

# The Alfalfa Leafcutting Bee, *Megachile rotundata*: The World's Most Intensively Managed Solitary Bee\*

Theresa L. Pitts-Singer and James H. Cane

USDA ARS Bee Biology & Systematics Laboratory, Utah State University, Logan, Utah 84322; email: Theresa.Pitts-Singer@ars.usda.gov, Jim.Cane@ars.usda.gov

Annu. Rev. Entomol. 2011. 56:221–37

First published online as a Review in Advance on August 30, 2010

The *Annual Review of Entomology* is online at [ento.annualreviews.org](http://ento.annualreviews.org)

This article's doi:  
10.1146/annurev-ento-120709-144836

Copyright © 2011 by Annual Reviews.  
All rights reserved

0066-4170/11/0107-0221\$20.00

\*This is a work of the U.S. Government and is not subject to copyright protection in the United States.

## Key Words

Apoidea, bivoltinism, chalkbrood, Fabaceae, pollination, Megachilidae

## Abstract

The alfalfa leafcutting bee (ALCB), *Megachile rotundata* F. (Megachilidae), was accidentally introduced into the United States by the 1940s. Nest management of this Eurasian nonsocial pollinator transformed the alfalfa seed industry in North America, tripling seed production. The most common ALCB management practice is the loose cell system, in which cocooned bees are removed from nesting cavities for cleaning and storage. Traits of ALCBs that favored their commercialization include gregarious nesting; use of leaves for lining nests; ready acceptance of affordable, mass-produced nesting materials; alfalfa pollination efficacy; and emergence synchrony with alfalfa bloom. The ALCB became a commercial success because much of its natural history was understood, targeted research was pursued, and producer ingenuity was encouraged. The ALCB presents a model system for commercializing other solitary bees and for advancing new testable hypotheses in diverse biological disciplines.

**Alfalfa a.k.a. lucerne:**

*Medicago sativa* L.  
(Fabaceae).  
Originating in southwest Asia, the tetraploid is now widely naturalized in Europe and North America

**Alfalfa leafcutting bee (ALCB):**

*Megachile*  
*(Eutricharaea)*  
*rotundata* Fab.  
(Apoidea:  
Megachilidae)

**INTRODUCTION**

Fifty-three years ago in the second volume of the *Annual Review of Entomology*, George “Ned” Bohart summarized the wild bees that pollinate the world’s leading forage legume, alfalfa (*Medicago sativa*). He concluded that only honey bees, although mediocre pollinators, could satisfy this crop’s need for an abundant bee (7). At that time, U.S. alfalfa seed production was shifting west to California, where fields packed with hived honey bees increased seed yields fivefold to 450 kg ha<sup>-1</sup>. Fifteen years later, Bohart again wrote for these pages about an unheralded new agricultural pollinator from Eurasia, the alfalfa leafcutting bee (ALCB) (8). Although detected in the United States without fanfare by the 1940s (122), the ALCB revolutionized the alfalfa seed industry, boosting yields to a remarkable 1300 kg ha<sup>-1</sup> (95). No other solitary bee is produced and managed so intensively, although several species are propagated to fill regional niche markets. What are this bee’s attributes that have made its management uniquely successful, but only where it and alfalfa are not native? Is ALCB management a model for other solitary bee pollinators or, like honey beekeeping, is it peculiar and unlikely to be replicated? In reviewing the ALCB’s life history, management, and ecological impacts, we highlight the factors that enable successful solitary bee management for crop pollination.

**TAXONOMY AND BIOGEOGRAPHY**

The bee genus *Megachile* is massive (ca. 1,478 described species, or one-third of all megachilids) and cosmopolitan (76); most cut leaf pieces to line their nests, a behavior to which they owe their common name (76, 144). This genus is subdivided into 52 subgenera (76), in part reflecting their collective diverse behaviors and morphologies. The largest subgenus, *Eutricharaea*, has more than 230 described species, all from the Eastern Hemisphere (37, 76). It includes the ALCB,

which has been introduced widely (e.g., the Americas, Australia) to pollinate alfalfa.

Fossil *Megachile* are unknown. However, trace fossils have been reported periodically for more than a century. These are leaf imprints that bear the characteristic semicircular marginal notches that remain after females clip leaf discs for nest building (see Life Cycle, below). Most examples come from the Eocene (34–55 Ma) (66); the oldest one dates to the Paleocene (55–65 Ma) (144).

**POLLINATION**

Owing to its use in alfalfa, the value of the ALCB is surpassed only by the honey bee for pollination of field crops. For example, ALCB use yielded 46,000 metric tons of alfalfa seed in North America in 2004, two-thirds of world production (80, 95). Planted worldwide for hay, alfalfa is fed to livestock, especially dairy cows. Alfalfa seed and resultant hay constitute one-third of the \$14 billion value ascribed to honey bees pollinating U.S. crops (139); managed ALCBs account for an additional 50% of alfalfa seed production in the northwestern United States (139) and central Canada. Paradoxically, the ALCB remains uncommon and irrelevant for pollinating alfalfa in its native range. It constituted just 0.03% of the 8,168 wild bees taken in 27 Hungarian alfalfa fields (78). In wild regions of Spain where alfalfa is also grown, the ALCB was absent from the 59 species sampled (82), and even in southern France, large populations are difficult to sustain for alfalfa (126).

Bees pollinate alfalfa flowers when they trip the staminal column, for which ALCBs are effective (29) (**Figure 1**). Rates of pod and seed set reflect primarily the frequencies of tripping (15), with lesser benefits from cross-pollination (124). Females of the ALCB and the alkali bee (*Nomia melanderi*; Halictidae) excel at pollinating alfalfa, tripping 80% of visited flowers (15), comparable to alfalfa’s effective but unmanaged European pollinators (23). In North America, diverse bee species—including native *Megachile* species—also pollinate alfalfa well (8, 45, 68),

but their abundances are never adequate to satisfy modern seed yield expectations.

Rates and distances of alfalfa gene flow cannot be reliably extrapolated from the flight range of ALCBs. ALCBs readily fly more than 100 m from the nest but then forage locally in a field, moving pollen only short distances of about 4 m (117). Among isolated alfalfa plots or plants, however, minor pollen-mediated gene flow occurs over much greater distances, 8% between plots 200 m apart (10) and 0.5% between plots 330 m apart (139).

The name alfalfa leafcutting bee belies its moderate foraging and pollination versatility. ALCBs have varying success in pollinating several other North American field crops. They are reportedly in extensive use for producing hybrid seed of canola (*Brassica napus*) in western Canada. In cage studies, they pollinated several annual clovers well (*Trifolium* spp.) (72, 104), but not vetches (*Vicia* spp.) (107). They pollinate some native legumes farmed for wildland restoration seed (16). ALCBs pollinate the small flowers of lowbush blueberries (*Vaccinium angustifolium*) grown commercially in the Canadian Maritimes (54) but are exposed to lethally cold nights there during bloom (113). ALCBs foraged at and pollinated cranberry in field cages (19), but in open bogs they foraged little and dispersed (69).

The ALCB can also be amenable to pollinating in confinement. Along with mason bees (*Osmia lignaria*) and honey bees, ALCBs are useful in cages to increase or regenerate germplasm accessions stored at crop seed repositories (21). Caged ALCBs effectively pollinated carrot (130) and canola (116) for hybrid seed, but they eschewed bloom of field-grown carrot (129). Greenhouse pollination of vegetables or fruits by ALCBs receives scant attention; they forage readily in glasshouses (15, 70, 136) but orient poorly under some plastic films (133).

## LIFE CYCLE

Adult ALCBs naturally emerge and nest during the hot days of summer. Females mate once, soon after emergence, and then consume

nectar and pollen as their first eggs mature (106); within a week females begin constructing and provisioning cells sequentially. Like most other *Megachile* species, they nest in existing holes above ground, fashioning nest walls, partitions, and plugs from strips and disks of leaves that are transported singly. Each nest cell requires 14–15 leaf pieces, both for wall strips and partition disks (58, 70) (Figure 1). They cut these pieces using the opposing beveled edges of their mandibles like scissors (Figure 1). At the nest, the female thoroughly chews the edges of each new leaf piece. The resultant sticky pulp binds the new leaf piece to the others (136). An ALCB can line and later cap one cell with leaf pieces on average in 81 min (70) to 2.5 h (58). Leaf piece dimensions and cell architecture represent precisely measurable physical manifestations of the bee's complex behaviors (44).

A female spends from 5 to 6 h per day foraging (58, 70), returning from flowers with both dry pollen in her scopa and nectar in her crop. The female enters the cavity headfirst to regurgitate her crop full of nectar. She then backs out, turns, and backs into the nest, using her metatarsal hair combs to sweep her abdominal scopa clean of its pollen load (32). Early in the provisioning sequence, the female carries mostly pollen (ca. 80%), but with each subsequent trip, she returns with proportionally more nectar (59, 136). On her final foraging trip, a female invariably returns with just nectar (58, 59, 70). Regurgitated atop the provision mass, this nectar constitutes the young larva's largely liquid diet (136). The final provision mass has a wet pasty consistency, weighs about 90–94 mg, and consists of 33%–36% pollen and 64%–67% nectar by weight (18, 59). It contains 1.3 million pollen grains and is 47% sugar by weight (18). Male progeny receive 17% less provision than do females (59). Larval provision masses of ALCBs contain diverse aerobic bacteria, filamentous fungi, and yeasts, but neither their removal by irradiation (50) nor individual restoration to sterile provisions (49) affected larval performance.

Under ideal greenhouse conditions with abundant sweetclover (*Melilotus officinalis*),

---


**Scopa:** a brush of setae (hairs) beneath the female bee's abdomen for transporting dry pollen

---

bloom for forage, ALCB females sometimes laid two eggs per day, and each female completed on average 57 cells with eggs over their 7–8 week life spans (71). Far fewer cells result when floral resources become limited (92, 96). Interference between females, or excessive male harassment of females, slows the pace of many

nesting activities and curtails reproductive output, at least in cages (70, 112).

The immature ALCB develops rapidly (**Figure 1**). Embryogenesis takes 2–3 days, followed by five larval instars (136). The first instar is spent inside the egg chorion from which the second instar larva hatches to begin feeding



(136). As with *A. mellifera*, young ALCB larvae consume a liquid diet, but they imbibe it from a shallow trough that the larva carves atop its provision mass (136). After 3–4 more days of development, the larva molts to its final fifth instar. In just the next three days, it eats the entire pollen provision. Few insects rival ALCB nitrogen and energy assimilation efficiencies (146); moreover, only 2% of the provision is left uneaten. Only when the larva is done feeding does it defecate (76). As with other megachilids and *Apis* species (among other long-tongued bees) (76), the larva then weaves a tough multilayered cocoon of secreted silk of unknown composition.

Typical of other *Megachile* and many summer-flying bees (76), the ALCB brood overwinters as a postfeeding, diapausing larva, the prepupa. Diapause terminates with warming conditions of late spring or early summer, or when they are artificially incubated (see Commercial Management, below). Alternatively, instead of diapause, up to half of the early ALCB summer brood completes development to yield a second generation before late summer (48, 62, 126). This facultative bivoltinism is irreversible (131). The cues and mechanism(s) that induce diapause or bivoltinism are unknown, but suspected factors that elicit summer emergence

include maternal and/or larval responses to long photoperiods, excessive heat, and poor larval nutrition, as well as maternal inheritance (57, 84, 86, 101, 127, 132).

## MATING

The courtship behaviors and mating biology of the ALCB resemble those of many solitary bees (24). Male ALCBs are usually smaller than females (83), having received smaller provisions as larvae (32). Females vary more in size than males (83), perhaps because daughters are more likely to receive less food when forage is sparse (92). Like most bees (111), the ALCB is protandrous. The first adult males emerge 1–3 days before females regardless of thermal regime (56, 109). Protandry facilitates an orderly exodus from a linear cavity nest whose male cells are nearer the entrance (**Figure 1**), as well as early access to receptive females. Sib-mating within intact ALCB nests is not reported. Confined to desktop cages, some female ALCBs mated and their spermathecae received sperm within hours of emergence (61). However, when newly emerged marked females were released in the field, only one-third of recaptured females were inseminated within 48 h;

---

### Facultative

**bivoltinism:** refers to some progeny emerging as a second generation in the same reproductive season

---

---

### Figure 1

Tripping alfalfa flower, alfalfa leafcutting bee (ALCB) life cycle, and ALCB field domicile. (a) Banner petal and keel of untripped alfalfa flower. (b) Staminal column snaps upward to banner petal when tripped by a bee. (c) The ALCB female lands on a flower and applies pressure to keel; the flower is tripped; pollen is exposed and released. (d) Extended proboscis of ALCB female while taking nectar. (e) ALCB female always probes flower for nectar. (f) ALCB egg laid atop provision mass of pollen and nectar that is enveloped in leaf pieces. (g) First instar usually remains inside the chorion; second instar then hatches to imbibe nectar pooled atop the provision. (h) Third and fourth instars continue feeding but do not defecate. (i) Fifth instar consumes final portions of the provision and defecates before spinning a cocoon. (j) In the cocoon stage, the postfeeding late fifth instar is called a prepupa. (k) ALCB cells are arranged linearly within a nest, with female cells made before male cells (left to right = back to front). (l) An X-radiograph of the same nest. (m) Reveals healthy diapausing prepupae, with larger females in the first (left) cells. (n) Scanning electron micrograph of female ALCB mandibles. Beveled, chisel-like opposing mandibular edges are used to cut leaf ovals and discs for lining nest cavities and capping individual nest cells, respectively. (Micrograph courtesy of N. Dakota State Univ., Fargo, North Dakota.) (o) ALCB domicile aside a blooming alfalfa field. (p) Crowded polystyrene nesting boards with thousands of ALCBs (black dots) seeking nest cavities. (q) Close-up view of ALCB nesting board surface showing a female completing a nest plug made of leaf discs; the face of another female peers out from a cavity from which second-generation bees chewed through the leaf plug and have already emerged. (Except where noted, all images are property of USDA ARS Bee Biology & Systematic Laboratory.)

### Loose cell bee

#### management system:

nest cells are removed from cavities of bees' nesting boards and then are separated, cleaned, and sorted using specialized equipment, rather than leaving cells within nests in boards

all were inseminated within a week (106). Mating is commonly seen at commercial nesting shelters (32), but its frequency at flowers is not known, particularly for bees emerging from intact natal nests at natural nesting densities. Territoriality is not evident, but males do pounce on perched females and other males.

ALCB males have simple genitalia (85), and copulation for ALCBs is apparently brief (8–10 s) (112). Copulation is preceded by male wing flipping and accompanied by pulsed buzzing and lateral sweeps of his antennae (32). Males firmly grip females in a stereotypical embrace that employs distinctive forecoxal spines and basal mandibular protrusions (147) but without the paddle-like foretarsi of some other *Megachile* species (147). Putative female sex pheromones of ALCBs are reported among their cuticular alkenes (88). Glandular regions of the male's foretarsi are pressed against the female's antennae during mating (147), but the possible exudates have not been isolated or studied for their functionality.

Males undoubtedly inseminate multiple females when they can (polygyny), as seems ubiquitous with non-*Apis* bees (24). As for many solitary bees (89), female ALCBs seem to be monogamous (31) and, in confinement at least, physically reject later suitors (61). Allozyme markers (74) and early genetic studies using RAPD markers indicated singular paternity among nestmates (6). Oogenesis can begin without mating in ALCBs, but as is the case for many other insects (145), eggs will not mature unless the female eats pollen (106). Overall, the mating biology of ALCBs seems unexceptional for *Megachile* (147) and solitary bees in general.

## COMMERCIAL MANAGEMENT

Several works over the decades have described ALCB management for alfalfa seed production (8, 28, 46, 47, 105, 119, 120). Unless otherwise cited, most of the following information is from comprehensive guides by Richards (105) and Frank (28) written for alfalfa pollination in Canada. U.S. bee managers tailor Canadian guidelines for their climate to maximize alfalfa

seed yield. Depending on the age, quality, and production potential of the alfalfa, growers seek to optimize the ALCB stocking rate for pollination. Because of their higher seed yield potential, U.S. producers use more bees (100,000–150,000 bees per hectare) than do Canadian producers (50,000–75,000 bees per hectare) (96). A loose cell bee management system is most commonly practiced in North America, consisting of four sequential phases: spring/summer incubation, summer brood production, fall/winter cleanup, and winter storage.

ALCB prepupae spend the winter in cold storage (at 4°C–5°C for 7–10 months), usually as cells removed from their nests (i.e., loose cells). In the spring, nest cells bearing prepupae are poured into large, shallow trays to incubate at constant 30°C and 50%–60% relative humidity for safe, synchronous, and timely adult emergence. Emergence is timed for when alfalfa bloom is expected to be at 25%–50% (3, 28). Prior to bee emergence, wasp parasitoids emerge and must be controlled (see Parasitoids and Predators, below). Male ALCBs emerge by day 17–20, and females emerge a few days later (97). Once ~75% of females have emerged, bees are taken to domiciles in blooming alfalfa fields for release from incubation trays. However, if weather is unfavorable, alfalfa bloom is delayed, or an insecticide application is needed for alfalfa pests, deliberate cooling (15°C–20°C) during incubation will slow adult emergence.

Prior to bee release, field domiciles containing nesting boards are readied. Occurring in many shapes, sizes, and materials, domiciles can be uniformly dispersed throughout the fields or placed around field borders. The domicile opening is oriented southeastward (121) so that bees and brood are warmed by the morning sun but shaded during the heat of the day. Nest materials are wood or polystyrene boards containing evenly spaced holes (hole length is 95–150 mm; diameter is 5–7 mm) (32) (**Figure 1**). Bees will only fly once the sun has risen and ambient temperature has warmed to 21°C; flight ceases at twilight, regardless of temperature (65, 125).

Most bees immediately fly out of the opened trays into the blooming field. However, bees that have not eclosed that continue to develop under field conditions emerge slowly and may even die. Nesting begins a few days after release and continues for about 11 weeks, although later activity may represent only the second-generation bees. After bee activity subsides, brood-filled nesting boards are moved from the field to a shop or incubator for storage.

Conscientious bee management using controlled conditions during the fall is practiced more in Canada than in the United States. Nests are allowed to dry in boards, with some of the larvae maturing to the prepupal stage if kept warm enough ( $>15^{\circ}\text{C}$ ); parasitoids may need to be controlled. Bees are gradually cooled for winter storage (about  $5^{\circ}\text{C}$ ), arresting development. During fall and winter, bee nests are removed from boards by stripping them from grooved wooden boards or “punching” them from polystyrene molded boards. This process is followed by cell tumbling, breaking, and separation, which are mechanized techniques for minimizing extraneous matter and helping to rid the bee stock of chalkbrood (see Chalkbrood Disease, below). During winter storage, stocks are sampled for the percent of live progeny and their sex ratio. Healthy, diseased, parasitized, and other dead bees can be diagnosed from X-radiographs of cells or manual cell dissections (123).

ALCB management is strikingly different from honey bee management. Migratory honey beekeepers move hives to follow bloom, but moving ALCB domiciles or boards disorients most nesting females and increases dispersal. Because ALCB adults are active for only about two months, most ALCB care focuses on prepupae in nest cells. Farmers can manage ALCBs amid regular duties or can contract with specialists to handle all ALCB management, in the same way farmers can rent honey bee hives.

To control *A. mellifera* hive pests, pesticides are placed in direct contact with adult bees. For ALCBs, most pests are controlled while larvae are protected in their cells and no adults are

present (see Parasitoids and Predators, below). Nevertheless, ALCB adults and brood may be exposed to pesticides and fungicides applied to the crop plant during the nesting season. Bees may be killed or sublethally affected if toxins are sprayed during daylight or if active residues persist (1, 110). Poisoned adult bees are evidenced by their corpses or irregular behavior, but it is difficult to ascertain if brood succumb to toxins that have reached them directly in their nests, as provision contaminants, or through transovarial transport to eggs from mother bees after their topical or oral exposure to active chemicals (79). Generally, pesticides toxic to honey bees are toxic to ALCBs, bumble bees, and wild bees, although  $\text{LD}_{50}$  levels can differ (64, 110, 134). Toxicological screening of new pesticide formulations and classes are vital for safekeeping all bees whose different life histories and foraging behaviors may render some more susceptible than others (e.g., ALCBs handling contaminated leaves) (1, 79, 90, 118, 134).

## CHALKBROOD DISEASE

ALCBs are attacked by disease pathogens just like other bees (5, 52). The most common pathogen that infects ALCB brood is the fungus *Ascospaera aggregata* (Ascomycete), which causes chalkbrood (34, 35). The related *A. apis* causes chalkbrood in honey bees (20), but the two *Ascospaera* species do not cross-infect. In the 1970s, chalkbrood disease was devastating to ALCB production; controls used today have lessened disease impact ( $\leq 20\%$  of U.S. brood) (36, 98). In Canada, chalkbrood is stringently controlled ( $\leq 2\%$ ) (28), and Canadian bees are sold to the United States, but not vice versa.

Larval cadavers filled with *A. aggregata* spores are the source of future disease. When emerging bees are trapped in nests behind chalkbrood cadavers, they must chew through the cadavers and thus are dusted with fungal spores. Spore-laden adults may then contaminate provisions of their own brood and also may contaminate other bees through physical contact or by depositing spores in tunnels when investigating future nest cavities

(102, 141). Although loose cell management keeps adults from chewing through cadavers (105), spores erupting from cadavers during cell removal and processing may contaminate loose cells (53). In addition, where ALCB bivoltinism is prevalent in the United States, the spread of chalkbrood disease is exacerbated by summer-emerging bees that must chew through any spore-filled siblings (140).

Canadian managers use paraformaldehyde fumigation to kill fungal spores on loose cells, trays, nesting boards, and any bee equipment (35). Paraformaldehyde is not registered for this use in the United States, so U.S. managers have tried to cleanse nesting boards with methyl bromide fumigation, heat (wood boards only), and chlorine dips (51). New chalkbrood controls are desired because paraformaldehyde is carcinogenic, methyl bromide is now banned, and heat and chlorine treatments are difficult and labor-intensive.

## PARASITIDS AND PREDATORS

ALCBs host many natural enemies native to both North America and Eurasia (8, 25). Fortunately, no pest that accompanied ALCBs from Europe has affected North American native bees. Most knowledge of ALCB parasitism and predation comes from managed populations. Where the ALCB has been used around the world, the same groups of parasitoids and predators have followed or adapted (63, 108, 126, 148).

Up to 20% of ALCB cells produced in U.S. western states can be parasitized by wasps (98), which are known natural enemies of other *Megachile* and *Osmia* spp. (25, 60) in their native ranges. The prevalent parasitoids are the European *Pteromalus venustus* (Pteromalidae) (25) and the native minute *Tetrastichus megachilidis* (Eulophidae). Also, the European *Monodontomerus aeneus* (Torymidae) (40, 60) once, but no longer, devastated ALCB populations. Adult parasitoids are typically killed with dichlorvos pest strips (73) that are placed in storage or incubator chambers, but that do not harm ALCBs enclosed in nest cells. Ultraviolet lamps above liquid traps also can be effective.

During nesting, parasitoids are physically deterred from nests by the thick walls of artificial cavities and by felt cloth tightly affixed to the back of nesting boards.

*Sapyga pumila* (Sapygidae) is a North American wasp that attacks native bees in several genera and quickly adopted the ALCB as a host (25, 60). A trap designed for this wasp (2, 25, 135) is seldom used because *S. pumila* parasitism is no longer a major concern. A trap also exists, but is rarely used, for the checkered flower beetle, *Trichodes ornatus* (Cleridae) (22, 67). Larvae of these beetles are pests during the nesting season and while bees are stored (25). They attack other megachilids, bees in other families, and some wasps (67).

Less persistent or problematic parasitoids, predators, and pests include the wasp parasitoids *Melittobia chalybii* and *M. acasta* (Eulophidae) (73), six native *Coelioxys* species (60), one *Stelis* species (cuckoo bees) (Megachilidae), the parasitic beetle *Nemognatha lutea* (Meloidae), and the beetles *Trogoderma* spp. (Dermestidae) and *Tribolium* spp. (Tenebrionidae), which eat immature bees and their provisions (25). Predators also include yellowjacket wasps (Vespidae: *Vespa* spp.), earwigs (Forficulidae: *Forficula auricularia*), ants (Formicidae), birds, and rodents (25).

As with honey bees, nest and brood destroyers can be problematic year-round. Unlike honey bees, ALCB adults are present only in summer with their brood sealed inside leafy capsules. Thus, pests can be treated with insecticides while ALCBs are protected within their cells.

## UNEXPLAINED MORTALITY

ALCB populations are difficult to sustain in the United States, so U.S. producers import ALCBs from Canada to supplement or completely supply their bees (95). Although all bee stocks suffer pestilence (chalkbrood, parasites, and predators), unexplained mortality occurs more often in U.S. than Canadian populations.

Poor management may account for some U.S. bee mortality. Compared to Canadian



samples, more U.S. ALCBs died over the winter and during incubation, with most bees dying as prepupae (97). Although optimal durations and temperatures for winter storage and incubation are reliable (56, 100, 109, 137, 138), less is known about potential bee losses due to conditions during the handling of completed nests prior to winter storage (57, 99). The duration of prewinter storage at the recommended temperature of 16°C can subtly affect bee survival to adulthood (99), but effects of excessive heat for various storage durations need study. Storage differences may underlie the superior survival of Canada-raised bees.

Unexplained brood mortality in the United States includes pollen ball (at times 40%–60% of brood), which are cells containing the pollen and nectar provision at a time when the provision should have been consumed by a larva. Many pollen ball cells lack eggs or larvae; others contain collapsed eggs, dead larvae, or fungi (93). Causes of pollen balls probably include unsuitable microclimates and overly dense bee populations (96, 98). Overstocked bees used in U.S. alfalfa fields quickly drain floral resources, suffer crowded nesting sites, and may increase the spread of disease (41, 52). Depleted resources constrain ALCB reproduction (91, 92, 96), and adults likely perish or depart crowded sites. Often under such conditions during the hot U.S. summers, less than half of ALCBs released into fields are replaced through reproduction. Other suspected causes of larval mortality include molds, viruses, other as yet identified pathogens, and pesticides (34, 35).

Second-generation emergence in the United States can reach 90% for the first nests made in the season, tapering to none for the season's last nests (55); over an entire season, summer emergence can total about 40% progeny loss. In contrast, summer adult emergence in Alberta, Canada, is quite low (e.g., <5%) (62). Reproduction by second-generation bees is constrained by sparse floral resources on farms and their progeny's race to become prepupae before cool weather arrests their development.

## VISUAL AND CHEMICAL CUES FOR ALFALFA LEAFCUTTING BEES

Just as honey bees orient to a hive, ALCBs must orient to their nests after foraging bouts. Large field domiciles can serve as long-distance visual cues for ALCBs flying over seed fields (32, 105, 121). Once at the domicile, a female bee finds her nest among thousands of holes using three-dimensionality, color contrast, and color patterns (27, 41, 42). If visual cues are manipulated, females become temporarily disoriented. Poor local orientation can result in wasted time, an increase in bee collisions, dropped nest-building materials, or further spread of disease that diminishes efficient, productive nesting.

Olfactory cues detected in close proximity or upon antennation seem to be important when initiating or recognizing nests. Females prefer once-used nesting boards over new ones, implicating olfactory attractants at nest initiation (14, 26, 87, 142). Isolating effective cues from the old nests has proven difficult. In laboratory Y-tube assays, ALCB females were attracted to a complete nest cell (after the adult emerged), larval feces, and solvent-extracted components from leaf pieces that once lined a cell (94). However, field bioassays were less conclusive, finding that only cavities cued with feces and a cocoon were somewhat more attractive than other components (142). Fractionation of chemicals from old nest components and testing them in field assays are needed for understanding chemical mediation of nest initiation and aggregation.

A nesting ALCB female distinguishes her own nest from others. ALCB females mark their nests by dragging the abdomen throughout the cavity (32, 43). Replacing an outer beemarked section of a nest cavity with a clean section causes a returning female to be confused and hesitant to enter, indicating an individual nest recognition pheromone. Cuticular waxes and/or the Dufour's gland may be the origin of the nest recognition cue (13). For honey bees, in which many individuals need to recognize the

---

**Pollen ball:** intact pollen and nectar provision mass remaining at end of the field season

---

colony odor to discern kin from nonkin, recognition requires them to learn odors both from cuticles of nestmates and from the colony's wax comb (11). Solitary bees need to recognize only their own nests.

## ALFALFA LEAFCUTTING BEES AS ECOLOGICAL DISRUPTORS

Exotic species can become ecologically disruptive invaders. Feral honey bees have penetrated far into wildlands (77), as have introduced European *Bombus terrestris* that are now escaping around the world (39). Among the few surveys of U.S. wildland bee communities where the ALCB might be expected, it has been rare or absent. In Wyoming short-grass prairie, none of the 13 species of *Megachile* (52 species of Megachilidae) were ALCBs (128). ALCB was similarly absent from the 125 bee species netted in tallgrass prairie remnants and restorations (103). In the shrub-steppe of eastern Oregon, 12 species of Megachilidae were trap-nested, but no ALCB was found (30). At a reserve amid intensive agriculture in central California, the ALCB was more common, but occupied only 3%–4% of available nesting cavities; it was outnumbered both by native cavity-nesters and by another escaped Eurasian species of *Eutricharaea*, *M. apicalis*, known to aggressively compete for nesting sites (4, 122). The ALCB is also rare or absent in southern Europe, even in alfalfa seed fields. The ALCB seems not to venture far from agricultural or otherwise disturbed sites in North America and so appears to be an inconsequential competitor with native bees for nesting sites or floral resources.

Some nonnative bees prefer and pollinate nonnative forbs, contributing to the invasions of weedy species that can outcompete native wildflowers for pollination services (38). Feral honey bees and feral bumble bees have been widely implicated in pollinating invasive Eurasian weeds in the Americas, New Zealand, Tasmania, and Australia (39). ALCBs also avidly visit several Eurasian weeds in North America, notably sweetclovers (*Melilotus alba* and *M. officinalis*) (58, 81) and purple loosestrife

(*Lythrum salicaria*), preferring both to alfalfa in choice tests (114). However, these and other exotic and invasive weeds are eagerly sought and pollinated by honey bees and bumble bees too (12).

## A MODEL SOLITARY BEE

Management of ALCBs is an exemplary system for solitary bee commercial pollination. Fortuitously, ALCBs are efficient crop pollinators that also thrive as aggregating cavity-nesters in anthropogenic habitats. In contrast, the solitary ground-nesting alkali bee is restricted in its use as a managed alfalfa pollinator because it requires specific soil moisture, temperature, and alkalinity. Few growing areas naturally or artificially can satisfy such requirements (95).

Where farming is intense and an abundance of pollinating bees is needed, managed cavity-nesting bees are advantageous. Cavity-nesters are transportable as packaged nests or loose cells; some species tolerate nesting in large aggregations. With progeny in nests or as loose cells, dead or parasitized cells can be culled, cells and management equipment can be treated for diseases, and the bee number and sex ratio can be determined. Other megachilids (e.g., *Osmia* spp. for orchard and berry crops) can be used as pollinators (9), but today they are not available at the scale of ALCBs (9, 17).

Just as honey bees and bumble bees represent the social bees for studies in evolution, genetics, behavior, learning, neurophysiology, and ecology (e.g., 33, 75), the readily available ALCB can serve as a model for solitary bees. ALCB laboratory maintenance is relatively simple, and adults can be kept to emerge through most of the year. Beyond management research, ALCB studies already include the effects of male harassment on reproductive success and the lack of response to proboscis extension reflex elicitation used in learning and conditioning experiments (112, 143). Recent reports have identified ALCB genes related to diapause regulation and bee immunity (149, 150). Future ALCB studies might reveal solitary bee commonalities in recognition, aggregation

behavior, mating strategies and courtship behaviors, and pheromone use. On the horizon are genome sequences of ALCB and other bees from which ancestral and shared traits in solitary, subsocial, and primitively eusocial bees may be found and thus contribute to the understanding of the mechanisms and evolutionary origins of bee sociality (115).

### SUMMARY POINTS

1. Accidentally introduced into the United States, the ALCB has become the world's most effectively used and intensely managed solitary bee. Its use transformed the alfalfa seed industry in North America. The most common management practice is the loose cell system.
2. Traits of ALCBs that contribute to their commercialization include their gregarious nature; philopatry; use of leaves for lining nests; ready acceptance of cheap, mass-produced nesting materials; and pollination efficacy at and emergence synchrony with alfalfa bloom.
3. The ALCB became a commercial success because its natural history was studied, targeted research was performed, and producer ingenuity was encouraged. ALCB management is a model system for commercializing other solitary bees and for advancing new testable hypotheses in diverse biological disciplines.

### DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

### ACKNOWLEDGMENTS

We thank Gordon Frank for insights into ALCB pollination of canola. Vincent Tepedino and James Pitts kindly provided helpful reviews.

### LITERATURE CITED

1. Abbott VA, Nadeau JL, Higo HA, Winston ML. 2008. Lethal and sublethal effects of imidacloprid on *Osmia lignaria* and clothianidin on *Megachile rotundata* (Hymenoptera: Megachilidae). *J. Econ. Entomol.* 101:784–96
2. Arnett WH. 1980. Alfalfa leafcutting bee parasite *Sapyga pumula* trap: effects of color and domicile placement on catch. *Environ. Entomol.* 9:783–84
3. Baird CR, Bitner RM. 1991. *Loose cell management of alfalfa leafcutting bees in Idaho*. Moscow: Univ. Idaho Coll. Agric., Curr. Inf. Ser. 588
4. Barthell JF, Frankie GW, Thorp RW. 1998. Invader effects in a community of cavity nesting megachilid bees (Hymenoptera: Megachilidae). *Environ. Entomol.* 27:240–47
5. Batra LR, Batra SWT, Bohart GE. 1973. The mycoflora of domesticated and wild bees (Apoidea). *Mycopathol. Mycol. Appl.* 49:13–44
6. Blanchetot A. 1992. DNA fingerprinting analysis in the solitary bee *Megachile rotundata*: variability and nest mate genetic relationships. *Genome* 35:681–88
7. Bohart GE. 1957. Pollination of alfalfa and red clover. *Annu. Rev. Entomol.* 2:355–80
8. Bohart GE. 1972. Management of wild bees for the pollination of crops. *Annu. Rev. Entomol.* 17:287–312
9. Bosch J, Sgolastra F, Kemp WP. 2008. Life cycle ecophysiology of *Osmia* mason bees used as crop pollinators. In *Bee Pollination in Agricultural Ecosystems*, ed. RR James, TL Pitts-Singer, 6:83–104. New York: Oxford Univ. Press. 232 pp.

10. Bradner NR, Frakes RV, Stephen WP. 1965. Effects of bee species and isolation distance on possible varietal contamination in alfalfa. *Agron. J.* 57:247–48
11. Breed MD. 1998. Chemical cues in kin recognition: criteria for identification, experimental approaches, and the honey bee as an example. In *Pheromone Communication in Social Insects*, ed. RK Vander Meer, MD Breed, ML Winston, KE Espelie, 3:57–78. Boulder, CO: Westview. 368 pp.
12. Brown BJ, Mitchell RJ, Graham SA. 2002. Competition for pollination between an invasive species (purple loosestrife) and a native congener. *Ecology* 83:2328–36
13. Buckner JS, Pitts-Singer TL, Guédot C, Hagen MM, Fatland CL, Kemp WP. 2009. Cuticular lipids of female solitary bees, *Osmia lignaria* Say and *Megachile rotundata* (F.) (Hymenoptera: Megachilidae). *Comp. Biochem. Physiol. B* 153:200–5
14. Buttery RG, Parker FD, Teranishi R, Mon TR, Ling LC. 1981. Volatile components of alfalfa leaf-cutter bee cells. *J. Agric. Food Chem.* 29:955–58
15. Cane JH. 2002. Pollinating bees (Hymenoptera: Apiformes) of U.S. alfalfa compared for rates of pod and seed set. *J. Econ. Entomol.* 95:22–27
16. Cane JH. 2006. An evaluation of pollination mechanisms for purple prairie-clover, *Dalea purpurea* (Fabaceae: Amorpheae). *Am. Midl. Nat.* 156:193–97
17. Cane JH. 2008. Pollinating bees crucial to farming wildflower seed for U.S. habitat restoration. In *Bee Pollination in Agricultural Ecosystems*, ed. RR James, TL Pitts-Singer, 4:48–64. New York: Oxford Univ. Press. 232 pp.
18. Cane JH, Gardner D, Harrison P. 2011. Nectar and pollen sugars constituting larval provisions of the alfalfa leaf-cutting bee (*Megachile rotundata*) (Hymenoptera: Apiformes: Megachilidae). *Apidologie*. In press
19. Cane JH, Schifffhauer D. 2003. Dose-response relationships between pollination and fruiting refine pollinator comparisons for cranberry (*Vaccinium macrocarpon* [Ericaceae]). *Am. J. Bot.* 90:1425–32
20. Christensen M, Gilliam M. 1983. Notes on the *Ascospaera* species inciting chalkbrood in honey bees. *Apidologie* 14:291–97
21. Cox RL, Abel C, Gustafson E. 1996. A novel use of bees: controlled pollination of germplasm collections. *Am. Bee J.* 136:709–12
22. Davis HG, George DA, McDonnough LM, Tamaki G, Burditt AK. 1983. Checkered flower beetle (Coleoptera: Cleridae) attractant: development of an effective bait. *J. Econ. Entomol.* 76:674–75
23. Dylewska M, Jablonski B, Sowa S, Bilinski M, Wrona S. 1970. An attempt of determination of the number of bees (*Hym., Apoidea*) needed for adequate pollination of alfalfa. *Pol. Pismo Entomol.* 40:371–98
24. Eickwort GC, Ginsberg HS. 1980. Foraging and mating behavior in Apoidea. *Annu. Rev. Entomol.* 25:421–46
25. Eves JD, Mayer DF, Johansen CA. 1980. Parasite, predators, and nest destroyers of the alfalfa leafcutting bee, *Megachile rotundata*. *West. Reg. Ext. Publ.* 32:1–15
26. Faurey DT, Lieveer JAC. 1986. Cell production by the alfalfa leafcutting bee (*Megachile rotundata*) in new and used wood and polystyrene nesting materials. *J. Appl. Entomol. Z. Angew. Entomol.* 102:148–53
27. Fauria K, Campan R, Grimal A. 2004. Visual marks learned by the solitary bee *Megachile rotundata* for localizing its nest. *Anim. Behav.* 67:523–30
28. Frank G. 2003. *Alfalfa Seed and Leafcutter Bee Production & Marketing Manual*. Brooks, Alberta, Can.: Irrig. Alfalfa Seed Prod. Assoc. 160 pp.
29. Free JB. 1993. *Insect Pollination of Crops*. New York: Academic. 684 pp.
30. Frohlich DR, Clark WH, Parker FD, Griswold TL. 1988. The xylophilous bees and wasps of a high, cold desert: Leslie Gulch, Oregon (Hymenoptera: Apoidea, Vespoidea). *Pan-Pac. Entomol.* 64:266–69
31. Gerber HS, Klostermeyer EC. 1970. Sex control by bees: a voluntary act of egg fertilization during oviposition. *Science* 167:82–84
32. Gerber HS, Klostermeyer EC. 1972. Factors affecting the sex ratio and nesting behavior of the alfalfa leafcutter bee. *Wash. Agric. Exp. Stn. Tech. Bull.* 73:1–11
33. Giurfa M. 2007. Behavioral and neural analysis of associative learning in the honeybee: a taste from the magic well. *J. Comp. Physiol. A* 193:801–24
34. Goerzen DW. 1989. *Foliar Moulds in Alfalfa Leafcutting Bees*. Can.: Saskatchewan Agric. Dev. Fund Ext. Bull. (ISBN 0-88656-534-0) 4 pp.

---

31. First documentation of female control of progeny sex in a solitary bee.

---

35. Goerzen DW. 1991. Microflora associated with the alfalfa leafcutting bee, *Megachile rotundata* (Fab.) (Hymenoptera: Megachilidae) in Saskatchewan, Canada. *Apidologie* 22:553–61
36. Goerzen DW, Watts TC. 1991. Efficacy of the fumigant paraformaldehyde for control of microflora associated with the alfalfa leafcutting bee, *Megachile rotundata* (Fabricius) (Hymenoptera: Megachilidae). *Bee Sci.* 1:212–18
37. Gonzalez VH, Engel MS, Hinojosa-Diaz IA. 2010. A new species of *Megachile* from Pakistan, with taxonomic notes on the subgenus *Eutricharaea* (Hymenoptera: Megachilidae). *J. Kans. Entomol. Soc.* 83:58–67
38. Goodell K. 2008. Invasive exotic plant-bee interactions. In *Bee Pollination in Agricultural Ecosystems*, ed. RR James, TL Pitts-Singer, 10:166–83. New York: Oxford Univ. Press. 232 pp.
39. Goulson D. 2003. Effects of introduced bees on native ecosystems. *Annu. Rev. Ecol. Syst.* 34:1–26
40. Grissell EE. 2007. Torymidae (Hymenoptera: Chalcidoidea) associated with bees (Apoidea), with a list of chalcidoid bee parasitoids. *J. Hymenopt. Res.* 16:234–65
41. Guédot C, Bosch J, James RR, Kemp WP. 2006. Effects of three-dimensional and color patterns on nest location and progeny mortality in alfalfa leafcutting bee (Hymenoptera: Megachilidae). *J. Econ. Entomol.* 99:626–33
42. Guédot C, Bosch J, Kemp WP. 2005. The relative importance of vertical and horizontal visual cues in nest location by *Megachile rotundata*. *J. Apic. Res.* 44:109–15
43. Guédot C, Pitts-Singer TL, Buckner JS, Bosch J, Kemp WP. 2005. Use of olfactory cues for individual nest recognition in the solitary bee *Megachile rotundata* (F.) (Hymenoptera: Megachilidae). Proc. Winter Seed Sch. Conf. Northwest Alfalfa Seed Growers Assoc. Meet., 30 Jan. – 1 Feb. pp. 45
44. Hasenkamp KR. 1974. Ökophysiologische und ökethologische Untersuchungen an Blattschneiderbienen. *Forma Funct.* 7:139–78
45. Hobbs GA. 1956. Ecology of the leaf-cutter bee *Megachile peribirta* Ckll. (Hymenoptera: Megachilidae) in relation to production of alfalfa seed. *Can. Entomol.* 87:625–31
46. Hobbs GA. 1965. Importing and managing the alfalfa leaf-cutter bee. *Publ. No. 1209*. Ottawa, Ontario: Can. Dep. Agric.
47. Hobbs GA. 1967. Domestication of alfalfa leaf-cutter bees. *Publ. No. 1313*. Ottawa, Ontario: Can. Dep. Agric.
48. Hobbs GA, Richards KW. 1976. Selection for a univoltine strain of *Megachile* (*Eutricharaea*) *pacifica* (Hymenoptera: Megachilidae). *Can. Entomol.* 108:165–67
49. Inglis GD, Goettel MS, Sigler L. 1993. Influence of microorganisms on alfalfa leafcutter bee (*Megachile rotundata*) larval development and susceptibility to *Ascospaera aggregata*. *J. Invertebr. Pathol.* 61:236–43
50. Inglis GD, Goettel MS, Sigler L, Borsari J. 1992. Effects of decontamination of eggs and  $\gamma$ -irradiation of provisions on alfalfa leafcutter bee (*Megachile rotundata*) larvae. *J. Apic. Res.* 31:15–21
51. James RR. 2005. Impact of disinfecting nesting boards on chalkbrood control in the alfalfa leafcutting bee. *J. Econ. Entomol.* 98:1094–100
52. James RR. 2008. The problem of disease when domesticating bees. In *Bee Pollination in Agricultural Ecosystems*, ed. RR James, TL Pitts-Singer, 8:124–41. New York: Oxford Univ. Press. 232 pp.
53. James RR, Pitts-Singer TL. 2005. *Ascospaera aggregata* contamination on alfalfa leafcutting bees in a loose cell incubation system. *J. Invertebr. Pathol.* 89:176–78
54. Javorek SK, MacKenzie KE, Vander Kloet SP. 2002. Comparative pollination effectiveness among bees (Hymenoptera: Apoidea) on lowbush blueberry (Ericaceae: *Vaccinium angustifolium*). *Ann. Entomol. Soc. Am.* 95:345–51
55. Johansen CA, Eves J. 1973. Effects of chilling, humidity and seasonal conditions on emergence of the alfalfa leafcutting bee. *Environ. Entomol.* 2:23–26
56. Kemp WP, Bosch J. 2000. Development and emergence of the alfalfa pollinator *Megachile rotundata* (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.* 93:904–11
57. Kemp WP, Bosch J. 2001. Postcocooning temperatures and diapause in the alfalfa pollinator *Megachile rotundata* (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.* 94:244–50
58. Klostermeyer EC, Gerber HS. 1969. Nesting behavior of *Megachile rotundata* (Hymenoptera: Megachilidae) monitored with an event recorder. *Ann. Entomol. Soc. Am.* 62:1321–26

59. Klostermeyer EC, Mech SJ Jr, Rasmussen WB. 1973. Sex and weight of *Megachile rotundata* (Hymenoptera: Megachilidae) progeny associated with provision weights. *J. Kans. Entomol. Soc.* 46:536–48
60. Krombein KV, Hurd PD, Smith DR, Burks BD. 1979. *Catalog of Hymenoptera in America North of Mexico*. Washington, DC: Smithsonian Inst. Press. 2731 pp.
61. Kronic MD. 1971. Mating of alfalfa leafcutter bee, *Megachile rotundata* (F.), in the laboratory. *Can. J. Zool.* 49:778–79
62. Kronic MD. 1972. Voltinism in *Megachile rotundata* (Megachilidae: Hymenoptera) in Southern Alberta. *Can. Entomol.* 104:185–88
63. Kronic MD, Tasei JN, Pinzauti M. 1995. Biology and management of *Megachile rotundata* Fabricius under European conditions. *Apicoltura* 10:71–97
64. Ladurner E, Bosch J, Kemp WP, Maini S. 2005. Assessing delayed and acute toxicity of five formulated fungicides to *Osmia lignaria* Say and *Apis mellifera*. *Apidologie* 36:449–60
65. Lerer H, Bailey WG, Mills PF, Pankiw P. 1982. Pollination activity of *Megachile rotundata* (Hymenoptera: Apoidea). *Environ. Entomol.* 11:997–1000
66. Lewis SE. 1994. Evidence of leaf-cutting bee damage from the Republic sites (Middle Eocene) of Washington. *J. Paleontol.* 68:172–73
67. Linsley EG, MacSwain JW. 1943. Observations on the life history of *Trichodes ornatus* (Coleoptera, Cleridae), a larval predator in the nests of bees and wasps. *Ann. Entomol. Soc. Am.* 36:589–601
68. Linsley EG, MacSwain JW. 1947. Factors influencing the effectiveness of insect pollinators of alfalfa in California. *J. Econ. Entomol.* 40:349–57
69. MacKenzie K, Javorek S. 1997. The potential of alfalfa leafcutter bees (*Megachile rotundata* L.) as pollinators of cranberry (*Vaccinium macrocarpon* Aiton). *Acta Hort.* 437:345–51
70. Maeta Y, Adachi K. 2005. Nesting behaviors of the alfalfa leaf-cutting bee, *Megachile (Eutricharaea) rotundata* (Fabricius) (Hymenoptera, Megachilidae). *Chugoku Kontyu* 18:5–21
71. Maeta Y, Kitamura T. 2005. On the number of eggs laid by one individual of females in the alfalfa leaf-cutting bee, *Megachile (Eutricharaea) rotundata* (Fabricius) (Hymenoptera, Megachilidae). *Chugoku Kontyu* 19:39–43
72. Maeta Y, Kitamura T. 2007. Efficiency of seed production in breeding of ladino clover by three species of wild bees, *Ceratina flavipes* Smith, *Megachile rotundata* (Fabricius) and *Megachile spissula* Cockerell (Hymenoptera, Apidae and Megachilidae). *Chugoku Kontyu* 21:35–53
73. Matthews RW, González JM, Matthews JR, Deyrup LD. 2009. Biology of the parasitoid *Melittobia* (Hymenoptera: Eulophidae). *Annu. Rev. Entomol.* 54:251–66
74. McCorquodale DB, Owen RE. 1997. Allozyme variation, relatedness among progeny in a nest, and sex ratio in the leafcutter bee, *Megachile rotundata* (Fabricius) (Hymenoptera: Megachilidae). *Can. Entomol.* 129:211–19
75. Menzel R. 2001 Behavioral and neural mechanisms of learning and memory as determinants of flower constancy. In *Cognitive Ecology of Pollination: Animal Behavior and Floral Evolution*, ed. L Chittka, JD Thomson, 2:21–40. Cambridge, UK: Cambridge Univ. Press. 343 pp.
76. Michener CD. 2007. *The Bees of the World*. Baltimore, MD: Johns Hopkins Univ. Press. 953 pp.
77. Minkley RL, Cane JH, Kervin L, Yanega D. 2003. Biological impediments to measures of competition among introduced honey bees and desert bees (Hymenoptera: Apiformes). *J. Kans. Entomol. Soc.* 76:306–19
78. Móczár L. 1961. The distribution of wild bees in the lucerne fields of Hungary (Hymenoptera: Apoidea). *Ann. Hist. Nat. Mus. Natl. Hungar. (Pars Zool.)* 53:451–61
79. Mommaerts V, Sterk G, Smagghe G. 2006. Hazards and uptake of chitin synthesis inhibitors in bumblebees *Bombus terrestris*. *Pest Manag. Sci.* 62:752–58
80. Morse RA, Calderone NW. 2000. The value of honey bees as pollinators of U.S. crops in 2000. *Bee Cult.* 128:1–15
81. O'Neill KM, O'Neill RP, Blodgett S, Fultz J. 2004. Composition of pollen loads of *Megachile rotundata* in relation to flower diversity (Hymenoptera: Megachilidae). *J. Kans. Entomol. Soc.* 77:619–25
82. Ortiz-Sanchez FJ, Aguirre-Segura A. 1991. Composition and seasonal patterns of a community of Apoidea in Almería. *EOS* 67:3–22

83. Owen RE, McCorquodale DB. 1994. Quantitative variation and heritability of postdiapause development time and body size in the alfalfa leafcutting bee (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.* 87:922–27
84. Pankiw P, Lieverse JAC, Siemens B. 1980. The relationship between latitude and the emergence of alfalfa leafcutter bees *Megachile rotundata* (Hymenoptera: Megachilidae). *Can. Entomol.* 112:555–58
85. Parker FD. 1978. An illustrated key to alfalfa leafcutter bees *Eutricharaca* (Hymenoptera: Megachilidae). *Pan-Pac. Entomol.* 54:61–64
86. Parker FD, Tepedino VJ. 1982. Maternal influence on diapause in the alfalfa leafcutting bee (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.* 75:407–10
87. Parker FD, Teranishi R, Olson AC. 1983. Influence of attractants on nest establishment by the alfalfa leafcutting bee (Hymenoptera: Megachilidae) in styrofoam and rolled paper. *J. Kans. Entomol. Soc.* 56:477–82
88. Paulmier I, Bagnères AG, Afonso CMM, Dusticier G, Rivière G, Clément JL. 1999. Alkenes as a sexual pheromone in the alfalfa leaf-cutter bee *Megachile rotundata*. *J. Chem. Ecol.* 25:471–90
89. Paxton RJ. 2005. Male mating behaviour and mating systems of bees: an overview. *Apidologie* 36:145–56
90. Peach ML, Alston DG, Tepedino VJ. 1995. Sublethal effects of carbaryl bran bait on nesting performance, parental investment, and offspring size and sex ratio of the alfalfa leafcutting bee (Hymenoptera, Megachilidae). *Environ. Entomol.* 24:34–39
91. Peterson JH, Roitberg BD. 2006. Impacts of flight distance on sex ratio and resource allocation to offspring in the leafcutter bee, *Megachile rotundata*. *Behav. Ecol. Sociobiol.* 59:589–96
92. Peterson JH, Roitberg BD. 2006. Impact of resource levels on sex ratio and resource allocation in the solitary bee, *Megachile rotundata*. *Environ. Entomol.* 35:1404–10
93. Pitts-Singer TL. 2004. Examination of ‘pollen balls’ in nests of the alfalfa leafcutting bee, *Megachile rotundata*. *J. Apic. Res.* 43:40–46
94. Pitts-Singer TL. 2007. Olfactory response of megachilid bees, *Osmia lignaria*, *Megachile rotundata*, and *M. pugnata*, to individual cues from old nest cavities. *Environ. Entomol.* 36:402–8
95. **Pitts-Singer TL. 2008. Past and present management of alfalfa bees. In *Bee Pollination in Agricultural Ecosystems*, ed. RR James, TL Pitts-Singer, 7:105–23. New York: Oxford Univ. Press. 232 pp.**
96. Pitts-Singer TL, Bosch J. 2010. Nest establishment, pollination efficiency, and reproductive success of *Megachile rotundata* (Hymenoptera: Megachilidae) in relation to resource availability in field enclosures. *Environ. Entomol.* 39:149–58
97. Pitts-Singer TL, James RR. 2005. Emergence success and sex ratio of commercial alfalfa leafcutting bees from the United States and Canada. *J. Econ. Entomol.* 98:1785–90
98. Pitts-Singer TL, James RR. 2008. Do weather conditions correlate with findings in failed, provision-filled nest cells of *Megachile rotundata* (Hymenoptera: Megachilidae) in western North America? *J. Econ. Entomol.* 101:674–85
99. Pitts-Singer TL, James RR. 2009. Prewinter management affects *Megachile rotundata* (Hymenoptera: Megachilidae) prepupal physiology and adult emergence and survival. *J. Econ. Entomol.* 102:1407–16
100. Rank GH, Goerzen DW. 1982. Effect of incubation temperatures on emergence of *Megachile rotundata* (Hymenoptera: Megachilidae). *J. Econ. Entomol.* 75:467–71
101. Rank GH, Rank FP. 1989. Diapause intensity in a French univoltine and a Saskatchewan commercial strain of *Megachile rotundata* (Fab.). *Can. Entomol.* 121:141–48
102. Rank GH, Rank FP, Watts R. 1990. Chalkbrood (*Ascosphaera aggregata* Skou.) resistance of a univoltine strain of the alfalfa leafcutting bee, *Megachile rotundata* F. *J. Appl. Entomol.* 109:524–27
103. Reed CC. 1995. Species richness of insects on prairie flowers in southeastern Minnesota. In *Proc. 14th Annu. N. Am. Prairie Conf.*, ed. DC Hartnett, pp. 103–15. Manhattan: Kans. State Univ.
104. Richards K. 1995. The alfalfa leafcutter bee, *Megachile rotundata*: a potential pollinator for some annual forage clovers. *J. Apic. Res.* 34:115–21
105. **Richards KW. 1984. Alfalfa leafcutter bee management in Western Canada. *Agric. Can.* 1495/E:1–53**
106. Richards KW. 1994. Ovarian development in the alfalfa leafcutter bee, *Megachile rotundata*. *J. Apic. Res.* 33:199–203

---

95. Review of alfalfa pollination by two managed solitary bees. Other chapters in book cover different aspects of solitary bees in agriculture.

---



---

105. A comprehensive guide to using ALCB for alfalfa seed production that includes ALCB life history and is largely based on author’s research. A needed update to this guide is Reference 28.

---

107. Richards KW. 1997. Potential of the alfalfa leafcutter bee, *Megachile rotundata* (F.) (Hym., Megachilidae) to pollinate hairy and winter vetches (*Vicia* spp.). *J. Appl. Entomol. Z. Angew. Entomol.* 121:225–29
108. Richards KW, Krunic MD. 1990. Introduction of alfalfa leafcutter bees to pollinate alfalfa in Yugoslavia. *Entomologist* 109:130–35
109. Richards KW, Whitfield GH. 1988. Emergence and survival of leafcutter bees, *Megachile rotundata*, held at constant incubation temperatures (Hymenoptera: Megachilidae). *J. Apic. Res.* 27:197–204
110. Riedl H, Johansen E, Brewer L, Barbour J. 2006. *How to reduce bee poisoning from pesticides. Rep. PWN 591.* Pac. Northwest Ext. Publ., Oregon State Univ., Corvallis
111. Robertson C. 1930. Proterandry and flight of bees (Hymen.: Apoidea). III. *Entomol. News* 41:331–36
112. Rossi BH, Nonacs P, Pitts-Singer TL. 2010. Sexual harassment by males reduces female fecundity in the alfalfa leafcutting bee, *Megachile rotundata*. *Anim. Behav.* 79:165–71
113. Sheffield CS. 2008. Summer bees for spring crops? Potential problems with *Megachile rotundata* (Fab.) (Hymenoptera: Megachilidae) as a pollinator of lowbush blueberry (Ericaceae). *J. Kans. Entomol. Soc.* 81:276–87
114. Small E, Brookes B, Lefkovitch LP, Fairey DT. 1997. A preliminary analysis of the floral preferences of the alfalfa leafcutting bee, *Megachile rotundata*. *Can. Field Nat.* 111:445–53
115. Smith CR, Toth AL, Suarez AV, Robinson GE. 2008. Genetic and genomic analyses of the division of labour in insect societies. *Nat. Rev. Genet.* 9:735–48
116. Soroka JJ, Goerzen DW, Falk KC, Bett KE. 2001. Alfalfa leafcutting bee (Hymenoptera: Megachilidae) pollination of oilseed rape (*Brassica napus* L.) under isolation tents for hybrid seed production. *Can. J. Plant Sci.* 81:199–204
117. St. Armand PC, Skinner DZ, Peaden RN. 2000. Risk of alfalfa transgene dissemination and scale-dependent effects. *Theor. Appl. Genet.* 101:107–14
118. Stark JD, Banks JE. 2003. Population-level effects of pesticides and other toxicants on arthropods. *Annu. Rev. Entomol.* 48:505–19
119. Stephen WP. 1961. Artificial nesting sites for the propagation of the leaf-cutter bee, *Megachile (Eutricharaea) rotundata*, for alfalfa pollination. *J. Econ. Entomol.* 54:989–93
120. Stephen WP. 1962. Propagation of the leaf-cutter bee for alfalfa seed production. *Agric. Exp. Stn. Bull.* 586:1–16. Oregon State Univ., Corvallis
121. Stephen WP. 1981. The design and function of field domiciles and incubators for leafcutting bee management (*Megachile rotundata* (Fabricius)). *Oregon State Coll. Agric. Exp. Stn. Bull.* 654:1–13
122. **Stephen WP. 2003. Solitary bees in North American agriculture: a perspective. In *For Non-native Crops, Whence Pollinators of the Future?*, ed. K Strickler, JH Cane, 3:41–66. Lanham, MD: Entomol. Soc. Am. 204 pp.**
123. Stephen WP, Undurraga JM. 1976. X-radiography, an analytical tool in population studies of the leaf-cutter bee *Megachile pacifica*. *J. Apic. Res.* 15:81–87
124. Strickler K. 1999. Impact of flower standing crop and pollinator movement on alfalfa seed yield. *Environ. Entomol.* 28:1067–76
125. Szabo TI, Smith MV. 1972. The influence of light intensity and temperature on the activity of the alfalfa leaf-cutter bee *Megachile rotundata* under field conditions. *J. Apic. Res.* 11:157–65
126. Tasei JN. 1975. Le problème de l'adaptation de *Megachile (Eutricharea) pacifica* Panz. (Megachilidae) Américain en France. *Apidologie* 6:1–57
127. Tasei JN, Masure MM. 1978. Sur quelques facteurs influençant le développement de *Megachile pacifica* Panz (Hymenoptera, Megachilidae). *Apidologie* 9:273–90
128. Tepedino VJ. 1982. Flower visitation and pollen collection records for bees of high altitude shortgrass prairie in southeastern Wyoming, USA. *Southwest. Entomol.* 7:16–25
129. Tepedino VJ. 1983. An open-field test of *Megachile rotundata* as a potential pollinator in hybrid carrot fields. *J. Apic. Res.* 22:64–68
130. Tepedino VJ. 1997. A comparison of the alfalfa leafcutting bee (*Megachile rotundata*) and the honey bee (*Apis mellifera*) as pollinators for hybrid carrot seed in field cages. *Acta Hort.* 437:457–61
131. Tepedino VJ, Parker FD. 1986. Effect of rearing temperature on mortality, second-generation emergence, and size of adult in *Megachile rotundata* (Hymenoptera: Megachilidae). *J. Econ. Entomol.* 79:974–77

---

122. Comprehensive review and history of establishment of solitary bees for pollination. Other chapters in book cover different aspects of alternative pollinators in agriculture.

---



132. Tepedino VJ, Parker FD. 1988. Alternation of sex ratio in a partially bivoltine bee, *Megachile rotundata* (Hymenoptera: Megachilidae). *Ann. Entomol. Soc. Am.* 81:467–76
133. Tezuka T, Maeta Y. 1993. Effect of UVA film on extranidal activities of three species of bees. *Jpn. J. Appl. Entomol. Zool.* 37:175–80
134. Torchio PF. 1973. Relative toxicity of insecticides to the honey bee, alkali bee, and alfalfa leafcutting bee (Hymenoptera: Apidae, Halictidae, Megachilidae). *J. Kans. Entomol. Soc.* 46:446–53
135. Torchio PF. 1979. An eight-year field study involving control of *Sapyga pumila* Cresson (Hymenoptera: Sapygidae), a wasp parasite of the alfalfa leafcutter bee, *Megachile pacifica* Panzer. *J. Kans. Entomol. Soc.* 52:412–19
- 136. Trostle G, Torchio PF. 1994. Comparative nesting behavior and immature development of *Megachile rotundata* (Fabricius) and *Megachile apicalis* Spinola (Hymenoptera, Megachilidae). *J. Kans. Entomol. Soc.* 67:53–72**
137. Undurraga JM, Stephen WP. 1980. Effects of temperature on development and survival in post-diapausing alfalfa leafcutting bee prepupae and pupae (*Megachile rotundata* (F.): Hymenoptera: Megachilidae). I. High temperatures. *J. Kans. Entomol. Soc.* 53:669–76
138. Undurraga JM, Stephen WP. 1980. Effects of temperature on development and survival in post-diapausing alfalfa leafcutting bee prepupae and pupae (*Megachile rotundata* (F.): Hymenoptera: Megachilidae). II. Low temperatures. *J. Kans. Entomol. Soc.* 53:677–82
139. Van Deynze AE, Fitzpatrick SM, Hammon B, McCaslin MH, Putnam DH, et al. 2008. Gene flow in alfalfa: biology, mitigation, and potential impact on production. *CAST Spec. Publ.* 28. Counc. Agric. Sci. Technol., Ames, IA. 30 pp.
140. Vandenberg JD, Fichter BL, Stephen WP. 1980. Spore load of *Ascospaera* spp. on emerging adults of the alfalfa leafcutting bee, *Megachile rotundata*. *Appl. Environ. Microbiol.* 39:650–55
141. Vandenberg JD, Stephen WP. 1982. Etiology and symptomatology of chalkbrood in the alfalfa leafcutting bee, *Megachile rotundata*. *J. Invertebr. Patbol.* 39:133–37
142. Vorel CA. 2010. *Learning ability and factors influencing nest establishment of the solitary bees Osmia lignaria and Megachile rotundata (Hymenoptera: Megachilidae)*. PhD thesis. Utah State Univ. 126 pp.
143. Vorel CA, Pitts-Singer TL. 2010. The proboscis extension reflex not elicited in megachilid bees. *J. Kans. Entomol. Soc.* 83:80–83
144. Wedmann S, Wappler T, Engel MS. 2009. Direct and indirect fossil records of megachilid bees from the Paleogene of Central Europe (Hymenoptera: Megachilidae). *Naturwissenschaften* 96:703–12
145. Wheeler D. 1996. The role of nourishment in oogenesis. *Annu. Rev. Entomol.* 41:407–31
146. Wightman JA, Rogers VM. 1978. Growth, energy and nitrogen budgets and efficiencies of the growing larvae of *Megachile pacifica* (Hymenoptera: Megachilidae). *Oecologia* 36:245–57
147. Wittmann D, Blochtein B. 1995. Why males of leafcutter bees hold the females' antennae with their front legs during mating. *Apidologie* 26:181–95
148. Woodward DR. 1994. Predators and parasitoids of *Megachile rotundata* (F.) (Hymenoptera: Megachilidae) in South Australia. *J. Aust. Entomol. Soc.* 33:13–15
149. Xu J, James R. 2009. Genes related to immunity, as expressed in the alfalfa leafcutting bee, *Megachile rotundata*, during pathogen change. *Insect Mol. Biol.* 18:785–94
150. Yocum GD, Kemp WP, Bosch J, Knoblett JN. 2005. Temporal variation in overwintering gene expression and respiration in the solitary bee *Megachile rotundata*. *J. Insect Physiol.* 51:621–29

---

136. Describes nesting behavior and immature development of *Megachile* species.

---



# Contents

<i>Bemisia tabaci</i> : A Statement of Species Status <i>Paul J. De Barro, Shu-Sheng Liu, Laura M. Boykin, and Adam B. Dinsdale</i> .....	1
Insect Seminal Fluid Proteins: Identification and Function <i>Frank W. Avila, Laura K. Sirot, Brooke A. LaFlamme, C. Dustin Rubinstein, and Mariana F. Wolfner</i> .....	21
Using Geographic Information Systems and Decision Support Systems for the Prediction, Prevention, and Control of Vector-Borne Diseases <i>Lars Eisen and Rebecca J. Eisen</i> .....	41
Salivary Gland Hypertrophy Viruses: A Novel Group of Insect Pathogenic Viruses <i>Verena-Ulrike Lietze, Adly M.M. Abd-Alla, Marc J.B. Vreysen, Christopher J. Geden, and Drion G. Boucias</i> .....	63
Insect-Resistant Genetically Modified Rice in China: From Research to Commercialization <i>Mao Chen, Anthony Shelton, and Gong-yin Ye</i> .....	81
Energetics of Insect Diapause <i>Daniel A. Hahn and David L. Denlinger</i> .....	103
Arthropods of Medicoveterinary Importance in Zoos <i>Peter H. Adler, Holly C. Tuten, and Mark P. Nelder</i> .....	123
Climate Change and Evolutionary Adaptations at Species' Range Margins <i>Jane K. Hill, Hannah M. Griffiths, and Chris D. Thomas</i> .....	143
Ecological Role of Volatiles Produced by Plants in Response to Damage by Herbivorous Insects <i>J. Daniel Hare</i> .....	161
Native and Exotic Pests of <i>Eucalyptus</i> : A Worldwide Perspective <i>Timothy D. Paine, Martin J. Steinbauer, and Simon A. Lawson</i> .....	181

Urticating Hairs in Arthropods: Their Nature and Medical Significance <i>Andrea Battisti, Göran Holm, Bengt Fagrell, and Stig Larsson</i> .....	203
The Alfalfa Leafcutting Bee, <i>Megachile rotundata</i> : The World's Most Intensively Managed Solitary Bee <i>Theresa L. Pitts-Singer and James H. Cane</i> .....	221
Vision and Visual Navigation in Nocturnal Insects <i>Eric Warrant and Marie Dacke</i> .....	239
The Role of Phytopathogenicity in Bark Beetle–Fungal Symbioses: A Challenge to the Classic Paradigm <i>Diana L. Six and Michael J. Wingfield</i> .....	255
Robert F. Denno (1945–2008): Insect Ecologist Extraordinaire <i>Micky D. Eubanks, Michael J. Raupp, and Deborah L. Finke</i> .....	273
The Role of Resources and Risks in Regulating Wild Bee Populations <i>T'ai H. Roulston and Karen Goodell</i> .....	293
Venom Proteins from Endoparasitoid Wasps and Their Role in Host-Parasite Interactions <i>Sassan Asgari and David B. Rivers</i> .....	313
Recent Insights from Radar Studies of Insect Flight <i>Jason W. Chapman, V. Alistair Drake, and Don R. Reynolds</i> .....	337
Arthropod-Borne Diseases Associated with Political and Social Disorder <i>Philippe Brouqui</i> .....	357
Ecology and Management of the Soybean Aphid in North America <i>David W. Ragsdale, Douglas A. Landis, Jacques Brodeur, George E. Heimpel, and Nicolas Desneux</i> .....	375
A Roadmap for Bridging Basic and Applied Research in Forensic Entomology <i>J.K. Tomberlin, R. Mohr, M.E. Benbow, A.M. Tarone, and S. VanLaerhoven</i> .....	401
Visual Cognition in Social Insects <i>Aurore Avarguès-Weber, Nina Deisig, and Martin Giurfa</i> .....	423
Evolution of Sexual Dimorphism in the Lepidoptera <i>Cerisse E. Allen, Bas J. Zwaan, and Paul M. Brakefield</i> .....	445
Forest Habitat Conservation in Africa Using Commercially Important Insects <i>Suresh Kumar Raina, Esther Kioko, Ole Zethner, and Susie Wren</i> .....	465
Systematics and Evolution of Heteroptera: 25 Years of Progress <i>Christiane Weirauch and Randall T. Schuh</i> .....	487