## Gratiola heterosepala

Scrophulariaceae Boggs Lake hedge hyssop






Virgina Crosby



best survey times
J|F|M|A|M|J|J|A|S|O|N|D





From: Threatened and Endangered Vascular Plants of Oregon (1982) R. J. Meinke

Erect, amphibious, annual herb. Stems 2-10 cm high, glabrous, stout and fistulose below, slender and glandular above. Leaves opposite, lower leaves linear-lanceolate, 1-2 cm long, becoming shorter and rounded above, 2-5 mm. Inflorescence glandular; pedicels 10-23 mm, slender, erect; bracts 0. Flowers few; sepals unequal, 4-6 mm long, lower two sepals distinct, upper three partially fused; upper corolla lobes yellow, three lower lobes white, 6-8 mm long; stigmas 2, unequal. Fruit a capsule, spheric, 4-6 mm in diameter; seeds oblong and ribbed.

#### Lookalikes Gratiola ebracteata Gratiola neglecta


### differs from featured plant by

its uniformly white corolla lobes and yellow tubes. its two bracts below the calyx.

# Gratiola heterosepala H. Mason & Bacig.

Boggs Lake hedge hyssop PLANTS symbol: GRHE

August 2019 status Federal:SOC; Oregon:LT; ORBIC: List 1



Distribution: Fresno Co. California north to extreme southern Lake Co. Oregon.

**Habitat:** Shallow water, mud or damp soil at edges of lakes and vernal pools.

Elevation: 1620 - 1650 m; less than 1200 m in California.

Best survey time (in flower): April – June

#### Associated species:

Marsilea vestita (Hairy waterclover) Downingia laeta (Great Basin calicoflower) Juncus (Rush)









