Kalmiopsis fragrans

Ericaceae Fragrant kalmiopsis




Melissa A. Carr

Bruce N. Newhouse







Plant is a perennial, evergreen, multi-stemmed shrub, may be erect or trailing and matted. **Stems** from a thickened base, glabrous, exfoliating, reddish-gray bark. **Leaves** mostly terminal, crowded, shiny, dark green and glabrous above, golden crystalline glands below, elliptic-obovate, short petiole. **Inflorescence** terminal corymbose raceme, 4-7 flowers, fragrant, calyx urn or bell-shaped, pale pink, sepals overlapping at the base, corolla 16-28 mm, reddish-purple to pink, lacking a defined throat, nearly flat as petal lobes reflex, stamens 10, spreading to erect, protruding from corolla, dense golden pubescence at base of filaments. **Fruit** a depressed capsule, shallowly five lobed and glandular-warty, 3-5 mm wide.

Lookalikes *Kalmiopsis leachiana*


differs from featured plant by

its smaller corolla, shorter stamens, lack of golden pubescence at the base of the filaments, well-defined corolla tube, and its lack of fragrance.

Kalmiopsis fragrans Meinke & Kaye

Fragrant kalmiopsis PLANTS symbol:

August 2019 status Oregon:C; ORBIC: List 1



Distribution: Douglas County, Oregon

Habitat: Rock outcrops and crevices, in sun or shady coniferous forests.

Elevation: 400-1600 m

Best survey time (in flower): Late April to late June

Associated species:

Pseudotsuga menziesii (Douglas fir)

Abies grandis (Grand fir)

Tsuga heterophylla (Western hemlock)

Calocedrus decurrens (Incense cedar)

Thuja plicata (Western redcedar)

Pinus lambertiana (Sugar pine)

Berberis nervosa (Cascade Oregon grape, Long leaved Oregon grape)

Gaultheria shallon (Salal)

Oxalis oregana (Oregon wood sorrel, Redwood sorrel)

Polystichum munitum (Common sword fern)

Linnaea borealis (Twinflower)

Rhododendron macrophyllum (Pacific rhododendron, Western rhododendron)

Acer circinatum (Vine maple)

Toxicodendron diversilobum (Poison oak)

