

Streptanthus howelliiBrassicaceae
Howell's streptanthus




ober Mansheld



best survey times
J|F|M|A|M|J|J|A|S|O|N|D





Plant is erect perennial, entire plant glabrous and glaucous. **Stems** stout and leafy, simple or branched, 30-70 cm tall, occasionally dilated upward. **Leaves** rather fleshy, obovate to spatulate, denticulate or entire, not clasping at the base. **Inflorescence** a long, loose raceme, sparsely flowered, bractless flowers have broad greenish sepals and oblong dark purple petals. **Fruit** is a flat and strongly curved silique.

LookalikesStreptanthus spp.


differs from featured plant by

Featured plant differs from other *Streptanthus* species by its non-clasping leaves, flat fruits, and bractless flowers.

Streptanthus howellii S. Watson

Howell's streptanthus PLANTS symbol: STHO

August 2019 status Oregon:C; ORBIC: List 1



Distribution: Josephine and Curry Counties, SW Oregon; and Del Norte County, California.

Habitat: Dry, serpentine slopes, mixed evergreen forests, open pine woods or brushy areas.

Elevation: 400-1700 m

Best survey time (in flower): Mid-June through August

Associated species:

Ceanothus cuneatus (Buckbrush)

Arctostaphylos viscida (White leaved manzanita, Whiteleaf manzanita)

Eriogonum pendulum (Waldo eriogonum, Waldo buckwheat)

Lomatium howellii (Howell's desert parsley)

Pinus jeffreyi (Jeffrey pine)

Quercus chrysolepis (Canyon live oak, Maul oak)

Rhamnus californica (California buckthorn)

