


Sullivantia oregana Saxifragaceae Sullivantia







Delicate, yellowish-green perennial spreading by long slender stolons, nearly or quite glabrous except for some glandular pubescence on the upper portions of the flowering stems and on the inflorescence, the hairs mostly purplish-tipped. Flowering stems 5-20(25) cm tall, with 1-3 leaves that are greatly reduced upward. Basal leaves long-petiolate with membranous stipules, blades reniform, 1-10 cm broad, lobed into 7-9 cuneate segments and again once or twice sharply toothed. Inflorescence a panicle. Flowers erect, but becoming sharply reflexed in fruit; calyx glabrous, pale green, 2.5-3.5 mm long, more or less campanulate; petals slightly longer than the calyx lobes, the blade oval to obovate-oblanceolate, narrowed to a very short, broad claw; stamens shorter than the sepals, the cordate anthers ca. equaling the slender filaments. Capsule ca. 4 mm long; seeds brown, ca. 1.5 mm long.




stolons present

differs from featured plant by stipules membranous, with dark red bristles on margins; petals longer (5-6 mm) and exceeding the calvx

Illustration by Jeanne R. Janish.
VASCULAR PLANTS OF THE
PACIFIC NORTHWEST (1961)
Hitchcock, Cronquist, & Ownbey,
courtesy of University of Washington Press.

small flowers




## Sullivantia oregana S. Watson

Sullivantia

PLANTS symbol: SUOR

August 2019 status Federal:SOC; Oregon:C; ORBIC: List 1



**Distribution:** Columbia River Gorge, northern Willamette Valley and Cascade Mountains; Skamania Co., southern Washington.

**Habitat:** Moist, shaded cliffs, especially near waterfalls; surrounding

forest dominated by Douglas fir.

**Elevation:** 30-365 (1370) m

Best survey time(in flower): late May-August

## Associated species:

Heracleum maximum (cow parsnip) Tolmiea menziesii (piggy back plant) Oxalis trilliifolia (trillium leaved oxalis)



