


Barbara L. Wilson, Richard Brainerd,  
Danna Lytjen, Bruce Newhouse, and Nick Otting  
of the *Carex* Working Group

# Field Guide to the Sedges of the Pacific Northwest

---

Second Edition



# Field Guide to the Sedges of the Pacific Northwest

*Second Edition*



The authors and publisher gratefully acknowledge the generous support of the following organizations, which made possible the writing, photography, and publication of the Field Guide to the Sedges of Oregon and Washington.



USDI Bureau of Land Management



USDA Forest Service



U.S. Environmental Protection Agency



USDA Natural Resources Conservation Service



Oregon State University Herbarium  
Department of Botany and Plant Pathology



Oregon Flora Project



Native Plant Society of Oregon

Field Guide to the Sedges of the Pacific Northwest  
*Second Edition*



*by*

Barbara L. Wilson  
Richard E. Brainerd  
Danna Lytjen  
Bruce Newhouse  
& Nick Otting

of the *Carex* Working Group

---

Oregon State University Press  
Corvallis



www.carexworkinggroup.com  
the place to go for updates to the identification key  
and for other news of Pacific Northwest sedges

The paper in this book meets the guidelines for permanence and durability of the Committee on Production Guidelines for Book Longevity of the Council on Library Resources and the minimum requirements of the American National Standard for Permanence of Paper for Printed Library Materials Z39.48-1984.

**Library of Congress Cataloging-in-Publication Data**

Field guide to the sedges of the Pacific Northwest / by Barbara L. Wilson ... [et al.].

p. cm.

Includes bibliographical references and index.

ISBN-13: 978-0-87071-197-8 (alk. paper)

1. Carex--Oregon--Identification. 2. Carex--Washington (State)--Identification. I. Wilson, Barbara L., 1951-

QK495.C997F54 2008

584'.8409795--dc22

2007039207

© 2008 *Carex* Working Group

All rights reserved. First published in 2008 by Oregon State University Press

Second edition 2014

Printed in China



**Oregon State University Press**

121 The Valley Library

Corvallis OR 97331-4501

541-737-3166 • fax 541-737-3170

<http://osupress.oregonstate.edu>



Field Guide to Sedges of the Pacific Northwest

*is dedicated to*

Danna Lytjen  
(1947 - 2006)

Danna was a charter member of the *Carex* Working Group and was instrumental in bringing this book into existence. She was an insightful observer of nature, an accomplished field biologist, a mother, and a grandmother. Her abilities as a scientist, her love of nature, and her kindness are greatly missed, but they live on in the hearts of those who knew her.

“I would like to know the grasses and sedges—and care. Then my least journey into the world would be a field trip, a series of happy recognitions.”—Annie Dillard

“I would like to go into perfectly new and wild country. I wish to lose myself amid reeds and sedges and wild grasses that have not been touched.”—Henry David Thoreau

Botanically, where there's sedge, there's often confusion.—Anon.

## Contents

10	Acknowledgments
12	Introduction
14	<i>Carex</i> Ecology
22	<i>Carex</i> Nomenclature and Classification
26	<i>Carex</i> Morphology
28	Sedge Parts (illustrations)
31	How to Use the Identification Key
34	Collecting Sedges
36	Abbreviations and Symbols
37	Identification Key to Sedges of the Pacific Northwest
38	Key A. Inflorescences consisting of a single spike
41	Key B. Inflorescences normally entirely staminate
42	Key C. Perigynia hairy
44	Key D. <i>Kobresia</i> ; perigynia open on one side; short cespitose plants of high elevations
44	Key E. Styles persistent; perigynia usually $\pm$ inflated
45	Key F. Stigmas 3; perigynia not pubescent; styles deciduous
51	Key G. Stigmas 2
55	Key H. Stigmas 2; perigynia unwinged; spikes androgynous, entirely pistillate or with staminate flowers mixed irregularly among the pistillate flowers
60	Key I. Stigmas 2; spikes gynecandrous; perigynia unwinged
62	Key J. Perigynia winged; stigmas 2; mainly section <i>Ovales</i>

71	Species Accounts
408	Sedges Rarely Found in Oregon or Washington
410	Sedges with Distinctive Traits or Habitats
413	Glossary
418	Illustration Credits
420	References
422	Index
432	Map of the counties of Oregon and Washington



## Acknowledgments

The *Carex* Working Group (CWG) began in 1993 when botanists intrigued by taxonomic and ecological issues in the genus *Carex* met at the Oregon State University Herbarium to exchange ideas. At that initial meeting, Dr. Dan Norris suggested that we produce an Atlas of Oregon *Carex*. Peter Zika became the informal leader of the CWG. This careful observer of details inspired us, showed us identification tricks, and led us on field trips. Other early members included Keli Kuykendall, Dr. Francisco (Ankie) Camacho, Jim Oliphant, and Dr. Kate Dwire.

The Native Plant Society of Oregon provided critically important early funding for the *Carex* Working Group's field work and databasing efforts. We are grateful to the society for publishing *The Atlas of Oregon Carex* (Wilson et al. 1999) as its first occasional paper.

Many people and organizations have helped make the Field Guide happen. Major financial support for the project was provided by the Bureau of Land Management. We especially thank Nora Taylor, who first arranged this financial support; Joan Seevers, without whom the project would have died; and Ron Exeter, our patient and effective project manager. BLM Botanists Jean Findley, Claire Hibler, Rick Hall, Ron Halvorson, Lucille Housely, Linda Mazzu, Mark Mousseaux, Bruce Rittenhouse, Tim Rodenkirk, Nancy Sawtelle, Jennifer Wheeler, and Lou Whiteaker made it possible for us to do much of the field work that underlies this book.

Major financial support was also provided by the USDA Forest Service. We thank Russ Holmes for making that possible. Botanists Paula Brooks, David Isle, Jennifer Lippert, Julie Kierstead Nelson, Teresa Ohlsen, Laura Potash Martin, Wayne Rolle, Marty Stein, Maria Ulloa, Gene Yates, and Joan Ziegltrum facilitated field work.

Significant financial support was provided by the U.S. Environmental Protection Agency, with help from Linda Storm. Significant financial support was also provided by the USDA Natural Resources Conservation Service. We thank Kathy Pendergrass for her assistance and Amy Bartow for information about propagation of sedges.

We could not have done this work without the aid of the Oregon State University Herbarium. We thank Dr. Aaron Liston, Director, for early and continued encouragement, and Dr. Richard Halse, Curator, for aid with loans and specimen storage. The Oregon Flora Project (OFP) provided databasing and mapping. The late Dr. Scott Sundberg provided much advice and arranged the initial databasing of *Carex* specimens and sightings. Dr. Linda Hardison and Thea Cook have provided continuing assistance. Katie Mitchell generated the range maps used in the book, managed the databases, assisted in the field, and supervised student workers. We are grateful for her quiet competence.

Our maps are based on databases of herbarium specimens and sightings provided by the Oregon Flora Project (Oregon State University Herbarium), the

herbaria of University of Washington and Washington State University, and the other herbaria which contributed data to the Oregon Flora Project.

Plant descriptions are modified from the *Carex* treatment in *Flora of North America* (Ball and Reznicek 2002). Tim Jones provided some of the descriptive material from the database for his on-line interactive key (Jones 2007).

University of Washington Press allowed us to use Jeanne R. Janish's fine drawings from the *Vascular Plants of the Pacific Northwest* (Hitchcock et al. 1969). New York Botanical Garden granted permission to use Harry Charles Creutzburg's drawings from *North American Cariceae* (Mackenzie 1940). Erin Stangel photographed perigynia and herbarium specimens at the Oregon State University Herbarium. Nancy Shaw made available photographs from the *Field Guide to Intermountain Sedges* (Hurd et al. 1998). We thank the following photographers for generously sharing their *Carex* photos with us: Gerald Carr (Oregon State University Herbarium), David Dister, Susan Farrington, Charles Feddema, Richard Heliwell (USDA Forest Service), Andrew Hipp (Morton Arboretum), Matthias Hoffmann (Martin-Luther-Universität Halle-Wittenberg), Ernst Horak, Emily Kapler, Tom Kaye (Institute for Applied Ecology), Natalie Kirchner, Steve Matson, Keir Morse, Julie Kierstead Nelson (USDA Forest Service), Anton A. Reznicek (University of Michigan Herbarium), Russ Schipper (Audubon Society of Kalamazoo), Forest and Kim Starr (Starr Environmental), Amadej Trnkoczy, Dana Visalli (Methow Biodiversity Project), Leanne Wallis, Fred Weinmann, Louise Wootton (Georgian Court University), Gene Yates (USDA Forest Service), and Peter Zika (University of Washington Herbarium). Stephen Meyers of the Oregon Flora Project formatted the new photos for the 2nd edition.

Dr. Anton A. Reznicek, our sedge guru, read parts of the manuscript and provided advice and insightful discussion. We thank Dr. Robert Naczi (Delaware State University) and Dr. Adolf Ceska (Royal British Columbia Museum) for information about sedges. Dr. Fred Weinmann not only provided information, but led highly productive field trips. Dana Visali and George Wooten provided information about unusual sedges of northern Washington. Dr. Karen Antell (Eastern Oregon University) provided records from northeast Oregon. Peter Lesica provided information about *C. idahoensis*. Gay Hunter (Olympic National Park) willingly took last-minute perigynium measurements for us. Dr. Lawrence Janeway (Herbarium at University of California, Chico) was involved in the discovery and description of *C. klamathensis*. Lynda Boyer (Heritage Seedlings, Inc.) provided information about *Carex* seed germination and propagation.

Dr. Mary O'Brien strengthened our early project proposals. Dr. Rhoda Love helped with both early proposals and late editing.

We are grateful to the participants in our sedge identification workshops, whose many suggestions greatly improved the identification key.

We thank Sami Gray for editing and research. We thank Mary Braun and Jo Alexander of OSU Press for their perseverance in assembling the book. Mary Harper of Access Points Indexing prepared the index.

## Introduction

Sedges are fun! Repeat this mantra whenever you find yourself facing the challenge of identifying a really difficult sedge. As with many of the more difficult plant groups, a positive attitude and a good sense of humor can smooth the way when you're stuck. Good identification tools help a lot too. Hopefully this book will assist you in becoming more comfortable with identifying sedges.

*Carex*—sedge—is the largest plant genus in the Pacific Northwest (PNW) with 164 species, and a total of 171 taxa when all subspecies and varieties are counted. These plants play essential roles in a variety of ecosystems and habitats, ranging from the seacoast to alpine slopes, from marshes, lakeshores, and river edges, to dry forests and sagebrush steppe. Particularly in wetlands, sedges often are community dominants, keystone species essential for stabilizing soil, maintaining stream quality, and providing habitat for fish, wildlife and plants. Because of their importance, sedges are frequently used for habitat restoration and enhancement projects. Knowledge of native ranges, habitat requirements, and ecological functions of sedges is essential if these projects are to be successful. Such knowledge begins with identifying the species that are present or that you want to reintroduce.

Sedges have a reputation for being devilishly difficult to identify. Much of the challenge comes from the fact that the plant parts used to identify most sedges are tiny, show great similarity between different species, and have unfamiliar names such as “perigynium.” But once you look carefully and master a bit of terminology and plant structure, it turns out that there are some useful “handles” you can use to make sedge identification a manageable, even enjoyable, activity. And having a name for your sedge is a starting point for learning about it and the places where it lives.

This book will be useful for professional botanists, ecologists, wetland scientists, wildlife biologists, and amateur plant enthusiasts. It contains an identification key and descriptions, range maps, photographs, and drawings of all *Carex* taxa known to occur naturally in Oregon and Washington. It also includes information on ecology, and morphology of sedges.

We hope this book will help you to identify most of the sedges that you encounter, and that you won't have to put unidentified “*Carex* sp.” on your species list ever again.

## Carex Ecology

Sedges perform important ecological functions in a variety of wetland and upland systems. They may be community dominants in an impressive diversity of habitats. They are most valued for stabilizing soil and as food for wild and domesticated animals. A few introduced species may spread as weeds, so understanding their methods of dispersal is important.

### Diversity

*Carex* reach their highest diversity in the north temperate and boreal zones. In the Pacific Northwest (PNW), *Carex* are especially diverse in the alpine zone. Sedges can achieve high species diversity in small areas because they tend to be microhabitat specialists. A few have wide ecological tolerances, and grow in a variety of habitats across the region, but most grow best in a very specific substrate, range of acidity, water and light regime, and elevation. Perhaps most importantly, each species has particular needs for water: deep water all year, deep water in spring and then dry, more or less dry all year, etc. Different species can occupy different zones across the hydrologic gradient around a pond, forming concentric rings from *C. utriculata* in the water through *C. pellita* and *C. scoparia* in seasonally wet marsh, to *C. pachystachya* on the mesic fringes and *C. tumulicola* on the dry slope above.

Climate change will seriously affect the distributions of the many PNW sedges that have limited habitat tolerances. The most vulnerable may be local populations of species such as *C. anthoxantha*, *C. circinata*, *C. pluriflora*, and *C. macrochaeta* that were probably widespread in the PNW at the end of the last ice age, 10,000 years ago. As the climate warmed, these species migrated north, leaving behind the few remnant PNW populations in small, isolated pockets of suitable habitat.

### Carex Below Ground

Sedge roots are probably much more interesting than we know. The roots of wetland species are adapted to maintain aerobic metabolism even in water-saturated soils. Air spaces called aerenchyma occur between the cells and allow diffusion of oxygen throughout *Carex* roots, stems, and leaves. This diffusion is efficient enough that excess oxygen may leak from *Carex* roots into adjacent saturated soils, where it oxidizes iron to produce “rust” surrounding the roots (Figure 1).

Most plants depend on an extensive network of mutualistic mycorrhizal fungi to obtain nutrients from the soil and water. However, about half the species in the sedge family lack mycorrhizae and many others are only facultatively mycorrhizal.



Figure 1. Roots of *Carex scoparia* in mud.

In the *Carex* that have mutualistic root fungi, arbuscular mycorrhizae (AM) may be the most common type. Ecotomycorrhizal associations occur in some sedges, such as *Kobresia myosuroides*. Dark septate fungi (DSF) occur in many sedges, especially in habitats where AM do not thrive, such as alpine tundra. The relationship of DSF and their *Carex* hosts needs more study, but appears to benefit both parties. A single plant may host both DSF and AM. Some sedge roots host other fungal species whose role is unclear.

*Carex* of wetlands frequently lack mycorrhizae because water-logged soil has too little oxygen to promote growth of mycorrhizal fungi. Several sedge adaptations contribute to this independence from mycorrhizal fungi (Muthukumar et al. 2004). Long rhizomes can explore the soil, producing roots in nutrient-rich patches and transporting the nutrients to other parts of the plant. Some sedges obtain nutrients unavailable to mycorrhizal fungi through root clusters in the uppermost soil layer, which contains much litter that could decompose to release nutrients, but which dries out too suddenly and thoroughly to provide good habitat for mycorrhizae. Each season, some sedges transport 80% or more of the nitrogen and phosphorus from their dying leaves to growing shoots. Arctic and alpine sedges need not wait for completion of the slow process of decomposition in cold soils because they can absorb small amino acids such as glycine and use them as a nitrogen source. These sedges also have enzymes on their root surfaces that remove phosphate ions from organic compounds in soil.

Not all below-ground sedge parts are roots. In fact, the most consistently perennial part of a *Carex* is its rhizome, an underground stem that gives rise to both roots and leafy shoots. Rhizomes explore the habitat, transfer nutrients from areas of concentrated resources to the shoots that need them, and determine how sedges compete with neighboring plants.

### Sedges, Streams, and Erosion

Sedges exert some of their most important effects on ecosystems by stabilizing soils in prairies, on streambanks, and in wetlands. In wet sedge meadows, 50–75% of the biomass is below ground. Rhizomatous, community-dominant sedges such as *C. nebrascensis*, *C. aquatilis*, *C. angustata*, and *C. utriculata* play an important role in stabilizing streamside soils east of the Cascades. Their long, strong



rhizomes and roots form a dense network that is tougher than that of dominant riparian zone grasses such as Kentucky Bluegrass (*Poa pratensis*) and Tufted Hairgrass (*Deschampsia cespitosa*). This network prevents both loss of small soil particles and mass wasting of streambanks. Along smaller streams, long rhizomes actually cross below the waterway, knitting the banks together. Even cespitose sedges can hold soil more effectively than one might expect (Figure 2).



Figure 2. Roots and soil-holding capacity of a small *Carex serratodens*.

Sedges are an important component of the streamside vegetation that provides habitat structure for diverse riparian invertebrates, which are part of the foundation of both aquatic and terrestrial food chains. Sedges also shade stream edges, providing cover for salmon, trout, and other cold-water species.

This is especially important in relatively treeless reaches east of the Cascades, such as parts of Oregon's John Day River where *C. nudata* may provide the only shade for fish.

Streambank stabilization by sedges can be seriously decreased by grazing livestock or elk. Ungrazed or lightly grazed sedges translocate adequate organic carbon to strengthen and extend rhizomes and roots, but more heavily grazed sedges cannot support the below-ground tissue, and the network of rhizomes and roots deteriorates. In one experiment, ungrazed wet sedge meadows had 60% more below-ground biomass and the soil infiltration rate was more than twice as high as in comparable grazed meadows. Reduction of below-ground biomass can result in stream incision, and the lowered water table further decreases the vigor of the sedge stands, which leads to additional loss of below-ground biomass. Not only does the reduced below-ground biomass lead to stream downcutting, but it also reduces forage production, leading to a spiral of riparian degradation when the animals graze the remaining stubble more intensely. With loss of herbaceous vegetation, grazers may resort to browsing willows with similar effects on their root systems, leading to additional erosion and stream incision. Grazing in sedge meadows while soils are wet can cause soil compaction and churning, directly damaging rhizomes and roots, and reducing their ability to penetrate and stabilize the soil.

Restoring the healthy riparian sedge meadows that stabilize streambanks can be difficult. Along narrow, deeply incised streams, the water table often drops too low to permit seedling establishment.

Too often, projects intended to improve stream habitat include planting of willows in healthy riparian sedge meadows. The planted willows usually die because they are poorly adapted for growth in the deep, anoxic soils. Instead, willows should be planted in disturbed soils to which they are adapted and which would benefit from stabilization by their roots.

### Sedges as Forage

Sedges provide nutritious forage for animals capable of digesting cellulose, and can be more nutritious or more easily digested than co-occurring grasses. They are especially important in early spring and late summer because many species green up earlier and remain green later than most grasses. Evergreen species such as *C. geyeri* are important in winter and early spring.

Grazing animals may selectively eat some sedge species, passing by other potential forage plants to do so. For example, cattle select *C. rossii* in sagebrush steppe in spring. Some bison herds consume mainly *C. atherodes* at all seasons of the year. However, many other sedges are tough or have serrate leaf margins that reduce palatability, and are consumed only when preferred forage plants are gone.

The animals that eat *Carex* are diverse. Caterpillars of some butterfly and moth species do so; most are thought to be generalists, eating several species of graminoids, but some are specialists, feeding only on *Carex*. Arctic geese and swans graze so intensively on *C. aquatilis* that they cause shifts in plant communities and in some places even denude the soil. Waterfowl, gallinaceous birds, and small seed-eating birds all eat *Carex* perigynia, sometimes in large quantities. *Carex* foliage is reported to be a significant food source for many mammals: voles, ground squirrels, prairie dogs, beavers, muskrats, pikas, rabbits, bears, bighorn sheep, bison, musk oxen, moose, elk, reindeer, deer, sheep, cattle, and horses. About a third of the diet of bison in Yellowstone National Park is wetland *Carex*; for some Canadian herds, *C. aquatilis* and *C. utriculata* form more than 70% of the winter diet, and for others, *C. atherodes* is the most important food plant at all seasons. Although bears are not specialized to digest cellulose, they may eat large quantities of wetland sedges, especially in spring, digesting mainly the shoot bases and starchy rhizomes. In uplands, bears eat *C. geyeri* in spring.

Effects of herbivory on *Carex* populations depend in part on the growth form of the plants. Cespitose species are dependent on seeds for reproduction. Therefore, although individual plants survive repeated grazing, populations eventually decline even on moderately grazed ranges. For example, *C. petasata*, a cespitose species of sagebrush steppe, declines with grazing and is considered an indicator of range condition, providing 2–3% cover on ranges in excellent condition, less under pressure of grazing. *Carex rossii* and its close relatives can maintain populations despite moderate grazing in part because many of their

## Carex Ecology

perigynia are produced on basal spikes too short to be eaten by large animals. Cespitose *C. filifolia* is a source of high-protein forage for livestock and wildlife. It increases under moderate grazing that reduces grass competition, but declines under heavy grazing that prevents seed set or favors monocultures of very short, highly grazing-tolerant grass species. In alpine sites, even very light grazing (one day per year) may cause declines in abundance and cover of *C. filifolia*.

Rhizomatous sedges have the potential to persist and spread in the near-absence of seed production, and therefore withstand grazing better than cespitose species. Some are important forage species (e.g., *C. nebrascensis* and *C. utriculata*), at least after their habitat dries seasonally so that animals have easy access to them. Grazing in fall tends to reduce nutrient storage in rhizomes and may cause harm. Grazing in moist springtime conditions may do less direct harm to plants, but must be balanced against trampling damage to soil and rhizomes. Although the rhizomatous *Carex* may persist under moderately heavy grazing, both above-ground and below-ground biomass are reduced, with potentially harmful effects on their wetland ecosystems.

A few short, wiry, rhizomatous sedges increase on degraded ranges where heavy grazing has reduced competition from grasses. For example, *Carex praegracilis* becomes a community dominant on heavily grazed, seasonally wet, alkaline ranges. In the northern Great Plains, *C. duriuscula* and *C. obtusata* respond the same way in heavily grazed prairies. Dominance by these unusually grazing-tolerant species is accompanied by a reduction of plant diversity, particularly of native species.

## Carex and Fire

Effects of fire have been studied in detail in few sedge species. Most *Carex* are well adapted to survive fire because of their growth form and, in many cases, their habitat. *Carex* foliage may burn but the growing points are protected below the soil surface. Survival and cover are greater after low-intensity fires than after high-intensity fires, with the qualification that the intensity that matters is at the rhizome level. Fire that consumes the duff layer can kill upland forest sedges with shallow rhizomes, such as *C. concinnoides*. Many *Carex* have long-lived seed banks that contribute to revegetation after fire and sedges can effectively stabilize soils in burned areas. For example, ten years after the North Waldo Lake fire in the Oregon Cascades, most of the ground cover in uplands is naturally regenerated *C. inops* and *C. rossii* and, in moist drainages, *C. kelloggii*.

Sedges are most susceptible to fire in late spring and early summer, when they are most physiologically active. Fall burns may harm sedges by interrupting storage of nutrients in rhizomes, but some species have responded better to fall burns than spring burns (e.g., *C. duriuscula* in northern prairies). Usually, early

spring fires do the least damage. Some species (e.g., *C. rossii*) seem unaffected by season of the fire. Prescribed burning of wet sedge meadows during drought years should be avoided because the dry litter layer and organic soil can burn, killing sedge rhizomes and roots.


Some upland species, such as *C. deflexa*, *C. geyeri*, *C. inops*, and *C. rossii*, thrive after fire because of reduced competition and increased soil nutrients. Both survival of mature plants and recruitment from the long-lived soil seed bank contribute to these post-fire sedge populations. Germination of *C. rossii* seeds in the seed bank may be aided by exposure to heat from fires.

*Carex geyeri* is an upland community dominant in open conifer forest. Prior to settlement, many of its habitats had a history of frequent, low-intensity fires. Its rhizomes survive such fires well and plants may flower the first year after the fire. In the Blue Mountains of Oregon, seed germination also contributes to an overall increase in *C. geyeri* cover after a fire. At some drier sites, competing bunchgrasses may benefit from low-intensity fire more than *C. geyeri*, leading to a reduction in *C. geyeri* cover. Rhizome survival and recruitment from the seed bank both decrease after high-intensity fires, delaying a return to pre-fire cover for seven years or more.

*Carex filifolia* is an important “bunchgrass” sedge that lives at the dry edge of the ecological niches *Carex* can occupy, in short-grass prairie and sagebrush steppe. It is especially vulnerable if it experiences two or more stressors such as fire, grazing pressure, competition, low precipitation, and poor soil. Severe fires may reduce cover and productivity for fifteen years or more. The effects of low- and moderate-intensity fires vary. If adequate moisture is available, plants recover in two to four years, but low rainfall delays recovery. *Carex filifolia* should not be burned during drought. Recovery after fire depends much more on rhizome survival than recruitment from the seed bank, though both occur. Usually, flowering is reduced the year after a fire, but lower competition and post-fire influx of nutrients may result in unusually high seed production if enough moisture is available. Fire may benefit *C. filifolia* by setting back competing shrubs or grasses, but it may also encourage weed invasion. Resting burned sites from grazing for at least one year facilitates recovery.

Wet sedge meadows may dry out enough to burn in summer or fall, but fire has little effect on deep-rooted wetland species such as *C. aquatilis*. This species may produce more foliage the year after a fire if competing vegetation is killed and additional nutrients are released into the soil. However, prescribed burns have damaged *C. aquatilis* populations already stressed by grazing. If prescribed burns are used as a management tool, pairing them with a rest from grazing will allow root reserves to build up. Ideally, the rest would extend from the season before the burn to two or three years after.

# Sedge Parts






## *Carex abrupta* Mack.

Common name: Abrupt-beak Sedge

Section: *Ovales*

Key: J



### Key features:

- Cespitose, with gynecandrous spikes & winged perigynia
- Perigynia brown and strongly veined on both faces
- Inflorescence a single dense head
- Meadows at moderate to high elevations in mountains

**Description:** **Habit:** Cespitose. **Culms:** 18-70 cm tall. **Leaves:** 1.5-3.7 (-4.9) mm wide. **Inflorescences:** Dense to slightly elongated heads 1.2-2.2 cm long, (6-) 9-20 mm wide. Spikes gynecandrous. ♀ **scales:** Reddish brown with lighter or green midstripe, 2.4-3.9 mm long, shorter than and as wide as or narrower than the perigynia. **Perigynia:** Reddish brown or dark brown, lacking metallic sheen, typically elliptic to lance-ovate, sometimes lanceolate and nearly wingless, usually planoconvex, (2.9-) 3.6-5.4 mm long, 1-2.1 mm wide, with 3+ conspicuous veins that reach or exceed the top of the achene on each face of the perigynium, with wing usually 0.2-0.3 mm wide and sometimes curved toward the ventral surface, making the perigynium boat-shaped. Beak tips coppery or reddish brown, usually unwinged, brown, and parallel-sided for the distal 0.5-0.8 mm; (1.6-) 2-2.3 mm from top of achene to tip of beak. Stigmas 2. **Achenes:** Lenticular, 1.2-1.8 mm long, 0.7-1.1 mm wide, (0.3-) 0.4-0.5 mm thick.

**Habitat and Distribution:** Moist meadows and streambanks at moderate to high elevations in mts; Siskiyou, Cascades, Wallowas, and Steens Mt., OR, and north Cascades, WA. OR to ID, S to CA and NV.

**Identification Tips:** *Carex abrupta* resembles *C. pachystachya* but has more strongly veined perigynia, especially on the ventral surface, and perigynium wings that may bend in toward the ventral surface more and narrow irregularly toward the beak. *Carex pachystachya* perigynia have a metallic sheen and usually lack veins on the ventral surface; if ventral veins are present, they end at or below top of the achene, or rarely 1 or 2 of them may extend onto the beak. *Carex microptera* has perigynia flat except over the achene, proportionately smaller achenes, and a green and black inflorescence. On Steens Mt., both *C. haydeniana* and *C. microptera* have unusually prominent ventral veins, similar to those of *C. abrupta*, but have either longer or more clearly defined beaks.

**Comments:** Many SW OR plants that have been identified as *C. abrupta* are odd. They may represent one of the NW CA species such as *C. subbracteata*, but more research is needed.

*Carex abrupta*



top left: pistillate scales, perigynia  
top right: inflorescence  
bottom: habit




*Carex albonigra* Mack.

Common name: Black-and-White Sedge

Section: *Ramosae*      Key: F

**Key features:**

- Inflorescence black or dark brown, the ♀ scales with hyaline margins
- Lower spikes on short peduncles
- Rocky alpine ridges and meadows



**Description:** **Habit:** Cespitose. **Culms:** 10-40 cm tall. **Leaves:** 2.5-5 mm wide. **Inflorescences:** 2-4 spikes, each 0.8-2 cm long, the lateral spikes ♀ and erect on short peduncles, the terminal spike gynecandrous. **♀ scales:** Brown to blackish, about equal to the perigynium, tip acute, the midrib dark like the rest of the scale, margins hyaline, contrasting with the rest of the inflorescence. **Perigynia:** Ovate, more or less flattened except over the achene, dark brown, 3-3.5 mm long, 2-2.5 mm wide, papillose, with a small beak 0.3-0.4 mm long. Stigmas 3. **Achenes:** Trigonous, nearly filling bodies of perigynia.

**Habitat and Distribution:** Dry, windblown alpine and subalpine rocky slopes and meadows where little snow accumulates in winter; mts of N WA. AK to NT, S to WA and NM.

**Identification Tips:** *Carex albonigra* inflorescences are dark except for pale margins of ♀ scales. Spikes are on stalks so short that they can superficially seem sessile, like the sessile spikes of *C. media* and *C. pelocarpa*. *Carex media* has more contrast between dark scales and greenish or lighter brown perigynia. *Carex pelocarpa* has a shorter, more uniformly dark inflorescence. *Carex atosquama* has obtuse ♀ scales that are shorter than the perigynia and less conspicuously pale-margined, and its perigynia are a golden brown except at the dark tips.

**Comments:** Plants previously called *C. albonigra* in the White Mountains and central Sierra Nevada of California have recently been described as a distinct species, *C. orestera* Zika.

*Carex albonigra*



top left: perigynia

top right: inflorescence

center left: inflorescence

bottom left and right:

habit





*Field Guide to the Sedges of the Pacific Northwest* is an illustrated guide to all 169 species, subspecies, and varieties in the genus *Carex* that grow in the wild in Oregon and Washington. Most of these species are found throughout the Pacific Northwest and California. This updated second edition includes eight additional species documented in the region since the guide was first published, along with an improved identification key, updated nomenclature and taxonomy, revised range maps, and better illustrations.

Sedges can be difficult to identify, with differences between species based on small, technical characters. This comprehensive guide contains identification keys, descriptions, more than 650 color photographs, and distribution maps for each species, providing users with helpful tools and tips for identifying the plants in this challenging group. Information about sedge ecology, habitat, management and restoration, ethnobotanical uses, and propagation enhances the guide's utility.

The *Field Guide* provides an invaluable resource for botanists, land managers, restoration ecologists, and plant enthusiasts. And, as the genus *Carex* becomes increasingly important amongst landscapers, nurseries, and gardeners, the guide will serve as a handy tool for choosing Northwest natives for the garden.

The *Carex* Working Group is made up of Oregon botanists fascinated by sedges, grasses, willows, and other difficult-to-identify plant groups. In addition to writing about and photographing sedges, CWG teaches plant identification workshops; completes botanical inventories, rare plant surveys, and natural resource planning projects; and conducts taxonomic research on plants of the western United States.

Oregon State University Press

Cover design by David Drummond  
Front cover: *Carex comosa* on the Klamath River,  
Klamath County, Oregon  
Back cover: *Carex capitata* and *Carex unilateralis*  
Photographs by Bruce Newhouse

ISBN 978-0-87071-728-4

