

Fossilium Catalogus

II: Plantae.

Editus a

W. Jongmans.

Pars 13:

H. N. Dixon

Muscineae.



W. Junk

Berlin W 15

5. X. 1927

Ds

115

Do 115 N)

Fossilium Catalogus


II: Plantae.

Editus a

W. Jongmans.

Pars 13:

H. N. Dixon



~~Wpisano do inwentarza
ZAKŁADU GEOLOGII~~

W. Junk
Berlin W 15

Dział B Nr 65
Data 9.10. 1946.

1927



3644



1927.574



Introductory Remarks.

By far the largest number of entries in the following list will be found to belong to quite recent geological periods; their claim to be considered "fossil" in the true sense being somewhat dubious. The number of undoubted mosses known from rocks earlier than the Tertiary is remarkably small. This is surprising in view of the presence in Carboniferous rocks of Hepaticae, very similar to existing species, having been clearly established (Walton, Carboniferous Bryophyta, I, in Ann. of Bot., XXXIX, p. 563, t. XIII).

One would have supposed that mosses with their firmer, often almost ligneous tissues would have stood a better chance of preservation than the more delicate stems of the foliose Hepaticae; but this does not appear to have been the case. Certain scale-like structures from the coal measures have been supposed to be moss leaves, but apart from their unusual dimensions the cell structure does not appear to be bryophytic, and they resemble so closely the stipes scales of some modern ferns that the conclusion seems to me inevitable that they must be considered to represent some similar organ of the Carboniferous Fern-allies. *Muscites polytrichaceus* Ren. and Zeiller, however, has strong claims to be considered a true Carboniferous moss, and the same may be said of *Muscites Bertrandii* Lignier.

Records of fossil or prehistoric mosses, scattered throughout botanical and geological literature, often unindexed, are exceedingly hard to trace, and it is fairly certain that I have overlooked numerous sources of information. The list of works given in the Bibliography will however show what papers have actually been consulted, and make it possible for such gaps as exist to be readily filled up.

It may be of some use to make reference to certain works which are of general importance or which have critical remarks on species already recorded. Schimper's *Traité de Paléontologie végétale* (1869) gives a valuable résumé of most of the records of fossil mosses up to the time of its publication. Neuweiler, *zur Interglazialflora der Schweizerischen Schieferkohlen* (1905), has some useful critical notes on the fossil *Hypna* described by Schimper in the above work; and Caspary, in Klebs, *Flora d. Bernsteins* (1906), does the same for many of the records of mosses by Goeppert in his great work on the Amber Flora. Some valuable notes are to be found in Britton and Hollick, *American fossil mosses, etc.* (1907); and Weyland, *Beitr. zur Kenntnis fossiler Moose* (1925) clears up many points left doubtful by Engelhardt und Kinkelin in their *Oberpliocäne Flora und Fauna des Untermaintals* (1908).

I have endeavoured to indicate in general terms the stratigraphical origin of the recorded mosses as far as possible, where such origin is mentioned in the title or elsewhere in the paper. For the most part I have used the terms employed by the au-

thors. These frequently employ different terms for the same geological epoch. Such terms as Pleistocene, Quaternary, Diluvial, Recent, etc., therefore, must not be taken as necessarily denoting different periods.

The taxonomic arrangement of the following catalogue is (with one or two very slight departures) that of Brotherus in the *Musci*, Engler & Prantl, *Pflanzenfamilien*, Ed. I, 1898—1909.

The species under each genus are arranged alphabetically; the entries under each species chronologically. An index of synonyms is given at the end.

My thanks are due for assistance in the compilation of this list in many directions, notably to the Botanical and Geological Departments of the British Museum, especially to Mr. W. N. Edwards of the Geological Department. Also to Mrs. Britton of the New York Botanical Garden, Mons. J. Cardot of Paris, Prof. I. Györffy of Szeged, Hungary, Herr Loeske of Berlin, Dr. Docturowsky of Moscow, and Dr. Jongmans, for information about and transcripts from papers published in their respective countries to which I was unable to obtain access.

17, St. Matthew's Parade,

Northampton, England.

H. N. Dixon.

Bibliography.

- A dám, Boros, Magyar Botanikai Lapok, Budapest; 1922, p. 71.
— Two fossil species of Mosses from the Diluvial Lime Tufa of Hungary. — Bryologist, XXVIII, p. 29; 1925.
- Andersson, Gunnar, Ueber das Fossile Vorkommen der *Brassenia purpurea* Mich. in Russland und Dänemark. — Bihang Svensk. Vet.-Akad. Handl., Bd. 22, Afd. III, No. 1; 1896.
- Studier öfver Finlands Torfmossar och fossila Kvartärflora. — Bull. de la Commission géologique de Finlande, Tome 2, No. 8. Helsingfors; 1898.
- Augé, L., Note sur la végétation des tufs quaternaires de Piécourt (Gard). — Bull. Soc. Etudes Sc. nat. Nîmes, vol. XXXV, p. 65; 1908.
- Baker, Frank C., The life of the Pleistocene or glacial period. — Univ. of Illinois Bull. 17; 500 pp. 56 plates; 1920.
- Baumberger, E., Die diluvialen Schieferkohlen der Schweiz. — Beitr. Geol. Schweiz. geotechn. Ser., Lief. 8; 1923. (cf. Gams, H., 1923.)
- Beck, R., und Weber, C. A., Ueber ein Torflager im älteren Diluvium des sächsischen Erzgebirges. — Zeitschr. d. D. geolog. Ges., Bd. XLIX, p. 662; Berlin; 1897.
- Bell, A. M., Implementiferous Sections at Wolvercote (Oxfordshire). — Quart. Journ. Geol. Soc., IX, p. 120; 1904.
- Behrendt, G. C. (& Goeppert), Die im Bernstein befindlichen organ. Reste der Vorwelt; Bd. 1, Abt. 1; Berlin; 1845.
- Bertrand, C. E., Description d'un échantillon de charbon papyracé ou Papierkohle trouvé à Prisches en 1859. — Ann. Soc. Géol. du Nord, XXVIII, p. 171; 1899.
- Beyle, M., Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Erster Teil. — Jahrb. der Hamburg. wiss. Anst. XXX, p. 83; 6 Beih.; 1913.
- Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. — Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 33; 1918.
- Ibid. Dritter Teil. — Mitteil. aus dem Mineralog.-Geologisch Staatsinstitut, Heft VI. Hamburg; 1924.
- Boulay, N., Flore pliocène du Mont-Dore. Paris; 1892.
- Braun, Al., Verzeichn. d. fossil. Pflanzen von Oeningen. — Stizenberger, Uebersicht d. Verstein. Badens. Freiburg; 1851.
- Britton, Elizabeth G., A new Tertiary Fossil Moss. — Bull. Torr. Bot. Club, XXVI: p. 79. New York; 1899.
- Britton, Elizabeth G., and Hollick, Arthur, American fossil mosses, with description of a new species from Florissant, Colorado. — Bull. Torr. Bot. Club, XXXIV: p. 139; 1907.
- A new American fossil moss. — Bull. Torr. Bot. Club, XLII: p. 9; 1915.

- Bronniart, Ad., Sur la Classification . . . des Végétaux Fossiles en Général. Chap. II. — Mém. du Mus. d'Histoire nat., viii; 1822.
 — Histoire des végétaux fossiles. — Paris, 1828—1838.
- Brotherus, V. F., Fossile Bryales. — Engler & Prantl, Pflanzenfam. Musci, II, p. 1239; 1909. Ibid. Ed. II; II, p. 523; 1925.
- Burrell, W. H., Pennine Peat. — The Naturalist, 1924, p. 145.
- Camus, F., Sur les mousses trouvées dans le contenu de l'estomac d'un mammouth. — Compt. Rend. Acad. Sci. Paris, CLX, p. 842; 1915.
- Candler, Chas. Observations on some Undescribed Lacustrine Deposits at Saint Cross, South Elmham, in Suffolk. — Quart. Journ. Geol. Soc., XLV, p. 504; 1889.
- Caspary, Robert. See Klebs, 1906.
- Clerici, E. Contribuzione alla conoscenza dei capisaldi per la geologia dei dintorni di Roma. — Rend. Lincei, Roma, X, p. 77; 1901.
- Cockerell, T. D. A. The fossil flora of Florissant, Colorado. — Bull. amer. Mus. nat. Hist. XXIV, p. 71; 1908.
 — A Redwood described as a Moss. — Torreya, VII: p. 203; 1907.
- Dawson, Sir Wm.: and Prof. D. P. Penhalilow. On the Pleistocene Flora of Canada. — Bull. Geol. Soc. Amer. I: pp. 31, 332; 1890. (See also Trans. Roy. Soc. Canada, VIII, 1890).
- Dixon, H. N. Plant remains in peat. — Journ. of Bot., XXXIII, p. 216; 1895.
 — See Lewis, F. J.; 1907, 1908 and 1911.
 — Some "Neolithic" Moss remains from Fort William. — Ann. Scottish Nat. Hist., 1910, p. 103.
 — See Warren, S. H.; 1912.
 — Note on a moss in amber. — Journ. of Bot., LX, p. 149; 1922.
 — Mnium antiquorum. — Bryologist, XIX, p. 51; 1916.
 — See Reid, Clement; 1915.
 — Moss Remains in Russian Peat. — Journ. of Bot., LXIII, p. 370; 1925. (& V. S. Docturowsky).
- Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. — Reports on Peat, 4; 1918—20.
 (1.) Les mousses participant à la formation des tourbières du Polessie (Gouv. de Moskva, etc.). — Rev. d'industr. de la tourbe, 3—4. Petrograd; 1916.
 (2) Ueber die Stratigraphie der russ. Torfmoore. — Geol. Foren. i Stockholm Foerh., XLVII, 1—2; 1925.
- Docturowsky, V. S. See Dixon, H. N. (1925).
- Douin, R. Les mousses et les hépatiques fossiles des tuis du Lautaret (Hautes Alpes). — Revue Génér. de Bot., XXXV, p. 113; 1926.
- Engel, Th. Geognostischer Wegweiser durch Württemberg. 2^o Aufl.; 1896. 3^o Aufl.; 1908.
- Engelhardt, H., and Kinkel, F. Oberpliocäne Flora und Fauna des Untermaintals, insbesondere des Frankfurter Klärbeckens. — Abhandl. Senckenb. Naturf. Ges., XXIX, p. 151; 1908.
- (N. B. The references to the Figures on Plate 22 are given incorrectly in the text, and partially in the Reference to Plates. The references in the Erläuterung der Abbildung are correct with the exception of Figures 22, 23, which certainly

- do not represent the same plant as Figs. 18—21, or indeed any species of *Neckera*.)
- Ettingshausen, C. von. Die Tertiärflora von Haering. — Abhandl. d. geol. Reichsanstalt, Bd. II; Wien; 1852.
- Beitr. zur Kenntniss der fossilen Flora von Parschlug in Steiermark. I. Theil. — Denkschr. d. K. Akad. Wiss. Wien, Bd. XXXVIII, Abth. I, p. 81; 1877.
- Farneti, R. Ricerche di briologia palaeontologica nelle torbe del sottosuolo pavese appartenenti al periodo glaciale. — Atti dell'Istituto Botanico della Reale Università di Pavia. Ser. II, vol. V; Milano; 1896.
- Fischer, Ed. Nachtrag zum Pflanzenverzeichniss des Interglacials von Pianico-Sellere. — Neues Jahrb. f. Mineralogie, Bd. II, p. 105; 1897.
- Fleischer, Max. Natürliches System der Laubmose. — Hedwigia, LXI, p. 390; 1920.
- Gadecau, E. Les Forêts submergées de Belle-Ile-en-mer. — Bull. Biologique de la France, etc., LIII, p. 276; (1919),
- Gams, H. See Heim, Arnold, 1918.
- Interglacial bryophytes, in Baumberger, etc. Die Schieferkohlen der Schweiz, B. 2, Geolog. Karte d. Schweiz, geotechn. Ser., Lief. 8; 1923.
- Geheeb, Adalbert. Ueber ein fossiles Laubmoos aus der Umgebung von Fulda. — Beihefte zum Bot. Centralbl., herausgeg. von Uhlwurm und Kohl, Bd. X, Heft 3, p. 125; Kassel; 1901.
- Geinitz, C., und Weber, C. A. Ueber ein Moostorflager der post-glacialen Foehrenzeit am Seestrande der Rostocker Heide. — Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 1; 1904.
- Gepp, A. Fossil plant-remains in peat. — Journ. of Bot., XXXIII, p. 180; 1895.
- Goeppert, H. R. See Berendt, G. C.; 1847.
- Ueber die Bernsteinflora. — Monatsber. d. Berliner Akad., 1853, p. 450. (Issued also as a reprint with entirely different pagination).
- Goodchild, H. H. See Whitehead, H.; 1912.
- Grout, A. J. A Fossil Camptothecium. — Bryologist, XX, p. 9; 1917.
- Halle, T. G. A fossil Sporogonium from the Lower Devonian of Röragen in Norway. — Bot. Not., 1916, p. 79.
- Lower Devonian plants from Röragen in Norway. — Kgl. Svensk. Vet.-Akad. Handl., LVII, No. 1; 1916.
- Hartmann, F. Die Fossile Flora von Ingaramsdorf. Inaug. Dissert. Breslau; 1907.
- Hartz, N. Bidrag til Danmarks senglaciale Flora og Fauna. Avec Résumé. — Danmarks geologiske Undersøgelse, II Raekke, No. 11; Kjobenhavn; 1902.
- Bidrag til Danmarks tertiaere og diluviale Flora. — Danmarks geologiske Undersøgelse, II Raekke, No. 20; Kjobenhavn; 1909.
- Hawkesworth, E. A pre-glacial Lake-bed near Northallerton. — Naturalist, 1912, p. 204.
- Hayden, F. V. Tertiary Flora of United States. — Rep. of U. S. A. Geol. Survey, VII; 1878. [Mosses by Lesquereux]. Ibid., VIII; 1883.
- Heer, Osw. Flora tertiaria Helvetiae; Zürich, 1855—1859.
- Die Urwelt der Schweiz. 1. Aufl. Zürich, 1865.

- Heim, Arnold & Gams, H. Interglaziale Bildungen bei Wildhaus (Kant. St. Gallen). — *Vierteljahrsschr. Naturf. Ges. Zürich*, LXIII: 19; 1918.
- Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. — *Danmarks geologiske Undersøgelse*, II Raekke, No. 25, p. 101; Kjobenhavn; 1910.
- Hollick, A. See Britton, E. G.; 1907; and 1915.
- Hollo, Dr. L., in Kecszemét Multja ás Jelene; Keczsemét; 1896.
- Holst, N. O. Preglaciala Dryasförande inneslutningar i den undre moränen vid Bjäresjöholms tegelbruk nära Ystad. — *Geol. Fören. Förh.*, XXIX, p. 228; 1907.
— Efterskörd från de senglaciala Lagren vid Toppeladugård. — *Sver. Geologiska Undersökning Arsbok*, II, No. 2; 1908.
- Holzinger, J. M. Some Fossil Mosses. — *Bryologist*, VI, 93; 1903.
- Jessen, Knud, og R. Rasmussen. Et Profil Gennem en Tørvemose paa Faerørne [Section of a Bog in the Faeroe Is.]. — *Danmarks geologiske Undersøgelse*, IV Raekke, No. 13; 1922.
- Jongmans, W. J., in Mededeelingen van de Ryks Opsporing van Delfstoffen, No. 3. 's Gravenhage; 1911.
- Kinkelin, F. See Engelhardt, H.; 1908.
- Kirchner, C. G. Walter. Contribution to the fossil flora of Florissant, Colorado. — *Trans. Acad. of Science of St. Louis*, VIII, p. 161; 1898.
- Klähn, Hans. Die Petrogenese der Kalktuffe nebst einigen sich daraus ergebenden geologischen Problemen. — *Geol. Arch.* 2: 298; 1923.
- Klebs, (and Caspary). Flora d. Bernsteins. — *Abhandl. K. preuss. Geol. Landesanstalt*, N. F., Heft 4; 1906. (With Atlas).
- Knowlton, F. H. Fossil Mosses. — *Plant World*, V, p. 243; 1902.
— A Review of the fossil plants in the U. S. National Museum from the Florissant lake beds at Florissant, Colorado, with description of new species and list of type specimens. — *Proc. United States Nat. Mus.* 51: 241; 1916.
- Koert, W., und Weber, C. Ueber ein neues interglaciales Torflager. — *Jahrb. d. Königl. preuss. geologisch. Landesanstalt* für 1899, p. 185. Berlin; 1900. [Mosses by Weber].
- Laurent, L. Flore plaisancienne des argiles cinéritiques de Mac (Cantal), avec une introduction Géol. — *Annales du Musée d'Histoire naturelle de Marseille*, — *Géologie*, Tome XII, p. 1; 1908.
— Flore pliocène des cinérites du Pas de la Mougado et de Saint-Vincent-la-Sabie (Cantal) — *Ann. du Mus. Hist. nat. Marseille, Géol. tom. IX*; 1904—1905. (Only a general reference to *Muscites*).
- Lesquereux, Léo. See Hayden, F. V.; 1878.
- Lewis, Francis J. Summary of Progress for 1907. — *Mem. Geol. Survey*. [Mosses by Dixon].
— The Plant Remains in the Scottish Peat Mosses. Part III. — The Scottish Highlands and the Shetland Islands. — *Trans. Roy. Soc. Edinb.* XLVI, p. 33; 1908. [Mosses by Dixon].
— Ibid, Part IV. — Op. cit. XLVII, p. 793; 1911. [Mosses by Dixon].
- Lignier, O. Sur une mousse houillère à structure conservée. — *Bull. Soc. Linn. Normandie*, 6 Sér., VII, p. 128; 1914.

- Lindberg, Harald. Finska torfmossar. — Finska Mooskulturför. Arsbok, Helsingfors; H. 2, p. 1—73; 1900.
[I have been unable to consult this paper.]
- Ludwig, R. Fossile Pflanzen aus der ältesten Abtheil. der Rheinisch-Wetterauer Tertiär-Formation. — Palaeontographica, VIII, pp. 39—181; 1859.
- Maffei, Luigi. Contributo allo studio della flora fossile del deposito lacustre de Pianico. — Atti Istit. Bot. Univ. Pavia, Ser. III, 1/2: p. 47; 1924.
- Marty, P. Flore miocène de Joursac (Cantal). — Rev. Haute Auvergne, Paris; 1903.
- Müller, G. und Weber, C. A. Ueber eine fröhdiuviale und vorglaziale Flora bei Lüneburg. — Abhandl. Kgl. preuss. geologisch. Landesanstalt und Bergakademie; Heft 40; Berlin; 1904. [Palaeontologischer Teil by Weber].
- Neuweiler, E. Die prähistorischen Pflanzenreste Mitteleuropas mit besonderer Berücksichtigung der schweizerischen Funde. — Vierteljahrsschr. Naturforsch. Ges. Zürich, L. Heft 6, p. 23; 1905.
- Zur Interglazialflora der schweizerischen Schieferkohlen. — Ber. d. schweizerisch. bot. Ges., Heft XV; 1905. Beilage. Neunter Ber. d. zürcherisch. bot. Ges., p. 93.
- Nørregaard, E. M. Et senglacialt, opfylt Vandøb fra Dejbjerg Bakker. — Meddel. fra Dansk geologisk Forening, No. 15, p. 317; 1909.
- Pax, F. Grundzüge der Pflanzenverbreitung in den Karpathen, II. Leipzig; 1908.
- Pelourde, F. Paléontologie végétale. Cryptogames cellulaires et crypt. vasculaires. — Paris, 1914.
- Penhallow, D. P. See Dawson, Sir Wm.; 1890.
- Canadian Pleistocene Flora of the Don Valley. — Report of 70th Meeting of Brit. Assoc., Bradford, Sept. 1900. Pt. II, p. 334.
- Report on Tertiary plants of British Columbia collected by Mr. Lawrence M. Lambe in 1906, together with a discussion of previously recorded Tertiary floras. — Canada Dept. of mines, Geol. Survey branch, No. 1013; 1908.
- Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone. — Zeitschr. f. Naturwiss., Bd. LXXVI, p. 161; Stuttgart; 1903.
- Rasmussen, R. See Jessen, Knud.; 1922.
- Reid, C., and Ridley, H. N. Fossil Arctic Plants from the Lacustrine Deposit at Hoxne, in Suffolk. — Geol. Mag., Decade III, Vol. V, p. 441; 1888.
- East Norfolk Geology. — Trans. Norf. & Norw. Naturalists' Soc., VII, p. 290; 1904.
- See Whitehead, H.; 1912.
- and E. M. The Pliocene Floras of the Dutch Prussian Border. — Med. van de Rijksopspor. van Delfstoffen, No. 6; 1915. [Mosses by Dixon].
- Reid, E. M. Two preglacial Floras from Castle Eden. — Quart. Journ. of Geol. Soc., LXXVI, p. 104; 1920.
- Renault, B. and Zeiller, R. Sur des Mousses de l'époque houillère. — Compt. Rend. Acad. Sci. Par., C, p. 660; 1885.
- Flore fossile du terrain houiller de Commentry. — Bull. Soc. Industr. Min.; Saint-Etienne; 1888.
- Rudolph, K. Untersuchungen über den Aufbau böhmischer Moore. — Abhandl. k. k. zoolog.-bot. Gesellsch. Wien, IX, p. 4; 1917.

- Saporta, le comte Gaston de. *Etudes sur la végétation du Sud-Est de la France*. — *Annales des Sci. nat. Par.*, 4^e sér. XVII, p. 191; 1862, do., XIX, p. 5; 1863. 5^e sér. III, p. 5; 1865. do., IV, p. 5; 1865. (Published later as a separate work, in 2 vols., with fresh pagination; cited in list below as *Etudes*, I, p. 24, etc.)
- Schilbersky, K., in *Kecszemét Multja ás Jelene*; Kecszemét; 1896.
- in *Term. tud. Kòzlöny*, XXX, p. 436; Budapest; 1898.
 - *Ein Laubmoos aus dem Pleistocän von Kecszemét*. — *Mathem. Termtud. Ertesítő*, Bd. XXX, p. 632; 1912. (Ungarisch.)
- Schimpfer, W. P. *Traité de Paléontologie Végétale*. I. Paris, 1869.
- Schuster, J. *Paläobotanische Notizen aus Bayern*. — *Ber. bayer. bot. Gesellsch.*, XII, I, p. 44; 1909;
- Seward, A. C. *Fossil Plants*. Vol. I. Cambridge; 1898.
- Sismonda, E. *Materiaux pour servir à la Paléontologie des terres tertiaires du Piémont*; 1865.
- Stark, P. *Beitr. zur Kenntniss der eiszeitlichen Flora und Fauna Badens*. — *Ber. Naturforsch. Gesellsch. Freiburg i. Baden*, XIX, pp. 153—272; 1912. (Separately paged, 1—115).
- Stoller, J. *Beitr. zur Kenntnis der diluvialen Flora (besonders Phanerogamen) Norddeutschlands*. — *Jahrb. d. Königl. preuss. geologisch. Landesanstalt*, XXIX, p. 102; 1908.
- *Die Pflanzenreste des altdiluvialen Torflagers in den Stuttgarter Anlagen*. — *Mitteil. der geofig. Abt. des K. württ. stat. Landesamts*, No. 6. Beilage zu *Jahresh. des Vereins für vaterl. Naturkunde in Württemberg*, Jahrg. LXV, 73; 1909.
 - *Die Flora der jungglazialen Ablagerungen Ostpreussens*. — *Jahrb. d. Königl. preuss. geologisch. Landesanstalt*, XXXI, T. II, H. 1, p. 120; 1911.
 - *Beitr. zur Kenntnis der diluvialen Flora (besonders Phanerogamen) Norddeutschlands*. II. Lauenburg a. Elbe (Kuhgrund). — *Jahrb. d. Königl. preuss. geologisch. Landesanstalt*, XXXII, T. 1, H. 1, p. 109; 1911.
- Sukačev, V. N. *Sur la trouvaille de la flore arctique fossile sur la rive du fleuve Irtyche près du village Demianskóe, gouv. Tobolsk*. — *Bull. Acad. Impér. Sci. St. Pétersbourg*, 1910, p. 457.
- Szafer, W. *Eine Dryasflora bei Krystynopol in Galizien*. — *Bull. internat. Acad. Sci. Cracovie, Sér. B*, 2, 1912, p. 1105.
- Travis, W. G. Plant remains in peat at Aintree. — *Trans. Liverpool Bot. Soc.*, I; 1909.
- On peaty bands in the Wallasey sandhills. — *Proc. Liverpool Geol. Soc.*, III, p. 207; 1922.
- Travis, C. B. and W. G. On plant remains in post-glacial gravels at Seaforth, Liverpool. — *Lancashire Naturalist*, 1913, p. 49.
- Unger, Fr. *Genera et Species Plantarum Fossilium*; 1850, p. 41.
- *Iconographia plantarum fossilium*. — *Denkschr. d. k. k. Akad. d. Wissenschaft.*; Wien, 1852.
- Vonderau, Joseph. *Pfahlbauten im Fuldathale*. — 1. Ergänzungsheft des Vereins für Naturkunde zu Fulda. Veröffentlichung des Fuldaer Geschichts-Vereins. Fulda, 1899. [I have not been able to see this paper].

- Warren, S. H. A late Glacial stage in the Lea Valley. — Quart. Journ. Geol. Soc., LXVIII, p. 213; 1912.
[Mosses by Dixon].
- Weber, C. See Koert, W., 1900.
- Weber, C. A. See Beck, R.; 1897.
- See Geinitz, O., 1904.
- See Müller, G.; 1904.
- Die Moostorfsschichten im Steilufer der Kurischen Nehrung zwischen Sarkau und Crantz. — Engler's Bot. Jahrb., XLII, p. 38; 1908.
- *Hypnum turgescens* Schimp. nicht auf der Kurischen Nehrung fossil. — Engler's Bot. Jahrb., XLII, p. 239; 1908.
- Die Mammutfloren von Borna. — Abhandl. Nat. Ver. Bremen, Bd. XXIII, Heft 1, p. 1; 1914.
- Die Pflanzenwelt des Rabutzer Beckentons und ihre Entwicklung unter Bezugnahme auf Klima und geologische Vorgänge. — Bot. Jahrb. Syst. LIV, Beibl. Nr. 120, p. 3; 1917.
- Weber, C. Otto. Die Tertiärfloren der niederrhein. Braunkohlenformation. — Palaeontographica, II, p. 115; Cassel; 1852.
- Wessei, P. & Weber, C. O. Neuer Beitrag zur Tertiärfloren der niederrhein. Braunkohlenformation. — Palaeontographica, IV, p. 111; 1856.
- Weyland, H. Beitrag zur Kenntnis fossiler Moose. I. Die Moose der oberpliocänen Flora des Frankfurter Klärbeckens. — Senckenbergiana, Bd. VII, Heft 1/2. Frankfurt a. M.; 1925.
- Whitehead, H., Goodchild, H. H., and Reid, C. Some notes on "moorlog" — a peaty deposit from the Dogger Bank in the North Sea, with report on the Plant remains. — Essex Naturalist, XVI, p. 51; 1912.
- Zeiller, R. See Renault, B. 1885, and 1888.
- Le Progrès de la Paléobotanique de l'ère des Gymnospermes. — Progressus Rei Botanicae, II, p. 179; 1908. "Quant aux Mousses, il n'en a toujours pas été signalé dans les couches secondaires, les Najadite die Rhétien d'Angleterre appartenir aux Lycopodiacées."
- Zmuda, A. J., in Szafer, Ueber eine alt-diluviale Flora in Krystynopol in Wolhynien. — Kosmos, XXXVI, p. 337; 1911.
- Fossile Flora des Krakauer Diluviums. — Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 209; 1914.

Sphagnaceae.

Sphagnum Ehrh.

Sphagnum acutifolium Ehrh.

- 1896 **Sphagnum acutifolium** aggr. Farneti, R. Ricerche di briologia palaeont. [Atti del Istit. Bot. della Reale Univ. di Pavia, Ser. II, vol. V, p. 58.] Recent
- 1904 **Sphagnum acutifolium** aggr. Weber, C. A. in Müller G. und Weber. Ueber eine fröhldiluviale... Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 15.] Preglacial
- 1909 **Sphagnum acutifolium**. Hartz, N. Bidr. til Danmarks tertiaere... Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] Interglacial
- 1909 **Sphagnum** cf. **acutifolium**. Schuster, J. Palaeobot. Not. aus Bayern. [Ber. bayer. bot. Gesellsch., XII, I, p. 58.]
- 1917 **Sphagnum acutifolium** aggr. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons... [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 13.]
- 1918 **Sphagnum** ? **acutifolium**. Beyle, M. Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. [Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 41 ter.] Recent

Sphagnum compactum De Cand.

- 1828—1844 **Sphagnum compactum**. Brongniart, Ad. No locality
Hist. des végétaux fossiles. Paris, p. 96, tab. X. or data
given

Sphagnum cuspidatum Ehrh.

- 1896 **Sphagnum cuspidatum**. Farneti, R. Ricerche di briologia palaeont. [Atti del Istit. Bot. della Reale Univ. di Pavia, Ser. II, vol. V, p. 58.] Recent
- 1904 **Sphagnum cuspidatum** aggr. Weber, C. A. in Müller G. und Weber. Ueber eine fröhldiluviale... Flora bei Lüneburg. [Abh. Kgl. preuss. geologisch. Landesanstalt und Bergakad.; Heft 40, p. 15.] Preglacial
- 1909 **Sphagnum** cf. **cuspidatum**. Schuster, G. Palaeobot. Not. aus Bayern. [Ber. bayer. bot. Gesellsch., XII, I, p. 58.]
- 1914 **Sphagnum cuspidatum**. Weber, C. A. Die Mammutflora von Borna. [Abh. nat. Ver. Bremen, XXIII, Heft 1, p. 14.] Glacial

- 1917 *Sphagnum cuspidatum*. Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Gesellsch. Wien, IX, p. 86.]
- 1924 *Sphagnum cuspidatum*. Burrell, W. H. Pennine Peat. [Naturalist, 1924, p. 148.]

Sphagnum cymbifolium Ehrh.

- 1869 *Sphagnum cymbifolium*. Schimper, W. P. Traité de Paléontologie Végétale. I, p. 253.
- 1895 *Sphagnum cymbifolium*. Gepp, A. Fossil plant-remains in peat. [Journ. of Bot., XXXIII, p. 181.]
- 1897 *Sphagnum cymbifolium* or near it. Beck, R., und Weber, C. A. Ueber ein Torflager.... des sächsisch. Erzgebirges. [Zeitschr. d. Deutsch. geolog. Gesellsch., XLIX, p. 666.]
- 1899 *Sphagnum cymbifolium*. Bertrand, C. E. Descr... de charbon papyracé à Prisches en 1859. Ann. Soc. géol. du Nord, XXVIII, p. 180.]
- 1904 *Sphagnum cymbifolium* aggr. Weber, C. A. in Müller, G. und Weber, Ueber eine fröhdiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geologisch. Landesanstalt und Bergakad., Heft 40, p. 15.]
- 1905 *Sphagnum cymbifolium*. Neuweiler, E. Die prähist. Pflanzenreste Mitteleuropas... [Vierteljahrs. Naturforsch. Ges. Zürich, I, Heft 6, p. 94.]
- 1909 *Sphagnum cymbifolium*. Hartz, N. Bidr. til Danmarks tertiaere.... Flora.... [Danmarks geol. Undersøgelse II Raekke, No. 20, p. 261.]
- 1912 *Sphagnum cf. cymbifolium*. Whitehead, H., Goodchild, H. H., and Reid, C. Some notes on "moorlog" [Essex Naturalist, XVI, p. 56.]
- 1913 *Sphagnum cymbifolium*. Beyle, M. Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Erster Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6 Beih., p. 83.]
- 1917 *Sphagnum palustre* Linn. Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Gesellsch. Wien, IX, p. 86.]
- 1923 *Sphagnum cymbifolium*. Baumberger, E., etc., Die diluv. Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz; geotechn. Ser., Lief. 8, pp. 83, 338.]
- 1924 *Sphagnum cymbifolium*. Burrell, W. H. Pennine Peat. [Naturalist, 1924, p. 148.]

Sphagnum fimbriatum Wils.

- 1908 *Sphagnum fimbriatum* (or teres). Stoller, J. Beitr. zur Kenntn. der diluv. Flora... Norddeutschlands. [Jahrb. d. Kgl. preuss. geolog. Landesanstalt, XXIX, p. 119.]

- 1910 *Sphagnum fimbriatum* (or *Girgensohnii*). Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 103.] Postglacial
- 1912 *Sphagnum fimbriatum*. Stark, P. Beitr. zur Kenntn. der eiszeitl. Fl. und Faun. Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.] Glacial

Sphagnum Girgensohnii Rüss.

cf. *S. fimbriatum* (1910).

Sphagnum imbricatum (Hornsch.) Rüss.

- 1914 *Sphagnum imbricatum*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 15.] Glacial

Sphagnum intermedium Hoffm.

cf. *S. recurvum*.

Sphagnum Lindbergii Schimp.

- 1918—20 ? *Sphagnum Lindbergii*. Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Interglacial

Sphagnum Ludwigi Schimp.

- 1869 *Sphagnum Ludwigi* Schimp., Traité de Paléontologie Végétale, p. 252. (*Gymnostomum ferrugineum* Ludw., Foss. Pflanzen aus der Rheinisch Tertiär-Formation, in Palaeontographica, VIII, p. 165, tab. 63; 1859.) Tertiary

Note. Schimper places it between *S. cymbifolium* and *S. subsecundum*. It is highly improbable that it is distinct from one of the existing species. Its origin is "dans l'hématite brune de Dernbach (Nassau)". Seward (Fossil Plants, I, p. 241) expresses a doubt as to whether the generic position can be satisfactorily determined. Schimper, however, had presumably examined the specimen.

Sphagnum magellanicum Brid.

- 1898 *Sphagnum medium* Limpr. Andersson, Gunnar Stud. öfver Finlands ... Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 138.] Quaternary
- 1904 *Sphagnum medium* Limpr. Weber, C. A. in Müller, G. und Weber, Ueber eine fröh diluviale.... Flora bei Lüneburg. [Abh. Kgl. preuss. geologisch. Landesanstalt und Bergakad., Heft 40, pp. 15, 18.] Preglacial
- 1917 *Sphagnum magellanicum*. Rudolph K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Gesellsch. Wien, IX, p. 86.] Recent

- 1917 *Sphagnum medium* Limpr. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons..... [Bot. Jahrb. Syst. LIV, Beibl. No. 120, p. 14.]

Sphagnum medium Limpr.

cf. *S. magellanicum*.

Sphagnum palustre Linn.

cf. *S. cymbifolium*.

Sphagnum papillosum Lindb.

- 1909 *Sphagnum papillosum*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1914 *Sphagnum papillosum*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 15.]
- 1924 *Sphagnum papillosum*. Burrell, W. H. Pennine Peat. [The Naturalist, 1924, p. 148, tab. X.]

Sphagnum recurvum P. Beauv.

- 1898 *Sphagnum recurvum*. Andersson. Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 138.]
- 1904 *Sphagnum cf. recurvum*. Weber C. A., in Müller G. und Weber, Ueber eine fröhiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geolog. Landesanstalt und Bergakad., Heft 40, p. 15.]
- 1911 *Sphagnum intermedium* Hoffm. Dixon, H. N. in Lewis, F. J., The Plant Remains in the Scottish Peat Mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.]
- 1917 *Sphagnum recurvum* var. *parvifolium* Warnst. (as *S. brevifolium* Röll). Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Gesellsch. Wien, IX, p. 87.]

Sphagnum rubellum Wils.

- 1898 *Sphagnum rubellum*. Andersson. Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 139.]
- 1917 *Sphagnum Wilsoni* Röll. Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 87.]

Sphagnum squarrosum Pers.

- 1896 *Sphagnum squarrosum*. Farneti, R. Ricerche di briologia paleoent. [Atti del Istit. Bot. della Reale Univ. di Pavia, Ser. II, vol. V, p. 57.]

Sphagnum subbicolor Hampe.

- 1923 *Sphagnum subbicolor*. Baumberger, E., etc. Die diluv. Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz; geotechn. Ser., Lief. 8, p. 83.]

Sphagnum subsecundum Nees aggr.

- 1895 *Sphagnum ? subsecundum*. Gepp, A. Fossil plant- Recent sub-
remains in peat. [Journ. of Bot., XXXIII, p. 181.] marine
peat.
- 1908 *Sphagnum subsecundum*. Stoller, J. Beitr. zur Diluvial
Kenntn. der diluv. Flora Norddeutschlands. [Jahrb.
d. Kgl. preuss. geol. Landesanstalt, XXIX, p. 119.]

Sphagnum teres Aongstr.

- 1904 *Sphagnum teres*. Weber, C. A., in Müller, G. Preglacial
und Weber, Ueber eine fröhldiluviale Flora
bei Lüneburg. [Abh. Kgl. preuss. geol. Landes-
anstalt und Bergakad., Heft 40, p. 15 bis.]
- 1908 *Sphagnum teres* (or fimbriatum). Stoller, J. Diluvial
Beitr. zur Kenntn. der diluv. Flora Norddeutsch-
lands. [Jahrb. d. Kgl. preuss. geol. Landesanstalt,
XXIX, p. 119.]
- 1912 *Sphagnum teres*. Stark, P. Beitr. zur Kenntn. Glacial
des eiszeitlich.... Badens. [Ber. naturforsch. Ge-
sellsch. Freiburg i. B., XIX, pp. 88, 116.]
- 1918—20 *Sphagnum teres*. Docturowsky, V. S., and Postglacial
Kudriashchev, V. Results of Russian studies on
peat moors. [Reports on Peat, 4.]
- 1925 *Sphagnum teres*. Dixon, H. N. Moss remains in Interglacial
Russian Peat. (Journ. of Bot., LXIII, p. 370.)

Sphagnum Wilsoni Röll.cf. *S. rubellum*.**Undetermined species of Sphagnum.**

- 1852 *Cryptothecium antediluvianum*. Hueben., MS., nov. gen. et sp., in Weber, C. Otto. Die Tertiär-
flora der niederrhein. Braunkohlenformation. [Pa-
laeontographica, 11, p. 228.] This is considered by
Schimper as without doubt a species of *Sphagnum*.
- 1903 *Sphagnum sp.* Range, Paul. Das Diluv. von Lübeck Diluvial
und seine Dryastone. [Zeitschr. f. Naturwiss.,
LXXVI, pp. 191, 256.]
- 1904 *Sphagnum sp.* Weber, C. A., in Müller, G. und Preglacial
Weber, Ueber eine fröhldiluviale.... Flora bei
Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt
und Bergakad., Heft 40, p. 15.]
- 1909 *Sphagnum sp.* Hartz, N. Bidr. til Danmarks ter- Interglacial
tiaere Flora. [Danmarks geol. Undersøgelse,
II Raekke, No. 20, p. 261.]
- 1910 *Sphagnum sp.* Hesselbo, A. Mosrester fra Diluviet Diluvial
ved Skaerumhede. [Danmarks geol. Undersøgelse,
II Raekke, No. 25, p. 106.]
- 1913 *Sphagnum sp.* Travis, C. B. and W. G. On plant Postglacial
remains in postglacial gravels at Seaforth, Liver-
pool. [Lancashire Naturalist, 1913, p. 20.]
- 1914 *Sphagnum sp.* Zmuda, A. J. Fossile Flora des Diluvial
Krakauer Diluviums. [Bull. int. Acad. Sci. Cra-

- covie d. sc. math. et nat., Sér. B. No. 2 B,
p. 248.]
- 1918 *Sphagnum* sp. Beyle, M. Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. [Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 39.]
- 1918—20 *Sphagnum* sp. (§ Subsecunda). Docturowsky, Interglacial V. S. and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.]
- 1920 *Sphagnum* sp. Baker, Frank C. The Life of the Pleistocene or Glacial period. [Univ. of Illinois, Bull., XVII, pp. 207, 211, 276, 280, 377.]
- 1923 *Sphagnum* sp. Baumberger, E., etc. Die diluv. Quaternary Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz; geotechn. Ser., Lief. 8, p. 338.]

Andreaeaceae.

Andreaea Ehrh.

Andreaea Huntii Limpr.

- 1912 *Andreaea Huntii*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlichen Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.]

Andreaea petrophila Ehrh.

1914. *Andreaea petrophila*. Zmuda, A. J. Fossile Flora Diluvial des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Ser. B., No. 2 B, p. 248.]

Andreaea Rothii Web. & Mohr.

- 1910 *Andreaea Rothii*. Dixon, H. N. Some "Neolithic" Postglacial Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 103.]

Andreaea sp.

cf. *Sporogonites exuberans*.

Dicranaceae.

Ditrichum Timm.

Ditrichum flexicaule (Schleich.) Hampe.

- 1902 *Ditrichum flexicaule*. Hartz, N. Bidr. til Danmarks senglacial Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.]
- 1909 *Ditrichum flexicaule*. Hartz, N. Bidr. til Danmarks tertiaere Flora [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]

- 1910 *Ditrichum flexicaule*. Hesselbo, A. Mosrester fra Diluvial Diluviet ved Skaerumhede. [Danmarks geoL Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Ditrichum flexicaule* var. *densum*. Dixon, H. N. Late glacial in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1918—20 *Ditrichum flexicaule*. Docturowsky, V. S., Postglacial and Kudriiashev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.]

***Ditrichum tortile* (Schrad.) Lindb.**

- 1915 *Ditrichum tortile*. Reid, C. and E. M. The Pliocene Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, p. 53.]

Note. With fragment of seta.

Distichium Bry. eur.

***Distichium capillaceum* (Sw.) Bry. eur.**

- 1900 *Distichium capillaceum*. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley. [Rep. of 70th Meeting of Brit. Assoc., Bradford, Sept., 1900, Pt. II, p. 335.]
- 1903 *Distichium capillaceum*. Range, Paul. Das Diluv. von Lübeck und seine Dryastone. [Zeitschr. f. Naturwiss., LXXVI, p. 253.]
- 1907 *Swartzia montana* Lindb. Host, N. O. Preglacial Dryasförande nära Ystad. [Geol. Fören. Förh., XXIX, p. 232.]
- 1909 *Swartzia montana* Lindb. Hartz, N. Bidr. til Interglacial Danmarks tertiaere Flora. [Danmarks geoL Undersøgelse, II Raekke, No. 20, p. 261.]
- 1910 *Swartzia montana* Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geoL Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Distichium capillaceum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *Distichium capillaceum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.]
- 1914 *Distichium capillaceum*. Weber, C. A. Die Mammutflora von Borna. [Abh. nat. Ver. Bremen, XXIII, Heft I, p. 15.]
- 1920 *Distichium capillaceum*. Baker, Frank C. The Glacial Life of the Pleistocene or Glacial period. [Univ. of Illinois, Bull. XVII, pp. 334, 335, 377.]
- Note. The text reads: —
- Dichelema capillaceum* (see *Distichium capillaceum*)
- Distichium (Dichelema) capillaceum*. As both species occur at the present day in the U. S. A., there is nothing to indicate which name is the species intended.

Ceratodon Brid.**Ceratodon purpureus** (L.) Brid.

- 1898 *Ceratodon purpureus*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.] Quaternary
- 1902 *Ceratodon purpureus*. Hartz, N. Bidrag til Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.] Late glacial
- 1909 *Ceratodon purpureus*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
- 1910 *Ceratodon purpureus*. Hesselbo, A. Mosreste fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 ? *Ceratodon purpureus*. Dixon, H. N. in Warren, S. H., A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *Ceratodon purpureus* var. *paludosus* Warnst. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.] Diluvial

Blindia Bry. eur.**Blindia acuta** (Huds.) Bry. eur.

- 1910 *Blindia acuta* var. *trichodes* Braithw. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 103.] Postglacial
- 1912 *Blindia acuta*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.] Glacial

Dicranella Schimp.**Dicranella cerviculata** (Hedw.) Schimp.

- 1914 *Dicranella cerviculata*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math., et nat., Sér. B, No. 2 B, p. 248.] Diluvial

Dicranella squarrosa (L.) Schimp.

- 1914 *Dicranella squarrosa*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 88.] Diluvial

Dicranella sp.

- 1853 *Dicranum simplex* Goepp. & Menge. Ueber die Amber Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 458.]
Note. Compared by the authors to *Dicranum Schreberi* (*Dicranella*). 2*

- 1907 *Dicranella* sp. Hartmann, F. Die Fossile Flora Diluvial von Ingramsdorf. Inaug. Dissert. Breslau, 1907, p. 14.

Oreoweisia De Not.

Oreoweisia serrulata (Funck) De Not.

- 1912 *Oreoweisia serrulata*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.]

Dichodontium Schimp.

Dichodontium pellucidum (L.) Schimp.

- 1904 *Dichodontium pellucidum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1910 *Dichodontium pellucidum*. Dixon, H. N. Some Postglacial "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 103.]
- 1923 *Dichodontium pellucidum* var. *fagimontanum* Schimp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz; geotechn. Ser., Lief. 8, p. 333.]

Dichodontium subpellucidum (Goepp. & Menge.)

- 1853 *Dicranum subpellucidum* Goepp. & Menge in Goepp., Amber H. R. Ueber die Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 458.]

Dicranoweisia Lindb.

Dicranoweisia crispula (Hedw.) Lindb.

- 1912 *Dicranoweisia crispula*. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitl. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.]

Oncophorus Brid.

Oncophorus virens (Sw.) Brid.

- 1910 *Oncophorus virens*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Oncophorus virens*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.]

Dicranum Hedw.

Dicranum Bergeri Bland.

- 1912 *Dicranum Bergeri* (as *Dicranella Bergeri*, lapsu calami). Stark, P. Beitr. zur Kenntn. der eis-

zeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Dicranum Bonjeani De Not.

- 1909 *Dicranum Bonjeani*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
 1910 *Dicranum Bonjeani*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 103.] Postglacial

Dicranum congestum Brid.

- 1910 *Dicranum congestum*. Hesselbo, A. Mosrester fra Diluvial Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

Dicranum fuscescens Turn.

- 1853 *Dicranum fuscescens*. Goeppert, H. R. Ueber die Amber Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 458.]
 Note. "Von D. fuscescens nicht zu unterscheiden". Goepp. l. c.

Dicranum Mühlenbeckii Bry. eur.

- 1912 *Dicranum Mühlenbeckii* (as *Dicranella lapsu calamii*). Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.] Glacial

Dicranum parietinum.

- 1909 *Dicranum parietinum*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] There is no moss known of this name, and an error is presumable; possibly *Hylocomium parietinum* is intended.

Dicranum Sauteri Bry. eur.

- 1912 *Dicranum Sauteri* (as *Dicranella lapsu calamii*). Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.] Glacial

Dicranum scoparium (L.) Hedw.

- 1909 *Dicranum scoparium*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
 1912 *Dicranum scoparium*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 116.] Glacial

- 1914 *Dicranum scoparium* var. *alpestre* Milde. Zmuda, Diluvial
A. J. Fossile Flora des Krakauer Diluviums. [Bull.
int. Acad. Sci. Cracovie d. sc. math. et nat.,
Sér. B., No. 2 B, p. 248.]

Dicranum Scottianum Turn.

- 1910 *Dicranum Scottianum*. Dixon, H. N. Some "Neo-
lithic" Moss remains from Fort William. [Ann.
Scott. Nat. Hist., 1910, p. 105.] Postglacial

Dicranum strictum Schleich.

- 1911 *Dicranum strictum*? Dixon, H. N. in Lewis, F. J. Iceland
The Plant Remains in the Scottish Peat Mosses. peat; recent
Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.]
Note. Almost certainly this species; but the
specimen was unfortunately lost in the post before
the determination was confirmed.

Dicranum subflagellare Goepp. & Menge.

- 1853 *Dicranum subflagellare*. Goepp. & Menge in Goepp., Amber
H. R. Ueber die Bernsteinflora. [Monatsber. d.
Berliner Akad., 1853, p. 458.]
Note. The authors say "cf. *D. flagellare*".
Syn. *Dicranites subflagellare* Caspary e Klebs,
Flora d. Bernsteins. [Abh. Kgl. preuss. geol.
Landesanst., N. F., Heft 4, p. 55, t. VIII, f. 47;
1903.] There seems no adequate ground for re-
moving this from *Dicranum*.

Dicranum subpellucidum Goepp. & Menge.

cf. *Dichodontium subpellucidum*.

Dicranum subscoparium Goepp. & Menge.

- 1853 *Dicranum subscoparium*. Goepp. & Menge in Goep- Amber
pert, H. R. Ueber die Bernsteinflora. [Monatsber.
d. Berliner Akad., 1853, p. 458.]
Note. The authors say "cf. *D. scoparium*".

Dicranum spp. undetermined.

- 1897 *Dicranum* sp. (Eudicranum). Beck, R. und Weber, Diluvial
C. A. Ueber ein Torflager des sächsisch.
Erzgebirges. [Zeitschr. d. Deutsch. geol. Gesellsch.,
XLIX, p. 666.]
- 1898 *Dicranum* sp. Andersson, Gunnar. Stud. öfver Quaternary
Finlands Kvartärflora. [Bull. de la Com-
mission géol. de Finlande, Tome 2, No. 8, p. 136.]

Dicranites Klebs.

- 1903 *Dicranites* gen. nov. Klebs in Caspary, Fl. d.
Bernsteins [Abh. Kgl. preuss. geol. Landesanst.,
N. F., Heft 4, p. 52; 1903.]

Klebs gives no diagnosis of this genus, but states that it consists of species not certainly referable to any existing species, while clearly belonging to Dicranaceae in the widest sense.

Dicranites Casparyi Klebs.

1903 *Dicranites Casparyi* Klebs, op. et loc. cit., t. VII, Amber f. 42—45.

Compared with such erect-leaved forms as *Dicranum Scottianum*, and *Campylopus flexuosus*. The leaf apex figured, with a few short teeth, does not suggest a species of this alliance at all.

Dicranites obtusifolius Casp. & Klebs.

1903 *Dicranites obtusifolius* Casp. & Klebs, op. cit., Amber p. 54, t. VII, f. 46.

The form of the leaf with toothed sheathing base strongly suggests *Eucladium verticillatum*; but the leaf apex is figured as rough, and described as "an der stumpfen Spitze sägeähnig".

Dicranites subflagellare Caspary.

cf. *Dicranum subflagellare*.

Dicranodontium Bry. eur.

Dicranodontium longirostre (Starke) Bry. eur.

1912 *Dicranodontium longirostre* var. *alpinum* Milde. Glacial Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Fissidentaceae.

Fissidens Hedw.

Fissidens adiantoides (L.) Hedw.

1910 *Schistophyllum adiantoides* Brid. Hesselbo, A. Mos- Quaternary rester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

Fissidens cristatus Wils.

1923 *Fissidens decipiens* De Not. Baumberger, E., etc. Quaternary Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz., geotechn. Ser., Lief. 8, p. 333.]

Fissidens julianus Sav.

1898 *Schistophyllum julianum* (Sav.) Lindb. Andersson, Quaternary Gunnar. Stud. öfver Finlands Kvartärfloren.

[Bull. de la Commission géol. de Finlande, Tome 2,
No. 8, p. 138.]

Fissidens osmundoides (Sw.) Hedw.

- 1910 *Fissidens osmundoides*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]
- 1912 *Fissidens osmundoides*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]
- 1924 *Fissidens osmundoides*. Beyle, M. Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Dritter Teil. [Mitteil. aus dem Mineralog.-Geol. Staatsinstitut, Hamburg, Heft VI, p. 7.]

Fissidens taxifolius (L.) Hedw.

- 1923 *Fissidens taxifolius*. Baumberger, E., etc. Die Quaternary diluv. Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 333.]

Pottiaceae.

Hymenostomum R. Br.

Hymenostomum microstomum (Hedw.) R. Br.

cf. *Muscites confertus*. (*Hymenostomum trichostomum* Goep., Douin, les Mousses et les Hepaticques Fossiles, p. 114, is a lapsus calami for *H. microstomum*).

Weisia Hedw.

Weisia conferta Schimp.

- 1845 *Muscites confertus* Goep. & Ber. Die im Bernstein befindlichen organ. Reste der Vorwelt, I, Abt. 1, p. 112; t. VI, figs. 29—31.
- 1869 *Weisia conferta* Schimper, Traité de Pal. Végét. 1, p. 243.
Note. Caspary writes that he has not examined the specimen, but from the description and figures he is inclined to think it the same thing as *Dicranites Casparyi* Klebs. The relation to *Weisia* seems very doubtful.

Gymnostomum Hedw.

Gymnostomum calcareum Bry. germ.

- 1923 *Gymnostomum calcareum*. Baumberger, E., etc. Die diluv. Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 333.]

Gymnostomum ferrugineum Ludw.

cf. *Sphagnum Ludwigii*.

Eucladium Bry. eur.**Eucladium verticillatum** (L.) Bry. eur.

- 1925 *Eucladium verticillatum*. Adam, Boros. Two fossil Recent tufa species of Mosses from the Diluvial Lime Tufa of Hungary. [Bryologist, XXVIII, p. 32.]

Trichostomum Hedw.**Trichostomum substrictum** Goepp. & Menge.

- 1853 *Trichostomum substrictum* Goepp. & Menge. Ueber Amber die Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 458.]
 Note. Compared by the authors with "*Trich. strictum*"; but, as Schimper remarks, they do not even say whether they refer to *T. strictum* Sw. or *T. strictum* Bry. eur. The affinity is very doubtful.

Trichostomum subpolystichum Goepp. & Menge.

- 1853 *Trichostomum subpolystichum* Goepp. & Menge op. Amber et loc. cit.
 Note. Compared by the authors with "*Trich. polystichum*". No such moss is known. Schimper suggests a slip for *T. polyphyllum* (*Ptychomitrium*); but a moss in amber is not likely to have a very close affinity with a purely rupestral species such as this.

Tortella Limpr.**Tortella fragilis** (Hook. f. & Wils.) Limpr.

- 1909 *Mollia fragilis* Lindb. Hartz, N. Bidrag til Danmarks tertiale Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]

Tortella inclinata (Hedw. f.) Limpr.

- 1904 *Tortella inclinata*. Weber, C. A., in Müller, G. Preglacial und Weber. Ueber eine fröhldiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 15.]

Tortella tortuosa (L.) Limpr.

- | | |
|--|--------------|
| 1902 <i>Mollia tortuosa</i> Lindb. Hartz, N. Bidr. til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.] | Late glacial |
| 1910 <i>Mollia tortuosa</i> Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] | Quaternary |
| 1914 <i>Tortella tortuosa</i> . Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.] | Diluvial |

Didymodon Hedw.**Didymodon alpigena Vent.**

- 1912 *Didymodon alpigena*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Didymodon rigidulus Hedw.

- 1911 *Didymodon rigidulus*. Stoller, J. Beitr. zur Kenntn. Diluvial der diluv. Flora Norddeutschlands. [Jahrb. d. Kgl. preuss. geol. Landesanst., XXIX, p. 123.]

Didymodon rubellus (Hoffm.) Bry. eur.

- 1909 *Barbula rubella* Lindb. Hartz, N. Bidr. til Danmarks tertiaere . . . Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]

- 1910 *Barbula rubella* Lindb. Hesselbo, A. Mosrester Quaternary fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

- 1912 *Barbula rubella* Lindb. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Didymodon tophaceus (Brid.) Jur.

- 1922 *Didymodon tophaceus*. Adám, Boros. Magyar Botanikai Lapok, Budapest, p. 71.

- 1925 *Didymodon tophaceus*. Adám, Boros. Two fossil species of Mosses from the Diluvial Lime Tufa of Hungary. [Bryologist, XXVIII, 29.]

Barbula Hedw.**Barbula fallax Hedw.**

- 1913 *Barbula fallax*. Travis, C. B. and W. G. On plant remains in post-glacial gravels at Seaforth, Liverpool. [Lancashire Naturalist, 1913, p. 50.]

Barbula icmadophila Schimp.

- 1912 *Barbula icmadophila*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Barbula paludosa Schleich.

- 1912 *Barbula paludosa*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Barbula subcanescens Goepp. et Ber.

cf. *Muscites apiculatus* Goepp. et Ber.

Phascum L.**Phascum cuspidatum Goepp.**

- 1853 *Phascum cuspidatum* Goepp., H. R. Ueber die Amber Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 458.]

Note. Schimper considers the identification as doubtful.

Desmatodon Brid.**Desmatodon latifolius** (Hedw.) Bry. eur.

- 1914 *Desmatodon latifolius* var. *muticus* Brid. Weber, Glacial
C. A. Die Mammutfloa von Borna. [Abh. Nat.
Ver. Bremen, XXIII, Heft 1, p. 16.]

Tortula Hedw.**Tortula norvegica** (Web.) Wahl.

- 1903 *Tortula aciphylla* Hartm. Range, Paul. Das Diluv. Diluvial
von Lübeck und seine Dryastone. [Zeitschr. f.
Naturwiss., LXXVI, p. 254.]
- 1910 *Tortula norvegica*. Hesselbo, A. Mosrester fra Quaternary
Diluviet ved Skaerumhede. [Danmarks geol. Un-
dersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Tortula aciphylla* var. *mucronata* Sendtn. Dixon, Late glacial
H. N., in Warren, S. H., A late Glacial stage in
the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII,
p. 231.]
- 1914 *Tortula aciphylla* Hartm. Weber, C. A. Die Mam- Glacial
mutfloa von Borna. [Abh. Nat. Ver. Bremen,
XXIII, Heft 1, p. 16.]

Tortula ruralis (L.) Ehrh.

- 1907 *Tortula ruralis*. Holst. N. O. Preglaciala Dryas- Preglacial
förande nara Ystad. [Geol. Fören. Förh.,
XXIX, p. 232.]
- 1910 *Tortula ruralis*. Hesselbo, A. Mosrester fra Quaternary
Diluviet ved Skaerumhede. [Danmarks geol. Under-
søgelse, II Raekke, No. 25, p. 106.]
- 1914 *Tortula ruralis*. Weber, C. A. Die Mammutfloa Glacial
von Borna. [Abh. Nat. Ver. Bremen, XXIII,
Heft 1, p. 16.]

Tortula species.

- 1914 *Tortula* sp. Weber, C. A. Die Mammutfloa von Glacial
Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1,
p. 16.]

Encalypta Schreb.**Encalypta alpina** Smith.

- 1910 *Leersia alpina* Lindb. Hesselbo, A. Mosrester fra Quaternary
Diluviet ved Skaerumhede. [Danmarks geol. Un-
dersøgelse, II Raekke, No. 25, p. 106.]

Encalypta rhabdocarpa Schwaegr.

- 1902 *Encalypta rhabdocarpa*. Hartz, N. Bidr. til Dan- Late glacial
marks senglaciale Flora og Fauna. [Danmarks
geol. Undersøgelse, II Raekke, No. 11, p. 11.]
- 1909 *Leersia rhabdocarpa* Lindb. Hartz, N. Bidr. til Interglacial
Danmarks tertiaere Flora. [Danmarks geol.
Undersøgelse, II Raekke, No. 20, p. 261.]

- 1910 *Encalypta rhabdocarpa?* Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Encalypta rhabdocarpa*, Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc. LXVIII, p. 231.]
- 1912 *Encalypta rhabdocarpa*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. u. Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Grimmiaceae.

Glyphomitrium Brid.

Glyphomitrium Cockerelleae Britt. et Hollick.

- 1907 *Glyphomitrium Cockerelleae* Britt. & Hollick, American Fossil Mosses, in Bull. Torr. Bot. Cl. 34: 140; tab. 9, figs. 6, 6a. Cum fructu. Tertiary Shales, Florissant, Colorado

Coscinodon Spreng.

Coscinodon humilis Milde.

- 1912 *Coscinodon humilis*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Faun. Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, 89.] Glacial

Grimmia Ehrh.

Grimmia alpicola Sw.

- 1912 *Schistidium alpicola* var. *rivulare* Limpr. Stark, P. Glacial Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Grimmia anodon Bry. eur.

- 1912 *Grimmia anodon*. Stark, P., op. et loc. cit. Glacial

Grimmia apocarpa (L.) Hedw.

- 1910 *Grimmia apocarpa*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumkede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

Grimmia elatior Bruch.

- 1912 *Grimmia elatior*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.] Glacial

Grimmia elongata Kaulf.

- 1912 *Grimmia elongata*. Stark, P., op. et loc. cit. Glacial

Grimmia ericoides Lindb.cf. *Rhacomitrium canescens*.**Grimmia funalis** (Schwaegr.) Schimp.1912 *Grimmia funalis*. Stark, P., op. et loc. cit. Glacial**Grimmia hypnoides** Lindb.cf. *Rhacomitrium lanuginosum*.**Grimmia incurva** Schwaegr.1912 *Grimmia incurva*. Stark, P., op. et loc. cit. Glacial**Grimmia subelongata** Goepp. et Menge.1853 *Grimmia subelongata* Goepp. & Menge in Goepp., Amber H. R., Ueber die Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 459.]

Note. Compared with *G. elongata* Kaulf. Schimper rightly remarks that it is very doubtful if any species allied to *G. elongata* would occur in the pine forests and under the conditions where amber was formed.

Grimmia torquata Hornsch.1912 *Grimmia torquata*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]**Grimmia** species.1908 *Grimmia* sp. Dixon, H. N., in Lewis, F. J. The Recent Plant Remains in the Scottish Peat Mosses. Part III. [Trans. Roy. Soc. Edinb., XLVI, p. 44.]1910 *Grimmia* sp. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]

Note. Of the *trichophylla* type, but with cells nearly all incrassate and sinuose to the base. Probably either *G. Mühlenbeckii* or *G. robusta*.

Rhacomitrium Brid.**Rhacomitrium affine** Lindb.1910 *Rhacomitrium heterostichum* var. Dixon, H. N. Postglacial Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]**Rhacomitrium canescens** (Weis) Brid.1910 *Rhacomitrium canescens*. Dixon, H. N., op. et Postglacial loc. cit.

- 1910 *Grimmia ericoides* Lindb. Hesselbo, A. Mosrester Quaternary fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raække, No. 25, p. 108.]
- 1922 *Rhacomitrium canescens*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faererne, p. 12.

Rhacomitrium fasciculare (Schrad.) Brid.

- 1922 *Rhacomitrium fasciculare*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faererne, p. 12.

Rhacomitrium heterostichum (Hedw.) Brid.

- 1922 *Rhacomitrium heterostichum*. Jessen, Knud, og Rasmussen, R., op. et loc. cit.

Rhacomitrium hypnoides (L.) Lindb.

- 1908 *Rhacomitrium lanuginosum* Brid. Dixon, H. N., Recent in Lewis, F. J. The Plant Remains in the Scottish Peat Mosses. Part III. [Trans. Roy. Soc. Edinb., XLVI, pp. 40, 48, 55.]
- 1910 *Grimmia hypnoides* Lindb. Hesselbo, A. Mosrester Diluvial fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raække, No. 25, p. 108.]
- 1914 *Rhacomitrium hypnoides* Lindb. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.]
- 1924 *Rhacomitrium lanuginosum* Brid. Burrell, W. H. Recent Pennine Peat. [The Naturalist, 1924, p. 148, tab. X.]

Rhacomitrium patens (Brid.)

- 1912 *Dryptodon patens* Brid. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Orthotrichaceae.

Anoectangium Hedw.

Anoectangium compactum Schwaegr.

- 1912 *Anoectangium compactum*. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 88.]

Amphidium Nees.

Amphidium lapponicum (Hedw.) Schimp.

- 1912 *Amphidium lapponicum*. Stark, P. op. cit., p. 89. Glacial

Zygodon Hook. & Tayl.**Zygodon viridissimus** (Dicks.) R. Br.

- 1913 *Zygodon viridissimus*. Travis, C. B. and W. G. Postglacial
On plant remains in post-glacial gravels at Sea-
forth, Liverpool. [Lancashire Naturalist, 1913,
p. 50.]

Ulota Mohr.**Ulota curvifolia** (Wahl.) Brid.

- 1912 *Ulota curvifolia*. Stark, P. Beitr. zur Kenntn. Glacial
der eiszeitlich. Fl. und Fauna Badens. [Ber. na-
turforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Orthotrichum Hedw.**Orthotrichum diaphanum** (Gmel.) Schrad.

- 1912 *Orthotrichum diaphanum*. Dixon, H. N., in Warren, Late glacial
S. H. A late Glacial stage in the Lea Valley.
[Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Orthotrichum species.

- 1907 *Orthotrichum* sp. Hartmann, F. Die Fossile Flora Diluvial
von Ingramsdorf. Inaug. Dissert. Breslau, p. 25.

Splachnaceae.**Splachnum** L.**Splachnum ampullaceum** L.

- 1912 *Splachnum ampullaceum*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.
[Ber. naturforsch. Gesellsch. Freiburg i. B., XIX,
p. 89.]

Splachnum sphaericum (L. f.) Sw.

- 1912 *Splachnum sphaericum*. Stark, P., op. et loc. cit. Glacial

Bryaceae.**Leptobryum** Wils.**Leptobryum pyriforme** (L.) Wils.

- 1910 *Leptobryum pyriforme*. Hesselbo, A. Mosrester Quaternary
fra Diluviet ved Skaerumhede. [Danmarks geol.
Undersøgelse, II Raekke, No. 25, p. 106.]

Pohlia Hedw.**Pohlia commutata (Schimp.) Lindb.**

- 1910 *Pohlia commutata*. Hesselbo, A. Mosrester fra Quaternary Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
 1912 *Webera commutata* Schimp. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Pohlia cruda (L.) Lindb.

- 1902 *Pohlia cruda*. Hartz, N. Bidr. til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.]
 1912 *Webera cruda* (L.) Bruch. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Pohlia cucullata (Schwaegr.) Bruch

- 1910 *Pohlia cucullata*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
 1912 *Webera cucullata* (Schwaegr.) Schimp. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Pohlia longicolla (Sw.) Lindb.

- 1912 *Webera longicolla* (Sw.) Hedw. Stark, P., op. et loc. cit. Glacial

Pohlia Ludwigii (Spreng.) Lindb.

- 1912 *Webera Ludwigii* Schimp. Stark, P., op. et loc. cit. Glacial

Pohlia nutans (Schreb.) Lindb.

- 1902 *Pohlia nutans*. Hartz, N. Bidr. til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.]
 1904 *Pohlia nutans*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
 1909 *Pohlia ?nutans*. Hartz, N. Bidr. til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
 1910 *Pohlia nutans*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
 1913 *Webera nutans* Hedw. Travis, C. B. and W. G. On plant remains in post-glacial gravels at Seaforth, Liverpool. [Lancashire Naturalist, 1913, p. 50.]
 1914 *Pohlia nutans*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracoviae d. sc. math. et nat., Sér. B, No. 2B, p. 248.]

- 1917 *Webera nutans* Hedw. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.]

Pohlia sphagnicola (Bry. eur.) Lindb.

- 1912 *Webera sphagnicola* Schimp. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Mniobryum Limpr.

Mniobryum albicans (Wahl.) Limpr.

- 1888 *Webera albicans* Schimp. Reid, C. and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Decade III, Vol. V, p. 444.]
- 1904 *Mniobryum albicans*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1910 *Pohlia albicans* Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Mniobryum albicans* var. *glaciale* Limpr. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]
- 1914 *Mniobryum albicans*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.]

Anomobryum Schimp.

Anomobryum concinnum (Spruce) Lindb.

- 1912 *Anomobryum concinnum*. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

Bryum Dill.

Bryum capillare L.

- 1912 *Bryum capillare*. Dixon, H. N. in Warren, S. H. A late Glacial Stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Bryum cirratum Hoppe et Hornsch.

- 1910 *Bryum cirratum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Bryum ?cirratum*. Dixon, H. N. in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Bryum crispulum Hampe.

- 1910 **Bryum crispulum**. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary

Bryum cyclophyllum (Schwaegr.) Bry. eur.

- 1910 **Bryum cyclophyllum**. Hesselbo, A., op. et loc. cit. Quaternary

Bryum Duvalii Voit.

- 1905 **Bryum ?Duvalii**. Neuweiler, E. Zur interglazial-flora der schweiz. Schieferkohlen. [Ber. d. Zürcher Bot. Gesellsch., XV, p. 98.] Interglacial
- 1914 **Bryum ?Duvalii**. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 17.] Glacial
- 1923 **Bryum Duvalii**. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 84.] Quaternary

Bryum elegans Nees.

- 1912 **Bryum elegans**. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.] Glacial

Bryum erythrocarpum Schwaegr.

- 1904 **Bryum erythrocarpum**. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial

Bryum intermedium Brid.

- 1912 **Bryum ?intermedium**. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial

Bryum lacustre Brid.

- 1914 **Bryum lacustre**. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracoviae d. sc. math. et nat., Sér. B. No. 2 B, p. 248.] Diluvial

Bryum microstegium Bry. eur.

- 1904 **Bryum cf. microstegium**. Weber, C. A., in Müller, G. und Weber. Ueber eine fröhdiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 16.] Preglacial

Bryum neodamense Itzigs.

- 1910 **Bryum neodamense** var. *ovatum* Lindb. and Arn. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary

- 1912 *Bryum neodamense*. Stark, P. Beitr. zur Kenntn. Glacial
der eiszeitlich. Fl. und Fauna Badens. [Ber.
naturforsch. Gesellsch. Freiburg i. B., XIX, p. 89.]

***Bryum pallens* Swartz.**

- 1888 *Bryum pallens*. Reid, C. and Ridley, H. N. Fossil Glacial
Arctic Plants from Hoxne, in Suffolk. [Geol.
Mag., Decade III, Vol. V, p. 444.]
- 1903 *Bryum pallens*. Range, Paul. Das Diluv. von Diluvial
Lübeck und seine Dryastone [Zeitschr. f.
Naturwiss., LXXVI, p. 253.]
- 1912 *Bryum pallens*. Dixon, H. N., in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1913 *Bryum pallens*. Travis, C. B. and W. G. On Postglacial
plant remains in post-glacial gravels at Seaforth,
Liverpool. [Lancashire Naturalist, 1913, p. 50.]
- 1923 *Bryum pallens*. Baumberger, E., etc. Die diluvia- Quaternary
len Schieferkohlen der Schweiz. [Beitr. Geol.
Schweiz, geotechn. Ser., Lief. 8, p. 521.]
- 1923 *Bryum pallens*. Gams, H. Interglacial bryophytes Interglacial
..... [B. 2, Geol. Karte d. Schweiz, Geotechn.
Ser., Lief. 8, p. 521.]

***Bryum Schleicheri* Schwaegr.**

- 1912 *Bryum Schleicheri* var. *latifolium* Schimp. Stark, P. Glacial
Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna
Badens. [Ber. naturforsch. Gesellsch. Freiburg i.
B., XIX, p. 89.]

***Bryum ventricosum* Dicks.**

- 1902 *Bryum ventricosum*. Hartz, N. Bidrag til Dan- Late glacial
marks senglacial Flora og Fauna. [Danmarks geol.
Undersøgelse, II Raekke, No. 11, p. 15.]
- 1903 *Bryum ?pseudo-triquetrum* Schwaegr. Range, Paul. Diluvial
Das Diluv. von Lübeck und seine Dryastone.
[Zeitschr. f. Naturwiss., LXXVI, p. 253.]
- 1909 *Bryum ventricosum*. Hartz, N. Bidr. til Dan- Interglacial
marks tertiaere Flora. [Danmarks geol. Un-
dersøgelse, II Raekke, No. 20, p. 260.]
- 1909 *Bryum ventricosum*. Nørregaard, E. M. Et sengla- Late glacial
cialt ... Vandløb fra Dejbjerg Bakker. [Meddel.
fra Dansk geol. Foren., No. 15, p. 325.]
- 1910 *Bryum ventricosum*. Dixon, H. N. Some "Neolithic" Postglacial
Moss remains from Fort William. [Ann.
Scott. Nat. Hist., 1910, p. 104.]
- 1910 *Bryum ventricosum*. Hesselbo, A. Mosrester fra Quaternary
Diluviet ved Skaerumhede. [Danmarks geol. Un-
dersøgelse, II Raekke, No. 25, p. 106.]
- 1912 *Bryum ventricosum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. Late glacial
[Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *Bryum pseudotriquetrum* Schwaegr. Weber, C. A. Diluvial
des Krakauer Diluviums. [Bull. int. Acad. Sci.
Cracovie d. sc. math. et nat., Sér. B, No. 2 B,
p. 248.]

- 1914 *Bryum pseudotriquetrum* Schwaegr. Weber, C. A. Glacial
Die Mammutflora von Borna. [Abh. Nat. Ver.
Bremen, XXIII, Heft 1, p. 17.]

Bryum species.

- 1890 *Bryum* sp. Dawson, Sir Wm., and Penhallow, D. Pleistocene
P. On the Pleistocene Flora of Canada. [Bull.
Geol. Soc. Amer., I, p. 315.]
- 1902 *Bryum* sp. Hartz, N. Bidrag til Danmarks sengla- Late glacial
cial Flora og Fauna. [Danmarks geol. Under-
søgelse, II Raekke, No. 11, p. 15.]
- 1904 *Bryum* sp. Weber, C. A. in Mueller, G. u. We- Preglacial
ber. Ueber einer fröh diluviale und vorglaciale
Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Lan-
desanstalt und Bergakad., Heft 40, p. 16.]
- 1910 *Bryum* sp. Hesselbo, A. Mosrester fra Diluviet Quaternary
ved Skaerumhede. [Danmarks geol. Undersøgelse,
II Raekke, No. 25, p. 106.]
- 1914 *Bryum* sp. Weber, C. A. Die Mammutflora von Glacial
Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1,
p. 17.]
- 1923 *Bryum* sp. Baumberger, E., etc. Die diluvialen Quaternary
Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz,
geotechn. Ser., Lief. 8, pp. 145, 333.]
- "*Bryon crispatum* Grand Eury ist Autophyllites fur-
catus Grand 'Eury." Jongmans, in Med. van de
Rijksopspor. van Delfstoffen, No. 3 (1911), p. 441.

Mniaceae.

Mnium L.

Mnium affine Bland.

- 1898 *Mnium affine*. Andersson, Gunnar. Stud. öfver Quaternary
Finlands Kvartärflora. [Bull. de la Com-
mission géol. de Finlande, Tome 2, No. 8, p. 137.]
- 1899 *Mnium affine*. Bertrand, C. E. Descr. d'un échan- Pleistocene
tillon de charbon papyracé [Ann. Soc.
géol. du Nord, XXVIII, p. 190.]
- 1902 *Astrophyllum cuspidatum* (L.) Lindb. Hartz, N. Late glacial
Bidrag til Danmarks senglaciale Flora og Fauna.
[Danmarks geol. Undersøgelse, II Raekke, No. 11,
p. 11.]
- 1904 *Mnium affine*. Bell, A. M. Implementiferous Late glacial
Sections at Wolvercote (Oxfordshire). [Quart.
Journ. Geol. Soc., LX, p. 124.]
- 1909 *Astrophyllum cuspidatum* (L.) Lindb. Hartz, N. Interglacial
Bidrag til Danmarks tertiaere Flora. [Dan-
marks geol. Undersøgelse, II Raekke, No. 20,
p. 260.]
- 1914 *Mnium affine* var. *integrifolium* Lindb. Zmuda, Diluvial
A. J. Fossile Flora des Krakauer Diluviums.
[Bull. int. Acad. Sci. Cravovie d. sc. math. et
nat., Sér. B., No. 2 B, p. 248.]

Mnium antiquorum Card et Dixon.

- 1915 **Mnium antiquorum** Card. & Dixon in Reid, Clement and E. M. The Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, p. 52, Fig. 3.] See also Bryologist, XIX, p. 51 (1916).

Mnium cinclidiooides (Blytt) Hüben.

- 1897 **Mnium cinclidiooides**. Beck, R., und Weber, C. A. Ueber ein Torflager des sächsisch. Erzgebirges. [Zeitschr. d. deutsch. geol. Ges., XLIX, p. 666.]
- 1898 **Mnium cinclidiooides**. Andersson, Gunnar. Stud. Finlands Kvartärfloara. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.]
- 1912 **Mnium cinclidiooides**. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. u. Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.]
- 1914 **Mnium cinclidiooides**. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 248.]

Mnium curvatum (Lindb.) Limpr.

- 1910 **Astrophyllum curvatum** Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

Mnium cuspidatum (L.) Leyss.

- 1909 **Astrophyllum silvaticum** Lindb. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
- 1910 **Astrophyllum silvaticum** Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

Mnium hornum L.

- 1895 **Mnium hornum**. Gepp, A. Fossil plant-remains in peat. [Journ. of Bot., XXXIII, p. 181.]
- 1898 **Mnium hornum**. Andersson, Gunnar. Stud. öfver Finlands Kvartärfloara. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.]
- 1910 **Mnium hornum**. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]
- 1922 **Mnium hornum**. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvmose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.]

Mnium hymenophylloides Hüben.

- 1910 **Mnium hymenophylloides**. Sukačev, V. N. Sur la flore arctique près du village Demianskoe, gouv. Tobolsk. [Bull. Acad. Impér. Sci. St. Pétersbourg, 1910, p. 460, figs. 1—3.]

- 1910 *Astrophyllum hymenophylloides* (Hueben.) Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]
- 1914 *Mnium hymenophylloides*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 16.]
- 1918—20 *Mnium hymenophylloides*. Docturowsky, V. S. and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.]

***Mnium marginatum* (Dicks.) P. Beauv.**

- 1910 *Astrophyllum marginatum* (Dicks.) Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

***Mnium medium* Bry. eur.**

- 1909 *Astrophyllum medium* (Bry. eur.) Lindb. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
- Note. The author gives no indication of the characters by which he distinguishes it from *M. affine*. It was presumably not fruiting.

***Mnium punctatum* (L.) Hedw.**

- 1888 *Mnium punctatum*. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Dec. III, Vol. V, p. 444.]
- 1898 *Mnium punctatum*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.]
- 1905 *Mnium punctatum*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas... [Vierteljahr. Naturforsch. Ges. Zurich, L, Heft 6, p. 40.]
- 1911 *Mnium punctatum*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish Peat Mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831 bis.]
- 1912 *Mnium punctatum* (or *M. subglobosum* Bry. eur.). Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *Mnium punctatum*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 17.]
- 1923 *Mnium punctatum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz., geotechn. Ser., Lief. 8, pp. 145, 333.]

***Mnium riparium* Mitten.**

- 1914 *Mnium riparium*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.]

Note. The identification, in the presumable absence of inflorescence, appears to me extremely doubtful.

Mnium rostratum Schrad.

- 1904 **Mnium rostratum.** Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
 1912 **Mnium rostratum.** Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial

Mnium rugicum Laur.

- 1904 **Mnium rugicum.** Weber, C. A., in Mueller, G. und Weber. Ueber eine frühdiluviale und vor-glaciale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 16.] Preglacial
 1912 **Mnium rugicum.** Stark, P. Beitr. zur Kennntn. der eiszeitlich. Flora und Fauna Badens. [Ber. Naturforsch. Ges. Freiburg i. B., XIX, p. 26.] Glacial
 1914 **Mnium rugicum.** Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.] Diluvial

Mnium subglobosum Br. Eur.

- 1909 **Astrophyllum subglobosum** (Bry. eur.) Lindb. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
 1912 **Mnium subglobosum** (cf. *M. punctatum*).

Mnium undulatum (L.) Weis.

- 1905 **Mnium undulatum.** Neuweiler, E., Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. Naturforsch. Ges. Zurich, I, Heft 6, p. 40.] Pleistocene
 1909 **Astrophyllum undulatum (L.) Lindb.** Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial

Mnium species.

- 1892 **Mnium sp.** Boulay, N. Flore pliocène du Mont-Dore, p. 44.
 Note. According to the author the generic position is very doubtful. *Fissidens* and *Hypnum* are also suggested.

Cinclidium Sw.

Cinclidium arcticum Schimp.

- 1910 **Cinclidium arcticum.** Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary

- 1914 *Cinclidium arcticum*. Weber, C. A. Die Mammut-flora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 17.] Glacial

Cinclidium latifolium Lindb.

- 1910 *Cinclidium latifolium*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary

Cinclidium stygium Sw.

- 1907 *Cinclidium stygium*. Dixon, H. N., in Lewis, F. J. Peat Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.]
 1914 *Cinclidium stygium*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.] Diluvial

Cinclidium species.

- 1925 *Cinclidium* sp. Dixon, H. N. Moss Remains in Recent Russian Peat. [Journ. of Bot., LXIII, p. 370.]

Aulacomniaceae.

Aulacomnium palustre (L.) Schwaegr.

- 1895 *Aulacomnium palustre*. Gepp, A. Fossil plant-remains in peat. [Journ. of Bot., XXXIII, p. 181.] Recent sub-marine peat
 1902 *Sphaerocephalus palustris* (L.) Lindb. Hartz, N. Bidrag til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.] Late glacial
 1903 *Aulacomnium palustre*. Range, Paul. Das Diluv. von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 269.] Diluvial
 1904 *A. palustre*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
 1907 *A. palustre*. Dixon, H. N., in Lewis, F. J. Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.] Peat
 1910 *Sphaerocephalus palustris* (L.) Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
 1912 *A. palustre*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
 1914 *A. palustre* var. *imbricatum* Bry. eur. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
 1917 *A. palustre*. Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 82.] Recent
 1917 *A. palustre*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.] Diluvial

- 1918—20 *A. palustre*. Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Interglacial
- 1922 *A. palustre*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faererne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] Recent
- 1923 *A. ?palustre*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.] Quaternary
- 1924 *A. palustre*. Burrell, W. H. Pennine Peat. [The Naturalist, 1924, p. 148.] Recent

Aulacomnium turgidum Schwaegr.

- 1910 *Aulacomnium turgidum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
- 1914 *A. turgidum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial

Meeseaceae.**Paludella** Ehrh.**Paludella squarrosa** (L.) Brid.

- 1896 *Paludella squarrosa*. Farneti, R. Ricerche di briol. palaeont. al periodo glaciale. [Atti del Istit. Bot. della Reale Univ. d. Pavia, Ser. 2, V, p. 53.] Glacial
- 1914 *P. squarrosa*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1918 *P. squarrosa*. Beyle, M. Ueber einige Ablagerungen fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. [Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 40.] Glacial
- 1925 *P. squarrosa*. Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot., LXIII, p. 370.] Recent

Amblyodon P. Beauv.**Amblyodon dealbatus** (Dicks.) Beauv.

- 1912 *Amblyodon dealbatus*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.] Glacial
- 1913 *A. dealbatus*. Travis, C. B. and W. G. On plant remains in postglacial gravels at Seaforth, Liverpool. [Lancashire Naturalist, 1913, p. 50.] Postglacial

Meesea Hedw.**Meesea Albertini** Bry. eur.

- 1912 *Meesea Albertini*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.] Glacial

Meesea longiseta Hedw.

- 1905 **Meesea longiseta**. Neuweiler, E. Zur Interglazialfl. der schweiz. Schieferkohlen. [Ber. d. schweiz. bot. Ges., Heft 15, Beilage. Neunter Ber. d. Zürich. bot. Ges., p. 96.] Interglacial
- 1909 **M. longiseta**. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] Interglacial
- 1912 **M. longiseta**. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.] Glacial
- 1923 **M. longiseta**. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 84, 334.] Quaternary

Meesea trichodes (L.) Spruce.

- 1910 **Meesea trichodes**. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
- 1912 **Meesea trichodes**. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.] Glacial

Meesea triquetra (L.) Aongstr.

- 1896 **Meesea triquetra**. Farneti, R. Ricerche di briol. palaeont. al periodo glaciale. [Atti del Istit. Bot. della Reale Univ. d. Padova, Ser. 2, V, p. 52.] Glacial
- 1898 **M. triquetra**. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.] Quaternary
- 1904 **M. triquetra**. Geinitz, C., und Weber, C. A. Ueber ein Moostorflager der Rostocker Heide. [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.] Postglacial
- 1904 **M. triquetra** var. *timmoides* Sanio. Geinitz und Weber, op. et loc. cit. Postglacial
- 1908 **Meesea tristicha** (Funck) Bry. eur. Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engler's Bot. Jahrb., XLII, p. 43.] Pleistocene
- 1908 **M. tristicha** var. *timmoides* Sanio. Weber, C. A. op. cit., pp. 41, 43. Pleistocene
- 1909 **M. tristicha** (Funck) Bry. eur. Schuster, J. Palaeobot. Not. aus Bayern. [Ber. bayér. bot. Ges., XI, 1, p. 58.] Schieferkohlen
- 1910 **M. triquetra**. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
- 1912 **M. tristicha** (Funck) Bry. eur. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.] Glacial
- 1914 **M. triquetra**. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracoviae d. sc. math. et nat., Sér. B., No. 2 B, p. 248.] Diluvial
- 1923 **M. triquetra**. Gams, H. Interglacial bryophytes. [Die Schieferkohlen der Schweiz, B. 2, Geol. Karte d. Schweiz, geotechn. Ser., Lief. 8, p. 521.] Interglacial

- 1923 *M. triquetra*. Baumberger, E., etc. Die diluvialen Quaternary Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 84, 334, 521.]
 1925 *M. triquetra*. Dixon, H. N. Moss Remains in Recent Russian Peat. [Journ. of Bot., LXIII, p. 370.]

Catoscopiaceae.

Catoscopium Brid.

Catoscopium nigritum (Hedw.) Brid.

- 1912 *Catoscopium nigritum*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.]

Bartramiaceae.

Plagiopodopsis Britton & Hollick

Plagiopodopsis Scudderii Britton et Hollick.

- in Bull. Torr. Bot. Cl. 42:10 (1915); Figs. 1, 2. Tertiary Shales of Florissant, Colorado, U. S. A.
Cum fructu.

Conostomum Sw.

Conostomum tetragonum (Dicks.) Lindb.

- 1914 *Conostomum boreale* Sw. Zmuda, A. J. Fossile Diluvial Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

Philonotis Brid.

Philonotis calcarea (Br. Eur.) Schimp.

- 1911 *Philonotis calcarea*. Dixon, H. N. in Lewis, F. J. Iceland The plant remains in the Scottish Peat Mosses. peat; recent Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831.]

Philonotis coespitosa Wils.

- 1918 *Philonotis coespitosa*. Heim, Arnold, and Gams, H. Interglacial Interglaz. Bild. bei Wildhaus. [Vierteljahrsschr. naturf. Ges. Zürich, LXIII, p. 27.]

Philonotis fontana (L.) Brid.

- 1888 *Philonotis fontana*. Reid, C. and Ridley, H. N. Glacial Fossil Arctic Plants from... Hoxne, in Suffolk. [Geol. Mag., Dec. III, Vol. V, p. 444.]

- 1904 *P. fontana* var. *ampliretis* Dixon. Bell, A. M. Late glacial
Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, 124.]
- 1909 *P. fontana*. Nørregaard, E. M. Et senglacialt ... Late glacial
fra Dejbjerg Bakker. [Meddel. fra Dansk geol. Förening, No. 15, p. 326.]
- 1910 *P. fontana*. Hesselbo, A. Mosrester fra Diluviet Diluvial
ved Skaerumhede. [Danmarks geol. Undersøgelse, II Rakke, No. 25, p. 106.]
- 1911 *P. fontana*. Dixon, H. N. in Lewis, F. J. The Iceland
plant remains in the Scottish Peat Mosses. Part peat, recent
IV. [Trans. Roy. Soc. Edinb., XLVII, pp. 830,
831.]
- 1912 *P. fontana*. Dixon, H. N. in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *P. fontana*. Zmuda, A. J. Fossile Flora des Diluvial
Krakauer Diluviums. [Bull. int. Acad. Sci. d.
sc. math. et nat., Sér. B, No. 2 B, p. 249.]
- 1915 *P. fontana*. Reid, Clement and E. M. The Pliocene Floras of the Dutch-Prussian Border. [Med.
van de Rijksopspor. van Delfstoffen, No. 6, p. 53.]
- 1923 *P. fontana* var. *glacialis*. Baumberger, E., etc. Quaternary
Die diluvialen Schieferkohlen der Schweiz. [Beitr.
Geol. Schweiz., geotechn. Ser., Lief. 8, p. 521.]

Philonotis seriata Mitt.

- 1912 *Philonotis seriata*. Stark, P. Beitr. zur Kenntn. Glacial
der eiszeitlich. Fl. und Fauna Badens. [Ber.
naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Philonotis tomentella Mol.

- 1912 *Philonotis alpicola* Jur. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 89.]
- 1914 *Philonotis tomentella*. Weber, C. A. Die Mammut- Glacial
flora von Borna. [Abh. Nat. Ver. Bremen, XXIII,
Heft 1, p. 18.]

Philonotis species.

- 1911 *Philonotis* sp. Dixon, H. N. in Lewis, F. J. Iceland
The plant remains in the Scottish Peat Mosses. peat, recent
Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.]
Note. Leaf with very stout nerve; much eroded.
May be an *adpressa* form.

Timmiaceae.

Timmia Hedw.

Timmia austriaca Hedw.

- 1912 *Timmia cf. austriaca*. Stark, P. Beitr. zur Kenntn. Glacial
der eiszeitlich. Fl. und Fauna Badens. [Ber. na-
turforsch. Ges. Freiburg i. B., XIX, p. 116.]

Timmia bavarica Hessl.

- 1910 **Timmia bavarica**. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Diluvial

Timmia megapolitana Hedw.

- 1912 **Timmia megapolitana**. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] Glacial

Timmia norvegica Zett.

- 1910 **Timmia norvegica**. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Diluvial
- 1912 **T. norvegica**. Dixon, H. N. in Warren, S. H. Late glacial A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, P. 231.]
- 1914 **T. norvegica**. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 18.] Glacial

Polytrichaceae.**Catharinaea** Ehrh.**Catharinaea subundulata** (Goepp. et Menge.)

Broth. in Engl. & Prantl, die natürl. Pflanzenfam., Ed. 2, Bd. XI, 2 Hälften, p. 523 (1925).

- 1853 **Polytrichum subundulatum** Goepp. & M. in Goepert, H. R. Ueber die Bernsteinflora, p. 459. Amber (Atrichum subundulatum Schimp., Traité de Pal. Végétale, I, p. 244).

Note. Neither Schimper nor any of the later writers seem to comment on this. It is presumably not specifically distinct from *C. undulata*.

Catharinaea undulata (L.) Web. et Mohr.

- 1922 **Catharinaea undulata**. Jessen, Knud og Rasmussen, Recent R. Et Profil Gennem en Tørvemose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.]

Oligotrichum Lam. et De Cand.**Oligotrichum hercynicum** (Huds.) Lindb.

- 1912 **Oligotrichum hercynicum** Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Pogonatum P. Beauv.**Pogonatum suburnigerum**
(Goepp. et Menge.) Dixon, comb. nov.

- 1853 **Polytrichum suburnigerum** Goepp. & M. in Goepert, H. R. Ueber die Bernsteinflora, p. 459. Amber

Note. Schimper (*Traité de Pal. Végétale*, I, p. 244) points out that the remains being solely leaves, the specific position must be doubtful.

Pogonatum urnigerum (L.) P. Beauv.

- | | | |
|------|---|---------------|
| 1902 | Pogonatum urnigerum. Hartz, N. Bidrag til Danmarks senglaciaile Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.] | Late glacial |
| 1904 | P. urnigerum. Weber, C. A. in Mueller, G. und Weber. Ueber eine fröhdiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanst. und Bergakad., Heft 40, p. 16.] | Early glacial |
| 1914 | P. urnigerum. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] | Diluvial |

Polytrichum L.
Polytrichum alpinum L.

- | | | |
|------|--|----------|
| 1912 | Polytrichum alpinum. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] | Glacial |
| 1914 | P. alpinum. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] | Diluvial |

Polytrichum attenuatum Menz.

- | | | |
|------|---|--------------|
| 1909 | Polytrichum attenuatum. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] | Interglacial |
| 1922 | P. attenuatum. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faerørne. Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] | Recent |

Polytrichum commune L.

- | | | |
|------|---|--------------|
| 1897 | Polytrichum commune. Beck, R. und Weber, C. A. Ueber ein Torflager des sächsisch. Erzgebirges. [Zeitschr. d. deutsch. geol. Ges., XLIX, p. 666.] | Diluvial |
| 1909 | P. commune. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] | Interglacial |
| 1910 | P. commune. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] | Quaternary |
| 1914 | P. commune. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] | Diluvial |
| 1917 | P. commune. Rudolph, K. Untersuch. über den Aufbau böhmisch. Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 85.] | Recent |
| 1922 | P. commune. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] | Recent |
| 1924 | P. commune. Burrell, W. H. Pennine Peat. [The Recent Naturalist, 1924, p. 148.] | Recent |

Polytrichum Florissanti Knowlton.

- 1916 (?) *Polytrichum Florissanti* Knowlton n. sp. A Review of the fossil plants from Florissant, Colorado. [Proc. United States Nat. Mus., Vol. 44, p. 245; t. 12.]

Note. A seta with capsule; no leaves found. The author compares with *P. juniperinum*, but there can hardly be any reason for selecting this species as the nearest ally. The reference to *Polytrichum* itself is not certain. From the figure it certainly looks like this genus.

Polytrichum gracile Menz.

- 1914 *Polytrichum gracile*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.]

Polytrichum juniperinum Willd.

- 1904 *Polytrichum juniperinum*. Weber, C. A., in Müller, G. Ueber eine frühdiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanst. und Bergakad., Heft 40, pp. 15, 16.]
- 1909 *P. juniperinum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1912 *P. juniperinum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, 116.]
- 1914 *P. juniperinum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.]

Tertiary.
Florissant shales

Polytrichum piliferum Schreb.

- 1910 *Polytrichum pilosum* Floercke (sic.). Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.]

Note. I presume that this refers to *P. pilosum* Neck., which is synonymous with *P. piliferum* Schreb.

Quaternary

Polytrichum sexangulare Floerke.

- 1914 *Polytrichum sexangulare*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.]
- 1915 *P. sexangulare*. Camus, F. Sur les mousses trouvées dans un mammouth. [Compt. Rend. Acad. Sci. Paris, CLX, p. 842.]

Diluvial

Glacial

Polytrichum strictum Banks.

- 1909 *Polytrichum strictum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]

Interglacial

- 1910 *P. strictum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
- 1912 *P. strictum* var. *alpestre* Limpr. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] Glacial
- 1923 *P. strictum*. Baumberger, S., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 84.] Quaternary

Polytrichum subseptentrionale Goepp. et Menge.

- 1853 *Polytrichum subseptentrionale* Goepp. & M. in Amber Goepert, H. R. Ueber die Bernsteinflora. [Monatsber. d. Berliner Akad., 1853, p. 459.] Note. Schimper (Traité de Pal. Végétale, I, p. 244) remarks on this that in view of the material available the affinity must be problematic, and even the genus may be doubtful.

Polytrichum species.

- 1898 *Polytrichum* sp. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.] Quaternary
- 1907 *Polytrichum* sp. Holst, N. O. Präglaciale Dryas-förande nära Ystad. [Geol. Fören. Förh., XXIX, p. 232.] Preglacial
- 1908 *Polytrichum* sp. Dixon, H. N. in Lewis, F. J. The plant remains in the Scottish Peat Mosses. Part III. [Trans. Roy. Soc. Edinb., XLVI, p. 48.] Quaternary
- 1910 *Polytrichum* sp. Sukačev, V. N. Sur la trouvaille de la flore arctique fossile près du village Demianskóe [Bull. Acad. Impér. Sc. St. Petersbourg, 1910, p. 460.] Quaternary
- 1910 *Polytrichum* sp. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 106.] Quaternary
- 1912 *Polytrichum* sp. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1918—20 *Polytrichum* sp. Docturowsky, V. S., and Kuriashev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Glacial

Hedwigiaceae.

Hedwigia Ehrh.

Hedwigia albicans (Web.) Lindb.

- 1914 *Hedwigia albicans*. Zmuda, A. J. Fossile Fl. des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 248.] Diluvial

Fontinalaceae.

Fontinalis (Dill.) L.

Fontinalis antipyretica L.

- 1914 **Fontinalis antipyretica**. Zmuda, A. J. Fossile Flora Diluvial des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]
- 1919 **F. antipyretica?** Gadeceau, E. Les Forêts submergées de Belle-Ile-en-mer. [Bull. biol. de la France, etc. LIII, p. 295.]
Note. Many fragments; almost certainly *F. antipyretica*.

Fontinalis hypnoides Hartm.

- 1898 **Fontinalis hypnoides**. Andersson, Gunnar. Stud. Quaternary öfver Finnlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.]

Fontinalis pristina Lesq.

- 1883 **Fontinalis pristina** Lesq. in Hayden, F. V. Tertiary Flora of United States. [Rep. of U. S. A. Geol. Survey, VIII, p. 135; t. 21, f. 9.] See also Britton & Hollick, Bull. Torr. Bot. Club, 34: 139; t. 9, f. 2, 2 a.
Note. Britton and Hollick say the determination is questionable, Knowlton (Rev. of the Fossil Plants from Florissant..., p. 245) states that precisely similar structures have proved to be avian feathers. It is, to say the least, exceedingly dubious whether it is a moss.

Fontinalis Sismondana Schimp.

- 1869 **Fontinalis Sismondana** Schimp., Traité de Pal. Végétale, I, p. 245. Figured by Sismonda, E. Matériaux pour servir à la Pal. d. terr. tert. du Piémont, 1865, t. XXXI bis.
Note. Schimper remarks that this is very close to *F. antipyretica*. He does not suggest wherein the difference consists.

? **Fontinalis Tournalii** (Brongn.) Schimp.

- 1828—38 **Muscites Tournalii** Brongn. Ad. Histoire des Miocene Végétaux Fossiles, p. 93, t. X. **Fontinalis** (?) *Tournalii* Schimp., Traité de Pal. Végétale, I, p. 245, t. VI, figs. 14, 15, 16 (nec Saporta, Végét. d. S. E. de la France, tab. X).

Note. Schimper says that the characters recall a *Fontinalis* or a *Dichelyma* rather than any other pleurocarpous moss; it is a very small plant.

Fontinalis species.

- 1890 *Fontinalis* sp. Dawson, Sir Wm., and Penhallow, D. P. On the Pleistocene Flora of Canada. [Bull. Geol. Soc. Amer., I, p. 315.] Pleistocene
- 1898 *Fontinalis* sp. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.] Quaternary
- 1900 *Fontinalis* sp. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley, [Rep. of 70th Meeting of Brit. Assoc., Bradford, 1900, Pt. 1, p. 335.] Pleistocene
- 1911 *Fontinalis* sp. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831 bis] Recent. Ice-land peat
- 1920 *Fontinalis* sp. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Ill. Bull., XVII, pp. 332, 335, 377.] Pleistocene

Dichelyma Myr.

D. capillaceum. cf. *Distichium capillaceum* (1920).
Dichelyma sp. cf. *Fontinalis Tournalii*.

Climaciaceae.**Climacium** Web. et Mohr**Climacium dendroides** (Dill.) Web. et Mohr

- 1902 *Climacium dendroides*. Hartz, N. Bidrag til Danmarks senglacial Fl. og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.] Late glacial
- 1904 *C. dendroides*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1910 *C. dendroides*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *C. dendroides*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *C. dendroides*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] Diluvial
- 1923 *C. dendroides*? Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 334.] Quaternary
- 1924 *C. dendroides*. Beyle, M. Ueber einige Ablag.... der Hamburger Gegend. Dritter Teil. [Mitteil. aus dem Mineralog.-geol. Staatsinstitut, Heft VI, p. 26.] Glacial

Leucodontaceae.**Leucodon** Schwaegr.**Leucodon sciurooides** (L.) Schwaegr.

- 1905 *Leucodon sciurooides*. Neuweiler, E. Die prähistor. Pleistocene

- Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]
- 1914 *L. sciuroides*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]
- 1923 *L. sciuroides*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]
- Note. A f. *falcata* is also recorded, with the type.

Antitrichia Brid.

Antitrichia curtipendula (Hedw.) Brid.

- 1898 *Antitrichia curtipendula*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 135.]
- 1905 *A. curtipendula*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, pp. 37, 39.] (The latter reference as *Anomodon curtipendula, lapsu calamis*).
- 1909 *A. curtipendula*. Hartz, N. Bidrag til Danmarks tertiaere.... Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
- 1910 *A. curtipendula*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]
- 1913 *A. curtipendula*. Beyle, M. Ueber einige Ablag... der Hamburger Gegend. Erster Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6 Beih., p. 83.]
- 1923 *A. curtipendula*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 334.]

Neckeraceae.

Neckera Hedw.

Neckera complanata (L.) Hüben,

- 1905 *Neckera complanata*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, pp. 38, 40.]
- 1909 *N. complanata*. Hartz, N. Bidrag til Danmarks tertiaere.... Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1910 *N. complanata*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]
- 1920 *N. complanata*. Reid, E. M. Two preglacial Floras from Castle Eden. [Quart. Journ. Geol. Soc., LXXVI, p. 111.]
- 1923 *N. complanata*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]



- 1925 *N. complanata*? Weyland, H. Beitr. zur Kenntn. Upper
fossiler Moose. I. [Senckenbergiana, VII, Heft Pliocene
1/2, p. 53, t. III.]

Neckera crispa (L.) Hedw.

- 1901 *Neckera crispa*. Clerici, E. Contrib. alla Recent
geologia dei dintorni di Roma. [Rend. Lincei,
Roma, X, p. 79.]
- 1905 *N. crispa*. Neuweiler, E. Die prähistor. Pflanzen- Pleistocene
reste Mitteleuropas.... [Vierteljahr. naturforsch.
Ges. Zürich, L, Heft 6, p. 38.]
- 1909 *N. crispa*. Hartz, N. Bidrag til Danmarks tertiaere Interglacial
.... Flora. [Danmarks geol. Undersøgelse, II
Raekke, No. 20, p. 261.]
- 1923 *N. crispa*. Baumberger, E., etc. Die diluvialen Quaternary
Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz,
geotechn. Ser., Lief. 8, p. 334.]
- 1924 *N. crispa*. Maffei, Luigi. Contrib. allo stud. d. Quaternary
flora fossile de Pianico. [Atti Istit. Bot.
Univ. Pavia, Ser. III, 1/2, p. 53, t. III.]

Neckera pennata (L.) Hedw.

- 1914 *Neckera pennata*. Zmuda, A. J. Fossile Flora Diluvial
des Krakauer Diluviums. [Bull. int. Acad. Sci.
Cracovie d. sc. math. et nat., Sér. B, No. 2 B,
p. 249.]

Neckera pumila Hedw.

- 1897 *Neckera pumila*. Fischer, Ed. Nachtr. zum Pflan- Interglacial
zenverzeichn. von Pianico-Sellere. [Neues
Jahrb. f. Mineralogie, II, p. 106.]
- 1910 *N. pumila*. Dixon, H. N. Some "Neolithic" Moss Postglacial
remains from Fort William. [Ann. Scott. Nat.
Hist., 1910, p. 104.]

Neckera species.

- 1908 *Neckera* sp. Engelhardt, H., and Kinkelin, F. Upper
Oberpliocene Fl. und Fauna. des Frank- Pliocene
furter Klärbeckens. [Abh. Senckenb. naturforsch.
Ges., XXIX, p. 188, t. XXII, figs. 18—21.]

Homalia (Bríd.) Bry. eur.

Homalia trichomanoides (Schreb.) Bry. eur.

- 1904 *Homalia trichomanoides*. Bell, A. M. Implementi- Late glacial
ferous Sections at Wolvercote (Oxfordshire).
[Quart. Journ. Geol. Soc., LX, p. 124.]
- 1914 *H. trichomanoides*. Zmuda, A. J. Fossile Flora des Diluvial
Krakauer Diluviums. [Bull. int. Acad. Sci. Cra-
covie d. sc. math. et nat., Sér. B, No. 2 B,
p. 249.]
- 1920 *H. trichomanoides*. Reid, E. M. Two preglacial Pliocene
Floras from Castle Eden. [Quart. Journ. Geol.
Soc., LXXVI, p. 111.]

Pinnatella (C. M.) Fleisch.**Pinnatella species.**

- 1915 *Pinnatella* sp. Reid, Clement and E. M. The Pliocene Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, pp. 51.] See also Bryologist, XIX, p. 51 (1916.) Note. Fragments only, showing traces of an intra-marginal border to the leaves, and in other respects agreeing with the § *Urocladium* of *Pinnatella*. Near to *P. alopecuroides* (Hook.) Fleisch., but probably specifically distinct.

Thamnium Bry. eur.**Thamnium alopecurum (L.) Bry. eur.**

- 1909 *Porotrichum alopecurum* (L.) Mitt. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1925 *Thamnium alopecurum*. Weyland, H. Beitr. zur Kenntn. fossiler Moose. I. [Senckenbergiana, VII, Heft 1/2, p. 11, fig. 1.]

Thamnium species.

- 1908 *Thamnium* sp. Engelhardt, H., and Kinkel, F. Oberpliocäne Fl. und Fauna des Frankfurter Klärbeckens. [Abh. Senckenb. naturforsch. Ges., XXIX, p. 189, t. 22, figs. 14, 15.] Note. I should say *T. alopecurum*, without doubt.

Lembophyllaceae.**Isothecium Brid.****Isothecium myosuroides (Dill.) Brid.**

- 1909 *Isothecium myosuroides*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1910 *Eurhynchium myosuroides* (Dill.) Schimp. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
- 1913 *Isothecium myosuroides*. Beyle, M. Ueber einige Ablagerungen der Hamburger Gegend. Erster Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6, Beih., p. 83.]

Isothecium viviparum (Neck.) Lindb.

- 1905 *Isothecium myurum* (Poll.) Brid. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]

- 1909 *I. myurum*. Hartz, N. Bidrag til Danmarks ter-
tiaere Flora. [Danmarks geol. Undersøgelse,
II Raekke, No. 20, p. 261.] Interglacial
- 1910 *Eurhynchium myurum* (Poll.) Dixon. Dixon, H. N.
Some "Neolithic" Moss remains from Fort
William. [Ann. Scott. Nat. Hist., 1910, p. 105.] Postglacial
- 1914 *Isothecium myurum*. Zmuda, A. J. Fossile Flora
des Krakauer Diluviums. [Bull. int. Acad. Sci.
Cracovie d. sc. math. et nat., Sér. B, No. 2 B,
p. 249.] Diluvial

Fabroniaceae.

Fabronia Raddi, cf. *Muscites serratus* Goepp. & Ber.

Entodontaceae.

Orthothecium Bry. eur.

Orthothecium chryseum (Schwaegr.) Bry. eur.

- 1910 *Stereodon chryseus* (Schwaegr.) Mitt.? Hesselbo, A. Diluvial
Mosrester fra Diluviet ved Skaerumhede. [Dan-
marks geol. Undersøgelse, II Raekke, No. 25,
p. 108.]

Orthothecium intricatum (Hartm.) Bry. eur.

- 1912 *Orthothecium intricatum*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX,
p. 90.]

Orthothecium rufescens (Dicks.) Bry. eur.

- 1912 *Orthothecium rufescens*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Pylaisia Bry. eur.

Pylaisia polyantha (Schreb.) Bry. eur.

- 1913 *Pylaisia polyantha*. Beyle, M. Ueber einige Ab-
lag. fossiler Pflanz. der Hamburger Gegend. Erster
Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6,
Beih., p. 83.] Glacial

Leskeaceae.

Heterocladium Bry. eur.

Heterocladium squarrosum (Voit) Lindb.

- 1914 *Heterocladium squarrosum*. Zmuda, A. J. Fossile Diluvial
Flora des Krakauer Diluviums. [Bull. int. Acad.
Sci. Cracovie d. sc. math. et nat., Sér. B, No.
2 B, p. 249.]

Heterocladium species.

- 1908 *Heterocladium* sp. Engelhardt, N., and Kinkelin, F. Oberpliocäne Fl. und Fauna des Frankfurter Klärbeckens. [Abh. Senckenb. naturforsch. Ges., XXIX, p. 189, t. 22, figs. 24—27.]

Note. Weyland [Senckenbergiana, VII, Heft 1/2, p. 13] states that no material of this was available. The figure of the leaf showing nerve reaching nearly to apex seems to preclude *Heterocladium*, though the cell stucture is not inconsistent with this genus.

Anomodon Hook. et Tayl.**Anomodon longifolius** (Schleich.) Bruch.

- 1923 *Anomodon longifolius*. Baumberger, E., etc. Die Quaternary diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]

Anomodon viticulosus (L.) Hook. et Tayl.

- 1905 *Anomodon viticulosus*. Neuweiler, E. Die prähistor. Pleistocene Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 37.]

- 1907 *A. viticulosus*. Dixon, H. N., in Lewis, F. J. Peat Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.]

- 1909 *A. viticulosus*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]

- 1914 *A. viticulosus*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

- 1923 *A. viticulosus*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]

- 1925 *A. viticulosus*. Weyland, H. Beitr. zur Kenntn. fossiler Moose. I. [Senckenbergiana, VII, Heft 1/2, p. 12, Fig. 4.]
= *Anomodon* sp. Engelhardt, H. & Kinkelin, F. Oberpliocäne Fl. und Fauna... des Frankfurter Klärbeckens. [Abh. Senckenb. naturforsch. Ges., XXIX, p. 189, t. 22, Figs. 16, 17.]

Note. As Weyland remarks, there is nothing to preclude this being *A. viticulosus*.

Lescuraea Bry. eur.**Lescuraea striata** (Schwaegr.) Bry. eur.

- 1912 *Lescuraea striata*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Leskea Hedw.**Leskea polycarpa** Ehrh.

- 1898 *Leskea polycarpa*. Andersson, Gunnar. Stud. öfver Quaternary

Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.]

Leskea species.

- 1908 *Leskea* sp. Engelhardt, H. & Kinkelin, F. Ober-pliocäne Fl. und Fauna des Frankfurter Klär-beckens. [Abh. Senckenb. naturforsch. Ges., XXIX, p. 188, t. 22, figs. 28, 29.] Upper Pliocene

Note. The determination as a *Leskea* cannot be justified; the upper half of the leaf shows no nerve. Weyland, H. (Beitr. zur Kenntn. fossiler Moose, in Senckenbergiana, VII, Heft 1/2) refers to this, and figures it (Fig. 6), remarking that in the condition of the fragment it must be left indeterminate.

Pseudoleskea Bry. eur.

Pseudoleskea filamentosa (Dicks.) Broth.

- 1912 *Pseudoleskea atrovirens*. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Pseudoleskea atrovirens (Dicks.) (P. patens Limpr.)

- 1915 *Pseudoleskea atrovirens*. Reid, Clement & E. M. Pliocene The Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, p. 52].

Note. A slight doubt attaches to the determination, owing to the remains being associated with such species as *Eurhynchium speciosum* Schimp., *Philonotis fontana* etc.; the objection being somewhat neutralized by the presence also of *Homalothecium sericeum* and *Ditrichum tortile*. Structurally the leaves quite agreed.

Thuidium Bry. eur.

Thuidium abietinum (Dill.) Bry. eur.

- 1903 *Thuidium abietinum*. Range, Paul. Das Diluv. Diluvial von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 269.]

- 1911 *T. abietinum*. Stoller, J. Die Flora der jung-glaziaalen Ablag. Ostpreussens. [Jahrb. d. Kgl. preuss. geol. Landesanst., XXXI, T. II, H. 1, p. 123.] Glacial

- 1914 *T. abietinum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial

- 1918—20 *T. abietinum*. Docturovsky, V. S., & Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Glacial

Thuidium antiquum Schimp.

- 1865 *Thuidium antiquum* Schimp. in Heer, Osw. Die Quaternary Urwelt der Schweiz, 1 Aufl., p. 497. lignites of Note. Schimper (Traité de Pal. Végét., I. Switzerland

p. 253) refers to this — without naming it — as apparently identical with *T. delicatulum*. Later, in Heer's work, he gives it the above name, and refers to it as being very close to *T. delicatulum*.

Thuidium delicatulum (Dill.) Mitt.

- | | | |
|---------|--|------------------------------------|
| 1869 | <i>Thuidium delicatulum</i> . Schimper, W. P. Traité de Pal. végétale, I, p. 253 (cf. <i>T. antiquum</i>). | Quaternary lignites of Switzerland |
| 1910 | <i>T. delicatulum</i> . Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.] | Postglacial |
| 1910 | <i>T. delicatulum</i> (or <i>T. recognitum</i> Lindb.). Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] | Diluvial |
| 1911 | <i>T. delicatulum</i> . Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands. [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXIX, p. 135.] | Diluvial |
| 1914 | <i>T. delicatulum</i> . Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] | Diluvial |
| 1918—20 | <i>T. delicatulum</i> . Docturovsky, V. S., & Kudriashchev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] | Postglacial |
| 1923 | <i>T. delicatulum</i> . Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.] | Quaternary |

Thuidium elatinum (Sap.) Schimp., Traité de pal. végétale, I, p. 250 (1869); t. VI, fig. 22—26.

- | | | |
|------|---|----------|
| 1865 | <i>Muscites elatinus</i> Saporta G. Etudes sur la vég. du Sud-Est de la France. [Ann. des Sc. nat. Par., 5 ^e sér., IV, p. 33; t. I, f. 2.] | Tertiary |
|------|---|----------|
- Note. Schimper places this "entre *T. abietinum* et *T. delicatulum*". The figures however show a clearly bipinnate branching, which precludes *T. abietinum*. The stem-leaves if correctly drawn are erect and short-pointed, which would suggest *T. delicatulum* or *T. recognitum*. There is apparently no reason to suppose it is distinct from one of our modern species.

Thuidium oeningense (Braun) Schimp. op. et loc. cit.

- | | | |
|------|--|----------|
| 1851 | <i>Muscites oeningensis</i> Braun Al. Verzeichn. d. foss. Pflanz. von Oeningen. [Stizenberger, Uebersicht d. Verstein. Badens, p. 72]. <i>Hypnum oeningense</i> Schimp. MS. in Heer, O., Flora tert. Helvetiae, I, p. 29; t. III, f. 8; II, t. XCIV, f. 20, e; III, t. CXLII, f. 26, 27. | Tertiary |
|------|--|----------|
- Note. Schimper states that this resembles *T. minutulum*. A specimen of the plant figured by Heer (I, t. III, f. 8) is in the Geological Department of the British Museum (No. 40884). The stem is distinctly pinnate, and it may well be a *Thuidium*, but it need by no means be compared with *T. minutulum*.

Thuidium Philiberti Limpr.

- 1905 *Thuidium Philiberti*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas.... [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]
 1910 *T. Philiberti*. Dixon, H. N. Some "Neolithic" Postglacial Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 104.]
 1923 *T. Philiberti*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]

var. pseudo-tamarisci (Limpr.) Ryan & Hagen

- 1905 *Thuidium pseudo-tamarisci*. Neuweiler, E., op. cit., pp. 38, 40.

Thuidium recognitum Lindb. cf. *T. delicatulum*, 1910.**Thuidium tamariscinum** (Hedw.) Bry. eur.

- 1905 *Thuidium tamariscinum*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]
 1908 *T. tamariscinum*. Pax, F. Grundzüge der Pflanzenverbreitung in den Karpathen, II, p. 44.
 1909 *T. tamariscinum*. Hartz, N. Bidrag til Danmarks tertiaære.... Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
 1910 *T. tamariscinum*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist. Soc., 1910, p. 104.]
 1913 *T. tamariscinum*. Beyle, M. Ueber einige Ablagerung.... der Hamburger Gegend. Erster Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6 Beih., p. 83.]
 1920 *T. tamariscinum*. Reid, E. M. Two pre-glacial Floras from Castle Eden. [Quart. Journ. Geol. Soc., LXXVI, p. 111.]
 1923 *T. tamariscinum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.]

Thuidium species.

- 1912 *Thuidium sp.* Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.]
 1912 *Thuidium sp.* Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Helodium (Sull.) Warnst.**Helodium lanatum** (Stroem) Broth.

- 1907 *Thuidium Blandovii* (Web. & Mohr) Bry. eur. Peat Dixon, H. N., in Lewis, F. J. Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.]
 1909 *Thuidium Blandovii*. Hartz, N. Bidrag til Danmarks tertiaære Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]

- 1910 *Thuidium lanatum* (Stroem) Hagen. Hesselbo, A. Diluvial
Mosrester fra Diluviet ved Skaerumhede. [Dan-
marks geol. Undersøgelse, II Raekke, No. 25,
p. 108.]
- 1914 *Helodium Blandovii* (Web. & Mohr). Zmuda, A. J. Diluvial
Fossile Flora des Krakauer Diluviums. [Bull. int.
Acad. Sci. Cracovie d. sc. math. et nat., Sér. B,
No. 2 B, p. 249.]
- 1918—20 *Helodium lanatum*. Docturowsky, V. S., and Postglacial
Kudriaspach, V. Results of Russian studies on
peat moors. [Reports on Peat, 4.]

Hypnaceae.

Amblystegium Bry. eur.

Amblystegium Kochii Bry. eur.

- 1904 *Amblystegium Kochii*. Bell, A. M. Implementi-
ferous Sections at Wolvercote (Oxfordshire). Late glacial
[Quart. Journ. Geol. Soc., LX, p. 124.]
- 1920 *A. Kochii*. Reid, E. M. Two pre-glacial Floras Pliocene
from Castle Eden. [Quart. Journ. Geol. Soc.,
LXXVI, p. 111.]
- 1924 *A. Kochii*. Beyle, M. Ueber einige Ablagerung... Pleistocene
der Hamburger Gegend. Dritter Teil. [Mitteil.
aus dem Mineralog.-geol. Staatsinstitut, Heft VI,
p. 26.]

Amblystegium riparium Bry. eur.

- 1912 *Amblystegium riparium*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlch. Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.]
- 1914 *Leptodictyum riparium* (L.) Warnst. Zmuda, A. J. Diluvial
Fossile Flora des Krakauer Diluviums. [Bull. int.
Acad. Sci. Cracovie d. sc. math. et nat., Sér. B,
No. 2 B, p. 249.]
- 1923 *A. riparium*. Douin, R. Les mousses des
tufs de Lautaret. [Rev. Génér. de Bot., XXXV,
p. 119.]

Amblystegium serpens Bry. eur.

- 1904 *Amblystegium serpens*. Bell, A. M. Implementi-
ferous Sections at Wolvercote (Oxfordshire). Late glacial
[Quart. Journ. Geol. Soc., LX, p. 124.]
- 1912 *A. serpens*. Dixon, H. N., in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *A. serpens*. Weber, C. A. Die Mammuthflora von Glacial
Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1,
p. 19.]
- 1920 *A. serpens*. Reid, E. M. Two pre-glacial Floras Pliocene
from Castle Eden. [Quart. Journ. Geol. Soc.,
LXXVI, p. 111.]

Amblystegium varium (Hedw.) Lindb.

- 1903 *Amblystegium varium*. Range, Paul. Das Diluv. Diluvial von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 253.]
- 1904 *A. varium*. Reid, Clement. East Norfolk Geology. Preglacial [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.]
- 1909 *A. radicale Bry. eur.* Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial

Amblystegium species.

- 1908 *Amblystegium* sp. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård. [Sver. geol. Undersökn. Årsbok, II, No. 2, p. 16.] Late glacial
- 1908 *Amblystegium* sp. Weber, C. A. Die Moostorf-schichten zwischen Sarkau und Cranz. [Engler's Bot. Jahrb., XLII, pp. 42, 43.] Pleistocene
- 1910 *Amblystegium* sp. Sukačev, V. N. Sur la flore arctique fossile près du village Demianskóe, gouv. Tobolsk. [Bull. Acad. Impér. Sci. St. Pétersbourg, 1910, p. 460, fig. g.] Glacial
- 1912 *Amblystegium* sp. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, 116.]
- 1918—20 *Amblystegium* sp. Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Postglacial
- 1923 *Amblystegium* sp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 521.] Quaternary

Amblystegiella Loeske**Amblystegiella confervoides (Brid.) Loeske**

- 1909 *Amblystegium* cfr. *confervoides*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial

Homomallium (Schimp.) Loeske**Homomallium incurvatum (Schrad.) Loeske**

- 1905 *Hypnum incurvatum* Schrad. Neuweiler, E. Die Pleistocene prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]

Hygroamblystegium Loeske**Hygroamblystegium curvicaule (Jur.) Loeske**

- 1914 *Amblystegium curvicaule* Jur. Weber, C. A. Die Glacial Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 19.]

Hygroamblystegium fallax (Brid.) Loeske

- 1903 *Amblystegium fallax* var. *spinifolium* (Schimp.) Diluvial Range, Paul. Das Diluv. von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 253.]

Hygroamblystegium filicinum (L.) Broth.

- 1901 *Amblystegium filicinum* (L.) De Not. Geheeb, A. Ueber ein fossiles Laubmoos aus der Umgebung von Fulda. [Beihefte zum Bot. Centralblatt, Bd. X, Heft 3, p. 125.]
- 1904 *A. filicinum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1908 *A. filicinum*. Holst, N. O. Efterskörd från de senglaciale Lagren vid Toppeladugård. [Sver. geol. Undersökn. Arsbok, II, No. 2, p. 19.]
- 1910 *A. filicinum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]
- 1912 *A. filicinum*, and var. *vallisclausae* (Wils.). Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *A. filicinum*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 19.]
- 1914 *Cratoneuron filicinum* (L.) Roth. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 429.]
- 1915 *A. filicinum*. Reid, Clement, and E. M. The Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, p. 53.]
- 1922 *A. filicinum*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.]
- 1922 *Hypnum filicinum* L. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faærerne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.]
- 1923 *A. filicinum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335, 521.]

Cratoneuron (Sulliv.) Roth**Cratoneuron commutatum (Hedw.) Roth**

- 1890 *Hypnum commutatum* Hedw. Dawson, Sir Wm., and Penhallow, D. P. On the Pleistocene Flora of Canada. [Bull. Geol. Soc. Amer., I, p. 315.]
- 1900 *H. commutatum*. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley. [Rep. of 70th Meeting of Brit. Assoc., Bradford. Pt. II, p. 335.]
- 1904 *Cratoneuron commutatum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1908 *H. commutatum*. Augé, L. Note sur tufs quaternaires de Piécourt. [Bull. Soc. Etudes Sc. nat. Nîmes, XXXV, p. 67.]
- 1909 *H. commutatum*. Schuster, J. Paläobot. Not. aus Bayern. [Ber. bayer. bot. Ges., XII, 1, p. 58.]

- 1920 *C. commutatum*. Baker, Frank C. The Life of Pleistocene the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 332, 335, 377.]
- 1923 *C. commutatum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 195, 335, 521.]
- 1923 *C. cfr. commutatum*. Baumberger, op. cit., p. 145, 335.]
- 1923 *C. commutatum*. Klähn, Hans. Die Petrogenese der Kalktuffe [Geol. Arch., 2, p. 305.]
- 1923 *H. commutatum*. Douin, R. Les mousses des tufs de Lautaret. [Rev. Génér. de Bot., XXXV, p. 120.]

Cratoneuron decipiens (De Not.) Broth.

- 1904 *Thuidium decipiens* De Not. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1910 *Th. decipiens*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undsøgelse, II Raekke, No. 25, p. 108.]

Cratoneuron falcatum (Brid.) Roth

- 1904 *Hypnum falcatum* Brid. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1910 *H. falcatum*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
- 1922 *H. falcatum*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.]
- 1923 *Cratoneuron falcatum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335.]
- 1923 *H. falcatum*. Douin, R. Les mousses des tufs de Lautaret. [Rev. Génér. de Bot., XXXV, p. 120.]

Cratoneuron irrigatum (Zett.) Roth

- 1912 *Hypnum irrigatum* Zett. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. Naturforsch. Ges. Freiburg i. B., XIX, p. 90.]
- 1923 *H. irrigatum*. Douin, R. Les mousses des tufs de Lautaret. [Rev. Génér. de Bot., XXXV, p. 120.]

Cratoneuron species.

- 1892 *Hypnum filicinum* or *commutatum*. Boulay, N. Pliocene Flore pliocène du Mont-Dore, p. 45.

Drepanocladus Roth (Hypnum § Harpidium auct.)

Drepanocladus aduncus (Hedw.) Moenk.

- 1869 *Hypnum aduncum* Hedw. Schimper, W. P. Traité de Pal. Végétale, I, p. 254.

- 1903 *H. aduncum* var. *sparsifolium* (Warnst.). Range, Diluvial
Paul. Das Diluvialgebiet von Lübeck und seine
Dryastone.... [Zeitschr. für Naturwiss., LXXVI,
p. 253.]
- 1904 *Drepanocladus aduncus*. Bell, A. M. Implementi-
ferous Sections at Wolvercote (Oxfordshire.)
[Quart. Journ. Geol. Soc., LX, p. 124.]
- 1904 *D. aduncus*? Weber, C. A. in Mueller, G. Ueber
eine fröhldiluviale Flora bei Lüneburg. [Abh.
Kgl. preuss. geol. Landesanstalt und Bergakad.
Heft 40, p. 16.]
- 1908 *H. aduncum*. Weber, C. A. Die Moostorfschicht.
.... zwischen Sarkau und Cranz. [Engl. Bot.
Jahrb., XLII, pp. 40, 43.]
- 1908 *H. aduncum*. Pax, F. Grundzüge der Pflanzen-
verbreitung in den Karpathen, II, p. 44.
- 1909 *H. aduncum*. Travis, W. G. Plant remains in Peat
peat at Aintree. [Trans. Liverpool Bot.
Soc., I, p. 5.]
- 1909 *H. aduncum*. Schuster, J. Paläobot. Not. aus Schiefer-
Bayern. [Ber. bayer. bot. Ges., XII, pp. 1, 58.] kohlen
- 1910 *Amblystegium aduncum* (Hedw.) Sanio. Hesselbo,
A. Mosrester fra Diluviet ved Skaerumhede. [Dan-
marks geol. Undersøgelse, II Raekke, No. 25,
p. 108.]
- 1912 *H. aduncum*. Dixon, H. N., in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1912 *Drepanocladus subaduncus* Warnst. Szafer, W. Eine Diluvial
Dryasflora bei Krystynopol in Galizien. [Bull.
int. Acad. Sci. Cracovie d. sc. math. et nat.,
Sér. B, No. 2, p. 1107.]
Note. The determination is said to be slightly doubtful.
- 1919 *D. aduncus*. Gadeceau, E. Les Forêts submergées Postglacial
de Belle-Ile-en-mer. [Bull. biol. de la France,
etc., LIII, p. 295.]
- 1920 *D. aduncus*. Baker, Frank C. The life of the Pleistocene
Pleistocene or glacial period. [Univ. of Illinois
Bull., XVII, pp. 245, 280, 377.]
- 1922 *H. aduncum*. Travis, W. G. On peaty bands in Peat
the Wallasey sandhills. [Proc. Liverpool Geol.
Soc., III, p. 209.]

***Drepanocladus aduncus* (Hedw.) Gr. Kneiffii**

- 1852 *Hypnum Noegerrathii* Weber C. O. Die Tertiär- Tertiary
flora der niederrhein. Braunkohlenformation. [Pa-
laeontographica, II, p. 227, t. XXV.]
Note. Schimper (Traité de Pal. Végét., I, 250)
refers this to *Hyp. aduncum* var. *Kneiffii*.
- 1896 *Hypnum aduncum* var. *groenlandicum* Heer. Engel,
Th. Geognost. Wegweiser durch Württemberg.
2^e Aufl., p. 436.]
- 1905 *H. aduncum* var. *Kneiffii groenlandicum*. Neuweiler, Pleistocene
E. Die prähistor. Pflanzenreste Mitteleuropas ...
[Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6,
p. 39.]
- 1909 *Amblystegium aduncum* (Hedw.) Sanio ? var. Interglacial
Kneiffii. Hartz, N. Bidrag til Danmarks ter-

- tiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
 1909 *Hypnum* cfr. *Kneiffii*. Schuster, J. Paläobot. Not. Schiefer-
 aus Bayern. [Ber. bayer. bot. Ges., XII, pp. 1, 58.] kohlen
 1910 *Amblystegium Kneiffii* Bry. eur. Hesselbo, A. Diluvial
 Mosrester fra Diluviet ved Skaerumhede. [Dan-
 marks geol. Undersøgelse, II Raekke, No. 25,
 p. 108.]
 1919 *Drepanocladus aduncus* var. *Kneiffii*. Gadeceau, E. Postglacial
 Les Forêts submergées de Belle-Ile-en-mer. [Bull.
 biol. de la France, etc., LIII, p. 295.]

Drepanocladus aduncus* (Hedw.) Gr. *pseudofluitans

- 1904 *D. aduncus* var. *pseudofluitans* (Sanio). Bell, A. M. Late glacial
 Implementiferous Sections at Wolvercote (Ox-
 fordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
 1909 *Hypnum aduncum* var. *paternum* Sanio. Travis, Peat
 W. G. Plant remains in peat at Aintree.
 [Trans. Liverpool Bot. Soc., I, p. 5.]
 1909 *Drepanocladus pseudofluitans* (Sanio). Stoller, J. Early
 Die Pflanzenreste in den Stuttgarter An-
 lagen. [Mitteil. der geol. Abt. des k. württ.
 stat. Landes-Amtes, No. 6. Beilage zu Jahresh. des
 Ver. für vaterl. Naturkunde in Würtemberg,
 Jahrg. LXV, p. 73, fig. 1.]
 1912 *H. aduncum* var. *pseudofluitans* Sanio. Dixon, H. Late glacial
 N., in Warren, S. H. A late Glacial stage in the
 Lea Valley. [Quart. Journ. Geol. Soc., LXVIII,
 p. 231.]
 1919 *D. pseudofluitans* (Sanio). Gadeceau, E. Les Postglacial
 Forêts submergées de Belle-Ile-en-mer. [Bull. biol.
 de la France, etc., LIII, p. 295.]

***Drepanocladus badius* (Hartm.) Roth**

- 1914 *Drepanocladus badius*. Zmuda, A. J. Fossile Flora Diluvial
 des Krakauer Diluviums. [Bull. int. Acad. Sci.
 Cracovie d. sc. math. et nat., Sér. B, No. 2 B,
 p. 250.]

***Drepanocladus capillifolius* Warnst.**

- 1904 *Drepanocladus capillifolius*. Bell, A. M. Implemen- Late glacial
 tiferous Sections at Wolvercote (Oxfordshire).
 [Quart. Journ. Geol. Soc., LX, p. 124.]
 1904 *Hypnum capillifolium* Warnst. Reid, Clement. East Preglacial
 Norfolk Geology. [Trans. Norf. & Norw. Nats'.
 Soc., VII, p. 293.]
 1910 *Amblystegium capillifolium* (Warnst.). Hesselbo, A. Diluvial
 Mosrester fra Diluviet ved Skaerumhede. [Dan-
 marks geol. Undersøgelse, II Raekke, No. 25,
 p. 108.]
 1911 *D. capillifolius*. Zmuda, A. J., in Szafer, W. Early
 Ueber eine alt-diluviale Flora in Krystynopol in diluvial
 Wolhynien. [Kosmos, XXXVI, p. 337.]
 1912 *D. capillifolius*. Szafer, W. Eine Dryasflora bei
 Krystynopol in Galizien. [Bull. int. Acad. Sci.

- Cracovie d. sc. math. et nat., Sér. B. No. 2,
p. 1107.]
- 1912 *H. capillifolium*. Dixon, H. N., in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1914 *D. capillifolius*. Zmuda, A. J. Fossile Flora des Diluvial
Krakauer Diluviums. [Bull. int. Acad. Sci. Cra-
covie d. Sc. math. et nat., Sér. B, No. 2B,
p. 250.]
- 1915 *H. capillifolium*. Reid, Clement and E. M. The Pliocene
Pliocene Floras of the Dutch-Prussian Border.
[Med. van de Rijksopspor. van Delfstoffen, No. 6,
p. 53.]

Drepanocladus exannulatus (Guemb.) Warnst.

- 1896 *Amblystegium exannulatum* (Guemb.) De Not. Pleistocene
Andersson, Gunnar. Ueber das Fossile Vorkommen clay
..... in Russland und Dänemark. [Bih. Svensk
Vet.-Akad. Handl., XXII, Afd. III, No. 1, p. 6.]
- 1902 *A. exannulatum*. Hartz, N. Bidrag til Danmarks Late glacial
senglaciiale Fl. og Fauna. [Danmarks geol. Un-
dersøgelse, II Raekke, No. 11, p. 15.]
- 1903 *Hypnum fluitans* var. *brachydictyon* Ren. Holzinger, J. M. Some Fossil Mosses. [Bryologist, VI,
p. 93.]
- 1904 *Drepanocladus exannulatus*. Weber, C. A., in Muel- Early glacial
ler, G. Ueber eine frühdiluviale Flora bei
Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt
und Bergakad., Heft 40, p. 16.]
- 1904 *H. exannulatum*. Bell, A. M. Implementiferous Late glacial
Sections at Wolvercote (Oxfordshire). [Quart.
Journ. Geol. Soc., LX, 124.]
- 1905 *H. fluitans-exannulatum*. Neuweiler, E. Die prähistor. Pleistocene
Pflanzenreste Mitteleuropas (Vier-
teljahr. naturforsch. Ges. Zürich, L, Heft 6,
p. 39.)
- 1907 *H. exannulatum* var. *brachydictyon* Ren. Dixon, H. Peat
N., in Lewis, F. J. Summary of Progress for
1907, p. 99. [Mem. Geol. Survey.]
- 1908 *H. exannulatum*. Weber, C. A. Die Moostorfsschichten. Peat
..... zwischen Sarkau und Cranz. [Engl.
Bot. Jahrb., XLII, pp. 33, 40, 43.]
- 1909 *Amblystegium exannulatum*. Hartz, N. Bidrag til Diluvial
Danmarks tertiaere Flora. [Danmarks geol.
Undersøgelse, II Raekke, No. 20, p. 260.]
- 1910 *A. exannulatum*. Hesselbo, A. Mosrester fra Dilu- Diluvial
viet ved Skaerumhede. [Danmarks geol. Under-
søgelse, II Raekke, No. 25, p. 108.]
- 1912 *H. exannulatum*. Dixon, H. N., in Warren, S. H. Late glacial
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.]
- 1912 *H. exannulatum* var. *orthophyllum* Milde. Dixon, Late glacial
H. N., in Warren, S. H., op. et loc. cit.
- 1912 *H. purpurascens* Limpr. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]
- 1912 *H. fluitans* Gr. *exannulatum* var. *pinnatum* Boul. f. Preglacial
Hawkesworth, E. A pre-glacial Lake-bed near
Northallerton. [Naturalist, 1912, p. 204.]

- 1914 *H. exannulatum*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 23.] Glacial
- 1914 *Warnstorffia exannulata* (Guemb.) Loeske. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1918 *D. exannulatus*. Heim, Arnold, & Gams, H. Inter-glaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXVIII, p. 27.] Interglacial
- 1918—20 *D. exannulatus*. Docturowsky, V. S., & Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Interglacial
- 1919 *D. exannulatus*. Gadeceau, E. Les Forêts submergées de Belle-Ile-en-mer. [Bull. biol. de la France, etc., LIII, p. 295.] Postglacial
- 1922 *H. exannulatum*. Jessen, Knud, og Rasmussen, R. Et Provil Gennem in Tørvemose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] Recent
- 1923 *D. exannulatus*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 521.] Quaternary

Drepanocladus fluitans (L.) Gr. Rotae (De Not.)

- 1903 *H. fluitans* var. *glaciale* Ren. Holzinger, J. M. Some Fossil Mosses. [Bryologist, VI, p. 94.] Pleistocene; below Kansas Drift
- 1904 *Hypnum fluitans* var. *falcifolium* Ren. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1908 *Drepanocladus fluitans* Gr. Rotae. Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands... [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXIX, p. 119.] Diluvial
- 1909 *Amblystegium Rotae* De Not. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Diluvial
- 1910 *Amblystegium Rotae*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *H. exannulatum* Gr. Rotae. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *H. purpurascens* Limpr. var. *Rotae* (De Not.). Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 23.] Glacial

Drepanocladus fluitans (L.) Warnst.

- 1869 *Hypnum fluitans* L. Schimper, W. P. Traité de Pal. Végét., I, p. 254. Glacial deposits in Württemberg
- 1888 *Amblystegium fluitans* (L.) De Not. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from.... Hoxne, in Suffolk. [Geol. Mag., Decade III, Vol. V, p. 444.] Glacial

- 1889 *A. fluitans*. Candler, Chas. Observations on some undescribed Lacustrine Deposits.... in Suffolk. [Quart. Journ. Geol. Soc., XLV, p. 507.]
- 1896 *H. fluitans* var. *tenuissimum* Schimp. Engel, Th. Ceognostisch. Wegweiser durch Württemberg. 2^e Aufl., p. 436.
- 1898 *A. fluitans*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome II, No. 8, p. 135.]
- 1900 *Hypnum fluitans*. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley. [Rep. of 70th Meeting of Brit. Assoc., Bradford, 1900. Pt. II, p. 335.]
- 1901 *H. fluitans* f. *fossilis*. Geheebe, A. Ueber ein fossiles Laubmoos aus der Umgebung von Fulda. [Beih. zum Bot. Centralbl., X, Heft 3, p. 127.]
- 1902 *A. fluitans*. Hartz, N. Bidrag til Danmarks sen-glaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 18.]
- 1904 *H. fluitans*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1904 *H. fluitans*. Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Natts. Soc., VII, p. 293.]
- 1905 *H. fluitans* var. *tenuissimum* Schimp. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 39.]
- 1908 *Drepanocladus fluitans*. Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXIX, p. 119.]
- 1908 *Amblystegium fluitans*. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård. [Sver. Geol. Undersökning Arsbok, II, No. 2, p. 19 bis.]
- 1909 *A. fluitans*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
- 1909 *H. fluitans*. Schuster, J. Paläobot. Not. aus Bayern. [Ber. bayer. bot. Ges., XII, 1, p. 58.]
- 1910 *H. fluitans*. Dixon, H. N. Some "Neolithic" Moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
- 1910 *A. fluitans*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]
- 1911 *D. fluitans*. Stoller, J. Beitr. zur Kenntn. der diluvialen Flora Norddeutschlands. [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXXII, T. 1, H. 1, p. 135.]
- 1911 *A. fluitans*. Stoller J., op. cit., p. 130.
- 1912 *H. fluitans*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1912 *H. fluitans* var. *falcatum* Schimp. Dixon, H. N., op. et loc. cit.
- 1912 *H. fluitans*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.]

- 1914 *H. fluitans*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 23.] Glacial
- 1914 *D. fluitans* var. *pseudostramineus* (C. M.) (as *D. pseudostramineus* (C. M.) Roth). Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] Diluvial
- 1917 *Hypnum* cfr. *fluitans*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons.... [Bot. Jahrb. Syst., LIV, Beibl. No. 120, p. 14.] Interglacial
- 1918 *D. fluitans*. Heim, Arnold, & Gams, H. Inter-glaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXVIII, p. 27.] Interglacial
- 1918—20 *D. fluitans*. Docturowsky, V. S., and Kudriashchev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Postglacial
- 1919 *D. fluitans*. Gadeceau, E. Les Forêts submergées de Belle-Ile-en-mer. [Bull. biol. de la France, etc., LIII, p. 295.] Postglacial
- 1920 *D. fluitans*. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 332, 335, 377.] Pleistocene
- 1923 *D. fluitans*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotech. Ser., Lief. 8, pp. 84, 336.] Quaternary

Drepanocladus Hollosianus (Schillb.)

- Györrfy MS., Broth. in Engl. & Prantl, Pflanzen-fam., Musci, Ed. 2, II, p. 523.
 Syn. *Hypnum* sp.? Hollós, L., in Kecszemét Multja és Jelene, 1896, p. 25, fig. 22.
Hypnum Hollósi Schilb. in Term. tud. Közlöny, XXX, p. 436; Budapest, 1898 (nomen solum).
 1912 *Hypnum Hollósianum* Schilb. in Math. és term. tud Ertesito, XXX, p. 636 (cf. pp. 646—649); figs. 1, 3, 4, 6, 8, 9.

Note. Full descriptions and figures are given in the 1912 paper. The author compares it with *Acrocladium cuspidatum* and *Hyp. Taramellianum* Farneti (proximum).

Drepanocladus intermedius (Lindb.) Warnst.

- 1898 *Amblystegium intermedium* (Lindb.) Lindb. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome II, No. 8, p. 135.] Quaternary
- 1904 *Hypnum intermedium* Lindb. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1904 *H. intermedium*. Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 193.] Preglacial
- 1905 *H. intermedium*. Neuweiler, E. Zur Interglazial-flora der schweiz. Schieferkohlen. [Ber. d. schweizerisch. bot. Ges., Heft XV. Beilage. Neunter Ber. d. zürch. bot. Ges., p. 98.] Interglacial
- 1909 *H. intermedium*. Schuster, J. Paläobot. Not. aus Bayern. [Ber. bay. bot. Ges., XII, 1, p. 58.] Schiefer-kohlen

- 1910 *A. intermedium*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1911 *H. intermedium*. Stoller, J. Die Flora der jungglaziaalen Ablag. Ostpreussens. [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXXI, II, p. 130.] Early glacial
- 1912 *H. intermedium*. Whitehead, Goodchild, and Reid. Some notes on "moorlog"..... [Essex Naturalist, XVI, p. 56.] Submarine peat deposit.
- 1914 *H. intermedium*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 21.] Glacial
- 1914 *Limprechtia intermedia* (Lindb.) Loeske. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 249.] Diluvial
- 1918 *Drepanocladus intermedius*. Heim, Arnold, & Gams, H. Interglaz. Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXVIII, p. 27 bis.] Interglacial
- 1920 *D. intermedius*. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 332, 335, 377.] Pleistocene
- 1922 *H. intermedium*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
- 1923 *D. intermedius*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 84, 145, 335, 336, 521.] Diluvial

***Drepanocladus latifolius* (Lindb. et Arn.) Broth.**

- 1910 *Amblystegium latifolium* (Lindb. & Arn.) Lindb. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial

***Drepanocladus lycopodioides* (Schwaegr.) Warnst.**

- 1904 *Hypnum lycopodioides* Schwaegr.? Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1914 *H. lycopodioides*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 22.] Glacial
- 1922 *H. lycopodioides*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat

***Drepanocladus revolvens* (Sw.) Warnst.**

- 1896 *Hypnum revolvens* Sw. Farneti, R. Ricerche di briol. palaeont al periodo glaciale. [Atti del Istit. Bot. della Reale Univ. d. Pavia, Ser. 2, V, p. 54.] Glacial
- 1890 *H. revolvens*. Dawson, Sir Wm., and Penhallow, D. P. On the Pleistocene Flora of Canada. [Bull. Geol. Soc. Amer., I, p. 315.] Pleistocene
- 1900 *H. revolvens*. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley. [Rep. of 70th Meeting, Brit. Assoc., Bradford. Pt. II, p. 336.] Pleistocene

- 1903 *H. revolvens*, Range, Paul. Das Diluv. von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 254.] Diluvial
- 1903 *H. revolvens*. Holzinger, J. M. Some Fossil Mosses. [Bryologist, VI, p. 94.] Pleistocene, below Kansas drift.
- 1904 *H. revolvens*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1904 *H. revolvens*. Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.] Preglacial
- 1909 *Amblystegium Cossoni* (Schreb.) Lindb. Nørregaard, E. M. Et senglaciale fra Dejbjerg Bakker. [Med. fra Dansk geol. Fören., No. 15, p. 326.] Late glacial
- 1910 *A. Cossoni*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *H. revolvens*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Flora und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] Glacial
- 1912 *H. Cossoni*. Stark, P., op. cit., p. 116. Glacial
- 1912 *H. revolvens*, Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *H. revolvens*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 22.] Glacial
- 1915 *H. revolvens sensu latiore*. Camus, F. Sur les mousses trouvées dans un mammouth. [Compt. Rend. Acad. Sci. Paris, CLX, p. 842.] Glacial
- 1917 *Hypnum cfr. revolvens*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. No. 120, p. 14.]
- 1918 *Drepanocladus revolvens*. Heim, Arnold, & Gams, H. Interglaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXIII, p. 27.] Interglacial
- 1920 *D. revolvens*. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 178, 184, 228, 241, 377.] Pleistocene
- 1922 *H. revolvens*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
- 1923 *D. revolvens*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335, 521.] Quaternary

Drepanocladus scorpioides (L.) Warnst.

- 1896 *Hypnum scorpioides* L. Farneti R. Ricerche di briol. palaeont al periodo glaciale. [Atti del Istit. Bot. della Reale Univ. d. Pavia, Ser. 2, V, p. 55.] Glacial
- 1903 *H. scorpioides*. Range, Paul. Das Diluv. von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 269.] Diluvial
- 1904 *H. scorpioides*. Geinitz, C., und Weber, C. A. Ueber ein Moostorflager der Rostocker Heide. [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.] Postglacial

- 1908 *Scorpidium scorpioides* (L.) Limpr. Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, pp. 41, 44.] Pleistocene
- 1908 *Drepanocladus scorpioides*. Weber, C. A. Hypnum turgescens Schimp. [Engl. Bot. Jahrb., XLII, p. 239.] Pleistocene
Note. This is a correction of *D. turgescens* to the present species, from p. 38 of the same volume.
- 1908 *Amblystegium scorpioides* (L.) Lindb. Holst, N.O. Efterskörd från de senglaciala Lagren vid Toppeledugård [Sver. geol. Undersökn. Arsbok, II, No. 2, p. 19 bis.] Late glacial
- 1909 *A. scorpioides*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Diluvial
- 1909 *H. scorpioides*. Travis, W. G. Plant remains in peat at Aintree. [Trans. Liverpool Bot. Soc., I, p. 5.] Peat
- 1909 *Sc. scorpioides*. Schuster, J. Paläobot. Not. aus Bayern. [Ber. bay. bot. Ges., XII, 1, p. 58 bis.] Schieferkohlen
- 1910 *A. scorpioides*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1911 *H. scorpioides*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.] Arctic bed in peat, West Shetland
- 1912 *H. scorpioides*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *Sc. scorpioides*. Szafer, W. Eine Dryasflora bei Krystynopol in Galizien. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2, p. 1109.] Glacial
- 1914 *Sc. scorpioides*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] Diluvial
- 1914 *H. scorpioides*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 24.] Glacial
- 1917 *H. scorpioides*. Weber, C. A. Die Pflanzenwelt des Ralutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.] Interglacial
- 1917 *Sc. scorpioides*. Rudolph, K. Untersuch. über den Aufbau böhmischer Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 83.] Recent
- 1918 *Sc. scorpioides*. Heim, Arnold, & Gams, H. Interglaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXIII, p. 27 bis.] Interglacial
- 1922 *H. scorpioides*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
- 1922 *Sc. scorpioides*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] Recent
- 1923 *Sc. scorpioides*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 146, 337.] Quaternary

Drepanocladus Sendtneri (Schimp.) Warnst.

- 1869 *Hypnum Sendtneri* Schimp. Traité de Pal. végét., Tertiary I, p. 250.
Syn. *H. Weberianum* Goepp. ex Weber, C. Otto.
Die Tertiärfloren der niederrhein. Braunkohlen-
formation, in Palaeontographica, II, p. 227, t.
XXV, f. 7, 10—13. (nec *H. Weberianum* Schimp.).
- 1896 *H. Sendtneri*. Farneti, R. Ricerche di briol. pa-
laeont al periodo glaciale. [Atti del Istit.
Bot. della Reale Univ. d. Pavia, Ser. 2, V, p. 55.] Glacial
- 1904 *H. Sendtneri* (with some slight doubt). Bell, A. M.
Implementiferous Sections at Wolvercote (Ox-
fordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1904 *H. Sendtneri* (or ? *H. Wilsoni*). Reid, Clement.
East Norfolk Geology. [Trans. Norf. & Norw.
Nats. Soc., VII, p. 293.] Preglacial
- 1905 *H. Sendtneri*. Neuweiler, E. Die prähistor. Pflan-
zenreste Mitteleuropas [Vierte Jahr. natur-
forsch. Ges. Zurich, L, Heft 6, p. 39.] Pleistocene
- 1905 *H. Sendtneri*? Neuweiler, E. Zur Interglazialflora
der schweiz. Schieferkohlen. [Ber. d. schweize-
risch. bot. Ges., Heft XV. Beilage. Neunter Ber.
d. zürcherisch. bot. Ges., p. 98.] Interglacial
- 1909 *H. Sendtneri*. Travis, W. G. Plant remains in peat at Aintree. [Trans. Liverpool Bot. Soc., I, p. 5.] Peat
- 1910 *Amblystegium Sendtneri* (Schimp.) De Not. Hes-
selbo, A. Mosrester fra Diluviet ved Skaerumhede.
[Danmarks geol. Undersøgelse, II Raekke, No. 25,
p. 108.] Diluvial
- 1912 *H. Sendtneri*. Dixon, H. N., in Warren, S. H.
A late Glacial stage in the Lea Valley. [Quart.
Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *Drepanocladus Sendtneri*. Weber, C. A. Die Mam-
mutflora von Borna. [Abh. Nat. Ver. Bremen,
XXIII, Heft 1, p. 22.] Glacial
- 1922 *H. Sendtneri*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol.
Soc., III, p. 209.] Peat
- 1923 *D. Sendtneri*. Baumberger, E., etc. Die diluv.
Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz,
geotechn. Ser., Lief. 8, pp. 84, 145, 336.] Quaternary
- 1925 *D. Sendtneri*. Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot., LXIII, p. 370.] Interglacial

Drepanocladus trifarius (Web. & Mohr) Broth.

- 1905 *Hypnum trifarium* (Web. & Mohr). Neuweiler, E. Interglacial
Zur Interglazialflora der schweiz. Schieferkohlen.
[Ber. d. schweizerisch. bot. Ges., Heft XV. Bei-
lage. Neunter Ber. d. zürcherisch. bot. Ges.,
p. 97.] Pleistocene
- 1908 *H. trifarium*. Weber, C. A. Die Moostorfschich-
ten zwischen Sarkau und Cranz. [Engl. Bot.
Jahrb., XLII, pp. 40, 41, 43.] Pleistocene
- 1909 *Calliergon trifarium* (Web. & Mohr) Kindb. Schuster,
J. Paläobot. Not. aus Bayern. [Ber. bay. bot. Ges.,
XII, 1, p. 58.] Schiefer-
kohlen

- 1912 *H. trifarium*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, pp. 90, 116.] Glacial
- 1914 *Pseudocalliergon trifarium* (Web. & Mohr) Loeske. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 250.] Diluvial
- 1917 *H. trifarium*. Rudolph, K. Untersuch. über den Aufbau böhmischer Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 83.] Recent
- 1918 *Calliergon trifarium*. Heim, Arnold, & Gams, H. Interglaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXIII, p. 27.] Interglacial
- 1923 *Calliergon trifarium*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 84, 145, 336, 521.] Quaternary

Drepanocladus Tundrae (Arn.).

- 1911 ?*Calliergon Tundrae* (Arn.). Zmuda, A. J. in Szafer, W. Ueber eine alt-diluviale Flora in Krystynopol in Wolhynien. [Kosmos, XXXVI, p. 338.] Early diluvial

Drepanocladus turgescens (Jens.) Broth.

- 1904 *Hypnum turgescens* Jens. Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.] Preglacial
- 1908 *H. turgescens*. Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, p. 38 sqq.] Pleistocene
Note. This was corrected later, in the same volume, to D. Sendtneri.
- 1909 *Amblystegium turgescens* (Jens.) Lindb. Nørregaard, E. M. Et senglacialt fra Dejbjerg Bakker. [Meddel. fra Dansk geol. Fören., No. 15, p. 325.] Late glacial
- 1910 *A. turgescens*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II. Raekke, No. 25.] Diluvial
- 1912 *H. turgescens*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *Drepanocladus turgescens*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] Glacial
- 1914 *Pseudocalliergon turgescens* (Jens.) Loeske. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 250.] Diluvial
- 1917 *H. turgescens*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. No. 120, p. 14.] Interglacial
- 1923 *Calliergon turgescens* (Jens.) Kindb. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 337, 521.] Quaternary

Drepanocladus vernicosus (Lindb.) Warnst.

- 1904 *Hypnum vernicosum* Lindb. Geinitz, E., und Weber, C. A. Ueber ein Moostorflager der Rostocker Heide. [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.] Postglacial
- 1908 *H. vernicosum?* Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, pp. 40, 41, 43.] Pleistocene
- 1911 **Drepanocladus vernicosus.** Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands. Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXXII, T. 1, H. 1, p. 135.] Diluvial
- 1912 *D. vernicosus.* Szafer, W. Eine Dryasflora bei Krystynopol in Galizien. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2, p. 1106.] Glacial
- 1914 *Limprichtia vernicosa* (Lindb.) Loeske. Zmuda, A. J., Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1914 *H. vernicosum.* Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 21.] Glacial
- 1917 *D. vernicosus.* Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.] Interglacial
- 1922 *H. vernicosum.* Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
- 1925 *D. vernicosus.* Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot., LXIII, p. 370.] Interglacial

Drepanocladus Wilsoni (Schimp.) Roth.

- 1904 *Hypnum Wilsoni* Schimp. ? Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.] Preglacial
Note. Possibly *D. Sendtneri* (Schimp.).
- 1908 **Drepanocladus Wilsoni.** Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands. [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXIX, p. 119.] Diluvial
- 1909 *H. Wilsoni.* Travis, W. G. Plant remains in peat at Aintree. [Trans. Liverpool Bot. Soc., I, p. 5.] Peat
- 1910 *D. Wilsoni.* Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *H. Wilsoni.* Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *H. Wilsoni.* Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 22.] Glacial
- 1918—20 *D. Sendtneri* var. *Wilsoni* (Schimp.). Doc-turowsky, V. S., and Kudriiashev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Interglacial

Drepanocadus, undetermined species.

- 1923 *Drepanocladus* sp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 336.] Quaternary
- 1925 *Drepanocladus* sp. Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot., LXIII, p. 370.] Novaya Zemlya peat

Calliergon (Sull.) Kindb.**Calliergon cordifolium** (Hedw.) Kindb.

- 1902 *Amblystegium cordifolium* (Hedw.) De Not. Hartz, N., Bidrag til Danmarks senglaciale Fl. og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.] Late glacial
- 1904 *Calliergon cordifolium*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1909 *A. cordifolium*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1910 *A. cordifolium*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial

Calliergon diluvianum (Schimp.) Dix., comb. nov.

- 1869 *Hypnum diluvianum* Schimp. Traité de Pal. Végét., I, p. 253. Quaternary lignites of Switzerland
Note. Schimper compares this with *C. giganteum*. Gams considers it may be *C. sarmentosum* (cfr. Baumberger, E., etc., in Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 520).
- 1923 *H. diluvianum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 521.] Quaternary

Calliergon giganteum (Schimp.) Kindb.

- 1898 *Amblystegium giganteum* (Schimp.) De Not. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 135.] Quaternary
- 1902 *A. giganteum*. Hartz, N. Bidrag til Danmarks senglaciale Fl. og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.] Late glacial
- 1903 *Hypnum giganteum* Schimp. Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 256.] Diluvial
- 1904 *Calliergon giganteum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1904 *H. giganteum*. Weber, C. A., in Müller, G. Ueber eine fröhldiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 16.] Preglacial
- 1904 *H. giganteum*. Geinitz, E., und Weber, C. A. Postglacial Ueber ein Moostorflager der Rostocker Heide.

- [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.]
- 1905 *H. giganteum*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 39 bis.] Pleistocene
- 1905 *H. giganteum*, var. Neuweiler, E. Zur Inter-glazialflora der schweizerisch. Schieferkohlen. [Ber. d. schweizerisch. bot. Ges., Heft XV. Beilage. Neunter Ber. d. zürcherisch. bot. Ges., p. 96.] Interglacial
- 1908 *A. giganteum*. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård [Sver. geol. Undersökning Arsbok, II, No. 2, p. 19.] Late glacial
- 1908 *H. giganteum*. Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, pp. 40, 41, 44.] Pleistocene
- 1909 *A. giganteum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Diluvial
- 1909 *C. giganteum*. Schuster, J. Palaeobot. Not. aus Bayern. [Ber. bayer. bot. Ges., XII, 1, p. 58.] Schiefer-kohlen
- 1910 *A. giganteum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1911 *H. giganteum*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.] Recent Ice-land peat
- 1912 *H. giganteum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *C. giganteum*. Szafer, W. Eine Dryasflora bei Krystynopol in Galizien. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2, p. 1108.] Glacial
Note. Zmuda gives as a synonym "*C. Richardsoni* Zmuda (nicht Kindb.) in Kosmos, XXXVI, p. 337."
- 1912 *H. giganteum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *C. giganteum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1914 *H. giganteum*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 24.] Glacial
- 1917 *C. giganteum*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons.... [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.] Interglacial
- 1917 *H. giganteum*. Rudolph, K. Untersuch. über den Aufbau böhmischer Moore. [Abh. k. k. zool.-bot. Ges., Wien, IX, p. 84.] Recent
- 1918 *C. giganteum*. Beyle, M. Ueber einige Ablag. fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. [Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 40.] Glacial
- 1918 *C. giganteum*. Heim, Arnold, & Gams, H. Inter-glaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXIII, p. 27 bis.] Interglacial

- 1922 *H. giganteum*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
- 1923 *C. giganteum*. Baumberger, E., etc., Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 84, 145, 336, 521.] Quaternary
- 1924 *H. giganteum*. Beyle, M. Ueber einige Ablag. fossiler Pflanzen der Hamburger Gegend. Dritter Teil. [Mitteil. aus dem Mineralog-geol. Staats-institut, Heft VI, p. 26.] Glacial

Calliergon Richardsoni (Mitt.) Kindb.

- 1903 *Hypnum Richardsoni* (Mitt.) Lesq. & James. Holzinger, J. M. On some fossil mosses. [Bryologist, VI, p. 94.] Pleistocene, below Kansas drift
- 1904 *H. Richardsoni*? Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.] Preglacial
- 1911 Calliergon Richardsoni. Zmuda, A. J., in Szafer, W. Ueber eine alt-diluv. Flora in Krystynopol in Wolhynien. [Kosmos, XXXVI, p. 337.] Early diluvial
Note. cfr. note on *C. giganteum* (1912, Szafer) above.
- 1912 *H. Richardsoni*. Whitehead, Goodchild, and Reid. Some notes on "moor-log" [Essex Naturalist, XVI, p. 56.] Marine peat deposit
- 1914 *H. Richardsoni*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 24.] Glacial
- 1914 *C. Richardsoni*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1917 *H. Richardsoni*. Weber, C. A. Die Pflanzenwelt des Rabutzer Beckentons [Bot. Jahrb. Syst., LIV, Beibl. Nr. 120, p. 14.] Interglacial
- 1920 *C. Richardsoni*. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 228, 241, 377.] Pleistocene
- 1925 *C. Richardsoni*. Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot., LXIII, p. 370.] Peat in Nova Zemlya

Calliergon sarmentosum (Wahl.) Kindb.

- 1869 *Hypnum sarmentosum* Wahl. Schimper, W. P. Traité de Pal. Végét., I, p. 253, bis. Quaternary lignites, Switzerland; and glacial deposits in Württemberg
- 1888 *Acroceratium sarmentosum* (Wahl.) Mitt. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Decade III, vol. V, p. 444.] Glacial
- 1896 *H. sarmentosum*. Farneti, R. Ricerche di briol. palaeont.... al periodo glaciale. [Atti del Istit. Bot. della Reale Univ. d. Pavia, Ser. 2, V, p. 57.] Glacial

- 1896 *H. sarmentosum*. Engel, Th. Geognost. Wegweiser durch Württemberg. 2^e Aufl., p. 436. Diluvial
- 1912 *H. sarmentosum*? Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *Hypnum cfr. sarmentosum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *Calliergon sarmentosum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1918 *C. sarmentosum*. Heim, Arnold, & Gams, H. Inter-glaziale Bildungen bei Wildhaus. [Vierteljahr. naturforsch. Ges. Zürich, LXIII, p. 28.] Interglacial
- 1923 *C. sarmentosum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 521.] Quaternary

Calliergon stramineum (Dicks.) Kindb.

- 1897 *Calliergon stramineum*. Beck, R. und Weber, C. A. Ueber ein Torflager des sächsischen Erzgebirges. [Zeitschr. d. D. geol. Ges., XLIX, p. 667.] Early diluvial
- 1898 *Amblystegium stramineum* (Dicks.) De Not. Andersson, Gunnar. Stud. öfver Finlands Kvar-tärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 135.] Quaternary
- 1904 *C. stramineum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
- 1907 *Hypnum stramineum* Dicks. Dixon, H. N., in Lewis, F. J. Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.] Peat
- 1908 *A. stramineum*. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård [Sver. geol. Undersökning Arsbok, II, No. 2, p. 19.] Late glacial
- 1910 *A. stramineum*. Hesselbo, A. Mosrester fra Diluviet ved. Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *H. stramineum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1912 *H. stramineum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *C. stramineum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1918—20 *C. stramineum*. Docturowsky, V. S. and Kudriashchev, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Postglacial
- 1922 *C. stramineum*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvmose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.] Recent

Calliergon undetermined species.

- 1869 *Hypnum stramineum* Dicks. or near it. Schimper, Traité de Pal. Végét., I, p. 253.] Quaternary lignites of Switzerland
1905 *Hypnum*, § *Calliergon* sp. Neuweiler, E. Zur Interglazialflora der schweizerisch. Schieferkohlen. [Ber. d. schweizerisch. bot. Ges., Heft XV. Beilage. Neunter Ber. d. zürcherisch. bot. Ges., p. 97.] Interglacial
1923 *Calliergon* sp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 337.] Quaternary

Acrocladium Mitt.**Acrocladium cuspidatum (L.) Lindb.**

- 1888 *Acroceratium cuspidatum* (L.) Mitt. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Decade III, vol. V, p. 444.] Glacial
1898 *Acrocladium cuspidatum*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 194.] Quaternary
1899 *A. cuspidatum*. Bertrand, C. E. Description d'un échantillon de charbon papyracé [Ann. Soc. Géol. du Nord, XXVIII, p. 186.] Pleistocene
1904 *A. cuspidatum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
1909 *A. cuspidatum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
1910 *Hypnum cuspidatum* L. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.] Postglacial
1911 *H. cuspidatum*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831.] Recent. Ice-land peat
1912 *H. cuspidatum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
1912 *H. cuspidatum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
1912 *A. cuspidatum*. Hawkesworth, E. A pre-glacial Lake-bed near Northallerton. [Naturalist, 1912, p. 204.] Preglacial
1922 *H. cuspidatum*. Travis, W. G. On peaty bands in the Wallasey sandhills. [Proc. Liverpool Geol. Soc., III, p. 209.] Peat
1923 *A. cuspidatum*? Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 146, 337, 521.] Quaternary
1924 *H. cuspidatum*. Beyle, M. Ueber einige Ablag. fossiler Pflanzen der Hamburger Gegend. Dritter Teil. [Mitteil. aus dem Mineralog.-geol. Staats-institut, Heft VI, pp. 23, 26.] Glacial

[*Acrocladium squarrosum* (L.) Lindb., Jessen & Rasmussen in Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12 (1922) is an unknown species, and probably a *lapsus calami*].

***Acrocladium* undetermined species.**

- 1908 *Acrocladium* sp. Holst, N. O. Efterskördf från Lateglacial de senglaciala Lagren vid Toppeladugård [Sver. geol. Undersökning Arsbok, II, p. 16.]

***Hygrohypnum* Lindb.**

(*Hypnum* § *Limnobium* auct.).

***Hygrohypnum alpestre* (Sw.) Loeske.**

- 1903 *Hypnum alpestre* Sw. Range, Paul. Das Diluvial- Diluvial gebiet von Lübeck und seine Dryastone [Zeitschr. für Naturwiss., LXXVI, p. 254.]

***Hygrohypnum dilatatum* (Wils.) Loeske.**

- 1910 *Amblystegium dilatatum* (Wils.) Lindb. Hesselbo, Diluvial A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

- 1912 *Hypnum dilatatum* Wils. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

***Hygrohypnum lignitorum* (Schimp.) Dix., comb. nov.**

- 1869 *Hypnum lignitorum* Schimp. Traité de Pal. Végét., I, p. 253.
Note. Neuweiler in Ber. d. Zürcher. bot. Ges. XV, p. 94 (1905) says "Zwischen H. palustre und H. ochraceum stehend". This is in agreement with Schimper's note; but Gams in his work on "Die Schieferkohlen der Schweiz", p. 28, cites Meylan as referring it to *Calliergon trifarium*, which does not seem probable.

***Hygrohypnum molle* (Dicks.) Loeske.**

- 1910 *Amblystegium molle* (Dicks.) Lindb. ? Hesselbo, A. Diluvial Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

- 1912 *Hypnum molle* Dicks. Dixon, H. N., in Warren, Late glacial S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

***Hygrohypnum ochraceum* (Turn.) Loeske.**

- 1869 *Hypnum ochraceum* Turn. Schimper, W. P. Traité de Pal. Végét., I, p. 253. Quaternary lignites,

- 1912 *Hyp. ochraceum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.] Glacial
 1914 *Hygr. ochraceum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 250.] Diluvial

Hygrohypnum palustre (Huds.) Loeske.

- 1905 *Hypnum palustre* Huds. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 39.] Pleistocene
 1910 *Hyp. palustre*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.] Postglacial
 Note. The species is a little doubtful.
 1923 *Hygr. palustre*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 336.] Quaternary
 1923 *Hygrohypnum* cfr. *palustre*. Baumberger, op. cit., pp. 145, 336. Quaternary

Hygrohypnum Smithii (Sw.) Broth.

- 1912 *Hypnum arcticum* Sommerf. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX. p. 90.] Glacial

Campylium (Sull.) Bryhn.

Campylium chrysophyllum (Brid.) Bryhn.

- 1904 *Campylium chrysophyllum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
 1910 *Amblystegium chrysophyllum* (Brid.) De Not. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
 1914 *Hypnum chrysophyllum* Brid. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 19.] Glacial

Campylium helodes (Spr.) Broth.

- 1907 *Hypnum elodes* Spr.? Hartmann, F. Die Fossile Flora von Ingaramsdorf, p. 25. Diluvial
 1911 H. elodes. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.] West Shetland peat, arctic bed.

Campylium insubricum (Farneti) Dix., comb. nov.

- 1896 *Hypnum insubricum* sp. nov. Farneti, R. Ricerche di briol. palaeont al periodo glaciale. [Atti del Istit. Bot. della Reale Univers. di Pavia, Ser. II, vol. V, p. 53, t. IV.] Glacial
 Note. The author refers this to *Campylium*.

Campylium polygamum (Bry. eur.) Bryhn.

- 1904 *Hypnum polygamum* (Bry. eur.) Wils. Reid, Clement. East Norfolk Geology. [Trans. Norf. & Norw. Nats. Soc., VII, p. 293.] Preglacial
- 1905 *H. polygamum*. Neuweiler, E. Zur Interglazialflora der schweizerisch. Schieferkohlen. [Ber. d. schweizerisch. bot. Ges., Heft XV. Beilage. Neunter Ber. d. Zürcherisch. bot. Ges., p. 98.] Interglacial
- 1909 *H. polygamum*. Travis, W. G. Plant remains in peat at Aintree. [Trans. Liverpool Bot. Soc., I, p. 5.] Peat
- 1909 *Amblystegium polygamum* Bry. eur. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial
- 1910 *A. polygamum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *A. polygamum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1914 *A. polygamum*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 20.] Glacial
- 1923 *Campylium polygamum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 335.] Diluvial

Campylium protensum (Brid.) Broth.

- 1902 *Amblystegium protensum* Lindb. Hartz, N. Bidrag til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.] Late glacial
- 1910 *A. protensum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *Hypnum protensum* Brid. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *H. protensum*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 20.] Glacial

Campylium Sommerfeltii (Myr.) Bryhn.

- 1909 *Campylium Sommerfeltii*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Interglacial

Campylium stellatum (Schreb.) Lang. & Jens.

- 1888 *Campylium stellatum*. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Decade III, Vol. V, p. 444.] Glacial
- 1902 *Amblystegium stellatum* (Schreb.) Lindb. Hartz, N. Bidrag til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 15.] Late glacial

- 1904 *C. stellatum*. Weber, C. A., in Müller G. Ueber eine fröhldiluviale Flora bei Lüneburg. [Abh. Kgl. preuss. geol. Landesanstalt und Bergakad., Heft 40, p. 16.] Preglacial
- 1904 *Hypnum stellatum* Schreb. Geinitz, E., und Weber, C. A. Ueber ein Moostorflager der Rostocker Heide. [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.] Postglacial
- 1907 *H. stellatum*? Hartmann, F. Die Fossile Flora von Ingamsdorf, p. 14. Diluvial
- 1908 *H. stellatum*. Weber, C. A. Die Moostorfschichten zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, p. 43.] Recent
- 1909 *C. stellatum*. Nørregaard, E. M. Et senglacialt fra Dejbjerg Bakker [Meddel. fra Dansk geol. Fören., No. 15, p. 326.] Late glacial
- 1912 *H. stellatum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *H. stellatum*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *H. stellatum*. Weber, C. A. Die Mammutflora von Bornia. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 20.] Glacial
- 1914 *C. stellatum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Glacial
- 1915 *H. stellatum* (with some slight doubt). Camus, F. Sur les mousses trouvées dans un mammouth. [Compt. Rend. Acad. Par., CLX, p. 843.] Glacial
- 1917 *H. stellatum*. Rudolph, K. Untersuch. über den Aufbau böhmischer Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 84.] Recent

Campylium undetermined species.

- 1923 *Campylium* sp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 335.] Quaternary

Hyocomium Bry. eur.

Hyocomium flagellare (Dicks.) Bry. eur.

- 1910 *Hyocomium flagellare*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.] Postglacial
- 1919 *H. flagellare*. Gadecau, E. Les Forêts submergées de Belle-Ile-en-mer. [Bull. biol. de la France, etc., LIII, p. 295.] Postglacial

Ctenidium (Schimp.) Mitt.

Ctenidium molluscum (Hedw.) Mitt.

- 1911 *Hypnum molluscum* Hedw. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831.] Recent. Ice-land peat

- 1923 *C. molluscum*. Baumberger, E., etc. Die diluvialen Quaternary Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335.]

Rhytidadelphus (Lindb.) Warnst.

Rhytidadelphus loreus (Dill.) Warnst.

- 1910 *Hylocomium loreum* (Dill.) Bry. eur. Dixon, H. N. Postglacial Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
 1924 *Hyl. loreum*. Burrell, W. H. Pennine Peat. [Naturalist, 1924, p. 148.]

Rhytidadelphus squarrosus (L.) Warnst.

- 1853 *Hypnum squarrosum* L. Goeppert, H. R. Ueber die Bernstein Flora. [Monatsber. d. Berliner Akad., 1853, p. 459.]
 1888 *Hylocomium squarrosum* (L.) Bry. eur. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. [Geol. Mag., Decade III, Vol. V, p. 444.]
 1909 *Hyl. squarrosum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raække, No. 20, p. 260.]
 1910 *Hyl. squarrosum*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
 1922 *Hyl. squarrosum*. Jessen, Knud, og Rasmussen, R. Et Profil Gennem en Tørvemose paa Faærerne. [Danmarks geol. Undersøgelse, IV Raække, No. 13, p. 12.]
 1924 *Hyl. squarrosum*. Burrell, W. H. Pennine Peat. Recent [Naturalist, 1924, p. 148.]

Rhytidadelphus triqueter (L.) Warnst.

- 1898 *Hylocomium triquetrum* (L.) Bry. eur. Andersson, Gunnar. Stud. öfver Finlands Kvartärfiora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.]
 1905 *Hyl. triquetrum*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, I, Heft 6, p. 38.]
 1911 *Hyl. triquetrum*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 831.]
 1914 *Rhytidadelphus triqueter*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 249.]
 1918—20 *R. triqueter*. Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.]
 1920 *Hyl. triquetrum*. Reid, E. M. Two preglacial Floras from Castle Eden. [Quart. Journ. Geol. Soc., LXXVI, p. 111.]
 1923 *Hyl. triquetrum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 146, 337.]

Rhytidium (Sull.) Kindb.**Rhytidium rugosum (Ehrh.) Kindb.**

- 1908 *Hylocomium rugosum* ((Ehrh.) Bry. eur. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård [Sver. geol. Undersökning Arsbok, II, p. 19.]
- 1910 *Hyl. rugosum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]
- 1914 *R. rugosum*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

Hylocomium Bry. eur.**Hylocomium brevirostre (Ehrh.) Bry. eur.**

- 1905 *Hylocomium brevirostre*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]
Note. This is said to have been determined as *Brachythecium rutabulum* by Uhlmann.
- 1910 *Hyl. brevirostre*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Hist., 1910, p. 105.]
- 1923 *Hyl. brevirostre*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 146, 338.]

Hylocomium proliferum (L.) Lindb.

- 1898 *Hylocomium proliferum*. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.]
- 1902 *Hyl. splendens* (Dill.) Bry. eur. Hartz, N. Bidrag til Danmarks senglaciale Fl. og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, pp. 11, 15, 18.]
- 1903 *Hypnum splendens* Dill. Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 269.]
- 1905 *Hypnum splendens*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 40.]
- 1909 *Hyl. proliferum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.]
- 1911 *Hyl. splendens*. Dixon, H. N., in Lewis, F. J. The plant remains in the Scottish peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.]
- 1912 *Hyl. splendens*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.]
- 1914 *Hyl. splendens*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

- 1922 *Hyl. proliferum*. Jessen, Knud, og Rasmussen, R. Recent
Et Profil Gennem en Tørvemose paa Faerørne.
Danmarks geol. Undersøgelse, IV Raekke, No. 13,
p. 12.]
- 1923 *Hyl. splendens*. Baumberger, E., etc. Die dilu- Quaternary
vialen Schieferkohlen der Schweiz. [Beitr. Geol.
Schweiz, geotechn. Ser., Lief. 8, pp. 146, 286, 338.]
- 1924 *Hyl. splendens*. Burrell, W. H. Pennine Peat. Recent
[Naturalist, 1924, p. 148.]

Hylocomium pyrenaicum (Spr.) Lindb.

- 1912 *Hylocomium pyrenaicum*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.
[Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Hylocomium undetermined species.

- 1909 *Hylocomium* sp. Travis, W. G. Plant remains Peat
in peat at Aintree. [Trans. Liverpool Bot.
Soc., I, p. 5.]

Hypnum Dill. ex. p.

Hypnum Schreberi Willd.

- 1898 *Hylocomium parietinum (L.) Lindb.* Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 136.]
- 1900 *Hypnum Schreberi*. Gepp, A. Fossil plant re- Recent; sub-
mains in peat. [Journ. of Bot., XXXIII, p. 181.] marine peat
- 1902 *Hyl. parietinum*. Hartz, N. Bidrag til Danmarks senglaciale Flora og Fauna. [Danmarks geol. Undersøgelse, II Raekke, No. 11, p. 11.] Late glacial
- 1905 *Hyp. Schreberi*. Neuweiler, E. Die prähistor. Pflanzenreste Mittereuropas [Vierteljahr. naturforsch. Ges. Zürich, I, Heft 6, p. 40.] Pleistocene
- 1908 *Hyl. parietinum*. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård [Sver. geol. Undersökning Arsbok, II, p. 19.] Late glacial
- 1909 *Hyl. parietinum*. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 260.] Diluvial
- 1912 *Hylocomium Schreberi (Willd.) De Not.* Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *Hyp. Schreberi*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
- 1922 *Hyl. parietinum*. Jessen, Knud, og Rasmussen, R. Recent
Et Profil Gennem en Tørvemose paa Faerørne.
[Danmarks geol. Undersøgelse, IV Raekke, No. 13,
p. 12.]
- 1923 *Hyp. Schreberi*. Baumberger, E., etc. Die dilu- Quaternary
vialen Schieferkohlen der Schweiz. [Beitr. Geol.
Schweiz, geotechn. Ser., Lief. 8, pp. 146, 338.]

Ptilium (Sull.) De Not.**Ptilium crista-castrensis (L.) De Not.**

- 1914 *Ptilium crista-castrensis*. Zmuda, A. J. Fossile Diluvial Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. Sc. math. et nat., Sér. B, No. 2B, p. 249.]

Stereodon (Brid.) Mitt.**Stereodon arcuatus Lindb.**

- 1907 *Stereodon cfr. arcuatus*. Holst, N. O. Preglaciala Preglacial Dryasförande nära Ystad. [Geol. Fören. Förh., XXIX, p. 232.]
- 1914 *Hypnum Lindbergii* Mitt. Weber, C. A. Die Mammutflora von Borl. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 21.]

Stereodon callichrous Brid.

- 1912 *Hypnum callichroum* (Brid.) C. M. Dixon, H. N. Late glacial in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]
- 1912 *H. callichroum*. Stark, P. Beitr. zur Kenntn. der eiszeitl. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Stereodon cupressiformis (L.) Brid.

- 1900 *Hypnum cupressiforme* L. Gepp, A. Fossil plant-remains in peat. [Journ. of Bot., XXXIII, p. 181.] Recent; sub-marine peat
- 1903 *H. cupressiforme*. Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone [Zeitschr. für Naturwiss., LXXVI, p. 254.] Diluvial
- 1905 *H. cupressiforme*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.] Pleistocene
- 1909 *Stereodon cupressiformis*. Hartz, N. Bidrag til Danmarks tertiære Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] Diluvial
- 1910 *H. cupressiforme*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.] Postglacial
- 1913 *S. cupressiformis*. Beyle, M. Ueber einige Ablag. der Hamburger Gegend. Erster Teil. [Jahrb. der Hamburg. wiss. Anst., XXX, 6, Beih., p. 83.] Quaternary
- 1914 *S. cupressiformis*. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 249.] Diluvial
- 1923 *S. cupressiformis*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 336.] Quaternary
- 1923 *S. cupressiformis* var. *filiformis* (Brid.). Baumberger, op. cit., p. 336. Quaternary
- 1924 *H. cupressiforme*. Burrell, W. H. Pennine Peat. [Naturalist, 1924, p. 148.] Peat

Stereodon fastigiatus Brid.

- 1912 *Hypnum fastigiatum* (Brid.) Hartm. Dixon, H. N., Late glacial in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Stereodon fertilis (Sendtn.) Lindb.

- 1912 *Hypnum fertile* Sendtn. Stark, P. Beitr. zur Glacial Kenntniss der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Stereodon hamulosus (Bry. eur.) Lindb.

- 1914 *Hypnum hamulosum* Bry. eur. Weber, C. A. Die Glacial Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 20.]

Stereodon pallescens (Hedw.) Lindb.

- 1912 *Stereodon pallescens*. Stark, P. Beitr. zur Kenntn. Glacial der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Stereodon pratensis (Koch) Warnst.

- 1911 *Hypnum pratense* Koch ? Dixon, H. N., in Lewis, Recent; Ice-F. J. The plant remains in the Scottish peat land peat mosses. Part IV. [Trans. Roy. Soc. Edinb., XLVII, p. 830.]

Note. A single fragment found, probably belonging here.

- 1911 *Hypnum pratense* Koch. Dixon, H. N., op. et loc. Recent; Ice-peat.

Note. From a different layer. Either this or *S. areuatus*, probably *S. pratensis*.

- 1914 *H. pratense*. Weber, C. A. Die Mammutflora von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 1, p. 21.]

Stereodon recurvans (Schwaegr.) Broth.

- 1900 *Hypnum recurvans* Schwaegr. Penhallow, D. P. Pleistocene Canadian Pleistocene Flora of the Don Valley. [Rep. of 70th Meeting, Brit. Assoc., Bradford. Part II, p. 335.]

- 1920 *Sematophyllum recurvans* (Schwaegr.) E. G. Britt. Pleistocene Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 334, 335, 377.]

Stereodon reptilis (Rich.) Mitt.

- 1907 *Hypnum reptile* (Rich.) ? Hartmann, F. Die Fossile Diluvial Flora von Ingamsdorf, p. 14.

- 1912 *H. reptile*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Stereodon revolutus Mitt.

- 1910 **Stereodon revolutus**. Hesselbo, A. Mosrester fra Diluvial Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

Stereodon Vaucherii (Lesq.) Lindb.

- 1912 **Hypnum Vaucherii** Lesq. Dixon, H. N., in Warren, Late glacial S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Isopterygium Mitt.**Isopterygium pulchellum** (Dicks.) Jaeg.

- 1912 **Plagiothecium pulchellum** (Dicks.) Bry. eur. Stark, Glacial P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Plagiothecium Bry. eur.**Plagiothecium latebricola** (Wils.) Bry. eur.

- 1912 **Plagiothecium latebricola**. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Plagiothecium Roeseanum (Hamp.) Bry. eur.

- 1920 **Plagiothecium denticulatum** (L.) Roeseanum(Hamp.) Pleistocene Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 126, 184, 377.]

Plagiothecium Ruthei Limpr.

- 1912 **Plagiothecium Ruthei** var. *rupincola* Limpr. Stark, Glacial P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Plagiothecium silvaticum (Huds.) Bry. eur.

- 1914 **Plagiothecium silvaticum**. Zmuda, A. J. Fossile Diluvial Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 249.]

Plagiothecium striatellum (Brid.) Lindb.

- 1912 **Plagiothecium Muchlenbeckii** (Schimp.) Bry. eur. Glacial Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Plagiothecium undulatum (L.) Bry. eur.

- 1910 **Plagiothecium undulatum**. Dixon, H. N. Some Postglacial "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]

Brachytheciaceae.**Homalothecium** Bry. eur.**Homalothecium sericeum** (L.) Bry. eur.

- 1909 **Hypnum sericeum** L. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] Diluvial
 1914 **Homalothecium sericeum**. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.] Diluvial
 1915 **Hom. sericeum**. Reid, Clement and E. M. The Pliocene Floras of the Dutch-Prussian Border. [Med. van de Rijksopspor. van Delfstoffen, No. 6, p. 53.] Pliocene
 1918—20 **Hom. sericeum**. Docturowsky, V. S., and Kudriashov, V. Results of Russian studies on peat moors. [Reports on Peat, 4.] Postglacial
 1923 **Hom. sericeum**. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.] Quaternary

Camptothecium Bry. eur.**Camptothecium nitens** (Schreb.) Schimp.

- 1899 **Camptothecium nitens**. Weber, C. in Koert, W., and Weber. Ueber ein neues interglaciales Torflager. [Jahrb. der Kgl. preuss. geol. Landesanstalt für 1899, pp. 190, 191.] Interglacial
 1900 **C. nitens**. Dixon, H. N. Plant-remains in peat. [Journ. of Bot., XXXIII, p. 216.] Peat
 1903 **Hypnum trichoides** Neck. Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 269.] Diluvial
 1904 **C. nitens**. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.] Late glacial
 1904 **C. nitens**. Geinitz, E., und Weber, C. A. Ueber ein Moostorflager ... der Rostocker Heide. [Arch. d. Ver. d. Fr. d. Naturgesch. in Mecklenburg, LVIII, p. 13.] Postglacial
 1907 **C. nitens**. Dixon, H. N., in Lewis, F. J. Summary of Progress for 1907, p. 99. [Mem. Geol. Survey.] Peat
 1908 **C. nitens**. Weber, C. A. Die Moostorfschichten ... zwischen Sarkau und Cranz. [Engl. Bot. Jahrb., XLII, p. 43.] Pleistocene
 1908 **Hypnum trichoides**. Holst, N. O. Efterskörd från de senglaciala Lagren vid Toppeladugård.... [Sver. geol. Undersökning Arsbok, II, No. 2, p. 19.] Late glacial

- 1909 *C. nitens*. Schuster, J. Paläobot. Not. aus Bay- ern. [Ber. bayer. bot. Ges., XII, 1, p. 58.] Schiefer- kohlen
- 1910 *H. trichoides*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.] Diluvial
- 1912 *C. nitens*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. [Quart. Journ. Geol. Soc., LXVIII, p. 231.] Late glacial
- 1912 *C. nitens*. Stark, P. Beitr. zur Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.] Glacial
- 1914 *C. nitens*. Weber, C. A. Die Mammutfloren von Borna. [Abh. Nat. Ver. Bremen, XXIII, Heft 2, p. 18.] Glacial
- 1914 *Tomentypnum nitens* (Schreb.) Loeske. Zmuda, A. J. Fossile Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2B, p. 247.] Diluvial
- 1917 *C. nitens*. Rudolph, K. Untersuch. über den Aufbau böhmischer Moore. [Abh. k. k. zool.-bot. Ges. Wien, IX, p. 82.] Pleistocene
- 1918 *C. nitens*. Beyle, M. Ueber einige Ablag. fossiler Pflanzen der Hamburger Gegend. Zweiter Teil. [Jahrb. der Hamburg. wiss. Anst., XXXVI, p. 40.] Glacial
- 1920 *C. nitens*. Baker, Frank C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 228, 241, 377.] Pleistocene
- 1923 *C. nitens*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335.] Quaternary

Camptothecium lutescens (Huds.) Bry. eur.

- 1905 *Camptothecium lutescens*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, I, Heft 6, pp. 37, 40.] Pleistocene
- 1909 *Hypnum lutescens* Huds. Hartz, N. Bidrag til Danmarks tertiale Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] Interglacial
- 1923 *C. lutescens*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 334.] Quaternary

Camptothecium Woldenii Grout.

- 1917 *C. Woldenii* Grout, A. J. A Fossil Camptothecium. [Bryologist, XX, p. 9, t. I.] Kansan Drift
Note. Grout states that this is nearest to *C. pinnatifidum* and *C. aureum*, but differs clearly from both.

Brachythecium Bry. eur.

Brachythecium albicans (Neck.) Bry. eur.

- 1898 *Hypnum near albicans* Neck. Andersson, Gunnar. Stud. öfver Finlands Kvartärflora. [Bull. de la Commission géol. de Finlande, Tome 2, No. 8, p. 137.] Quaternary

- 1910 *H. albicans* Neck. Hesselbo, A. Mosrester fra Diluvial Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

Brachythecium glareosum (Bruch) Bry. eur.

- 1904 *Brachythecium glareosum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). Late glacial [Quart. Journ. Geol. Soc., LX, p. 124.]

Brachythecium Mildeanum (Schimp.) Schimp.

- 1912 *Brachythecium Mildeanum*. Dixon, H. N., in Warren, S. H. A late Glacial stage in the Lea Valley. Late glacial [Quart. Journ. Geol. Soc., LXVIII, p. 231.]

Brachythecium plumosum (Sw.) Bry. eur.

- 1889 *Brachythecium plumosum*. Candler, Chas. Observations on some Deposits at Saint Cross in Suffolk. [Quart. Journ. Geol. Soc., XLV, p. 507.]
 1910 *B. plumosum*. Dixon, H. N. Some "Neolithic" Postglacial moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
 1910 *B. plumosum*. Hesselbo, A. Mosrester fra Diluviet ved Skaerumhede. [Danmarks geol. Undersøgelse, II Raekke, No. 25, p. 108.]

Brachythecium reflexum (Stark) Bry. eur.

- 1912 *Brachythecium reflexum*. Stark, P. Beitr. zur Glacial Kenntn. der eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 90.]

Brachythecium rivulare (Bruch) Bry. eur.

- 1923 *Brachythecium cfr. rivulare*. Baumberger, E., etc. Quaternary Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, pp. 145, 335.]

Brachythecium rutabulum (L.) Bry. eur.

- 1888 *Brachythecium rutabulum*. Reid, Clement, and Ridley, H. N. Fossil Arctic Plants from Hoxne, in Suffolk. Geol. Mag., Decade III, Vol. V, p. 444.]
 1904 *B. rutabulum*. Bell, A. M. Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
 1905 *B. rutabulum*. Neuweiler, E. Die prähistor. Pflanzreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 40.]
 1909 *Hypnum rutabulum* L. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
 1910 *B. rutabulum*. Dixon, H. N. Some "Neolithic" Postglacial moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]

Brachythecium salebrosum (Hoffm.) Bry. eur.

- 1909 *Hypnum salebrosum* Hoffm. Hartz, N. Bidrag til Interglacial Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]

Brachythecium turgidum (Hartm.) Hartm. f.

- 1914 *B. turgidum*. Zmuda, A. J. Fossile Flora des Diluvial Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

Brachythecium velutinum (L.) Bry. eur.

- 1909 *Hypnum velutinum* L. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1914 *Brachythecium velutinum*. Zmuda, A. J. Fossile Diluvial Flora des Krakauer Diluviums. [Bull. int. Acad. Sci. Cracovie d. sc. math. et nat., Sér. B, No. 2 B, p. 249.]

Brachythecium undetermined species.

- 1920 *Brachythecium* sp. Reid, E. M. Two preglacial Pliocene Floras of Castle Eden. [Quart. Journ. Geol. Soc., LXXVI, p. 111.]
- 1922 *Brachythecium* sp. Jessen, Knud, og Rasmussen, R. Recent Et Profil Gennem en Tørvemose paa Faerørne. [Danmarks geol. Undersøgelse, IV Raekke, No. 13, p. 12.]
- 1923 *Brachythecium* sp. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 521.]
- 1925 *Brachythecium* sp. Dixon, H. N. Moss Remains in Russian Peat. [Journ. of Bot. LXIII, p. 370.]
- 1925 *Brachythecium* sp. Weyland, H. Beitr. zur Kenntn. fossiler Moose. I. [Senckenbergiana, VII, Heft 1/2, p. 52, t. III.]

Scleropodium Bry. eur.**Scleropodium purum** (L.) Limpr.

- 1909 *Hypnum purum* L. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1909 *H. purum*. Schuster, J. Paläobot. Not. aus Bayern. [Ber. bayer. bot. Ges., XII, 1, p. 58.]
- 1911 *H. purum*. Stöller, J. Die Flora der jungglazialen Ablag. Ostpreussens. [Jahrb. d. Kgl. preuss. geol. Landesanstalt, XXXI, t. II, Heft 1, p. 123.]

Scorpiurium Schimp.**Scorpiurium deflexifolium** (Solms) Schimp.

- 1923 *Eurhynchium circinatum* (Brid.) var. *deflexifolium* (Solms) Boul. Douin, R. Les mousses fossiles

des tufs du Lautaret. [Rev. Génér. de Bot., XXXV, p. 121.]

Cirriphyllum Grout.

Cirriphyllum cirrosum (Schwaegr.) Grout.

- 1910 *Hypnum cirrosum* Schwaegr. Hesselbo, A. Mos- Diluvial
rester fra Diluviet ved Skaerumhede. [Danmarks
geol. Undersøgelse, II Raekke, No. 25, p. 108.]

Cirriphyllum Vaucheri (Bry. eur. ex p.) Loeske & Fleisch.

- 1925 *Eurhynchium* sp. ? *Tommasinii* (Sendtn.) Ruthe. Upper
Weyland, H. Beitr. zur Kenntn. fossiler Moose. I. Pliocene.
[Senckenbergiana, VII, Heft 1/2, p. 52, t. III.]

Note. This is the plant recorded as *Eurhynchium* sp. by Engelhardt & Kinkelin, Oberpliocâne Flora des Frankfurter Klärbeckens, in Abh. Senckenb. naturforsch. Ges., XXIX, p. 188, t. XXII, figs. 9—13. The figures there show an entire leaf, apparently nerveless, very concave, and with recurved margins; and strongly suggest *Plagiothecium piliferum*. Weyland however who has examined the specimens figures and describes the piliform acumen as clearly toothed, and states that the margin is frequently *incurved*; the nerve is stated by him to be not present in the upper part of the leaf, while the lower part of the leaf does not permit of observation. He considers it may be *C. Tommasinii* (Vaucheri) or possibly *C. cirrosum*.

Oxyrrhynchium Warnst.

Oxyrrhynchium praelongum (Bry. eur.) Warnst.

[Note. There is considerable confusion about this and *Eurhynchium Stokesii*. The plant described as *Eurh. praelongum* (L.) in the Bry. eur. is not *Hypnum praelongum* L., which is the plant usually known as *Eurh. Stokesii*. It is therefore not always clear to which of these two plants a record of "*H. praelongum*" or "*Eurh. praelongum*" refers.]

- 1904 *Eurhynchium Swartzii* (Turn.) Hobk. Bell, A. M. Late glacial
Implementiferous Sections at Wolvercote (Oxfordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1905 *Eurhynchium praelongum* Bry. eur. Neuweiler, E. Pleistocene
Die prähistor. Pflanzenreste Mitteleuropas
[Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6,
p. 38.]
- 1912 *E. praelongum*. Stark, P. Beitr. zur Kenntn. der Glacial
eiszeitlich. Fl. und Fauna Badens. [Ber. naturforsch. Ges. Freiburg i. B., XIX, p. 116.]
- 1913 *E. praelongum*. Beyle, M. Ueber eine Ablag. fossiler Pflanzen der Hamburger Gegend. [Jahrb. Hamburg. wiss. Anstalt, XXX, 6, Beih., p. 83.] Pleistocene
- 1918—20 *O. praelongum*. Docturowsky, V. S., and Kudriashchev, V. Results of Russian studies on peat Postglacial
moors. [Reports on Peat, 4.]

- 1920 *E. Swartzii* (Turn.) Hobk. Reid, E. M. Two pre- Pliocene
glacial Floras from Castle Eden. [Quart. Journ.
Geol. Soc., LXXVI, p. 111.]
- 1922 *E. praelongum*. Jessen, Knud, og Rasmussen, R. Recent
Et Profil Gennem en Tørvemose paa Faerørne.
[Danmarks geol. Undersøgelse, IV Raekke, No. 13,
p. 12.]
- 1923 *E. praelongum*. Baumberger, E. etc. Die dilu- Quaternary
vialen Schieferkohlen der Schweiz. [Beitr. Geol.
Schweiz, geotechn. Ser., Lief. 8, p. 335.]

Oxyrrhynchium rusciforme (Neck.) Warnst.

- 1910 *Eurhynchium rusciforme* (Neck.) Milde. Dixon, H. Postglacial
N. Some "Neolithic" moss remains from Fort Wil-
liam. [Ann. Scott. Nat. Hist., 1910, p. 105.]

Oxyrrhynchium speciosum (Brid.) Warnst.

- 1904 *Eurhynchium speciosum* (Brid.) Milde. Bell, A. M. Late glacial
Implementiferous Sections at Wolvercote (Ox-
fordshire). [Quart. Journ. Geol. Soc., LX, p. 124.]
- 1915 *E. speciosum*. Reid, Clement and E. M. The Plio- Pliocene
cene Floras of the Dutch-Prussian Border. [Med.
van de Rijksopspor. van Delfstoffen, No. 6, p. 53.]

Oxyrrhynchium Swartzii (Turn.) Warnst.

- 1909 *Hypnum Swartzii* Turn. Hartz, N. Bidrag til Interglacial
Danmarks tertiaere Flora. [Danmarks geol.
Undersøgelse, II Raekke, No. 20, p. 261.]
- 1919 *Eurhynchium Swartzii* (Turn.) Bry. eur. Gadeceau, Postglacial
E. Les Forêts submergées de Belle-Ile-en-mer.
[Bull. Biol. de la France, etc., LIII, p. 295.]
- 1923 *E. Swartzii*. Baumberger, E., etc. Die diluvialen Quaternary
Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz,
geotechn. Ser., Lief. 8, p. 335.]

Eurhynchium Bry. eur.

Eurhynchium Stokesii (Turn.) Bry. eur.

- 1909 *Hypnum praelongum* L. Hartz, N. Bidrag til Interglacial
Danmarks tertiaere Flora. [Danmarks geol.
Undersøgelse, II Raeke, No. 20, p. 261.]
- 1910 *Eurhynchium praelongum* (L.) Hobk. Dixon, H. Postglacial
N. Some "Neolithic" moss remains from Fort
William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
- 1920 *E. praelongum*. Reid, E. M. Two preglacial Floras Pliocene
from Castle Eden. [Quart. Journ. Geol. Soc.,
LXXVI, p. 111.]
- 1922 *E. Stokesii* (*Stockesii, false*). Jessen, Knud, og Recent
Rasmussen, R. Et Profil Gennem en Tørvemose paa
Faerørne. [Danmarks geol. Undersøgelse, IV
Raekke, No. 13, p. 12.]

Eurhynchium striatum (Spr.) Bry. eur.

- 1912 *Eurhynchium* cfr. *striatum*. Stark, P. Beitr. zur Glacial
Kenntn. der eiszeitlich. Fl. und Fauna Badens.

- [Ber. naturforsch. Ges. Freiburg i. B., XIX,
p. 116.]
- 1923 *E. striatum*. Baumberger, E., etc. Die diluvialen Schieferkohlen der Schweiz. [Beitr. Geol. Schweiz, geotechn. Ser., Lief. 8, p. 335.]
- 1923 *E. striatum* var. *integrifolium*. Baumberger, op. et loc. cit.

***Eurhynchium striatum* (Schreb.) Schimp.**

- 1905 *Eurhynchium striatum*. Neuweiler, E. Die prähistor. Pflanzenreste Mitteleuropas [Vierteljahr. naturforsch. Ges. Zürich, L, Heft 6, p. 38.]
- 1909 *Hypnum striatum* Schreb. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]
- 1910 *E. striatum*. Dixon, H. N. Some "Neolithic" moss remains from Fort William. [Ann. Scott. Nat. Hist., 1910, p. 105.]
- 1911 *H. striatum*. Stoller, J. Beitr. zur Kenntn. der diluv. Flora Norddeutschlands. [Jahrb. Kgl. preuss. geol. Landesanstalt, XXXII, Teil 1, Heft 1, p. 135.]

***Eurhynchium strigosum* (Hoffm.) Bry. eur.**

- 1909 *Hypnum strigosum* Hoffm. Hartz, N. Bidrag til Danmarks tertiaere Flora. [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.]

***Eurhynchium* spec. Eng. et Kink.
cf. *Cirriphyllum* Vaucheri Loeske et Fl.**

***Rhynchosstegium* Bry. eur.**

***Rhynchosstegium Knowltoni* E. G. Britt.**

- 1899 *Rhynchosstegium Knowltoni* E. G. Britt. A new Tertiary fossil Moss, in Bull. Torr. Bot. Club, Upper XXVI, p. 79, with text figure (repeated op. cit., XXXIV, (1907), t. 9, fig. 5).

***Hypnodendron* (C. M.) Lindb.**

- 1922 *Hypnodendron arborescens* (Mitt.) Lindb., or *H. Reinwardtii* (Hornsch.) Lindb. Dixon, H. N. Note on a moss in amber. [Journ. of Bot., LX, p. 149, with text-figure.]

Note. The fragment of branch might very well belong to either of the above species. *H. arborescens* is known from the Malay Peninsula, but *H. Reinwardtii* has not been recorded from continental Asia.

Excluded and doubtful genera and species.

***Cryptothecium antediluvianum* Hueb.**

Cryptothecium antediluvianum nov. gen. & sp. Tertiary Huebener MS. in Weber, C. O. Die Tertiärfloren der niederrhein. Braunkohlenformationen, in Palaeontographica, II, p. 228 (1852).

This is stated by Schimper to be a species of *Sphagnum*.

Hypnites haeringianus Ett.

Hypnites haeringianus nov. gen. & sp. Ett. Tertiär; tingshausen, C. von. Die Tertiärfloren von Haer- schiste cal- ring, in Abh. d. geol. Reichsanstalt, Wien, II, caire bitu- p. 27, t. IV (1852).

Syn. *Hypnum haeringianum* Schimp., Traité de Pal. Végét., I, p. 249. Schimper considers this to belong to the *Hypna adunca*.

Hypnum armissanense Schimp. cf. *Muscites pulvinatus*.**Hypnum Brownii** Kirchner.

1898 *Hypnum Brownii* Kirchner, C. G. Walter. Contrib. to the fossil flora of Florissant, Colorado. [Trans. Acad. Sci. St. Louis, VIII, p. 178, t. XII]. cfr. also Knowlton, F. H. A review of the fossil plants in the U. S. Nat. Mus. in Proc. U. S. Nat. Mus., LI (1916), p. 245. See also Britton & Hollick, Bull. Torr. Bot. Club, XXXIV, p. 139—140, t. IX. The author compares it with *Brachythecium populeum*, but the figures do not suggest that, the leaves being broader and shorter. "Brachythecoid" seems as near as it seems safe to go.

Hypnum carbonarium Ludw.

1859 *Hypnum carbonarium* Ludwig, R. Fossile Pflanzen aus der Rheinisch-Wetterauer-Tertiär-Formation. [Palaeontographica, VIII, p. 62, t. IX and XXIV]. The figures rather suggest a *Fontinalis*. Schimper thinks a *Hypnum* (Traité de Pal. Végét., I, p. 248).

Hypnum columbianum Penhallow.

1890 *Hypnum columbianum* Penhallow, D. P., in Dawson, Sir Wm., Trans. Roy. Soc. Canada, VIII, Pt. 4, p. 77, fig. 3. Britton and Hollick, Bull. Torr. Bot. Club, XXXIV, p. 140, t. IX, fig. 4, consider this to be a conifer, near to *Widdringtonia helvetica* Heer, or *Glyptostrobus Ungerii* Heer.

Hypnum Haydenii Lesq.

1878 *Hypnum Haydenii* Lesq. in Hayden, F. V. Tertiary Flora of U. S. [Rep. of U. S. A. Geol. Survey, VII, p. 44]. Cockerell, T. D. A. [A Redwood described as a Moss, *Torreya*, VII, p. 203 (1907),] shows this to be a conifer, *Sequoia Haydenii* (Lesq.) Cock.

Hypnum Heerii Schimp.

1855—59 *Hypnum Schimperi* Heer. Flora tertiaria Helveticae, I, p. 28, t. III (nec *H. Schimperianum* Fossilium Catalogus II. 13.

Lor. 1864; nec *H. Schimperi* (Ung.) Schimp.
Schimper (*Traité de Pal. Végét.*, I, p. 247) alters the name to *H. Heerii*, in view of the earlier *H. Schimperianum* and *Muscites Schimperi* Ung. He compares it with *Amblystegium serpens*, but the resemblance cannot be very close; the leaves being longer and narrower, with an apparently percurrent nerve.

Hypnum Heppii Heer.

1855—59 *Hypnum Heppii* Heer. *Fl. tertiaria Helveticæ*, I, p. 28, t. III.

Syn. *Hygroamblystegium Heppii* (Heer) Broth. in Engler, *Die Natürliche Pflanzenfam., Musci*, II, p. 523 (1925).

Schimper figures this, *Traité de Pal. Végét.*, t. VI, figs. 20, 21 (not 17, 18, 19, as in the text). He considers it to be nearest to *Amblystegium irriguum* var. *fallax*, but the leaves are, as he says, too narrow. It suggests to me a *Drepanocladus*, of the *fluitans* group, but the acumen is perhaps too broad, and the dimensions rather small.

Hypnum Mastodontum Boul.

1892 *Hypnum Mastodontum*. Boul, N., *Flore pliocène du Mont-Dore*, p. 45, f. 9.

Boulay suggests *Amblystegium riparium* as the nearest affinity, but agrees that it cannot be that. The upper cells are shortly linear; no distinct alar cells visible.

1903 *Hypnum cfr. Mastodontum* Boul. Marty, P. *Flore Miocène de Joursac. Rev. Haute Auvergne*, 1903, p. 18, t. I, figs. 7—10].

This does not appear to have been critically examined. All one can say is that it is a pleurocarpous moss, with lanceolate, single-nerved leaves.

Hypnum plumula Schimp.

1869 *Hypnum plumula* Schimp., *Traité de Pal. Végét.*, I, p. 248.

According to Schimper this is "une forme exotique", not allied to any European moss. It is not figured.

Etage ton-grien, près d'Allauch

Hypnum priscum Schimp.

1865 *Hypnum priscum* Schimp. in Heer, O. *Die Urwelt der Schweiz*, I. Aufl., p. 497.

Schimper says this is very close to *Calliergon sarmentosum*. Gams, in *Vierteljahrsschr. naturforsch. Ges. Zürich*, LXIII, p. 28 (1913) states that according to Meylan it is between *Calliergon giganteum* and *Scorpidium scorpioides*.

Quaternary lignites of Dürnten, Switzerland

Hypnum Saportanum Schimp. cfr. *Muscites setosus*.**Hypnum Schimperi** (Ung.) Schimp.

- 1869 *Hypnum Schimperi* (Ung.) Schimp. *Traité de Pal. Végét.*, 1, p. 246. *Muscites Schimperi* Ung. *Genera et Spec. Plant. Fossil.*, p. 42 (1850); and *Iconograph. plant. fossil.*, in *Denkschr. d. k. k. Akad. d. Wissensch.*, Wien, 1852, p. 10, t. IV; and p. 82, t. XXVII, Figs. 1, 2.

Unger compares it with *Campylium stellatum* ("admodum convenit"). Ettingshausen, in *Denkschr. d. k. k. Akad. d. Wissensch.*, Wien, XXXVIII, Abth. 1, figures on t. V, figs. 7—9 a separate capsule, associated with the vegetative fragments. I have examined the specimen on which this is based, in the Geol. Dept. of the British Museum (No. 592). It may be the impression of a moss capsule, but if so it is much more like *Catoscopium* than *Hypnum*!

Schimper considers it of doubtful affinity.

Schiste calcaire argileux de Paschlug

Hypnum Schimperi Heer. cfr. *Hypnum Heerii*.**Hypnum Taramellianum** Farneti.

- 1896 *Hyp. Taramellianum* Farn. *Ricerche di briol. paleont. al periodo glaciale.* [Atti dell' Istit. Bot. della Reale Univ. di Pavia, Ser. II, vol. V, p. 56, t. IV] cf. also Schilbersky, K. Ein Laubm. aus dem Pleistocän von Kecskemét [Math. Termtud. Ertesítő, XXX (1912) p. 632] text and figures. The author says it recalls *Scleropodium purum*, *Drepanocladus turgescens*, *Acrocladium euspidatum*, etc., and does not venture to place it. The figure looks to me more like a *Hygrohypnum* (e. g. *H. molle* or *H. dilatatum*) than any of the above.

Pleistocene

Hypnum Ungerii Schimp.

- 1869 *Hyp. Ungerii* Schimp. *Traité de Pal. Végét.*, I, p. 247. *Muscites hypnoides* Ung. *Iconograph. plant. fossil.*, in *Denkschr. d. k. k. Akad. d. Wissenschaft.*, Wien, 1852, p. 11, t. IV.

Probably, according to Schimper, a submerged or floating form of a riparian moss.

Schiste calcaire marneux de Radoboj

Hypnum viride.

- 1909 *Hypnum viride*. Hartz, N. *Bidrag til Danmarks tertiaere Flora.* [Danmarks geol. Undersøgelse, II Raekke, No. 20, p. 261.] This may refer either to *H. viride* Lam. (*Brachythecium populeum*), or to *H. viride* Schleich. (*Amblystegium serpens*).

Hypnum Weberianum Schimp.

1856 **Hypnum Weberianum** Schimp., Traité de Pal. Végét., I, p. 249.

Hyp. lycopodioides Weber, C. O. in Wessel & Weber, Neuer Beitrag zur Tertiärfloren der niederrhein. Braunkohlförderung. [Palaeontographica, IV, p. 126, t. XXX, f. 12] (nec H. lycopodioides Brid.).

Schimper compares it to *Brachythecium salebrosum* or *Antitrichia curtipendula*, but with leaves narrowly linear-lanceolate. He does not suggest any closer affinity. The figure shows a well preserved moss with regularly plumose stems; it suggests to me an aquatic *Drepanocladus*, e. g. a *fluitans* form, rather than either of the above plants. The name was given by Weber in reference to a supposed resemblance to certain Lycopods, and has no connection with *Drepanocladus lycopodioides* (Brid.). The name was altered by Schimper to *H. Weberianum*, rather unfortunately, in view of *H. Weberianum* Goepf., which latter Schimper referred to *Drepanocladus Sendtneri*.

Hypnum Spp.

1892 **Hypnum** sp. Boulay, N. Flore pliocène du Mont Dore, p. 45.

The author compares with *Hygrohypnum filicinum* and *Cratoneuron commutatum*.

1900 **Hypnum** sp. Penhallow, D. P. Canadian Pleistocene Flora of the Don Valley. [Rep. of Brit. Assoc., 1900, Bradford. Pt. II, p. 336.]

1903 **Hypnum** sp. Range, Paul. Das Diluvialgebiet von Lübeck und seine Dryastone [Zeitschr. f. Naturwiss., LXXVI, p. 191 bis].

1903 **Hypnum** sp. Range, Paul, op. cit., p. 257.
1905 **Hypnum** sp. Neuweiler, E. Zur Interglazialflora der schweizerisch. Schieferkohlen. [Ber. d. schweizerisch. bot. Ges., Heft XV. Beilage. Neunter Ber. d. zürcherisch. bot. Ges., p. 94].

The author comments "entspricht dem *H. sarmatosum* und *trifarium*".

1907 **Hypnum** sp. Hartmann, F. Die fossile Flora von Ingramsdorf, p. 14.

1908 **Hypnum** sp. Dixon, H. N., in Lewis, F. J. The Plant Remains in the Scottish Peat Mosses. Part III. [Trans. Roy. Soc. Edinb., XLVI, p. 40.]

1920 **Hypnum** sp. Baker, Frank, C. The life of the Pleistocene or glacial period. [Univ. of Illinois Bull., XVII, pp. 27, 87, 101, 329.]

1923 **Hypnum** sp. Douin, R. Les mousses fossiles des tufs du Lautaret. Rev. Gén. Bot., XXXV, p. 120.]

Lycopodites Meekii Lesq.

Lycopodites Meekii Lesq. It has been suggested by Solms-Laubach (Fossil Botany, p. 186, 1891) that this fossil, figured by Lesquereux

in 2nd Geol. Surv. Penn., Rep. of Progress P. Harrisburg, 1879, t. LXII, f. 1, may be a moss. Seward however shows that this is untenable, and remarks that it resembles closely *Selaginella Oregana*. It has no real claim to be considered a Bryophyte.

Muscites

Muscites Brongn., Hist. des Végétaux Fossiles, p. 93 (1828).

The diagnosis is preceded by a note that the author designated under this genus such fossil mosses as could not be determined as belonging to any living genus.

Muscites apiculatus Goepp. & Ber.

1845 **Muscites apiculatus** Goepp. & Ber. Die im Bernstein befindlichen organ. Reste der Vorwelt, I, Abt. 1, p. 111, t. VI, fig. 32.

Syn. *Barbula subcanescens* Goepp. & Ber. in Goeppert, H. R. Ueber die Bernsteinflora, p. 458.

Schimper, Traité de Pal. Végét., I, p. 243, says of this "plus que problématique". Caspary (Flora d. Bernsteins, in Abh. Kgl. preuss. geol. Landesanstalt, N. F., Heft 4,) says of this that it "ist auch kein Moos".

Schimper adds that it can in no case belong to *Barbula*, and says that a single leaf suggests a floral leaf from *Polytrichum*. This appears to me by no means impossible; cfr. Goeppert & Berendt's fig. 32 with the ♂ bract e. g. of *Pol. gracile* (Bry. eur., t. 421, fig. 7), or of *P. juniperinum* (ibid., t. 423, fig. 10).

Muscites Bertrandii Lignier.

1914 **Muscites Bertrandii** Lignier, V. Sur une mousse houillère à structure conservée. [Bull. Soc. Linn. Normandie, 6 Sér., VII, p. 128.]

Palaeozoic;
Stephanien;
silex de
Grand Croix

This is described as showing a partial transverse section of a stem, exhibiting parenchymatous tissue, the unistratose outermost layer with thickened walls, and rhizoids with *oblique* partitions. The evidence for its belonging to the Bryophytes seems rather slight, though I am not aware of any other suggestion having been made for it; and the figure certainly indicates a moss stem.

Muscites confertus Goepp. et Ber.

cfr. *Weisia conferta* Schimp.

Muscites dubius Goepp. & Ber.

1845 **Muscites dubius** Goepp. & Ber. Die im Bernstein befindlichen organ. Reste der Vorwelt, I, Abt. 1, p. 112, t. VI, figs. 25, 26; and Goeppert, Ueber die Bernsteinflora, p. 459.

Schimper in the Traité de Pal. Végét. makes no reference to this; from the figure I should



feel doubtful if it is a moss; possibly plumose feathers of a bird, or pappus hairs of a higher plant.

Muscites elatinus Sap.

cf. *Thuidium elatinum* (Sap.) Schimp.

Muscites elegans Goepp.

1853 *Muscites elegans* Goepp. Ueber die Bernstein-flora. [Monatsber. d. Berliner Akad., 1853, p. 459.]

Of doubtful affinity. Goeppert says of it "Ein höchst zierliches, 3 Linien langes Pflänzchen, ähnlich manchen *Mnium* Arten hinsichtlich der Insertion der fast gestielten, etwas $\frac{1}{2}$ Lin. langen Blättchen."

Schimper (Traité de Pal. Végét., I, p. 244) suggests that it may be a small Hookeriaceous moss.

Muscites fontinaloides Ung.

1852 *Muscites fontinaloides* Unger. Iconograph. plant. fossil. [Denkschr. d. k. k. Akad. d. Wissensch., Wien, 1852, p. 82, t. XXVII, figs. 3, 4.]

Schimper makes no reference to this. From the figure it must, I think, if a moss at all, which seems probable, almost certainly be a *Fontinalis*.

Muscites Hauchecornei Casp. & Klebs.

1906 *Muscites Hauchecornei* Caspary & Klebs, in Caspary, Flora d. Bernsteins. [Abh. Kgl. preuss. geol. Landesanstalt, N. F., Heft 4, p. 56, t. VIII, fig. 48.]

Syn. *Muscites serratus* Goepp. in sched., non *M. serratus* Goepp. & Ber.

Caspary can suggest no relationship. The figures strongly suggest a *Mnium* or *Mnioid* moss (cfr. *M. antiquorum* Card. & Dix.); the cells are described as isodiametric and irregular, —.0085 —.0414 mm. (? .014.).

Muscites hirsutissimus Goepp. & Ber.

1845 *Muscites hirsutissimus* Goepp. & Ber. Die im Amber Bernstein befindlichen organ. Reste der Vorwelt, I, Abt. 1, p. 112, t. VI, figs. 23, 24; and Goepp. Ueber die Bernsteinflora, p. 459.

Schimper (Traité de Pal. Végét., I, p. 245) says of this that it is evidently a phanerogamous plant.

Muscites hypnoides Ung.

cf. *Hypnum Ungerii* Schimp.

Muscites intricatus Sap.

1863 *Muscites intricatus* Saporta. Etudes sur la végétation du Sud-Est de la France. [Ann. des Sc. nat. Par., 4^e sér., XIX, p. 30].

Saporta compares this with *Philonotis fontana*. The position must be considered extremely doubtful.

Muscites journacensis Marty.

1903 **Muscites journacensis** Marty, P. Flore miocène Miocene de Journac. [Rev. Haute Auvergne, 1903, p. 18, t. I, figs. 4—6.]

Marty says that it suggests a *Fissidens* or *Conomitrium*; but this from the branching is quite out of the question. The thickness of the stem as shown in the enlarged fig. 5 appears to me to preclude a moss altogether, and I am inclined to think it more likely a phanerogam.

Muscites oeningensis Braun

ct. *Thuidium oeningense* (Braun) Schimp.

Muscites pereger Sap.

1862 **Muscites pereger** Saporta, Etudes sur la végétation du Sud-Est de la France. [Ann. des Sc. nat. Par., 4^e sér., XVII, p. 207.] Gypse d'Aix

Schimper makes no reference to this. Saporta compares it at once with *Polytrichum commune* and *Dicranum scoparium*!

Muscites polytrichaceus Ren. & Zeill.

1885 **Muscites polytrichaceus** Renault, B. & Zeiller, R. Sur des Mousses de l'époque houillère. [Compt. Rend. Acad. Sci. Paris, C, p. 660]. Also in Ren. & Zeill., Flore Fossile de Commentry, in Bull. Soc. Industr. Min., 1888, p. 34, t. XLI. Described and figured by Jongmans, in Med. van de Rijksopspor. van Delfstoffen, No. 3, p. 11. Palaeozoic, Stéphanien de Commentry

Supposed to be an Acrocarpous moss. Numerous stems with clearly defined, uninerved leaves, "alternate", e descr., and so figured. The texture and size seemed to leave no doubt of its being a moss. It is compared with *Rhizogonium* and *Polytrichum*; it has certainly nothing to do with the latter, and very doubtfully with the former.

It is figured also by Seward, Fossil Plants, vol. I, p. 239, fig. 51.

Muscites pulvinatus Sap.

1865 **Muscites pulvinatus** Saporta, Etudes sur la végétation du Sud-Est de la France. [Ann. des Sci. nat. Par., 5^e sér., IV, p. 32, t. I, f. 3.] Tertiary lig-

Hypnum armisanense Schimp., Traité de Pal. Végét., I, 248. nites of Armissan

Schimper compares it with *Eurhynchium Stokesii*.

Muscites redivivus Sap.

1862 **Muscites redivivus** Saporta, op. cit., 4^e sér., XVII, p. 207, t. II, f. 2.

A Hypnaceous moss of doubtful affinity. According to Schimper the figure recalls somewhat *Eurhynchium Teesdalei*, but he considers it impossible to determine the genus.

Schistes
marneux
dans les
gypses
d'Aix

Muscites Schimperi Ung.

cf. *Hypnum Schimperi* (Ung.) Schimp.

Muscites serratus Goepp. & Ber.

1845 **Muscites serratus** Goeppert & Berendt, Die im Bernstein befindlichen organ. Reste der Vorwelt, I, Abt. 1, p. 111, t. VI, figs. 27, 28. Also Goepert, Ueber die Bernsteinflora, p. 459.

Schimper, Traité de Pal. Végét., I, 250, suggests an affinity with the exotic genus, *Trichosteleum*, e. g. *T. Boschii* (Doz. & Molk.) This affinity is based solely on the ground of the appearance of papillae on the cells of the enlarged figure 28, — "les feuilles à bord denté auraient le dos couvert d'aspérités spiniformes". He adds that he thinks the figure inexact, the asperities being probably papillae, and being associated with rhomboidal cells, lax, and without a median nerve, they are distinct from all European species with papillose cells.

A careful examination of this figure, however, reveals no such asperities. What Schimper no doubt took for these are, I have no doubt, simply the marginal denticulations drawn so as to appear dorsal in one of the leaves. Moreover the plant from its size could not possibly have had leaves large enough for *Trichosteleum* or the allied genera; fig. 27, of a branch apex, containing about 20 leaves, is less than 1.5 mm. in length! Schimper's own suggestion of a *Fabronia* is without doubt the correct one; the size, the areolation, and the long and fine denticulations of the leaf, agree exactly with species of this genus, e. g. *F. octoblepharis* Schleich.; and fig. 28 might well have been enlarged slightly from the plate in *Bry. eur.*, Tab. 451, of this species.

Muscites setosus Sap.

1895 **Muscites setosus** Saporta, Etudes sur la végétation Tertiary du Sud-Est de la France. [Ann. des sci. nat. lignites of Paris, 5^e Série, IV, p. 30, t. 1, f. 1]. Armissan *Hypnum Saportanum* Schimp., Traité de Pal. Végét., I, p. 247.

Muscites squamatus Brongn.

1828—1844 *Muscites ?squamatus* Brongn. Ad. Hist. des végétaux fossiles, p. 95, t. X.

Syn. *Lycopodites squamatus* Brongn. in Cuvier & Brongn. Descr. géol. des env. de Paris, p. 359, t. XI, fig. 3 (1822). And Brongn., Sur la Classification des Végétaux Fossiles en Général; Chap. II; in Mém. du Mus. d'Histoire nat., VIII, p. 304; t. VI, fig. 1.

This is in all probability a Lycopod rather than a moss.

Muscites Sternbergianus Dunker is not a moss, but a *Sphenolepidium*.

Muscites tortifolius Casp. & Klebs.

1906 *Muscites tortifolius* Caspary & Klebs, in Caspary, Flora d. Bernsteins. [Abh. Kgl. preuss. geol. Landesanstalt, N. F., Heft 4, p. 57; t. VIII, fig. 49.]

Caspary can suggest no relationship. It is clearly a pleurocarpous moss. The leaves are figured as strongly toothed, the acumen strongly half twisted; approximately two-thirds of a millimetre in length, and deeply plicate — an unusual feature in leaves of such small size.

Muscites Tournalii Brongn.

cf. *Fontinalis Tournalii* (Brongn.) Schimp.

Najadita Buckman

1850 *Najadita* gen. nov. Buckman, J. On some Fossil plants of the Lower Lias, in Quart. Journ. Geol. Soc., VI, p. 415.

Gardner, on Mesozoic Angiosperms, Geol. Mag. III, 1886, p. 203, refers the three species here described to Bryophytes; but Miss Sollas (Quart. Journ. Geol. Soc., LVII, p. 307) places them with the Lycopods, and this seems to be generally accepted. Cf. Zeiller, in Progressus Rei Botanicae, II, p. 179 (1908).

Schlotheimia tenuifolia Sternb.

Jongmans in Med. van de Rijksopspor. van Delfstoffen, No. 3, p. 214, refers to this. It is a genus of Calamites alliance, and has nothing to do with the moss genus *Schlotheimia* Brid.

Sporogonites Halle**Sporogonites** Halle.

Sporogonites Halle, gen. nov. A fossil Sporogonium from the Lower Devonian of Röragen in Norway; Bot. Not. 1916, p. 79; with text figure.

Sporogonites exuberans Halle.

- 1916 *Sporogonites exuberans* Halle, T. G. Lower Devonian plants from Röragen in Norway. [Kgl. Svensk. Vet.-Akad. Handl., LVII, p. 27; t. III, figs. 11—32]. Cfr. also Fleischer, Max, in Hedwig., LXI, p. 400.

This has been referred with some confidence to *Andreaea*, but it certainly belongs to no existing genus of mosses. The affinity with *Andreaea* is suggested from the longitudinal dehiscence of the capsule as suggested by vertical clefts in the wall of the sporogonium, but the resemblance ends there. It is in all probability — while possibly indicating the ancestry of the Bryophyta — not a true Bryophyte, but rather related — perhaps closely — to the genera *Rhynia* and *Hornea* recently described by Kidston & Lang (cf. Trans. Roy. Soc. Edinb. L1, p. 761, LII, pp. 603, 643, 831). The vegetative plant of *Sporogonites* is unknown, the large sporogonium, 6—9 × 3—4 mm., on a stalk at least 50 mm. high, resembles that of *Hornea* (rather than *Rhynia*) in the presence of a large columella, rendering the spore-cavity dome-shaped; the spores measure about 20—25 μ , those of the two genera in question being about 40 μ & 50 μ respectively; while *Sporogonites* differs apparently from both in the presence of longitudinal slits; though the capsule of *Hornea* appears to exhibit normally certain irregularities of structure on the capsule wall. It seems in the present state of our knowledge impossible to determine the position of *Sporogonites* more definitely than this.



Index.

Heavy types indicate those places where the synonymy is given.

- | | |
|---|--|
| Acroceratium cuspidatum (L.)
Mitt. 79.
A. sarmentosum (Wahl.) Mitt.
77.
Acrocladium Mitt. 79—80.
A. cuspidatum (L.) Lindb. 68,
79.
A. squarrosum (L.) Lindb. 80.
A. species 80.
Amblyodon P. Beauv. 41.
A. dealbatus (Dicks.) Beauv. 41.
Amblystegiella Loeske 60.
A. confervoides (Brid.) Loeske
60.
Amblystegium Bry. eur. 59-60.
A. aduncum (Hedw.) Sanio 63.
A. aduncum var. Kneiffii 63.
A. capillifolium (Warnst.) 64.
A. chrysophyllum (Brid.) De
Not. 81.
A. confervoides (Brid.) 60.
A. cordifolium (Hedw.) De Not.
75.
A. Cossoni (Schreb.) Lindb. 70.
A. curvicaule Jur. 60.
A. dilatatum Wils. 80.
A. exannulatum (Guemb.) De
Not. 65.
A. fallax var. spinifolium
Schimp. 60.
A. filicinum (L.) De Not. 61.
A. fluitans (L.) De Not. 66, 67.
A. giganteum (Schimp.) De Not.
75, 76.
A. intermedium Lindb. 68.
A. Kneiffii Bry. eur. 64.
A. Kochii Bry. eur. 59.
A. latifolium (Lindb. et Arn.)
Lindb. 69.
A. molle (Dicks.) Lindb. 80.
A. polygamum Bry. eur. 82.
A. protensum Lindb. 82.
A. radicale Bry. eur. 60.
A. riparium Bry. eur. 59. | A. Rotae De Not. 66.
A. scorpioides (L.) Lindb. 71.
A. Sendtneri (Schimp.) De Not.
72.
A. serpens Bry. eur. 59.
A. stellatum (Schreb.) Lindb.
82.
A. stramineum (Dicks.) De Not.
78.
A. turgescens (Jens.) Lindb. 73.
A. varium (Hedw.) Lindb. 60.
A. species 60.
Amphidium Nees 30.
A. lapponicum (Hedw.) Schimp.
30.
Andreaea Ehrh. 17, 106.
Andreaeaceae 17.
A. Huntii Limpr. 17.
A. petrophila Ehrh. 17.
A. Rothii Web. et Mohr. 17.
A. species 17.
Anoectangium Hedw. 30.
A. compactum Schwaegr. 30.
Anomobryum Schimp. 33.
A. concinnatum (Spruce) Lindb.
33.
Anomodon Hook. et Tayl. 55.
A. curtipendula (Neuweiler) 51.
A. longifolius (Schleich.) Bruch
55.
A. viticulosus (L.) Hook. et
Tayl. 55.
A. species Eng. et Kink. 55.
Antitrichia Brid. 51.
A. curtipendula (Hedw.) Brid.
51.
Astrophyllum curvatulum
Lindb. 37.
A. cuspidatum (L.) Lindb. 36.
A. hymenophylloides (Hueben.)
Lindb. 38.
A. marginatum (Dicks.) Lindb.
38.
A. medium (Bry. eur.) Lindb. 38. |
|---|--|

- A. silvicum Lindb. 37.
 A. subglobosum (Bry. eur.)
 Lindb. 39.
 A. undulatum (L.) Lindb. 39.
Atrichum subundulatum Schimp.
 45.
Aulacomniaceae 40-41.
Aulacomnium Schwaegr. 40-41.
 A. palustre (L.) Schwaegr. 40-
 41.
 A. palustre var. imbricatum
 Bry. eur. 40.
 A. turgidum Schwaegr. 41.
Autophyllum furcatus Grand-
 Eury 36.
Barbula Hedw. 26.
 B. fallax Hedw. 26.
 B. icmadophila Schimp. 26.
 B. paludosa Schleich. 26.
 B. rubella Lindb. 26.
 B. subcanescens Goepp. et Ber.
 26, 101.
Bartramia 43-44.
Blindia Bry. eur. 19.
 B. acuta (Huds.) Bry. eur. 19.
 B. acuta var. trichodes Braithw.
 19.
Brachytheciaceae 90-96.
Brachythecium Bry. eur. 91-93.
 B. albicans (Neck.) Bry. eur.
 91.
 B. glareosum (Bruch) Bry. eur.
 92.
 B. Mildeanum (Schimp.)
 Schimp. 92.
 B. plumosum (Sw.) Bry. eur. 92.
 B. reflexum (Stark) Bry. eur.
 92.
 B. rivulare (Bruch) Bry. eur.
 92.
 B. rutabulum (L.) Bry. eur. 92.
 B. salebrosum (Hoffm.) Bry.
 eur. 93.
 B. turgidum (Hartm.) Hartm.
 f. 93.
 B. velutinum (L.) Bry. eur. 93.
 B. species 93.
Bryaceae 31-36.
Bryum Dill. 33-36.
Bryum capillare L. 33.
B. cirratum Hoppe et Hornsch.
 33.
B. crispulum Hampe 34.
B. cyclophyllum (Schwaegr.)
 Bry. eur. 34.
B. Duvalii Voit. 34.
B. elegans Nees 34.
B. erythrocarpum Schwaegr. 34.
B. intermedium Brid. 34.
 B. lacustre Brid. 34.
 B. microstegium Bry. eur. 34.
 B. neodamense Itzigs. 34.
 B. pallens Swartz 35.
 B. pseudotriquetrum Schwaegr.
 35, 36.
 B. Schleicheri Schwaegr. 35.
 B. Schleicheri var. latifolium
 Schimp. 35.
 B. ventricosum Dicks. 35.
 B. species 36.
Bryon crispatum Grand'Eury
 36.
Calliergon (Sull.) Kindb. 75-79.
C. cordifolium (Hedw.) Kindb.
 75.
C. diluvianum (Schimp.) Dixon
 75.
C. giganteum (Schimp.) Kindb.
 75.
C. Richardsoni (Mitt.) Kindb.
 77.
C. Richardsoni Zmuda (not
 Kindb.) 76, 77.
C. sarmentonum (Wahl.) Kindb.
 77-78.
C. stramineum (Dicks.) Kindb.
 78.
C. trifarium (Web. et Mohr)
 Kindb. 72, 80.
C. tundrae (Arn.) 73.
C. turgescens (Jens.) Kindb.
 73.
C. species 79.
Camptothecium Bry. eur. 90-91.
C. lutescens (Huds.) Bry. eur.
 91.
C. nitens (Schreb.) Schimp.
 90-91.
C. Woldenii Grout 91.
Campylium (Sull.) Bryhn 81-83.
C. chrysophyllum (Brid.) Bryhn
 81.
C. helodes (Spr.) Broth. 81.
C. insubricum (Farneti) Dixon
 81.
C. polygamum (Bry. eur.) Bryhn
 82.
C. protensum (Brid.) Broth. 82.
C. Sommerfeltii (Myr.) Bryhn
 82.
C. stellatum (Schreb.) Lang. et
 Jens. 82-83.
C. species 83.
Campylopus flexuosus (L.)
 Brid. 23.
Catharinaea Ehrh. 45.
C. subundulata (Goepp. et
 Menge) 45.

- | |
|---|
| <i>C. undulata</i> (L.) Web. et Mohr. 45.
<i>Catascopiaceae</i> 43.
<i>Catoscopium</i> Brid. 43.
<i>C. nigritum</i> (Hedw.) Brid. 43.
<i>Ceratodon</i> Brid. 19.
<i>C. purpureus</i> (L.) Brid. 19.
<i>C. purpureus</i> var. <i>paludosus</i> Warnst. 19.
<i>Cinclidium</i> Sw. 39, 40.
<i>C. arcticum</i> Schimp. 39, 40.
<i>C. latifolium</i> Lindb. 40.
<i>C. stygium</i> Sw. 40.
<i>C. species</i> 40.
<i>Cirriphyllum</i> Grout 94.
<i>C. cirrosum</i> (Schwaegr.) Grout 94.
<i>C. Vaucherii</i> (Bry. eur. ex p.) Loeske et Fleisch. 94.
<i>Climaciaceae</i> 50.
<i>Climacium</i> Web. et Mohr 50.
<i>C. dendroides</i> (Dill.) Web. et Mohr 50.
<i>Conostomum</i> Sw. 43.
<i>C. boreale</i> Sw. 43.
<i>C. tetragonum</i> (Dicks.) Lindb. 43.
<i>Coscinodon</i> Spreng. 28.
<i>C. humilis</i> Milde 28.
<i>Cratoneuron</i> (Sulliv.) Roth 61-62.
<i>C. commutatum</i> (Hedw.) Roth 61-62.
<i>C. decipiens</i> (De Not.) Broth. 62.
<i>C. falcatum</i> (Brid.) Roth 62.
<i>C. irrigatum</i> (Zett.) Roth 62.
<i>C. species</i> 62.
<i>Cryptothecium</i> Hueb. 96.
<i>C. antediluvianum</i> Hueb. 96.
<i>Otenidium</i> (Schimp.) Mitt. 83.
<i>C. molluscum</i> (Hedw.) Mitt. 83.
<i>Desmatodon</i> Brid. 27.
<i>D. latifolius</i> (Hedw.) Bry. eur. 27.
<i>D. latifolius</i> var. <i>muticus</i> Brid. 27.
<i>Dichelyma</i> Myr. 50.
<i>D. capillaceum</i> 18, 50.
<i>D. species</i> 49, 50.
<i>Dichodontium</i> Schimp. 20.
<i>D. pellucidum</i> (L.) Schimp. 20.
<i>D. subpellucidum</i> (Goepp. et Menge) 20.
<i>Dicranaceae</i> 17-23.
<i>Dicranella</i> Schimp. 19-20.
<i>D. Bergeri</i> (Stark) 20.
<i>D. cerviculata</i> (Hedw.) Schimp. 19.
<i>D. Mühlenbeckii</i> (Stark) 21.
<i>D. squarrosa</i> (L.) Schimp. 19.
<i>D. Sauteri</i> (Stark) 21.
<i>D. species</i> 19, 20.
<i>Dicranites</i> Klebs 22, 23.
<i>D. Casparyi</i> Klebs 23, 24.
<i>D. obtusifolius</i> Casp. et Klebs 23.
<i>D. subflagellare</i> Casp. 22, 23.
<i>Dicranodontium</i> Bry. eur. 23.
<i>D. longirostre</i> (Starke) Bry. eur. 23.
<i>Dicranoweisia</i> Lindb. 20.
<i>D. crispula</i> (Hedw.) Lindb. 20.
<i>Dicranum</i> Hedw. 20-22.
<i>D. Bergeri</i> Bland. 20.
<i>D. Bonjeani</i> De Not. 21.
<i>D. congestum</i> Brid. 21.
<i>D. fuscessens</i> Turn. 21.
<i>D. Mühlenbeckii</i> Bry. eur. 21.
<i>D. parietinum</i> Hartz 21.
<i>D. Sauteri</i> Bry. eur. 21.
<i>D. scoparium</i> (L.) Hedw. 21, 22.
<i>D. scoparium</i> var. <i>alpestre</i> Milde 22.
<i>D. Scottianum</i> Turn. 22, 23.
<i>D. simplex</i> Goepp. et Menge 19.
<i>D. strictum</i> Schleich. 22.
<i>D. subflagellare</i> Goepp. et Menge 22.
<i>D. subpellucidum</i> Goepp. et Menge 20, 22.
<i>D. subskoparium</i> Goepp. et Menge 22.
<i>D. species</i> 22.
<i>Didymodon</i> Hedw. 26.
<i>D. alpigena</i> Vent. 26.
<i>D. rigidulus</i> Hedw. 26.
<i>D. rubellus</i> (Hoffm.) Bry. eur. 26.
<i>D. tophaceus</i> (Brid.) Jur. 26.
<i>Distichium</i> Bry. Eur. 18.
<i>D. capillaceum</i> (Sw.) Bry. eur. 18.
<i>Ditrichum</i> Timm 17.
<i>D. flexicaule</i> (Schleich.) Hampe 17.
<i>D. flexicaule</i> var. <i>densum</i> 18.
<i>D. tortile</i> (Schrad.) Lindb. 18, 56.
<i>Drepanocladus</i> Roth 62-75.
<i>D. aduncus</i> (Hedw.) Moenk. 62-63.
<i>D. aduncus</i> (Hedw.) Gr. Kneiffii 63-64.
<i>D. aduncus</i> (Hedw.) Gr. pseudofluitans 64.
<i>D. badius</i> (Hartm.) Roth 64.
<i>D. capillifolius</i> Warnst. 64. |
|---|

- D. exannulatus (Guemb.) Warnst. **65 - 66.**
 D. fluitans (L.) Warnst. **66-68,** 100.
 D. fluitans (L.) Gr. Rotae (De Not.) **66.**
 D. fluitans (L.) Warnst. var. pseudostramineus (C. M.) **68.**
 D. Hollosianus (Schilb.) Györrfy **68.**
 D. intermedius (Lindb.) Warnst. **68-69.**
 D. latifolius (Lindb. et Arn.) Broth. **69.**
 D. lycopodioides (Schwaegr.) Warnst. **69.**
 D. pseudofluitans (Sanio) **64.**
 D. pseudostramineum (C. M.) Roth **68.**
 D. revolvens (Sw.) Warnst. **69-70.**
 D. scorpioides (L.) Warnst. **70-71.**
 D. Sendtneri (Schimp.) Warnst. **72.**
 D. Sendtneri var. Wilsoni (Schimp.) **74.**
 D. subaduncus Warnst. **63.**
 D. trifarius (Web. et Mohr) Broth. **72-73.**
 D. Tundrae (Arn.) **73.**
 D. turgescens (Jens.) Broth. **73.**
 D. vernicosus (Lindb.) Warnst. **74.**
 D. Wilsoni (Schimp.) Roth **74.**
 D. species **75.**
Dryptodon patens Brid. **30.**
Elodium see *Helodium.*
Encalypta Schreb. **27-28.**
E. alpina Smith **27.**
E. rhabdocarpa Schwaegr. **27, 28.**
 Entodontaceae **54.**
Eucladium Bry. eur. **25.**
E. verticillatum (L.) Bry. eur. **23, 25.**
Eurhynchium Bry. eur. **95-96.**
E. circinatum Brid. var. *deflexifolium* (Solms) Boul. **93.**
E. myosuroides (Dill.) Schimp. **53.**
E. myurum (Poll.) Dixon **54.**
E. praelongum Bry. eur. **94, 95.**
E. rusciforme (Neck.) Milde **95.**
E. speciosum (Brid.) Milde **56, 95.**
E. Stokesii (Turn.) Bry. eur. **94, 95.**
E. striatulum (Spr.) Bry. eur. **95.**
E. striatulum var. *integrifolium* **96.**
E. striatum (Schreb.) Schimp. **96.**
E. strigosum (Hoffm.) Bry. eur. **96.**
E. Swartzii (Turn.) Hobk. **94, 95.**
E. Tommasinii (Sendtn.) Ruthe **94.**
E. species Eng. et Kink. **94.**
Fabronia Raddi **54.**
F. octoblepharis Schleich. **104.**
 Fabroniaceae **54.**
Fissidens Hedw. **23-24.**
F. adiantoides (L.) Hedw. **23.**
F. cristatus Wils. **23.**
F. decipiens De Not. **23.**
F. julianus Sav. **23.**
F. osmundoides (Sw.) Hedw. **24.**
F. taxifolius (L.) Hedw. **24.**
 Fissidentaceae **23-24.**
Fontinalaceae **49-50.**
Fontinalis (Dill.) L. **49-50.**
F. antipyretica L. **49.**
F. hypnoides Hartm. **49.**
F. pristina Lesq. **49.**
F. Sismondiana Schimp. **49.**
F. Tournalii (Brongn.) Schimp. **49, 105.**
F. species **50.**
 Glyptomitrium Brid. **28.**
G. Cockerelleae Brid. et Hollick **28.**
Glyptostrobus Unger. Heer **97.**
Grimmia Ehrh. **28-29.**
G. alpicola Sw. **28.**
G. anodon Bry. eur. **28.**
G. apocarpa (L.) Hedw. **28.**
G. elatior Bruch **28.**
G. elongata Kaulf. **28, 29.**
G. ericoides Lindb. **29, 30.**
G. funalis (Schwaegr.) Schimp. **29.**
G. hypnoides Lindb. **29, 30.**
G. incurva Schwaegr. **29.**
G. Mühlenbeckii Schimp. **29.**
G. robusta **29.**
G. subelongata Goepp. et Menge **29.**
G. torquata Hornsch. **29.**
G. trichophylla Grev. **29.**
G. species **29.**
 Grimmiaceae **28-29.**
Gymnostomum Hedw. **24.**
G. calcareum Bry. germ. **24.**
G. ferrugineum Ludw. **14, 24.**

- | | |
|---|---|
| <i>Hedwigiaceae</i> 48.
<i>Hedwigia</i> Ehrh. 48.
<i>H. albicans</i> (Web.) Lindb. 48.
<i>Helodium</i> (Sull.) Warnst. 58-59.
<i>H. Blandovii</i> (Web. et Mohr) 59.
<i>H. lanatum</i> (Stroem) Broth. 58.
<i>Heterocladium</i> Bry. eur. 54.
<i>H. squarrosum</i> (Voit) Lindb. 54.
<i>H. species</i> 55.
<i>Homalia</i> (Brid.) Bry. eur. 52.
<i>H. trichomanoides</i> (Schreb.) Bry. eur. 52.
<i>Homalothecium</i> Bry. eur. 90.
<i>H. sericeum</i> (L.) Bry. eur. 56, 90.
<i>Homomallium</i> (Schimp) Loeske 60.
<i>H. incurvatum</i> (Schrad.) Loeske 60.
<i>Hornea</i> Kidst. et Lang 106.
<i>Hygroamblystegium</i> Loeske 60-61.
<i>H. curvicaule</i> (Jur.) Loeske 60.
<i>H. fallax</i> (Brid.) Loeske 60.
<i>H. filicinum</i> (L.) Broth. 61.
<i>H. Heppi</i> (Heer) Broth. 98.
<i>Hygrohypnum</i> Lindb. 80-81.
<i>H. alpestre</i> (Sw.) Loeske 80.
<i>H. dilatatum</i> (Wils.) Loeske 80.
<i>H. lignitorum</i> (Schimp.) Dixon 80.
<i>H. molle</i> (Dicks.) Loeske 80.
<i>H. ochraceum</i> (Turn.) Loeske 80-81.
<i>H. palustre</i> (Huds.) Loeske 81.
<i>H. Smithii</i> (SW.) Broth. 81.
<i>Hylocomium</i> Bry. eur. 85-86.
<i>H. brevirostre</i> (Ehrh.) Bry. eur. 85.
<i>H. loreum</i> (Dill.) Bry. eur. 84.
<i>H. parietinum</i> (L.) Lindb. 21, 86.
<i>H. proliferum</i> (L.) Lindb. 85.
<i>H. pyrenaicum</i> (Spr.) Lindb. 86.
<i>H. rugosum</i> (Ehrh.) Bry. eur. 85.
<i>H. Schreberi</i> (Willd.) De Not. 86.
<i>H. splendens</i> (Dill.) Bry. eur. 85.
<i>H. squarrosum</i> (L.) Bry. eur. 84.
<i>H. triquetrum</i> (L.) Bry. eur. 84.
<i>H. species</i> 86.
<i>Hymenostomum</i> R. Br. 24.
<i>H. microstomum</i> (Hedw.) R. Br. 24.
<i>H. trichostomum</i> Goepp. 24.
<i>Hyocomium</i> Bry. eur. 83. | <i>H. flagellare</i> (Dicks.) Bry. eur. 83.
<i>Hypnaceae</i> 59-90.
<i>Hypnites</i> Ett. 97.
<i>H. Haeringianus</i> Ett. 97.
<i>Hypnodendron</i> (C. M.) Lindb. 96.
<i>H. arborescens</i> (Mitt.) Lindb. 96.
<i>H. Reinwardtii</i> (Hornschr.) Lindb. 96.
<i>Hypnum</i> auct. 62.
<i>Hypnum</i> Dill. ex p. 86.
<i>H. aduncum</i> Hedw. 62, 63.
<i>H. aduncum</i> var. <i>groenlandicum</i> Heer 63.
<i>H. aduncum</i> var. <i>Kneiffii</i> <i>groenlandicum</i> 63.
<i>H. aduncum</i> var. <i>paternum</i> Sanio 64.
<i>H. aduncum</i> var. <i>pseudofluitans</i> Sanio 64.
<i>H. aduncum</i> var. <i>sparsifolium</i> (Warnst.) 63.
<i>H. albicans</i> Meck. 91, 92.
<i>H. alpestre</i> Sw. 80.
<i>H. arcticum</i> Sommert. 81.
<i>H. amrisanense</i> Schimp. 97, 103.
<i>H. Brownii</i> Kirchner 97.
<i>H. callichroum</i> (Brid.) C. M. 87.
<i>H. capillifolium</i> Warnst. 64.
<i>H. carbonarium</i> Ludw. 97.
<i>H. chrysophyllum</i> Brid. 81.
<i>H. cirrosum</i> Schwaegr. 94.
<i>H. columbianum</i> Penh. 97.
<i>H. commutatum</i> Hedw. 61, 62.
<i>H. cypressiforme</i> L. 87.
<i>H. cuspidatum</i> L. 79.
<i>H. dilatatum</i> Wils. 80.
<i>H. diluvianum</i> Schimp. 75.
<i>H. elodes</i> Spr. 81.
<i>H. exannulatum</i> Guemb. 65, 66.
<i>H. exannulatum</i> var. <i>brachydictyon</i> Ren. 65.
<i>H. exannulatum</i> var. <i>orthophyllum</i> Milde 65.
<i>H. exannulatum</i> Gr. Rotae 66.
<i>H. falcatum</i> Brid. 62.
<i>H. fastigiatum</i> (Brid.) Hartm. 88.
<i>H. fertile</i> Sendtn. 88.
<i>H. filicinum</i> L. 61, 62.
<i>H. fluitans</i> L. 66, 67, 68.
<i>H. fluitans</i> var. <i>brachydictyon</i> Ren. 65.
<i>H. fluitans</i> <i>exannulatum</i> 65.
<i>H. fluitans</i> <i>exannulatum</i> var. <i>pinnatum</i> Boul. f. 65.
<i>H. fluitans</i> var. <i>falcatum</i> Schimp. 67. |
|---|---|

- H. fluitans* var. *falcitolium* Ren. 66.
H. fluitans t. *fossilis* Geheeb 67.
H. fluitans var. *glaciale* Ren. 66.
H. fluitans var. *tenuissimum* Schimp. 67.
H. giganteum Schimp. 75, 76.
H. haeringianum Schimp. 97.
H. hamulosum Bry. eur. 88.
H. Haydenii Lesq. 97.
H. Heerii Schimp. 97, 98.
H. Heppii Heer 98.
H. Hollosianum Schilb. 68.
H. incurvatum Schrad. 60.
H. insubricum Farneti 81.
H. intermedium Lindb. 68, 69.
H. irrigatum Zett. 62.
H. Kneiffii (Bry. eur.) Schimp. 64.
H. lignitorum Schimp. 80.
H. Lindbergii Mitt. 87.
H. lutescens Huds. 91.
H. lycopodioides Brid. 69.
H. lycopodioides Weber 100.
H. Mastodontum Boul. 98.
H. moile Dicks. 80.
H. molluscum Hedw. 83.
H. Noegerrathii Weber 63.
H. ochraceum Turn. 80.
H. oeningense Schimp. 57.
H. palustre Huds. 81.
H. plumula Schimp. 98.
H. polygamum (Bry. eur.) Wils. 82.
H. praelongum L. 94, 95.
H. pratense Koch 88.
H. priscum Schimp. 98.
H. protensum Brid. 82.
H. purpurascens Limpr. 65.
H. purpurascens var. *Rotae* (De Not.) 66.
H. purum L. 93.
H. recurvans Schwaegr. 88.
H. reptile (Rich.) 88.
H. revolvans Sw. 69, 70.
H. Richardsoni (Mitt.) Lesq. et James 77.
H. rutabulum L. 92.
H. saiebrosum Hoffm. 93.
H. Saportanum Schimp. 99, 104.
H. sarmentosum Wahl. 77, 78.
H. Schimperi Heer 97, 99.
H. Schimperi (Ung.) Schimp. 99, 104.
H. Schreberi Willd. 86.
H. scorpioides L. 70, 71.
H. Sendteri Schimp. 72.
H. sericeum L. 90.
H. splendens Dill. 85.
H. squarrosum L. 84.
H. stellatum Schreb. 83.
H. stramineum Dicks. 78.
H. striatum Schreb. 96.
H. strigosum Hoffm. 96.
H. Swartzii Turn. 95.
H. Taramellianum Farneti 68, 99.
H. trichoides Neck. 90.
H. trifarium (Web. et Mohr) 72, 73.
H. turgescens Jens. 73.
H. Ungerii Schimp. 99, 102.
H. Vaucherii Lesq. 89.
H. velutinum L. 93.
H. vernicosum Lindb. 74.
H. viride Hartz 99.
H. Weberianum Goepp. 72, 100.
H. Wilsonii Schimp. 74.
H. species? Hollos 68.
H. species 100.
Isopterygium Mitt. 89.
I. pulchellum (Dicks.) Jaeg. 89.
Isothecium Brid. 53-54.
I. myosuroides (Dill.) Brid. 53.
I. myurum (Poli.) Brid. 53.
I. viviparum (Neck.) Lindb. 53.
Leersia alpina Lindb. 27.
L. rhabdocarpa Lindb. 27.
Lembophyllaceae 53-54.
Leptobryum Wils. 31.
L. pyriforme (L.) Wils. 31.
Leptodictyum riparium (L.) Warnst. 59.
Lescuraea Bry. eur. 55.
L. striata (Schwaegr.) Bry. eur. 55.
Leskea Hedw. 55-56.
L. polycarpa Ehrh. 55.
L. species 56.
Leucodon Schwaegr. 50.
L. sciuroides (L.) Schwaegr. 50-51.
Leucodontaceae 50-51.
Limprichtia intermedia (Lindb.) Loeske 69.
L. vernicosa (Lindb.) Loeske 74.
Lycopodites Meekii Lesq. 100.
L. squamatus Brongn. 105.
Meesea Hedw. 41-43.
M. Albertini Bry. eur. 41.
M. longiseta Hedw. 42.
M. trichodes (L.) Spruce 42.
M. triquetra (L.) Aongstr. 42, 43.
M. triquetra var. *timmiooides* Sanio 42.

M. tristicha (Funck) Bry. eur. 42.
 Meesaceae 41-43.
 Mniaceae 36-40.
Mniobryum Limpr. 33.
M. albicans (Wahl.) Limpr. 33.
M. albicans var. *glaciale* Limpr. 35.
Mnium L. 36-39.
M. affine Bland. 36, 38.
M. affine var. *integritolium* Lindb. 36.
M. antiquorum Card. et Dixon 37, 102.
M. cinctidiooides (Blytt) Hüben. 37.
M. curvatum (Lindb.) Limpr. 37.
M. cuspidatum (L.) Leyss. 37.
M. hornum L. 37.
M. hymenophylloides Hüben. 37, 38.
M. marginatum (Dicks.) P. Beauv. 38.
M. medium Bry. eur. 38.
M. punctatum (L.) Hedw. 38.
M. riparium Mitten. 38.
M. rostratum Schrad. 39.
M. rugicium Laur. 39.
M. subglobosum Bry. eur. 38, 39.
M. undulatum (L.) Weis 39.
M. species 39.
Mollia tortuosa Lindb. 25.
Muscites Brongn. 101-104.
M. apiculatus Goepp. et Ber. 26, 101.
M. Bertrandii Lignier 101.
M. confertus Goepp. et Ber. 24, 101.
M. dubius Goepp. et Ber. 101.
M. elatinus Sap. 57, 102.
M. elegans Goepp. 102.
M. fontinaloides Ung. 102.
M. Hauchecornei Casp. et Klebs 102.
M. hirsutissimus Goepp. et Ber. 102.
M. hypnoides Ung. 99, 102.
M. intricatus Sap. 102.
M. journacensis Marty 103.
M. oenigenensis Braun 57, 103.
M. pereger Sap. 103.
M. polytrichaceus Ren. et Zeili. 103.
M. pulvinatus Sap. 103.
M. redivivus Sap. 104.
M. Schimperi Ung. 99, 104.
M. serratus Goepp. et Ber. 54, 104.
M. setosus Sap. 104.
Fossilium Catalogus II. 13

M. squamatus Brongn. 105.
M. Sternbergianus Dunker 105.
M. tortifolius Casp. et Klebs 105.
M. Tournalii Brongn. 49, 105.
Najadita Bruckm. 105.
Neckera Hedw. 51-52.
N. complanata (L.) Hüben. 51.
N. crispa (L.) Hedw. 52.
N. pennata (L.) Hedw. 52.
N. pumila Hedw. 52.
N. species 52.
Neckeraceae 51-53.
Oligotrichum Lam. et De Cand. 45.
O. hercynicum (Huds.) Lindb. 45.
Oncophorus Brid. 20.
O. virens (Sw.) Brid. 20.
Oreoweisia De Not. 20.
O. serrulata (Funck) De Not. 20.
Orthothecium Bry. eur. 54.
O. chrysseum (Schwaegr.) Bry. eur. 54.
O. intricatum (Hartm.) Bry. eur. 54.
O. rufescens (Dicks.) Bry. eur. 54.
Orthotrichaceae 30-31.
Orthotrichum Hedw. 31.
O. diaphanum (Gmel.) Schrad. 31.
O. species 31.
Oxyrrhynchium Warnst. 94-95.
O. praelongum (Bry. eur.) Warnst. 94-95.
O. rusciforme (Neck.) Warnst. 95.
O. speciosum (Brid.) Warnst. 95.
O. Swartzii (Turn.) Warnst. 95.
Paludella Ehrh. 41.
P. squarrosa (L.) Brid. 41.
Phascum L. 26.
P. cuspidatum Goepp. 26.
Philonotis Brid. 43-44.
P. alpicola Jur. 44.
P. calcarea (Bry. eur.) Schimp. 43.
P. coespitosa Wils. 43.
P. fontana (L.) Brid. 43-44, 56, 103.
P. fontana var. *ampliretis* Dixon 44.
P. fontana var. *glacialis* 44.
P. seriata Mitt. 44.
P. tomentella Mol. 44.

- P. species 44.
Pinnatella (C. M.) Fleisch. 53.
P. alopecuroides (Hook.)
Fleisch. 53.
P. species 53.
Plagiopodopsis Britton et Hollick 43.
P. Scudderi Britton et Hollick 43.
Plagiothecium Bry. eur. 89-90.
P. denticulatum L. *Roeseanum*
Hamp. 89.
P. latebricola (Wils.) Bry. eur.
89.
P. Muehlenbeckii (Schimp.)
Bry. eur. 89.
P. pulchellum (Dicks.) Bry. eur.
89.
P. Roeseanum (Hamp.) Bry. eur.
89.
P. Ruthei Limpr. 89.
P. Ruthei var. *rupinicola*
Limpr. 89.
P. silvaticum (Huds.) Bry. eur.
89.
P. striatellum (Brid.) Lindb. 89.
P. undulatum (L.) Bry. eur. 90.
Pogonatum P. Beauv. 45-46.
P. suburnigerum (Goepp. et
Menge) Dixon 45.
P. urnigerum (L.) P. Beauv. 46.
Pohlia Hedw. 32-33.
P. albicans Lindb. 33.
P. commutata (Schimp.) Lindb.
32.
P. cruda (L.) Lindb. 32.
P. cucullata (Schwaegr.) Bruch
32.
P. longicolla (Sw.) Lindb. 32.
P. Ludwigii (Spreng.) Lindb. 32.
P. nutans (Schreb.) Lindb. 32.
P. sphagnicola (Bry. eur.)
Lindb. 33.
Polytrichaceae 45-48.
Polytrichum L. 46-48.
P. alpinum L. 46.
P. attenuatum Menz. 46.
P. commune L. 46.
P. Florissanti Knowlton 47.
P. gracile Menz. 47, 101.
P. juniperinum Wild. 47, 101.
P. piliferum Schreb. 47.
P. pilosum Floercke 47.
P. sexangulare Floercke 47.
P. strictum Banks 47, 48.
P. strictum var. *alpestre*
Limpr. 48.
P. subseptentriionale Goepp. et
Menge 48.
P. subundulatum Goepp. et
Menge 45.
- P. suburnigerum* Goepp. et
Menge 45.
P. species 48.
Porotrichum alopecurum (L.)
Mitt. 53.
Pottiaceae 24-28.
Pseudocaliergon trifarium (Web.
et Mohr) Loeske 73.
P. turgescens (Jens.) Loeske 73.
Pseudoleskea Bry. eur. 56.
P. atrovirens (Dicks.) Bry. eur.
56.
P. filamentosa (Dicks.) Broth.
56.
P. patens Limpr. 56.
Ptilium (Sull.) De Not. 87.
P. crista-castrensis (L.) De Not.
87.
Pylaisia Bry. eur. 54.
P. polyantha (Schreb.) Bry. eur.
54.
Rhacomitrium Brid. 29-30.
R. affine Lindb. 29.
R. canescens (Weis) Brid. 29, 30.
R. fasciculare (Schrad.) Brid. 30.
R. heterostichum (Hedw.) Brid.
30.
R. heterostichum var. 29.
R. hypnoides (L.) Lindb. 30.
R. lanuginosum Brid. 30.
R. patens (Brid.) 30.
Rhynchosporium Bry. eur. 96.
R. Knowltoni E. G. Britt. 96.
Rhynia Kidst. et Lang 106.
Rhytidadelphus (Lindb.)
Warnst. 84.
R. lorenus (Dill.) Warnst. 84.
R. squarrosus (L.) Warnst. 84.
R. triquetus (L.) Warnst. 84.
Rhytidium (Sull.) Kindb. 85.
R. rugosum (Ehrh.) Kindb. 85.
Schistidium alpicola var. *rivulare* Limpr. 28.
Schistophyllum adiantoides
Brid. 23.
S. juleanum (Sav.) Lindb. 23.
Schlotheimia tenuifolia Sternb.
105.
Scleropodium Bry. eur. 93.
S. purum (L.) Limpr. 93.
Scorpidium scorpioides (L.)
Limpr. 71.
Scorpiurium Schimp. 93.
S. deflexifolium (Solms) Schimp.
93.
Sematophyllum recurvans
(Schwaegr.) E. G. Britt. 88.
Sequoia Haydenii (Lesq.) Cock.
97.

- Sphaerocephalus palustris* (L.)
Lindb. 40.
Sphagnaceae 12-17.
Sphagnum Erhrh. 12-17.
S. acutifolium Ehrh. 12.
S. brevifolium Röll 15.
S. compactum De Cand. 12.
S. cuspidatum Ehrh. 13, 14.
S. cymbifolium Ehrh. 13, 14.
S. fimbriatum Wils. 13.
S. Girgensohnii Rüss. 14.
S. imbricatum (Hornsch.) Rüss. 14.
S. intermedium Hoffm. 14, 15.
S. Lindbergii Schimp. 14.
S. Ludwigii Schimp. 14.
S. magellanicum Brid. 14.
S. medium Limpr. 14, 15.
S. palustre Linn. 13, 15.
S. papillosum Lindb. 15.
S. recurvum P. Beauv. 15.
S. recurvum var. *parvifolium* Warnst. 15.
S. rubellum Wils. 15.
S. squarrosum Pers. 15.
S. subbicolor Hampe 15.
S. subsecundum Nees 14, 16.
S. teres Aongstr. 13, 16.
S. Wilsoni Röll. 15, 16.
S. species 16, 17.
Sphenolepidium Sternbergianum (Dunk.) 105.
Splachnaceae 31.
Splachnum L. 31.
S. ampullaceum L. 31.
S. sphaericum (L. f.) Sw. 31.
Sporogonites Halle 105-106.
S. exuberans Halle 17, 106.
Stereodon (Brid.) Mitt. 87-89.
S. arcuatus Lindb. 87.
S. callichrous Brid. 87.
S. chrysanthus (Schwaegr.) Bry. eur. 54.
S. cupressiformis (L.) Brid. 87.
S. cupressiformis var. *filiformis* (Brid.) 87.
S. fastigiatus Brid. 88.
S. fertilis (Sendtn.) Lindb. 88.
S. hamulosus (Bry. eur.) Lindb. 88.
S. paflescens (Hedw.) Lindb. 88.
S. pratensis (Koch) Warnst. 88.
S. recurvans (Schwaegr.) Broth. 88.
S. reptilis (Rich.) Mitt. 88.
S. revolutus Mitt. 89.
S. Vaucheri (Lesq.) Lindb. 89.
Swartzia montana Lindb. 18.

Thamnium Bry. eur. 53.
- Th. alopecurum* (L.) Bry. eur. 53.
Th. species 53.
Thuidium Bry. eur. 56-58.
Th. abietinum (Dill.) Bry. eur. 56.
Th. antiquum Schimp. 56.
Th. Blandowii (Web. et Mohr)
Bry. eur. 58.
Th. decipiens De Not. 62.
Th. delicatulum (Dill.) Mitt. 57.
Th. elatinum (Sap.) Schimp. 57,
102.
Th. lanatum (Stroem.) Hagen 59.
Th. minutulum (Hedw.) Bry.
eur. 57.
Th. oeningense (Braun) Schimp.
57.
Th. Philiberti Limpr. 58.
Th. Philiberti var. *pseudo-tama-*
risci (Limpr.) Ryan et Ha-
gen 58.
Th. recognitum Lindb. 57, 58.
Th. tamariscinum (Hedw.) Bry.
eur. 58.
Th. species 58.
Timmia Hedw. 44-45.
T. austriaca Hedw. 44.
T. bavarica Hessl. 45.
T. megapolitana Hedw. 45.
T. norvegica Zett. 45.
Timmiceae 44 - 45.
Tomentypnum nitens (Schreb.)
Loeske 911.
Tortella Limpr. 25.
T. fragilis (Hook. f. et Wils.)
Limpr. 25.
T. inclinata (Hedw. f.) Limpr.
25.
T. tortuosa (L.) Limpr. 25.
Tortula Hedw. 27.
T. aciphylla Hartm. 27.
T. aciphylla var. *mucronata*
Sendtn. 27.
T. norvegica (Web.) Wahl. 27.
T. ruralis (L.) Ehrh. 27.
T. species 27.
Trichosteleum Boschi (Doz. et
Molk.) 104.
Trichostomum Hedw. 25.
T. polyphyllum 25.
T. polystichum (?) in Goepp. et
Menge 25.
T. strictum (Sw. or Bry. eur.)
26.
T. subpolystichum Goepp. et
Menge 25.
T. substrictum Goepp. et Menge
25.

Uliota Mohr 31.
U. curvifolia (Wahl.) Brid. 31.

- | | |
|--|---|
| <i>Warnstorffia exannulata</i> | W. nutans Hedw. 32, 33. |
| (Guemb.) Loeske 66. | <i>W. sphagnicola</i> Schimp. 33. |
| <i>Webera albicans</i> Schimp. 33. | <i>Weisia</i> Hedw. 24. |
| <i>W. commutata</i> Schimp. 32. | <i>W. conferta</i> Schimp. 24, 101. |
| <i>W. cruda</i> (L.) Bruch 32. | <i>Widdringtonia helvetica</i> Heer 97. |
| <i>W. cucullata</i> (Schwaegr.)
Schimp. 32. | <i>Zygodon</i> Hook. et Tayl. 31. |
| <i>W. longicolla</i> (Sw.) Hedw. 32. | <i>Z. viridissimus</i> (Dicks.) R. Br. |
| <i>W. Ludwigii</i> Schimp. 32. | 31. |



