

Chemical Constituents of *Eremostachys macrophylla* Montbr. & Auch. Aerial Parts

Parina Asgharian^{1,2}, Abbas Delazar^{1,2}, Solmaz Asnaashari^{3*}

¹Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

²Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

³Biotechnology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

Article Info

Article History:

Received: 1 June 2019

Accepted: 7 October 2019

ePublished: 27 June 2020

Keywords:

-*Eremostachys macrophylla*

-Iridoid

-Phenylethanoid

-Flavonoid

Abstract

Background: *Eremostachys macrophylla* Montbr. & Auch. is one of the wild growing species of herbs found in East Azerbaijan province of Iran. These species are used in folk medicine for the healing of wound, treatment of snake bites, rheumatism and joint pains. The primary aim of this study was to obtain natural pure compounds and this was done by subjecting the aerial parts of *Eremostachys macrophylla* Montbr. & Auch. to phytochemical analysis.

Methods: The air-dried and crushed aerial parts were respectively extracted with n-hexane, dichloromethane (DCM) and methanol (MeOH) solvents using a soxhlet apparatus. The 10%, 20% and 40% of MeOH in water Sep-Pak fractions of the MeOH extract were subjected to a preparative reversed-phase high performance liquid chromatography (RP-HPLC). Also, the isolated pure compounds were identified by one-dimensional nuclear magnetic resonance (1D-NMR) spectroscopic technique

Results: The results obtained in this study showed the presence of seven pure components; (1) Lamalbide, (2) Sesamoid, (3) Phlomiol, (4) Verbascoside, (5) Luteolin-7-O- glucoside, (6) Apigenin-7-O- rutinoside and (7) Kaempferol-3-O- glucoside with iridoid, phenylethanoid and flavonoid structures.

Conclusion: The results from the study demonstrated that the aerial parts of *E. macrophylla* could be a good source of iridoids, phenylethanoids and flavonoids.

Introduction

The genus *Eremostachys* belongs to the Lamiaceae family. It is made up of about 60 species of perennial herbs, which originate generally in Central, Middle-East and South-West Asian countries.^{1,2} This genus is presented by fifteen species in Iran flora.³ Some plants of this genus are traditionally used in making local analgesia, for treating inflammation, allergy, for wound healing, for treating snake bites, rheumatism, joint pains, headache and liver diseases.^{1,3,4} Moreover, previous studies have shown the wide range of biological activities from the genus *Eremostachys*; for example, according to published reports, *E. laciniata* has possessed analgesic, antinociceptive, antidepressant, antioxidant, antibacterial and anti-inflammatory effects.^{2,3,5-8}

In pursuant to previous literatures, phytochemical studies about the essential oils of some *Eremostachys* species revealed the presence of terpenoids, linear and branched hydrocarbons and derivatives as the main constituents in different stages of growth. The analysis of the volatile oil of *E. adenatha* using Gas Chromatography Mass Spectroscopy (GCMS) demonstrated the presence of

dodecanal, hexadecanoic acid and 6, 10, 14-trimethyl-2-pentadecanone as major components; whereas, 6, 10, 14-trimethyl-2-pentadecanone, 1, 8-cineole and α -pinene were the main compound of the essential oil of *E. labiosa* aerial parts and the chief constituents identified from the stem oil were α -phellandrene, β -phellandrene, α -pinene and tetradecane.^{9,10} The analysis of the *E. laevigata* aerial parts essential oil represented dodecanal, germacrene-D, β -caryophyllene and caryophyllene oxide structures as the main components.¹¹ Although, the major constituent of *E. laciniata* aerial parts was dodecanol,¹² the chief components of the *E. azerbaijanica* oil were hexahydrofarnesyl acetone, 2-methyl-6-propyl-dodecane and tricosane.¹³ *Eremostachys macrophylla* Montbr. & Auch or "Sonbole biabani" is a species which is distributed in Turkey, Iraq, Iran and Central Asia. This wild-growing species have been used for treatment of snake bites, healing of wounds, rheumatism and joint pains.^{14,15} There are several studies about phytochemical analysis of *E. macrophylla* essential oil. According to Javidnia et al. report, spathulenol, hexadecanoic acid and caryophyllene oxide were the main

*Corresponding Author: Solmaz Asnaashari, E-mail: asnaasharisolmaz@gmail.com

©2020 The Author(s). This is an open access article and applies the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited.

compounds of this species.⁹ Likewise, in the other study by Nori-shargh et al., the oil of *E. macrophylla* aerial parts consisted essentially of germacrene-D, germacrene-B and γ -elemene. Rustaiyan et al. assessed different parts of *E. macrophylla* separately and their reports demonstrated that 1,8-cineol and germacrene D-4-ol were the major structures in the flower oil, while the leaf oil contained α -pinene, 1, 10-di-epi cubenol, elemol and bornyl acetate. The oil of the stem was dominated also by 1, 10-di-epi cubenol and elemol.¹⁰

No study has ever been performed on the non-volatile phyto-constituents of *E. macrophylla*, except for our previous evaluations on the phyto-constituents of *E. macrophylla* rhizomes, which represented two iridoid structures (Lamalbide and 6-Hydroxyl Loganin) in 2018.¹⁶ Thus, the aim of this study was to extract, purify and identify natural constituents of *E. macrophylla* aerial parts grown in East Azerbaijan province, Iran.

Materials and Methods

Materials

All the solvents that were used for extraction and fractionation were bought from Caledon (Canada), D₂O and d₆-DMSO for NMR spectroscopy were purchased from Sigma (USA).

Plant material

The aerial parts of *E. macrophylla* Montbr. & Auch. were collected in the month of July, 2012 from Sahand mountains in East Azerbaijan province of Iran on 37.759 (37° 45' 32.4" N) latitude, 45.9783 (45° 58' 41.9" E) longitude and an altitude of 1950m above sea level. It was taxonomically identified and deposited by Dr. Hossein Nazemiyeh and Dr. Atefeh Ebrahimi in the Herbarium of Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran (Voucher Nos. TBZ-fph-739).

Extraction, separation and identification of compounds

The air-dried and ground aerial parts of *E. macrophylla* (50g) were extracted with n-hexane, dichloromethane (DCM) and MeOH, with a soxhlet apparatus (500mL, throughout 8 hours for each solvent), sequentially (solvents were from Caledon, Canada).

The MeOH extract (2g) was subjected to solid phase extraction (SPE) using a C₁₈ Sep-Pak cartridge (Waters, USA), eluting with a step gradient of MeOH/water mixture (10:90, 20:80, 40:60, 60:40, 80:20 and 100:0). All these extracts and fractions were separately concentrated using a rotary evaporator at a maximum temperature of 45°C. The 10%, 20% and 40% MeOH in water Sep-Pak fraction were also subjected to preparative reversed-phase HPLC (prep-HPLC) conducted on a Knauer HPLC (preparative pump 1800) fitted with a C₁₈ column (250mm length, 20mm i.d, 10 μ m particle size, Dr. Maisch, Germany) system.

The mobile phase time program for each fraction was set as below:

The 10% fraction: linear gradient of 0-36% MeOH in water

during 0-50 min, isocratic gradient of 36% MeOH in water during 50-62.5 min, linear gradient of 36-0% during 62.5-67.5 min, isocratic gradient of 0% MeOH in water during 67.5-75 min to yield compound (1) (8.4mg, 51.1% of 10% fraction) and compound (2) (4.3mg, 21.8% of 10% fraction).

The 20% fraction: linear gradient of 20-40% MeOH in water during 0-50 min, isocratic gradient of 40% MeOH during 50-62.5 min, linear gradient of 40-20% MeOH in water during 62.5-67.5 min, isocratic gradient of 20% MeOH in water during 67.5-75 min to isolate compound (3) (3mg, 3.3% of 20% fraction).

The 40% fraction: linear gradient of 30-60% MeOH in water during 0-50 min, isocratic gradient of 60% MeOH during 50-62.5 min, linear gradient of 60-30% MeOH in water during 62.5-67.5 min, isocratic gradient of 30% MeOH in water during 67.5-75 min to isolate compound (4) (4.7mg, 6.1% of 40% fraction), compound (5) (9.1mg, 46.5% of 40% fraction), compound (6) (4.5mg, 1.8% of 40% fraction) and compound (7) (3.1mg, 1.7% of 40% fraction).

Flow rate was 8 ml/min, volume of each injection was 1 ml and the detector set at 220 nm was used to monitor chromatogram. Elucidation of structures was performed using ¹H-NMR and ¹³C-NMR (Bruker, Germany) and the spectroscopic data of the known compounds were also compared with the respective published data.¹³

Results and Discussion

The reversed-phase preparative HPLC analysis of 10%, 20% and 40% fraction of the MeOH extract of *E. macrophylla* aerial parts afforded three iridoid glycoside, one phenylethanoid glycoside and three flavonoid structures, which were identified unequivocally as (1) Lamalbide (White amorphous, 8.4mg, 51.1%, Rt: 51.35min), (2) Sesamoside (White amorphous, 4.3mg, 21.8%, Rt: 53.13min), (3) Phlomiol (White amorphous, 3mg, 3.3%, Rt: 53.65min), (4) Verbascoside (Yellow amorphous, 4.7mg, 6.1%, 23.38min), (5) Luteolin-7-O- glucoside (Brown amorphous, 9.1mg, 46.5%, 32.22min), (6) Apigenin-7-O- rutinoside (Brown amorphous, 4.5mg, 1.8%, Rt.36.50min) and (7) Kaempferol-3-O- glucoside (Brown amorphous, 3.1mg, 1.7%, Rt: 39.81min) on the extensive 1D H-NMR and C-NMR data analyses. The spectroscopic data of the known compounds were also compared with the respective published data.

All three compounds (1), (2) and (3) showed that UV, ¹H-NMR and ¹³C-NMR signals were consistent with iridoid glycoside skeletons.¹⁷

Compound (1):

λ_{\max} : 237 nm

¹H-NMR and ¹³C-NMR signals of compound (1) presented a methyl group at C₈ ($\sigma_{\text{H}10}$: 1.11ppm, $\sigma_{\text{C}10}$: 19.09ppm), a methoxy group ($\sigma_{\text{H}(\text{OCH}_3)}$: 3.66ppm, $\sigma_{\text{C}(\text{OCH}_3)}$: 50.35ppm), an olefinic methine at C₃ ($\sigma_{\text{H}3}$: 7.36ppm, $\sigma_{\text{C}3}$: 150.18ppm), oxymethines at C₁ ($\sigma_{\text{H}1}$: 5.54ppm, $\sigma_{\text{C}1}$: 92.28ppm), C₆ ($\sigma_{\text{H}6}$: 3.84ppm, $\sigma_{\text{C}6}$: 75.96ppm) and C₇ ($\sigma_{\text{H}7}$: 3.56 ppm, $\sigma_{\text{C}7}$: 75.96ppm).

σ_{C_7} : 73.97ppm), two methine at C5 (σ_{H_5} : 2.85ppm, σ_{C_5} : 33.70ppm) and C9 (σ_{H_9} : 2.74ppm, σ_{C_9} : 45.55ppm) and a β -glucose unit (σ_{H_1} : 4.67ppm, σ_{C_1} : 96.60ppm).

Compound (2):

λ_{\max} : 239 nm

The $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ data of compound (2) showed the presence of a methyl group at C₈ ($\sigma_{H_{10}}$: 1.39ppm, $\sigma_{C_{10}}$: 14.36ppm), a methoxy group ($\sigma_{H(\text{OCH}_3)}$: 3.67ppm, $\sigma_{C(\text{OCH}_3)}$: 50.61ppm), an olefinic methine at C₃ (σ_{H_3} : 7.60ppm, σ_{C_3} : 153.73ppm), oxymethines at C₁ (σ_{H_1} : 5.62ppm, σ_{C_1} : 93.31ppm), C₆ (σ_{H_6} : 4.30ppm, σ_{C_6} : 74.73ppm) and C₇ (σ_{H_7} : 3.53ppm, σ_{C_7} : 70.88ppm), a methine at C₉ (σ_{H_9} : 2.56ppm, σ_{C_9} : 50.42ppm) and a β -glucose unit (σ_{H_1} : 4.70ppm, σ_{C_1} : 96.99ppm).

Compound (3):

λ_{\max} : 229 nm

The $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ data of compound (3) indicated the existence of a methyl group at C₈ ($\sigma_{H_{10}}$: 1.06ppm, $\sigma_{C_{10}}$: 18.99ppm), a methoxy group ($\sigma_{H(\text{OCH}_3)}$: 3.67ppm, $\sigma_{C(\text{OCH}_3)}$: 53.94ppm), an olefinic methine at C₃ (σ_{H_3} : 7.51ppm, σ_{C_3} : 155.25ppm), oxymethines at C₁ (σ_{H_1} : 5.79ppm, σ_{C_1} : 94.20ppm), C₆ (σ_{H_6} : 4.03ppm, σ_{C_6} : 80.46ppm) and C₇ (σ_{H_7} : 3.65ppm, σ_{C_7} : 74.34ppm), a methine at C₉ (σ_{H_9} : 2.51ppm, σ_{C_9} : 58.10ppm) and a β -glucose unit (σ_{H_1} : 4.66ppm, σ_{C_1} : 100.40ppm).

These three structures spectroscopic data were compared with the former published data and the presence of Lamalbid (1), Sesamoside (2) and Phlomiol (3) were established.¹⁷⁻¹⁹

Lamalbid and Sesamoside were reported previously from *E. azerbaijanica* rhizomes, *E. moluccelloides* and *E. laciniata* aerial parts. In addition, Phlomiol was isolated from *E. laciniata* rhizomes.^{3,20-22} In addition, our earlier study on *E. macrophylla* rhizomes showed the presence of Lamalbid as an iridoid structure.¹⁶ Previous literature revealed a broad spectrum of biological activities of the compounds with iridoid structure, such as neuroprotective, hypoglycemic, anti-inflammatory, hypolipidemic, immunomodulator, choleric, hepatoprotective, cardioprotective, anti-cancer, anti-oxidant, anti-microbial, anti-spasmodic and purgative effects.²³ Moreover, several studies have been mentioned to antioxidant effect of Lamalbid and anti-tumor effect of Phlomiol.^{24,25}

Compound (4):

UV λ_{\max} (MeOH): 220, 290, 330nm.

The $^1\text{H-NMR}$ spectrum of structure (4) indicated the presence of a tri-substituted phenyl moiety characterized by signals appearing at σ 7.02(bs, 1H, H-2'), 6.96 (d, 1H, $J = 8.22\text{Hz}$, H-6') and 6.76 (d, 1H, $J = 8.08$, H-5'), a 3,4 di-hydroxyphenethyl alcohol moiety with proton resonances at σ 6.64 (s, 1H, H-2), 6.62 (bs, 1H, H-5), 6.50 (d, 1H, $J = 6.62\text{Hz}$, H-6), 3.72 (m, 2H, H- α) and 2.70 (bs, 2H, H- β) and olefinic protons of caffeic acid derivative at 6.21 (d, 1H, H- α') and 7.47 (d, 1H, H- β'). Anomeric proton

signals were presented at σ 4.36ppm (d, 1H, $J = 7.82\text{Hz}$, H-1') and σ 5.02ppm (bs, 1H, H-1'') that were consistent with the β -glucopyranose unit and α -rhamnopyranosyl moiety, respectively. The results identified the compound (4) as phenylethanoid glycoside, acteoside or verbascoside. The data were in agreement with published spectral data.¹³ This phenylethanoid structure had been previously reported from other species of the genus *Eremostachys* such as the aerial parts and rhizomes of *E. azerbaijanica* and *E. laciniata*.^{7,13,22,26} Early reports described several pharmacological benefits for human health of Verbascoside, including anti-inflammatory, anti-oxidant, anti-neoplastic, wound healing and neuroprotective effects.^{27,28}

The compounds (5), (6) and (7) from 40% MeOH fraction with flavonoid structures were confirmed with the UV and $^1\text{H-NMR}$ spectrum.

Compound (5):

UV λ_{\max} (MeOH): 207, 255, 350 nm

The $^1\text{H-NMR}$ data of compound (5) showed characteristic signals of 1, 2, 4- trisubstituted phenyl unit at σ 7.42ppm (bs, 1H, H-2'), 6.90ppm (d, 1H, $J = 8.26\text{Hz}$, H-5'), 7.45ppm (d, 1H, $J = 8.39\text{Hz}$, H-6'). Other signals at σ 6.75, 6.44 and 6.78ppm mentioned to the H-3, H-6 and H-8 in flavone, respectively. In addition, the anomeric proton signal appeared at σ 5.09ppm (d, 1H, $J = 7.25\text{Hz}$, H-1'').

Compound (6):

UV λ_{\max} (MeOH): 205, 265, 345 nm

In the same manner, The $^1\text{H-NMR}$ data of compound (6) presented the 1, 4- disubstituted phenyl unit at σ 7.94ppm (d, 2H, $J = 8.82\text{Hz}$, H-2' and H-6'), 6.92ppm (d, 2H, $J = 8.60\text{Hz}$, H-3' and H-5'). The signals of H-3, H-6 and H-8 were displayed at 6.78ppm, 6.44ppm and 6.84ppm, respectively. Two anomeric protons were observed at 4.86ppm and 5.04ppm, which related to rutinoid group.

Compound (7):

UV λ_{\max} (MeOH): 205, 265, 345 nm

The $^1\text{H-NMR}$ spectrum of compound (7) showed two doublet signals at σ 7.92ppm ($J = 8.62\text{Hz}$) and 6.87ppm ($J = 8.87\text{Hz}$) with orientation of two ortho situation protons for H-2', 6' and H-3', 5'. Two aromatic protons at σ 6.42ppm and 6.81ppm referring to H-6 and H-8 and a doublet signal at σ 5.07ppm ($J = 7.13\text{Hz}$) were related to an anomeric proton.

These spectral data revealed the existence of (5) Luteolin-7-O- glucoside, (6) Apigenin-7- o- rutinoid and (7) Kaempferol 3-O-glucoside structures. (25-27) According to previous literatures, Luteolin-7-O- glucoside was reported from the aerial parts of *E. laciniata*.²²

Flavonoids as a group of natural compounds with variable phenolic structures exhibit a wide range of bioactivities including anti-oxidant, anti-tumor, anti-inflammatory, anti-viral, anti-fungal, anti-bacterial and platelet aggregation inhibitory actions.²⁹ According to Ozcan and Matthaus, Luteolin-7-glucoside and Apigenin-7-glucoside

showed significant anti-oxidant activities as the phenolic compounds of olive leaves.³⁰ Moreover, anti-inflammatory effects of these two flavonoid structures were established previously.^{31,32} In other previous study, Parveen *et al.* presented significant anti-inflammatory and analgesic effects of Kaempferol 3-O-glucoside.³³

The focus on the phytochemical studies of a number of species of genus *Eremostachys* showed the presence

of various chemical structures. In some studies, iridoid glycosides, flavonoids, phenylethanoid glycosides and phytosterols were reported from the rhizomes or aerial parts of *E. laciniata*.^{6,7} Ferulic acid derivatives, furanolabdane-type diterpenoids, and phenylethanoid glycosides were identified from the rhizomes of *E. glabra* in other studies.^{1,34,35} Furthermore, several other studies reported iridoid glycoside, flavonoid, phenylethanoid, fatty acid




Figure 1. The suggested structures of seven pure compounds from *E. macrophylla* aerial parts: (1) Lamalbide, (2) Sesamoside, (3) Phloriol, (4) Verbascoside, (5) Luteolin-7-O-glucoside, (6) Apigenin-7-O-rutinoside and (7) Kaempferol-3-O-glucoside.

and steroid structures from the rhizome or aerial parts of *E. azerbaijanica*,^{13,36,37} iridoids and flavonoids from *E. loasifolia*,³⁸⁻⁴⁰ iridoid glycosides from the aerial parts of *E. moluccelloides*²¹ and various flavonoid derivatives from *E. vicaryi*⁴¹ were isolated previously.

This is the first report on the purification and identification of iridoid, phenylethanoid and flavonoid structure from the aerial parts of *E. macrophylla*.

Conclusion

The present study has shown that the aerial parts of *E. macrophylla* are a source of iridoid, phenylethanoids and flavonoids. Since the anti-oxidant and anti-inflammatory effects of phenylethanoids and flavonoids,⁴²⁻⁴⁶ analgesic, anti-inflammatory and anti-arthritic properties of iridoid glycosides have previously been confirmed by several *in vitro* studies,⁴⁷ it is therefore reasonable to conclude that *E. macrophylla* can be a good candidate for the studies on inflammatory diseases.

Acknowledgments

This assay was financially supported by the Drug Applied Research Center of Tabriz University of Medical Sciences.

Conflict of Interests

The authors claim that there is no conflict of interest.

References

- Delazar A, Shoeb M, Kumarasamy Y, Byres M, Nahar L, Modarresi M, et al. Two bioactive ferulic acid derivatives from *eremostachys glabra*. *Daru*. 2004;12(2):49-53.
- Nisar M, Khan S, Dar A, Rehman W, Khan R, Jan I. Antidepressant screening and flavonoids isolation from *eremostachys laciniata* (l) bunge. *S Afr J Sci*. 2011;10(9):1696-9.
- Modarresi M, Delazar A, Nazemiyeh H, Fathi-Azad F, Smith E, Rahman MM, et al. Antibacterial iridoid glucosides from *eremostachys laciniata*. *Phytother Res*. 2009;23(1):99-103. doi:10.1002/ptr.2568
- Asgharian P, Delazar A, Vatankhah A, Javadzadeh M, Asnaashari S. In vitro bioactivity and phytochemical evaluation of extracts from aerial parts of *Eremostachys macrophylla* montbr. & Auch. Growing in Iran. *Res J Pharmacogn*. 2017;4(2):65-73.
- Delazar A, Habibi Asl B, Mohammadi O, Heshmati Afshar F, Nahar L, Modarresi M, et al. Evaluation of analgesic activity of *Eremostachys laciniata* in mice. *J Nat Rem*. 2009;9(1):1-7. doi:10.18311/jnr/2009/213
- Khan S, Nisar M, Simjee SU, Rehman W, Khan R, Jan I, et al. Evaluation of micronutrients level and antinociceptive property of *Eremostachys laciniata* (L) bunge. *Afr J Biotechnol*. 2010;9(5):775-7.
- Delazar A, Sarker SD, Nahar L, Jalali SB, Modaresi M, Hamedeyazdan S, et al. Rhizomes of *Eremostachys laciniata*: Isolation and structure elucidation of chemical constituents and a clinical trial on inflammatory diseases. *Adv Pharm Bull*. 2013;3(2):385-93. doi:10.5681/apb.2013.062
- Delnavazi MR, Baba-ali F, Soufiabadi S, Sherafatmand M, Ghahremani F, Tavakoli S, et al. Essential oil composition, antioxidant activity and total phenolic content of some *Lamiaceae taxa* growing in northwest of Iran. *Pharm Sci*. 2014;20(1):22-8.
- Javidnia K, Miri R, Soltani M, Khosravi A. Essential oil composition of two species of *eremostachys* from Iran (*E. Adenantha jaub.* Et spach and *E. Macrophylla montbr.* Et auch.). *J Essent Oil Res*. 2008;20(3):226-8. doi:10.1080/10412905.2008.9699998
- Rustaiyan A, Masoudi S, Ezzatzadeh E, Akhlaghi H, Aboli J. Composition of the aerial part, flower, leaf and stem oils of *eremostachys Macrophylla montbr.* & auch. And *Eremostachys labiosa* bunge. From Iran. *J Essent Oil Bear Pl*. 2011;14(1):84-8. doi:10.1080/0972060X.2011.10643904
- Amiri H, Meshkat Al Sadat M, Lari Yazdi H. Chemical composition of the essential oil of *Eremostachys laevigata* bung. *Daru*. 2007;15(1):34-40.
- Navaei MN, Mirza M. Chemical composition of the oil of *Eremostachys laciniata* (l.) bunge from Iran. *Flavour Fragr J*. 2006;21(4):645-6. doi:10.1002/ffj.1635
- Asnaashari S, Afshar FH, Ebrahimi A, Moghadam SB, Delazar A. Chemical composition and radical scavenging activity of essential oil and methanolic extract of *Eremostachys azerbaijanica* rech. F. From Iran. *Res Pharm Sci*. 2016;11(2):113-9.
- Mosaddegh M, Naghibi F, Moazzeni H, Pirani A, Esmaeili S. Ethnobotanical survey of herbal remedies traditionally used in kohghiluyeh va boyer ahmad province of iran. *J Ethnopharmacol*. 2012;141(1):80-95.
- Jamzad Z. Flora of Iran. Tehran: Research Institute of Forests and Rangelands; 2012. p.333-8.
- Delazar A, Asnaashari S. Two iridoid structures from *Eremostachys macrophylla* montbr. & auch. Rhizomes. *J Rep Pharm Sci*. 2018;7(2):221-6.
- Boros CA, Stermitz FR. Iridoids. An updated review, part ii. *J Nat Prod*. 1991;54(5):1173-246. doi:10.1021/np50077a001
- Ersöz T, Schühly W, Popov S, Handjieva N, Sticher O, Çaliş I. Iridoid and phenylethanoid glycosides from *phlomis longifolia* var. *Longifolia*. *Nat Prod Lett*. 2001;15(5):345-51. doi:10.1080/10575630108041302
- Pan Z, Fan G, Yang Rp, Luo Wz, Zhou Xd, Zhang Y. Discriminating *lamiophlomis rotata* according to geographical origin by 1h-nmr spectroscopy and multivariate analysis. *Phytochem Anal*. 2015;26(4):247-52. doi:10.1002/pca.2557
- Modarresi M, Foadnia M, Rafiee Z, Jafari A, Zarzasangan K. Iridoid glycosides from *eremostachys azerbaijanica* rech. f. Root. *J Med Plants*. 2013;2(46):66-77.
- Çaliş İ, Güvenç A, Armağan M, Koyuncu M, Gotfredsen CH, Jensen SR. Iridoid glucosides from *eremostachys moluccelloides*bunge. *Helv Chim Acta*. 2007;90(8):1461-6. doi:10.1002/hlca.200790150

22. Calis I, Güvenc A, Armagan M, Koyuncu M, Gotfredsen CH, Jensen SR. Secondary metabolites from *eremostachys laciniata*. *Nat Prod Commun*. 2008;3(2):117-24. doi:10.1177/1934578x0800300202
23. Tundis R, Loizzo MR, Menichini F, Statti GA, Menichini F. Biological and pharmacological activities of iridoids: Recent developments. *Mini Rev Med Chem*. 2008;8(4):399-420. doi:10.2174/138955708783955926
24. Vidyalakshmi KS, Nagarajan S, Vasanthi HR, Rajamanickam V. Hepatoprotective and antioxidant activity of two iridoids from *mussaenda 'dona aurora'*. *Z Naturforsch C*. 2009;64(5-6):329-34. doi:10.1515/znc-2009-5-604
25. Xie W, Zhu J, Wan Z, Zhao Y. Anti-tumor effect of phlomiol and its mechanism study. *China Pharm*. 2010;39:1-7.
26. Fouladnia M, Modarresi M. Phenylethanoid glycosides from *Eremostachys azerbaijanica* rech. *F. Res Pharm Sci*. 2012;7(5):760-1.
27. Alipieva K, Korkina L, Orhan IE, Georgiev MI. Verbascoside—a review of its occurrence, (bio) synthesis and pharmacological significance. *Biotechnol Adv*. 2014;32(6):1065-76. doi:10.1016/j.biotechadv.2014.07.001
28. Carrillo-Ocampo D, Bazaldúa-Gómez S, Bonilla-Barbosa J, Aburto-Amar R, Rodríguez-López V. Anti-inflammatory activity of iridoids and verbascoside isolated from *castilleja tenuiflora*. *Molecules*. 2013;18(10):12109-18. doi:10.3390/molecules181012109
29. Kosmider B, Osiecka R. Flavonoid compounds: A review of anticancer properties and interactions with cis-diamminedichloroplatinum (ii). *Drug Dev Res*. 2004;63(4):200-11. doi:10.1002/ddr.10421
30. Özcan MM, Matthäus B. A review: Benefit and bioactive properties of olive (*olea europaea* L.) leaves. *Eur Food Res Technol*. 2017;243(1):89-99. doi:10.1007/s00217-016-2726-9
31. Palombo R, Savini I, Avigliano L, Madonna S, Cavani A, Albanesi C, et al. Luteolin-7-glucoside inhibits il-22/stat3 pathway, reducing proliferation, acanthosis, and inflammation in keratinocytes and in mouse psoriatic model. *Cell Death Dis*. 2016;7(8):e2344. doi:10.1038/cddis.2016.201
32. Fuchs J, Milbradt R. Skin anti-inflammatory activity of apigenin-7-glucoside in rats. *Arzneimittelforschung*. 1993;43(3):370-2.
33. Parveen Z, Deng Y, Saeed MK, Dai R, Ahamad W, Yu YH. Antiinflammatory and analgesic activities of thesium chinense turcz extracts and its major flavonoids, kaempferol and kaempferol-3-o-glucoside. *Yakugaku Zasshi*. 2007;127(8):1275-9. doi:10.1248/yakushi.127.1275
34. Delazar A, Modarresi M, Shoeb M, Nahar L, Reid RG, Kumarasamy Y, et al. Eremostachiin: A new furanolabdane diterpene glycoside from *eremostachys glabra*. *Nat Prod Res*. 2006;20(2):167-72. doi:10.1080/13518470500047082
35. Delazar A, Gibbons S, Kumarasamy Y, Nahar L, Shoeb M, Sarker SD. Antioxidant phenylethanoid glycosides from the rhizomes of *Eremostachys glabra* (lamiaceae). *Biochem Syst Ecol*. 2005;33(1):87-90. doi:10.1016/j.bse.2004.05.011
36. Asnaashari S, Heshmati Afshar F, Moghadam SB, Delazar A. Evaluation of in vitro antimalarial activity of different extracts of *Eremostachys azerbaijanica* rech. *F. Iran J Pharm Res*. 2016;15(3):523-9.
37. Asnaashari S, Delazar A, Asgharian P, Lotfipour F, Moghaddam SB, Heshmati Afshar F. In-vitro bioactivity and phytochemical screening of extracts from rhizomes of *Eremostachys azerbaijanica* rech. f. growing in Iran. *Iran J Pharm Res*. 2017;16(1):306-14.
38. Mughal UR, Fatima I, Malik A, Bakhsh Tareen R. Loasifolin, a new flavonoid from *Eremostachys loasifolia*. *J Asian Nat Prod Res*. 2010;12(4):328-30. doi:10.1080/10286021003627379
39. Mughal UR, Fareed G, Zubair A, Malik A, Versiani MA, Afza N, et al. Loasins a and b, new flavonoids from *Eremostachys loasifolia*. *Nat Prod Res*. 2013;27(20):1906-10. doi:10.1080/14786419.2013.784872
40. Ali B, Mehmood R, Mughal UR, Malik A, Safder M, Hussain R, et al. Eremosides a-c, new iridoid glucosides from *Eremostachys loasifolia*. *Helv Chim Acta*. 2012;95(4):586-93. doi:10.1002/hlca.201100316
41. Imran M, Mehmood R, Mughal UR, Ali B, Malik A. Vicarin, a new isoflavone from *Eremostachys vicaryi*. *J Asian Nat Prod Res*. 2012;14(3):293-6. doi:10.1080/10286020.2011.652951
42. Nan ZD, Zeng KW, Shi SP, Zhao MB, Jiang Y, Tu PF. Phenylethanoid glycosides with anti-inflammatory activities from the stems of *cistanche deserticola* cultured in tarim desert. *Fitoterapia*. 2013;89:167-74. doi:10.1016/j.fitote.2013.05.008
43. He ZD, Lau KM, Xu HX, Li PC, But PPH. Antioxidant activity of phenylethanoid glycosides from *brandisia hancei*. *J Ethnopharmacol*. 2000;71(3):483-6. doi:10.1016/S0378-8741(00)00189-6
44. Asnaashari S, Delazar A, Alipour SS, Nahar L, Williams AS, Pasdaran A, et al. Chemical composition, free-radical-scavenging and insecticidal activities of the aerial parts of *stachys byzantina*. *Arch Biol Sci*. 2010;62(3):653-62. doi:10.2298/ABS1003653A
45. Serafini M, Peluso I, Raguzzini A. Flavonoids as anti-inflammatory agents. *Proc Nutr Soc*. 2010;69(3):273-8. doi:10.1017/s002966511000162x
46. Pietta PG. Flavonoids as antioxidants. *J Nat Prod*. 2000;63(7):1035-42. doi:10.1021/np9904509
47. Dinda B, Debnath S, Harigaya Y. Naturally occurring secoiridoids and bioactivity of naturally occurring iridoids and secoiridoids. A review, part 2. *Chem Pharm Bull*. 2007;55(5):689-728. doi:10.1248/cpb.55.689