



Acta Scientiarum. Biological Sciences

ISSN: 1679-9283

eduem@uem.br

Universidade Estadual de Maringá

Brasil

Felisberto, Sirlene Aparecida; Rodrigues, Liliana
Periphytic algal community in artificial and natural substratum in a tributary of the Rosana reservoir
(Corvo Stream, Paraná State, Brazil)
Acta Scientiarum. Biological Sciences, vol. 32, núm. 4, 2010, pp. 373-385
Universidade Estadual de Maringá
.png, Brasil

Disponibile en: <http://www.redalyc.org/articulo.oa?id=187115378007>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica
Red de Revistas Científicas de América Latina, el Caribe, España y Portugal
Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto

Periphytic algal community in artificial and natural substratum in a tributary of the Rosana reservoir (Corvo Stream, Paraná State, Brazil)

Sirlene Aparecida Felisberto^{1*} and Liliana Rodrigues²

¹Universidade Federal de Goiás, Câmpus Samambaia, Cx. Postal 131, 74001-970, Goiânia Goiás, Brazil. ²Universidade Estadual de Maringá, Maringá, Paraná, Brazil. *Author for correspondence. Email: sirfe@hotmail.com

ABSTRACT. Periphytic algal community in artificial and natural substratum in a tributary of the Rosana reservoir (Corvo Stream, Paraná State, Brazil). This study evaluated the heterogeneity in periphytic algal community, under the influence of time colonization on artificial substratum. We also examined which abiotic variables most influenced the community in natural and artificial substratum. *Egeria najas* Planchon was used as natural substratum, and a plastic plant, as artificial. This experiment was carried out in a lateral arm from Rosana Reservoir, formed by Corvo Stream (Paranapanema river basin), in the period from November 21st to December 12nd, 2003, characterized as a warm and rainy period. Changes in species composition were assessed using the similarity indices. 495 taxa were registered in the phycoperiphytic community, distributed in 133 genera and 11 classes. Zygnemaphyceae, Bacillariophyceae, Chlorophyceae and Cyanophyceae presented higher species number, in both substrata. *Staurodesmus*, *Closterium*, *Staurastrum* and *Cosmarium* (Desmidiales); *Gomphonema* and *Eunotia* (Pennales); *Characium*, *Scenedesmus* and *Desmodesmus* (Chlorococcales); *Anabaena* and *Aphanocapsa* (Nostocales and Chroococcales, respectively) were the most species-rich genera. The epiphytic community reached the highest species richness in the 15th successional day. Regardless the substratum type, the number of species was probably related to the high concentrations of nutrients (phosphorus and nitrogen).

Key words: ecology, colonization, phycoperiphyton, Zygnemaphyceae, similarity.

RESUMO. Comunidade de algas perifíticas em substrato artificial e natural em um tributário do reservatório de Rosana (Ribeirão do Corvo, Estado do Paraná, Brasil).

Este trabalho objetivou analisar a composição da comunidade de algas perifíticas sob influência do tempo de colonização em substrato artificial. Ainda procurou responder quais variáveis abióticas mais influenciaram na comunidade em substrato artificial e natural. Utilizou-se *Egeria najas* Planchon como substrato natural e planta de plástico semelhante à *Egeria* como artificial. O experimento foi realizado no rio do Corvo, tributário do reservatório de Rosana (bacia do Paranapanema). As amostragens foram realizadas de 21 de novembro a 12 de dezembro de 2003, quando o clima é caracterizado como quente e chuvoso. As alterações na composição específica do ficoperifiton foram analisadas utilizando-se o índice de similaridade. A comunidade de algas perifíticas compreendeu 495 táxons, distribuídos em 133 gêneros e 11 classes. Zygnemaphyceae, Bacillariophyceae, Chlorophyceae e Cyanophyceae apresentaram maior número de espécies, em ambos os substratos. *Staurodesmus*, *Closterium*, *Staurastrum* e *Cosmarium* (Desmidiales); *Gomphonema* e *Eunotia* (Pennales); *Characium*, *Scenedesmus* e *Desmodesmus* (Chlorococcales); *Anabaena* e *Aphanocapsa* (Nostocales e Chroococcales, respectivamente) foram os gêneros mais representativos. A comunidade perifítica do substrato artificial atingiu maior número de espécies no 15^o dia sucessional. O número de espécies independentemente do tipo de substrato foi relacionado à maior disponibilidade de nutrientes (fósforo e nitrogênio).

Palavras-chave: ecologia, colonização, ficoperifiton, Zygnemaphyceae, similaridade.

Introduction

The species diversity plays an important role in the processes of production, consumption, respiration and decaying, mainly at littoral regions, since the aquatic vegetation is generally associated to different communities. The high availability of habitats provided by the diversity of aquatic vegetation favors the establishment and development of several organisms,

mostly the periphyton (WETZEL, 1981; WETZEL; LIKENS, 1991).

The periphyton is a sessile community that responds punctually to environmental conditions. This community presents short life cycle (3 - 10 days) which allows quick answers to environmental changes (HAMBROOK, 2002; RODRIGUES et al., 2003; WU et al., 2009). The heterogeneity in the

structure of phycoperiphytic community may be understood through the species composition in different successional stages. The studies about the distribution and mobility of algae from this community, when analyzed in short time scale, are more suitable due to population changes (WETZEL, 1983). This distribution extremely homogeneous of periphyton and its interaction with natural substrata, coupled to the need to standardize the community development time and the substratum nature, have motivated the use of artificial substrata. The review made by Cattaneo and Amireault (1992) indicated that 60% from published papers had used different types of artificial substrata.

In Brazil, among the published researches with artificial substratum reproducing a natural one are those using glass tubes (MOSCHINI-CARLOS et al., 2000), glass slides (RODRIGUES; BICUDO, 2001, 2004) and plastic tubes (FERNANDES; ESTEVES, 2003) to compare with petioles of emerging macrophytes, whereas other studies had used polyamide strings (OLIVEIRA et al., 2001) and glass slides (VERCELLINO; BICUDO, 2006) to analyze the structure and dynamic of periphytic algae community. Although Brazil presents one of the major hydroelectric parks of the world, the knowledge about periphyton in these environments is still scarce.

This study hypothesized that the composition of periphytic algae varies in function to the type of substratum (artificial and natural) and/or due to abiotic conditions of the system. The prediction is that in the course of time the phycoperiphytic community from an artificial substratum will be similar to that found in natural one, considering the species composition, and that both communities will be mainly influenced by nutrient concentrations. Therefore, we analyzed (a) the floristic composition from the community of periphytic algae in both substrata of one tributary of Rosana reservoir, Corvo Stream, during a warm and rainy period; (b) the algae similarity, between artificial and natural substratum; and (c) the principal abiotic factors influencing the species composition and richness of phycoperiphytic community, in both substrata.

Material and methods

Study area

The samplings location is one tributary that flows into the lacustrine region from Rosana

reservoir (Corvo Stream, 22°39'S; 052°46'W, Figure 1), near the dam. This river is situated in Paraná State, between the counties of Diamante do Norte and Terra Rica. The sampling station was 4.95 km far from the lacustrine region of the reservoir.




Figure 1. Map with the location of Corvo Stream, tributary of Rosana reservoir (Paranapanema river watershed).

Corvo Stream presents, in the sampling local, approximately 250 m of width and 5.8 m of depth. The banks practically did not present arborous vegetation, and the littoral region present several ecological types of aquatic macrophytes: emerged (*Typha* sp., *Sagittaria* sp. and *Eichhornia azurea* Kunth), floating (*Eichhornia crassipes* (Mart.) Solms, *Nymphaea* sp. and *Salvinia* sp.) and rooted-submersed (*Cabomba furcata* (Schult.) Schult., *Egeria densa* Planchon, *Egeria najas* Planchon, *Miriophyllum* sp. and *Utricularia foliosa* Lineau).

Samplings and analyses

Periphyton samplings were performed from November 21st to December 12nd, 2003, a period characterized as warm and rainy. Two types of substrata were chosen for the samplings: *Egeria najas* (natural substratum) and a plastic plant (artificial substratum), similar to natural substratum (Figure 2).

For the natural substratum, a fragment of *Egeria najas* was removed from the environment, and a median part was detached (about 7 to 8 cm of length), excluding the apex and base. Afterwards, this median part was transferred to previously moistened flasks, and kept in ice, and later the periphyton was removed for analysis. The samplings of both substrata were carried out in the lacustrine region of the environment. For

the artificial substratum, each one was washed in running water and after this; each branch was attached to a weight (pebbles wrapped in plastic) using a nylon line, and Styrofoam pieces as buoys. These sets were put in Corvo Stream, close to the left bank, about 60 cm of depth, in November 21st.



Figure 2. Types of substrata, *Egeria najas* Planchon (natural) and a plastic plant, sampled in Corvo Stream (tributary of Rosana reservoir), during the study period.

Samplings of artificial substratum occurred every three days, between November 24th (1st sampling and 3rd successional days) and December 12nd, 2003 (last sampling and 21st day), while for the natural substratum, the samplings began at the first day of installation of artificial substratum (11/21), and during alternate days (11/26; 01, 06, 09 and 12/12), totaling 13 samples (six for the natural substratum, and seven for the artificial one). The periphyton was washed from the substrata using brush and distilled water, and then transferred to transparent flasks. The samples were preserved with Transeau solution (BICUDO; MENEZES, 2006).

Abiotic data evaluated in the study were furnished by Limnology Laboratory, from Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura – Nupélia. Water temperature (YSI 55 portable oximeter), turbidity (LaMotte portable turbidimeter) and wind (anemometer) were measured during the samplings. The concentrations of total nitrogen, total phosphorus and total dissolved phosphorus were determined based on Valderrama (1981), nitrate, ammonium and orthophosphate were estimated according to Mackereth et al. (1978), Solorzano (1969) and Golterman et al. (1978), respectively.

The taxonomical study of periphytic algae was accomplished using approximately 15 temporary slides, by sample. For this procedure, we used optical microscope with micrometric ocular. Some genera of green filamentous algae, as *Bulbochaete*, *Oedogonium*, *Mougeotia* and *Spirogyra*, were distinguished only in vegetative groups, based on the cells diameter and length, due to the absence of reproductive structures in the analyzed individuals.

The algae identification was based on classical literature, as Croasdale and Flint (1986, 1988), Dillard (1990, 1991), Förster (1982), Komárek and Anagnostidis (1999, 2005), Krammer and Lange-Bertalot (1986, 1988, 1991), Patrick and Reimer (1966, 1975), Prescott (1982), Prescott et al. (1981, 1982) among others. Regarding the classification system of algae, we adopted Round (1965, 1971), following the recommendation made by Bicudo and Menezes (2006).

The species richness, expressed in number of taxa for both substrata, was obtained from qualitative samples complemented with quantitative samples.

The similarity of periphytic algae from artificial and natural substratum, between the sampled periods, was analyzed using the Jaccard similarity index (species presence/absence data), using NTSYS software, version 2.1 (ROHLF, 2000) and unweighted average (UPGMA).

The values of total richness of algae classes from both substrata were correlated with the abiotic variables, through Pearson correlation analysis, using Statistica software, version 7.1 (STATSOFT, 2005).

Results

Floristic composition

The community of periphytic algae comprised 495 taxa distributed in 133 genera and 11 classes, considering both substrata (natural and artificial) (Table 1). Among these taxa, 403 occurred in the natural substratum, and 401 in the artificial one. In the table 1, the complete list of species is presented, and, from this total, 86 taxa occurred exclusively in the natural substratum, and 98 in the artificial. The most representative classes in the natural and artificial substratum were Zygnemaphyceae (37.2 and 36.4%, respectively) Bacillariophyceae (21.6 and 24.7%, respectively), Chlorophyceae (17.4 and 17.2%, respectively) and Cyanophyceae (13.2 and 12%, respectively).

Amongst Zygnemaphyceae, *Stauroidesmus*, *Closterium*, *Staurastrum* and *Cosmarium* (66.5% from the total of Desmidiaceae) were the best represented genera, in both substrata. Bacillariophyceae, Order Pennales,

presented *Gomphonema* and *Eunotia* as the most representative genera regarding the number of taxa. Among Chlorophyceae, *Characium*, *Desmodesmus* and *Scenedesmus* were the best represented (Table 1).

Table 1. Taxa of periphytic algae surveyed in Corvo Stream (tributary of Rosana reservoir), in the study period.

	Bacillariophyceae
<i>Achnanthydium exiguum</i> (Grun.) Czarn.	<i>Gomphonema augur</i> Ehr.**
<i>A. minutissimum</i> (Kütz.) Czarn.	<i>Gomphonema augur</i> Ehr. var. <i>turris</i> (Ehr.) Lang.-Bert.
<i>Amphipleura lindheimeri</i> Grun.	<i>Gomphonema</i> cf. <i>auritum</i> Braun ex Kütz.
<i>Amphora copulata</i> (Kütz.) Schoem. & Arch.	<i>Gomphonema brasiliense</i> Grun.
<i>Amphora</i> cf. <i>ovalis</i> (Kütz.) Kütz.**	<i>Gomphonema gracile</i> Ehr.
<i>Amphora</i> sp.	<i>Gomphonema parvulum</i> Kütz.
<i>Aulacoseira granulata</i> (Ehr.) Sim	<i>Gomphonema subtile</i> Ehr.**
<i>Aulacoseira granulata</i> (Ehr.) Sim. var. <i>angustissima</i> (O. Müll.) Sim. **	<i>Gomphonema</i> cf. <i>truncatum</i> Ehr.*
<i>Aulacoseira</i> sp. *	<i>Gomphonema</i> sp.
<i>Brachysira neoxilis</i> Hor. Lang.-Bert. & Mos.	<i>Gomphonema</i> sp.1*
<i>Capartogramma crucicula</i> (Grun. & Cleve)**	<i>Hantzschia amphioxys</i> (Ehr.) Grun.
<i>Cyclotella stelligera</i> (Cleve & Grun.) V. Heur.	<i>Navicula aikenensis</i> Patr.
<i>Cymbella acuta</i> (A. Schm.) Cleve	<i>Navicula constans</i> Hust.
<i>Cymbella affinis</i> Kütz.	<i>Navicula cryptocephala</i> Kütz.
<i>Cymbella cuspidata</i> Kütz.	<i>Navicula cryptonella</i> Lang.-Bert.
<i>Cymbella</i> cf. <i>moreirae</i> (Rod.) Ludw.	<i>Navicula</i> cf. <i>drouetiana</i> Patr.**
<i>Cymbella</i> cf. <i>myrocephala</i> Grun.*	<i>Navicula</i> cf. <i>menisculus</i> Schum.**
<i>C. naviculiformes</i> Auer. ex Heib.	<i>Navicula</i> cf. <i>tenera</i> *
<i>Diatoma</i> sp.	<i>Navicula</i> sp.*
<i>Diploneis</i> cf. <i>subovalis</i> Cleve	<i>Nitzschia amphibia</i> Grun.
<i>Encyonema lunatum</i> (Smith) V. Heur.*	<i>Nitzschia frustulum</i> Grun.**
<i>Encyonema mesianum</i> (Chol.) Mann*	<i>Nitzschia</i> cf. <i>intermedia</i> Hantz. ex Cleve & Grun.
<i>Encyonema minutum</i> (Hil.) Mann*	<i>Nitzschia</i> cf. <i>linearis</i> Grun.
<i>Encyonema</i> cf. <i>perpusillum</i> Cleve & Mann*	<i>Nitzschia palea</i> (Kütz.) Smith
<i>Encyonema silesiacum</i> (Bleis.) Mann*	<i>Nitzschia</i> sp.*
<i>Encyonema</i> sp.	Penales unidentified 1
<i>Encyonema</i> sp.1	Penales unidentified 2*
<i>Eunotia bilunaris</i> Ehr.	<i>Pinnularia</i> cf. <i>braunii</i> Grun.*
<i>Eunotia camelus</i> Ehr.	<i>Pinnularia divergens</i> Smith
<i>Eunotia curvata</i> (Kütz.) Lagerst.	<i>Pinnularia gibba</i> Ehr.
<i>Eunotia didyma</i> Hust.	<i>Pinnularia interrupta</i> Smith
<i>Eunotia</i> cf. <i>faba</i> (Ehr.) Grun.*	<i>Pinnularia luculenta</i> A. Schm.
<i>Eunotia flexuosa</i> Bréb.	<i>Pinnularia maior</i> (Kütz.) Rab.
<i>Eunotia formica</i> Ehr.	<i>Pinnularia mesolepta</i> (Ehr.) Smith
<i>Eunotia indica</i> Grun.	<i>Pinnularia</i> cf. <i>microstauron</i> (Ehr.) Cleve*
<i>Eunotia lineolata</i> Hust.	<i>Pinnularia subcapitata</i> Greg. *
<i>Eunotia maior</i> (Smith) Rab.	<i>Pinnularia viridis</i> (Nitz.) Ehr.
<i>Eunotia</i> cf. <i>minor</i> (Kütz.) Grun.*	<i>Pinnularia</i> sp.
<i>Eunotia monodon</i> Ehr.*	<i>Placoneis disparilis</i> (Hust.) Lang.-Bert.
<i>Eunotia pectinalis</i> (Dillw.) Rab.	<i>Planothidium lanceolatum</i> (Bréb.) Round & Bukh.*
<i>Eunotia pectinalis</i> (Dillw.) Rab. var. <i>ventricosa</i> Grun.	<i>Pleurosira laevis</i> (Ehr.) Comp.
<i>Eunotia sudetica</i> O. Müll.	<i>Rhopalodia</i> cf. <i>brebissoni</i> Kram.*
<i>Eunotia zygodon</i> Ehr.*	<i>Rhopalodia gibberula</i> (Ehr.) O. Müll.**
<i>Fragilaria capucina</i> Desm.	<i>Sellaphora pupulla</i> Kütz.
<i>Fragilaria capucina</i> Desm. var. <i>gracilis</i> (Oestrup) Hust.	<i>Stauroneis phoenicenteron</i> (Nitz.) Ehr.
<i>Fragilaria delicatissima</i> (Smith) Lang.-Bert.	<i>Stenopterobia delicatissima</i> (Lew.) Bréb. ex V. Heur.
<i>Fragilaria javanica</i> Hust.*	<i>Stenopterobia intermedia</i> (Lew.) V. Heur.
<i>Fragilaria pinnata</i> Ehr.*	<i>Surirella bisseriata</i> Bréb.
<i>Frustulia rhomboides</i> (Ehr.) De Toni var. <i>rhomboides</i>	<i>Surirella linearis</i> Smith
<i>Frustulia rhomboides</i> (Ehr.) De Toni var. <i>crassinervia</i> (Bréb. ex Smith) Ross	<i>Surirella</i> cf. <i>robusta</i> Hust.**
<i>Frustulia rhomboides</i> (Ehr.) De Toni var. <i>saxonica</i> (Rab.) De Toni	<i>Surirella sublinearis</i> Hust.
<i>Frustulia</i> cf. <i>vitrea</i> Oestrup*	<i>Surirella</i> sp.
<i>Frustulia vulgaris</i> (Thwait.) De Toni*	<i>Synedra acus</i> Kütz. var. <i>angustissima</i> Grun.*
<i>Frustulia</i> sp.*	<i>Synedra gouldarii</i> Bréb.*
<i>Gomphonema affine</i> Kütz.	<i>Ulnaria ulna</i> (Nitz.) Ehr.
<i>Gomphonema</i> cf. <i>apicatum</i> Ehr.*	<i>Talassiosira</i> sp.**

Continuac...

..continuation

Ankistrodesmus falcatus (Corda) Ralfs
Ankistrodesmus fusiformis Corda
Ankistrodesmus spiralis (Turn.) Lemm.
Aphanochaete repens Braun
Bicuspidela cf. *ineus* Pasch.
Botryococcus sp.
Chaetosphaeridium globosum (Nord.) Kleb.
Chaetosphaeridium sp.*
Characiellopsis skujae (Fott) Kom.
Characium cf. *acuminatum* Braun**
Characium ambiguum Herm.
Characium cf. *cucurbitinum* (Biss.) Teil.*
Characium ensiforme Herm.
Characium guttula Play.
Characium ornithocephalum (Braun) var. *ornithocephalum*
Characium ornithocephalum (Braun) var. *hapochytriforme* Prin.
Characium ornithocephalum (Braun) var. *pringsheimii* (Braun) Kom.
Characium sp.**
Chlamydomonas cf. *globosa* Snow
Chlamydomonas sp.
Closteriopsis acicularis (Smith) Belc. & Swale
Coelastrum indicum Turner
Coleochaete cf. *irregularis* Prings.**
Coleochaete orbicularis Prings.**
Coleochaete sp.
Crucigenia sp.
Desmodesmus cf. *abundans* (Kirch.) Hegew.
Desmodesmus armatus (Chod.) Hegew.
Desmodesmus armatus (Chod.) var. *bicaudatus* (Gugl.) Hegew.
Desmodesmus cf. *armatus* (Chod.) var. *spinulosus* (Frit. & Rich) Hegew.**
Desmodesmus communis (Hegew.) Hegew.
Desmodesmus denticulatus (Lagerh.) Friedl & Hegew.
Desmodesmus denticulatus (Lagerh.) Friedl & Hegew. var. *linearis* (Hansg.) Hegew.
Desmodesmus dispar Bréb.
Desmodesmus maximus (W. & G.S. West) Hegew.
Desmodesmus opoliensis (Rich.) Hegew. var. *carinatus* (Lemm.) Hegew.*

Desmodesmus perforatus (Lemm.) Hegew.
Desmodesmus quadricauda (Turp.) Hegew.
Desmodesmus serratus (Corda) Friedl & Hegew.
Desmodesmus spinosus (Chod.) Hegew.
Dictyosphaerium pulchellum Wood*
Dictyosphaerium sp.*

Anabaena cf. *affinis* Lemm.**
Anabaena cf. *circinalis* Rab.**
Anabaena spiroides Kleb.
Anabaena sp.
Anabaena sp.1**
Aphanocapsa elachista W. & G.S. West
Aphanocapsa delicatissima W. & G.S. West**
Aphanocapsa grevillei Raben.
Aphanocapsa cf. *pulchra* (Kütz.) Raben.*
Aphanocapsa sp.
Calothrix sp.*
Chamaesiphon incrustans Grun.
Chamaesiphon sp.
Chroococcus minor (Kütz.) Näg.
Chroococcus minimus (Keis.) Lemm.
Chroococcus sp.

Chlorophyceae

Dimorphococcus lunatus Braun*
Drepanochloris uherkovichii Marvan
Gloeoecystis cf. *gigas* (Kütz.) Lagerh.**
Golenkinia radiata (Chod.) Wille**
Kirchneriella lunaris (Kirch.)*
Kirchneriella obesa (West) West & G.S. West*
Monoraphidium arcuatum (Korsh.) Hind.
Monoraphidium caribeum Hind.
Monoraphidium circinale (Nyg.) Nyg.
Monoraphidium contortum (Thur.) Kom.-Legn.
Monoraphidium griffithii (Berk.) Kom.-Legn.**
Monoraphidium sp.*
Oocystis lacustris Chod.*
Pediastrum duplex Meyen
Pediastrum duplex Meyen var. *subgranulatum* Racib.
Pediastrum tetras (Ehr.) Ralfs
Planktosphaeria gelatinosa Smith
Radiococcus sp.*
Scenedesmus antennatus Bréb.
Scenedesmus acuminatus (Lagerh.) Chod.
Scenedesmus cf. *acumae* Comas**
Scenedesmus acutus (Meyen) Chod.
Scenedesmus acutus (Meyen) Chod. var. *alternans* Hortob.
Scenedesmus acutus (Meyen) Chod. var. *globosus* Hortob.
Scenedesmus bijugus (Turp.) Kütz.
Scenedesmus javanensis Chod.*
Scenedesmus linearis Kom.*
Scenedesmus cf. *regularis* Svir.*
Scenedesmus wisconsinensis (Smith) Chod.**
Scenedesmus sp.
Selenastrum gracile (Rein.)
Selenastrum rinoi Kom. et Com.*
Sphaerocystis schroeteri Chod.**
Stigeodolium sp.
Tetraedron caudatum (Corda) Hans.**
Tetraedron minimum (Braun) Hans.*
Tetraedron regulare Kütz.**
Tetrastrum sp.**
Westella sp.
 Chlorococcales unidentified
 Chlorococcales unidentified 1
 Chlorococcales unidentified 2

Cyanophyceae

Lyngbya maior Men.
Merismopedia glauca (Ehr.) Kütz.**
Merismopedia tenuissima Lemm.
Merismopedia sp.*
Microcrocis cf. *irregularis* (Lagerh.) Geitler
Microcrocis cf. *pulchella* (Buell) Geitler**
Microcystis aeruginosa (Kütz.) Lemm.
Microcystis firma (Bréb. et Len.) Schm.
Oscillatoria cf. *curviceps* Agardh.**
Oscillatoria principes Vaucher
Oscillatoria sancta (Kütz.) Gom.
Planktolynghya limnetica (Lemm.) Kom.-Legn.
Planktolynghya sp.
Phormidium cf. *chlorinum* (Kütz.) Umez. & Wat.
Phormidium irriguum (Kütz. ex Gom.) Anag. & Kom.
Planktothrix agardhii Anag. & Kom.*

Continue...

...continuation

- Coelosphaerium aeruginum* Lemm.*
Geitleribactron periphyticum Kom.**
Geitlerinema jasorvense (Vouk) Anag.**
Geitlerinema splendidum (Grev.) Anag.
Gloeocapsa cf. *rupestris* Kütz.**
Gloeocapsa sp.
Gloeocapsopsis sp.**
Jaaginema quadripunctulatum (Brühl. & Bisw.) Anag. & Kom.
Komvophoron crassum (Voz.) Anag. & Kom.
Komvophoron minutum (Skuja) Anag. & Kom.*
Komvophoron schmidlei (Jaag) Anag. & Kom.
Leptolyngbya fragilis (Gom.) Anag. & Kom.**
Leptolyngbya lagerheimii (Gom.) Anag. & Kom.
Leptolyngbya perelegans (Lemm.) Anag. & Kom.
Lynngbya comperei Senna
- Bicoeca* cf. *synoica* Skuja
Dinobryon divergens Imhof
Dinobryon sertularia Ehr.
Mallomonas sp.
Salpingoeca marsonii Lemm.*
- Cryptomonas ovata* Ehr.**
- Peridinium* cf. *cinctum* O. Müll.*
Peridinium cf. *volzii* Lemm.
- Colacium* sp.
Euglena cf. *acus* Ehr.**
Euglena cf. *gracilis* Klebs
Euglena cf. *spirogyra* Ehr.*
Euglena viridis Ehr.**
Euglena sp.
Lepocinclis sp.
Phacus acuminatus Stokes
Phacus curvicauda Swir.*
Phacus cf. *polytrophos* Poch.*
- Bulbochaete* sp.*
Bulbochaete sp.1
Bulbochaete sp.2
Bulbochaete sp.3 *
Bulbochaete sp.4
- Ulothrix* sp.
Uronema elongatum Hodg.
- Characiopsis acuta* (Braun) Borzi**
Characiopsis aquilonaris Skuja
Characiopsis elegans Ettl
Characiopsis minuta (Braun) Lemm.**
- Actinotaenium diploporum* (Lund.) Teil.**
Actinotaenium globosum (Bulnh.) Teil.
Actinotaenium lagenarioides (Roy) Teil.*
Closterium acutum Ehr.*
C. acutum Ehr. var. *variabile* (Lemm.) W. Krieg.
Closterium closterioides (Ralfs) Louis & Pect.*
Closterium closterioides (Ralfs) Louis & Pect. var. *intermedium* (Roy & Biss.) Ruz.*
Closterium cornu Ehr. ex Ralfs
Closterium cynthia de Not.
Closterium diana Ehr. ex Ralfs var. *minus* Hieron.
- Pseudanabaena catenata* Laut.
Pseudanabaena lonchoides Anag.
Pseudanabaena moniliformes Kom. & Kling
Pseudanabaena mucicula (Hub.-Pest. & Naum.) Schw.
Radiocystis fernandoi Kom. & Kom.-Legn.
Rhabdoderma irregulare (Naum.) Geitler**
Rhabdoderma lineare Schm. & Laut.
Rhabdoderma vermiculare Fott
Rhabdoderma sp.**
Snowella sp.
Spirulina laxa Smith
Synechocystis sp.*
Woronichinia elorantae Kom. & Kom.-Legn.*
 Chroococcales unidentified*
- Chrysophyceae**
Salpingoeca serpettei Bourr.**
Salpingoeca urceolata Kent
Salpingoeca sp.
Synura sp.*
- Cryptophyceae**
Cryptomonas sp.
- Dinophyceae**
Peridinium sp.*
- Euglenophyceae**
Phacus cf. *pusillus* Lemm.**
Phacus cf. *raciborskii* Drez.**
Phacus sp.**
Strombomonas sp.**
Trachelomonas cf. *bernardi* Wolos.**
Trachelomonas crista Balech
Trachelomonas hispida (Perty) Stein.*
Trachelomonas intermedia Dang.
Trachelomonas klebsii Ehr.**
Trachelomonas sp.
- Oedogoniophyceae**
Oedogonium sp.
Oedogonium sp.1
Oedogonium sp.2
Oedogonium sp.3
Oedogonium sp.4
- Ulothricophyceae**
Uronema sp.**
- Xanthophyceae**
Characiopsis sphagnicola Pascher**
Isthmochloron lobulatum (Näg.) Skuja**
Stipitococcus vasiformis Tiff.
- Zygnemaphyceae**
Haplotaenium minutum (Ralfs) Bando**
H. minutum (Ralfs) Bando var. *attenuatum* (W. West) Bando**
Hyalotheca dissiliens (Smith) Bréb. ex Ralfs
Micrasterias abrupta West & G.S. West*
Micrasterias furcata Ralfs
Micrasterias lateiceps Nordst. var. *acuminata* W. Krieg.**
Micrasterias lateiceps Nordst. var. *lateiceps*
Micrasterias mahabuleshwariensis Hobs.
Micrasterias radiosa Ralfs var. *radiosa*
Micrasterias radiosa Ralfs var. *elegantior* (G.S. West) Croasd.

Continue...

...continuation

<i>Closterium ehrenbergii</i> Menegh. ex Ralfs**	<i>Micrasterias rotata</i> (Grev.) Ralfs ex Ralfs **
<i>Closterium exiguum</i> West & G.S. West*	<i>Micrasterias truncata</i> (Corda) Bréb. ex Ralfs var. <i>pusilla</i> G.S. West
<i>Closterium gracile</i> Bréb.	<i>Mougeotia</i> sp.
<i>Closterium incurvum</i> Bréb.	<i>Mougeotia</i> sp.1
<i>Closterium jenneri</i> Ralfs	<i>Mougeotia</i> sp.2
<i>Closterium laterale</i> Nordst.**	<i>Mougeotia</i> sp.3
<i>Closterium leibleinii</i> Kütz. ex Ralfs	<i>Mougeotia</i> sp.4
<i>Closterium moniliferum</i> (Bory) Ehr. ex Ralfs	<i>Mougeotia</i> sp.5**
<i>Closterium navicula</i> (Bréb.) Lütik.	<i>Netrium digitus</i> (Ehr.) Itzigs. & Rothe
<i>Closterium nematodes</i> Josh. var. <i>proboscideum</i> Turn.**	<i>Netrium digitus</i> (Ehr.) Itzigs. & Rothe var. <i>parvum</i> (Borge) Krieg.**
<i>Closterium pusillum</i> Hantzsch**	<i>Netrium oblongum</i> (De Bary) Lütik.*
<i>Closterium setaceum</i> Ehr. ex Ralfs	<i>Octacanthium mucronulatus</i> (Nordst.) Comp.
<i>Closterium tortum</i> B.M. Griffiths **	<i>Onychonema laeve</i> Nordst.
<i>Closterium toxon</i> W. West	<i>Penium exiguum</i> West *
<i>Closterium tumidum</i> Johns.	<i>Penium margaritaceum</i> (Ehr.) ex Bréb.
<i>Closterium tumidum</i> Johns. var. <i>nylandicum</i> Grönbl.	<i>Pleurotaenium ehrenbergii</i> (Bréb.) de Bary
<i>Closterium venus</i> Kütz. ex Ralfs*	<i>Pleurotaenium nodosum</i> (Bail.) Lund.**
<i>Closterium</i> sp.	<i>Sphaerozosma aubertianum</i> W. West*
<i>Cosmarium abbreviatum</i> Racib. var. <i>minus</i> (West & G. S. West) W. Krieg. & Gerl.	<i>Spirogyra</i> sp.
<i>Cosmarium</i> cf. <i>abruptum</i> Lund.*	<i>Spirogyra</i> sp.1
<i>Cosmarium anisochondrum</i> Nordst. var. <i>tetrachondrum</i> Scott & Grönbl.*	<i>Spirogyra</i> sp.2*
<i>Cosmarium bayleyi</i> Wolle*	<i>Spirogyra</i> sp.3**
<i>Cosmarium bitriangulum</i> Grönbl. var. <i>groenbladii</i> Grönbl.	<i>Spirogyra</i> sp.4**
<i>Cosmarium blyty</i> Wille	<i>Spondylosium moniliforme</i> Lund.
<i>Cosmarium brasiliense</i> (Wille) Nordst.	<i>Spondylosium panduriforme</i> (Heimerl) Teil. var. <i>limneticum</i> (West & G. S. West) Först.
<i>Cosmarium</i> cf. <i>devei</i> (Lund.) Lütik.**	<i>Spondylosium planum</i> (Wolle) West & G.S. West
<i>Cosmarium comisurale</i> (Bréb.) Ralfs var. <i>crassum</i> Nordst.	<i>Spondylosium pulchellum</i> Arch.**
<i>Cosmarium contractum</i> Kirch.	<i>Spondylosium pulchrum</i> Bail.
<i>Cosmarium excavatum</i> Nordst.	<i>Spondylosium pygmaeum</i> (Cooke) W. & G.S. West*
<i>Cosmarium galeritum</i> Nordst. var. <i>borgei</i> Krieg. & Gerl.	<i>Staurastrum ambiguum</i> Turner
<i>Cosmarium granatum</i> Bréb. ex Ralfs	<i>Staurastrum anatinum</i> Cooke & Wills
<i>Cosmarium impressulum</i> Eلف.	<i>Staurastrum bineanum</i> Rabenh.**
<i>Cosmarium laeve</i> Rabenh. var. <i>laeve</i>	<i>Staurastrum boreale</i> West & G.S. West
<i>Cosmarium laeve</i> Rabenh. var. <i>uestii</i> Krieg. & Gerl.*	<i>Staurastrum brasiliense</i> Nordst.**
<i>Cosmarium lagoense</i> (Nordst.) Nordst. var. <i>amoebum</i> Först. & Eck.**	<i>Staurastrum brebissonii</i> Arch. var. <i>brasiliense</i> Grönbl.
<i>Cosmarium mamiliferum</i> Nordst.**	<i>Staurastrum claviferum</i> West & G.S. West
<i>Cosmarium margaritatum</i> (Lund.) Roy & Biss. var. <i>margaritatum</i> f. <i>minor</i> (Boldt) West & G.S. West	<i>Staurastrum cyclacanthum</i> West & G.S. West
<i>Cosmarium minimum</i> var. <i>subrotundatum</i> W. & G.S. West	<i>Staurastrum dilatatum</i> (Ehr.) Ralfs
<i>Cosmarium moerliani</i> Lütik. var. <i>brasiliense</i> Borge	<i>Staurastrum hagnmannii</i> Grönbl.
<i>Cosmarium naegelianum</i> Bréb.	<i>Staurastrum hantzschii</i> Reinsch
<i>Cosmarium norimbergense</i> Reinsch var. <i>depressum</i> (West & G. S. West) W. Krieg. & Gerl.	<i>Staurastrum leptacanthum</i> Nordst. var. <i>borgei</i> Först.*
<i>Cosmarium panamense</i> Presc.*	<i>Staurastrum leptocladum</i> Nordst. var. <i>leptocladum</i>
<i>Cosmarium phaseolus</i> var. <i>phaseolus</i> Bréb. ex Ralfs f. <i>minus</i> Boldt.	<i>Staurastrum leptocladum</i> Nordst. var. <i>cornutum</i> Wille**
<i>Cosmarium portianum</i> Arch.	<i>Staurastrum margaritaceum</i> (Ehr.) Ralfs
<i>Cosmarium protractum</i> (Näg.) de Bary	<i>Staurastrum muticum</i> (Bréb.) Ralfs
<i>Cosmarium pseudobroomei</i> Wolle	<i>Staurastrum nudibrachiatum</i> Borge
<i>Cosmarium pseudocannatum</i> Nordst.	<i>Staurastrum orbiculare</i> (Ehr.) Ralfs var. <i>orbiculare</i>
<i>Cosmarium pseudoexiguum</i> Racib.	<i>Staurastrum orbiculare</i> (Ehr.) Ralfs var. <i>depressum</i> Roy & Biss.
<i>Cosmarium</i> cf. <i>pseudopyramidatum</i> Lund.**	<i>Staurastrum</i> cf. <i>polymorphum</i> Bréb.*
<i>Cosmarium punctulatum</i> Bréb.	<i>Staurastrum</i> cf. <i>pseudotetracerum</i> (Nordst.) West & G. S. West
<i>Cosmarium</i> cf. <i>pyramidatum</i> Bréb. (Ralfs)*	<i>Staurastrum quadrangulare</i> Bréb. ex Ralfs var. <i>quadrangulare</i>
<i>Cosmarium quadrum</i> Lund. var. <i>minus</i> Nordst.	<i>Staurastrum quadrangulare</i> Bréb. ex Ralfs var. <i>contectum</i> (Turn.)*
<i>Cosmarium quadrum</i> Lund. var. <i>sublatum</i> (Nordst.) West & G.S. West	<i>Staurastrum quadricornutum</i> Roy & Biss.
<i>Cosmarium quinarium</i> Lund.*	<i>Staurastrum rotula</i> Nordst.**
<i>Cosmarium regnellii</i> Wille**	<i>Staurastrum seboldi</i> Reinsch var. <i>ornatum</i> Nordst.
<i>Cosmarium regnesii</i> Reinsch	<i>Staurastrum minesotense</i> Wolle
<i>Cosmarium regnesii</i> Reinsch var. <i>montanum</i> Schm.	<i>Staurastrum setigerum</i> Cleve var. <i>pectinatum</i> West & G.S. West
<i>Cosmarium reniforme</i> (Ralfs) Arch. var. <i>reniforme</i>	<i>Staurastrum subavicula</i> West & G.S. West
<i>Cosmarium reniforme</i> (Ralfs) Arch. var. <i>compressum</i> Nordst.	<i>Staurastrum tetracerum</i> (Kütz.) Ralfs

Continue...

...continuation

<i>Cosmarium sublobatum</i> (Bréb.) Arch. var. <i>brasiliense</i> Borge	<i>Staurastrum trifidum</i> Nordst. var. <i>inflexum</i> West & G.S. West
<i>Cosmarium subspeciosum</i> Nordst. var. <i>subspectiosum</i>	<i>Staurastrum</i> sp.*
<i>Cosmarium subspeciosum</i> Nordst. var. <i>validius</i> Nordst. *	<i>Staurodesmus brevispina</i> (Bréb.) Croas.
<i>Cosmarium subtumidum</i> Nordst.	<i>Staurodesmus clepsydra</i> (Nordst.) Teil.
<i>Cosmarium trilobulatum</i> Reinsch	<i>Staurodesmus corniculatus</i> (Lund.) Teil. var. <i>spinigerum</i> W. West
<i>Cosmarium vexatum</i> W. West	<i>Staurodesmus cuspidatus</i> (Bréb. ex Ralfs) Teil.
<i>Cosmarium</i> sp.*	<i>Staurodesmus dejectus</i> (Bréb.) Teil. var. <i>dejectus</i>
<i>Cosmarium</i> sp.1	<i>Staurodesmus dejectus</i> (Bréb.) Teil. var. <i>apiculatus</i> (Bréb.) Teil.
<i>Cylindrocystis brebissonii</i> (Menegh. ex Ralfs) de Bary	<i>Staurodesmus dejectus</i> (Bréb.) Teil. var. <i>brevispinus</i> (Nygaa.) Coescl *
<i>Desmidium aptogonium</i> Bréb. *	<i>Staurodesmus dickiei</i> (Ralfs) S. Lill.
<i>Desmidium baileyi</i> (Ralfs) Nordst.**	<i>Staurodesmus glaber</i> (Ehr. ex Ralfs) S. Lill.*
<i>Desmidium grevillii</i> (Kütz.) de Bary**	<i>Staurodesmus grandis</i> Teil. var. <i>parvus</i> W. & G.S. West**
<i>Desmidium pseudostreptonema</i> West & G.S. West**	<i>Staurodesmus lobatus</i> (Börger.) Bourr.**
<i>Desmidium swartzii</i> Agardh**	<i>Staurodesmus mamillatus</i> (Nordst.) Teil.
<i>Euastrum abruptum</i> Nordst.	<i>Staurodesmus patens</i> (Nordst.) Croas.
<i>Euastrum denticulatum</i> (Kirchn.) Gay	<i>Staurodesmus</i> cf. <i>spencerianus</i> (Mask.) Teil.*
<i>Euastrum evolutum</i> (Nordst.) W. & G.S. West**	<i>Staurodesmus</i> sp.
<i>Euastrum monocylum</i> (Nordst.) Rabic. var. <i>borgei</i> Grönb. l.	<i>Teilingia granulata</i> (Roy et Biss.) Bourr.
<i>Groenbladia undulata</i> Nordst.*	<i>Teilingia quadrispinata</i> (Scott et Grönb. l.) Bourr.*
<i>Gonatozygon aculeatum</i> Hast.	<i>Zygnema</i> sp.
<i>Gonatozygon monotaenium</i> de Bary	<i>Zygnema</i> sp.1
<i>Gonatozygon pilosum</i> Wolle*	Desmidiaceae unidentified

*taxa exclusive to artificial substratum; **taxa exclusive to natural substratum.

Therefore, independently of substratum type, we verified the predominance of Zygnemaphyceae (desmids), followed by Bacillariophyceae (diatoms), Chlorophyceae and Cyanophyceae (Table 1, Figure 4).

The number of species, in natural substratum, ranged from 156, in December 12nd (final of the experiment, 6th sampling) to 261 taxa, in November 26th (beginning of the experiment, 2nd sampling), thus, we observed a sudden decrease in the number of species during the final phase (Figure 3). In relation to artificial substratum, this number varied between 169, in December 9th (18th day), and 259 taxa, in December 6th (15th day).




Figure 3. Number of periphytic species in natural (11/21 = 1st, 11/26 = 2nd, 12/01 = 3rd, 12/06 = 4th, 12/09 = 5th, 12/12 = 6th sampling) and artificial substratum (11/24 = 1st, 11/27 = 2nd, 11/30 = 3rd, 12/03 = 4th, 12/06 = 5th, 12/09 = 6th, 12/12 = 7th sampling), sampled in Corvo Stream (tributary of Rosana reservoir).

In the natural substratum, there was a change in the number of desmids taxa, in December 6th (4th sampling), when we registered an expressive contribution of diatoms. The same was observed for the artificial substratum, however, in the last two samplings (December 9th and 12th, 18th and 21st successional days, Figure 4).




Figure 4. Relative contribution of taxonomic classes in Corvo Stream (tributary of Rosana reservoir), during the study period. (Other classes: Chrysophyceae, Cryptophyceae, Dinophyceae, Euglenophyceae, Oedogoniophyceae, Ulothricophyceae and Xanthophyceae).

Taxonomic similarity in periphytic community

In the diagram from the grouping analysis performed with periphytic algae from both substrata, there was a separation of groups, for the distinct substrata. Two groups were distinguished for artificial and natural substratum (Figure 5). For the artificial substratum, the assemblages of species referring to 3rd and 6th successional days (1st and 2nd samplings) were distinguished amongst themselves, and from the other days; a second group was formed, between the 9th and 12nd days (3rd and 4th samplings). The 15th successional day (5th sampling) was distinct from the others, however presented higher similarity with the final phase of the experiment, between the 18th and 21st days (Figure 5).

For the natural substratum, the results pointed higher similarity among the samples from the three

last samplings (December 6th, 9th and 12th), otherwise, the sample of the beginning of the experiment (November, 21st) was grouped with intermediary experimental phase (November 26th, and December 1st; Figure 5).




Figure 5. Dendrogram, resulting from Jaccard similarity index (UPGMA), among species of periphytic algae in Corvo Stream, from 13 sampling units. Mantel Test with $r = 0.78$. Substratum: Natural = N; Artificial = S.

The values referring to physical and chemical parameters of the water, electric conductivity, dissolved oxygen, water temperature and pH presented more constant values, while the other variables as nutrient concentrations and turbidity presented higher variation (Table 2).

Relationship between abiotic variables and phycoperiphytic community

In Paranapanema river watershed, as well as in Corvo Stream (lateral arm from Rosana reservoir), here is a precipitation regime, with concentration of rainfall between November to February, period characterized as warm and rainy. In the study

period (November-December), there was higher volume of precipitation at the end of November (11/28, 1st study week) and beginning of December (12/01, Figure 6), two days before the 4th sampling, 12nd successional day at artificial substratum.




Figure 6. Precipitation values in Corvo Stream (tributary of Rosana reservoir), in the period of 21/11 to 12/12/03.

This high precipitation along with the strong winds during the study period was responsible for the high quantity of material carried into the reservoir, contributing to increase the turbidity and nutrients concentration, which influenced the species composition of periphytic algae. In this way, the number of taxa recorded in the natural substratum was positively correlated to NT ($r = 0.95$), PT ($r = 0.88$), PO_4^{-3} ($r = 0.93$), PDT ($r = 0.84$) during the 1st week. This community attribute was also positively correlated to PO_4^{-3} ($r = 0.92$) during the 2nd week, and also with the turbidity ($r = 0.93$) in the 3rd week.

Considering the artificial substratum, during the 1st week, the species richness of phycoperiphytic species was positively correlated to PT ($r = 0.93$), NH_4^+ ($r = 0.97$), turbidity ($r = 0.91$), and wind ($r = 0.99$).

Table 2. Values of abiotic variables in Corvo Stream (Rosana reservoir), during the period of 11/21 to 12/12/03. (DO = dissolved oxygen, PT = total phosphorus, PDT = total dissolved phosphorus, PO_4^{-3} = orthophosphate, NT = total nitrogen, NO_3^- = nitrate, NH_4^+ = ammonium).

Weeks	1 st week			2 nd week			3 rd week			
Variable	11-21	11-24	11-26	11-27	11-30	12-01	12-03	12-06	12-09	12-12
Wind (m s ⁻¹)	2.8	3.1	1.5	1.0	5.6	1.0	3.4	3.0	6.7	2.3
Conductivity ($\mu S\ cm^{-1}$)	36.1	40.0	44.8	43.9	37.0	37.6	34.8	40.5	42.6	46.2
Water temperature (°C)	26.0	28.2	27.5	26.8	26.7	26.5	27.0	26.9	26.7	28.6
DO (mg L ⁻¹)	7.0	7.6	6.4	6.6	6.4	6.6	6.0	6.6	5.8	7.8
pH	6.9	7.0	7.0	6.7	6.7	6.5	6.3	6.6	6.0	6.2
Turbidity (NTU)	2.9	2.8	2.5	3.0	4.2	6.6	11.9	4.1	27.0	18.7
PT ($\mu g\ L^{-1}$)	5.7	9.6	13.1	11.1	18.0	17.8	22.4	22.7	21.6	26.3
PDT ($\mu g\ L^{-1}$)	3.1	3.2	5.2	4.4	9.0	8.8	9.2	9.5	8.2	16.6
PO_4^{-3} ($\mu g\ L^{-1}$)	2.3	3.3	4.3	4.2	7.0	6.9	3.7	4.7	4.5	3.8
NT ($\mu g\ L^{-1}$)	322.1	409.0	392.6	346.8	300.0	299.8	236.5	310.0	300.6	339.5
NO_3^- ($\mu g\ L^{-1}$)	185.3	215.1	206.3	176.6	120.0	119.0	114.4	136.0	134.0	97.6
NH_4^+ ($\mu g\ L^{-1}$)	18.0	20.9	31.7	65.1	66.0	29.2	35.8	75.0	73.9	44.0

During the 2nd week, positive correlations were observed between the species richness and NO₃⁻ ($r = 0.95$), and wind ($r = 0.88$), whereas negative correlations were verified with PDT ($r = -0.97$), and turbidity ($r = -0.93$). In the 3rd week, a negative correlation was registered with the turbidity ($r = -0.98$).

Discussion

The increase in species richness observed during the first days of colonization in the artificial substratum, was possibly due to the propagules present in the environment, since the habitat was not yet colonized, the migration processes favor the colonization and succession of species, allowing the composition of not only colonizing and opportunistic species of diatoms, as well as Chlorococcales, Desmidiaceae and filamentous algae, as *Bulbochaete* and *Oedogonium*. The immigration is an important process in the composition, colonization and increase of periphytic algae in artificial substratum (HILLEBRAND; SOMMER, 2000; PETERSON, 1996; STEVENSON; PETERSON, 1989).

The decrease in the number of species in the natural substratum over the experiment occurred during the days with higher pluviometric intensity. On the other hand, in artificial substratum, there was a different result, possibly, because this substratum is under succession process, since the increase in the number of species was gradate until November 30th (9th successional day and 3rd sampling), with a decrease in December 3rd (12nd day and 4th sampling). This reduction during the 4th sampling for both substrata may be related to the high precipitation that occurred in November 28th and mainly during the night of December 1st, i.e., two days before the sampling. From this period, the environmental conditions stabilized and hence, the number of species reached the maximum values during the 5th sampling (December 6th, 15th successional day), decreasing drastically again from the 6th sampling (December 9th, 18th day).

The oscillation in the species heterogeneity may be explained by the predominance of resilient species in phycoperiphytic community. These species can recover from disturbing events, as fluctuations in water level and large movement of the water column, caused by rainfall and constant winds. Based on studies about the succession of periphytic algae, the assemblages would be more resilient in more heterogeneous environments (more diversified environmental conditions in face of disturbing events, e.g., rainfall, wind), than in

homogeneous ones, because the species diversity in one habitat, would increase the efficient use of available resources (STEVENSON, 1997). The high resistance of many diatoms to wave's effect, even flooding, shows competitive advantages (BIGGS; THOMSEN, 1995; PETERSON, 1996; SABATER et al., 1998; STEVENSON, 1996b), and allows greater representativeness in rivers with high frequency of flooding. *A. minutissimum*, e.g., (species present in all samplings from both substrata), is widely recorded as intermediary colonizer, and highly resistant to disturbances (BIGGS; THOMSEN, 1995; PETERSON, 1996).

The differentiation of phycoperiphytic community in artificial substratum, was evidenced by the grouping analysis, in three phases: the initial, mainly formed when the community receives a gradate increase of species; the intermediate, when there is still an increase in the number of species, reaching a maximum; and the final phase, when there is a severe decrease in number of species, probably due to the replacement and addition of species in different successional stages. The exposure time necessary to periphyton community reaches the maturity stage (maximum in the number and density of species) may vary from two weeks to longer periods, depending on the type of environment, water temperature and the type substratum (LOBO; BUSELATO-TONIOLL, 1985; PATRICK; REIMER, 1975; SABATER et al., 1998). In Corvo Stream, this phase was achieved until the 15th

colonization day, as also verified in other studies (CATTANEO et al., 1975; LAM; LEI, 1999; RODRIGUES; BICUDO, 2001).

Moreover, the environmental processes of each habitat, exert a direct influence on the development, composition, and distribution of phycoperiphytic community, specifically concerning the algal flora of each location (CASCO; TOJA, 1994; STEVENSON, 1996a). Consequently, the high variation in nutrient concentrations (especially phosphorus and nitrogen), mainly from rainfall and constant winds that cause greater displacement of allochthonous material into the environment, besides the resuspension of sediment for the water column, elevating the turbidity, indicate a possible influence on the number and composition of species. On the one hand, when these events are more intense (high precipitation, strong winds) there may be a negative influence, as seen during the 4th sampling, both in natural and artificial substrata, when a sharp decrease in species number was detected. Otherwise, the positive effects of environmental variables previously mentioned may

be corroborated by the correlations between the species richness with nutrients (especially phosphorus and nitrogen), turbidity, wind and rainfall.

The highest number of Zygnemaphyceae species, followed by Bacillariophyceae, in both substrata, leads to a greater representativeness of these unicellular algae. Unicellular organisms are fast colonizers, due to their high reproductive rate; while the colonial or filamentous organisms are slower in propagation process (HILLEBRAND; SOMMER, 2000). These traits should influence the phycoperiphytic communities in the upper Paraná river floodplain, considering the high predominance of these non-flagellate unicellular forms, in the studied environments (RODRIGUES; BICUDO, 2004).

The contribution of diatoms, with several species of *Eunotia* (higher number of species in Penales), forming large filaments of cells aggregate in chains (*E. pectinalis*, *E. sudetica* and *E. camelus*), or composing extensive 'arborescent colonies' connected by the edges (*E. flexuosa* and *E. lineolata*), among others, may be associated to morphological and adaptive traits from each taxa, since these algae are better adapted to periphytic habit, besides being grouped and involved in mucilage sheaths that favor the better attachment to the substrata.

The great representativeness of Zygnemaphyceae (desmids) may be related in part to the amount of available substratum for colonization (diversity of aquatic vegetation), and the high values of nutrients and to the water temperature; and also due to the unicellular condition with mucilage sheaths around the cells. In a study of colonization by periphytic algae in glass slides close to macrophyte stands, Rodrigues and Bicudo (2001) related the high richness of desmids in the periphytic community to the presence of macrophyte stands. Diversified flora of desmids presents faster growth at warmer temperatures, with optimum ranging from 25°C to 30°C (COESEL, 1996; COESEL; WARDENAAR, 1990; FELISBERTO; RODRIGUES, 2005a and b). In this way, Corvo Stream, tropical environment, with temperatures varying between 26 and 28°C, with high nutrient concentrations and diversity of aquatic vegetation, is a favorable local to the development of this algal flora.

Regarding the Chlorococcales, *Desmodesmus* and *Scenedesmus* are exceptionally common in any environment, oligo, meso or eutrophic (BICUDO; MENEZES, 2006; LÜRLING, 2003), and are among the first to colonize the environment (BICUDO; MENEZES, 2006). The high variation in nutrient concentrations of the studied

environment allowed the development of species from this Order. Additionally, *Characium* species, possibly due to the development stage of periphytic biofilm, were favored by the increase of filamentous green algae (as *Oedogonium* and *Bulbochaete*) and also by the quantity of species from the *Eunotia* genus, to which they were fixed through attaching discs.

In summary, the results of taxonomical composition of periphytic algae in natural (*Egeria najas* Planchon) and artificial (plastic plant) substratum were similar, revealing that the development of these algae were also related to environmental conditions, over time. The grouping analysis enabled to register three growth phases of the community, with a maximum peak at 15^o successional day, which allow suggesting that the community achieved the maturity stage. Thus, we may state that the heterogeneity in the structure of phycoperiphytic community may be understood through species composition in different succession stages. Furthermore, the number of species, regardless the substratum type, was positively related to the higher nutrient availability, higher values of turbidity, especially during the first week; and negatively related to the turbidity, during the second and third weeks.

Acknowledgements

We wish to thank the Núcleo de Pesquisas em Limnologia, Ictiologia e Aqüicultura – Nupélia, and PEA (Programa de Pós-Graduação em Ambientes Aquáticos Continentais) from the State University of Maringá, for the logistic, technical and scientific support during the accomplishment of this study; to Capes for the scholarship. We also thank to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for productivity scholarship to LR. This research is inserted in PRONEX Project 'Produtividade em Reservatório: relações com o estado trófico e predação', developed by Nupélia.

Referências

- BICUDO, C. E. M.; MENEZES, M. **Gêneros de algas de águas continentais do Brasil (Chave de identificação e descrições)**. São Carlos: Rima, 2006.
- BIGGS, B. J. F.; THOMSEN, H. A. Disturbance of stream periphyton by perturbations in shear stress: time to structural failure and differences in community resistance. **Journal of Phycology**, v. 31, n. 2, p. 233-241, 1995.
- CASCO, M. A.; TOJA, J. The distribution and interaction of algal communities in reservoir. **Archiv für Hydrobiologie Beih Ergebn Limnologie**, v. 40, p. 85-96, 1994.
- CATTANEO, A.; AMIREAULT, M. C. How artificial are artificial substrata for periphyton? **Journal of the**

- North American Benthological Society**, v. 11, n. 2, p. 244-256, 1992.
- CATTANEO, A.; GHITTORI, S.; VENDEGNA, V. The development of benthonic phytoecoenosis on artificial substrates in the Ticiano River. **Oecologia**, v. 19, n. 4, p. 315-327, 1975.
- COESEL, P. F. M. Biogeography of desmids. **Hydrobiologia**, v. 336, n. 1, p. 41-53, 1996.
- COESEL, P. F. M.; WARDENAAR, K. Growth responses of planktonic desmid species in a temperature: light gradient. **Freshwater Biology**, v. 23, n. 3, p. 551-560, 1990.
- CROASDALE, H.; FLINT, E. A. **Flora of New Zealand: freshwater algae, Chlorophyta, Desmids**. Wellington: Government Printer, 1986.
- CROASDALE, H.; FLINT, E. A. **Flora of New Zealand: freshwater algae, Chlorophyta, Desmids**. Wellington: Government Printer, 1988.
- DILLARD, G. E. **Freshwater algae of the southeastern United States. Part 3. Chrolophyceae: Zygnematales: Mesotaeniaceae and Desmidiaceae (Section 1)**. Berlin: J. Cramer, 1990. (Bibliotheca phycologica, v. 85).
- DILLARD, G. E. **Freshwater algae of the southeastern United States. Part 4. Chrolophyceae: Zygnematales: Desmidiaceae (Section 2)**. Berlin: J. Cramer, 1991. (Bibliotheca phycologica, v. 89).
- FELISBERTO, S. A.; RODRIGUES, L. Influência do gradiente longitudinal (rio-barragem) na similaridade das comunidades de desmídias perifíticas. **Revista Brasileira de Botânica**, v. 28, n. 2, p. 241-254, 2005a.
- FELISBERTO, S. A.; RODRIGUES, L. Comunidade de Algas Perifíticas em reservatórios de diferentes latitudes. In: RODRIGUES, L.; THOMAZ, S. M.; AGOSTINHO, A. A.; GOMES, L. C. (Ed.). **Biocenoses em reservatórios: padrões espaciais e temporais**. São Carlos: Rima, 2005b. p. 97-114.
- FERNANDES, V. O.; ESTEVES, F. A. The use of indices for evaluating the periphytic community in two kinds of substrate in imboassica lagoon, Rio de Janeiro, Brazil. **Brazilian Journal of Biology**, v. 63, n. 2, p. 233-243, 2003.
- FÖRSTER, K. Conjugatophyceae: Zygnematales und Desmidiales (excl. Zygnemataceae). In: HUBER-PESTALOZZI, G. (Ed.). **Das Phytoplankton des Süßwassers: systematik und biologie**. Stuttgart: Schweizerbart'sche Verlagsbuchhandlung, 1982. p. 1-543.
- GOLTERMAN, H. L.; CLYMO, R. S.; OHMSTAD, M. A. M. **Methods for physical and chemical analysis of freshwaters**. Oxford: Blackwell Scientific Publication, 1978.
- HAMBROOK, J. A. Bioassessment of stream-water quality using benthic and planktonic algae collected along an urban intensity gradient in the Eastern Cornbelt Plains Ecoregion, Ohio, USA. **Journal of Phycology**, v. 38, n. 1, p. 14-15, 2002.
- HILLEBRAND, H.; SOMMER, U. Diversity of benthic microalgae in response to colonization time and eutrophication. **Aquatic Botany**, v. 67, n. 3, p. 221-236, 2000.
- KOMÁREK, J.; ANAGNOSTIDIS, K. Cyanoprocaryota 1 – Chroococcales. In: Ettl, H.; Gerloff, J.; Heynig, H.; Mollenhauer, D. (Ed.). **Süßwasserflora von Mitteleuropa**. Band 19/1. Stuttgart: Gustav Fischer, 1999. p. 1-548.
- KOMÁREK, J.; ANAGNOSTIDIS, K. Cyanoprocaryota 2 – Oscillatoriales. In: Ettl, H.; Gerloff, J.; Heynig, H.; Mollenhauer, D. (Ed.). **Süßwasserflora von Mitteleuropa**. Band 19/2. Heidelberg, Elsevier: 2005. p. 1-759.
- KRAMMER, K.; LANGE-BERTALOT, H. Bacillariophyceae: Naviculaceae. In: Ettl, H.; Gerloff, J.; Heynig, H.; Mollenhauer, D. (Ed.). **Süßwasserflora von Mitteleuropa**. Stuttgart: Gustav Fischer, 1986. v. 2, p. 1-596.
- KRAMMER, K.; LANGE-BERTALOT, H. Bacillariophyceae: Bacillariaceae, Epithemiaceae, Surirellaceae. In: Ettl, H.; Gerloff, J.; Heynig, H.; Mollenhauer, D. (Ed.). **Süßwasserflora von Mitteleuropa**. Stuttgart: Gustav Fischer, 1988. v. 2, p. 1-596.
- KRAMMER, K.; LANGE-BERTALOT, H. Bacillariophyceae: Centrales, Fragilariaceae, Eunotiaceae. In: Ettl, H.; Gerloff, J.; Heynig, H.; Mollenhauer, D. (Ed.). **Süßwasserflora von Mitteleuropa**. Stuttgart: Gustav Fischer, 1991. v. 2, p. 1-576.
- LAM, P. K. S.; LEI, A. Colonization of periphytic algae on artificial substrates in a tropical stream. **Diatom Research**, v. 14, n. 2, p. 307-322, 1999.
- LOBO, E.; BUSELATO-TONIOLLI, T. C. Tempo de exposição de um substrato artificial para o estabelecimento da comunidade do perifíton no curso inferior do rio Caí, Rio Grande do Sul, Brasil. **Rickia**, v. 12, p. 35-51, 1985.
- LÜRLING, M. Phenotypic plasticity in the green algae *Desmodesmus* and *Scenedesmus* with special reference to the induction of defensive morphology. **Annales de Limnologie**, v. 39, n. 2, p. 85-101, 2003.
- MACKERETH, F. J. H.; HERON, J.; TALLING, J. F. **Water analysis: some revised methods for limnologists**. Ambleside: Scientific Publication, 1978.
- MOSCHINI-CARLOS, V.; HENRY, R.; POMPÊO, M. L. M. Seasonal variation of biomass and productivity of the periphytic community on artificial substrata in the Jurumirim Reservoir (São Paulo, Brazil). **Hydrobiologia**, v. 434, n. 1-3, p. 35-40, 2000.
- OLIVEIRA, M. A.; TORGAN, L. C.; LOBO, E. A.; SCHARZBOLD, A. Association of periphytic diatom species of artificial substrate in lotic environments in the Arroio Sampaio basin, RS, Brazil: relationships with abiotic variables. **Brazilian Journal of Biology**, v. 61, n. 4, p. 523-540, 2001.
- PATRICK, R.; REIMER, C. W. **The diatoms of the United States, exclusive of Alaska and Hawaii**. Philadelphia: Livingstgen Company, 1966. v. 1, pt. 1.
- PATRICK, R.; REIMER, C. W. **The Diatoms of United States: exclusive of Alaska and Hawaii**. Philadelphia: Philadelphia Academy of Natural Sciences, 1975. v. 2, part. 1, n. 13, p. 1-214.
- PETERSON, C. G. Response of Benthic Algal communities to natural physical disturbance. In: STEVENSON, R. J.; BOTHWELL, M. L.; LOWE, R. L. (Ed.). **Algal ecology: freshwater benthic ecosystems**. San Diego: Academic Press, 1996. cap. 13, p. 375-402.

- PRESCOTT, G. W. **Algae of de western great lakes area**. Koenigstein: Otto Koeltz Science, 1982.
- PRESCOTT, G. W.; BICUDO, C. E. M.; VINYARD, W. C. A Synopsis of North American Desmids. Part II. Desmidiaceae: Placodermae. Section 4. In: PRESCOTT, G. W. (Ed.). **Desmidiales**. Lincoln: University Nebraska Press, 1982. p. 1-700.
- PRESCOTT, G. W.; CROASDALE, H. T.; VINYARD, W. C.; BICUDO, C. E. M. A Synopsis of North American Desmids. Part II. Desmidiaceae: Placodermae. Section 3. In: PRESCOTT, G. W. (Ed.). **Desmidiales**. Lincoln: University Nebraska Press, 1981. p. 1-720.
- RODRIGUES, L.; BICUDO, D. C. Similarity among periphyton algal communities in a lentic-lotic gradient of the upper Paraná river floodplain, Brazil. **Revista Brasileira de Botânica**, v. 24, n. 3, p. 235-248, 2001.
- RODRIGUES, L.; BICUDO, D. C. Periphytic Algae. In: THOMAZ, S. M.; AGOSTINHO, A. A.; HAHN, N. S. (Ed.). **The upper Paraná river and its floodplain: physical aspects, ecology and conservation**. Leiden: Backhuys, 2004. p. 125-143.
- RODRIGUES, L.; BICUDO, D. C.; MOSCHINI-CARLOS, V. O papel do perifiton em áreas alagáveis e nos diagnósticos ambientais. In: THOMAZ, S. M.; BINI, L. M. (Ed.). **Ecologia e manejo de macrófitas aquáticas**. Maringá: Eduem, 2003. cap. 10, p. 211-230.
- ROHLF, F. J. **NTSYS-PC: numerical taxonomy and multivariate analysis system: version 2.1**. New York: Exeter Publications, 2000.
- ROUND, F. E. **The biology of the algae**. London: Edward Arnold, 1965.
- ROUND, F. E. The taxonomy of the Chlorophyta, 2. **British Phycological Journal**, v. 6, n. 2, p. 235-264, 1971.
- SABATER, S.; GREGORY, S. V.; SEDELL, J. R. Community dynamics and metabolism of benthic algae colonizing wood and rock substrata in a forest stream. **Journal of Phycology**, v. 34, n. 4, p. 561-567, 1998.
- SOLORZANO, L. Determination of ammonia in natural waters by the phenolhypochlorite method. **Limnology and Oceanography**, v. 14, n. 5, p. 799-801, 1969.
- STATSOFT, Inc. **Statistica** (data analysis software system): version 7.1. Disponível em: <www.statsoft.com>. Acesso em: 2005.
- STEVENSON, R. J. An introduction to algal ecology in freshwater benthic habitats. In: STEVENSON, R. J.; BOTHWELL, M. L.; LOWE, R. L. (Ed.). **Algal ecology: freshwater benthic ecosystems**. San Diego: Academic Press, 1996a. cap. 1, p. 3-30.
- STEVENSON, R. J. The stimulation and drag of current. In: STEVENSON, R. J.; BOTHWELL, M. L.; LOWE, R. L. (Ed.). **Algal ecology: freshwater benthic ecosystems**. San Diego: Academic Press, 1996b. cap. 11, p. 321-340.
- STEVENSON, R. J. Scale-dependent determinants and consequences of benthic algal heterogeneity. **Journal of the North American Benthological Society**, v. 16, n. 1, p. 248-262, 1997.
- STEVENSON, R. J.; PETERSON, C. G. Variation in benthic diatom (Bacillariophyceae) immigration with habitat characteristics and cell morphology. **Journal of Phycology**, v. 25, n. 1, p. 120-129, 1989.
- VALDERRAMA, J. C. The simultaneous analysis of total nitrogen and total phosphorus in natural waters. **Marine Chemistry**, v. 10, n. 2, p. 109-122, 1981.
- VERCELLINO, I. S.; BICUDO, D. C. Sucessão da comunidade de algas perifíticas em reservatório oligotrófico tropical (São Paulo, Brasil): comparação entre período seco e chuvoso. **Revista Brasileira de Botânica**, v. 29, n. 3, p. 363-377, 2006.
- WETZEL, R. G. **Limnologia**. Barcelona: Omega, 1981.
- WETZEL, R. G. Recommendations for future research on periphyton. In: WETZEL, R. L. (Ed.). **Periphyton of freshwater ecosystems developments in hidrobiology**. The Hague: Dr. W. Junk, 1983. p. 339-346.
- WETZEL, R. G.; LIKENS, G. E. **Limnological analyses**. 2nd ed. New York: Springer Verlag, 1991.
- WU, N.; TANG, T.; ZHOU, S.; JIA, X.; LI, D.; LIU, R.; CAI, Q. Changes in benthic algal communities following construction of a run-of-river dam. **Journal of the North American Benthological Society**, v. 28, n. 1, p. 69-79, 2009.

Received on July 28, 2008.

Accepted on July 15, 2009.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.