



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Understanding Acid-Base Disorders

Maxwell, A., Hamilton, P. K., Connolly, G. M., & Morgan, N. (2017). Understanding Acid-Base Disorders. The Ulster Medical Journal.

**Published in:**  
The Ulster Medical Journal

**Document Version:**  
Publisher's PDF, also known as Version of record

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

Copyright 2017 the authors.

This is an open access article published under a Creative Commons Attribution-NonCommercial-ShareAlike License (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits use, distribution and reproduction for non-commercial purposes, provided the author and source are cited and new creations are licensed under the identical terms.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

# Understanding Acid-Base Disorders

Paul K. Hamilton<sup>1</sup>, Neal A. Morgan<sup>2</sup>, Grainne M. Connolly<sup>3</sup> and Alexander P. Maxwell<sup>4</sup>

Accepted: 8th January 2017

Provenance: externally peer-reviewed

<i>Case for consideration</i>			
<i>A 25 year old mechanic is admitted in a very confused and drowsy state. His initial laboratory results are listed below. What diagnoses should be considered?</i>			
	<i>Reference Range</i>		<i>Reference Range</i>
<i>pH 7.1</i>	<i>7.35-7.45</i>	<i>Na<sup>+</sup> 136 mmol/L</i>	<i>136-145</i>
<i>PaO<sub>2</sub> 13.4 kPa</i>	<i>11.0-14.0</i>	<i>K<sup>+</sup> 3.8 mmol/L</i>	<i>3.5-5.3</i>
<i>PaCO<sub>2</sub> 2.5 kPa</i>	<i>4.5-6.0</i>	<i>Cl<sup>-</sup> 95 mmol/L</i>	<i>95-108</i>
<i>Bicarbonate 5.3 mmol/L</i>	<i>22-29</i>	<i>Total CO<sub>2</sub> 6 mmol/L</i>	<i>22-29</i>
<i>Lactate 1.2 mmol/L</i>	<i>0.6-2.4</i>	<i>Urea 4.2 mmol/L</i>	<i>2.5-7.8</i>
<i>Ketones &lt;1 mmol/L</i>	<i>Depends on context</i>	<i>Creatinine 92 µmol/L</i>	<i>40-110</i>
		<i>eGFR &gt;60 mL/min/1.73m<sup>2</sup></i>	<i>&gt;60</i>

## INTRODUCTION

The accurate interpretation of laboratory tests in patients with acid-base disorders is critical for understanding pathophysiology, making a diagnosis, planning effective treatment and monitoring progress. This is an important topic particularly for junior medical staff who may encounter acid-base problems outside normal working hours when patients become acutely unwell. These clinical situations may be a source of confusion particularly because of the variety of terms used to describe and classify acid-base disorders. In this article, we aim to provide the reader with an overview of the key concepts necessary for developing a good working understanding of acid-base disorders that commonly present in clinical medicine. We start with some acid-base disorder definitions and then provide a series of case vignettes to illustrate the key points.


## DEFINITIONS

- Acidaemia** An arterial pH below the normal range (pH<7.35).
- Alkalaemia** An arterial pH above the normal range (pH>7.45).
- Acidosis** A process lowering pH. This may be caused by a fall in serum bicarbonate and/or a rise in the partial pressure of carbon dioxide (PaCO<sub>2</sub>).
- Alkalosis** A process raising pH. This may be caused by a rise in serum bicarbonate and/or a fall in PaCO<sub>2</sub>.

## ACID-BASE HOMEOSTASIS

Like temperature, blood pressure, osmolality and many other physiological parameters, the human body strives to keep its acid-base balance within tightly controlled limits. It is not the aim of this article to review in detail the physiology of acid-base homeostasis, but to provide a working knowledge of some key concepts that will help in the interpretation of results encountered commonly in clinical practice. More detailed free text reviews of acid-base homeostasis are available<sup>1-5</sup>.

A buffer is a solution that resists a change in pH. There are many different buffer systems in the body, but the key one for understanding most acid-base disorders is the bicarbonate system present in the extracellular fluid. Like any buffer, this system comprises a weak acid (in this case carbonic acid, H<sub>2</sub>CO<sub>3</sub>) and its conjugate base (the bicarbonate ion, HCO<sub>3</sub><sup>-</sup>), which exist in a dynamic equilibrium as shown in Equation 1<sup>6</sup>:



Equation 1

1 Specialty Registrar in Chemical Pathology (Metabolic Medicine), Belfast Health and Social Care Trust; Honorary Lecturer, Queen's University Belfast. 2 Consultant Nephrologist, Southern Health and Social Care Trust. 3 Consultant Chemical Pathologist, Belfast Health and Social Care Trust. 4 Consultant Nephrologist, Belfast Health and Social Care Trust


a.p.maxwell@qub.ac.uk

Correspondence to Prof AP Maxwell.

Address: Regional Nephrology Unit, Belfast City Hospital, Belfast, BT9 7AB, United Kingdom.



The acidity of a solution is governed by the concentration of hydrogen ions ( $H^+$ ) present. If a disease process results in an increase in the concentration of hydrogen ions, one would expect the body to become more acidic. However, the bicarbonate buffer system resists this change because the excess of hydrogen ions drives the reaction in Equation 1 to the right: hydrogen ions react with and “consume” bicarbonate ions and any change in acidity is minimised. This process requires an adequate supply of bicarbonate ions. The kidneys are vital organs in acid-base balance as they can both generate “new” bicarbonate buffer and reclaim filtered bicarbonate in the proximal tubules (Figure 1).



**Fig 1.** Acid-base balance is maintained by effective renal and respiratory homeostatic mechanisms

By rearranging and simplifying the above acid-base reaction, it is possible to derive the useful relationship shown in Equation 2:

$$H^+ \text{ is proportional to } \frac{PaCO_2}{HCO_3^-}$$


*Equation 2*

Equation 2 helps to illustrate how the body’s hydrogen ion concentration can be regulated by altering the ratio of  $CO_2$  to bicarbonate. Ventilation controls the  $PaCO_2$  level and the kidneys regulate the bicarbonate level (Figure 1).

This makes it easy to see that the concentration of hydrogen ions increases in two settings: an increase in  $PaCO_2$  or a reduction in plasma bicarbonate. One of the functions of ventilation is the elimination of  $CO_2$  during exhalation. If a patient is tachypnoeic, they will tend to lose  $CO_2$ , while patients with a reduced respiratory drive will retain  $CO_2$ . An increased concentration of hydrogen ions (an acidosis) stimulates the respiratory centre to increase the rate of breathing (exhaling more  $CO_2$ ). This mechanism is another key physiological response that helps to maintain acid-base balance.

Acid-base disorders are broadly classified into problems involving metabolic and/or respiratory processes. Metabolic processes primarily direct change in the level of bicarbonate and respiratory processes primarily direct changes in  $PaCO_2$  (Figure 2).

The body adapts, or compensates where there is an acid-base disturbance in an attempt to maintain homeostasis<sup>7</sup>.



**Fig 2.** Changes in  $PaCO_2$  level and bicarbonate concentration  $[HCO_3^-]$  can help identify the nature of the acid-base disorder.

If the primary acid-base problem is metabolic, then the compensatory mechanism is respiratory. The respiratory rate is altered, usually within minutes, in an attempt to keep the hydrogen ion concentration normal. If the primary acid-base problem is respiratory, then the kidneys adapt to counteract the change by altering their handling of hydrogen ions. This process in the kidneys usually takes place over several days.

### CAUSES OF ACID-BASE DISORDERS

Acid-base disorders are classified according to whether there is acidosis or alkalosis present (see pH section for details), and whether the primary problem is metabolic or respiratory (Figure 2). Bear in mind that there may be more than one problem occurring simultaneously and that the body may be compensating for the derangement. Table 1 outlines, with some clinical examples, acid-base disorders that are commonly encountered.

TABLE 1

#### Definitions and main causes of acid-base disorders

Metabolic acidosis	Process that primarily reduces bicarbonate: Excessive $H^+$ formation e.g. lactic acidosis, ketoacidosis Reduced $H^+$ excretion e.g. renal failure Excessive $HCO_3^-$ loss e.g. diarrhoea
Metabolic alkalosis	Process that primarily raises bicarbonate: Extracellular fluid volume loss e.g. due to vomiting or diuretics Excessive potassium loss with subsequent hyperaldosteronism
Respiratory acidosis	Process that primarily causes elevation in $PaCO_2$ : Reduced effective ventilation e.g. many chronic respiratory diseases or drugs depressing the respiratory centre
Respiratory alkalosis	Process that primarily causes reduction in $PaCO_2$ : Increased ventilation e.g. in response to hypoxia or secondary to a metabolic acidosis

Remember, metabolic processes primarily direct changes in bicarbonate and respiratory processes primarily direct changes in  $PaCO_2$  (Figure 2).

### MEASURED AND DERIVED INDICES

Some potentially confusing terminology is often used when discussing acid-base disorders. These terms include  $PaCO_2$ ,



total bicarbonate, total  $\text{CO}_2$ , standard bicarbonate and base excess<sup>8</sup>. It is useful to know what these terms mean and how they are derived. Most blood gas analysis is carried out on point-of-care blood gas analysers, and these generally only measure two substances when it comes to acid-base reports: hydrogen ions (from which pH is calculated – see below) and  $\text{PaCO}_2$ . The ‘bicarbonate’ results that are given from such analysers are generally calculated using Equation 2.

Most laboratories measure total  $\text{CO}_2$  concentration as part of the standard electrolyte profile. The reason behind this is that it is technically difficult to measure bicarbonate ions in isolation, but relatively straightforward to measure total  $\text{CO}_2$ . Total  $\text{CO}_2$  represents the total amount of bicarbonate ions, dissolved  $\text{CO}_2$  and other  $\text{CO}_2$ -containing substances in a solution. Since bicarbonate normally constitutes the majority of this, total  $\text{CO}_2$  is normally used as a convenient surrogate measure of bicarbonate. The total  $\text{CO}_2$  on the electrolyte profile may provide the first clue to the presence of an acid-base disturbance in a patient and should not be overlooked when reviewing electrolyte results. One cannot, however, diagnose acid-base disturbances from an isolated total  $\text{CO}_2$  measurement. In order to characterise an acid-base disturbance, measures of pH,  $\text{PaCO}_2$ , total  $\text{CO}_2$  or bicarbonate are required, as well as measurement of the anion gap.

Standard bicarbonate is a calculated index that attempts to provide information on what the bicarbonate concentration would be if the respiratory components of the disorder were eliminated. Base excess is another calculated index which will be elevated in the setting of metabolic alkalosis and reduced in metabolic acidosis. We will not consider the use of these calculated indices further in this article.

## UNDERSTANDING ACID-BASE DISORDERS – A FOUR STEP APPROACH

In order to understand the nature of an acid-base problem, we recommend a structured approach during which the following four questions should be asked.

### Question 1: What is the pH?

The first step in interpreting an acid-base problem is to look at the pH (or  $[\text{H}^+]$ ) and decide if you are dealing with acidosis, alkalosis or normality. The concept of pH as a measure of acidity will already be familiar. With most human enzymes favouring physiologically neutral conditions, acidaemia is deemed to be present when the pH is less than 7.35 and alkalaemia when the pH exceeds 7.45. It is becoming increasingly common to directly quote the concentration of hydrogen ions ( $[\text{H}^+]$ ) present in a solution. pH and  $[\text{H}^+]$  are directly related using Equation 3:

$$\text{pH} = -\log_{10} [\text{H}^+], \text{ where } [\text{H}^+] \text{ is in mol/L}$$

Equation 3

Thus, pH 6.8 corresponds to  $1.6 \times 10^{-7}$  mol/L  $[\text{H}^+]$ , pH 7.4 to  $4.0 \times 10^{-8}$  mol/L, and pH 7.6 to  $2.5 \times 10^{-8}$  mol/L, i.e. pH falls as  $[\text{H}^+]$  rises.

Because the body compensates for acid-base disorders, it is possible that a disorder might be present even if the pH is normal. It should also be borne in mind that the body never over-compensates.

### Question 2: What is the bicarbonate?

The second step in interpreting an acid-base disorder is to consider the bicarbonate concentration relative to the normal reference range (which will vary from laboratory to laboratory, but is typically in the range 22-29 mmol/L).

A reduced bicarbonate concentration could mean that the body’s main buffer is being used up buffering excess acid (hydrogen ion) production e.g. in lactic acidosis or ketoacidosis. Alternatively the reduced bicarbonate concentration could indicate a problem related to loss of bicarbonate from the gastrointestinal tract e.g. diarrhoea or a kidney problem i.e. failure to generate new bicarbonate or reclaim bicarbonate filtered into the renal tubules. **A reduced bicarbonate concentration is a hallmark of metabolic acidosis.**

An increased bicarbonate concentration may indicate that there have been substantial losses of acidic fluid e.g. loss of gastric fluid from persistent vomiting or prolonged nasogastric aspiration. Alternatively an increased bicarbonate concentration may be a chronic adaptation by the kidney to high  $\text{PaCO}_2$  levels in persons with chronic respiratory diseases associated with  $\text{CO}_2$  retention (see Equation 1 where elevated  $\text{CO}_2$  levels drives the equation to the left producing more bicarbonate). **An elevated bicarbonate concentration is a feature of metabolic alkalosis.**

### Question 3: What is the $\text{PaCO}_2$ ?

The third step in assessing an acid-base problem is to measure the  $\text{PaCO}_2$ . This is helpful in determining whether the respiratory system is responding normally to an acid load and reducing the  $\text{PaCO}_2$  to compensate for an acidosis i.e. the primary acid-base disturbance is a metabolic acidosis and this is compensated by an increased respiratory rate resulting in a secondary respiratory alkalosis. **A decreased  $\text{PaCO}_2$  is a feature of respiratory alkalosis.**

Alternatively, if there is a primary respiratory problem, e.g. respiratory failure associated with chronic obstructive pulmonary disease, the retained  $\text{CO}_2$  results in an elevated  $\text{PaCO}_2$  (and will drive Equation 1 to the left) and produce a respiratory acidosis. It is also possible to develop a respiratory acidosis if drugs, such as opiate analgesics, depress the respiratory centre resulting in a critical reduction in the rate of ventilation resulting in  $\text{CO}_2$  retention. **An elevated  $\text{PaCO}_2$  is a feature of respiratory acidosis.**

*One can see that by examining the pH, bicarbonate and  $\text{PaCO}_2$  it is possible to deduce the nature of the primary acid-base disorder present and the compensatory response.*

### Question 4: What is the anion gap?



The final step in assessing an acid-base disorder is to calculate the anion gap. Bodily fluids are electrically neutral, meaning that the number of positive charges (cations) present equals the number of negative charges (anions). The most abundant anions are chloride and bicarbonate; numerous other anions are not routinely quantitated, for example proteins and sulphate ions. Sodium is by far the most abundant plasma cation; other cations present in much lower quantities include potassium, calcium and magnesium. If it were feasible to measure all charged substances in blood, it could be shown that the sum of the positively charged particles is exactly balanced by the number of those substances carrying negative charges. It is routine practice to measure only four charged particles: sodium, potassium, chloride and bicarbonate ions. As discussed earlier, total CO<sub>2</sub> on the electrolyte profile may be considered as a convenient surrogate measure of bicarbonate and can be used in the calculation of the anion gap. When the numbers of cations (sodium and potassium) are added, one will always find that they outnumber the anions (chloride and bicarbonate). This difference is what is meant by the term ‘anion gap’ and reflects the unmeasured anions in Equation 4 or Equation 5. An anion gap may be low, normal or high, and can be conveniently calculated using Equation 4:

$$\text{Anion Gap} = [\text{Na}^+] + [\text{K}^+] - [\text{Cl}^-] + [\text{HCO}_3^-], \text{ where all concentrations are in mmol/L.}$$

*Equation 4*

Since the extracellular fluid potassium concentration is very much lower than the sodium, chloride or bicarbonate concentrations and because it can only vary by a few mmol/L, it is often ignored making the anion gap calculation simpler as shown in Equation 5:

$$\text{Anion gap} = [\text{Na}^+] - [\text{Cl}^-] + [\text{HCO}_3^-]$$

*Equation 5*

The reference interval (normal range) for anion gap varies from laboratory to laboratory, and is inherently imprecise because of the number of measurements required for its calculation. An anion gap greater than 20 mmol/L is always considered to be abnormally elevated and a gap of less than 10 mmol/L abnormally low. There is some debate in the literature about the significance of anion gaps in the range 10-20 mmol/L, but a pragmatic approach would be to actively seek out causes of a high anion gap in patients with gaps exceeding 14 mmol/L (or 18 mmol/L if potassium is included in the equation).

TABLE 2

*Causes of metabolic acidosis (common causes are in bold)*

Normal anion gap	High anion gap
Gastrointestinal losses of bicarbonate	Renal failure
Renal tubular acidosis	Ketoacidosis
Treatment with carbonic anhydrase inhibitors	Lactic acidosis
Urinary diversion procedures	Salicylate poisoning
Excessive administration of 0.9% saline	Glycol ingestion (ethylene glycol, propylene glycol)
	Methanol ingestion





Fig 3

(a) Illustration of the “normal” anion gap



(b) High anion gap present in a metabolic acidosis

Consider the following normal electrolyte profile: Na<sup>+</sup> 136 mmol/L, K<sup>+</sup> 4.0 mmol/L, Cl<sup>-</sup> 100 mmol/L, HCO<sub>3</sub><sup>-</sup> (or total CO<sub>2</sub>) 25 mmol/L. The anion gap is calculated as 140 – 100 – 25 = 11 mmol/L (or 15 mmol/L if potassium is included in the calculation). The Anion Gap is illustrated in Figure 3a.

TABLE 3

*Mnemonics for high anion gap metabolic acidosis*

Mnemonic	Causes of a high anion gap metabolic acidosis
<b>GOLD MARK</b>	Glycols, Oxoprolin, L-lactate, D-lactate, Methanol, Aspirin, Renal failure, and Ketoacidosis
<b>KARMEL</b>	Ketoacidosis, Aspirin, Renal Failure, Methanol, Ethylene Glycol, and Lactic Acidosis
<b>MUDPILES</b>	Methanol, Uraemia, Diabetes, Paraldehyde, Iron/Isoniazid, Lactate, Ethylene glycol and Salicylate
<b>KUSMALE</b>	Ketoacidosis, Uraemia, Salicylate, Methanol, Aldehyde, Lactate and Ethylene glycol

Calculation of the anion gap is particularly useful in cases of metabolic acidosis since it can help in formulating a differential diagnosis<sup>10</sup>. There are two main categories of metabolic acidosis: high anion gap metabolic acidosis (HAGMA) and normal anion gap metabolic acidosis (NAGMA). A HAGMA is illustrated in Figure 3b. Common causes of HAGMA and NAGMA are detailed in Table 2.



Several mnemonics for common causes of HAGMA have been developed<sup>11</sup>, and some of the more useful examples are included in Table 3.

From a clinical perspective, if a HAGMA is identified then the simplest approach to establishing a cause is to consider if the patient has one (or more) of the three common aetiologies (lactic acidosis, ketoacidosis or kidney failure)<sup>12</sup>.

If these conditions are not present then the HAGMA may be linked to ingestion of a toxin e.g. methanol or ethylene glycol, or be due to the build-up of another acid such as 5-oxoproline (also known as pyroglutamic acid) which may accumulate with chronic paracetamol use in susceptible individuals<sup>13</sup>.

As the laboratory tests for toxic alcohols are not rapidly available it can be useful in a patient with an unexplained HAGMA to assess the “osmolal gap”<sup>14</sup>. This “gap” is the difference between the calculated serum osmolality and the laboratory measurement of serum osmolality (from a U&E sample). The calculated osmolality can be simply derived by using Equation 6. A high osmolal gap suggests the presence of toxic alcohols such as methanol or ethylene glycol.

$$\text{Calculated osmolality (mmol/L)} = 2 \times [\text{Na}^+] + [\text{glucose}] + [\text{urea}]$$

Equation 6

Rarely, patients with short bowel syndrome or following bariatric surgery can develop severe D-lactic acidosis and an associated encephalopathy<sup>15</sup>. Unabsorbed carbohydrates act as a substrate for colonic bacteria to produce D-lactate. This will result in a high anion gap metabolic acidosis but the standard laboratory measured lactate (L-lactate) will be normal<sup>15</sup>.

Calculated anion gaps that are low (below the reference interval) are uncommon. Causes include laboratory error or hypoalbuminaemia but rarely may be found in association with a paraproteinaemia or intoxication with lithium, bromide, or iodide<sup>10</sup>.

## ILLUSTRATIVE CASES

### Case 1

An elderly man is admitted with septic shock. Shortly after admission, blood tests reveal the following:

pH 7.18, PO<sub>2</sub> 34.2 kPa, PaCO<sub>2</sub> 2.1 kPa, HCO<sub>3</sub><sup>-</sup> 7 mmol/L

Na<sup>+</sup> 138 mmol/L, K<sup>+</sup> 3.9 mmol/L, Cl<sup>-</sup> 95 mmol/L, Total CO<sub>2</sub> 8 mmol/L, Urea 8.2 mmol/L, Creatinine 102 µmol/L, eGFR >60 mL/min/1.73m<sup>2</sup>

Question 1: What is the pH? The pH is low indicating an acidosis.

Question 2: What is the bicarbonate? Bicarbonate is low, indicating that the acidosis is metabolic in nature.

Question 3: What is the PaCO<sub>2</sub>? The PaCO<sub>2</sub> is low, reflecting a respiratory alkalosis. The low level seen here is a reflection

of the body's compensation in an attempt to correct the pH, i.e. a compensatory respiratory alkalosis is present.

Question 4: What is the anion gap? The anion gap is high, indicating HAGMA.

The most likely cause for this acid-base disorder is lactic acidosis due to poor tissue perfusion as a result of septic shock.

### Case 2

A woman is being treated for congestive cardiac failure on the coronary care unit. After several days of treatment, the following results are returned:

pH 7.49, PO<sub>2</sub> 11.6 kPa, PaCO<sub>2</sub> 5.8 kPa, HCO<sub>3</sub><sup>-</sup> 42 mmol/L

Na<sup>+</sup> 142 mmol/L, K<sup>+</sup> 3.0 mmol/L, Cl<sup>-</sup> 85 mmol/L, Total CO<sub>2</sub> 44 mmol/L, Urea 9.3 mmol/L, Creatinine 84 µmol/L, eGFR >60 mL/min/1.73m<sup>2</sup>

Question 1: What is the pH? The pH is high indicating an alkalosis.

Question 2: What is the bicarbonate? Bicarbonate is high, in keeping with a metabolic alkalosis

Question 3: What is the PaCO<sub>2</sub>? The result towards the higher end of the reference range reflects a degree of respiratory compensation for the metabolic alkalosis.

Question 4: What is the anion gap? The anion gap is 13 mmol/L which is normal.

The most likely cause for this acid-base abnormality is extracellular fluid volume loss and hypokalaemia due to treatment with diuretics.

### Case 3

An elderly woman with chronic obstructive pulmonary disease (COPD) is admitted with increasing confusion. Shortly after admission, blood tests reveal the following:

pH 7.21, PO<sub>2</sub> 8.2 kPa, PaCO<sub>2</sub> 11.1 kPa, HCO<sub>3</sub><sup>-</sup> 35 mmol/L

Na<sup>+</sup> 140 mmol/L, K<sup>+</sup> 4.7 mmol/L, Cl<sup>-</sup> 94 mmol/L, Total CO<sub>2</sub> 34 mmol/L, Urea 8.2 mmol/L, Creatinine 66 µmol/L, eGFR >60 mL/min/1.73m<sup>2</sup>

Question 1: What is the pH? The pH is low indicating an acidosis.

Question 2: What is the bicarbonate? Bicarbonate is high, indicating that a metabolic alkalosis is present. The pH is low so the primary problem is an acidosis and is likely to be respiratory in nature.

Question 3: What is the PaCO<sub>2</sub>? The PaCO<sub>2</sub> is very high and indicates a respiratory acidosis is present. The very high PaCO<sub>2</sub> level seen here is typical of a person with respiratory disease that results in retention of CO<sub>2</sub> i.e. the primary clinical problem is respiratory failure due to COPD.

Question 4: What is the anion gap? The calculated anion gap is 12 mmol/L i.e. normal.

The most likely cause for this acid-base abnormality is an acute exacerbation of COPD.

#### Case 4

An elderly man developed profuse diarrhoea following antibiotic treatment of a chest infection. He is thirsty and light headed. Shortly after admission, blood tests reveal the following:

pH 7.25, PO<sub>2</sub> 13.2 kPa, PaCO<sub>2</sub> 4.2 kPa, HCO<sub>3</sub><sup>-</sup> 17 mmol/L

Na<sup>+</sup> 134 mmol/L, K<sup>+</sup> 3.4 mmol/L, Cl<sup>-</sup> 104 mmol/L, Total CO<sub>2</sub> 18 mmol/L, Urea 9.3 mmol/L, Creatinine 102 µmol/L, eGFR >60 mL/min/1.73m<sup>2</sup>

Question 1: What is the pH? The pH is low indicating an acidosis.

Question 2: What is the bicarbonate? Bicarbonate is low, indicating that a metabolic acidosis is present.

Question 3: What is the PaCO<sub>2</sub>? The PaCO<sub>2</sub> level is just below the lower end of the normal range indicating a respiratory alkalosis is present. The pH is low so the primary problem is an acidosis (metabolic acidosis). The respiratory alkalosis therefore represents partial compensation of the metabolic acidosis.

Question 4: What is the anion gap? The anion gap is 12 mmol/L, indicating that this is a normal anion gap metabolic acidosis.

The most likely cause for this acid-base disorder is bicarbonate loss from the gastrointestinal tract due to diarrhoea.

#### Returning to our initial case...

Applying the four question approach to this case, it should now be apparent that the patient has a high anion gap metabolic acidosis with respiratory compensation. The common causes for this presentation can be quickly eliminated since his renal function is normal, and lactate and ketone levels are not elevated. A more unusual explanation for the presentation should be sought (see Table 2). In this case, the patient was subsequently found to have ingested 500 mL of screenwash containing ethylene glycol (antifreeze) in an attempt to end his life.

Prompt recognition of the likely cause of this patient's high anion gap metabolic acidosis helps inform further investigation and management. This would include quantitation of ethanol and toxic alcohol concentrations to confirm the type of ingested poison. Ethylene glycol and methanol are metabolised by alcohol dehydrogenase to very toxic metabolites. If this diagnosis seems likely it is important to urgently seek senior help. Fomepizole is an alcohol dehydrogenase inhibitor which is easy to administer and prevents metabolism of these alcohols to their toxic

metabolites. Haemodialysis will rapidly clear ethylene glycol, methanol and their metabolites and should be started if the patient is severely acidaemic or has evidence of end organ damage e.g. renal failure or visual loss.

#### CONCLUSION

Acid-base disorders are commonly encountered in clinical practice and a structured approach to assessment includes taking a history, performing a physical examination and careful interpretation of routine biochemical tests and arterial blood gas analysis. Additional investigations such as lactate, glucose, ketones or toxicology testing may be needed to more fully characterise a metabolic acidosis.

Answering four questions will help determine the problems present in the clinical scenario: What is the pH? What is the bicarbonate? What is the PaCO<sub>2</sub>? What is the anion gap? Using this approach will help guide further investigations and management of the patient.

There are no conflicts of interest.

#### REFERENCES

1. Adrogué HJ, Madias NE. Management of life-threatening acid-base disorders. First of two parts. *N Engl J Med.* 1998; **338**(1): 26-34.
2. Adrogué HJ, Madias NE. Management of life-threatening acid-base disorders. Second of two parts. *N Engl J Med.* 1998; **338**(2): 107-11.
3. Haber RJ. A practical approach to acid-base disorders. *West J Med.* 1991; **155**(2): 146-51.
4. Hamm LL, Nakhoul N, Hering-Smith KS. Acid-Base Homeostasis. *Clin J Am Soc Nephrol.* 2015; **10**(12): 2232-42.
5. Scott MG, LeGrys VA, Hood JL. Blood gases. Chapter 28. In: Burtis CA, Ashwood ER, Bruns DE, editors. *Tietz Textbook of clinical chemistry and molecular diagnostics.* 5<sup>th</sup> ed. St Louis: Elsevier; 2012. p. 807-35
6. Rose DB, Post TW. *Clinical physiology of acid-base and electrolyte disorders.* 5<sup>th</sup> ed. New York: McGraw Hill. 2001.
7. Adrogué HJ, Madias NE. Secondary responses to altered acid-base status: the rules of engagement. *J Am Soc Nephrol.* 2010; **21**(6): 920-3.
8. Marshall WJ, Bangert SK, Lapsley ML. Hydrogen ion homeostasis and blood gases. Chapter 3. In: Marshall WJ, Bangert SK, Lapsley ML, editors. *Clinical chemistry 7<sup>th</sup> ed.* Edinburgh: Mosby; 2012. p. 41-62.
9. Badrick T, Hickman PE. The anion gap: a reappraisal. *Am J Clin Pathol.* 1992; **98**(2): 249-52.
10. Kraut JA, Madias NE. Serum anion gap: its uses and limitations in clinical medicine. *Clin J Am Soc Nephrol.* 2007; **2**(1): 162-74.
11. Mehta AN, Emmett JB, Emmett M. GOLD MARK: an anion gap mnemonic for the 21<sup>st</sup> century. *Lancet.* 2008; **372**(9642): 892.
12. Kraut JA, Madias NE. Lactic acidosis: current treatments and future directions. *Am J Kidney Dis.* 2016; **68**(3): 473-82
13. Fenves AZ, Kirkpatrick HM 3rd, Patel VV, Sweetman L, Emmett M. Increased anion gap metabolic acidosis as a result of 5-oxoproline (pyroglutamic acid): a role for acetaminophen. *Clin J Am Soc Nephrol.* 2006; **1**(3): 441-7
14. Choy KW, Wijeratne N, Lu ZX, Doery JC. Harmonisation of Osmolal Gap - Can We Use a Common Formula? *Clin Biochem Rev.* 2016; **37**(3): 113-9.
15. Kowlgi NG, Chhabra L. D-lactic acidosis: an underrecognized complication of short bowel syndrome. *Gastroenterol Res Pract.* 2015; **2015**: 476215

