

# A new subspecies of *Seseli gummiferum* (Apiaceae) from Ilgaz Mountain National Park, northern Turkey

Özlem Çetin<sup>1</sup>, Meryem Öztürk Şeker<sup>2</sup>, Ahmet Duran<sup>2</sup>

**1** Department of Biotechnology, Selçuk University, Selçuklu, Konya, Turkey **2** Department of Biology, Selçuk University, Konya, Turkey

Corresponding author: Özlem Çetin (ozlemcetin8419@gmail.com)

Academic editor: Peter de Lange | Received 28 July 2014 | Accepted 4 September 2015 | Published 6 October 2015

**Citation:** Çetin Ö, Şeker MÖ, Duran A (2015) A new subspecies of *Seseli gummiferum* (Apiaceae) from Ilgaz Mountain National Park, northern Turkey. *PhytoKeys* 56: 99–110. doi: 10.3897/phytokeys.56.5755

## Abstract

A new subspecies *Seseli gummiferum* Pall. ex Sm. subsp. *ilgazense* A.Duran, Ö.Çetin & M.Öztürk, **subsp. nov.** (Apiaceae) is described from Kastamonu province, Turkey. It was collected from the open *Pinus sylvestris* L. and *Abies nordmanniana* (Steven) É.Spach. mixed forest in the northern Anatolian region. An endemic apparently confined to the Ilgaz Mountain National Park, the new taxon is closely related to *Seseli gummiferum* subsp. *gummiferum*. Diagnostic morphological characters for closely similar taxa are discussed, and a key to the subspecies of *Seseli gummiferum* is presented. ITS (Internal Transcribed Spacer) region of the nuclear ribosomal DNA of closely related *Seseli* L. taxa and *Pimpinella* is used to construct phylogenetic tree by using BioEdit and Seaview Programme.

## Keywords

New taxon, *Seseli gummiferum*, Turkey, Umbelliferae

## Introduction

The Apiaceae comprise approximately 450 genera and 3700 species worldwide (Pimenov and Leonov 1993). However, the distribution of species among the genera vary, with almost half of the genera monotypic and 26% consisting of only two or three species. Sixty percent of the species in the family assign to just a few genera, which genera encompass over 20 species have been noted as polyphyletic (Spalik et al. 2004). Asian countries with the greatest biodiversity for the Apiaceae include China,

Asian Turkey, Iran, Asian Russia, and Kazakhstan, with the most species given for the Chinese flora (677 species in 108 genera) (Pimenov and Leonov 2004). Turkey, with a considerably smaller geographic area, is second only to China in its diversity for the Apiaceae, with 450 species in 109 genera. There are four endemic genera in Turkey, with 140 species among 42 genera. This suggests that the Asiatic region in Turkey has the highest known species-level diversity for the Apiaceae in Asia, if not in the world (Pimenov and Leonov 2004).

*Seseli* L. is one of the largest genera in the Apiaceae with 125 to 140 taxa ascribed to the genus worldwide. Represented by both intraspecific and interspecific diversity, *Seseli* is distributed in Europe, Asia, Africa, North America and Australia (Pimenov and Leonov 2004). The first revision of *Seseli* in Turkey was made by Hedge and Lamond (1972), who recognized 10 infrageneric taxa. *Seseli gummiferum* subsp. *gummiferum* was reported as a new record from Turkey (Duman 2000). After that *Seseli ramosissimum* replaced with *Seseli hartvigii* by Parolly (Parolly and Nordt 2001). Latest revision of *Seseli* in Turkey was made by Doğan Güner and Duman (2013). So the total number of these taxa is 13 now.

## Material and methods

In 2008, during a field trip in the Black Sea region of Turkey, an unusual specimen of *Seseli* was collected by the authors. Study of the descriptions in Hedge and Lamond (1972), Davis et al. (1988), Duman (2000), Ball (1968), Rechinger (1987), Parolly and Nordt (2001), Shishkin (1950), Özhatay et al. (2009), Doğan Güner and Duman (2013) as well as comparison with herbarium material in GAZI, HUB, KNYA and ANK revealed that the specimens indeed represented a new taxon. In particular, the new taxon was compared with the closely similar taxa *Seseli gummiferum* Pall. ex Sm. subsp. *gummiferum* and *S. corymbosum* Boiss. & Heldr. In the morphological description below, each numerical value is the average of ten measurements from different specimens. The abbreviations of the authors of plant names were checked from Brummitt and Powell (1992).

**DNA isolation:** Total DNA was obtained from 50–75 mg leaf and fruit from six different individuals. DNAs are isolated with CTAB method and after concentrations were determined by Nanodrop. Sample DNAs were diluted 25 ng/μl. Stock DNAs were kept at -86 °C.

**ITS amplifications:** ITS region of studied taxa were amplified using ITS4 (5' TCC TCC GCT TAT TGA TAT GC 3') and ITS5 (5' GGA AGG AGA AGT CGT AAC AAG 3') primers. PCR condition is 95 °C for 5 min initial denaturation, 35 cycles of 94 °C for 30 s denaturation, 50 °C for 30 s annealing, and 72 °C for 1 min extension, 72 °C for 10 min final extension.

**Data collection and cluster of phylogenetic analysis:** PCR products were visualised by agarose gel. The amplified fragments were sequenced using the same primers used for amplification. ITS sequences of the taxa were aligned via Bioedit and were used to construct phylogenetic trees by using Seaview.

## Taxonomic treatment

*Seseli gummiferum gummiferum* Pall. ex Sm., *Exot. Bot.* [Smith] ii, 121 (1807).

Replaced synonym: *Bubon rigidus* Spreng., *Syst. Veg.* (ed. 16) [Sprengel] 1: 900 1825 [1824], nom. illeg., non *Bubon rigidus* (Waldst. & Kit.) Spreng. *Pl. Min. Cogn. Pug.* 2: 53. 1815. Type: [London] Cult. in Oxford Bot. Garden and Hort. Lady Hume.

*Seseli gummiferum crithmifolium* (Boiss.) P.H.Davis, *Notes Roy Bot. Gard. Edinb.* 21: 120 (1953).

Basionym: *Seseli crithmifolium* Boiss., *Fl. Orient.* [Boissier] 2: 962 (1872). Type: Greece, Insulae maris Aegaei, *Tournefort* 324 (holotype P, photo!, E!).

*Seseli gummiferum ilgazense* A.Duran, Ö.Çetin & M.Öztürk, *subsp. nov.*

urn:lsid:ipni.org:names:77150275-1

Figs 1–6

*Affinis* *S. gummiferum* Pall. ex Sm. *subsp. gummiferum* sed *umbellis centralibus radiis* 13–21 (*nec* 25–30), *bracteis* 2–7 (*nec plerumque* 8–15), *umbellis lateralibus radiis* 7–13 (*nec* 15–20), *ovariis glabris (nec cum pilis)*, *fructibus glabris (nec cum pilis)* differt.


**Type.** TURKEY. A4 Kastamonu: Ilgaz Mountain Natural Park, Kastamonu road, from Çatören village to Büyük Hacet Hill, 6 km, in open *Pinus sylvestris* L. and *Abies nordmanniana* (Stev.) Spach. mixed forest, serpentine stony slopes, 41°06'344"N, 33°48'628"E, 1465 m, 22 August 2008, A.Duran 8135, Ö.Çetin & M.Öztürk (holotype KNYA! isotypes ANK!, GAZI!, HUB!).

**Description.** Plants monocarpic, 15–30 cm tall; rootstock thickened, cylindrical, oblong, ± vertical, 8–15 mm diameter; stems 3 to 5, terete, finely ridged, mostly glabrous, sparsely puberulent above, with a developed fibrous collar 2.5–7 cm; stems mostly branching from the base, below and rarely at the middle part, green to purplish green. Basal leaves ovate to oblong-ovate, 2-pinnate, 6–20 × 3–10 cm, glabrous; ultimate segments lanceolate, linear-oblong, 5–15 × 0.5–1.5 mm, acute to acuminate; sheaths developed, distinctly widened at base, upper sheath surface sulcate, margin membranous; cauline leaves similar to basal leaves, partly reduced, amplexicaule, middle and upper portions of the stem leafless. Flowers hermaphroditic; the central umbel stout, 7–11 cm diam., equal to or longer than lateral umbels, rarely shorter, with 13 to 21 rays, 1.8–5.5 cm long, puberulent above, unequal, each central umbel with by 2 to 7 bracts; bracts lanceolate, rarely widened at base, 10–17 mm long, margin ± membranous, sometimes bifid to trifid, glabrous or puberulent; umbellules 9–15 mm diam, each comprising



**Figure 1.** *Seseli gummiferum* subsp. *ilgazense* A.Duran, Ö.Çetin & M.Öztürk, subsp. nov.






**Figure 2.** Cross section of fruit: emb = embriyo; dvb = dorsal vascular bundle; ex = exocarp; cdv = central dorsal vittae; es = endosperm; lvb = lateral vascular bundle; ldv = lateral dorsal vittae; ae = aerenchyma; cv = commissural vittae; cvb = carphophore vascular bundle; covb = commissural vascular bundle. Scale bar = 0.2 mm.

55 to 70 flowers; bracteoles 15–20, connate at base, 5–8 mm long, linear-lanceolate, acuminate, entire or bifid to trifid, green to partly purple, margin distinctly membranous, especially ciliate at apex, out side glabrous to puberulent; lateral umbels 3.5–6 cm diam. with 7 to 13 rays, 1–3.5 cm long, with bracts or without. Flowers subsesile, glabrous, sepals ±purplish, broadly lanceolate, glabrous, persistent, ca. 1 mm; petals white, glabrous with deflexed apex; filament white, anther clearly purple; ovary glabrous. Fruit oblong-elliptic, 3.5–4 × 1.5–2 mm, glabrous, with 5 ribs prominent, obtuse; stylopodium short-conical; styles deflexed in fruit, distinctly purple.


**Distribution and IUCN red list category.** The new subspecies is known only from the type locality, with specimens collected only from Ilgaz Mountain National Park (Kastamonu province) in Turkey, where the species seems to be very rare. This area is ca. 1.5 km<sup>2</sup>, and mature individuals of the type population number approximately 125. The location is very close to the road side and near forest management. The population is going to be negatively affected from cars, trucks and people in the future. The habitat of this subspecies is clearly under threat of destruction, and therefore, the taxon should be considered Critically Endangered (CR), according to IUCN Red List Criteria (IUCN 2001).

**Habitat and ecology.** This new subspecies grows at 1450–1470 m with *Lapsana communis* L., *Abies nordmanniana* (Stev.) Spaach, *Centaurea drabifolia* Sm., *Erysimum thyrsoideum* Boiss., *Pteridium aquilinum* (L.) Kuhn, *Sorbus umbellata* (Desf.) Fritsch, *Valeriana alliariifolia* Adams, *Eryngium giganteum* M.Bieb., *Bupleurum falcatum* L. subsp. *persicum* (Boiss.) Koso-Pol., *Salvia verticillata* L., *Teucrium chamaedrys* L., *Dactylis glomerata* L., *Asyneuma rigidum* (Willd.) Grossh. subsp. *rigidum*.




**Figure 3.** *Seseli gummiferum* subsp. *ilgazense*. **A** trifid and entire bracteoles **B** lateral umbel **C** general aspects **D** habitat and general view of type locality. Photo by A.Duran.






**Figure 4.** Comparison of *Seseli gummiferum* fruits. *S. corymbosum* fruits. **A** general view **B** surface details. *S. corymbosum*: **C** general view **D** surface details. *S. gummiferum* subsp. *ilgazense*: **E** general view **F** surface details.

**Discussion and conclusion.** Turkey is the most complex country in the Middle East with regard to geographic structure and landforms. It's comprised of comparatively narrow and long, variously oriented mountain chains, separated by deep valleys and also high- and medium-elevational plateaus. The geological composition and physical direction, exposure and altitude of these mountains are here largely influential not only upon the diversity of vegetation, but also on the richness of the flora (Zohary 1973). Ilgaz Mountain, which is situated in a transitional zone in central and North of Anatolia and is generally composed of serpentine, schist and volcanic rocks. The mountain is orogenically interesting, with the quite active north Anatolian fault found along the southern slopes of Ilgaz Mountain (Kuter 2008). One of the more important reasons for protecting the Ilgaz Mountain is the richness and endemism of its flora. Approximately 100 endemic plants occur within the boundaries of the National Park and the type localities of 19 endemic taxa are found on Ilgaz Mountain. *Delphinium ilgazense*



**Figure 5.** Distribution of *Seseli gummiiferum* subsp. *ilgazense* (◆), *S. gummiiferum* subsp. *gummiiferum* (●) and *S. corymbosum* (■) in Turkey.



**Figure 6.** The neighbour joining tree generated using nrITS DNA sequences of some *Seseli* taxa and *Pimpinella saxifraga*.

P.H.Davis, *Arabis abietina* Bornmüller, *Draba anatolica* A.Duran & Dinç, *Astragalus nabelekii* Czechtz, *Heracleum paphlagonicum* Czechtz, *Hieracium macrogonum* (Zahn) P.D.Sell & C.West and *H. tuberculatum* Freyn, *Festuca ilgazensis* Markgr.-Dann. are all noteworthy endemic taxa confined to the national park (Davis 1965–1985, Davis et al. 1988, Duman 2000, Duran et al. 2008).

*Seseli gummiiferum* subsp. *ilgazense* is closely related to two other subspecies found in Turkey. *S. gummiiferum* subsp. *gummiiferum* is distributed in Crimea, Central Anatolia (Shishkin 1950, Ball 1968, Duman 2000). The new subspecies differs from *S. gummiiferum* subsp. *gummiiferum*, based on its glabrous fruits (not with indument), fewer central umbels, with 13 to 21 rays (not 25 to 30), fewer lateral umbels, with seven to 13 rays (not 15 to 20), as well as the bracteoles 15 to 20 (not 11 to 16) (Hedge and Lamond 1972, Duman 2000).

*Seseli gummiiferum* subsp. *ilgazense* also differs from *S. gummiiferum* subsp. *crithmifolium* (DC.) P.H.Davis, which is distributed in west and east Crete, Folegandros, Sikinos,



Amargos, Karpathos, Saria and neighbouring islands in Aegean Sea. The new taxon is not so broadly distributed, endemic and found only in northern Anatolia (Ilgaz Mountain). *Seseli gummiferum* subsp. *crithmifolium* principally differs from subsp. *ilgazense* by its puberulent fruits, the central umbels with 20–45 rays, and the oblong leaf segments (6–) 10–30 × 2–5 mm (Ball 1968, Hedge and Lamond 1972).

### Key to closely related *Seseli gummiferum* subspecies

- 1 Fruit glabrous; the central umbel 13 to 21 rays; lateral umbels with 7 to 13 rays ..... **subsp. *ilgazense***
- Fruit with hairs; the central umbel 22 to 70 rays; lateral umbels with 13 to 47 rays .....
- 2 Basal leaves glabrous ..... **subsp. *crithmifolium***
- Basal leaves puberulent or finely pubescent ..... **subsp. *gummiferum***

Since Morison's (1672), *Plantarum umbelliferarum*, fruit morphology and anatomy have been regarded as essential to the taxonomy of Apiaceae (Drude 1898, Constance 1971, Spalik et al. 2001). Details of the fruits have been traditionally viewed as rich sources of taxonomic characters, exhibiting some, but not excessive variation in features such as fruit shape, the degree and direction of mericarp compression, modifications of the pericarp ribs (e.g. wings or spines), and the shape of mericarp commissural faces. Thus, most traditional classifications of Apiaceae have relied almost exclusively on fruit characters (Plunkett and Downie 1999, Pimenov et al. 2004). *Seseli gummiferum* subsp. *ilgazense* has some distinctive characteristics in terms of carpological features. Mericarps of *Seseli gummiferum* subsp. *ilgazense* have five large vascular bundles situated beneath the each rib. Large vittae present in mesocarp layer adjacent to the endocarp. There are also two large vittae in the commissure. Differing from *Seseli gummiferum* subsp. *gummiferum*, the new subspecies does not have short secretory ducts around vascular bundles. In addition, the vittae around the endocarp are always large and elliptical. The endosperm is round shaped. Commissure width is approximately equal to mericarp width.

Diagnostic characters of *Seseli gummiferum* subsp. *ilgazense* with the three related taxa are provided in Table 1.

Five *Seseli* taxa and *Pimpinella* were evaluated in the phylogenetic analysis. PCR amplification with ITS 4/ITS 5 primers generated bands ranging from 595 to 665 bp. Alignment of the ITS sequences was done using Bioedit. Neighbour-Joining (NJ) tree were constructed using Seaview programme (Figure 6). According to NJ analyses, the taxa of *Pimpinella* placed in first clade. This species was used as outgroup. The second clade includes *Seseli libanotis*, which is found in Section *Libanotis*. *Seseli libanotis* is separated from *S. gummiferum* and *S. petraeum* by having highly wide leaf segment. The second clade includes *Seseli petraeum*, *S. corymbosum*, *S. gummiferum* in sect. *Seseli*. *Seseli petraeum* is close to *Seseli gummiferum*. It mainly differs from *S. gummiferum*

**Table 1.** Diagnostic characters of *Seseli gummiferum* subsp. *ilgazense*, *S. gummiferum* subsp. *gummiferum*, *S. gummiferum* subsp. *crithmifolium* and *S. corymbosum*.

Characters	subsp. <i>ilgazense</i>	subsp. <i>gummiferum</i>	subsp. <i>crithmifolium</i>	<i>S. corymbosum</i>
Stems per plant	3 to 5	1 to 4	solitary	solitary
Stem branching	mostly below, rarely in middle portion	below the middle	above the middle	above the middle
Central umbel rays	13 to 21	25 to 30	22 to 45	30 to 70
Lateral umbel rays	7 to 13	15 to 20	17 to 35	13 to 47
Bracts	2 to 7	mostly 8 to 15	mostly absent, rarely up to 2	0 to 1
Bracteoles	15 to 20	11 to 16	19 to 26	19 to 23
Petals	glabrous	glabrous	pubescent	pubescent
Ovary	glabrous	pubescent	puberulent	pubescent
Fruit	glabrous	pubescent	pubescent	pubescent

because it has free bracteol from base and main umbel shorter than lateral umbels. *Seseli corymbosum* (Boiss. & Heldr.) P.H.Davis which only occurs in South Anatolia, is readily distinguished from *S. gummiferum*, by its solitary stem branching above, bracts either lacking or one below the central umbel, more numerous bracteoles, 19 to 23, larger central umbels with 30 to 70 rays and the lateral umbels with 13 to 47 rays, and finally by the pubescent petals and fruits (Hedge and Lamond 1972). Although *Seseli gummiferum* subsp. *ilgazense* is closely related to *S. gummiferum* subsp. *gummiferum*, this taxa is treated as different subspecies.

## Acknowledgements

The authors express their thanks to the Scientific Investigation Projects Coordination Office of the Selçuk University (project no: 08201037) for financial support.

## References

- Ball PW (1968) *Seseli* L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) *Flora Europaea*. Vol 2, Cambridge University Press, Cambridge, 334–338.
- Brummitt RK, Powell CE (1992) *Authors of Plant Names*. Royal Botanic Gardens, Kew, 736 pp.
- Constance L (1971) History of the classification of Umbelliferae (Apiaceae). In: Heywood VH (Ed.) *The Biology and Chemistry of the Umbelliferae*. Academic Press, New York, 1–8.
- Davis PH (1965–1985) *Flora of Turkey and the East Aegean Islands*. Vols 1–9. Edinburgh University Press, Edinburgh, 567 pp.

- Davis PH, Mill RR, Tan T (1988) *Seseli* L. In: Davis PH, Mill RR, Tan T (Eds) Flora of Turkey and the East Aegean Islands – 11. Edinburgh University Press, Edinburgh, 150 pp.
- Doğan Güner E, Duman H (2013) The revision of genus *Seseli* (Umbelliferae) in Turkey. Turkish Journal of Botany 37: 1018–1037. doi: 10.3906/bot-1302-41
- Drude CGO (1898) Umbelliferae. In: Engler A, Prantl K (Eds) Die Natürlichen Pflanzenfamilien – Volume 3. Leipzig, 63–250.
- Duman H (2000) *Seseli* L. In: Güner A, Özhatay N, Ekim T, Başer KHC (Eds) Flora of Turkey and the East Aegean Islands (Suppl. 2). Edinburgh University Press, Edinburgh, 141 pp.
- Duran A, Dinç M, Martin E (2008) A new species of *Draba* (Brassicaceae) from North Anatolia, Turkey. Novon 18: 464–468. doi: 10.3417/2007023
- Hedge IC, Lamond JM (1972) *Seseli* L. In: Davis PH (Ed.) Flora of Turkey and the East Aegean Islands – Volume 4. Edinburgh University Press, Edinburgh, 367–372.
- IUCN (2001) IUCN red list categories, ver. 3.1. IUCN Species Survival Commission.
- Kuter N (2008) Ilgaz Dağı milli Parkının orman peyzajı ve estetiği açısından değerlendirilmesi. Süleyman Demirel Üniversitesi Orman Fakültesi Dergisi 1: 36–47.
- Morison R (1672) Plantarum umbelliferarum distributio nova. Oxford.
- Özhatay N, Kültür Ş, Aslan S (2009) Check-list of Additional Taxa to the Supplement Flora of Turkey, IV. Turkish Journal of Botany 33: 191–226.
- Parolly G, Nordt B (2001) *Seseli hartwigii* (Apiaceae), a new name for *S. ramosissimum* Hartvig & Strid., with carpological and ecological notes on this species. Willdenowia 31: 87–93. doi: 10.3372/wi.31.31107
- Pimenov MG, Valiejo-Roman CM, Terentieva EI, Samigullin TH, Mozaffarian V (2004) Enigmatic genus *Hausknechtia* (Umbelliferae): systematic relationships based on molecular and carpological data. Nordic Journal of Botany 24(5): 555–565. doi: 10.1111/j.1756-1051.2004.tb01640.x
- Pimenov MG, Leonov MV (2004) The Asian Umbelliferae biodiversity database (ASIUM) with particular reference to South-West Asian taxa. Turkish Journal of Botany 28: 139–145.
- Pimenov MG, Leonov MV (1993) *Seseli* L. The genera of the Umbelliferae: a nomenclature. Kew Royal Botanic Gardens, Kew.
- Plunkett GM, Downie SR (1999) Major lineages within Apiaceae subfamily Apioideae: A comparison of chloroplast restriction site and DNA sequence data. American Journal of Botany 86: 1014–1026. doi: 10.2307/2656619
- Rechinger KH (1987) Umbelliferae. Flora Iranica. Akademische Druck-u., Verlagsanstalt, Graz-Austria, 347–349.
- Shishkin BK (1950) *Seseli* L. In: Shishkin BK (Ed.) Flora of the USSR, Vol. 16, Izdatel'stvo Akademii Nauk SSSR, Moskva-Leningrad, 349–378. [translated from Russian, Israel Prog. Sci. Transl., 1973, Jerusalem]
- Spalik K, Wojewódzka A, Downie SR (2001) The evolution of fruit in Scandiceae subtribe Scandicinae (Apiaceae). Canadian Journal of Botany 79: 1358–1374. doi: 10.1139/cjb-79-11-1358
- Spalik K, Reduron J-P, Downie SR (2004) The phylogenetic position of *Peucedanum* sensu lato and allied genera and their placement in tribe Selineae (Apiaceae, subfamily Apioideae). Plant Systematics and Evolution 243: 189–210. doi: 10.1007/s00606-003-0066-2
- Zohary M (1973) Geobotanical foundations of the Middle East, Vol. 1. Gustav Fischer, City.



## Appendix

**Representative specimens examined** (\*which used for DNA samples): – *Seseli corymbosum*: Turkey, C3 Antalya: Akseki, Pınarbası village, S of Gidefi Mountain, *A.Duran* 2970 (GAZI); Akseki, Gidefi Mountain, *A.Duran* 1847 (GAZI); C5 Niğde; Ulukışla, between Alihoca-Maden villages, *A.Duran* 6078\* (KNYA). – *S. gummiferum* subsp. *gummiferum*: Turkey, A4 Ankara: Hasanoğlu, İdris Mountain, above Adilahmet village, *M.Koyuncu* 16348 (GAZI); Hasanoğlu, İdris Mountain, 1500–1600 m., 14.09.2002, *E.Doğan* 1650\* (GAZI). – *S. gummiferum* subsp. *critmifolium*: Greece, Dodecanese, Karpathos, side of Kali Limni, 10 m, 21.07.1950, *Davis* 18010 (E). – *Seseli petraeum*: Turkey, Trabzon, between Araklı-Bayburt, 27.07.2002, *A.Duran* 6059\* (KNYA). – *Seseli libanotis*: Turkey, Erzurum, between İspir-Ovit, 1700 m, 18.08.2013, *A.Duran* 9778\* (KNYA).