

---

# MYCOTAXON

<http://dx.doi.org/10.5248/119.439>

Volume 119, pp. 439–443

January–March 2012

---

## *Herpothallon weii*, a new lichen from China

YU-LIANG CHENG<sup>1</sup>, JING NING<sup>1</sup>, HOU-PING XU<sup>2</sup>, LU-LU ZHANG<sup>1</sup>,  
HAI-YING WANG<sup>1A\*</sup> & ZUN-TIAN ZHAO<sup>1B\*</sup>

<sup>1</sup>College of Life Sciences, Shandong Normal University, Jinan, 250014, P. R. China

<sup>2</sup>Jinan Institute of Metrological Verification, Jinan, 250002, P. R. China

\* CORRESPONDENCE TO: <sup>a</sup>lichenwhy@yahoo.com.cn, <sup>b</sup>ztzhao@sohu.com

**ABSTRACT** — A new species, *Herpothallon weii*, is described from China. This species is characterized by a byssoid whitish green thallus tightly attached to the substratum, pinkish pseudisidia, and the presence of psoromic acid and unknown substances.

**KEY WORDS** — biodiversity, *Arthoniaceae*, *Cryptothecia*, taxonomy

### Introduction

*Herpothallon* (*Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, *Ascomycota*) was established by Tobler in 1937 and reinstated by Aptroot et al. (2009). The lichen genus is characterized by a crustose thallus without upper cortex, a byssoid prothallus, and a trentepohlioid photobiont. *Herpothallon* species often have pseudisidia or pseudisidioid structures, calcium oxalate crystals, pigments, and various lichen substances, but usually lack ascocarps. *Cryptothecia* species, which are morphologically similar to *Herpothallon* species, have an I+ blue medulla. Of *Herpothallon*, only six species (*H. biacidum*, *H. flavominutum*, *H. granulosum*, *H. isidiatum*, *H. philippinum*, and *H. rubroechinatum*) have an I+ blue medulla (Aptroot et al. 2009, Frisch et al. 2010a,b, Jagadeesh & Sinha 2009, Jagadeesh et al. 2009).

*Herpothallon* includes 35 species worldwide (Aptroot et al. 2009, Frisch et al. 2010a,b, Jagadeesh & Sinha 2009, Jagadeesh et al. 2009). In China, three *Herpothallon* species, *H. echinatum*, *H. granulare*, and *H. philippinum* have been reported (Aptroot et al. 2009). During our study of the lichen flora of Mt. Leigong in Guizhou province, China, a species of *Herpothallon* new to science was found.

### Materials & methods

The specimens studied were collected from Guizhou, China, and are preserved in SDNU (Lichen Section of Botanical Herbarium, Shandong Normal University).

The morphological and anatomical characters of the specimens were examined using a stereo microscope (COIC XTL7045B2) and a polarizing microscope (OLYMPUS CX41). Lichen substances in the specimen cited were identified using the standardized thin layer chromatography techniques (TLC) with C system (Orange et al. 2010). Photos of the thalli were taken using a Olympus SZX12 with DP72.

## Taxonomic description

*Herpothallon weii* Y.L. Cheng & H.Y. Wang, sp. nov.

FIGS 1–2

MYCOBANK MB 563539

Differs from *Herpothallon albidum* by pinkish pseudisidia and lack of pigmentosin D.

TYPE: China. Guizhou province, Mt. Leigong, on bark, alt. 1000 m, 2 Nov. 2009, Q. Tian 20103366. (Holotype in SDNU).

ETYMOLOGY: In honour of the 80<sup>th</sup> birthday (6 Nov. 2011) of our dear teacher, Prof. Jiang-Chun Wei, who as the ‘father of lichenology in China’ has made a preeminent contribution to the development of lichenology and the education of Chinese talent in this field.

Thallus corticolous, crustose, tightly attached,  $\leq 1.5$  cm in diam., without upper cortex, distinctly byssoid, whitish green, with irregular calcium oxalate crystals. Prothallus distinct,  $\leq 0.5$  mm wide, white, byssoid, composed of radiating hyphae. Pseudisidia (also called pseudisidioid structure) numerous, pinkish, irregularly cushion-shaped, fluffy-felty with erect hyphae,  $\leq 1 \times 0.5$  mm. Photobiont cells single or a few cells aggregated. Apothecia and pycnidia not seen.

CHEMISTRY — Thallus K<sup>+</sup> yellow, C $\pm$  yellow, KC<sup>+</sup> yellow, P<sup>+</sup> yellow, I<sup>-</sup>, UV<sup>-</sup>. Psoromic acid and unknown substances present.

DISTRIBUTION AND SUBSTRATE — *Herpothallon weii* is known only from the type locality; on bark.

ADDITIONAL SPECIMEN EXAMINED — CHINA. GUIZHOU PROVINCE, Mt. Leigong, on bark, alt. 1000 m, 2 Nov. 2009, Q. Tian, 20103174 (SDNU).

COMMENTS — Other *Herpothallon* species with pseudisidia and psoromic acid include *H. australasicum*, *H. echinatum*, *H. flavominutum*, *H. globosum*, *H. rubroechinatum*, and *H. albidum*. In contrast to *H. weii*, *H. australasicum* has a red thallus with concolorous compact pseudisidia; *H. echinatum* has soredioid fragments and concolourous pseudisidia and thallus; *H. flavominutum* has a greyish yellow thallus, granular pseudisidia, and lichexanthone, norlichexanthone, and psoromic acid; *H. globosum* has a red thallus, globose pseudisidia, and chiodectonic acid (red pigment); and *H. rubroechinatum* has an orange to red thallus, pseudisidia with red hairs, and common pycnidia. The more closely related *H. albidum* possesses very similar pseudisidia (also called pseudisidioid structures) but has white, rather than pinkish, pseudisidia and possesses pigmentosin D.




FIG. 1. *Herpothallon weii* (holotype). A. Thallus; B. Prothallus.

#### Acknowledgements

This research was supported by the City Science Foundation of Guiyang (20101Zhong05), the National Natural Science Foundation of China (31070010, 31000008), and Natural Science Foundation of Shandong Province (2R2010CQ038). We thank Prof. A. Aptroot (ABL Herbarium, Soest, The Netherlands) and Prof. Shou-Yu Guo (Institute of Microbiology, Chinese Academy of Sciences, Beijing, China) for presubmission reviews.




FIG. 2. *Herpotherallon weii* (holotype).

A. Pseudisidia; B. TLC of *H. weii* (U: unknown substance; P: psoromic acid).

**Literature cited**

- Aptroot A, Thor G, Lücking R, Elix JA, Chaves JL. 2009. The lichen genus *Herpothallon* reinstated. *Bibliotheca Lichenologica* 99: 19–66.
- Frisch A, Elix JA, Thor G. 2010a. *Herpothallon biacidum*, a new lichen species from tropical Australia. *Lichenologist* 42(3): 285–289. <http://dx.doi.org/10.1017/S0024282909990697>
- Frisch A, Thor G, Elix J. 2010b. *Herpothallon rubroechinatum* (Arthoniaceae), a new species from tropical and subtropical America. *Bryologist* 113(1): 144–148. <http://dx.doi.org/10.1639/0007-2745-113.1.144>
- Jagadeesh Ram TAM, Sinha GP. 2009. New species and new records of *Herpothallon* (lichenized *Ascomycota*) from India. *Mycotaxon* 110: 37–42. <http://dx.doi.org/10.5248/110.37>
- Jagadeesh Ram TAM, Sinha GP, Singh KP. 2009. New species and new records of *Cryptothecia* and *Herpothallon* (Arthoniales) from India. *Lichenologist* 41(6): 605–613. <http://dx.doi.org/10.1017/S0024282909008123>
- Orange A, James PW, White FJ. 2010. *Microchemical methods for the identification of lichens*. 2nd edition. London: British Lichen Society.