

# A new species of *Cyanea* (Campanulaceae, Lobelioideae), from the Ko‘olau Mountains of O‘ahu, Hawaiian Islands

Margaret J. Sporck-Koehler<sup>1,2</sup>, Tobias B. Koehler<sup>4</sup>, Sebastian N. Marquez<sup>3</sup>,  
Mashuri Waite<sup>5</sup>, Adam M. Williams<sup>1</sup>

**1** *Hawai‘i Department of Land and Natural Resources, Division of Forestry and Wildlife, 1151 Punchbowl St. Rm. 325, Honolulu, HI 96813 USA* **2** *Department of Botany, University of Hawai‘i at Mānoa, 3190 Maile Way, Honolulu, HI 96822, USA* **3** *Papahana Kuaola 46-403 Haiku Road, Kaneohe, Hawai‘i 96744 USA* **4** *Koehler Enterprises LLC., Honolulu, Hawai‘i 96813 USA* **5** *Ko‘olau Mountains Watershed Partnership, 2551 Waimano Home Rd, Pearl City, HI 96782, USA*

Corresponding author: Margaret J. Sporck-Koehler (margaret.j.sporck-koehler@hawaii.gov)

Academic editor: Vicki Funk | Received 6 October 2014 | Accepted 22 December 2014 | Published 5 February 2015

**Citation:** Sporck-Koehler MJ, Koehler TB, Marquez SN, Waite M, Williams AM (2015) A new species of *Cyanea* (Campanulaceae, Lobelioideae), from the Ko‘olau Mountains of O‘ahu, Hawaiian Islands. *PhytoKeys* 46: 45–60. doi: 10.3897/phytokeys.46.8694

## Abstract

*Cyanea konahuanuiensis* Sporck-Koehler, M. Waite, A.M. Williams, **sp. nov.**, a recently documented, narrowly endemic species from the Hawaiian Island of O‘ahu, is described and illustrated with photographs from the field. The closest likely relatives to the species, current conservation needs, and management future are discussed. It is currently known from 20 mature plants from two subpopulations and is restricted to a drainage below the Kōnāhua-nui summit (K1), the highest summit of the Ko‘olau Mountains, located on Windward O‘ahu. It differs from all other *Cyanea* species by its combination of densely pubescent leaves, petioles, and flowers; sparsely pubescent to glabrous stems, long calyx lobes, and staminal column being adnate to the corolla.

## Keywords

Campanulaceae, conservation, *Cyanea*, endemic, Hawai‘i, Hawaiian Islands, IUCN Red List, Kōnāhua-nui, Ko‘olau Mountains, Lobeliads, Lobelioideae, O‘ahu, rare plants, *Rollandia*


## Introduction

The Campanulaceae is a large, diverse, and cosmopolitan plant family with representatives inhabiting a wide range of ecosystems including tropical, subtropical, temperate, and even frigid zones with exceptional diversification in South Africa and Hawai'i (Lammers 2007). One of the most exceptional adaptive radiation events known in the family is the monophyletic lobelioid group in Hawai'i (Givnish et al. 2008). The Hawaiian lobelioids are a group of woody eudicots comprised of six genera, *Brighamia* A. Gray, *Clermontia* Gaudich., *Cyanea* Gaudich., *Delissea* Gaudich., *Lobelia* L., and *Trematolobelia* A. Zahlbr. Together they account for roughly ten percent of Hawaiian angiosperm diversity (Wagner et al. 1999, 2005) with a total of 128 taxa, 78 of which are currently recognized as Threatened and Endangered (T&E; USFWS <http://www.fws.gov/endangered/>). The lobelioids in Hawai'i represent the largest adaptive radiation from a single colonization event known from any plant group restricted to an oceanic island chain (Oppenheimer and Lorence 2012, Givnish et al. 2008). According to the most current treatment of the Hawaiian Lobelioids (Lammers 2007), *Cyanea* (including the merger of the genus *Rollandia*) is the most species-rich genus in the radiation, comprised of 79 currently recognized species (54 T&E). A possible explanation for the impressive speciation in this genus is that the fleshy fruits are poorly distributed by Hawai'i's native forest birds, which do not typically travel long distances, leading to parallel speciation events on multiple islands (Givnish 2008). *Cyanea* occurs in mesic to wet forests across the Hawaiian archipelago and includes many taxa with restricted distributions, most of which are single island endemics.

In September 2012, the Kōnāhua-nui summit area of the Ko'olau Mountains on the island of O'ahu was surveyed (Figure 1). The target species was *Cyanea humboldtiana* (Gaudich.) Lammers, Givnish & Sytsma, a species federally listed as Endangered, and endemic to the Ko'olau Mountains. It was hoped to locate additional individuals to monitor and manage as a part of species recovery efforts. We summited K1 (the highest of the two peaks of Kōnāhua-nui) and descended into a stream drainage and once near the bottom of the gulch, directly adjacent to the stream, several plants were discovered of a *Cyanea* with hairy leaves and petioles, glabrous stems, and long, hairy calyx lobes. There were no flowers present on the plants and only one immature infructescence. The specific taxon could not be confirmed in the field and it was decided to take photos and collect a dropped and decaying leaf for further investigation.

Preliminary research in the following weeks consisted of examining the plants using photos and the one fallen, partially decayed leaf collected from the field. After reviewing the current treatments of *Cyanea*, sharing the photos with local experts, and looking at specimens at the Bishop Museum, it was concluded that the species in question was most likely undescribed. To be certain, fertile specimen was required.

Thereafter, trips to the remote population were made every two to three months. These subsequent expeditions surveyed the surrounding area and revealed the presence of additional individuals. In June 2013, an automated game camera was installed




**Figure 1.** Distribution map of *Cyanea konahuanuiensis* showing site of population on South East O'ahu in the Ko'olau Mountain Range.

connected to cellular phone service that transmitted a photo of one of the plants three times per 24 hour period (Figure 2). This camera allowed us to monitor the flower development and visit the plants again when we were certain the flowers would be fully mature.

In addition, type specimens of all of the Hawaiian Lobeliads were examined in the Herbarium Pacificum of the Bernice Pauahi Bishop Museum (BISH) collection, with special attention given to *Clermontia*, *Cyanea*, *Delissea*, and the formerly recognized generic group, *Rollandia* Gaudich. None of the Hawaiian Campanulaceae was a match to the *Cyanea* under study.

There are several species that share some characteristics with *C. konahuanuiensis* including: *Cyanea crispa* Gaudich., *Cyanea calycina* Gaudich., *Cyanea humboldtiana* Gaudich., and *Cyanea pilosa* A. Gray (Table 1). For example, *C. crispa*, *C. calycina*, and *C. humboldtiana* all share a staminal column that is adnate to the corolla tube, similar overall stature and leaf shape, presence of some leaf pubescence (though not to the same degree as *C. konahuanuiensis*), and similar number of flowers in each inflorescence (*C. konahuanuiensis* also shares the characteristic of pendant inflorescences but to a lesser degree than *C. humboldtiana*). However, they all differ in the degree of leaf pubescence (*C. konahuanuiensis* being significantly more hairy), calyx lobe length, and corolla surface characteristics.



**Figure 2.** *Cyanea konahuanuiensis*. Game camera time lapse series illustrating flower and early fruit development over a nearly two month time frame. Panels **A–D** each illustrate a different stage of development. Red arrows point to developing flower and fruit features.

The Hawai'i Island endemic *C. pilosa* shares a comparable leaf shape and is similarly pubescent (though dissection scope inspection revealed that hair structure differs) to *C. konahuanuiensis*, but the latter lacks hairy stems and has more abundant hair on both leaf surfaces. Also, flower color and calyx lobe length differ radically, *C. pilosa* usually having white flowers and much shorter calyx lobes (2–5 mm long) and *C. konahuanuiensis* having brilliant purple flowers. Lastly, unlike *C. konahuanuiensis*, the staminal column of *C. pilosa* is not adnate to the corolla tube (Figures 3, 4, 5, 6, 7, 8 and Table 1).

Because so few individuals of *C. konahuanuiensis* are known, collections were limited to two fertile vouchers, each including one inflorescence and two leaves. Flowers were also collected from three individual plants and preserved in alcohol for precise floral analysis (see Specimens Examined section). In order to ensure limited damage to the population and to further conserve the species, detailed measurements were taken of five, reproductively mature and flowering, live plants in the wild for most of the descriptive information in this paper. For corolla tube lobe length and width, surface characteristics, and anther length and surface characteristic data from the preserved flowers were utilized. Seed data are taken from living material deposited at the University of Hawai'i Harold L. Lyon Arboretum micropropagation facility. Selected measurements from the dried holotype specimen are included parenthetically in the description.

**Table 1.** Selected traits for *Cyanea konahuaniensis* compared to four *Cyanea* taxa sharing some similar morphological characteristics. Abbreviations: O = O‘ahu, H = Hawai‘i Island.

Species	Plant height (m)	Leaf length (cm)	Leaf width (cm)	Leaf shape	Adaxial leaf surface char.	Abaxial leaf surface char.	Peduncle Length (mm)	Number of flowers in inflorescence	Calyx lobe shape	Calyx lobe length (mm)	Hypanthium length (mm)	Flower Color	Corolla surface char.	Island found on
<i>C. konahuaniensis</i>	0.57–0.69	20–33	10–16	elliptic–oblong	densely hirsute, juveniles subdeltate muriccate	densely hirsute, juveniles subdeltate muriccate	50–122	3–12	linear to linear-oblong	16–18	10–15	dark magenta/purple w/ some lighter streaking as flowers age	densely pubescent	O
<i>C. calycina</i>	1–3	15–60	5.5–14	elliptic–oblanccolate	glabrous, juveniles muriccate	densely pubescent w/ branched & clustered hairs	20–100	4–16	oblong-ovate	4–10	6–12	pale–dark magenta, rarely pale greenish w/ lighter or darker longitudinal stripes	pubescent	O
<i>C. crispa</i>	0.3–1.3	30–75	9–16	broadly obovate	glabrous	glabrous or pubescent	20–30	3–8	ovate–oblong	6–12	8–12	pale magenta w/ darker longitudinal stripes	pubescent	O
<i>C. humboldtiana</i>	1–2	18–45	7–16	obovate–broadly elliptic	glabrous	pubescent	80–250	5–12	oblong	4–10	8–12	dark magenta, rarely white	pubescent	O
<i>C. pilosa</i>	0.8–2	15–42	8–15	broadly elliptic–broadly obovate	pubescent	whitish green, densely pubescent	15–110	6–28	narrowly triangular	2–5	4–6	white	pubescent	H

## Systematics

### *Cyanea konahuanuiensis* Sporck-Koehler, M. Waite & A.M. Williams, sp. nov.


urn:lsid:ipni.org:names:77145079-1

Figures 3–7

**Note.** Species believed to be allied to *Cyanea humboldtiana* (Gaudich.) Lammers, Givnish & Sytsma, but primarily differs in its longer calyx lobes (16–18 mm long); and its more densely pubescent leaves, petioles, flowers, and differing flowering period. *C. humboldtiana* has leaves, petioles, and flowers that are more sparsely pubescent with shorter trichomes; leaf margins are callose-crenulate, floral bracts have acute apices, and its calyx lobes are ovate, acuminate and considerably shorter (4–10 mm long). The two species are reproductively isolated from each other due to lack of overlapping flowering periods (*C. konahuanuiensis* flowers from June–August and *C. humboldtiana* from October–December), and they are not known to intergrade.

**Type.** USA, Hawai'i, O'ahu Island, Ko'olau Range: Kōnāhua-nui, near summit, 912m (2991 ft), 9 July 2013, M. Sporck, T. Koehler, & M. Waite MJS 0019 (holotype: BISH 1049136), (Figure 9).

**Description.** *Unarmed shrubs* 57–69 cm high, with 1–6 stems originating at the base; stems light green and darkening to a light brownish gray closer to the base, erect to decumbent, 58–119 cm in length, some branches partially resting on the ground, occasionally rooting when in contact with the soil or moss producing aerial roots, leaf scars subcircular, 9.5–12 × 6.5–11.8 mm, upper end of leaf scar depressed, basal portion slightly raised; leaf scar with a protuberance; latex white. *Leaves* clustered distally near end of stems, petiolate; petioles 2–4.2 cm long, pubescent; blades elliptic to oblong, in adult plants 20–33 × 10–16 cm, base cuneate to rounded, occasionally slightly truncate, apex acute to sub-obtuse, margins serrate to serrate-dentate, dull grayish-green on adaxial surface and dull greenish white on abaxial surface, stiff, slightly fleshy, both surfaces densely hirsute and minutely muricate; in juvenile plants leaves are less stiff, margins dentate, and hairs softer. *Inflorescences* axillary just above the leaf, up to 4 per stem, young inflorescences roughly perpendicular to stem, larger and more developed inflorescences pendant, 3–12 flowered, peduncles 5–12.2 cm long (dried specimen 7.7 cm long), pubescent. *Flowers* on pedicels 7–14 mm long (5–10 mm when dried), pubescent, subtended by linear bracts 7–18 × 2–6 mm, apex obtuse, margins entire, densely pubescent; hypanthium 10–15 mm (9–14 mm when dried) × 7–10 mm (5–7 mm when dried), obovoid to cylindrical, pubescent; calyx lobes linear to linear-oblong, 16–18 × 5–7 mm (13–17 × 3–5 mm when dried), apex acute to subobtuse, retained on immature (green) fruits (no mature fruit seen); corolla dark purple with some lighter streaks developing with age, tubular, laterally compressed, curved, 86–99 mm long (80–95 mm when dried) × 12–13 mm wide medially (9–12 mm when dried), externally densely pubescent, internally glabrous, the lobes linear-subulate, 10–16 mm long × 5 mm wide at the base, reflexed, c. 1/4–1/3 as long as the tube; staminal column glabrous, adnate to corolla for half its




**Figure 3.** *Cyanea konabuanuiensis*. **A** habit, photo with Adam M. Williams crouched down next to plant for scale **B** Seedlings growing in test tube in micropropagation lab **C** juvenile plant in the field **D** flowering stem with pendent inflorescence captured (photos by Margaret J. Sporck-Koehler).

length, anthers 9–10 mm long, scantily pubescent, the lower two with apical tufts of white hairs 2–3 mm long. *Fruits* berries (mature fruits not seen), immature fruits densely pubescent, with calyx lobes persistent. *Seeds* from immature fruits numerous, embedded in green pulp, obovoid,  $0.74\text{--}0.84 \times 0.58\text{--}0.64$  mm, testa medium to dark brown, shiny and smooth.

**Distribution.** Known only from the Kōnāhua-nui summit area in the Ko‘olau Mountains of O‘ahu, Hawaiian Islands. The population is on land owned by the State of Hawai‘i, Department of Land and Natural Resources (DLNR), and is part of the Honolulu Watershed Forest Reserve (Figure 1).

**Habitat and ecology.** *Cyanea konabuanuiensis* occurs in wet forest sites at elevations from 884 to 932 m. The associated native Hawaiian plant species include *Broussaisia arguta* Gaudich., *Cheirodendron platyphylla* (Hook & Arn.) Seem., *Dubautia laxa* (Hook. & Arn.) *Machaerina angustifolia* (Gaudich.) T. Koyama, *Metrosideros rugosa* A. Gray, *Perrottetia sandwicensis* A. Gray, *Phyllostegia grandiflora* (Gaudich.) Benth., *Polyscias gymnocarpa* (Hillebr.) Lowry & G. M. Plunkett, *Sadleria pallida* Hook. & Arn., *Scaevola mollis* Hook. & Arn., and species of *Cibotium* Kaulf., *Kadua* Cham. & Schldtl., *Labordia* Gaud., *Melicope* J.R. Forst. & G. Forst., and *Wikstroemia* Endl. Introduced naturalized plant species include: *Ageratina adenophora* (Spreng.) R.M. King & H. Rob., *Clidemia hirta* (L.) D.



**Figure 4.** *Cyanea konahuanuiensis*. **A** detail of corolla tube in profile. Dense pubescence and long calyx lobes apparent **B** close up view of buds **C** view of plant habit depicting single-stalked individual **D** view of plant habit depicting multi-stalked individual (photos by Tobias B. Koehler).

Don, *Hedychium gardnerianum* Sheppard ex Ker Gawl., *Rubus rosifolius* Sm., and *Setaria palmifolia* (J. König) Stapf. Bryophyte species are prevalent and include *Distichophyllum freycinetii* (Schwägr.) Mitt., *Plagiochila diflexa* Mont. & Gottsche, and species of *Bazzania* S. Gray. The native arthropod, *Megalagrion oahuense* Blackburn (a Hawaiian endemic damselfly) has been observed on *C. konahuanuiensis*. Soil is of basaltic origin and typical of wet forest sites on O‘ahu and the average annual rainfall is approximately 2600 mm (Giambelluca et al. 2013). The plants occupy a gulch both at the base and middle of relatively steep slopes which results in direct sunlight exposure occurring for a few hours close to midday and varying seasonally. Plants occur on both northwest and south facing slopes of the gulch, mostly along the banks of intermittent streams, but can also be found several meters from the stream with no apparent preference. Small stem fragments that are detached from plant have been observed to take root, forming new clones.

**Phenology.** *Cyanea konahuanuiensis* has been observed flowering from June–August with fruit developing from August–October. All observed fruits have aborted, been eaten, or decomposed before maturity. The lifespan and time to maturity of the species is unknown. Immature, nearly aborted fruits have been collected when all others had aborted. Fruits have been submitted to the Lyon Arboretum Micropropaga-





**Figure 5.** *Cyanea konahuanuiensis*. Inflorescence/flowers illustrating pendant inflorescence, dense pubescence of flowers, and long calyx lobes. Pubescent petioles and hairless stems are also apparent (photo by Tobias B. Koehler).

tion lab where some of the seeds have germinated and it is interesting to note that the seedlings have densely pubescent leaves (Figure 3).

**Etymology.** The specific name pays homage to the twin-peaked (946 m and 960 m) Kōnāhua-nui Pu‘u (summit), the tallest peaks in the Ko‘olau Mountain range on windward (east) O‘ahu. *Lit.* Large fat innards (Pukui et al. 1974), + Latin suffix *-ensis*, indicating a place of origin or belonging. “In one story a giant threw his great testicles (*Kona hua nui*) at a woman who escaped him.” (Pukui et al. 1974). Kōnāhua-nui has significance not only because it is the highest peak in the Ko‘olau Range, but because the summit area is a largely intact native ecosystem in relative close proximity to Honolulu, the largest city in the State of Hawai‘i. To our knowledge, there has never before been a plant species named after this beautiful and biologically important locality. After seeking counsel with Hawaiian cultural practitioners Kaua Neumann and Kihei Nahale-a (pers. comm. 2014), it is proposed to give the species a Hawaiian name of *Hāhā mili‘ohu*, meaning “The *Cyanea* that is caressed by the mist”.

**Conservation.** *Cyanea konahuanuiensis* is a critically imperiled species (see Conservation status below) due to its low population numbers and exceptionally narrow endemism. Some of the conservation obstacles to overcome include probable loss of



**Figure 6.** *Cyanea konahuanuiensis*. Inflorescence/flowers illustrating dense pubescence of flowers, long calyx lobes. Pubescent leaves also apparent (photo by Chris A. Johns).

most, if not all, of its native avian pollinators and dispersers, and suspected herbivory by introduced taxa such as rats, terrestrial gastropod mollusks (slugs), and feral pigs (*Sus scrofa*). Invasive plant species are becoming increasingly common even in relatively hard to access sites along and near mountain summits in Hawai'i. Species such as *Ageratina adenophora*, *Clidemia hirta*, *Erigeron karvinskianus*, *Hedychium gardnerianum*, *Rubus rosifolius* and *Setaria palmifolia* are competing with *C. konahuanuiensis* and other native species for space and resources. It is conceivable that stochastic events such as landslides, hurricanes, and flash-flooding could obliterate the majority or all of the currently known plants with a single event. Approximately 20 mature plants and several immature plants have been observed in total. Plants that were observed ranged from seedlings to reproductively mature individuals. Seedlings are scarce, however, which suggests that the population may be declining.

The Hawai'i DLNR, Division of Forestry and Wildlife (DOFAW) has largely funded this research by providing staff time to further investigate this species. The Hawai'i Plant Extinction Prevention (PEP) Program focuses on conserving and restoring plants with less than 50 known wild individuals. Because *C. konahuanuiensis* falls within that threshold, O'ahu PEP is working closely with DOFAW staff to protect this critically rare taxon. The first goal of the PEP Program is to secure seeds or propagules from each individual mature plant for *ex situ* germplasm banking. The long-term goal for the PEP Program and DOFAW will be to collaborate in the effort to grow and out-



**Figure 7.** *Cyanea konahuaniensis*. Solitary stem illustrating typical leaf arrangement clustered distally near the stem terminus (photo by Chris A. Johns).

plant nursery stock into appropriate restoration sites. Currently there are no protected (fenced) areas in similar habitat with comparable elevation, rainfall, humidity, and species composition on O‘ahu. Our recommendation is that additional fenced out-planting sites be established in appropriate areas of the Ko‘olau Mountains in order to establish multiple populations of this species. The authors emphasize the importance of prioritizing staff time to carry out further vegetation surveys in areas that have not been explored in recent history as this exciting new find shows that even seemingly well botanized areas in Hawai‘i may yet yield new discoveries.

In October 2013, immature fruits from two plants were collected and have since germinated at the UH Harold L. Lyon Arboretum after being directly sown on an agar medium. This is valuable information since at this time all observed fruits seem to be aborting prior to maturity. We recommend collecting immature fruit (or mature fruit if possible) from all reproductive individuals during future fruiting seasons in order to secure genetic representation from all reproductively mature individuals in *ex situ* collections.



**Figure 8.** *Cyanea konahuanuiensis*. Photo of leaf illustrating dense pubescence on both adaxial and abaxial leaf surfaces (photo by Tobias B. Koehler).

The use of cellular phone-connected game cameras is recommended for monitoring of rare plants in remote locations. This novel use of game camera technology saved time and resources, optimizing the timing of visits and increasing the likelihood of making successful and representative observations required for the species description and fruit collections. This rapidly improving technology could have many positive impacts on monitoring rare plants for flower development, fruit development, herbivory impacts, and the effects of various seasonal events.

**Specimens examined.** USA. Hawaiian Islands O‘ahu [East O‘ahu]: Paratypes: dried herbarium specimen BISH 1049144; and spirit collections: BISH 1059013, BISH 1059014, and BISH 1059015.

**Discussion.** For over a century the taxa that currently comprise the genus *Cyanea* were recognized as two separate genera, *Cyanea* and *Rollandia*. The genus *Rollandia* was distinguished from *Cyanea* based on the single character of staminal column adnation to corolla in the former (Gaudichaud-Beaupré 1844; Hillebrand 1888; Lammers 1990; Rock 1919). Phylogenetic investigations of molecular data revealed that the taxa of *Rollandia* are embedded within the genus *Cyanea* (Lammers, 1993). Together with two species of *Cyanea* (*C. acuminata* and *C. grimesiana*), they form a clade referred to as the “*acuminata* clade” (Givnish et al. 1995). *Cyanea konahuanuiensis* most likely



**Figure 9.** *Cyanea konahuaniensis*. Photo of holotype (photo courtesy of Bishop Museum).

belongs in this clade based on the staminal column being adnate to the corolla tube. It is noteworthy to mention that *C. konahuanuiensis* shares close geographic proximity to several taxa of the *acuminata* clade. Eight out of the nine previously recognized *Rollandia* taxa from this clade are endemic to O‘ahu (the ninth taxon, *C. parvifolia* C. N. Forbes is only known from the type specimen collected on Kaua‘i) and six out of those eight are even more restricted, occurring only in the Ko‘olau Range. Of those six, one taxon (*C. humboldtiana*) is known to occur on the summit ridges near *C. konahuanuiensis*.

### Key to species

There is currently no published taxonomic key for *Cyanea* that includes the merger of the *Rollandia* clade. The following couplets can be inserted into the most recent revision of *Rollandia* (Lammers 1990 in Wagner et al. 1990) to separate *C. konahuanuiensis* from *C. humboldtiana*.

- 2(1) Inflorescence pendent, peduncles 50–250 mm long.....2'  
 2'(2) Upper leaf surface glabrous, calyx lobes 4–10 mm long..... *C. humboldtiana*  
 2' Upper leaf surface densely pubescent, calyx lobes 16–18 mm long.....  
 ..... *C. konahuanuiensis*

### Conservation status

Using the IUCN Red List criteria (IUCN 2012), *Cyanea konahuanuiensis* falls into the Critically Endangered (CR) category, a rank given to species facing the highest threat of extinction in the wild, fitting the following criteria defined by IUCN Red List: B1) Extent of occurrence estimated to be < 100 km<sup>2</sup> and D) number of mature individuals < 50. This species is currently known from only one population in two sub-gulches of a single stream drainage. It is our recommendation that *C. konahuanuiensis* be evaluated by the United States Fish and Wildlife Service to be added as a Candidate for listing as Endangered under the Endangered Species Act of 1973. A recovery plan should be written and implemented.

### Acknowledgements

The authors extend our appreciation to the Department of Land and Natural Resources, Division of Forestry and Wildlife (DLNR/DOFAW) for access to the Honolulu Watershed Forest Reserve (FR) and use of staff time for this project. We thank Randall W. Kennedy, Ryan K. Peralta, David G. Smith, and Marigold S. Zoll from DLNR/DOFAW for encouraging us to write this species description and for technical support with accessing the FR. Sincere thanks go to Keahi M. Bustamente, Marian M. Chau, William P. Haines, Susan Ching Harbin, J. Christopher Havran, Clyde T. Imada,

Matthew J. Keir, Timothy I. Kroessig, Joel Q. C. Lau, Clifford W. Morden, Kihei Nahale-a, Kaua Neumann, Douglas O. Okamoto, Hank L. Oppenheimer, Richard J. Pender, Tom A. Ranker, Lara S. Reynolds, and Lauren A. Weisenberger for botanical expertise, assistance, encouragement, and thought-provoking discussions. Thank you to Yoshiko Akashi for making the species distribution map and thanks to Chris A. Johns for making a special trip to take descriptive photos. We thank Barbara H. Kennedy, the collection manager of the Bernice P. Bishop Museum herbarium (BISH) for access to specimens and assistance with preparing the Type specimens for this study and we thank the Lyon Arboretum for assistance with managing propagules that were obtained during the course of this study. This manuscript was greatly improved by suggestions from David H. Lorence (PTBG) who also provided encouragement and mentorship throughout the process from investigating the plants to writing the paper. Vicki A. Funk and Warren L. Wagner (US) are thanked for valuable suggestions on the manuscript.

## References

- Gaudichaud-Beaupré (1844) Voyage autour du monde, exécuté pendant les années 1836 et 1837 sur la corvette la *Bonite*, commandée par M. Vaillant...Botanique...Atlas. Arthus Bertrand, Paris, pl. 1–150 (*Rollandia humboldtiana*, pl. 76).
- Giambelluca TW, Chen Q, Frazier AG, Price JP, Chen Y-L, Chu P-S, Eischeid JK, Delparte DM, (2013) Online Rainfall Atlas of Hawai'i. Bull. Amer. Meteor. Soc. 94: 313–316. doi: 10.1175/BAMS-D-11-00228.1
- Givnish TJ, Millam KC, Mast AR, Paterson TB, Theim TJ, Hipp AL, Henss JM, Smith JF, Wood KR, Sytsma KJ (2008) Origin, adaptive radiation and diversification of the Hawaiian lobeliads, (Asterales: Campanulaceae). Proceedings of the Royal Society of London, Series B 276: 407–416. doi: 10.1098/rspb.2008.1204
- Givnish TJ, Sytsma KJ, Smith JF, Hahn WJ (1995) Molecular evolution, adaptive radiation, and geographic speciation in *Cyanea* (Campanulaceae, Lobelioideae). In: Wagner WL, Funk VA. Hawaiian biogeography, evolution on a hot spot archipelago. Smithsonian Institution Press, Washington and London, 288–337.
- Hillebrand W (1888) Flora of the Hawaiian Islands. 1965 reprint, Hafner Publishing Co., New York & London, 1–673.
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland, Switzerland and Cambridge, UK, iv + 32 pp.
- Lammers TG (1990) Campanulaceae. In: Wagner WL, Herbst DR, Sohmer SH. Manual of the Flowering Plants of Hawai'i. University of Hawai'i Press & Bishop Museum Press, Honolulu, 420–489.
- Lammers TG, Givnish TJ, Sytsma KJ (1993) Merger of the Endemic Hawaiian Genera *Cyanea* and *Rollandia* (Campanulaceae, Lobelioideae). Novon 3: 437–441. doi: 10.2307/3391391
- Lammers TG (1999) Campanulaceae. In: Wagner WL, Herbst DR, Sohmer SH. Manual of the Flowering Plants of Hawai'i. University of Hawai'i Press & Bishop Museum Press, Honolulu, 420–489.

- Lammers TG (2007) World Checklist and Bibliography of Campanulaceae. Royal Botanic Gardens, Kew, 675 pp.
- Oppenheimer H, Lorence DH (2012) A new species of *Cyanea* (Campanulaceae, Lobelioideae) from Maui, Hawaiian Islands. *PhytoKeys* 13: 15–23 doi: 10.3897/phytokeys.13.3447
- Pukui MK, Elbert SH, Mookini ET (1974) Place Names of Hawaii rev. ed. University of Hawaii Press, Honolulu.
- Rock JF (1919) A monographic study of the Hawaiian species of the tribe Lobelioideae, Family Campanulaceae. Bishop Museum Press, Honolulu. 1977 reprint, Krauss Reprint Co., Millwood, New York.
- Wagner WL, Herbst DR, Lorence DH (2005) Flora of the Hawaiian Islands website. <http://botany.si.edu/pacificislandbiodiversity/hawaiianflora/index.htm> [accessed July 2014]
- Wagner WL, Herbst DR, Sohmer SH (1999) Manual of the Flowering Plants of Hawai`i. 2 vols. University of Hawai`i Press & Bishop Museum Press, Honolulu, 1–1853.