



# A phylogenetic and taxonomic revision of sequestrate Russulaceae in Mediterranean and temperate Europe

J.M. Vidal<sup>1\*</sup>, P. Alvarado<sup>2\*</sup>, M. Loizides<sup>3</sup>, G. Konstantinidis<sup>4</sup>, P. Chachula<sup>5</sup>,  
P. Mleczko<sup>6</sup>, G. Moreno<sup>7</sup>, A. Vizzini<sup>8</sup>, M. Krakhmalnyi<sup>9</sup>, A. Paz<sup>10</sup>, J. Cabero<sup>11</sup>,  
V. Kaounas<sup>12</sup>, M. Slavova<sup>13</sup>, B. Moreno-Arroyo<sup>14</sup>, J. Llistosella<sup>15</sup>

## Key words

angiocarpic  
*Arcangiella*  
*Elasmomyces*  
Europe  
*Gymnomyces*  
*Lactarius*  
*Macowanites*  
*Martellia*  
pseudoangiocarpic  
taxonomy  
*Zelleromyces*

**Abstract** A comprehensive morphological and genetic study of type material and new collections of sequestrate *Russulales* species formerly belonging to the genera *Arcangiella*, *Elasmomyces*, *Gymnomyces*, *Hydnangium*, *Hymenogaster*, *Macowanites*, *Martellia*, *Secotium* and *Zelleromyces* is here undertaken, for the purpose of providing a complete taxonomical revision of sequestrate *Russulaceae* species in the Mediterranean and temperate regions of Europe. As a result, seven distinct taxa in the genus *Lactarius* and 18 in the genus *Russula* are identified. Six of them are new species: *L. populicola*, *L. subgiennensis*, *R. bavarica*, *R. candidissima*, *R. hobartiae* and *R. mediterraneensis*, and seven represent new combinations: *L. josserandii* ( $\equiv$  *Zelleromyces josserandii*), *L. soehneri* ( $\equiv$  *Hydnangium soehneri*), *R. candida* ( $\equiv$  *Hydnangium candidum*), *R. cerea* ( $\equiv$  *Hydnangium cereum*), *R. messapica* var. *messapicoides* ( $\equiv$  *Macowanites messapicoides*), *R. meridionalis* ( $\equiv$  *Zelleromyces meridionalis*) and *R. neuhoffii* ( $\equiv$  *Hydnangium neuhoffii*). Twenty-two of the 25 taxa are illustrated, while descriptions, microscopy images, as well as extensive information on the ecology, chorology and phylogeny for all taxa are provided. A key is further included to facilitate their identification.

**Article info** Received: 2 July 2018; Accepted: 4 December 2018; Published: 5 April 2019.

## INTRODUCTION

Sequestrate *Russulaceae* produce pseudoangiocarpic ('seco-toid') basidiomata, where the stipe is external and more or less reduced, to angiocarpic ('gasteroid') basidiomata, where the stipe is internal, reduced to a columella, or altogether absent. Sequestrate *Russulaceae* seem to be widespread throughout the globe, especially in arid and semi-arid regions of Australia and New Zealand (Bouger 1997, Lebel 1998, 2001, 2002, 2003a, b, Lebel & Trappe 2000, Bouger & Lebel 2001, Lebel & Castellano 2002, Trappe & Claridge 2003, Lebel & Tonkin 2007) and North America (Zeller & Dodge 1919, 1937, Singer

& Smith 1960, Smith 1963, Miller & Lebel 1999, Fogel & States 2001, Desjardin 2003, Smith et al. 2006). However, several sequestrate species have also been documented in tropical forests in Africa (Dring & Pegler 1978, Eberhardt & Verbeken 2004, Verbeken & Walleyn 2010, Beenken et al. 2016) and Asia (Corner & Hawker 1953, Heim 1959, Zhang & Yu 1990, Tao et al. 1993, Verbeken et al. 2014a, b), as well as in temperate *Nothofagus* forests in Patagonia (Trierweiler-Pereira et al. 2015).

The generic organization of sequestrate taxa within the *Russulaceae* has long been a subject of debate (Singer & Smith 1960, Pegler & Young 1979, Beaton et al. 1984, Zhang & Yu 1990). The subtle differences between *Gymnomyces*, *Martellia*, *Cystangium*, *Elasmomyces* and *Macowanites*, as well as the latex-bleeding *Arcangiella* and *Zelleromyces*, led to different taxonomic arrangements to accommodate an increasing number of emerging species (Lebel & Trappe 2000, Trappe et al. 2002, Vidal 2004a). Early authors hypothesized a single evolutionary lineage from the gymnocarpic to the angiocarpic genera in the *Russulales* (Malençon 1931, Heim 1938, Singer & Smith 1960, Oberwinkler 1977), but genetic studies revealed that sequestrate russuloid lineages are polyphyletic and nest within the genera *Russula* and *Lactarius* (Calonge & Martín 2000, Miller et al. 2001, Larsson & Larsson 2003, Eberhardt & Verbeken 2004, Nuytinck et al. 2003, Shimono et al. 2004, Smith et al. 2006, Lebel & Tonkin 2007). The split of *Lactarius* due to genetic evidence (Buyck et al. 2008, 2010) led to the restoration of the old genus *Lactifluus* and the erection of *Multfurca*, although no sequestrate species are yet known to occur in either of these two lineages.

The genus *Gymnomyces*, typified by *G. pallidus* (Massee 1898), a species found in myrtaceous forests whose current name is *R. paneeroides* (Lebel 2017), was characterized by the absence of a stipe-columella and the presence of sphaerocytes in the hymenophoral trama. The European species *G. xanthosporus*

<sup>1</sup> C/ Massaballs 10, E-17118 Sant Sadurní de l'Heura, Girona, Spain.

<sup>2</sup> ALVALAB, La Rochela 47, E-39012 Santander, Spain;  
corresponding author e-mail: pablo.alvarado@gmail.com.

<sup>3</sup> P.O. Box 58499, 3734 Limassol, Cyprus.

<sup>4</sup> Agiou Kosma 25, 51100 Grevena, Greece.

<sup>5</sup> Pieniny National Park, Jagiellońska 107b, PL-34-450 Krościenko nad Dunajcem, Poland.

<sup>6</sup> Institute of Botany, Jagiellonian University in Kraków, Gronostajowa 3, PL-30-387 Kraków, Poland.

<sup>7</sup> Dept. de Ciencias de la Vida (Botánica), Facultad de Biología, Universidad de Alcalá, E-28871 Alcalá de Henares, Spain.

<sup>8</sup> Dipt. di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Viale P.A. Mattioli 25, I-10125 Torino, Italy.

<sup>9</sup> Institute of Evolution and Department of Evolutionary and Environmental Biology, University of Haifa, 199 Abba Khoushy Ave, Haifa, 3498838, Israel.

<sup>10</sup> C/ Vall Ter 791, Urbanització Llac del Cigne, E-17455 Caldes de Malavella, Girona, Spain.

<sup>11</sup> C/ El Sol 6, E-49800 Toro, Zamora, Spain.

<sup>12</sup> Sokratous 40, TK-19016 Artemis Attika, Greece.

<sup>13</sup> Krivolak 4, 1164 Sofia, Bulgaria.

<sup>14</sup> Dept. de Biología Vegetal, Facultad de Biología, Universidad de Córdoba, Colonia San José 4, Campus de Rabanales, E-14014 Córdoba, Spain.

<sup>15</sup> Dept. de Biología Evolutiva, Ecología i Ciències Ambientals, Universitat de Barcelona, Av. Diagonal 643, E-08028 Barcelona, Spain.

\* These authors contributed equally.

was originally described by Hawker (1952) as a variety of *Hydnangium carneum*, to accommodate some specimens found in the UK, and subsequently re-combined into *Gymnomyces* by Smith (1962), because of its amyloid spores and sphaerocytes. Several species were described later from the Mediterranean basin or re-combined into this genus, such as *G. ilicis* (Llistosella & Vidal 1995), *G. dominguezii* (Moreno-Arroyo et al. 1999, and subsequently validated in Calonge 2000), *G. sublevisporus* (Moreno-Arroyo et al. 2002) and *G. meridionalis* (Vidal 2004a), the latter originally treated in the genus *Zelleromyces* by Moreno-Arroyo et al. (1998b, and subsequently validated in Calonge 2000).

The genus *Martellia*, typified by *M. mistiformis* (Mattirollo 1900), was first described from several collections found in Sardinia (Italy) under *Quercus suber*. It was characterized by the absence of a stipe-columella and lack of sphaerocytes in the hymenophoral trama (Singer & Smith 1960). In the late 20th century, new species and combinations were proposed: *M. mediterranea* (Moreno et al. 1991), *M. pila* (Vidal 1991b), *M. monospora* (Astier & Pacioni 1998), the last two originally described from France in the genus *Hydnangium* by Patouillard (1910) and Boudier & Patouillard (1888) respectively, and *M. aurantiaca* (Astier & Pacioni 1998), originally described from Spain in the genus *Hydnangium* by Heim et al. (1934) and later re-combined into the genus *Stephanospora* by Vidal (2004c). After comparing the types of *Gymnomyces* and *Martellia*, Lebel & Trappe (2000) considered that the presence or absence of sphaerocytes in the hymenophoral trama was of no taxonomic relevance in keeping the two genera separate, and subsumed *Martellia* into *Gymnomyces* (Trappe et al. 2002).

The genus *Elasmomyces*, typified by *E. mattiroloanus* (Cavara 1897), was proposed to accommodate some stipitate collections obtained under *Abies alba* in Italy, that lacked sphaerocytes in the hymenophoral trama. Saccardo & Saccardo (1905) proposed the combinations *E. krjukowensis* and *E. michailowskianus* for two species of the genus *Secotium*, originally found in Russia by Bucholtz (1901, 1903).

The genus *Macowanites*, typified by *M. agaricinus* (Kalchbrenner 1882), which was originally placed in the genus *Macowania* (Kalchbrenner 1876), was first proposed for a stipitate fungus found in Eastern Cape (South Africa), exhibiting sphaerocytes in the hymenophoral trama. Much later, some Mediterranean species were added or re-combined into this genus, namely *M. galileensis* from Mount Carmel, Israel (Moser et al. 1977) and *M. messapicoides* (Llistosella & Vidal 1995), *M. ammophilus* (Vidal et al. 2002), originally described as *Gymnomyces* by Calonge & Vidal (1999), and *M. vinaceodorus* (Calonge & Vidal 2001), all from Spain. Recently, *M. candidus* (Vidal 2004b), a species originally described from Northern France in the genus *Hydnangium* (Tulasne & Tulasne 1843), was also placed into this genus. Same as for the angiocarpic genera *Gymnomyces* and *Martellia*, the pseudoangiocarpic taxa *Elasmomyces* and *Macowanites* were also considered synonyms by Lebel & Trappe (2000), following comparison of their type collections.

The genus *Cystangium*, typified by *C. sessile* (Singer & Smith 1960), is another russuloid sequestrate genus, but appears to be confined to Australia and South America, with no known representatives in Europe. It is characterized by an epithelial pileipellis, in contrast to *Gymnomyces*, *Martellia*, *Elasmomyces* and *Macowanites*, all of which have a trichodermal or undifferentiated pileipellis. The stipe in this genus can be external, reduced to a columella, or completely absent (Lebel & Trappe 2000).

The latex-bleeding genus *Arcangeliella* was typified by *A. borziana* (Cavara 1900), and proposed for collections found under *Abies alba* in central Italy. *Arcangeliella borziana* is a sessile

species that sometimes presents a poorly developed or residual stipe and a percurrent columella, especially in young basidiomata. Despite these features, *Arcangeliella* was chosen by Singer & Smith (1960) to describe several non-European sequestrate species with a well-developed stipe-columella, whereas Vidal (2004a) maintained this genus only for angiocarpic, sessile species, keeping *Gastrolactarius* for pseudoangiocarpic, stipitate species. American mycologists (Dodge 1931, Zeller & Dodge 1935) combined several European taxa into *Arcangeliella*, e.g., *A. asterosperma* var. *asterosperma*, *A. asterosperma* var. *depauperata*, *A. asterosperma* var. *hololeuca*, *A. laevis* or *A. stephensii*; the first four, however, belonging to the order *Boletales*, have since been transferred to the genus *Octaviania* (Paz et al. 2016). Finally, Mader & Mader (1992) proposed *A. volemoides* for a collection made under *Picea abies* in the Austrian Alps, but this species was considered a synonym of *A. borziana* by Nuytinck et al. (2003) and Vidal (2004a).

The latex-bleeding genus *Zelleromyces*, typified by *Z. cinnabarinus* (Singer & Smith 1960), was proposed for angiocarpic species devoid of a stipe-columella. Several Mediterranean and Central European species were later described or re-combined into this genus: *Z. josserandii* from Morocco and France (Malençon 1975), *Z. hispanicus* (Calonge & Pegler 1998) and *Z. giennensis* (Moreno-Arroyo et al. 1998a) from Spain, and *Z. soehneri* (Trappe et al. 2002), originally described from Germany in the genus *Hydnangium* by Zeller & Dodge (1935). Recently, several taxa previously placed in the sequestrate genera *Cystangium*, *Elasmomyces*, *Gymnomyces*, *Macowanites*, and *Martellia*, were re-combined into *Russula* by Lebel (2017) and Elliott & Trappe (2018). About 30 sequestrate species of *Russulaceae* have been described from Mediterranean and temperate regions of Europe, but the generic placement for many of these still remains unclear. In the present work, all of them are reviewed and revised in the light of extensive genetic data. New sequences were produced from herbarium specimens and new collections of pan-European origins, in order to evaluate the phylogenetic basis of characters traditionally employed to discriminate among sequestrate taxa, and propose the most appropriate supraspecific affiliation of these species within *Lactarius* and *Russula*.

## MATERIAL AND METHODS

### Fungal collections

Morphological studies include type material and old representative collections of selected species placed in the genera *Arcangeliella*, *Elasmomyces*, *Gymnomyces*, *Hydnangium*, *Hymenogaster*, *Macowanites*, *Martellia*, *Octaviania*, *Secotium* and *Zelleromyces* kept in the public herbaria of BPI (Beltsville, USA), FH (Cambridge, USA), IB (Innsbruck, Austria), K (Kew, UK), M (Munich, Germany), MA (Madrid, Spain), MPU (Montpellier, France), NICE (Nice, France), NY (New York, USA), PC (Paris, France), PRM (Praha, Czech Republic), UPS (Uppsala, Sweden) and WU (Wien, Austria), as well as recent collections from different localities in Europe and the Middle East (Bulgaria, Cyprus, France, Germany, Greece, Hungary, Israel, Italy, Poland, Portugal and Spain) kept in the public herbaria of AH (Alcalá de Henares, Spain), BCN (Barcelona, Spain), HAI (Haifa, Israel), KRA (Kraków, Poland), MCVE (Venezia, Italy), or in private herbaria of the Asociación Vallisoletana de Micología (AVM), B. Moreno Arroyo (BM), G. Konstantinidis (GK), A. Paz Conde (IC), J. Cabero (JC), J.M. Vidal (JMV), M. Loizides (ML), M. Slavova (MSL), V. Kaounas (VK) and other contributors to this paper (see acknowledgements). Newly collected specimens have been deposited in the Herbarium of the Universidad de Alcalá (AH), in the Herbarium of

the CeDocBiV of the Universitat de Barcelona (BCN-myc), in the Herbarium of the Jagiellonian University in Kraków (KRA), and in the Mycological Collection of the Bulgarian Academy of Sciences (SOMF).

### Morphological study

Most species are fully described, with the exception of the recently described species, for which only a summary description is provided. Detailed macromorphological notes and images were taken from the majority of collections, and the presence/absence of latex and columella, as well as ecological data (altitude, associated vegetation, putative ectomycorrhizal partner, substrate) were annotated. Colour of basidiomata, hymenophore, and spore mass observed in locules was described according to Kornerup & Wanscher (1978). Methuen code colour of fresh specimens was obtained from digital images. For microscopical features, a Carl Zeiss Jena Jenaval microscope with a DIC device, a Leica BM E binocular, and a Leica DMRB with condenser P 1.40 OIL S1, provided with planapochromatic optics 25 $\times$ , 40 $\times$ , 50 $\times$ , 63 $\times$  and 100 $\times$ , were employed, coupled with Reflex Pentax K-20 (14.6 Mpx) cameras with remote control. Amyloid reaction was checked from mounts in Melzer's reagent, while ammoniacal (10 %) or potassium hydroxide (2–5 %) Congo red was employed to enhance contrast in microscopical observations.

The terminology of Heilmann-Clausen et al. (1998) is adopted for naming the pileipellis structure. For the different types of secretory hyphae, the terminology of Cléménçon (2004) is adopted, using the terms: 1) 'laticifera' for non-septate, latex-containing hyphae of granular content (crystalloid in exsiccata), staining violet in contact with sulfovainillin, typical of *Lactarius* (Fig. 12c, f); 2) 'gloeoplera' for the septate hyphae not bleeding latex, also of granular or crystalloid content, not staining violet with sulfovainillin, which are found in the trama and context of *Russula* and generate both fertile (basidia) and sterile elements (macrocystidia and dermatocystidia) (Fig. 15f); and 3) 'thromboplera' for the tortuous hyphae of homogeneous, non-granular, yellow to brownish content, found in both *Lactarius* and *Russula* (Fig. 10b–c, 12c). We use the term 'cystidole' (Romagnesi 1944, Josserand 1983) for sterile hymenial elements similar to dermatocystidia first developing in external locules at the same level as basidioles, and sometimes remaining in the mature hymenium as in *Russula mistiformis* (Fig. 25j–l). We apply the term 'paraphysoid cells' (Hawksworth et al. 1995) to cylindrical or slightly clavate and sometimes septate sterile elements without granular content, present in the hymenium of many of the sequestrate species studied, e.g., sterile cells similar to septate basidioles found in the hymenium of *Lactarius josserandii* (Fig. 8g–h), called 'paracystidia' by Malençon (1975), or those in the hymenium of *L. borzianus* and *L. stephensii* (Fig. 11b–c), called 'hymenial hairs' or 'paracystidia' in Vidal (2004a).

The measurements of spores and other elements were obtained with the aid of 'Mycometre VA' software (G. Fannechère). Spores were measured in side view, always excluding ornamentation and hilar appendix, and at least 20 spores from each specimen were studied, indicating the minimum and maximum average and the Q-value. Scanning electron microscopes (SEM) Hitachi S-4100 and Zeiss DSM-960A (Unitat de Microscòpia dels Serveis Tècnics de Recerca de la Universitat de Girona, UdG), Hitachi S-4700 (Laboratory of Scanning Microscopy, Institute of Geological Sciences, Jagiellonian University in Kraków, UJ) and Zeiss DSM-950 (Servicio de Microscopía Electrónica de la Universidad de Alcalá de Henares, AH), were also employed for imaging spores. Light microscopy images of spores were stacked digitally with the aid of 'Helicon Focus' (Helicon Soft Ltd.).

### Molecular analysis

#### DNA extraction, amplification and sequencing

DNA extraction and PCR amplification were performed as described by Alvarado et al. (2012). Primers ITS1F and ITS4 (White et al. 1990, Gardes & Bruns 1993) were used for the ITS region; primers LR0R and LR5 (Vilgalys & Hester 1990) were used for the 28S rDNA ribosomal region (28S rDNA); bRPB2-6F2 (reverse from bRPB2-6R2) and bRPB2-7.1R2 (modified from bRPB2-7.1R) for the DNA-directed second largest subunit two of RNA polymerase II (*rpb2*) gene (Matheny 2005, Matheny et al. 2007, Gelardi et al. 2015); and in some samples, EF1-983F and EF1-1567R (Rehner & Buckley 2005) for the translation elongation factor 1α (*tef1*) gene for both, PCR and sequencing. Sequences were edited for errors in MEGA v. 5 (Tamura et al. 2011). New sequences are listed in Table 1 highlighted in **bold** characters, and newly sequenced specimens are marked with an asterisk symbol (\*) in the paragraph of material studied. Attempts to obtain genetic data from some old specimens were conducted without success.

#### Phylogenetic analyses

Newly generated sequences were aligned with the most similar sequences in the International Nucleotide Sequence Database Collaboration (INSDC) identified through BLASTn searches (Altschul et al. 1997). A multigenic ITS – 28S rDNA – *rpb2* alignment of Russulaceae was built including all lineages identified in the most recent phylogenetic works (Kong et al. 2015, Looney et al. 2016), as well as all sequences available originated from sequestrate species. Reference publications for the sequences employed are provided in Table 1. *Stereum hirsutum*, *Auriscalpium vulgare*, *Amylostereum laevigatum* and *Echinodontium tinctorium* were employed as outgroups for the general alignment because these are known to be the closest lineages outside the Russulaceae clade (Buyck et al. 2008). Sequences were first aligned in MEGA v. 5 (Tamura et al. 2011) with Clustal W (Higgins et al. 1994) and the resulting alignments edited manually. First, a full query for the best-scoring maximum likelihood tree was performed on the general alignment in RAxML (Stamatakis 2006) using the standard search algorithm (ITS, 28S rDNA, *rpb2* data partitioned, 2000 bootstrap replications). Four independent alignments were taken from the general one: 1) *Lactarius*; 2) *Russula* sect. *Ingratae*; 3) *Russula* sect. *Rigidae*; and 4) *Russula* s.str., using *Lactifluus piperatus* as outgroup for all of them. GBlocks v. 0.91b (Castresana 2000) was employed with the most permissive conditions to remove ambiguously aligned positions from ITS rDNA data. Bayesian analyses were performed in MrBayes v. 3.2.6 (Ronquist & Huelsenbeck 2003) employing optimal models determined for ITS, 28S rDNA and *rpb2* partitions in MrModeltest v. 2.3 (Nylander 2004) package loaded in PAUP v. 4.0b10, two simultaneous runs, six chains, temperature set to 0.2, and sampling every 100th generation until convergence parameters (< 0.01) were met after 9.35 M generations in *Lactarius*, 10.4 M in *Russula* sect. *Ingratae*, 0.74 M in *Russula* sect. *Rigidae* and 8.64 M generation in *Russula* s.str. The first 25 % trees were discarded as burn-in. Fig. 1 represents the phylogenetic tree resulting from the ML analysis of all sequences produced in the present work and retrieved from public databases with the major clades condensed. Fig. 2–5 were obtained from the previous one, collapsing all nodes except a single major clade: *Lactarius* (Fig. 2), *Russula* sect. *Ingratae* (Fig. 3), *Russula* sect. *Rigidae* (Fig. 4) and *Russula* s.str. (Fig. 5). Nodes were annotated with support values from Maximum Likelihood and Bayesian analyses if at least one of these was considered significant, or annotated in brackets when at least one of these was considered subsignificant. Support values were considered significant when ML bootstrap (BP) values were above 70 % or posterior probability (PP) values

**Table 1** Specimens used in molecular phylogenetic studies and their GenBank accession numbers.

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<b>Eurussulloid clade</b>							
<i>Lamellostereaceae</i>							
<i>Amylostereum laevigatum</i>	<i>Amylostereum laevigatum</i>	France	olrim409, 623.84 (CBS)	Wang et al. (2004)	AY781246	AF287843	AY218469
<i>Auriscalpiaceae</i>					DQ911613	AF287843	AY218469
<i>Auriscalpium vulgare</i>	<i>Auriscalpium vulgare</i>	USA	PBM944 (WTU)	Lutzoni et al. (2004)	AY854088	AF393056	AY218482
<i>Echinodontiaceae</i>							
<i>Echinodontium tinctorium</i>	<i>Echinodontium tinctorium</i>	USA	16666 (DAOM)	Bindler & Hibbett (2002)			
<i>Irussulaceae/lactarius</i>							
<i>Lactarius</i> sp.							
<i>Lactarius</i> sp.*							
<i>Lactarius akahatsu</i>	<i>Lactarius akahatsu</i>	Tasmania	BH2148F, T991 (HO)	Kohout (unpubl.)	JF960610	—	—
<i>Lactarius angiocarpus*</i>	<i>Lactarius angiocarpus*</i>	Tasmania	BH1714P (HO)	Horton (2011)	JF960854	—	—
<i>Lactarius atroviridis</i>	<i>Lactarius atroviridis</i>	Tasmania	BH3528R (HO)	Horton (2011)	JF960852	—	—
<i>Lactarius aurilla</i>	<i>Lactarius aurilla</i>	Germany	BH2174F (HO)	Horton (2011)	JF960853	—	—
<i>Lactarius acris</i>	<i>Lactarius acris</i>	Thailand	EU014 (UPS)	Buyck et al. (2008)	DQ421988	DQ421922	DQ421922
<i>Lactarius albovires</i>	<i>Lactarius albovires</i>	Zambia	AV2004-141 (GENT)	Verbeken et al. (2014b)	KF133269	KF133301	KF133333
<i>Lactarius angiocarpus*</i>	<i>Lactarius angiocarpus*</i>	USA	DA00-448 (GENT)	Eberhardt & Verbeken (2004), Stubbe et al. (2010)	AY608942	DQ421981	GU258310
<i>Lactarius atroviridis</i>	<i>Lactarius atroviridis</i>	Sweden	AV05-306 (GENT)	Verbeken et al. (2014b)	KF133270	KF133302	KF133334
<i>Lactarius aurilla</i>	<i>Lactarius aurilla</i>	Belgium	RW1601 (GENT)	Verbeken et al. (2014b)	KF133257	KF133288	KF133321
<i>Lactarius azonites</i>	<i>Lactarius azonites</i>	Malawi	DS08-520 (GENT)	Stubbe & Verbeken (2012)	JQ446168	JQ446231	JQ446231
<i>Lactarius ballophaeus</i>	<i>Lactarius ballophaeus</i>	Thailand	AV05-155 (GENT)	Verbeken et al. (2014b)	GU258277	GU265576	GU258312
<i>Lactarius bisporus*</i>	<i>Lactarius bisporus*</i>	France	FH12-160 (GENT)	Verbeken et al. (2014a)	KF856293	—	—
<i>Lactarius bozianus*</i>	<i>Lactarius bozianus*</i>	Italy	JMV800279b (BCN)	<b>This work</b>	<b>MK105610</b>	<b>MK102740</b>	<b>MK105696</b>
<i>Arcangelia boziana</i>	<i>Arcangelia boziana</i>	Switzerland	16944 (MCE/E)	Osmundson et al. (2013)	JF908775	—	—
<i>Arcangelia boziana</i>	<i>Arcangelia boziana</i>	Sweden	FA96-05-3344 (WSL)	Peer et al. (2001)	AF286204	AF286203	—
<i>Lactarius camphoratus</i>	<i>Lactarius camphoratus</i>	Italy	UE04-09-2004 (UPS)	Verbeken et al. (2014b)	DQ422009	DQ421933	DQ421933
<i>Lactarius chrysorheus</i>	<i>Lactarius chrysorheus</i>	Sweden	UE04-10-2002-8 (UPS)	Verbeken et al. (2014b)	KF133261	KF133293	KF133325
<i>Lactarius citriolens</i>	<i>Lactarius citriolens</i>	Thailand	UE20-09-2004-03 (UPS)	Verbeken et al. (2014b)	DQ422003	DQ421931	DQ421931
<i>Lactarius crassiusculus</i>	<i>Lactarius crassiusculus</i>	USA	HTL369 (GENT)	EF560684	KF133303	KF133335	KF133335
<i>Lactarius crassus*</i>	<i>Lactarius crassus*</i>	USA	T17996	KT9688636	KT9688581	KT9688636	—
<i>Arcangelia crassa</i>	<i>Arcangelia crassa</i>	USA	41826 (OSC)	KT968863	KT9688615	KT9688615	—
<i>Lactarius cyanescens</i>	<i>Lactarius cyanescens</i>	Malaysia	DS06-058 (GENT)	DQ422009	JN375602	JN375602	JN375602
<i>Lactarius cyathiformis</i>	<i>Lactarius cyathiformis</i>	Sweden	UE04-09-2004-2 (UPS)	KF133278	GU258278	GU258278	GU258278
<i>Lactarius deliciosus</i>	<i>Lactarius deliciosus</i>	Slovakia	JN2001-046 (GENT)	KF133266	KF133266	KF133266	KF133266
<i>Lactarius echinellus</i>	<i>Lactarius echinellus</i>	Sri Lanka	AV07-169 (GENT)	KF133272	KF133272	KF133272	KF133272
<i>Lactarius echinus*</i>	<i>Lactarius echinus*</i>	Sri Lanka	AV07-168 (GENT)	KF133273	KF133273	KF133273	KF133273
<i>Lactarius evosmus</i>	<i>Lactarius evosmus</i>	Sweden	536 (UP)	DQ658882	—	—	—
<i>Lactarius fulacetus*</i>	<i>Lactarius fulacetus*</i>	Thailand	KVP08-038 (GENT)	KF133274	KF133307	KF133339	KF133339
<i>Lactarius flavopalustris</i>	<i>Lactarius flavopalustris</i>	Finland	JV23334 (TURA)	KR090904	KR090982	KR090982	KR090982
<i>Lactarius flexuosus</i>	<i>Lactarius flexuosus</i>	Sweden	UE06-09-2002-1 (UPS)	DQ421992	DQ421925	DQ421925	DQ421925
<i>Lactarius fuliginosus</i>	<i>Lactarius fuliginosus</i>	Sweden	MTB97-24 (GENT)	JQ446111	JQ446180	JQ446180	JQ446180
<i>Lactarius fulvissimus</i>	<i>Lactarius fulvissimus</i>	Germany	JKLAC10082002 (GENT)	KF432970	KR025662	KR025662	KR025661
<i>Lactarius fumosibrunneus</i>	<i>Lactarius fumosibrunneus</i>	Mexico	JN2012-025 (GENT)	KR025576	—	—	—
<i>Lactarius fumosus</i>	<i>Lactarius fumosus</i>	Canada	EG10 (XAL)	JN797633	—	—	—
<i>Zelleromyces gardneri</i>	<i>Zelleromyces gardneri</i>	USA	HRL0894	KJ705224	—	—	—
<i>Zelleromyces giennensis*</i>	<i>Zelleromyces giennensis*</i>	USA	1537 (SOC)	JN022500	—	—	—
<i>Zelleromyces giennensis*</i>	<i>Zelleromyces giennensis*</i>	Spain	513 (SOC)	DQ453696	—	—	—
<i>Lactarius fumosus</i>	<i>Lactarius fumosus</i>	Spain	Fungi 38674 (MA)	AF230900	<b>MK105697</b>	<b>MK105697</b>	<b>MK105697</b>
<i>Lactarius fumosus</i>	<i>Lactarius fumosus</i>	Spain	AVM 1615, JMV800629 (BCN)	—	<b>MK105612</b>	<b>MK105612</b>	<b>MK105612</b>
<i>Lactarius gardneri*</i>	<i>Lactarius gardneri*</i>	USA	JC20061020 (pers. herb.)	KR090988	—	—	—
<i>Lactarius giennensis*</i>	<i>Lactarius giennensis*</i>	USA	20923 (DGB)	KT583642	—	—	—
<i>Lactarius haugiae</i>	<i>Lactarius haugiae</i>	Mexico	LM4994 (XAL)	KT583649	KT736507	KT736507	KT736507
<i>Lactarius helvus</i>	<i>Lactarius helvus</i>	Sweden	UE08-09-2004-1 (UPS)	KF133263	KF133295	KF133295	KF133295
<i>Lactarius hispidulus</i>	<i>Lactarius hispidulus</i>	Guinea	AB152 (GENT)	KF133258	KF133322	KF133322	KF133322

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Lactarius horakii</i> <sup>*</sup>	<i>Lactarius horakii</i> <sup>*</sup>	Indonesia	EH8336 (ZT)	Nuytinck et al. (2007)	EF685069	—	—
<i>Lactarius josserandii</i> <sup>*</sup>	<i>Lactarius josserandii</i>	Spain	JMV800621 (BCN)	<b>MK105613</b>	AF231911	—	<b>MK102742</b>
<i>Zelleromyces hispanicus</i>	<i>Zelleromyces hispanicus</i>	Spain	Fungi 37497 (MA)	<b>This work</b>	AF231912	—	—
<i>Zelleromyces hispanicus*</i>	<i>Zelleromyces hispanicus*</i>	Spain	Fungi 37498 (MA)	Calonge & Martin (2003)	AJ555267	—	—
<i>Lactarius kabansus</i>	<i>Lactarius kabansus</i>	Zimbabwe	Fungi 53339 (MA)	Calonge & Martin (2003)	KF133295	KF133323	—
<i>Lactarius lanceolatus</i>	<i>Lactarius lanceolatus</i>	USA	AV98-205 (GENT)	Verbeken et al. (2014b)	KR090915	KR090989	—
<i>Lactarius lignyotus</i>	<i>Lactarius lignyotus</i>	USA	F 4239 (VPI)	Barge et al. (2016)	DQ221107	DQ408128	—
<i>Lactarius lignyotus</i>	<i>Lactarius lignyotus</i>	Sweden	PBM2424 (CULW)	Matheny et al. (2007)	DQ421993	DQ421926	—
<i>Lactarius lignyotus</i>	<i>Lactarius lignyotus</i>	Belgium	UE06-09-2003-5 (UPSG)	Buijck et al. (2008)	KF133275	KF133340	—
<i>Lactarius lilacinus</i>	<i>Lactarius lilacinus</i>	USA	RW3774 (GENT)	Verbeken et al. (2014b)	KR090922	KR090994	—
<i>Lactarius luculentus</i> var. <i>laetus</i>	<i>Lactarius luculentus</i> var. <i>laetus</i>	Belgium	F 024643 (DBG)	Barge et al. (2016)	KF1241547	—	—
<i>Lactarius luteolus</i>	<i>Lactarius luteolus</i>	Sweden	OB11-011 (GENT)	Verbeken et al. (2014b)	KF133297	KF133329	—
<i>Lactarius mammosus</i>	<i>Lactarius mammosus</i>	India	UE09-09-2004-5 (UPSG)	Verbeken et al. (2014b)	KF133265	KF133342	—
<i>Lactarius montoyae*</i>	<i>Lactarius montoyae*</i>	USA	KD1065 (BSD)	Stibbe et al. (2010)	KU265641	KU258380	—
<i>Lactarius nanus</i>	<i>Lactarius nanus</i>	France	EB106-13 (MONT)	Barge et al. (2016)	KR091000	KF133341	—
<i>Lactarius necator</i>	<i>Lactarius necator</i>	USA	AV04-231 (GENT)	Verbeken et al. (2014b)	KF133276	KR364320	—
<i>Lactarius olympianus</i>	<i>Lactarius olympianus</i>	USA	ED08-018 (GENT)	De Crop et al. (2017)	KR364089	KR091009	—
<i>Lactarius pallidomarginatus</i>	<i>Lactarius pallidomarginatus</i>	USA	CLC1470 (MONT)	Barge et al. (2016)	KR090939	KF133310	—
<i>Lactarius peckii</i>	<i>Lactarius peckii</i>	USA	JN2004-020 (GENT)	Verbeken et al. (2014b)	KF133277	KF133342	—
<i>Lactarius pomoliensis*</i>	<i>Lactarius pomoliensis*</i>	Sri Lanka	AV07-159 (GENT)	Verbeken et al. (2014b)	KF133282	KF133315	—
<i>Lactarius populicola*</i>	<i>Lactarius populicola*</i>	Belgium	RW2930 (GENT)	Nuytinck et al. (2003)	AY331012	—	—
<i>Lactarius psammincola</i>	<i>Lactarius psammincola</i>	Greece	GK4831, JMV800648 (BCN)	<b>MK105700</b>	<b>MK102743</b>	<b>KY848507</b>	<b>KY848507</b>
<i>Lactarius pterosporus</i>	<i>Lactarius pterosporus</i>	USA	BPL869 (TENN)	Looney et al. (2016)	JO446153	—	—
<i>Lactarius quieticolor</i>	<i>Lactarius quieticolor</i>	France	PAM061-00705 (GENT)	Stibbe & Verbeken (2012)	DQ42275	DQ421930	—
<i>Lactarius quietus</i>	<i>Lactarius quietus</i>	Sweden	UE10-09-2004-1 (UPSG)	Buijck et al. (2008)	KF133296	KF133328	—
<i>Lactarius representaneus</i>	<i>Lactarius representaneus</i>	USA	UE16-09-2004 (UP-S)	Verbeken et al. (2014b)	KR091020	KR091020	—
<i>Lactarius romagnesi</i>	<i>Lactarius romagnesi</i>	France	CLC2318 (MONT)	Barge et al. (2016)	DQ421989	DQ421923	—
<i>Lactarius ruginosus</i>	<i>Lactarius ruginosus</i>	Italy	UE29-09-2002-6 (UPSG)	Buijck et al. (2008)	KJ446150	KJ446216	—
<i>Lactarius salicis-reticulatae</i>	<i>Lactarius salicis-reticulatae</i>	Finland	DS09-09-2002-13 (GENT)	Stibbe & Verbeken (2012)	KR090959	KR091030	—
<i>Lactarius salmonicolor</i>	<i>Lactarius salmonicolor</i>	Germany	CLC2885 (MONT)	Barge et al. (2016)	AF325284	—	—
<i>Lactarius saturnisporus*</i>	<i>Lactarius saturnisporus*</i>	Germany	UE158/FO46879 (TUB)	Eberhardt et al. (2000)	KF133316	KF133348	—
<i>Lactarius scrobiculatus</i>	<i>Lactarius scrobiculatus</i>	Italy	AV07-170 (GENT)	Verbeken et al. (2014b)	KF08281	KF133311	—
<i>Lactarius Shoreae*</i>	<i>Lactarius Shoreae*</i>	Sri Lanka	732 (MCVE)	Osmundson et al. (2013)	EU834192	—	—
<i>Lactarius salmonicolor</i>	<i>Lactarius salmonicolor</i>	USA	AV07-164 (GENT)	Verbeken et al. (2014b)	EU652366	—	—
<i>Lactarius saturnisporus*</i>	<i>Lactarius saturnisporus*</i>	USA	64481 (OSC)	Gordon (unpubl.)	<b>MK105701</b>	<b>MK105615</b>	<b>MK102744</b>
<i>Lactarius silvæ*</i>	<i>Lactarius silvæ*</i>	USA	74231 (OSC)	Gordon (unpubl.)	<b>MK105616</b>	<b>MK105702</b>	<b>MK105702</b>
<i>Lactarius soehneri</i>	<i>Lactarius soehneri</i>	Spain	39272 (AH)	<b>This work</b>	KF133268	KF133300	KF133332
<i>Lactarius sphagneti</i>	<i>Lactarius sphagneti</i>	UK	46013 (AH)	<b>This work</b>	KY270490	KY270494	—
<i>Lactarius spinosporus*</i>	<i>Lactarius spinosporus*</i>	China	PL2805 (pers. herb.)	<b>This work</b>	KF133262	KF133294	KF133326
<i>Lactarius spinulosus</i>	<i>Lactarius spinulosus</i>	Sweden	FAN445 (BJTC)	<b>This work</b>	<b>MK105617</b>	<b>MK105618</b>	<b>MK105620</b>
<i>Lactarius stephensii</i>	<i>Lactarius stephensii</i>	Poland	AT2013068 (UPS)	<b>This work</b>	EU784439	KF133312	<b>MK105621</b>
<i>ArcangelIELLA stephensii</i>	<i>ArcangelIELLA stephensii</i>	Belgium	F-2014-17 (KRA)	<b>This work</b>	KF133279	KF133344	<b>MK105624</b>
<i>Lactarius subdulcis</i>	<i>Lactarius subdulcis</i>	Cyprus	JMV20010609-1 (BCN)	<b>This work</b>	<b>MK105703</b>	<b>MK105704</b>	<b>MK105705</b>
<i>Lactarius subgennensis*</i>	<i>Lactarius subgennensis*</i>	Cyprus	ML21152E (JMV800627 (BCN))	<b>This work</b>	—	—	<b>MK102747</b>
<i>Lactarius subgennensis</i>	<i>Lactarius subgennensis</i>	Cyprus	ML611322 (pers. herb.)	<b>This work</b>	DQ422201	DQ421934	DQ421934
<i>Lactarius subsericeus</i>	<i>Lactarius subsericeus</i>	Cyprus	ML411272 (pers. herb.)	<b>This work</b>	KF133313	KF133345	KF133345
<i>Lactarius tenellus</i>	<i>Lactarius tenellus</i>	Sweden	UE11-10-2004-8 (UP-S)	Buijck et al. (2008)	KF133280	KF133281	KF133336
<i>Lactarius thyinos</i>	<i>Lactarius thyinos</i>	Benin	ADK3598 (BR)	Verbeken et al. (2014b)	KF133271	KF133304	KF133344
<i>Lactarius torrinosus</i>	<i>Lactarius torrinosus</i>	Canada	Avoik23-08-2004 (GENT)	Verbeken et al. (2014b)	KF133314	KF133346	KF133346
<i>Lactarius torrinosus</i>	<i>Lactarius torrinosus</i>	Czech Republic	RW3183 (GENT)	Verbeken et al. (2014b)	—	—	—

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Lactarius trivialis</i>	<i>Lactarius trivialis</i>	Sweden	UE27.08.2002-17a (UPS)	Buyck et al. (2008)	DQ421991	DQ421924	
<i>Lactarius uvidus</i>	<i>Lactarius uvidus</i>	Estonia	185041 (TAA)	Tedersoo et al. (2003)	AJ534936	—	
<i>Lactarius vietus</i>	<i>Lactarius vietus</i>	Sweden	UE11.19.2004-1 (UPS)	Verbeeken et al. (2014b)	KF133331	—	
<i>Lactarius zonarius</i>	<i>Lactarius zonarius</i>	France	UE27.09.2002-4 (UPS)	Buyck et al. (2008)	EU278678	EU278679	
<i>Lactarius zonarius</i> var. <i>riparius</i>	<i>Lactarius zonarius</i> var. <i>riparius</i>	USA	CLC2933 (MONT)	Barge et al. (2016)	KX394302	—	
<i>Zelleromyces daucinus*</i>	<i>Zelleromyces daucinus*</i>	Australia	T18311 (OSC)	Miller et al. (2001)	—	AF265543	—
<i>Zelleromyces stratus*</i>	<i>Zelleromyces stratus*</i>	Australia	T18858 (OSC)	Miller et al. (2001)	—	AF265544	—
<i>Lactifluus edulis</i>	<i>Lactarius edulis</i>	Zimbabwe	AV99-041 (GENT)	Buyck et al. (2007)	DQ421977	DQ421916	
<i>Lactifluus emergens</i>	<i>Lactifluus emergens</i>	Zimbabwe	AV99-012 (GENT)	De Crop et al. (2017)	KR364021	KR364276	
<i>Lactifluus longisporus*</i>	<i>Lactarius longisporus*</i>	Zimbabwe	AV94-557 (GENT), BB00.1519 (PC)	Buyck et al. (2007), De Crop et al. (2017)	KR364118	KR364244	
<i>Lactifluus nodosicystidiosus</i>	<i>Lactarius nodosicystidiosus*</i>	Madagascar	BB97-072 (PC)	Buyck et al. (2007)	DQ421976	DQ421915	
<i>Lactifluus phlebophyllus</i>	<i>Lactarius phlebophyllus</i>	Madagascar	BB00-1388 (PC)	Buyck et al. (2007)	DQ421979	DQ421918	
<i>Lactifluus piperatus</i>	<i>Lactarius piperatus</i>	Sweden	UE09.08.2004-6 (UPS)	Buyck et al. (2008)	DQ422035	DQ421937	
<i>Lactifluus velleoreus</i>	<i>Lactarius velleoreus</i>	Sweden	UE20.09.2004-22 (UPS)	Buyck et al. (2007)	DQ422034	DQ421936	
<i>Lactifluus velutissimus</i>	<i>Lactarius velutissimus</i>	Zimbabwe	AV99-185 (GENT)	Buyck et al. (2007)	DQ421973	DQ421912	
<i>Lactifluus volentus</i>	<i>Lactarius volentus</i>	Sweden	UE09.08.2004-5 (UPS)	Buyck et al. (2008)	DQ422008	DQ421932	
<i>Multifurca furcata</i>	<i>Lactarius furcatus</i>	Costa Rica	RH7804 (NY)	DQ421994	DQ421927	DQ421940	
<i>Multifurca ochrocompacta</i>	<i>Russula ochrocompacta</i>	USA	BB02.107 (PC)	DQ421984	DQ421984	DQ421940	
<i>Multifurca stenophylla</i>	<i>Multifurca stenophylla</i>	Australia	CWD584 (MEL)	JX266628	JX266633	—	
<i>Multifurca zonaria*</i>	<i>Russula zonaria*</i>	Thailand	DED7442 (PC)	DQ421990	DQ421942	DQ421942	
<i>Russula</i> basal clades							
<i>Russula acrifolia</i>		USA	r-05065	JF834363	—	JF834460	
<i>Russula albonigra</i>		Sweden	AT2002064 (UPS)	DQ422029	DQ422029	DQ421966	
<i>Russula archaea</i>		USA	BB2004-255 (PC)	EU598165	—	—	
<i>Russula cf. brevipes</i>		Canada	F30230 (UBC)	KX812841	KX813647	KX813647	
<i>Russula camarophylla</i>		France	PAM01081108 (PC)	DQ421982	DQ421982	DQ421938	
<i>Russula cascadensis</i>		Canada	F23910 (UBC)	KJ146726	JF834478	JF834427	
<i>Russula aff. chloroides</i>		USA	r-01002	JF834332	KT933890	KT933890	
<i>Russula compacta</i>		USA	BPL242 (TENN)	KT933860	KX812861	KX812861	
<i>Russula crassutinata</i>		Canada	F30159 (UBC)	FH12-272 (GENT)	KF432955	KR364340	
<i>Russula delica</i>		Belgium	WCR00-412 (PC)	DQ422025	DQ422025	DQ421963	
<i>Russula earlei</i>		USA	UE28.09.2002-4 (UPS)	DQ421983	DQ421983	DQ421939	
<i>Russula farinipes</i>		France	12221S87, PAM93071601 (pers. herb.)	AY061702	—	—	
<i>Russula litoralis</i>		Europe	MCA1856 (pers. herb.)	JN168745	—	JN168745	
<i>Russula metachromatica</i>		French Guiana	UE20.09.2004-07 (UPS)	DQ422010	DQ422010	DQ421952	
<i>Russula nigricans</i>		Sweden	146/2002 (TUR)	DQ421987	DQ421987	DQ421941	
<i>Russula pallens</i>		Norway	JV02-218 (C)	DQ422032	DQ422032	—	
<i>Russula pallidospora</i>		Denmark	BB07.134 (PC)	KP033486	KP033497	KP033508	
<i>Russula polypyilla</i>		USA	T14771 (MEL, OSC)	EU019931	EU019931	—	
<i>Russula pumicoides*</i>		Australia	H4755 (HO)	EU019943	—	—	
<i>Russula sinuata*</i>		Australia	—	—	—	—	
<i>Ingrateirigidae russula</i> s.str.							
<i>Russula</i> sp.		Iran	FR852097	—	—	—	
<i>Russula</i> sp.		Estonia	FN669244	—	—	—	
<i>Russula</i> sp.		China	JF273537	—	—	—	
<i>Russula</i> sp.		China	JN129407	—	—	—	
<i>Russula</i> sp.		Pakistan	KR082870	—	—	—	
<i>Russula</i> sp.		USA	KU886598	—	—	—	
<i>Russula</i> sp.		USA	JF834493	JF834493	JF834493	JF834493	

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Russula</i> sp. *	<i>Gymnomyces</i> sp.	USA	T13003 (OSC)	Whitbeck (2003)	AY239337	—	—
	<i>Gymnomyces</i> sp.	USA	T13006 (OSC)	Whitbeck (2003)	AY239340	—	—
	<i>Gymnomyces</i> sp.	USA	1059 (SOC)	Gladish et al. (2010)	FJ789600	—	—
	<i>Gymnomyces</i> sp.	USA	T35827 (OSC)	Frank (unpubl.)	KF386763	—	—
	<i>Martellia</i> sp.	USA	TK1719, SNF35	Izzo et al. (2005)	AY558757	—	—
	<i>Russula adsphearocephalifaris</i> *	China	TK165	Izzo et al. (2005)	AY558788	KY270493	—
	<i>Russula aeruginosa</i>	China	FAN492 (BJTC)	Sang et al. (2016)	KY270488	—	—
	<i>Russula aeruginosa</i>	Sweden	FAN448 (BJTC)	Sang et al. (2016)	KY270489	—	—
	<i>Russula albobrunnea</i>	Australia	AT200317 (UPS)	Buyck et al. (2008)	DQ421999	DQ421946	—
	<i>Russula alutacea</i>	Italy	TL2136 (MEI)	Lebel & Tonkin (2007)	EU019933	EU019933	—
	<i>Russula amethystina</i>	Germany	6472 (MCVE)	Osmundson et al. (2013)	JF908676	—	—
	<i>Macowanites ammophilus</i>	Portugal	hue215 (TUB)	Eberhardt (2002)	AY606971	—	—
	<i>Gymnomyces ammophilus</i>	Portugal	42956 (AH)	This work	<b>MK105622</b>	—	—
	<i>Macowanites ammophilus</i>	Portugal	Fungi 40132 (MA)	Calonge & Martin (2000)	AF230890	—	—
	<i>Macowanites ammophilus</i>	Portugal	Fungi 40137 (MA)	Calonge & Martin (2000)	AF230891	—	—
	<i>Macowanites ammophilus</i>	Spain	Fungi 51167 (MA)	Vidal et al. (2002)	AJ438036	—	—
	<i>Macowanites ammophilus</i>	Spain	Fungi 51165 (MA)	Vidal et al. (2002)	AJ438038	—	—
	<i>Macowanites ammophilus</i>	Spain	Fungi 51166 (MA)	Vidal et al. (2002)	AJ438037	—	—
	<i>Macowanites ammophilus</i>	Spain	43950 (AH)	This work	<b>MK105623</b>	—	—
	<i>Macowanites ammophilus</i>	Spain	46370 (AH)	This work	<b>MK105624</b>	<b>MK105706</b>	<b>MK102748</b>
	<i>Russula ammophila</i>	Europe	IC09010703 (BCN)	This work	<b>MK112566</b>	<b>MK108033</b>	—
	<i>Russula amoenicolor</i>	USA	311IX76, BBB9.823 (PC)	Miller & Buyck (2002)	AY061655	—	—
	<i>Russula cf. amoenolens</i>	USA	CDW122	Avis (2012)	JQ622327	—	—
	<i>Russula cf. amoenolens</i>	USA	PRL4149	Bazzicalupo et al. (2017)	KF245512	—	—
	<i>Russula cf. amoenolens</i>	USA	12794 (MICH)	Loney et al. (2016)	KT933954	KT933884	—
	<i>Russula cf. amoenolens</i>	France	BPL232 (TENN)	Bazzicalupo & Berbee (unpubl.)	KF245510	—	—
	<i>Russula amoena</i>	Spain	12838 (MICH)	This work	<b>MK105625</b>	<b>MK102749</b>	<b>MK102750</b>
	<i>Russula amoena</i>	Spain	46371 (AH)	This work	<b>MK105626</b>	<b>MK105707</b>	<b>MK102264</b>
	<i>Russula amoena</i>	New Zealand	46372 (AH)	This work	<b>MK105627</b>	<b>MK105708</b>	<b>MK102751</b>
	<i>Gymnomyces sublevisporus</i> *	Spain	77763 (PDD)	This work	<b>MK105628</b>	<b>MK105709</b>	<b>MK102752</b>
	<i>Gymnomyces sublevisporus</i>	Spain	BM360 (pers. herb.)	This work	<b>MK105629</b>	<b>MK105709</b>	<b>MK102753</b>
	<i>Russula aquosa</i>	Spain	39198 (AH)	This work	<b>MK105629</b>	<b>MK105709</b>	<b>MK102753</b>
	<i>Gymnomyces fragrans</i>	Spain	39239 (AH)	Bazzicalupo et al. (2017)	KX812873	KX812873	KX812873
	<i>Russula atrorubens</i>	USA	101708 (TU)	Whitbeck (2003)	AY239331	—	—
	<i>Russula atroglauca</i>	Estonia	PNW 5607 (OSC)	Bazzicalupo et al. (2017)	KX579812	—	—
	<i>Russula atroglauca</i>	Poland	101718 (TU)	Trocha & Rudy (unpubl.)	KM085418	—	—
	<i>Russula atrovirens</i>	New Zealand	ID PAN 248	Johnston & Park (unpubl.)	GU222260	—	—
	<i>Russula aurantiolammas</i>	Europe	77744 (PDD)	Adamčík et al. (2016)	KU928167	KX812878	KX812878
	<i>Russula aurea</i>	Estonia	r3245	Bazzicalupo et al. (2017)	JN944002	JN940591	JN940591
	<i>Russula aurea</i>	Italy	BB537/08.668 (PC)	Schoch et al. (2012)	JF834520	JF834520	JF834520
	<i>Russula aff. betularum</i>	USA	r-09003	Davis (unpubl.)	KY270487	KY270487	KY270487
	<i>Russula brevilocystidiata</i> *	China	FAN45 (BJTC)	Sang et al. (2016)	JF834341	JF834489	JF834489
	<i>Russula bruneolea</i>	USA	r-03034	Davis (unpubl.)	EU019945	—	—
	<i>Russula brunneonigra</i> *	Australia	H5813 (DAR)	Lebel & Tonkin (2007)	AF325297	AF325297	AF325297
	<i>Russula caerulea</i>	Germany	hue146, 5041S77 (TUB)	Eberhardt (2002)	AY239333	AY239333	AY239333
	<i>Martellia californica</i>	USA	T16027 (OSC)	Whitbeck (2003)	<b>MK105630</b>	<b>MK105709</b>	<b>MK102752</b>
	<i>Elasmomyces mattoianus</i>	Germany	GG171 (M), JM799894 (BCN)	This work	<b>MK105631</b>	<b>MK105710</b>	<b>MK105711</b>
	<i>Russula candida</i>	Spain	JC1710011N (pers. herb.)	This work	<b>MK105632</b>	<b>MK105711</b>	<b>MK105712</b>
	<i>Russula candida</i>	Spain	JMV20100724b (BCN)	This work	<b>MK105633</b>	<b>MK105712</b>	<b>MK105712</b>
	<i>Macowanites candidus</i>	Italy	ELG930704-1, JM/V800185 (BCN)	This work	<b>MK105634</b>	<b>MK105712</b>	<b>MK105712</b>
	<i>Russula candidissima</i>	Poland	F-2009-57 (KRA)	This work	<b>MK105634</b>	<b>MK105712</b>	<b>MK105712</b>

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Russula candidissima</i> * (cont.)	<i>Russula candidissima</i>	Poland	F-2017-2 (KRA)	This work	MK105635	—	—
	<i>Russula candidissima</i>	Spain	JMV20110906-6a (BCN)	This work	MK105636	<b>MK105713</b>	<b>MK102753</b>
	<i>Russula cerea</i>	Spain	JMV20000817-1 (BCN)	This work	MK105637	<b>MK105714</b>	—
	<i>Russula cerea</i>	Germany	JMV800660 (BCN)	This work	MK105638	<b>MK105715</b>	—
	<i>Russula cerea</i>	Poland	F-2012-28 (KRA)	This work	MK105639	—	<b>MK102754</b>
	<i>Gymnomyces xanthosporus</i>	Spain	JMV961003-8 (BCN)	This work	MK105640	—	—
	<i>Russula cerea</i>	Spain	JMV201160705 (BCN)	This work	MK105641	—	—
	<i>Russula cerea</i>	Spain	JMV800656 (BCN)	This work	MK105642	—	—
	<i>Gymnomyces</i> sp.	U.K.	LM1492	Suz et al. (2014)	KM576488	—	—
	<i>Russula cerolens</i>	China	HE2720	Sun et al. (unpubl.)	KC505578	—	—
	<i>Russula cerolens</i>	China	C3	Chai (unpubl.)	JX434671	—	—
	<i>Russula cessans</i>	Slovakia	BB52507-219 (PC)	Schoch et al. (2012)	JN940595	JN993601	—
	<i>Russula cheelii</i>	Australia	JE11002 (MEL)	Lebel et al. (2013)	JX266638	JX266623	—
	<i>Russula chiuif</i> *	China	250410 (HMAS)	Li et al. (2015)	KF225491	—	—
	<i>Macowanites chlorinosmus</i>	USA	36339 (OSC)	Gordon (unpubl.)	KT968633	KT968633	—
	<i>Macowanites chlorinosmus</i>	USA	51028 (OSC)	Gordon (unpubl.)	KT968632	KT968632	—
	<i>Russula clariana</i>	Europe	492RUS26, BB2733 (PC)	Miller & Buyck (2002)	AY061664	—	—
	<i>Russula clarolava</i>	Germany	FH12-212 (GENT)	Looney et al. (2016)	KT933985	KT933929	—
	<i>Russula consobrina</i>	Estonia	101714 (TU)	Bazzicalupo et al. (2017)	KX812874	KX813655	—
	<i>Russula corallina</i>	USA	BB229/06-324 (PC)	Schoch et al. (2012)	JN940605	JN993609	—
	<i>Russula cremeolilacina</i>	Guadeloupe (FR)	BA02-11-04	Sene et al. (2015)	KT933963	KT933822	—
	<i>Russula crustosa</i>	USA	BPI251 (TENN)	Looney et al. (2016)	KT933971	KT933942	—
	<i>Russula cuprea</i>	Germany	FH12-250 (GENT)	Looney et al. (2016)	KT933955	KT933927	—
	<i>Russula curtipes</i>	Germany	FH12-206 (GENT)	Looney et al. (2016)	KT933976	KT933908	—
	<i>Russula cf. cyanoxantha</i>	USA	BPI280 (TENN)	De Crop et al. (2017)	KR364093	KR364225	—
	<i>Russula cyanoxantha</i>	Germany	FH12-201 (GENT)	Looney et al. (2016)	KT933992	KT933924	—
	<i>Russula decolorans</i>	Germany	FH12-196 (GENT)	Osmundson et al. (2013)	JF908710	—	—
	<i>Russula dyadicola</i>	France	16243 (MCVE)	Adamčík et al. (2016)	KU928146	KY616724	—
	<i>Russula dyadicola</i>	Finland	151632 (TURA)	Whitbeck (2003)	AY239304	—	—
	<i>Gymnomyces ellipsosporus</i>	USA	T61608 (OSC)	Whitbeck (2003)	AY239306	DQ421943	—
	<i>Gymnomyces ellipsosporus</i>	USA	58973 (OSC)	Eberhardt (2002)	AF323506	—	—
	<i>Russula emetica</i>	Sweden	UE05-10-2003-11 (UPS)	Qiao et al. (unpubl.)	KX655858	DQ421997	—
	<i>Russula exalticans</i>	Germany	n179/3 (TUB)	Loney et al. (2016)	KT933989	KT933921	—
	<i>Russula faustiana</i>	China	XJ2013-11-00408	Buyck et al. (2008)	DQ422017	DQ421958	—
	<i>Russula felaea</i>	Germany	FH12-185 (GENT)	Kong et al. (2015)	EU598162	EU598162	—
	<i>Russula firmula</i>	Sweden	AT2004142 (UPS)	Eberhardt (2002)	AF418613	AF322599	—
	<i>Russula cf. flavisiccans</i>	USA	BB2004-254 (PC)	Loney et al. (2016)	KT933864	KT933935	—
	<i>Russula foetens</i>	Germany	hue124 (TUB)	Lebel & Tonkin (2007)	EU019936	EU019936	—
	<i>Russula fontqueri</i>	Australia	FH12-223 (GENT)	This work	<b>MK105643</b>	<b>MK105716</b>	<b>MK102755</b>
	<i>Russula galbana</i> *	Israel	H4667 (MEL)	This work	<b>MK105644</b>	<b>MK105717</b>	—
	<i>Macowanites galileensis</i>	Israel	G-83 (HA)	G-201 (HA)	<b>MK105645</b>	—	—
	<i>Macowanites galileensis</i>	Israel	G-166 (HA)	G-202 (HA)	<b>MK105646</b>	—	—
	<i>Macowanites galileensis</i>	Israel	G-209 (HA)	G-209 (HA)	<b>MK105647</b>	—	—
	<i>Cystangium</i> sp.	Chile	T26316 (OSC)	Trievelier-Pereira et al. (2015)	KF819810	KF819810	—
	<i>Gymnomyces monosporus</i>	USA	117360 (OSC)	Gordon & Zych (unpubl.)	EU669222	EU669222	—
	<i>Russula gillkeyae*</i>	USA	T2572 (OSC)	Whitbeck (2003)	AY239346	AY239346	—
	<i>Russula globispora</i>	Germany	FH2007-BT111 (GENT)	Adamčík et al. (2016)	KU928144	KU928144	—
	<i>Russula gracilina</i>	Germany	FH2007-BT121 (GENT)	Adamčík et al. (2016)	KU886594	KU886594	—
	<i>Russula granulata</i>	Sweden	UE23-08-2004-14 (PC)	Buyck et al. (2008)	DQ422004	DQ422004	—
		USA	BPL272 (TENN)	Looney et al. (2016)	KT933832	KT933832	—

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher id (Herbarium) <sup>2</sup>	GenBank accession codes			
				ITS rDNA	28S rDNA	RPB2	
<i>Russula grisea</i>	<i>Russula grisea</i>	Sweden	UE2005_08_16-01 (UPS)	Buyck et al. (2008)	DQ422030	DQ421968	
<i>Russula heterophylla</i>	<i>Russula heterophylla</i>	Sweden	UE20_08_2004-2 (UPS)	Buyck et al. (2008)	DQ422006	DQ421951	
<i>Russula hemisphaerica</i>	<i>Russula hemisphaerica</i>	Cameroun	12031(S83, SLM (pers. herb.)	Kleine et al. (2013)	JQ902080	—	
<b><i>Russula hobartiae*</i></b>	<i>Russula hobartiae</i>	Cyprus	ML4193GY (pers. herb.)	<b>This work</b>	<b>MK105648</b>	<b>MK102756</b>	
<i>Russula hobartiae</i>	<i>Russula hobartiae</i>	Cyprus	ML4116GY (pers. herb.)	<b>This work</b>	<b>MK105649</b>	<b>MK102757</b>	
<i>Russula hobartiae*</i>	<i>Russula hobartiae*</i>	Cyprus	ML10131GY, JMV800628 (BCN)	<b>This work</b>	<b>MK105650</b>	<b>MK102758</b>	
<i>Russula ilicis</i>	<i>Russula ilicis</i>	Cyprus	GK5889, JMV800647 (BCN)	Miller & Buyck (2002)	<b>MK105651</b>	<b>MK102759</b>	
<i>Russula illoota</i>	<i>Russula illoota</i>	Europe	5631C52, MS1018/99 (pers. herb.)	Buyck et al. (2008)	AY061682	DQ422024	DQ421967
<i>Russula inamoena</i>	<i>Russula inamoena</i>	Sweden	UE26_07_2002-3 (UPS)	Buyck et al. (2008)	DQ422024	DQ421967	
<i>Russula inflata</i>	<i>Russula inflata</i>	Italy	107	Mejera et al. (2017)	KJ834597	—	
<i>Russula insignis</i>	<i>Russula insignis</i>	Madagascar	SLM00/344	Kleine et al. (2013)	JQ902062	—	
<i>Russula integra</i>	<i>Russula integra</i>	Europe	12231S85, BB00_2149 (PC)	Miller & Buyck (2002)	AY061700	KT933916	
<i>Russula intervenosa*</i>	<i>Russula intervenosa*</i>	Germany	FH12-172 (GENT)	Looney et al. (2016)	KT933984	—	
<i>Russula kanadii</i>	<i>Russula kanadii</i>	India	AM273 (CUH), 1272 (CAL)	Crous et al. (2016)	KT824241	—	
<i>Russula kermesina*</i>	<i>Russula kermesina*</i>	India	AM086 (CUH)	Dutta et al. (2015)	KJ866936	—	
<i>Russula larinicina</i>	<i>Russula larinicina</i>	New Zealand	92049 (PDD)	Johnston & Park (unpubl.)	GU223233	—	
<i>Russula larinocaffinis</i>	<i>Russula larinocaffinis</i>	Italy	BB575/08_681 (PC)	Schoch et al. (2012)	JN940593	KU237846	
<i>Russula laurocerasi</i>	<i>Russula laurocerasi</i>	Italy	662 (MCVE)	Osmundson et al. (2013)	JF908646	—	
<i>Russula lepida</i>	<i>Russula lepida</i>	Chile	T26311 (OSC)	Trienveiler-Pereira et al. (2015)	KF198111	KT933920	
<i>Russula liliacea</i>	<i>Russula liliacea</i>	Germany	FH12-178 (GENT)	Looney et al. (2016)	KT933988	—	
<i>Russula livescens</i>	<i>Russula livescens</i>	Belgium	HAB9990 (UPS)	Buyck et al. (2008)	DO422013	DQ421954	
<i>Russula longistigmata*</i>	<i>Russula longistigmata*</i>	Slovakia	BB435/07_213 (PC)	Schoch et al. (2012)	JN940592	JN993610	
<i>Russula maculata</i>	<i>Russula maculata</i>	China	F0177	Xie et al. (2010)	GU371295	—	
<i>Russula mariae</i>	<i>Russula mariae</i>	China	T262665 (OSC)	Trienveiler-Pereira et al. (2015)	KF813808	—	
<i>Russula mariae</i>	<i>Russula mariae</i>	Chile	FH2010-B1784 (GENT)	Adamäk et al. (2016)	KU928154	KT933945	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Germany	FH12-262 (GENT)	Looney et al. (2016)	KT933874	—	
<i>Russula mariae</i>	<i>Russula mariae</i>	Korea	19111 (HCCN)	Park et al. (2013)	KF361762	KF361712	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Korea	20120922-08 (SFC)	Park et al. (2013)	KF361778	KF361728	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Greece	GK3901, JMV800638 (BCN)	<b>This work</b>	<b>MK105652</b>	<b>MK102758</b>	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Greece	GK8136, JMV800644 (BCN)	<b>This work</b>	<b>MK105653</b>	<b>MK102758</b>	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Poland	F-2012-153 (KRA)	<b>This work</b>	<b>MK105654</b>	<b>MK102758</b>	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Poland	F-2017-1 (KRA)	<b>This work</b>	<b>MK105655</b>	<b>MK102758</b>	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Poland	F-2018-1 (KRA)	<b>This work</b>	<b>MK105656</b>	<b>MK102758</b>	
<i>Russula matitroiana</i>	<i>Russula matitroiana</i>	Poland	F-2018-2 (KRA)	<b>This work</b>	<b>MK105657</b>	<b>MK102758</b>	
<i>Gymnomyces compactus</i>	<i>Gymnomyces compactus</i>	USA	T13171 (OSC)	Whitbeck (2003)	AY238303	—	
<i>Russula mediterraneensis</i>	<i>Russula mediterraneensis</i>	USA	T13565 (OSC)	Whitbeck (2003)	AY238342	—	
<i>Russula mediterraneensis</i>	<i>Russula mediterraneensis</i>	Greece	GK3930, JMV800639 (BCN)	<b>This work</b>	<b>MK105658</b>	<b>MK102755</b>	
<i>Russula megaseudocyathidata</i> *	<i>Russula megaseudocyathidata</i> *	Greece	GK7286, JMV800642 (BCN)	<b>This work</b>	<b>MK105659</b>	<b>MK102756</b>	
<i>Russula matsmili**</i>	<i>Russula matsmili**</i>	Australia	GM6710, JMV800641 (BCN)	<b>This work</b>	<b>MK105660</b>	<b>MK102760</b>	
<b><i>Russula mediterraneensis*</i></b>	<i>Russula mediterraneensis*</i>	Italy	MG630, 29085 (MCVE)	<b>This work</b>	<b>MK105661</b>	<b>MK102761</b>	
<i>Russula megaseudocyathidata</i> *	<i>Russula megaseudocyathidata</i> *	China	MG636, 29086 (MCVE)	<b>This work</b>	<b>MK105662</b>	<b>MK102769</b>	
<i>Gymnomyces megasporus</i>	<i>Gymnomyces megasporus</i>	Australia	FAN454 (BJTC)	Sang et al. (2016)	KY270488	KY270491	
<i>Zelleromyces meridionalis*</i>	<i>Zelleromyces meridionalis*</i>	Spain	T18717 (OSC)	Miller et al. (2001)	AF265535	—	
<i>Russula meridionalis</i>	<i>Russula meridionalis</i>	Spain	BM410 (pers. herb.)	<b>This work</b>	<b>MK105663</b>	<b>MK102762</b>	
<i>Russula meridionalis</i>	<i>Russula meridionalis</i>	Spain	IC20051417 (BCN)	<b>This work</b>	<b>MK105664</b>	<b>MK102762</b>	
<i>Russula meridionalis</i>	<i>Russula meridionalis</i>	Spain	IC24051506 (BCN)	<b>This work</b>	<b>MK105665</b>	<b>MK102768</b>	
<i>Russula messapica</i>	<i>Russula messapica</i>	Italy	JC180617NR (pers. herb.)	<b>This work</b>	<b>MK105667</b>	<b>MK102769</b>	
<i>Russula messapica</i>	<i>Russula messapica</i>	Spain	5621C52, MS61189 (pers. herb.)	<b>This work</b>	<b>MK105668</b>	<b>MK102763</b>	
<i>Russula messapica</i>	<i>Russula messapica</i>	Spain	46373 (AH)	<b>This work</b>	<b>MK105669</b>	<b>MK102764</b>	
<i>Russula messapica</i>	<i>Russula messapica</i>	Spain	JL20111182 (BCN)	<b>This work</b>	<b>MK105730</b>	<b>MK102764</b>	
<i>Russula messapica</i>	<i>Russula messapica</i>	Spain	IC28110613 (BCN)	<b>This work</b>	<b>MK105666</b>	<b>MK102764</b>	

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Russula messapica</i> var. <i>messapicoides</i> *	<i>Russula messapica</i> var. <i>messapicoides</i>	Greece	VK2998 (pers. herb.)	This work	<b>MK105670</b>	<b>MK105731</b>	<b>MK102765</b>
<i>Russula messapica</i> var. <i>messapicoides</i>	<i>Russula messapica</i>	Greece	VK3368 (pers. herb.)	This work	<b>MK105671</b>	<b>MK105732</b>	<b>MK102766</b>
<i>Russula messapica</i> var. <i>messapicoides</i>	<i>Russula messapica</i>	Greece	VK3411, JMV800682 (BCN)	This work	<b>MK105672</b>	–	–
<i>Russula messapica</i> var. <i>messapicoides</i>	<i>Russula messapica</i>	Greece	GK934-1, JMV800645 (BCN)	This work	<b>MK105673</b>	–	–
<i>Russula messapica</i> var. <i>messapicoides</i>	<i>Macowanites messapicoides*</i>	Greece	JL1493 (BCN)	This work	<b>MK105674</b>	–	–
<i>Russula messapica</i> var. <i>messapicoides</i>	<i>Russula messapicoides*</i>	Spain	JMV800652 (BCN)	This work	<b>MK105675</b>	–	–
<i>Russula misiformis</i>	<i>Gymnomyces misiformis</i>	Greece	AM1653 (pers. herb.)	Whitbeck (2003)	AY472079	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Italy	JMV800125 (BCN)	This work	<b>MK105676</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Italy	JC170305NR, JMV800661 (BCN)	This work	<b>MK105677</b>	–	<b>MK102767</b>
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JC131117NR, JMV800662 (BCN)	This work	<b>MK105678</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JC101206BT, JMV800663 (BCN)	This work	<b>MK105679</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JM8-H-69, JMV971127 (BCN)	This work	<b>MK105680</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JC110-102NR (pers. herb.)	This work	<b>MK105681</b>	–	–
<i>Martellia pila</i>	<i>Russula misiformis</i>	Spain	JMV920626-1 (BCN)	This work	<b>MK105682</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JMV20140607-3 (BCN)	This work	<b>MK105683</b>	–	–
<i>Russula misiformis</i>	<i>Russula misiformis</i>	Spain	JMV20160524-1 (BCN)	This work	<b>MK105684</b>	–	–
<i>Martellia mediterranea*</i>	<i>Russula misiformis</i>	Spain	GM-RG11057 (AH)	This work	<b>MK105685</b>	–	–
<i>Russula monospora</i>	<i>Russula monospora</i>	Spain	46459 (AH)	This work	<b>MK105686</b>	–	–
<i>Russula monospora</i>	<i>Russula monospora</i>	Spain	JMV800671 (BCN)	This work	<b>MK105687</b>	–	–
<i>Russula monospora</i>	<i>Russula monospora</i>	Spain	JMV800672 (BCN)	This work	<b>MK105688</b>	–	–
<i>Russula monospora</i>	<i>Russula monospora</i>	Spain	JMV800673 (BCN)	This work	<b>MK105689</b>	–	–
<i>Gymnomyces monticola</i>	<i>Gymnomyces monticola</i>	USA	56167 (OSC)	Whitbeck (2003)	AY239313	JF834500	JF834449
<i>Russula murrillii</i>	<i>Russula murrillii</i>	USA	r-04075	Davis (unpubl.)	KT934005	KX578009	KT933937
<i>Russula mustelina</i>	<i>Russula mustelina</i>	Germany	FH-12-226 (GENT)	Looney et al. (2016)	KT933866	KX578009	KT933937
<i>Russula nana</i>	<i>Russula nana</i>	Germany	101701 (TU)	Wood et al. (unpubl.)	–	–	–
<i>Russula nauseosa</i>	<i>Russula nauseosa</i>	Australia	FH12-173 (GENT)	Looney et al. (2016)	KT933985	KT933986	KT933917
<i>Russula neerimea</i>	<i>Russula neerimea</i>	USA	2101871 (MEL)	Lebel & Tonkin (2007)	EU019915	EU019915	–
<i>Russula nitida</i>	<i>Russula nitida</i>	USA	BB2004-272 (PC)	Hughes & Buyck (unpubl.)	EU598164	EU598164	EU598164
<i>Russula nitida</i>	<i>Russula nitida</i>	Germany	FH12-218 (GENT)	Looney et al. (2016)	KT934001	KT933862	KT933933
<i>Gymnomyces nondistincta</i>	<i>Gymnomyces nondistincta</i>	USA	62139 (OSC)	Gordon (unpubl.)	KP859276	KP859294	–
<i>Cystangiolum sp.</i>	<i>Cystangiolum sp.</i>	USA	T26350 (OSC)	Trievilleir-Pereira et al. (2015)	KU928159	KU928159	–
<i>Russula ochroleuca</i>	<i>Russula ochroleuca</i>	Chile	HM-R-9702 (M)	Adamicik et al. (2016)	KT933996	KT933987	KT933928
<i>Russula nympharum</i>	<i>Russula nympharum</i>	Germany	FH12-211 (GENT)	Looney et al. (2016)	KT933953	KT933812	KT933883
<i>Russula cf. ochrophylla</i>	<i>Russula cf. ochrophylla</i>	USA	BPL231 (TENN)	Buyck et al. (2008)	DQ422012	DQ421953	DQ421945
<i>Russula ochrospora</i>	<i>Russula ochrospora</i>	Italy	GD20-07-2004 (UPS)	Schoch et al. (2012)	JN944010	JN993603	JN940607
<i>Russula odorata</i>	<i>Russula odorata</i>	Slovakia	BB526-07-186 (PC)	Smith et al. (2006)	AF265536	–	–
<i>Gymnomyces redolens</i>	<i>Gymnomyces redolens</i>	New Zealand	SLM4162 (pers. herb.)	Looney et al. (2016)	KT934001	KT933861	KT933932
<i>Russula paludosa</i>	<i>Russula paludosa</i>	Sweden	FH12-216 (GENT)	Buyck et al. (2008)	DQ422007	DQ422007	DQ421945
<i>Russula parazurea</i>	<i>Russula parazurea</i>	USA	BW06-09-2002-16/MF01.10.2003 (UPS)	Whitbeck (2003)	–	–	–
<i>Gymnomyces parksii</i>	<i>Gymnomyces parksii</i>	India	T14997 (OSC)	Das et al. (2017)	MF535185	–	–
<i>Russula pauperiensis</i>	<i>Russula pauperiensis</i>	USA	1395 (CAL)	Looney et al. (2016)	KT933830	KT933901	KT933901
<i>Russula peckii</i>	<i>Russula peckii</i>	USA	BPL270 (TENN)	Buyck et al. (2008)	DQ422019	DQ422019	DQ421960
<i>Russula pectinatoides</i>	<i>Russula pectinatoides</i>	USA	BPL276 (TENN)	Whitbeck (2003)	–	–	–
<i>Russula pericina</i>	<i>Russula pericina</i>	Sweden	UE21.09.2003-01 (UPS)	Looney et al. (2016)	<b>MK105690</b>	<b>MK105691</b>	<b>MK102768</b>
<i>Russula pilosa</i> *	<i>Russula pilosa</i> *	Spain	JMV970816-8 (BCN)	Buyck et al. (2008)	–	–	–
<i>Russula pilosa</i>	<i>Russula pilosa</i>	Spain	JMV800654 (BCN)	This work	–	–	–
<i>Russula aff. pilosella*</i>	<i>Russula aff. pilosella*</i>	Australia	H4784 (MEL)	Lebel & Tonkin (2007)	EU019932	EU019941	EU019941
<i>Russula pilosella*</i>	<i>Russula pilosella*</i>	Australia	H5974 (BR)	Lebel & Tonkin (2007)	–	–	–

Table 1 (cont.)

Species (*) sequestrate <sup>1</sup>	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Russula praeterita</i>	<i>Russula praeterita</i> *	Italy	1997/0812 (IB)	Sannari & Eberhardt (unpubl.)	UDB019331	—	—
<i>Russula pseudocarminea</i>	<i>Russula pseudocarminea</i>	Burundi	BB5401 (PC)	Kleinie et al. (2013)	JQ902068	—	—
<i>Russula pseudopectinatoidea</i>	<i>Russula pseudopectinatoidea*</i>	China	251523 (HMAS)	Li et al. (2015)	KM269077	—	—
<i>Russula puelaria</i>	<i>Russula puelaria</i>	Estonia	101839 (TU)	Bazzicalupo et al. (2017)	UDB015695	KX812887	KX813668
<i>Russula puelaria</i>	<i>Russula puelaria</i>	Europe	2-11301876, RW27065 (E)	Miller & Buyck (2002)	AY061710	—	KY616697
<i>Russula puligera</i>	<i>Russula puligera</i>	Slovakia	F-3107 (SAV)	Cabon et al. (unpubl.)	KY582704	KJ786689	—
<i>Russula pulverulenta</i>	<i>Russula pulverulenta</i>	French Guiana	G3130	Roy et al. (2016)	KJ786592	AY061736	—
<i>Russula punicea</i>	<i>Russula punicea</i>	USA	4-11441879, RF566 (pers. herb.)	Miller & Buyck (2002)	JF834441	JF834441	—
<i>Russula purpureoflava</i>	<i>Russula purpureoflava</i>	USA	r-03043	Davis (unpubl.)	JF834344	JX266641	—
<i>Russula putida</i>	<i>Russula putida</i> *	Australia	JET1128 (MEL)	Lebel et al. (2013)	HG798527	HG798526	—
<i>Russula risigallina</i>	<i>Russula risigallina</i>	Italy	1987/0791 (IB)	Borovicka (unpubl.)	DQ422022	DQ422022	—
<i>Russula romellii</i>	<i>Russula romellii</i>	Sweden	UE03/07.2003-08 (UPS)	Buyck et al. (2008)	KT933887	KT933919	—
<i>Russula rosea</i>	<i>Russula rosea</i>	Germany	FH12-177 (GENT)	Looney et al. (2016)	JN944003	JN993613	—
<i>Russula roseipes</i>	<i>Russula roseipes</i>	France	BB430/07.780 (PC)	Schoch et al. (2012)	UDB015972	KX812884	KX813665
<i>Russula rostratycystidea</i>	<i>Russula rostratycystidea*</i>	Estonia	101806 (TU)	Bazzicalupo et al. (2017)	EU019838	—	—
<i>Russula rubellipes</i>	<i>Russula rubellipes</i>	Australia	H6165 (BRI)	Lebel & Tonkin (2007)	KT933858	KT933888	—
<i>Russula rubroluteus</i>	<i>Russula rubroluteus*</i>	USA	BP1240 (TENN)	Looney et al. (2016)	KY019940	EU019940	—
<i>Russula rubropunctatissima</i>	<i>Russula rubropunctatissima</i>	Australia	T12610 (PDD, OSC)	Lebel & Tonkin (2007)	KY087583	KY087583	—
<i>Russula rugulosa</i>	<i>Russula rugulosa</i>	Brazil	IDuque55	Duque et al. (unpubl.)	KY848516	KY701373	—
<i>Russula sanguinea</i>	<i>Russula sanguinea</i>	USA	BP1654 (TENN)	Looney et al. (2016)	KT934008	KT933940	—
<i>Russula sarnarii</i>	<i>Russula sarnarii*</i>	Germany	FH12-240 (GENT)	Looney et al. (2016)	KY284154	—	—
<i>Russula seminudum</i>	<i>Cystangium seminudum</i>	India	1395 (CAL)	Ghosh et al. (2017)	KY019947	AF265533	—
<i>Russula sessilis</i>	<i>Russula sessilis</i> *	Australia	H5346 (MEI, OSC)	Lebel et al. (unpubl.)	KY848516	KY848516	—
<i>Russula segeria</i>	<i>Russula segeria</i> *	Brazil	H5038 (OSC, MEI)	Miller et al. (2007), Lebel & Tonkin (2007)	KT934008	KT933940	—
<i>Russula shafferi</i>	<i>Russula shafferi</i> *	USA	29622 (OSC)	Whitbeck (2003)	KY284154	—	—
<i>Russula shingtaensis</i>	<i>Russula shingtaensis</i>	USA	51046 (OSC)	Whitbeck (2003)	KY284154	—	—
<i>Russula sichuanensis</i>	<i>Russula sichuanensis</i>	India	KD11-094 (BSHC)	Das et al. (2014)	KY284154	—	—
<i>Russula similis</i>	<i>Russula similis</i> *	Pakistan	MSM0025 (LAH)	Saba & Khalid (2015)	KY284154	—	—
<i>Russula cf. sonora</i>	<i>Russula cf. sonora</i>	Mexico	GO2009-239 (MEYL)	Kong et al. (2015)	KY284154	—	—
<i>Russula stewartii</i>	<i>Russula stewartii</i> *	Korea	ASIS22640 (HCCN)	Lee et al. (2017)	KY284154	—	—
<i>Russula cf. subfoetens</i>	<i>Russula cf. subfoetens</i>	Korea	2208ASI (HCCN)	Lee et al. (2017)	KX574703	—	—
<i>Russula subfulva</i>	<i>Russula subfulva</i> *	USA	611382 (OSC)	Gordon (unpubl.)	KP859259	KP859279	—
<i>Russula subioculata</i>	<i>Russula subioculata*</i>	USA	611637 (OSC)	Gordon (unpubl.)	KP859261	KT968617	—
<i>Russula subochracea</i>	<i>Russula subochracea*</i>	Spain	JMV94/1022.7 (BCN)	<b>This work</b>	<b>MK105692</b>	—	—
<i>Russula suecica</i>	<i>Russula suecica</i>	USA	T14998 (OSC)	Whitbeck (2003)	AY239321	—	—
<i>Russula tapawera</i>	<i>Macowanites tapawera*</i>	USA	SLM480MC (pers. herb.)	Miller et al. (2001)	GU222263	—	—
<i>Russula tawai</i>	<i>Russula tawai</i>	USA	T13386 (OSC)	Whitbeck (2003)	DQ403804	—	—
<i>Russula theodorou</i>	<i>Cystangium theodorou</i>	Norway	JR7994 (KUO)	Vauras et al. (2016)	KX888014	EU019935	—
<i>Russula tricholomopsis</i>	<i>Russula tricholomopsis</i>	New Zealand	T12607 (OSC), 83696 (PDD)	Lebel & Tonkin (2007)	KX812891	KX813672	—
<i>Russula turci</i>	<i>Russula turci</i>	New Zealand	77760 (PDD)	Johnston & Park (unpubl.)	AY239338	EU019934	—
<i>Russula unicolor</i>	<i>Gymnomyces cinnamomeus</i>	Australia	SLM4384 (pers. herb.)	Smith et al. (2006)	KY582701	KY616695	—
<i>Russula unicolor</i>	<i>Russula variispora*</i>	Finland	77749 (PDD)	Johnston & Park (unpubl.)	KT339269	JF834447	—
<i>Russula venenovskii</i>	<i>Russula venenovskii</i>	Slovakia	101874 (TU)	Bazzicalupo et al. (2017)	JF834498	JN940594	—
<i>Russula venezueliana</i>	<i>Russula venezueliana</i>	French Guiana	T19930 (OSC)	Whitbeck (2003)	KY616695	KY616695	—
<i>Russula versicolor</i>	<i>Russula versicolor</i>	USA	H5855 (DAR)	Lebel & Tonkin (2007)	JF834350	JN940594	—
<i>Russula versicolor</i>	<i>Russula versicolor</i>	Slovakia	F-2921 (SAV)	Cabon et al. (unpubl.)	DQ422018	DQ422018	—
<i>Russula vesca</i>	<i>Russula vesca</i>	Spain	TH7874 (pers. herb.)	Smith et al. (2017)	<b>MK105693</b>	<b>MK105738</b>	—
<i>Russula vidalii</i>	<i>Russula vidalii</i>	Spain	r-04039	Davis (unpubl.)	<b>MK105694</b>	<b>MK105739</b>	—
<i>Russula vinaceodora</i>	<i>Macowanites vinaceodorus</i>	Spain	BB569/07.288 (PC)	Schoch et al. (2012)	Fungi 46524 (MA)	Fungi 46524 (MA)	—

Table 1 (cont.)

Species (*) sequestrate	Original identification (*) type	Origin	Voucher Id (Herbarium) <sup>2</sup>	Reference	GenBank accession codes		
					ITS rDNA	28S rDNA	RPB2
<i>Russula vinaceodora*</i> (cont.)	<i>Macowanites vinaceodorus</i>	Spain	Fungi 47416 (MA)	Vidal et al. (2002)	AJ438034	–	–
	<i>Macowanites vinaceodorus</i>	Spain	46374 (AH)	<b>This work</b>	<b>MK105695</b>	<b>MK105740</b>	<b>MK102770</b>
<i>Russula vinoso</i>	<i>Russula violeipes</i>	Sweden	F124791 (UPS)	Looney et al. (2016)	KX812900	KX8113680	
<i>Russula violeipes</i>	<i>Russula violeipes</i>	Korea	2012/10/06 (SFC)	Park et al. (2013)	KF361808	KF3617758	
	<i>Russula violeipes</i>	Korea	16735 (HCCN)	Park et al. (2013)	KF361783	KF361833	KF361733
	<i>Russula violeipes</i>	Belgium	HU9989 (UPS)	Buyck et al. (2008)	DQ422014	DQ421955	
	<i>Russula violeipes</i>	Canada	F 16576 (UBC)	Berbee et al. (unpubl.)	FJ627039	–	
	<i>Russula viscida</i>	Italy	1997/07/86 (IB)	Eberhardt (unpubl.)	DQ422021	DQ422021	
	<i>Russula wernerii*</i>	Australia	2238232 (MEL)	Lebel & Tonkin (2007)	EU019921	EU019921	
	<i>Russula wollumbina</i>	Canada	OUC97303 (DAV/FP)	Durrall et al. (2006)	DQ367916	DQ367916	
	<i>Russula xerampelina</i>	USA	82212-8, SRC672 (OSC)	Smith et al. (2006)	DQ028473	–	
	<i>Gymnomyces xerophilus*</i>	USA	82212-9, SRC648 (OSC)	Smith et al. (2006)	DQ028476	–	
	<i>Gymnomyces xerophilus</i>	USA					UDP028475

卷之三

STEREACEAE/STEREUM

*Stereum hirsutum* Binder & Hibbett (2002) **FPL 8805**

New species, new combinations and sequences produced from this study are in **bold**. Sequestrate taxa are marked with an asterisk (\*) symbol.

- : Collection sources: AB = Ananad Bâ (BR); AM = Amer Monteiro (pers. herb.); AT = Andrew Taylor (UPS); AV = Annemieke Verbeeken (GENT); AVoik = Andrus Voik (GENT); BH = Bryony M. Horton (HO); BB = Bart Buyck (PO); BM = Baldomero Moreno-Aroyo (pers. herb.); BPL = Brian P. Looney (TENN); BDW = Birgitta Wästberg (UPPS); CLC = Cathly L. Cripps (MONT); DA = David Christopher W. Dunk (ME); DE = Dennis E. Desjardin (PC); DFSU = David G. Barge (MONT); DS = Dirk Stubbe (GENT); EB = Edward G. Barge (MONT); EG = Edith Georges Konstantinidis (pers. herb.); Garay-Serano (XAL); EH = Egon Horak (ZT); EU = Elisabeth Uhlmann (UPS); FA = François Ayer (WSL); FAN = Li Fan (BJTC); FH = Felix Hampe (GENT); FC = Franz Oberwinkler (TUB); GD = Giuseppe Donelli (UPS); GG = Gerhard Gross (M); GK = Georges Konstantinidis (pers. herb.); GM-RG = Gabriel Moreno and Ricardo Galán (AH); GO = Roberto Garbay Oriol (MEXU); H-B = Henrique Marxmüller (M); HRL = Raneče Lebeuf (pers. herb.); HTL = Huyen Than Le (CMU, GENT); IC = Aurelia Pia Conde (BCN); JC = Julio Cabero (pers. herb.); JET = Jennifer E. Tonkin (ME); JL = Jaime Litostrobo (BCN); JMV = Josep M. Vidal (BCN); JN = József Nyitrik (ME); JR = Juhani Ruotsalmi (KLUO); JV = Jan Verheyen (GENT); KAD = Kanad Verheyen (C, GENT); KD = Kodebe Van de Putte (GENT); LM = Leticia Montoya (XAL); MAM = Makela Saba (LAH); MCA = M. Catherine Aime (pers. herb.); MTB = Marco Floriani (pers. herb.); MTB = Michael Illoizoides (pers. herb.); MS = Mauro Samarti (pers. herb.); MTB = Maria Teresa Bassi (TUR); MU = Michael Illoizoides (pers. herb.); PAM = Pierre-Arthur Moreau (GENT); PAM = Pierre-Arthur Moreau (ME); PBM = P. Brandon Matherly (CLW); PL = Patrick Leonard (pers. herb.); RF = Raymond M. Fatto (pers. herb.); RH = Roy E. Halling (NY); RW (GENT) = Ruben Wallenius (GENT); RW (NY) = Roy Walling (E); SLM = Steven L. Miller (pers. herb.); T = James M. Trappe (ME), OSC PC; PBM = P. Brandon Matherly (CLW); WTU = Wayne T. Tamm (WTU); Y = Yves (pers. herb.)

AY218520  
AY218520  
AY854063 AF9393078

*Stereum hirsutum* Binder & Hibbett (2002) **FPL 8805**

New species, new combinations and sequences produced from this study are in **bold**. Sequestrate taxa are marked with an asterisk (\*) symbol.


- : Collection sources: AB = Ananad Bâ (BR); AM = Amer Monteiro (pers. herb.); AT = Andrew Taylor (UPS); AV = Annemieke Verbeeken (GENT); AVoik = Andrus Voik (GENT); BH = Bryony M. Horton (HO); BB = Bart Buyck (PO); BM = Baldomero Moreno-Aroyo (pers. herb.); BPL = Brian P. Looney (TENN); BDW = Birgitta Wästberg (UPPS); CLC = Cathie L. Cripps (MONT); DA = David Christopher W. Dunk (ME); DE = Dennis E. Desjardin (PC); DFSU = David G. Barge (MONT); DS = Dirk Stubbe (GENT); EB = Edward G. Barge (MONT); EG = Edith Georges Konstantinidis (pers. herb.); Garay-Serano (XAL); EH = Egon Horak (ZT); EU = Elisabeth Uhlmann (UPS); FA = François Ayer (WSL); FAN = Li Fan (BJTC); FH = Felix Hampe (GENT); FC = Franz Oberwinkler (TUB); GD = Giuseppe Donelli (UPS); GG = Gerhard Gross (M); GK = Georges Konstantinidis (pers. herb.); GM-RG = Gabriel Moreno and Ricardo Galán (AH); GO = Roberto Garbay Oriol (MEUX); H-B = Henrique Becker (UPS); HMR = Hermann Marxmüller (M); HRL = Raneče Lebeuf (pers. herb.); IC = Aurelia Pia Conde (BCN); JC = Julio Cabero (pers. herb.); JET = Jennifer E. Tonkin (ME); JL = Jaime Llristosa (BCN); JMV = Josep M. Vidal (BCN); JN = József Nyitrik (ME); JR = Juhani Ruotsimäki (KLUO); JV = Jan Verheyen (GENT); KAD = Kanad Verheyen (C); KDV = Koeke Van de Putte (GENT); LM = Leticia Montoya (XAL); MAM = Makela Saba (LAH); MCA = M. Catherine Aime (pers. herb.); MTB = Marco Floriani (pers. herb.); MTB = Michael Illoizoides (pers. herb.); MS = Mauro Samarti (pers. herb.); MTB = Maria Teresa Bassi (TUR); MU = Michael Illoizoides (pers. herb.); PAM = Pierre-Arthur Moreau (GENT); PAM = Pierre-Arthur Moreau (ME); PBM = P. Brandon Matherly (CLW); PL = Patrick Leonard (pers. herb.); RF = Raymond M. Fatto (pers. herb.); RH = Roy E. Halling (NY); RW (GENT) = Ruben Wallenius (GENT); RW (NY) = Roy E. Halling (E); SLM = Steven L. Miller (pers. herb.); T = James M. Trappe (ME), OSC PC; PBM = P. Brandon Matherly (CLW); PL = Patrick Leonard (pers. herb.); RF = Raymond M. Fatto (pers. herb.); RH = Roy E. Halling (E); SLM = Steven L. Miller (pers. herb.); T = James M. Trappe (ME), OSC

were above 0.95, and subsignificant if BP values were above 60 % or PP values were above 0.90.


## RESULTS

Maximum-likelihood analysis of the general ITS – 28S rDNA – *rpb2* dataset including all new sequences and those recovered from public databases (Fig. 1) produced a phylogenetic tree consisting of four significantly supported clades, those of genera *Lactarius*, *Lactifluus*, *Multifurca* and *Russula*, in accordance with previous reconstructions of the *Russulaceae* lineage (Buyck et al. 2008, Lebel et al. 2013, Verbeken et al. 2014b, Kong et al. 2015, Looney et al. 2016, Song et al. 2016, De Crop et al. 2017).


*Lactarius* (Fig. 2) was composed of five main clades, namely *Lactarius* subg. *Plinthogalus*, */L. hispidulus*, */L. bisporus*, */L. kabansus*, and a lineage including *Lactarius* subg. *Lactarius* (= *Lactarius* subg. *Piperites*) and *Lactarius* subg. *Russularia*, all of them receiving significant or subsignificant support, in agreement with previous analyses of *Lactarius* (Eberhardt & Verbeken 2004, Buyck et al. 2008, Verbeken et al. 2014a, b, Barge et al. 2016), except for subg. *Lactarius* (PP 0.88, BP 58). All European taxa of sequestrate *Lactarius* nested within the monophyletic clade formed by *L.* subg. *Lactarius* and *L.* subg. *Russularia*. Seven significantly supported sequestrate species were identified, two of which belong to *L.* subg. *Russularia* and five fall outside and are therefore considered part of *L.* subg. *Lactarius*. *Lactarius* subg. *Russularia* received subsignificant support (PP 0.94, BP 47),




**Fig. 1** Consensus phylogram of the family *Russulaceae* obtained in RAxML after 2000 bootstrap iterations of a combined alignment of ITS rDNA, 28S rDNA and rpb2. Major clades were collapsed and rooting branch shortened for publication. Nodes were annotated if supported by > 70 % ML BP, but non-significant support values are exceptionally represented inside parentheses.



**Fig. 2** Extended phylogram of genus *Lactarius* obtained from Fig. 1. Rooting branch was shortened for publication. Nodes were annotated if supported by > 0.95 bayesian PP or > 70 % ML BP, but non-significant support values are exceptionally represented inside parentheses. Major lineages are highlighted with coloured boxes. Sequestrate species are marked with black circle (●) symbols. Specimens sequenced in the present work are highlighted in bold.



**Fig. 3** Extended phylogram of *Russula* sect. *Ingratae* obtained from Fig. 1. Rooting branch was shortened for publication. Nodes were annotated if supported by > 0.95 bayesian PP or > 70 % ML BP, but non-significant support values are exceptionally represented inside parentheses. Major lineages are highlighted with coloured boxes. Sequestrate species are marked with black circle (•) symbols. Specimens sequenced in the present work are highlighted in **bold**.




**Fig. 4** Extended phylogram of *Russula* sect. *Rigidiae* obtained from Fig. 1. Rooting branch was shortened for publication. Nodes were annotated if supported by > 0.95 bayesian PP or > 70 % ML BP, but non-significant support values are exceptionally represented inside parentheses. Major lineages are highlighted with coloured boxes. Sequestrate species are marked with black circle (•) symbols. Specimens sequenced in the present work are highlighted in **bold**.

a result similar to that of Verbeken et al. (2014b), probably due to the insufficient phylogenetic signal in the species analyzed, gaps in the phylogenetic diversity because of an incomplete sampling, or too much phylogenetic noise in the dataset employed. However, a significantly supported clade formed by *L. borzianus*, *L. josserandii* (= *Z. hispanicus*) and some gymnocarpic species, such as *L. fulvissimus* and *L. subsericatus*, in accordance with results of Verbeken et al. (2014b), Liu et al. (2015), or Barge et al. (2016). The remaining European sequestrate Russulaceae species included in *L.* subg. *Lactarius* were identified as the existing species *L. giennensis*, *L. stephensii* and *Z. soehneri*, or otherwise accommodated into the new taxa *L. populicola* and *L. subgiennensis*.

*Russula* was divided in eight significantly supported monophyletic clades (Fig. 1, 3–5):

- 1) an isolated branch for */R. compacta*;
- 2) a clade formed by *Russula* subsect. *Pallidosporinae* (Bon 1988), as well as the lineages of */R. cascadiensis* and */R. metachromatica*;
- 3) *Russula* subsect. *Lactarioideae* (including *R. delica*);
- 4) *Russula* sect. *Nigricantinae*;
- 5) *Russula* sect. *Archaeinae*;
- 6) *Russula* subsect. *Farinipedes* (including *R. crassotunicata* and *R. pallescens*);
- 7) a monophyletic lineage formed by *Russula* sect. *Ingratae* (≡ subg. *Ingratula*) and *Russula* sect. *Rigidiae* (≡ subg. *Heterophyllidia*); and
- 8) a large *Russula* s.str. clade including the remaining species.



**Fig. 5** Extended phylogram of *Russula* s.str. obtained from Fig. 1. Rooting branch was shortened for publication. Nodes were annotated if supported by > 0.95 bayesian PP or > 70 % ML BP, but non-significant support values are exceptionally represented inside parentheses. Major lineages are highlighted with coloured boxes. Sequestrate species are marked with black circle (●) symbols. Specimens sequenced in the present work are highlighted in bold.

These clades were also significantly supported by Kong et al. (2015), Looney et al. (2016) and Bazzicalupo et al. (2017). All European sequestrate Russulaceae taxa nested within the *Ingratae-Rigidae* clade or *Russula* s.str.

*Russula* sect. *Ingratae* (Fig. 3) received significant support (PP 1.00, BP 99). It was composed of *R.* subsect. *Subvelatae*, *Pectinatinae*, and *Foetentinae*, as well as the lineages of /*R. granulata*, /*R. amoenolens* and /*R. pilosella*, all of them significantly supported except *Foetentinae* (PP 0.84, BP 29), probably because of incomplete data from many American sequestrate taxa. These results agree with those obtained by other researchers (Kong et al. 2015, Li et al. 2015, Lee et al. 2017, Melera et al. 2017). Sequestrate Russulaceae taxa are present in most of these lineages, but all currently known European sequestrate species were related to subsect. *Foetentinae* or the /*R. amoenolens* lineage. *Russula ammophila* was not significantly different from *R. amoenolens* employing ITS, 28S rDNA and *rpb2* data, but significant differences were found in *tef1* gene-data (data not shown).

*Russula* sect. *Rigidae* (Fig. 4) was split into two clades. The first one was composed of *R.* subsect. *Cyanoxanthinae* and *Pluviales* (Singer 1986, Buyck & Ovrebo 2002, Cheype & Campo 2012), as well as the /*R. cheelii* lineage (PP 0.99, BP 55). The second clade was formed by the remaining species of *Rigidae* (PP 0.98, BP 62). This clade was composed of subsect. *Ilicinae* (BP 100, PP 1.00) on the one side, and the remaining subsections of *Rigidae* (PP 0.87, BP 77) on the other, including subsections *Griseinae* (divided in two clades, those of /*R. ochrospora* and /*R. grisea*), *Virescentinae*, *Pseudoepitheiosinae*, *Heterophyllae* and *Amoeninae*. Most of these clades were also significantly supported by the phylogenetic reconstructions of *Rigidae* conducted by Dutta et al. (2015) and Zhao et al. (2015). The only European sequestrate species related to sect. *Rigidae* was *R. andaluciana*, a replacement name for *Gymnomyces sublevisporus*, whose isotype was significantly related with subsect. *Amoeninae*.

*Russula* s.str. (Fig. 5) was composed of a basal clade which did not receive significant support (PP 0.73, BP 50), including *Russula* subsect. *Russula*, as well as subsect. *Sardoninae*, subsect. *Viscidinae*, and the lineages of *R. consobrina* (subsect. *Consobrinae*) and *R. fellea* (subsect. *Felleinae*). A second large

clade with significant support clustered all other species of *Russula*, and had itself two basal clades: subsections *Lepidinae* and *Firmiores*, the latter including gymnocarpic, pseudoangiocarpic and angiocarpic species. The remaining species of *Russula* grouped into several distinct clades with significant support: subsections *Cupreinae*, *Laricinae*, *Lilaceinae*, *Maculatinae*, *Paludosinae*, *Puellinae*, *Roseinae*, a monophyletic clade containing subsections *Amethystinae* and *Chamaeleontinae*, and finally the so-called 'crown' clade (Looney et al. 2016), containing species from subsections *Decolorantinae*, *Insidiolinae*, *Integrae*, *Olivaceinae*, *Vinosinae* and *Xerampeliniae*, as well as a lineage including the type species of genus *Cystangium* (*C. sessile* ≡ *Russula sessilis*), and many sequestrate and gymnocarpic species from Australia (Lebel & Tonkin 2007) and South America (Trieveiler-Pereira et al. 2015). Most European sequestrate taxa were related to subsections *Firmiores*, *Laricinae*, *Maculatinae* and *Puellinae*, but there were also two independent lineages: one of them, *R. monospora*, was subsignificantly related with *R. consobrina*, the type species of subsect. *Consobrinae* (PP 0.94, BP 53), while the other, *R. hobartiae*, represented a new species from Cyprus significantly related with a putative specimen of *Russula ochrophylla*. Lineages matching known species of sequestrate Russulaceae are listed below or re-described, and new names are proposed with complete descriptions for those representing new taxa.

## TAXONOMY

***Lactarius borzianus* (Cavara) Verbeken & Nuytinck, Belg. J. Bot. 136, 2: 151. 2003 — Fig. 6**

*Basionym.* *Arcangeliella borziana* Cavara, Nuovo Giorn. Bot. Ital., Nuov. Ser. 7, 2: 126. 1900.

*Synonyms.* *Octaviania borziana* (Cavara) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 740. 1958.

*Arcangeliella stephensi* var. *borziana* (Cavara) Krieglst., Z. Mykol. 57, 1: 19. 1991.

*Arcangeliella volemoides* K. Mader & A. Mader, Österr. Z. Pilzk. 1: 5. 1992.

*Basidiomata* 1–4 cm wide, angiocarpic, subglobose to tuberiform, often bilobed, sessile or subsessile, with a residual stipe; radially alveolate in the base. *Pileus* dry, first smooth, then areo-



**Fig. 6** *Lactarius borzianus*. a. Reproduction of Cavara's plate of *Arcangeliella borziana* (Cavara 1900, tav. VII). — b. MSL1962F0412. Basidiomata. — c—e. Spores in Melzer; d—e. SEM images of spores. — Scale bars: b = 1 cm, c = 10 µm; d—e = 5 µm. — Photos: b. M. Slavova; c. J.M. Vidal; d—e. UdG.

late, basally closed, or open exposing the hymenophore; initially yellowish white (2A2) to pale yellow (3A3–4A5), then brownish orange (7C7) with reddish brown to dark brown (8E8–F8) maculae. *Hymenophore* loculate, labyrinthoid, yellowish white (2A2), pale yellow (4A4) to pale brown (7D7); locules irregularly arranged, 0.6–1.8 × 0.15–3 mm (1–2 per mm), elongated, sinuous, empty; fresh spore mass in locules yellowish white (4A1–A2). *Columella* percurrent or branched, apparent only in young specimens. *Latex* scant, more abundant in columella and pileal context, colourless to white, changing to yellow. *Odour* of wild bug (*Nazara viridula*) or musky; taste unpleasant, bitterish after a while.

*Spores* 9.5–13(–15) × 8–10(–11) µm, Q = 1.16–1.28, heterotropic, subglobose to broadly ellipsoid, echinate; warts 1–1.5 µm high, isolated, amyloid; suprahilar plage absent; hilar appendix 1–1.5 µm long, sometimes with a short sterigmal fragment 1–3 µm long. *Basidia* 3–4-spored, 30–65 × 10–15 µm, cylindrical to subclavate; sterigmata conical, curved, 5–8 µm long. *Macrocytidia* absent. *Cystidioles* infrequent, 20–50 × 4–6 µm, septate, cylindrico-rostrate to ventricose-rostrate, acuminate, more abundant in sterile locules. *Paraphysoid cells* abundant, 15–40 × 5–12 µm, cylindrical to slightly clavate, sometimes subglobose, often 1–2-septate. *Pseudocystidia* not observed. *Subhymenium* cellular, formed by prismatic cells 8–25 µm diam. *Hymenophoral trama* homomerous, formed by septate hyphae 2–6.5 µm diam, with subglobose thickenings up to 12 µm wide in tramal anastomoses. *Pileipellis* and *context* 100–250 µm thick; suprapellis trichodermal to oedotrichodermal, formed by cylindrical, sinuous, septate hairs 10–40 × 3–6 µm, and lanceolate dermatocystidia similar to those of the sterile locules, that soon collapses in a brown mass; subpellis made of prismatic hyphae up to 15 µm diam. *Pileal context* a cutis with the same composition of the hymenophoral trama, plenty of laticifera. *Laticifera* 2.5–10 µm diam, scarce in the hymenophoral trama and more abundant in the pileal context. *Thromboplera* 2–8 µm diam, abundant in young specimens. Some specimens present isolated sphaerocytes in the hymenophoral trama and pileal context, 15–30 µm diam, being more abundant and grouped in rosettes or chains in the columella.

**Habitat, Distribution & Season** — Gregarious, hypogeous or semi-hypogeous under needles, in subalpine conifer forests of *Abies* and *Picea*, on siliceous soil. Summer and autumn. So far known from the Alps, Apennine (Italy), and Rhodope mountains (Greece), at 1000–1800 m altitude.

**Material studied.** AUSTRIA, Lower Austria, Gföhl, near Dorf Brunn, under *Picea abies*, 12 Sept. 1970, A. & K. Mader (WU 10875, holotype of *A. volemoides*). — BULGARIA, Smolyan, Pamporovo, Southern Rhodope mountains, 1650–1750 m, under *Picea abies*, on siliceous soil, 15 Aug. 2017, M. Slavova (MSL1962F0412, duplicate BCN JMV800685). — FRANCE, Rhône-Alpes, Savoie, Héry-sur-Ugine, Réserve Naturelle du Nant Pareu-Merdassier, 1450 m, under *Picea abies* in the vicinity of *Lactarius aurantioculus*, on siliceous soil, 13 Aug. 2002, P.-A. Moreau as *A. borziana* (PAM02081306, duplicate BCN JMV800279b). — ITALY, Emilia-Romagna, Reggio Emilia, Civago, Abetina Reale forest, Tuscan-Emilian Apennines, under *Abies alba*, on siliceous soil, 5 Aug. 1999, A. Montecchia as *A. borziana* (AM1973, duplicate BCN JMV800239); Tuscany, Firenze, Reggello, Vallombrosa Forest, Tuscan-Emilian Apennines, 'In silvis abiegnis Vallisumbrosae (Etruria), aestate, F. Cavara' (NY, herb. S.M. Zeller, labelled 'Elasmomyces mattiroloanus, Italy, Vallombrosa, coll. F. Cavara, semiepigeous in fallen needles under *Abies pectinata*, autumn 1896, sent to Zeller by Cavara, type'; isotype of *A. borziana*).

**Notes** — *Arcangeliella borziana* was collected in the summer of 1898 in Tuscany (Italy) by Cavara (1900), growing hypogaeously under *Abies alba*. According to Zeller & Dodge (1937), collections of *A. borziana* preserved in their personal herbaria are duplicates of type material preserved at the Herbarium of the Università degli Studi di Napoli (Naples, Italy), but these were apparently not studied in detail by Zeller & Dodge, who just included in their works a brief description of *A. borziana*.

taken from Cavara (Zeller & Dodge 1919). Later descriptions by Singer & Smith (1960), Pegler & Young (1979), Lebel & Trappe (2000), probably based on the isotypes of Zeller & Dodge (C.W. Dodge 2087 at FH, and S.M. Zeller 1671 at NY), highlighted the same group of features: presence of a percurrent stipe-columella, hymenophoral trama and pileal context containing nests of sphaerocytes and laticifera, hymenium with abundant macrocystidia, and large ochraceous globose spores measuring 9–15 × 10–13 µm. Vidal (2004a), conducted a detailed study of these specimens, and observed that these microscopical features do not match the protologue of this species, but are more compatible with that of *Elasmomyces mattiroloanus*, suggesting that they do not represent the original concept of *A. borziana* sensu Cavara, and probably both species were erroneously labelled. This is all the more likely if laticifera had been mistaken for gloeoplera. The original concept of *Lactarius borzianus* is of a sessile or subsessile species that may have a rudimentary stipe-columella, a hymenophoral trama and pileal context rarely containing sphaerocytes (sometimes just represented by scattered globose elements, but never forming nests), a fertile hymenium deprived of macrocystidia (with only cystidioles present in the sterile hymenium), and whose spores are hyaline and broadly ellipsoid. It is a silicolous species growing in wet, subalpine conifer forests of *Abies* and *Picea*, receiving large amounts of precipitation (about 2000 mm).


Genetic studies by Peter et al. (2001) conducted on specimens collected in the Swiss Alps (F. Ayer 96-05-3344 WSL) show that this species is significantly related to members of *L. subg. Russularia*. Nuytinck et al. (2003) compared sequences of *Z. stephensii* with those of *A. borziana* obtained by Peter et al. (2001), and found both species to be genetically distinct, recombining them into *Lactarius*. Neither Nuytinck et al. (2003) nor Vidal (2004a) were able to observe any significant macro- or microscopical differences between the type collection of *A. volemoides* and *A. borziana*, and both taxa were therefore considered to be synonyms by these authors. Another specimen (MCVE 16944) collected at Abetina Reale (Reggio Emilia, Italy) by Amer Montecchi was sequenced by Osmundson et al. (2013) and found to be genetically identical with the Swiss collection already mentioned. Species of *Lactarius* subg. *Russularia* are characterized by reddish brown or orange pilei sometimes producing remarkable smells, bleeding white or transparent latex not changing in colour upon exposure or slowly changing to yellow, globose to ellipsoid spores ornamented with a more or less complete reticulum and an inamyloid or distally amyloid plage, as well as dermatocystidia (Hesler & Smith 1979, Wisitrassameewong et al. 2014). Other sequestrate species in *Lactarius* subg. *Russularia* include the European *L. josserandii* (= *Z. hispanicus*), the North American *L. silviae* (≡ *Elasmomyces camphoratus*) and the Australian *Zelleromyces striatus* and *Z. daucinus*, all of them matching the characteristic features of *Lactarius* subg. *Russularia* (Singer & Smith 1960, Beaton et al. 1984, Miller et al. 2001, Verbeken et al. 2014b). The North American species *L. paulus* (≡ *Arcangeliella parva*) and *L. variegatus* (≡ *A. variegata*) could also be related to subgenus *Russularia*, but their phylogenetic affiliation is still dubious because of the scarce genetic data available (Miller et al. 2001).

#### *Lactarius giennensis* (Mor.-Arr. et al.) Pierotti, Index Fungorum 254: 1. 2015 — Fig. 7

**Basionym.** *Zelleromyces giennensis* Mor.-Arr. et al., Cryptog. Mycol. 19, 1–2: 108. 1998.

**Synonym.** *Arcangeliella giennensis* (Mor.-Arr. et al.) J.M. Vidal, Rev. Catalana Micol. 26: 74. 2004.

**Basidiomata** 1–2.5 cm wide, angiocarpic, subglobose to tuberiform, somewhat lobate, sessile. **Pileus** whitish to pale yellowish,



**Fig. 7** *Lactarius giennensis*. a. JC20061020. Basidiomata. — b–c. JMV20010204-1. b. Pileipellis and context; c. cystidioles of external immature locule. — d–f. JMV931123-1. d. Basidiomata; e. basidia; f. spores in Melzer. — g–i. JMV20010204-1. SEM images of spores. — Scale bars: a, d = 1 cm; b = 20 µm; c, e–h = 10 µm; i = 5 µm. — Photos: a. J. Cabero; b–f. J.M. Vidal; g–i. UdG.

membranous, smooth, partially evanescent, indistinctly scrobiculate or with some scattered minute openings. *Hymenophore* loculate, whitish to pale cream or pinkish. *Columella* branched, inconspicuous. *Latex* not observed. *Odour* mild.

*Spores* 10–13 × 7–9 µm, Q = 1.15–1.35, ovoid to ellipsoid, orthotropic; reticulum 1–2 µm high, complete, amyloid. *Basidia* 1-spored, 30–45 × 6–10 µm, cylindrical, clavate or sublageniform. *Macrocytidia* absent. *Cystidioles* 30–42 × 4–8 µm, lageniform-urticiform. *Subhymenium* rameose to cellular. *Hymenophoral trama* homoiomerous, made of septate hyphae 3–6 µm diam. *Pileipellis* and *context* 150–200 µm thick; suprapellis a trichoderm of septate hyphae. *Pileal context* a prosenchyma of intricate hyphae 3–6 µm diam. *Laticifera* 5–8 µm diam, present in the hymenophoral trama and pileal context.

**Habitat, Distribution & Season** — Gregarious, hypogeous, associated with *Cistus* and *Halimium*, often accompanied by *Pinus*, on siliceous soil. Autumn to spring. Xerophytic. So far restricted to the western Mediterranean region, in Central Spain, between 600–1000 m altitude.

**Material studied.** SPAIN, Andalusia, Jaén, La Aliseda, under *Pinus halepensis*, 28 Feb. 1994, J. Gómez & B. Moreno-Arroyo (MA-Fungi 38674, holotype of *Z. giennensis*); Castilla and Leon, León, Lugar, under *Cistus* sp. and *Eucalyptus* sp., 25 Sept. 1993, T. Pérez-Jarauta (BCN JMV930925); Salamanca, Casarito, under *Halimium alyssoides*, on siliceous soil, 6 Dec. 1992, T. Pérez-Jarauta (BCN JMV921206); Salamanca, Miranda del Castañar, under *Cistus ladanifer*, 18 Apr. 2003, A. García-Blanco, M. Sanz-Carazo & J.B. Del Val as *Z. giennensis* (AVM 1615, duplicate BCN JMV800629)\*; Zamora, Tábara, 825 m, under *Cistus ladanifer* and *Pinus pinaster*, on siliceous soil, 4 Feb. 2001, P. Juste & F. García (BCN JMV20010204-1); ibid., 20 Oct. 2006, J. Cabero as *Z. giennensis* (JC20061020)\*; Castilla-La Mancha, Ciudad Real, El Viso del Marqués, under *Halimium ocymoides*, on siliceous soil, 23 Nov. 1993, T. Pérez-Jarauta (BCN JMV931123-1).

**Notes** — *Lactarius giennensis* is morphologically characterized by its whitish to pinkish basidiomata, with a rather thin,

membranous pileus with perforations opening to the hymenophoral locules, not bleeding latex when cut. Spores are ovoid or ellipsoid, ornamented with a closed reticulum composed of ridges up to 2 µm high. Abundant laticifera and thromboplera were observed in the hymenophoral trama, as well as lageniform-urticiform cystidioles in the hymenium of immature external locules.

Genetic data from the holotype specimen of *L. giennensis* (MA-Fungi 38674, GenBank AF230900, incorrectly published as AF230800 by Calonge & Martín 2000), evidenced that this species is related with other lactarioid taxa. In the present work this species is subsignificantly related with the lineages of subg. *Lactarius* and *Russularia*, so a hypothetical affiliation with subg. *Lactarius* is suggested. The clade containing *L. giennensis* has an unusually long branch, suggesting it could represent a highly evolved or ancestral lineage, or else evidence an incomplete sampling. The holotype of *L. giennensis* was found at Jaén (Andalusia, Spain) growing subhypogaeously under *Pinus halepensis*, but the collections analyzed in the present work suggest a broader distribution, being present in central Spain, associated with *Cistaceae* hosts in acidic soils.

***Lactarius josserandii* (Malençon) J.M. Vidal & P. Alvarado, comb. nov.** — MycoBank MB828496; Fig. 8

**Basionym.** *Zelleromyces josserandii* Malençon, Rev. Mycol. (Paris) 39: 303. 1975.


**Synonyms.** *Arcangeliella josserandii* (Malençon) J.M. Vidal, Rev. Catalana Micol. 26: 75. 2004.

*Zelleromyces hispanicus* Calonge & Pegler, Cryptog. Mycol. 19, 1–2: 100. 1998.

**Basidiomata** 2–5 cm wide, angiocarpic, irregular, obpyriform to subglobose, tuberiform to lobulate, sessile, with a depressed

or conical base immersed into the soil. *Pileus* dry, smooth, continuous, with minute circular depressions in the surface, but not exposing the locules; initially pale orange (5A5), then orange-red (8B8), brownish red (8C8) or reddish brown (9D8), like in *Lactarius fulvissimus*, remaining pale orange in unexposed areas. *Hymenophore* loculate, labyrinthoid, pale, initially yellowish white (4A2), then orange-white (5A2); locules 0.5–1.5 × 0.15–0.3 mm (1–2 per mm), elongated, sinuous, empty; fresh spore mass in locules whitish; yellowish white (4A2) in exsiccata. *Columella* absent or rudimentary, with thin whitish veins. *Latex* scant, white, unchanging, more abundant in the pileal context. *Odour* fruity, taste sweetish, later astringent. Spores 8.5–12.5(–13.5) × 7–9.5(–10.5) µm, Q = 1.1–1.3, sub-

globose to ellipsoid, heterotrophic, initially hyaline then yellow, reticulated; reticulum amyloid, complete or with some interruptions, formed by 0.5–1 µm high ridges and warts; hilar appendix 0.7–1.4 × 0.7–1.1 µm, short, cylindrical to conical; suprahilar plage not seen. *Basidia* typically 4-spored, but also 1–3-spored, 40–50 × 7.5–10 µm, cylindrical, sinuous, with granular content; sterigmata 5–8 µm long, conical. *Basidiobles* 20–30 × 7–10 µm, cylindrical. *Macrocystidia* absent. *Cystidioles* 42–44 × 7–8 µm, lageniform, scarce, present only in external immature locules. *Paraphysoid* cells abundant, 10–30 × 4–11 µm, entire or 1–2-septate. *Pseudocystidia* not observed. *Subhymenium* rameous to cellular, formed by chains of prismatic cells 8–10 µm diam. *Hymenophoral* trama 80–125 µm wide,



**Fig. 8** *Lactarius josserandii*. a–b. Reproduction of Malençon's plates of *Zelleromyces josserandii* corresponding to the sample n° 3242 conserved in the herbarium of Montpellier (MPU C03565, ©Université de Montpellier-Herbier). — c. MPU 0310529 (holotype of *Z. josserandii*). Spores in Melzer. — d–m. JMV800621. d–e. Basidiomata; f. pileipellis and context; g–h. basidia, basidiobles, paraphysoid cells and subhymenium; i. cystidiole of an external immature locule; j. spores in Melzer; k. spores in ammonia; l–m. SEM images of spores. — Scale bars: c, i–k = 10 µm; d–e = 1 cm; f–h = 20 µm; l–m = 5 µm. — Photos: c, f–k. J.M. Vidal; d–e. F. García; l–m. UdG.

homoiomericous, formed by septate, tortuous hyphae 2–7 µm diam, some of them inflated, but lacking sphaerocytes, with abundant laticifera 2–8 µm diam, and some thromboplera 2–6 µm diam. *Pileipellis* and *context* 175–300 µm thick; pileipellis 55–65 µm thick, arranged in an oedotrichoderm resembling that of *L. fulvissimus*, formed by: 1) a suprapellis of cylindrical and sinuous septate hairs, soon collapsing in a brown mass; and 2) a subpellis composed of prismatic hyphae 8–10 µm diam. *Pileal context* a prosenchyma, with the same composition of the hymenophoral trama, with plenty of laticifera and some thromboplera.

**Habitat, Distribution & Season** — Gregarious, hypogeous to semi-hypogeous, growing under needles in montane conifer forests of *Pinus* and *Cedrus*, on siliceous soil, frequently in the vicinity of *Lactarius aurantiacus*. Autumn. Occurring in submediterranean regions, in Morocco, France and Spain, between 1250–1900 m altitude.

**Material studied.** MOROCCO, Taza-Al Hoceima-Taounate, Ketama, Llano Amarillo, 'Rif cristallin, 1600 m, parmi les jeunes cèdres mêlés de *Cistus laurifolius*', 16 Nov. 1957, G. Malençon (MPU 0310528, herb. G. Malençon 3242, paratype of *Z. josserandii*); ibid., 'peu de distance de la 1er (env. 300 m)', 18 Nov. 1957, G. Malençon (MPU 0310529, herb. G. Malençon 3242 Bis, holotype of *Z. josserandii*); ibid., 'sous les cèdres', 29-X-1972, R. Bertault (MPU 0310527, herb. G. Malençon 7252, paratype of *Z. josserandii*). — SPAIN, Castilla and Leon, Segovia, Navafria, Puerto de Navafria, El Nevero, Sierra de Guadarrama, 1800–1900 m, under *Pinus sylvestris*, on siliceous soil, 6 Oct. 2006, F. García (BCN JMV800621)\*; Community of Madrid, Cercedilla, Dehesas, under *Pinus sylvestris*, on siliceous soil, 27 Nov. 1996, F.D. Calonge & C. García-Ruz (MA-Fungi 37498, holotype of *Z. hispanicus*); ibid., 23 Nov. 1996, J. Daniel & J.M. Santos (MA-Fungi 37497, paratype of *Z. hispanicus*); ibid., 16 Nov. 1997, R. Cifuentes (MA-Fungi 38311, paratype of *Z. hispanicus*).

**Notes** — Malençon (1975) proposed the name *Zelleromyces josserandii* for an orange hypogeous fungus, bleeding white latex when cut, and displaying reticulate spores, found in 1957 and 1972 in Morocco and France (five collections made by G. Malençon in the Rif mountains, one found in the same area by R. Bertault, and two more collections found by R. Ruffier-Lanche and J. Mornand in the Aitone forest in Corsica under *Pinus nigra* subsp. *laricio* at 1250 m altitude). Later on, Calonge & Pegler (1998) studied several collections found under *Pinus sylvestris* at Sierra de Guadarrama (Madrid, Spain), with morphological features closely resembling those of *Z. josserandii*, which they named *Z. hispanicus*.

Vidal (2004a) proposed a synonymy between *Z. josserandii* and *Z. hispanicus* after observing identical macro- and microscopical features in the type specimens, but all attempts to sequence the type material of *Z. josserandii* have been unsuccessful. Calonge & Pegler (1998) first suggested a close relationship between *Z. hispanicus* and *Lactarius aurantiacus*, and later Calonge & Martín (2000, 2003) highlighted the slight genetic differences (37/634 bp) between ITS sequences of these species, both belonging to *Lactarius* subg. *Russularia*. However, phylogenetic analyses conducted by Verbeken et al. (2014b), as well as present results, suggest a closer affinity to *L. borzianus* and *L. fulvissimus*, which share a similar oedotrichodermal pileipellis.

***Lactarius populicola* J.M. Vidal, Konstantin., Setkos & Slavova, sp. nov.** — MycoBank MB828497; Fig. 9

Basidiomata similar to *L. stephensii*, 1–4 cm wide, angiocarpic, sessile, subglobose or tuberiform, lactescent, smooth, at first pale orange, then reddish brown to violet brown. Hymenophore loculate, deeply coloured, reddish yellow to orange red. Columella absent or inconspicuous. Latex watery-white to white, of mild taste. Odour fruity, of pears. Microscopy as in *L. stephensii*, except the spores which are subglobose to broadly ellipsoid, measuring 13–15 × 11–13 µm in size, covered by strongly amyloid, conical or tooth-like warts, 1–2 µm long. Hypogeous or semi-hypogeous under *Populus alba*.

**Etymology.** From Latin, *populus* = poplar, and -cola = inhabitant of, who lives in, in reference to its association to poplars (*Populus*).


**Holotype.** GREECE, Central Macedonia, Serres, Sidirokastro, 50 m, under *Populus alba*, 13 Mar. 2010, G. Setkos (BCN JMV800648)\*; isotype in herb. pers. G. Konstantinidis (GK4831).

**Basidiomata** 1–4 cm wide, angiocarpic, subglobose or irregular, lobate, reniform, tuberiform, sessile or with a minute sterile base, attached to soil by thin mycelial threads. *Pileus* smooth, viscid, persistent, open basally and exposing the locules in old specimens; at first pale orange (6A3), brownish orange (7C6) or orange-red (8A6) and finally reddish brown (8D6–9E5) to violet-brown (10E6). *Hymenophore* loculate, labyrinthoid, deeply coloured, reddish yellow (4A7), orange (5A6–6B7) to reddish orange (7A6), or orange-red (8A6–B6); locules very small, 0.2–0.5 × 0.02–0.1 mm (3–4 per mm), irregularly arranged, elongated, sinuous; septa 250–500 µm thick; fresh spore mass in locules pale yellow (4A4–A5); pale orange (5A3) to brownish orange (6C5) in exsiccata. *Columella* absent or inconspicuous. *Latex* watery-white in immature basidiomata and white in mature basidiomata, of mild taste. *Odour* strong, aromatic and fruity, reminiscent of pears.

**Spores** 13–15 × 11–13 µm, Q = 1.07–1.2, subglobose to broadly ellipsoid, orthotropic, echinulated, unigutulate, deep orange; warts deeply amyloid, robust, 1–2 µm long, conical or tooth-like, usually curved, with some verrucae among them; suprahilar plage absent; hilar appendix minute, usually united to a fragment of sterigma up to 4 × 2.5 µm. *Basidia* 1-spored, 35–50 × 8–10 µm, cylindrical, straight or sinuous, soon collapsed; initially hyaline and filled with oleiferous guttules, then of dark orange colour; sterigmata central or eccentric, 2–4 µm long. *Basidioles* similar to basidia. *Macrocytidia* and *cystidiolles* absent. *Paraphysoid cells* 20–30 × 6–8 µm, usually with 1–2 septa, cylindrical or with a clavate apex. *Pseudocystidia* not observed. *Subhymenium* ramos, formed by septate, cylindrical hyphae, 10–30 × 3–9 µm, perpendicular to the tramal hyphae. *Hymenophoral trama* 60–90 µm thick, homoiomericous, formed by hyaline, thin-walled, subgelatinized hyphae, 3–6 µm diam, finally dark yellow, completely gelatinized, with a prosenchymatous aspect; laticifera abundant, 8–10 µm diam, with thinner branches 3–5 µm diam; sometimes penetrating into the hymenium and terminating as pseudocystidia; thromboplera yellow, 3–5 µm diam. *Pileipellis* and *context* 300–500 µm thick; pileipellis 50–90 µm thick composed of: 1) a trichodermal suprapellis of septate hairs and dermatocystidia, 20–40 × 2.5–4 µm, straight to sinuous, rounded to acute at the apex, with yellow granular content, that soon collapses in a brown granular mass; and 2) a prosenchymatous subpellis 20–50 µm thick, of densely entangled hyphae 1.5–3 µm diam. *Pileal context* an ixocutis 250–400 µm thick, made by subgelatinized, entangled hyphae 2–4 µm diam, crossed by some thromboplera and abundant laticifera.

**Habitat, Distribution & Season** — Gregarious, hypogeous to semi-hypogeous in riparian forests under *Populus*, on alluvial soils. Spring to autumn. Distributed in temperate and Mediterranean regions, from Western to Southeastern Europe, from sea level up to 1000 m altitude.

**Additional material studied.** BULGARIA, Stara Zagora, Parvomai, Mirovo, 130 m, in a riparian forest under *Populus alba* and *Fraxinus ornus*, 12 Dec. 2015, T. Chokova MSL1629F6516 (SOMF 29973, duplicate BCN JMV800687). — FRANCE, Franche-Comté, Jura, Hérimoncourt, Aug. 1892, *L. Quélet* as 'Hydnangium galatheum' (UPS F013405). — GREECE, Central Macedonia, Serres, Sidirokastro, 50 m, under *Populus alba*, 24 Apr. 2010, G. Setkos (GK4969, duplicate BCN JMV800649); West Macedonia, Kastoria, Maniakoi, 650 m, under *Populus alba*, *Corylus avellana* and *Alnus glutinosa*, 7 Oct. 2010, G. Setkos (GK5223, duplicate BCN JMV800650); ibid., 25 Nov. 2011, G. Setkos (GK5904, duplicate BCN JMV800651). — ITALY, Emilia-Romagna, Reggio Emilia, Febbio, 1050 m, under *Populus alba*, 30 Oct. 1991, A. Montecchi as 'A. borziana' (AM1042, duplicate BCN JMV800036); Tuscany, Lucca, Montuolo, 15 m, under *Populus alba*, 1 Nov. 1993, G. Bernardini & L. Gori as 'A. stephensii' (ELG931101-2, duplicate BCN JMV800171); ibid., 28 Nov. 1993, G. Bernardini & L. Gori as 'A. stephensii' (ELG931128-2, duplicate BCN JMV800169); Lucca, Nozzano Castello, 20 m, under *Populus alba*,



**Fig. 9** *Lactarius populicola*. a–c. MSL1629F6516. a. Mature basidiomata and cut of hymenophore bleeding white latex; b. pileipellis and context; c. section of a septum with gelatinized hymenophoral trama. — d. GK4969. Old basidiomata. — e–i. GK4831 (BCN JMV800648, holotype). e. Locules of hymenophore; f. spores in Melzer; g–i. SEM images of spores. — j–l. GK5904. j–k. Young basidiomata; l. cut of hymenophore bleeding watery-white latex. — m. UPS F013405, herb. L. Quélet (as '*Hydnangium galatheum*'). Spores in Melzer. — Scale bars: a, d, j–k = 1 cm; b–c = 20 µm; e, l = 1 mm; f, m = 10 µm; g–i = 5 µm. — Photos: a. M. Slavova; b–c, f, m. J.M. Vidal; d–e, j–l. G. Konstantinidis; g–i. UdG.

31 July 1993, G. Bernardini & L. Gori as 'A. stephensi' (ELG930731-1, duplicate BCN JMV800170).

**Notes** — *Lactarius populicola* has been confused with *L. stephensi* (Nuytinck et al. 2003, Gori 2005) because of their multiple similarities. Microscopically, *L. populicola* has the same pileal and hymenial structure, and identical spore shape with *L. stephensi*, but spores are intensely amyloid and ornamented with robust conical or tooth-like warts, while *L. stephensi* has weakly amyloid spiny spores. Macroscopically, *L. populicola* is distinguished by the violaceous coloration of basidiomata in mature specimens and smaller locules in the hymenophore. While *L. stephensi* is probably associated with *Fagaceae* and *Betulaceae* hosts, *L. populicola* is linked to *Salicaceae* plants. Genetically, *L. populicola* is a monophyletic taxon significantly related with *L. stephensi*, both forming a significantly monophyletic clade with a third species, *L. soehneri*. These angiocarpic species were also related (PP 1.00, BP 88) with a clade formed by samples identified as *L. evosmus* and *L. zonarius*. The only sequenced specimens of *L. populicola* are by now the type proposed here, and a sample from Belgium (RW2930 GENT) treated as *L. stephensi* by Nuytinck et al. (2003).

***Lactarius soehneri* (Zeller & C.W. Dodge) J.M. Vidal & G. Moreno, comb. nov. — MycoBank MB828498; Fig. 10**

*Basionym.* *Hydnangium soehneri* Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 22: 372. 1935.


*Synonyms.* *Octaviania soehneri* (Zeller & C.W. Dodge) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 200. 1958.

*Martellia soehneri* (Zeller & C.W. Dodge) Singer & A.H. Sm., Mem. Torrey Bot. Club 21, 3: 31. 1960.

*Zelleromyces soehneri* (Zeller & C.W. Dodge) Trappe et al., Mycotaxon 81: 205. 2002.

*Hydnangium soehneri* var. *ettenbergii* Soehner, Z. Pilzk., N.F. 20, 3–4: 110. 1941 (nom. inval., Art. 39.1).

**Basidiomata** 1–2 cm wide, angiocarpic, subglobose or irregular, reniform, tuberiform, sessile; sterile base not observed. **Pileus** smooth, viscid, at first pale orange (5A3), then greyish red (9B4) and finally reddish brown (9D4) to violet-brown (10E7), basally open and alveolate in old specimens. **Hymenophore** loculate, labyrinthoid, initially pinkish white (7A2), then deeply coloured, pale red (9A3), dull red (9B4) and finally dull violet (15D4); locules large, 0.4–1.5 × 0.1–0.2 mm (1–2 per mm), irregularly arranged, elongated, sinuous, empty; septa 250–500 µm thick; fresh spore mass in locules greyish ruby (12D6);



**Fig. 10** *Lactarius soehneri*. — a. AH 39272. Basidiomata. — b–d. AH 46013. b. Pileipellis and context; c. hymenium and trama; d. spores in Melzer. — e. IC21091207. Basidiomata. — f. M, herb. E. Soehner 1041 (holotype of *Hydnangium soehneri* var. *ettenbergii*). Spores in Melzer. — g–k. M, herb. E. Soehner 1081 (holotype of *Hydnangium soehneri*). g. Basidia; h. spores in Melzer; i–k. SEM images of spores. — l. JMV800677. SEM image of a spore. — Scale bars: a, e = 1 cm; b–c = 20 µm; d, f–i = 10 µm; j–l = 5 µm. — Photos: a. G. Moreno; b–d, f–h. J.M. Vidal; e. C. Lavoise; i–l. UdG.

brownish orange (6C5) to brown (6D7) in exsiccata. *Columella* absent. *Latex* watery, scant or absent. *Odour* fruity. *Taste* earthy. *Spores* 14–18.5 × 12–15 µm, Q = 1.1–1.3, broadly ellipsoid to ellipsoid, orthotropic, echinate, unigutulate, pale orange; warts deeply amyloid, dense, 1–2 µm long, cylindrical, usually curved, sometimes fused forming short ridges, with numerous verrucae among them; suprahilar plage absent; hilar appendix 1.5–3 × 2 µm, usually attached to a fragment of sterigma. *Basidia* 1-spored, 50–70 × 6–10 µm, cylindrical, straight or sinuous, soon collapsed, initially hyaline, with a good number of oil drops inside, later filled with a dark orange substance that finally saturates all the hymenium; sterigmata 3–6 µm long. *Basidioles* similar to basidia. *Macrocytistidia* and *cystidioles* absent. *Paraphysoid* cells 30–60 × 2.5–7 µm, 1–3-septate, difficult to observe. *Pseudocystidia* not observed. *Subhymenium* ramoso. *Hymenophoral* trama 60–90 µm thick, homoiomerous, devoid of sphaerocytes, formed by hyaline, thin-walled, subgelatinized hyphae, 2–5 µm diam, displaying several yellow-brown thromboplera, 2–5 µm diam, and some laticifera, 3–10 µm diam, which sometimes penetrate into the hymenium terminating as pseudocystidia. *Pileipellis* and *context* 125–250 µm thick; pileipellis 70–150 µm composed of: 1) a trichodermal suprapellis of similar structure to that in *L. stephensii*, constituted by fragile,

septate hairs and dermatocystidia, 20–30 × 3–6 µm, rounded to acute at the apex, sometimes mucronate, with dark yellow granular contents and walls up to 0.5 µm, soon collapsing into a brown granular mass formed by remnants of dermatocystidia and yellow granules; and 2) a subpellis 50–80 µm thick of prosenchymatous aspect, composed of subgelatinized hyphae 2–5 µm diam. *Pileal* *context* an ixocutis 60–120 µm thick, composed of subgelatinized, entangled hyphae, 3–6 µm diam, intersected by abundant thromboplera and some laticifera.

**Habitat, Distribution & Season** — Gregarious, hypogeous to semi-hypogeous in montane forests of conifers (*Abies*, *Pinus*) or under broadleaved trees (*Corylus*, *Quercus*), on calcareous soil. Spring to autumn. Distributed in temperate regions of Germany, Italy and Spain, between 300–1400 m altitude.

**Material studied.** GERMANY, Bavaria, 'Ettenberg bei Berchtesgaden', 1100 m, under *Abies alba*, Aug. 1925, E. Soehner (M, herb. E. Soehner 1041, holotype of *H. soehneri* var. *ettenbergii*); 'Pupplinger Heide bei Wolfratshausen', 19 Aug. 1919, E. Soehneras 'Hydnangium monosporum' (M, herb. E. Soehner 1374); 'Pupplinger Heide bei Wolfratshausen, Föhrenwald', 21 Oct. 1928, E. Soehner as 'Hydnangium carneum' (M, herb. E. Soehner 1081, holotype of *H. soehneri*; FH, herb. C.W. Dodge, isotype; NY, herb. S.M. Zeller, isotype). — ITALY, Lombardy, Bergamo, San Pellegrino Terme, 350 m, under *Corylus avellana*, on calcareous soil, Sept. 2016, M. Berbenni, comm. M. Morara

(BCN JMV800677); Umbria, Perugia, Gubbio, 340 m, under *Quercus* sp. and *Corylus avellana*, in a truffle plantation of *Tuber melanosporum*, on calcareous soil, 21 Sept. 2012, A. Paz & C. Lavoise (BCN IC21091207). — SPAIN, Castilla-La Mancha, Guadalajara, Cantalojas, 1300–1400 m, in humus of *Pinus sylvestris*, on calcareous soil, 3 Apr. 2011, M.A. Sanz (AH 39272)\*; ibid., 9 Apr. 2011, M.A. Sanz (AH 46013, duplicate BCN JMV800633)\*.

**Notes** — *Lactarius soehneri* was considered a macrosporic form of *L. stephensii* by several authors (Mader & Mader 1992, Vidal 2004a), because of the very similar microscopic features, especially the presence of laticiferous hyphae and monosporic basidia documented in the type collection by Vidal (2004a). However, *L. soehneri* differs from *L. stephensii* and *L. populicola* by its scarce, watery latex, by the larger hymenophoral locules and by the ellipsoid spores up to  $18.5 \times 15 \mu\text{m}$ . The dark orange substance of hymenium that impregnates basidia and spores is another differentiating feature. While *L. stephensii* and *L. populicola* occur exclusively under broadleaved trees, *L. soehneri* also occurs under conifers.

Genetically, *L. soehneri* is significantly related with *L. populicola* and *L. stephensii*, and to a lesser extent to the gymnocarpic species *L. evosmus* and *L. zonarius*, and probably belongs to *Lactarius* subg. *Lactarius*. Unfortunately, no samples from the region where the type was found (Bavaria, Germany) or the same habitat (*Abies* or *Picea* forests) could be analyzed, and therefore the identification made here relies entirely on morphological similarities between the newly collected samples and the type specimen.

***Lactarius stephensii* (Berk.) Verbeken & Walleyn, Belg. J. Bot. 136, 2: 151. 2003 — Fig. 11**

**Basionym.** *Hydnangium stephensii* Berk., Ann. Mag. Nat. Hist., Ser. I, 13: 352. 1844.

**Synonyms.** *Octaviania stephensii* (Berk.) Tul. & C. Tul., Fung. Hypog.: 78. 1851.

*Arcangiella stephensii* (Berk.) Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 18: 463. 1931.

*Zelleromyces stephensii* (Berk.) A.H. Sm., Mycologia 54: 635. 1962.

*Martellia stephensii* (Berk.) K. Mader & A. Mader, Österr. Z. Pilzk. 1: 4. 1992.

*Hydnangium galatheum* Quél., Enchir. Fung.: 247. 1886 ('galatheum').

*Octaviania galathea* (Quél.) De Toni in P. A. Saccardo, Syll. Fung. 7, 1: 491. 1888 ('galathea').

**Basidiomata** 1–3.5 cm wide, angiocarpic, subglobose to tuberiform, lobate, sessile, sometimes with a minute sterile base; basally closed, or open and radially alveolate in old specimens. **Pileus** smooth, viscid, at first white, then maize yellow (4A6), oxide yellow (5C7) to carrod red (6B7), finally reddish brown (7D8–E8), with the same colour as *L. rufus*. **Hymenophore** loculate, labyrinthoid, initially white or ochraceous, acquiring a reddish brown colour like the pileus; locules small,  $0.3–1.5 \times 0.1–0.5 \text{ mm}$  (2–3 per mm), irregularly arranged, elongated, sinuous; septa  $150–300 \mu\text{m}$  thick; fresh spore mass in locules greyish orange (5B5). **Columella** absent or inconspicuous. **Latex** scant and hyaline, changing to citrine yellow in young specimens, white, abundant, and almost immutable in old basidiomata (orange in contact with KOH), with a bitter and more or less hot taste as in *L. chrysorrhoeus*. **Odour** mild in young basidiomata, but intensely fruity when mature.

**Spores**  $(11.5–)12–14.5(–15) \times (10–)11–13 \mu\text{m}$ , Q = 1.1–1.2, subglobose to broadly ellipsoid, orthotropic, echinate; warts 0.5–2  $\mu\text{m}$  high, isolated, inamyloid or weakly amyloid; suprahilar plage



**Fig. 11** *Lactarius stephensii*. a–d. JMV970830-1. a. Basidiomata and hymenophore of immature and mature specimens; b–c. hymenium and rameous subhymenium; d. weakly amyloid spores. — e. JMV951012-1. Basidiomata. — f. JMV20010609-1. Pileipellis. — g. K(M)69330 (lectotype of *Hydnangium stephensii*). Spores in Melzer. — h–j. KRA F-2014-147. h. Spores in Melzer; i–j. SEM images of spores. — Scale bars: a, e = 1 cm; b–c, f = 20  $\mu\text{m}$ ; d, g–h = 10  $\mu\text{m}$ ; i–j = 5  $\mu\text{m}$ . — Photos: a–h. J.M. Vidal; i–j. P. Mleczko, UJ.

absent; hilar appendix inconspicuous, usually united to a fragment of sterigma up to 4 µm long. *Basidia* 1-spored, 35–70 × 6–10 µm, cylindrical, straight to sinuose, soon collapsed; initially hyaline, with a good number of oil drops inside, that become filled with a dark orange substance; sterigmata central or lateral, 4–6 µm long. *Macrocytidia* absent. *Cystidioles* only present in sterile locules, of the same shape and size of dermatocystidia. *Paraphysoid cells* plenty, 20–60 × 6–8 µm, cylindrical or clavate, usually with 1–3 septa. *Pseudocystidia* occasionally observed. *Subhyphmenium* ramos, formed by cylindrical hyphae 10–30 × 3–9 µm, perpendicular to the tramal hyphae. *Hymenophoral trama* homoiomericous, formed by subgelatinized hyphae 2–5 µm diam, with abundant laticifera. *Pileipellis* and *context* 250–500 µm thick; suprapellis a trichoderm formed by yellow, straight to sinuous, septate dermatocystidia 20–60 × 3–6 µm, with rounded to acute, sometimes mucronate apex, that soon collapses in a brown granular mass. *Pileal context* formed by intricate gelatinized hyphae 2–6 µm diam, crossed by some thromboplera and abundant laticifera 3–10(–12.5) µm diam.

**Habitat, Distribution & Season** — Gregarious, hypogeous to semi-hypogeous among plant debris, in montane woods of *Corylus*, *Carpinus*, *Fagus*, *Quercus* accompanied by *Acer*, *Buxus*, *Cornus*, *Fraxinus*, *Sambucus*, *Tilia*, *Ulmus*, on calcareous soil. Summer to autumn. Widely distributed in temperate and submediterranean regions, from almost sea level in the British Isles to 1400 m altitude in Southern Europe.

**Material studied.** CZECH REPUBLIC, Central Bohemia, Karlštejn, in a leafy forest, under *Fagus sylvatica* and *Carpinus*, 10 July 1949, V. Vacek as '*Hydnangium monosporum*' (PRM 685987); ibid., in a leafy forest, under *Quercus*, *Carpinus*, *Acer campester* and *Tilia*, 17 July 1949, V. Vacek as '*Hydnangium monosporum*' (PRM 619116); South Moravia, Dolní Věstonice, Děvičky, in Acereto-Carpinetum, 28 Aug. 1955, K. Kříž, det. M. Svrček as *O. stephensi* (PRM 719216); Veverská Bítýška, Krnovc, under *Carpinus betulus* and *Cornus mas*, 10 Aug. 1955, K. Kříž, det. M. Svrček as *O. stephensi* (PRM 719217); Žarošice, Aug. 1937, V. Vacek as '*Hydnangium carneum*' (PRM 154169, coll. J. Velenovský); ibid., in a leafy forest, 31 Aug. 1948, VI. Vacek & V. Vacek as *H. stephensi* (PRM 685988); ibid., in a leafy forest, under *Tilia*, *Quercus* and *Larix*, 25 Aug. 1949, V. Vacek as *H. galatheum* (PRM 619118); ibid., in a leafy forest, under *Tilia*, *Quercus* and *Larix*, 25 Aug. 1949, V. Vacek as *H. galatheum* (PRM 685993); Ždánský les, near Žarošice, in a leafy forest, 31 Aug. 1950, V. Vacek as *H. galatheum* (PRM 685990); ibid., in a leafy forest, under *Tilia*, *Quercus* and *Fraxinus*, 9 Sept. 1950, V. Vacek as *H. galatheum* (PRM 685992); Zdravá Voda, near Žarošice, in a leafy forest, under *Tilia*, *Carpinus*, *Fagus* and *Cornus mas*, 24 Aug. 1949, V. Vacek as *H. galatheum* (PRM 685996); ibid., in a leafy forest, under *Carpinus*, *Quercus* and *Tilia*, 29 Aug. 1949, V. Vacek as *H. galatheum* (PRM 685994). — HUNGARY, Central Hungary, Budapest, Normafa, under *Tilia* sp., 25 Aug. 1995, I. Király (BCN JMV800071). — ITALY, Lombardy, Bergamo, San Pellegrino Terme, 350 m, under *Corylus avellana*, on calcareous soil, 30 Nov. 2016, M. Berbenni, comm. M. Morara (BCN IC30111603). — POLAND, Silesia, Pogórze Śląskie region, Machowa mountain, Western Carpathians, 15 June 2014, R. Rutkowski (KRA F-2014-147, duplicate BCN JMV800668)\*. — SPAIN, Asturias, Somiedo, Parque Natural de Somiedo, under *Corylus avellana*, 8 Apr. 2009, A. Paz (BCN IC08040913); Catalonia, Girona, Albanyà, Serra de Corsavell, Can Padern, 800 m, under *Corylus avellana*, on calcareous soil, 9 June 2001, J.M. Vidal as *A. stephensi* (BCN JMV20010609-1)\*; Girona, La Vall de Bianya, Vall del Bac, Bac de Mariner, 900 m, under *Corylus avellana*, *Fagus sylvatica*, *Quercus humilis* and *Buxus sempervirens*, on calcareous soil, 30 Aug. 1997, J.M. Vidal as *A. stephensi* (BCN JMV970830-1); Girona, Montagut, Sant Miquel de Pera, 700 m, under *Corylus avellana*, *Quercus humilis*, *Populus tremula* and *Buxus sempervirens*, on calcareous soil, 15 Oct. 1995, J.M. Vidal as *A. stephensi* (BCN JMV951012-1). — UNITED KINGDOM, England, Somerset, Clifton, near Bristol, Leigh Woods, 6 Aug. 1843, C.E. Broome (K(M)69330, lectotype of *H. stephensi*); ibid., Bristol, 1844, H.O. Stephens (K(M)69331, original material of *H. stephensi*); ibid., Bristol, sine dat., ex M.J. Berkeley (PRM 719218, herb. A.C.J. Corda, original material of *H. stephensi*).

**Notes** — *Hydnangium stephensi* is a lactescent angiocarpic fungus described by Berkeley (1844) from Leigh Woods, North Somerset (UK), which was recombined in the genus *Zelleroomyces* by Smith (1962) based on the presence of amyloid spores and laticifera. It is macroscopically characterized by maize yellow to reddish brown basidiomata bleeding white latex

(scant and hyaline when young) that turns yellow in contact with air, and microscopically, by its monosporic basidia and warty subglobose to broadly ellipsoid spores, which are inamyloid to weakly amyloid (Vidal 2004a, Fraiture & Derboven 2009, all as *A. stephensi*).

Nuytinck et al. (2003) suggested *L. stephensi* belongs to *Lactarius* subg. *Lactarius*, and considered it a priority synonym of *Hydnangium monosporum*, while Vidal (2004a) considered *L. stephensi* (treated as *Arcangiella stephensi*) to be a priority synonym of *Hydnangium soehneri*. Following careful studying of the types and other collections in the present study, *L. stephensi*, *H. soehneri* and *H. monosporum* are here considered to be three distinct species. *Hydnangium monosporum* differs by its perfectly rounded and pink-coloured spores when mature, and *H. soehneri* by its bigger, strongly amyloid and ellipsoid spores. It was not possible to locate original material of *H. galatheum* (Quélet 1875 as '*stephensi*', 1886 as '*galatheum*'), but only a collection made later by Quélet in August 1892 at Hérimoncourt (UPS F013405), which we have identified as *L. populicola*, and another collection without date from Abbévillers (UPS F013936), which we have identified as *Russula monospora*. However, based on descriptions and illustrations provided by Quélet (1875: 446, pl. I, f. 9, as *H. stephensi*) and (1886: 247, as *H. galatheum*), we consider, like Patouillard (1910, 1914) and Zeller & Dodge (1937), *H. galatheum* to be a synonym of *L. stephensi*. The only genetic sequence available in GenBank before the present work from *L. stephensi*, was an ITS rDNA sequence of K(M)64067 specimen (Brock et al. 2009), collected by Brian Spooner in 1999 at North Somerset (UK), probably close to the locality where the type was found. This sequence clearly differs from that produced from specimen RW2930 (GENT) by Nuytinck et al. (2003), which is here considered to represent *L. populicola*.


### *Lactarius subgiennensis* Loizides, J.M. Vidal & P. Alvarado, sp. nov. — MycoBank MB828499; Fig. 12

**Basidiomata** 0.5–2.5 cm wide, angiocarpic, sessile, globose to subglobose or lobate, lactescent, scrobiculate-porate, ivory white. *Hymenophore* loculate, whitish to faintly pinkish. *Latex* white. *Spores* 8.5–11 × 6.5–8 µm, ovoid, reticulated; reticulum 0.5–1 µm high, partially interrupted. *Basidia* 1-spored, 30–42 × 5–9 µm, lageniform-urticiform. *Cystidioles* 32–40 × 5–8 µm, lageniform-urticiform, present only in external locules. *Macrocytidia* absent. *Hymenophoral trama* homoiomericous, prosenchymatous, gelatinized, with laticifera 3–9 µm diam. *Suprapellis* formed by repent to semi-erect, septate hyphae 4–6 µm wide. *Pileal context* prosenchymatous, gelatinized, with abundant laticifera and some thromboplera 4–14 µm diam. Semi-hypogeous in *Cistus* maquis on the island of Cyprus.

**Etymology.** From Latin, *sub* = nearly, close to, and *-giennensis*, referring to its close phylogenetic and morphological affinity to *Lactarius giennensis*.

**Holotype.** CYPRUS, Larnaca District, Kalavasos, 170 m, under *Cistus salviifolius* and *Cistus creticus*, 25 Nov. 2012, M. Loizides (BCN JMV800627)\*; isotype in herb. pers. M. Loizides (ML211152E).

**Basidiomata** 0.5–2.5 cm wide, angiocarpic, irregularly globose, subglobose or tuberiform to weakly lobulate, occasionally more distinctly lobulate to almost irregularly-shaped, sessile, lactescent, loosely attached to the substrate by rhizoids. *Pileus* smooth, silky, subtomentose in parts, moderately to distinctly scrobiculate and porate, ivory white (5A2–6A2), often with ochraceous stains at maturity (5A4–A5), overripe basidiomata sometimes brownish buff (6C6); greyish orange (5B5) in excise-cata. *Hymenophore* loculate, labyrinthoid, white when young, at maturity becoming cream (6A2–A3) or pale ochraceous salmon (7A2); locules 0.5–1 mm long (1–3 per mm), polygonal to somewhat elongated, irregular, mostly empty. *Columella* absent or rudimentary, but thin, sterile veins frequently intersecting the hymenophore. *Latex* abundant in humid conditions, white, unchanging. *Odour* acidic, slightly fruity.



**Fig. 12** *Lactarius subgiennensis*. a. ML61132Z. Basidiomata. — b. ML211192Z. Basidiomata. — c–l. ML211152E (BCN JMV800627, holotype). c. Pileipellis and context with one thromboplera and some laticifera; d. detail of the pileipellis; e–f. hymenium (basidia, basidioles and paraphysoid cells), subhymenium and hymenophoral trama with some laticifera; g. cystidioles of external immature locule; h. emerging necks of basidia; i. basidia; j. spores in Melzer; k–l. SEM images of spores. — Scale bars: a–b = 1 cm; c–f = 20 µm; g–j = 10 µm; k–l = 5 µm. — Photos: a–b. M. Loizides; c–j. J.M. Vidal; k–l. UdG.

Spores  $8.5–11 \times 6.5–8 \mu\text{m}$ ,  $Q = 1.2–1.4$ , ovoid, orthotropic, reticulated; reticulum strongly amyloid, formed by well-developed crests projecting  $0.5–1 \mu\text{m}$  in side view, partially interrupted to nearly complete at full maturity; hilar appendix indistinct or rudimentary. *Basidia* 1-spored (rarely 2-spored),  $30–42 \times 5–9 \mu\text{m}$  (neck included), lageniform-urticiform, promptly collapsed, with a long and sinuous neck measuring  $20–25 \times 2.5–4 \mu\text{m}$ , protruding the hymenium; sterigmata  $\pm 1 \mu\text{m}$  long, inconspicuous. *Basidioles*  $22–24 \times 4.5–6 \mu\text{m}$ , ventricose-rostrate. *Macro-cystidia* absent. *Cystidioles*  $32–40 \times 5–8 \mu\text{m}$ , lageniform-urticiform, observed only in immature external locules. *Paraphysoid cells*  $9–25 \times 5–8.5 \mu\text{m}$ , cylindrical to subclavate, entire or 1-septate. *Subhymenium* rameous to cellular, composed of chains of rounded, polygonal or irregularly elongated cells  $5–10 \mu\text{m}$  diam. *Hymenophoral trama*  $90–120 \mu\text{m}$  wide, homoiomerous, prosenchymatous, formed by intricate, tortuous, thin-walled hyphae  $3–6 \mu\text{m}$  diam, with plenty of lipidic guttules, finally of pseudoparenchymatous aspect, moderately to strongly gelatinized, but without the presence of sphaerocytes; laticifera  $3–9 \mu\text{m}$  diam, thick-walled, tortuous, often branching, sometimes forming projecting pseudocystidia. *Pileipellis* and *context* thin,  $50–100 \mu\text{m}$  thick; pileipellis composed of: 1) an easily

collapsible trichodermal suprapellis, comprised of terminal, septate hyphae  $4–6 \mu\text{m}$  diam; and 2) a prosenchymatous subpellis of gelatinized, tightly packed, tortuous, septate hyphae  $3–6 \mu\text{m}$  diam. *Pileal context* undifferentiated from the subpellis with abundant laticifera and some thromboplera  $4–11 \mu\text{m}$  diam, locally enlarged up to  $14 \mu\text{m}$ .

**Habitat, Distribution & Season** — Gregarious to caespitose, growing semi-hypogaeously in thick litter, exclusively associated with *Cistus*, on siliceous soil. Autumn and winter. Xerophytic. So far restricted in the eastern Mediterranean region on the island of Cyprus, at altitudes below 400 m.

**Additional material studied.** CYPRUS, Larnaca District, Kalavasos, 170 m, under *Cistus salvifolius* and *Cistus creticus*, 29 Nov. 2012, M. Loizides (ML211192Z); ibid., 200 m, 23 Jan. 2016, M. Loizides (ML61132Z)\*; Nicosia District, Lythrodontas, 400 m, under *Cistus salvifolius* and *Cistus creticus*, 27 Nov. 2014, M. Loizides (ML411172Z)\*.

**Notes** — *Lactarius subgiennensis* differs from the closely related *L. giennensis* in its markedly scrobiculate basidiomata, lageniform-urticiform basidia, and smaller, subreticulated spores with a partially incomplete or discontinuous reticulum, not exceeding  $1 \mu\text{m}$  height in side view. Both species are found in xerophytic environments of Southern Europe in association

with *Cistaceae* plants, but seem to be geographically isolated. Additional collections from intermediate localities (Greece, Italy) would be needed to confirm the distribution boundaries between both taxa. Genetically, *L. subgiennensis* seems to be the sister-species of *L. giennensis*. Both taxa seem to have some intraspecific variability, but are significantly different.

**Russula ammophila** (J.M. Vidal & Calonge) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 231. 2018 — Fig. 13

*Basionym.* *Gymnomyces ammophilus* J.M. Vidal & Calonge, Bol. Soc. Micol. Madrid 24: 66. 1999.

*Synonym.* *Macowanites ammophilus* (J.M. Vidal & Calonge) J.M. Vidal & Calonge, Rev. Catalana Micol. 24: 70. 2002.

*Basidiomata* russuloid, angiocarpic to pseudoangiocarpic, stipitate. *Pileus* 2–7 cm wide, globose to hemispherical, orange white to pale orange, with brownish orange and dark brown maculae; margin open, alveolate to sublamellate. *Hymenophore* loculate, orange-white to pale orange. *Stipe-columella* 1–4 × 0.7–2.5 cm, with brownish orange dots; context white. *Odour* fruity; taste sweetish to acrid.

*Spores* 7–9 × 5.5–7.5 µm, Q = 1.16–1.4, subglobose to ovoid, heterotrophic; warts 0.25–0.75 µm high, amyloid, some forming short ridges or connected by short lines. *Basidia* 2–4-spored, 35–45 × 8–11 µm, clavate. *Macrocytidia* 45–60 × 6–10 µm. *Hymenophoral trama* with sphaerocytes 6–35 µm diam. *Pileipellis* 125–200 µm thick; suprapellis an intricate trichoderm made of sinuous, clavate, lanceolate or mucronate tips of hyphae and some poorly differentiated dermatocystidia 15–40 × 2–5 µm; subpellis an intricate cutis of gelatinized hyphae 2.5–7.5 µm diam. *Pileal* and *stipe-columella* context heteromerous.


*Habitat, Distribution & Season* — Gregarious, hypogeous to semi-hypogeous in sandy soils, associated with *Pinus*. Spring and autumn. Common in sandy pine forests of the Atlantic coast of Southern Spain and Portugal, more rarely in Northern and Central Spain.

*Material studied.* PORTUGAL, Setubal, road from Grândola to Santa Margarita do Sado, stabilized dunes, under *Pinus pinea* and *Pinus pinaster* with *Halimium halimifolium* and *Corema*, 26 Mar. 1998, P.P. Daniels, J.M. Vidal & F.D. Calonge as *G. ammophilus* (MA-Fungi 40132; duplicate BCN JMV980326-11b); Castelo Ventoso, road to Alcácer do Sal, stabilized dunes, under *Pinus pinea* with *Corema*, 26 Nov. 1999, J.M. Vidal & F.D. Calonge as *M. ammophilus* (MA-Fungi 51167; duplicate BCN JMV991126-1); without loc., without date, as *M. ammophilus* (AH 42956). — SPAIN, Andalusia, Cádiz, Barbate, Loma

del Teniente, 120 m, stabilized dunes, under *Pinus pinea* with *Halimium*, 15 Dec. 2012, M. Becerra & E. Robles (AH 43950, as *M. ammophilus*)\*; Cádiz, Sanlúcar de Barrameda, Pinar de Algaida, stabilized dunes, under *Pinus pinea*, 30 Oct. 2014, M. Becerra as *M. ammophilus* (AH 46370)\*; Huelva, Matalascañas, stabilized dunes, under *Pinus pinea* with *Halimium*, *Corema* and *Juniperus phoenicea*, 27 Nov. 1999, J.M. Vidal & F.D. Calonge as *M. ammophilus* (MA-Fungi 51165; duplicate BCN JMV991127-8); Mazagón, stabilized dunes, under *Pinus pinea* with *Halimium* and *Corema*, 27 Nov. 1999, J.M. Vidal & F.D. Calonge as *M. ammophilus* (MA-Fungi 51166; duplicate BCN JMV991127-2); Castilla and Leon, Palencia, Dehesa de Montejo, Tosande, under *Pinus nigra* subsp. *salzmannii* and *Quercus rotundifolia*, in sandy basic soil, 9 Jan. 2007, A. Paz (BCN IC09010703)\*.

*Notes* — *Gymnomyces ammophilus* was proposed to accommodate specimens of a fully angiocarpic basidiomycete without any trace of a stipe-columella, found in South-Western Spain and Portugal (Calonge & Vidal 1999). However, new specimens with an evident stipe-columella were later collected by the same authors (Vidal et al. 2002) and DNA data suggested they all represent a single species, so it was concluded that the holotype material was not fully mature and the taxon was re-combined as *Macowanites ammophilus*. Since its formal description, this species has been regularly collected in coastal sand dunes of the South-Western Iberian Peninsula and a high morphological variability has been observed. Recently, this species was re-combined into genus *Russula* by Elliott & Trappe (2018).

A high genetic variability (5.8 %) among ITS sequences from these specimens can be observed (Fig. 3). This could be due in part to sequencing errors in older GenBank accessions (M.P. Martín, pers. comm.), although some variability was found also among newly produced sequences. Most interestingly, ITS sequences did not support significant differences between *R. ammophila* and sequences of the gymnocarpic species *R. amoena* produced in the present work and those present in public databases, including one obtained from a specimen identified by Romagnesi (MICH12838, KF245510, Bazzicalupo & Berbee, unpubl. data). These European sequences probably represent the original concept of *R. amoena*, and are significantly different from other lineages identified with the same name (Fig. 3). In addition, 28S rDNA and *rpb2* data failed to support any significant difference between *R. ammophila* and *R. amoena*, but translation elongation factor 1-alpha (*tef1*) data contains diagnostic differences that can be employed to discriminate them (GenBank MK102724–MK102727). Owing to



**Fig. 13** *Russula ammophila*. a. MA-Fungi 51167 (as *Macowanites ammophilus*). Basidiomata. — b–d. IC09010703. b. Pileipellis; c–d. hymenium. — e. JMV991127-2. Spores in Melzer. — f–g. JMV991126-1. SEM images of spores. — Scale bars: a = 1 cm; b–c = 20 µm; d–e = 10 µm; f–g = 5 µm. — Photos: a–e. J.M. Vidal; f–g. UdG.

genetic differences in *tef1* gene and the lack of a stabilized concept of *R. amoenolens*, we here still accept that *R. ammophila* represents an independent species. Both taxa belong to *R. sect. Ingratae*, and are nested within the clade of /*R. amoenolens*, closely related with *R. cerolens*, *R. sororia*, and several other lineages identified as *R. amoenolens* not matching Romagnesi's concept of this species. In *R. amoenolens* spores are ellipsoid, measuring 7–8.5(–9) × 5–6.7 µm and ornamented with warts up to 0.75 µm high (Romagnesi 1985), or 6.5–8.4 × 5.2–6.4 µm and with warts up to 0.8 µm high (Sarnari 1998), some warts interconnected with short ridges. Spores of *R. ammophila* are subglobose to ovoid, measuring 7–9 × 5.5–7.5 µm and ornamented with warts up to 0.75 µm high, some of them fused and connected by short ridges.

*Russula amoenolens* Romagn., Bull. Mens. Soc. Linn. Lyon 21: 111. 1952.

*Material studied.* SPAIN, Cantabria, Castro Urdiales, Prado Maya, under *Quercus* sp., 23 July 2015, J.A. Cadiñanos (AH 46371)\*; Castro Urdiales, under *Quercus* sp., 23 July 2015, J.A. Cadiñanos (AH 46372)\*.

***Russula andaluciana*** T.F. Elliott & Trappe, Fungal Systematics and Evolution 1: 239. 2018 — Fig. 14

Replaced synonym. *Gymnomycetes sublevisporus* Mor.-Arr., Llistos. & L. Romero, Rev. Catalana Micol. 24: 179. 2002.


*Basidiomata* 0.5–2 cm wide, angiocarpic, globose to subglobose, somewhat lobate, sessile. *Pileus* smooth, pruinose, whitish, with brownish red maculae. *Hymenophore* loculate, whitish at first, finally brownish red. *Columella* percurrent, sometimes branched, inconspicuous. *Odour* of bitter almonds; taste mild, slightly bitter.

*Spores* (6.5)–7.5–9.5(–11.5) × (6)–7–9(–11) µm, Q = 1.0–1.1, globose to subglobose, orthotropic; warts up to 0.3 µm high, amyloid, some connected with low ridges. *Basidia* 4-spored, 36–50 × 7.5–11 µm, clavate. *Macrocystidia* and cystidioles absent. *Hymenophoral trama* prosenchymatous, of dense interwoven, tortuous hyphae about 2–6 µm diam, with some inflated ampullaceous cells up to 10 µm diam. *Pileipellis* and *context* 125–200 µm thick; suprapellis a trichoderm formed by short, cylindrical to clavate hairs 9–18 × 5–7 µm, and long, subulate, sometimes 1–2-septate hairs 30–49 × 2–7 µm; dermatocystidia absent; subpellis made by ampullaceous cells 8–10 µm diam. *Pileal context* with the same structure as the hymenophoral trama.

**Habitat, Distribution & Season** — Gregarious, hypogeous, associated with *Cistus ladanifer*, often accompanied by *Quercus rotundifolia* or *Q. suber*, on siliceous soil. Spring. Located in the Western Mediterranean region, in Central Spain, between 400–800 m altitude.

*Material studied.* SPAIN, Andalusia, Huelva, Cortelazor, finca Galindo, 540 m, under *Cistus ladanifer* with *Cistus crispus*, *Cistus salvifolius* and *Quercus rotundifolia*, 11 Mar. 1995, L. Romero de la Osa (BCN JL5101, holotype of *G. sublevisporus*; BM360, isotype)\*; Extremadura, Badajoz, Mirandilla, Sierra Bermeja, 400 m, under *Cistus ladanifer* with *Quercus suber*, on siliceous soil, 25 Mar. 2017, J.L. Becerra (BCN IC25031720); Cáceres, Cuesta de Jaraiz, under *Cistus ladanifer*, on siliceous soil, 2 Apr. 2011, C. Gelpí as *G. sublevisporus* (AH 39239)\*; Cáceres, Cuesta de las Veguillas, under *Cistus ladanifer* with *Eucalyptus camaldulensis*, on siliceous soil, 2 Mar. 2011, C. Gelpí & J. Muñoz as *G. sublevisporus* (AH 39198)\*.

**Notes** — As already mentioned above, *R. andaluciana* is a replacement name proposed by Elliott & Trappe (2018) to re-combine *G. sublevisporus* into *Russula*. This species is distinguished by its whitish to yellowish basidiomata, and by its globose spores, ornamented with very small verrucae not



**Fig. 14** *Russula andaluciana*. a. AH 39239. Basidiomata. — b. AH 39198. Basidiomata — c–i. BM360 (isotype of *Gymnomycetes sublevisporus*). c. Pileipellis; d. hymenium (basidia and paraphysoid cells); e–f. spores in Melzer; g–i. SEM images of spores. — Scale bars: a–b = 1 cm; c–d = 20 µm; e–f = 10 µm; g–i = 5 µm. — Photos: a–b. C. Gelpí; c–f. J.M. Vidal; g–i. UdG.

exceeding 0.3 µm high. Sphaerocytes were not found in our observations, neither in the hymenophoral trama nor in the pileal context. Genetically, *R. andaluciana* is the only known European sequestrate species in *Russula* sect. *Rigidae*. It is nested within *R. subsect. Amoeninae*, but shows no clear relationship with any other taxon (Fig. 4).

***Russula bavarica* J.M. Vidal, sp. nov.** — MycoBank MB828500; Fig. 15

**Basidiomata** 1–2 cm wide, angiocarpic, sessile, subglobose to tuberiform, subtomentose, pale orange with brown maculae. **Hymenophore** pale orange, forming small cells or locules. Spores 13–15(–15.5) × 12.5–14.5(–15) µm, globose to subglobose, orthotropic, intensely yellow, weakly amyloid, ornamented with dense, curved warts, 1.5–3 µm long. **Basidia** 1-spored, 24–36 × 8–12 µm, clavate, hollow; sterigmata 3–5 µm long, conical. **Basidioles** 15–28 × 6–10 µm, clavate. **Macrocystidia** 30–70 × 8–16 µm, clavate, abundant, containing refringent crystals (in exsiccata). **Subhymenium** ramoso. **Hymenophoral trama** homoiomerous. **Suprapellis** a trichoderm of repent to semi-erect hyphae and dermatocystidia 50–100 × 2–4 µm; pileal context a subgelatinized cutis.


**Etymology.** From Latin, *bavaricus* = refers to the German state of Bavaria, where mycologist Ert Soehner collected this sequestrate fungus.

**Holotype.** GERMANY, Bavaria, 'Wolfratshausen bei München, Pupplinger Heide in der Nähe der Aumühle', 8 July 1923, E. Soehner #724 as '*Hydnangium carneum*' (M); isotype in FH (herb. C.W. Dodge).

**Basidiomata** 1–2 cm wide, angiocarpic, subglobose to tuberiform, sessile. **Pileus** finely tomentose, white to pale orange (5A4) in young specimens, partially covered with brownish maculae (7E6) in mature basidiomata. **Hymenophore** loculate, labyrinthoid, pale orange in colour (5A4–A5); locules 0.4–1 × 0.1–0.4 mm, elongated, sinuous; spore mass in locules pale orange (5A3). **Columella** absent. **Odour** and taste unknown. (Description based on herbarium material).

**Spores** 13–15(–15.5) × 12.5–14.5(–15) µm, Q = 1.0–1.1, globose to subglobose, orthotropic, echinate, hyaline when young, yellowish when mature; warts inamyloid or weakly amyloid, 1.5–3 µm long, dense, curved, sharp, with some verrucae among them; hilar appendix short, 0.5–1 µm long, often with remnants of the sterigmata attached. **Basidia** 1-spored, 24–36 × 8–12 µm, clavate, hollow; sterigmata 3–5 µm long, conical. **Basidioles** 15–28 × 6–10 µm, clavate. **Macrocystidia** 30–70 × 8–16 µm, clavate, abundant, filled with highly refringent hyaline needles. **Pseudocystidia** not observed. **Subhymenium** very reduced, ramoso, formed by short chains of 1–3 cylindrical elements, measuring 3–10 × 3–7 µm. **Hymenophoral trama** 20–40 µm wide, homoiomerous, formed by an entanglement of branched, tortuous, septate, subgelatinized hyphae about 2–6 µm diam, including scattered endomacrocystidia; sphaerocytes absent. **Pileipellis** and **context** thin, 100–150 µm thick; pileipellis 25–50 µm thick, arranged as a trichoderm formed by hyphae and septate, repent or semi-erect dermatocystidia 50–100 × 2–4 µm, finally collapsing into a brownish mass. **Pileal context** 75–100 µm thick, formed by a cutis of intricate, subgelatinized hyphae 2–5 µm diam, connecting with the hymenophoral trama. **Laticifera** not observed. **Gloeoplera** abundant, 2–5 µm diam, present in trama and context.

**Notes** — Collection Soehner num. 724 was cited by Dodge & Zeller (1937) and Soehner (1941), under the name *Hydnangium carneum*. It was found by Soehner in July 1923 near Munich (Bavaria, Germany). After studying the specimens preserved nowadays in M and FH herbaria, amyloid spores, hymenial macrocystidia, dermatocystidia and gloeoplera were observed, suggesting this species does not belong to genus *Hydnangium*. These specimens resemble *R. monospora*, because of the presence of monosporic basidia, echinate and



**Fig. 15** *Russula bavarica*. a–i. M, herb. E. Soehner 724 as '*Hydnangium carneum*' (holotype). a. Pileipellis and context; b. suprapellis; c. macrocystidia; d–e. basidia, macrocystidia and ramoso subhymenium; f. hymenophoral trama with some gloeoplera; g. spores in Melzer; h–i. SEM images of spores. — Scale bars: a–f = 20 µm; g = 10 µm; h–i = 5 µm. — Photos: a–g. J.M. Vidal; h–i. UdG.

weakly amyloid spores, rameose subhymenium, homoiomerous hymenophoral trama, and trichodermal pileipellis. Although it has not been possible to obtain recent collections matching these specimens, we believe it necessary to accommodate the collection Soehner 724 in a new species of *Russula* because of its unique combination of microscopical features, very similar to those of *R. monospora*, but differing and being characterized by the presence of clavate macrocystidia and larger spores ornamented with longer and more acute and dense spines. Further collections are needed to confirm the ecology of this species, which is probably the same as *Lactarius soehneri*, whose holotype was also collected by Soehner in Pupplinger Heide (south of Munich), and also determined as *H. carneum*.

***Russula candida*** (Tul. & C. Tul.) J.M. Vidal, comb. nov. — MycoBank MB828501; Fig. 16

Basionym. *Hydnangium candidum* Tul. & C. Tul., Ann. Sci. Nat., Bot., Sér. 2, 19: 376. 1843.

Synonyms. *Octaviania candida* (Tul. & C. Tul.) Lloyd, Mycol. Writings 7, Letter 67: 1142. 1922.

*Sclerogaster candidus* (Tul. & C. Tul.) Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 23: 570. 1937.

*Macowanites candidus* (Tul. & C. Tul.) J.M. Vidal, Rev. Catalana Microl. 26: 84. 2004.

*Secotium krjukowense* Bucholtz, Hedwigia 40: 314. 1901.

*Elasmomyces krjukowensis* (Bucholtz) Sacc. & D. Sacc. in P.A. Saccardo, Syll. Fung. 17: 218. 1905.

*Bucholtzia krjukowensis* (Bucholtz) Lohwag, Oesterr. Bot. Z. 73, 7–9: 173. 1924.

*Arcangelia krjukowensis* var. *krjukowensis* (Bucholtz) Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 22: 368. 1935.

*Hydnangium krjukowense* var. *krjukowense* (Bucholtz) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 206. 1958.

*Macowanites krjukowensis* (Bucholtz) Singer & A.H. Sm., Mem. Torrey Bot. Club 21, 3: 77. 1960.

*Russula krjukowensis* (Bucholtz) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 235. 2018.

*Secotium michailowskianum* Bucholtz, Hedwigia 40: 315. 1901.

*Elasmomyces michailowskianus* (Bucholtz) Sacc. & D. Sacc. in P.A. Saccardo, Syll. Fung. 17: 218. 1905 ('*michailowskjanus*').

*Arcangelia krjukowensis* var. *michailowskiana* (Bucholtz) Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 22: 368. 1935.

*Secotium krjukowense* forma *pleurotopsis* Bucholtz, Bull. Soc. Imp. Naturalistes Moscou 4: 463. 1908 (1907).


*Octaviania moravica* Velen., Opera Bot. Cech. 4: 95. 1947.

*Hydnangium krjukowense* var. *moravicum* (Velen.) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 208. 1958.

*Basidiomata* russuloid, angiocarpic, stipitate. *Pileus* 0.5–3.5 cm wide, globose to subglobose, sometimes lobate, smooth, pruinose, white to yellowish white (4A2), with yellowish orange (4A6) to dark orange (5A8) maculae; margin closed or laterally open, alveolate. *Hymenophore* loculate, labyrinthoid, pale orange (5A3–A4); locules 0.7–3 × 0.1–0.4 mm (1–2 per mm), elongated, irregular, sinuous; fresh spore mass in locules pale yellow (3A3–A4). *Stipe-columella* 0.6–1.5 × 0.15–0.4 cm, usually half stipe, half columella, cylindrical to clavate, central or lateral, sometimes deeply lateral and thus barely visible, pure white, pruinose; context white, not changing on exposure to air. *Odour* fruity; taste mild.

*Spores* 8.5–11(–12.5) × 7–9(–11) µm, Q = 1.08–1.25(–1.3), subglobose to broadly ellipsoid, heterotropic to subheterotropic, echinate; warts amyloid, 0.5–1.5 µm high, isolated, cylindrical and sometimes curved, with a rounded apex; hilar appendix 1–2.5 × 1.2 µm, straight, cylindrical to conical, occasionally retaining a sterigmal appendage; suprahilar plage amyloid; immature spores sometimes with a non-functional apicular drop. *Basidia* 1–3(–4)-spored, 28–42 × 14–18 µm, clavate. *Basidioles* clavate, 20–24 × 10–16 µm. *Macrocytidia* 45–70 × 8–12 µm, lanceolate or fusiform, rostrate, narrowly acute or mucronate, easily collapsing. *Paraphysoid cells* absent. *Subhymenium* cellular, formed by subglobose cells of 8–16 µm diam. *Hymenophoral trama* 40–50 µm wide, formed by septate hyphae 1.5–10 µm diam, with frequent enlargements up to 16 µm and abundant sphaerocytes 15–40 µm diam. *Pileipellis* and *context* 80–250 µm thick, thinning in the perimarginal zone; suprapellis a trichoderm of cylindrical dermatocystidia 15–70 × 3–8 µm, with tips of gelatinized hyphae soon collapsing into a brown, granular slimy mass; mediopellis 60–100 µm thick, that consists of an ixocutis of intricate hyphae 2–6 µm diam, with numerous gloeoplera 4–5 µm diam terminating in dermatocystidia; subpellis 50–120 µm thick, formed by an intricate cutis of subgelatinized hyphae 2–6 µm diam, connecting with the pileus and hymenophoral trama. Context of pileus and stipe-columella heteromerous. *Stipitipellis* a turf of repent to erect hyphal tips mixed with dermatocystidia. *Thromboplera* 3–5 µm diam, present in the trama and context.

*Habitat, Distribution & Season* — Solitary to gregarious, hypogeous to semi-hypogeous among plant debris, in montane broadleaved woods of *Carpinus* and *Corylus* mixed with *Fagus*, *Populus*, *Quercus*, *Tilia*, *Betula*, *Acer*, *Fraxinus*, *Sambucus*, *Ulmus*, on calcareous soil. Summer and autumn. Widely dis-



**Fig. 16** *Russula candida*. a–e. JMV20100724b. a. Basidiomata; b. hymenium and subhymenium; c. spores in Melzer; d–e. SEM images of spores. — f–g. M, herb. G. Gross 171 (as '*Elasmomyces mattiroloanus*'). f. Pileipellis; g. spores in Melzer. — Scale bars: a = 1 cm; b, f = 20 µm; c–d, g = 10 µm; e = 5 µm. — Photos: a–c, f–g. J.M. Vidal; d–e. UdG.

tributed in temperate regions of Europe, from 100 m altitude in Northern Europe to 1100 m around the Mediterranean basin.

**Material studied.** BULGARIA, Vraca, Chiren, under *Tilia* sp., 23 June 2018, M. Slavova & O. Ligabue (MSL2177F3265, duplicate BCN JMV800751). – CZECH REPUBLIC, Central Bohemia, Černošice, under *Carpinus*, *Tilia* and *Acer*, 9 July 1950, V. Vacek as *E. krukowensis* (PRM 619106); South Moravia, Ždanice, near Bučovice, in forest, July 1941, F. Neuwirth (PRM 151648, holotype of *O. moravica*); Ždanice, near Zdravá Voda, in a leafy forest, July 1960, K. Kríž, det. M. Svrček as *E. krukowensis* (PRM 620134). – FRANCE, Poitou-Charentes, Vienne, Couhé-Vérac, solitary under *Carpinus*, Oct. 1841, Tulasne (PC, herb. Tulasne, holotype of *H. candidum*). – GERMANY, Saarland, Gerlfangen, under *Fagus*, *Carpinus*, *Quercus* and *Acer*, 26 July 1968, G. Gross as '*E. mattioloanus*' (M, herb. G. Gross 171; duplicate BCN JMV799894)\*; Honzrath, under *Fagus*, *Carpinus* and *Corylus*, on calcareous soil, 27 June 1968, G. Gross as '*E. mattioloanus*' (M, herb. G. Gross 176). – HUNGARY, Nógrád, Litke, under *Carpinus*, on calcareous soil, 20 Aug. 1996, Z. Lukács (BCN JMV800073); Tötökmező, 250 m, under *Carpinus*, on calcareous soil, 1 Aug. 1996, Z. Lukács (BCN JMV800074). – ITALY, Lombardy, Monza e Brianza, Monza, Viale Reale, under *Carpinus*, 15 Oct. 1989, M. Sarasinias as '*E. mattioloanus*' (MS394, duplicate BCN JMV800003); Monza e Brianza, Monza, Viale Romano, under *Carpinus*, 15 Oct. 1992, M. Sarasinias as '*E. mattioloanus*' (MS467, duplicate BCN JMV800004); ibid., 19 Sept. 1994, M. Sarasinis & G. Imperatori as '*E. mattioloanus*' (MS619, duplicate BCN JMV800005). – RUSSIA, Moscow, Kryukovo, Kryukovo station, 18 July 1899, F. Bucholtz (FH, herb. F. Bucholtz, original material of *S. krukowense*); Podolsk, Michailowskoje station, 20 July 1899, F. Bucholtz (FH, herb. F. Bucholtz 290, holotype of *S. michailowskianum*). – SPAIN, Cantabria, Saja, Reserva Natural del Saja, 658 m, under *Fagus sylvatica*, 1 Oct. 2017, J. Cabero (JC171001N)\*; Catalonia, Girona, Molló, Coll de la Boixeda, El Querol, 1100 m, under *Corylus avellana*, *Quercus pubescens*, *Tilia platyphyllos* and *Populus tremula*, on calcareous soil, 24 July 2010, M.A. Pérez-De-Gregorio (BCN JMV20100724b)\*; ibid., 6 Sept. 2011, J.M. Vidal & F. Rodríguez (BCN JMV20110906-6b).

**Notes** — Tulasne & Tulasne (1843) collected in the French region of Vienne a fungus of angiocarpic appearance devoid of a stipe-columella, which they named *Hydnangium candidum*. Vidal (2004b) studied the holotype of *H. candidum* preserved in Tulasne's herbarium at PC, as well as the holotype of *Octaviania moravica* preserved in Velenovský's herbarium at PRM and part of the original material of *Secotium krukowense* and *S. michailowskianum* preserved in Bucholtz's herbarium at FH, and concluded that all these taxa represent a single species, which he re-described as *Macowanites candidus*. *Russula candida* is characterized by its whitish angiocarpic basidiomata, which are maculated with yellowish orange, sometimes developing a minute stipe and a conspicuous columella. Spores are warty, subglobose to broadly elliptical, measuring 8.5–11(–12.5) × 7–9(–11) µm, many of them with a minute suprahilar plage in the base of hilar appendix and conserving a non-functional apical drop, a character that is also present in other pseudo-angiocarpic species of the same clade (*R. mattioloana* and *R. mediterraneensis*).

This species is genetically nested within *Russula* sect. *Russula*, subsect. *Maculatinae*, and shows a significant relationship with *R. maculata* and *R. nympharum*, although it seems to represent an independent lineage not identified in previous studies of this clade (Adamčík et al. 2016). The complex around *R. maculata* was found by Adamčík et al. (2016) to be characterized by spores which are ornamented with low or medium-sized warts frequently connected and forming small ridges. However, *R. candida* has spiny spores not forming a reticulum or ridges, suggesting that this feature could be variable, or else restricted to the lineage formed by *R. maculata* and *R. nympharum*.

***Russula candidissima* J.M. Vidal, Pasabán & Chachuła, sp. nov.** — MycoBank MB828502; Fig. 17

**Basidiomata** russuloid, angiocarpic to partially pseudoangiocarpic, stipitate. **Pileus** 1.4–2.8 cm wide, globose to lobate, subtomentose, pure white, later featuring pale yellow maculae; margin closed or laterally open, sublamellate. **Hymenophore** loculate, pale yellow to yellow. **Stipe-columella** 1–2 ×

0.2–0.4 cm, percurrent, white. **Odour** fruity. **Spores** 9.5–12.5 × 8.5–10.5 µm, subglobose to broadly ellipsoid, heterotropic, amyloid; warts 1.2–2 µm high, isolated or in groups of 2–4, commonly interconnected by low ridges. **Basidia** 2–4-spored, 37–55 × 14–18 µm, broadly clavate. **Macrocytidia** 45–65 × 6–14 µm. **Hymenophoral trama** with sphaerocytes measuring 20–36 µm diam. **Suprapellis** a cutis of repent hyphae 3–8 µm diam, and some dermatocystidia measuring 20–40 × 3–7 µm; **pileal context** a cutis with some sphaerocytes up to 15 µm diam. Hypogeous or semi-hypogeous between plant debris, in fir and broadleaved montane woods.


**Etymology.** From Latin, *candidissima* = superlative for *candidus* (white), meaning very white because of its white colour more persistent than that of *Russula candida*.

**Holotype.** SPAIN, Catalonia, Girona, Molló, Coll de la Boixeda, El Querol, 1100 m, gregarious under *Corylus avellana*, *Quercus pubescens*, *Tilia platyphyllos* and *Populus tremula*, occurring in the same habitat as *Russula candida*, on calcareous soil, 6 Sept. 2011, J.M. Vidal & F. Rodríguez (BCN JMV20110906-6a)\*.

**Basidiomata** similar to *Russula candida*, russuloid, angiocarpic to partially hemiangiocarpic, stipitate. **Pileus** 1.4–2.8 cm wide, initially globose or subglobose, finally lobate, with a flattened to depressed apex, laterally open and sublamellate at the base, exposing the hymenophore; initially finely tomentose, then smooth, pure white for a long time, staining pale yellow with manipulation (4A4); greyish orange (5B5) to pale brown in exsiccata (6D7). **Hymenophore** loculate, labyrinthoid, pale yellow to yellow (4A3–A6); pale brown (6D6) to brown in exsiccata (6E7); locules 0.7–1.8 × 0.15–0.3 mm (1–3 per mm), irregular, elongated, sinuous; fresh spore mass in locules pale yellow to yellow (4A3–A5); pale orange (5A4) to greyish orange (5B6) in exsiccata. **Stipe-columella** 1–2 × 0.2–0.4 cm, well developed, cylindrical, attenuate, central or eccentric, straight or curved, percurrent, not branched, pure white; context white, unchanging upon exposure to air. **Odour** fruity, taste mild.

**Spores** 9.5–12.5(–15) × (8–)8.5–10.5(–13) µm, Q = 1.1–1.2 (–1.3), subglobose to broadly ellipsoid, heterotropic, echinate, yellow; warts amyloid, 1.2–2 µm high, cylindrical and straight, with a truncate apex, in groups of 2–4, interconnected with low inamyloid ridges forming a basal reticulum; hilar appendix 1–1.8 × 1.2 µm, conical or cylindrical, straight; suprahilar plage amyloid; immature spores with a non-functional apical drop. **Basidia** (1–)2–4-spored, 37–55 × 14–18 µm, broadly clavate; sterigmata 4–5 µm long. **Basidioles** 30–35 × 10–15 µm, clavate. **Macrocytidia** scarce, 45–65(–100) × 6–14 µm, cylindrical, lanceolate or fusiform, rostrate. **Paraphysoid cells** 20–30 × 5–10 µm, 1-septate. **Subhymenium** composed of isodiametric cells 10–20 µm diam. **Hymenophoral trama** 30–50 µm wide, composed of hyaline hyphae 3–8 µm diam, with nests and columns of sphaerocytes 20–36 µm diam. **Pileipellis** and **context** 150–250 µm thick; pileipellis composed of: 1) a suprapellis soon collapsing into a yellow slimy mass, consisting of a cutis of upright to repent hyphae 2–4 µm diam, and some hyaline, cylindrical or clavate dermatocystidia 20–40 × 3–7 µm; and 2) a subpellis 100–130 µm thick, consisting of a subixocutis of densely intricate, hyaline hyphae 2–3 µm diam. **Pileal context** 100–110 µm thick, forming a cutis of intricate hyaline hyphae 2–4 µm diam, with some inflated hyphae and sphaerocytes up to 15 µm diam. **Stipitipellis** a turf of repent to erect hyphal tips mixed with dermatocystidia; context of stipe-columella heteromerous, composed of densely interwoven narrow hyphae mixed with nests of sphaerocytes. **Gloeoplera** 2–5 µm diam, present in trama and context.

**Habitat, Distribution & Season** — Solitary to gregarious, hypogeous to semi-hypogeous among plant debris, in broad-leaved montane woods of *Carpinus*, *Corylus*, *Fagus*, *Quercus*, accompanied by *Acer*, *Betula*, *Castanea*, *Fraxinus*, *Populus*, *Tilia*, but also under *Abies*, on calcareous soil, frequently found together with *Russula candida*. Summer and autumn. Widely distributed in temperate regions of Europe, from 100 m altitude in Northern countries to 1100 m in Southern countries.



**Fig. 17** *Russula candidissima*. a–e. JMV20110906-6a (BCN, holotype). a. Basidiomata; b. pileipellis; c. stipitipellis; d–e. SEM images of spores. — f. JMV800675. Basidiomata. — g–i. JMV20100724a. g–h. Basidia, basidioles and macrocystidium; i. spores in Melzer. — j. KRA F-2013-94. SEM image of spores. — k. KRA F-2009-57. SEM image of sporal ornamentation. — l. PRM 619108. Spores in Melzer. — Scale bars: a, f = 1 cm; b, i, l = 10 µm; c, g–h = 20 µm; d–e, j = 5 µm; k = 1 µm — Photos: a–c, g–i, l. J.M. Vidal; d–e. UdG; f. P.M. Pasabán; j–k. P. Mleczko, UJ.

*Additional material studied.* AUSTRIA, Lower Austria, Wien-Neustadt, Näu-dorf, 10 June 1937, M. Jacob, det. E. Soehner as '*Hydnangium cereum*' (M, herb. E. Soehner 1894). — CZECH REPUBLIC, Central Bohemia, Černošice, in a leafy forest, under *Carpinus*, *Crataegus* and *Cornus*, 4 July 1950, V. Va-cek, det. M. Svrček as '*Elasmomyces krjukowensis*' (PRM 619107); South Moravia, Súchov, under *Carpinus*, *Quercus* and *Fagus*, 8 July 1953, K. Kríž, det. M. Svrček as '*Hydnangium krjukowense* var. *moravicum*' (PRM 619108, herb. F. Šmarda). — GERMANY, Baden-Württemberg, Altbach/Neckar, in a *Carpinetum*, 23 Oct. 1961, H. Steinmann as '*Hydnangium krjukowense*' (PRM 616208); Saarland, Ballweiler, under *Fagus*, *Carpinus* and *Quercus*, on calcareous soil, 19 July 1968, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 161); Bebelsheim, under *Fagus*, *Carpinus* and *Quercus*, on calcar-eous soil, 27 Aug. 1967, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 82); ibid., 12 July 1968, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 157); ibid., 26 Aug. 1969, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 323); Eimersdorf, under *Carpinus*, *Fagus* and *Quercus*, on calcareous soil, 15 Aug 1968, Derbsch & G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 188); ibid., 6 Sept. 1969, J. Schmitt, det. G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 335); Eschingen, under *Fagus*, *Quercus* and *Fraxinus*, 3 Sept. 1967, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 85); Gerlfangen, under *Fagus*, on calcareous soil, 20 July 1968, J. Schmitt, det. G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 164); ibid., under *Fagus*, *Carpinus* and *Quercus*, 20 July 1968, G. Korn, det. G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 164a); ibid., under *Betula* and *Fagus*, 10 Aug. 1968, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 179); Sitterswald, under *Fagus sylva-tica*, on calcareous soil, 4 July 1967, G. Gross as '*E. mattiroloanus*' (M, herb.

G. Gross 79'); ibid., 31 Aug. 1967, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 84); ibid., 18 July 1968, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 159); ibid., 10 July 1969, G. Gross as '*E. mattiroloanus*' (M, herb. G. Gross 303). — ITALY, Tuscany, Lucca, Vagli di Sotto, Monte Roggio, 900 m, under *Corylus avellana*, gregarious, 4 July 1993, G. Bernardini & L. Gori as '*M. candidus*' (ELG930704-1, duplicate BCN JMV800185)\*. — POLAND, Lesser Poland, Krościenko nad Dunajcem, Pieniny Nat. Park, near Gródek, Western Carpathians, 480 m, under *Abies alba*, on calcareous soil, 19 Sept. 2009, P. Chachula (KRA F-2009-57, duplicate BCN JMV800664)\*; ibid., yellow route from Sromowce Niżne to Szopka pass, 640 m, under *Abies alba*, on calcareous soil, 13 July 2011, P. Chachula (KRA F-2013-94, duplicate BCN JMV800665); Silesia, Cieszyn, Pogórze Śląskie region, Western Carpathians, 296 m, under *Carpinus betulus* and *Quercus* sp., on calcareous soil, 22 Aug. 2017, P. Chachula (KRA F-2017-2, duplicate BCN JMV800669)\*. — SPAIN, Basque Country, Guipúzcoa, Villabona, under *Corylus avellana*, *Castanea sativa* and *Acer* sp., on calcareous soil, 19 Sept. 2004, P.M. Pasabán as '*M. candidus*' (BCN JMV800674); Catalonia, Girona, Molló, Coll de la Boixeda, El Querol, 1100 m, under *Corylus avellana*, *Quercus pubescens*, *Tilia platyphyllos* and *Populus tremula*, on calcareous soil, occurring in the same habitat as *Russula candida*, 17/24 July 2010, M.A. Pérez-De-Gregorio (BCN JMV20100724a); Navarre, Aizarotz, under *Corylus avellana*, on calcareous soil, 26 Sept. 2006, P.M. Pasabán as '*M. candidus*' (BCN JMV800675).

*Notes* — After carefully studying all basidiomata present in collection JMV20100724 (*Russula candida* from Catalonia, Spain), some of them were found to have spores with intercon-

nected warts forming low ridges, different from the spores with isolated warts found in the remaining specimens of *R. candida*. These basidiomata also had a well-developed stipe-columella. After revisiting all collections of *Macowanites candidus* studied by Vidal (2004b), similar spore ornamentation was found also in Czech, German and Italian collections. The same German collections had been previously described by Gross (1969), under the name of *Elasmomyces mattioloanus*. Similar spore ornamentation was also depicted in SEM images obtained from Swiss specimens by Pegler & Young (1979). More recently, Calonge & Pasabán (2005), and Gori (2005), described and illustrated new collections (identified by them as *M. candidus*) found in Spain and Italy, respectively, with the same spore morphology.

*Russula candidissima* can be discriminated macroscopically from *R. candida* because of its more persistently white external colour, its pileus that turns slightly yellowish with handling, and its mature basidiomata, which are opened in the base exposing the sublamellate hymenium, while in *R. candida* pileus becomes intensely yellow or yellowish orange with handling and its hymenium is rarely exposed. Microscopically, *R. candidissima* has a suprapellis composed of more or less parallel hyphae with scattered dermatocystidia and spore warts are interconnected by low ridges, while *R. candida* has a trichodermal suprapellis and spore warts are isolated. *Russula mattioloana* differs microscopically from *R. candida* and *R. candidissima* by its bigger macrocystidia up to 130 µm long, and by its subglobose to globose spores up to 15.5(–18) × 15(–17) µm, ornamented with isolated warts up to 2.5–3 µm long.

Genetically, *R. candidissima* is significantly related to several sequestrate species from North America such as *R. xerophila*, *R. ellipsospora*, *R. matthsmithii* (a replacement name for *Gymnomyces compactus*), *R. stewartii* (a replacement name for *G. monosporus*), and one from New Zealand, *R. ospranticarpa* (a replacement name for *Gymnomyces redolens*). All these sequestrate species constitute a significantly supported clade related to subsect. *Firmiores*, in concordance with previous results (Smith et al. 2006). Several species of this subsection are characterized by the yellow or yellowish orange colour of their lamellae and spores, a cuticular structure with short emergent hyphal tips, and spores ornamented with a spiny reticulum (Lebel 2002, 2003b, Smith et al. 2006).

### *Russula cerea* (Soehner) J.M. Vidal, comb. nov. — MycoBank MB828503; Fig. 18

Basionym. *Hydnangium cereum* Soehner, Kryptog. Forsch. 1, 6: 394. 1924.

Synonyms. *Octaviania cerea* (Soehner) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 197. 1958.

*Hydnangium carneum* var. *xanthosporum* Hawker, Trans. Brit. Mycol. Soc. 35, 4: 281. 1952. (syn. nov.)

*Gymnomyces xanthosporus* (Hawker) A.H. Sm., Mycologia 54: 635. 1962.

*Russula xanthospora* (Hawker) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 240. 2018.

Lectotype of *Hydnangium cereum* (here designated MBT384232): GERMANY, Bavaria, 'Erharting bei Mühldorf', under oaks, 16 Aug. 1921, E. Soehner (M, herb. E. Soehner 527); isolectotype in FH (herb. C.W. Dodge).

**Basidiomata** 1–3 cm wide, angiocarpic, subglobose, slightly lobate, sessile, sometimes presenting an inconspicuous sterile base and attached to the substrate by a mycelial strand. **Pileus** dry, smooth, membranous, translucent, displaying hymenophoral locules, basally open and alveolate in old specimens; initially white, then greyish orange (5B6) to pale brown (6D6) when handled or upon exposure to the air, finally reddish brown (8E8). **Hymenophore** loculate, labyrinthoid, brown (7D7) to reddish brown (8D8); locules minute, 0.4–1.4 × 0.1–0.3 mm (2–3 per mm),

elongated, sinuous; external locules always fertile; fresh spore mass in locules reddish brown (9D6); brown (6E7) in exsiccata. **Columella** absent or rudimentary, but thin sterile veins can be observed in some specimens. **Odour** fruity, taste mild.

**Spores** (8–)9.5–12.5(–14) µm, Q = 1, variable in size, globose, orthotropic, echinate, orange to reddish in KOH; warts of irregular length, 1–3 µm high, intense yellow, deeply amyloid, dense, robust, conical or cylindrical, obtuse, longitudinally striate, isolated or fused in groups, often interconnected by low ridges; hilar appendix short, inconspicuous. **Basidia** typically 2-spored, 40–50 × 18–24 µm, pyriform to clavate, filled with minute, hyaline to dark reddish droplets, soon collapsing; sterigmata 8–10(–12) µm long, cylindrical to conical. **Basidioles** 22–36 × 10–13 µm, clavate. **Macrocytidia** abundant, (25–)30–50 × (5–)7–12(–16) µm, cylindrical to cylindro-clavate, wall up to 1 µm, with a dark yellow refractive content, rarely exceeding the basidia in length, resembling pseudocystidia because many of them originate at the deep subhymenium or hymenophoral trama. **Paraphysoid cells** abundant, 20–42 × 10–15 µm, vesiculose, 1-septate. **Subhymenium** very thick, pseudoparenchymatous, formed by 2–3 layers of prismatic to globose cells 8–20(–25) µm diam. **Hymenophoral trama** reduced, 10 µm thick, formed by hyphae 3–6 µm diam, and some sphaerocytes 10–20 µm diam, only present in tramal anastomoses. **Pileipellis** and **context** 70–125 µm thick; pileipellis thin, 50–100 µm, consisting of: 1) a trichodermal suprapellis of subulate to lageniform, capitulate dermatocystidia, measuring 15–30 × 3–5 µm, that soon collapses into a yellowish mass; and 2) a prosenchymatous subpellis 20–30 µm thick, formed by a dense mesh of interwoven, subgelatinized yellowish hyphae, measuring only 2–2.5 µm diam. **Peridial context** formed of interwoven hyaline hyphae 2–5 µm diam, lacking sphaerocytes. **Gloeoplera** and **thromboplera** present in the trama and context.

**Habitat, Distribution & Season** — Solitary to gregarious, hypogeous, in montane conifer (*Abies*, *Picea*, *Pinus*) or broad-leaved (*Carpinus*, *Castanea*, *Corylus*, *Fagus*, *Quercus*) forests, commonly on siliceous soil. Summer and autumn. Widely distributed in temperate and submediterranean regions of Europe, from almost sea level in Northern countries to 1100 m altitude in Southern countries.

**Additional material studied.** GERMANY, Saxony, Chemnitz, Park der OdF, 300 m, under *Castanea sativa*, 20 June 2009, B. Mühlner & G. Hensel (GH20090620, duplicate BCN JMV800660)\*. — HUNGARY, Veszprém, Zirc, under *Fagus* and *Carpinus*, 15 Sept. 2000, Z. Lukács (BCN JMV800164). — ITALY, Emilia-Romagna, Forlì-Cesena, Santa Sofia, Foresta di Campigna, under *Abies alba*, 11 Oct. 1991, M. Sarasini as *G. xanthosporus* (MS419, duplicate BCN JMV800006); Tuscany, Lucca, Vagli di Sotto, Monte Roggio, 800 m, under *Corylus avellana*, 4 Dec. 1994, G. Bernardini & L. Gori as *G. xanthosporus* (ELG941204-3, duplicate BCN JMV800172). — POLAND, Lower Silesia, Góry Złote region, Sudety mountains, under *Quercus* sp., *Fagus sylvatica*, *Picea abies*, *Abies alba* and *Acersp.*, 9 Aug. 2012, M. Kozak & P. Mleczko (KRA F-2012-28, duplicate BCN JMV800667)\*. — PORTUGAL, Aveiro, Luso, bosque de Buçaco, 547 m, under *Castanea sativa*, on siliceous soil, 11 Nov. 2015, A. Paz (BCN IC11111504). — SPAIN, Asturias, Perán, under *Castanea sativa*, 17 Aug. 2000, F. García (BCN JMV20000817-1)\*; Catalonia, Girona, Campelles, Torrent de Prat de Jou, 1100 m, under *Corylus avellana*, *Fraxinus*, *Acer* and *Buxus*, on siliceous soil, 3 Oct. 1996, J.M. Vidal as *G. xanthosporus* (BCN JMV961003-8)\*; Girona, Espinelves, crossroad to Viladrau, 740 m, in a *Picea abies* plantation, on siliceous soil, 5 July 2016, F. Rodríguez (BCN JMV20160705)\*; ibid., 4 Sept. 2018, F. Rodríguez & J.M. Vidal (BCN JMV20180904-3); Navarre, Eratsun, under *Pinus pinaster*, 10 June 2015, P.M. Pasabán & F. Sáinz (BCN JMV800656)\*. — UNITED KINGDOM, Wales, Caernarvon, 'Bettws-y-Coed, North Wales, inter acibus emortius Piceae', 13 Sept. 1951, L.E. Hawker (K(M)69329, herb. L. Hawker H251, holotype of *H. carneum* var. *xanthosporum*).

**Notes** — Soehner (1924) described an angiocarpic fungus found under oaks in the German region of Bavaria, which he called *Hydnangium cereum*, but did not select a type collection. In a later paper, Soehner (1941) provided new data and reference




**Fig. 18** *Russula cerea*. — a–b. GH20090620. a. Basidiomata; b. basidia and macrocystidia. — c. MS419. Pileipellis. — d. JMV20180904-3. Basidiomata. — e. JMV20160705. Basidia and macrocystidia. — f. ELG941204-3. Spores in Melzer. — g–h. M, herb. E. Soehner 527 (original material of *Hydnangium cereum*). g. Basidium and macrocystidia; h. spores in KOH 5 %. — i. JMV20160705. SEM image of spores. — j–k. K(M)69329 (holotype of *Hydnangium carneum* var. *xanthosporum*). j. Basidia and macrocystidia; k. spores in KOH 5 %. — l. KRAF-2012-28. SEM image of spores. — Scale bars: a, d = 1 cm; b–c, e, g, j = 20 µm; f, h, k = 10 µm; i, l = 5 µm. — Photos: a. G. Hensel; b–h, j–k. J.M. Vidal; i. UdG; l. P. Mleczko, UJ.

codes for several collections of this species, indicating that duplicates of a collection labelled 'Summer 1921' were sent to several mycologists such as E. Fischer, O. Mattiolo and C.W Dodge. Dodge & Zeller (1937) considered this collection, which they labelled 'E. Soehner 527', as the type of *H. cereum*. After studying several original collections of *H. cereum* from E. Soehner's herbarium at München, we found that 'Soehner no. 527' is the only collection consisting of mature basidiomata, so it is here designated as lectotype.

Later on, Hawker (1952) described from Wales, UK, an angiocarpic fungus of similar characteristics that she called *Hydnangium carneum* var. *xanthosporum*. Type collections of *H. cereum* and *H. carneum* var. *xanthosporum* were compared in the present work, and found to be probably conspecific because of their bisporic basidia, cylindrical thick-walled macrocystidia, and conspicuously ornamented globose spores, with *H. cereum* being the priority name. Interestingly, the presence of hymenial macrocystidia was not observed by Soehner (1924, 1941) in *H. cereum*, neither by Hawker (1952, 1954), Smith (1962), nor by Pegler et al. (1993) in *H. carneum* var. *xanthosporum*,

despite them being very conspicuous. Morphological traits as well as genetic data from specimens originating from Central and Southern Europe suggest that this taxon is another sequestrate species of *Russula*, and therefore the combination *R. cerea* is here proposed.

Phylogenetic inferences suggested that *R. cerea* belongs to sect. *Ingratae*, and probably subsect. *Foetentinae*. *Russula cerea* is significantly related to other European sequestrate taxa such as *R. pila* and *R. mistiformis*, being macroscopically very similar to both species, but differing because of its thinner membranous pileus, bisporic basidia and presence of thick walled, clavate macrocystidia. All samples of *R. cerea* seemed genetically similar to each-other, except for JMV20000817-1 from Asturias (Spain), which showed significant differences in ITS (15/564 bp = 2.65 %, including a 7 bp insertion) and 28S rDNA (4/851 bp = 0.5 %). After microscopically studying this specimen, we have been unable to find any distinct features, so it could represent a cryptic species or maybe a hybrid specimen with either *R. pila* or *R. mistiformis*.



**Fig. 19** *Russula galileensis*. a. IB M72.340 (isotype of *Macowanites galileensis*). Spores in Melzer. — b. HAI-G-83. Suprapellis. — c–d. HAI-G-201. c. Hymenium and subhymenium; d. spores in Melzer. — e–g. HAI-G-83. SEM images of spores. — Scale bars: a, d–e = 10 µm; b–c = 20 µm; f–g = 1 µm. — Photos: a–d. J.M. Vidal; e–g. UdG.

***Russula galileensis* (M.M. Moser et al.) Trappe & T.F. Elliott  
Fungal Systematics and Evolution 1: 234. 2018 — Fig. 19**

*Basionym.* *Macowanites galileensis* M.M. Moser et al., Trans. Brit. Mycol. Soc. 68, 3: 371. 1977.

*Basidiomata* russuloid, angiocarpic, stipitate. *Pileus* 2–6 cm wide, irregular and tuberous, sometimes depressed at the apex, white with cream to pale umber maculae; margin radially alveolate when mature, but not open. *Hymenophore* loculate, cream-ochre. *Stipe-columella* 0.5–3.5 × 0.5–1.7(–3) cm, white. *Odour* and taste mild.

Spores (9)–10–11.5(–15) × (7.5)–9.5–10.5(–14) µm, Q = 1.05–1.15, subglobose, heterotropic; reticulum 0.5 µm high, amyloid, composed of crests and warts. *Basidia* 4-spored, 26–37 × 11–13 µm, broadly clavate. *Macrocytidia* 40–60 × 10–15 µm, sometimes 1–2-septate. *Hymenophoral trama* heteromerous. *Pileipellis* and *context* 200–250 µm thick; suprapellis an intricate trichoderm of repent, to semi-erect hyphae and dermatocystidia 2–7 µm diam.

*Habitat, Distribution & Season* — Subepigaeous under fallen leaves, in sclerophyllous woods of *Quercus*, on calcareous soil. From late autumn to spring. So far located only in the eastern Mediterranean region, in Israel, found at low altitudes.

*Material studied.* ISRAEL, Carmel Mount, Mt Carmel National Park, near Haifa, Shajarat Al Arba'in (Horshat Ha'arba'im), under *Quercus calliprinos*, 12 Dec. 1972, M. Moser, N. Binyamini & Z. Avizohar-Hershzon 72/401 (IB M72340, isotype of *M. galileensis*); ibid., near *Pinus*, 13 Jan. 2012, M. Krakhmalnyi as *M. galileensis* (HAI-G-91); Haifa, Technion, Dec. 2011, M. Krakhmalnyi as *M. galileensis* (HAI-G-59); Upper Galilee, Goren Park, Dec. 2011, Z. Shafranov, det. M. Krakhmalnyi as *M. galileensis* (HAI-G-83; duplicate BCN JMV800634)\*; ibid., Hanita Forest, 22 Dec. 2012, Z. Shafranov, det. M. Krakhmalnyi as *M. galileensis* (HAI-G-166)\*; Mount Meron National Park, 7 Jan. 2006, I. Shams, det. S.P. Wasser as *M. galileensis* (HAI-G-202)\*; Nahal Kziv Nature Reserve, 'Evolution Canyon-II', under *Quercus*, 3 Jan. 2001, S. Reshetnikov, det. S.P. Wasser as *M. galileensis* (HAI-G-45); ibid., under *Quercus*, 3 Jan. 2001, T. Pavlichek & I. Shams, det. S.P. Wasser as *M. galileensis* (HAI-G-201; duplicate BCN JMV800635)\*; ibid., under *Laurus*, 7 Jan. 2001, I. Duckman, det. S.P. Wasser as *M. galileensis* (HAI-G-46); Nahal Zalmon National Park, under *Quercus*, 1 Febr. 2003, Y. Ur, det. S.P. Wasser as *M. galileensis* (HAI-G-209)\*.

*Notes* — *Macowanites galileensis* was originally described and illustrated from the specimens collected on Mt Carmel in 1972 (Moser et al. 1977, 1994). Interestingly, the authors initially regarded their collection to be *M. krjukowensis*, due to white

colour of basidiomata. Following more careful examination, however, it became evident that new collections possessed a number of distinctive macro- and micromorphological characters, such as a very robust pileus and stipe-columella, as well as warts of spores connected by ridges and crests, features which separated them from the latter. We could not obtain images to illustrate the studied collections. Recently, this species was re-combined into genus *Russula* by Elliott & Trappe (2018). Current molecular results indicate that *R. galileensis* belongs to *Russula* subsect. *Laricinae*, being significantly related to the *R. laricinoaffinis* clade. Subsection *Laricinae* is characterized by ochre to yellowish spore deposits, short basidia, rare pleurocystidia and multisepitate dermatocystidia (Romagnesi 1985, Li et al. 2013). Other European sequestrate taxa also related to *Laricinae* include *R. vinaceodora* and *R. vidalii* (a replacement name for *Gymnomyces ilicis*), both species apparently endemic to the Mediterranean region (Moser et al. 1977, Llistosella & Vidal 1995, Calonge & Vidal 2001, Cabero 2011). *Russula galileensis* and *R. vinaceodora* produce large basidiomata (up to 5–6 cm across), with a conspicuous stipe-columella that is sublamellate in the lower part, and spores with warts forming small interconnecting ridges, while *R. vidalii* is a smaller fungus (1–2.5 cm diam), with a poorly developed columella, a loculate hymenophore, and spores which are ornamented with more or less isolated warts.

***Russula hobartiae* Loizides & J.M. Vidal, sp. nov. — MycoBank MB828504; Fig. 20**

*Basidiomata* 2–7 cm wide, angiocarpic, turbinate to pyriform, weakly lobate, tomentose, whitish to ochraceous brown, firmly rooted into the substrate. *Hymenophore* loculate, whitish to ochraceous orange, becoming vinaceous in FeSO<sub>4</sub>; locules minute, rounded to elongate. *Columella* absent or rudimentary, sometimes forming short white veins. *Odour* slightly fruity, with a vague hint of chocolate. Spores 7–9.5(–10.5) × 7–9(–10) µm, globose; reticulum complete to incomplete, up to 0.5 µm high. *Basidia* 2–4-spored, 20–35 × 8–12 µm, clavate. *Macrocytidia* 25–40 × 8–10 µm, clavate, scarce. *Hymenophoral trama* prosenchymatosus, with scarce sphaerocytes. *Suprapellis* a palisado-trichoderm of clavate dermatocystidia 25–60 × 5–10 µm; pileal context a cutis of hyaline hyphae 1.5–3 µm diam, lacking sphaerocytes. Hypogeous or semi-hypogeous in montane woods of *Pinus nigra* subsp. *pallasiana* on the island of Cyprus.

*Etymology.* *hobartiae* = in honor of mycologist Caroline Hobart, for her contribution to the study of hypogeous and sequestrate fungi in Cyprus.

*Holotype.* CYPRUS, Nicosia District, Prodromos, 1450 m, under *Pinus nigra* subsp. *pallasiana*, 13 Oct. 2011, M. Loizides (BCN JMV800628)\*; isotype in herb. pers. M. Loizides (ML110131GY).

*Basidiomata* 2–7 cm wide, angiocarpic, turbinate, oblate-globose to pyriform and somewhat applanate at the apex, weakly lobate, distinctly tapering at base, longitudinally furrowed and firmly rooted into the substrate. *Pileus* dry, finely tomentose under the lens, often cracked, dirty white, cream-white (5A2–A4) or ochraceous cream (6A2), usually with darker ochraceous or brownish stains (6C5). *Hymenophore* loculate, labyrinthoid, cream at first, at maturity becoming deep ochraceous yellow to ochraceous orange (5A6); instantly turning vinaceous in FeSO<sub>4</sub>; locules minute, 0.2–1 × 0.1–0.3 mm (3–5 per mm), irregularly rounded to elongated, empty or filled; fresh spore mass in locules pale yellow to yellow (4A3–A4); pale orange (5A4–A5) in exsiccata. *Columella* absent or rudimentary, sometimes forming a sterile base or short white veins, but small sterile patches often scattered throughout the hymenophore. *Odour* slightly fruity, sometimes with a vague hint of chocolate when cut; taste astringent.


*Spores* 7–9.5(–10.5) × 7–9(–10) µm, Q = 1.0–1.1, globose, orthotropic, reticulated; reticulum complete to incomplete, up to 0.5 µm high, formed by strongly amyloid crests and warts; hilar appendix usually rudimentary, exceptionally up to 2–3 µm long. *Basidia* 2–4-spored, 20–35 × 8–12 µm, clavate, thick-walled; sterigmata 4–6 µm long. *Macrocytidia* 25–40 × 8–10 µm, clavate, scarce, very scattered and immersed in the hymenium. *Paraphysoid cells* 11–34 × 7–12 µm, cylindrical to slightly clavate, 1–2-septate. *Subhymenium* rameose, composed of chains of 1–2 globose to irregularly polygonal or elongated cells 5–15 µm diam. *Hymenophoral trama* 40–70 µm wide, prosenchymatous, formed by septate, tortuous, interwoven hyaline hyphae 3–6(–8) µm diam, with some inflated elements up to 12 µm

diam and scattered sphaerocytes or nests of sphaerocytes 6–20 µm diam, especially in tramal anastomoses. *Pileipellis* and *context* 120–250(–400) µm thick; pileipellis consisting of: 1) a palisadotrichodermal suprapellis of yellow, clavate dermatocystidia 25–60 × 5–10 µm; and 2) a prosenchymatous subpellis 30–40 µm thick of branched, densely interwoven, septate hyphae 3–6 µm diam. *Pileal context* 80–200 µm thick, arranged in a cutis of septate, hyaline hyphae 1.5–3 µm diam; sphaerocytes absent. Gloeoplera abundant, up to 8 µm diam.

**Habitat, Distribution & Season** — Solitary to gregarious, hypogeous to semi-hypogeous, in montane woods of *Pinus nigra* subsp. *pallasiana*, rarely also with *Pinus brutia*, on siliceous, well-drained serpentine soils. Summer and autumn. Xerophytic. Known from the eastern Mediterranean region, where it is probably endemic to the island of Cyprus, between 1100–1800 m altitude.

**Additional material studied.** CYPRUS, Limassol District, Troodos, 1800 m, under *Pinus nigra* subsp. *pallasiana*, 7 Oct. 2008, M. Loizides (ML8017GY); ibid., 1700 m, 28 Sept. 2009, M. Loizides (ML90982GY); ibid., 1600 m, 4 Nov. 2012, M. Loizides (ML21114GY); ibid., 26 Aug. 2014, M. Loizides (ML41862GY); ibid., 1650 m, 3 Sept. 2014, M. Loizides (ML4193GY)\*; ibid., 1700 m, 20 Sept. 2016, M. Loizides (ML61902GY); Nicosia District, Prodromos, 1450 m, under *Pinus nigra* subsp. *pallasiana*, 7 Oct. 2011, M. Loizides (ML11701GY); Platania, 1100 m, under *Pinus brutia*, 16 Nov. 2011, M. Tordelli (ML111161GY); ibid., 1120 m, 20 Nov. 2011, G. Konstantinidis & D. Klisiari (GK5889, duplicate BCN JMV800647)\*; ibid., 16 Nov. 2014, M. Loizides (ML411161GY)\*.

**Notes** — *Russula hobartiae* is characterized by the typically turbinete shape of its basidiomata, with a flattened upper side and a conical base that is deeply rooted into the substrate and breaks easily. Hymenial chambers are very small, only visible under a lens, cream at first but turning into an intensely orange colour when mature. Microscopically, it is characterised by its



**Fig. 20** *Russula hobartiae*. a. ML110131GY (BCN JMV800628, holotype). Basidiomata. — b. ML411161GY. Basidiomata. — c. ML4193GY. Basidiomata. — d–i. ML110131GY (BCN JMV800628, holotype). d. Pileipellis; e. hymenium, subhymenium and hymenophoral trama; f. hymenium and subhymenium; g. spores in Melzer; h–i. SEM images of spores. — Scale bars: a–c = 1 cm; d–f = 20 µm; g = 10 µm; h–i = 5 µm. — Photos: a–c. M. Loizides; d–g. J.M. Vidal; h–i. UdG.

globose or subglobose spores ornamented with a reticulum, its infrequent and very scattered clavate macrocystidia, not exceeding the hymenial layer, and the pileipellis formed by abundant clavate dermatocystidia forming a palisade.

Genetically, *R. hobartiae* is significantly related to a North American specimen identified as *Russula* cf. *ochrophylla* (BPL231, Looney et al. 2016). Currently, there are no other records of *R. ochrophylla* in public databases, so the identity of this sample cannot be confirmed until additional collections are studied. *Russula ochrophylla* is characterized by a purple pileus, ochraceous yellow lamellae, mild taste, echinulate spores, and preference for broadleaved trees (Peck 1897, Adamčík et al. 2017), features which are remarkably diverse from those of *R. hobartiae*. This morphological disparity could perhaps be explained if: 1) specimen BPL231 does not actually represent *R. ochrophylla*; 2) the features of one of both concerned species are not typical of their lineage; or 3) *R. ochrophylla* and *R. hobartiae* actually belong to different subsections.


***Russula mattiroloana* (Cavara) T. Lebel, Muelleria 36: 11. 2017**

— Fig. 21

*Basionym. Elasmomyces mattiroloanus* Cavara, Malpighia 11, 1–3: 426. 1897 ('mattirolianus').

*Synonym. Macowanites mattiroloanus* (Cavara) T. Lebel & Trappe, Mycologia 92, 6: 1194. 2000.

**Basidiomata** russuloid, pseudoangiocarpic, stipitate. **Pileus** 1.2–3.8 cm wide, subglobose to hemispherical, open in the base and exposing a marginally sublamellate hymenophore; initially finely tomentose, then smooth, whitish, pale yellow (4A4) to pale orange (5A4), maculated with brownish orange (6C8), finally brownish red (8D7) maculated with dark reddish brown (8E7–F6); greyish orange (5B5) to brown in exsiccatum (7E8). **Hymenophore** loculate, labyrinthoid-daedaleoid, initially with an enclosed stipe-columella, soon expanding outwards and exposing the sublamellate hymenium underneath; initially whitish, yellowish white to pale yellow (4A2–A3), then pale yellow (4A5) to pale orange (5A4), finally titian red (7D6); pale orange (5A4) to dark brown in exsiccatum (6F7); locules large, 1.5–3 × 0.2–0.8 mm (1–2 per mm), radially arranged,



**Fig. 21** *Russula mattiroloana*. a. Reproduction of Cavara's plate of *Elasmomyces mattiroloanus* (Cavara 1897, tav. VII). — b. GK8136. Basidiomata. — c. KRA F-2018-1. Basidiomata. — d–l. KRA F-2012-153. d. Pileipellis; e. stipitipellis; f. hymenium and subhymenium; g–h. spores in Melzer; i–l. SEM images of spores. i. two ellipsoid young spores; j. globose mature spore; k. detached apical drop over the ornamental warts; l. non-functional apical drop united to the base of hilar appendix. — Scale bars: b–c = 1 cm; d–f = 20 µm; g–h = 10 µm; i–k = 5 µm; l = 1 µm. — Photos: b. G. Konstantinidis; c. P. Chachula; d–h. J.M. Vidal; i–l. P. Mleczko, UJ.

elongated, irregular, sinuous; fresh spore mass in locules pale yellow (4A4); pale orange (5A3) to brown (6E6) in exsiccata. *Stipe-columella* 0.8–2.5 × 0.3–1 cm, well developed, percurrent, stipe 2–3 times longer than the columella, cylindrical, generally central, often curved, pruinose, pure white; context white, not changing upon exposure to air, but yellow in the cortical zone. Odour and taste mild.

Spores (9–)10.5–15.5(–18) × (8–)10–15(–17) µm, Q = 1.0–1.15, very variable in size, globose to subglobose, some ellipsoid when immature (Q = 1.1–1.2), heterotrophic to subheterotrophic, echinate; warts (0.7–)1.5–2.5(–3) µm high, isolated, amyloid, conical and straight, with an acute apex; hilar appendix 1.5–2 × 1–1.5 µm, cylindrical, straight, with a minute, inconspicuous, amyloid suprahilar plage in its base and conserving a non-functional apical drop. *Basidia* 2–4-spored, (25–)45–65 × 17–23 µm, ventricose to broadly clavate; sterigmata 3–5 µm long. *Macrocytidia* abundant, 70–125(–180) × 15–18(–25) µm, cylindrical, lanceolate or fusiform, obtuse or rostrate, with a wall 1 µm thick. *Paraphysoid cells* scarce, 18–28 × 6–8 µm, cylindrical or clavate, aseptate or 1-septate. *Subhymenium* of pseudoparenchymatous aspect, consisting of 3–4 strata of globose to prismatic cells 14–30 µm diam. *Hymenophoral trama* 20–30 µm wide, with scarce sphaerocytes 15–35 µm diam, more abundant in tramal anastomoses. *Pileipellis* and *context* 100–300 µm thick; pileipellis composed of: 1) a trichodermal suprapellis consisting of upright to repent hyphae 2–6.5 µm diam, and abundant, long, cylindrical or clavate, usually 1-septate, yellow dermatocystidia, 60–140 × 3–8 µm; and 2) a subpellis 100–180 µm thick, comprised of a subixocutis of intricate, hyaline hyphae 1.5–5 µm diam, with frequent enlargements 10–20 µm at hyphal ramifications. *Pileal context* heteromerous, 200–1000 µm thick, formed by hyaline hyphae 3–6 µm diam, and abundant nests of sphaerocytes up to 40 µm diam. *Stipitipellis* a turf of repent to erect, septate hyphae, 3–5 µm diam, and clavate dermatocystidia, 50–100 × 5–12 µm, similar in shape and content to those of pileipellis; context of stipe-columella heteromerous, composed of densely interwoven narrow hyphae mixed with nests of sphaerocytes. *Gloeoplera* 2–5 µm diam, present in trama and context where they terminate as dermatocystidia.

**Habitat, Distribution & Season** — Solitary to gregarious, subepigaeous among needles, in forests of *Abies* and *Picea*, on siliceous or calcareous soils. Summer and autumn. Distributed in temperate and submediterranean regions, from Central to Southern Europe (Carpathian, Pindus and Apennine mountains), between 500–1600 m altitude.

**Material studied.** GREECE, Thessaly, Trikala, Pertouli, Koziakas mountain, Pindus mountains, 1180 m, under *Abies borisi-regis*, 5 July 2009, G. Konstantinidis as *M. mattiroloanus* (GK3901, duplicate BCN JMV800638)\*; ibid., 1200 m, under *Abies borisi-regis*, 11 July 2015, G. Konstantinidis as *M. mattiroloanus* (GK8136, duplicate BCN JMV800644)\*. – ITALY, Emilia-Romagna, Reggio Emilia, Civago, Abetina Reale Forest, Tuscan-Emilian Apennines, 1500 m, under *Abies alba* and *Fagus sylvatica*, on siliceous soil, 5 Oct. 1983, A. Montecchi as '*M. ellipsospora*' (AM1616, duplicate BCN JMV800245); ibid., 1400 m, 10 Aug. 1985, A. Montecchi as *E. mattiroloanus* (AM65, duplicate BCN JMV800241); ibid., 16 Aug. 1985, A. Montecchi as *E. mattiroloanus* (AM64, duplicate BCN JMV800240); ibid., 1600 m, under *Abies alba* and *Fagus sylvatica*, on siliceous soil, 5 Aug. 1988, A. Montecchi as *E. mattiroloanus* (AM825, duplicate BCN JMV800038); Tuscany, Firenze, Reggello, Vallombrosa Forest, Tuscan-Emilian Apennines, 'ad terram sub acibus *Abietis pectinatae* prope S. Giovanni Gualberto Vallumbrosae (Florentia), F. Cavara' (FH, herb. C.W. Dodge 2087; NY, herb. S.M. Zeller 1671; all labelled '*Arcangeliella borziana*', Italy, Etruria, Vallombrosa, forest of firs, autumn 1898, Instituto Botanico di Napoli, coll. F. Cavara, type'; isotypes of *E. mattiroloanus*); Tuscany, 24 Nov. 1899, O. Mattirolo as *E. mattiroloanus*, conf. F. Cavara (FH, herb. F. Bucholtz 275). – POLAND, Lesser Poland, Pieniny Nat. Park, Western Carpathians, Sokolica massif, close to Hukowa Skała rock, 490 m, under *Abies alba*, on calcareous soil, 13 July 2012, P. Chachula (KRA F-2012-153, duplicate BCN JMV800666)\*; ibid., Bajków Groń, 710 m, under *Abies alba*, on calcareous soil, 3 July 2018, P. Chachula (KRA F-2018-1,

duplicate BCN JMV800713)\*; ibid., Lasek, 660 m, under *Abies alba*, on calcareous soil, 21 July 2018, P. Chachula (KRA F-2018-3, duplicate BCN JMV800715); ibid., Ula mountain, 620 m, under *Abies alba*, on calcareous soil, 8 Nov. 2017, P. Chachula (KRA F-2017-1, duplicate BCN JMV800670)\*; ibid., 8 July 2018, P. Chachula (KRA F-2018-2, duplicate BCN-JMV800714)\*.

**Notes** — *Elasmomyces mattiroloanus* was collected under *Abies alba* at the beginning of September 1896 in Tuscany (Italy) by Cavara (1897), who reported it as a russuloid fungus with a pseudoangiocarpic habit, characterized by its globose spores, 14–15 µm diam, and abundant macrocystidia, 70–72 µm long. Singer & Smith (1960) observed microscopical differences between the type material kept at New York Botanical Garden (NY) and the descriptions provided by Cavara: spores were subglobose or ellipsoid, measuring 10–15 × 9.5–13.5 µm, lacked macrocystidia and showed cystidiole 'sterile bodies', measuring 22–27 × 11–12 µm. Lebel & Trappe (2000) studied the same material at NY and confirmed the subglobose to broadly ellipsoid spore shape (although they report different spore measures, 10–12 × 9–11.5 µm), and further discriminated between two different kinds of cystidia: ventricose to clavate, 28–37 × 7–11 µm, and clavate to cylindrical, 8–14 × 3–5 µm. Vidal (2004b) also studied the same type material kept at S.M. Zeller's herbarium in NY (without voucher number) and concluded that microscopical features of these specimens do not match those reported in the protologue of *E. mattiroloanus*, but are compatible with those of *Arcangeliella borziana*, concluding that both species were probably erroneously labelled. The original concept of *E. mattiroloanus* has spores of unequal size and shape, 10.5–15.5(–18) × 10–15(–17) µm, typically globose or subglobose, sometimes ellipsoid when immature, resembling those of *Martellia ellipsospora* sensu Montecchi & Sarasini (2000), and numerous macrocystidia measuring 70–125(–180) × 15–18 µm. In mature spores, it is interesting to observe the intensely amyloid, non-functional apical drop, which remains located in a basal position in the hilar appendix, immediately above a minute and residual suprahilar plage, which is not always visible, and finally becomes detached on top of ornamental warts, or fused with the suprahilar plage, also observed in spores of *R. mediterraneensis*. Recently, *E. mattiroloanus* was re-combined into genus *Russula* by Lebel (2017).

Genetically, *R. mattiroloana* belongs to the *R. globispora* complex of subsect. *Maculatinae* (Adamčík et al. 2016). Despite ITS rDNA sequences of *R. mattiroloana* sharing a 20 bp deletion not present in any other species of *Maculatinae*, its clade collapsed or received non-significant support (Fig. 5), evidencing the need of additional samples and genetic markers. Subsection *Maculatinae* was originally characterized by species with reddish pilei, acrid taste and a yellow spore print (Romagnesi 1985), but these features are also shared by species not directly related (Adamčík et al. 2016). Interestingly, Sarnari (1998) subsumed *Maculatinae* within subsect. *Urentes*, and highlighted the presence in this group of an amyloid suprahilar spot in spores. The clade formed by *R. globispora* and *R. dryadicola* was reported by Adamčík et al. (2016) to be characterized by spores ornamented with large isolated spines, a feature also observed in *R. mattiroloana*, although also present in species not directly related with this clade, such as *R. candida*. *Russula mattiroloana* seems to be present in *Abies* sp. and *Picea* sp. forests of the alpine Mediterranean regions (Cavara 1897, Montecchi & Lazzari 1986, Montecchi & Sarasini 2000, Konstantinidis 2009) and Central Europe. *Russula globispora* has large, globose spores (Bon 1986, Llistosella 1998, Lejeune 2005) similar to those of *R. mattiroloana*, but spores of *R. dryadicola* and the new species *R. heilongjiangensis* (Li et al. 2018) do not share these features, suggesting that they are not representative of the entire genetic lineage.

**Russula mediterraneensis** Konstantin., J.M. Vidal, Gelardi, Papadimitriou, Tulli, Angeli & Vizzini, sp. nov. — MycoBank MB828505; Fig. 22

*Basidiomata* russuloid, pseudoangiocarpic, stipitate. *Pileus* 2.3–4.3(–5.2) cm wide, hemispherical to spherical, sometimes lobate, smooth, whitish to yellowish with brown maculae. *Hymenophore* internally daedaleoid and externally lamellate, cream to ochraceous orange. *Stipe-columella* 1.6–4.2 × 0.7–1.8 cm, well developed, percurrent, white. *Odour* of pears or yoghurt and fruit. *Spores* (8–)10–15(–17) × (7–)9–14(–15) µm, subglobose to broadly ellipsoid, heterotropic; warts 0.6–1(–1.5) µm high, isolated or in few groups. *Basidia* 2–3-spored, 30–55 × 11–18 µm, clavate. *Macrocystidia* 70–130 × 12–20 µm. *Hymenophoral trama* heteromerous. *Suprapellis* a trichoderm of erected hyphae 2–6 µm diam, and clavate, septate dermatocystidia 30–160 × 5–10 µm; pileal context heteromerous with nests and columns of sphaerocytes 21–47 µm diam. Found in Greece and Italy, subepigaeous under *Quercus* and *Castanea*.

**Etymology.** From Latin, *Mediterraneum (mare)* = Mediterranean Sea, and *-ensis* = from, found in, for its occurrence in Mediterranean localities.

**Holotype.** GREECE, West Macedonia, Kozani, Voucherina, 700 m, under *Quercus robur*, 23 July 2013, G. Konstantinidis (BCN JMV800641)\*; isotype in herb. pers. G. Konstantinidis (GK6710).

*Basidiomata* russuloid, small to medium sized, pseudoangiocarpic, stipitate. *Pileus* 2.3–4.3(–5.2) cm wide, initially globose to subglobose, sometimes lobate, almost completely closed and covering the whole stipe except its base, later convex to plane, opened basally and exposing a lamellate hymenium, with an involute, dentate margin; rugulose, finely tomentose, white or whitish at first, sometimes with translucent spots, especially at the periphery and margin, later yellowish with ochraceous areas (4A2–A3) and finally mostly bicoloured, yellowish buff (4A4) to ochraceous buff (5A4), with brown maculae (6C8–7D8). *Hymenophore* irregularly loculate inside, daedaleoid, lamellate in

the most external parts; locules large, 2–5 × 0.2–1 mm (0.5–1 per mm), labyrinthoid or flexuous in cross-section; lamellae irregular, more or less sinuous, frequently anastomosing and bifurcate, adnate, initially white, later cream, yellowish to ochraceous (4A3–A7), and finally ochre-orange or ochre-buff (6C8), with concolourous, acute or obtuse edges; white remnants of a partial veil can be observed in lamellae edges of some young basidiomata; fresh spore mass in locules yellowish white (4A2). *Stipe-columella* 1.6–4.2 × 0.7–1.8 cm, percurrent, well developed, cylindrical, straight or rarely flexuous, central or eccentric, white, initially solid and later medullary, usually broadening but rarely also tapering at the base; context white, not changing after contact with air. *Odour* pleasantly fruity, of pear or reminiscent of yoghurt and fruit; taste mild to slightly acrid.

*Spores* (8–)10–15(–17) × (7–)9–14(–15) µm, Q = (1.0–)1.10–1.25(–1.3), very variable in form and size, globose, subglobose, broadly ellipsoid to ovoid, heterotropic, echinate, some with one large, central oil drop; warts 0.6–1(–1.5) µm high, amyloid, irregularly conical or truncated, scattered or packed in few groups, with some verrucae among them; hilar appendix 1.2–1.8 µm long, cylindrical, with a distinct amyloid suprahilar plage in its base and conserving a non-functional apicular drop as in *R. mattioloana*. *Basidia* (1–)2–3(–4)-spored, 30–55 × 11–18 µm, clavate, some of them with one or more oil droplets and amorphous yellowish content; sterigmata (4–)6.5–7.5 µm long. *Basidioles* 30–35 × 10–15 µm, clavate. *Macrocystidia* abundant (40–)70–130(–180) × 12–20 µm, subcylindrical, lanceolate to fusiform, thin-walled or slightly thick-walled in the central portion, 1–1.6 µm thick, with abundant amorphous yellowish content, and an obtuse to acute or subulate apex, sometimes rostrate. *Paraphysoid cells* scarce, 12–26 × 6–9 µm, cylindrical, entire or 1-septate. *Subhymenium* cellular,



**Fig. 22** *Russula mediterraneensis*. a. MG630. Basidiomata. — b–i. GK6710 (BCN JMV800641, holotype). b. Basidiomata; c. external lamellar hymenium; d. daedaleoid hymenophore; e. pileipellis; f. hymenium and subhymenium; g. spores in Melzer; h–i. SEM images of spores. — j. GK7286. Spores in Melzer. — Scale bars: a–b = 1 cm; c–d = 5 mm; e–f = 20 µm; g, j = 10 µm; h–i = 5 µm. — Photos: a. M. Gelardi; b–d. G. Konstantinidis; e–g, j. J.M. Vidal; h–i. UdG.

consisting of 3–4 strata of globose to prismatic cells 9–28 µm diam. *Hymenophoral trama* heteromerous, made of hyaline, loosely interwoven septate hyphae 3–9 µm wide, with abundant sphaerocytes 21–47 µm diam, and some gloeoplera 2–7 µm diam. *Pileipellis* and *context* 100–300 µm thick; pileipellis composed of: 1) a trichodermal suprapellis made of erect to semi-erect, septate hyphae 2–6 µm diam, and abundant clavate, septate dermatocystidia 30–160 × 5–10 µm; and 2) a prosenchymatous subpellis of intricate hyphae 2–6 µm diam. *Pileal context* heteromerous, composed of septate hyphae 2–6 µm diam, mixed with several inflated hyphae and abundant nests and columns of sphaerocytes 21–47 µm diam. *Stipitipellis* a texture of interwoven, branched hyphae similar in shape and content to pileipellis dermatocystidia.

**Habitat, Distribution & Season** — Gregarious, subepigaeous among plant debris, in broadleaved woods of *Quercus* and *Castanea*, on siliceous soil. Summer. Occurring in the central and east-Mediterranean regions, in Greece and Italy, from sea level up to 1000 m altitude.

**Additional material studied.** GREECE, Central Macedonia, Kilkis, Koupa, 590 m, under *Quercus* sp., 8 July 2009, G. Konstantinidis (GK3930, duplicate BCN JMV800639)\*; North Aegean, Lesbos Island, Agiasos, under *Castanea sativa*, 8 June 2014, sine leg. (GK7286, duplicate BCN JMV800642)\*; Thessaly, Trikala, Logga, 1040 m, under *Quercus petraea*, 16 June 2012, G. Konstantinidis (GK6072, duplicate BCN JMV800640). — ITALY, Lazio, Roma, Nettuno, Torre Astura, 5 m, under *Quercus robur*, *Quercus frainetto* and *Quercus cerris* with presence of *Pinus pinea*, 2 Aug. 2014, M. Gelardi, M. Tulli & R. Polverini (MG630, duplicate MCVE 29085)\*; ibid., 9 Aug. 2014, M. Gelardi, M. Tulli & R. Polverini (MG636, duplicate MCVE 29086)\*.

**Notes** — *Russula mediterraneensis* belongs to the *R. globispora* complex of subsect. *Maculatinae*, and is therefore related with *R. mattioloana*, but differs because of its larger basidiomata, its hymenophore comprised of a daedaleoid, loculate inner part and a clearly lamellate external part, its subglobose to ellipsoid spores with lower warts and a distinct amyloid suprahilar plage, its trama and context featuring abundant sphaerocytes, and its ecological association with broadleaved trees. The large spore size (up to 17 µm) is similar to that observed in *R. globispora* and *R. mattioloana*, but different from those found in *R. dryadicola* and *R. heilongjiangensis*.

***Russula meridionalis* (Calonge et al.) J.M. Vidal, Mor.-Arr. & A. Paz, comb. nov.** — MycoBank MB828506; Fig. 23

**Basionym.** *Zelleromyces meridionalis* Calonge et al., Bol. Soc. Micol. Madrid 25: 302. 2000.

**Synonyms.** *Gymnomyces meridionalis* (Calonge et al.) J.M. Vidal, Rev. Catalana Micol. 26: 78. 2004.

*Gymnomyces dominguezii* Mor.-Arr. et al., Bol. Soc. Micol. Madrid 25: 301. 2000. (syn. nov.)

*Russula dominguezii* (Mor.-Arr. et al.) Trappe & T.F. Elliott, Fungal Systematics and Evolution 2: 361. 2018.

**Basidiomata** 1–2 cm wide, angiocarpic, sessile, subglobose to lobate or irregular. *Pileus* smooth, pale cream to ochraceous, drying dark reddish brown, intense red in contact with KOH. *Hymenophore* loculate, pale cream to ochraceous. *Columella* and sterile base present or absent. *Odour* fruity. Spores 8–11 × 7–10 µm, Q = 1.03–1.12, globose to subglobose, orthotropic; reticulum 0.4–0.6 µm high, composed of isolate warts and ridges, amyloid. *Basidia* 4-spored, 25–40 × 10–16 µm, clavate. *Macrocytidia* absent. *Cystidioles* like in var. *messapicoides*. *Hymenophoral trama* heteromerous. *Suprapellis* a trichoderm of septate hairs less developed than in var. *messapicoides*.

**Habitat, Distribution & Season** — Gregarious, hypogeous in continental sclerophyllous woods of *Quercus rotundifolia*, commonly on siliceous soil. Autumn and spring. Located in the western Mediterranean region, in Central Spain, between 400–1100 m altitude.


**Material studied.** SPAIN, Andalusia, Córdoba, Cabra, under *Quercus rotundifolia*, 21 June 1997, B. Moreno-Arroyo & J. Gómez (MA-Fungi 38502, holotype of *Z. meridionalis*; BM410, isotype)\*; Córdoba, Cabra, La Nava, 1070 m, under *Quercus rotundifolia*, 24 May 2015, C. Lavoise, A. Paz & R. Molina (BCN IC24051506)\*; Córdoba, Priego de Córdoba, Dehesilla Carcabuey, 700 m, under *Quercus rotundifolia*, 18 May 1993, J. Gómez (MA-Fungi 32069, paratype of *Z. meridionalis*); ibid., 20 May 2014, J. Gómez & A. Paz (BCN IC20051417)\*; Córdoba, carretera de Los Villares, under *Quercus rotundifolia* with *Cistus crispus* and *Pistacia lentiscus*, Feb. 1996, B. Moreno-Arroyo & J. Gómez BM412 (MA-Fungi 38572, holotype of *G. dominguezii*); Castilla and Leon, Valladolid, Urueña, montes Torozos, 830 m, under *Quercus rotundifolia*, on sandy soil, 17 June 2018, J. Cabero (JC-180617NR)\*; Extremadura, Cáceres, Torrejón el Rubio, Parque Nacional de Monfragüe, 370 m, under *Quercus rotundifolia*, on siliceous soil, 15 Apr. 2007, A. Paz (BCN IC15040721).

**Notes** — The holotype of *Z. meridionalis* is genetically very similar to the sequences of *R. messapica* and *M. messapicoides* analyzed in the present work. The three taxa seem to represent a monophyletic lineage, where no significant differences in ITS, 28S rDNA or *rpb2* data were found between *R. messapica* and *M. messapicoides*. Both taxa were in turn significantly different from the holotype of *Z. meridionalis* and other samples of this species, although no significant relationship was found between other collections identified as *Z. meridionalis*. Therefore, *M. messapicoides* is subsumed under the priority name *R. messapica*, while *Z. meridionalis* is re-combined into *Russula* but retained as an independent species due to the small differences found in ITS rDNA (2/715 bp different from *R. messapica* in sequences MK105663–MK105665 and MK105667) and *rpb2* (6/639 bp different in MK102762). These differences did not produce a significant phylogenetic support for a monophyletic *R. meridionalis*, but this was probably caused by the lack of data, especially *rpb2*, from all specimens tested. *Russula meridionalis* has a restricted geographic distribution (under *Quercus rotundifolia* in Southern and Central Spain), and constitutes at least a partially supported distinct genetic lineage, while *R. messapica* var. *messapicoides* occurs in the same areas as *R. messapica* var. *messapica* (under *Quercus ilex* in the Mediterranean basin, from Eastern Spain to Greece), but did not receive any significant support as a distinct genetic lineage employing ITS, 28S rDNA, *rpb2* and *tef1* data.

Basidiomata of *R. meridionalis* are angiocarpic and sessile or subsessile, with a completely loculate hymenophore, generally with traces of a columella and a small stipe or sterile base. The trichodermis (which is part of the universal veil) is reduced or partially lost in the substrate due to the hypogeous mode of growth, so the pileus does not always react intensely in contact with KOH 10 % as it does in *R. messapica*. Microscopical features of *R. meridionalis* are very similar to those of *R. messapica*, viz. tetrasporic basidia, subreticulate spores ornamented with an incomplete reticulum, and a more or less developed trichodermal suprapellis made of septate hairs with golden yellow encrusted pigment, which turns red in contact with KOH. Microscopical characters of *Gymnomyces dominguezii* (Moreno-Arroyo et al. 1999) also match those of *R. meridionalis*, viz. suprapellis a trichoderm, basidia tetrasporic and globose to subglobose spores 8–10 µm diam, ornamented with an incomplete reticulum 0.4–0.6 µm high. Despite the fact that attempts to produce genetic data from *G. dominguezii* yielded no useful results, this species is here considered a synonym of *R. meridionalis* because of their morphological similarities and identical geographical distribution.

***Russula messapica* Sarnari var. *messapica* in Sarnari, Boll. Assoc. Micol. Ecol. Romana 5, 18: 12. 1989 — Fig. 24a left**

**Material studied.** ITALY, Apulia, Taranto, Avetrana, Bosco Modunato, under *Quercus ilex*, on calcareous soil, 15 Dec. 2013, C. Agnello (AH 46373)\*. — SPAIN, Balearic Islands, Menorca, Alaior, Binixems, on calcareous soil, 120 m, under *Quercus ilex*, 18 Nov. 2011, J. Llistosella (BCN JL20111182)\*; Catalonia, Girona, Rupià, Bosc Geltrú, 120 m, under *Quercus ilex*, close to *R. messapica* var. *messapicoides*, 22 May 1993, J.M. Vidal (BCN JMV930522-12a).



**Fig. 23** *Russula meridionalis*. a. JC180617NR. Basidiomata. — b–c. IC15040721. b. Basidiomata; c. spores in Melzer. — d–e. IC20051417. Red reaction of the hairs of the suprapellis in contact with KOH 5 %. — f. IC24051506. Red reaction of the suprapellis in contact with KOH 5 %. — g–k. MA-Fungi 38502 (holotype of *Zelleromyces meridionalis*). g. Suprapellis in KOH 5 % and red Congo; h. spores in Melzer; i–k. SEM images of spores. — l. MA-Fungi 38572 (holotype of *Gymnomyces dominguezi*). Spores in Melzer. — Scale bars: a–b = 1 cm; c, h–i, l = 10 µm; d–g = 20 µm; j–k = 5 µm. — Photos: a. J. Cabero; b. A. Paz; c–h, l. J.M. Vidal; i–k. UdG.

#### *Russula messapica* var. *messapicoides* (Llistos. & J.M. Vidal)

J.M. Vidal, Llistos., Kaounas & P. Alvarado, comb. & stat. nov. — MycoBank MB828507; Fig. 24

Basionym. *Macowanites messapicoides* Llistos. & J.M. Vidal, Rivista Micol. AMB 38, 2: 155. 1995.

Synonym. *Russula messapicoides* (Llistos. & J.M. Vidal) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 236. 2018.


**Basidiomata** russuloid, pseudoangiocarpic, stipitate. **Pileus** 0.5–2 cm wide, rounded or occasionally bi- or trilobate, areolate, papillose, pale yellow to orange-yellow, intensely red in contact with KOH 10%; margin laterally open, alveolate to sublamellate. **Hymenophore** loculate to sublamellate, pale yellow to pale orange. **Stipe-columella** 0.3–0.7 × 0.15–0.2 cm, concolourous with pileus; context white. **Odour** fruity, similar to fermented fruits of *Sorbus domestica*.

**Spores** 8–10 × 7.5–9.5 µm, Q = 1.0–1.1, globose to subglobose, subheterotrophic to orthotropic; reticulum 0.6–1.2 µm high, amyloid, made by isolate warts forming ridges. **Basidia** 4-spored, 40–56 × 12–15 µm, clavate. **Macrocytidia** absent. **Cystidioles** 44–67 × 7–14 µm, scarce. **Hymenophoral trama** heteromerous. **Suprapellis** a trichoderm arranged in tufts of septate hairs 40–125 × 3–6 µm, with golden yellow encrusted

pigment (soon dissolved in KOH) and sometimes also intracellular pigment (reddish brown in KOH); basal elements ampullaceous, 8–30 µm diam; dermatocystidia absent.

**Habitat, Distribution & Season** — Gregarious, semi-hypogeous under fallen leaves, in sclerophyllous woods of *Quercus ilex* and *Q. coccifera*, commonly on calcareous soil, occasionally coexisting with *R. messapica* var. *messapica*. Spring to autumn. Widely distributed along the Mediterranean littoral, from Greece to Northern Spain, between 100–650 m altitude.

**Material studied.** GREECE, Central Greece, Attica, Katsimidi, 650 m, under *Quercus ilex* with *Pinus halepensis* and *Quercus coccifera*, on calcareous soil, 5 June 2013, V. Kaounas (VK2998)\*; ibid., 8 May 2014, V. Kaounas (VK3368)\*; ibid., 28 May 2014, V. Kaounas (VK3411, duplicate BCN JMV800682)\*; Crete, Retymnon, Arkadi, 520 m, under *Quercus ilex*, on calcareous soil, 27 Mar. 2016, V. Ramoutsakis (GK9341, duplicate BCN JMV800645)\*. — SPAIN, Castilla and Leon, León, Santibáñez de la Isla, under *Quercus rotundifolia*, *Quercus faginea* and *Fraxinus angustifolia*, on siliceous soil, 28 Nov. 2006, A. Paz as *M. messapicoides* (BCN IC28110613)\*; Catalonia, Girona, Rupià, Bosc Geltrú, 120 m, under *Quercus ilex*, in the same habitat as *Tuber aestivum*, 28 Apr. 1992, J.M. Vidal (BCN JMV920428-1, paratype of *M. messapicoides*); ibid., 22 May 1993, J.M. Vidal (BCN JMV930522-12b, paratype of *M. messapicoides*); ibid., 26 May 1993, J.M. Vidal & J. Llistosella (BCN JL1493, holotype of *M. messapicoides*)\*; ibid., 17 Oct. 2001, J.M. Vidal (BCN JMV20011017-1).



**Fig. 24** *Russula messapica* var. *messapicoides*. a. JMV930522-12a/b. Basidiomata of *R. messapica* var. *messapica* (left) and var. *messapicoides* (right). — b. VK2998b/3368b. Red reaction of the pileus in contact with KOH 10 %. — c. VK3411. Basidiomata. — d–e. JMV920428-1 (paratype of *M. messapicoides*). d. Basidiomata; e. spores in Melzer. — f. GK9341. Spores in Melzer. — g–m. JMV20011017-1. g. Basidiomata; h. red reaction of the hairs of the suprapellis in contact with KOH 5 %; i. detail of suprapellis and subpellis; j. hymenium with one cystidiole and subhymenium; k–m. SEM images of spores. — Scale bars: a–d, g = 1 cm; e–f = 10 µm; h–j = 20 µm; k–m = 5 µm. — Photos: a, d–j. J.M. Vidal; b–c. V. Kaounas; k–m. UdG.

**Notes** — *Macowanites messapicoides* was first proposed as a new species more or less identical with *Russula messapica*, except for its sequestrate habit (Llistosella & Vidal 1995). Martín et al. (1999) found that RFLP profiles of mt-LrDNA, mt-SrDNA and ITS nrDNA regions were identical in *R. messapica* and *M. messapicoides*, and only the highly variable IGS region could be used to discriminate between them. However, Martín et al. (1999) only analyzed a single specimen from each species, and these were found growing at the same time only a few meters apart from one-another (Llistosella & Vidal 1995). Martín et al. (1999) suggested that *M. messapicoides* was recently derived from *R. messapica*, but noted that the status of both taxa as independent species could not be confirmed by their results. Later, Miller et al. (2006) referred to this case as an example of an angiocarpic taxon being conspecific with a gymnocarpic species of *Russula*, even though they did not formally propose the synonymy. In the present work, four DNA markers (ITS, 28S rDNA, *rpb2*, *tef1*) were sequenced from several specimens of *M. messapicoides* and *R. messapica*, originating from different countries of the Mediterranean basin. Results show that no significant differences exist between the two taxa (*tef1* GenBank accessions MK102731–MK102735),

but both taxa are here kept as distinct varieties because of their different habit. The rank of variety is the first infraspecific category recommended by Art. 4 of the International Code of Nomenclature (Turland et al. 2018), and there is currently no consensus about the application of other ranks, such as subspecies or forma (Turner & Nesom 2000). *Russula messapica* var. *messapica*, *R. messapica* var. *messapicoides*, and *R. meridionalis* were significantly related to subsect. *Puellarinae* (PP 1.00, BP 90), in concordance with previous phylogenetic reconstructions of *Russula* (Miller & Buyck 2002, Bazzicalupo et al. 2017). The species most closely related are *R. cessans*, *R. odorata*, *R. puellaris* and *R. versicolor*.

Morphologically, *R. messapica* var. *messapicoides* is a sequestrate miniature of *R. messapica* var. *messapica*, where lamellae and stipe are progressively transformed into alveoli and columella. In the specimens from Attica (Greece) studied in the present work, the pileus is barely closed, the hymenophore is daedaleoid-sulphureous, and the stipe is well developed. On the other hand, the specimens from Catalonia (Spain) present an almost closed pileus, loculate hymenophore, and a reduced stipe. In all specimens, the suprapellis is a yellowish trichoderm that turns red in contact with KOH, the hymenium is formed by

tetrasporic basidia and scattered macrocystidia, and spores are subreticulated. The spores are ellipsoid and heterotrophic in *R. messapica* var. *messapica*, showing a suprahilar plage, and subglobose to globose, orthotropic, without a suprahilar plage in *R. messapica* var. *messapicoides*, which produces statismospores, except in Greek collections from Attica where the spores are subheterotrophic and the suprahilar plage is still present. Both taxa seem to occur under *Quercus ilex* across the entire Mediterranean region.

***Russula mistiformis* (Mattir.) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 236. 2018 — Fig. 25**

Basionym. *Martellia mistiformis* Mattir., Malpighia 14: 81. 1900.

Synonyms. *Hydnangium mistiforme* (Mattir.) Zeller & C.W. Dodge, Ann. Missouri Bot. Gard. 22: 372. 1935.

*Octaviania mistiformis* (Mattir.) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 743. 1958.

*Gymnomyces mistiformis* (Mattir.) T. Lebel & Trappe, Mycologia 92, 6: 1199. 2000.

*Martellia mediterranea* G. Moreno et al., Mycotaxon 42: 227. 1991. (syn. nov.)

*Russula mediterranea* (G. Moreno et al.) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 236. 2018.

**Basidiomata** 0.6–2.2 cm wide, angiocarpic, subglobose to obovoid or tuberiform, sessile, lacking a sterile base, but occasionally attached to the soil by a fragile mycelial strand the same colour as the pileus. **Pileus** at first finely tomentose, white, then smooth, with olivaceous, pastel yellow (3A4) or cream, pale orange (5A3) tones, finally displaying brown (7E8) maculae because of contact with air and handling; basally open and alveolate in old specimens. **Hymenophore** loculate, labyrinthoid, at first pale yellow (4A3) or pale orange (6A3), finally brown (6C6–7D7–7E8); locules 0.4–1.2 × 0.1–0.4 mm (2–4 per mm), radially arranged, elongated, sinuous; fresh spore mass in locules pale yellow (4A5) to brownish red (8C7); dark brown (6F8) in exsiccata. **Columella** absent. **Odour** in young specimens resembling that of an apple, very intense in mature specimens, similar to *Tuber melanosporum*.

**Spores** (8.5–)9.5–11(–12.5) × (8–)8.5–10(–10.5) µm, Q = 1.1–1.2, subglobose to ovoid, orthotropic, uniguttulate, echinate, orange to reddish in KOH; warts of regular length, 0.8–1.6(–3) µm high, yellow, deeply amyloid, cylindrical with an obtuse tip, sometimes interconnected with low ridges and with some isolated verrucae among them; hilar appendix about 2 µm long. **Basidia** scarce, typically 4-spored, but sometimes also 1–3-spored, 30–50 × 12–18 µm, clavate to broadly clavate, filled with small dark yellowish or dark reddish oil droplets, soon collapsing, originating in the deep subhymenium; sterigmata 3–7(–10) µm long, conical. **Basidioles** scarce, 21–35 × 7.5–12.5 µm, clavate. **Macrocystidia** absent. **Cystidioles** 27–40 × 6.5–9 µm; abundant in immature external locules, similar to dermatocystidia, cylindrical to fusoid, septate, sinuous, acute, capitulate, moniliform, filled with granular hyaline or yellowish content; rare in mature locules, cylindrical to fusoid or lageniform, typically capitate, not exceeding basidia in length. **Paraphysoid cells** abundant, 20–40 × 8–16 µm, commonly 1-septate. **Subhymenium** pseudoparenchymatous, formed of 2–3 layers of polygonal cells, 7–20(–25) µm diam. **Hymenophoral trama** 10–20 µm wide, formed of hyphae about 3–8.5 µm, connecting with those of pileal context; inflated elements up to 12 µm can be found sometimes in tramal anastomoses as well as sphaerocytes 10–20 µm diam. **Pileipellis** and context 80–240 µm thick, separable from the hymenophore; pileipellis thin, consisting of: 1) a trichodermal suprapellis, soon collapsing into a brownish mass, formed of very fragile, tapered dermatocystidia 20–40 × 3–6 µm, frequently with a mucronate apex; and 2) a prosenchymatous subpellis 20–30 µm thick, formed of interwoven subgelatinized hyphae 2–5.5 µm diam, slightly


differing from those of the pileal context. **Pileal context** 70–220 µm thick, prosenchymatous, formed by interwoven subgelatinized hyphae about 2–5.5 µm thick, lacking sphaerocytes. **Thromboplera** abundant in trama and context.

**Habitat, Distribution & Season** — Solitary to gregarious, hypogeous, associated with *Quercus*, *Castanea* and *Pinus*, on siliceous soil. From spring to late autumn. Found throughout the Mediterranean region, from Greece to Spain, from sea level up to 1200 m altitude.

**Material studied.** GREECE, North Aegean, Lesbos Island, 500 m, under *Castanea sativa* with *Crataegus monogyna* and *Quercus coccifera*, on siliceous soil, 29 Jan. 2017, G. Fransouas (BCN JMV800652)\*. — ITALY, Sardinia, Nuoro, Orune, under *Quercus suber*, May 1900, U. Martelli (FH, herb. N. Patouillard, original material of *M. mistiformis*); Oristano, Pau, Is Lottus, 550 m, under *Quercus ilex*, on siliceous soil, 1 May 1989, P. Fantini (BCN JMV800124); Oristano, Santu Lussurgiu, Rio e Messi, 650 m, under *Quercus pubescens* and *Alnus glutinosa*, on siliceous soil, 28 Oct. 1999, P. Fantini (BCN JMV800125)\*. — SPAIN, Castilla and Leon, Segovia, La Granja de San Ildefonso, 1200 m, under *Quercus pyrenaica*, on siliceous soil, 27 Nov. 1997, F. García (AMC H-69, duplicate BCN JMV971127)\*; Zamora, Tábara, under *Pinus sylvestris* with *Quercus rotundifolia* and *Erica arborea*, on siliceous soil, 6 Dec. 2010, J. Cabero (JC101206BT, duplicate BCN JMV800663)\*; ibid., 2 Jan. 2011, J. Cabero (JC110102NR)\*; ibid., 17 Nov. 2013, J. Cabero (JC131117NR, duplicate BCN JMV800662)\*; Zamora, Villar del Buey, under *Quercus pyrenaica*, on siliceous soil, 5 Mar. 2017, J. Cabero (JC170305NR, duplicate BCN JMV800661)\*; Catalonia, Girona, Quart, Sant Mateu de Montnegre, Les Gavarres, 350 m, under *Quercus ilex*, *Quercus suber* and *Pinus pinaster*, 7 June 2014, F. Rodríguez (BCN JMV20140607-3)\*; Girona, Sant Sadurní de l'Heura, Can Torrent, Les Gavarres, 150 m, under *Pinus radiata*, 23 May 1991, J.M. Vidal as '*M. pila*' (BCN JMV910523-3); ibid., 11 May 1992, J.M. Vidal as '*M. pila*' (BCN JMV920511-3); ibid., 26 June 1992, J.M. Vidal as '*M. pila*' (BCN JMV920626-1)\*; ibid., Pla de Banyeres, Les Gavarres, 80 m, under *Pinus pinea*, on siliceous soil, 24 May 2016, F. Rodríguez & J.M. Vidal (BCN JMV20160524-1)\*; Girona, Santa Cristina d'Aro, Romanà, Can Pons, Les Gavarres, 365 m, under *Pinus pinaster*, 26 Dec. 1990, J.M. Vidal as '*M. pila*' (BCN JMV901226-6); ibid., 20 Nov. 1993, J.M. Vidal, A. Montecchi & M.P. Martín (BCN JMV931120-9); ibid., 26 Dec. 1996, J.M. Vidal (BCN JMV961226-1); Extremadura, Cáceres, Parque Nacional de Monfragüe, under *Quercus suber* with *Cistus ladanifer*, 9 Nov. 1987, A. Montecchi et al. (AH GM-RG11057, holotype of *M. mediterranea*; BCN JMV800632, isotype)\*; Cáceres, Jarandilla de la Vera, under *Quercus pyrenaica* and *Pinus* sp., 15 Dec. 2012, A. Paz (BCN IC15121207/IC15121208/IC15121209).

**Notes** — *Martellia mistiformis* was described and illustrated by Mattiolo (1900) to accommodate several collections found by U. Martelli under *Quercus ilex* and *Q. suber* in the island of Sardinia (Italy), as well as additional collections from the island of Sicily. Mattiolo highlighted the hyphal hymenophoral trama, the predominantly tetrasporic basidia and the umber-coloured, globose to subglobose or slightly ellipsoid, echinate spores of 10 µm diam. This species was redescribed by Lebel & Trappe (2000) in a paper dedicated to the study of the generic types of sequestrate Russulales, and recombined into the genus *Gymnomyces* after observing sphaerocytes in the hymenophoral trama of the isotypic material at FH and NY herbaria. Recently, this species was re-combined into genus *Russula* by Elliott & Trappe (2018).

The first genetic data of *R. mistiformis* obtained by Calonge & Martín (2000) from specimens collected and illustrated by Vidal (1991a) as *Octaviania pila*, and subsequently described by Vidal (1991b) as *Martellia pila*, were found to have a close relationship with *R. foetens*. These early sequences (AF230893 and AF230894) contained some sequencing errors and so new data from the same collections, GM-RG11057 (AH) and JMV920626-1 (BCN), were produced in this study (Table 1). Whitbeck (2003) obtained additional sequences from a specimen collected by A. Montecchi in Italy (AM1653). In the present work, new collections from Italy and Spain were tested and shown to be genetically identical to one-another. ITS and 28S rDNA obtained from the holotype of *Martellia mediterranea* (AH GM-RG11057), were identical to those of *R. mistiformis*, and both species have also very similar morphology, ecological preferences and geographi-



**Fig. 25** *Russula mistiformis*. a. Reproduction of Mattioli's plate of *Martellia mistiformis* (Mattioli 1900, tav. 1). — b. FH, herb. N. Patouillard (original material of *M. mistiformis*). Spores in KOH 5 %. — c–f. JMV920626-1. c. Spores in Melzer; d. basidiomata; e–f. SEM images of spores. — g–k. JMV961226-1. g. Basidiomata; h. pileipellis and context; i. suprapellis; j–k. cystidioles of external immature locules. — l–n. JMV20160524-1. l. Cystidiole of mature hymenium; m. basidia, basidioles, paraphysoid cells and trama; n. spores in KOH 5 %. — o. AH GM-RG11057 (holotype of *Martellia mediterranea*). Spores in KOH 5 %. — Scale bars: b–c, i, n–o = 10 µm; d, g = 1 cm; e–f = 5 µm; h, j–k, m = 20 µm. — Photos: b–d, g–o. J.M. Vidal; e–f. UdG.

cal distribution, so they are here considered synonyms. Trappe (in Moreno et al. 1991), studied the holotype of *M. mistiformis* at TO and the isotype at FH, concluding that the spores of *M. mediterranea* are smaller than those of *M. mistiformis*, and the spore ornamentation thinner. These differences in spore size and ornamentation are here considered merely as part of the intraspecific variability of *R. mistiformis*.

*Russula mistiformis* belongs to sect. *Ingratae* subsect. *Foetentinae*, and is closely related to other European sequestrate russuloid taxa, such as *R. cerea* and *R. pila*. It differs from these species due to the thickness of pileipellis and context (80–240 µm in *R. mistiformis*, 70–125 µm in *R. cerea*, 150–500 µm

in *R. pila*), bisporic or tetrasporic basidia (typically bisporic in *R. cerea*, tetrasporic in *R. pila*), hymenium with capitate cystidioles and abundant paraphysoid cells, but lacking macrocystidia (macrocystidia thick walled and clavate in *R. cerea*, macrocystidia and cystidioles absent in *R. pila*) and subglobose to ovoid spores (globose in *R. cerea* and *R. pila*). In addition, the three species seem to be ecologically isolated, with *R. mistiformis* being apparently endemic to the Mediterranean region under *Quercus* and *Pinus*, while *R. cerea* is associated with lowland and montane trees (conifers and broadleaved trees) in Mediterranean and Central Europe, and *R. pila* shows preference for *Fagus sylvatica* stands of the Mediterranean subalpine range.

**Russula monospora** (Boud. & Pat.) Trappe & T.F. Elliott,  
Fungal Systematics and Evolution 1: 236. 2018 — Fig. 26

Basionym. *Hydnangium monosporum* Boud. & Pat., J. Bot., Paris 2, 24: 445. 1888.

Synonyms. *Octaviania monospora* (Boud. & Pat.) Lloyd, Mycol. Writings 7, Letter 67: 1141. 1922.

*Martellia monospora* (Boud. & Pat.) Astier & Pacioni, Doc. Mycol. 28, 109–110: 9. 1998.

**Basidiomata** 1.5–5.5 cm wide, angiocarpic, subglobose to ovate-pyriform or tuberiform, often caespitose and deformed, irregularly depressed at the top, sometimes longitudinally plicate and basally alveolate, attached to the soil by a minute sterile base that often remains in the substrate when the basidiomata are extracted. **Pileus** dry, finely tomentose to minutely papillate-squamulose under the lens, evanescent (old specimens nude), pale orange (5A2) to melon (5A6) or greyish orange (5B6), covered by fulvous yellowish to ferruginous brown hairs, with wine-red (11D7) and olivaceous (3C3) maculae when handled; pale orange (5A2) to dark brown (7F7) in exsiccata. **Hymenophore** firm, minutely loculate, labyrinthoid, unevenly ripening, at first pure white to ochraceous, becoming pink (13A3) to purplish red (13A6) at maturity; greyish orange (5B4) to brown (7E5) in exsiccata; locules minute, 0.4–1 × 0.1–0.3 mm (2–4 per mm), elongated, compressed, flexuose, empty or filled of spores; fresh spore mass in locules initially white to yellowish, finally pink (12A3–A6) when mature; yellowish white (4A2) in immature exsiccata, and greyish red (10D5–11D5) to brownish red (10D6–11D6) in mature exsiccata. **Columella** absent. **Odour** pleasant, fruity, strongly of pineapple according to Boudier & Patouillard (1888).


**Spores** (9–)10–13 µm, Q = 1, perfectly spherical, orthotropic, uniguttulate, echinate, hyaline, colourless for a long time, then yellowish with purplish pink hues, and finally dark pink; in contact with Melzer's reagent the immature hyaline spores stain dark yellow and the mature pink spores stain orange; warts 0.4–1.4(–1.6) µm high, isolated, inamyloid or weakly amyloid, conical, with some verrucae among them; hilar appendix 1.5–3 × 1.5–2 µm long, sometimes retaining a sterigmal appendix. **Basidia** 1-spored (rarely 2-spored), 30–60 × 7–12 µm, cylindrical to oblong-clavate when young, then sinuous, constricted, lageniform-urticiform with a long neck, filled of yellow oleiferous guttules, soon collapsed; sterigmata 4–10 µm long, tapering above. **Basidioles** clavate, 17–38 × 7–17 µm, with numerous oleiferous guttules inside. **Macrocytidia** and **cystidioles** absent. **Paraphysoid cells** abundant, aseptate or 1-septate, similar to basidioles, but internally empty. **Subhymenium** ramosa, composed of chains of 1–3 elongated cells measuring 7–15 × 4–9 µm. **Hymenophoral trama** 30–50 µm wide, homoiomerous, composed of loosely interwoven, hyaline, subgelatinized hyphae, 2–6 µm diam, filled of oleiferous guttules; sphaerocytes absent. **Pileipellis** and **context** 150–350 µm thick, lacking in old specimens; pileipellis consisting of: 1) an intricate trichodermal suprapellis of dark yellow, septate hairs and dermatocystidia, 40–100 × 2–4 µm, with a thick wall up to 1 µm, finally collapsing into a brownish mass; and 2) an undifferentiated prosenchymatous subpellis. **Pileal context** 125–250 µm thick, densely prosenchymatous, composed of subgelatinized hyaline hyphae 2–5 µm diam; sphaerocytes absent. **Gloeoplera** up to 5 µm diam, abundant in the trama and context. Thromboplera also present.

**Habitat, Distribution & Season** — Hypogeous to semi-hypogeous, solitary to gregarious, sometimes in compact, caespitose groups, under conifers (*Pinus nigra*) or deciduous trees (*Quercus*), on calcareous soil. Summer to winter. Found in Mediterranean and submediterranean regions of Southern Europe (Bulgaria, France and Spain), between 300–1150 m altitude.

**Material studied.** BULGARIA, Blagoevgrad, Ilindentsi, Struma Valley, 550 m, under *Pinus nigra* subsp. *nigra* with *Corylus avellana*, *Fraxinus ornus* and *Ulmus* sp., on calcareous soil, 16 Jan. 2014, M. Slavova (MSL0932F1017); ibid., 31 Jan. 2016, M. Slavova (MSL1689F7395); ibid., 11 Dec. 2017, M. Slavova MSL2032F2679 (SOMF 29974, duplicate BCN JMV800686). — FRANCE, Franche-Comté, Jura, Abbévillers, sine dat., L. Quélet as 'H. galatheum' (UPS F013936, herb. L. Quélet); Provence-Alpes-Côte d'Azur, Nice, July 1885, 'dedit D. Barla' (PC, herb. E. Boudier, lectotype of *H. monosporum*); ibid., sine dat., ex E. Boudier as *H. monosporum* (BPI 712228, coll. C.G. Lloyd 7201); ibid., sine dat., 'fragmentum spec. orig.', ex E. Boudier as *H. monosporum* (UPS F016541, ex herb. G. Bresadola); Nice, Drap, Grand Bois, 12 July 1886, J.-B. Barla as 'H. candidum' (NICE 2012-0-06711/06712); ibid., 23 June 1887, J.-B. Barla as 'H. candidum' (NICE 2012-0-06714); 'environ de Nice', July 1887, as *H. monosporum* (FH, herb. N. Patouillard); Nice, Drap, La Bauma, 8 Aug. 1887, J.-B. Barla as 'H. candidum' (NICE 2012-0-06713); Nice, Drap, Oct. 1890, J.-B. Barla as 'H. candidum' (NICE 2012-0-06715-1); Nice, Drap, Grand Bois, 30 June 1891, J.-B. Barla as *H. monosporum* (NICE 2012-0-06715-2/6716/6717/6718). — SPAIN, Castilla and Leon, Soria, San Leonardo de Yagüe, Sistema Ibérico, under *Quercus rotundifolia*, on calcareous soil, 15 June 2001, A. Sanz-Becerra & N. Redondo (AH 46459)\*; Castilla-La Mancha, Cuenca, Área Recreativa de Los Lagunillos, Serranía de Cuenca, Sistema Ibérico, 1150 m, in *Pinus nigra* subsp. *salzmannii*/forest with *Buxus sempervirens* and *Crataegus monogyna*, on calcareous soil, 10 Nov. 2017, J.A. Martínez (BCN JMV800671)\*; ibid., 1120 m, in *Pinus nigra* subsp. *salzmannii* forest with *Quercus faginea* and *Juniperus communis*, on calcareous soil, 16 Dec. 2017, A. Carreres & J.A. Martínez (BCN JMV800672)\*; ibid., 16 Dec. 2017, A. Carreres & J.A. Martínez (BCN JMV800673, duplicate AH 50141)\*.

**Notes** — This rare species was collected by J.-B. Barla between 1885 and 1891 in Nice (Southern France), in the vicinity of Drap, in the places called Grand Bois and La Bauma (Trimbach 1996), and initially identified as *Hydnangium candidum*. Parts of these collections were sent to Boudier and also to Patouillard who, after studying them and observing the presence of monosporic basidia in all the specimens, published this species as *Hydnangium monosporum* (Boudier & Patouillard 1888), which was later illustrated by Boudier (1906: pl. 193) in his *Icones Mycologicae*. The lectotype of *H. monosporum* (PC) designated by Nuytinck et al. (2003), as well as original and authentic collections deposited in various public herbaria (BPI, FH, NICE, UPS), were checked for the purpose of this study. Despite being completely or partially immature, or in some cases parasitized by moulds, several microscopic features were noted, such as a trichodermal pileipellis, monosporic basidia, and globose, echinate spores measuring 9–13 µm diam, still hyaline or yellowish, but showing an intense yellow reaction in contact with Melzer's reagent. Therefore, the specimens collected by Barla were probably too immature and led Boudier (1906) to depict them as yellowish. The sample collected by Quélet (1886) in the Jura region of France displays slightly more mature spores, with purplish pink tones in the spore wall under ammonium 10 %. Modern Spanish and Bulgarian collections studied in the present work show an intense pink colour in mature spores and hymenophore, a unique character discriminating this species from all other European sequestrate Russulaceae taxa. Recently, this species was re-combined into genus *Russula* by Elliott & Trappe (2018). *Russula monospora* can be confused with *Lactarius stephensii*, which also has monosporic basidia, a ramosa subhymenium, a homoiomerous hyphal trama, and echinate spores. However, in *L. stephensii*, spores are distinctly subglobose instead of spherical, while numerous laticifera are present in the trama and context, which are completely absent in *R. monospora*.

Genetically, *R. monospora* shows no significant relationships, but is probably close to *R. consobrina* (PP 0.94, BP 53) and other lineages such as subsections *Russula*, *Viscidinae* and *Sardoninae* (PP 0.73, BP 50). The only other sequestrate species in these clades is *R. gilkeyae* (Trappe 2572, listed as 'Gymnomyces gilkeyae', and OSC 117360, as 'G. monosporus'), which probably belongs to subsect. *Sardoninae* (Trendel et al. 2017).



**Fig. 26** *Russula monospora*. a. Reproduction of Boudier's plate of *Hydnangium monosporum* (Boudier 1906: pl. 193). — b–c. PC, herb. E. Boudier (lectotype of *H. monosporum*). b. Pileipellis; c. immature spores in Melzer. — d–e. UPS F013936, herb. L. Quélet (as '*Hydnangium galatheum*'). d. Immature spores in Melzer; e. immature spores in ammonia showing the purplish pink hues of the spore wall. — f–h. MSL2032F2679. f. Basidiomata; g. hymenium and trama; h. immature spores in Melzer. — i–j. JMV800672. i. Basidiomata initiating maturation; j. pileipellis and context. — k–q. JMV800673. k. Basidiomata showing the non-uniform ripening of hymenophore; l. detail of monosporic basidium; m. emerging necks of basidia; n. mature spores in Melzer; o. mature spores in ammonia; p–q. SEM images of spores. — Scale bars: b, g, j, l–m = 20 µm; c–e, h, n–o = 10 µm; f, i, k = 1 cm; p–q = 5 µm. — Photos: b–e, g–h, j, l–o. J.M. Vidal; f. M. Slavova; i, k. A. Carreres & J.A. Martínez; p–q. UdG.

**Russula neuhoffii** (Soehner) J.M. Vidal, comb. nov. — MycoBank MB828508; Fig. 27

Basionym. *Hydnangium neuhoffii* Soehner, Z. Pilzk., N.F. 20, 3–4: 111. 1941.

Synonyms. *Octaviania neuhoffii* (Soehner) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 202. 1958.

*Hymenogaster pisiformis* Velen., Opera Bot. Cech. 4: 96. 1947. (syn. nov.)

Lectotype of *Hydnangium neuhoffii* (here designated MBT384233): GERMANY, Brandenburg, 'Liebenthaler Wäldchen bei Marienwerder', July 1926, W. Neuhoff (M, herb. E. Soehner 1060).

**Basidiomata** 1–3 cm wide, angiocarpic, globose, ovoid to irregular, slightly sulcate and tuberous, sessile, with a minute sterile base. **Pileus** pruinose, initially pure white, later orange-white to pale orange (5A2–A4), with brown maculae (7D7). **Hymenophore** loculate, labyrinthoid, initially yellow to orange-yellow, then ochre; pale orange in exsiccata (5A4); spore mass in locules pale orange (5A4). **Stipe-columella** not observed. **Odour** mild. (Description based on Soehner 1941, Velenovský 1947 and herbarium material).


**Spores** 9.5–12.5 × 8.5–11 µm, Q = 1.0–1.1, globose to subglobose, orthotropic to subheterotropic, statismosporic, echinate, intensely yellow; warts 0.5–1 µm high, amyloid, densely packed and short, cylindrical to dentiform, or rounded at the apices, with numerous verrucae among them; hilar appendix 1–2 × 1.5 µm, straight, cylindrical to conical, sometimes retaining a sterigmal appendix; suprahilar plage present, inamyloid. **Basidia** 2–3-spored, 27–40 × 9–15 µm, clavate to broadly clavate; sterigmate 3.5–6.5 µm long. **Macrocystidia** scarce, 55–70 × 7–9 µm, cylindrical, lanceolate, fusiform, or rostrate, with amorphous content, soon collapsed. **Paraphysoid cells** absent. **Subhymenium** made of a layer of globose to prismatic cells about 5–15 µm diam. **Hymenophoral trama** 40–50 µm wide, homoiomerous, made of septate, branched, tortuous hyaline hyphae 1.5–7 µm

diam; inflated elements and sphaerocytes rare or absent. **Pileipellis** and **context** 100–150 µm thick; pileipellis a trichoeplithelium formed by: 1) a trichodermal suprapellis made of septate hairs and dermatocystidia 15–30 × 2–5 µm, that soon collapses in a yellow granular mass; and 2) a pseudoparenchymatous subpellis 50–80 µm thick, made of ampullaceous cells up to 17 µm wide and globose cells 6–30(–40) µm diam. **Pileal context** 30–100 µm thick, formed by tortuous and intricate hyaline hyphae 2–5 µm diam, with numerous ampullaceous inflated elements up to 14 µm diam and some sphaerocytes up to 16 µm diam. **Gloeoplera** 3–5 µm diam, present in trama and context.

**Habitat, Distribution & Season** — Solitary to gregarious, hypogeous under broadleaved trees (*Quercus*, *Carpinus*, *Betula*), on siliceous soil. Summer and autumn. Found in temperate regions of Central Europe.

**Additional material studied.** CZECH REPUBLIC, Central Bohemia, Jidášky, near Mnichovice, 'in humo nigro ca 10 cm profundo, supra foliis quercinis et carpineis tecto in duobus speciminiibus in betuleto, statio quarcitica', 12 July 1945, J. Velenovský (PRM 153797, coll. J. Velenovský, holotype of *H. pisiformis*). — GERMANY, Bavaria, Planegg, near Munich, in oak forest, 28 Sept. 1941, E. Soehner as *H. neuhoffii* (M, herb. E. Soehner 1626).

**Notes** — Soehner (1941) described an angiocarpic fungus found by W. Neuhoff near Berlin, naming it *Hydnangium neuhoffii*, characterized by whitish basidiomata, yellow to fulvous loculated hymenophore, 1–3-spored basidia and echinate spores. Some years later, Velenovský (1947) described a similar hypogeous species collected in the south of Prague, which he named *Hymenogaster pisiformis*. Type collections of *H. neuhoffii* and *H. pisiformis* were compared in the present work and found to represent a single species of *Russula* characterized by the lack of laticifera, a pileipellis arranged in trichoeplithelium, and globose to subglobose amyloid spores provided with an inamyloid plage and ornamented with densely packed short



**Fig. 27** *Russula neuhoffii*. a–g. M, herb. E. Soehner 1060 (lectotype of *Hydnangium neuhoffii*). a. Pileipellis and context; b. suprapellis and subpellis; c. cells of the subpellis; d. hymenium; e. spores in Melzer; f–g. SEM images of spores. — h–i. PRM 153797 (holotype of *Hymenogaster pisiformis*). h. Hymenium and subhymenium; i. spores in Melzer. — Scale bars: a–b, d, h = 20 µm; c, e, i = 10 µm; f–g = 5 µm. — Photos: a–e, h–i. J.M. Vidal; f–g. UdG.

spines ( $< 1 \mu\text{m}$ ), with rounded apices. The type material of *H. neuhoffii* and *H. pisiformis* are parasitized by moulds and, unfortunately, we were unable to obtain recent collections of these species to sequence.

**Russula pila** (Pat.) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 237. 2018 — Fig. 28


*Basionym.* *Hydnangium pila* Pat., Bull. Soc. Mycol. France 26: 201. 1910.  
*Synonyms.* *Octaviania pila* (Pat.) Svrček in Pilát, Flora ČSR B1, Gasteromycetes: 199. 1958.

*Martellia pila* (Pat.) J.M. Vidal, Butll. Soc. Catalana Micol. 14–15: 172. 1991.

*Gymnomyces pila* (Pat.) Trappe et al., Mycotaxon 81: 200. 2002.

*Basidiomata* 1–3 cm wide, angiocarpic, subglobose to tuberiform, frequently caespitose, sessile, with an inconspicuous sterile base, occasionally attached to the soil by a basal mycelial strand. *Pileus* initially densely tomentose, greyish, then smooth, pale orange (5A4) to orange (6B6), finally dark brown (8F8) after handling or in contact with air, with aromatic viscid exudations in mature basidiomata; basally open and alveolate in old specimens. *Hymenophore* loculate, labyrinthoid, at first orange-white (6A2) or greyish orange (6B4), finally brownish orange (6C6); locules 0.3–1 × 0.1–0.3 mm (2–4 per mm), elongated, sinuous, minute; fresh spore mass in locules pale orange

(5A4) to reddish brown (5A4–8D8); brown (6E7) in exsiccata. *Columella* absent. *Odour* fruity or similar to tuber, taste mild. *Spores* (9–)10–12(–13)  $\mu\text{m}$ , Q = 1, globose, orthotropic, echinate, reddish in KOH; warts of irregular length, 1–2.5  $\mu\text{m}$  high, hyaline to yellow, deeply amyloid, cylindrical with obtuse tips, with some verrucae among them; hilar appendix short, inconspicuous. *Basidia* abundant, typically 4-spored, but also 2–3-spored, 35–45 × 11–17  $\mu\text{m}$ , broadly clavate, filled with granulose or homogeneous dark reddish content, originating deep in the subhymenium, persistent; sterigmata conical, 3–5  $\mu\text{m}$  long in tetrasporic basidia, up to 10  $\mu\text{m}$  in bisporic basidia. *Basidioles* abundant, 25–40 × 10–14  $\mu\text{m}$ , with tiny, hyaline to yellowish oily droplets. *Macrocytidia* absent. *Cystidioles* 30–60 × 3–6  $\mu\text{m}$ , similar to dermatocystidia, 1–3-septate, hyaline, cylindrical to fusoid, sinuous, acute, capitate, moniliform, present only in immature external hymenial locules. *Paraphysoid cells* scarce, 20–30 × (5–)8–14  $\mu\text{m}$ , commonly 1-septate. *Subhymenium* formed by 2–3 layers of prismatic cells about (6–)10–20  $\mu\text{m}$  diam. *Hymenophoral trama* 10–20  $\mu\text{m}$  wide, formed by branched hyphae (2–)4–6  $\mu\text{m}$  diam, with occasional inflated elements up to 12  $\mu\text{m}$  and chains of sphaerocytes in trmal anastomoses 6–20  $\mu\text{m}$  diam. *Pileipellis* and *context* 150–500  $\mu\text{m}$  thick, homogeneous, separable from the hymenophore; pileipellis consisting of: 1) a trichodermal suprapellis formed



**Fig. 28** *Russula pila*. a–k. JMV970816-8. a–b. Basidiomata; c. pileipellis and context; d. suprapellis; e–f. septate cystidioles of external immature locule; g. section of a septum; h. spores in KOH 5 %; i. spores in Melzer; j–k. SEM images of spores. — l–m. FH, herb. N. Patouillard (original material of *Hydnangium pila*). SEM images of spores. — Scale bars: a–b = 1 cm; c–g = 20  $\mu\text{m}$ ; h–i = 10  $\mu\text{m}$ ; j–m = 5  $\mu\text{m}$ . — Photos: a–i. J.M. Vidal; j–k. UdG; l–m. AH.

of subulate to lageniform, capitulate dermatocystidia 20–40 × 3–6 µm, that soon collapses into a yellowish mass; and 2) a prosenchymatous subpellis 50–250 µm thick, formed by a dense mesh of interwoven, subgelatinized, yellowish hyphae 2–4 µm diam. Pileal context 100–300 µm thick, formed by interwoven, subgelatinized, hyaline hyphae 2–4 µm diam, lacking sphaerocytes. Thromboplera present in trama and context.

**Habitat, Distribution & Season** — Commonly gregarious, caespitose, hypogeous to semi-hypogeous, in montane broad-leaved woods of *Quercus* and *Fagus*, in different types of substrates. Summer to winter. Found in temperate regions of Southern Europe, between 200–900 m altitude.

**Material studied.** FRANCE, Rhône-Alpes, Ain, Lepinay (Cras-sur-Reysouze), Jura mountains, under *Quercus* sp., Aug. 1909, N. Patouillard (FH, herb. N. Patouillard, original material of *H. pila*). — SPAIN, Catalonia, Girona, Vall de Bianya, Capsacosta, Coll Pregon, 900 m, under *Fagus sylvatica*, on sandstone soil, 16 Aug. 1997, J.M. Vidal (BCN JMV970816-8)\*; ibid., 18 Aug. 1997, J.M. Vidal (BCN JMV970818-2); Navarre, Lekunberri, road to Aralar, km. 6.5, under *Fagus sylvatica* and *Corylus avellana*, on calcareous soil, 23 Nov. 2011, P.M. Pasabán & F. Sáinz (BCN JMV800654)\*.

**Notes** — *Hydnangium pila* was proposed by Patouillard (1910), to name some collections of a semi-hypogeous fungus found in oak forests of the calcareous mountain range of French Jura. He reported it as sessile, at first white becoming reddish with age, with a dense and pubescent pileipellis easily separable from the hymenophore, tetrasporic clavate basidia and echinate globose spores measuring 10–12 µm diam, suggesting a close relationship of this species with *Martellia*. *Hydnangium pila* was therefore re-combined into *Martellia* by Vidal (1991b), because of these features, as well as its amyloid spores and the hyphal structure of the hymenophoral trama, and later re-combined into *Gymnomyces* by Trappe et al. (2002), following the synonymy of the genera *Martellia* and *Gymnomyces* proposed by Lebel

& Trappe (2000). Recently, this species was placed in genus *Russula* by Elliott & Trappe (2018).

*Russula pila* exhibits features reminiscent of both *R. mistiformis* and *R. cerea*. *Russula mistiformis* also has tetrasporic basidia, but differs in its subglobose to ovoid spores, thinner pileipellis and context and presence of cystidioles in the mature hymenium. *Russula cerea* also has globose spores, but differs in its abundant hymenial macrocystidia, bisporic basidia, and a thinner membranous pileipellis and context. Finally, *R. pila* and *R. cerea* are commonly found in montane woods, while *R. mistiformis* is found in Mediterranean woods. The three species are genetically related, but *R. cerea* and *R. pila* seem to be significantly closer to one-another. All of them probably belong to *Russula* sect. *Ingratae* subsect. *Foetentinae*.

#### *Russula vidalii* Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 234. 2018 — Fig. 29

Replaced synonym. *Gymnomyces ilicis* J.M. Vidal & Llistos., Rivista Micologica 38, 2: 160. 1995.

**Basidiomata** 0.5–3.5 cm wide, angiocarpic, subglobose to turbinete, bi-lobate, sessile or subsessile, with a residual stipe. **Pileus** pruinose, pure white, with pale orange to reddish brown maculae; old specimens nude, completely alveolate. **Hymenophore** loculate, yellowish white, then yellowish orange, pale orange, and finally deep orange. **Columella** white, branched. **Odour** fruity.

**Spores** 9–11(–13) µm, Q = 1.0–1.1, globose to subglobose, orthotropic; warts 0.5–1 µm high, isolated, amyloid, some forming short ridges or even an incomplete reticulum. **Basidia** 2-spored, 30–50 × 10–15 µm, clavate. **Macrocystidia** 43–85 × 6–10 µm. **Hymenophoral trama** developing nests of sphaerocytes. **Pileipellis** and **context** 175–300 µm thick, evanescent; supra-



**Fig. 29** *Russula vidalii*. a–b. JMV20160517-1. a. Basidiomata; b. hymenium and subhymenium. — c–g. JMV920401-1 (paratype of *Gymnomyces ilicis*). c. Basidiomata; d. suprapellis; e. spores in Melzer's reagent; f–g. SEM images of spores. — h–i. JC100508BT01. h. Basidiomata; i. SEM image of spores. — Scale bars: a, c, h = 1 cm; b, d = 20 µm; e = 10 µm; f–g, i = 5 µm. — Photos: a–e. J.M. Vidal; f–g. UdG; h–i. J. Cabero.

pellis a palisadotrichoderm of cylindrical to fusiform, erect dermatocystidia 5–12 µm thick; subpellis a cutis.

**Habitat, Distribution & Season** — Gregarious, semi-hypogeous under leaf litter of *Quercus ilex* and *Quercus rotundifolia*, sometimes mixed with *Pinus*, on calcareous soil. Spring and autumn. So far known from the western Mediterranean region, in France and Spain, from sea level up to 800 m altitude.

**Material studied.** FRANCE, Provence-Alpes-Côte D'azur, Bouches du Rhône, Saint Rémy de Provence, under *Quercus ilex* and *Pinus halepensis*, 17 Nov. 1993, L. Riousset (BCN JL1501, paratype of *G. ilicis*). — SPAIN, Castilla and Leon, Zamora, Toro, 800 m, under *Quercus rotundifolia*, on calcareous soil, 9 May 2010, J. Cabero as *G. ilicis* (JC100508BT01, duplicate BCN JMV800688)\*; Catalonia, Barcelona, Sant Just Desvern, Can Fetjó, 220 m, under *Quercus ilex*, 13 Apr. 1992, J. Llistosella, A. Rocabruna & J. Vila (BCN JL1497, holotype of *G. ilicis*); Girona, Rupià, Can Candell, under *Quercus ilex*, 3 Apr. 2005, F. Rodríguez (BCN JMV20050403-2); Girona, Sant Sadurní de l'Horta, Can Barris, 90 m, under *Quercus ilex*, on sandy basic soil, 1 Apr. 1992, J.M. Vidal (BCN JMV920401-1, paratype of *G. ilicis*); ibid., 17 May 2016, J.M. Vidal & F. Rodríguez (BCN JMV20160517-1)\*; Girona, Viladamat, Gorners, 40 m, under *Quercus ilex*, 10 Apr. 1991, J.M. Vidal (BCN JMV910410-3, paratype of *G. ilicis*); Girona, Viladamat, Palau Borrell, under *Quercus ilex*, 11 May 1996, J.M. Vidal (BCN JMV960511-2).

**Notes** — *Russula vidalii* is a replacement name for *Gymnomyces ilicis* (Elliott & Trappe 2018), an epithet originating from the apparent association of this taxon with *Quercus ilex* in the Mediterranean region. However, collections BCN JL1501 (Llistosella & Vidal 1995) and JC100508BT01 (Cabero 2011) were found in mixed stands of *Quercus* and *Pinus*, so it is possible that *R. vidalii* has a wider host range than originally thought. It is characterized by an evanescent pileipellis and context that gives mature basidiomata an alveolate look. Genetically, *R. vidalii* belongs to subsect. *Laricinae*, displaying close affinities with *R. laricinaaffinis* and *R. galileensis*. Basidiomata of *R. vidalii* are relatively small (1–2.5 cm), have a poorly developed columella, a residual stipe, and a loculate hymenophore, while *R. galileensis* produces larger basidiomata (up to 5–6 cm), with a conspicuous stipe-columella and a sublamellate hymenium. Spore warts of *R. vidalii* are isolated or forming short ridges, while basidia are bisporic, and the suprapellis is a palisadotrichoderm. By contrast, spore warts of *R. galileensis* are interconnected by small ridges, basidia are tetrasporic, and the suprapellis is an intricate trichoderm.

***Russula vinaceodora* (Calonge & J.M. Vidal) Trappe & T.F. Elliott, Fungal Systematics and Evolution 1: 240. 2018 — Fig. 30**

**Basionym.** *Macowanites vinaceodorus* Calonge & J.M. Vidal, Mycotaxon 79: 2. 2001.


**Basidiomata** russuloid, pseudoangiocarpic, stipitate. **Pileus** 3–8.5 cm wide, convex to plano-convex and depressed, smooth, viscid, initially pale yellow, then pinkish white to purplish brown, darker at the center, with pale orange maculae; margin open, lamellate. **Hymenophore** loculate in the upper zone and sub-lamellate in the lower, pale orange. **Stipe-columella** 1.5–4 × 0.8–2.5 cm, white; context white. **Odour** vinaceous, intense; taste sweetish to slightly acrid.

**Spores** 7–11 × 6–9.5 µm, Q = 1.05–1.17, globose to broadly ellipsoid, heterotrophic; reticulum 0.7–1.5 µm high, amyloid, made of crests and isolate warts. **Basidia** 2–4-spored, 30–45 × 11–16 µm, broadly clavate. **Macrocystidia** 50–75 × 10–16 µm, fusiform. **Hymenophoral trama** with large sphaerocytes up to 50 µm diam. **Subhymenium** cellular. **Pileipellis** and **context** 150–250 µm thick; suprapellis arranged as an intricate ixotrichoderm of erect hyphae and cylindrical to clavate dermatocystidia 30–75 × 2.5–11 µm; subpellis an intricate ixocutis. **Pileal** and **stipe-columella context** heteromerous.

**Habitat, Distribution & Season** — Gregarious, hypogeous to semi-hypogeous in sandy substrates, associated with *Pinus*, in fixed coastal dunes. Autumn. Located in the Atlantic coast of Southern Spain.

**Material studied.** SPAIN, Andalusia, Cádiz, Sanlúcar de Barrameda, Pinar de Algaida, littoral stabilized sand dunes, under *Pinus pinea*, 30 Oct. 2014, M. Becerra as *M. vinaceodorus* (AH 46374)\*; Huelva, Mazagón, littoral stabilized sand dunes, under *Pinus pinea* with *Corema*, *Halimium* and *Helichrysum*, 27 Nov. 1999, J.M. Vidal & F.D. Calonge (MA-Fungi 47416, holotype of *M. vinaceodorus*; BCN JMV991127-4, isotype); ibid., three young specimens growing fasciculate among *Macowanites ammophilus*, 27 Nov. 1999, J.M. Vidal & F.D. Calonge as *M. vinaceodorus* (MA-Fungi 46524; duplicate BCN JMV991127-5).

**Notes** — Recently, this species was re-combined into genus *Russula* by Elliott & Trappe (2018). *Russula vinaceodora* is characterized by its russuloid, pseudoangiocarpic basidiomata, with a pinkish white to purplish brown pileal colour, especially in the centre. Mature basidiomata have a very characteristic odour of fermented wine.



**Fig. 30** *Russula vinaceodora*. a. MA-Fungi 47416 (holotype of *Macowanites vinaceodorus*). Basidiomata. — b–f. JMV991127-4 (isotype of *M. vinaceodorus*). b. Suprapellis; c. hymenium and subhymenium; d. spores in Melzer; e–f. SEM images of spores. — g. MA-Fungi 46524 (as *M. vinaceodorus*). Loculated hymenophore of young basidioma. — Scale bars: a, g = 1 cm; b–c = 20 µm; d = 10 µm; e–f = 5 µm. — Photos: a–d, g. J.M. Vidal; e–f. UdG.

Genetically, *R. vinaceodora* belongs to subsect. *Laricinae*, and is closely related to *R. laricina*, *R. murrillii* and *R. nauseosa*. The most closely related sequestrate taxon is *R. sichuanensis* (Li et al. 2013), another pseudoangiocarpic species found in *Picea* forests of Xizang and Sichuan provinces of China, which has similar cream to yellowish lamellae, globose or subglobose spores ornamented with an incomplete reticulum, and short basidia. However, *R. vinaceodora* has a pinkish to violet pileus, different from the cream-coloured one of *R. sichuanensis*, a distinctive odour of wine, and a different habitat under *Pinus pinea*, in Mediterranean coastal sandy soils.

## IDENTIFICATION KEY TO EUROPEAN SPECIES OF SEQUESTRATE RUSSULACEAE

1. Hymenophore lactescent or with laticifera. Hymenophoral trama homoiomerous, lacking nests of sphaerocytes . . . . . *Lactarius* 2
1. Hymenophore not lactescent, lacking laticifera. Hymenophoral trama heteromerous, with nests of sphaerocytes, especially in tramal anastomoses . . . . . *Russula* 8
2. Spores echinate . . . . . 3
2. Spores reticulate . . . . . 6
3. Basidia 3–4-spored. — Spores 9.5–13(–15) × 8–10(–11) µm, subglobose to broadly ellipsoid; warts 1–1.5 µm high, isolated. Basidiomata 1–4 cm, subglobose to tuberiform, with a residual stipe, pale yellow with reddish brown maculae. Hymenophore loculate, pale yellow to pale brown. Latex scant, colourless to white, changing to yellow. In subalpine conifer woods (*Abies*, *Picea*). Temperate (Alps to Rhodopes) . . . . . *L. borzianus*
3. Basidia 1-spored . . . . . 4
4. Spores weakly amyloid. — Spores (11.5–)12–14.5(–15) × (10–)11–13 µm, subglobose to broadly ellipsoid; warts 0.5–2 µm high, isolated. Basidiomata 1–3.5 cm, subglobose to tuberiform, maize yellow to reddish brown. Hymenophore loculate, ochraceous to reddish brown. Latex scant and hyaline, changing to citrine yellow in young specimens, white, abundant, and almost immutable in old basidiomata. In montane woods of *Carpinus*, *Corylus*, *Fagus*, *Quercus*, *Tilia*. Temperate to submediterranean (British Isles to Southern Europe) . . . . . *L. stephensii*
4. Spores strongly amyloid . . . . . 5
5. Spores subglobose to broadly ellipsoid. — Spores 13–15 × 11–13 µm; warts 1–2 µm high, isolated or tooth-like fused. Basidiomata 1–4 cm, subglobose to tuberiform, at first pale orange then reddish brown to violet brown. Hymenophore loculate, deeply coloured, reddish yellow to orange red. Latex white. Under *Populus*. Temperate to Mediterranean (Belgium to Bulgaria) . . . . . *L. populicola*
5. Spores broadly ellipsoid to ellipsoid. — Spores 14–18.5 × 12–15 µm; warts 1–2 µm high, cylindrical, isolated or forming short ridges. Hymenium completely embedded in a dark orange substance. Basidiomata 1–2 cm, subglobose to tuberiform, orange to reddish brown or violet brown. Hymenophore loculate, deeply coloured, dull red to dull violet. Latex watery, scant. In montane conifer woods (*Abies*, *Pinus*) or under broadleaved trees (*Corylus*, *Quercus*). Temperate (Germany, Italy, Spain) . . . . . *L. soehneri*
6. Basidia 2–4-spored. — Spores 8.5–12.5(–13.5) × 7–9.5(–10.5) µm, subglobose to ellipsoid; reticulum 0.5–1 µm high, incomplete. Basidiomata 2–5 cm, obpyriform to tuberiform, pale orange to reddish brown, with minute depressions. Hymenophore loculate, yellowish white to orange-white. Latex scant, white, immutable. Taste sweetish, later astringent. In montane conifer woods (*Cedrus*, *Pinus*). Submediterranean (France, Morocco, Spain) . . . . . *L. josserandii*
6. Basidia 1-spored . . . . . 7
7. Reticulum 1–2 µm high, complete. — Spores 10–13 × 7–9 µm, ovoid to ellipsoid. Basidiomata 1–2.5 cm, subglobose to tuberiform, whitish to pale yellowish, pileus membranous, partially evanescent, indistinctly scrobiculate or with some scattered minute openings. Hymenophore loculate, whitish to pale cream or pinkish. Latex not observed. Under *Cistus* and *Halimium*. Mediterranean (Central Spain) . . . . . *L. giennensis*
7. Reticulum 0.5–1 µm high, incomplete. — Spores 8.5–11 × 6.5–8 µm, ovoid. Basidiomata 0.5–2.5 cm, globose to tuberiform, whitish to brownish buff, pileus membranous, partially evanescent, distinctly scrobiculate, with abundant large openings. Hymenophore loculate, whitish to cream or faintly ochraceous pink. Latex white. Under *Cistus*. Mediterranean (Cyprus) . . . . . *L. subgiennensis*
8. Basidiomata stipitate, pseudoangiocarpic. Spores heterotrophic. Amyloid suprahilar plage present . . . . . 9
8. Basidiomata sessile, angiocarpic. Spores orthotropic. Amyloid suprahilar plage absent . . . . . 16
9. Spores verrucose to echinate. Warts isolated or connected by low ridges . . . . . 10
9. Spores subreticulated . . . . . 14
10. Macrocytidia 45–70 µm long . . . . . 11
10. Macrocytidia 70–130 µm long . . . . . 13
11. Spores subglobose to ovoid. — Spores 7–9 × 5.5–7.5 µm; warts 0.25–0.75 µm high, some forming short ridges or connected by short lines. Pileus 2–7 cm, orange white to pale orange, with brownish orange and dark brown maculae; margin open, alveolate to sublamellate. Hymenophore loculate, orange-white to pale orange. Stipe-columella 1–4 × 0.7–2.5 cm, with brownish orange dots. Common in coastal sand dunes, under *Pinus*. Mediterranean (Southern Portugal and Spain). . . . . *R. ammophila*
11. Spores globose to broadly ellipsoid . . . . . 12
12. Warts interconnected with low ridges. — Spores 9.5–12.5 × 8.5–10.5 µm; warts 1.2–2 µm high, in groups of 2–4. Pileus 1.4–2.8 cm, pure white, belatedly maculated of pale yellow; margin closed or laterally open, sublamellate. Hymenophore loculate, pale yellow to yellow. Stipe-columella 1–2 × 0.2–0.4 cm. In montane broadleaved woods (*Carpinus*, *Corylus*, *Fagus*, *Quercus*) or conifer woods (*Abies*). Temperate (Eastern to Southern Europe) . . . . . *R. candidissima*
12. Warts isolated. — Spores 8.5–11(–12.5) × 7–9(–11) µm; warts 0.5–1.5 µm high. Pileus 0.5–3.5 cm, white to yellowish white, with yellowish orange maculae; margin closed or laterally open, alveolate. Hymenophore loculate, pale orange. Stipe-columella 0.6–1.5 × 0.15–0.4 cm. In montane broadleaved woods (*Carpinus*, *Corylus*). Temperate (Eastern to Southern Europe) . . . . . *R. candida*
13. Warts 0.6–1(–1.5) µm high. — Spores (8–)10–15(–17) × (7–)9–14(–15) µm, subglobose to broadly ellipsoid. Pileus 2.3–5 cm, yellowish to yellowish buff, with dark brown maculae; margin open, lamellate. Hymenophore sublamellate-daedaleoid, cream to ochre-orange. Stipe-columella 1.6–4.2 × 0.7–1.8 cm. Under broadleaved trees (*Castanea*, *Quercus*). Mediterranean (Greece and Italy) . . . . . *R. mediterraneensis*
13. Warts (0.7–)1.5–2.5(–3) µm high. — Spores (9–)10.5–15.5(–18) × (8–)10–15(–17) µm, globose to subglobose, some ellipsoid when immature. Pileus 1.2–3.8 cm, pale yellow to pale orange with dark brown maculae; margin open, sublamellate. Hymenophore loculate, pale yellow

- to titian red. Stipe-columella  $0.8\text{--}2.5 \times 0.3\text{--}1$  cm. In montane conifer woods (*Abies*, *Picea*). Temperate to submediterranean (Southern Poland, Greece and Italy) ..... *R. mattiroloana*
14. Reticulum  $0.5 \mu\text{m}$  high. — Spores  $(9\text{--})10\text{--}11.5\text{--}(15) \times (7.5\text{--}9.5\text{--}10.5\text{--}14) \mu\text{m}$ , subglobose; reticulum made of crests and warts. Pileus  $2\text{--}6$  cm, white with cream to pale umber maculae; margin radially alveolate when mature, but not open. Hymenophore loculate, cream-ochre. Stipe-columella  $0.5\text{--}3.5 \times 0.5\text{--}1.7\text{--}(3)$  cm. In sclerophyllous woods of *Quercus*. Mediterranean (Israel) ..... *R. galileensis*
14. Reticulum  $0.7\text{--}1.5 \mu\text{m}$  high ..... 15
15. Macrocytidia present. — Spores  $7\text{--}11 \times 6\text{--}9.5 \mu\text{m}$ , globose to broadly ellipsoid; reticulum made of crests and isolate warts. Pileus  $3\text{--}8.5$  cm, pinkish white to purplish brown; margin open, lamellate. Hymenophore loculate to sublamellate, pale orange. Stipe-columella  $1.5\text{--}4 \times 0.8\text{--}2.5$  cm. Odour intense, vinaceous. In littoral sand dunes, under *Pinus*. Mediterranean (Atlantic coast of Southern Spain) ..... *R. vinaceodora*
15. Macrocytidia absent. — Spores  $8\text{--}10 \times 7.5\text{--}9.5 \mu\text{m}$ , globose to subglobose; reticulum made of isolate warts and ridges. Pileus  $0.5\text{--}2$  cm, rounded or bi-trilobate, areolate, papillose, pale yellow to orange-yellow, intense red in contact with KOH; margin laterally open, alveolate to sublamellate. Hymenophore loculate to sublamellate, pale yellow to pale orange. Stipe-columella  $0.3\text{--}0.7 \times 0.15\text{--}0.2$  cm, concolourous with pileus. In littoral sclerophyllous woods of *Quercus ilex*. Mediterranean (Greece to Spain) ..... *R. messapica* var. *messapicoides*
16. Spores reticulated ..... 17
16. Spores echinate or verrucose ..... 18
17. Macrocytidia absent. — Spores  $8\text{--}11 \times 7\text{--}10 \mu\text{m}$ , globose to subglobose; reticulum  $0.4\text{--}0.6 \mu\text{m}$  high, made of isolate warts and ridges. Basidiomata  $1\text{--}2$  cm, subglobose to lobate or irregular, smooth, pale cream to ochraceous, drying dark reddish brown, intense red in contact with KOH. Hymenophore loculate, pale cream to ochraceous. In continental sclerophyllous woods of *Quercus rotundifolia*. Mediterranean (Central Spain) ..... *R. meridionalis*
17. Macrocytidia present but scarce. — Spores  $7\text{--}9.5\text{--}(10.5) \times 7\text{--}9\text{--}(10)$ , globose; reticulum  $0.5 \mu\text{m}$  high, complete to incomplete, made of low ridges and warts. Basidiomata  $2\text{--}7$  cm, turbinate, firmly rooted into the substrate, often cracked, cream-white to ochraceous cream, with ochraceous to brownish stains. Hymenophore loculate, ochraceous yellow to ochraceous orange, vinaceous in  $\text{FeSO}_4$ . In montane woods of *Pinus*. Mediterranean (Cyprus) ..... *R. hobartiae*
18. Pileipellis a trichoepithelium or an oedotrichoderm ..... 19
18. Pileipellis a trichoderm ..... 20
19. Pileipellis a trichoepithelium. — Spores  $9.5\text{--}12.5 \times 8.5\text{--}11 \mu\text{m}$ , globose to subglobose; warts dense  $0.5\text{--}1 \mu\text{m}$  high, isolated. Basidiomata  $1\text{--}3$  cm, globose to irregular, whitish. Hymenophore loculate, yellow, orange yellow to ochre. Under broadleaved trees (*Carpinus*, *Betula*, *Quercus*). Temperate (Central Europe) ..... *R. neuhoffii*
19. Pileipellis an oedotrichoderm. — Spores  $(6.5\text{--})7.5\text{--}9.5\text{--}(11.5) \times (6\text{--})7\text{--}9\text{--}(11) \mu\text{m}$ , globose to subglobose; warts up to  $0.3 \mu\text{m}$  high, some connected with low ridges. Basidiomata  $0.5\text{--}2$  cm, globose to subglobose, pruinose, whitish, with brownish red maculae. Hymenophore loculate, whitish at first, finally brownish red. Under *Cistus*. Mediterranean (Central Spain) ..... *R. andaluciana*
20. Basidia 1-spored ..... 21
20. Basidia 2–4-spored ..... 22
21. Macrocytidia present. — Spores  $13\text{--}15\text{--}(15.5) \times 12.5\text{--}14.5\text{--}(15) \mu\text{m}$ , globose to subglobose, weakly amyloid, yellow; warts dense,  $1.5\text{--}3 \mu\text{m}$  high, isolated. Macrocytidia numerous,  $30\text{--}70 \times 8\text{--}16 \mu\text{m}$ , clavate. Basidiomata  $1\text{--}2$  cm, subglobose to tuberiform, finely tomentose, pale orange with brown maculae. Hymenophore loculate, pale orange. Temperate (Germany) ..... *R. bavarica*
21. Macrocytidia absent. — Spores  $(9\text{--})10\text{--}13 \mu\text{m}$ , spherical, weakly amyloid, intense pink at maturity; warts  $0.4\text{--}1.4\text{--}(1.6) \mu\text{m}$  high, isolated. Basidia clavate to lageniform-urticiform, sometimes 2-spored. Basidiomata  $1.5\text{--}5.5$  cm, subglobose to tuberiform, finely tomentose to papillate-squamulose, pale orange to greyish orange with wine red and olivaceous maculae when rubbing. Old specimens nude. Hymenophore loculate, pink to purplish red at maturity. Under *Pinus* and *Quercus*. Mediterranean to submediterranean (Bulgaria, France and Spain) ..... *R. monospora*
22. Warts  $0.5\text{--}1 \mu\text{m}$  high. — Spores  $9\text{--}11\text{--}(13) \mu\text{m}$ , globose to subglobose; warts isolated, some forming short ridges or even an incomplete reticulum. Basidiomata  $0.5\text{--}3.5$  cm, subglobose to turbinate, with a residual stipe, pruinose, pure white, with pale orange to reddish brown maculae. Old specimens nude, completely alveolate. Hymenophore loculate, yellowish white to yellowish orange or deep orange. In sclerophyllous woods of *Quercus*. Mediterranean (France and Spain) ..... *R. vidalii*
22. Warts  $0.8\text{--}3 \mu\text{m}$  high ..... 23
23. Basidia 2-spored. Macrocytidia present. — Spores  $(8\text{--})9.5\text{--}12.5\text{--}(14) \mu\text{m}$ , globose; warts variable in length,  $1\text{--}3 \mu\text{m}$  high, isolated. Macrocytidia  $(25\text{--})30\text{--}50 \times (5\text{--})7\text{--}12\text{--}(16) \mu\text{m}$ , cylindrical to cylindro-clavate, thick walled. Basidiomata  $1\text{--}3$  cm, subglobose, smooth, greyish orange to pale brown, maculated of reddish brown. Hymenophore loculate, brown to reddish brown. In montane conifer woods (*Abies*, *Picea*, *Pinus*) or broadleaved woods (*Carpinus*, *Castanea*, *Corylus*, *Fagus*, *Quercus*). Temperate to submediterranean (British Isles to Southern Europe) ..... *R. cerea*
23. Basidia 1–4-spored. Macrocytidia absent ..... 24
24. Spores globose. — Spores  $(9\text{--})10\text{--}12\text{--}(13) \mu\text{m}$ ; warts variable in length,  $1\text{--}2.5 \mu\text{m}$  high, isolated. Basidiomata  $1\text{--}3$  cm, subglobose to tuberiform, caespitose, pubescent, greyish to pale orange or orange, maculated of dark brown and producing aromatic exudations. Hymenophore loculate, orange-white to greyish orange or brownish orange. In montane broadleaved woods (*Fagus*, *Quercus*). Temperate (Southern Europe) ..... *R. pila*
24. Spores subglobose to ovoid. — Spores  $(8.5\text{--})9.5\text{--}11\text{--}(12.5) \times (8\text{--})8.5\text{--}10\text{--}(10.5) \mu\text{m}$ ; warts of regular length,  $0.8\text{--}1.6\text{--}(3) \mu\text{m}$  high, isolated. Basidiomata  $0.6\text{--}2.2$  cm, subglobose to tuberiform, finely tomentose, pastel yellow to pale orange, maculated of brown. Hymenophore loculate, pale yellow, pale orange to brown. In woods of *Castanea*, *Pinus* and *Quercus*. Mediterranean (Greece to Spain) ..... *R. mistiformis*

## DISCUSSION

The multigene phylogenetic analysis and morphological revision of European sequestrate *Russulaceae* taxa carried out in this work, allowed us to clarify several taxonomic issues at specific and supraspecific levels. The overall topology of the multigenic analyses was not different from those obtained by other studies focused on gymnocarpic species (Shimono et al. 2004, Verbeken et al. 2014b, Kong et al. 2015, Looney et al. 2016, Buyck et al. 2017), except for lower support values for

some supraspecific lineages of *Lactarius*. These differences could be due to phylogenetic noise introduced by some sequences analyzed in the present work, or differences in the alignment of the highly variable ITS rDNA region. In contrast to *Lactarius*, multiple significant supraspecific clades were found within *Russula*, in concordance with previous phylogenetic reconstructions of this genus (Kong et al. 2015, Looney et al. 2016). Most of these clades include type species of subgenera, sections and/or subsections proposed by different classical authors on the basis of morphological evidence (Romagnesi 1985, Singer 1986, Bon 1988, Sarnari 1998, 2005), but until now, no formal re-arrangement had been proposed to incorporate the current genetic knowledge into the often overlapping nomenclature available. Kong et al. (2015) informally named the genetic lineages of *Russula* with a representative species or a supraspecific name without rank, and discussed the different supraspecific taxa in classical literature putatively matching these lineages. In the present work, eight major clades were found in *Russula* (Fig. 1). The largest ones (*Ingratae*, *Rigidae*, *Russula* s.str.) have often been treated as subgenera or sections, while the remaining were usually considered as sections or subsections. Treating all of them as sections could be a natural solution requiring few nomenclatural changes, with the major phylogenetic clades within these sections deserving the status of subsection.

Most European sequestrate Russulaceae species tested in the present work nested within previously known supraspecific clades, except two: *Russula monospora* and *R. hobartiae*. *Russula monospora* nested within a poorly supported clade formed by subsections *Russula* s.str. (= *Emeticinae*), *Sardoniinae* (= *Firmeae*) and *Viscidinae*. This clade was recovered also in the analyses of Kong et al. (2015), but received no significant support (PP 0.87). *Russula monospora* represents a distinct lineage within this clade, and so it could deserve its own subsection. However, a deeper study of the whole group and the other isolated lineages (e.g., those of *R. consobrina* and *R. fellea*), would be necessary to propose a stable nomenclature at the supraspecific level. The other taxon studied in the present work not matching any known supraspecific clade, is the new species *R. hobartiae*, which is significantly related with a putative specimen of *R. ochrophylla*. The identity of this specimen needs to be compared with the type collection (Adamčík et al. 2017), in order to support the introduction of a supraspecific name for the monophyletic lineage shared with *R. hobartiae*. All other European sequestrate Russulaceae taxa belong to existing supraspecific taxa, except for *R. ammophila*, which belongs to a significantly supported clade lacking a formal name, related to subsections *Foetentinae*, *Pectinatinae* and *Subvelatae*, within sect. *Ingratae*. This clade is often referred to as the /*R. amoena* clade, but genetic data support a formal name at the same rank of the other clades.

European sequestrate Russulaceae taxa have unequal intraspecific variability in the DNA markers analyzed. Some species were found to be genetically homogeneous (e.g., *R. mistiformis*), while others presented conspicuous random differences between most individuals (e.g., *R. candidissima*), or had a genetically homogeneous core and one or few individuals showing some differences (e.g., *R. mediterraneensis*). In addition, there was one species recently evolved from a gymnocarpic taxon, which could only be discriminated from the former using *tef1* marker (*R. ammophila*). The diverse phylogenetic status of these species suggests differences in their evolutive past and present, but these can be due to very different factors, such as their life-cycles, mating systems, metapopulation structure, hybridization and introgression events, or selection pressure, to name a few. The future of these lineages remains, of course, an open question, and so predictions based on their eventual

status are risky based on current evidence. Partially or recently isolated lineages may thus:

- 1) genetically diverge from each other and lead to fully isolated species;
- 2) merge again if reproductive barriers disappear; or
- 3) become extinct after the conditions that favored diversification change.

Two varieties of *Russula messapica* are here recognized: the original gymnocarpic phenotype (var. *messapica*) and a pseudoangiocarpic phenotype (var. *messapicoides*) that cannot be phylogenetically discriminated even using up to four DNA markers. This is a very rare case, but a few other examples can be found: for instance, a broad morphological variation from gymnocarpic to angiocarpic habit can be observed in *Hydnangium sublamellatum* (Bougher et al. 1993), or *Setchellliogaster tenuipes* var. *rheophyllus* (Martín & Rocabruna 1999, Lago et al. 2000, Martín & Moreno 2001). Hibbett et al. (1994) demonstrated that a pseudoangiocarpic forma can be triggered by a recessive allele at a single locus in *Lentinus tigrinus*, and so it is possible that specific matings or even environmental factors can be responsible for the apparition of sequestrate phenotypes. However, despite the evidence provided here by multigenic data, we cannot exclude the possibility that none of the markers selected reflect a hypothetical reproductive isolation between *R. messapica* var. *messapica* and *R. messapica* var. *messapicoides*. In addition, a closely related species with an exclusively angiocarpic habit (*R. meridionalis*) was found to have at least a partial genetic isolation, probably linked to its restricted geographic distribution. Despite the fact that ITS and 28S rDNA cannot be employed to significantly discriminate between *R. messapica* and *R. meridionalis*, the latter taxon is retained as an independent species because: 1) this is the most conservative decision in nomenclatural terms; and 2) the only *rpb2* sequence successfully obtained from samples of *R. meridionalis* (MK102762) seems different enough from those of *R. messapica* (MK102763–MK102766), with 6/634 bp varying between both species, vs only 1/634 variable sites among *rpb2* sequences of *R. messapica*. Therefore, a more complete *rpb2* dataset would likely support the separation between *R. meridionalis* and *R. messapica*.

Angiocarpic species are present in most lineages of *Lactarius* and *Russula*, confirming that transitions from gymnocarpic ancestors occurred multiple times. Such transitions were often explained as an evolutive adaption to adverse and arid environmental conditions (Thiers 1984, Buyck 1995, Bougher & Lebel 2001, Trappe & Claridge 2005, Smith et al. 2006), although angiocarpic species could be far more frequent in tropical climates than previously assumed (Verbeken et al. 2014b). Wilson et al. (2011) showed that angiocarpic species in Agaricomycetes appeared rather recently, but evolve at similar or even faster rates than the gymnocarpic lineages they have derived from, eventually predominating over them. In the present phylogenetic reconstructions, most angiocarpic species seem to be isolated in gymnocarpic lineages, but small groups of angiocarpic species can be found too (e.g., *R. candidissima* and related species within subsect. *Firmiores*). Interestingly, the clade named /*R. tapawera* is composed mainly of Australian and South American angiocarpic species with scattered gymnocarpic taxa, such as *R. purpureoflava*, *R. tawai* or *R. tricholomopsis* (Lebel & Tonkin 2007, Trierweiler-Pereira et al. 2015). Such an abundance of sequestrate species in the Southern Hemisphere could be due to: 1) very frequent transitions from gymnocarpic to angiocarpic states and/or a faster divergence rate of angiocarpic lineages; 2) an angiocarpic ancestor of the entire lineage with scattered reversions to a gymnocarpic state; or 3) a fragmented picture of the biogeographical distribution of gymnocarpic russuloid taxa in the Southern Hemisphere. Additional information is critically needed to evaluate these hypotheses.

**Table 2** Comparison of the different European sequestrate species of *Lactarius* and *Russula* ordered by systematic sections.

Basidiomata type	Columella	Suprapellis	Hymenophoral trama	Cystidia	Basidia	Spores ( $\mu\text{m}$ )	Ornamentation	Host
<i>Lactarius</i> subgen. <i>Lactiferus</i>								
<i>L. gennensis</i>	angiocarpic, sessile	branched, inconspicuous	trichoderm	homoiomericous homoiomericous	cystidioles absent	1-spored 1-spored	10–13 $\times$ 7–9 13–15 $\times$ 11–13	reticulum isolated and tooth-like fused warts
<i>L. populincola</i>	angiocarpic, sessile	absent or inconspicuous	trichoderm	homoiomericous	absent	1-spored	14–18.5 $\times$ 12–15	isolated warts and short ridges
<i>L. soehneri</i>	angiocarpic, sessile	absent	trichoderm	homoiomericous	absent	1-spored	12–14.5 $\times$ 11–13 (weakly amyloid)	isolated warts
<i>L. stephensi</i>	angiocarpic, sessile	absent or inconspicuous	trichoderm	homoiomericous	cystidioles	1-spored	8.5–11 $\times$ 6.5–7.8	incomplete reticulum
<i>L. subgiliennensis</i>	angiocarpic, sessile	branched, inconspicuous	trichoderm	homoiomericous	cystidioles	3–4-spored	9.5–13 $\times$ 8–10	isolated warts
<i>Lactarius</i> subgen. <i>Russularia</i>								
<i>L. borzianus</i>	angiocarpic, sessile	percurrent, branched	trichoderm to cedotrichoderm	homoiomericous	cystidioles	2–4-spored	8.5–12.5 $\times$ 7–9.5	incomplete reticulum
<i>L. josserandii</i>	angiocarpic, sessile	absent or inconspicuous	cedotrichoderm	homoiomericous	macrocystidia	2–4-spored	7–9 $\times$ 5.5–7.5	isolated warts and short ridges
<i>Russula</i> sect. <i>Ingratae</i>								
<i>R. ammophila</i>	angiocarpic to pseudoangiocarpic, stipitate	stipe-columnella	intricate trichoderm	heteromericous	macrocystidia	2-spored	9.5–12.5	isolated warts
<i>R. cerea</i>	angiocarpic, sessile	absent or inconspicuous	trichoderm	sphaerocysts only in trama anastomoses	macrocystidia	1–4-spored	9.5–11 $\times$ 8.5–10	isolated warts
<i>R. mistiformis</i>	angiocarpic, sessile	absent	trichoderm	sphaerocysts only in trama anastomoses	cystidioles	3–4-spored	10–12	isolated warts
<i>R. pila</i>	angiocarpic, sessile	absent	trichoderm	sphaerocysts only in trama anastomoses	proschnymatos	absent	7.5–9.5 $\times$ 7–9	isolated warts and low ridges
<i>Russula</i> sect. <i>Rigidae</i>								
<i>R. andaluciana</i>	angiocarpic, sessile	inconspicuous	cedotrichoderm	proschnymatos	macrocystidia	1–3-spored 2–4-spored	8.5–11 $\times$ 7–9 9.5–12.5 $\times$ 8.5–10.5	isolated warts
<i>Russula</i> sect. <i>Russula</i>								
<i>R. candida</i>	angiocarpic, stipitate	stipe-columnella	trichoderm	heteromericous	macrocystidia	4-spored	10–11.5 $\times$ 9.5–10.5	groups of 2–4 warts interconnected by low ridges
<i>R. candidissima</i>	angiocarpic to pseudoangiocarpic, stipitate	stipe-columnella	intricate trichoderm	heteromericous	macrocystidia	2–4-spored	7–9.5 $\times$ 7–9	reticulum of crests and warts
<i>R. galileensis</i>	angiocarpic, stipitate	stipe-columnella	palisadotrichoderm	proschnymatos	macrocystidia	2–4-spored	10.5–15.5 $\times$ 10–15	reticulum of low ridges and warts
<i>R. horbariae</i>	angiocarpic, sessile	absent or inconspicuous	trichoderm	heteromericous	macrocystidia	2–4-spored	10–15 $\times$ 9–14	isolated warts
<i>R. matitiroiana</i>	pseudoangiocarpic, stipitate	stipe-columnella	trichoderm	heteromericous	macrocystidia	2–3-spored	8–11 $\times$ 7–10	reticulum of ridges and isolated warts
<i>R. mediterraneensis</i>	pseudoangiocarpic, stipitate	stipe-columnella	trichoderm	heteromericous	cystidioles	4-spored	8–10 $\times$ 7.5–9.5	reticulum of ridges and isolated warts
<i>R. meridionalis</i>	angiocarpic, sessile	absent or present	trichoderm	heteromericous	absent	1-spored	10–13 (weakly amyloid)	isolated warts
<i>R. messapica</i>	pseudoangiocarpic, stipitate	stipe-columnella	intricate trichoderm	homoiomericous	macrocystidia	2-spored	9–11	short ridges
<i>R. var. messepicoides</i>	angiocarpic, sessile	absent	palisadotrichoderm	heteromericous	macrocystidia	2–4-spored	7–11 $\times$ 6–9.5	reticulum of crests and isolated warts
<i>R. monospora</i>	angiocarpic, sessile	percurrent, branched	intricate ixotrichoderm	heteromericous	macrocystidia	1-spored	13–15 $\times$ 12.5–14.5 (weakly amyloid)	isolated warts
<i>R. vidali</i>	angiocarpic, sessile	stipe-columnella	trichoderm	homoiomericous with endomacrocystidia	macrocystidia	2–3-spored	9.5–12.5 $\times$ 8.5–11	isolated warts
<i>R. vinaceodora</i>	pseudoangiocarpic, stipitate	absent	trichoderm	homoiomericous	macrocystidia	1-spored	13–15 $\times$ 12.5–14.5 (weakly amyloid)	isolated warts
<i>Russula</i> not classified								
<i>R. bavarica</i>	angiocarpic, sessile	absent						
<i>R. neuhoffii</i>	angiocarpic, sessile	absent	trichopithelium					

Speciation processes seem to be quite diverse among European sequestrate russuloid taxa. For example, *L. giennensis* and *L. subgiennensis* seem to represent independent genetic lineages maybe diverging as a result of allopatric speciation. Both species can be found associated with *Cistaceae* hosts in acidic soils in the Mediterranean basin, with *L. giennensis* apparently restricted to Western and Southern Spain, while *L. subgiennensis* is so far known only from the island of Cyprus at the opposite end of the Mediterranean basin. This clade apparently represents one of the few lineages within *Russulaceae* associated with *Cistaceae* hosts, although comprehensive phylogenetic assessments of fungal diversity within *Cistaceae* ecosystems are generally lacking (Fellner & Biber 1990, Lavorato 1991, Vila & Llimona 1999, 2002, 2006, 2009, Comandini et al. 2006, Torrejón 2007, 2009, Campos et al. 2010, Loizides & Kyriakou 2011, Malloch & Thorn 2011, Zambonelli et al. 2014, Leonardi et al. 2016, Loizides 2016). *Russulaceae* species associated with *Cistaceae* are not monophyletic, suggesting that multiple events of host specialization took place in the evolutionary history of this family. Moreover, speciation processes through biogeographic isolation (*L. josserandii* vs *L. borzianus*), host specialization (*L. populicola* vs *L. stephensi*), or climatic regions (*R. mistiformis* vs *R. cerea*), seem to be unstable in time, as no apomorphic features based solely on these factors could be identified in any section or subsection. Despite the loss of aerial spore dispersal, however, the majority of lineages do not appear to be endemic to specific continents, such as the widespread European species *R. candidissima*, which nests within a clade otherwise composed exclusively of American species. These observations are consistent with the 'generalized diversification rate' pattern found in *Russula* by Looney et al. (2016).

Morphological features traditionally employed to discriminate between sequestrate *Russulaceae* species seem to have a phylogenetic basis, although some of them were in need of re-interpretation. In Table 2, all European species studied are compared according to:

- 1) basidiomata development (pseudoangiocarpic, angiocarpic) and presence or absence of a stipe (stipitate, sessile);
- 2) presence or absence of a columella (percurrent, branched, stipe-columella);
- 3) suprapellis structure (trichoderm, intricate trichoderm, palisadotrichoderm, oedotrichoderm, trichoepithelium);
- 4) hymenophoral trama composition (heteromerous or homoiomerous);
- 5) presence or absence of cystidia (cystidioles, macrocystidia);
- 6) number of spores in basidia;
- 7) size of spores;
- 8) spore ornamentation (warts, reticulum, ridges, crests); and finally
- 9) family of putative symbiont plants (*Cistaceae*, *Fagaceae*, *Pinaceae*, *Salicaceae*).

The degree of development of the sequestrate syndrome (gasteromycetation) is indeed not only poorly correlated with classical taxonomical concepts, as already evident in other groups (Hibbett et al. 1997, Peintner et al. 2001, Hosaka et al. 2006), but probably variable within some species, and maybe even between basidiomes from the same mycelium. Hymenophoral trama, suprapellis structure, and number of spores per basidia seem to be features characteristic of supraspecific clades, in concordance with observations by Calonge & Martín (2000) and Lebel & Trappe (2000), and so is ectomycorrhizal morphology (Beenken 2004), while spore size, shape, and ornamentations, ecological traits, as well as classical macroscopic features (size, colour, odor, spore print), are still useful at the species level.

**Acknowledgements** The authors want to express their gratitude to Antoni Sánchez-Cuxart, curator of Herbarium BCN (Barcelona University, CeDocBiV, Barcelona), for his help with loan requests. To the directors and curators of the following herbaria: Universidad de Alcalá, Alcalá de Henares (AH), Asociación Vallisoletana de Micología, Valladolid (AVM), U.S. National Fungus Collections, Systematic Mycology and Microbiology Laboratory, Beltsville (BPI), Harvard University, Farlow Reference Library and Herbarium of Cryptogamic Botany, Cambridge (FH), University of Haifa, Institute of Evolution, Haifa (HAI), Leopold Franzens Universität, Institut für Mikrobiologie, Innsbruck (IB), Royal Botanic Gardens, Kew (K), Jagiellonian University, Institute of Botany, Kraków (KRA), Botanische Staatssammlung, München (M), Real Jardín Botánico, CSIC, Madrid (MA), Museo di Storia Naturale di Venezia, Venezia (MCVE), Institut de Botanique, Montpellier (MPU), Muséum d'Histoire Naturelle de Nice, Nice (NICE), New York Botanical Garden, New York (NY), Muséum National d'Histoire Naturelle, Laboratoire de Cryptogamie, Paris (PC), National Museum, Mycological Department, Praha (PRM), Uppsala Universitet, Museum of Evolution, Botany, Uppsala (UPS) and University of Vienna, Institute of Botany, Wien (WU).

The authors also thank the following collaborators for kindly providing collections and/or images or valuable information data: Carlo Agnello for the Italian collection of *R. messapica*, José Luis Becerra for the Spanish collections and photographs of *R. andaluciana*, Manuel Becerra for the Spanish collections of *R. ammophila* and *R. vinaceodora*, Alyona Biketova, Zohar Shafranov, Anatoliy Fedorenko, T. Pavlichek, S. Reshetnikov, I. Duckman, I. Shams and Y. Ur for providing collections, photographs and data of *R. galileensis*, José Antonio Cadiñanos for the Spanish collections of *R. amoena*, T. Chokova for the Bulgarian collection of *L. populicola*, Paolo Fantini for the Italian collections of *R. mistiformis*, Giorgos Fransouas for the Greek collection and photographs of *R. mistiformis*, Aurelio García-Blanco, Miguel Sanz-Carazo and J.B. Del Val for the Spanish collection of *L. giennensis*, Faustino García for the Spanish collections and photographs of *L. josserandii*, *R. cerea* and *R. mistiformis*, Matteo Gelardi (MG), María Tulli and R. Polverini for the Italian collections and photographs of *R. mediterraneensis*, Celestino Gelpí and Julián Muñoz for the Spanish collections and photographs of *R. andaluciana*, Javier Gómez for helping in collecting *R. meridionalis*, Lamberto Gorit (ELG) and Giampaolo Bernardini for the Italian collections of *L. populicola*, *R. candidissima* and *R. cerea*, Gerhard Gross† for the German collection of *R. candida*, Gunnar Hensel (GH) and B. Mühlér for the German collection and photographs of *R. cerea*, Pablo Juste for the Spanish collection of *L. giennensis*, Istvan Király for the Hungarian collection of *L. stephensi*, Maciej Kozak for the Polish collection of *R. cerea*, Claude Lavoise for the photographs of *L. soehneri*, Zoltán Lukács for the Hungarian collections of *R. candida*, José Angel Martínez and Andreu Carreres for the Spanish collections and photographs of *R. monospora*, Amer Montecci (AM) for the Italian collections of *L. borzianus*, *L. populicola* and *R. mattiroloana*, Marco Morara and Maurizio Berbenni for the Italian collections and photographs of *L. soehneri* and *L. stephensi*, Pierre-Arthur Moreau (PAM) for the French collections and images of *L. borzianus* and for his help in consulting the original material of *H. monosporum* from Nice, Pedro María Pasabán and Francisco Sáinz for the Spanish collections and photographs of *R. candidissima*, *R. cerea* and *R. pila*, Miquel Angel Pérez-De-Gregorio for the Spanish collections of *R. candida* and *R. candidissima*, Tomás Pérez-Jarauta for the Spanish collections of *L. giennensis*, Vasilis Ramoutsakis for the Greek collection of *R. messapica* var. *messapicoides*, Fernando Rodríguez for his friendship and invaluable help in many Catalan collections, Ryszard Rutkowski for the Polish collection of *L. stephensi*, A. Sanz-Becerra and Natalio Redondo for the Spanish collection of *R. monospora*, Miguel Ángel Sanz for the Spanish collections of *L. soehneri*, Mario Sarasin (MS) for the Italian collections of *R. candida* and *R. cerea*, George Setkos for the Greek collections of *L. populicola*, Josep Leonard Squier for images and information of the Spanish collections of *R. messapica* and M. Tordelli and Despina Klisiari for the Cypriot collections of *R. hobartiae*. Finally the authors thank Caroline Loup, curator of the Herbier of the Université de Montpellier (SPH, MPU) for kindly allowing us to reproduce the original plates of *Zelleromyces josserandii* of G. Malençon, Olivier Gerriet, curator of the Muséum d'Histoire Naturelle de Nice (NICE), for kindly providing us information, images and original material of *Hydnangium monosporum* from Barla's Herbarium, Prof. Solomon P. Wasser for providing access to HAI Herbarium and Carles Roqué and the Unitat de Microscòpia dels Serveis Tècnics de Recerca of the Universitat de Girona (STR, UdG) for their invaluable help with SEM imaging. Last but not least, the authors are grateful to Caroline Hobart for her help and valuable advice on sequestrate material from Cyprus, particularly *R. hobartiae* and also to Angelos Papadimitriou for translating into English the Greek description of *L. populicola*.

## REFERENCES

- Adamčík S, Caboň M, Eberhardt U, et al. 2016. A molecular analysis reveals hidden species diversity within the current concept of *Russula maculata* (Russulaceae, Basidiomycota). *Phytotaxa* 270: 71–88.
- Adamčík S, Jančovičová S, Buyck B. 2017. The Russulas described by Charles Horton Peck. *Cryptogamie, Mycologie* 39: 3–108.
- Altschul SF, Madden TL, Schäffer AA, et al. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389–3402.
- Alvarado P, Moreno G, Manjón JL. 2012. Comparison between *Tuber gennadii* and *T. oligospermum* lineages reveals the existence of the new species *T. cistophilum* (Tuberaceae, Pezizales). *Mycologia* 104: 894–910.
- Astier J, Pacioni G. 1998. Le genre *Martellia* Matt. en Europe. *Documents Mycologiques* 109–110: 7–9.
- Avis PG. 2012. Ectomycorrhizal iconoclasts: the ITS rDNA diversity and nitrophilic tendencies of fetid *Russula*. *Mycologia* 104: 998–1007.
- Bahram M, Polme S, Koljalg U, et al. 2011. A single European aspen (*Populus tremula*) tree individual may potentially harbour dozens of *Cenococcum geophilum* ITS genotypes and hundreds of species of ectomycorrhizal fungi. *FEMS Microbiology Ecology* 75: 313–320.
- Bahram M, Polme S, Koljalg U, et al. 2012. Regional and local patterns of ectomycorrhizal fungal diversity and community structure along an altitudinal gradient in the Hircanian forests of northern Iran. *New Phytologist* 193: 465–473.
- Bandala VM, Montoya L, Ramos A. 2016. Two new *Lactarius* species in a subtropical cloud forest in eastern Mexico. *Mycologia* 108: 967–980.
- Barge EG, Cripps CL, Osmundson TW. 2016. Systematics of the ectomycorrhizal genus *Lactarius* in the Rocky Mountain alpine zone. *Mycologia* 108: 414–440.
- Bazzicalupo AL, Buyck B, Saar I, et al. 2017. Troubles with mycorrhizal mushroom identification where morphological differentiation lags behind barcode sequence divergence. *Taxon* 66: 791–810.
- Beaton G, Pegler DN, Young TWK. 1984. Gasteroid Basidiomycota of Victoria State, Australia. 2. Russulales. *Kew Bulletin* 39: 669–698.
- Beenken L. 2004. Les ectomycorrhizes du genre *Russula*. *Bulletin Trimestriel de la Société Mycologique de France* 120: 293–333.
- Beenken L, Saine MN, Kocyan A. 2016. *Lactarius megalopterus*, a new angiocarpous species from tropical rainforest in Central Africa, shows adaptations to endozoochorous spore dispersal. *Mycological Progress* 15: 1–10.
- Berkeley MJ. 1844. Notices of British fungi (257–322). *Annals and Magazine of Natural History*, Series 1: 340–360.
- Binder M, Hibbett DS. 2002. Higher-level phylogenetic relationships of Homobasidiomycetes (mushroom-forming fungi) inferred from four rDNA regions. *Molecular Phylogenetics & Evolution* 22: 76–90.
- Bon M. 1986. Novitaires. Validations et taxons nouveaux. *Documents Mycologiques* 65: 51–56.
- Bon M. 1988. Clé monographique des russules d'Europe. *Documents Mycologiques* 70–71: 1–120.
- Boudier E. 1906. *Icônes mycologicae ou iconographie des champignons de France, principalement Discomycètes*. Tome 1. Librairie des Sciences Naturelles Paul Klincksieck, Paris, France.
- Boudier E, Patouillard N. 1888. Note sur deux nouvelles espèces de Champignons des environs de Nice. *Journal de Botanique*, Paris 2: 445–446.
- Bougher NL. 1997. Three new sequestrate Basidiomycetes from Western Australia. *Mycotaxon* 63: 37–48.
- Bougher NL, Lebel T. 2001. Sequestrate (truffle-like) fungi of Australia and New Zealand. *Australian Systematic Botany* 14: 439–484.
- Bougher NL, Tommerup IC, Malajczuk N. 1993. Broad variation in developmental and mature basidiome morphology of the ectomycorrhizal fungus *Hydnangium sublamellatum* sp. nov. bridges morphologically based generic concepts of *Hydnangium*, *Podothyridium* and *Laccaria*. *Mycological Research* 97: 613–619.
- Brock PM, Doring H, Bidartondo MI. 2009. How to know unknown fungi: the role of a herbarium. *New Phytologist* 181: 719–724.
- Bucholtz F. 1901. Hypogaeen aus Russland. *Hedwigia* 40: 304–322.
- Bucholtz F. 1903. Zur Morphologie und Systematik der Fungi hypogaei. *Annales Mycologici* 1: 152–174, taf. IV, V.
- Buyck B. 1995. A global integrated approach on the taxonomy of Russulales. *Russulales News* 3: 2–17.
- Buyck B, Duhem B, Das K, et al. 2017. Fungal Biodiversity Profiles 21–30. *Cryptogamie, Mycologie* 38: 101–146.
- Buyck B, Hofstetter V, Eberhardt U, et al. 2008. Walking the thin line between *Russula* and *Lactarius*: the dilemma of *Russula* subsect. *Ochricompactae*. *Fungal Diversity* 28: 15–40.
- Buyck B, Hofstetter V, Verbeken A, et al. 2010. Proposal 1919: To conserve *Lactarius* nom. cons. (Basidiomycota) with a conserved type. *Taxon* 59: 295–296.
- Buyck B, Ovrebo CL. 2002. New and interesting *Russula* species from Panama. *Mycologia* 94: 888–901.
- Buyck B, Verbeken A, Eberhardt U. 2007. The genus *Lactarius* in Madagascar. *Mycological Research* 111: 787–798.
- Cabero J. 2011. Segunda cita para la península ibérica de *Gymnomyces ilicis*. *Boletín Micológico de la FAMCAL* 6: 63–68.
- Calonge FD. 2000. Validation or confirmation of some new taxa recently published. *Boletín de la Sociedad Micológica de Madrid* 25: 301–302.
- Calonge FD, Martín MP. 2000. Morphological and molecular data on the taxonomy of *Gymnomyces*, *Martellia* and *Zelleromyces* (Russulales). *Mycotaxon* 76: 9–15.
- Calonge FD, Martín MP. 2003. *Zelleromyces hispanicus*, the gasteroid phase of *Lactarius aurantiacus*. *Boletín de la Sociedad Micológica de Madrid* 27: 231–236.
- Calonge FD, Pasabán PM. 2005. *Macowanites candidus* (Russulales), nuevo para el catálogo micológico español. *Boletín de la Sociedad Micológica de Madrid* 29: 87–90.
- Calonge FD, Pegler DN. 1998. *Zelleromyces hispanicus* sp. nov. (Russulales, Elasmomycetaceae), an orange-red species possibly related to *Lactarius aurantiacus*. *Cryptogamie, Mycologie* 19: 99–105.
- Calonge FD, Vidal JM. 1999. *Gymnomyces ammophilus* Vidal & Calonge, sp. nov., encontrado en Portugal. *Boletín de la Sociedad Micológica de Madrid* 24: 65–70.
- Calonge FD, Vidal JM. 2001. *Macowanites vinaceodorus* sp. nov. (Russulales), a new gasteroid fungus from coastal dunes of Spain. *Mycotaxon* 79: 1–6.
- Campos JC, Zamora JC, Vila J. 2010. Estudio de la micobiota de las comunidades de Cistaceae en el centro de la Península Ibérica. *Boletín de la Sociedad Micológica de Madrid* 34: 257–270.
- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540–552.
- Cavara F. 1897. Contributo alla conoscenza delle Podaxineae: *Elasmomyces mattirolianus* nov gen. et sp. *Malpighia* 11: 414–428, tav. VII.
- Cavara F. 1900. *Arcangiella borziana* nov. gen. nov. sp. Nuova Imenogastera delle abetine di Vallombrosa. *Nuovo Giornale Botanico Italiano, Nuova Serie*, 7: 117–128, tav. VII.
- Cheype J-L, Campo E. 2012. *Russula rubropunctatissima* Cheype & Campo une nouvelle russule découverte en Guyane Française. *Bulletin Trimestriel de la Société Mycologique de France* 128: 127–135.
- Cléménçon H. 2004. Cytology and plectology of the Hymenomycetes. *Biblioteca Mycologica* 199. Cramer, Berlin-Stuttgart, Germany.
- Comandini O, Contu M, Rinaldi AC. 2006. An overview of *Cistus* ectomycorrhizal fungi. *Mycorrhiza* 16: 381–395.
- Corner EJH, Hawker LE. 1953. Hypogeous fungi from Malaya. *Transactions of the British Mycological Society* 36: 125–137.
- Crous PW, Wingfield MJ, Richardson DM, et al. 2016. Fungal Planet description sheets 400–468. *Persoonia* 36: 316–458.
- Das K, Dowie NJ, Li GJ, et al. 2014. Two new species of *Russula* (Russulales) from India. *Mycosphere* 5: 612–622.
- Das K, Ghosh A, Chakraborty D, et al. 2017. Fungal Biodiversity Profiles 31–40. *Cryptogamie, Mycologie* 38: 353–406.
- De Crop E, Nyuytinck J, Van de Putte K, et al. 2017. A multi-gene phylogeny of *Lactifluus* (Basidiomycota, Russulales) translated into a new infrageneric classification of the genus. *Persoonia* 38: 58–80.
- Desjardin D. 2003. A unique ballistosporic hypogeous sequestrate *Lactarius* from California. *Mycologia* 95: 148–155.
- Dodge CW. 1931. Alpova, a new genus of Rhizopogonaceae, with further notes on *Leucogaster* and *Arcangiella*. *Annals of the Missouri Botanical Garden* 18: 457–464.
- Dodge CW, Zeller SM. 1937 '1936'. *Hydnangium* and related genera. *Annals of the Missouri Botanical Garden* 23: 565–598.
- Dring DM, Pegler DN. 1978. New and noteworthy gasteroid relatives of the Agaricales from tropical Africa. *Kew Bulletin* 32: 563–569.
- Durrall DM, Gamiet S, Simard SW, et al. 2006. Effects of clearcut logging and tree species composition on the diversity and community composition of epigeous fruit bodies formed by ectomycorrhizal fungi. *Canadian Journal of Botany* 84: 966–980.
- Dutta AK, Paloi S, Pradhan P, et al. 2015. A new species of *Russula* (Russulaceae) from India based on morphological and molecular (ITS sequence) data. *Turkish Journal of Botany* 39: 850–856.
- Eberhardt U. 2002. Molecular kinship analyses of the agaricoid Russulaceae: correspondence with mycorrhizal anatomy and sporocarp features in the genus *Russula*. *Mycological Progress* 1: 201–223.
- Eberhardt U, Oberwinkler F, Verbeken A, et al. 2000. *Lactarius ectomycorrhizae* on Silver fir (*Abies alba*): morphological description, molecular characterization and taxonomic remarks. *Mycologia* 92: 860–873.

- Eberhardt U, Verbeken A. 2004. Sequestrate Lactarius species from tropical Africa: *L. angiocarpus* nov. sp. and *L. dolichocaulis* comb. nov. Mycological Research 108: 1042–1052.
- Elliott TF, Trappe JM. 2018. A worldwide nomenclature revision of sequestrate Russula species. Fungal Systematics and Evolution 1: 229–242.
- Fellner R, Bibel J. 1990. Helianthemum and some agaricales: unusual case of ectomycorrhizal symbiosis. Agriculture, Ecosystems & Environment 28: 121–125.
- Fogel R, States J. 2001. Materials for a hypogeous mycoflora of the Great Basin and adjacent cordilleras of the Western United States IV: Zelleromyces rogersonii, sp. nov. (Basidiomycota, Elasmomycetaceae). Mycotaxon 80: 321–326.
- Fraiture A, Derboven P. 2009. Deux nouvelles récoltes belges d'*Arcangeliella stephensii*, espèce tubéroïde proche des lactaires. Revue du Cercle de Mycologie de Bruxelles 9: 27–42.
- Frank JL, Barry S, Southworth D. 2006. Mammal mycophagy and dispersal of mycorrhizal inoculum in Oregon white oak woodlands. Northwest Science 80: 264–273.
- Garay-Serrano E, Bandala VM, Montoya L. 2012. Morphological and molecular identification of the ectomycorrhizal association of *Lactarius fumosibrunneus* and *Fagus grandifolia* var. *mexicana* trees in eastern Mexico. Mycorrhiza 22: 583–588.
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for Basidiomycetes — Application to the identification of mycorrhizae and rusts. Molecular Ecology 2: 113–118.
- Ge ZW, Smith ME, Zhang QY, et al. 2012. Two species of the Asian endemic genus *Keteleeria* form ectomycorrhizas with diverse fungal symbionts in southwestern China. Mycorrhiza 22: 403–408.
- Gelardi M, Vizzini A, Ercol E, et al. 2015. Circumscription and taxonomic arrangement of *Nigroboletus roseonigrescens* gen. et sp. nov., a new member of Boletaceae from tropical South-Eastern China. PLoS One 10: e0134295.
- Ghosh A, Das K, Bhatt RP. 2017. *Russula sarnarii* sp. nov. (Russulaceae, Basidiomycota) from Indian Himalaya. Current Research in Environmental and Applied Mycology 7: 64–72.
- Gladish S, Frank J, Southworth D. 2010. The serpentine syndrome below ground: ectomycorrhizas and hypogeous fungi associated with conifers. Canadian Journal of Forest Research 40: 1671–1679.
- Gori L. 2005. Funghi Ipogeici della Lucchesia, di altre province italiane e dall'estero. Maria Pacini Fazzi Editore, Lucca, Italy.
- Gross G. 1969. Ein Saarländischer fund von Elasmomyces mattirolianus Cav. Zeitschrift für Pilzkunde, N.F., 34: 27–32.
- Hawker LE. 1952. Hypogeous fungi. II. A new variety of *Hydnangium carneum* Wallr. from North Wales. III. Three new British records: *Gautieria morchellaeformis* Vitt., *Hymenogaster hessei* Soehner, and *Elaphomyces aculeatus* Vitt. Transactions of the British Mycological Society 35: 279–284.
- Hawker LE. 1954. British hypogeous fungi. Philosophical Transactions of the Royal Society London, Series B, 237: 429–546.
- Hawksworth DL, Kirk PM, Sutton BC, et al. 1995. Ainsworth & Bisby's dictionary of the fungi. 8th edition. CAB International, Oxon, U.K.
- Heilmann-Clausen J, Verbeken A, Vesterholz J. 1998. The genus *Lactarius*. Fungi of Northern Europe. Vol. 2. The Danish Mycological Society, Copenhagen, Denmark.
- Heim R. 1938. Les Lactario-russulés du domaine oriental de Madagascar. Essai sur la classification et la phylogénie des Astérosporales. Prodrome à une flore mycologique de Madagascar et dépendances. I. Laboratoire de Cryptogamie du Muséum National d'Histoire Naturelle de Paris, Paris, France.
- Heim R. 1959. Une espèce nouvelle de Gastrolactarié en Thaïlande. Revue de Mycologie, Paris 24: 93–102.
- Heim R, Font-Quer P, Codina J. 1934. Fungi Iberici. Observations sur la flore mycologique Catalane. Treballs del Museu de Ciències Naturals de Barcelona, Sèr. Bot. 15: 1–146, 4 pl.
- Hesler LR, Smith AH. 1979. North American species of *Lactarius*. The University of Michigan Press, Ann Arbor, Michigan, USA.
- Hibbett DS, Pine EM, Langer E, et al. 1997. Evolution of gilled mushrooms and puffballs inferred from ribosomal DNA sequences. Proceedings of the National Academy of Sciences 94: 12002–12006.
- Hibbett DS, Tsuneda A, Murakami S. 1994. The secotoid form of *Lentinus tigrinus*: genetics and development of a fungal morphological innovation. American Journal of Botany 81: 466–478.
- Higgins D, Thompson J, Gibson T, et al. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Research 22: 4673–4680.
- Horton BM. 2011. Eucalypt decline and ectomycorrhizal fungal community ecology of *Eucalyptus delegatensis* forest, Tasmania, Australia. PhD thesis, University of Tasmania.
- Hosaka K, Bates ST, Beever RE, et al. 2006. Molecular phylogenetics of the gomphoid-phalloid fungi with an establishment of the new subclass Phallomycetidae and two new orders. Mycologia 98: 949–959.
- Izzo AD, Meyer M, Trappe JM, et al. 2005. Hypogeous ectomycorrhizal fungal species on roots and in small mammal diet in a mixed-conifer forest. Forest Science 51: 243–254.
- Josserand M. 1983. La description des champignons supérieurs. 2e éd. Editions Lechevalier, Paris, France.
- Kalchbrenner C. 1876. Zwei neue Pilzgattungen. Hedwigia 15: 115–116.
- Kalchbrenner C. 1882. Fungi Macowaniani. Grevillea 10: 104–109.
- Kleine CS, McClean T, Miller SL. 2013. Genetic divergence among disjunct populations of three *Russula* spp. from Africa and Madagascar. Mycologia 105: 80–89.
- Kong A, Cifuentes J, Estrada-Torres A, et al. 2015. Russulaceae associated with mycoheterotroph *Monotropa uniflora* (Ericaceae) in Tlaxcala, Mexico: a phylogenetic approach. Cryptogamie, Mycologie 36: 479–512.
- Konstantinidis G. 2009. Μανύρια, φωτογραφικός οδηγός μανιταροσυλλέκτη (Mushrooms, a photographic collector's guide). Published by the author, Athens, Greece.
- Kornerup A, Wanscher JH. 1978. Methuen handbook of colour. 3 ed. Eyre Methuen, London, UK.
- Lago M, Bouger NL, Castro ML. 2000. Morphological variability and implications for definition of taxa in the Descolea-Setchellio-gaster-Descomyces complex. Mycotaxon 18: 37–57.
- Larsson E, Larsson KE. 2003. Phylogenetic relationships of russuloid basidiomycetes with emphasis on aphylophoraean taxa. Mycologia 95: 1037–1065.
- Lavorato C. 1991. Chiave analitica e note bibliografiche della micoflora del cisto. Bollettino dell'Associazione Micologica ed Ecologica Romana 24: 16–45.
- Lebel T. 1998. Taxonomic revision of the sequestrate relatives of *Russula* from Australia and New Zealand. PhD Thesis, Oregon State University, USA.
- Lebel T. 2001. A new species of Zelleromyces (Russulales) from Australia. Australasian Mycologist 20: 4–8.
- Lebel T. 2002. Sequestrate Russulales of New Zealand: Gymnomycetes and Macowanites. New Zealand Journal of Botany 40: 489–509.
- Lebel T. 2003a. Australian sequestrate (truffle-like) fungi. XIII. *Cystangium* (Russulales, Basidiomycota). Australian Systematic Botany 16: 371–400.
- Lebel T. 2003b. Australian sequestrate (truffle-like) fungi. XIV. *Gymnomyces* (Russulales, Basidiomycota). Australian Systematic Botany 16: 401–426.
- Lebel T. 2017. Nomenclatural changes and corrections for some previously described Australasian truffle-like fungi (Basidiomycetes). Muelleria 36: 8–14.
- Lebel T, Castellano MA. 2002. Type studies of sequestrate Russulales II. Australian and New Zealand species related to *Russula*. Mycologia 94: 327–354.
- Lebel T, Dunk CW, May TW. 2013. Rediscovery of *Multifurca stenophylla* (Berk.) Lebel, C.W. Dunk & T.W. May comb. nov. (Russulaceae) from Australia. Mycological Progress 12: 497–504.
- Lebel T, Tonkin JE. 2007. Australasian species of Macowanites are sequestrate species of *Russula* (Russulaceae, Basidiomycota). Australian Systematic Botany 20: 355–381.
- Lebel T, Trappe JM. 2000. Type studies of sequestrate Russulales. I. Generic type species. Mycologia 92: 1188–1205.
- Lee H, Park MS, Jung PE, et al. 2017. Re-evaluation of the taxonomy and diversity of *Russula* section Foetentinae (Russulales, Basidiomycota) in Korea. Mycoscience 58: 351–360.
- Lejeune C. 2005. *Russula straminea* ou *Russula globispora*? Annales Confederationis Europeae Mycologiae Mediterranensis 2003: 61–72.
- Leonardi M, Comandini O, Rinaldi AC. 2016. Peering into the Mediterranean black box: *Lactifluus rugatus* ectomycorrhizas on *Cistus*. IMA Fungus 7: 275–284.
- Li G-J, Zhang C-L, Zhao R-L, et al. 2018. Two new species of *Russula* from Northeast China. Mycosphere 9: 431–443.
- Li G-J, Zhao D, Li S-F, et al. 2015. *Russula chuii* and *R. pseudopectinatoides*, two new species from southwestern China supported by morphological and molecular evidence. Mycological Progress 14: 33.
- Li G-J, Zhao Q, Zhao D, et al. 2013. *Russula atroeruginea* and *R. sichuanensis* spp. nov. from southwest China. Mycotaxon 124: 173–188.
- Liu JK, Hyde KD, Jones EBG, et al. 2015. Fungal diversity notes 1–110: Taxonomic and phylogenetic contributions to fungal species. Fungal Diversity 72: 1197.
- Llistosella J. 1998. Algunes espècies del gènere *Russula* de Catalunya i les Illes Balears. 2a Contribució. Revista Catalana de Micología 21: 75–92.
- Llistosella J, Vidal JM. 1995. Due nuove specie di Russulales gasteroidi della regione mediterranea. Rivista di Micologia AMB 38: 149–162.
- Loizides M. 2016. Macromycetes within Cistaceae-dominated ecosystems in Cyprus. Mycotaxon 131: 255–256.

- Loizides M, Kyriakou T. 2011. Fungi of the Cistus maquis. *Field Mycology* 12: 14–22.
- Looney BP, Ryberg M, Hampe F, et al. 2016. Into and out of the tropics: global diversification patterns in a hyper-diverse clade of ectomycorrhizal fungi. *Molecular Ecology* 25: 630–647.
- Lutzoni F, Kauff F, Cox CJ, et al. 2004. Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *American Journal of Botany* 91: 1446–1480.
- Mader K, Mader A. 1992. Ein Beitrag zur Kenntnis der sternsporigen Hypogäen. *Österreichische Zeitschrift für Pilzkunde* 1: 3–10.
- Malençon G. 1931. La série des Astérosporés. Recueil de travaux dédiés à L. Mangin: 377–396, 1 pl.
- Malençon G. 1975. Champignons hypogés du Nord de l'Afrique II. Basidiomycetes. *Revue de Mycologie*, Paris 39: 279–306.
- Malloch D, Thorn RG. 2011. The occurrence of ectomycorrhizae in some species of Cistaceae in North America. *Canadian Journal of Botany* 63: 872–875.
- Martín MP, Höglberg N, Llistosella J. 1999. Macowanites messapicoides, a hypogeous relative to Russula messapica. *Mycological Research* 103: 203–208.
- Martín MP, Moreno G. 2001. Molecular data confirm Setchellio-gaster tenuipes and S. rheophyllus as Cortinariales. *Mycotaxon* 78: 257–263.
- Martín MP, Rocabruna A. 1999. The taxonomic boundaries between Naucoria rheophylla and Setchellio-gaster tenuipes based on morphology and molecular data. *Mycotaxon* 71: 141–148.
- Massee G. 1898. Fungi Exotici, I. *Bulletin of Miscellaneous Information, Royal Botanic Gardens, Kew* 1898: 113–136.
- Matheny PB. 2005. Improving phylogenetic inference of mushrooms with RPB1 and RPB2 nucleotide sequences (Inocybe; Agaricales). *Molecular Phylogenetics and Evolution* 35: 1–20.
- Matheny PB, Wang Z, Binder M, et al. 2007. Contributions of rpb2 and tef1 to the phylogeny of mushrooms and allies (Basidiomycota, Fungi). *Molecular Phylogenetics and Evolution* 43: 430–451.
- Mattioli O. 1900. Gli ipogei di Sardegna e di Sicilia. *Malpighia* 14: 39–110, tav. I.
- Melera S, Ostellari C, Roemer N, et al. 2017. Analysis of morphological, ecological and molecular characters of Russula pectinatoides Peck and Russula praetervisa Sarnari, with a description of the new taxon Russula recondita Melera & Ostellari. *Mycological Progress* 16: 117–134.
- Miller SL, Buyck B. 2002. Molecular phylogeny of the genus Russula in Europe with a comparison of modern infrageneric classifications. *Mycological Research* 106: 259–276.
- Miller SL, Larsson E, Larsson KH, et al. 2006. Perspectives in the new Russulales. *Mycologia* 98: 960–970.
- Miller SL, Lebel T. 1999. Hypogeous fungi from the South-eastern United States. II. The genus Zelleromyces. *Mycotaxon* 72: 15–25.
- Miller SL, McClean TM, Walker JF, et al. 2001. A molecular phylogeny of the Russulales including agaricoid, gasteroid and pleurotoid taxa. *Mycologia* 93: 344–354.
- Montecchi A, Lazzari G. 1986. Elasmomyces mattirolianus Cav. (= Macowanites krikowensis (Buch.) Sing. & Smith), forme di transizione tra Gasteromiceti e Russulaceae. *Bollettino del Gruppo Micologico G. Bresadola Trento* 29: 157–163.
- Montecchi A, Sarasini M. 2000. Funghi Ipogei d'Europa. *Associazione Micologica Bresadola, Fondazione Centro Studi Micologici, Trento-Vicenza*, Italy.
- Moreno G, Galán R, Montecchi A. 1991. Hypogeous fungi from peninsular Spain. II. *Mycotaxon* 42: 201–238.
- Moreno-Arroyo B, Gómez J, Calonge FD. 1998a. Zelleromyces giennensis sp. nov. (Russulales), a gasteroid fungus from the South of Spain. *Cryptogamie, Mycologie* 19: 107–111.
- Moreno-Arroyo B, Gómez J, Calonge FD. 1998b. Zelleromyces meridionalis (Russulales, Elasmomycetaceae), a new species from Spain. *Mycotaxon* 69: 467–471.
- Moreno-Arroyo B, Gómez J, Calonge FD. 1999. Gymnomyces dominguezii sp. nov. from Spain. *Mycological Research* 103: 215–218.
- Moreno-Arroyo B, Llistosella J, Romero De La Osa L. 2002. Gymnomyces sublevisporus (Russulales), una nueva especie de la región mediterránea. *Revista Catalana de Micología* 24: 179–186.
- Moser M, Binyamin N, Avizohar-Hershenson Z. 1977. New and noteworthy Russulales from Israel. *Transactions of the British Mycological Society* 68: 371–377.
- Moser M, Jülich W, Furrer-Ziegas C. 1994. IV Macowanites 1. In: Moser M, Jülich W (eds), *Farbatlas der Basidiomyceten* 12. Gustav Fischer Verlag, Stuttgart, Germany.
- Nuytinck J, Verbeken A, Delarue S, et al. 2003. Systematics of European sequestrate lactarioid Russulaceae with spiny spore ornamentation. *Belgian Journal of Botany* 136: 145–153.
- Nuytinck J, Verbeken A, Miller SL. 2007. Worldwide phylogeny of Lactarius section Deliciosi inferred from ITS and glyceraldehyde-3-phosphate dehydrogenase gene sequences. *Mycologia* 99: 820–832.
- Nygren CM, Edqvist J, Elfstrand M, et al. 2007. Detection of extracellular protease activity in different species and genera of ectomycorrhizal fungi. *Mycorrhiza* 17: 241–248.
- Nylander JAA. 2004. MrModeltest v2. Program distributed by the autor. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.
- Oberwinkler F. 1977. Das neue System der Basidiomyceten. In: Frey W, Hurka H, Oberwinkler F (eds), *Beiträge zur Biologie der Niederen Pflanzen. Systematik, Stammesgeschichte, Ökologie*: 59–105. Gustav Fischer Verlag, Stuttgart, Germany.
- Osmundson TW, Robert VA, Schoch CL, et al. 2013. Filling gaps in biodiversity knowledge for macrofungi: contributions and assessment of an herbarium collection DNA barcode sequencing project. *PLoS ONE* 8: e62419.
- Park MS, Fong JJ, Lee H, et al. 2013. Delimitation of Russula subgenus Amoenula in Korea using three molecular markers. *Mycobiology* 41: 191–201.
- Patouillard N. 1910. Note sur trois espèces d'Hydnangium de la flore du Jura. *Bulletin Trimestriel de la Société Mycologique de France* 26: 199–204.
- Patouillard N. 1914. Contribution à la flore mycologique du Jura. *Bulletin Trimestriel de la Société Mycologique de France* 30: 347–354.
- Paz A, Vidal JM, Lavoise C, et al. 2016. Revisión taxonómica del género Octaviania (Boletales) en Europa. *Boletín Micológico de la FAMCAL* 11: 101–138.
- Peck CH. 1897. Report of the State Botanist (1896). *Annual Report on the New York State Museum of Natural History* 50: 77–159.
- Pegler DN, Spooner BM, Young TWK. 1993. British truffles. A revision of British hypogeous fungi. Royal Botanic Gardens, Kew, UK.
- Pegler DN, Young TWK. 1979. The gasteroid Russulales. *Transactions of the British Mycological Society* 72: 353–388, 134 fig.
- Peintner U, Bouger NL, Castellano MA, et al. 2001. Multiple origins of sequestrate fungi related to Cortinarius (Cortinariaceae). *American Journal of Botany* 88: 2168–2179.
- Peter M, Buechler U, Ayer F, et al. 2001. Ectomycorrhizas and molecular phylogeny of the hypogeous russuloid fungus Arcangelia borziana. *Mycological Research* 105: 1231–1238.
- Quélet L. 1875. Les champignons du Jura et des Vosges. IIIe Partie. Mémoires de la Société d'Émulation de Montbéliard 5: 429–556, pl. I–IV.
- Quélet L. 1886. Enchiridion Fungorum in Europa media et praesertim in Gallia Vigentium. Lutetiae, Doin, France.
- Rehner SA, Buckley E. 2005. A Beauveria phylogeny inferred from nuclear ITS and EF-1 $\alpha$  sequences: evidence for cryptic diversification and links to Cordyceps teleomorphs. *Mycologia* 97: 84–98.
- Romagnesi H. 1944. La cystide chez les Agaricacées. *Supplément à la Revue de Mycologie*, Paris 9: 4–21.
- Romagnesi H. 1985. Les Russules d'Europe et d'Afrique du Nord. Reprint of 1967 ed. by Cramer, Vaduz, Germany.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Roy M, Schimann H, Braga-Neto R, et al. 2016. Diversity and distribution of ectomycorrhizal fungi from Amazonian lowland white-sand forests in Brazil and French Guiana. *Biotropica* 48: 90–100.
- Saba M, Khalid AN. 2015. Russula sichuanensis and its ectomycorrhizae from Himalayan moist temperate forests of Pakistan. *Mycotaxon* 130: 629–639.
- Saccardo PA, Saccardo D. 1905. *Hymenomycetace-Laboulbeniomycetace*. Supplementum Universale, Pars VI. In: Saccardo PA, *Sylloge Fungorum* 17. Padova, Italy.
- Sang X-Y, Li X-D, Wang Y-W, et al. 2016. Four new sequestrate species of Russulaceae found in China. *Phytotaxa* 289: 101–117.
- Sarnari M. 1998. Monografia illustrata del genere Russula in Europa. Tomo Primo. A.M.B. Fondazione Centro Studi Micologici, Trento, Italy.
- Sarnari M. 2005. Monografia illustrata del genere Russula in Europa. Tomo Secondo. A.M.B. Fondazione Centro Studi Micologici, Trento, Italy.
- Schoch CL, Seifert KA, Huhndorf S, et al. 2012. Fungal Barcoding Consortium. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences, USA* 109: 6241–6246.
- Sene S, Avril R, Chaintreuil C, et al. 2015. Ectomycorrhizal fungal communities of *Coccocloba uvifera* (L.) L. mature trees and seedlings in the neotropical coastal forests of Guadeloupe (Lesser Antilles). *Mycorrhiza* 25: 547–559.
- Shimono Y, Kato M, Takamatsu S. 2004. Molecular phylogeny of Russulaceae (Basidiomycetes, Russulales) inferred from the nucleotide sequences of nuclear large subunit rDNA. *Mycoscience* 45: 303–316.
- Singer R. 1986. The Agaricales in modern taxonomy. Fourth edition. Koeltz Scientific Books, Koenigstein, Germany.
- Singer R, Smith AH. 1960. Studies on secoziaceous fungi. IX. The Astrogastraceous series. *Memoirs of the Torrey Botanical Club* 21: 1–212.

- Smith AH. 1962. Notes on Astrogastraceous fungi. *Mycologia* 54: 626–639.
- Smith AH. 1963. New Astrogastraceous fungi from the Pacific northwest. *Mycologia* 55: 421–441.
- Smith ME, Henkel TW, Aime CM, et al. 2011. Ectomycorrhizal fungal diversity and community structure on three co-occurring leguminous canopy tree species in a Neotropical rainforest. *New Phytologist* 192: 699–712.
- Smith ME, Henkel TW, Williams GC, et al. 2017. Investigating niche partitioning of ectomycorrhizal fungi in specialized rooting zones of the monodominant leguminous tree *Dicymbe corymbosa*. *New Phytologist* 215: 443–453.
- Smith ME, Trappe JM, Rizzo DM, et al. 2006. *Gymnomyces xerophilus* sp. nov. (sequestrate Russulaceae), an ectomycorrhizal associate of *Quercus* in California. *Mycological Research* 110: 575–582.
- Soehner E. 1924. Prodromus der Fungi hypogaei Bavariae. Kryptogamische Forschungen, Bayerischen Botanischen Gesellschaft, München 1: 390–398.
- Soehner E. 1941. Deutsche Hydnangiaceae. *Zeitschrift für Pilzkunde*, N.F. 20: 108–119.
- Song J, Chen J-J, Wang M, et al. 2016. Phylogeny and biogeography of the remarkable genus *Bondarzewia* (Basidiomycota, Russulales). *Scientific Reports* 6: 34568.
- Southworth D. 2016. Distribution and abundance of rare sequestrate fungi in Southwestern Oregon. FY10-15 ISSSSP Project Final Report. Southern Oregon University, Ashland.
- Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690.
- Stubbe D, Nuytinck J, Verbeken A. 2010. Critical assessment of the *Lactarius gerardii* species complex (Russulales). *Fungal Biology* 114: 271–283.
- Stubbe D, Verbeken A. 2012. *Lactarius* subg. *Plinthogalus*: the European taxa and American varieties of *L. lignyotus* re-evaluated. *Mycologia* 104: 1490–1501.
- Suz LM, Barsoum N, Benham S, et al. 2014. Environmental drivers of ectomycorrhizal communities in Europe's temperate oak forests. *Molecular Ecology* 23: 5628–5644.
- Tamura K, Peterson D, Peterson N, et al. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28: 2731–2739.
- Tao K, Chang M-C, Liu B. 1993. New species and new record of hypogeous fungi from China IV. *Acta Mycologica Sinica* 12: 103–106.
- Tedersoo L, Hallenberg N, Larsson KH, et al. 2003. Fine scale distribution of ectomycorrhizal fungi and roots across substrate layers including coarse woody debris in a mixed forest. *New Phytologist* 159: 153–165.
- Thiers HD. 1984. The secotoid syndrome. *Mycologia* 76: 1–8.
- Torrejón M. 2007. Contribución al estudio de los hongos del parque natural de la Serra Calderona y su área de influencia: Castelló-València (España). I. Jarales (Cistion). *Revista Catalana de Micología* 29: 17–28.
- Torrejón M. 2009. A contribution to the study of fungi associated with *Cistus* spp. in the Sierra Calderona Nature Reserve, Castellón-Valencia, Spain. II. *Mycologia Balcanica* 6: 111–122.
- Trappe JM, Claridge AW. 2003. Australasian sequestrate (truffle-like) fungi. XV. New species from tree line in the Australian Alps. *Australasian Mycologist* 22: 27–38.
- Trappe JM, Claridge AW. 2005. Hypogeous fungi: evolution of reproductive and dispersal strategies through interactions with animals and mycorrhizal plants. In: Dighton J, White JF, Oudemans P (eds), *The fungal community: its organization and role in the ecosystem*: 613–623. 3rd ed. Taylor & Francis, Boca Raton, USA.
- Trappe JM, Lebel T, Castellano MA. 2002. Nomenclatural revisions in the sequestrate russuloid genera. *Mycotaxon* 81: 195–214.
- Trendel JM, Hampe F, Verbeken A. 2017. *Russula vinosoflavescens* sp. nov., from deciduous forests of Northern Alsace, France. *Mycotaxon* 132: 707–721.
- Trieveriler-Pereira L, Smith ME, Trappe JM, et al. 2015. Sequestrate fungi from Patagonian Nothofagus forests: *Cystangium* (Russulaceae, Basidiomycota). *Mycologia* 107: 90–103.
- Trimbach J. 1996. Barla et la mycologie. *Annales du Muséum d'Histoire Naturelle de Nice* 11: 179–256.
- Tulasne L-R, Tulasne C. 1843. Champignons hypogés de la famille des Lycopéracées, observés dans les environs de Paris et les départements de la Vienne et d'Indre-et-Loire. *Annales des Sciences Naturelles, Botanique*, 2e Série, 19: 373–381, pl. 17.
- Turland NJ, Wiersema JH, Barrie FR, et al. (eds). 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Glashütten, Koeltz Botanical Books.
- Turner BL, Nesom GL. 2000. Use of variety and subspecies and new varietal combinations for *Styrax platanifolius* (Styraceae). *SIDA, Contributions to Botany* 19: 257–262.
- Vauras J, Ruotsalainen J, Liimatainen K. 2016. *Russula suecica*, a new red species from Northern Fennoscandia. *Karstenia* 56: 5–12.
- Velenovský J. 1947. Novitates mycologicae novissimae. *Opera Botanica Čechica* 4: 1–167, 2 pl.
- Verbeken A, Hampe F, Wissitrasameewong K, et al. 2014a. A new angiocarpous *Lactarius* species from Thailand. *Phytotaxa* 181: 163–170.
- Verbeken A, Stubbe D, Van De Putte K, et al. 2014b. Tales of the unexpected: angiocarpous representatives of the Russulaceae in tropical South East Asia. *Persoonia* 32: 13–24.
- Verbeken A, Walley R. 2010. *Lactarius*. Fungus Flora of Tropical Africa. Nationale Plantentuin België, Meise.
- Vidal JM. 1991a. *Octaviania pila* (Pat.) Svrček. *Bolets de Catalunya* 10: lám. 486.
- Vidal JM. 1991b. Contribución al conocimiento de la flora micológica del Baix Empordà y zonas limítrofes (Cataluña). IV. Hongos hipogeos (Zygomycota, Ascomycota y Basidiomycota). *Butletí de la Societat Catalana de Micología* 14–15: 143–194.
- Vidal JM. 2004a. *Arcangeliella borziana* and *A. stephensii*, two gasteroid fungi often mistaken. A taxonomic revision of *Lactarius*-related sequestrate fungi. *Revista Catalana de Micología* 26: 59–82.
- Vidal JM. 2004b. *Macowanites candidus*, a new combination for *Hydnangium candidum* Tul. et C. Tul. *Revista Catalana de Micología* 26: 83–96.
- Vidal JM. 2004c. The genus *Stephanospora* Pat., two new combinations. *Revista Catalana de Micología* 26: 97–111.
- Vidal JM, Calonge FD, Martín MP. 2002. *Macowanites ammophilus* (Russulales), a new combination based on new evidences. *Revista Catalana de Micología* 24: 69–74.
- Vila J, Llimona X. 1999. Els fongs del Parc Natural del Cap de Creus i Serra Verdera (Girona). II. Aproximació al component fúngic del Cistion. *Revista Catalana de Micología* 22: 95–114.
- Vila J, Llimona X. 2002. Noves dades sobre el component fúngic de les comunitats de Cistus de Catalunya. *Revista Catalana de Micología* 24: 74–121.
- Vila J, Llimona X. 2006. Noves dades sobre el component fúngic de les comunitats de Cistus de Catalunya. II. *Revista Catalana de Micología* 28: 167–207.
- Vila J, Llimona X. 2009. Noves dades sobre el component fúngic de les comunitats de Cistus de Catalunya. III. Addicions, correccions i claus d'identificació. *Revista Catalana de Micología* 31: 103–137.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246.
- Wang Z, Binder M, Dai Y-C, et al. 2004. Phylogenetic relationships of *Sparassis* inferred from nuclear and mitochondrial ribosomal DNA and RNA polymerase sequences. *Mycologia* 96: 1015–1029.
- Whitbeck KL. 2003. Systematics of Pacific Northwestern species of the genus *Gymnomyces* inferred from nuclear ribosomal DNA internal transcribed spacer sequences. Masters thesis, Oregon State University, Corvallis.
- White TJ, Bruns TD, Lee SB, et al. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, et al. (eds), *PCR protocols: a guide to methods and applications*: 315–322. Academic Press Inc, New York-San Diego, USA.
- Wilson AW, Binder M, Hibbett DS. 2011. Effects of gasteroid fruiting body morphology on diversification rates in three independent clades of fungi estimated using binary state speciation and extinction analysis. *Evolution* 65: 1305–1322.
- Wissitrasameewong K, Nuytinck J, Hyde KD, et al. 2014. *Lactarius* subgenus *Russularia* (Russulaceae) in Southeast Asia: 1. Species with very distant gills. *Phytotaxa* 158: 23–42.
- Xie X-D, Liu P-G, Yu F-Q. 2010. Species diversity of russuloid mycorrhiza-forming fungi on *Pinus yunnanensis* seedlings and the mycorrhizal morphology. *Acta Botanica Yunnanica* 32: 211–220.
- Zambonelli A, Donnini D, Rana GL, et al. 2014. Hypogeous fungi in Mediterranean maquis, arid and semi-arid forests. *Plant Biosystems* 148: 392–401.
- Zeller SM, Dodge CW. 1919. *Arcangeliella*, *Gymnomyces* and *Macowanites* in North America. *Annals of the Missouri Botanical Garden* 6: 49–59.
- Zeller SM, Dodge CW. 1935. New species of Hydnangiaceae. *Annals of the Missouri Botanical Garden* 25: 365–373.
- Zeller SM, Dodge CW. 1937 '1936'. *Elasmomyces*, *Arcangeliella* and *Macowanites*. *Annals of the Missouri Botanical Garden* 23: 599–638.
- Zhang B-C, Yu Y-N. 1990. Two new species of gasteroid Russulales from China, with notes on taxonomy of *Gymnomyces*, *Martellia* and *Zelleromyces*. *Mycological Research* 94: 457–462.
- Zhao Q, Li Y-K, Zhu X-T, et al. 2015. *Russula nigrovirens* sp. nov. (Russulaceae) from southwestern China. *Phytotaxa* 236: 249–256.