

# Inventory of the ichthyofauna from the Jordão and Areia river basins (Iguaçu drainage, Brazil) reveals greater sharing of species than thought

Augusto Frota<sup>1,2\*</sup>, Eduardo Vieira Real Gonçalves<sup>2</sup>, Gabriel de Carvalho Deprá<sup>1,2</sup> and Weferson Júnio da Graça<sup>2,3</sup>

1 Universidade Estadual de Maringá. Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais. Av. Colombo, 5790, 87020-900 Maringá, PR, Brazil

2 Coleção Ictiológica do Núcleo de Pesquisas em Limnologia Ictiologia e Aquicultura (Nupélia), Universidade Estadual de Maringá. Av. Colombo, 5790, 87020-900 Maringá, PR, Brazil

3 Universidade Estadual de Maringá, Centro de Ciências Biológicas, Departamento de Biologia, Núcleo de Pesquisas em Limnologia Ictiologia e Aquicultura (Nupélia), Universidade Estadual de Maringá. Av. Colombo, 5790, 87020-900 Maringá, PR, Brazil

\* Corresponding author. E-mail: [frota.augusto@gmail.com](mailto:frota.augusto@gmail.com)

**Abstract:** Recent sampling on fish from the headwaters of the Jordão and Areia rivers, combined with collection databases and specialized literature reports, yielded the first species inventory. Both basins reflect the high endemism rate associated with the Iguaçu River. However, four species previously thought to be restricted to the Jordão River basin are now known to also occur in the basin of the Areia River: *Astyanax jordanensis* Vera Alcaraz, Pavanelli & Bertaco, 2009; *Characidium travassosi* Melo, Buckup & Oyakawa, 2016; *Jenynsia diphyes* Lucinda, Ghedotti & Graça, 2006; and *Trichomycterus igobi* Wosiacki & de Pinna, 2008. This suggests an ancestral connection between the tributaries of the Jordão and Areia rivers, with putative drainage rearrangements resulting from the uplift of the Serra da Esperança. The endemic fauna of this region is presently threatened by invasive species, the construction of dams, and other human activities.

**Key words:** drainage rearrangement; endemism; vicariance; river capture; Serra da Esperança; Paraná state

## INTRODUCTION

South America has a diverse freshwater fish fauna, with 20 orders, 69 families, 739 genera and 5,160 valid species (Reis et al. 2016). No less than 3,300 of these occur in Brazilian territory (Froese and Pauly 2016), but distributional data for them are scarce and for some species limited to the original descriptions. Biogeographical knowledge of freshwater fishes of Paraná state shows a clear bias toward economic

interests. Ichthyological inventories are mostly the result of ecological surveys on impounded waterbodies by hydroelectric companies that are obliged to fund such studies. Thus, large headwater areas with lesser hydroelectric profitability but with an expected rate of high endemism remain under sampled, although already threatened by intense human occupation (Baumgartner et al. 2012). There is the risk of losing species before becoming well known.

About 75% of the species of fishes in the Iguaçu basin are endemic (Julio Jr. et al. 1997; Zawadzki et al. 1999). Among the species inventories available (Severi and Cordeiro 1994; Garavello et al. 1997; Ingenito et al. 2004; Bifi et al. 2006; Baumgartner et al. 2006; Abilhoa et al. 2008; Baumgartner et al. 2012), the majority focused on areas influenced by dams built on the main channel of the Iguaçu River. Thus the biogeography of several of the river's subbasins remain poorly understood, and the existing inventories are insufficient. The Jordão and Areia rivers are right-bank tributaries of the lower stretch of the Iguaçu River. The headwaters of these two tributaries are adjacent to one another for several kilometers (Figure 1). Endemic species of fishes have been reported from both rivers, but mostly from the basin of the Jordão River (Lucinda et al. 2006; Vera Alcaraz et al. 2009; Baumgartner et al. 2012).

There are currently no species lists available for either of these two basins. Our inventory helps fill the knowledge gap in the biological and ecological understanding of fish species in the Neotropics (Vari and Malabarba 1998) and may aid in the formulation of conservation programs. As part of a project designed specifically to understand the biogeography of the Serra

da Esperança, we sampled several localities within the two basins. It became clear to us that the Jordão and Areia river basins have quite similar species compositions and share species that are possibly absent from all other subbasins of the Iguaçu drainage. Included are species that were thought to be restricted to the Jordão River basin but are actually also present in the Areia River basin. Thus, the objective of this paper is to provide a species inventory that accounts for both systems.

## MATERIALS AND METHODS

### Study area

The Jordão and Areia rivers are important right-bank tributaries of the lower stretch of the Iguaçu River in Paraná state, Brazil (Figure 1). The main characteristics of these river basins are presented in Table 1.

### Data collection

The specimens were collected under a permanent license to collect zoological material, number 14028-1, granted 10 December 2008 by the Ministério do Meio Ambiente (MMA), Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA), Instituto Chico Mendes de Conservação da Biodiversidade (ICMbio), and Sistema de Autorização e Informação em Biodiversidade (SISBIO) to Weferson J. da Graça.


Three collection trips were carried out: the first, in January 2014, with nine sampling sites; the second, in October 2014, with 13 sampling sites; and the third, in October 2015, with eight sampling sites, totalling 30 different sampling sites. Of these, 14 are located in the Jordão River basin and 16 in the Areia River basin (Figures 1; Appendix Figures A1 and A2; Table A1). A 100-m stretch was defined at each site,

georeferenced (Appendix Table A1), and sampled by electrofishing. This technique employs two electrified dip nets, producing an electric discharge between 200 and 400 V and current of 2 A, that stun the fish temporarily (Lobón-Cervia 1991). The specimens were anesthetized in benzocaine hydrochloride and fixed in 10% formalin in the field. After a few days, the fishes were transferred to 70% ethanol in the laboratory and deposited in the Coleção Ictiológica do Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (Nupélia). Species determinations were accomplished by comparison of the specimens with original descriptions or with type specimens of species reported from the lower portion of the Iguaçu River basin by Baumgartner et al. (2012).

In addition to those species that we collected for this study, other species have been reported for the area in the literature. The literature includes recent species descriptions (Casciotta et al. 2004; Wosiacki and Garavello 2004; Garavello 2005; Pereira 2005; Lucinda et al. 2006; Garavello and Shibatta 2007; Casciotta and Almirón 2008; Ingenito et al. 2008; Lucinda 2008; Wosiacki and de Pinna 2008a, 2008b; Pavanelli and Bifi 2009; Bifi et al. 2009; Vera Alcaraz et al. 2009; Garavello and Sampaio 2010; Řičan et al. 2011; Garavello et al. 2012) as well as ecological or ecomorphological studies and inventories covering the study area (Garavello et al. 1997; Lucinda 2005; Loureiro-Crippa and Hahn 2006; Kantec et al. 2007; Hahn and Fugi 2008; Pavanelli and Oliveira 2009; Garavello et al. 2012; Baumgartner et al. 2012; Mise et al. 2013). Species not captured by us but mentioned in previous studies were included in the list, as were species recorded in the Species Link on-line database

**Table 1.** Summary of the characteristics of the study area.

Character	Jordão River basin (Figure 1)	Areia River basin (Figure 1)
Extent	185 km (Maack 2012)	90 km (Maack 2012)
Area	4,780 km <sup>2</sup> (Júlio Jr et al. 1997)	2,290 km <sup>2</sup> (Júlio Jr et al. 1997)
Headwaters	In the Serra da Esperança, in the reverse slope of the Ivaí and Piquiri rivers (and of the Areia River), at altitudes over 1,200 m (Maack 2012)	In the Serra da Esperança, in the reverse slope of the Ivaí River (and of the Jordão River)
Mouth	At the Iguaçu River 2.5 km downstream of the dam of the Governador Ney Aminthas de Barros Braga hydroelectric plant, which is commonly referred to as the Segredo hydroelectric plant	At the reservoir of the Governador Bento Munhoz da Rocha Neto hydroelectric plant, commonly known as Foz do Areia hydroelectric plant
Largest tributaries	Cascavel, Coutinho, Campo Real and Caracu Rivers on the right margin (Paraná 2008) Capão Grande, Capivara, Pinhãozinho and Pinhão Rivers on the left margin (Paraná 2008)	Pimpão and Lageado Feio rivers on the right margin (Lactec 2009) da Praia River on the left margin (Lactec 2009)
Stretches	The upper stretch, with about 50 km of extension and low average steepness (0.9 m/km), is situated upstream of the Salto Curuaca and drains the Guarapuava grasslands (Paraná 2008). The middle stretch, also with 50 km of extension but a slightly higher average steepness (1.5 m/km), is comprised between the Salto Curuaca and the Santa Clara hydroelectric plant downstream (Paraná 2008) The lower stretch, with 40 km and a much higher average steepness (6.0 m/km), comprises watercourses situated in steep valleys and marked by several cascades (Paraná 2008)	There are no references There are no references There are no references
Current use	Water supply, industrial water use, irrigation and electricity generation (Paraná 2008); although drains areas of agriculture, it also drains areas of preserved native forests and reforestation (Pachechenik and Souza 2005; Paraná 2008)	The landscape in the vicinities of the Foz do Areia reservoir are heavily anthropized (Lactec 2009) by large pine plantations



**Figure 1.** Map of the study area showing (A) the location of the Iguaçú River basin within Brazil and Paraná state and (B) the 30 survey sites within the Jordão (grey area) and Areia (green area) subbasins. Yellow circles represent sampling sites, with numbers corresponding to Appendix Table A1. The red and orange stars represent the Governor Ney Aminthas de Barros Braga hydroelectric plant and the Governor Bento Munhoz da Rocha Neto hydroelectric plant, respectively (both of them located on the main channel of the Iguaçú River).


(CRIA 2016). However, in this case, species with dubious determination or distribution were disregarded. Whenever possible, determinations were checked by specialists of each taxonomic group.

Fishes were categorized based on their distribution in the Iguaçú River basin, following Baumgartner et al. (2012), as endemic indigenous (species with distribution restricted to the Iguaçú River basin), native indigenous (species native from the Iguaçú River basin, but not restricted to it), and non-indigenous (species introduced to the Iguaçú River basin).

**RESULTS**

We collected from our 30 sampling sites 1,557 specimens, of which 401 are from the Jordão River basin and 1,156 from the Areia River basin. The number of species we collected (27) combined with those reported in the literature and the Species Link database is 67. Six orders and 18 families of Actinopterygii are represented. In each of the basins, 54 and 48 species are present in the Jordão and Areia basins, respectively (Table 2). Siluriformes and Characiformes are the most species-rich orders that together comprise about

79.6% and 77.1% of the species in the Jordão and Areia basins, respectively (Figure 2). In the Jordão basin the families Characidae and Trichomycteridae represent a combined 38.9% of ichthyofauna, and in the Areia basin,



**Figure 2.** Specific richness of each of the six orders recorded in the Jordão and Areia basins, shown as percentages of the total richness in each basin.



**Table 2.** List of the species recorded in the Jordão and Areia basins, Paraná state, Brazil. The presence of each species in each of the two basins is signalled with an “X” in the respective column. When a species is present in both, two voucher numbers are provided: the first representing specimens from the Jordão River basin, and the second representing specimens from the Areia River basin. Eni = endemic indigenous, Nai = native indigenous, Noi = non-indigenous.

Classification	Basin		Origin	Voucher numbers
	Jordão	Areia		
<b>ACTINOPTERYGII</b>				
<b>CHARACIFORMES</b>				
<b>Anostomidae</b>				
<i>Leporinus octofasciatus</i> Steindachner, 1915		X	Noi	NUP 12787
<b>Characidae</b>				
<i>Astyanax bifasciatus</i> Garavello & Sampaio, 2010	X	X	Eni	NUP 16106 and NUP 15913
<i>Astyanax dissimilis</i> Garavello & Sampaio, 2010	X	X	Eni	NUP 9996 and NUP 15885
<i>Astyanax gymnodontus</i> (Eigenmann, 1911)	X	X	Eni	NUP 2460 and NUP 2423
<i>Astyanax gymnogenys</i> Eigenmann, 1911	X	X	Eni	NUP 1576 and NUP 2037
<i>Astyanax jordanensis</i> Vera Alcaraz, Pavanelli & Bertaco, 2009	X	X	Eni	NUP 17373 and NUP 15922
<i>Astyanax lacustris</i> (Lütken, 1875)	X		Nai	NUP 1314
<i>Astyanax longirhinus</i> Garavello & Sampaio, 2010	X	X	Eni	NUP 10152 and NUP 11858
<i>Astyanax minor</i> Garavello & Sampaio, 2010	X	X	Eni	NUP 11258 and NUP 15884
<i>Astyanax serratus</i> Garavello & Sampaio, 2010		X	Eni	NUP 11859
<i>Bryconamericus ikaa</i> Casciotta, Almirón & Azpelicueta, 2004	X	X	Eni	NUP 1649 and NUP 15725
<i>Bryconamericus</i> sp.	X	X	Eni	NUP 1654 and NUP 15911
<i>Hyphessobrycon reticulatus</i> Ellis, 1911		X	Nai	NUP 15797
<i>Oligosarcus longirostris</i> Menezes & Géry, 1983	X	X	Eni	MHNCI 8346 and NUP 15881
Undetermined genus and species ( <i>sensu</i> Baumgartner et al. 2012: 86)	X		Eni	NUP 11752
<b>Crenuchidae</b>				
<i>Characidium</i> sp. 1	X	X	Eni	NUP 1655 and NUP 15876
<i>Characidium travassosi</i> Melo, Buckup & Oyakawa, 2016	X	X	Eni	NUP 17799 and NUP 15935
<b>Erythrinidae</b>				
<i>Hoplias</i> sp.	X	X	Nai	NUP 10848 and MHNCI 8668
<b>Parodontidae</b>				
<i>Apareiodon vittatus</i> Garavello, 1977	X		Eni	NUP 12097
<b>Prochilodontidae</b>				
<i>Prochilodus lineatus</i> (Valenciennes, 1837)		X	Noi	NUP 3010
<b>CYPRINIFORMES</b>				
<b>Cyprinidae</b>				
<i>Cyprinus carpio</i> Linnaeus, 1758	X	X	Noi	NUP 725 and NUP 3062
<i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844)		X	Noi	NUP 2383
<b>CYPRINODONTIFORMES</b>				
<b>Poeciliidae</b>				
<i>Cnesterodon carnegiei</i> Haseman, 1911		X	Eni	NUP 15936
<i>Cnesterodon omorgmatus</i> Lucinda & Garavello, 2001	X		Eni	MCP 22741
<i>Phalloceros harpagos</i> Lucinda, 2008	X	X	Nai	NUP 16103 and NUP 15983
<b>Anablepidae</b>				
<i>Jenynsia diphyes</i> Lucinda, Ghedotti & Graça, 2006	X	X	Eni	NUP 608 and NUP 15912
<b>GYMNOTIFORMES</b>				
<b>Gymnotidae</b>				
<i>Gymnotus inaequilabiatus</i> (Valenciennes, 1839)	X	X	Noi	NUP 3752 and NUP 3043
<b>PERCIFORMES</b>				
<b>Cichlidae</b>				
<i>Australoheros angiru</i> Řičan, Piálek, Almirón & Casciotta, 2011	X		Nai	NUP 2430
<i>Cichlasoma dimerus</i> (Heckel, 1840)		X	Noi	NUP 9758
<i>Coptodon rendalli</i> (Boulenger, 1897)	X		Noi	NUP 3749
<i>Crenicichla iguassuensis</i> Haseman, 1911	X	X	Eni	NUP 3683 and NUP 2954
<i>Crenicichla tesay</i> Casciotta & Almirón, 2009	X	X	Eni	NUP 11288 and NUP 11428
<i>Geophagus</i> aff. <i>brasiliensis</i> (Quoy & Gaimard, 1824)	X	X	Nai	NUP 16107 and MCP 49248
<i>Oreochromis niloticus</i> (Linnaeus, 1758)	X	X	Noi	NUP 11438 and NUP 3063
<b>SILURIFORMES</b>				
<b>Auchenipteridae</b>				
<i>Glanidium ribeiroi</i> Haseman, 1911	X		Eni	NUP 11191
<b>Callichthyidae</b>				
<i>Callichthys callichthys</i> (Linnaeus, 1758)		X	Noi	NUP 5490

Continued

Table 1. Continued.

Classification	Basin		Origin	Voucher numbers
	Jordão	Areia		
<i>Corydoras ehrhardti</i> Steindachner, 1910	X	X	Nai	NUP 17790 and NUP 15802
<i>Corydoras aff. paleatus</i> (Jenyns, 1842)	X	X	Eni	NUP 13206 and NUP 2960
<i>Hoplosternum littorale</i> (Hancock, 1828)		X	Noi	NUP 11201
<b>Heptapteridae</b>				
<i>Heptapterus</i> sp.		X	Eni	NUP 15925
" <i>Pariolius</i> " <i>hollandi</i> (Haseman, 1911)	X		Eni	NUP 2976
" <i>Pariolius</i> " sp.	X	X	Eni	NUP 3710 and NUP 15888
<i>Rhamdia branneri</i> Haseman, 1911	X	X	Eni	NUP 10849 and NUP 569
<i>Rhamdia voulezi</i> Haseman, 1911	X	X	Eni	NUP 3741 and NUP 2763
<i>Rhamdia</i> sp.	X		Nai	NUP 2428
<b>Loricariidae</b>				
<i>Ancistrus abelhoi</i> Bifi, Pavanelli & Zawadzki, 2009		X	Eni	NUP 7502
<i>Ancistrus agostinhoi</i> Bifi, Pavanelli & Zawadzki, 2009	X		Eni	MZUSP 104118
<i>Hypostomus albopunctatus</i> (Regan, 1908)	X		Nai	NUP 593
<i>Hypostomus commersoni</i> Valenciennes, 1836	X	X	Nai	NUP 596 and NUP 2541
<i>Hypostomus derbyi</i> (Haseman, 1911)	X	X	Nai	NUP 585 and NUP 2541
<i>Hypostomus myersi</i> (Gosline, 1947)	X	X	Nai	MHNCI 7633 and NUP 5749
<i>Hypostomus nigropunctatus</i> Garavello, Britski & Zawadzki, 2012	X		Eni	NUP 5082
<i>Pareiorhaphis cf. parmula</i> Pereira, 2005		X	Eni	NUP 15928
<i>Rineloricaria maacki</i> Ingenito, Ghazzi, Duboc & Abilhoa, 2008		X	Eni	NUP 2540
<i>Neoplecostomus</i> sp.	X		Eni	NUP 4069
<b>Pimelodidae</b>				
<i>Pimelodus britskii</i> Garavello & Shibatta, 2007	X	X	Eni	NUP 598 and NUP 2381
<i>Pimelodus ortmanni</i> Haseman, 1911	X	X	Eni	NUP 3747 and NUP 3007
<b>Trichomycteridae</b>				
<i>Trichomycterus castroi</i> de Pinna, 1992	X		Eni	NUP 3127
<i>Trichomycterus crassicaudatus</i> Wosiacki & de Pinna, 2008	X		Eni	MZUSP 88518
<i>Trichomycterus davisii</i> (Haseman, 1911)	X	X	Nai	NUP 16104 and NUP 15927
<i>Trichomycterus igobi</i> Wosiacki & de Pinna, 2008	X	X	Eni	NUP 9866 and NUP 15882
<i>Trichomycterus mboycy</i> Wosiacki & Garavello, 2004	X		Eni	NUP 3716
<i>Trichomycterus papilliferus</i> Wosiacki & Garavello, 2004	X	X	Eni	NUP 10828 and NUP 17364
<i>Trichomycterus plumbeus</i> Wosiacki & Garavello, 2004	X		Eni	NUP 10829
<i>Trichomycterus stawiarski</i> (Miranda Ribeiro, 1968)	X	X	Eni	NUP 10830 and NUP 15909
<i>Trichomycterus taroba</i> Wosiacki & Garavello, 2004	X		Eni	NUP 3152
<b>Ictaluridae</b>				
<i>Ictalurus punctatus</i> (Rafinesque, 1818)	X		Noi	NUP 584
Total	54	48	-	-





Figure 3. Specific richness of each of the 16 families recorded in the Jordão and Areia basins, shown as percentages of the total richness in each basin.

the Characidae and Loricariidae have a combined 37.5% of the species (Figure 3).

All sampling sites were dominated by small-sized species such as *Astyanax bifasciatus*, *A. jordanensis*, and *Trichomycterus davisii*. Those were the most abundant, and in all but one site (23; Figure 1; Appendix Figure A2, Table A1) at least one of the three was captured. In the Jordão River basin, 68.5% of the species are endemic indigenous, 22.2% are native indigenous and 9.3% are non-indigenous. In the Areia River basin, 62.4% are endemic indigenous, 18.8% are native indigenous and other 18.8% are non-indigenous. Four species, *Astyanax jordanensis*, *Characidium travassosi*, *Jenynsia diphyes*, and *Trichomycterus igobi*, previously thought to be restricted to the Jordão River basin, were also captured in the Areia River (Figure 4). Short descriptions of those species are provided below.



**Figure 4.** Representative individuals of species previously thought to be endemic to the Jordão basin (images on the right) but now known from the Areia basin (images on the left). Species, voucher numbers and SL are as follows: *Astyanax jordanensis*, (A) NUP 15922, 68.3 mm, (B) NUP 17373, 64.2 mm; *Characidium travassosi*, (C) NUP 15935, 54.4 mm, (D) NUP 17799, 55.2 mm; *Jenynsia diphyes*, (E) NUP 15912, 50.8 mm, (F) NUP 608, 57.0 mm; *Trychomycterus igobi*, (G) NUP 15882, 92.4 mm; (H) NUP 9866, 125.9 mm. Photographs by Celso Ikedo.

***Astyanax jordanensis*** Vera Alcaraz, Pavanelli & Bertaco, 2009

*Astyanax jordanensis* Vera Alcaraz, Pavanelli & Bertaco (2009): 185, 186, 187, 188, 189. — Baumgartner et al. (2012): 74, 80.

**Material examined:** Table 1; Appendix Table A1; Figure 4 (A and B).

The presence of two tooth rows in the premaxilla, the inner with five teeth; lateral line complete; adipose fin present; caudal fin naked, with scales only to its base; absence of teeth on the palate; dentary teeth abruptly decreasing in size posterior to fifth tooth and pelvic bone not prolonged anteriorly into a spine distinguishes *A. jordanensis* from other characiform genera occurring in the surveyed basins and adjacent areas. Among other congeners in the Iguaçu River basin, *A. jordanensis* is distinguished by having 14–17 branched anal-fin rays; 35–38 perforated lateral-line scales; premaxillary symphyseal tooth tetracuspid, anteriorly convex; inner series of premaxilla with slightly convex teeth; only one humeral spot, which is dark grey, vertically elongate; posteroventral margin of third infraorbital close to

preopercle or touching it, leaving but a narrow naked area in the cheek or none at all. This species belongs in the *A. scabripinnis* species group.

***Characidium travassosi*** Melo, Buckup & Oyakawa, 2016

*Characidium travassosi* Melo, Buckup & Oyakawa (2016): Figures 1 A–E, 3 A–C.

*Characidium* sp. 2 — Baumgartner et al. (2012): 70, 71.

*Characidium* sp. B — Severi & Cordeiro (1994): 47, 48.

**Material examined:** Table 1; Appendix Table A1; Figure 4 (C and D).

The three anterior most pectoral-fin rays unbranched, instead of only the first ray unbranched, distinguishes *Characidium travassosi* from other characiform genera occurring in the surveyed basins and adjacent areas. Among other congeners in the Iguaçu River basin, *Characidium travassosi* is distinguished by lacking adipose fin.

***Jenynsia diphyes*** Lucinda, Ghedotti & Graça, 2006

*Jenynsia diphyes* Lucinda, Ghedotti & Graça (2006): 616, 617, 619, 620. — Baumgartner et al. (2012): 161, 162.

**Material examined:** Table 1; Appendix Table A1; Figure 4 (E and F).

The presence of scales on the top of the head anteriorly extending to about the vertical through the anterior margin of the eye; only one dorsal fin; gonopodium shorter than the head; caudal fin truncate; mouth terminal; inner dentary teeth tricuspid; and vertical through base of first dorsal fin ray anterior to vertical through base of first anal fin ray even in males distinguishes *J. diphyes* from other genera occurring in the surveyed basins and adjacent areas. A midlateral stripe along the flank, usually discontinuous anteriorly to vertical through first dorsal fin ray and 10 anal fin rays distinguishes it from the other congener in the basin.

***Trichomycterus igobi*** Wosiacki & de Pinna, 2008

*Trichomycterus igobi* Wosiacki & de Pinna (2008): 17, 18, 19, 20, 21, 22. — Baumgartner et al. (2012): 103, 106.

**Material examined:** Tables 1 and A1; Figure 4 (G and H).

The body completely naked, with no bony plates; opercle and interopercle with posteriorly directed odontodes; presence of a nasal barbel; and terminal mouth distinguishes *T. igobi* from other siluriform genera occurring in the surveyed basins and adjacent areas. It is distinguished from other congeners from the Iguaçu River basin by having the caudal fin truncate, without a black bar close to its distal margin; body with randomly distributed dark-brown spots; head long, comprised 4.2 times or less in the SL; and eight pectoral fin rays.

## DISCUSSION

In the most recent inventory of fishes of the Iguaçu River basin, Baumgartner et al. (2012) recorded 106 species in the lower Iguaçu River. Of these, 50.9% and 45.3% are present in the Jordão and Areia basins, respectively. The high species richness of Siluriformes and Characiformes in comparison to other orders is a pattern widespread throughout the Neotropics (Castro 1999; Lowe-McConnell 1999; Buckup et al. 2007) and this pattern has already been shown for the upper Paraná River basin (Graça and Pavanelli 2007; Langeani et al. 2007; Maier et al. 2008; Cunico et al. 2009; Gubiani et al. 2010; Araújo et al. 2011; Delariva and Silva 2013; Frota et al. 2016) and even for the Iguaçu River basin itself (Severi and Cordeiro 1994; Garavello et al. 1997; Ingenito et al. 2004; Baumgartner et al. 2006; Bifi et al. 2006; Abilhoa et al. 2008; Baumgartner et al. 2012).

The dominance of Characidae, Loricariidae and Trichomycteridae, comprising small and medium-sized species (15 cm or less), is characteristic of localities such as the ones sampled (Appendix Figures A1 and A2). This dominance of these families in the Iguaçu basin was also found by Baumgartner et al. (2012). Environments with rocky substrates, and grassy margins with series of pools (Súarez et al. 2007; Ferreira et al. 2010; Araújo et al. 2011; Pagotto et al. 2011) favour fast swimmers (e.g., *Astyanax* spp. and *Bryconamericus* spp.) and bottom dwellers that

fixate to rocks with suckermouths, odontodes, and spines (e.g., loricariids and trichomycterids) (Ferreira et al. 2010).

Endemism in the two river basins inventoried here is high, and is thought to be associated with the formation of Iguaçu Falls in the Miocene (about 22 Mya), which isolated populations in the Iguaçu River (Mello et al. 2015). Further explanations to the high rate of endemism in the Iguaçu basin have been developed lately. Garavello and Sampaio (2010) explain that the juvenile aspect of several regions within the basin of Iguaçu River is a probable consequence of a tectonic reactivation, with several “breaks” in the drainage. This would have isolated waterbodies and thus compartmentalized different subbasins.

The close relationship between the geomorphological and morphodynamic characteristics and the hydrography of the Iguaçu River developed rugged terrain with several waterfalls, which influenced the geographic distribution of several fish species (Baumgartner et al. 2012). Lucinda et al. (2006), Vera Alcaraz et al. (2009), and Baumgartner et al. (2012) considered the following species to be endemic to the Jordão basin: *Ancistrus agostinhoi*, *Astyanax jordanensis*, *Characidium travassosi*, *Cnesterodon omorgmatus*, *Jenynsia diphyes*, *Trichomycterus igobi*, *T. crassicaudatus*, *T. plumbeus*, and *T. taroba*. This was explained by the existence, until recently, of a 15-m waterfall near the river’s mouth that is now submerged by the Usina Hidrelétrica da Derivação do rio Jordão reservoir (Lucinda et al. 2006; Vera Alcaraz et al. 2009). We expand the geographic distributions of *Astyanax jordanensis*, *Characidium travassosi*, *Jenynsia diphyes*, and *Trichomycterus igobi* to include the Areia River basin and show that the rate of endemism in the Jordão River is not as great as previously thought. The similarity between the two basins is possibly the result of a common geological past, because before the uplift of the Serra da Esperança their headwaters lay closer together (Ribeiro 2006) suggesting a past connection.

In modern times, the rate of endemism in the Iguaçu basin has decreased, mostly due to anthropic actions and particularly because of species introduction. Considering introduced species, Baumgartner et al. (2012) observed an endemism rate of only 50% for the lower Iguaçu basin. In aquatic environments, invasive species can cause negative effects on diversity and perpetrate extreme changes in the structure of communities (Pellice and Agostinho 2009). According to Johnson et al. (2008) biological invasions have been a major problem to freshwater ecosystems together with hydrological changes, which on a global scale seriously threaten the aquatic biota. The main mechanisms by which species introduction takes place in the Iguaçu basin are their use as bait by sport fishers (*Callichthys callichthys*, *Gymnotus inaequilabiatus* and *Hoplosternum littorale*) and escapees



from aquaculture farms (*Cyprinus carpio*, *Hypophthalmichthys molitrix*, *Ictalurus punctatus*, *Oreochromis niloticus*, *Prochilodus lineatus* and *Coptodon rendalli*).

The portion of the Iguaçú River basin upstream of Iguaçú Falls represents an ecoregion (or biogeographic region) considered different from that of the Paraná River, in spite of emptying into the latter (Abell et al. 2008). That is because the Iguaçú River has a simplified ichthyofauna, lacking several genera and species present in the Paraná River, and a high rate of endemism. In addition to introduced fish species that threaten native species, the Jordão and Areia basins are also plagued by urban expansion, untreated sewage, and pesticides as well as other agrichemicals. Thus, Baumgartner et al. (2012) pointed out the need for new species inventories and taxonomic studies on ichthyofauna of the Iguaçú River. Recent studies have shown an increasing number of species restricted to this drainage that justify the need for conservation of sites in the Iguaçú River basin.

## ACKNOWLEDGEMENTS

We are deeply grateful to Steven Grant for English review, Fundação Araucária (Seti-PR) and Nupélia-UEM for financial support, Francisco Alves Teixeira, Wladimir Marques Domingues, and Rodrigo Júnio da Graça for assistance in the field, Claudimar Jean dos Santos for help in fish cataloging, Cláudio Zawadzki, Luiz Fernando Caserta Tencatt, Fagner de Souza, and Carlos Alexandre Miranda Oliveira for help in fish identification, and Celso Ikedo for the photography. AF and WJG received grants from Fundação Araucária, and GCD, from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

## LITERATURE CITED

- Abell, R., M.L. Thieme, C. Revenga, M. Bryer, M. Kottelat, N. Bogutskaya, B. Coad, N. Mandrak, S. Contreras-Balderas, W. Bussing, M.L.J. Stiassny, P. Skelton, G. R. Allen, P. Unmack, A. Naseka, N.G. Rebecca, N. Sindorf, J. Robertson, E. Armijo, J.V. Higgins, T.J. Heibel, E. Wikramanayake, D. Olson, H.L. López, R. E. Reis, J.G. Lundberg, M.H. Sabaj Pérez and P. Petry. 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience* 58(5): 403–414. doi: [10.1641/B580507](https://doi.org/10.1641/B580507)
- Abilhoa, V., L.F. Duboc and D.P. de Azevedo Filho. 2008. A comunidade de peixes de um riacho de Floresta com Araucária, alto rio Iguaçú, sul do Brasil. *Revista Brasileira de Zoologia* 25(2): 238–246. doi: [10.1590/S0101-81752008000200012](https://doi.org/10.1590/S0101-81752008000200012)
- Araújo, M.I., R.L. Delariva, K.O. Bonato and J.C. Silva. 2011. Fishes in first order stream in Ivaí River drainage basin, upper Paraná River basin, Paraná state, Brazil. *Check List* 7(6): 774–777. doi: [10.15560/11023](https://doi.org/10.15560/11023)
- Baumgartner, D., G. Baumgartner, C.S. Pavanelli, P.R.L. da Silva, V.A. Frana, L.C. de Oliveira and M.R. Michelon. 2006. Fish, Salto Osório Reservoir, Iguaçú River basin, Paraná state, Brazil. *Check List* 2(1): 1–4. doi: [10.15560/2.1.1](https://doi.org/10.15560/2.1.1)
- Baumgartner, G., C.S. Pavanelli, D. Baumgartner, A.G. Bifi, T. Debona and V.A. Frana 2012. Peixes do baixo rio Iguaçú. Maringá: Eduem. 203 pp.
- Bifi, A.G., C.S. Pavanelli and C.H. Zawadzki. 2009. Three new species of Ancistrus Kner, 1854 (Siluriformes: Loricariidae) from the Rio Iguaçú basin, Paraná state, Brazil. *Zootaxa* 2275: 41–59. <http://peixe.nupelia.uem.br/uploads/pdf/64.pdf>
- Bifi, A.G., D. Baumgartner, G. Baumgartner, V.A. Frana and T. Debona. 2006. Composição específica e abundância da ictiofauna do rio dos Padres, bacia do rio Iguaçú, Brasil. *Acta Scientiarum. Biological Sciences* 28(3): 203–211. doi: [10.4025/actascibiolsci.v28i3.193](https://doi.org/10.4025/actascibiolsci.v28i3.193)
- Buckup, P.A., N.A. Menezes and M.S. Ghazzi. 2007. Catálogo das espécies de peixes de água doce do Brasil. Rio de Janeiro: Museu Nacional (UFRJ). 195 pp.
- Casciotta, J. and A. Almirón. 2008. *Crenicichla tesay*, a new species of cichlid (Perciformes: Labroidei) from the rio Iguazú basin in Argentina. *Revue Suisse de Zoologie* 115(3): 651–660. doi: [10.5962/bhl.part.80451](https://doi.org/10.5962/bhl.part.80451)
- Casciotta, J.R., A.E. Almirón and M.M. Azpelicueta. 2004. *Brycon-americanus ikaa*, a new species from tributaries of the rio Iguazú in Argentina (Characiformes, Characidae). *Ichthyological Exploration of Freshwaters* 15(1): 61–66.
- Castro, R.M.C. 1999. Evolução da ictiofauna de riachos sul americanos: padrões gerais e possíveis processos casuais; pp. 139–155, in: E.P. Caramaschi, R. Mazzoni and P.R. Peres-Neto (eds.). *Ecologia de peixes de riachos: estado atual e perspectivas*. Rio de Janeiro: UFRJ.
- Cunico, A.M., W.J. Graça, A.A. Agostinho, W.M. Domingues and J.D. Latini. 2009. Fish, Maringá urban streams, Pirapó River drainage, upper Paraná River basin, Paraná state, Brazil. *Check List* 5(2): 273–280. doi: [10.15560/5.2.273](https://doi.org/10.15560/5.2.273)
- CRIA [Centro de Referência em Informação Ambiental]. 2016. SpeciesLink. Accessed at <http://splink.cria.org.br>, 17 October 2016.
- Delariva, R.L. and J.C. da Silva. 2013. Fish fauna of headwater streams of Perobas Biological Reserve, a conservation unit in the Atlantic Forest of the northwestern Paraná state, Brazil. *Check List* 9(3): 549–554. doi: [10.15560/9.3.549](https://doi.org/10.15560/9.3.549)
- Ferreira, F.C., U.P. Souza and M. Petrere Jr. 2010. Zonacção longitudinal da ictiofauna em ambientes lóticos. *Boletim da Sociedade Brasileira de Limnologia* 38(1): 1–17. [http://www.ablimno.org.br/boletins/pdf/bol\\_38\(1-4\).pdf](http://www.ablimno.org.br/boletins/pdf/bol_38(1-4).pdf)
- Froese, R. and D. Pauly. 2016. FishBase. Worldwide web electronic publication. 2014. Accessed at <http://www.fishbase.org/>, 13 March 2016.
- Frota, A., G.C. Deprá, L.M. Petenucci and W.J. Graça. 2016. Inventory of the fish fauna from Ivaí River basin, Paraná state, Brazil. *Biota Neotropica* 16(3): e20150151. doi: [10.1590/1676-0611-BN-2015-0151](https://doi.org/10.1590/1676-0611-BN-2015-0151)
- Garavello, J.C. 2005. Revision of genus *Steindachneridion* (Siluriformes: Pimeloidae). *Neotropical Ichthyology* 3(4): 607–623. doi: [10.1590/S1679-62252005000400018](https://doi.org/10.1590/S1679-62252005000400018)
- Garavello, J.C. and F.A.A. Sampaio. 2010. Five new species of genus *Astyanax* Bird & Girard, 1854 from Rio Iguaçú, Paraná, Brazil (Ostariophysi, Characiformes, Characidae). *Brazilian Journal of Biology* 70(3): 847–865. doi: [10.1590/S1519-69842010000400016](https://doi.org/10.1590/S1519-69842010000400016)
- Garavello, J.C. and O.A. Shibatta. 2007. A new species of the genus *Pimelodus* La Cépède, 1803 from the rio Iguaçú basin and a reappraisal of *Pimelodus ortmanni* Haseman, 1911 from the rio Paraná system, Brazil (Ostariophysi: Siluriformes: Pimelodidae). *Neotropical Ichthyology* 5(3): 285–292. doi: [10.1590/S1679-62252007000300008](https://doi.org/10.1590/S1679-62252007000300008)
- Garavello, J.C., C.S. Pavanelli, H.I. Suzuki. 1997. Caracterização da ictiofauna do rio Iguaçú; pp. 61–84, in: A.A. Agostinho and L.C. Gomes (Eds.). *Reservatório de Segredo: bases ecológicas para o manejo*. Maringá: EDUEM.
- Garavello, J.C., H.A. Britski and C.H. Zawadzki. 2012. The cascudos of the genus *Hyposotomus* Lacépède (Ostariophysi: Loricariidae)



- from the Rio Iguaçu basin. *Neotropical Ichthyology* 10(2): 263–283. doi: [10.1590/S1679-62252012000200005](https://doi.org/10.1590/S1679-62252012000200005)
- Graça, W.J. and C.S. Pavanelli. 2007. Peixes da planície de inundação do alto rio Paraná e áreas adjacentes. Maringá: EDUEM. 241 pp.
- Gubiani, E.A., V.S. Daga, V.A. Frana and W.J. Graça. 2010. Fish, Toledo urban streams, São Francisco Verdadeiro River drainage, upper Paraná River basin, state of Paraná, Brazil. *Check List* 6(1): 45–48. <http://www.checklist.org.br/getpdf?SL055-09>
- Hahn, N.S. and R. Fugi. 2008. Environmental changes, habitat modifications and feeding ecology of freshwater fish; pp. 35–65, in: J.E.P. Cyrino, D.P. Bureau and B.G. Kapoor (eds.). *Feeding and digestive functions of fishes*. Enfield: Science Publishers.
- Ingenito, L.F.S., L.F. Duboc and V. Abilhoa. 2004. Contribuição ao conhecimento da ictiofauna da bacia do alto rio Iguaçu, Paraná, Brasil. *Arquivos de Ciências Veterinárias e Zoológicas da UNIPAR* 7(1): 23–36. <http://www.revistas.unipar.br/?journal=veterinaria&page=article&op=view&path%5B%5D=540&path%5B%5D=479>
- Ingenito, L.F.S., M.S. Ghazzi, L.F. Duboc and V. Abilhoa. 2008. Two new species of *Rineloricaria* (Siluriformes: Loricariidae) from the Rio Iguaçu basin, southern Brazil. *Neotropical Ichthyology* 6(3): 355–366. doi: [10.1590/S1679-62252008000300009](https://doi.org/10.1590/S1679-62252008000300009)
- Johnson, P.T.J., J.D. Olden and M.J.V. Zanden. 2008. Dam invaders: impoundments facilitate biological invasions into freshwaters. *Frontiers in Ecology and the Environment* 6(7): 357–363. doi: [10.1890/070156](https://doi.org/10.1890/070156)
- Julio Jr, H.F., C.C. Bonecker and A.A. Agostinho. 1997. Reservatório de Segredo e sua inserção na bacia do rio Iguaçu; pp. 1–17, in: A.A. Agostinho and L.C. Gomes (eds.). *Reservatório de Segredo: bases ecológicas para o manejo*. Maringá: EDUEM.
- Kantec, D.L.Z., R.R. Cipriano, V. Abilhoa, R.F. Artoni and M.M. Cestari. 2007. Cytotaxonomic and evolutionary considerations about karyotypic data of fishes from the Iguaçu River basin in south of Brazil. *Brazilian Archives of Biology and Technology* 50(5): 793–802. doi: [10.1590/S1516-89132007000500007](https://doi.org/10.1590/S1516-89132007000500007)
- Lactec [Instituto de Tecnologia para o Desenvolvimento]. 2009. Plano Ambiental de Conservação e Uso do Entorno de Reservatório Artificial UHE Governador Bento Munhoz da Rocha Neto (Atualização). Curitiba. Accessed at [http://www.iap.pr.gov.br/arquivos/File/EIA\\_RIMA/UHE\\_GNB\\_SEGREDO\\_PLANO\\_AMBIENTAL/CAPA\\_volume\\_II.pdf](http://www.iap.pr.gov.br/arquivos/File/EIA_RIMA/UHE_GNB_SEGREDO_PLANO_AMBIENTAL/CAPA_volume_II.pdf), 16 March 2016.
- Langeani, F., R.M.C. Castro, O.T. Oyakawa, O.A. Shibatta, C.S. Pavanelli and L. Casatti. 2007. Diversidade da ictiofauna do Alto Rio Paraná: composição atual e perspectivas futuras. *Biota Neotropica* 7(3): 1–17. doi: [10.1590/S1676-06032007000300020](https://doi.org/10.1590/S1676-06032007000300020)
- Lobón-Cerviá, J. 1991. Dinámica de poblaciones de peces en rios: pesca eléctrica y métodos de capturas sucesivas en la estima de abundancias. Monografía. Museo Nacional de Ciencias Naturales. Consejo Superior de Investigaciones Científicas. 157 pp.
- Loureiro-Crippa, V.E. and N.S. Hahn. 2006. Use of food resources by the fish fauna of a small reservoir (rio Jordão, Brazil) before and shortly after its filling. *Neotropical Ichthyology* 4(3): 357–362. doi: [10.1590/S1679-62252006000300007](https://doi.org/10.1590/S1679-62252006000300007)
- Lowe-McConnell, R.H. 1999. *Estudos ecológicos de comunidades de peixes tropicais*. São Paulo: EDUSP. 534 pp.
- Lucinda, P.H.F. 2005. Systematics of the genus *Cnesterodon* Garman, 1895 (Cyprinodontiformes: Poeciliidae: Poeciliinae). *Neotropical Ichthyology* 3(2): 259–270. doi: [10.1590/S1679-62252005000200003](https://doi.org/10.1590/S1679-62252005000200003)
- Lucinda, P.H.F. 2008. Systematics and biogeography of the genus *Phalloceros* Eigenmann, 1907 (Cyprinodontiformes: Poeciliidae: Poeciliinae), with the description of twenty-one new species. *Neotropical Ichthyology* 6(2): 113–158. doi: [10.1590/S1679-62252008000200001](https://doi.org/10.1590/S1679-62252008000200001)
- Lucinda, P.H.F., M.J. Ghedotti and W.J. Graça. 2006. A new *Jenynsia* species (Teleostei, Cyprinodontiformes, Anablepidae) from southern Brazil and its phylogenetic position. *Copeia* 4: 613–622. doi: [10.1643/0045-8511\(2006\)6\[613:ANJSTC\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2006)6[613:ANJSTC]2.0.CO;2)
- Maack, R. 2012. *Geografia física do Estado do Paraná*. 4 edition. Ponta Grossa: Editora UEPG. 526 pp.
- Maier, A., C.H. Zawadzki, W.J. Graça and A.G. Bifi. 2008. Fish, Barra Bonita River, Ivaí River drainage, upper Paraná basin, Paraná State, Brazil. *Check List* 4(3): 336–340. doi: [10.15560/4.3.336](https://doi.org/10.15560/4.3.336)
- Mello, R., T.C. Maniglia, S.M.A. P. Prioli and A.J. Prioli. 2015. Genetic and biogeographical relationships among species of *Astyanax* (Teleostei, Characidae) in Brazilian river basins. *Genetics and Molecular Research* 14(4): 15356–15364. doi: [10.4238/2015.November.30.13](https://doi.org/10.4238/2015.November.30.13)
- Melo, M.R.S., P.A. Buckup and O.T. Oyakawa. 2016. A new species of *Characidium* Reinhardt, 1867 (Characiformes: Crenuchidae) endemic to the Atlantic Forest in Paraná State, southern Brazil. *Neotropical Ichthyology* 14(3): e160014. doi: [10.1590/1982-0224-20160014](https://doi.org/10.1590/1982-0224-20160014)
- Mise, F.T., L.F.C. Tencatt and F. de Souza. 2013. Ecomorphological acustress between *Rhamdia* (Bleeker, 1858) populations from the Iguaçu River Basin. *Biota Neotropica* 13(4): 1–6. <http://www.biotaneotropica.org.br/v13n4/pt/abstract?article+bn01513042013>
- Pachechenik, P.E. and J.L.M. de Souza. 2005. Estudo do processo chuva-vazão na bacia do rio das Pedras, Guarapuava, Estado do Paraná, com o modelo *HYCYMODEL*. *Acta Scientiarum. Agronomy* 27(4): 677–686. doi: [10.4025/actasciagron.v27i4.1689](https://doi.org/10.4025/actasciagron.v27i4.1689)
- Pagotto, J.P.A., E. Goulart, E.F. Oliveira and C.B. Yamamura. 2011. Trophic ecomorphology of Siluriformes (Pisces, Osteichthyes) from a tropical stream. *Brazilian Journal of Biology* 71(2): 469–479. <http://www.scielo.br/pdf/bjb/v71n2/a17v71n2>
- Paraná. Secretaria do Estado do Meio Ambiente e Recursos Hídricos (SEMA). 2008. Bacia Hidrográfica do Rio Jordão. Fase 1—Diagnóstico. Curitiba: SEMA. Accessed at [http://www.aguasparana.pr.gov.br/arquivos/File/jordao/plano\\_de\\_bacia/diagnostico\\_bhj.pdf](http://www.aguasparana.pr.gov.br/arquivos/File/jordao/plano_de_bacia/diagnostico_bhj.pdf), 14 March 2016.
- Pavanelli, C.S. and A.G. Bifi. 2009. A new *Tatia* (Ostariophysi: Siluriformes: Auchenipteridae) from the Rio Iguaçu basin, Paraná state, Brazil. *Neotropical Ichthyology* 7(2): 199–204. doi: [10.1590/S1679-62252009000200010](https://doi.org/10.1590/S1679-62252009000200010)
- Pavanelli, C.S. and C.A.M. Oliveira. 2009. A redescription of *Astyanax gymnodontus* (Eigenmann, 1911), new combination, a polymorphic characid fish from the Rio Iguaçu basin, Brazil. *Neotropical Ichthyology* 7(4): 569–578. doi: [10.1590/S1679-62252009000400003](https://doi.org/10.1590/S1679-62252009000400003)
- Pellicice, F.M. and A.A. Agostinho. 2009. Fish fauna destruction after the introduction of a non-native presator (*Cichla kelberi*) in a Neotropical reservoir. *Biological Invasions* 11(8): 1789–1801. doi: [10.1007/s10530-008-9358-3](https://doi.org/10.1007/s10530-008-9358-3)
- Pereira, E.H.L. 2005. Resurrection of *Pareiorhaphis* Miranda Ribeiro, 1918 (Teleostei: Siluriformes: Loricariidae), and description of a new species from the Rio Iguaçu basin, Brazil. *Neotropical Ichthyology* 3(2): 271–276. doi: [10.1590/S1679-62252005000200004](https://doi.org/10.1590/S1679-62252005000200004)
- Reis, R.E., J.S. Albert, F. Di Dario, M.M. Mincarone, P. Petry and L.A. Rocha. 2016. Fish biodiversity and conservation in South America. *Journal of Fish Biology* 89(1): 12–47. doi: [10.1111/jfb.13016](https://doi.org/10.1111/jfb.13016)
- Ribeiro, A.C. 2006. Tectonic history and the biogeography of the freshwater fishes from the coastal drainages of eastern Brazil: an example of faunal evolution associated with a divergent continental margin. *Neotropical Ichthyology* 4(2): 225–246. doi: [10.1590/S1679-62252006000200009](https://doi.org/10.1590/S1679-62252006000200009)
- Říčan, O., L. Piálek, A. Almirón and J. Casciotta. 2011. Two new species of *Australoheros* (Teleostei: Cichlidae), with notes on diversity of the genus and biogeography of the Río de la Plata basin. *Zootaxa* 2982: 1–26. <http://www.mapress.com/zootaxa/2011/f/zt02982p026.pdf>
- Severi, W. and A.A.M. Cordeiro. 1994. *Catálogo de peixes da bacia do rio Iguaçu*. Curitiba: IAP GTZ. 118 pp.

- Suárez, Y.R., S.B. Valério, K.K. Tondato, A.C. Florentino, T.R.A. Felipe, L.Q.L. Ximenes and L.S. Lorenço. 2007. Fish species diversity in headwaters streams of Paraguay and Paraná basins. *Brazilian Archives of Biology and Technology* 50(6): 1033–1042. doi: [10.1590/S1516-89132007000700014](https://doi.org/10.1590/S1516-89132007000700014)
- Vari, R.P. and L.R. Malabarba. 1998. Neotropical Ichthyology: an overview; pp. 1–11, in: L.R. Malabarba and R.P. Reis (eds.). *Phylogeny and classification of Neotropical fishes*. Porto Alegre: EDIPUCRS.
- Vera Alcaraz, H.S., C.S. Pavanelli and V.A. Bertaco. 2009. *Astyanax jordanensis* (Ostariophysi: Characidae), a new species from the Rio Iguaçú basin, Paraná, Brazil. *Neotropical Ichthyology* 7(2): 185–190. doi: [10.1590/S1679-62252009000200008](https://doi.org/10.1590/S1679-62252009000200008)
- Wosiacki, W.B. and M. de Pinna. 2008a. A new species of the Neotropical catfish genus *Trichomycterus* (Siluriformes: Trichomycteridae) representing a new body shape for the family. *Copeia* 2008(2): 273–278. doi: [10.1643/CI-06-237](https://doi.org/10.1643/CI-06-237)
- Wosiacki, W.B. and M. de Pinna. 2008b. *Trichomycterus igobi*, a new catfish species from the Rio Iguaçú drainage: the largest head in Trichomycteridae (Siluriformes: Trichomycteridae). *Neotropical Ichthyology* 6(1): 17–23. doi: [10.1590/S1679-6225008000100003](https://doi.org/10.1590/S1679-6225008000100003)
- Wosiacki, W.B. and J.C. Garavello. 2004. Five new species of *Trichomycterus* from the Rio Iguaçú (Rio Paraná basin), southern Brazil (Siluriformes: Trichomycteridae). *Ichthyological Exploration of Freshwaters* 15(1): 1–16.
- Zawadzki, C.H., E. Renesto and L.M. Bini. 1999. Genetic and morphometric analysis of three species of the genus *Hypostomus* Lacépède, 1803 (Osteichthyes: Loricariidae) from the Rio Iguaçú basin (Brazil). *Revue Suisse de Zoologie* 106(1): 91–105.

**Author contributions:** AF and WJG collected data, verified the specimens, and wrote the text. GCD verified the specimens and wrote the text. EVRG filtered data and wrote the text.

**Received:** 15 April 2016

**Accepted:** 30 September 2016

**Academic editor:** Mariangeles Arce H.

## APPENDIX

**Table A1.** Geographic descriptions of the 30 sampling sites, with a list of species captured in each of them.


Sample point	Locality	County	Geographic Coordinates	Basin	Species caught
1	Unnamed river	Pinhão	25°37'39.2" S 051°28'23.6" W	Jordão	<i>Astyanax bifasciatus</i> , <i>A. jordanensis</i> , <i>Geophagus</i> aff. <i>brasiliensis</i> , <i>Phalloceros harpagos</i> , <i>Rhamdia voulezi</i> , <i>Trichomycterus davis</i> and <i>T. igobi</i>
2	Unnamed river	Pinhão	25°44'35.5" S 051°35'14.5" W	Jordão	<i>Astyanax bifasciatus</i> and <i>Trichomycterus davis</i>
3	Santa Terezinha River	Pinhão	25°44'1.5" S 051°36'0.4" W	Jordão	<i>Astyanax bifasciatus</i> , <i>Geophagus</i> aff. <i>brasiliensis</i> , <i>Hoplias</i> sp. and <i>Trichomycterus davis</i>
4	Tapejara River	Pinhão	25°41'22.3" S 051°44'33.2" W	Jordão	<i>Rhamdia branneri</i> and <i>Trichomycterus davis</i>
5	Unnamed river	Reserva do Iguaçú	25°48'40.6" S 052°00'9.8" W	Jordão	<i>Astyanax jordanensis</i> and <i>Trichomycterus davis</i>
6	Unnamed river	Reserva do Iguaçú	25°52'13.9" S 051°54'30.9" W	Jordão	<i>Astyanax jordanensis</i> and <i>Trichomycterus davis</i>
7	Lageado Capão Cortado River	Reserva do Iguaçú	25°47'6.3" S 051°54'30.4" W	Jordão	<i>Trichomycterus davis</i>
8	Da Arca Brook	Guarapuava	25°33'11.2" S 051°29'25" W	Jordão	<i>Astyanax bifasciatus</i> and <i>Rhamdia branneri</i>
9	Banana Brook	Guarapuava	25°26'32.6" S 051°25'44.4" W	Jordão	<i>Astyanax bifasciatus</i> , <i>Corydoras ehrhardti</i> , <i>Geophagus</i> aff. <i>brasiliensis</i> , <i>Hoplias</i> sp., <i>Phalloceros harpagos</i> , <i>Rhamdia branneri</i> and <i>Trichomycterus davis</i>
10	Unnamed river	Guarapuava	25°28'38.2" S 051°39'15.5" W	Jordão	<i>Astyanax bifasciatus</i> and <i>Geophagus</i> aff. <i>brasiliensis</i>
11	Unnamed river	Guarapuava	25°21'34" S 051°21'27.7" W	Jordão	<i>Astyanax bifasciatus</i> and <i>Trichomycterus davis</i>
12	Unnamed river	Guarapuava	25°28'4.6" S 051°14'36.3" W	Jordão	<i>Phalloceros harpagos</i> , <i>Rhamdia branneri</i> , <i>Trichomycterus davis</i> , <i>T. igobi</i> and <i>T. stawiarski</i>
13	Unnamed river	Inácio Martins	25°28'26.7" S 051°14'14" W	Jordão	<i>Astyanax bifasciatus</i> , <i>A. dissimilis</i> and <i>Geophagus</i> aff. <i>brasiliensis</i>
14	Lajeado Brook	Inácio Martins	25°30'5.3" S 051°10'9.7" W	Jordão	<i>Astyanax bifasciatus</i> , <i>Characidium travassosi</i> and <i>Trichomycterus davis</i>
15	Unnamed river	Mallet	25°5'0'10.1" S 050°52'21.9" W	Areia	<i>Astyanax bifasciatus</i> , <i>Bryconamericus ikaa</i> , <i>Corydoras ehrhardti</i> , <i>Hyphessobrycon reticulatus</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> , <i>Phalloceros harpagos</i> and <i>Rineloricaria maacki</i>
16	Unnamed river	Mallet	25°56'24.3" S 051°00'31" W	Areia	<i>Astyanax bifasciatus</i> , <i>A. minor</i> , <i>Bryconamericus ikaa</i> , <i>Corydoras ehrhardti</i> , <i>Hyphessobrycon reticulatus</i> , <i>Oligosarcus longirostris</i> , <i>Phalloceros harpagos</i> , <i>Rhamdia voulezi</i> and <i>Rineloricaria maacki</i>
17	Louro River	Cruz Machado	25°51'1.0" S, 050°50'30.3" W	Areia	<i>Characidium travassosi</i> , <i>Cnesterodon carnegiei</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>
18	Pedrinho River	Cruz Machado	25°56'48.2" S, 051°02'45.9" W	Areia	<i>Astyanax jordanensis</i> , <i>Characidium travassosi</i> , <i>Heptapterus</i> sp., <i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>

Continued



**Table A1.** *Continued.*

Sample point	Locality	County	Geographic Coordinates	Basin	Species caught
19	Jacutinga River	Cruz Machado	25°56'12.7" S 051°13'46.6" W	Areia	<i>Astyanax bifasciatus</i> , <i>Bryconamericus ikaa</i> , <i>Bryconamericus</i> sp., <i>Jenynsia diphyes</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> , <i>Trichomycterus davis</i> and <i>T. stawiarski</i>
20	Unnamed river	Cruz Machado	25°58'27.4" S 051°22' 13.7" W	Areia	<i>Astyanax bifasciatus</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> , <i>Rineloricaria maacki</i> and <i>Trichomycterus davis</i>
21	Unnamed river	Cruz Machado	25°56'47.8" S 051°23' 44.9" W	Areia	<i>Ancistrus abilhoai</i> , <i>Astyanax bifasciatus</i> , <i>A. dissimilis</i> , <i>A. minor</i> , <i>Bryconamericus ikaa</i> , <i>Characidium</i> sp. 1, <i>Oligosarcus longirostris</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> , " <i>Pariolius</i> " sp., <i>Rhamdia voulezi</i> , <i>Trichomycterus davis</i> , <i>T. igobi</i> and <i>T. papilliferus</i>
22	Pimpãozinho River	Pinhão	25°50'58.3" S, 051°27' 26.5" W	Areia	<i>Jenynsia diphyes</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>
23	Unnamed river	Pinhão	25°54'45.9" S, 051°34'49.6" W	Areia	<i>Phalloceros harpagos</i>
24	Unnamed river	Pinhão	25°56'7.0" S, 051°33'4.6" W	Areia	<i>Astyanax bifasciatus</i> , <i>Heptapterus</i> sp., <i>Pareiorhaphis</i> cf. <i>parmula</i> , <i>Trichomycterus davis</i> and <i>T. papilliferus</i>
25	Unnamed river	Pinhão	25°50'24.6" S, 051°32'1.1" W	Areia	<i>Astyanax jordanensis</i> and <i>Trichomycterus davis</i>
26	Unnamed river	Inácio Martins	25°31'34.6" S, 051°10'45.6" W	Areia	<i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>
27	Unnamed river	Inácio Martins	25°32'37.4" S, 051°12'42.7" W	Areia	<i>Astyanax bifasciatus</i> , <i>Cnesterodon carnegiei</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>
28	Unnamed river	Inácio Martins	25°35'20.1" S, 051°06'59.8" W	Areia	<i>Astyanax bifasciatus</i> and <i>Trichomycterus davis</i>
29	Unnamed river	Inácio Martins	25°35'16.3" S, 051°06'19.9" W	Areia	<i>Astyanax bifasciatus</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> and <i>Trichomycterus davis</i>
30	Unnamed river	Inácio Martins	25°37'12.6" S, 051°07'21.2" W	Areia	<i>Astyanax bifasciatus</i> , <i>Geophagus</i> aff. <i>brasiliensis</i> , <i>Pareiorhaphis</i> cf. <i>parmula</i> , <i>Trichomycterus davis</i> and <i>T. stawiarski</i>



**Figure A1.** Sampling sites in the Jordão River basin (labelled in accordance to Table A1).





**Figure A2.** Sampling sites within the Areia River basin (labelled in accordance to Table A1).