



# Bryophytes of the cloud forest of Pico do Marumbi State Park, Paraná, Brazil

Emanuelle L. dos Santos,<sup>1</sup> Dimas M. do Carmo,<sup>2</sup> Denilson F. Peralta<sup>2</sup>

<sup>1</sup> Universidade Federal do Paraná, Setor de Ciências Biológicas, Departamento de Botânica, Caixa Postal 19031, 81531-980, Curitiba, PR, Brazil.

<sup>2</sup> Instituto de Botânica, Núcleo de Pesquisa em Briologia, Caixa Postal 68041, 04045-972 São Paulo, SP, Brazil.

**Corresponding author:** Emanuelle L. dos Santos, [emanuellelais.s@gmail.com](mailto:emanuellelais.s@gmail.com)

## Abstract

Bryophyte diversity in tropical forests is closely related to elevation. A survey was undertaken of the bryoflora of a poorly known cloud forest environment from 1,000 to 1,500 m.a.s. in Pico do Marumbi State Park, state of Paraná, Brazil. The aim was to analyze the geographical range of the bryoflora and significance of this elevational zone to bryophyte diversity. We found 364 bryophyte species, including 187 species of mosses, 175 liverworts and two hornworts. This diversity represents 40% of the bryoflora of Paraná and 24% of that of Brazil. Forty-one Brazilian endemic species were found, representing a highly relevant feature for the preservation of the area. These results emphasize the importance of Pico do Marumbi State Park to the conservation of bryophyte diversity and to understanding the species of the state of Paraná, as well as demonstrating the importance of bryophyte surveys in the Atlantic Forest of Southeast Brazil.

## Keywords

Mosses; liverworts; Neotropics; cloud forest.

**Academic editor:** Denise Pinheiro da Costa | Received 8 September 2016 | Accepted 26 June 2017 | Published 15 December 2017

**Citation:** dos Santos EL, do Carmo DM, Peralta DF (2017) Bryophytes of the cloud forest of Pico do Marumbi State Park, Paraná, Brazil. *Check List* 13 (6): 959–986. <https://doi.org/10.15560/13.6.959>

## Introduction

Bryophytes are avascular terrestrial plants whose sexual reproduction is dependent on water because of their flagellated anterozoids. In their lifecycle, the gametophytic generation (haploid) is predominant over their sporophytic generation (diploid) (Gradstein et al. 2001). They are the second largest group of terrestrial plants (Frahm 2003), and currently, they comprise 3 divisions, the Anthocerothophyta (hornworts), the Marchantiophyta (liverworts), and the Bryophyta (mosses) (Shaw and Goffinet 2009).

There are 1548 species of bryophytes reported from Brazil, with 541 species occurring in the state of Paraná (Yano 2013, 2014, Ristow et al. 2015), 570 species in the state of Rio Grande do Sul, and 542 in the state of Santa Catarina (Costa and Peralta 2017).

The first studies of bryophytes of Paraná were those of Angely (1961, 1965, 1968), who produced a list of taxa. Kummrow and Prevedello (1982) published a list of samples deposited in the herbarium MBM (Municipal Botanical Museum, Curitiba). Hirai et al. (1998) undertook a floristic survey in a small secondary *Araucaria* forest, and Yano and Colletes (2000) investigated the

bryophytes of an area of Atlantic Forest. Recently Yano (2013, 2014) published a compilation of the literature on bryophyte taxa of t Paraná and listed 706 taxa. Since then, Ristow et al. (2015) added 203 species records for the state.

Studies of bryophytes in Brazilian cloud forests remain scarce and are non-existent for the state of Paraná. The studies of Santos and Costa (2010) and Costa et al. (2015) represent the only bryophyte investigations conducted above 1000 m above sea level in Brazil. These authors indicated that bryophytes were the richest and most diverse above this elevation. The results of several other authors are consistent with this finding (Enroth 1990, Frahm 1990, Gradstein 1991, Kessler 2000, Santos and Costa 2010, Costa et al 2015).

Pico do Marumbi State Park (PMSP) is located in southern Brazil, and is 1 of the 10 centers of diversity and endemism in tropical America (Tan and Pócs 2000), and thus, it offers an excellent environment for the study of bryophyte diversity. PMSP is completely within the Atlantic Forest biome and comprises Dense High Mountain Ombrophylous Forest phytophysiognomy. These unique high-elevation montane forests possess high endemism and, in general, are subjected to minimal anthropogenic disturbance due their difficult access (Scheer et al. 2011).

The forest vegetation of PMSP decreases in size with increasing elevation and because soil is replaced by rocky outcrops near 1400–1500 m elevation. At elevations of 1000–1400 m the vegetation in the park is cloud (IBGE 2012). forest. The term “cloud forest” derives of the high condensation of evaporated water in the air, which forms a haze and maintains high precipitation. The features of cloud forest correspond to an arboreal mesophanerophytic formation (reaching >30 m), which is typically located on the summits of higher mountains (1000–1500 m elevation) (Roderjan et al. 2002, Scheer et al. 2011).

Upper Montane Forests and High-Altitude Meadows are fundamental to the conservation of bryophyte diversity in the Atlantic Forest, and Santos and Costa (2010) and Costa et al. (2015) emphasized the importance of their protection and conservation. The objective of our study is to record and analyze the floristic composition of bryophytes occurring in the cloud forest of Pico do Marumbi State Park, as well as to evaluate the geographic distributions of bryophyte species found.

## Methods

**Study area.** PMSP (Fig. 1) is located between the cities of Morretes, Quatro Barras, and Piraquara, and covers an area of 8745 ha (IAP 2012). It is within the Atlantic Forest domain (IAP 1996) and includes areas of Dense Ombrophylous Submountain Forest, Dense Mountain Ombrophylous Forest, and Dense High Mountain Ombrophylous Forest in good states of preservation. The floristic composition and tree layer structure varies along the elevational gradient (Figs 2–7).

**Floristic surveys.** Sampling was permitted by the Instituto Ambiental do Paraná (IAP), and we followed the methodology of Frahm (2003). Random collects were performed using pre-existent trails (approximate coordinates plotted in Fig. 1) with an effort made to cover the range of habitats and microhabitats available in cloud forests between 1000 and 1400 m above sea level. Eleven expeditions were taken from March 2014 to June 2015. The samples were deposited in the Maria Eneyda P. K. Fidalgo (SP) herbarium with duplicates in Departamento de Botânica da Universidade Federal do Paraná (UPCB) herbarium.

**Bryophyte identification.** Bryophytes were identified from the preparation of slides and their observation with stereo- and optical microscopes using specialized literature according to each family (Frahm 1991, Sharp et al. 1994, Yano and Carvalho 1995, Buck 1998, Gradstein et al. 2001, Gradstein and Costa 2003, Yano and Peralta 2009, Yano and Peralta 2011) and comparisons with herbarium specimens.

The classification systems follow Söderström et al. (2016) for Anthocerotophyta and Marchantiophyta and Goffinet et al. (2009) for Bryophyta. Author abbreviations for the specific epithets followed Brummitt and Powell (1992). The species list provided here is in alphabetical order by division, family and species. The geographic distributions in Brazil follow the standardization of Valente and Porto (2006), who classify species as Endemic, Rare, Moderate or Wide, while the occurrences in states followed Costa and Peralta (2015) and Gradstein and Costa (2003).


## Results

We found 364 bryophyte species, including 187 species of mosses (35 families, 85 genera), 175 liverworts (22 families, 58 genera) and two hornworts (1 family, 2 genera). The richest families of Marchantiophyta were Lejeuneaceae with 66 species (18%) and Lepidoziaceae with 16 species (4%). For Bryophyta, the most speciose families were Leucobryaceae with 19 species (5%), Pilotrichaceae with 16 species, Sematophyllaceae with 15 species, Dicranaceae with 14 species, Fissidentaceae and Sphagnaceae with 13 species each, Orthotrichaceae with 10 species and Bryaceae with nine species (Table 1).

Here we provide, in alphabetical order, brief characterizations of the 41 Brazilian endemic species found in PMSP. This list follows the characterization of cloud forest by Costa et al. (2015), with species tolerant to wide environmental variation, such as low temperatures, high humidity, and high light intensity.

### *Aptychopsis pyrrophylla* (Müll. Hal.) Wijk & Margad.

Comments: distinguished by ovate leaves with cuspidate apices. Habitat: corticolous. Geographic distribution: Amazonian and Atlantic Forest at various elevations but more common above 1500 m. Brazilian geographic distribution: AM, BA, ES, MG, RJ, SP.



**Figure 1.** Location map of Pico do Marumbi State Park in Paraná state.

***Atractylocarpus brasiliensis* (Müll. Hal.) R.S. Williams**

Comments: differs from *Atractylocarpus nanus* R.S. Williams, an Andean species, by having a less rugose dorsal surface of costa; shorter exotechial cells; and peristome teeth divided from the base. Habitat: soil. Geographic distribution: Atlantic Forest (high montane). Brazilian geographic distribution: BA, RJ.

***Barbula riograndensis* E.B. Bartram**

Comments: distinguishable from any other known Brazilian species of *Barbula* by costa with dorsal and ventral bands of stereids in cross section, and short, yellow and strongly dentate apiculi. Habitat: soil, rocks and streams. Geographic distribution: Atlantic Forest (lowlands). Brazilian geographic distribution: PR, RS.

***Bazzania heterostipa* (Steph.) Fulford**

Comments: forked, terminal branching scarce and plants without terminal branches may be mistaken with species of *Calypogeia*; the presence of a vitta in the leaves immediately separates this species from *Calypogeia*. Habitat: soil, rocks and corticolous. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: BA, ES, MG, PE, PR, RJ, RS, SC, SP.

***Brachythecium podadelphus* Müll. Hal. (Figure 10)**

Comments: recognized by its medium to large size; leaves approximately 2.0 × 0.8 mm; distinctly different

weak alar cells; dioicous. Habitat: corticolous. Geographic distribution: Atlantic Forest (montane). Brazilian geographic distribution: MG, RJ, SP.

***Campylopus dichrostis* (Mull. Hal.) Paris**

Comments: characterized by leaf lamina reaching the apex; percurrent costa with lamellae; dorsal and ventral stereids in transverse section. Habitat: soil and rocks. Geographic distribution: Atlantic Forest and Cerrado. Brazilian geographic distribution: BA, GO, MG, RJ, RS, SC, SP.

***Campylopus fragilis* (Brid.) Bruch & Schimp.**

Comments: characterized by masses of sessile brood leaves; hyaline basal leaf cells grading evenly into the quadrate median leaf cells; alar cells absent. Habitat: soil, decomposing tree trunks and rocks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: MG, RJ, SC, SP.

***Cheilolejeunea caducifolia* (Gradst. & Schäf.-Verw.) W. Ye & R.L. Zhu**

Comments: distinguished by caducous leaves. Habitat: corticolous. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: BA, ES, MG.

***Cololejeunea manaosensis* (Herzog) O. Yano (Fig. 42)**

Comments: characterized by a smooth cuticle and opaque lobe cells (not transparent). Habitat: epiphyllous.



**Figures 2–7.** Studied area. **2.** Marumbi peaks group. **3–6.** Overview of the Pico do Marumbi State Park. **7.** Taquaral River.

Geographic distribution: Amazon and Atlantic Forest. Brazilian geographic distribution: AM, MT, PA, SP.

***Cololejeunea microscopica* (Taylor) Schiffn.**

Comments: conical protuberances restricted to the keel and sometimes a few apical cells of the lobe, never exceeding 10  $\mu\text{m}$  in the height. Habitat: epiphyllous and decomposing tree trunks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: SP.

***Cololejeunea papilliloba* (Steph.) Steph.**

Comments: characterized by elongate-ovate leaves with obtuse apexes; perianth elongate-obconoidal, with

well-developed beak; female bracts as long as, or longer than, perianth. Habitat: corticolous. Geographic distribution: Atlantic Forest and Pampa. Brazilian geographic distribution: MG, RS, SP.

***Fissidens pseudoplurisetus* Bordin, Pursell & O. Yano**

Comments: characterized by limbate leaves, limbidia bi- or tri-stratose, unipapillose cells and 1–4 perichaetial sporophytes. Habitat: corticolous. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: SP.

***Fissidens yanoae* Pursell.**

Comments: characterized by terminal sporophytes

with short seta and long perichaetial leaves. Habitat: corticolous. Geographic distribution: Atlantic Forest (montane). Brazilian geographic distribution: RS, SP.

***Holomitrium nitidum* Herzog.**

Comments: characterized by elongate laminal cells and long perichaetial leaves exceeding the capsule (Herzog 1925). Habitat: corticolous. Geographic distribution: Atlantic Forest (submontane and montane). Brazilian geographic distribution: ES, MG, PR, RJ, SC, SP.

***Isotachis inflata* Steph.**

Comments: recognized by bordered leaves. Habitat: wet soil. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: ES, PR, RJ.

***Leiomela bartramioides* (Hook. f.) Paris (Fig. 9)**

Comments: smaller stem and perichaetial leaves, 5 and 15 mm, respectively, usually half the size of *Leiomela piligera* (Hampe); always found with sporophyte; globose capsule present totally covered with leaves. Habitat: corticolous, generally in bracken fern-like caudex. Geographic distribution: Atlantic Forest (submontane and high montane). Brazilian geographic distribution: MG, RJ, SP.

***Lejeunea cristulata* (Steph.) E. Reiner & Goda**

Comments: characterized by perianth keel without lacinia, occasionally with cilia; plants 0.7–0.9 mm wide; lobule inflated throughout, free margin involute, apex tooth cell 20–25 µm long. Habitat: corticolous. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: BA, ES, MG, PE, PR, RJ, RS, SC, SP.

***Lejeunea oligoclada* Spruce**

Comments: at first glance resembles a small species of *Cheilolejeunea* due to dull greenish color, long lobule tooth, and thick leaf cell walls; also resembles *Lejeunea phyllobola* Nees & Mont., but the latter is larger (0.6–1.1 mm wide), and with the leaf lobe apex widely rounded; plants pale and yellowish or light green. Habitat: corticolous and decomposing tree trunks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: AL, BA, ES, MG, PE, PR, RJ, SC, SP.

***Lepidopilidium brevisetum* (Hampe) Broth.**

Comments: distinguished by leaves with acute apexes, margin serrate at the apex and entire toward the base; costa reaching approximately ½ of the leaf length. Habitat: corticolous. Geographic distribution: Atlantic Forest (lowlands to montane) Brazilian geographic distribution: AL, ES, MG, RJ, RS, SC, SP.

***Lepidopilum caudicaule* Müll. Hal.**

Comments: distinguished by serrulate leaf margin at the apex and entire to sinuous toward the base. Habitat: moist rocks. Geographic distribution: Atlantic Forest (lowlands to high montane). Brazilian geographic distribution: PE, RJ, RS, SC.

***Leptodontium viticulosoides* (P. Beauv.) Wijk & Margad. (Fig. 30)**

Comments: characterized by serrate leaves with small laminal papillae; generally hemispherical and homogeneous spores. Habitat: soil, corticolous, decomposing tree trunks and rocks. Geographic distribution: Atlantic Forest (lowlands to high montane). Brazilian geographic distribution: BA, ES, MG, PR, RJ, RS, SC, SP.

***Leskeodon aristatus* (Geh. & Hampe) Broth. (Fig. 13)**

Comments: characterized by complanate, elliptical to orbicular oblong leaves, slightly revolute when dry, oval-lanceolate, 2.0–3.0 mm, margin entire or denticulate above and apex apiculate. Habitat: corticolous, sometimes epiphyllous. Geographic distribution: Atlantic Forest (submontane and high montane). Brazilian geographic distribution: MG, PR, RJ, RS, SC, SP.

***Leucoloma trifforme* (Mitt.) A. Jaeger**

Comments: characterized by predominantly rectangular to elongated, obliquely-oriented alar cells. Habitat: rocks. Geographic distribution: Atlantic Forest (lowlands to montane). Brazilian geographic distribution: ES, PR, RJ, SP.

***Metzgeria bahiensis* Schiffn.**

Comments: similar to *M. convoluta* Steph., but differs by costa formed by 4–6 rows of ventral epidermal cells; verrucous cuticle; male branch smaller than female branch; external development of female casing absent, stalk vegetative; *M. brasiliensis* differs by costa formed by 2 or 3 rows of ventral epidermal cells; 10–19 lamina formed by cells; occasional rhizoids on ventral surface of lamina; 6–7(–8) medulla cells; cuticle verrucous; marginal cells differentiated; male branch with rhizoids; external development of the female casing absent; stalk vegetative. Habitat: corticolous. Geographic distribution: Atlantic Forest and Pampa. Brazilian geographic distribution: BA, RS, SP.

***Metzgeria brasiliensis* Schiffn.**


Comments: similar to *Metzgeria aurantiaca* Steph., *Metzgeria convoluta* Steph. and *Metzgeria cratoneura* Schiffn. by having cells of verrucous lamina and medulla with thickened walls. *Metzgeria aurantiaca* differs by possessing rhizoids on the ventral surface of the lamina and on the margin, non-differentiated margin of the cells and male stem smaller than the female; *Metzgeria convoluta* differs by costa formed by 4 (–6) rows of ventral epidermal cells, rhizoids on uniform margin, male stem smaller than the female stem and male branch without rhizoids. Habitat: corticolous, epiphyllous and rocks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: AL, BA, PR, RJ, RS, SC, SP.

***Paranapiacabaea paulista* W.R. Buck & Vital.**

Comments: can only be identified with sporophyte; sterile resembles *Donnellia*, when fertile resembles *Pterogoniopsis cylindrical* Müll. Hal., but this latter genus has a higher basal membrane, various super-alar cells and leaves with a flat edge. Habitat: corticolous. Geographic distribution: Atlantic Forest (submontane to montane). Brazilian geographic distribution: PR, SP.



**Figures 8–22.** 8. *Dendroceros crispus* (Sw.) Nees. 9. *Leiomela bartramioides* (Hook. f.) Paris. 10. *Brachythecium poadelphus* Müll. Hal. 11. *Rosulabryum densifolium* (Brid.) Ochyra. 12. *Syrrhopodon prolifer* Schwägr. 13. *Leskeodon aristatus* (Geh. & Hampe) Broth. 14. *Holomitrium arboreum* Mitt. 15. *Leucoloma serrulatum* Brid. 16. *Fissidens asplenioides* Hedw. 17. *Chryso-hypnum diminutivum* (Hampe) W.R. Buck. 18. *Hypopterygium tamariscinum* (Hedw.) Brid. 19. *Orthostichella pachygastrella* (Müll. Hal. ex Ångstr.) B.H. Allen & Magill. 20. *Campylopus heterostachys* (Hampe) A. Jaeger. 21. *Leucobryum albicans* (Schwägr.) Lindb. 22. *Toloxia imponderosa* (Taylor) W.R. Buck.



**Figures 23–37.** **23.** *Porotrichum substriatum* (Hampe) Mitt. **24.** *Thamnomalia glabella* (Hedw.) S. Olsson et al. **25.** *Macrocoma tenuis* (Hook. f. & Grev.) Vitt. **26.** *Schlotheimia tecta* Hook. f. & Wilson. **27.** *Lepidopilum muelleri* (Hampe) Mitt. **28.** *Thamniopsis langsdorffii* (Hook. f.) W.R. Buck. **29.** *Polytrichum commune* Hedw. **30.** *Leptodontium viticulosoides* (P. Beauv.) Wijk & Margad. **31.** *Acroporium estrellae* (Müll. Hal.) W.R. Buck & Schäf.-Verw. **32.** *Sematophyllum galipense* (Müll. Hal.) Mitt. **33.** *Sphagnum palustre* L. **34.** *Aneura pinguis* (L.) Dumort. **35.** *Isotachis aubertii* (Schwägr.) Mitt. **36.** *Odontoschisma falcifolium* Steph. **37.** *Frullania brasiliensis* Raddi.



**Figures 38–52.** **38.** *Syzygiella anomala* (Lindenb. & Gott.) Steph. **39.** *Chiloscyphus martianus* (Nees) J.J. Engel & R.M. Schust. **40.** *Bryopteris filicina* (Sw.) Nees. **41.** *Cheilolejeunea clausa* (Nees & Mont.) R.M. Schuster. **42.** *Cololejeunea manaoensis* (Herzog) O.Yano. **43.** *Diplasiolejeunea brunnea* Steph. **44.** *Lejeunea flava* (Sw.) Nees. **45.** *Microlejeunea epiphylla* Bischl. **46.** *Taxilejeunea pterigonia* (Lehm. & Lindb.) Schiffn. **47.** *Bazzania hookeri* (Lindb.) Trevis. **48.** *Lepidozia inaequalis* Lehm. & Lindb. **49.** *Metzgeria albinea* Spruce. **50.** *Symphyogyna brasiliensis* Nees. **51.** *Plagiochila adiantoides* Dumort. **52.** *Radula recubans* Taylor.



***Polytrichum angustifolium* Mitt.**

Comments: distinguished by pyriform lamella of cells in cross section, vaginate sometimes with orange flat-edged sheath. Habitat: soil, rocks and humus. Geographic distribution: Atlantic Forest (submontane to high montane). Brazilian geographic distribution: ES, MG, PR, RJ, RS, SC, SP.

***Porotrichum thieleanum* (Müll. Hal.) Mitt.**

Comments: characterized by stipe with squarrose leaves with recurved margin and stem leaves orbicular. Habitat: corticolous. Geographic distribution: Atlantic Forest (submontane). Brazilian geographic distribution: MG, PR, RJ, RS, SP.

***Prionolejeunea scaberula* (Spruce) Steph.**

Comments: margins of caducous leaves can be strongly crenate with marginal cells evolving to rhizoids. Habitat: corticolous, epiphyllous, and rocks. Geographic distribution: Amazonian and Atlantic Forest. Brazilian geographic distribution: AM, BA, MA, SP.

***Radula brasilica* Yamada**

Comments: characterized by recurved free margin of lobules; leaf cells with large trigones. Habitat: decomposing tree trunks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: SP.

***Rhacocarpus inermis* (Mull. Hal.) Lindb.**

Comments: distinguished by oblong-lanceolate leaves with short acute apexes that are cuspidate or when piliferous with the upper margin not bordered. Habitat: exposed, moist or soil covered rocks. Geographic distribution: Atlantic Forest (montane to high montane). Brazilian geographic distribution: ES, MG, RJ, RS, SC.

***Riccardia emarginata* (Steph.) Hell.**

Comments: related to *R. regnelli* but differs by having a thicker axis (9–12 cells thick; 5–7 cells thick in *R. regnelli*) and a subepidermis in the axis (subepidermis lacking in *R. regnelli*). Habitat: soil and corticolous. Brazilian geographic distribution: BA, MG, RJ, SP.

***Riccardia regnellii* (Ångstr.) Hell.**

Comments: related to *R. emarginata*. Habitat: soil, decomposing tree trunks and corticolous. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: BA, ES, MG, MT, PE, RJ, RN, RS, SC, SP.

***Sphagnum divisum* H.A. Crum**

Comments: with 1 or 2 fasciculate branches, large leaves of branches with annular pores commissurally arranged in continuous rows on both surfaces, and leucocyst stem leaves consistently divided with more numerous pores on internal than external surface. Habitat: soil and rocks. Geographic distribution: Atlantic Forest (montane and high montane) and Cerrado. Brazilian geographic distribution: BA, GO, MG, RJ, SC.

***Sphagnum exquisitum* H.A. Crum**

Comments: similar to *S. ramulinum*, but possessing smaller leaves, oval branches, and fibrilous leucocysts

only on upper portions of both stem leaves and branches. Habitat: soil and rocks. Geographic distribution: Atlantic Forest (submontane to high montane). Brazilian geographic distribution: MG, PR, RJ, SP.

***Sphagnum globicephalum* Müll. Hal. ex Warnst.**

Comments: characterized by possessing a simple stem and sparse branches with terminal globular structures formed by groups of imbricate leaves; stem leaves and branches are similar with few or numerous pores on both surfaces. Habitat: dry soil. Geographic distribution: Atlantic Forest (montane to high montane). Brazilian geographic distribution: RJ, SC.

***Sphagnum multiporosum* H.A. Crum**

Comments: characterized by branches in fascicles of three; leaves broad-elliptic, broad-rounded apex, apex smooth, sometimes fringing or bordered by a reabsorption furrow; leucocyst of leaves of branches on outer surface with group of 3 conspicuous pores at adjacent angles; chlorocyst elliptical. Habitat: soil and rocks. Geographic distribution: Atlantic Forest and Pampa. Brazilian geographic distribution: BA, MG, RJ, RS, SP.

***Sphagnum pseudoramulinum* H.A. Crum**

Comments: similar to *S. ramulinum* in Southeast Brazil, however, the pores of the external surface of the stem leaves and branches are large and arranged in a commissural row, leucocysts of stem leaves are divided and secondary branches grow near base. Habitat: soil and rocks. Geographic distribution: Atlantic Forest (high montane). Brazilian geographic distribution: RJ, RS, SP.

***Sphagnum sucrei* H.A. Crum**

Comments: stem leaves large, oblong, with no divided hyaline cells and numerous pores in commissural rows on external surface, but absent on internal surface, contrasting in size and shape with leaves of the branches, which possess a similar pore distribution; the chlorocyst of the leaves of the branches are narrowly trapezoidal in cross-section, more widely exposed on the internal surface. Habitat: rocks. Geographic distribution: Atlantic Forest. Brazilian geographic distribution: MG, RJ.

***Trichocolea argentea* Herzog.**

Comments: characterized by lamina of the leaf with scattered cilia on the surface. Habitat: soil, rocks, and decomposing tree trunks. Geographic distribution: Amazonia, Caatinga, Cerrado, Atlantic Forest, Pampa and Pantanal. Brazilian geographic distribution: PR, RJ, SC, SP.

***Trichosteleum glaziovii* (Hampe) W.R. Buck**

Comments: is distinguished by falcate leaves. Habitat: corticolous and decomposing tree trunks. Geographic distribution: Atlantic Forest (lowlands to high montane). Brazilian geographic distribution: MG, PE, PR, RJ, SC, SP.

**Table 1.** List of bryophyte species found in cloud forest of Pico do Marumbi State Park with their geographic distribution in Brazil. Dist. Brazil = distribution in Brazil, EN = Endemic, RA = Rare, MO = Moderate and AM = Wide. Voucher information. DFP = Denilson Fernandes Peralta; DMC = Dimas Marchi do Carmo; ELS = Emanuele Lais dos Santos; GTF = G.T. Ferreira; LAA = Leandro de Almeida Amelio; RR = Rony Ristow; and WTF = W.T. Ferreira.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<b>Anthocerothphyta</b>				
Dendrocerotaceae				
<i>Dendroceros crispus</i> (Sw.) Nees (Swartz 1788: 146, Nees 1846: 581) (Fig. 8)	MO	BA, MG, PR, RJ, SP	DFP 16094 (SP452520)	25°26'11"S, 48°55'14"W
<i>Nothoceros minarum</i> (Nees) J.C. Villarreal (Nees 1838: 340, Villarreal A. 2014: 34)	MO	PE, RJ, RS, SP	DFP 16016 (SP452439)	25°26'11"S, 48°55'14"W
<b>Bryophyta</b>				
Bartramiaceae				
<i>Bretulia subtomentosa</i> (Hampe) A. Jaeger (Hampe 1872: 49, Jaeger 1875: 94)	MO	ES, MG, PR, RJ, RS, SC, SP	DFP 17925 (SP461467)	25°30'55"S, 48°58'58"W
<i>Leiomela bartramioides</i> (Hook. f.) Paris (Hooker 1836: 71, Paris 1905: 132) (Fig. 9)	RA/EN	MG, RJ, SP	DFP 16120 (SP 452546)	25°26'11"S, 48°55'14"W
<i>Philonotis hastata</i> (Duby) Wijk & Argent (Duby 1846: 132, Wijk & Margadant 1959: 74)	AM	AM, BA, CE, GO, MA, MG, MS, MT, PA, PI, PR, RJ, RO, RS, SP	DFP 15946 (SP452368)	25°26'55"S, 48°54'54"W
<i>Philonotis sphaerocarpa</i> (Hedw.) Brid. (Hedwig 1801: 197, Bridel 1827: 25)	MO	AM, BA, CE, MT, RJ, SC, SP	ELS 379 (SP477549)	25°15'57"S, 48°32'42"W
Brachytheciaceae				
<i>Brachythecium poadelphus</i> Müll. Hal., 1901 (Fig. 10)	RA/EN	MG, RJ, SP	DFP 16823 (SP460357)	25°26'15"S, 48°55'09"
<i>Meteorioidium remotifolium</i> (Müll. Hal.) Manuel (Müller 1846: 216, Manuel 1977: 49)	AM	BA, ES, GO, MG, MT, PB, PE, PR, RJ, RR, RS, SC, SP	DFP 16804 (SP460338)	25°27'00"S, 48°55'05"W
<i>Squamidium brasiliense</i> (Hornsch.) Broth. (Horschuch 1840: 52, Brotherus 1906: 809)	MO	BA, ES, MG, PR, RJ, RS, SC, SP	DFP 16819 (SP460353)	25°26'15"S, 48°55'09"W
<i>Squamidium leucotrichum</i> (Taylor) Broth. (Taylor 1848: 196, Brotherus 1906: 809)	AM	AC, AL, AM, BA, CE, ES, MG, PA, PE, PR, RJ, RO, RR, RS, SC, SP	WTF 35 (SP437089)	25°27'14"S, 48°55'12"W
<i>Squamidium nigricans</i> (Hook. f.) Broth. (Hooker 1822: 64, Brotherus 1906: 808)	MO	AP, CE, ES, PE, PR, RJ, RS, SC	DFP 16249 (SP454827)	25°26'11"S, 48°55'14"W
<i>Zelometeorium patens</i> (Hook. f.) Manuel (Hooker 1818: 56, Manuel 1977: 116)	MO	ES, MG, MT, RJ, SP	DFP 16254 (SP454832)	25°26'11"S, 48°55'14"W
<i>Zelometeorium patulum</i> (Hedw.) Manuel (Hedwig 1801: 279, Manuel 1977: 116)	AM	AC, AL, AM, AP, BA, CE, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RO, RR, RS, SC, SP, TO	RR 2852 (SP438301)	25°27'14"S, 48°55'12"W
Bryaceae				
<i>Brachymerium patulum</i> (Müll. Hal.) Schimp. (Müller 1851: 579, Schimper in Bescherelle 1872: 195)	RA	ES, MG, SP	DFP 15991 (SP452413)	25°26'55"S, 48°54'54"W
<i>Bryum alpinum</i> Huds. ex With., 1801	RA	RS, SP	ELS 260 (SP459068)	25°30'49"S, 48°59'20"W
<i>Bryum limbatum</i> Müll. Hal., 1851	MO	DF, ES, MG, MS, PR, RJ, RS, SC, SP	RR s.n. (SP462188)	25°26'57"S, 48°54'50"W
<i>Rhodobryum roseolum</i> (Müll. Hal.) Paris (Müller 1879: 287, Paris 1898: 1120)	MO	RJ, SC, SP	ELS 539 (SP477711)	25°18'20"S, 48°35'09"W
<i>Rhodobryum roseum</i> (Hedw.) Limpr. (Hedwig 1801: 194-195, Limpricht 1892: 445)	RA	PE, RJ, RN	ELS 263 (SP459071)	25°30'49"S, 48°59'20"W
<i>Rhodobryum subverticillatum</i> Broth., 1924	MO	BA, MG, PA, PE, RJ, SC, SP	DFP 17921 (SP461463)	25°30'55"S, 48°58'58"W
<i>Rosulabryum billardieri</i> (Schwär) J.R. Spence (Schwägrichen 1826: 76, Spence 1996: 223)	AM	AM, BA, DF, ES, GO, MA, MG, MS, MT, PA, PE, PR, RJ, RO, RR, RS, SC, SP	DFP 16000 (SP452422)	25°26'11"S, 48°55'14"W
<i>Rosulabryum densifolium</i> (Brid.) Ochyra (Bridel 1827: 855, Ochyra 2003: 162) (Fig. 11)	AM	BA, DF, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 16240 (SP454818)	25°26'11"S, 48°55'14"W
<i>Rosulabryum huillense</i> (Welw. & Duby) Ochyra (Welwitsch & Duby in Duby 1872: 220, Ochyra 2003: 162)	RA	MG, RJ, SP	ELS 342 (SP477512)	25°15'57"S, 48°32'42"W
Calymperaceae				
<i>Syrrhopodon elongatus</i> Sull., 1861	MO	BA, PR, RJ, RS, SC, SP	ELS 269 (SP459077)	25°30'50"S, 48°59'16"W
<i>Syrrhopodon gardneri</i> (Hook. f.) Schwägr. (Hooker 1819: 146, Schwägrichen 1824: 110)	MO	AM, BA, GO, MG, MT, RJ	ELS 247 (SP459055)	25°30'49"S, 48°59'20"W
<i>Syrrhopodon incompletus</i> Schwägr., 1824	AM	AC, AM, AP, BA, DF, GO, MG, MS, MT, PA, PE, PR, RJ, RO, RR, SC, SP, TO	DFP 16063 (SP452487)	25°26'11"S, 48°55'14"W
<i>Syrrhopodon ligulatus</i> Mont., 1856	AM	AC, AM, AP, BA, DF, GO, MG, MS, MT, PA, PE, RJ, RO, RR, SP	DFP 16248 (SP454826)	25°26'11"S, 48°55'14"W
<i>Syrrhopodon prolifer</i> Schwägr., 1827 (Fig. 12)	AM	AL, AM, AP, BA, CE, DF, ES, GO, MG, MT, PA, PE, PI, PR, RJ, RO, RS, SC, SE, SP, TO	ELS 264 (SP459072)	25°30'50"S, 48°59'16"W
<i>Syrrhopodon prolifer</i> var <i>cinninnatus</i> (Hampe) W.D. Reese (Hampe 1874: 131, Reese 1978: 200)	MO	BA, DF, ES, GO, MG, MT, PA, PI, PR, RJ, RO, RS, SC, SP	RR s.n. (SP462176)	25°26'49"S, 48°54'53"W
<i>Syrrhopodon prolifer</i> var <i>tenuifolius</i> (Sull.) W.D. Reese (Sullivant 1861: 280, Reese 1978: 199)	MO	BA, ES, MG, RJ, RS, SC, SP	RR s.n. (SP462179)	25°26'49"S, 48°54'53"W
<i>Syrrhopodon tortilis</i> Hampe, 1872	MO	MG, RJ, RS, SC, SP	ELS 294 (SP477464)	25°15'57"S, 48°32'42"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
Catagoniaceae				
<i>Catagonium nitens</i> (Brid.) Cardot (Bridel 1812: 50, Cardot 1915: 469)	RA	AM, MG, SC	DFP 17933 (SP461475)	25°30'55"S, 48°58'58"W
Daltoniaceae				
<i>Daltonia bilimbata</i> Hampe, 1863	RA	MG, PR, RJ, SP	DFP 16267 (SP454846)	25°26'11"S, 48°55'14"W
<i>Daltonia splachnoides</i> (Sm.) Hook. & Tayl. (Smith 1813: 36, Hooker & Taylor 1818: 80)	AM	BA, ES, MG, PE, SP	DFP 17851 (SP461393)	25°30'55"S, 48°58'58"W
<i>Leskeodon aristatus</i> (Geh. & Hampe) Broth. (Geheeb & Hampe 1879, Brotherus 1907: 926) (Fig. 13)	MO/EN	MG, PR, RJ, RS, SC, SP	DFP 16057 (SP452481)	25°26'11"S, 48°55'14"W
Dicranaceae				
<i>Dicranella exigua</i> (Schwägr.) Mitt. (Schwägrichen 1827: 93, Mitten 1869: 30)	MO	AM, ES, MG, MT, PA, PR, RJ, RS, SC, SP	DFP 17743 (SP461284)	25°27'10"S, 48°55'11"W
<i>Dicranella hilariana</i> (Mont.) Mitt. (Montagne 1839: 52, Mitten 1860: 31)	AM	AC, AM, AP, BA, CE, ES, GO, MA, MG, MS, MT, PA, PB, PE, RJ, RO, RR, RS, SC, SP, TO	ELS 285 (SP477455)	25°15'57"S, 48°32'42"W
<i>Dicranella lindigiana</i> Hampe, 1869	MO	CE	DFP 15721 (SP452143)	25°26'55"S, 48°54'54"W
<i>Dicranella varia</i> (Hedw.) Schimp. (Hedwig 1801: 133, Schimper 1855: 13)	MO	CE, GO, SP	ELS 383 (SP477553)	25°15'57"S, 48°32'42"W
<i>Dicranodontium pulcholare</i> Broth subsp. <i>brasiliense</i> (Herzog) J.-P. Frahm (Herzog 1927: 254, Frahm 1997: 196)	RA	RJ	ELS 283 (SP477453)	25°15'57"S, 48°32'42"W
<i>Dicranoloma billardieri</i> (Brid. ex Anon) Paris (Bridel 1802: 214, Paris 1904: 24)	RA	RS, SC	ELS 449 (SP 477620)	25°15'57"S, 48°32'42"W
<i>Holomitrium arboreum</i> Mitt., 1869 (Fig. 14)	AM	AM, BA, ES, GO, MG, MT, PA, PE, PR, RJ, RO, RR, RS, SP	RR 2849 (SP438300)	25°27'14"S, 48°55'12"W
<i>Holomitrium crispulum</i> Mart., 1834	AM	AM, BA, ES, GO, MG, PE, PR, RJ, RS, SC, SP	ELS 354 (SP477524)	25°15'57"S, 48°32'42"W
<i>Holomitrium nitidum</i> Herzog, 1924	MO/EN	ES, MG, PR, RJ, SC, SP	DFP 17665 (SP461206)	25°27'10"S, 48°55'11"W
<i>Holomitrium offersianum</i> Hornsch., 1840	AM	ES, MG, MT, PR, RJ, RS, SC, SP	LAA 59 (SP464601)	25°26'55"S, 48°54'54"W
<i>Leucoloma cruegerianum</i> (Müll. Hal.) A. Jaeger (Müller 1851: 588, Jaeger 1872: 412)	MO	AL, BA, ES, GO, MG, PE, PR, RJ, SP	ELS 287 (SP477457)	25°15'57"S, 48°32'42"W
<i>Leucoloma serrulatum</i> Brid., 1827 (Fig. 15)	MO	AL, BA, DF, ES, MG, PE, PR, RJ, SP	ELS 290 (SP477460)	25°15'57"S, 48°32'42"W
Dicranaceae				
<i>Leucoloma trifforme</i> (Mitt.) A. Jaeger (Mitten 1869: 94, Jaeger 1872: 413)	RA/EN	ES, PR, RJ, SP	ELS 286 (SP477456)	25°15'57"S, 48°32'42"W
Diphysciaceae				
<i>Diphyscium longifolium</i> Griff. (Griffith 1842: 477)	MO	AM, MT, RJ, RO, SP	DFP 15791 (SP452213)	25°26'55"S, 48°54'54"W
Ditrichaceae				
<i>Rhamphidium dicranoides</i> (Müll. Hal.) Paris (Müller 1851: 612, Paris 1905: 164)	RA	RJ	DFP 16073 (SP452497)	25°26'11"S, 48°55'14"W
Fissidentaceae				
<i>Fissidens asplenioides</i> Hedw., 1801 (Fig. 16)	MO	BA, MG, MT, PB, PR, RJ, RS, SC, SP	DFP 15950 (SP452372)	25°26'55"S, 48°54'54"W
<i>Fissidens bryoides</i> Hedw., 1801	RA	PE, PR, SP	RR s.n. (SP462175)	25°26'13"S, 48°55'19"W
<i>Fissidens elegans</i> Brid., 1806	AM	AC, AM, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PB, PE, PI, PR, RJ, RO, RR, RS, SC, SP	ELS 492 (SP477664)	25°16'16"S, 48°33'04"W
<i>Fissidens guianensis</i> Mont., 1840	AM	AC, AL, AM, BA, CE, ES, MA, MG, MS, MT, PA, PB, PE, PI, RO, RR, RS, SP, TO	DFP 17625 (SP461165)	25°26'08"S, 48°55'16"W
<i>Fissidens hornsuschii</i> Mont., 1840	AM	AM, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PB, PE, PI, RJ, RO, RS, SC, SP	DFP 17750 (SP461291)	25°27'10"S, 48°55'11"W
<i>Fissidens oediloma</i> Müll. Hal. ex Broth., 1895	MO	ES, MG, PR, RJ, RS, SC, SP	DFP 16176 (SP452602)	25°26'11"S, 48°55'14"W
<i>Fissidens pellucidus</i> Hornsch., 1840	AM	AC, AM, BA, CE, DF, ES, GO, MG, MT, PA, PB, PE, PR, RJ, RO, RR, RS, SC, SP, TO	DFP 17690 (SP461231)	25°27'10"S, 48°55'11"W
<i>Fissidens pseudoplurisetus</i> Bordin et al., 2011	RA/EN	SP	DFP 15988 (SP452410)	25°26'55"S, 48°54'54"W
<i>Fissidens scariosus</i> Mitt., 1869	AM	BA, ES, MA, MG, PA, PB, PE, PR, RJ, RO, RS, SC, SP	DFP 16238 (SP454816)	25°26'11"S, 48°55'14"W
<i>Fissidens serratus</i> Müll. Hal., 1847	AM	AM, BA, CE, ES, GO, MG, MT, PE, PI, RJ, RS, SC, SP	DFP 16268 (SP454847)	25°26'11"S, 48°55'14"W
<i>Fissidens submarginatus</i> Bruch in Krauss, 1846	AM	AC, AM, BA, CE, DF, ES, GO, MA, MG, MT, PA, PB, PE, PI, RJ, RN, RO, RS, SC, SP	RR s.n. (SP462187)	25°26'57"S, 48°54'50"W
Fissidentaceae				
<i>Fissidens weirii</i> Mitt., 1869	MO	GO, MG, PR, RJ, SC, SP	DFP 15928 (SP452350)	25°26'55"S, 48°54'54"W
<i>Fissidens yanoae</i> Pursell, 1994	RA/EN	RS, SP	DFP 16815 (SP460349)	25°26'15"S, 48°55'09"W
Funariaceae				
<i>Entosthodon bonplandii</i> (Hook. f.) Mitt. (Hooker 1816: 1B, Mitten 1869: 245)	MO	ES, GO, MG, PE, PI, RJ, RS, SC, SP	DFP 16793 (SP460327)	25°27'00"S, 48°55'05"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
Hedwigiaceae				
<i>Braunia subincana</i> Broth., 1900	RA	PR, RS	RR 2835 (SP437116)	25°27'14"S, 48°55'12"W
<i>Hedwigidium integrifolium</i> (P. Beauv.) Dixon (Palisot de Beauvois 1805: 60, Dixon 1939: 369)	MO	ES, MG, PE, PR, RJ, RS, SC	DFP 17868 (SP461410)	25°30'55"S, 48°58'58"W
Hypnaceae				
<i>Chryso-hypnum diminutivum</i> (Hampe) W.R. Buck (Hampe 1847: 86-87, Buck 1984: 182) (Fig. 17)	AM	AC, AM, AP, BA, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RO, RR, RS, SC, SP, TO	DFP 16766 (SP460300)	25°27'00"S, 48°55'05"W
<i>Mittenothamnium reduncum</i> (Mitt.) Ochyra (Mittentr 1869: 509, Ochyra 1999: 256)	MO	ES, MG, PR, RJ, RS, SC	ELS 479 (SP477651)	25°16'16"S, 48°33'04"W
<i>Phyllocladon truncatulus</i> (Müll. Hal.) W.R. Buck (Müller 1851: 263, Buck 1987: 521)	MO	AC, AM, ES, MG, MT, PR, RJ, SC, SP	DFP 16196 (SP454774)	25°26'11"S, 48°55'14"W
<i>Puiggariopsis aurifolia</i> (Mitt.) M. Menzel (Mitt 1869: 509, Menzel 1992: 239)	RA	SP	ELS 527 (SP477699)	25°18'20"S, 48°35'09"W
<i>Taxiphyllum ligulaefolium</i> (E.B. Bartram) W.R. Buck (Bartram 1946: 123, Buck 1990: 42)	AM	AM, GO	ELS 378 (SP477548)	25°15'57"S, 48°32'42"W
Hypopterygiaceae				
<i>Hypopterygium tamariscinum</i> (Hedw.) Brid. (Hedwig 1801: 212-214, Bridel 1851: 8) (Fig. 18)	MO	BA, ES, MG, PR, RJ, RS, SC, SP	DFP 16022 (SP452445)	25°26'11"S, 48°55'14"W
<i>Lopidium concinnum</i> (Hook. f.) Wilson (Hooker 1818: 34, Wilson 1854: 119)	MO	ES, MG, PA, PR, RJ, RS, SC, SP	DFP 16195 (SP454773)	25°26'11"S, 48°55'14"W
Lembophyllaceae				
<i>Orthostichella pachygastralla</i> (Müll. Hal. ex Ångström) B.H. Allen & Magill (Müller 1876: 33, Allen & Magill 2007) (Fig. 19)	MO	MG, PR, RJ, RS, SC, SP	RR 4242 (SP458461)	25°26'57"S, 48°55'53"W
<i>Orthostichella versicolor</i> (Müll. Hal.) B.H. Allen & W.R. Buck (Müller 1850: 127, Allen & Buck 2003: 140)	AM	AM, ES, MG, PE, PR, RJ, RO, RS, SC, SP	DFP 16818 (SP460352)	25°26'15"S, 48°55'09"W
<i>Pilotrichella flexilis</i> (Hedw.) Ångstr. (Hedwig 1801: 234, Ångström 1876: 34)	AM	AP, BA, ES, MG, MS, MT, PE, PR, RJ, RS, SC, SP	ELS 349 (SP477519)	25°15'57"S, 48°32'42"W
Leucobryaceae				
<i>Attractylacarpus brasiliensis</i> (Müll. Hal.) R.S. Williams (Müller 1898: 39, Williams 1928: 110)	RA/EN	BA, RJ	DFP 15841 (SP452263)	25°26'55"S, 48°54'54"W
<i>Campylopus arctocarpus</i> (Hornsch.) Mitt. (Hornschuch 1840: 12, Mitten 1869: 87)	AM	BA, ES, GO, MG, MT, PE, PI, PR, RJ, RS, SC, SP	ELS 295 (SP477465)	25°15'57"S, 48°32'42"W
<i>Campylopus cryptopodioides</i> Broth., 1900	AM	AP, DF, ES, GO, MG, MS, MT, PR, RJ, RS, SC, SP	DFP 15735 (SP452157)	25°26'55"S, 48°54'54"W
<i>Campylopus cuspidatus</i> (Hornsch.) Mitt. (Hornschuch 1840: 13, Mitten 1869: 90)	MO	BA, MG, RJ	RR s.n. (SP462186)	25°26'57"S, 48°54'50"W
<i>Campylopus dichrostis</i> (Müll. Hal.) Paris. (Müller 1900: 255, Paris 1901: 333)	MO/EN	BA, GO, MG, RJ, RS, SC, SP	ELS 319 (SP477489)	25°15'57"S, 48°32'42"W
<i>Campylopus filifolius</i> (Hornsch.) Mitt. (Hornschuch 1840: 12, Mitt 1869: 76)	AM	BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 16165 (SP452591)	25°26'11"S, 48°55'14"W
<i>Campylopus flexuosus</i> (Hedw.) Brid. (Hedwig 1801: 38, Bridel 1819: 71)	RA	PE, RJ, SP, RS	ELS 303 (SP477473)	25°15'57"S, 48°32'42"W
<i>Campylopus fragilis</i> (Brid.) Bruch & Schimp. (Bridel 1801: 296, Buch & Schimper 1847: 164)	RA/EN	MG, RJ, SC, SP	ELS 267 (SP459075)	25°30'50"S, 48°59'16"W
<i>Campylopus heterostachys</i> (Hampe) A. Jaeger (Hampe 1865: 581, Jaeger 1872: 421) (Fig. 20)	AM	BA, CE, ES, GO, MA, MG, MT, PE, PI, PR, RJ, RR, RS, SP	ELS 252 (SP459060)	25°30'49"S, 48°59'20"W
<i>Campylopus julaceus</i> A. Jaeger, 1880	MO	BA, MG, PR, RJ, RS, SC, SP	DFP 17948 (SP461490)	25°30'55"S, 48°58'58"W
<i>Campylopus lamellinervis</i> (Müll. Hal.) Mitt. (Müller 1848: 390, Mitten 1869: 82)	AM	BA, ES, MG, PE, PI, PR, RJ, RS, SC, SP	ELS 248 (SP459056)	25°30'49"S, 48°59'20"W
<i>Campylopus occultus</i> Mitt., 1869	AM	AP, BA, DF, ES, GO, MA, MG, MS, MT, PA, PE, PR, RJ, RR, RS, SC, SP	DFP 17906 (SP461448)	25°30'55"S, 48°58'58"W
<i>Campylopus pilifer</i> Brid., 1819	AM	AL, AM, BA, CE, DF, ES, MG, MT, PA, PE, PR, RJ, RR, RS, SP	ELS 282 (SP477452)	25°15'57"S, 48°32'42"W
<i>Campylopus richardii</i> Brid., 1819	AM	AM, BA, CE, ES, MG, PE, PR, RJ, RR, RS, SC, SP	ELS 373 (SP477543)	25°15'57"S, 48°32'42"W
<i>Campylopus savannarum</i> (Müll. Hal.) Mitt. (Müller 1851: 596, Mitten 1869: 85)	AM	AM, BA, CE, ES, GO, MA, MG, MS, MT, PA, PE, PI, PR, RJ, RO, RR, SE, SP, TO	ELS 291 (SP477461)	25°15'57"S, 48°32'42"W
<i>Campylopus subcuspidatus</i> (Hampe) A. Jaeger (Hampe 1870: 273, Jaeger 1872: 441)	RA	RJ, SP	ELS 255 (SP459063)	25°30'49"S, 48°59'20"W
<i>Campylopus thwaitesii</i> (Mitt.) A. Jaeger (Mitten 1859: 19, Jaeger 1872: 417)	AM	AM, BA, DF, MG, RJ, RS, SC, SP	ELS 534 (SP477706)	25°18'20"S, 48°35'09"W
<i>Leucobryum albicans</i> (Schwägr.) Lindb. (Schwägrichen 1827: 122, Lindberg 1863: 402) (Fig. 21)	AM	BA, CE, DF, ES, MG, MT, PA, PE, PR, RJ, RS, SC, SP	ELS 284 (SP477454)	25°15'57"S, 48°32'42"W
<i>Leucobryum crispum</i> Müll. Hal., 1848	AM	AM, AP, BA, CE, DF, ES, GO, MG, MT, PA, PR, RJ, RO, RR, RS, SC, SP, TO	ELS 301 (SP477471)	25°15'57"S, 48°32'42"W
<i>Leucobryum giganteum</i> Müll. Hal., 1848	AM	AL, AM, BA, ES, MG, PE, PR, RJ, RS, SC, SP	ELS 310 (SP477480)	25°15'57"S, 48°32'42"W
Leucomiaceae				
<i>Leucomium strumosum</i> (Hornsch.) Mitt. (Hornschuch 1840: 69, Mitten 1869: 502)	AM	AC, AL, AM, AP, ES, MG, PA, PE, RJ, RO, RR, SC, SP	DFP 16066 (SP452490)	25°26'11"S, 48°55'14"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<b>Meteoriaceae</b>				
<i>Toloxis imponderosa</i> (Taylor) W.R. Buck (Taylor 1846: 62, Buck 1994: 436) (Fig. 22)	RA	MG, RJ, RS	DFP 15753 (SP452175)	25°26'55"S, 48°54'54"W
<i>Trachypus bicolor</i> Reinw. & Hornsch., 1829	RA	RJ	DFP 15784 (SP452206)	25°26'55"S, 48°54'54"W
<b>Mniaceae</b>				
<i>Epipterygium immarginatum</i> Mitt., 1869	RA	SC	DFP 16186 (SP454764)	25°26'11"S, 48°55'14"W
<i>Pohlia tenuifolia</i> (Mitt.) A. Jaeger (Jaeger 1875: 137, Brotherus 1903: 549)	RA	RJ, SP	DFP 16778 (SP460312)	25°27'00"S, 48°55'05"W
<b>Neckeraceae</b>				
<i>Thamnobryum fasciculatum</i> (Sw. ex Hedw.) I. Sastre (Swartz 1801: 245, Sastre-de Jesús 1993: 232)	AM	ES, MG, PR, RJ, RS, SC, SP	RR s.n. (SP462182)	25°26'59"S, 48°54'51"W
<i>Homaliodendron piniforme</i> (Brid.) Enroth (Bridel 1827: 260, Enroth 1990: 551)	MO	BA, PE, RJ, SC, SP	DFP 16771 (SP460305)	25°27'00"S, 48°55'05"W
<i>Isodrepanium lentulum</i> (Wilson) E. Britton (Wilson 1847: 379, Britton 1914: 28)	RA	BA, RJ, RO, SP	DFP 16216 (SP454794)	25°26'11"S, 48°55'14"W
<i>Neckera villae-ricae</i> Besch., 1877	MO	MS, PR, RJ, RS, SC, SP	DFP 16043 (SP452467)	25°26'11"S, 48°55'14"W
<i>Porotrichodendron superbum</i> (Taylor) Broth. (Taylor 1846: 61, Brotherus 1916: 120)	RA	PR, RS, SC	DFP 15818 (SP452240)	25°26'55"S, 48°54'54"W
<i>Porotrichum filiferum</i> Mitt., 1869	RA	ES, MG, MT, RJ, SP	DFP 15873 (SP452295)	25°26'55"S, 48°54'54"W
<i>Porotrichum korthalsianum</i> (Dozy & Molke.) Mitt. (Dozy & Monkenboer 1854: 42, Mitten 1869: 463)	MO	MG, PE, RJ, RS, SP	DFP 15939 (SP452361)	25°26'55"S, 48°54'54"W
<i>Porotrichum lancifrons</i> (Hampe) Mitt. (Hampe 1863: 158, Mitten 1869: 462)	MO	ES, MG, PR, RJ, RS, SC, SP	DFP 16070 (SP452494)	25°26'11"S, 48°55'14"W
<i>Porotrichum longirostre</i> (Hook. f.) Mitt. (Hooker 1818: 1, Mitten 1869: 461)	MO	MG, MS, MT, PE, RJ, RS, SC, SP	DFP 15777 (SP452199)	25°26'55"S, 48°54'54"W
<i>Porotrichum mutabile</i> Hampe, 1862	MO	MG, PE, RJ, RS, SC	DFP 17844 (SP461386)	25°30'55"S, 48°58'58"W
<i>Porotrichum substriatum</i> (Hampe) Mitt. (Hampe 1866: 340, Mitten 1869: 463) (Fig. 23)	AM	AC, AL, AM, BA, MT, PA, PE, RJ, RO, RR, RS, SC, SP	DFP 15729 (SP452151)	25°26'55"S, 48°54'54"W
<i>Porotrichum thieleanus</i> (Müll. Hal.) Mitt. (Müller 1851: 227, Mitten 1869: 465)	MO/EN	MG, PR, RJ, RS, SP	DFP 15750 (SP452172)	25°26'55"S, 48°54'54"W
<i>Thamnocalia glabella</i> (Hedw.) S. Olsson et al. (Hedwig 1801: 235, Olsson et al. 2011: 46) (Fig. 24)	RA	BA, RJ, SC, SP	DFP 15778 (SP452200)	25°26'55"S, 48°54'54"W
<b>Orthotrichaceae</b>				
<i>Groutiella tumidula</i> (Mitt.) Vitt (Mitten 1869: 201, Vitt 1979: 9)	AM	AC, AM, BA, CE, ES, MT, PA, PR, RJ, RO, SP	GTF 6 (SP438161)	25°26'55"S, 48°55'03"W
<i>Macrocoma tenuis</i> (Hook. f. & Grev.) Vitt (Hooker & Greville 1824: 120, Vitt 1973: 217) (Fig. 25)	MO	GO, PR, RS, SP	DFP 17875 (SP461417)	25°30'55"S, 48°58'58"W
<i>Macromitrium argutum</i> Hampe, 1849	MO	MG, PR, RJ, RS, SC, SP	DFP 17918 (SP461460)	25°30'55"S, 48°58'58"W
<i>Macromitrium cirrosomum</i> (Hedw.) Brid. (Hedwig 1801: 42, Bridel 1826: 316)	AM	AM, AP, BA, CE, MG, PA, PR, RJ, RS, SC, SP	DFP 16727 (SP460261)	25°27'00"S, 48°55'05"W
<i>Macromitrium microstomum</i> (Hook. f. & Grev.) Schwägr. (Hooker & Greville 1824: 114, Schwägrichen 1827: 130)	RA	MG, PR, RJ, SP	DFP 16785 (SP460319)	25°27'00"S, 48°55'05"W
<i>Macromitrium richardii</i> Schwägr., 1826	AM	AM, BA, DF, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 16164 (SP452590)	25°26'11"S, 48°55'14"W
<i>Schlotheimia jamesonii</i> (Arn.) Brid. (Arnott 1823: 349, Bridel 1826: 742)	AM	AC, BA, DF, ES, GO, MA, MG, MS, PE, PR, RJ, RS, SC, SP	DFP 17945 (SP461487)	25°30'55"S, 48°58'58"W
<i>Schlotheimia rugifolia</i> (Hook. f.) Schwägr. (Hooker 1819: 128, Schwägrichen 1824: 150)	AM	AC, AM, BA, CE, DF, ES, GO, MG, MT, PA, PE, PR, RJ, RO, RS, SC, SP	LAA 13 (SP464555)	25°27'10"S, 48°55'11"W
<i>Schlotheimia tecta</i> Hook. f. & Wilson (Fig. 26)	MO	CE, ES, MG, PR, RJ, RS, SC, SP	ELS 254 (SP459062)	25°30'49"S, 48°59'20"W
<i>Schlotheimia torquata</i> (Sw. ex Hedw.) Brid. (Swartz 1801: 246, Bridel 1812: 16)	AM	AM, BA, MG, PE, PR, RJ, RR, RS, SC, SP	ELS 258 (SP459066)	25°30'49"S, 48°59'20"W
<b>Phyllogoniaceae</b>				
<i>Phyllogonium viride</i> Brid., 1827	AM	AL, BA, CE, ES, MG, PE, PR, RJ, RS, SC, SP	ELS 328 (SP477498)	25°15'57"S, 48°32'42"W
<b>Pilotrichaceae</b>				
<i>Callicostella merkelii</i> (Hornsch.) A. Jaeger (Hornschuch 1840: 62, Jaeger 1877: 255)	AM	AC, AM, BA, CE, GO, MG, PA, PE, RJ, RR, SC, SP	DFP 17672 (SP461213)	25°27'10"S, 48°55'11"W
<i>Callicostella pallida</i> (Hornsch.) Ångstr. (Hornschuch 1840: 64, Ångström 1876: 27)	AM	AC, AL, AM, AP, BA, CE, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RN, RO, RR, RS, SC, SE, SP, TO	DFP 16003 (SP452425)	25°26'11"S, 48°55'14"W
<i>Callicostella rufescens</i> (Mitt.) A. Jaeger (Mitten 1869: 352, Jaeger 1877: 355)	MO	AL, AM, PA, PE, RJ	DFP 16036 (SP452460)	25°26'11"S, 48°55'14"W
<i>Cyclodictyon varians</i> (Sull.) Kuntze (Sullivant 1861: 285-286, Kuntze 1891: 835)	MO	MS, PR, RJ, SC, SP	DFP 16184 (SP454762)	25°26'11"S, 48°55'14"W
<i>Hypnella pilifera</i> (Hook. f. & Wilson) A. Jaeger (Hooker 1844: 160, Jaeger 1877: 270)	MO	ES, MG, PB, PR, RJ, RS, SC, SP	ELS 382 (SP477552)	25°15'57"S, 48°32'42"W
<i>Lepidopilidium brevisetum</i> (Hampe) Broth. (Hampe 1878: 266, Brotherus 1907: 944)	MO/EN	AL, ES, MG, RJ, RS, SC, SP	DFP 15986 (SP452408)	25°26'55"S, 48°54'54"W
<i>Lepidopilidium nitens</i> (Hornsch.) Broth. (Hornschuch 1840: 65, Brotherus 1907: 944)	MO	MG, PR, RJ, RN, RS, SC, SP	DFP 17651 (SP461192)	25°27'10"S, 48°55'11"W
<i>Lepidopilidium caudicaule</i> Müll. Hal., 1900	RA/EN	PE, RJ, RS, SC	ELS 377 (SP477547)	25°15'57"S, 48°32'42"W
<i>Lepidopilidium muelleri</i> (Hampe) Hampe (Hampe 1847: 84, Hampe 1865: 365) (Fig. 27)	MO	AL, MG, PE, RJ, RS, SP	DFP 17682 (SP461223)	25°27'10"S, 48°55'11"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Lepidopilum ovalifolium</i> (Duby) Broth. (Duby 1880: 172, Brotherus 1895: 34)	MO	PR, RJ, RS, SC, SP	DFP 15900 (SP452322)	25°26'55"S, 48°54'54"W
<i>Thamniopsis cruegeriana</i> (Müll. Hal.) W.R. Buck (Müller 1851: 208, Buck 1987: 218)	RA	AM, MT	DFP 16111 (SP452537)	25°26'11"S, 48°55'14"W
<i>Thamniopsis incurva</i> (Hornsch.) W.R. Buck (Hornschuch 1840: 65, Buck 1987: 218)	AM	AM, BA, ES, MG, PA, PB, PE, PR, RJ, RS, SC, SP	DFP 16243 (SP454821)	25°26'11"S, 48°55'14"W
<i>Thamniopsis langsdorffii</i> (Hook. f.) W.R. Buck (Hooker 1819: 17, Buck 1987: 218) (Fig. 28)	MO	CE, ES, MG, PR, RJ, RN, RS, SC, SP	ELS 330 (SP477500)	25°15'57"S, 48°32'42"W
<i>Thamniopsis undata</i> (Hedw.) W.R. Buck (Hedwig 1801: 52, Buck 1987: 219)	AM	BA, ES, MG, MT, PR, RJ, RN, SC, SP	DFP 17897 (SP461439)	25°30'55"S, 48°58'58"W
<i>Trachyxiphium guadalupense</i> (Spreng.) W.R. Buck (Sprengel 1812: 96, Buck 1987: 220)	MO	ES, MG, MS, PR, RJ, RS, SC, SP	ELS 329 (SP477499)	25°15'57"S, 48°32'42"W
<i>Trachyxiphium saxicola</i> (R.S. Williams) Vaz-Imbassahy & D.P. Costa (Williams 1909: 248, Vaz-Imbassahy & Costa 2009: 472)	MO	AL, MG, MT, RJ, RN, RS	DFP 15860 (SP452282)	25°26'55"S, 48°54'54"W
Plagiomniaceae				
<i>Plagiomnium rhynchophorum</i> (Harv.) T.J. Kop. (Harvey 1836: 20, Koponen 1971: 57)	MO	AP, ES, GO, MG, PR, RJ, RS, SC, SP	DFP 16812 (SP460346)	25°26'15"S, 48°55'09"W
Polytrichaceae				
<i>Polytrichum angustifolium</i> Mitt., 1869	MO/EN	ES, MG, PR, RJ, RS, SC, SP	DFP 16222 (SP454800)	25°26'11"S, 48°55'14"W
<i>Polytrichum commune</i> Hedw., 1801 (Fig. 29)	AM	AM, BA, DF, ES, GO, MG, PR, RJ, RO, RR, RS, SC, SP	RR 4253 (SP458470)	25°26'57"S, 48°55'53"W
<i>Polytrichum juniperinum</i> Hedw., 1801	AM	BA, DF, ES, GO, MG, PR, RJ, RO, RR, RS, SC, SP, TO	DFP 16253 (SP454831)	25°26'11"S, 48°55'14"W
Pottiaceae				
<i>Barbula riograndensis</i> E.B. Bartram, 1952	RA/EN	PR, RS	DFP 15907 (SP452329)	25°26'55"S, 48°54'54"W
<i>Leptodontium viticulosoides</i> (P. Beauv.) Wijk & Margad. (Palisot de Beauvois 1805: 78, Wijk & Margadant 1960: 51) (Fig. 30)	MO/EN	BA, ES, MG, PR, RJ, RS, SC, SP	DFP 17924 (SP461466)	25°30'55"S, 48°58'58"W
<i>Tortella humilis</i> (Hedw.) Jenn. (Hedwig 1801: 116-117, Jennings 1913: 96)	AM	BA, DF, ES, GO, MA, MG, MS, PE, PR, RJ, RS, SC, SP	DFP 16075 (SP452499)	25°26'11"S, 48°55'14"W
Pterobryaceae				
<i>Orthostichopsis praetermissa</i> W.R. Buck, 1991	RA	AM, BA, MA	DFP 15775 (SP15775)	25°26'55"S, 48°54'54"W
Pterobryaceae				
<i>Orthostichopsis tortipilis</i> (Müll. Hal.) Broth. (Brotherus 1925: 150, Müller 1855: 768)	MO	AM, AP, BA, ES, MG, PE, RJ, SP	DFP 15793 (SP452215)	25°26'55"S, 48°54'54"W
<i>Pterobryon densum</i> Hornsch., 1840	MO	BA, ES, MG, PR, RJ, RS, SC, SP	DFP 15965 (SP452387)	25°26'55"S, 48°54'54"W
Ptychomniaceae				
<i>Ptychomnion cygnisetum</i> (Müll. Hal.) Kindberg (Müller 1885: 425, Kindberg 1888: 31)	MO	PR, RJ, RS, SC	ELS 528 (SP477700)	25°18'20"S, 48°35'09"W
Pylaisiadelphaceae				
<i>Isopterygium tenerifolium</i> Mitt., 1869	AM	AM, BA, CE, DF, GO, MG, MT, PA, PR, RJ, RO, RS, SC, SP	DFP 16237 (SP454815)	25°26'11"S, 48°55'14"W
Rhacocarpaceae				
<i>Rhacocarpus inermis</i> (Müll. Hal.) Lindb. (Müller 1862: 382, Lindberg 1891: 22)	MO/EN	ES, MG, RJ, RS, SC	RR s.n. (SP462191)	25°26'57"S, 48°54'50"W
<i>Rhacocarpus purpurascens</i> (Brid.) Paris (Bridel 1812: 121, Paris 1900: 292)	MO	ES, MG, PR, RJ, RS, SC	ELS 314 (SP477484)	25°15'57"S, 48°32'42"W
Rhizogoniaceae				
<i>Hymenodon aeruginosus</i> (Hook.f. & Wilson) Müll. Hal. (Hooker & Wilson 1844: 154, Müller 1847: 804)	MO	ES, MG, PR, RJ, RS, SC, SP	RR s.n. (SP462183)	25°26'59"S, 48°54'51"W
<i>Pyrrhobryum spiniforme</i> (Hedw.) Mitt. (Hedwig 1801: 236-237, Mitten 1868: 174)	AM	AM, BA, ES, MG, MT, PE, PR, RJ, RO, RR, RS, SC, SP	ELS 526 (SP477698)	25°18'20"S, 48°35'09"W
Sematophyllaceae				
<i>Acroporium caespitosum</i> (Hedw.) W.R. Buck (Hedwig 1802: 233, Buck 1983: 310)	MO	DF, MT, PB, PR, RS	DFP 16006 (SP452428)	25°26'11"S, 48°55'14"W
<i>Acroporium estrellae</i> (Müll. Hal.) W.R. Buck & Schäf.-Verw. (Müller 1851: 275, Buck & Schäfer-Verwimp 1991: 646) (Fig. 31)	AM	BA, DF, GO, MG, PA, PR, RJ, RS, SC, SP	DFP 15773 (SP452195)	25°26'55"S, 48°54'54"W
<i>Aptychella prolifera</i> (Broth.) Herzog (Brotherus 1908: 1115, Herzog 1916: 157)	RA	MG, RJ, SC	DFP 16764 (SP460298)	25°27'00"S, 48°55'05"W
<i>Aptychopsis pyrrophylla</i> (Müll. Hal.) Wijk & Margad. (Müller 1851: 344, Wijk & Margadant 1959: 71)	MO/EN	AM, BA, ES, MG, RJ, SP	DFP 17693 (SP461234)	25°27'10"S, 48°55'11"W
<i>Aptychopsis pungifolia</i> (Hampe) Broth. (Hampe 1879: 152, Brotherus 1925: 411)	MO	BA, MG, PE, RJ, SP	DFP 15744 (SP452166)	25°26'55"S, 48°54'54"W
<i>Paranapiacabaea paulista</i> W.R. Buck & Vital, 1992	RA/EN	PR, SP	DFP 16109 (SP452535)	25°26'11"S, 48°55'14"W
<i>Sematophyllum beyrichii</i> (Hornsch.) Broth. (Hornschuch 1840: 81, Brotherus 1925: 431)	AM	BA, DF, ES, GO, MA, MG, PE, PR, RJ, SP	DFP 17860 (SP461402)	25°30'55"S, 48°58'58"W
<i>Sematophyllum galipense</i> (Müll. Hal.) Mitt. (Müller 1848: 780, Mitten 1869: 480) (Fig. 32)	AM	BA, CE, DF, ES, GO, MA, MG, MT, PA, PR, RJ, RS, SC, SP, TO	ELS 316 (SP477486)	25°15'57"S, 48°32'42"W
<i>Sematophyllum lithophilum</i> (Hornsch.) Ångstr. (Hornschuch 1840: 84, Ångström 1876: 42)	RA	MG, MT, RJ, RS	DFP 17735 (SP461276)	25°27'10"S, 48°55'11"W
<i>Sematophyllum subdepressum</i> (Hampe) Broth. (Hampe 1878: 270, Brotherus 1925: 433)	MO	MT, PR, RJ, RS, SC, SP	DFP 16730 (SP460264)	25°27'00"S, 48°55'05"W
<i>Sematophyllum subsimplex</i> (Hedw.) Mitt. (Hedwig 1801: 69, Mitten 1869: 494)	AM	AC, AL, AM, AP, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PE, PI, PR, RJ, RO, RR, RS, SC, SE, SP, TO	ELS 257 (SP459065)	25°30'49"S, 48°59'20"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Trichosteleum glaziovii</i> (Hampe) W.R. Buck (Hampe 1874: 174, Buck 1998: 243)	MO/EN	MG, PE, PR, RJ, SC, SP	ELS 318 (SP477488)	25°15'57"S, 48°32'42"W
<i>Trichosteleum hornsuschuchii</i> (Hampe) A. Jaeger (Hampe 1844: 9, Jaeger 1878: 418)	AM	AM, BA, MT, PA, PI, PR, RJ, SP	ELS 375 (SP477545)	25°15'57"S, 48°32'42"W
<i>Trichosteleum sentosum</i> (Sull.) A. Jaeger (Sullivant 1861: 288, Jaeger 1878: 415)	MO	AM, BA, PA, PE, PR, RJ	DFP 17666 (SP461207)	25°27'10"S, 48°55'11"W
<i>Wijkia flagellifera</i> (Broth.) H.A. Crum (Brotherus 1895: 54, Crum 1971: 172)	MO	BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 16077 (SP452502)	25°26'11"S, 48°55'14"W
<b>Sphagnaceae</b>				
<i>Sphagnum cyclophyllum</i> Sull. & Lesq. in Gray, 1856	MO	BA, ES, GO, MG, PR, RJ, SC, SP	DFP 15762 (SP452184)	25°26'55"S, 48°54'54"W
<i>Sphagnum divisum</i> H.A. Crum, 1992	MO/EN	BA, GO, MG, RJ, SC	DFP 16786 (SP460320)	25°27'00"S, 48°55'05"W
<i>Sphagnum exquisitum</i> H.A. Crum, 1992	RA/EN	MG, PR, RJ, SP	DFP 15866 (SP452288)	25°26'55"S, 48°54'54"W
<i>Sphagnum globicephalum</i> Müll. Hal. ex Warnst., 1911	RA/EN	RJ, SC	DFP 15899 (SP452321)	25°26'55"S, 48°54'54"W
<i>Sphagnum multiporosum</i> H.A. Crum, 1987	MO/EN	BA, MG, RJ, RS, SP	DFP 17721 (SP461262)	25°27'10"S, 48°55'11"W
<i>Sphagnum ovalifolium</i> Warnst., 1891	RA	GO, MG	DFP 15738 (SP452160)	25°26'55"S, 48°54'54"W
<i>Sphagnum palustre</i> L., 1753 (Fig. 33)	AM	AM, AP, BA, CE, ES, GO, MG, MS, PA, PB, PE, PR, RJ, RO, RR, RS, SC, SE, SP	RR s.n. (SP462180)	25°26'49"S, 48°54'53"W
<i>Sphagnum perichaetiale</i> Hampe, 1848	AM	AM, AP, CE, DF, ES, MG, MS, MT, PA, PB, PE, PR, RJ, RO, RR, RS, SC, SE, SP	DFP 15864 (SP452286)	25°26'55"S, 48°54'54"W
<i>Sphagnum pseudoramulinum</i> H.A. Crum, 1987	RA/EN	RJ, RS, SP	DFP 15819 (SP452241)	25°26'55"S, 48°54'54"W
<i>Sphagnum strictum</i> Sull., 1846	RA	BA, ES	DFP 17659 (SP461200)	25°27'10"S, 48°55'11"W
<i>Sphagnum subsecundum</i> Nees in Sturm, 1819	AM	BA, DF, ES, GO, MG, MT, PR, RJ, RO, RS, SC, SP, TO	RR 2834 (SP437115)	25°27'14"S, 48°55'12"W
<i>Sphagnum sucrei</i> H.A. Crum, 1987	RA/EN	MG, RJ	RR s.n. (SP462185)	25°26'57"S, 48°54'50"W
<i>Sphagnum tenellum</i> (Brid.) Brid., 1819	RA	AM, RJ	DFP 15854 (SP452276)	25°26'55"S, 48°54'54"W
<b>Symphyodontaceae</b>				
<i>Symphyodon imbricatifolius</i> (Mitt.) S.P. Churchill (Mitten 1869: 372, Churchill 1994: 194)	RA	GO, MG, RJ	DFP 16256 (SP454834)	25°26'11"S, 48°55'14"W
<b>Thuidiaceae</b>				
<i>Thuidiopsis furfurosa</i> (Hook. f. & Wilson) M. Fleisch. (Hooker & Wilson 1854: 10, Fleischer 1923: 1497)	AM	ES, MG, PR, RJ, RS, SC, SP	DFP 17848 (SP461390)	25°30'55"S, 48°58'58"W
<i>Thuidium brasiliense</i> Mitt., 1869	MO	MG, PR, RJ, RS, SC, SP	DFP 16817 (SP460351)	25°26'15"S, 48°55'09"W
<i>Thuidium delicatulum</i> (Hedw.) Bruch & Schimp. (Hedwig 1801: 26-261, Schimper 1852: 164)	AM	AM, BA, GO, MG, MT, PA, PR, RJ, RS, SC, SP	ELS 530 (SP477702)	25°18'20"S, 48°35'09"W
<i>Thuidium tomentosum</i> Besch., 1872	AM	AL, AM, BA, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RO, RR, RS, SC, SP	DFP 16203 (SP454781)	25°26'11"S, 48°55'14"W
<b>Marchantiophyta</b>				
<b>Acrobolbaceae</b>				
<i>Tylimanthus laxus</i> (Lehm. & Lindb.) Spruce (Lindenberg 1841: 68, Stephani 1885: 502)	MO	ES, MS, MT, PR, RJ, SP	DFP 17755 (SP461297)	25°27'10"S, 48°55'11"W
<b>Adelanthaceae</b>				
<i>Adelanthus decipiens</i> (Hook.) Mitt. (Hooker 1813: 50, Mitten 1864: 244)	MO	BA, ES, MG, PR, RJ	RR 2803 (SP437094)	25°27'14"S, 48°55'12"W
<b>Aneuraceae</b>				
<i>Aneura pinguis</i> (L.) Dumort. (Linnaeus 1753: 1136, Dumortier 1831: 86) (Fig. 34)	AM	AL, AM, ES, MG, MS, PA, PR, RJ, SC, SP	DFP 16118 (SP452544)	25°26'11"S, 48°55'14"W
<i>Riccardia chamedryfolia</i> (With.) Grolle (Withering 1776: 699, Grolle 1969: 772)	MO	DF, ES, GO, MG, MT, PR, RJ, RS, SP	ELS 338 (SP477508)	25°15'57"S, 48°32'42"W
<i>Riccardia digitiloba</i> (Spruce ex Steph.) Pagán (Spruce 1888: 276, Pagán 1939: 6)	AM	AC, AM, BA, CE, ES, MG, MS, MT, PE, RJ, SP	ELS 546 (SP477718)	25°18'20"S, 48°35'09"W
<i>Riccardia emarginata</i> (Steph.) K.G. Hell (Stephani 1893: 20, Hell 1969: 100)	RA/EN	BA, MG, RJ, SP	ELS 250 (SP459058)	25°30'49"S, 48°59'20"W
<i>Riccardia fucoides</i> (Sw.) C. Massal. (Swartz 1788: 145, Massalongo 1885: 256)	RA	BA, ES, RJ, SP	DFP 17660 (SP461201)	25°27'10"S, 48°55'11"W
<i>Riccardia glaziovii</i> (Spruce) Meenks (Spruce 1889: 201, Meenks 1987: 173)	MO	AP, BA, ES, PA, PR, RJ, RS, SC, SP	ELS 365 (SP477535)	25°15'57"S, 48°32'42"W
<i>Riccardia regnellii</i> K.G. Hell, 1969	AM/EN	BA, ES, MG, MT, PE, RJ, RN, RS, SC, SP	DFP 16072 (SP16072)	25°26'11"S, 48°55'14"W
<b>Balantiopsidaceae</b>				
<i>Isotachis aubertii</i> (Schwägr.) Mitt. (Schwägrichen 1814: 19, Mitten 1887: 322) (Fig. 35)	MO	ES, MG, PR, RJ, RS, SC, SP	RR 2806 (SP438273)	25°27'14"S, 48°55'12"W
<i>Isotachis inflata</i> Steph., 1903	RA/EN	ES, PR, RJ	ELS 488 (SP477660)	25°16'16"S, 48°33'04"W
<i>Isotachis multiceps</i> (Lindb. & Gottsche) Gottsche (Lindenberg & Gottsche 1847: 687, Gottsche 1863: 105)	MO	ES, PR, RJ, RS, SP	DFP 17697 (SP461238)	25°27'10"S, 48°55'11"W
<i>Neesioscyphus bicuspidatus</i> (Nees) Grolle (Stephani 1901: 471, Grolle 1966: 182)	RA	MG	DFP 16239 (SP454817)	25°26'11"S, 48°55'14"W
<i>Neesioscyphus homophyllus</i> (Nees) Nees (Nees von Esenbeck in Martius 1833: 336, Grolle 1964: 188)	RA	BA, MG, RJ, SP	DFP 17891 (SP461433)	25°30'55"S, 48°58'58"W
<b>Calypogeiaceae</b>				
<i>Calypogeia peruviana</i> Nees & Mont., 1838	AM	AC, AL, AM, BA, DF, GO, MG, PE, PR, RJ, RS, SC, SP	DFP 16058 (SP452482)	25°26'11"S, 48°55'14"W
<i>Mnioloma caespitosum</i> (Spruce) R.M. Schuster (Spruce 1885: 412, Schuster 1995: 839)	RA	AM	DFP 16777 (SP460311)	25°27'00"S, 48°55'05"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Mnioloma cyclostipum</i> (Spruce) R.M. Schuster (Spruce 1885: 411, Schuster 1995: 843)	RA	RJ, SC	DFP 15820 (SP452242)	25°26'55" S, 48°54'54" W
<b>Cephaloziaceae</b>				
<i>Fuscocephaloziopsis crassifolia</i> (Lindenb. & Gottsche) Vána & L. Söderstr.	RA	BA, ES, MG, PR, RJ, RS, SP	DFP 17889 (SP461431)	25°30'55" S, 48°58'58" W
<i>Odontoschisma denudatum</i> (Nees) Dumort. (Nees von Esenbeck in Martius 1817: 14, Dumortier 1835: 19)	AM	AM, BA, MT, PE, RJ, SC, SP	ELS 251 (SP459059)	25°30'49" S, 48°59'20" W
<i>Odontoschisma falcifolium</i> Steph., 1908 (Fig. 36)	RA	AM, BA, ES, GO, MG, MT, RJ, SP	DFP 15761 (SP452183)	25°26'55" S, 48°54'54" W
<i>Odontoschisma longiflorum</i> (Taylor) Steph. (Taylor 1846: 281, Trevisan 1877: 419)	MO	BA, ES, GO, MG, MT, PR, RJ, SP	ELS 246 (SP459054)	25°30'49" S, 48°59'20" W
<b>Herbertaceae</b>				
<i>Herbertus acantheilus</i> Spruce, 1885	MO	MG, RJ, RS, SC	WTF 39 (SP437099)	25°27'14" S, 48°55'12" W
<i>Herbertus bivittatus</i> Spruce, 1885	MO	BA, CE, PE, RJ	DFP 17756 (SP461298)	25°27'10" S, 48°55'11" W
<i>Herbertus pensilis</i> Spruce, 1885	RA	RJ	DFP 17729 (SP461270)	25°27'10" S, 48°55'11" W
<b>Frullaniaceae</b>				
<i>Frullania atrata</i> (Sw.) Dumort. (Swartz 1788: 144, Dumortier 1835: 13)	AM	AM, BA, MG, PA, PE, PR, RJ, RS, SC, SP	ELS 327 (SP477497)	25°15'57" S, 48°32'42" W
<i>Frullania beyrichiana</i> (Lehm. & Lindenb.) Lehm. & Lindenb. (Lehmann & Lindenberg in Lehmann 1833: 25, Lehmann & Lindenberg 1845: 460)	MO	AC, BA, ES, GO, MG, MT, PA, PE, RJ, RS, SP	RR s.n. (SP462184)	25°26'59" S, 48°54'51" W
<i>Frullania brasiliensis</i> Raddi, 1822 (Fig. 37)	AM	BA, CE, DF, ES, GO, MG, PE, RJ, RS, SC, SP	ELS 274 (SP459082)	25°30'50" S, 48°59'16" W
<i>Frullania breuteliana</i> Gottsche, 1845	MO	BA, PE, RJ, RS, SP	ELS 326 (SP477496)	25°15'57" S, 48°32'42" W
<i>Frullania caulisequa</i> (Nees) Nees (Nees von Esenbeck 1833: 373, Montagne 1839: 51)	AM	AC, AL, BA, CE, DF, ES, GO, MG, MT, PA, PB, PE, RJ, RR, RS, SC, SE, SP	DFP 16139 (SP452565)	25°26'11" S, 48°55'14" W
<i>Frullania curvilobula</i> Schäf.-Verw. et al., 2012	RA	PR, SP	DFP 16126 (SP452552)	25°26'11" S, 48°55'14" W
<i>Frullania flexicaulis</i> Spruce, 1884	RA	SC, SP	DFP 17650 (SP461191)	25°27'10" S, 48°55'11" W
<i>Frullania kunzei</i> (Lehm. & Lindb.) Lehm. & Lindb. (Lehmann & Lindenberg 1834: 50, Lehmann & Lindenberg 1845: 449)	AM	AC, AM, BA, CE, DF, ES, GO, MG, MT, PA, PB, PE, PR, RJ, RR, RS, SC, SE, SP	DFP 16127 (SP452553)	25°26'11" S, 48°55'14" W
<i>Frullania setigera</i> Steph., 1894	MO	ES, MG, PR, RJ, RS, SC, SP	DFP 16733 (SP460267)	25°27'00" S, 48°55'05" W
<b>Geocalycaceae</b>				
<i>Saccogynidium caldense</i> (Ångstr.) Grolle (Ångström 1876: 80, Grolle 1960: 44)	MO	GO, RJ, SP	ELS 541 (SP477713)	25°18'20" S, 48°35'09" W
<b>Jamesoniellaceae</b>				
<i>Syzygiella anomala</i> (Lindenb. & Gott.) Steph. (Lindenberg & Gottsche 1844: 646, Stephani 1902: 190) (Fig. 38)	RA	MG, PR, RJ, SP	DFP 15732 (SP452154)	25°26'55" S, 48°54'54" W
<i>Syzygiella contigua</i> (Gottsche) Steph. (Gottsche 1864: 118, Stephani 1902: 470)	MO	RJ, SP	ELS 485 (SP477657)	25°16'16" S, 48°33'04" W
<i>Syzygiella integerrima</i> Steph., 1917	RA	ES, MG, RJ	DFP 17656 (SP461197)	25°27'10" S, 48°55'11" W
<i>Syzygiella liberata</i> Inoue, 1974	MO	BA, MG, RJ, SC, SP	LAA 73 (SP464615)	25°27'00" S, 48°55'05" W
<i>Syzygiella rubricaulis</i> (Nees) Steph. (Nees von Esenbeck in Martius 1833: 34, Stephani 1902: 187)	MO	BA, MG, RJ, RS, SC, SP	ELS 348 (SP477518)	25°15'57" S, 48°32'42" W
<b>Lophocoleaceae</b>				
<i>Chiloscyphus muricatus</i> (Lehm.) J.J. Engel & R.M. Schust. (Lehmann 1829: 363, Engel & Schuster 1984: 419)	MO	ES, MG, RJ, RS, SC, SP	DFP 15947 (SP452369)	25°26'55" S, 48°54'54" W
<i>Chiloscyphus martianus</i> (Nees) J.J. Engel & R.M. Schust. (Nees von Esenbeck 1845: 152, Engel & Schuster 1984: 419) (Fig. 39)	AM	AM, AP, BA, CE, ES, GO, MG, MT, PA, PE, PR, RJ, RS, SC, SE, SP	DFP 16784 (SP460318)	25°27'00" S, 48°55'05" W
<i>Chiloscyphus proteus</i> (Herzog) J.J. Engel & R.M. Schust. (Herzog 1955: 164, Engel & Schuster 1984: 421)	RA	RJ, SP	RR s.n. (SP462182)	25°26'59" S, 48°54'51" W
<i>Heteroscyphus valdiviensis</i> (Mont.) Schiffn. (Montagne 1845: 351, Schiffner 1910: 172)	RA	SP	DFP 17674 (SP461215)	25°27'10" S, 48°55'11" W
<i>Leptoscyphus amphibolius</i> (Nees) Grolle (Nees von Esenbeck 1833 in Martius: 334, Grolle 1962: 54)	MO	BA, ES, GO, MG, RJ, SP	DFP 15935 (SP452357)	25°26'55" S, 48°54'54" W
<i>Leptoscyphus porphyrius</i> (Nees) Grolle (Nees von Esenbeck 1845: 185, Grolle 1969: 3)	MO	ES, MG, PA, PE, PR, RJ, SP	DFP 16726 (SP460260)	25°27'00" S, 48°55'05" W
<b>Lejeuneaceae</b>				
<i>Anoplolejeunea conferta</i> (Meissn. ex Spreng.) A. Evans (Spreng 1827: 325, Evans 1908: 175)	AM	AL, BA, ES, MG, PA, PB, PE, PR, RJ, RR, RS, SC, SP	DFP 16207 (SP454785)	25°26'11" S, 48°55'14" W
<i>Archilejeunea ludoviciana</i> (De Not. ex Lehm.) Gradst. & Geissler (Lehmann 1857: 11, Gradstein & Geissler 1994: 58)	RA	AM	DFP 15922 (SP452344)	25°26'55" S, 48°54'54" W
<i>Brachiolejeunea phyllorhiza</i> (Nees) Kruijt & Gradst. (Nees von Esenbeck 1833 in Martius: 348, Kruijt & Gradstein 1986: 299)	MO	AM, BA, ES, MG, PE, RJ, SP	DFP 15953 (SP452375)	25°26'55" S, 48°54'54" W
<i>Bryopteris diffusa</i> (Sw.) Nees (Swartz 1788: 144, Nees von Esenbeck 1845: 286)	AM	AL, AM, BA, CE, ES, MG, MT, PA, PB, PE, PR, RJ, RS, SC, SP	RR 4268 (SP458541)	25°26'57" S, 48°55'53" W
<i>Bryopteris filicina</i> (Sw.) Nees (Swartz 1788: 145, Nees von Esenbeck 1845: 284) (Fig. 40)	AM	AL, AM, CE, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RR, RS, SC, SP	DFP 15751 (SP452173)	25°26'55" S, 48°54'54" W
<i>Ceratolejeunea confusa</i> R.M. Schuster, 1956	RA	AM, BA, PA, PE, SP	DFP 15713 (SP452135)	25°26'55" S, 48°54'54" W
<i>Ceratolejeunea cornuta</i> (Lindb.) Steph. (Lindenberg 1829: 23, Stephani 1895: 65)	MO	AC, AM, AP, CE, MG, PA, PE, PR, RJ, RO, RR, SC, SP	ELS 452 (SP477623)	25°15'57" S, 48°32'42" W
<i>Cheilolejeunea acutangula</i> (Nees) Grolle (Nees von Esenbeck 1833 in Martius: 357, Grolle 1979: 173)	AM	AL, AM, BA, DF, ES, GO, MG, MT, PA, PE, RJ, RR, RS, SC, SP	ELS 481 (SP477653)	25°16'16" S, 48°33'04" W



Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Cheilolejeunea adnata</i> (Kutze.) Grolle (Kunze 1834: 46, Grolle 1977: 529)	AM	AC, AL, AM, AP, BA, ES, MT, PA, PE, PR, RN, RR, SC, SP	DFP 16087 (SP452513)	25°26'11"S, 48°55'14"W
<i>Cheilolejeunea caducifolia</i> (Gradst. & Schäf.-Verw.) W. Ye & R.L. Zhu (Gradstein & Schäfer-Verwimp 1993: 64, Ye & Zhu 2010: 280)	RA/EN	BA, ES, MG	DFP 17736 (SP461277)	25°27'10"S, 48°55'11"W
<i>Cheilolejeunea clausa</i> (Nees & Mont.) R.M. Schuster (Nees von Esenbeck in Montagne 1840: 337, Schuster 1980: 863) (Fig. 41)	AM	AL, AM, BA, CE, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RO, SP	DFP 15801 (SP452223)	25°26'55"S, 48°54'54"W
<i>Cheilolejeunea holostipa</i> (Spruce) Grolle & R.L. Zhu (Spruce 1884: 171, Grolle & Zhu 2001: 1071)	MO	BA, ES, MG, PA, PE, PR, RJ, SP	DFP 15957 (SP452379)	25°26'55"S, 48°54'54"W
<i>Cheilolejeunea oncophylla</i> (Ångstr.) Grolle & E. Reiner (Ångström 1876: 86, Grolle & Reiner 1997: 781)	MO	AP, BA, MG, PA, PR, RJ, RR, SC, SP	DFP 15861 (SP452283)	25°26'55"S, 48°54'54"W
<i>Cheilolejeunea rigidula</i> (Mont.) R.M. Schuster (Montagne 1840: 336, Schuster 1971: 102)	AM	AC, AL, AM, AP, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PB, PE, PR, RJ, RR, SC, SE, SP, TO	DFP 17903 (SP461445)	25°30'55"S, 48°58'58"W
<i>Cheilolejeunea uncioloba</i> (Lidenb.) Malombe (Lindenberg 1845: 331, Malombe 2009: 325)	MO	BA, CE, ES, MG, RJ, RS, SP	ELS 538 (SP477710)	25°18'20"S, 48°35'09"W
<i>Cheilolejeunea xanthocarpa</i> (Lehm. & Lindenberg) Malombe (Lehman & Lindenberg in Lehmann 1833: 8, Malombe 2009: 326)	MO	BA, CE, ES, MG, SP	ELS 540 (SP477712)	25°18'20"S, 48°35'09"W
<i>Cololejeunea diaphana</i> A. Evans, 1905	AM	AM, ES, GO, MT, PA, PE, RJ, RS, SC, SP	DFP 16233 (SP454811)	25°26'11"S, 48°55'14"W
<i>Cololejeunea gracilis</i> (Ast.) Pócs (Jovet-Ast 1947: 21, Pócs 2008: 233)	MO	AM, ES, MG, MT, PA, RJ, SP	DMC 1122 (SP455270)	25°26'11"S, 48°55'14"W
<i>Cololejeunea manaosensis</i> (Herzog) O.Yano (Herzog 1931: 349, Yano 1894: 513) (Fig. 42)	RA/EN	AM, MT, PA, SP	DFP 17685 (SP461226)	25°27'10"S, 48°55'11"W
<i>Cololejeunea microscopica</i> (Taylor) Schiffn. (Taylor in MacKay 1836: 59, Schiffner in Engler & Prantl 1893: 122)	RA/EN	SP	ELS 489 (SP477661)	25°16'16"S, 48°33'04"W
<i>Cololejeunea papilliloba</i> (Steph.) Steph. (Stephani 1890: 73, Stephani 1890: 135)	RA/EN	MG, RS, SP	DFP 17861 (SP461403)	25°30'55"S, 48°58'58"W
<i>Colura calyptrifolia</i> (Hook. f.) Dumort. (Hooker 1813: 1013, Dumortier 1835: 12)	RA	MG, RJ	DFP 17757 (SP461299)	25°27'10"S, 48°55'11"W
<i>Cyclolejeunea convexistipa</i> (Lehm. & Lindb.) A. Evans (Lehmann & Lindenberg 1834: 43, Evans 1904: 198)	AM	AL, AM, AP, BA, CE, MA, PA, PE, RJ, RN, RO, SP	DFP 16185 (SP454763)	25°26'11"S, 48°55'14"W
<i>Cyclolejeunea luteola</i> (Spruce) Grolle (Spruce 1884: 205, Grolle 1984: 761)	MO	AM, BA, MG, MT, PA, PE, RJ, RR, SP	DFP 15800 (SP452222)	25°26'55"S, 48°54'54"W
<i>Dicranolejeunea axillaris</i> (Nees & Mont.) Schiffn. (Nees von Esenbeck & Montagne 1936: 59, Schiffner in Engler & Prantl 1893: 128)	RA	PR, RJ, SP	DFP 16813 (SP460347)	25°26'15"S, 48°55'09"W
<i>Diplasiolejeunea brunnea</i> Steph., 1916 (Fig. 43)	MO	AC, AL, AM, BA, CE, ES, MT, PA, RJ, RO, SC, SP	DFP 17699 (SP461240)	25°27'10"S, 48°55'11"W
<i>Diplasiolejeunea inermis</i> P. Tixier, 1985	RA	RJ, SC	DFP 15771 (SP452193)	25°26'55"S, 48°54'54"W
<i>Drepanolejeunea anoplantha</i> (Spruce) Steph. (Spruce 1884: 189, Stephani 1913: 325)	MO	AM, BA, CE, ES, PB, RJ, RS, SP	DFP 15972 (SP452394)	25°26'55"S, 48°54'54"W
<i>Drepanolejeunea araucariae</i> Steph., 1896	MO	MG, RJ, RS, SC, SP	DFP 15881 (SP452303)	25°26'55"S, 48°54'54"W
<i>Drepanolejeunea campanulata</i> (Spruce) Steph. (Spruce 1884: 192, Stephani 1913: 328)	RA	RJ, SC, SP	DFP 17890 (SP461432)	25°30'55"S, 48°58'58"W
<i>Drepanolejeunea capulata</i> (Taylor) Steph. (Taylor 1846: 394, Stephani 1913: 328)	MO	AM, SP	DFP 17723 (SP461264)	25°27'10"S, 48°55'11"W
<i>Drepanolejeunea fragilis</i> Bischl., 1964	AM	AL, AM, AP, BA, CE, ES, MG, PA, PE, RJ, RR, SP	DFP 17745 (SP461286)	25°27'10"S, 48°55'11"W
<i>Drepanolejeunea mosenii</i> (Steph.) Bischl. (Stephani 1913: 372, Bischer 1967: 118)	MO	AM, BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 17752 (SP461294)	25°27'10"S, 48°55'11"W
<i>Drepanolejeunea orthophylla</i> (Nees & Mont.) Bischl. (Nees von Esenbeck & Montagne 1843: 265, Bischler 1967: 102)	MO	AM, PA, RJ, SC, SP	DFP 17696 (SP461237)	25°27'10"S, 48°55'11"W
<i>Harpalejeunea oxyphylla</i> (Nees & Mont.) Steph. (Nees von Esenbeck 1843: 262, Stephani 1890: 76)	MO	AM, BA, PA, PB, PE, RJ, RR, SP	DFP 15909 (SP452331)	25°26'55"S, 48°54'54"W
<i>Harpalejeunea stricta</i> (Lindenb. & Gottsche) Steph. (Lindenberg & Gottsche 1847: 756, Stephani 1913: 259)	MO	AL, BA, MG, PA, PE, RJ, SP	ELS 336 (SP477506)	25°15'57"S, 48°32'42"W
<i>Harpalejeunea tridens</i> (Besch. & Spruce) Steph. (Bescherelle & Spruce 1890: 36, Stephani 1913: 263)	RA	SP, PE	DFP 17670 (SP461211)	25°27'10"S, 48°55'11"W
<i>Lejeunea bermudiana</i> (A. Evans) R.M. Schuster (Evans 1906: 132, Schuster 1980: 1105)	MO	AC, ES, PA, RJ, SP	DMC 1143 (SP455291)	25°26'11"S, 48°55'14"W
<i>Lejeunea caulicalyx</i> (Steph.) M.E. Reiner & Goda (Stephani 1913: 237, Reiner & Goda 2000: 13)	AM	AC, AL, BA, ES, MS, MT, PA, PE, PR, RJ, RR, SP	DFP 17853 (SP461395)	25°30'55"S, 48°58'58"W
<i>Lejeunea cerina</i> (Lehm. & Lindb.) Gottsche & et al. (Lehmann & Lindenberg in Lehmann 1833: 16, Gottsche et al. 1845: 391)	MO	AC, BA, ES, PE, RJ, SP	DFP 17728 (SP461269)	25°27'10"S, 48°55'11"W
<i>Lejeunea controversa</i> Gottsche & Rabenh., 1872	MO	AC, AL, BA, MS, PA, PE, RJ, SP	DFP 16251 (SP454829)	25°26'11"S, 48°55'14"W
<i>Lejeunea cristulata</i> (Steph.) M.E. Reiner & Goda (Stephani 1896: 75, Reiner & Goda 2000: 89)	MO/EN	BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 15709 (SP452131)	25°26'55"S, 48°54'54"W
<i>Lejeunea deplanata</i> Ness, 1845	MO	AC, AL, BA, ES, MS, PA, RJ	DFP 16004 (SP452426)	25°26'11"S, 48°55'14"W
<i>Lejeunea flava</i> (Sw.) Nees (Swartz 1788: 144, Nees von Esenbeck 1838: 277) (Fig. 44)	AM	AC, AL, AM, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PB, PE, PR, RJ, RR, RS, SC, SE, SP, TO	ELS 347 (SP477517)	25°15'57"S, 48°32'42"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Lejeunea glaucescens</i> Gottsche, 1845	AM	AC, AM, BA, CE, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RR, RS, SC, SP	RR 2867 (SP437142)	25°27'14"S, 48°55'12"W
<i>Lejeunea grossitexta</i> (Steph.) E. Reiner & Goda (Steph 1913: 240, Reiner & Goda 2000)	MO	AL, BA, CE, ES, MG, PR, RJ, SC, SP	DFP 16119 (SP452545)	25°26'11"S, 48°55'14"W
<i>Lejeunea laeta</i> (Lehm. & Lindenb.) Lehm. & Lindenb. (Lehmann & Lindenberg 1834: 45, Lehmann & Lindenberg 1845: 380)	MO	BA, CE, MG, PR, RJ, SC, SP	RR s.n. (SP462189)	25°26'20"S, 48°55'11"W
<i>Lejeunea laetevirens</i> Nees & Mont., 1842	AM	AC, AL, AM, AP, BA, CE, DF, ES, GO, MA, MS, MT, PA, PB, PE, PR, RJ, RN, RR, RS, SC, SE, SP	DFP 17953 (SP462404)	20°45'39"W, 41°32'05"W
<i>Lejeunea oligoclada</i> Spruce, 1889	MO/EN	AL, BA, ES, MG, PE, PR, RJ, SC, SP	DFP 17688 (SP461229)	25°27'10"S, 48°55'11"W
<i>Lopholejeunea nigricans</i> (Lindb.) Schiffn. (Lindenberg 1845: 316, Stephani 1898: 293)	AM	AC, AL, AM, BA, CE, DF, ES, GO, MA, MG, MS, MT, PA, PE, PR, RJ, RS, SC, SP	DFP 16011 (SP452434)	25°26'11"S, 48°55'14"W
<i>Marchesinia bongardiana</i> Trevis., 1877	AM	AM, RJ, SC, SP	DFP 16213 (SP454791)	25°26'11"S, 48°55'14"W
<i>Marchesinia brachiata</i> (Sw.) Schiffn. (Swartz 1788: 144, Schiffner in Engler & Prantl 1893: 128)	AM	BA, CE, ES, MG, MT, PE, PR, RJ, RR, SC, SE, SP	DFP 15929 (SP452351)	25°26'55"S, 48°54'54"W
<i>Microlejeunea acutifolia</i> Steph., 1896	RA	PA	DMC 1129 (SP455277)	25°26'11"S, 48°55'14"W
<i>Microlejeunea cystifera</i> Herzog, 1950	RA	BA, SP	DFP 15764 (SP452186)	25°26'55"S, 48°54'54"W
<i>Microlejeunea epiphylla</i> Bischl., 1963 (Fig. 45)	AM	AL, AP, BA, CE, ES, GO, MA, MG, MS, PA, PB, PE, RJ, SE, SP, TO	DFP 15996 (SP452418)	25°26'11"S, 48°55'14"W
Lejeuneaceae				
<i>Microlejeunea globosa</i> (Spruce) Steph. (Spruce 1889: 193, Stephani 1915: 821)	MO	ES, PA, PR, RS, SC, SE, SP	DFP 17714 (SP461255)	25°27'10"S, 48°55'11"W
<i>Myriocoleopsis minutissima</i> (Sm.) R.L. Zhu et al. (Smith 1806: 1633, Zhu et al. 2014: 293)	AM	AC, AM, BA, ES, MG, MS, MT, PE, PR, RJ, RR, SC	DFP 16822 (SP460356)	25°26'15"S, 48°55'09"W
<i>Neurolejeunea breutelii</i> (Gottsche) A. Evans (Gottsche 1845: 324, Evans 1907: 13)	MO	BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 17746 (SP461287)	25°27'10"S, 48°55'11"W
<i>Odontolejeunea lunulata</i> (F. Weber) Schiffn. (Weber 1815: 33, Schiffner in Engler & Prantl 1893: 128)	AM	AC, AM, AP, BA, CE, ES, MG, MT, PA, PE, PR, RJ, RR, RS, SP	DFP 16021 (SP452444)	25°26'11"S, 48°55'14"W
<i>Omphalanthus filiformis</i> (Sw.) Nees (Swartz 1788: 144, Nees 1845: 304)	AM	AM, BA, CE, ES, MG, PE, RJ, RS, SE, SP	ELS 337 (SP477507)	25°15'57"S, 48°32'42"W
<i>Physantholejeunea huctumalcensis</i> (Lindenb. & Gottsche) Heinrichs & Schäf.-Verw. (Lindenberg & Gottsche 1847: 762, Heinrichs et al. 2013: 387)	RA	AM, BA, PA	DFP 16007 (SP452429)	25°26'11"S, 48°55'14"W
<i>Prionolejeunea aemula</i> (Gottsche) A. Evans (Gottsche 1845: 338, Evans 1904: 219)	MO	AM, BA, MT, PA, PE, RJ, RR, SE, SP	ELS 339 (SP477509)	25°15'57"S, 48°32'42"W
<i>Prionolejeunea scaberula</i> (Spruce) Steph. (Spruce 1884: 159, Stephani 1913: 214)	RA/EN	AM, BA, MA, SP	DFP 16024 (SP452447)	25°26'11"S, 48°55'14"W
<i>Taxilejeunea isocalycina</i> (Nees) Steph. (Nees von Esenbeck in Martius 1833: 356, Stephani 1903: 118)	MO	AM, ES, MG, PA, PR, RJ, SC, SP	DFP 17880 (SP461422)	25°30'55"S, 48°58'58"W
<i>Taxilejeunea pterigonia</i> (Lehm. & Lindb.) Schiffn. (Lehmann & Lindenberg 1834: 44, Schiffner in Engler & Prantl 1893: 125) (Fig. 46)	MO	ES, MG, RJ, RS, SC, SP	DFP 17691 (SP461232)	25°27'10"S, 48°55'11"W
<i>Xylolejeunea crenata</i> (Nees & Mont.) X.-L. He & Grolle (Montagne 1838: 48, He & Grolle 2001: 36)	AM	AL, AM, AP, BA, ES, MA, MG, PA, PE, RJ, RO, RR, SC, SP	DFP 16086 (SP452512)	25°26'11"S, 48°55'14"W
Lepicoleaceae				
<i>Lepicolea ochroleuca</i> (Spreng.) Spruce (Sprengel 1827: 325, Spruce 1885: 345)	RA	RS, SC	DFP 16747 (SP460281)	25°27'00"S, 48°55'05"W
Lepidoziaceae				
<i>Bazzania arcuata</i> (Lindb. & Gottsche) Trevis. (Lindenberg & Gottsche 1863, Trevisan 1877: 414)	AM	AM, MG, RJ, SP	ELS 323 (SP477493)	25°15'57"S, 48°32'42"W
<i>Bazzania aurescens</i> Spruce, 1885	MO	AM, BA, ES, GO, MG, PR, RJ, SC, SP	ELS 451 (SP477622)	25°15'57"S, 48°32'42"W
<i>Bazzania cuneistipula</i> (Gottsche et al.) Trevis. (Gottsche et al. 1845: 225, Trevisan 1877: 414)	RA	MG, RJ, SP	DFP 17907 (SP461449)	25°30'55"S, 48°58'58"W
<i>Bazzania gracilis</i> (Hampe & Gottsche) Stephani (Hampe & Gottsche 1852: 346, Stephani 1888: 279)	MO	AM, BA, ES, MG, PE, RJ, RR, SP	DFP 17653 (SP461194)	25°27'10"S, 48°55'11"W
<i>Bazzania heterostipa</i> (Steph.) Fulford (Stephani 1909: 532, Fulford 1959: 410)	MO/EN	BA, ES, MG, PE, PR, RJ, RS, SC, SP	DFP 15726 (SP452148)	25°26'55"S, 48°54'54"W
Lepidoziaceae				
<i>Bazzania hookeri</i> (Lindb.) Trevis. (Lindenberg 1845: 166, Trevisan 1877: 414) (Fig. 47)	AM	AM, BA, ES, MG, PR, RJ, RR, RS, SC, SP	ELS 325 (SP477495)	25°15'57"S, 48°32'42"W
<i>Bazzania jamaicensis</i> (Lehm. & Lindb.) Trevis. (Lehmann & Lindenberg 1838: 7, Trevisan 1877: 414)	MO	MG, PR, RJ, SC, SP	DFP 17713 (SP461254)	25°27'10"S, 48°55'11"W
<i>Bazzania pallide-virens</i> (Steph.) Fulford (Stephani 1908: 473, Fulford 1946: 42)	MO	AM, CE, GO, MT, RJ, RR	RR s.n. (SP462181)	25°26'49"S, 48°54'53"W
<i>Bazzania taleana</i> (Gottsche) Fulford (Gottsche 1863: 131, Fulford 1946: 54)	RA	ES, RJ, SC, SP	ELS 551 (SP477723)	25°18'20"S, 48°35'09"W
<i>Kurzia capillaris</i> (Sw.) Grolle (Swartz 1788: 144, Grolle 1963: 173)	AM	AM, BA, CE, DF, ES, GO, MG, MT, PA, PR, RJ, SC, SP	ELS 322 (SP477492)	25°15'57"S, 48°32'42"W
<i>Lepidozia coilophylla</i> Taylor, 1846	RA	MG, RJ, SC, SP	DFP 16147 (SP452573)	25°26'11"S, 48°55'14"W
<i>Lepidozia cupressina</i> (Sw.) Lindb. (Swartz 1788: 144, Lindenberg 1845: 207)	RA	BA, PE, RJ, SP	DFP 17843 (SP461385)	25°30'55"S, 48°58'58"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Lepidozia inaequalis</i> Lehm. & Lindb., 1845 (Fig. 48)	MO	BA, MG, PR, RJ, SC, SP	DFP 15942 (SP452364)	25°26'55"S, 48°54'54"W
<i>Paracromastigum pachyrhizum</i> (Nees) Fulford (Nees von Esenbeck in Martius 1833: 339, Fulford 1968: 390)	MO	DF, ES, GO, MG, RJ, SC, SP	ELS 384 (SP477554)	25°15'57"S, 48°32'42"W
<i>Telaranea diacantha</i> (Mont.) J.J. Engel & G.L. Merrill (Montagne 1845: 349, Engel & Merrill 2004: 145)	AM	AC, AM, BA, DF, ES, GO, PA, PE, PR, RJ, RS, SP	DFP 16263 (SP454842)	25°26'11"S, 48°55'14"W
<i>Telaranea nematodes</i> (Austin) M. Howe (Austin 1879: 302, Howe 1902: 284)	AM	AC, AM, BA, CE, DF, ES, GO, MG, MS, MT, RJ, RR, RS, SC, SE, SP	DFP 15944 (SP452366)	25°26'55"S, 48°54'54"W
<b>Marchantiaceae</b>				
<i>Dumortiera hirsuta</i> (Sw.) Nees (Swartz 1788: 145, Nees von Esenbeck in Martius 1833: 307)	AM	AC, AM, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RS, SC, SP	DFP 16013 (SP452436)	25°26'11"S, 48°55'14"W
<i>Marchantia chenopoda</i> L., 1753	AM	AM, DF, ES, GO, MG, MS, MT, PR, RJ, RS, SC, SP	DFP 16168 (SP452594)	25°26'11"S, 48°55'14"W
<i>Marchantia papillata</i> Raddi, 1822	MO	MG, MS, MT, PR, RJ, RS, SC, SP	DFP 16002 (SP452424)	25°26'11"S, 48°55'14"W
<b>Metzgeriaceae</b>				
<i>Metzgeria albinea</i> Spruce, 1889 (Fig. 49)	AM	BA, CE, ES, GO, MG, PE, PR, RJ, RS, SC, SP	DFP 16227 (SP454805)	25°26'11"S, 48°55'14"W
<i>Metzgeria bahiensis</i> Schiffn., 1911	RA/EN	BA, RS, SP	DFP 15770 (SP452192)	25°26'55"S, 48°54'54"W
<i>Metzgeria brasiliensis</i> Schiffn., 1964	MO/EN	AL, BA, PR, RJ, RS, SC, SP	DFP 16765 (SP460299)	25°27'00"S, 48°55'05"W
<i>Metzgeria conjugata</i> Lindb., 1875	MO	ES, MG, PR, RJ, RS, SP	DFP 15767 (SP452189)	25°26'55"S, 48°54'54"W
<i>Metzgeria consanguinea</i> Schiffn., 1893	RA	PR, RJ, SP	RR 4254 (SP458471)	25°26'57"S, 48°55'53"W
<i>Metzgeria fruticosa</i> Spruce, 1885	RA	AL, RJ, RS	RR s.n. (SP462177)	25°26'49"S, 48°54'53"W
<i>Metzgeria hegewaldii</i> Kuwah., 1981	RA	RS	DFP 17862 (SP461404)	25°30'55"S, 48°58'58"W
<i>Metzgeria myriopoda</i> Lindb., 1877	MO	DF, ES, GO, MG, PE, RJ, RS, SC, SP	DFP 15886 (SP452308)	25°26'55"S, 48°54'54"W
<i>Metzgeria uncigera</i> A. Evans, 1910	MO	ES, MG, PE, PR, RJ, RS, SC, SP	DFP 17917 (SP461459)	25°30'55"S, 48°58'58"W
<b>Pallaviciniaceae</b>				
<i>Symphogyna aspera</i> Steph., 1914	AM	AM, BA, CE, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RS, SC, SE, SP	DFP 16085 (SP452511)	25°26'11"S, 48°55'14"W
<i>Symphogyna brasiliensis</i> Nees, 1836 (Fig. 50)	AM	BA, CE, DF, ES, GO, MG, MT, PR, RJ, RO, RR, RS, SC, SP	DFP 15741 (SP452163)	25°26'55"S, 48°54'54"W
<i>Symphogyna brongniartii</i> Mont., 1843	MO	AC, AM, MG, MS, RJ, RS, SC, SP	DFP 16808 (SP460341)	25°26'15"S, 48°55'09"W
<i>Symphogyna podophylla</i> (Thunb.) Mont. & Nees (Thunberg 1800: 174, Montagne & Nees von Esenbeck 1846: 481)	MO	BA, CE, ES, MG, PR, RJ, SP	ELS 535 (SP477707)	25°18'20"S, 48°35'09"W
<b>Pelliaceae</b>				
<i>Notoclada confluens</i> (Hook. f. & Taylor) Spruce (Hooker & Taylor 1844: 478, Spruce 1885: 531)	MO	DF, ES, GO, MG, PR, RJ, RS, SC, SP	DFP 16810 (SP460343)	25°26'15"S, 48°55'09"W
<b>Plagiochilaceae</b>				
<i>Plagiochila adianthoides</i> (Sw.) Lindenb. (Swartz 1788: 142, Lindenberg 1840: 77) (Fig. 51)	MO	BA, ES, GO, MG, RJ, SP	DFP 15736 (SP452158)	25°26'55"S, 48°54'54"W
<i>Plagiochila aerea</i> Taylor, 1846	RA	BA, PE, RJ	DFP 17738 (SP461279)	25°27'10"S, 48°55'11"W
<i>Plagiochila bifaria</i> (Sw.) Lindb. (Swartz 1788: 145, Lindenberg 1843: 127)	MO	AM, BA, ES, MG, PA, RJ, SP	ELS 249 (SP459057)	25°30'49"S, 48°59'20"W
<i>Plagiochila corrugata</i> (Nees) Nees & Mont. (Nees von Esenbeck in Martius 1833: 378, Nees von Esenbeck & Montagne 1836: 52)	AM	AC, BA, CE, DF, ES, GO, MG, PE, PR, RJ, RS, SC, SE, SP	ELS 313 (SP477483)	25°15'57"S, 48°32'42"W
<i>Plagiochila cristata</i> (Sw.) Lindb. (Swartz 1788: 143, Lindenberg 1839: 33)	MO	AM, BA, ES, MG, PA, PR, RJ, RN, SP	ELS 447 (SP477618)	25°15'57"S, 48°32'42"W
<i>Plagiochila exigua</i> (Taylor) Taylor (Taylor 1843: 179, Taylor 1846: 264-265)	RA	BA, MG, RJ, SP	DFP 17668 (SP461209)	25°27'10"S, 48°55'11"W
<i>Plagiochila gymnocalycina</i> (Lehm. & Lindenb.) Mont. (Lehmann & Lindenberg in Lehmann 1833: 28, Montagne 1839: 2581)	MO	AL, BA, ES, MG, PE, RJ, SC, SP	ELS 312 (SP477482)	25°15'57"S, 48°32'42"W
<i>Plagiochila macrostachya</i> Lindb., 1840	MO	ES, MG, PA, PR, RJ, SC, SP	DFP 15960 (SP452382)	25°26'55"S, 48°54'54"W
<i>Plagiochila martiana</i> (Nees) Lindb. (Nees von Esenbeck 1831: 617, Lindenberg 1839: 12)	AM	AC, AL, BA, CE, DF, ES, GO, MG, MS, MT, PA, PE, PR, RJ, RS, SC, SP	ELS 259 (SP459067)	25°30'49"S, 48°59'20"W
<i>Plagiochila patentissima</i> Lindb., 1841	AM	BA, CE, ES, MG, PB, PE, PR, RJ, RS, SC, SP	DFP 16056 (SP452480)	25°26'11"S, 48°55'14"W
<i>Plagiochila patula</i> (Sw.) Lindb. (Swartz 1806: 1844, Lindenberg 1839: 21)	MO	AC, BA, MG, PA, PE, PR, RJ, SP	ELS 355 (SP477525)	25°15'57"S, 48°32'42"W
<i>Plagiochila rutilans</i> Lindb., 1840	AM	AC, AM, AP, BA, CE, ES, MG, MT, PA, PE, PR, RJ, RR, RS, SC, SP	ELS 359 (SP477529)	25°15'57"S, 48°32'42"W
<i>Plagiochila simplex</i> (Sw.) Lindb. (Swartz 1788: 145, Lindenberg 1840: 54)	AM	AM, BA, ES, MG, MT, PA, PE, PR, RJ, RS, SC, SP	ELS 525 (SP477697)	25°18'20"S, 48°35'09"W
<i>Plagiochila subplana</i> Lindb., 1840	AM	AM, AP, BA, ES, MG, MS, PA, PE, RJ, RR, SC, SP	DFP 15792 (SP452214)	25°26'55"S, 48°54'54"W
<b>Radulaceae</b>				
<i>Radula angulata</i> Steph., 1884	MO	ES, GO, MG, PE, PR, RJ, SC, SP	DFP 15834 (SP452256)	25°26'55"S, 48°54'54"W
<i>Radula brasílica</i> Yamada, 1993	RA/EN	SP	DFP 15765 (SP452187)	25°26'55"S, 48°54'54"W

Table 1. Continued.

Taxon	Dist. Brazil	Brazilian states	Voucher	Coordinates
<i>Radula elliotii</i> Castle, 1959	RA	SP	DFP 15817 (SP452239)	25°26'55"S, 48°54'54"W
<i>Radula gottscheana</i> Taylor, 1846	RA	AM, RJ	DFP 15731 (SP452153)	25°26'55"S, 48°54'54"W
<i>Radula javanica</i> Gottsche, 1845	AM	AC, AM, AP, BA, ES, MG, MS, MT, PA, PE, PR, RJ, RS, SC, SP	DFP 16809 (SP460342)	25°26'15"S, 48°55'09"W
<i>Radula mammosa</i> Spruce, 1890	RA	AM, RJ, SP	DFP 15870 (SP452292)	25°26'55"S, 48°54'54"W
<i>Radula mexicana</i> Lindb. & Gottsche, 1863	MO	BA, ES, PA, PE, RJ, RS, SP	ELS 376 (SP477546)	25°15'57"S, 48°32'42"W
<i>Radula nudicaulis</i> Steph., 1910	MO	ES, MG, PR, RJ, RS, SP	DFP 16018 (SP452441)	25°26'11"S, 48°55'14"W
<i>Radula recubans</i> Taylor, 1846 (Fig. 52)	AM	AL, BA, ES, MG, PA, PB, PE, PR, RJ, RS, SC, SP	ELS 262 (SP459070)	25°30'49"S, 48°59'20"W
<i>Radula schaefer-verwimpii</i> Yamada, 1990	RA	ES, MG, RJ, SP	DFP 17704 (SP461245)	25°27'10"S, 48°55'11"W
<i>Radula tenera</i> Mitt. ex Steph., 1884	RA	PE, PR, RJ, SP	DFP 17683 (SP461224)	25°27'10"S, 48°55'11"W
Scapaniaceae				
<i>Scapania portoricensis</i> Hampe & Gottsche, 1852	MO	ES, MG, PR, RJ, RR, SC, SP	ELS 490 (SP477662)	25°16'16"S, 48°33'04"W
Trichocoleaceae				
<i>Trichocolea argentea</i> Herzog, 1824	RA/EN	PR, RJ, SC, SP	RR 2804 (SP437095)	25°27'14"S, 48°55'12"W
<i>Trichocolea brevifissa</i> Steph., 1909	MO	ES, MG, PR, RJ, RS, SC, SP	ELS 553 (SP477725)	25°18'20"S, 48°35'09"W
<i>Trichocolea flaccida</i> (Spruce) Spruce (Spruce 1885: 349, Spruce 1888: 34)	RA	BA, PR, RJ, SP	ELS 261 (SP459069)	25°30'49"S, 48°59'20"W
<i>Trichocolea tomentosa</i> (Sw.) Gottsche (Swartz 1788: 145, Gottsche 1863: 119)	MO	ES, MG, PE, PR, RJ, RS, SC, SP	DFP 16757 (SP460291)	25°27'00"S, 48°55'05"W

## Discussion

**Bryophyte species richness.** The bryophytes found in our study of PMSP correspond to 40% of all the species recorded from the state of Paraná, although we found fewer species than indicated for this elevational range by Santos and Costa (2010). However, we found 41 Brazilian endemic species, including mosses and liverworts, which is near twice the total of endemic species (21) reported by Santos and Costa (2010). A possible explanation for this difference in species richness and endemism at PMSP is the high incidence of mist, as also observed by Montfoort and Ek (1990) in lowland tropical forests.

Moss species richness at PMSP was greater than that of liverworts, which is in accordance with findings by Santos (2011) in Rio de Janeiro, where liverworts had a greater species richness at lower elevations (between 50 m and 950 m) and mosses had a greater species richness at higher elevations (above 950 m). This pattern is well described in literature, and Gradstein et al. (2001) related the morphology of mosses to the colonization of open sites, like rock cliffs and exposed soils.

Some genera are better represented on our species list than others, and some are well represented by Brazilian endemics. Such genera include *Campylopus* with 16 species (2 endemic), *Fissidens* with 13 species (2 endemic), *Sphagnum* with 13 species (6 endemic), *Plagiochila* with 15 species, *Lejeunea* with 15 species (7 endemic), *Radula* with 12 species (1 endemic), *Frullania* with 10

species, and *Bazzania* with 9 species (1 endemic).

In the PMSP, the family Sphagnaceae had the greatest richness of endemic moss species found. The genus *Sphagnum* was found to be dominant in the cloud forests that we surveyed, and this dominance was earlier noted by IAP (2006). Sphagnaceae was among the families with the greatest contribution to the high species richness in High Mountain Forest habitats and elevational meadows of Itatiaia National Park, state of Rio de Janeiro (Costa et al. 2015). Species of the genus *Sphagnum* were found growing in dry habitats, such as on such rocks and tree trunks, as well as moist habitats, such as springs and bogs.

With 66 species, the family Lejeuneaceae represented 18% of our list. Citing 83 species from sea level to 1200 m above sea level, this family was identified by Santos (2011) as the most representative of the Atlantic Forest; Sonnleitner et al. (2001) recorded 59 species in a cloud forest in Monte Verde, Costa Rica but at a lower altitude, below 285 m. Lejeuneaceae is the richest family of the recent lineage of Porellales, with about 1000 species (Bechteler et al. 2016) and includes specialists at colonizing bark, branches, twigs, and leaves of trees. Thus, this family is frequently abundant in humid tropical forests (Lücking 1995, Pócs 1996, Gradstein 1997, Cornelissen and Ter Steege 1989, Sonnleitner et al. 2009).

The family Leucobryaceae was represented by 19 species (2 Brazilian endemics) in our study, and Costa et al. (2015) also observed this in Rio de Janeiro state.

Table 2. Bryophyte species number comparison among the Pico do Marumbi State Park with different geographic regions.

Area	Mosses	Liverworts	Reference
Marumbi	189	176	Present research
Paraná	591	309	Yano (2013); Yano (2014); Ristow et al. (2015)
Atlantic Forest	756	464	Costa (2009)
Brazil	880	633	Costa & Peralta (2015)
Tropical America	2600	1350	Gradstein et al. (2001)
World	13000	5000	Gradstein et al. (2001)

Among the Brazilian endemics found in PMSP, *Campylopus occultus* Mitt. (Frahm 1991), has its type locality in the state of Paraná. (Table 1).

**Conservation.** Our findings reinforce the importance of PMSP for the preservation of the bryophyte flora of Paraná and the recognition of diversity hot spots for prioritizing conservation efforts in areas with high species diversity and endemism (Kier et al. 2009, Costa et al. 2015). Furthermore, our findings expand the known distributions of several Brazilian endemic bryophytes. The bryophyte flora of PMSP has great importance to the conservation of these species with narrow to moderate distributions in Brazil. Of the species that we list, 95 are rare (50 liverworts and 45 mosses), 151 moderate (75 liverworts, 74 mosses, and 2 hornworts), and 118 widespread (60 mosses and 50 liverworts) (Table 1).

According to Costa and Peralta (2015), Brazil has 298 endemic species of bryophytes, with 73 species occurring in Paraná (24% of the total). In PMSP, 41 endemic species were found (25 mosses and 16 liverworts; see Results). Table 2 compares bryophytes richness in various regions of the world, and shows the significance of the PMSP for bryoflora.

Our study, based on new surveys, is a pioneering contribution to the knowledge of the bryoflora of the state of Paraná and Brazil.

## Acknowledgements

We thank IAP, the director of the park Mr Lothário Horst Stoltz Júnior, and other park staff for logistics and assistance during sampling. We greatly appreciate the contributions of Sionara Eliasaro, Juçara Bordin, Christopher Thomas Blum, and the reviewers.

## Authors' Contributions

ELS collected the data, wrote the text, and made the analysis; DMC and DFP also collected the data and made the analysis.

## References

- Allen BH, Magill RE (2003) A revision of *Pilotrichella* (Lembophylaceae, Musci). Acta Academiae Paedagogicae Agriensis, Sectio Biologiae, Nova Series 24: 43–83.
- Allen BH, Magill RE (2007) A revision of *Orthostichella* (Neckeraaceae). The Bryologist 110: 1–45. <http://doi.org/b68qqp>
- Angely J (1961) Musgos paranaenses: contribuição para o estudo e conhecimento da flora briológica do Paraná. Revista do Instituto Paranaense de Botânica 20: 1–7.
- Angely J (1965) Bryophytos Paranaenses. In: Angley J (Ed.) Flora Analítica do Paraná 1. Editora Phytos, São Paulo, 55–91.
- Angely J (1968) Bryophytos paranaenses. In: Angley J (Ed.) Flora Analítica do Paraná 7. Editora Phytos, São Paulo, 1–728.
- Angström J (1876) Primae lineae muscorum Cognoscendorum, qui ad Caldas Brasiliae sunt collecti. Öfersigt af kongliga vetenskaps-akademiens förhandlingar 33 (4): 2–50.
- Arnot W (1823) Notice sur quelques mousses de Rio-Janeiro. Mémoires de la Société d'histoire naturelle de Paris 1: 346–352.
- Austin CF (1879) Notes on hepaticology. Bulletin of the Torrey Botanical Club 6: 301–306.
- Bartram EB (1946) New species and new combinations of Guatemalan mosses. The Bryologist 49: 109–125.
- Bartram EB (1952) New mosses from southern Brazil. Journal of the Washington Academy of Sciences 42: 178.
- Bechteler J, Lee GE, Schafer-Verwimp A, Pócs TC, Peralta DF, Renner MAM, Schneider H, Heinrichs J (2016) Towards a monophyletic classification of Lejeuneaceae IV: reinstatement of *Allorgella*, transfer of *Microlejeunea aphanella* to *Vitalianthus* and refinements of the subtribal classification. Plant Systematics and Evolution 302: 187–201. <https://doi.org/10.1007/s00606-015-1252-8>
- Bescherelle É (1872) Prodromus bryologiae Mexicanae ou énumération des mousses de Mexique. Mémoires de la Société nationale des sciences naturelles de Cherbourg 16: 144–256.
- Bescherelle É, Spruce R (1890) Hepatiques nouvelles des colonies francaises. Bulletin de la Société botanique de France 36: 177–206.
- Bischler H (1964) Le genre *Drepanolejeunea* Steph. en Amérique Centrale et Méridionale. I. Revue bryologique et lichénologique 33: 15–179.
- Bischler H (1967) Le genre *Drepanolejeunea* Steph. en Amérique Centrale et Méridionale. II. Revue bryologique et lichénologique 35: 135–137.
- Bischler H, Bonner CEB, Miller HA (1963) Studies in Lejeuneaceae VI: The genus *Microlejeunea* Steph. in Central and South America. Nova Hedwigia 5: 359–411.
- Bordin J, Pursell RA, Yano O (2011) *Fissidens pseudophurisetus* sp. nov. Fissidentaceae, subgenus *Aloma*, from the Atlantic forest, Brazil. The Bryologist 114 (4): 785–789. <http://doi.org/xfx5bm>
- Bridel S (1801) Animadversiones in Muscologiae Recentiorum Tomum secundum, ab ipso auctore propositae. Journal of Botany (Schrader) 1800 (2): 268–299.
- Bridel S (1806-1812) Muscologiae recentiorum supplementum, Vol. 1. C.G. Ettinger, Gotha, 167 pp. <http://doi.org/cg3v>
- Bridel S (1819) Muscologia recentiorum supplementum. Pars IV seu Mantissa generum specierum que muscorum frondosorum universa. Gothae, apud A. Ukertum 4 (18): 1–220. <http://doi.org/cg3v>
- Bridel S (1827) Bryologia universa seu systematica ad novam methodum dispositio: historia et descriptio omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis. Pars II. J.A. Barth, Leipzig, 856 pp.
- Brotherus VF (1895) Nouvelles contributions à la Flore bryologique du Brésil. Bihang til kongliga svenska vetenskaps-akademiens handlingar 21 (3): 3–76.
- Brotherus VF (1900) Die Laubmoose der ersten Regnellschen Expedition. Bihang till kongliga svenska vetenskaps akademien handlingar. Stockholm 26: 1–65.
- Brotherus VF (1903) Musci (Laubmoose). III Unterklasse Bryales: II. Spezieller Teil, Gruppe I. Acrocarpi. In: A. Engler & K. Prantl (ed.), Die natürlichen Pflanzenfamilien. Teil 1, Abt. 3, Hälfte 1, Lief. 216. Leipzig: Engelmann, 481–528.
- Brotherus VF (1906) Musci Amazonici et subandini Uleani. Nova Hedwigia 45: 260–288.
- Brotherus VF (1916) Descriptions of some new species of Australian, Tasmanian and New Zealand mosses. Proceedings of the Linnean Society of New South Wales 41: 575–596.
- Brotherus VF (1925) Musci (Laubmoose) 1. Ergebnisse der botanische Expedition der Kaiserlichen Akademie der Wissenschaften nach Sudbrasilien 1901 (Thallophyta et Bryophyta). W. Engelmann, Leipzig, 251–358.
- Brotherus VF (1925) Musci (Laubmoose) 2. Hälfte. In: Engler A, Prantl K (Eds) Die natürlichen Pflanzenfamilien. Teil 1, Abt. 3, Hälfte 2, Lief. 216. Engelmann, Leipzig, 1–542.
- Bruggeman Nannenga, M. and R.A. Pursell. 1994. A re-evaluation of *Fissidens* subgenus *Serridium* section *Amblyothalia*. The Journal of the Hattori Botanical Laboratory 77: 255–271.
- Brummitt RK, Powell CE (1992) Authors of Plant Names. Royal Botanic Gardens, Kew, 732 pp.

- Buck WR (1983) New species and new combinations in the *Sematophyllum subpinnatum* complex Sematophyllaceae. *Brittonia* 35: 327–330.
- Buck WR (1984) Taxonomic and nomenclatural notes on West Indian Hypnaceae. *Brittonia* 36: 178–183.
- Buck WR (1987) Notes on Asian Hypnaceae and associated taxa. *Memoirs of The New York Botanical Garden* 45: 519–527.
- Buck WR (1987) Taxonomic and nomenclatural rearrangement in the Hookeriales with notes on West Indian taxa. *Brittonia* 39(2): 210–224.
- Buck WR (1990) Biogeography of the Greater Antillean mosses. *Tropical Bryology* 2: 35–48.
- Buck WR (1991) Notes of Neotropical Pterobryaceae. *Brittonia* 43 (2): 96–101.
- Buck WR (1994) The resurrection of *Orthostichella*. *The Bryologist* 97 (4): 434–435.
- Buck WR (1998) New combination and new synonymy in Brazilian Sematophyllaceae. *Nova Hedwigia* 66 (1): 241–246.
- Buck WR (1998) Pleurocarpous mosses of the West Indies. *Memoirs of the New York Botanical Garden* 1: 1–401.
- Buck WR, Vital DM (1992) *Paranapiacabaea paulista*, a new genus and species of Sematophyllaceae from southeastern Brazil. *Brittonia* 44: 339–343.
- Buck WR, Schäfer-Verwimp A (1993) A reassessment of *Schradobryum* (Sematophyllaceae). *Boletim do Museu Paraense Emílio Goeldi, nova série Botânica* 7 (2): 645–654.
- Cardot J (1915) Histoire physique, naturelle et politique de Madagascar, mousses. L'Imprimerie nationale, Paris, 469 pp.
- Castle H (1959) A revision of the genus *Radula*. Part II. Subgenus *Acroradula*. Section 4. Marginatae. *Revue bryologique et lichénologique* 28: 290–296.
- Churchill SP (1994) Catalog of Amazonian Mosses. *The Journal of the Hattori Botanical Laboratory* 85: 191–238.
- Churchill SP (1994) The mosses of Amazonian Ecuador. *Aarhus Universitetsforlag* 35: 1–211.
- Cornelissen JHC, Steege HT (1989) Distribution and ecology of epiphytic bryophytes and lichens in dry evergreen forest of Guyana. *Journal of Tropical Ecology* 5: 131–150.
- Costa DP (2009) Briófitas. In: Stehlmann JR, Forzza RC, Salino A, Cabral M, Costa DP, Kamino LHY (Eds) *Plantas da Floresta Atlântica*. Jardim Botânico do Rio de Janeiro, Rio de Janeiro, 13–17.
- Costa DP, Peralta DF (2017) Briófitas. Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. <https://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB128472>. Accessed on: 2017-2-25.
- Costa DP, Almeida JSS, Dias NS, Gradstein SR, Churchill SP (2010) *Manual de Briologia*. Editora Interciência, Rio de Janeiro, 222 pp.
- Costa DP, Imbassahy CAA, Silva VPAV (2005) Diversidade e importância das espécies de briófitas na conservação dos ecossistemas do Estado do Rio de Janeiro. *Rodriguésia* 56: 13–49.
- Costa DP, Pôrto KC, Luiz-Ponzo AP, Ilkiu-Borges AL, Bastos CJP, Câmara PEAS, Peralta DF, Bôas-Bastos SBV, Imbassahy CAA, Henriques DK, Gomes HCS, Rocha LM, Santos ND, Siviero TS, Vaz-Imbassahy TF, Churchill SP (2011) Synopsis of the Brazilian moss flora: checklist, distribution and conservation. *Nova Hedwigia* 93: 277–334.
- Costa DP, Santos ND, Rezende MA, Buck RW, Schäfer-Verwimp A (2015) Bryoflora of the Itatiaia National Park along an elevation gradient: diversity and conservation. *Biodiversity and Conservation* 24 (9): 2199–2212. <http://doi.org/b2ndn3>
- Crandall-Stotler B, Stotler RE, Long DG (2009) Morphology and classification of the Marchantiophyta. In: Goffinet B, Shaw AJ (Eds) *Bryophyte Biology* 2. Cambridge University Press, Cambridge, 1–54.
- Crum HA (1971) Nomenclatural changes in the Musci. *The Bryologist* 74: 165–174.
- Crum HA (1987) New species of *Sphagnum* from South America. *The Journal of the Hattori Botanical Laboratory* 63: 77–97.
- Crum HA (1992) Miscellaneous notes on the genus *Sphagnum*. 3. New species from Brazil. *The Bryologist* 95: 419–429. <https://doi.org/10.2307/3243565>
- Crum HA (1994) Miscellaneous notes on the genus *Sphagnum*. 5. New and notable species of South America. *The Journal of the Hattori Botanical Laboratory* 77: 233–253.
- Crum HA (1997) Miscellaneous notes on the genus *Sphagnum* 10. Contributions of the University of Michigan Herbarium 21: 147–159.
- Czumay A, Dong S, Scheben A, Schäfer-Verwimp A, Feldberg K, Heinrichs J (2013) Transfer of *Lejeunea huctumalcensis* to *Physoantholejeunea* (Lejeuneaceae, Porellales). *Australian Systematic Botany* 26: 386–392. <https://doi.org/10.1071/SB13039>
- Dixon HN (1939) High alpine mosses from Sumatra. *Annales Bryologiques* 12: 48–56.
- Dozy F, Molkenboer HJ (1854) Musci. In: Zollinger H (Ed.) *Systematisches Verzeichnis der im indischen Archipel in den Jahren 1842–1848 gesammelten sowie der aus Japan empfangenen Pflanzen* 1, E. Kiesling, Zurich, 22–33.
- Duby JE (1846) Musci. In: Zollinger H (Ed.) *Systematisches Verzeichnis der im indischen archipel in den Jahren 1842–1848 gesammelten sowie der aus Japan empfangenen Pflanzen* 1, E. Kiesling, Zurich, 130–135.
- Duby JE (1872) Choix de cryptogames exotiques nouvelles ou mal connues. I. Mousses (3me suite. Musci Welwitschiani), A. Acrocarpi. *Memoires de la Société de physique et d'histoire naturelle de Genève* 21: 215–227.
- Duby JE (1880) Aliquot diagnose muscorum novarum aut non rite cognitorum ab J.E. Duby. *Flora* 63: 168–174.
- Dumortier BCJ (1831) *Sylloge Jungermannidearum Europae indigenarum*. *Commentationes botanicae: observationes botanicae dédiées a la Société d'horticulture de Tournay* par B.-C. Ghent University, Ghent, 100 pp.
- Dumortier BCJ (1835) *Recueil d'observations sur les Jungermanniacées*. I. Révision des genres. *Commentationes botanicae: observationes botanicae dédiées a la Société d'horticulture de Tournay* par B.-C. Blanquart, Tournay, 27 pp.
- Engel JJ, Merrill GLS (2004) Austral Hepaticae. A taxonomic and phylogenetic study of *Telaranea* (Lepidoziaceae), with a monograph of the genus in temperate Australasia and commentary on extra-Australasian taxa. *Fieldiana Botany* 44: 1–265.
- Engel JJ, Schuster RM (1984) An overview and evaluation of the genera of Geocalycaceae subfamily Lophocoleoideae (Hepaticae). *Nova Hedwigia* 39: 385–463.
- Engler A, Prantl K (1893) Laubmooses. In: Engler A, Prantl K (Eds) *Die Natürlichen Pflanzenfamilien nebst ihren Gattungen und wichtigeren Arten, insbesondere den Nutzpflanzen, unter Mitwirkung zahlreicher hervorragender Fachgelehrten*. 37(7). W. Engelmann, Leipzig, 248–250.
- Enroth J (1990) Altitudinal zonation of bryophytes on the Huon Peninsula, Papua New Guinea. A floristic approach, with phytogeographic considerations. *Tropical Bryology* 2: 61–90.
- Enroth J (1990) Notes on the Neckeraceae (Musci) 3–7. *Homaliodendron piniforme* comb. nov. and new synonyms in *Porotrichum*, *Himantochalidum* and *Neolindbergia*. *Nova Hedwigia* 51 (3–4): 551–559.
- Evans AW (1904) Hepaticae of Puerto Rico. IV. *Odontolejeunea*, *Cyclolejeunea* and *Prionolejeunea*. *Bulletin of the Torrey Botanical Club* 31: 183–226.
- Evans AW (1905) Hepaticae of Puerto Rico V. *Ceratolejeunea*. *Bulletin of the Torrey Botanical Club* 32: 273–290.
- Evans AW (1906) Hepaticae of Puerto Rico. VI. *Cheilelejeunea*, *Rectolejeunea*, *Cystolejeunea*, and *Pycnolejeunea*. *Bulletin of the Torrey Botanical Club* 33: 1–25.
- Evans AW (1907) Hepaticae of Puerto Rico. VII. *Stictolejeunea*, *Neurolejeunea*, *Omphalanthus*, and *Lopholejeunea*. *Bulletin of the Torrey Botanical Club* 34: 1–34.
- Evans AW (1908) Hepaticae of Puerto Rico. IX. *Brachiolejeunea*, *Ptychocoleus*, *Archilejeunea*, *Leucolejeunea* and *Anoplolejeunea*. *Bulletin of the Torrey Botanical Club* 35: 155–179.

- Evans AW (1910) Notes on North American Hepaticae. I. The Bryologist 13: 33–36.
- Fleischer M (1923) Beitrag zur Laubmoosflora der Vulkaninsel Krakatau. Annales du Jardin botanique de Buitenzorg 33: 105–109.
- Frahm JP (1990) The altitudinal zonation of bryophytes on Mt. Kinabalu. Nova Hedwigia 51: 133–149.
- Frahm JP (1991) Dicranaceae: Campylopodioideae, Paraleucobryodeae. Flora Neotropica 54: 1–237.
- Frahm JP (1997) A taxonomic revision of *Dicranodontium* (Musci). Annales Botanici Fennici 34(3): 179–204.
- Frahm JP (2003) Manual of tropical bryology. Tropical Bryology 23: 1–196.
- Frahm JP, Gradstein SR (1991) An altitudinal zonation of tropical rain forests using bryophytes. Journal of Biogeography 18: 669–678. <https://doi.org/10.2307/2845548>
- Fulford M (1946) The genus *Bazzania* in Central and South America. Annales Cryptogamici et Phytopathologici 3: 1–175.
- Fulford M (1959) Studies on American Hepaticae. VII–XI. A supplement to the genus *Bazzania* in Central and South America. Part II. Tridentatae (3–5). Bulletin of the Torrey Botanical Club 86: 394–412.
- Fulford M (1968) Manual of the leafy Hepaticae of Latin America. Part III. Memoirs of the New York Botanical Garden 11(3): 277–392.
- Goffinet B, Shaw AJ (2009) Bryophyte Biology, 2nd edition. Cambridge University Press, Cambridge, 565 pp.
- Goffinet B, Buck WR, Shaw AJ (2009) Morphology, anatomy and classification of the Bryophyta. In: Goffinet B, Shaw AJ (Eds) Bryophyte Biology, 2nd edition. Cambridge University Press, Cambridge, 56–138.
- Gottsche CM (1863) De Mexikanske Levermooser, efter Prof. Fr. Liebmanns Samling. Kongelige Danske Videnskabernes Selskabs Skrifter. Naturvidenskabelige og Mathematisk Afdeling 6: 97–381.
- Gottsche CM (1864) Hepaticae. In: Triana J, Planchon JE (Eds) Prodromus Florae Novo-Granatensis. Annales des sciences naturelles, botanique 1: 95–198.
- Gottsche CM, Rabenhorst L (1872) Hepaticae Europaeae. Die Lebermoose Europa's unter Mitwirkung Mehrerer Namhafter Botaniker. Decades 53–55: 521–550.
- Gottsche CM, Lindenbergh JBG, Nees von Esenbeck SG (1843) Hepaticae. In: Meyen FJF (Ed.) Observationes botanicas in itinere circum terram institutas. Novorum Actorum Academia Caesarea Leopoldinae-Carolinae Naturae Curiosorum 19 (1): 469–477.
- Gottsche CM, Lindberg JBG, Nees von Esenbeck SG (1844) Synopsis Hepaticarum. Universidade de Ghent, Ghent. 835 pp.
- Gottsche CM, Lindberg JBG, Nees von Esenbeck SG (1845) Synopsis Hepaticarum. 1(2): 45–304. Universidade de Ghent, Ghent.
- Gottsche CM, Lindberg JBG, Nees von Esenbeck SG (1845) Synopsis Hepaticarum. 1(3): 305–464. Universidade de Ghent, Ghent.
- Gottsche CM, Lindberg JBG, Nees von Esenbeck SG (1846) Synopsis Hepaticarum. 1(4): 465–624. Universidade de Ghent, Ghent.
- Gottsche CM, Lindberg JBG, Nees von Esenbeck SG (1847) Synopsis Hepaticarum. 1 (5): 625–834. Universidade de Ghent, Ghent.
- Gradstein SR (1994) Lejeuneaceae: Ptychantheae, Brachirolejeuneae. Flora Neotropica, Monograph 61: 1–216.
- Gradstein SR (1997) The taxonomic diversity of epiphyllous bryophytes. Abstracta Botanica 21(1): 15–19.
- Gradstein SR, Costa DP (2003) The Hepaticae and Anthocerotae of Brazil. Memoirs of the New York Botanical Garden 87: 1–318.
- Gradstein SR, Grolle R, Schäfer-Verwimp A (1993) Two interesting species of Lejeuneaceae from Brazil. The Journal of the Hattori Botanical Laboratory 74: 39–70.
- Gradstein, S.R., D. Griffin III, M.I. Morales and N.M. Nadkarni 2001. Diversity and habitat differentiation of mosses and liverworts in the cloud forest of Monteverde, Costa Rica. Caldasia 23 (1) 203–212.
- Gradstein SR, Churchill SP, Salazar-Allen N (2001) Guide to the bryophytes of tropical America. Memoirs of the New York Botanical Garden 1: 1–577.
- Griffith W (1842) Muscologia itineris Assamici [I]; or, a description of mosses, collected during the journey of the Assam Deputation, in the years 1835 and 1836. Calcutta Journal of Natural History, and Miscellany of the Arts and Sciences in India 2: 465–514.
- Grolle R (1960) Zur Nomenklatur von *Riccardia pinguis*. Transactions of the British Bryological Society 3: 746–747.
- Grolle R (1962) *Goebelobryum*, eine neue marsupiale Lebermoosgattung. Journal of the Hattori Botanical Laboratory 25: 135–144.
- Grolle R (1963) [1964] Über *Kurzia* v. Martens. Revue bryologique et lichénologique 32: 166–180.
- Grolle R (1966) Miscellanea Hepaticologica 61–70. Journal of Japanese Botany 41: 225–232.
- Grolle R (1969) Miscellanea Hepaticologica 101–110. Österreichische Botanische Zeitschrift 117: 1–6.
- Grolle R (1969) Miscellanea Hepaticologica 91–100. Transactions of the British Bryological Society 5: 766–774.
- Grolle R (1977) Miscellanea hepaticologica 161–170. Journal of Bryology 9: 529–538.
- Grolle R (1979) Miscellanea hepaticologica 181–190. Journal of the Hattori Botanical Laboratory 45: 173–183.
- Grolle R (1984) Zur Kenntnis der Lejeuneoideae in Cuba (1): *Cyclolejeunea*. Mathematisch-naturwissenschaftliche Reihe 33: 759–764.
- Grolle R, Reiner ME (1997) *Cheilolejeunea oncophylla* (Aongstr.) Grolle & Reiner, comb. nov. (Lejeuneaceae), from the Neotropics. Journal of Bryology 19: 781–785.
- Grolle, R., R.-L. Zhu and Gradstein SR. 2001. On *Cyrtolejeunea* A. Evans (Lejeuneaceae, Hepaticae). Taxon 50: 1067–1074.
- Inoue H, Schuster RM (1971) A monograph of New Zealand and Tasmanian Plagioclilaceae. Journal of the Hattori Botanical Laboratory 34: 1–225.
- Hampe E (1844) Icones Muscorum novorum vel minus Cognitorum. Henry & Cohen, Bonn, 88 pp.
- Hampe E (1847) Berichte über die Hepaticae, welche Hr. Moritz in Columbien sammelte und dem königlichen Herbarium in Schönberg überlieferte, nach Synopsis Hepaticarum und den Moritschen Nummern aufgeführt. Linnaea 20: 321–326.
- Hampe E (1849) Musci frondosi in Plantae Regnellianae. Linnaea 22: 581.
- Hampe E (1852) Hepaticae Oerstedianae. Linnaea 24: 300–304.
- Hampe E (1862) Species novas muscorum ab Dr. Alexandro Lindigio in Nova-Granada mensibus Julio et Augusto a. 1861 collectas. Linnaea 31: 518–532.
- Hampe E (1863) Species novas muscorum ab Alexandro Lindigio in Nova-Granada collectas. Linnaea 32: 127–164.
- Hampe E (1865) Prodromus florae Novo-granatensis ou énumération des plantes de la Nouvelle-grenade. Annales des sciences naturelles, botanique 4 (5): 324–378.
- Hampe E (1870) Symbolae ad floram Brasiliae centralis cognoscendam. Videnskabelige meddelelser fra den naturhistoriske forening i Kjöbenhavn 8 (10): 267–296.
- Hampe E (1872). Musci frondosi. Symbolae ad floram Brasiliae centrales cognoscendam. Videnskabelige meddelelser fra dansk naturhistoriske Forening i Kjöbenhavn (serie 3) 10: 36–59.
- Hampe E (1874) Symbolae ad floram Brasiliae centralis cognoscendam. Videnskabelige meddelelser fra den naturhistoriske forening i Kjöbenhavn 19 (9): 129–178.
- Hampe E (1879) Enumeratio muscorum hactenus in provinciis Brasilienisub Rio de Janeiro et São Paulo detectorum. Videnskabelige meddelelser fra den naturhistoriske forening i Kjöbenhavn 1879: 73–164.
- Harvey WH (1836) Musci Indici. Icones plantarum. 1(1): 17–24.
- He XL, Grolle R (2001) *Xylolejeunea*, a new genus of the Lejeuneaceae (Hepaticae) from the Neotropics, Madagascar and the Seychelles. Annales Botanici Fennici 38: 25–44.
- Hedwig J (1801) Species muscorum frondosorum: descriptae et tabulis aeneis LXXVII coloratis illustratae. Barthius 1 (18): 1–353.
- Hell K (1969) Briófitas talosas dos Arredores da Cidade de São Paulo (Brasil). Ph.D. thesis, Universidade de São Paulo, São Paulo, 225 pp.

- Herzog T (1916). Die bryophyten meiner zweiten Reise durch Bolivia. *Bibliotheca Botanica* 87 (1): 1–168.
- Herzog T (1924). Contribuições ao conhecimento da flora bryologica do Brasil. *Archivos de Botanica do Estado de São Paulo* 1 (2): 27–105.
- Herzog T (1927). Zwei Bryophytensammlungen aus Süd-Amerika. *Hedwigia* 67: 1–249.
- Herzog T (1931). Die Bryophyten der Lützelburgschen Reisen in Nordbrasilien. *Hedwigia* 71: 332–350.
- Herzog T (1950). *Miscellanea bryologica. I. Neotropica. Memoranda Societatis pro Fauna et Flora Fennica* 25: 43–72.
- Herzog T (1955). Hepaticae aus Columbia und Peru. *Feddes Repertorium Specierum Novarum Regni Vegetabilis* 57: 156–203.
- Hirai RY, Yano O, Ribas MEG (1998). Musgos da mata residual do Centro Politécnico (Capão da Educação Física), Curitiba, Paraná, Brasil. *Boletim do Instituto de Botânica* 11: 81–118.
- Hooker JD, Thomson T (1854). On *Maddenia* and *Diplarche*, new genera of Himalayan plants. *Hooker's Journal of Botany and Kew Garden Miscellany* 4: 1–4.
- Hooker WJ (1816). *British Jungermanniae: being a History and Description, with Figures, of each Species of the Genus, and Microscopical Analysis of the Parts*. Longman, Hurst, Rees, Orme, and Brown; Sherwood, Neely, and Jones, London, 986 pp.
- Hooker WJ (1816). *Plantae Cryptogamicae, quas in plagra orbis novi aequinoctali collegerunt Alexander de Humboldt et Amat. Bonpland*. IDC, Leiden, 1–4.
- Hooker WJ (1818–1819). *Musci Exotici*. Longmans, London, 1: 1–96.
- Hooker WJ, Greville RK (1824). Sketch of the characters of the species of Mosses, belonging to the Genera *Orthotrichum* (Including *Schlotheimia*, *Macromitrium* and *Ulota*), *Glyphomitrium* and *Zygodon*. *Edinburgh Journal of Science* 1: 110–132.
- Hooker WJ, Wilson W (1844). Enumeration of the mosses and hepaticae, collected in Brazil by George Gardner, Esq. drawn by Sir W.J. Hooker and W. Wilson, Esq. *Journal of Botany* 3: 149–167.
- Hornsusch F (1840). Musci. In: Martius CFP (Ed.) *Flora Brasiliensis enumeratio plantarum in Brasilia hactenus detectarum quas suis aliorumque botanicorum studiis descriptas et methodo naturali digestas partim icone illustratas*. *Flora Brasiliensis* 1(2): 1–172.
- Howe M (1902). Notes on American Hepaticae. *Bulletin of the Torrey Botanical Club* 29: 281–289.
- IBGE (2012). *Manual Técnico da Vegetação Brasileira*, 2nd edition. *Manuais Técnicos em Geociências*. Ministério do Planejamento, Orçamento e Gestão Instituto Brasileiro de Geografia e Estatística, Diretoria de Geociências Coordenação de Recursos Naturais e Estudos Ambientais, Rio de Janeiro, 271 pp.
- IAP (1996). Plano de manejo Parque Estadual Pico do Marumbi, 1996. [https://www.iap.pr.gov.br/arquivos/File/Plano\\_de\\_Manejo/Parque\\_Estadual\\_Pico\\_do\\_Marumbi/PM\\_PE\\_Marumbi.pdf](https://www.iap.pr.gov.br/arquivos/File/Plano_de_Manejo/Parque_Estadual_Pico_do_Marumbi/PM_PE_Marumbi.pdf). Accessed on: 2014-4-7.
- IAP (2012). Parque Estadual Pico do Marumbi. Cadastro Nacional de Unidades de Conservação. <https://sistemas.mma.gov.br/portalcnucrel/index.php?fuseaction=portal.exibeUc&idUc=540>. Accessed on: 2016-2-9.
- Inoue H (1974). Two new species of *Syzygiella* Spruce from Colombia, South America. *Bulletin of the National Science Museum, Tokyo* 17: 301–305.
- Jaeger A (1872). *Adumbratio flore muscorum totius orbis terrarum. Part 2. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft während des Vereinsjahres 1870–1871*: 357–451.
- Jaeger A (1875). *Adumbratio flore muscorum totius orbis terrarum. Part 5. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft während des Vereinsjahres 1873–1874*: 53–278.
- Jaeger A, Sauerbeck F (1877). *Adumbratio flore muscorum totius orbis terrarum. Part 7. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft während des Vereinsjahres 1875–1876*: 201–371.
- Jaeger A Sauerbeck F (1878). *Adumbratio flore muscorum totius orbis terrarum. Part 8. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft während des Vereinsjahres 1876–1878*: 211–454.
- Jaeger A Sauerbeck F (1880). *Adumbratio flore muscorum totius orbis terrarum. Index. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft während des Vereinsjahres 1878–1879*: 213–252.
- Jennings OE (1913). *Brachythecium pacificum*, new species. *The Bryologist* 16: 95–96.
- Jovet-Ast S (1947). Hépatiques des Antilles françaises récoltées par P. et V. Allorge en 1936. *Revue bryologique et lichénologique* 16: 17–46.
- Kessler M (2000). Altitudinal zonation of Andean cryptogam communities. *Journal of Biogeography* 27: 275–282. <http://doi.org/cgcnb5>
- Kindberg NC (1888). *Enumeratio Bryinearum Exoticarum*. *Officina Corresp. Ostrogoth, Linköping*, 84 pp.
- Koponen T (1971). A monograph of *Plagiomnium* sect. *Rosulata* (Mniaceae). *Annales Botanici Fennici* 8: 305–367.
- Krauss F (1846). *Pflanzen des Cap- und Natal-Landes, geammelt und zusammengestellt von Dr. Ferdinand Krauss*. *Flora* 29: 129–138.
- Kruijt RC, Gradstein SR (1986). Studies on Lejeuneaceae subfam. Ptychanhoideae X. On *Brachiolejeunea phyllorhiza* (Nees) Kruijt & Gradstein comb. nov. (Hepaticae). *Nova Hedwigia* 43: 299–309.
- Kummrow RP, Prevedello SM (1982). Lista de musgos paranaenses do MBM. *Boletim do Museu Botânico Municipal* 54: 1–36.
- Kuntze O (1891). *Revisio generum plantarum vascularium omnium atque cellularium multarum leges nomenclaturae internationales cum enumeratione plantarum exoticarum in itinere mundi collectarum. 2: 375–1011*. A. Felix, Leipzig.
- Kunze G (1834). *Synopsis plantarum cryptogamicarum ab E. Poeppig*. *Linnaea* 1–757.
- Kury AB (2006). Diretrizes e estratégias para a modernização de coleções biológicas brasileiras e a consolidação de sistemas integrados de informação sobre biodiversidade. Brasília: Centro de Gestão e Estudos Estratégicos. Ministério da Ciência e Tecnologia. 43 pp. [https://www.dpi.inpe.br/referata/arq/12\\_candinha/11194.pdf](https://www.dpi.inpe.br/referata/arq/12_candinha/11194.pdf). Accessed on: 2016-10-16.
- Kuwahara Y (1981). Studies on Peruvian collections of the genus *Metzgeria* made by P. & E. Hegewald in 1973 and 1977. *Nova Hedwigia* 34: 769–815.
- Lehmann JGC (1829). *Hepaticarum capensium a C.F. Ecklon*. *Linnaea* 4: 357–371.
- Lehmann JGC (1833). *Novarum et Minus Cognitarum Stirpium Pugillum V addita enumeratione plantarum omnium in his pugillis descriptarum*. Meissner, Hamburg, 28 pp.
- Lehmann JGC (1834). *Novarum et Minus Cognitarum Stirpium Pugillum VI addita enumeratione plantarum omnium in his pugillis descriptarum*. Meissner, Hamburg, 66 pp.
- Lehmann JGC (1838). *Novarum et Minus Cognitarum Stirpium Pugillum VII addita enumeratione plantarum omnium in his pugillis descriptarum*. Meissner, Hamburg, 41 pp.
- Lehmann JGC (1857). *Novarum et Minus Cognitarum Stirpium Pugillum X addita enumeratione plantarum omnium in his pugillis descriptarum*. Meissner, Hamburg, 40 pp.
- Limpricht KG (1892). *Die Laubmoose Deutschlands, Oesterreichs und der Schweiz* 2. E. Kummer, Leipzig, 445 pp.
- Lindberg GA (1891). *Zum Kakteenstudium*. *Monatsschrift für Kakteenkunde* 1: 62–71.
- Lindberg SO (1863). *Bidrag till mossornas synonymi. Bihang till Kongl. Svenska vetenskaps-akademiens handlingar* 20 (7): 385–418.
- Lindberg SO (1875). *Hepaticae in Hibernia mense Julii 1873 lectae*. *Acta Societatis Scientiarum Fennicae* 10: 465–559.
- Lindberg SO (1877). *Monographia Metzgeriae*. *Societatis Litterariae Fennicae, Helsinki*, 47 pp.
- Lindenberg JBG (1829). *Synopsis Hepaticarum Europaeorum, Adnexis Observationibus et Adnotationibus Criticis Illustrata, Auctore I. B. G. Lindenberg, Ditionis Bergedorfensis Praefecto. Cum Tabulis*



- Duabus. Bonn. Nova Acta Academiae Caesareae Leopoldino-Carolinae Germanicae Naturae Curiosorum 14: 1–133.
- Lindenberg JBG (1839–1844) Species hepaticarum 1–5. Henry et Cohen, Bona, 359 pp.
- Linnaeus C (1753) Species Plantarum, ed. 1. Impensis G.C. Nauk, Stockholm, 1200 pp.
- Lücking A (1995) Diversität und Mikrohabitatpräferenzen epiphyller Moose in einem tropischen Regenwald in Costa Rica unter besonderer Berücksichtigung der Lejeuneaceae. Fakultät für Naturwissenschaften der Universität Ulm, Ulm, Germany 6: 1–211.
- Malombe I (2009) Studies on African *Cheilolejeunea* (Lejeuneaceae) I: new species and new combinations. Acta Botanica Hungarica 51: 315–328.
- Manuel M (1977) A monograph of the genus *Zelometeorium* Manuel, gen. nov. (Bryopsida: Meteoriaceae). The Journal of the Hattori Botanical Laboratory 43: 107–126.
- Martius KFP (1817) Flora Cryptogamica Erlangensis. J.L. Schrag, Norinbergae, 512 pp.
- Martius KFP (1833) Flora Brasiliensis seu Enumeratio Plantarum 1. Amsterdam: sumptibus J. G. Cottae. 390 pp.
- Martius KFP (1834) Icones plantarum cryptogamicarum, quas in itinere annis 1817–1820 per Brasilien instituto collegit et descripsit. Monachii: Impensis auctoris. 152 pp.
- Massalongo C (1885) Epatiche raccolte alla Tierra del Fuoco del Dott. C. Spegazzini nell' anno 1882. Nuovo Giornale Botanico Italiano 17: 201–277.
- Meenks JLD (1987) Studies on Colombian cryptogams XXVIII. A guide to the tropical Andean species of *Riccardia* (Hepaticae). The Journal of the Hattori Botanical Laboratory 62: 161–182.
- Menzel M (1992) Preliminary checklist of the mosses of Peru (Studies on Peruvian Bryophytes IV). The Journal of the Hattori Botanical Laboratory 71: 175–254.
- Mitten W (1859) Musci Indiae Orientalis, an enumeration of the mosses of the East Indies. Journal of the Proceedings of the Linnean Society, Botany 1: 1–96.
- Mitten, W. 1860. Hepaticae Indiae Orientalis; an enumeration of the species of the Hepaticae of East Indies. Journal of the Proceedings Linnean Society of London, Botany 5: 89–108.
- Mitten W (1864) On some species of Musci and Hepaticae, additional to the floras of Japan and the coast of China. Journal of the Proceedings of the Linnean Society, Botany 8: 148–162.
- Mitten W (1868) A list of the Musci collected by the Rev. Thomas Powell in the Samoa or Navigator's Islands. Journal of the Proceedings of the Linnean Society, Botany 10: 166–195.
- Mitten W (1869) Musci austro-americi. Journal of the Proceedings of the Linnean Society, Botany 12: 1–659.
- Mitten W (1887) Musci. In W. B. Hemsley, Report on the vegetation of Diego Garcia. Journal of the Proceedings of the Linnean Society, Botany 22: 339–340.
- Montagne JFC (1838) Centurie I. de plantes cellulaires exotiques nouvelles (suite). Annales des sciences naturelles, botanique 9 (4): 38–57.
- Montagne JFC (1839) Cryptogamae Brasilienses seu Plantae cellulares, quas in itinere per Brasiliam a celeb. Auguste de Sainte-Hilaire collectas, recensuit observationibusque nonnullis illustravit. Annales des sciences naturelles, botanique 12 (4): 42–55.
- Montagne JFC (1840) Seconde Centurie de plants cellulaires exotiques nouvelles. Décades VI, VII et VIII. Journal of the Proceedings of the Linnean Society, Botany 14: 331–337.
- Montagne JFC (1843) Troisième centurie de plantes cellulaires exotiques nouvelles. Annales des sciences naturelles, botanique 19: 241–282.
- Montagne JFC (1845) Cinquième centurie de plantes cellulaires exotiques Nouvelles Décades I à VI. Annales des sciences naturelles, botanique sér. 3 4: 86–123.
- Montagne JFC (1856) Huitième centurie de plantes cellulaires nouvelles tant indigènes qu'exotiques, decades I à III. Annales des sciences naturelles, botanique, sér. 4 6: 179–199.
- Müller C (1846) Nachtragliche Bemerkungen über die von Gardner in Brasilien gesammelten Laubmoose. Botanische Zeitung 3: 105–111.
- Müller C (1847) De muscis nonnullis novis vel minus cognitis exoticis. Botanische Zeitung 5: 801–806, 825–830.
- Müller C (1848) Plantae Kegelinae Surinamenses: musci frondosi. Linnaea 21: 181–200.
- Müller C (1850–1851) Synopsis muscorum frondosorum omnium hucusque cognitorum 2: musci vegetation plerocarpicae. Sumptibus A. Foerstner, Berlin, 772 pp.
- Müller C (1855) De muscis novis, incomplete descriptis, neglectis criticis. Botanische Zeitung 13: 745–789.
- Müller C (1862) Addimenta ad synopsis muscorum nova. Botanische Zeitung 20: 327–393.
- Müller C (1876) Musci Hildebrandtiani in Archipelago Comorense et in Somalia littoris Africani anno 1875 ab I. M. Hildebrandt lecti. Linnaea 40: 225–300.
- Müller C (1879) Musci Africae Orientali-tropicae Hildebrandtiani. Flora 62: 376–380.
- Müller C (1885) Bryologia Fuegiana. Flora 68: 391–429.
- Müller C (1898) Bryologia Serrae Itatiaiae (Minas Gerais Brasiliae) adjectus nonnullis speciosis affinis regionum vicinarum. Bulletin de le Herbarier Boissier 6(2): 18–48.
- Müller C (1900) Symbolae ad bryologiam Brasiliae et regionum vicinarum. Hedwigia 39: 235–289.
- Müller C (1901) Genera muscorum frondosorum. Gattungen und Gruppen Der Laubmoose inhistorischer und systematischer Beziehung sowie nach ihrer geographischen Verbreitung unter Berücksichtigung Der Arten. Leipzig: Kummer. (Handschriftlicher Nachlaß hrsg. v. Schliephacke). 474 pp.
- Nees von Esenbeck CG, Montagne JFC (1842) Familia X Hepaticae, Juss. In: Ramon de la Sagra (Ed.) Histoire physique, politique et naturelle de l'Île de Cuba. Arthus Bertrand, Paris, 427–492.
- Nees von Esenbeck CG (1831) Berichtigungen zur Enumeratio plantarum cryptogamicarum Javae. Linnaea 6: 602–623.
- Nees von Esenbeck CG (1833) Beiträge zur Naturgeschichte der deutschen Lebermoose. Flora 16: 401–412.
- Nees von Esenbeck CG (1836) Excursus de solano wightii quem arnotii collegae observationibus. Nova acta Academiae Caesareae Leopoldino-Carolinae Germanicae Naturae Curiosorum 357–360.
- Nees von Esenbeck CG (1838) Naturgeschichte der Europäischen Lebermoose mit besonderer Beziehung auf Schlesien und de Örtlichkeiten des Riesengebirge. Breslau 4: 1–540.
- Nees von Esenbeck CG (1838) Naturgeschichte der Europäischen Lebermoose mit besonderer Beziehung auf Schlesien und de Örtlichkeiten des Riesengebirge. Breslau 3: 1–504.
- Nees von Esenbeck CG, Montagne JFC (1836) Jungermanniearum Herbarii Montagneani species. Annales des sciences naturelles, botanique 5 (4): 52–72.
- Ochyra R (1999) New combinations in Neotropical mosses. Fragmenta Floristica et Geobotanica 44 (2): 255–259.
- Ochyra R, Zarnowiec J, Bednarek-Ochyra H (2003) Census catalogue of Polish mosses. Biodiversity on Poland 3: 1–372.
- Olsson S, Enroth J, Buchbender V, Hedenäs L, Huttunen SM, Quandt D (2011) *Neckera* and *Thamnobryum* (Neckeraceae, Bryopsida): paraphyletic assemblages. Taxon 60 (1): 36–50.
- Pagán FM (1939) A preliminary list of the hepaticae of Puerto Rico including Vieques and Mona Island. The Bryologist 42: 1–12.
- Palisot de Beauvois AMFJ (1805) Prodromus of Aethogamia, or of a treatise on those families of plants whose fructification is extraordinary. Annales Botanici 2: 218–251.
- Paris EG (1898) Index Bryologicus sive enumeratio muscorum hucusque cognitorum adjunctis synonyma distributioneque geographica locupletissimis quem conscripsit. Paul Klincksieck, Paris, 1379 pp.
- Paris EG (1900) Muscinées du Tonkin et de Madagascar (suite). Revue bryologique 27: 76–80.
- Paris EG (1901) Muscinées de la Côte de l'Ivoire et du Quang Tcheou Wan. Revue bryologique 28: 15–17.

- Paris EG (1904) Index Bryologicus sive enumeratio muscorum ad diem ultimam anni 1900 cognitorum adjunctis synonymia distributioneque geographica locupletissimis. 2nd edition. Hermann, Paris, 375 pp.
- Paris EG (1905) Muscinees de l'Afrique occidentale française. Revue bryologique 32: 101–104.
- Peralta DF, Yano O (2008) Briófitas do Parque Estadual da Ilha Anchieta, Ubatuba, estado de São Paulo. Iheringia, série botânica 63: 101–127.
- Peralta DF, Yano O (2012) Briófitas da Serra do Itapeti. Aspectos históricos, sociais e naturalísticos. Canal 6: 75–86.
- Pócs TC (1996) Epiphyllous liverwort diversity at worldwide level and its threat and conservation. Anales del Instituto de Biología de la Universidad Nacional Autónoma de México, botánica 67: 109–127.
- Pócs TC (2008) Contributions to the bryoflora of Australia, III. The genus *Jubula* Dumort., with the description of *J. hutchinsiae* Hook. subsp. nov. *australiae* (Jubulaceae, Jungermanniopsida. Nova Hedwigia 86: 231–236.
- Pursell RA (2007) Fissidentaceae. Flora Neotropica Monograph 101. New York Botanical Garden, New York, 278 pp.
- Raddi G (1822) Di Alcune specie nuove di rettili, e piante Brasiliane. Memorie della Società Italiana delle Scienze residente in Modena 18: 3–39.
- Reese WD (1978) The genus *Syrrophodon* in the Americas II. The limbate species. The Bryologist 81: 189–225.
- Reiner-Drehwald ME, Goda A (2000) Revision of the genus *Crossotolejeunea* (Lejeuneaceae, Hepaticae). The Journal of the Hattori Botanical Laboratory 89: 1–54.
- Reinwardt CGC, Hornschuch CF (1829) Musci frondosi Javanici, redditi coniunctis studiis et opera. Nova Acta Physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum Exhibentia Ephemerides sive Observationes Historias et Experimenta 14 (2): 697–731.
- Renzaglia KS, Villarreal JC, Duff RJ (2009) New insights into morphology, anatomy and systematics of hornworts. In: Goffinet B, Shaw AJ (Eds) Bryophyte Biology. 2nd edition. Cambridge University Press, Cambridge, 139–171.
- Roderjan CV, Galvão F, Kuniyoshi YS, Hatschbach GG (2002) As unidades fitogeográficas do estado do Paraná, Brasil. Ciência & Ambiente 24: 75–92.
- Santos ND (2011) Distribuição espacial de briófitas na floresta Atlântica Ph.D. thesis, Universidade Estadual de Campinas, Campinas, 149 pp.
- Santos ND, Costa DP (2008) A importância de Reservas Particulares do Patrimônio Natural para a conservação da brioflora da Mata Atlântica: um estudo em El Nagual, Magé, RJ, Brasil. Acta Botanica Brasilica 22: 359–372.
- Santos ND, Costa DP (2010) Altitudinal zonation of liverworts in the Atlantic Forest, southeastern, Brazil. The Bryologist 113: 631–645. <https://doi.org/10.1639/0007-2745-113.3.631>
- Sastre-de Jesús I, Buck WR (1993) Annotated checklist of the mosses of Puerto Rico. Caribbean Journal of Science 29: 226–234.
- Schäfer-Verwimp A (1998) Zur Kenntnis der *Sphagnum*. Flora Brasiliensis mit Beschreibung von *Sphagnum crumii* sp. nov. und *Sphagnum delamboyense* sp. nov. Nova Hedwigia 67 (3): 409–419.
- Schäfer-Verwimp A, Peralta DF, Siqueira SMC (2012) *Frullania curvibulba* (Frullaniaceae, Marchantiophyta), a new species from Brazil. Phytotaxa 57: 27–30. <https://doi.org/10.11646/phytotaxa.57.1.5>
- Scheer MB, Moco-chinski AY, Roderjan CV (2011) Estrutura arbórea da Floresta Ombrófila Densa Altomontana de serras do sul do Brasil. Acta Botanica Brasilica 25 (4): 735–750.
- Schiffner V (1893) Ueber Exotische Hepaticae, hauptsächlich Aus Java, Amboina und Brasilien, nebst Einigen morphologischen und kritischen Bemerkungen über Marchantia. Nova acta physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosum 60 (2): 217–316.
- Schiffner V (1910) Über die Gattungen *Chiloscyphus* und *Heteroscyphus* n. gen. Österreichische Botanische Zeitschrift 60: 169–173.
- Schiffner V (1911) Über einige neotropische Metzgeria-Arten. Österreichische botanische Zeitschrift 61: 183–187.
- Schiffner V (1964) Ergebnisse der Botanischen Expedition der kaiserlichen Akademie der Wissenschaften nach Südbrasilien 1901. II. Thalophyta und Bryophyta. Hepaticae (Lebermoose. Österreichische Akademie der Wissenschaften). Mathematisch-Naturwissenschaftliche Klasse, Denkschriften 111: 1–156.
- Schimper WP (1852) Thedenia, Anisodon, Pseudoleskea, Heterocladium, Thuidium, Hylocomium, Thamnium, Rhynchostegium. Bryologia Europaea 5:49–51.
- Schimper WP (1856) Corollarium Bryologiae Europaeae, Conspectum Diagnosticum Familiarum, Generum et Specierum, Adnotationes Novas at que Emendationes Complectens, scriptis. Sumptibus Librariae E. Schweizerbart, Stuttgartiae, 164 pp.
- Schuster RM (1956) North American Lejeuneaceae. V. Schizostipae: *Ceratolejeunea*. Journal of the Elisha Mitchell Scientific Society 72: 292–316.
- Schuster RM (1980) The Hepaticae and Anthocerotae of North America. IV. Columbia University Press, New York, 1133 pp.
- Schuster RM (1995) Phylogenetic and taxonomic studies of Jungermanniidae, III. Calypogeiaceae. Fragmenta Floristica et Geobotanica 40: 825–888.
- Schwägrichen CF (1814) Historiae muscorum hepaticorum prodromus: commentatio qua hortum botanicum lipsiensem. I.A. Barth, Leipzig, 1–39.
- Sehnem A (1969) Musgos sul brasileiros. Pesquisas, série Botânica 27: 1–36.
- Sehnem A (1970) Musgos sul brasileiros 2. Pesquisas, série Botânica 28: 1–96.
- Sehnem A (1972) Musgos sul brasileiros 3. Pesquisas, série Botânica 29: 1–70.
- Sehnem A (1976) Musgos sul brasileiros 4. Pesquisas, série Botânica 30: 1–79.
- Sehnem A (1978) Musgos sul brasileiros 5. Pesquisas, série Botânica 32: 1–170.
- Sehnem A (1979) Musgos sul brasileiros 6. Pesquisas, série Botânica 33: 1–149.
- Sehnem A (1980) Musgos sul brasileiros 7. Pesquisas, série Botânica 34: 1–121.
- Sharp AJ, Crum H, Eckel P (1994) The Mosses flora of Mexico. Memoirs of the New York Botanical Garden 69: 1–1113.
- Shaw AJ, Goffinet B (2000) Bryophyte Biology. Cambridge University Press, Cambridge, 368 pp.
- Smith JE (1806) English Botany 23: 1585–1650.
- Smith JE (1813) English Botany 36: 1–150.
- Sonnleitner M, Dullinger S, Wanek W, Zechmeister H (2009) Microclimatic patterns correlate with the distribution of epiphyllous bryophytes in a tropical lowland rain forest in Costa Rica. Journal of Tropical Ecology 25: 321–330. <https://doi.org/10.1017/S0266467409006002>
- Söderström L, Hagborg A, von Konrat M, Bartholomew-Began S, Bell D, Briscoe L, Brown E, Cargill DC, Cooper ED, Costa DP, Crandall-Stotler BJ, Dauphin G, Engel JJ, Feldberg K, Glenn D, Gradstein SR, He X, Heinrichs J, Hentschel J, Ilkiu-Borges AL, Katagiri T, Konstantinova NA, Larrain J, Long DG, Nebel M, Pócs T, Puche F, Reiner-Drehwald ME, Renner MAM, Sass-Gyarmati A, Schäfer-Verwimp A, Segarra-Moragues JG, Stotler RE, Sukkharak P, Thiers BM, Uribe J, Váña J, Villarreal JC, Wigginton M, Zhang L, Zhu R-L (2016) World checklist of hornworts and liverworts. PhytoKeys 59: 1–828. <https://doi.org/10.3897/phytokeys.59.6261>
- Spence J (1996) *Rosulabryum* genus novum (Bryaceae). The Bryologist 99 (2): 221–225.
- Sprengel C (1827) Caroli Linnaei Systema vegetabilium 16 ed. 4 (1): 1–592. Dieterich, Göttingen.
- Spruce R (1884) Hepaticae of the Amazon and the Andes of Peru and Ecuador. I. Transactions and Proceedings of the Botanical Society of Edinburgh 15: 1–308.
- Spruce R (1885) Hepaticae of the Amazon and the Andes of Peru and

- Ecuador. II. Transactions and Proceedings of the Botanical Society of Edinburgh 15: 309–588.
- Spruce R (1888) Hepaticae in prov. Rio de Janeiro a Glaziou lectae, a R. Spruce determinatae. Revue bryologique et lichénologique 15: 33–34.
- Spruce R (1889) Hepaticae novae americanae tropical et aliae. Bulletin de la Société botanique de France 36: 188–224.
- Spruce R (1890) Hepaticae Novae Americanae Tropicae. Bulletin de la Société botanique de France 11: 202–202.
- Stephani F (1884) Die Gattung *Radula*. Hedwigia 23: 113–116.
- Stephani F (1885) Hepaticarum species novae vel minus cognitae II–VIII. Hedwigia 24: 166–168.
- Stephani F (1888) Westindische Hepaticae. Hedwigia 27: 276–302.
- Stephani F (1890) Die Gattung *Lejeunea* im Herbarium Lindenberg 2. Hedwigia 29: 68–99.
- Stephani F (1893) Hepaticarum species novae. I. Hedwigia 32: 17–29.
- Stephani F (1895) Hepaticarum species novae VII. Hedwigia 34: 43–65.
- Stephani F (1896) Hepaticarum species novae IX. Hedwigia 35: 73–140.
- Stephani F (1898–1900) Species Hepaticarum I. George & Cia, Libraires-Éditeurs, Genève, 1–413.
- Stephani F (1901) Hepaticae novae Dussianae. Urban. Symbolae Antillanae 2: 469–472.
- Stephani F (1903) Species Hepaticarum. Bulletin de le Herbar Boissier (série 2) 3 (1): 140–177.
- Stephani F (1908) Species Hepaticarum. Bulletin de le Herbar Boissier (série 2) 8 (1): 49–64.
- Stephani F (1909) Species Hepaticarum. Bulletin de le Herbar Boissier (série 2) 9: 1–99.
- Stephani F (1910) Species Hepaticarum IV. George & Cia, Libraires-Éditeurs, Genève, 100–119.
- Stephani F (1911) Species Hepaticarum IV. George & Cia, Libraires-Éditeurs, Genève, 465–736.
- Stephani F (1913) Species Hepaticarum V. George & Cia, Libraires-Éditeurs, Genève, 178–480.
- Stephani F (1914) Species Hepaticarum V. George & Cia, Libraires-Éditeurs, Genève, 481–705.
- Stephani F (1915) Species Hepaticarum V. George & Cia, Libraires-Éditeurs, Genève, 706–848.
- Stephani F (1916) Species Hepaticarum V. George & Cia, Libraires-Éditeurs, Genève, 849–1006.
- Stephani F (1917) Species Hepaticarum V. George & Cia, Libraires-Éditeurs, Genève, 1009–1044.
- Struminski E (2001) Parque Estadual Pico do Marumbi. Série Pesquisa, n. 55. Editora UFPR, Curitiba, 179 pp.
- Sturm J (1819) Deutschlands Flora, Abtheilung II, Cryptogamie vol. 17. Kosten der Verfassers, Nürnberg, 16 pp.
- Sullivant WS (1846) Musci Alleghanensis: sive enumeratio muscorum atque hepaticarum quos in itinere a Marylandia usque ad Georgiam per tractus montium A.D. M DCCC XLIII. Typis Metcalf et Socorum, Columbus, 87 pp.
- Sullivant WS (1861) Musci Cubensis, or mosses collected by Charles Wright in the eastern part of the Island of Cuba during the years 1856, 1857, and 1858. Proceedings of the American Academy of Arts and Sciences 5: 273–290.
- Swartz O (1788) Nova Genera & Species Plantarum seu Prodrum descriptionum Vegetabilium, maximam partem incognitorum quae sub itinere in Indiam Occidentalem annis 1783–87. Uppsala & Aboae: In Bibliopoliis Acad. M. Swederi, Holmiae, 152 pp. <https://doi.org/10.5962/bhl.title.4400>
- Swartz O (1801) [1802] Observationes species nonnullas muscorum minus cognitae illustrantes. Journal of Botany (Schrader) 1800 (1–2): 171–185.
- Swartz O (1806) Flora Indiae occidentalis aucta atque illustrata, sive descriptiones plantarum in Prodrumo recensitarum. Vol. 3: 1231–2018. J. Palm, Erlingen.
- Tan BC, Pócs TC (2000) Bryogeography and conservation of bryophytes. In: Shaw AJ, Goffinet B (Eds) Bryophyte Biology. Cambridge University Press, Cambridge, 403–408.
- Taylor T (1843) Descriptions of two species of British Jungermanniae. Annals and Magazine of Natural History 12: 172–173.
- Taylor T (1846) The distinctive characters of some new species of Musci, collected by Professor William Jameson, in the vicinity of Quito, and by Mr. James Drummond at Swan River. London Journal of Botany 5: 41–67.
- Taylor T (1848) Some new musci, collected by Professor W. Jameson on Pichincha. London Journal of Botany 7: 187–199.
- The Plant list (2015). Briófitas. In: The Plant List. <http://www.theplantlist.org>. Accessed on: 2015-9-1.
- Thunberg CP (1800) Prodrum plantarum Capensium, quas in Promontorio Bonae Spei Africes, annis 1772–1775, pars Posterior. J. Edman, Uppsala, 191 pp. <https://doi.org/10.5962/bhl.title.84>
- Tixier P (1985) Contribution à la Connaissance des Cololejeunoideae. Bryophytorum Bibliotheca 27: 1–44.
- Trevisan de Saint-Léon VBATD (1877) Schema di una nuova classificazione delle Epatiche. Memorie del Reale Istituto Lombardo de Scienze e Lettere. Classe di Scienze Matematiche e Naturali 4 (3): 383–451.
- Valente EB, Pôrto KC (2006) Hepáticas (Marchantiophyta) de um fragmento de Mata Atlântica na Serra da Jibóia, Município de Santa Teresinha, BA, Brasil. Acta Botânica Brasileira 20: 433–439.
- Vaz-Imbassahy TF, Costa DP (2009) New combinations and new synonyms in Pilotrichaceae (Bryophyta) II. Nova Hedwigia 88 (3–4): 465–474. <https://doi.org/10.1127/0029-5035/2009/0088-0465>
- Villarreal JC, Renner SS (2014) A review of molecular-clock calibrations and substitution rates in liverworts, mosses, and hornworts, and a timeframe for a taxonomically cleaned-up genus *Nothoceros*. Molecular Phylogenetics and Evolution 78: 25–35. <https://doi.org/10.1016/j.ympev.2014.04.014>
- Vitt D (1973) A revisionary study of the genus *Macrococma*. Revue bryologique et lichénologique 39 (2): 205–220.
- Vitt D (1979) New taxa and new combinations in the Orthotrichaceae of Mexico. The Bryologist 82 (1): 1–19.
- Warnstorff C (1891) Beiträge zur Kenntniss Exotischer Sphagna. II. Hedwigia 30 (1): 12–60.
- Warnstorff C (1911) Sphagnales–Sphagnaceae (Sphagnologia Universalis). Engelmann, Leipzig, 550 pp.
- Weber F (1815) Historiae Muscorum hepaticorum Prodrum. Kiliae: Academiae Bibliopolae. 160 pp.
- Wijk RVD, Margadant WD (1959) New combinations in mosses II. Taxon 8: 70–75.
- Wijk RVD, Margadant WD (1960) New combinations in mosses IV. Taxon 9: 50–52.
- Williams RS (1909) Bolivian mosses. Part II. Bulletin of the New York Botanical Garden 6 (21): 227–261.
- Williams RS (1928) Some apparently undescribed mosses from Peru, also new combinations. The Bryologist 31: 109–122.
- Williams RS, Britton EG (1914) Central American mosses. Torreya 14: 24–31.
- Wilson W (1847) Description of two mosses from Jamaica. Transactions and Proceedings of the Botanical Society of Edinburgh 20: 21–22.
- Wilson W (1854) Flora of the Isthmus of Panama, Musci. In: Seeman B (Ed.) The Botany of the Voyage of H.M.S. Herald under the command of Captain Henry Kellett, R.N., C.B., during the years 1845–51. Lovell Reeve, London, 244–245. <https://doi.org/10.5962/bhl.title.425>
- Withering W (1776) A Botanical Arrangement of All the Vegetables Naturally Growing in Great Britain. T. Cadel, P. Elmsley, G. Robinson, London, 838 pp.
- Withering W (1801) A Systematic Arrangement of British Plants; with an Easy Introduction to the Study of Botany, 3rd edition. Musci. Vol. 3. Baldwin and Son, London, 824 pp.
- Yamada K (1990) Two new species of *Radula* (Hepaticae) from Australia and Brazil. Journal of Japanese Botany 65 (1): 1–6.
- Yamada K (1993) Four new species of *Radula* from Neotropics. The

- Journal of the Hattori Botanical Laboratory 74: 35–44.
- Yano O (1995) A new additional annotated checklist of Brazilian bryophytes. The Journal of the Hattori Botanical Laboratory 78: 137–182.
- Yano O (1981) A checklist of Brazilian mosses. The Journal of the Hattori Botanical Laboratory 50: 279–456.
- Yano O (1984) Checklist of Brazilian liverworts and hornworts. The Journal of the Hattori Botanical Laboratory 56: 481–548.
- Yano O (1989) An additional checklist of Brazilian bryophytes. The Journal of the Hattori Botanical Laboratory 66: 371–434.
- Yano O (1992) Novas localidades de musgos nos estados do Brasil. Acta Amazonica 22: 197–218. <http://doi.org/cg3w>
- Yano O (2006a) Novas adições ao catálogo de briófitas brasileiras. Boletim do Instituto de Botânica 17: 1–142.
- Yano O (2006b) Novas adições ao catálogo de briófitas brasileiras. Boletim do Instituto de Botânica 18: 229–233.
- Yano O (2010) Levantamento de novas ocorrências de briófitas brasileiras. Publicação on line do Instituto de Botânica. <http://www.santoandre.sp.gov.br/pesquisa/ebooks/342354.pdf>. Accessed on: 2017-11-16.
- Yano O (2013) Catálogo das Briófitas (Antóceros, Hepáticas e Musgos) do Estado do Paraná, Brasil. Pesquisas, Série Botânica 64: 347–420.
- Yano O (2014) Ocorrências novas de Briófitas para o Estado do Paraná, Brasil. Pesquisas, Série Botânica 65: 67–122.
- Yano O, Carvalho AB (1995) Briófitas da Serra da Piedade, Minas Gerais, Brasil. In: Noronha MRP (Ed.) Anais 9º Congresso da Sociedade Botânica de São Paulo. Sociedade Botânica de São Paulo, São Paulo, 15–25.
- Yano O, Colletes AG (2000) Briófitas do Parque Nacional de Sete Quedas, Guaira, PR, Brasil. Acta Botanica Brasílica 14: 215–242.
- Yano O, Peralta DF (2009) Flora de Grão-Mogol, Minas Gerais. Briófitas (Bryophyta e Marchantiophyta). Boletim de Botânica da Universidade de São Paulo 27: 1–26. <https://doi.org/10.11606/issn.2316-9052.v27i1p1-26>
- Yano O, Peralta DF (2011) Flora da Serra do Cipó, Minas Gerais: Briófitas (Anthocerotophyta, Bryophyta e Marchantiophyta). Boletim de Botânica da Universidade de São Paulo 29: 135–211. <https://doi.org/10.11606/issn.2316-9052.v29i2p135-299>
- Ye W, Zhu R-L (2010) *Leucolejeunea*, a new synonym of *Cheilolejeunea* (Lejeuneaceae), with special reference to new combinations and nomenclature. Journal of Bryology 32 (4): 279–282. <https://doi.org/10.1179/037366810x12814321877507>
- Yu Y, Pócs TC, Zhu R-L (2014) Notes on early land plants today. 62. A synopsis of *Myriocoleopsis* (Lejeuneaceae, Marchantiophyta) with special reference to transfer of *Cololejeunea minutissima* to *Myriocoleopsis*. Phytotaxa 183 (4): 293–297. <https://doi.org/10.11646/phytotaxa.183.4.11>