



Diversity and distribution of microlichens in the state of Arunachal Pradesh, Eastern Himalaya, India

Pushpi Singh^{1*}, Krishna Pal Singh¹ and Ajay Ballabh Bhatt²

¹ Botanical Survey of India, Central Regional Centre, Allahabad – 211 002, India

² Hemwati Nandan Bahuguna Garhwal University, Srinagar, Uttarakhand

* Corresponding author. E-mail: pushpisingh08@gmail.com

Abstract: The paper reports the occurrence of 404 species of microlichens belonging to 105 genera and 39 families known so far, from the state of Arunachal Pradesh, a part of the Himalaya biodiversity hotspot. Twelve species, namely *Arthopyrenia saxicola*, *Arthothelium subbessale*, *Diorygma macgregorii*, *D. pachygraphum*, *Graphis nuda*, *G. oligospora*, *G. paraserpens*, *G. renschiana*, *Herpothallon japonicum*, *Megalospora atrorubicans*, *Porina tijucana* and *Rhabdodiscus crassus*, are new distributional records for India. *Astrothelium meghalayense* (Makhija & Patw.) Pushpi Singh & Kr. P. Singh and *Astrothelium subnitidiusculum* (Makhija & Patw.) Pushpi Singh & Kr. P. Singh are proposed as new combinations and 66 species marked by an asterisk (*) are new distributional records for the state.

Key words: microlichens, diversity, Eastern Himalaya, new records

INTRODUCTION

The Indian state of Arunachal Pradesh, one of the most important parts of the Himalaya biodiversity hotspot (Mittermeier et al. 2005) covers an area of 83,743 km² (2.54 % total area of India) and is at altitudes ranging between 200 and 7,000 m above the mean sea level. Arunachal Pradesh lies between 26°28' and 29°30' N latitude and 091°30' and 097°30' E longitude. It harbours rich and unique diversity of lichen flora in northeastern India due to varied climate and topography. Various types of substrata, such as bark, twigs, leaves, soil, and rocks, provide suitable conditions for the rich growth of lichens from tropical to alpine regions. Lichenological exploration in the state was first made by Rolla Seshagiri Rao and Gopinath Panigrahi of Botanical Survey of India during 1956–1958. These collections were studied by Awasthi (1961) who reported 42 species of macrolichens. Subsequently, based on new collections in the state, some additional publications on foliicolous lichens, new species, and new records for India have

been made (e.g., Pinokiyo et al. 2004; Dubey et al. 2007, 2010; Pinokiyo et al. 2008; Singh and Pinokiyo 2008; Singh and Swarnlatha 2011a, 2011b; Jagadeesh Ram and Sinha 2011; Upreti et al. 2011; Singh and Singh 2012a, 2012b, 2012c, 2014; Singh et al. 2013; Joshi et al. 2014). Recently, a publication on foliicolous lichens of India (Singh and Pinokiyo 2014) recorded 98 species from Arunachal Pradesh. However, the microlichens of upper northern regions of the state could not be fully explored because of rugged and inaccessible hilly terrain. In the present study, the microlichens collected so far, have been investigated, identified and enumerated together with published reports in the tabular form (Appendix, Table A1) for future users.

MATERIALS AND METHODS

Arunachal Pradesh borders the states of Assam and Nagaland to the south, and shares international borders with the countries of Myanmar in the east, Bhutan in the west, and Tibet (People's Republic of China) in the north. Several extensive collection expeditions were undertaken in different localities in the state (Figure 1). Collected specimens were deposited in the herbaria of Botanical Survey of India, Central Regional Centre, Allahabad (BSA), and Eastern Regional Centre, Shillong (ASSAM). Also studied were specimens loaned from the National Botanical Research Institute, Lucknow (LWG). Morphological observations were made using a stereomicroscope (Nikon SMZ 1500). Thin, hand-cut sections, of thalli and ascocarps were mounted in water, lactophenol cotton blue (LPCB), 10% KOH and Lugol's iodine solution. All anatomical measurements were taken in water mounts and examined under a compound microscope (Nikon Eclipse 50i). Secondary metabolites were identified by thin layer chromatography (TLC) following Orange et al. (2001) and White and James (1985). All the specimens were identified with the help of authenticated specimens available in the various herbaria and published literature. Current names of the




Figure 1. Map of Arunachal Pradesh showing areas of exploration (stippled) of microlichens in various districts.

species, habit and distribution in districts along with selected voucher specimen numbers are also provided.

RESULTS

This study of microlichens of Arunachal Pradesh revealed the occurrence of 404 species (Appendix, Table A1), belonging to 105 genera and 39 families (Figure 2), of which 12 species are newly recorded for the Indian lichen flora and 66 species (marked with an asterisk) are new distributional records for Arunachal Pradesh.

As far as the microlichen diversity is concerned, the family Graphidaceae comprises 123 species and shows maximum diversity, followed by Porinaceae (35 species), Pyrenulaceae (34 species), Pilocarpaceae (31 species), Arthoniaceae (30 species), Lecanoraceae (22 species), Physciaceae (16 species), Pertusariaceae (12 species), etc. At the generic level, *Graphis* comprises 46 species and shows maximum species diversity, followed by *Porina* 34 species and *Lecanora* 21 species.

New distributional records

Arthopyrenia saxicola A. Massal. *Symmict. Lich.*: 107. 1855. (Figure 3A–C)

Notes: This species is characterized by its saxicolous habit (on cemented rock), lichenized, superficial to endolithic, whitish or pale brownish or greyish thallus; black solitary ascomata, branched and anastomosing pseudoparaphyses with simple periphysoids; 8-spored ascii; colorless, transversely 1-septate, 18–19 (–21) × 7–9 µm ascospores with upper cell slightly broader than the lower cell and lacking lichen substances. Sometimes this species is confused with *Porina linearis* (Leight.) Zahlbr., which has 3-septate ascospores and ± simple paraphyses (*fide* Orange 2013). Earlier, this species was erroneously




Figure 2. Total number of genera and species of per microlichen family in Arunachal Pradesh.




Figure 3. A–C: *Arthopyrenia saxicola*, (A) habit, (B) transverse section of ascoma, (C) ascus with ascospores. D–E: *Arthothelium subbessale*, (D) habit, (E) ascus and ascospores. F–G: *Diorygma macgregorii*, (F) habit, (G) ascospore. H–I: *D. pachygraphum*, (H) habit, (I) ascospore. Scale bars: A, D, F, H = 1 mm; B=100 µm; C, E=20 µm; G=25 µm; I=50 µm.

identified as *Anisomeridium calcicolum* Upreti & Nayaka (Pinokiyo et al. 2008). It grows between 300 and 400 m altitude in shady exposed places in tropical forests. The species is distributed in Britain, China, Ireland and Hong Kong.

Specimen examined: Arunachal Pradesh, Lower Dibang Valley district, Mehao Wildlife Sanctuary, Sally Lake, on cemented stones, alt. 390–400 m, K.P. Singh & P.K. Dixit 476 (BSA).

Arthothelium subbessale (Nyl.) Makhija & Patw., *Tropical Bryology* 10: 210. 1995. *Arthonia subbessalis* Nyl., *Sert. Lich. Trop. Labuan Singapore*: 23. 1891
(Figure 3D–E)

Notes: *Arthothelium subbessale* is characterized by its endophloeodal to slightly epiphloeodal, irregular, brownish grey thallus; blackish-brown semi-immersed substellate to irregular, blackish-brown ascocarps; branched and anastomosing paraphysoids; 8-spored ascii; muriform, transversely 6–7-septate and longitudinally 1–3-septate, 30–35 (–36) × 14–16 µm ascospores with a large undivided apical cell and absence of lichen substances. In ascospores character, it closely resembles *Arthothelium bessale* (Nyl.) Zahlbr., which has rounded, large ascocarps and large ascospores (36–44 µm long, *fide* Makhija and Patwardhan 1995). The species is distributed in Singapore.

Specimen examined: Arunachal Pradesh, Upper Subansiri district, Taliha, Subansiri River bed, on bark, K.P. Singh 10502 (ASSAM).

Diorygma macgregorii (Vain.) Kalb, Staiger & Elix, *Symb. Bot. Ups.* 34(1): 159–160. 2004. *Cyclographina macgregorii* (Vain.) D.D. Awasthi & M. Joshi, *Norw. J. Bot.* 26(3): 172. 1979.
(Figure 3F–G)

Notes: *Diorygma macgregorii* is characterized by its pale brown or grayish brown, rugose ecorcinate thallus; straight to incurved, flexuous and branched, prominently raised lirellae; wide open whitish pruinose disc; divergent, non-carbonized exciple; clear, completely I+ blue hymenium; colorless, oblong, densely muriform, 110–180 × 40–50 µm ascospores and presence of norstictic and connorstictic acids. This species resembles *Diorygma pachygraphum* (Nyl.) Kalb, Staiger & Elix, but later species differs in having closed to slightly open disc and strongly raised apothecia. This species is widely distributed in China, Philippines and Papua New Guinea.

Specimens examined: Arunachal Pradesh, West Kameng district, near Dedza bridge, Bhalukpong-Tenga road, on bark, alt. ca. 1216 m, K.P. Singh & G. Swarnlatha 4833, 4826 (BSA).

Diorygma pachygraphum (Nyl.) Kalb, Staiger & Elix,

Symb. Bot. Upsal. 34 (1): 163. 2004. *Graphis pachygrapha* Nyl., *Acta Soc. Sci. fenn.* 7(2): 472. 1863.
(Figure 3H–I)

Notes: This species is characterized by its greenish or whitish grey slightly fissured to rimose ecorcinate thallus; flexuous, elongated, prominently raised constricted lirellae; closed or slightly opened whitish pruinose disc; convergent to divergent, uncarbonized exciple; hyaline, weakly I+ blue-violet hymenium mostly in lateral part; 1-spored ascii with large densely muriform, 164–195 × 34–50 µm ascospores and presence of norstictic and connorstictic acids. In chemistry and ascospores characters, it closely resembles *Diorygma macgregorii* (Vain.) Kalb, Staiger & Elix, which has whitish pruinose exposed disc. The species is distributed in China, Colombia, Philippines and Tanzania.

Specimens examined: Arunachal Pradesh, West Siang district, Basar to Bame, alt. ca. 850 m, K.P. Singh 2445 (ASSAM); West Kameng district, Bhalukpong-Tenga road, near Dedza Bridge, on bark, alt. ca. 1216 m, K.P. Singh & G. Swarnlatha 4931, 4835, 4845, 4832 (BSA).

Graphis nuda (Magn.) Staiger & Lücking, in Lücking, Chaves, Sipman, Umaña & Aptroot, *Fieldiana Bot.* 38 (no. 1549): 93. 2008. *Graphina nuda* H. Magn., *Ark. Bot.* 3 (no. 10): 266. 1955.
(Figure 4A–B)

Notes: This species is characterized by the greyish or whitish grey thallus; very short, sessile lirellae lacking thalline margin; concealed disc; entire labia; completely carbonized exciple; clear hymenium; 8-spored ascii with colorless, muriform, 28–44 (–47) × 14–19 µm ascospores and absence of lichen substances. Morphologically, it closely resembles *Graphis ruiziana* (Fée) A. Massal., and *G. subruiziana* Sipman, Chaves & Lücking, but both later species have larger 35–65 µm and 55–80 µm long ascospores respectively (*fide* Lücking et al. 2009) ascospores. The species is widely distributed in Neotropics and Eastern Palaeotropics.

Specimen examined: Arunachal Pradesh, Lower Dibang Valley district, Mehao Wildlife Sanctuary, Sally lake, on bark, alt. 390–400 m, K.P. Singh & P.K. Dixit 448 (BSA).

Graphis oligospora Zahlbr. apud Handel-Mazzetti, *Symbol. Sinic. Pars 3*: 45. 1930.
(Figure 4C–D)

Notes: This species is characterized by its pale brown, rugose thallus; short and sparsely branched, emergent lirellae covered with lateral thalline margin; concealed disc; entire labia; apically to peripherally carbonized exciple; clear hymenium; 2–4 (–6)-spored ascii with colorless, fusiform, transversely 8–10-septate, (26–) 36–42 × (7–) 8–10 µm ascospores and absence of lichen substances. Anatomically, it closely resembles *Graphis*




Figure 4. A–B. *Graphis nuda*, (A) habit, (B) ascospore. C–D: *G. oligospora*, (C) habit, (D) ascospores. E–F: *G. paraserpens*, (E) habit, (F) ascospores. G–H: *G. renschiana*, (G) habit, (H) ascospores. Scale bars: A, C, E, G = 1 mm; B, D, F, H = 20 µm.

intermediella Stirt., which has very long and radiately branched lirellae. The species is widely distributed in eastern palaeotropics.

Specimen examined: Arunachal Pradesh, Lower Dibang Valley district, Mehao Wildlife Sanctuary, Mehao lake, on bark, alt. 1550–1600 m, K.P. Singh & P. Barua 10871 (ASSAM).

Graphis paraserpens Lizano & Lücking, in Lücking, Chaves, Sipman, Umaña & Aptroot, *Fieldiana Bot.* 38 (1549): 96. 2008.

(Figure 4E–F)

Notes: *Graphis paraserpens* is characterized by its greyish brown to pale brown thallus; elongate, straight to flexuose, partly branched, immersed-erumpent lirellae with concealed disc and striate labia; apically to peripherally carbonized exciple with apically thin complete thalline margin; 2–6-spored ascospores with colorless, muriform, 22–35 × 8.5–10 µm ascospores and absence of lichen substances. Morphologically, it resembles *Graphis symplecta* Nyl., which has broader ascospores (15–20 µm, *fide* Lücking et al. 2009). It also closely resembles *Graphis puiggarii* (Müll. Arg.) Lücking in morphological characters and ascospores size but the latter species differs in having lateral carbonized exciple. The species is distributed in Costa Rica.

Specimen examined: Arunachal Pradesh, West Kameng district, Sessa, on bark, alt. 1110–1500 m, Pinokiyo 64 (BSA).

Graphis renschiana (Müll. Arg.) Stizenb., *Bericht. Über die Thatigk. St. Gallisch. Naturw. Gesellsch.* 184. 1891.

Graphina renschiana Müll. Arg., *Flora Regensburg* 68(28):

512. 1885.

(Figure 4G–H)

Notes: This species is characterized by its greyish brown to reddish brown thallus; elongate and irregularly branched, erumpent lirellae with concealed disc and entire labia; laterally carbonized exciple; clear hymenium; 4–6-spored ascospores with colorless, muriform, 22–35 × 10–15 µm ascospores and presence of norstictic acid. Morphologically and in chemistry, it resembles *Graphis norstictica* A.W. Archer & Lücking and *G. borealis* (A.W. Archer) A.W. Archer, but both latter species have larger ascospores. Morphologically, it also resembles *G. deserrens* Vain, which has stictic acid. This species is widely distributed in China, Florida, Madagascar and the Philippines (Pantropical).

Specimen examined: Arunachal Pradesh, West Kameng district, Tipi, Festival ground, on bark, alt. ca. 160 m, K.P. Singh & G. Swarnlatha 4741 (BSA).

Herpothallon japonicum (Zahlbr.) G. Thor in Aptroot et al. *Biblioth. Lichnol.* 99: 44. 2009. *Chiodection japonicum* Zahlbr., *Annals mycol.* 29(1/2): 77. 1931.

(Figure 5A–B)

Notes: This species is characterized by the whitish to greenish-grey, minutely felty thallus lacking calcium oxalate crystals, I-/KI-; numerous, dense, cylindrical, unbranched, byssoid-felty, 0.8–1 × 0.2 mm pseudoisidia; whitish byssoid hypothallus; pale brown to whitish byssoid prothallus and presence of gyrophoric acid as major substance. It closely resembles *H. philippinum* (Vain.) Aptroot & Lücking in morphology but later species differs in I+/KI+ blue reaction and presence of calcium oxalate crystals in the thallus. Earlier this species was reported from Japan.

Specimen examined: Arunachal Pradesh, East Kameng district, Phakui Wildlife Sanctuary, Ditchu-Julley Nala, on bark, alt. 300–400 m, K. P. Singh 10091 A, B (ASSAM).

Megalospora atrorubricans (Nyl.) Zahlbr. *Cat. Lich. Univ.* 4: 86. 1927. *Lecidea marginiflexa* var. *atrorubricans* Nyl., *Flora Regensburg* 49: 132. 1866.

(Figure 5C–D)

Notes: This species is characterized by its greyish green, rugulose thallus; sessile, concave to strongly convex, dark brown to black, epruinose apothecia; inspersed hymenium; brown epiphymenium; 1-spored ascospores with colorless, ellipsoid to oblong, straight, transversely 1-septate, 88–102 × 30–35 µm ascospores and presence of usnic acid and zeorin. In ascospores character and chemistry, it closely resembles *Megalospora sulphurata* Meyen, which has 2–8-spored ascospores and ±curved (*sulphurata*-type) ascospores. This species is widely distributed in the Hawaiian Islands, Indonesia, the Mascarene Islands, Papua New Guinea, the Philippines and New Caledonia.

Specimen examined: Arunachal Pradesh, West Kameng district, Shergoan river side, on bark, alt. ca. 1,600 m, G.P. Sinha & T.A.M. Jagadeesh Ram 11416 (ASSAM).

Porina tijucana Vain., *Étude Lich. Brésil* 2: 220. 1890.

(Figure 5E–F)

Notes: *Porina tijucana* is characterized by its siccicolous habit, grey brown to greenish brown thallus; emergent 0.3–0.65 µm diam. perithecia with broad black perithecial cap; well-developed involucellum and fusiform, transversely 5–9 (-11)-septate, 69–80 × 8.5–10 (-13) µm ascospores with gelatinous sheath. It resembles *P. guaranitica* Malme, which has a highly reduced involucellum. It is also close to *P. mastoidea* (Ach.) Müll. Arg., but latter species differs in having smaller ascospores (32–65 × 6–12 µm, *fide* McCarthy 2001). The species is widely distributed in Brazil, Costa Rica, Colombia, Panama and Thailand.

Specimens examined: Arunachal Pradesh, Dibang Valley district, Ryali-Anini forest, on rock, alt. ca. 900 m, K. P. Singh 4783 (ASSAM); Lower Subansiri district, Yazali river side forest, on rock, alt. 480–494 m, K. P.




Figure 5. A–B: *Herpothallon japonicum*, (A) habit, (B) pseudoisidia (close-up). C–D: *Megalospora atrorubicans*, (C) habit, (D) ascus with ascospore. E–F: *Porina tijucana*, (E) habit, (F) ascospore. G–H: *Rhabdodiscus crassus* (G) habit, (H) ascospores. Scale bars: A, B, C, E, G = 1 mm; D, H= 20 μ m; F= 25 μ m.

Singh 5547 (ASSAM).

Rhabdodiscus crassus (Müll. Arg.) Rivas Plata, Lücking & Lumbsch, *Taxon* 61: 1175. 2012. *Leptotrema crassum* Müll. Arg., *Flora, Regensburg* 65(21): 332. 1882. (Figure 5G–H)

Notes: This species is characterized by its greenish grey to brownish thallus; emergent to prominent apothecia with broad-stump shaped carbonized columella with whitish pruinosity at the tip and open disc; lateral carbonized exciple with clear hymenium; 8-spored ascospores with fusiform, 5–7 × 0–1-septate, 15–20 × 7–9 µm ascospores and presence of psoromic acid (major). Morphologically, it closely resembles *Ocellularia papillata* (Leighton) Zahlbr., which has a simple columella, 3–5-septate ascospores and lacks psoromic acid. The species is widely distributed in Australia, Costa Rica, Indonesia (Java), Japan and USA.

Specimen examined: Arunachal Pradesh, Lower Dibang Valley district, Mehao Wildlife Sanctuary, Mehao lake, alt. ca. 1,550 m, K.P. Singh & P.K. Dixit 547 C, F (BSA).

New combinations

Recently, phylogenetic studies on Trypetheliaceae (Nelsen et al. 2014) have shown that the species with

rounded (not diamond-shaped) lumina in ascospores cluster around the *Trypethelium* group, while species with diamond-shaped lumina form another cluster around the *Astrothelium conicum*-group, regardless of their ascoma organization. Accordingly following two new combinations are proposed.


Astrothelium meghalayense (Makhija & Patw.) Pushpi Singh & Kr. P. Singh comb. nov.; *Trypethelium meghalayense* Makhija & Patw., *J. Hattori Bot. Lab.* 73: 201. 1993. (Figure 6A–B)

Mycobank: MB 814410

Notes: *Astrothelium meghalayense* is characterized by its greenish grey thallus; multicarpic, thallus colored, raised pseudostromata with creamish colored top; inspersed hamathecium; 8-spored ascospores with diamond-shaped lumina.

Specimen examined: Arunachal Pradesh, Lower Dibang Valley district, Mehao Wildlife Sanctuary, Mehao Lake, on bark, alt. 1,550–1,700 m, K. P. Singh & P. K. Dixit 589 E (ASSAM).

Astrothelium subnitidiusculum (Makhija & Patw.) Pushpi Singh & Kr. P. Singh comb. nov.; *Trypethelium*



Figures 6. A–B: *Astrothelium meghalayense*, (A) habit, (B) ascus with ascospores. **C–D:** *Astrothelium subnitidiusculum*, (C) habit, (D) ascospore. Scale bars: A, C = 1 mm; B, D = 20 µm.

subnitidiusculum Makhija & Patw., *J. Hattori Bot. Lab.* 73:

207. 1993.

(Figure 6C–D)

Mycobank: MB 814411

Notes: *A. subnitidiusculum* is characterized by its brownish-yellow thallus; monocarpic, thallus colored, raised pseudostromata with pale top; inspersed hamathecium; 8-spored asci and colorless, transversely 7-septate, 30–58 × 10–13 µm ascospores with diamond-shaped lumina.

Specimen examined: Arunachal Pradesh, Namsai district, Madhuban Reserve Forest, alt. ca. 250 m, K.P. Singh 4196 (ASSAM).

DISCUSSION

Geographically, Arunachal Pradesh is the largest state among the eight northeastern states (Arunachal Pradesh, Assam, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim and Tripura) of India and harbours a wide spectrum of lichen diversity. The present study provides an idea of the diversity and distribution of microlichens in the state and provides baseline data for future research in this region. Arunachal Pradesh has characteristic lichen genera such as *Coccotrema* (Singh and Singh 2012c), *Erythrodection* (Singh et al. 2009) and *Mycoblastus* (Singh and Singh 2015), which are hitherto unknown from the other parts of India. Forty-eight species (marked with an “E” in Appendix, Table A1) are endemic to India. Thus, Arunachal Pradesh has a distinct and unique lichen diversity and justifies its inclusion as a hotspot of biodiversity in the Indian Himalaya. Habitat-wise, 265 species are corticolous (67%), 98 species are foliicolous (25%), 27 species are saxicolous (7%), and four species are terricolous (1%). It is observed that the corticolous and foliicolous species are dominant and commonly distributed in almost all localities where as saxicolous and terricolous ones are scarce.

A comparison of microlichen diversity with other Indian Himalayan states indicates that the Arunachal Pradesh has the highest number of microlichen species known so far (Figure 7). The state has 404 species, but Uttarakhand has 282 species, Sikkim has 265 species, Himachal Pradesh has 155 species, Jammu and Kashmir has 149 species, and West Bengal-Darjeeling has 114 species (Singh and Sinha 2010; Upreti et al. 2010; Sinha and Jagadeesh Ram 2011; Jagadeesh Ram and Sinha 2011a, 2011b; Joseph and Sinha 2012, 2015; Goni et al. 2015). Similarly, a comparison among the north eastern states the Arunachal Pradesh also shows high percentage of microlichen diversity (Figure 8).

The rugged, hilly, and largely inaccessible terrain, which is cut by many rivers and streams originating from higher Himalayas, has made lichen surveys in the region extremely difficult. As a result, many areas in the north at higher elevations are still lichenologically




Figure 7. Microlichen diversity in Indian Himalayan states.




Figure 8. Microlichen diversity in northeastern states of India.

unexplored; future exploration of these areas may provide many interesting finds. The present assessment of microlichens will be helpful in biomonitoring and climate change studies, conservation research, and sustainable utilization of lichen resources.

ACKNOWLEDGEMENTS

The authors thank Dr. Paramjit Singh, Director, Botanical Survey of India, Kolkata, for encouragement and Dr. G.P. Sinha, Head of Office, Botanical Survey of India, Central Regional Centre, Allahabad for facilities. One of the authors (PS) also thanks the authorities of Botanical Survey of India for financial assistance under the “Flora of India Project”.

LITERATURE CITED

- Awasthi, D.D. 1961. Some foliose and fruticose lichens from Assam and North-East Frontier Agency of India. *Proceedings of the Indian Academy of Sciences, Section B*, 54: 24–44.
- Dubey, U., D.K. Upreti and J. Rout. 2007. Lichen flora of Along, West Siang district, Arunachal Pradesh. *Phytotaxonomy* 7: 21–26.
- Dubey, U., D.K. Upreti and N. Sanjeeva. 2010. A new species of *Leiorreuma* Eschw. from India. *The Lichenologist* 42: 711–713. doi: [10.1017/S002428291000040X](https://doi.org/10.1017/S002428291000040X)
- Goni, R., A.K.P. Raina, R. Magotra and N. Sharma. 2014. Lichen

- flora of Jammu and Kashmir State, India: An updated checklist. *Tropical Plant Research* 2(1): 64–71.
- Jagadeesh Ram, T.A.M. and G.P. Sinha. 2011. A new species and a new record of *Herpothallon* (lichenized Ascomycota) from India. *Mycotaxon* 116: 313–316. doi: [10.5248/116.313](https://doi.org/10.5248/116.313)
- Jagadeesh Ram, T.A.M. and G.P. Sinha. 2011a. Additional records of lichens for Darjeeling hills. *Annals of Forestry* 19(2): 264–265.
- Jagadeesh Ram, T.A.M. and G.P. Sinha. 2011b. Additions to lichen flora of Himalaya from Darjeeling and Sikkim. *Geophytology* 40(1–2): 21–28.
- Joseph, S. and G.P. Sinha. 2012. *Lecanora bicincta* Ramond, a new record to Indian lichen flora and some additions to the lichen flora of Sikkim, India. *Geophytology* 42(1): 71–75.
- Joseph, S. and G.P. Sinha. 2015. Contributions to the genus *Synarthonia* (lichenized Ascomycota, Arthoniaceae). *The Lichenologist* 47(2): 123–130. doi: [10.1017/s002428291500002x](https://doi.org/10.1017/s002428291500002x)
- Joshi, Y., T.A.M. Jagadeesh Ram, P. Singh and G.P. Sinha. 2014. *Caloplaca indica*, a new lichenized Ascomycetes (Teloschistaceae) from Eastern Himalaya, India. *National Academy Science Letter* 37(6) 517–519. doi: [10.1007/s40009-014-0270-3](https://doi.org/10.1007/s40009-014-0270-3)
- Lücking, R., A.W. Archer and A. Aptroot. 2009. A world wide key to the genus *Graphis* (Ostropales: Graphidaceae). *The Lichenologist* 41: 363–452. doi: [10.1017/S0024282909008305](https://doi.org/10.1017/S0024282909008305)
- Makhija, U. and P.G. Patwardhan. 1995. The Lichen genus *Arthothelium* (Family Arthoniaceae) in India. *Journal, Hattori Botanical Laboratory* 78: 189–235.
- McCarthy, P.M. 2001. *Porina*; pp. 106–153, in: P.M. McCarthy (ed.). *Flora of Australia*. 58A, Lichens 3. Canberra: Australia Government Publishing Service.
- Mittermeier, R.A., P.R. Gil, M. Hoffman, J. Pilgrim, T. Brooks, C.G. Mittermeier, J. Lamoreux and G.A.B. de Fonseca (eds.). 2005. Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions. Mexico: CEMEX. 392 pp.
- Nelsen, M.P., R. Lücking, A. Aptroot, C.J. Andrew, M.E.S. Cáceres, E. Rivas Plata, C. Gueidan, L. DA Silva Canez, A. Knight, L.R. Ludwig, J. Mercadi-Diaz., S. Parnmens and H.T. Lumbsch. 2014. Elucidating phylogenetic relationships and genus-level classification within the fungal family Trypetheliaceae (Dothideomycets: Ascomycota). *Taxon* 65: 974–992.
- Orange, A. 2013. British and other pyrenocarpous lichens. Cardiff, Wales: Department of Biodiversity and Systematic Biology, National Museum of Wales. 250 pp.
- Orange, A., P.W. James and F.J. White. 2001. Microchemical Methods for the Identification of Lichens. London: British Lichen Society. 101 pp.
- Pinokiyo, A., K.P. Singh and S.K. Borthakur. 2004. Folioicolous species of *Porina* (Lichens) from Arunachal Pradesh, India. *Indian Journal of Forestry* 27(4): 407–416.
- Pinokiyo, A., K.P. Singh and J.S. Singh. 2008. Diversity and distribution of lichens in relation to altitude in a protected area within a biodiversity hot spot of North-east India. *The Lichenologist* 40(1): 47–62. doi: [10.1017/S0024282911000721](https://doi.org/10.1017/S0024282911000721)
- Singh, K.P. and A. Pinokiyo. 2008. New taxa of lichens from eastern India. *The Lichenologist* 40(1): 23–29. doi: [10.1017/S0024282908006889](https://doi.org/10.1017/S0024282908006889)
- Singh, K.P. and A. Pinokiyo. 2014. Folioicolous lichens of India. Dehradun: Bishen Singh Mehendra Pal Singh. 335 pp.
- Singh, K.P. and G. Swarnlatha. 2011a. A new species of *Schistophoron* from India. *The Lichenologist* 43(3): 209–212. doi: [10.1017/S0024282910000800](https://doi.org/10.1017/S0024282910000800)
- Singh, K.P. and G. Swarnlatha. 2011b. A note on Graphidaceous lichens from Arunachal Pradesh. *Indian Journal of Forestry* 34: 353–360.
- Singh, K.P. and G.P. Sinha. 2010. Indian lichens: Annotated Checklist. Kolkata: Botanical Survey of India. 571 pp.
- Singh, K.P. and P. Singh. 2012a. A new species of *Pyrgillus* from India. *The Lichenologist* 44(6): 773–776. doi: [10.1017/S002428291200002X](https://doi.org/10.1017/S002428291200002X)
- Singh, K.P. and P. Singh. 2012b. Genus *Pyrgillus* Nyl. (Lichenized: Ascomycota: Pyrenulaceae) in India. *Taiwania* 57(4): 391–395.
- Singh, K.P., T.A.M. Jagadeesh Ram and G.P. Sinha. 2009. *Erythrodection* G. Thor (Lichenized Ascomycetes), a new generic record for India. *Indian Journal of Forestry* 32(2): 335–336.
- Singh, P. and K.P. Singh. 2012c. *Coccotrema* (Lichenized: Ascomycota) a new generic record for India. *Indian Journal of Forestry* 35(1): 131–132.
- Singh, P. and K.P. Singh. 2014. *Cyphelium inquinans* (Sm.) Trev. (lichenized: Ascomycota): a new record for India. *Check List* 10(1): 178–179. doi: [10.15560/10.1.178](https://doi.org/10.15560/10.1.178)
- Singh, P. and K.P. Singh. 2015. *Mycoblastus* Norman (Mycoblastaceae): A new generic record for India from Arunachal Pradesh. *Indian Journal of Forestry* 38(3): 249–251.
- Singh, P., K.P. Singh and A.B. Bhatt. 2013. New species and new records of theloremoid Graphidaceae (lichenized Ascomycota) from Arunachal Pradesh (India). *Taiwania* 58(4): 246–250. doi: [10.6165/tai.2013.58.246](https://doi.org/10.6165/tai.2013.58.246)
- Sinha, G.P. and T.A.M. Jagadeesh Ram. 2011. Lichen diversity in Sikkim; pp. 13–29, in: M.L. Arrawatia and S. Tambe (eds.). *Biodiversity of Sikkim exploring and conserving a global hotspot*. Gantok: Department of Information and Public Relations, Government of Sikkim.
- Upreti, D.K., S. Nayaka and S. Chatterjee. 2010. Lichen diversity of Uttarakhand Himalaya; pp. 79–196, in: P.L. Uniyal, B.P. Chamola and D.P. Semwal (eds.). *The plant wealth of Uttarakhand*. New Delhi: Jagdamba Publishing.
- Upreti, D.K., U. Dubey and N. Sanjeeva. 2011. A new isidiate species of *Hemithecium* Trevis. (Ascomycota, Graphidaceae) from India. *The Lichenologist* 43(5): 483–485. doi: [10.1017/S0024282911000430](https://doi.org/10.1017/S0024282911000430)
- White F.J. and P.W. James. 1985. A new guide to microchemical techniques for the identification of lichen substances. *British Lichen Society Bulletin* 57(Suppl.): 1–41.

Author contributions: KPS collected the specimens, PS and KPS did identification and photography and wrote the text and ABB attended the review processes.

Received: 2 July 2015

Accepted: 16 September 2015

Academic editor: Sandra Costa-Böddeker

APPENDIX.

Table A1. List of microlichens of Arunachal Pradesh, India. Abbreviations: + Present; C, corticolous; F, foliicolous; S, saxicolous; T, tericolous; E, endemic.

Species	Family	Habitat	Altitudinal range (m)	Distribution in districts															
				Tawang	Papampu	West Kameng	East Kameng	Upper Subansiri	Lower Subansiri	West Siang	East Siang	Upper Siang	Lower Dieng Valley	Lohit	Ajiaw	Changlang	Trishap	Namsai	Voucher Number
<i>Aderomyces albostrigosus</i> (R. Sant.) Lücking, Sérus. & Vézda	Graphidaceae	CF	2500–2750																397(B) (BSA)
<i>Anisomeridium bifforme</i> (Borrer) R.C. Harris	Monoblastiaceae	C	ca. 1600																10561, 10569 (ASSAM)
<i>Anthracothecium prasinum</i> (Eschw.) R.C. Harris	Pyrenulaceae	C	300–400																10021 (ASSAM)
<i>Anthracothecium macrosporum</i> (Heppe) Müll. Arg.		C	200–700	+															2835, 9620 (ASSAM)
<i>Arthonia antillarum</i> (Fée) Nyf.	Arthoniaceae	C	ca. 400																474B (BSA)
<i>Arthonia cinnabarinina</i> (DC.) Wall.	"	C	1000–1200																10309, 10665 (ASSAM)
<i>Arthonia collective</i> Stirn.	"	"	"	C	—													Dubey et al. 2007	
<i>Arthonia inconspicua</i> Stirn.				C	ca. 250													4233 (ASSAM)	
<i>Arthonia palmulacea</i> (Müll. Arg.) R. Sant.				F	200–800													12410A (ASSAM)	
<i>Arthonia recedens</i> Stirn.				C	390–400													592A, 514 (BSA)	
<i>Arthonia subgyrosa</i> Nyf.				C	300–400													Dubey et al. 2007	
<i>Arthonia trilocularis</i> Müll. Arg.				F	200–800													12706E (ASSAM)	
<i>Arthonia</i> sp. 1				C	400–500													2939, 2945 (ASSAM)	
<i>Arthonia</i> sp. 2				C	ca. 1800													9941 (ASSAM)	
<i>Arthopyrenia claviformis</i> (Stirt.) D. Hawksw.	Arthopyreniaceae	C	E	350–850														3023 (ASSAM)	
<i>Arthopyrenia grisea</i> (Schleich. ex Schaefer), Körb.	"	C	ca. 300															3047 (ASSAM)	
* <i>Arthopyrenia maguscula</i> (Nyf.) Zahlbr.	"	C	ca. 202															5195 (BSA)	
<i>Arthopyrenia saxicola</i> A. Massal.		S	390–400															476 (BSA)	
* <i>Arthothelium abnorme</i> (Ach.) Müll. Arg.	Arthoniaceae	C	115–200															10120, 10165 (ASSAM)	
<i>Arthothelium chiodectoides</i> (Nyf.) Zahlbr.	"	C	—														Dubey et al. 2007		
<i>Arthothelium subsessile</i> (Nyf.) Makhija & Patw.	"	C	600–800														10502 (ASSAM)		
<i>Arthothelium</i> sp. 1	"	C	1350–1500														9874 (ASSAM)		
<i>Aspicilia ewalensis</i> Räsänen	Hymeneliaceae	S	E	ca. 2400													508B (BSA)		
<i>Aspidothelium scutellifolium</i> Lücking var. <i>indicum</i> Kr.P. Singh & Pinokiyo	Aspidotheliaceae	F	E	1600–800													12972E (CAL)		
<i>Asterothrygium decipiens</i> (Rehm) R. Sant.	Graphidaceae	F	500–600														12856D (ASSAM)		
<i>Asterothrygium rotuliforme</i> (Müll. Arg.) Sérus.	"	F	1100–1200														12864D (ASSAM)		
<i>Astrothelium megalayense</i> (Makhija & Patw.) Pushpi Singh & Kr. P. Singh	Tryptotheliaceae	C	E	1550–1700													589E (BSA)		
<i>Astrothelium subnitidisculum</i> (Makhija & Patw.) Pushpi Singh & Kr. P. Singh	"	C	ca. 250														2676 (ASSAM)		
<i>Aulaxina quadrangula</i> (Stirt.) R. Sant.	Graphidaceae	F	200–300														11819C (ASSAM)		
<i>Aulaxina uniseptata</i> R. Sant.	"	F	550–950														12850D (ASSAM)		
* <i>Bacidia heterochroa</i> (Müll. Arg.) Zahlbr.	Ramalinaceae	C	ca. 400														2912 (ASSAM)		
* <i>Bacidia laurocerasi</i> (Delise ex Duby) Vain.	"	C	300–600														4029 (BSA)		
* <i>Bacidia medialis</i> (Tuck. ex Nyf.) Zahlbr.	"	C	1100–1200														11282 (ASSAM)		
* <i>Bacidia millegiana</i> (Taylor) Zahlbr.	"	C	ca. 1575														5060B (BSA)		
<i>Bacidia nigrotusca</i> (Müll. Arg.) Zahlbr.	"	C	ca. 1400														619E (BSA)		
<i>Bacidia olivacea</i> (Vain.) Zahlbr.	"	F	200–300														11810 (ASSAM)		
* <i>Bacidia rubella</i> (Hoffm.) A. Massal.	"	C	750–1500														10592 (ASSAM)		
<i>Bacidia submedialis</i> (Nyf.) Zahlbr.	"	C	500–1700														284B (BSA), 7-013645 (LWG)		
<i>Bacidia apidatica</i> (Müll. Arg.) Vezda	"	F	225–900														12894P (ASSAM)		
<i>Bacidia mastothallina</i> (Vain.) Vezda	"	F	300–800														12977I (ASSAM)		

Continued

Table A1. *Continued.*

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts											
				Tawang		West Kameng		Lower Subansiri		Upper Subansiri		West Siang		Upper Siang	
		Altitudinal range (m)													
<i>Chroodiscus coccineus</i> (Leight.) Müll. Arg.	Graphidaceae	F	ca.250												
<i>Chroodiscus mirificus</i> (Kremp.) R. Sant.	"	F	200-800												
* <i>Chrysotrichia candelaris</i> (L.) J.R. Laundon	Chrysotrichaceae	C	200-1645												
<i>Coccocarpia palmicola</i> (Spreng.) Arn. & D.J.Galloway	Coccocarpicaceae	F	ca.325												
<i>Coccotremma cucurbitula</i> (Mont.) Müll. Arg.	Coccotremataceae	C	ca.2125												
<i>Coenogonium luteum</i> (Dicks.) Kalb & Lücking	Coenogoniaceae	CF	ca.1500												
<i>Coenogonium subluteum</i> (Rehm) Kalb & Lücking	CF	300-1340													
<i>Coenogonium zonatum</i> (Müll. Arg.) Kalb & Lücking	"	F	ca.1100												
* <i>Crespoa plurilocularis</i> Egea & Torrente	Roccellaceae	C	400-950												
<i>Cryptothecia candida</i> (Kremp.) R. Sant.	Arthoniaceae	C	ca.500												
<i>Cryptothecia effusa</i> (Müll. Arg.) R. Sant.	"	C	ca.200												
* <i>Cryptothecia faveolata</i> Makhija & Patw.	"	C	ca.750												
<i>Cryptothecia lunulata</i> (Zahlbr.) Makhija & Patw.	"	C	ca.900												
* <i>Cryptothecia punctulata</i> Makhija & Patw.	"	C	300-400												
* <i>Cryptothecia scripta</i> G. Thor	"	C	ca.200												
<i>Cryptothecia</i> sp.1	"	C	ca.500												
<i>Cyphellium inquinans</i> (Sm.) Trev.	Physciaceae	C	3200-3650												
<i>Diorymia heteroglyphium</i> (Pers.) Staiger & Kalb	Graphidaceae	C	300-1216												
<i>Diorymia junghuhnii</i> (Mont. & Bosch.) Kalb, Staiger & Elix	"	C	350-500												
<i>Diorymia longirellumatum</i> B.O. Sharma & Makhija	"	C	E	ca.1216											
<i>Diorymia macgeorgei</i> (Vain.) Staiger & Elix	"	C	ca.1216												
<i>Diorymia megaspodium</i> Kalb, Staiger & Elix	"	C	202-1700												
<i>Diorymia pachygraphum</i> (Nyl.) Kalb, Staiger & Elix.	"	C	850-1216												
<i>Diorymia pruinosum</i> (Eschw.) Kalb, Staiger & Elix	"	C	ca.300												
<i>Diorymia radiatum</i> (D.D.Awasthi & S.R. Singh) K.P.Singh & Swanalatha	"	S	E	ca.500											
* <i>Diorymia soozanum</i> (Zahlbr.) M. Nakan. & Kashiw.	"	C	ca.202												
* <i>Diploschistes caesiopubescens</i> (Nyl.) Vain.	Graphidaceae	S	ca.900												
* <i>Diploschistes cinereoracemosus</i> (Sw. ex Ach.) Vain.	"	T	1050-1750												
* <i>Diploschistes diacapsis</i> (Ach.) Lumbsch	"	S	ca.1000												
* <i>Diploschistes epiphylla</i> Fée	"	C	ca.150												
<i>Echinoplaca pellicula</i> (Müll. Arg.) R. Sant.	"	F	750-1100												
<i>Echinoplaca</i> sp.1	"	F	800-2000												
<i>Erythrodection malacum</i> (Kremp.) G. Thor	Roccellaceae	C	550-1200												
<i>Fellhanera bouteillei</i> (Desm.) Vezda	Pilocarpaceae	F	200-1500												
<i>Fellhanera fuscata</i> (Müll. Arg.) Vezda	"	F	500-1300												
<i>Fellhanera thapidophylli</i> (Rehm) Vezda	"	F	230-500												
<i>Fellhanera semicarpi</i> (Vain.) Vezda	"	F	1300-1500												
<i>Fissurina subcontexta</i> Ach.	Graphidaceae	C	ca.400												
<i>Fissurina</i> sp. 1	"	C	ca.136												
<i>Glyphis cicatrica</i> Ach.	"	C	202-1100												
<i>Glypis scyphulifera</i> (Ach.) Staiger	"	C	220-229												

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts	
				Altitudinal range (m)	Voucher Number
<i>Graphidula byssoides</i> (Müll. Arg.) G. Thor	Roccellaceae	C	1033–1170		08-009421 (LWG)
<i>Graphis glareolaria</i> Pat. & C.R.Kulk	Graphidaceae	C	ca.202		5257 (BSA)
<i>Graphis anfractuosa</i> (Eschw.) Eschw.	"	C	ca.500m		Dubey et al. 2007
<i>Graphis caesiella</i> Vain.	"	C	1000–1575		5068 (BSA)
<i>Graphis capillacea</i> Stirn.	"	C	300–400		10035 (ASSAM)
<i>Graphis chlorotica</i> A. Massal.	"	C	1350–1650		4698 (BSA)
<i>Graphis cincta</i> (Pers.) Aptroot	"	C	ca.1095		823 (ASSAM)
<i>Graphis contortuplicata</i> Müll. Arg.	"	C	ca.300		40159 (ASSAM)
<i>Graphis crebra</i> Vain.	"	C	—		10696D (ASSAM)
<i>Graphis cycasicola</i> A.W. Archer & Ellix	"	C	1950–2050		5683 (ASSAM)
<i>Graphis dianthensis</i> (A.W. Archer) A.W. Archer	"	C	ca.300		4640 (BSA)
<i>Graphis dimidiata</i> Vain.	"	C	1700–1900		2011 (ASSAM)
<i>Graphis duplicita</i> Ach.	"	C	680–1601		+
<i>Graphis farinulenta</i> Müll. Arg.	"	C	ca.500		282C (BSA)
<i>Graphis filiformis</i> Adaw. & Makhija	"	C	ca.1216		4872,4866 (BSA)
<i>Graphis furcata</i> Fée	"	C	400–1184		1116A (ASSAM)
* <i>Graphis galactodera</i> (Zahlbr.) Lücking	"	C	ca.200		10173 (BSA)
<i>Graphis glaucescens</i> Fée	"	C	136–200		10164 (BSA)
<i>Graphis handelii</i> Zahlbr.	"	C	ca.2250		10696D (ASSAM)
<i>Graphis intermedia</i> Stirn.	"	C	200–300		659 (ASSAM)
<i>Graphis japonica</i> (Müll.Arg.) A.W. Archer & Lücking	"	C	300–600		2136 (ASSAM)
<i>Graphis kousyuensis</i> (Horik. & M. Nakan.) Lücking	"	C	220–229		5313, 5294 (BSA)
<i>Graphis leptoclada</i> Müll. Arg.	"	C	ca.500		11038 (ASSAM)
<i>Graphis librata</i> C. Knight	"	C	500–1184		516 (BSA)
<i>Graphis longiramea</i> Müll. Arg.	"	C	250–900		967 (ASSAM)
<i>Graphis longispora</i> D.D.Awasthi & S.R.Singh	"	C	ca.900		4786 (ASSAM)
<i>Graphis marginata</i> Raddi	"	C	ca.400		2611 (ASSAM)
<i>Graphis nanodes</i> Vain.	"	C	250–300		55, 5305 (BSA)
<i>Graphis nuda</i> (Magn.) Staiger & Lücking	"	C	390–400		448 (BSA)
<i>Graphis oligospora</i> Zahlbr.	"	C	ca.500		10871 (ASSAM)
<i>Graphis paraserpens</i> Lizano & Lücking	"	C	1100–1500		64 (BSA)
<i>Graphis parilis</i> Krempp.	"	C	300–440		10084 (ASSAM)
<i>Graphis pavoniana</i> Fée	"	C	ca.202		5256 (ASSAM)
<i>Graphis perticosa</i> (Krempp.) A.W. Archer	"	C	1184–1800		1897 (ASSAM)
<i>Graphis pinicola</i> Zahlbr.	"	C	ca.202		5244 (BSA)
<i>Graphis prospersens</i> Vain.	"	C	2500–2750		376A (BSA)
<i>Graphis pyrhocheiloidea</i> Zahlbr.	"	C	ca.600		4277A (ASSAM)
<i>Graphis retschiana</i> (Müll. Arg.) Stizenb.	"	C	ca.160		4741 (BSA)
<i>Graphis scripta</i> (L.) Ach.	"	C	300–400		10690 (ASSAM)
<i>Graphis sitapurensis</i> Makhija & Adaw.	"	C	ca.1500		551A/2 (BSA)
<i>Graphis strebiocarpa</i> (Bel) Nyf.	"	C	750–900		4835 (ASSAM)
<i>Graphis striatula</i> (Ach.) Spreng.	"	C	550–1050		4787 (BSA)

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts												
				Altitudinal range (m)	Tawang	West Kameng	Lower Subansiri	Upper Subansiri	West Siang	Upper Siang	Lower Dibang Valley	Upper Dibang Valley	Changlang	Ajlaw	Namsai	Voucher Number
<i>Graphis subassimilis</i> Müll. Arg.	Graphidaceae	C	300–500	"	C	1625–1800										3002 (ASSAM)
<i>Graphis subserpentina</i> Nyf.		"		C	1550–1700											1900A (ASSAM)
<i>Graphis tenella</i> Ach.		"		C	400–1500											608 (BSA)
<i>Graphis valbaraiensis</i> Adáw. & Makhija		"		C	ca.400											5180 (ASSAM)
<i>Graphis vittata</i> Müll. Arg.		"		F	200–300											331 (ASSAM)
<i>Gyalectidium filicinum</i> Müll. Arg.		"		C	ca.1600											11830C (ASSAM)
* <i>Hematomma puniceum</i> (Sw.) A. Massal.	Haematommataceae	"		C	ca.2350											5286 (BSA)
<i>Haematomma wattii</i> (Stern.) Zahlbr.		"		C	2500–2700											298A (BSA)
<i>Haematomma sp. 1</i>		"		C	300–1600											394B (BSA)
<i>Hafellia curatellae</i> (Malme) Marbach	Physciaceae	C		C	ca.500											9752 (ASSAM)
<i>Hemitrichium amboliense</i> Makhija & A. Dube	Graphidaceae	C		C	ca.160											504F (ASSAM)
<i>Hemitrichium aphanes</i> (Mont. & Bosch.) M.Nakan. & Kashiw.		"		C	300–440											4757 (BSA)
<i>Hemitrichium isidiatum</i> Upadhyay & U. Dubey		"		C	ca.200											4901 (ASSAM)
<i>Hemitrichium nagalandicum</i> (K.R.P.Singh & G.P.Sinha)		"		C	1400–1250											06-006534 (LWG)
<i>Hemitrichium nakanishianum</i> (Patw. & C.R.Kulk.) Makhija & A.Dube		"		C	ca.136											11225 (ASSAM)
<i>Hemitrichium noristicum</i> Makhija & A.Dube		"		C	250–300											4652 (BSA)
* <i>Herpothallon cinerarium</i> G. Thor	Arthoniaceae	C		C	ca.1300											11292 (ASSAM)
* <i>Herpothallon echinatum</i> Aptroot		"		C	ca.750											7139 (ASSAM)
* <i>Herpothallon isidiatum</i> Jagadeesh Ram & G.P.Sinha		"		C	160–229											10472 (ASSAM)
<i>Herpothallon japonicum</i> (Zahlbr.) G. Thor		"		C	300–440											4234 (BSA)
<i>Herpothallon philippinum</i> (Vain.) Aptroot & Lücking		"		C	136–400											10091 (ASSAM)
<i>Herpothallon sticticum</i> Jagadeesh Ram & G.P.Sinha		"		E	ca.1200											346 (ASSAM)
<i>Ionaspis acutris</i> (Wirth) Lutzoni	Hymeneliaceae	C		ca.2400												11331 (ASSAM)
<i>Lasioloma arachnoidium</i> (Kemp.) R. Sant.	Pilocarpaceae	F		300–400												428C (BSA)
<i>Lasioloma phycophilum</i> (Vain.) R. Sant.		"		F	ca.325											12431B (ASSAM)
* <i>Laurea megasperma</i> (Mont.) Riddle	Trypetheliaceae	C		ca.1200												12884C (ASSAM)
<i>Laurea meristospora</i> (Mont. & Bosch.) Zahlbr.		"		ca.1550												583E (BSA)
<i>Lecanora achaia</i> Nyl.	Lecanoraceae	C		1200–1500												9908 (ASSAM)
<i>Lecanora alba</i> Lumbsch		"		ca.2050												9882,9875 (ASSAM)
<i>Lecanora aff. albella</i> (Pers.) Ach.		"		ca.1575												4773,4485 (ASSAM)
<i>Lecanora austointumescens</i> Lumbsch & Elix		"		ca.1250												10567,10605 (ASSAM)
<i>Lecanora conciliolanda</i> Vain.		"		ca.1500												4921 (BSA)
<i>Lecanora fimbriatula</i> Siltz.		"		C	600–900											4411,4389 (ASSAM)
<i>Lecanora helva</i> Sizzenb.		"		C	ca.1600											715 (ASSAM)
<i>Lecanora hennenniae</i> Vänska		"		S	1500–2300											Phonkiyo et al. 2008
<i>Lecanora impudens</i> Degel.		"		C	ca.1100											4212,4095 (ASSAM)
<i>Lecanora inshaqii</i> Brodo		"		C	600–800											388A (BSA)
<i>Lecanora interjecta</i> Müll. Arg.		"		C	ca.200											4969 (BSA)
<i>Lecanora perplexa</i> Brodo		"		C	2500–2750											
<i>Lecanora phaeocarpa</i> Vain.		"		C	ca.2125											

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts											
				Tawang		West Kameng		Lower Subansiri		Upper Subansiri		West Siang		Upper Siang	
<i>Lecanora streimanii</i> Lumbsch	Lecanoraceae	C	ca. 500												
<i>Lecanora subalbellina</i> Vain.	"	C	ca. 1600												
<i>Lecanora subimmersa</i> (Fée) Vain.	"	C	600–800												
<i>Lecanora subapponical</i> Lü & H.Y. Wang	"	C	ca. 2250												
<i>Lecanora tropica</i> Zahlbr.	"	C	ca. 2100												
<i>Lecanora wilsonii</i> Müll. Arg.	"	S	1500–1550												
<i>Lecanora</i> sp. 1	"	C	2306–2344												
<i>Lecidea lapicida</i> (Ach.) Ach.	Lecideaceae	S	600–800												
<i>Lecidea</i> sp. 1	"	C	500–950												
<i>Leiorreuma exaltatum</i> (Mont. & Bosch) Staiger	Graphidaceae	C	400–900												
<i>Leiorreuma lyellii</i> (Sm.) Staiger	"	C	ca. 500												
<i>Leiorreuma subpatellulum</i> Dubey, Upadhyay & Nayaka	"	C	E	900											
<i>Lepraria aff. casieoalba</i> (B. de Lesd.) J.R. Laundon	Stereocaulaceae	C	ca. 1100												
* <i>Lepraria lobifrons</i> Nyf.	"	C	ca. 200												
<i>Letrouitia transgressa</i> (Malme) Hafellner & Bellerm	Letrouitiaceae	C	150–900												
<i>Lithothelium decumbens</i> (Müll. Arg.) Aptroot	Pyrenulaceae	C	—												
* <i>Lithothelium hyalosporum</i> (Nyf.) Aptroot	"	C	ca. 136												
<i>Lithothelium obectatum</i> (Müll. Arg.) Aptroot	"	C	ca. 500												
<i>Lofiamnia gabrielis</i> (Müll. Arg.) Vezda	Pilocarpaceae	F	ca. 800												
<i>Lofiamnia intermedia</i> (R. Sant.) Vezda	"	F	230–500												
<i>Malmidea granifera</i> (Ach.) Kalb, Rivas Plata & Lumbsch	Malmideaceae	C	500–700												
<i>Maronea manipurensis</i> H. Magn.	Fuscideaceae	C	ca. 2350												
<i>Mazosia bambusae</i> (Vain.) R. Sant.	Roccellaceae	F	500–800												
<i>Mazosia melanophthalma</i> (Müll. Arg.) R. Sant.	"	F	200–800												
<i>Mazosia phyllospema</i> (Nyf.) Zahlbr.	"	F	200–800												
<i>Mazosia rotula</i> (Mont.) A. Massal.	"	F	200–650												
<i>Mazosia tumidula</i> (Stern.) Müll. Arg.	"	F	200–850												
<i>Megalaria laurenii</i> (Hép.) ex Th. Fr. Hafellner	Megalariaeae	C	ca. 1500												
<i>Megalospora atrorubicans</i> (Nyf.) Zahlbr.	Megalosporaceae	C	375–1600												
<i>Megalospora tuberosculosa</i> (Fée) Sipman	"	C	ca. 1600												
<i>Mycobilimbia humana</i> (Zahlbr.) D.D. Awasthi	Lecideaceae	T	ca. 1800												
<i>Mycobilimbia philippina</i> (Vain.) D.D. Awasthi	"	S	ca. 2350												
<i>Mycoblastus affinis</i> (Schaeff.) Schaefer	Mycoblastaceae	C	3200–3600												
<i>Myconicrothelia conothelena</i> (Nyf.) D. Hawksw.	Arthopyreniaceae	C	ca. 2344												
* <i>Myriotrema microporum</i> (Mont.) Hale	Graphidaceae	C	1500–1700												
* <i>Myriotrema rugiferum</i> (Harm.) Hale	"	C	ca. 1550												
* <i>Ocellularia allosporoides</i> (Nyf.) Patw. & C.R. Kulk.	Graphidaceae	C	ca. 300												
<i>Ocellularia neopertusiformis</i> Hale	"	C	200–300												
<i>Ocellularia subgranulosa</i> (Homchanta & Coppins) Lumbsch & Papong	Ochrolechiaceae	C	ca. 340												
* <i>Ochrolechia hamannii</i> Verseghy	"	C	2360–2344												
* <i>Ochrolechia subpalleicensis</i> Verseghy	Roccellaceae	F	350–600												
<i>Opegrapha filicina</i> Mont.															

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts											
				Tawang		West Kameng		Upper Subansiri		Lower Subansiri		Upper Siang		Lower Dieng Valley	
					Altitudinal range (m)										
* <i>Opegrapha vulgata</i> (Ach.) Ach.	Roccellaceae	C	ca. 500												
<i>Ophioparma ventosa</i> (L.) Norman	Ophioparmaceae	C	3000–3470												
* <i>Pallidogramme bengalense</i> B.O.Sharma & P.Khadilkar	Graphidaceae	C	E 1250–1500												
<i>Pallidogramme chrysenteron</i> (Mont.) Staiger, Kalb & Lücking	"	C	ca. 900												
<i>Pallidogramme sp. 1</i>	Pertusariaceae	"	1250–1500												
* <i>Pertusaria alpina</i> Hepp.	Pertusariaceae	C	ca. 1601												
* <i>Pertusaria ceylonica</i> Müll. Arg.	"	C	ca. 800												
<i>Pertusaria coccodes</i> (Ach.) Nyl.	"	C	1100–1300												
* <i>Pertusaria colorata</i> Awasthi & Srivastava	"	C	E ca. 1360												
<i>Pertusaria composita</i> Zahlbr.	"	C	ca. 2123												
<i>Pertusaria indica</i> Preeti Srivast. & D.D. Awasthi	"	S	E ca. 400												
<i>Pertusaria leucosordes</i> Nyl.	"	C	ca. 2123												
<i>Pertusaria multipunctata</i> (Turn.) Nyl.	"	C	1800–3200												
* <i>Pertusaria pertusa</i> (L.) Tuck.	"	C	ca. 1200												
<i>Pertusaria quassiae</i> (Fee) Nyl.	"	C	ca. 750												
<i>Pertusaria submultipunctata</i> Nyl.	"	C	200–300												
<i>Pertusaria sp. 1</i>	"	C	2600–2750												
<i>Phaeographis caesiioradians</i> (Leight.) A.W. Archer	Graphidaceae	C	300–600												
<i>Phaeographis dendroides</i> (Leight.) Müll. Arg.	"	C	ca. 500												
<i>Phaeographis divaricoides</i> Räsänen	"	C	E 1500–1550												
<i>Phaeographis intricans</i> (Nyl.) Staiger	"	C	1500–1700												
<i>Phaeographis sculptata</i> (Ach.) Staiger	"	C	300–800												
<i>Phaeographopsis indica</i> Patw. & Nagarkar Sipman & Aptroot	"	C	ca. 202												
* <i>Phylctis karnatakana</i> S. Joshi & Upadhyay	Phlyctidaceae	C	E 750–1500												
<i>Phylctis subhimalayensis</i> S. Joshi & Upadhyay	"	C	E –												
<i>Phyllobathelium indicum</i> G.P. Sinha & Kr. P. Singh	Phlyctidaceae	F	E 200–300												
<i>Playgramme caesiopruinosa</i> (Fee) Fee	"	C	1625–1800												
<i>Playgramme discurrenti</i> (Nyl.) Staiger	"	C	ca. 900												
* <i>Playgramme muelleri</i> (A.W. Archer) Staiger	"	C	300–800												
<i>Pilaria montagnei</i> (Bosc.) A. Massal.	"	C	–												
<i>Porina albicera</i> (Kremp.) Overeem	Porinaceae	F	200–800												
<i>Porina andamanica</i> Makhija & al.	"	C	E 1100–1200												
<i>Porina applanata</i> Vain.	"	F	ca. 500												
<i>Porina articeps</i> (Vain.) Vain.	"	F	200–1300												
<i>Porina atrocoerulea</i> Müll. Arg.	"	F	950–1150												
<i>Porina atroperiostola</i> Makhija, Adawa & Patw.	"	C	E 400–500												
<i>Porina belanospora</i> (Müll. Arg.) F. Schill.	"	C	300–400												
<i>Porina cupreola</i> (Fee) Fee	"	F	200–300												
<i>Porina epiphylla</i> Müll. Arg.	"	F	200–1550												
<i>Porina fulvella</i> Müll. Arg.	"	F	500–800												

Continued

Table A1. Continued.

Species	Family	Endemic status	Distribution in districts	
			Altitudinal range (m)	Voucher Number
* <i>Porina glaucoflava</i> Makhija & al.	Porinaceae	C	300-400	9838 (ASSAM)
* <i>Porina halei</i> Makhija & al.		C	1100-1200	31 (BSA)
<i>Porina imitatrix</i> Müll. Arg.		"	200-1150	11856A (ASSAM)
<i>Porina innata</i> (Nyl.) Müll. Arg.		"	ca.650	4205 (ASSAM)
<i>Porina intermigrans</i> (Nyl.) Müll.		C	ca.500	9936 (ASSAM)
<i>Porina kamerunensis</i> F. Schill.		"	200-300	11816A (ASSAM)
<i>Porina kannatakensis</i> Makhija & al.		F	200-800	11816B (ASSAM)
<i>Porina limbulata</i> (Kremp.) Vain.		F	200-800	11557A (ASSAM)
<i>Porina lucida</i> R. Sant.		F	200-800	11811C (ASSAM)
<i>Porina mastoidella</i> (Nyl.) Müll. Arg.		C	ca.600	10788 (ASSAM)
<i>Porina monocarpa</i> (Kremp.) F. Schill.		F	200-1100	11795B (ASSAM)
<i>Porina napensis</i> Lücking		F	1200	13276G (ASSAM)
<i>Porina nitidula</i> Müll. Arg.		F	200-800	11799B (ASSAM)
<i>Porina rufula</i> (Kremp.) Vain.		F	200-1550	11812B (ASSAM)
<i>Porina subalpina</i> Upreti		C	600-800	4210 (ASSAM)
<i>Porina subinterstes</i> (Nyl.) Müll. Arg.		"	ca.1500	630A (ASSAM)
<i>Porina tetracerata</i> (Afz.) Müll. Arg.		C/F	650-1500	615A (BSA), 4209 (ASSAM)
<i>Porina iijiciana</i> Vain.		S	ca.900	4783 (ASSAM)
<i>Porina trichothelioides</i> R. Sant.		F	200-800	11802B (ASSAM)
<i>Porina</i> sp. 1		S	ca.500	5547 (ASSAM)
<i>Porina</i> sp. 2		C	ca.930	3040 (ASSAM)
<i>Porina virescens</i> (Kremp.) Müll. Arg.		"	200-800	11829B (ASSAM)
* <i>Porpidia albocoerulescens</i> (Wulfen) Hertel & Knoph	Lecideaceae	S	950-1150	10521, 8031 (ASSAM)
* <i>Porpidia macrocarpa</i> (DC.) Hertel & Knoph		S	ca.1100	4453 (ASSAM)
<i>Pseudopyrenula diluta</i> (Fée) Müll. Arg.	Trypethelaceae	C	ca.400	505B (BSA)
* <i>Pseudopyrenula subnudata</i> Müll. Arg.		C	ca.136	4691 (BSA)
<i>Pyrenula aggregata</i> (Fée) Fée	Pyrenulaceae	C	1500-1700	10729 (ASSAM)
<i>Pyrenula anomala</i> (Ach.) Vain.		C	400-500	1111 (ASSAM)
<i>Pyrenula aspista</i> (Ach.) Ach.		"	ca.136	4689 (BSA)
<i>Pyrenula balia</i> (Kremp.) R.C. Harris		"	-	Upreti 1998 (LWG)
<i>Pyrenula breutelli</i> (Müll. Arg.) Aptroot		C	ca.400	1146 (ASSAM)
<i>Pyrenula brunnea</i> Fée		C	300-400	10080 (ASSAM)
<i>Pyrenula castanea</i> (Eschw.) Müll. Arg.		C	100-1500	4014, 4039 (ASSAM)
<i>Pyrenula complanata</i> (Mont.) Trevis.		C	ca.1000	10820 (ASSAM)
<i>Pyrenula duplicans</i> (Nyl.) Aptroot		C	ca.400	1145 (ASSAM)
<i>Pyrenula festivica</i> (Kremp.) Müll. Arg.		C	ca.400	10490 (ASSAM)
<i>Pyrenula globifera</i> (Eschw.) Aptroot		C	ca.400	10384 (ASSAM)
<i>Pyrenula immissa</i> (Sitt.) Zahlbr.		C	ca.1216	10541 (ASSAM)
<i>Pyrenula interducta</i> (Nyl.) Zahlbr.		"	-	Dubey et al. 2007
<i>Pyrenula leucotrysma</i> (Nyl.) Upreti		C	ca.500	102A (BSA)
<i>Pyrenula maravalensis</i> Vain.		C	100-1400	11219 (ASSAM)

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts	
				Altitudinal range (m)	Voucher Number
<i>Pyrenula oculata</i> Ajay Singh & Upreti	Pyrenulaceae	C	ca. 1000	"	3054 (ASSAM)
<i>Pyrenula papillifera</i> (Nyl.) Aptroot	"	C	1500–1550	"	615C (BSA)
<i>Pyrenula playstoma</i> (Müll. Arg.) Aptroot	"	C	ca. 1600	"	9762 (ASSAM)
<i>Pyrenula punctella</i> (Nyl.) Trevis.	"	C	600–800	"	10490 (ASSAM)
<i>Pyrenula quassiaecola</i> Féé	"	C	1550–1700	"	87C (BSA)
<i>Pyrenula subelliptica</i> (Tuck.) R.C. Harris	"	C	ca. 1575	"	5060 (BSA)
<i>Pyrenula sublaevigata</i> (Patw. & Makhija) Upreti	"	C	ca. 1000	"	10744 (ASSAM)
<i>Pyrenula thailandica</i> Aptroot	"	C	ca. 300	"	81724 (LWG)
<i>Pyrenula zeylanica</i> Upreti & Ajay Singh	"	C	E	"	20150A (LWG)
<i>Pyrailus cubanus</i> Nyl.	"	C	1500–1700	"	587C (BSA)
<i>Pyrailus javanicus</i> (Mont. & Bosch.) Nyl.	"	C	1500–1850	"	9316 (ASSAM)
<i>Pyrailus tibetii</i> Kr.P. Singh & Pushpi Singh	"	C	600–800	"	10495, 10507 (ASSAM)
* <i>Ramboldia tessula</i> (Ach.) Lumbsch & Elix	Lecanoraceae	C	ca. 340	"	9616 (ASSAM)
* <i>Rhabdodiscus asiaticus</i> (Vain.) Rivas Plata, Lücking & Lumbsch	Graphidaceae	C	ca. 1645	"	5038 (BSA)
* <i>Rhabdodiscus auberianus</i> (Nyl.) Vain.	"	C	600–800	"	10549 (ASSAM)
<i>Rhabdodiscus cossus</i> (Müll. Arg.) Rivas Plata, Lücking & Lumbsch	"	C	ca. 500	"	323 (ASSAM)
<i>Rhabdodiscus indicus</i> Pushpi Singh & Kr. P. Singh	"	C	ca. 1550	"	547F (BSA)
* <i>Rhizocarpon geographicum</i> (L.) DC.	Rhizocarpaceae	S	600–800	"	10512 (ASSAM)
<i>Rhizocarpon sp. 1</i>	"	S	3200–3650	"	7829, 7747 (ASSAM)
<i>Rinodina intrusa</i> (Kemp.) Malme	"	S	3000–3470	"	9524 (ASSAM)
<i>Rinodina mackenziei</i> Räsänen	"	S	ca. 500	"	591E (ASSAM)
<i>Rinodina sophodes</i> (Ach.) A. Massal.	"	S	1704–1900	"	334 (BSA)
<i>Rinodina sp. 1</i>	"	S	—	"	Phnokiyo et al. 2008
<i>Sarcographa heteroclitia</i> (Mont.) Zahlbr.	Physciaceae	C	1800–2400	"	11441 (ASSAM)
<i>Sarcographa glyphiza</i> (Nyl.) Kr.P. Singh & G.P. Sinha	"	C	ca. 500	"	Dubey et al. 2007
<i>Sarcographa labyrinthica</i> (Ach.) Müll. Arg.	"	C	300–600	"	10090 (ASSAM)
<i>Sarcographa medusulina</i> (Nyl.) Müll. Arg.	"	C	250–600	"	761, 977 (ASSAM)
<i>Sarcographa subtorquescens</i> (Nyl.) Zahlbr.	"	C	—	"	Dubey et al. 2007
<i>Sarcographa tricosa</i> (Ach.) Müll. Arg.	"	C	400–900	"	Dubey et al. 2007
<i>Schistophoron indicum</i> Kr.P.Singh & Swarnalatha	"	C	500–700	"	969, 11117 (ASSAM)
<i>Sporopodium argillaceum</i> (Müll. Arg.) Zahlbr.	Pilocarpaceae	F	200–300	"	2858 (BSA)
<i>Sporopodium awasthiicum</i> Kr.P.Singh & Pinokiyo	"	F	300–400	"	11833 (ASSAM)
<i>Sporopodium phyllocharris</i> (Mont.) A. Massal.	"	F	230–850	"	12818C (ASSAM)
<i>Sporopodium xantholeucum</i> (Müll. Arg.) Zahlbr.	"	F	200–1300	"	12931 (ASSAM)
<i>Stauropelta clopima</i> (Vahlent.) Th. Fr.	"	C	ca. 1000	"	11798A (ASSAM)
<i>Strigula antillarum</i> (Fée) Müll. Arg.	"	F	325–850	"	10757 (ASSAM)
<i>Strigula concreta</i> (Fée) R. Sant.	"	F	200–500	"	12883P (ASSAM)
<i>Strigula Janeirensis</i> (Müll. Arg.) Lücking	"	F	ca. 500	"	11760A (ASSAM)
<i>Strigula maculata</i> (Cook & Masse) R. Sant.	"	F	750–1150	"	12930A (ASSAM)
<i>Strigula melanobapha</i> (Kemp.) R. Sant.	"	F	ca. 325	"	12630B (ASSAM)
<i>Strigula multipunctata</i> (G. Merr. ex R. Sant.) R.C. Harris	"	F	ca. 800	"	12883C (ASSAM)
					12336C (ASSAM)

Continued

Table A1. Continued.

Species	Family	Habitat	Endemic status	Distribution in districts	
				Altitudinal range (m)	Voucher Number
<i>Strigula nemathora</i> Mont. f. <i>hypothelia</i> (Nyl.) Lücking	Strigulaceae	F	500-1300	"	11780E (ASSAM)
<i>Strigula nemathora</i> Mont. f. <i>nemathora</i> Mont.		F	ca. 800	"	77A (ASSAM)
<i>Strigula nitidula</i> Mont.		F	200-300	"	11810G (ASSAM)
<i>Strigula orbicularis</i> Fr.		F	200-800	"	12984F (ASSAM)
<i>Strigula phyllogena</i> (Müll. Arg.) R.C. Harris		F	200-800	"	11970A (ASSAM)
<i>Strigula smaragdula</i> Fr.		F	500-1300	"	12307 (ASSAM)
<i>Strigula subelegans</i> Vain.		F	235-800	"	12322 (ASSAM)
<i>Strigula subtilissima</i> (Fée) Müll. Arg.		F	200-800	"	12635D (ASSAM)
<i>Tapellaria bilimboides</i> R. Sant.	Pilocarpaceae	F	500-1300	"	11770A (ASSAM)
<i>Tapellaria epiphylla</i> (Müll. Arg.) R. Sant.		F	500-1300	"	11786 (ASSAM)
<i>Tapellaria molleri</i> (Henrik) R. Sant.		F	500-1300	"	11747B (ASSAM)
<i>Tapellaria nana</i> (Fée) R. Sant.		F	ca. 325	"	12883 (ASSAM)
<i>Tapellaria nigra</i> (Müll. Arg.) R. Sant.		F	950-1150	"	12961F (ASSAM)
* <i>Thecaria austroindica</i> (D.D. Awasthi & Upreti) Kr.P. Singh & G.P. Sinha	Graphidaceae	C	ca. 400	"	520 (ASSAM)
<i>Thecaria quassiicola</i> Fee		C	400-500	"	1141 (ASSAM)
<i>Tricharia santessonii</i> D. Hawksw.		F	ca. 800	"	12623F (ASSAM)
<i>Tricharia vainioi</i> R. Sant.		F	700-1500	"	13085C (ASSAM)
<i>Trichothelium epiphyllum</i> Müll. Arg.	Porinaceae	F	200-1550	"	11886B (ASSAM)
<i>Trypethelium abpruinoseum</i> Makhija & Patw.	Trypeteliaceae	C	E	"	Dubey et al. 2007
<i>Trypethelium dichroum</i> Makhija & Patw.		C	E	"	
<i>Trypethelium elatior</i> Spreng.		C	E	"	
<i>Trypethelium inamoenum</i> Müll. Arg.		C	E	"	
<i>Trypethelium tropicum</i> (Ach.) Müll. Arg.		C	E	"	
* <i>Tylophoron moderatum</i> Nyl.		C	E	"	
* <i>Tylophoron nidulans</i> Strit.		C	E	"	
<i>Verrucaria cethabola</i> (Wahlb.)	Verrucariaceae	S	-	"	7139, 10797 (ASSAM)
<i>Verrucaria coerulea</i> (Ramond) DC.		S	ca. 1000	"	2871 (ASSAM)
<i>Verrucaria transiliensis</i> Arnold		S	2306-2344	"	Dubey et al. 2007
				"	4861 (ASSAM)
				"	7829 (ASSAM)