

## LISTS OF SPECIES

### Angiosperms, Climbing plants in tropical forests of southern Eastern Ghats, Tamil Nadu, India

Chellam Muthumperumal  
Narayanaswamy Parthasarathy

*Department of Ecology and Environmental Sciences, Pondicherry University.  
Puducherry – 605 014, India. E-mail: parthapu@yahoo.com*

#### Abstract

We provide a check list of angiosperm climbing plant species, along with their climbing modes, enumerated from a total of one hundred and fifty grids in tropical forests of southern Eastern Ghats, peninsular India. The Eastern Ghats constitute an important biodiversity area in India and have been studied earlier mainly for the floristics, and that too confined to a few prioritized sites. Lianas, the woody vines contribute substantially to the diversity and structure of most tropical forests. Yet, little is known about the importance of habitat specialization in maintaining tropical liana diversity. A total of 175 climbing plant species that belong to 100 genera and 40 families are included in this enumeration.

#### Introduction

Lianas, the woody vines are an important, but understudied growth-form, common to most forests throughout the world, particularly in the tropics (Schnitzer and Bongers 2002). The past few decades have brought increasing awareness of the importance of lianas to species diversity (Gentry and Dodson 1987; Schnitzer and Carson 2001; Burnham 2002). Lianas constitute ~ 25 % of the woody stem density and species diversity in tropical forests (Gentry 1991). They are often a large component of the canopy in tropical forests, often one-third or more of the entire leaf area, though only a small component of the basal area and biomass (Schnitzer and Bongers 2002). Recent studies on lianas have focused on the significant contribution of this life-form to the overall density and species diversity of tropical forests (Mascaro et al. 2004; Rice et al. 2004), mechanisms by which lianas alter the tropical forest diversity and regeneration (Schnitzer and Carson 2001), harming some shade tolerant species, while promoting some pioneer species (Schnitzer et al. 2000; Toledo-Aceves and Swaine 2007), and their significant contribution to carbon sequestration (Restom and Nepstad 2001; Schnitzer and Bongers 2002).

The mean abundance, diversity and taxonomic composition of lianas in lowland tropical moist and wet forests are similar among tropical regions, although liana abundance is higher in Africa

(Schnitzer and Bongers 2002). Liana density ( $\geq 1.6$  cm diameter) decreased with increasing altitude, whereas richness was highest at intermediate elevations (Parthasarathy et al. 2004). Reports suggest that lianas are now increasing in tropical forests as a result of the recent climatological trend (Malhi and Wright 2004). The higher frequency of lianas is not only caused by declining rainfall (Swaine and Grace 2007) but also several other factors are known to favour them such as an increase in disturbance (Laurance et al. 2001; Ibarra-Manriquez and Martinez-Ramos 2002; Londre and Schnitzer 2006), or an especially high responsiveness to elevated CO<sub>2</sub> (Grandos and Körner 2002; Zotz et al. 2006). More recently (Schnitzer 2005) examined floristic data from 69 tropical forests worldwide and found a negative correlation between mean annual precipitation and liana abundance.

#### Materials and methods


The Eastern Ghats constitute an important biogeographic region, as one of the nine floristic zones in India and form a broken chain of mountains spreading along the states of Orissa, Karnataka, Andhra Pradesh and Tamil Nadu and lie between Mahanadhi and Vaigai rivers. The present study was carried out in southern-most Eastern Ghats which falls in Tamil Nadu (10° 93'-12° 18' N) (Figure 1) in six major hill complexes viz., Chitteris, Kolli hills, Pachaimalais,

## LISTS OF SPECIES

Shervarayans, Kalrayans and Sathyamangalam forests (Figures 2 and 3). These sites harbor five distinct forest types -tropical evergreen, semi-evergreen, mixed deciduous, dry deciduous and thorn forests.

The overall climate in the Eastern Ghats is dry and salubrious. The climatic conditions favour both the south-west and the north-east monsoons. The months of April to June are hottest, while

November to February is relatively cooler. The average summer temperature in the plains can reach up to 40 °C particularly in some of the semi-arid areas in interior Tamil Nadu, while the average minimum temperature normally stays around 18-20 °C and the relative humidity is quite high throughout the year 70-75 %. Much of the region's rainfall occurs between October and December and ranges from 635-1,905 mm a year (Areendran and Rao 2006).



**Figure 1.** Map: (a) India, (b) Eastern Ghats stretch, (c) Southern Eastern Ghats.

Enumeration of angiosperm climbing plants was carried in a total of 150 grids in forests of southern Eastern Ghats. The entire stretch of southern Eastern Ghats ( $10^{\circ}93' - 12^{\circ}18'$  N) was divided into  $6.25\text{ km} \times 6.25\text{ km}$  grids and within each grid a 0.5 ha transect ( $5\text{ m} \times 1\text{ km}$ ) was established. Each transect was subdivided into fifty  $5\text{ m} \times 20\text{ m}$  units. All lianas - 1.5 cm dbh (5gbh, girth at breast height) were enumerated in the whole transect, and those of herbaceous vines of  $<1.5\text{ cm dbh}$  only in the beginning and end of the transect. Voucher specimens were collected

and identified using regional floras (Hooker 1872-1885; Gamble and Fischer 1921-1935; Matthew 1991), confirmed with the collections lodged in Department of Ecology and Environmental Sciences (DEES), Pondicherry University and Botanical Survey of India, Coimbatore (MH) and deposited in the herbarium of DEES.

### Results and Discussion

A total of 175 angiosperm climbing plants in 100 genera and 40 families were recorded from the total of 150 grids of southern Eastern Ghats

## LISTS OF SPECIES

forests. Jongkind and Hawthorne (2005) reported 746 species of climbers in Upper Guinean forests and the total number of climbing plants species that are found in Puerto Rico and the Virgin Islands amounts to 386 (Acevedo-Rodriguez 2005). The most speciose families in our sites include Asclepiadaceae (18 species), Convolvulaceae and Papilionaceae (13 species each), Apocynaceae (12 species), Vitaceae (11 species) and Menispermaceae (9 species) etc. The most abundant liana species include the thorny stragglers *Pterolobium hexapetalum* (Caesalpiniaceae), *Lantana camara* (Verbenaceae), *Ziziphus oenoplia* (Rhamnaceae), and the twiners *Jasminum angustifolium* (Oleaceae), *Gymnema sylvestre*, *Secamone emetica* (Asclepiadaceae) and *Aganosma cymosa* var. *cymosa* (Apocynaceae). The rare taxa include *Aganosma cymosa* var. *elegans*, *A. cymosa* var. *lanceolata*, *Ellertonia rheedii* (Apocynaceae), *Argyreia sericea* (Convolvulaceae), *Butea parviflora* (Papilionaceae), *Caesalpinia cucullata* (Caesalpiniaceae), *Capparis shevaroyensis*, *C. sepiaria* var. *retusella* (Capparaceae), *Cissus gigantea* (Vitaceae), *Cocculus pendulus* (Menispermaceae), *Dalbergia congesta* (Papilionaceae), *Rubus niveus* (Rosaceae), *Salacia oblonga* (Celastraceae) and *Ventilago goughii* (Rhamnaceae), all of which occurred in any one of the 150 study transects. Even though *Olax*

*scandens* (Olacaceae), *Chilocarpus atrovirens* (Apocynaceae), *Artobotrys zeylanicus* (Annonaceae) and *Calamus gamblei* (Arecaceae) were reported as most abundant species in the Western Ghats and *Strychnos minor* (Loganiaceae) in the tropical dry evergreen forests on the Coromandel Coast of India (Parthasarathy et al. 2004) these species did not occur in our sites. Pictures of selected liana species are featured in Figures 3-13. Plant binomial, family and voucher number of the 175 liana species are listed in Table 1. The enumerated climbing plants were classified into woody vines, the lianas (145 species) and herbaceous vines (30 species). Six climbing modes of lianas were recognized: stem twiners (46 %) followed by stragglers-armed (22 %), stragglers-unarmed (17 %), tendril climbers (13 %), root climbers (1 %) and hook climber (0.5 %). Only one climbing mode, the grapple-like climber (rattans) which was reported from Indian Western Ghats sites (Muthuramkumar and Parthasarathy 2000; Padaki and Parthasarathy 2000), did not occur in our study transects. Presently, many forest sites of southern Eastern Ghats are subjected to various anthropogenic pressures and the data of plant diversity, such as this on lianas will be useful in highlighting the importance of these forests in species conservation and forest management.

**Table 1.** Angiosperm climbing plants enumerated from 150 grids in tropical forests of southern Eastern Ghats, Tamil Nadu, peninsular India with their binomial, family, category, climbing mode and voucher number. WV: Woody vines; HV: Herbaceous vines; ST: Stem twiners; Str-A: Stragglers-armed; Str-UA: Stragglers-unarmed; TC: Tendril climbers; RC: Root climbers; and HC-1: Hook climber.

Species / Family	Category	Climbing mode	Voucher No.
Annonaceae			
<i>Desmos viridiflora</i> (Bedd.) Safford	WV	Str-UA	6056
<i>Uvaria narum</i> (Dunal) Wall. ex Wight & Arn.	WV	Str-UA	6324
Apocynaceae			
<i>Aganosma cymosa</i> (Roxb.) G. Don var. <i>cymosa</i>	WV	ST	6037
<i>Aganosma cymosa</i> (Roxb.) G. Don var. <i>elegans</i> Hook. f.	WV	ST	6454
<i>Aganosma cymosa</i> (Roxb.) G. Don var. <i>lanceolata</i> Hook. f.	WV	ST	6262
<i>Anodendron paniculatum</i> A. DC.	WV	ST	6176
<i>Carissa carandas</i> L.	WV	Str-A	6020
<i>Carissa gangetica</i> Stapf	WV	Str-A	6271
<i>Carissa paucinervia</i> A. DC.	WV	Str-A	6273
<i>Carissa salicina</i> Lam.	WV	Str-A	6263
<i>Carissa spinarum</i> L.	WV	Str-A	6076

## LISTS OF SPECIES

Species / Family	Category	Climbing mode	Voucher No.
<i>Ellerntonia rheedii</i> Wight	WV	ST	6038
<i>Ichnocarpus frutescens</i> (L.) R. Br.	HV	ST	6067
<i>Ichnocarpus ovatifolius</i> A. DC.	WV	ST	6041
Araceae			
<i>Rhaphidophora laciniata</i> (Burm.f.) Merr.	WV	RC	6178
Aristolochiaceae			
<i>Aristolochia indica</i> L.	HV	ST	6432
<i>Aristolochia tagala</i> Cham.	WV	ST	6168
Asclepiadaceae			
<i>Cosmostigma racemosum</i> (Roxb.) Wight	WV	ST	6025
<i>Cryptolepis buchanani</i> Roemer & Schultes	WV	ST	6018
<i>Cynanchum callialatum</i> Buch.-Ham. ex Wight & Arn.	WV	ST	6241
<i>Decalepis hamiltonii</i> Wight & Arn.	WV	ST	6060
<i>Gymnema hirsutum</i> Wight & Arn.	WV	ST	6321
<i>Gymnema montanum</i> (Roxb.) Hook. f. var. <i>beddomei</i> Hook. f.	HV	ST	6334
<i>Gymnema sylvestre</i> (Retz.) R.Br.ex Roemer & Schultes	WV	ST	6006
<i>Gymnema tingens</i> (Roxb.) Wight & Arn.	WV	ST	6246
<i>Hemidesmus indicus</i> (L.) R. Br. var. <i>indicus</i>	HV	ST	6083
<i>Hemidesmus indicus</i> (L.) R. Br. var. <i>pubescens</i> (Wight & Arn.) Hook. f.	HV	ST	6428
<i>Marsdenia brunoniana</i> Wt. & Arn.	WV	ST	6244
<i>Marsdenia tenacissima</i> (Roxb.) Moon	WV	ST	6030
<i>Pergularia daemia</i> (Forssk.) Chiov.	HV	ST	6224
<i>Sarcostemma acidum</i> (Roxb.) Voigt	WV	Str-UA	6147
<i>Secamone emetica</i> (Roxb.) R. Br. ex Schultes	WV	ST	6010
<i>Tylophora capparidifolia</i> Wight & Arn.	WV	ST	6087
<i>Tylophora indica</i> (Burm.f) Merr.	HV	ST	6128
<i>Wattakaka volubilis</i> (L.f.) T. Cooke	WV	ST	6043
Basellaceae			
<i>Basella alba</i> L.	HV	ST	6234
Caesalpiniaceae			
<i>Caesalpinia crista</i> L.	WV	Str-A	6027
<i>Caesalpinia cucullata</i> Roxb.	WV	Str-A	6259
<i>Pterolobium hexapetalum</i> (Roth) Sant. & Wagh	WV	Str-A	6012
Capparaceae			
<i>Capparis brevispina</i> DC.	WV	Str-A	6110
<i>Capparis divaricata</i> Lam.	WV	Str-A	6138
<i>Capparis sepiaria</i> L. var. <i>sepiaria</i>	WV	Str-A	6057
<i>Capparis sepiaria</i> L. var. <i>retusella</i> Thwaites	WV	Str-A	6204
<i>Capparis shevaroyensis</i> Sun.-Ragh.	WV	Str-A	6321
<i>Capparis zeylanica</i> L.	WV	Str-A	6059
<i>Maerua oblongifolia</i> (Forsk.) A.Rich.	WV	Str-UA	6050
Celastraceae			
<i>Celastrus paniculatus</i> Willd.	WV	Str-UA	6014
<i>Loeseneriella obtusifolia</i> (Roxb.) A.C. Smith	WV	Str-UA	6137
<i>Maytenus heyneana</i> (Roth) Raju & Babu	WV	Str-A	6195
<i>Maytenus royleanus</i> (Wallich ex M. Lawson) M.A. Rau	WV	Str-A	6013
<i>Reissantia indica</i> (Willd.) Halle	WV	Str-UA	6286
<i>Salacia chinensis</i> L.	WV	Str-UA	6106

LISTS OF SPECIES

Species / Family	Category	Climbing mode	Voucher No.
<i>Salacia oblonga</i> Wall. ex Wight & Arn.	WV	Str-UA	6273
<b>Combretaceae</b>			
<i>Combretum acuminatum</i> Lam.	WV	TC	6055
<i>Combretum albidum</i> G. Don	WV	ST	6024
<b>Convolvulaceae</b>			
<i>Argyreia cuneata</i> (Willd.) Ker-Gawl.	WV	ST	6328
<i>Argyreia elliptica</i> (Roth) Choisy	WV	ST	6015
<i>Argyreia involucrata</i> Clarke	WV	ST	6321
<i>Argyreia kleiniana</i> (Roem. & Schultes) Raizada	WV	ST	6319
<i>Argyreia pilosa</i> Arn.	WV	ST	6032
<i>Argyreia sericea</i> Dalz.	WV	ST	6149
<i>Ipomoea asarifolia</i> (Desr.) Roem. & Schultes	HV	ST	6034
<i>Ipomoea complanata</i> L.	WV	ST	6474
<i>Ipomoea eriocarpa</i> R. Br.	WV	ST	6257
<i>Ipomoea quamoclit</i> L.	HV	ST	6332
<i>Ipomoea staphylina</i> Roem & Schultes	WV	ST	6149
<i>Merremia vitifolia</i> (Burm. f.) Hall. f.	HV	ST	6276
<i>Rivea hypocrateriformis</i> (Desr.) Choisy	WV	ST	6274
<b>Cucurbitaceae</b>			
<i>Coccinia grandis</i> (L.) J. Voigt	WV	TC	6146
<i>Gymnopetalum cochinchinense</i> Kurz	WV	TC	6323
<i>Kedrostis courtallensis</i> (Arn.) Jeffrey	WV	TC	6264
<i>Trichosanthes anaimalaiensis</i> Bedd.	WV	TC	6343
<b>Dioscoreaceae</b>			
<i>Dioscorea oppositifolia</i> L.	HV	ST	6064
<i>Dioscorea pentaphylla</i> L.	HV	ST	6068
<i>Dioscorea tomentosa</i> J. Koenig ex Sprengel	HV	ST	6066
<b>Elaeagnaceae</b>			
<i>Elaeagnus indica</i> Servettaz	WV	Str-UA	6209
<b>Euphorbiaceae</b>			
<i>Phyllanthus reticulatus</i> Poir.	WV	Str-UA	6426
<i>Tragia involucrata</i> L.	WV	ST	6047
<i>Tragia pluknetii</i> R. Smith	HV	ST	6262
<b>Gnetaceae</b>			
<i>Gnetum ula</i> Brongn.	WV	ST	6188
<b>Liliaceae</b>			
<i>Asparagus racemosus</i> Willd.	HV	Str-A	6222
<b>Linaceae</b>			
<i>Hugonia mystax</i> L.	WV	HC	6031
<b>Malpighiaceae</b>			
<i>Hiptage benghalensis</i> (L.) Kurz	WV	Str-UA	6074
<b>Menispermaceae</b>			
<i>Anamirta cocculus</i> (L.) Wight & Arn.	WV	ST	6253
<i>Cissampelos pareira</i> L. var. <i>hirsuta</i> (DC.) Forman	HV	ST	6069
<i>Cocculus hirsutus</i> (L.) Diels	WV	ST	6135
<i>Cocculus pendulus</i> (Forst.) Diels	WV	ST	6452
<i>Cyclea peltata</i> (Lam.) Hook.f. & Thoms.	HV	ST	6318
<i>Diploclisia glaucescens</i> (Blume) Diels	WV	ST	6188
<i>Pachygone ovata</i> (Poir.) Miers ex Hook.	WV	ST	6035


## LISTS OF SPECIES

Species / Family	Category	Climbing mode	Voucher No.
<i>Stephania japonica</i> (Thunb.) Miers	HV	ST	6070
<i>Tinospora cordifolia</i> (Willd.) Hook.f. & Thoms.	WV	ST	6033
<b>Mimosaceae</b>			
<i>Acacia caesia</i> (L.) Willd.	WV	Str-A	6026
<i>Acacia canescens</i> Grah.	WV	Str-A	6272
<i>Acacia intsia</i> Willd. var. <i>intsia</i>	WV	Str-A	6229
<i>Acacia pennata</i> (L.) Willd	WV	Str-A	6071
<i>Acacia sinuata</i> (Lour.) Merr.	WV	Str-A	6192
<i>Acacia torta</i> (Roxb.) Craib	WV	Str-A	6271
<i>Entada pursaetha</i> DC.	WV	Str-UA	6197
<i>Mimosa intsia</i> L.	WV	Str-A	6327
<b>Moraceae</b>			
<i>Plecospermum spinosum</i> Trecul.	WV	Str-A	6017
<b>Myrsinaceae</b>			
<i>Embelia basaal</i> (Roem. ex Schultes) A. DC.	WV	Str-UA	6179
<i>Embelia ribes</i> Burm.f.	WV	Str-UA	6192
<b>Nyctaginaceae</b>			
<i>Pisonia aculeata</i> L.	WV	Str-A	6226
<b>Oleaceae</b>			
<i>Jasminum angustifolium</i> (L.) Willd.	WV	ST	6004
<i>Jasminum auriculatum</i> Vahl	WV	ST	6022
<i>Jasminum azoricum</i> L. var. <i>azoricum</i>	WV	ST	6195
<i>Jasminum cuspidatum</i> Rottl.	WV	ST	6009
<i>Jasminum malabaricum</i> Wight	WV	ST	6073
<i>Jasminum multiflorum</i> (Burm. f.) Andr.	WV	ST	6188
<i>Jasminum sessiliflorum</i> Vahl	WV	ST	6001
<i>Jasminum trichotomum</i> Heyne ex Roth	WV	ST	6437
<b>Opiliaceae</b>			
<i>Cansjera rheedii</i> J. Gmelin	WV	Str-A	6021
<b>Papilionaceae</b>			
<i>Abrus precatorius</i> L.	WV	ST	6079
<i>Butea parviflora</i> Roxb.	WV	Str-UA	6308
<i>Canavalia virosa</i> (Roxb.) Wight & Arn.	WV	ST	6228
<i>Dalbergia congesta</i> Graham ex Wight & Arn.	WV	ST	6029
<i>Dalbergia rubiginosa</i> Roxb.	WV	ST	6045
<i>Derris scandens</i> (Roxb.) Benth.	WV	ST	6042
<i>Dolichos trilobus</i> L.	HV	ST	6086
<i>Galactia villosa</i> Wight & Arn.	HV	ST	6126
<i>Mucuna atropurpurea</i> DC.	WV	ST	6236
<i>Mucuna monosperma</i> DC. ex Wight	WV	ST	6184
<i>Mucuna pruriens</i> (L.) DC.	WV	ST	6023
<i>Neonotonia wightii</i> (Wight & Arn.) Lackey	WV	ST	6165
<i>Pseudarthria viscosa</i> (L.) Wight & Arn.	HV	ST	6039
<b>Passifloraceae</b>			
<i>Passiflora edulis</i> Sims	HV	TC	6267
<i>Passiflora foetida</i> L.	HV	TC	6421
<i>Passiflora subpeltata</i> Ortega	HV	TC	6077
<b>Piperaceae</b>			
<i>Piper nigrum</i> L.	WV	RC	6273
<b>Ranunculaceae</b>			
<i>Clematis gouriana</i> Roxb. ex DC.	WV	Str-UA	6124
<i>Naravelia zeylanica</i> (L.) DC.	WV	TC	6019
<b>Rhamnaceae</b>			

## LISTS OF SPECIES

Species / Family	Category	Climbing mode	Voucher No.
<i>Sageretia filiformis</i> (Schultes) Don	WV	Str-A	6007
<i>Scutia myrtina</i> (Burm.f.) Kurz	WV	Str-A	6008
<i>Ventilago goughii</i> Gamble	WV	ST	6325
<i>Ventilago maderaspatana</i> Gaertner	WV	ST	6003
<i>Ziziphus horrida</i> Roth	WV	Str-A	6028
<i>Ziziphus oenoplia</i> (L.) Miller	WV	Str-A	6005
<i>Ziziphus rugosa</i> Lam.	WV	Str-A	6228
Rosaceae			
<i>Rubus ellipticus</i> Smith	WV	Str-A	6214
<i>Rubus niveus</i> Thunb.	WV	Str-A	6329
Rubiaceae			
<i>Morinda umbellata</i> L.	WV	ST	6040
<i>Mussaenda hirsutissima</i> (Hook.f.) Hutch. ex Gamble	WV	ST	6224
<i>Pseudaida speciosa</i> (Bedd.) Tirveng.	WV	Str-UA	6178
<i>Rubia cordifolia</i> L.	HV	Str-UA	6156
Rutaceae			
<i>Paramignya beddomei</i> Tanaka	WV	Str-A	6142
<i>Toddalia asiatica</i> (L.) Lam.	WV	Str-A	6011
<i>Zanthoxylum ovalifolium</i> Wight	WV	Str-A	6176
<i>Zanthoxylum tetraspermum</i> Wight & Arn.	WV	Str-A	6185
Sapindaceae			
<i>Cardiospermum canescens</i> Wall.	HV	TC	6239
<i>Cardiospermum halicacabum</i> L. var. <i>luridum</i> (Blume) Adelb.	HV	TC	6094
Smilacaceae			
<i>Smilax zeylanica</i> L.	HV	TC	6157
Solanaceae			
<i>Solanum seaforthianum</i> Andr.	HV	ST	6159
Tiliaceae			
<i>Grewia flavescens</i> Juss.	WV	Str-UA	6075
<i>Grewia heterotricha</i> Mast.	WV	Str-UA	6269
<i>Grewia obtusa</i> Wall.	WV	Str-UA	6443
<i>Grewia oppositifolia</i> Buch-Ham.	WV	Str-UA	6148
<i>Grewia rhamnifolia</i> Heyne ex Roth	WV	Str-UA	6016
Verbenaceae			
<i>Lantana camara</i> L.	WV	Str-A	6124
<i>Premna coriacea</i> Clarke	WV	Str-UA	6176
<i>Premna corymbosa</i> (Burm.f.) Rottler & Willd.	WV	Str-UA	6002
<i>Premna flavescens</i> Ham.	WV	Str-UA	6234
<i>Premna villosa</i> Clark	WV	Str-UA	6204
<i>Premna wightiana</i> Schauer	WV	Str-UA	6324
Vitaceae			
<i>Ampelocissus araneosa</i> (Dalz. & Gibbs.) Planch.	WV	TC	6327
<i>Ampelocissus arnottiana</i> Planch.	WV	TC	6109
<i>Ampelocissus tomentosa</i> (Heyne ex Roth) Planch.	WV	TC	6090
<i>Cayratia japonica</i> (Thunb.) Gagnep.	WV	TC	6172
<i>Cayratia pedata</i> (Lour.) A.L. Juss. ex Gagnep.	WV	TC	6146
<i>Cayratia roxburghii</i> (Wight & Arn.) Gagnep.	WV	TC	6166
<i>Cissus gigantea</i> (Bedd.) Planch.	WV	TC	6251
<i>Cissus glyptocarpa</i> (Thw.) Planch.	WV	TC	6174
<i>Cissus heyneana</i> (Wall. ex Lawson) Planch.	WV	TC	6149
<i>Cissus quadrangularis</i> L.	WV	TC	6128
<i>Cissus vitiginea</i> L.	WV	TC	6078

LISTS OF SPECIES



**Figure 2.** a) Overview of Chitteri hills; b) Inner stand view of Chitteri hills; c) Overview of Kolli hills; d) Inner stand view of Kolli hills; e) Overview of Pachaimalais; f) Overview of Shervarayans g) Inner stand view of Shervarayans; h) Overview of Kalrayans.

LISTS OF SPECIES



**Figure 3.** a) Inner stand view of Kalrayans; b) Overview of Sathyamangalam forests; c) Stand view of Sathyamangalam forests; d) *Abrus precatorius* (Papilionaceae); e) *Acacia caesia* (Mimosaceae); f) *Acacia intsia* var. *intsia* (Mimosaceae); g) *Acacia pennata* (Mimosaceae); h) *Acacia sinuata* (Mimosaceae).

LISTS OF SPECIES



**Figure 4.** a) *Acacia torta* (Mimosaceae); b) *Aganosma cymosa* (Apocynaceae); c) *Argyreia cuneata* (Convolvulaceae); d) *Argyreia kleiniana* (Convolvulaceae); e) *Argyreia pilosa* (Convolvulaceae); f) *Argyreia involucrata* (Convolvulaceae); g) *Aristolochia indica* (Aristolochiaceae); h) *Aristolochia tagala* (Aristolochiaceae); i) *Caesalpinia crista* (Caesalpiniaceae).

LISTS OF SPECIES



**Figure 5.** a) *Canavalia virosa* (Papilionaceae); b) *Cansjera rheedii* (Opiliaceae); c) *Capparis shevaroyensis* (Capparaceae); d) *Capparis divaricata* (Capparaceae); e) *Capparis sepiaria* (Capparaceae); f) *Capparis zeylanica* (Capparaceae); g) *Cardiospermum halicacabum* var. *luridum* (Sapindaceae); h) *Carissa carandas* (Apocynaceae)

LISTS OF SPECIES




**Figure 6.** a) *Carissa paucinervia* (Apocynaceae); b) *Carissa salicina* (Apocynaceae); c) *Carissa spinarum* (Apocynaceae); d) *Cayratia pedata* (Vitaceae); e) *Cissus heyneana* (Vitaceae); f) *Cissus quadrangularis* (Vitaceae); g) *Cissus vitiginea* (Vitaceae); h) *Clematis gouriana* (Ranunculaceae); g) *Coccinia grandis* (Cucurbitaceae).

LISTS OF SPECIES



**Figure 7.** a) *Cocculus hirsutus* (Menispermaceae); b) *Combretum albidum* (Combretaceae); c) *Cosmostigma racemosum* (Asclepiadaceae); d) *Decalepis hamiltonii* (Asclepiadaceae); e) *Derris scandens* (Papilionaceae); f) *Diploclisia glaucescens* (Menispermaceae); g) *Embelia basal* (Myrsinaceae); h) *Entada pursaetha* (Mimosaceae); i) *Gnetum ula* (Gnetaceae).

LISTS OF SPECIES




**Figure 8.** a) *Grewia flavescentia* (Tiliaceae); b) *Grewia rhamnifolia* (Tiliaceae); c) *Gymnema sylvestre* (Asclepiadaceae); d) *Hemidesmus indicus* var. *indicus* (Asclepiadaceae); e) *Hiptage benghalensis* (Malpighiaceae); f) *Hugonia mystax* (Linaceae); g) *Ichnocarpus frutescens* (Apocynaceae); h) *Ipomoea asarifolia* (Convolvulaceae).

LISTS OF SPECIES



**Figure 9.** a) *Ipomoeacompanulata* (Convolvulaceae); b) *Ipomoeastaphylina* (Convolvulaceae); c) *Jasminum angustifolium* (Oleaceae); d) *Jasminumauriculatum* (Oleaceae); e) *Jasminumazoricum* var. *azoricum* (Oleaceae); f) *Jasminummalabaricum* (Oleaceae); g) *Jasminumsessiliflorum* (Oleaceae); h) *Lantana camara* (Verbenaceae).

LISTS OF SPECIES



**Figure 10.** a) *Loeseneriella obtusifolia* (Celastraceae); b) *Maerua oblongifolia* (Capparaceae); c) *Marsdenia brunoniana* (Asclepiadaceae); d) *Marsdenia tenacissima* (Asclepiadaceae); e) *Maytenus heyneana* (Celastraceae); f) *Mimosa intsia* (Mimosaceae); g) *Naravelia zeylanica* (Ranunculaceae); h) *Neonotonia wightii* (Papilionaceae).

LISTS OF SPECIES



**Figure 11.** a) *Pachygone ovata* (Menispermaceae); b) *Piper nigrum* (Piperaceae); c) *Plecospermum spinosum* (Moraceae); d) *Premna villosa* (Verbenaceae); e) *Pseudaida speciosa* (Rubiaceae); f) *Pterolobium hexapetalum* (Caesalpiniaceae); g) *Rhaphidophora laciniata* (Araceae); h) *Rubus ellipticus* (Rosaceae); i) *Rubus niveus* (Rosaceae).

LISTS OF SPECIES



**Figure 12.** a) *Sageretia filiformis* (Rhamnaceae); b) *Sarcostemma acidum* (Asclepiadaceae); c) *Scutia myrtina* (Rhamnaceae); d) *Secamone emetica* (Asclepiadaceae); e) *Smilax zeylanica* (Smilacaceae); f) *Solanum seaforthianum* (Solanaceae); g) *Toddalia asiatica* (Rutaceae); h) *Uvaria narum* (Annonaceae).

## LISTS OF SPECIES



**Figure 13.** a) *Ventilago maderaspatana* (Rhamnaceae); b) *Wattakaka volubilis* (Asclepiadaceae); c) *Zanthoxylum ovalifolium* (Rutaceae); d) *Zanthoxylum tetraspermum* (Rutaceae); e) *Ziziphus oenoplia* (Rhamnaceae)

### Acknowledgements

We thank the Department of Biotechnology (DBT), New Delhi for financial support through a project (BT/PR6603/NDB/51/089/2005) and Tamil Nadu forest department – both the Headquarters and Divisional offices of southern Eastern Ghats region for site permission.

## LISTS OF SPECIES

### Literature cited

- Acevedo-Rodriguez, P. 2005. Vines and climbing plants of Puerto Rico and the Virgin Islands Contributions from the United States National Herbarium 51: 1-483.
- Areendran, G. and P. Rao. 2006. Vegetation types of the southern Eastern Ghats A Remote Sensing perspective. New Delhi. World Wide Fund for Nature-India. 1-92.
- Burnham, R. J. 2002. Dominance, diversity and distribution of lianas in Yasuni. Ecuador: Who is on top? Journal of Tropical Ecology 18: 845-864.
- Gamble, J. S. and C. E. C. Fischer. 1915-1935. Flora of the Presidency of Madras. London. Adlard & Son, Limited. vols. I to III.
- Gentry, A. H. 1991. The distribution and evaluation of climbing plants; p. 3-49 In F.E. Putz and H. A. Mooney (ed.), The Biology of vines. Cambridge, Cambridge University Press.
- Gentry, A. H. and C. Dodson. 1987. Contribution of non-trees to species richness of a tropical rain forest. Biotropica 19: 149-155.
- Grandos, J. and C. Körner. 2002. In deep shade, elevated CO<sub>2</sub> increases the vigour of tropical climbing plants. Global Change Biology 8: 1109-1117.
- Hooker, J. D. 1872-1897. The Flora of British India. London. L. Reeve & Co. part 1-7.
- Ibarra-Manríquez, G. and M. Martínez-Ramos. 2002. Landscape variation of liana communities in a Neotropical rain forest. Plant Ecology 160: 91-112.
- Jongkind, C. C. H. and W. D. Hawthorne. 2005. A botanical synopsis of the lianes and other forest climbers; p. 19-39 In Bongers, F., M. P. E. Parren and D. Traore (ed.), Forest climbing plants of West Africa diversity, ecology and management. Cambridge. CABI Publishing.
- Laurance, W. F., D. Perez-Salicrup and P. Delamonica et al. 2001. Rain forest fragmentation and the structure of Amazonian liana communities. Ecology 82: 105-116.
- Londre, R. A. and S. A. Schnitzer. 2006. The distribution of lianas and their change in abundance in temperate forests over the past 45 years. Ecology 87: 2973-2978.
- Matthew, K. M. 1991. An excursion flora of central Tamil Nadu. India. New Delhi. Oxford & IBH Publishing Co. Pvt. Ltd.
- Malhi, Y. and J. Wright. 2004. Spatial patterns and recent trends in the climate of tropical rainforest regions. Philosophical Transactions of the Royal Society: Biological Sciences 359: 311-329.
- Mascaro, J. S., S. A. Schnitzer, and W. P. Carson. 2004. Liana diversity, abundance and mortality in a tropical wet forest in Costa Rica. Forest Ecology and Management 190: 3-14.
- Muthuramkumar, S. and N. Parthasarathy. 2000. Alpha diversity of lianas in a tropical evergreen forest in the Anamalais, Western Ghats, India. Diversity and Distributions 6: 1-14.
- Padaki, A. and N. Parthasarathy. 2000. Abundance and distribution of lianas in tropical lowland evergreen forest of Agumbe, central Western Ghats, India. Tropical Ecology 41: 143-154.
- Parthasarathy, N., S. Muthuramkumar, and M. S. Reddy. 2004. Patterns of liana diversity in tropical evergreen forests of peninsular India. Forest Ecology and Management 190: 15-31.
- Restom, T. G. and D. C. Nepstad. 2001. Contribution of vines to the evapotranspiration of a secondary forest in eastern Amazon. Plant and Soil 236: 155-163.
- Rice, K., N. Brokaw, and J. Thompson. 2004. Liana abundance in a Puerto Rican forest. Forest Ecology and Management 190: 33-41.
- Schnitzer, S. A. 2005. A mechanistic explanation for global patterns of liana abundance and distribution. American Naturalist 166: 262-276.
- Schnitzer, S. A. and F. Bongers. 2002. The ecology of lianas and their role in forests. Trends in Ecology and Evolution 17: 223-230.
- Schnitzer, S. A. and W. P. Carson. 2001. Tree fall gaps and the maintenance of species diversity in a tropical forest. Ecology 82: 913-919.
- Schnitzer, S. A., J. W. Dalling, and W. Carson. 2000. The importance of lianas on tree regeneration in tropical forest canopy gaps: evidence for an alternative pathway of gap-phase regeneration. Journal of Ecology 88: 655-666.
- Swaine, M. D. and J. Grace. 2007. Liana may be favoured by low rainfall: evidence from Ghana. Plant Ecology 192: 271-276.
- Toledo-Aceves, T. and M. D. Swaine. 2007. Effects of the species of climbers on the performance of *Ceiba pentandra* seedling in gaps in a tropical forest in Ghana. Journal of Tropical Ecology 23: 45-52.
- Zotz, G., N. Cueni and C. Körner. 2006. *In situ* growth stimulation of a temperate zone liana (*Hedera helix*) in elevated CO<sub>2</sub>. Functional Ecology 20: 763-769.

Received September 2008

Accepted January 2009

Published online March 2009