

# Flavonoidal Constituents, Antioxidant, Antimicrobial, and Cytotoxic Activities of *Dipterygium glaucum* Grown in Kingdom of Saudi Arabia

Usama Shaheen<sup>1,2</sup>, Nagwa Abdelkader Shoeib<sup>1,3</sup>, Abeer Temraz<sup>1,2</sup>, Mohamed I. S. Abdelhady<sup>1,4</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, Umm Al-Qura University, Makkah Al Mukarramah 21955, Saudi Arabia, <sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Cairo, <sup>3</sup>Department of Pharmacognosy, Faculty of Pharmacy, Tanta University, Tanta 31527, <sup>4</sup>Department of Pharmacognosy, Faculty of Pharmacy, Helwan University, Cairo, Egypt

Submitted: 21-01-2016

Revised: 11-07-2016

Published: 11-10-2017

## ABSTRACT


**Background:** *Dipterygium glaucum* Decne. herb is one of the common traditional plants with multiple medicinal uses. **Objective:** To isolate the major constituents and to investigate the antioxidant, antimicrobial, and cytotoxic activities of this herb. **Materials and Methods:** Methanolic extract of *D. glaucum* herb was fractionated using *n*-hexane, dichloromethane, and *n*-butanol. Butanol fraction was chromatographed using column chromatography and preparative thin layer chromatography to isolate the major constituents. Isolated compounds were elucidated by means of spectroscopic methods, including 1D, 2D NMR (<sup>1</sup>H, <sup>13</sup>C, DEPT, COSY, HSQC, HMBC, NEOSY) and MS analysis. Total phenolic content using Folin–Ciocalteu reagent and antioxidant activity using 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay of the total methanolic extract were evaluated. Cytotoxic potential of both methanolic extract and butanol fraction was tested using a crystal violet viability assay. Antimicrobial activities of both extracts were investigated using diffusion agar technique. **Results:** Apigenin 6, 8-di-C-glucopyranoside (vicenin-2), quercetin-3'-O-methyl-3-O-glucopyranoside, quercetin-3'-O-methyl-3-O-galactopyranoside, quercetin-3-O-β-D-glucopyranoside, and quercetin-3-O-β-D-galactopyranoside were isolated and elucidated. Total phenolic content was (83.89 mg gallic acid equivalent/g extract). The EC<sub>50</sub> value of scavenging DPPH radical was 152.0 ± 2 μg/mL. Butanol fraction showed the highest cytotoxic activity against cervical and breast carcinoma cells (IC<sub>50</sub> 3.6 and 6.1 μg/mL, respectively). Both methanolic extract and butanol fraction showed wide spectrum antimicrobial activity against Gram-positive and Gram-negative bacteria and some fungi. The highest activity was from methanolic extract against *Enterococcus faecalis* (83.25%) and against *Candida tropicalis* (77.03%) as compared to reference antibiotics. **Conclusion:** Data obtained from this study demonstrate that *D. glaucum* possesses significant antioxidant, cytotoxic, and antimicrobial activities which could be ascribed to its flavonoidal content.

**Key words:** Antimicrobial, antioxidant, cytotoxic, *Dipterygium glaucum*, flavonoids

## SUMMARY

- Dipterygium glaucum* Decne. herb is one of the common traditional plants with multiple medicinal uses in KSA

- Five flavonoidal glycosides were isolated and elucidated
- This study demonstrated that *D. glaucum* possesses significant antioxidant, cytotoxic, and antimicrobial activities.



**Abbreviations used:** KSA: Kingdom of Saudi Arabia; TLC: Thin Layer Chromatography; DPPH: 2,2'-diphenyl-1-picrylhydrazyl; EC<sub>50</sub>: Half maximal effective concentration; IC<sub>50</sub>: Half maximal inhibitory concentration; DMSO: dimethyl sulfoxide; NMR: Nuclear Magnetic Resonance; ESIMS: Electrospray ionization mass spectrometry; MeOH: Methyl alcohol.

## Correspondence:

Dr. Nagwa Abdelkader Shoeib,  
Department of Pharmacognosy,  
Faculty of Pharmacy, Umm Al-Qura University,  
Makkah Al Mukarramah 21955,  
Saudi Arabia.  
E-mail: nashoeib@uqu.edu.sa; na2001eg@yahoo.  
com  
**DOI:** 10.4103/pm.pm\_44\_16

## Access this article online

**Website:** www.phcog.com

## Quick Response Code:



## INTRODUCTION

The use of medicinal plants as an essential component for traditional healthcare systems is the oldest and the most varied of all therapeutic systems. The World Health Organization estimated that 80% of the world population currently uses herbal medicine for some aspects of primary health care.<sup>[1]</sup>

In Saudi Arabia, traditional medicine is based on herbal remedies and spiritual healing. It is not possible to find a city or village where traditional herbal medicines are not used.<sup>[2]</sup>

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

**Cite this article as:** Shaheen U, Shoeib NA, Temraz A, Abdelhady MI. Flavonoidal constituents, antioxidant, antimicrobial, and cytotoxic activities of *Dipterygium glaucum* grown in Kingdom of Saudi Arabia. Phcog Mag 2017;13:S484-8.

*Dipterygium glaucum* Decne. herb (*Capparidaceae*) is a branched under shrub plant. It is a monotypic genus with one species, widely distributed in Saudi Arabia, Egypt, Sudan, and Pakistan.<sup>[3,4]</sup>

It is one of the common plants that are used by the Bedouins in Saudi Arabia. It has multiple medicinal uses; it is popular for the treatment of miss-breathing troubles as trachea dilating agent.<sup>[5]</sup>

It is also used traditionally in Pakistan as decoction and infusion for the treatment of jaundice, blood purifier, psoriasis, and ringworm infestation and as an antiasthma drug.<sup>[4,6]</sup>

Phytochemical studies on *D. glaucum* revealed the presence of alkaloids, cardiac glycoside, bound anthraquinones, saponins, terpenoids, and sterols.<sup>[3,7]</sup>

Previous biological studies on the plant revealed that ethanolic plant extract has antioxidant and antispasmodic activities while *n*-hexane, ethyl acetate, and butanol extracts have phytotoxic activity.<sup>[8]</sup>

Oxidative stress is the leading cause for the development of many recent diseases such as arteriosclerosis, inflammation, autoimmune diseases, cancer, and aging. Antioxidants deactivate and stabilize reactive oxygen species and other oxidants before they affect biological cells.<sup>[9]</sup>

Food and herbal plants contain many classes of antioxidant constituents such as flavonoids, phenolics, proanthocyanidins, and tannins. These constituents have free radical scavenging abilities and contribute to the protection of human body from diseases.<sup>[10]</sup>

Microbial resistance is considered serious threat to public health as infections caused by resistant microorganisms often fail to respond to the standard treatment. Natural products, in the form of pure compounds or standardized extracts, offer excellent chances for new drug discovery due to the chemical diversity of naturally derived compounds.<sup>[11]</sup>

As phytochemicals are important compounds that play an essential role for restoring normal functioning of human body and amelioration of diseases, this study was designed to isolate the major phytochemical constituents and to investigate the antioxidant, antimicrobial, and cytotoxic activities of the wild *D. glaucum* collected from Makkah deserts, Kingdom of Saudi Arabia (KSA).

## MATERIALS AND METHODS

### Plant material

The aerial parts of *D. glaucum* herb were collected in February 2013 from Makkah, KSA, and was identified by Mohamed Meelad Late, Professor of Plant Taxonomy, Faculty of Applied Sciences, Umm Al-Qura University. A voucher specimen (DG 013) is deposited at Department of Pharmacognosy, College of Pharmacy, Umm Al-Qura University.

### Extraction and isolation

The dried powdered aerial parts of *D. glaucum* (1.5 kg) were exhaustively extracted with methanol by maceration at room temperature (3 × 6 L). Methanolic extract was evaporated under reduced pressure using rotary evaporator (Buchi Co., Switzerland) till dryness to give 368.5 g dry extract. Dry extract (200 g) was suspended in distilled water (350 mL) and was fractionated using *n*-hexane, dichloromethane, and *n*-butanol (3 × 500 mL). The fractions were collected and evaporated till dryness to give the following weights, 25 g, 12 g, and 78 g. Butanol fraction (78 g) was chromatographed on silica gel column (Sigma-Aldrich, USA) (5 cm × 120 cm, flow rate 2.5 mL/min) and was eluted by chloroform/methanol mixtures of increasing polarities by the addition of methanol in 10% increments till 100% methanol. Fractions of 50 mL were collected, monitored by thin layer chromatography (TLC) (aluminum-coated silica gel 60

F<sub>254</sub>, Macherey-Nagel, Germany) using chloroform-methanol-water (80:20:2, v/v/v) and chloroform-methanol-water-acetic acid (60:32:7:1 v/v/v/v) as mobile phases, and spots were detected on the plates using methanol/sulfuric acid (20% v/v) spray reagent; then, similar fractions were grouped into 18 major fractions (A-R).

Fraction L (3.99 g) was chromatographed on Sephadex LH-20 column (Pharmacia, Uppsala, Sweden) (2.5 cm × 30 cm, flow rate 1.0 mL/min) using methanol as the eluent to give nine fractions. Fraction 3 was rechromatographed on Sephadex LH-20 column (1.5 cm × 20 cm, flow rate 0.5 mL/min) using methanol as the eluent to give three subfractions. Subfraction 1 was subjected to silica gel column (1.0 cm × 15 cm, flow rate 0.5 mL/min) using chloroform-methanol-water (8:20:2 v/v/v) as starting eluent, then increasing the polarity to 70:30:3 v/v/v which affords compounds 4 and 5.

Fraction M (230 mg) was chromatographed on Sephadex LH-20 column (2.5 cm × 30 cm, flow rate 1.0 mL/min) using methanol as eluent, which gave four subfractions. Subfraction number 4 was subjected to preparative TLC using chloroform-methanol-water-acetic acid (60:32:7:1 v/v/v/v), and bands at R<sub>f</sub> 0.5 and 0.55 were scratched, extracted with methanol, and purified using small Sephadex column; this affords compounds 2 and 3.

Fraction R (600 mg) was applied to Sephadex LH-20 column (2.5 cm × 30 cm, flow rate 1.0 mL/min) and was eluted using methanol, this afforded six subfractions. Subfraction number 5 was separated by silica gel column (1.5 cm × 15 cm, flow rate 0.5 mL/min) using ethyl acetate-methanol mixture (80:20 v/v) as eluent, which gave seven fractions. Fraction number 5 was subjected to preparative TLC using chloroform-methanol-water-acetic acid (60:32:7:1 v/v/v/v) as mobile phase, band at R<sub>f</sub> 0.3 was scratched, extracted with methanol, and purified using sephadex column to give compound 1.

### Compound 1 (6,8-Di-C-glucosylapigenin or vicanin-2)

Yellow amorphous powder; UV λ<sub>max</sub> MeOH 232, 282, 344 nm. ESI MS (m/z): 593 [M-H]<sup>-</sup>, 503, 473, 383, and 353. <sup>1</sup>H nuclear magnetic resonance (NMR) (500 MHz, dimethyl sulfoxide [DMSO]-d<sub>6</sub>) δ: 8.03 (2H, d, J = 8.4 Hz, H-2', 6'), 6.93 (2H, d, J = 8.4 Hz, H-3', 5'), 6.82 (1H, s, H-3); 6-C-β-Glc: 4.81 (1H, d, J = 9.7 Hz, H-1''); 8-C-β-Glc:




Figure 1: Structure of compounds 1-5 isolated from *Dipterygium glaucum*

4.76 (<sup>1</sup>H, d, *J* = 9.8 Hz, H-1''). <sup>13</sup>C NMR (125MHz, DMSO-d<sub>6</sub>) δ<sub>c</sub> 182.8 (C-4), 164.5 (C-2), 161.7 (C-3), 161.69 (C-4'), 159.0 (C-5), 155.5 (C-9), 129.5 (C-6'), 129.5 (C-2'), 122.0 (C-1'), 116.3 (C-5'), 116.3 (C-3'), 107.9 (C-6), 105.7 (C-10), 104.3 (C-8), 103.0 (C-3), 82.3 (C-5''), 82.3 (C-5'''), 79.3 (C-3''), 78.3 (C-3'''), 74.5 (C-1''), 73.8 (C-1'''), 72.4 (C-2''), 71.4 (C-2'''), 71.0 (C-4''), 69.5 (C-4'''), 61.7 (C-6''), 60.2 (C-6''').

### Compound 2 (quercetin-3'-O-methyl-3-O-glucopyranoside)

It was obtained as a light yellow amorphous powder; UV λ<sub>max</sub> MeOH: 258, 305, 361 nm. ESI-MS: m/z 479.0 [M + H]<sup>+</sup>. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ: 7.55 (<sup>1</sup>H, brs, H-2'), 7.50 (<sup>1</sup>H, d, *J* = 8.4 Hz, H-6'), 6.70 (<sup>1</sup>H, d, *J* = 7.9 Hz, H-5'), 6.37 (<sup>1</sup>H, d, *J* = 1.8 Hz, H-8), 6.17 (<sup>1</sup>H, d, *J* = 1.8 Hz, H-6), 3.80 (3H, s, OCH<sub>3</sub>-3'); 3-O-β-Glc: 5.32 (<sup>1</sup>H, d, *J* = 7.3 Hz, H-1'). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>) δ<sub>c</sub> 175.2 (C-4), 160.1 (C-5), 156.2 (C-2), 156.2 (C-9), 144.9 (C-3'), 133.1 (C-3), 121.8 (C-1'), 121.8 (C-6'), 116.4 (C-2'), 116.2 (C-5'), 103.3 (C-10), 101.5 (C-1''), 99.9 (C-6), 94.9 (C-8), 77.8 (C-3''), 77.1 (C-5''), 74.6 (C-2''), 70.4 (C-4''), 61.4 (C-6''), 56.03(OCH<sub>3</sub>).

### Compound 3 (quercetin-3'-O-methyl-3-O-galactopyranoside)

Yellow powder, UV λ<sub>max</sub> MeOH: 353 and 254 nm; ESI-MS: m/z 477.0 [M<sup>-</sup> H]<sup>-</sup>. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ: 7.54 (<sup>1</sup>H, brs, H-2'), 7.53 (<sup>1</sup>H, d, *J* = 8.4 Hz, H-6'), 6.73 (<sup>1</sup>H, d, *J* = 8.4 Hz, H-5'), 6.04 (<sup>1</sup>H, d, *J* = 2 Hz, H-8), 5.86 (<sup>1</sup>H, d, *J* = 2 Hz, H-6), 3.82 (3H, s, OCH<sub>3</sub>-3'); 3-O-β-Gla: 5.22 (<sup>1</sup>H, d, *J* = 7.8 Hz, H-1'). <sup>13</sup>C NMR (125MHz, DMSO-d<sub>6</sub>) δ<sub>c</sub> 176.3 (C-4), 161.2 (C-5), 157.3 (C-2), 157.3 (C-9), 146.0 (C-3'), 133.3 (C-3), 121.8 (C-1'), 121.8 (C-6'), 115.9 (C-2'), 115.8 (C-5'), 103.3 (C-10), 102.3 (C-1''), 99.9 (C-6), 94.9 (C-8), 74.0 (C-3''), 76.2 (C-5''), 71.7 (C-2''), 68.3 (C-4''), 60.5 (C-6''), 56.26 (OCH<sub>3</sub>).

### Compound 4 (quercetin 3-O-β-D-glucopyranoside)

Yellow amorphous powder; EI-MS m/z 463.08 [M<sup>-</sup> H]<sup>-</sup>; <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 500 MHz) δ: 7.54 (<sup>1</sup>H, d, *J* = 1.3, H-2'), 7.65 (<sup>1</sup>H, dd, *J* = 6.8, 1.3 Hz, H-6'), 6.80 (<sup>1</sup>H, d, *J* = 6.8 Hz, H-5'), 6.37 (<sup>1</sup>H, d, *J* = 2 Hz, H-8), 6.17 (<sup>1</sup>H, d, *J* = 2 Hz, H-6); 3-O-β-Glc: 5.45 (<sup>1</sup>H, d, *J* = 7.5 Hz, H-1'). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>) δ<sub>c</sub> 177.7 (C-4), 165.9 (C-7), 161.6 (C-5), 156.5 (C-2), 156.8 (C-9), 149.1 (C-4'), 145.4 (C-3'), 133.8 (C-3), 121.4 (C-1'), 122.3 (C-6'), 116.4 (C-2'), 115.6 (C-5'), 104.0 (C-10), 101.4 (C-1''), 99.4 (C-6), 94.1 (C-8), 77.9 (C-3''), 77.0 (C-5''), 74.5 (C-2''), 70.4 (C-4''), 61.4 (C-6'').

### Compound 5 (quercetin-3-O-β-D-galactopyranoside)

Yellow powder, UV λ<sub>max</sub> MeOH: 353 and 254 nm; EI-MS m/z 463.07 [M<sup>-</sup> H]<sup>-</sup>; <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 500 MHz) δ: 7.56 (<sup>1</sup>H, brs, H-2'), 7.65 (<sup>1</sup>H, d, *J* = 8.5 Hz, H-6'), 6.81 (<sup>1</sup>H, d, *J* = 8.5 Hz, H-5'), 6.38 (<sup>1</sup>H, d, *J* = 1.71 Hz, H-8), 6.17 (<sup>1</sup>H, d, *J* = 1.71 Hz, H-6); 3-O-β-Gla: 5.36 (<sup>1</sup>H, d, *J* = 7.5 Hz, H-1'). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>) δ<sub>c</sub> 177.7 (C-4), 165.9 (C-7), 161.6 (C-5), 156.5 (C-2), 156.8 (C-9), 149.1 (C-4'), 145.3 (C-3'), 133.8 (C-3), 121.4 (C-1'), 122.3 (C-6'), 116.3 (C-2'), 115.6 (C-5'), 103.9 (C-10), 102.4 (C-1''), 99.4 (C-6), 94.1 (C-8), 73.6 (C-3''), 76.2 (C-5''), 71.6 (C-2''), 68.3 (C-4''), 60.5 (C-6'') [Figure 1].

### Quantitative estimation of total phenolic content

Total phenolic content in the plant methanolic extract was determined by Folin-Ciocalteu method.<sup>[12]</sup> Calibration curve was constructed as




Figure 2: Standard calibration curve of gallic acid

follows: 1 mL of 30, 60, 90, 120, 150, and 180 µg/mL ethanolic gallic acid solutions was added to 5 mL Folin-Ciocalteu reagent (diluted 10-fold) and 4 mL sodium carbonate (75 g/L). The absorbance was measured after 30 min at 765 nm using a spectrophotometer (UV2550, Shimadzu, Japan) and the calibration curve was plotted. 0.2 mL methanolic extract (1 mg/mL) was mixed with the same reagents as described above, and the absorbance was measured. All determinations were done in triplicate.

The total phenolic content was calculated using the following linear equation based on the calibration curve of gallic acid [Figure 2].

$$A = 0.008X - 0.019.$$

where A is the absorbance and X is the amount of gallic acid.

The result was expressed as gallic acid equivalent (GAE)/g.

### In vitro determination of antioxidant activity

*In vitro* free radical scavenging activity of the methanolic extract was measured using 2,2'-diphenyl-1-picrylhydrazyl (DPPH) assay according to the published method.<sup>[13]</sup> Extract solution was prepared by dissolving 10 mg of dry extract in 10 mL methanol. The solution of DPPH (Sigma-Aldrich, KSA) in methanol ( $6 \times 10^{-5}$  M) was freshly prepared daily. Three milliliters of DPPH solution was mixed with 20, 40, 60, 80, 100, and 150 µl extract solution in 1 cm path length microcuvettes. The samples were kept in the dark at room temperature for 15 min, and then, the decrease in the absorbance was measured using the spectrophotometer. Blank sample containing the same amount of methanol and DPPH solution was prepared and measured daily. The experiment was carried out in triplicate. Radical scavenging activity was calculated by the following formula:

$$\% \text{ inhibition} = [(AB - AA)/AB] \times 100$$

where AB is the absorbance of DPPH (*t* = 0 min) and AA is the absorbance of plant extract solution + DPPH (*t* = 15 min).

### In vitro cytotoxic activity against human cell lines

The cytotoxicity of methanolic extract and butanol fraction against hepatocellular (HepG2), breast (MCF-7), pharynx (HEP-2), colon (HCT), and cervical (HeLa) carcinoma human cell lines was tested by a crystal violet viability assay,<sup>[14,15]</sup> and IC<sub>50</sub> values are determined and illustrated in Table 1.

### Antimicrobial activity

Diffusion agar technique was applied using trypticase soy agar (Difco) medium containing bacterial or fungal suspension of the test organisms.<sup>[16]</sup> Impregnated discs with plant extracts, DMSO as negative control and amphotericin B, ampicillin, gentamicin as standard

antibiotics, were placed on the surface of the cultured agar and were incubated at 35°C–37°C for 24–48 h in case of bacteria and at 25°C for 48 h in case of fungi. After incubation, zones of inhibition diameters were measured in millimeters and the results are shown in Table 2.

## RESULTS

Phytochemical investigation of *D. glaucum* led to the isolation of one C-flavone glycoside (compound 1) and four O-flavonol glycosides, compounds (2-5). They were elucidated by means of spectroscopic methods including 1D, 2D NMR (<sup>1</sup>H, <sup>13</sup>C, DEPT, COSY, HSQC, HMBC), MS analysis and by the comparison of the physical and spectral data with those published before as apigenin 6, 8-di-C-glucopyranoside (vicenin-2),<sup>[17]</sup> quercetin-3'-O-methyl-3-O-glucopyranoside,<sup>[18]</sup> quercetin-3'-O-methyl-3-O-galactopyranoside,<sup>[19]</sup> quercetin-3-O-β-D-glucopyranoside,<sup>[20]</sup> quercetin-3-O-β-D-galactopyranoside<sup>[21]</sup> [Figure 1].

### Quantitative estimation of total phenolic content

Phenolic content of the methanolic extract of the studied plant was 83.89 mg/g expressed as GAEs.

### Antioxidant activity

Methanolic extract of *D. glaucum* showed concentration-dependent scavenging effect on DPPH radical [Figure 3]. The EC<sub>50</sub> value of scavenging DPPH radical was found to be 152.0 ± 2 μg/mL.

### Cytotoxic activity

Methanolic extract and butanol fraction of *D. glaucum* exhibited cytotoxic activity against tested cell lines. Butanol fraction showed the highest cytotoxic activity against cervical and breast carcinoma cells (IC<sub>50</sub> 3.6 and 6.1 μg/mL, respectively).

### Antimicrobial activity

Methanolic extract and butanol fraction of *D. glaucum* showed comparable antibacterial activity against Gram-positive bacteria; the

**Table 1:** Results of cytotoxicity of methanolic extract and butanol fraction of *Dipterygium glaucum*

Extract	IC <sub>50</sub> (μg/ml)				
	HCT	HEP-2	HeLa	HepG2	MCF-7
Butanol fraction	35.7	24.6	3.6	22.3	6.1
Methanolic extract	18.6	14.4	24.7	28.3	35.2

highest activity was obvious from methanolic extract against *Enterococcus faecalis* (83.25%), while both extracts showed weak activity against tested Gram-negative bacteria. Both extracts showed moderate antifungal activity. Methanolic extract showed the highest activity against *Candida tropicalis* (77.03%) [Table 2].

## DISCUSSION


This study reported the isolation of C- and O-flavonoidal glycosides from *D. glaucum* for the first time.

During the present investigation of *D. glaucum*, the methanolic extract showed concentration-dependent antioxidant activity against DPPH radical by colored solution bleaching with EC<sub>50</sub> 152.0 ± 2 μg/mL. This inhibition percentage could be correlated positively to phenolic content of the plant.

*D. glaucum* butanol fraction demonstrated a promising anticancer activity against cervical (HeLa) and breast (MCF-7) carcinoma cells, and this could be attributed to flavonoidal content of the plants.

Flavonoids are naturally occurring plants secondary metabolites; they have positive impact on human health. They are highly effective scavengers against most oxidizing molecules as free radicals and singlet oxygen species are responsible for many diseases as cancer.<sup>[22]</sup>

Previous study was conducted to determine synergistic effect of vicenin-2 in combination with docetaxel in prostate cancer treatment. The authors reported that vicenin-2 is effective against



**Figure 3:** 1,1-diphenyl-2-picrylhydrazyl radical scavenging activity of *Dipterygium glaucum* methanolic extract

**Table 2:** Antimicrobial activities of methanolic extract and butanol fraction of *Dipterygium glaucum*

Tested organism	Sample		
	Butanol fraction (%)	Methanolic extract (%)	Standard
<b>Gram-positive</b>			<b>ampicillin</b>
<i>Streptococcus pneumonia</i> (RCB 010010)	14.8±0.34 (62.18)	15.3±0.34 (64.28)	23.8±0.2
<i>Enterococcus faecalis</i> (RCMB 010068)	16.4±0.19 (78.84)	16.9±0.62 (83.25)	20.3±0.3
<i>Staphylococcus aureus</i> (RCMB 010028)	13.2±0.67 (46.64)	15.4±0.38 (54.42)	28.3±0.1
<b>Gram-negative</b>			<b>gentamicin</b>
<i>Escherichia coli</i> (RCMB 010052)	9.6±0.34 (47.05)	11.4±0.44 (55.88)	20.4±0.6
<i>Pseudomonas aeruginosa</i> (RCMB 010043)	NA	NA	18.2±0.1
<b>Fungi</b>			<b>amphotericin B</b>
<i>Aspergillus fumigatus</i> (RCMB 02568)	12.6±0.23 (53.16)	14.5±0.35 (61.18)	23.7±0.1
<i>Candida tropicalis</i> (RCMB 05239)	15.8±0.51 (75.59)	16.1±0.38 (77.03)	20.9±0.3
<i>Cryptococcus neoformans</i> (RCMB 05642)	14.1±0.55 (64.38)	16.2±0.46 (73.97)	21.9±0.12

Results are expressed by mean zone of inhibition (in mm)±SD (potency %). RCMB: Regional Center for Mycology and Biotechnology; NA: No activity; SD: Standard deviation

prostate carcinoma progression when coadministered with docetaxel than either of the single agents in androgen-independent prostate cancer.<sup>[23]</sup>

Isorhamnetin is found in many plants and is a metabolite of quercetin. Lee *et al.*<sup>[24]</sup> demonstrated the anticancer effect of isorhamnetin on lung cancer cell culture and on mice diseased with lung cancer. *In vitro* study confirmed that isorhamnetin induces apoptosis, which is mediated by mitochondria-dependent caspase activation. *In vivo* test proved the anticancer effect of isorhamnetin; weights of tumors significantly reduced by injecting mice with isorhamnetin.<sup>[24]</sup>

In this study, the isolation of vicenin-2 (compound 1) together with isorhamnetin and quercetin glycosides (compounds 2-5) from butanol fraction suggested that flavonoids could be the major group of constituents that may be responsible for the anticancer activity.

Diseases resulting from microbial infection represent an important cause of morbidity and mortality among human and animals, particularly in developing countries. Therefore, searching for new antimicrobial drugs from medicinal plants has greatly increased in recent years, especially due to the constant rise of microorganisms resistant to conventional medications.

According to Alves *et al.*,<sup>[25]</sup> inhibition zones > 9 mm correspond to active antimicrobial samples. In the present study, *D. glaucum* methanolic extract and butanol fraction showed wide spectrum antimicrobial activity against Gram-positive and Gram-negative bacteria and fungi. The highest activity was from methanolic extract against *E. faecalis* (83.25%) and against *C. tropicalis* (77.03%) as compared to reference antibiotics.

In previous work,<sup>[3]</sup> it was reported that *D. glaucum* has no antimicrobial activity, and in the same study, they reported the absence of flavonoids in the plant extract. Herein, we report the isolation of flavonoidal glycosides which may be the responsible constituents for the reported activity and the slight increase in the activity of methanolic extract over butanol could be attributed to synergistic activity of flavonoidal glycosides with other secondary metabolites in the plant extract.

## CONCLUSION

The present finding showed that *D. glaucum* has antimicrobial, cytotoxic, and antioxidant activities, thus demonstrating its potential benefits in ameliorating many diseases. Further studies are required to identify the mechanisms, whereby *D. glaucum* exerts its beneficial effects.

## Financial support and sponsorship

The authors would like to thank the Deanship of Scientific Research and the Institute of Scientific Research and Revival of Islamic Heritage at Umm Al-Qura University (Project ID: 4331009) for the financial support.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Ghasemzadeh A, Jaafar HZ, Rahmat A. Phytochemical constituents and biological activities

- of different extracts of *Strobilanthes crispus* (L.) Bremek leaves grown in different locations of Malaysia. BMC Complement Altern Med 2015;15:422.
- Mahomoodally MF. Traditional medicines in Africa: An appraisal of ten potent African medicinal plants. Evid Based Complement Alternat Med 2013;2013:617459.
- Mehmood K, Mehmood S, Ramzan M, Arshad M, Yasmeen F. Biochemical and phytochemical analysis of *Dipterygium glaucum* collected from Cholistan desert. J Sci Res 2010;XXX:13-8.
- Ahmad S, Wariss HM, Alam K, Anjum S, Mukhtar M. Ethnobotanical studies of plant resources of Cholistan desert, Pakistan. Int J Sci Res 2014;3:1782-8.
- Moussa SA, Taia WK, Al-Ghamdy FG. Acclimation of *Dipterygium glaucum* Decne. Grown in the Western Coastal part of Saudi Arabia to different water supplies. Int J Res Chem Environ 2012;2:301-9.
- Rahman MA, Mossa JS, Al-Said MS, Al-Yahya MA. Medicinal plant diversity in the flora of Saudi Arabia 1: A report on seven plant families. Fitoterapia 2004;75:149-61.
- Abdel-Mogib M, Ezmirly ST, Basaif SA. Phytochemistry of *Dipterygium glaucum* and *Capparis decidua*. J Saudi Chem Soc 2000;4:103-8.
- Ahmed S, Ashraf M, Jabbar A, Janbaz KH, Khan MS, Rahman N, *et al.* Biological activity of *Dipterygium glaucum*. Pak J Biol Sci 2006;9:1173-4.
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. Pharmacogn Rev 2010;4:118-26.
- Gülçin I. Antioxidant activity of food constituents: An overview. Arch Toxicol 2012;86:345-91.
- Mothana RA, Abdo SA, Hasson S, Althawab FM, Alaghabari SA, Lindequist U. Antimicrobial, antioxidant and cytotoxic activities and phytochemical screening of some Yemeni medicinal plants. Evid Based Complement Alternat Med 2010;7:323-30.
- Jayaprakasha G, Patil BS. *In vitro* evaluation of the antioxidant activities in fruit extracts from citron and blood orange. Food Chem 2007;101:410-8.
- Singh RP, Chidambara Murthy KN, Jayaprakasha GK. Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using *in vitro* models. J Agric Food Chem 2002;50:81-6.
- Mosmann T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. J Immunol Methods 1983;65:55-63.
- Al-Salahi R, Alswaidan I, Marzouk M. Cytotoxicity evaluation of a new set of 2-aminobenzo[de]iso-quinoline-1,3-diones. Int J Mol Sci 2014;15:22483-91.
- Bauer AW, Kirby WM, Sherris JC, Turk M. Antibiotic susceptibility testing by a standardized single disk method. Am J Clin Pathol 1966;45:493-6.
- Lu Y, Foo LY. Flavonoid and phenolic glycosides from *Salvia officinalis*. Phytochemistry 2000;55:263-7.
- Wang DM, Pu WJ, Wang YH, Zhang YJ, Wang SS. A new isorhamnetin glycoside and other phenolic compounds from *Callianthemum taipaicum*. Molecules 2012;17:4595-603.
- Kim DW, Cho HI, Kim KM, Kim SJ, Choi JS, Kim YS, *et al.* Isorhamnetin-3-O-galactoside protects against CCl4-induced hepatic injury in mice. Biomol Ther (Seoul) 2012;20:406-12.
- Liu H, Mou Y, Zhao J, Wang J, Zhou L, Wang M, *et al.* Flavonoids from *Halostachys caspica* and their antimicrobial and antioxidant activities. Molecules 2010;15:7933-45.
- Pereira C, Barreto Júnior CB, Kuster RM, Simas NK, Sakuragui CM, Porzel A, *et al.* Flavonoids and a neolignan glucoside from *Guarea macrophylla* (Meliaceae). Quim Nova 2012;35:1123-6.
- Bravo L. Polyphenols: Chemistry, dietary sources, metabolism, and nutritional significance. Nutr Rev 1998;56:317-33.
- Nagaprasanthan LD, Vatsyayan R, Singhal J, Fast S, Roby R, Awasthi S, *et al.* Anti-cancer effects of novel flavonoid vicenin-2 as a single agent and in synergistic combination with docetaxel in prostate cancer. Biochem Pharmacol 2011;82:1100-9.
- Lee HJ, Lee HJ, Lee EO, Ko SG, Bae HS, Kim CH, *et al.* Mitochondria-cytochrome C-caspase-9 cascade mediates isorhamnetin-induced apoptosis. Cancer Lett 2008;270:342-53.
- Alves TM, Silva AF, Brandão M, Grandi TS, Smânia E, Smânia Júnior A, *et al.* Biological screening of Brazilian medicinal plants. Mem Inst Oswaldo Cruz 2000;95:367-73.