

Okinawan pit viper genome reveals evolution of snake venom

October 4 2017



The habu (*Protobothrops flavoviridis*), native to Okinawa, can reach lengths of 2.5 meters. Credit: Okinawa Institute of Science and Technology Graduate University

A bite from a pit viper, locally known as habu, can cause permanent

disability and even death. Yet, much about its venom remains an enigma. Highly variable in composition, even between littermates, this toxic cocktail keeps changing over generations.

A recent study in *Genome Biology and Evolution* sheds light on the evolution of snake venoms. For the first time, researchers have sequenced a habu genome, that of the Taiwan habu (*Protobothrops mucrosquamatus*), and compared it to that of its sister species, the Sakishima habu (*Protobothrops elegans*).

More than 50 instances of snake bites were recorded in the past year on Okinawa alone, prefectural government figures show. Globally, snake bites cause between 81,000 and 138,000 mortalities per year, according to the World Health Organization. In developing countries and rural areas with high exposure to venomous species and scant medical resources, [snake bites](#) can be especially devastating. For such places, creating effective antivenom can be a matter of life or death.

"For many years it was known that snake venoms evolve very rapidly, and the most common explanation for this has been natural selection," said Alexander Mikheyev, senior author on the paper and head of the Ecology and Evolution Unit at the Okinawa Institute of Science and Technology (OIST), "but there are reasons to suspect that this might not be the only evolutionary force at work."



The Taiwan habu (*Protobothrops mucrosquamatus*) is an invasive species that has become well established in Okinawa. Credit: Okinawa Institute of Science and Technology Graduate University

By taking samples of venoms and soft tissues from more than 30 specimens of the Taiwan and Sakishima habus, invasive but well-established species on Okinawa, researchers from OIST and the Okinawa Prefectural Institute of Health and Environment were able to map entire sequences of [venom](#) genes. Their study shows more than one factor at play in the evolution of this venom.

To understand how the chemical composition of a snake bite evolves, it

is crucial to understand its redundancy. Like multiple engines that allow a plane to fly if one of them should fail, venom targets multiple systems, assuring the snake's success. This complex mixture of proteins and small organic molecules attacks crucial prey physiological systems, such as blood pressure or blood coagulation, at several points. Even if one venom component does not prove optimally effective, various others do.

Typically, a habu injects a small amount of venom, a drop the size of a pinhead. Yet, it is more than strong enough to paralyze a rodent. Evolutionary biologists call this surplus power, which prevents prey from injuring or killing a [snake](#), "overkill."

Over time, as snakes reproduce, advantageous traits of venom are passed on to offspring in the process of [natural selection](#). However, the offspring can also inherit other traits—not necessarily beneficial ones. Because the average dose of venom is so high—in some cases killing prey almost instantaneously—it can mask inefficiencies in the venom's chemical makeup. These inefficiencies can be passed down from generation to generation with relatively little effect on the venom's function.



A Sakishima habu (*Protobothrops elegans*), endemic to the southwestern Ryukyus, has now become established in southern Okinawa Island. This specimen, encountered on Iriomote Island, was eating a tree lizard (*Japalura polygonata*). Credit: Okinawa Institute of Science and Technology Graduate University

"You can think of venoms evolving over two axes," said Mikheyev. "One of those is pushing them to be more effective, but another axis actually pushes them to be less effective."

"We're only now coming up with analytical methods to look at venoms comprehensively," said Steven Aird, first author on the paper. "There's a tremendous amount we can learn."

The researchers' work opens the door to new avenues of study as well as medical applications.

More information: Steven D. Aird et al. Population genomic analysis of a pitviper reveals microevolutionary forces underlying venom chemistry, *Genome Biology and Evolution* (2017). [DOI: 10.1093/gbe/evx199](https://doi.org/10.1093/gbe/evx199)

Provided by Okinawa Institute of Science and Technology Graduate University

Citation: Okinawan pit viper genome reveals evolution of snake venom (2017, October 4) retrieved 26 April 2024 from <https://phys.org/news/2017-10-okinawan-pit-viper-genome-reveals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.