


Research team reveals cellular and molecular basis of spur development in Impatiens uliginosa

February 29 2024



IuABP and IuEXT silencing led to phenotypic changes in spurs. Credit:



Horticulture Research

As an important reproductive organ of angiosperms, flowers have a clear purpose and adaptive significance in their various characteristics. As a typical floral evolutionary feature, the floral spur is a tubular structure extending from the petal, which has undergone several independent evolutions in angiosperms (e.g., Impatiens, Aquilegia, Linaria, etc.).

It plays a vital role in the pollination process because of its properties of secreting and storing nectar. In addition, the morphology (length, diameter, degree of distortion), color, and internal structure of spur exhibit a wide range of <u>species diversity</u>.

In the process of interacting with pollinators, spur improves pollination efficiency and ecological adaptability through the evolution of characteristics such as length and curvature. Plant lineages with spur often exhibit astonishing species diversity and faster species formation rates, so spur is considered a key innovation and has been the focus of numerous studies such as evolution and ecological hypotheses.


The study on the spur development process and the mechanism is helpful to further understand the species diversity and evolutionary mechanism of plant lineages. However, research on spur development has only been conducted in a few species, and its mechanism is not yet clear, showing significant species differences.

In January 2024, *Horticulture Research* published research titled "<u>The</u> <u>cellular and molecular basis of the spur development in Impatiens</u> <u>uliginosa</u>."

In this study, Impatiens uliginosa was used to analyze the characteristics



and rules of spur development from the perspectives of morphology, histocytology, physiology and <u>molecular biology</u>. The spur development in I. uliginosa is composed of early <u>cell division</u> and later anisotropic cell elongation.



The variation in cell number and cell morphology during spur development. Credit: *Horticulture Research*

During the entire development process, there were two imbalanced cell division events on the inner and outer sides of the spur, which



respectively led to the unique upward growth of the spur in the early stage and the formation of spur curve in the middle stage. In addition, substances such as auxin, gibberellin, and jasmonic acid exhibit significant differences in different parts and stages of spur development, indicating that hormones are involved in regulating spur development.

The regulation of IuABP and IuEXT genes on spur development was clarified by using RNA-seq and transient expression system in I. uliginosa. The downregulation of IuABP in spur leads to an increase in the number of spur curves and morphological distortion, suggesting that it may regulate cell division or elongation at different stages through the mediation of auxin action, and coordinate the transition between them.

The downregulation of IuEXT significantly shortened the spur length by inhibiting cell elongation, leading to the disappearance of spur curve and the inner epidermal papillae.

This study provides new insights into the cellular and molecular mechanisms of spur development in angiosperms.

More information: Yang Li et al, The cellular and molecular basis of the spur development in Impatiens uliginosa, *Horticulture Research* (2024). DOI: 10.1093/hr/uhae015

Provided by NanJing Agricultural University

Citation: Research team reveals cellular and molecular basis of spur development in Impatiens uliginosa (2024, February 29) retrieved 27 April 2024 from <u>https://phys.org/news/2024-02-team-reveals-cellular-molecular-basis.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.