



**Entry for TAVERNIERA abyssinica A. Rich. [family  
LEGUMINOSAE-PAPILIONOIDEAE]**

<http://plants.jstor.org/flora/flota001983>

<http://www.jstor.org>

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the contributing partner regarding any further use of this work. Partner contact information may be obtained at <http://plants.jstor.org/page/about/plants/PlantsProject.jsp>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [plants@jstor.org](mailto:plants@jstor.org).

**Entry for TAVERNIERA abyssinica A. Rich. [family LEGUMINOSAE-PAPILIONOIDEAE]**

Herbarium	Royal Botanic Gardens, Kew (K)
Collection	Flora of Tropical Africa
Resource Type	Reference Sources
Entry from	Flora of Tropical Africa, Vol 2, page 1 (1871) Author: Papilionaceae by Mr. J. G. Baker; Caesalpinieae and Mimoseae by Prof. Oliver)
Names	TAVERNIERA abyssinica A. Rich. [family LEGUMINOSAE-PAPILIONOIDEAE], Fl. Abyss. i. 206. Onobrychis simplicifolia R. Br. [family LEGUMINOSAE-PAPILIONOIDEAE], in Salt, App. lxxv.?
Information	A much branched shrub with terete pubescent branches. Leaves simple, small, short-stalked, obovate or obovate-oblong, glabrous above, paler and grey-silky beneath, the stipules as in the preceding. Flowers reddish, in few-flowered short-stalked axillary racemes. Calyx silky, the teeth narrow, acute. Pod linear, with 2–3 joints, clothed with white silky hairs.
Distribution	Nile Land Abyssinia: Petit.

<http://plants.jstor.org/flora/flota001983>