

Species	I. Abundance			IIa. Length statistics (mm)							IIb. Length frequency M = mean length		Growth characteristics: variations with length												
	Total in split	Perfect in split	Total in sample	Forms measured	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	Number of pairs of chambers	III. Number of chambers				IV. Width				V. Flare			
														Length (mm)				Length (mm)				Length (mm)		Length (mm)	
<i>Bolivina (Bolivina) pacifica</i>	19	9	2432	Total	106	S.R. 0.13-0.50 O.R. 0.16-0.48	0.31 ± 0.01	0.05	0.06 ± 0.004	16.8 ± 0.81		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.017$ mm				0.25	0.25 0.50 0.75 1.00				4	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) interjuncta bicostata</i>	13	11	1664	Total	237	S.R. 0-1.13 O.R. 0.17-1.04	0.52 ± 0.01	0.17	0.20 ± 0.01	38.7 ± 1.8		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = < 0.017$ mm $\times = 0.05-0.084$ mm				0.50	0.25 0.50 0.75 1.00				3	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) striatula</i>	230	204	29,440	Living forms	53	O.R. 0.16-0.45	0.32	0.06	0.07	20.3		0.25 0.50 0.75 1.00	6	Proloculus diameter: $\cdot = 0.017-0.05$ mm				0.50	0.25 0.50 0.75 1.00				4	0.25 0.50 0.75 1.00	
				Dead forms	204	S.R. 0.23-0.55 O.R. 0.20-0.48	0.39 ± 0.002	0.04	0.04 ± 0.003	13.4 ± 0.66		0.25 0.50 0.75 1.00	30	Proloculus diameter: $\cdot = 0.017-0.033$ mm				0.25	0.25 0.50 0.75 1.00				1	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) tongi flaccostata</i>	17	17	2162	Total	87	S.R. 0.22-0.46 O.R. 0.23-0.42	0.34 ± 0.004	0.03	0.04 ± 0.003	11.9 ± 0.9		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.017-0.033$ mm				0.25	0.25 0.50 0.75 1.00				3	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) acuminata</i>	6	4	768	Total	83	S.R. 0.04-0.55 O.R. 0.15-0.50	0.29 ± 0.01	0.07	0.08 ± 0.01	26.5 ± 2.2		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.017$ mm $\times = 0.025-0.059$ mm				0.25	0.25 0.50 0.75 1.00				3	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) vaughani</i>	3	3	384	Total	53	S.R. 0.12-0.22 O.R. 0.13-0.20	0.17 ± 0.003	0.01	0.02 ± 0.001	10.2 ± 0.99		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = < 0.017$ mm				0.25	0.25 0.50 0.75 1.00				3	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) inflata</i>	2	2	256	Total	40	S.R. 0.12-0.32 O.R. 0.17-0.28	0.22 ± 0.01	0.03	0.03 ± 0.003	14.8 ± 1.7		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.017$ mm				0.25	0.25 0.50 0.75 1.00				2	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) bradyi</i>	2	2	256	Total	18	S.R. 0.22-0.90 O.R. 0.26-0.74	0.56 ± 0.03	0.08	0.11 ± 0.02	20.3 ± 3.4		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.025-0.042$ mm				0.25	0.25 0.50 0.75 1.00				4	0.25 0.50 0.75 1.00	
<i>Suggrunda eckisi</i>	11	10	1408	Total	55	S.R. 0.13-0.36 O.R. 0.18-0.33	0.25 ± 0.01	0.03	0.04 ± 0.003	14.9 ± 1.4		0.25 0.50 0.75 1.00	15	Proloculus diameter: $\cdot = 0.017$ mm				0.25	0.25 0.50 0.75 1.00				2	0.25 0.50 0.75 1.00	

SAMPLE C-17 (82 METERS), QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens