

A new miniature species of *Priocharax* (Teleostei: Characiformes: Characidae) from the Rio Madeira drainage, Brazil, with comments on the adipose fin in characiforms

GEORGE M. T. MATTOX^{1,*}, CAMILA S. SOUZA², MÔNICA TOLEDO-PIZA³, RALF BRITZ⁴ & CLAUDIO OLIVEIRA²

¹ Laboratório de Ictiologia de Sorocaba, Departamento de Biologia, Universidade Federal de São Carlos – campus Sorocaba, São Paulo, Brazil — ² Departamento de Biologia Estrutural e Funcional, Instituto de Biociências, Universidade Estadual Paulista – campus Botucatu, São Paulo, Brazil — ³ Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil — ⁴ Museum of Zoology, Senckenberg Dresden, A. B. Meyer Building, 01109 Dresden, Germany — * Corresponding author; email: gmattox@ufscar.br

Submitted June 5, 2020.

Accepted July 24, 2020.

Published online at www.senckenberg.de/vertebrate-zoology on August 11, 2020.

Published in print Q3/2020.

Editor in charge: Uwe Fritz

Abstract

A new miniature species of the freshwater fish genus *Priocharax* is described from the Rio Jamari, Rio Madeira drainage, Amazon basin. *Priocharax varii* sp. n. is the fourth species currently recognized in the genus and shares with the other three the presence of a conspicuous larval pectoral fin in adults, a fully toothed maxilla, a triangular pseudotympanum situated anterior to the fifth vertebra and a tiny size combined with a translucent body. The new species is diagnosed from congeners in having an adipose fin, a unique feature within *Priocharax*. Molecular identification of the new species and the other valid *Priocharax* species supports the morphological findings. A brief discussion on the presence/absence of the adipose fin in Characiformes is provided.

Key words

Amazon River basin, barcoding, biodiversity, osteology, taxonomy.

Introduction

Priocharax is a genus originally described by WEITZMAN & VARI (1987) to include two miniature species (*sensu* WEITZMAN & VARI, 1988): *Priocharax ariel* Weitzman & Vari and *Priocharax pygmaeus* Weitzman & Vari. These species were collected from the upper Río Orinoco in Venezuela and in the region of Leticia, Río Amazonas Colombia, respectively. The authors proposed the presence in adults of *Priocharax* of a peculiar, larval-like, rayless pectoral fin as the main character to diagnose the genus. They also listed a set of other diagnostic features to distinguish *Priocharax* from most New World characids except those included in the Characinae and Cynopotaminae (=Characinae): dentary with approximately

30–50 conical teeth in somewhat irregular row, upper jaw (premaxilla and maxilla) with approximately 45–90 conical teeth in somewhat irregular row, and adult body size of 11–17 mm SL, a feature otherwise unknown in the Characinae and Cynopotaminae (=Characinae) at that time. Another feature that according to WEITZMAN & VARI (1987) distinguishes *Priocharax* from the remaining characines is the number of i,5 pelvic-fin rays, with all the remaining members of the subfamily having i,7 pelvic-fin rays.

Based on the presence of numerous conical teeth in the upper and lower jaws and on the elongate maxilla, WEITZMAN & VARI (1987) hypothesized that the relation-

ships of *Priopcharax* would be among the Characinae, pending further studies. Twenty-five years later, MATTOX & TOLEDO-PIZA (2012) carried out a phylogenetic study of this subfamily and found out that the Characinae as previously proposed were not monophyletic. Rather, they restricted the Characinae to a small group of seven genera (*Acanthocharax* Eigenmann, *Acstrocephalus* Eigenmann, *Charax* Scopoli, *Cynopotamus* Valenciennes, *Galeocharax* Fowler, *Phenacogaster* Eigenmann and *Roeboides* Günther) and transferred the other genera of more diminutive fishes to the Heterocharacinae (*Gnathocharax* Fowler, *Heterocharax* Eigenmann, *Hoplocharax* Géry, *Lonchogenys* Myers and *Priopcharax*), together with *Gilbertolus* Eigenmann and *Roestes* Günther. Similar results were obtained by other morphological (i.e. MIRANDE, 2010) and molecular approaches (i.e., OLIVEIRA *et al.*, 2011). A recent dataset (MIRANDE, 2018) combining both sources of information also partially corroborated the results, albeit not including exactly the same taxa, but none of those studies except that of MATTOX & TOLEDO-PIZA (2012) included *Priopcharax* in their analyses. In the same year, BETANCUR-R *et al.* (2018) included *Priopcharax ariel* in a molecular analysis of characiforms based on exons and recovered the genus among genera of the Characinae, but did not discuss this result any further.

Since WEITZMAN & VARI's (1987) publication, a third species of *Priopcharax* was discovered in the Rio Negro: *P. nanus* Toledo-Piza *et al.* (2014). *Priopcharax nanus* was diagnosed from *P. ariel* and *P. pygmaeus* by the unique presence of i,6 pelvic-fin rays, presence of the claustrum and two post-cleithra. Although still a miniature, *P. nanus* was interpreted as the least ontogenetically truncated species in the genus. Additional specimens of *Priopcharax* were reported in the literature from new localities (e.g., QUEIROZ *et al.*, 2013; VIEIRA *et al.*, 2016) or collected during recent expeditions. Upon closer study, samples from the Rio Jamari, Rio Madeira drainage, Amazon basin, Brazil were found to represent a new species, which is described in this paper.

Material & Methods

Morphological analysis

Counts and measurements follow FINK & WEITZMAN (1974), Weitzman & Vari (1987) and Menezes & Weitzman (1990) and were taken on the left side of each specimen whenever possible. All measurements other than standard length (SL) are expressed as percentages of SL, except for subunits of the head which are expressed as percentages of head length (HL). Caudal-peduncle depth is also expressed as percentage of caudal-peduncle length. Measurements were taken point to point with a precision of 0.1 mm from digital photographs of specimens taken under the stereomicroscope. Counts of vertebrae, teeth, gill-rakers and procurrent caudal-fin rays were obtained

from five specimens cleared and double stained for cartilage and bone following the protocol of TAYLOR & VAN DYKE (1985). Total vertebral number includes the four vertebrae of the Weberian apparatus as separated elements. The compound ural centrum was counted as a single vertebra. The gill-raker at the junction of the ceratobranchial and epibranchial is considered as the posteriormost gill-raker on the lower branch of the gill arch. Photographs were made with a Zeiss Discovery V20 stereomicroscope with a Zeiss AxioCam digital camera attached. Osteological terminology follows WEITZMAN (1962) except for inner arm of the *os suspensorium* instead of *os suspensorium*, and outer arm of the *os suspensorium* instead of rib of fourth vertebra, following CONWAY & BRITZ (2007), and other updates summarized in MATTOX *et al.* (2014). In the description, the frequency of each count is provided in parentheses after the respective count, with the count of the holotype indicated by an asterisk. Information on meristic and morphometric data of *Priopcharax ariel* and *P. pygmaeus* were taken from WEITZMAN & VARI (1987). Specimens examined for this study are deposited in the Coleção de Peixes da Universidade Federal de Rondônia (UFRO-ICT), Instituto Nacional de Pesquisas da Amazônia (INPA), Laboratório de Biologia e Genética de Peixes, Departamento de Biologia Estrutural e Funcional, Universidade Estadual Paulista, Botucatu (LBP), Museu de Zoologia da Universidade de São Paulo (MZUSP), and National Museum of Natural History – Smithsonian (USNM). Sampling for this study was authorized by Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) through permit number SISBIO/MMA 45429.

Molecular analysis

Voucher specimens for the molecular study are deposited in the collection of Laboratório de Biologia e Genética de Peixes (LBP), Departamento de Biologia Estrutural e Funcional, Instituto de Biociências, Universidade Estadual Paulista, Botucatu, São Paulo, Brazil. This study is in accordance with the National Council for the Control of Animal Experimentation (CONCEA) approved by UNE-SP Ethics Committee on Use of Animals (CEUA), protocol number 1058. Twenty specimens of *Priopcharax* were included in the analysis: one specimen of *P. nanus* and six specimens of *P. ariel* from the Rio Negro, nine specimens of *P. pygmaeus* from the Río Amazonas and four specimens of the new species from the Rio Madeira, described herein. A single specimen of *Acstrocephalus sardina* (Characiformes, Characidae, Characinae) was used to root the trees. Voucher data are summarized in Table 1.

DNA extraction followed IVANOVA *et al.* (2006), and partial sequences of the cytochrome c oxidase subunit I (COI) gene were amplified by polymerase chain reaction (PCR) with primers FishF1, FishR1 and FishF2, FishR2 (WARD *et al.*, 2005) or L6252-Asn, H7271-COXI (MELO *et al.*, 2011). PCR amplifications were performed in a total volume of 12.5 µl that included 1.25 µl of 10X buffer, 0.25 µl of MgCl₂ (50 mM), 0.2 µl dNTPs (2 mM), 0.5 µl of

each primer (5 mM), 0.1 µl of PHT Taq DNA polymerase (*Phoneutria*), 1.0 µl of genomic DNA (200 ng) and 8.7 µl ddH₂O. The thermocycling profile consisted of an initial denaturation (5 min at 94°C), followed by 30 cycles of chain denaturation (40s at 94°C), primer hybridization (30s at 50–54°C) and nucleotide extension (1 min at 68°C), and a final extension (8 min at 68°C). All PCR products were checked on 1% agarose gels and then purified with ExoSap-IT (USB Corporation) following the manufacturer's instructions. The subsequently purified PCR products were submitted to sequencing reactions using BigDye Terminator v 3.1 Cycle Sequencing Ready Reaction Kit (Applied Biosystems) and purified again by ethanol precipitation. Products were loaded onto an ABI 3130 DNA Analyzer automatic sequencer (Applied Biosystems).

Raw sequences were assembled to consensus using GENEIOUS v7.1.9 (KEARSE *et al.*, 2012) and subsequently aligned with MUSCLE (EDGAR, 2004) under default parameters. To evaluate the occurrence of substitution saturation, we estimated whether the Iss (index of substitution saturation) is significantly lower than Iss.cAsym (asymmetrical topology) and Iss.Sym (symmetrical topology) using the method of XIA *et al.* (2003) in DAMBE v5.3.38 (XIA, 2013). Nucleotide variation, substitution patterns and best-fit model of nucleotide evolution were estimated in MEGA v6.0 (TAMURA *et al.*, 2013).

The overall mean distance (among all specimens), the intraspecific distances (among specimens of each species group) and interspecific distances (among species groups) were estimated using the Kimura-2-parameter model (K2P; KIMURA, 1980) in MEGA v6.0 (TAMURA *et al.*, 2013). The Maximum-likelihood analysis (ML) was generated with 1000 bootstrap replicates in MEGA v6.0 (TAMURA *et al.*, 2013). Species delimitation analyses included two approaches: (a) the Automatic Barcode Gap Discovery analysis (ABGD; PUILLANDRE *et al.*, 2012) available in the ABGD webserver (<https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html>), excluding the root *Acestrocephalus sardina* and under default parameters; and (b) Poisson Tree Processes (PTP; ZHANG *et al.*, 2013) using the ML tree and other parameters at default in the PTP webserver (<http://species.h-its.org/ptp/>). Groups were ordered based on ABGD and PTP results.

Results

Priocharax variii, sp. nov.

ZOOBANK: urn:lsid:zoobank.org:act:811F685B-BBB7-45B2-8294-C1BC837565DB

Figs. 1–4, Table 2

Holotype. MZUSP 125786, 12.2 mm SL, Brazil, Rondônia, Candeias do Jamari, Rio Preto approximately 12km South from road BR-364, before bridge crossing the river, Rio Jamari drainage, 08°52'53.5"S, 63°37'50.8"W, G. Mattox & S. Souza, 02 Sep 2018.

Paratypes. LBP 28495, 5, 12.5–12.7 mm SL, collected with holotype. MZUSP 125787, 44, 11.8–14.0 mm SL (5 c&s, 11.8–13.7 mm SL), collected with holotype. UFRO-ICT 027656, 5, 12.2–13.3 mm SL, collected with holotype.

Diagnosis. *Priocharax variii* is distinguished from all congeners by the presence of an adipose fin (vs adipose fin absent). *Priocharax variii* can be further diagnosed from *P. ariel* and *P. pygmaeus* by the presence of i,6 pelvic-fin rays (i,5 in one out of 50 specimens examined) (vs i,5) and of two postcleithra (vs postcleithra absent). *Priocharax variii* can be further diagnosed from *P. ariel* by having a shorter caudal peduncle (14–18 % SL vs 18–24 % SL).

Description. For overall appearance, see Figure 1. Morphometric data of holotype and paratypes are presented in Table 2. Body laterally compressed, greatest depth at vertical through dorsal-fin origin. Dorsal-fin origin approximately at midbody, at vertical slightly anterior to anal-fin origin. Pectoral-fin bud at vertical through anterior portion of pseudotympanum. Pelvic-fin origin approximately midway between posterior margin of opercle and anal-fin origin. Dorsal profile of head and body slightly convex from tip of snout to dorsal-fin origin. Dorsal profile of body along dorsal-fin base nearly straight, gently sloping posteroventrally; straight and posteroventrally sloping from latter point to caudal peduncle. Dorsal profile of caudal peduncle slightly concave to base of dorsal procurrent rays. Ventral profile of head and body slightly convex from symphysis of lower jaw to vertical through pectoral-fin origin; straight to slightly convex from latter point to pelvic-fin origin. Ventral profile of body posteroventrally sloping from pelvic-fin to anal-fin origin; straight and posterodorsally rising along anterior one-half of anal-fin base, gently concave from latter point to base of ventral procurrent rays. Caudal peduncle short. Pseudotympanum located anterior to rib of fifth vertebra.

Snout blunt in lateral view. Infraorbitals 1 to 6 and supraorbital absent, antorbital present. Mouth terminal with lower jaw slightly shorter than upper jaw. Tip of maxilla elongate, posterior border reaching vertical through posterior border of pupil. Premaxilla with 23(1), 25(2), 26(1) or 29(1) teeth in single series. Maxilla with 34(1), 39(2) or 42(2) teeth. Dentary with 37(2), 40(1), 42(1) or 46(1) teeth. Dentary teeth in single series, with few anterior teeth slightly displaced anteriorly. Conspicuous elongate foramen at anterior portion of dentary. All jaw teeth small, conical and lingually curved to a moderate extent (Figure 2).

Dorsal-fin rays ii,9*(50). Endoskeletal part and some dermal bones of pectoral fin with structure similar to larval specimens of most characiforms (Figure 3). Cartilaginous pectoral-radial plate with incomplete longitudinal middle fissure leaving upper and lower halves connected at base and tip; base articulating with vertically elongated scapulocoracoid cartilage and round distal margin with larval-like pectoral-fin fold supported solely by actinotrichia. Pectoral-fin rays absent. All bones of endoskeletal pectoral girdle absent, exoskeletal part with posttem-



Fig 1. *Priocarax varii*, a) holotype, MZUSP 125786, 12.2 mm SL; Brazil, Rondônia, Candeias do Jamari, Rio Preto approximately 12km South from road BR-364, before bridge crossing the river, Rio Jamari drainage; scale bar = 2mm. b) live specimen photographed right after capture. Arrow points to adipose fin.

poral, supracleithrum, cleithrum and two postcleithra. Cleithrum with posteriorly directed, curved process immediately below ventral tip of supracleithrum. Pelvic-fin rays i,5 (1) or i,6*(49). Posterior tip of pelvic fin falling short of origin of anal fin but extending slightly beyond vent. Anal-fin rays ii, 18(1), 19(1), 20(1), 21(4), 22(11), 23*(14), 24(14) or 25(2). Anal-fin distal margin concave with anterior lobe formed by elongated fin rays and posterior portion with shorter rays. Caudal-fin rays i,8,7,1 (1), i,8,8,i (1) or i,9,8,i*(42), dorsal procurrent rays 9(5), ventral procurrent rays 7(4) or 8(1). Caudal fin forked. Adipose fin small and situated midway between terminus of dorsal fin and anteriormost dorsal procurrent caudal-fin ray ($n=50$).

Squamation present in almost all specimens, but scales highly deciduous and easily lost during handling. Scales cycloid, very thin, with no obvious circuli or radii. Scales in midlateral row 26(2), 27*(14), 28(14), 29(9), 30(6), 31(1) or 32(2); no canal bearing lateral-line scales on body. Horizontal scale rows between dorsal-fin origin and pelvic-fin origin 7(3) or 8*(7). Horizontal scale rows around caudal peduncle 8*(5) or 9(5). Predorsal scales typically absent with one or two scales just anterior to dorsal fin in few specimens. Scales restricted to base of caudal-fin rays, not covering caudal-fin lobes.

Total vertebrae 33(5); abdominal vertebrae 14(1) or 15(4); caudal vertebrae 18(4) or 19(1). Total number of gill-rakers on first branchial arch 8(1), 10(1), 11(2) or 12(1), upper limb gill-rakers 1(1) or 3(4), lower limb gill-rakers 7(2), 8(2) or 9(1). Weberian apparatus well devel-

oped, all components ossified except for claustrum (Figure 4), absent in all but one specimen with tiny, vestigial claustrum. Large gap between neural arches 3 and 4, with gap partially filled by dorsally projecting pointed process from vertebral centrum 3. Inner arm of *os suspensorium* large, projecting forward to vertical through middle of second centrum.

Colour in alcohol. Overall ground coloration pale yellow (Fig. 1a). Patch of melanophores on dorsal portion of head and scattered melanophores on opercle. Head with dark stripe radiating anteriorly from eye to tip of snout. Line of melanophores along dentary with concentration of melanophores on articulation between lower jaw and quadrate. Iridophores present in eye of most specimens. Longitudinal line of melanophores along midlateral body, from vertical through dorsal-fin origin to middle portion of caudal peduncle. Triangular to circular patch of melanophores at base of caudal fin forming inconspicuous spot. Bases of anal-fin rays with series of spots forming irregular line along fin. Second dark line, dorsal and more superficial, extending along ventral margin of hypaxial myomeres posteriorly from vertical through fifth branched anal-fin ray. Both lines more evident and well separated anteriorly but approaching each other posteriorly. Two patches of melanophores ventrally on body anterior to pelvic fin: anteriormost in form of an inconspicuous small spot on isthmus and posteriormost patch elongated just anterior to basipterygium. Few melanophores present in region around vent. Melanophores at




Fig 2. *Priocnemis varii*, paratype, MZUSP 125787, 13.6 mm SL, c&s; jaws in lateral view. Ana = anguloarticular; Cm = coronomeckelian; De = dentary; MC = Meckel's cartilage; Mx = maxilla; Pmx = premaxilla; Ra = retroarticular. Scale bar = 0.2 mm.




Fig 3. *Priocnemis varii*, paratype, MZUSP 125787, 13.6 mm SL, c&s; shoulder girdle in lateral view. Cl = cleithrum; PecRdC = pectoral-fin radial cartilage; Pcl = unidentified postcleithrum; Pcl3 = postcleithrum 3; Pt = posttemporal; ScCoC = scapulocoracoid cartilage; Sucl = supracleithrum. Arrow points to middle fissure on pectoral-fin radial cartilage. Tip of larval finfold is bent. Scale bar = 0.5 mm.




Fig 4. *Priocnemis varii*, paratype, MZUSP 125787, 13.6 mm SL, c&s; Weberian apparatus in lateral view. Exoc = Exoccipital; Int = intercalarium; NA3–4 = neural arches 3–4; NS4 = neural spine 4; OsS = os suspensorium; Sca = scaphium; Sn3 = supraneural 3; Soc = supraoccipital; T = tripus. Scale bar = 0.2 mm.

origin of dorsal, pelvic, and anal fins. All fins except pectoral and adipose with scattered melanophores along margins of fin rays. Caudal fin with two faint dark blotches on bases of four dorsal- and four ventralmost rays.

Colour in life. Body mostly translucent (Fig. 1b). Pattern of melanophores generally as described above for alco-

hol specimens. Melanophores also scattered on dorsal surface of swim bladder and along neural cord. Melanophores in line along base of dorsal-fin rays. Melanophores arranged along neural and haemal spines deep inside body forming few oblique lines. Dorsal portion of head and snout predominantly golden yellow. Xanthophores along dorsal profile of body, especially along dorsal midline and dorsal-fin base. Few scattered xanthophores on dorsal flanks of body. Xanthophores also present and abundant along neural chord and ventral margin of vertebral column, dorsal surface of swim bladder, anal-fin base and forming irregular patches on dorsal and ventral bases of caudal-fin lobes. Scattered melanophores on dorsal and caudal fins, all other fins hyaline including adipose fin. Eye silvery, dorsal half predominantly red and dorsal margin covered with melanophores.

Sexual dimorphism. Hooks, typical of many mature male characids, absent in dorsal-, pectoral-, pelvic-, and anal-fins of all examined specimens ($n=50$), but gonads not checked for maturity and sex.

Genetics. The alignment of the 21 terminals contained 652 bp with 208 variable sites. The nucleotide frequencies were 23.1% adenine, 17.8% guanine, 31.8% thymine, and 27.2% cytosine. DAMBE indicated no saturation in either transitions or transversions in both asymmetrical

Table 1. Vouchers, locality information, and GenBank accession numbers of the analyzed specimens of *Priobarax*.

| Lot | Voucher | Species | Basin | Locality | Coordinates | GenBank n. |
|-----------|-----------|--------------------------------|--------------|--|------------------------------|------------|
| LBP 28495 | LBP 96981 | <i>Priobarax variii</i> | Rio Madeira | Rio Preto, affluent of Rio Jamari, Candeias do Jamari, RO, Brazil | 08°52'53.5"S 63°37'50.8"W | MT754786 |
| LBP 28495 | LBP 96982 | <i>Priobarax variii</i> | Rio Madeira | Rio Preto, affluent of Rio Jamari, Candeias do Jamari, RO, Brazil | 08°52'53.5"S 63°37'50.8"W | MT754785 |
| LBP 28495 | LBP 96984 | <i>Priobarax variii</i> | Rio Madeira | Rio Preto, affluent of Rio Jamari, Candeias do Jamari, RO, Brazil | 08°52'53.5"S 63°37'50.8"W | MT754783 |
| LBP 28495 | LBP 96985 | <i>Priobarax variii</i> | Rio Madeira | Rio Preto, affluent of Rio Jamari, Candeias do Jamari, RO, Brazil | 08°52'53.5"S 63°37'50.8"W | MT7547864 |
| LBP 28442 | LBP 98284 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Tibarrá on left side of Rio Negro, Santa Isabel do Rio Negro, AM, Brazil | 00°26'28.1"S 64°56'57.5"W | MT754780 |
| LBP 28442 | LBP 98285 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Tibarrá on left side of Rio Negro, Santa Isabel do Rio Negro, AM, Brazil | 00°26'28.1"S 64°56'57.5"W | MT754781 |
| LBP 27704 | LBP 98286 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Tapage, Rio Urubaxi, approximately 1 hour from mouth of river, S. I. Rio Negro, AM, Brazil | 00°30'05.3"S 64°49'11.7"W | MT754778 |
| LBP 27704 | LBP 98287 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Tapage, Rio Urubaxi, approximately 1 hour from mouth of river, S. I. Rio Negro, AM, Brazil | 00°30'05.3"S 64°49'11.7"W | MT754782 |
| LBP 27704 | LBP 98288 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Tapage, Rio Urubaxi, approximately 1 hour from mouth of river, S. I. Rio Negro, AM, Brazil | 00°30'05.3"S 64°49'11.7"W | MT754779 |
| LBP 25858 | LBP 96383 | <i>Priobarax ariel</i> | Rio Negro | Igarapé Uacatuna, São Gabriel da Cachoeira, AM, Brazil | 00°03'38.0"S 67°05'45.0"W | MT754777 |
| LBP 28490 | LBP 98283 | <i>Priobarax nanus</i> | Rio Negro | Igarapé Tibarrá on left side of Rio Negro, Santa Isabel do Rio Negro, AM, Brazil | 00°26'28.1"S 64°56'57.5"W | MT754766 |
| LBP 22464 | LBP 96986 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada La Ponderosa, Letícia, Colombia | 04°08'24.4"S 69°56'53.4"W | MT754771 |
| LBP 22464 | LBP 96987 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada La Ponderosa, Letícia, Colombia | 04°08'24.4"S 69°56'53.4"W | MT754774 |
| LBP 22464 | LBP 96988 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada La Ponderosa, Letícia, Colombia | 04°08'24.4"S 69°56'53.4"W | MT754769 |
| LBP 22464 | LBP 96998 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada La Ponderosa, Letícia, Colombia | 04°08'24.4"S 69°56'53.4"W | MT754768 |
| LBP 22739 | LBP 96989 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada Pichuna, Leticia, Colombia | 04°07'33.8"S 70°00'28.9"W | MT754772 |
| LBP 22739 | LBP 96990 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada Pichuna, Leticia, Colombia | 04°07'33.8"S 70°00'28.9"W | MT754773 |
| LBP 22739 | LBP 96991 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada Pichuna, Leticia, Colombia | 04°07'33.8"S 70°00'28.9"W | MT754776 |
| LBP 22739 | LBP 96992 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada Pichuna, Leticia, Colombia | 04°07'33.8"S 70°00'28.9"W | MT754770 |
| LBP 22739 | LBP 96993 | <i>Priobarax pygmaeus</i> | Río Amazonas | Quebrada Pichuna, Leticia, Colombia | 04°07'33.8"S 70°00'28.9"W | MT754775 |
| LBP 17833 | LBP 70161 | <i>Acestrocephalus sardina</i> | Rio Negro | Santa Isabel do Rio Negro, AM, Brazil | 00°30'5.3"S 64°49'12.2"W | MT754767 |

(Iss.cAsym) and symmetrical (Iss.cSym) topologies. The overall mean of K2P genetic distances without outgroup was 18.1%. The values of interspecific distances among the four species ranged from 20.7% between *Priobarax variii* and *P. nanus* to 30.5% between *P. ariel* and *P. pygmaeus* (Table 3). Intraspecific genetic variation ranged from 0% within *Priobarax pygmaeus* to 0.2% within *P. ariel* and *P. variii* (Table 3). The convergence of initial and recursive partitions of the ABGD analysis suggested four species, *Priobarax ariel*, *P. nanus*, *P. pygmaeus*,

and *P. variii* (Fig. 5; Supplementary Fig. 1). The ML solution of the PTP analysis returned identical results (Fig. 5; Supplementary Fig. 2).

Distribution. *Priobarax variii* is known to date from the type locality in the Rio Preto, a tributary of the Rio Jamari, itself an affluent of the Rio Madeira (Fig. 6). Specimens from several other localities in the Madeira system in Brazil, including some of its large tributaries (e.g., Jaciparaná, Jamari, Marmelos) as well as some are-

Table 2. Morphometric data of *Priobarax variii* sp. nov. (n=50, except for anal-fin length and upper jaw length with n=49, range includes holotype); SD = Standard Deviation.

| | Holotype | Range | Mean | SD |
|--------------------------------------|----------|-----------|------|-----|
| Standard length (SL) (mm) | 12.2 | 11.8–14.0 | — | — |
| Percentages of SL | | | | |
| Depth at dorsal-fin origin | 24 | 18–31 | 23.5 | 1.7 |
| Snout to dorsal-fin origin | 54 | 49–58 | 54.3 | 1.5 |
| Snout to pelvic-fin origin | 41 | 39–44 | 40.5 | 0.9 |
| Snout to anal-fin origin | 53 | 52–57 | 53.8 | 1.0 |
| Dorsal-fin length | 25 | 23–29 | 25.6 | 1.1 |
| Dorsal-fin base | 13 | 10–14 | 12.0 | 1.1 |
| Pelvic-fin length | 11 | 9–13 | 11.5 | 0.8 |
| Anal-fin length | 35 | 32–38 | 35.5 | 1.3 |
| Anal-fin base | 32 | 29–36 | 32.4 | 1.2 |
| Caudal-peduncle depth | 8 | 7–9 | 8.0 | 0.4 |
| Caudal-peduncle length | 16 | 14–18 | 15.6 | 0.9 |
| Head length (HL) | 26 | 24–27 | 25.3 | 0.6 |
| Percentages of HL | | | | |
| Orbital diameter | 36 | 32–57 | 34.8 | 3.5 |
| Interorbital distance | 38 | 34–47 | 39.1 | 2.9 |
| Snout length | 21 | 20–27 | 23.0 | 1.3 |
| Upper jaw length | 54 | 48–62 | 54.7 | 3.5 |
| Percentage of caudal-peduncle length | | | | |
| Caudal-peduncle depth | 51 | 43–63 | 51.4 | 4.0 |

Table 3. Pairwise K2P genetic distances among species of *Priobarax*. Intraspecific genetic variations are highlighted in bold. Numbers below diagonal are values of interspecific distances. Values are shown in percentages, followed by standard deviation. Number of sequences: *P. ariel* = 6, *P. nanus* = 1, *P. pygmaeus* = 9, *P. variii* = 4.

| | 1 | 2 | 3 | 4 |
|-----------------------------|----------------|----------|----------|----------------|
| 1 <i>P. ariel</i> | 0.2±0.1 | | | |
| 2 <i>P. pygmaeus</i> | 30.5±3.6 | 0 | | |
| 3 <i>P. nanus</i> | 21.9±2.8 | 21.1±2.9 | — | |
| 4 <i>P. variii</i> | 22.7±2.9 | 23.5±2.8 | 20.7±2.8 | 0.2±0.1 |

as near the main channel (e.g., Lago Cuniã) were examined herein and identified as *P. ariel*. Figure 6 also shows the known distribution of all congeners.

Ecological notes. Specimens were collected between 14:00 and 17:00 in a small tributary of the Rio Jamari (Fig. 7). They were all collected close to a bush of submerged macrophytes in a slow flowing part of the Rio Preto, approximately 200m above rapids. *Priobarax variii* was collected with *Apistogramma resticulosa* Kullander 1980, *Axelrodia lindeae* Géry 1973, *Axelrodia stigmatias* (Fowler 1913), *Hemigrammus bellottii* (Steindachner 1882), *Hyphessobrycon* aff. *swelesi* (Géry 1961), *Leporinus* sp., *Nannostomus trifasciatus* Steindachner 1876, *Odontocharacidium aphanes* (Weitzman & Kanazawa 1977) and *Satanoperca* sp. Three of these species are considered miniature (*sensu* Weitzman & Vari 1988): the characids *A. lindeae*, *A. stigmatias*

and the crenuchid *O. aphanes*. In addition to *Priobarax variii* and those three miniatures which are mostly transparent, also the larger *H. bellottii* has a predominantly translucent body in life, lending support to the idea of an “almost invisible league” (Fig. 8) for the Rio Negro proposed by CARVALHO *et al.* (2006).

Etymology. *Priobarax variii* honours the late Richard P. Vari, our dear friend and mentor, and one of the most influential researchers in Neotropical ichthyology of the past 40 years. A noun in genitive. In addition to being one of the authors of the genus *Priobarax*, he continuously encouraged the first author to study this taxon further and in more detail.

Discussion

WEITZMAN & VARI (1987) referred to the retention of the larval pectoral-fin structure in adults of *Priobarax* as the main diagnostic feature of genus (e.g., Figure 3). Based on the hypothesis of phylogenetic relationships of *Priobarax* with *Gnathocharax*, *Heterocharax*, *Hoplocharax* and *Lonchogenys* (MATTOX & TOLEDO-PIZA, 2012:880) this feature was corroborated as autapomorphic for the genus (TOLEDO-PIZA *et al.* 2014:234). In addition to retaining the larval rayless structure of the pectoral fin, *Priobarax variii* possesses most other features listed by WEITZMAN & VARI (1987) to diagnose *Priobarax*, except for the number of pelvic-fin rays.




Fig. 5. Maximum likelihood tree of four known species of *Priobarax*, based on the COI gene (652 pb). Values < 50% are not shown. Bars represent the number of species obtained by the ABGD and bPTP analyses. Numbers near nodes represent bootstrap support.

Among the diagnostic characters of *Priobarax* listed by WEITZMAN & VARI (1987), *Priobarax nanus*, described years later (TOLEDO-PIZA *et al.*, 2014), has a claustrum (vs. claustrum absent) and has six branched pelvic-fin rays (vs. five). Five to six branched pelvic-fin rays are still diagnostic for *Priobarax* among the Heterocharacinae, which invariably have seven. We recognize *Priobarax variii* as a member of the genus *Priobarax* based on the following features which are in accordance with the diagnostic features of the genus mentioned by WEITZMAN & VARI (1987) and discussed by TOLEDO-PIZA *et al.* (2014): the retention of larval pectoral fin in adults (Figure 3), upper and lower jaws with somewhat irregular single rows of conical teeth (57–71 and 37–46, respectively) (Figure 2), and low number of branched anal-fin (18–25) and branched pelvic-fin rays (6 – one specimens out of 50 with 5 branched rays).

Like the other species of the genus, *Priobarax variii* also has a number of reductive anatomical characters associated with miniaturization: e.g. loss of the laterosensory canal system on the head and body, loss of the bones of the infraorbital series and the presence of a gap in the Weberian apparatus between neural arches 3 and 4 (WEITZMAN & VARI, 1987: fig. 3; MATTOX *et al.*,

2013:313, TOLEDO-PIZA *et al.*, 2014: fig. 4). However, other anatomical complexes present different degrees of reduction among *Priobarax* species. For example, *Priobarax variii* and *P. nanus* have two postcleithra in the pectoral girdle, while in *P. ariel* and *P. pygmaeus* all postcleithra are absent. *Priobarax variii* and *P. nanus* have six branched pelvic-fin rays (with the exception of one specimen of *P. variii* with five) and *P. ariel* and *P. pygmaeus* have only five. On the other hand, the claustrum in the Weberian apparatus is absent in most specimens of *Priobarax variii* ($n=4$ out of 5), and in all specimens of *P. ariel* and *P. pygmaeus* (WEITZMAN & VARI, 1987) while it is present in *P. nanus* (TOLEDO-PIZA *et al.*, 2014).

Priobarax ariel, *P. nanus* and *P. pygmaeus* all lack the adipose fin, a feature interpreted herein as a developmental truncation since the adipose fin does not develop from the larval fin fold, but forms *de novo* in the ontogeny of Characiformes (FUIMAN, 1984; BENDER & MORITZ, 2014; MARINHO, 2017). *Priobarax variii* can be distinguished from all three congeners by the presence of a small adipose fin located midway between the dorsal-fin terminus and the anterior most caudal-fin ray (Figure 1). Presence of the adipose fin has been used as one of the




Fig. 6. Map of South America with a close-up of upper and central Amazon basin, Brazil, illustrating the distribution of the four species of *Priocnemis*: *P. variii* (yellow star) described herein from the Rio Madeira basin, *P. ariel* (green dots), *P. nanus* (red dot), and *P. pygmaeus* (blue dots). Colours are same as in Figure 5. Some dots may represent more than one lot.



Fig. 7. Rio Preto approximately 12km South from road BR-364, Candeias do Jamari, Rondônia, Brazil, the type locality of *Priocnemis variii*.

characters to diagnose species in other genera of Characiformes composed of miniature species that otherwise lack such structure: for example, *Elachocharax pulcher*

Myers is the only one of four miniatures in the crenuchid genus *Elachocharax* Myers that has an adipose fin (BUCKUP, 1993) and *Xenurobrycon pteropus* Weitzman &



Fig. 8. Examples of fishes found with *Priocarax variii* that are translucent in life: a) *Axelrodia stigmatias*; b) *Axelrodia lindeae*. Live specimens photographed right after capture.

Fink is the only one of six miniatures in the genus that has an adipose fin (MENDONÇA *et al.*, 2016). In addition to *Elachocharax*, *Priocarax* and *Xenurobrycon* Myers & Miranda-Ribeiro of which the majority of species lack an adipose fin, the following miniature genera are currently composed solely of species lacking the adipose fin: *Dundocharax* Poll, *Iotabrycon* Roberts and *Scopaeocharax* Weitzman & Fink, and two other genera of Characidae are composed exclusively of miniature species, some of them also lacking the adipose fin: *Tytobrycon* Géry and *Tytocharax* Fowler (see Table 4 for details).

In some other genera that are not composed solely of miniature species *sensu* WEITZMAN & VARI (1988), the adipose-fin is absent in both miniature and some non-miniature taxa. Such is the case in *Carnegiella* Eigenmann, *Characidium* Reinhardt, *Nannostomus* Günther, and *Neolebias* Steindachner (Table 4). Moreover, the lack of the adipose fin is not exclusive to genera with miniature species. There are small to medium sized characiforms that also lack this fin such as the two species of the crenuchid *Poecilocharax* Eigenmann, the distichodontid *Nannocharax* Günther, the lebiasinids *Copeina* Fowler, *Copella* Myers, *Pyrrhulina* Valenciennes and some species of *Lebiasina* Valenciennes (Table 4). Among characids, the adipose fin is lacking in species of *Coptobrycon* Géry, *Corynopoma* Gill, *Grundulus* Valenciennes, *Gymnocharacinus* Steindachner, *Hasemania*

Ellis, *Hyphessobrycon* Durbin, *Monotocheirodon* Eigenmann, *Nematobrycon* Eigenmann, *Phenagoniates* Eigenmann & Wilson, and *Spintherobolus* Eigenmann (Table 4). Finally, the adipose fin is absent in all 17 species of the Erythrinidae (Table 4), including some of the largest characiforms (MATTOX *et al.*, 2006; OYAKAWA & MATTOX, 2009; 2018). This is evidence that the loss of the adipose fin is not necessarily related to small size.

Intraspecific variation in the presence of the adipose fin and its development was mentioned for some characiforms. WEITZMAN & COBB (1975) cited intraspecific variation in *Nannostomus eques* and *N. trifasciatus*, TRAVASSOS (1949) in *Characidium lauroi*, BUCKUP & HAHN (2000) in *C. vestigipinne* and MELO *et al.* (2016) in *C. oiticicai*. GÉRY (1977) reported variation in the presence/absence of the adipose fin in the African *Neolebias trilineatus* and *N. unifasciatus*. Among the Characidae, intraspecific variation in the size or presence of the adipose fin was reported for *Hasemania nana* (GÉRY, 1977:518; DAGOSTA *et al.*, 2014) *Hyphessobrycon diastatos* (DAGOSTA *et al.*, 2014), *H. negodagua* (LIMA & GERHARD, 2001), *H. eilyos* (LIMA & MOREIRA, 2003), and *H. uaiso* (CARVALHO & LANGEANI, 2013). All examined specimens of *Priocarax variii* ($n=50$) consistently have a small adipose fin and no intraspecific variation was observed.

The molecular analysis from samples of all four taxa supports the conclusion that each represents a species

Table 4. List of species of Characiformes that lack the adipose fin, organized in families according to Mirande (2018). Families and species are listed in alphabetical order. The second column refer to number of species lacking the adipose fin in relation to the total number of species in the genus. References are those that summarize information on the lack of the adipose fin for a species or group of species. Species in bold are miniature according to Weitzman & Vari (1987), with Neotropical fishes updated according to Toledo-Piza *et al.* (2014) or described after that. Species with intraspecific variation in the absence/presence of the adipose fin are not listed here, but are mentioned in the text.

| TAXON | | REFERENCE |
|--|-------------|---------------------------------------|
| Characidae | | |
| <i>Coptobrycon bilineatus</i> (Ellis, 1911) | 1/1 | GÉRY, 1977 |
| <i>Corynopoma riisei</i> Gill, 1858 | 1/1 | GÉRY, 1977 |
| <i>Grundulus bogotensis</i> Humboldt, 1821 | 3/3 | ROMÁN-VALENCIA <i>et al.</i> , 2003 |
| <i>Grundulus cochae</i> Román-Valencia <i>et al.</i> , 2003 | 3/3 | ROMÁN-VALENCIA <i>et al.</i> , 2003 |
| <i>Grundulus quitoensis</i> Román-Valencia <i>et al.</i> , 2005 | 3/3 | ROMÁN-VALENCIA <i>et al.</i> , 2005 |
| <i>Gymnocharacinus bergii</i> Steindachner, 1903 | 1/1 | GÉRY, 1977 |
| <i>Hasemania crenuchooides</i> Zarske & Géry, 1999 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania hansenii</i> (Fowler, 1949) | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania kalunga</i> Bertaco & Carvalho, 2010 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania maxillaris</i> Ellis, 1911 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania melanura</i> Ellis, 1911 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania nambiquara</i> Bertaco & Malabarba, 2007 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania piatan</i> Zanata & Serra, 2010 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hasemania uberaba</i> Serra & Langeani, 2015 | 8/9 | SERRA & LANGEANI, 2015 |
| <i>Hypessobrycon taurocephalus</i> Ellis, 1911 | 1/161 | LIMA & GERHARD, 2001 |
| <i>Iotabrycon praecox</i> Roberts, 1973 | 1/1 | ROBERTS, 1973 |
| <i>Monotocheirodon drilos</i> Menezes <i>et al.</i> , 2013 | 3/3 | MENEZES <i>et al.</i> , 2013 |
| <i>Monotocheirodon kontos</i> Menezes <i>et al.</i> , 2013 | 3/3 | MENEZES <i>et al.</i> , 2013 |
| <i>Monotocheirodon pearsoni</i> Eigenmann, 1924 | 3/3 | MENEZES <i>et al.</i> , 2013 |
| <i>Nematobrycon lacortei</i> Weitzman & Fink, 1971 | 2/2 | GÉRY, 1977 |
| <i>Nematobrycon palmeri</i> Eigenmann, 1911 | 2/2 | GÉRY, 1977 |
| <i>Phenagoniates macrolepis</i> (Meek & Hildebrand, 1913) | 1/1 | GÉRY, 1977 |
| <i>Priocharax ariel</i> Weitzman & Vari, 1987 | 3/4 | WEITZMAN & VARI, 1987 |
| <i>Priocharax nanus</i> Toledo-Piza <i>et al.</i> , 2014 | 3/4 | TOLEDO-PIZA <i>et al.</i> , 2014 |
| <i>Priocharax pygmaeus</i> Weitzman & Vari, 1987 | 3/4 | WEITZMAN & VARI, 1987 |
| <i>Scopaeocharax atopodus</i> (Böhlke, 1958) | 2/2 | WEITZMAN & FINK, 1985 |
| <i>Scopaeocharax rhinodus</i> (Böhlke, 1958) | 2/2 | WEITZMAN & FINK, 1985 |
| <i>Spintherobolus ankoseion</i> Weitzman & Malabarba, 1999 | 4/4 | WEITZMAN & MALABARBA, 1999 |
| <i>Spintherobolus broccae</i> Myers, 1925 | 4/4 | WEITZMAN & MALABARBA, 1999 |
| <i>Spintherobolus leptoura</i> Weitzman & Malabarba, 1999 | 4/4 | WEITZMAN & MALABARBA, 1999 |
| <i>Spintherobolus papilliferus</i> Eigenmann, 1911 | 4/4 | WEITZMAN & MALABARBA, 1999 |
| <i>Tyttocharax hamatus</i> Géry, 1973 | 2/6 | MARINHO <i>et al.</i> , 2013 |
| <i>Tyttocharax spinosus</i> Géry, 1973 | 2/6 | MARINHO <i>et al.</i> , 2013 |
| <i>Tyttocharax metae</i> Román-Valencia <i>et al.</i> , 2012 | 2/4 | ROMÁN-VALENCIA <i>et al.</i> , 2012 |
| <i>Tyttocharax tambopatensis</i> Weitzman & Ortega, 1995 | 2/4 | WEITZMAN & ORTEGA, 1995 |
| <i>Xenurobrycon coracorainae</i> Moreira, 2005 | 5/6 | MENDONÇA <i>et al.</i> , 2016 |
| <i>Xenurobrycon heterodon</i> Weitzman & Fink, 1985 | 5/6 | MENDONÇA <i>et al.</i> , 2016 |
| <i>Xenurobrycon macropus</i> Myers & Miranda Ribeiro, 1945 | 5/6 | MENDONÇA <i>et al.</i> , 2016 |
| <i>Xenurobrycon polyancistrus</i> Weitzman, 1987 | 5/6 | MENDONÇA <i>et al.</i> , 2016 |
| <i>Xenurobrycon variii</i> Mendonça <i>et al.</i> , 2016 | 5/6 | MENDONÇA <i>et al.</i> , 2016 |
| | | |
| Crenuchidae | | |
| <i>Characidium chicoi</i> Graça <i>et al.</i>, 2019 | 9/73 | GRAÇA <i>et al.</i>, 2019 |
| <i>Characidium helmeri</i> Zanata <i>et al.</i> , 2015 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Characidium mirim</i> Netto-Ferreira <i>et al.</i>, 2013 | 9/73 | MELO <i>et al.</i>, 2016 |
| <i>Characidium nana</i> Mendonça & Netto-Ferreira, 2015 | 9/73 | MELO <i>et al.</i>, 2016 |
| <i>Characidium nupelia</i> Graça <i>et al.</i> , 2008 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Characidium sterbai</i> Zarske, 1997 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Characidium stigmosum</i> Melo & Buckup, 2002 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Characidium travassosi</i> Melo <i>et al.</i> , 2016 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Characidium xavante</i> Graça <i>et al.</i> , 2008 | 9/73 | MELO <i>et al.</i> , 2016 |
| <i>Elachocharax geryi</i> Weitzman & Kanazawa, 1978 | 3/4 | BUCKUP, 1993 |

Table 4 continued.

| Taxon | | Reference |
|---|-------|--------------------------------------|
| Crenuchidae | | |
| <i>Elachocharax junki</i> (Géry, 1971) | 3/4 | BUCKUP, 1993 |
| <i>Elachocharax mitopterus</i> Weitzman, 1986 | 3/4 | BUCKUP, 1993 |
| <i>Poecilocharax bovaliorum</i> Eigenmann, 1909 | 2/2 | BUCKUP & VAN DER SLEEN, 2018 |
| <i>Poecilocharax weitzmani</i> Géry, 1965 | 2/2 | BUCKUP & VAN DER SLEEN, 2018 |
| Distichodontidae | | |
| <i>Dundocharax bidentatus</i> Poll, 1967 | 1/1 | GÉRY, 1977 |
| <i>Nannocharax machadoi</i> | 1/41 | GÉRY, 1977 |
| <i>Neolebias ansorgii</i> Boulenger, 1912 | 5/12 | GÉRY, 1977 |
| <i>Neolebias axelrodi</i> Poll & Gosse, 1963 | 5/12 | GÉRY, 1977 |
| <i>Neolebias lozii</i> Winemiller & Kelson-Winemiller, 1993 | 5/12 | WINEMILLER & KELSON-WINEMILLER, 1993 |
| <i>Neolebias philippei</i> Poll & Gosse, 1963 | 5/12 | GÉRY, 1977 |
| <i>Neolebias powelli</i> Teugels & Roberts, 1990 | 5/12 | TEUGELS & ROBERTS, 1990 |
| Erythrinidae | | |
| <i>Erythrinus erythrinus</i> (Bloch & Schneider, 1801) | 2/2 | OYAKAWA & MATTOX, 2018 |
| <i>Erythrinus kessleri</i> Steindachner, 1877 | 2/2 | OYAKAWA & MATTOX, 2018 |
| <i>Hopleriethrinus cinereus</i> (Gill, 1858) | 3/3 | OYAKAWA & MATTOX, 2018 |
| <i>Hopleriethrinus gronovii</i> (Valenciennes, 1847) | 3/3 | OYAKAWA & MATTOX, 2018 |
| <i>Hopleriethrinus unitaeniatus</i> (Spix & Agassiz, 1829) | 3/3 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas aimara</i> (Valenciennes, 1847) | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas argentinensis</i> Rosso <i>et al.</i> , 2018 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas australis</i> Oyakawa & Mattox, 2009 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas brasiliensis</i> (Spix & Agassiz, 1829) | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas curupira</i> Oyakawa & Mattox, 2009 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas intermedius</i> (Günther, 1864) | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas lacerdae</i> Miranda Ribeiro, 1908 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas malabaricus</i> (Bloch, 1794) | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas mbigua</i> Azpelicueta <i>et al.</i> , 2015 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas microlepis</i> (Günther, 1864) | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas misionera</i> Rosso <i>et al.</i> , 2016 | 12/12 | OYAKAWA & MATTOX, 2018 |
| <i>Hoplitas teres</i> (Valenciennes, 1847) | 12/12 | OYAKAWA & MATTOX, 2018 |
| Gasteropelecidae | | |
| <i>Carnegiella marthae</i> Myers, 1927 | 4/4 | WEITZMAN, 1960 |
| <i>Carnegiella myersi</i> Fernández-Yépez, 1950 | 4/4 | WEITZMAN, 1960 |
| <i>Carnegiella schereri</i> Fernández-Yépez, 1950 | 4/4 | WEITZMAN, 1960 |
| <i>Carnegiella strigata</i> (Günther, 1864) | 4/4 | WEITZMAN, 1960 |
| Lebiasinidae | | |
| <i>Copeina guttata</i> (Steindachner, 1876) | 2/2 | NETTO-FERREIRA, 2018 |
| <i>Copeina osgoodi</i> Eigenmann, 1912 | 2/2 | NETTO-FERREIRA, 2018 |
| <i>Copella arnoldi</i> (Regan, 1912) | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Copella callolepis</i> (Regan, 1912) | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Copella compita</i> (Myers, 1927) | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Copella eigenmanni</i> (Regan, 1912) | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Copella nattereri</i> (Steindachner, 1876) | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Copella vilmae</i> Géry, 1963 | 6/6 | MARINHO & MENEZES, 2017 |
| <i>Lebiasina bimaculata</i> Valenciennes, 1847 | 7/26 | NETTO-FERREIRA, 2010 |
| <i>Lebiasina marilynae</i> Netto-Ferreira, 2012 | 7/26 | NETTO-FERREIRA, 2012 |
| <i>Lebiasina melanoguttata</i> Netto-Ferreira, 2012 | 7/26 | NETTO-FERREIRA, 2012 |
| <i>Lebiasina minuta</i> Netto-Ferreira, 2012 | 7/26 | NETTO-FERREIRA, 2012 |
| <i>Lebiasina multimaculata</i> Boulenger, 1911 | 7/26 | NETTO-FERREIRA, 2010 |
| <i>Lebiasina taphorni</i> Ardila Rodriguez, 2004 | 7/26 | ARDILA RODRIGUEZ, 2004 |
| <i>Lebiasina uruyensis</i> Fernández-Yépez, 1967 | 7/26 | NETTO-FERREIRA, 2010 |
| <i>Nannostomus anduzei</i> Fernandez & Weitzman, 1987 | 6/20 | FERNANDEZ & WEITZMAN, 1987 |
| <i>Nannostomus beckfordi</i> Günther, 1872 | 6/20 | WEITZMAN & COBB, 1975 |
| <i>Nannostomus bifasciatus</i> Hoedeman, 1954 | 6/20 | WEITZMAN & COBB, 1975 |
| <i>Nannostomus limatus</i> Weitzman, 1978 | 6/20 | WEITZMAN, 1978 |

Table 4 continued.

| TAXON | | REFERENCE |
|---|-------------|----------------------------------|
| <i>Nannostomus marginatus</i> Eigenmann, 1909 | 6/20 | WEITZMAN & COBB, 1975 |
| <i>Nannostomus minimus</i> Eigenmann, 1909 | 6/20 | WEITZMAN & COBB, 1975 |
| <i>Pyrrhulina australis</i> Eigenmann & Kennedy 1903 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina beni</i> Pearson 1924 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina brevis</i> Steindachner 1876 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina capim</i> Vieira & Netto-Ferreira 2019 | 19/19 | VIEIRA & NETTO-FERREIRA 2019 |
| <i>Pyrrhulina eleanorae</i> Fowler 1940 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina elongata</i> Zarske & Géry 2001 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina filamentosa</i> Valenciennes 1847 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina laeta</i> (Cope 1872) | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina lugubris</i> Eigenmann 1922 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina marilynae</i> Netto-Ferreira & Marinho 2013 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina maxima</i> Eigenmann & Eigenmann 1889 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina melanostomus</i> (Cope 1870) | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina obermulleri</i> Myers 1926 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina rachoviana</i> Myers 1926 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina semifasciata</i> Steindachner 1876 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina spilota</i> Weitzman 1960 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina stoli</i> Boeseman 1953 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina vittata</i> Regan 1912 | 19/19 | NETTO-FERREIRA, 2018 |
| <i>Pyrrhulina zigzag</i> Zarske & Géry 1997 | 19/19 | NETTO-FERREIRA, 2018 |

(Figure 5). The genetic distance between the four species ranges from 20.7% between *P. nanus* and *P. varii* to 30.5% between *P. ariel* and *P. pygmaeus*, a high genetic divergence compared with other Neotropical fishes which have an average of 6.8% (e.g., CARVALHO *et al.*, 2011; PEREIRA *et al.*, 2011a; 2013). Considering only characiforms, the range in genetic divergence between species is 1–22% (BENINE *et al.*, 2009; MELO *et al.*, 2011; PEREIRA *et al.*, 2011b; BELLAFRONTI *et al.*, 2013; CASTRO PAZ *et al.*, 2014; SILVA *et al.*, 2013; RAMIREZ & GALETTI JR, 2015; ROSSINI *et al.*, 2016).

Two hypotheses could explain the higher genetic differences in *Priocharax*: either its species have been evolving separately for a long time and have accumulated a large amount of mutations, or they have an accelerated evolutionary rate due to intrinsic features (e.g., CASTRO PAZ *et al.*, 2014). The first hypothesis predicts that *Priocharax* has a long evolutionary and biogeographic history, something that needs to be tested in the future when more data on the taxonomy and phylogeny of the genus are available. The second hypothesis predicts that *Priocharax* has biological features that would increase its evolutionary rate. In fact, smaller organisms tend to have higher nucleotide substitution rates (GILLOOLY *et al.*, 2005) because of their shorter generation time and therefore accumulate more genetic divergence compared to larger organisms in the same time period. ROBERTS (1984) already pointed out that miniature fishes could reproduce year-round continuously increasing the number of generations per time unit and, therefore, their mutation rate. The hypothesis raised by ROBERTS (1984) is particularly interesting and should be tested for *Priocharax* and other miniature taxa. The results for the miniatur-

ized and highly developmentally truncated cypriniforms *Danionella* and *Paedocypris* which both sit at the end of very long branches in molecular phylogenetic trees (see Supplementary figure 7 in BRITZ *et al.*, 2014) is a further example of a higher mutation rate in highly miniaturized fish taxa.

Acknowledgements

Most of this study was conducted at Departamento de Biologia, UFSCar – Universidade Federal de São Carlos, which provided space and access to facilities. The molecular analyses were conducted at Departamento de Biologia Estrutural e Funcional, UNESP Botucatu. The authors are thankful for S. C. P. Souza (DBio-So/UFSCar) who provided valuable help during fieldwork and actually found the first specimens of *Priocharax varii*. W. M. Ohara (UFRO) provided information for the fieldwork. R. G. M. Ferreira (FIOCRUZ) gave valuable suggestions to the locality where we sampled the new species. C. R. C. Doria (UFRO) sent specimens on loan and provided formalin and alcohol for the expedition, to whom the authors are very grateful. B. M. Corazza, E. Silva, K. E. Reis (UFSCar) and M. M. F. Marinho (UFPB) aided in species identification. M. Gianeti and O. T. Oyakawa (MZUSP), and A. Andriolo (UFRO) provided curatorial assistance. F. C. T. Lima (ZUEC), H. Ortega (MHN-UNMSM), M. R. S. Melo (IO-USP) and R. R. Ota (UEM) sent publications and/or information regarding the adipose fin. GMTM acknowledges financial support from FAPESP (Proc 2017/01970-4), CS acknowledges financial support from FAPESP (Proc 2017/06551-0), and CO acknowledges financial support from FAPESP (Proc 2018/20610-1, 2016/09204-6, 2014/26508-3) and CNPq (Proc 306054/2006-0).

References

- ARDILA RODRÍGUEZ, C. A. (2004). *Lebiasina taphorni* (Pisces: Characiformes: Lebiasinidae), una nueva especie. *Dahlia*, **7**, 57–65.
- BELAFRONTÉ, E., MARIGUELA, T. C., PEREIRA, L. H. G., OLIVEIRA, C. & MOREIRA-FILHO, O. (2013). DNA barcode of Parodontidae species from the La Plata river basin – applying new data to clarify taxonomic problems. *Neotropical Ichthyology*, **11**, 497–506.
- BENDER, A. & MORITZ, T. (2013). Developmental residue and developmental novelty – different modes of adipose-fin formation during ontogeny. *Zoosystematics and Evolution*, **89**, 209–214.
- BENINE, R. C., MARIGUELA, T. C. & OLIVEIRA, C. (2009). New species of *Moenkhausia* Eigenmann, 1903 (Characiformes: Characidae) with comments on the *Moenkhausia oligolepis* species complex. *Neotropical Ichthyology*, **7**, 161–168.
- BETANCUR-R, R., ARCILA, D., VARI, R. P., HUGHES, L. C., OLIVEIRA, C., SABAJ, M. H. & ORTÍ, G. (2018). Phylogenomic incongruence, hypothesis testing, and taxonomic sampling: The monophyly of characiform fishes. *Evolution*, **73**, 329–345.
- BRITZ, R., CONWAY, K. W. & RÜBER, L. (2014). Miniatures, morphology and molecules: *Paedocypris* and its phylogenetic position (Teleostei, Cypriniformes). *Zoological Journal of the Linnean Society*, **172**, 556–615.
- BUCKUP, P. A. (1993). Review of the characidiini fishes (Teleostei, Characiformes), with descriptions of four new genera and ten new species. *Ichthyological Exploration of Freshwaters*, **4**, 97–154.
- BUCKUP, P. A. & HAHN, L. (2000). *Characidium vestigipinne*: a new species of Characidiinae (Teleostei, Characiformes) from Southern Brazil. *Copeia*, **2000**, 150–155.
- BUCKUP, P. A. & VAN DER SLEEN, P. (2018). Family Crenuchidae – South American darters, pp. 142–148 in: VAN DER SLEEN, P. & ALBERT, J. S. (eds) *Field guide to the fishes of the Amazon, Orinoco & Guianas*. Princeton, NJ: Princeton University Press.
- CARVALHO, D. C., OLIVEIRA, D. A. A., POMPEU, P. S., LEAL, C. G., OLIVEIRA, C. & HANNER, R. (2011). Deep barcode divergence in Brazilian freshwater fishes: the case of the São Francisco River basin. *Mitochondrial DNA*, **22**, 80–86.
- CARVALHO, F. R. & LANGEANI, F. (2013). *Hypessobrycon uaiso*: a new characid fish from the rio Grande, upper rio Paraná basin, Minas Gerais State (Ostariophysi: Characidae), with a brief comment about some types of *Hypessobrycon*. *Neotropical Ichthyology*, **11**, 525–536.
- CARVALHO, L. N., ZUANON, J. & SAZIMA, I. (2006). The almost invisible league: crypsis and association between minute fishes and shrimps as a possible defence against visually hunting predators. *Neotropical Ichthyology*, **4**, 219–224.
- CASTRO PAZ, F. P., BATISTA, J. S. & PORTO, J. I. R. (2014). DNA barcodes of Rosy Tetras and allied species (Characiformes: Characidae: *Hypessobrycon*) from the Brazilian Amazon basin. *PLoS ONE*, **9**, e98603.
- CONWAY, K. W. & BRITZ, R. (2007). Sexual dimorphism of the Weberian apparatus and pectoral girdle in *Sundadanio axelrodi* (Ostariophysi: Cyprinidae), a miniature cyprinid from South East Asia. *Journal of Fish Biology*, **71**, 1562–1570.
- DAGOSTA, F. C. P., MARINHO, M. M. F. & CAMELIER, P. (2014). A new species of *Hypessobrycon* Durbin (Characiformes: Characidae) from the middle rio São Francisco and upper rio Tocantins basins, Brazil, with comments on its biogeographic history. *Neotropical Ichthyology*, **12**, 365–375.
- FERNANDEZ, J. M. & WEITZMAN, S. H. (1987). A new species of *Nannostomus* (Teleostei: Lebiasinidae) from near Puerto Ayacucho, Río Orinoco drainage, Venezuela. *Proceedings of the Biological Society of Washington*, **100**, 164–172.
- FINK, W. L. & WEITZMAN, S. H. (1974). The so-called cheirodontin fishes of Central America with descriptions of two new species (Pisces: Characidae). *Smithsonian Contributions to Zoology*, **172**, 1–46.
- FUIMAN, L. A. (1984). Ostariophysi: development and relationships, pp. 126–137 in: MOSER, H. G., RICHARDS, W. J., COHEN, D. M., FAHAY, M. P., KENDALL JR., A. W. & RICHARDSON, S. L. (eds) *Ontogeny and Systematics of Fishes*. Lawrence, KS: Allen Press.
- GÉRY, J. (1977). *Characoids of the world*. Neptune City, NJ: T.F.H. Publications, Inc.
- GILLOOLY, J. F., ALLEN, A. P., WEST, G. B. & BROWN, J. H. (2005). The rate of DNA evolution: effects of body size and temperature on the molecular clock. *Proceedings of the National Academy of Sciences*, **102**, 140–145.
- GRAÇA, W. J., OTA, R. R. & DOMINGUES, W. M. (2019). A new species of miniature *Characidium* (Characiformes: Characidae) from the upper Rio Paraguay basin, Mato Grosso state, Brasil. *Journal of Fish Biology*, **95**, 1480–1485.
- IVANOVA, N. V., DEWAARD, J. R. & HEBERT, P. D. (2006). An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Molecular Ecology Notes*, **6**, 998–1002.
- KEARSE, M., MOIR, R., WILSON, A., STONES-HAVAS, S., CHEUNG, M., STURROCK, S., et al. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, **28**, 1647–1649.
- KIMURA, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, **16**, 111–120.
- LIMA, F. C. T. & GERHARD, P. (2001). A new *Hypessobrycon* (Characiformes: Characidae) from Chapada Diamantina, Bahia, Brazil, with notes on its natural history. *Ichthyological Exploration of Freshwaters*, **12**, 105–114.
- LIMA, F. C. T. & MOREIRA, C. R. (2003). Three new species of *Hypessobrycon* (Characiformes, Characidae) from the upper Rio Araguaia basin in Brazil. *Neotropical Ichthyology*, **1**, 21–33.
- MARINHO, M. M. F. (2017). Comparative development in *Moenkhausia pittieri* and *Paracheirodon inesi* (Ostariophysi, Characiformes) with comments on heterochrony and miniaturization in the Characidae. *Journal of Fish Biology*, **91**, 851–865.
- MARINHO, M. M. F. & MENEZES, N. A. (2017). Taxonomic review of *Copella* (Characiformes: Lebiasinidae) with an identification key for the species. *PLoS ONE*, **12**, e0183069.
- MARINHO, M. M. F., BASTOS, D. A. & MENEZES, N. A. (2013). New species of miniature fish from Marajó Island, Pará, Brazil, with comments on its relationships (Characiformes: Characidae). *Neotropical Ichthyology*, **11**, 739–746.
- MATTOX, G. M. T. & TOLEDO-PIZA, M. (2012). Phylogenetic study of the Characinae (Teleostei: Characiformes: Characidae). *Zoological Journal of the Linnean Society*, **165**, 809–915.
- MATTOX, G. M. T., BRITZ, R. & TOLEDO-PIZA, M. (2014). Skeletal development and ossification sequence of the characiform *Salminus brasiliensis* (Teleostei: Ostariophysi: Characidae). *Ichthyological Exploration of Freshwaters*, **25**, 103–158.
- MATTOX, G. M. T., BRITZ, R., TOLEDO-PIZA, M. & MARINHO, M. M. F. (2013). *Cyanogaster noctivaga*, a remarkable new genus and species of miniature fish from the Rio Negro, Amazon basin (Ostariophysi: Characidae). *Ichthyological Exploration of Freshwaters*, **23**, 297–318.
- MATTOX, G. M. T., TOLEDO-PIZA, M. & OYAKAWA, O. T. (2006). Taxonomic study of *Hoplias aimara* (Valenciennes, 1846) and *Hoplias macrourhthalmus* (Pellegrin, 1907) (Ostariophysi, Characiformes, Erythrinidae). *Copeia*, **2006**, 516–528.
- MELO, B. F., BENINE, R. C., MARIGUELA, T. C. & OLIVEIRA, C. (2011). A new species of *Tetragonopterus* Cuvier, 1816 (Characiformes: Characidae: Tetragonopterinae) from the rio Jari, Amapá, northern Brazil. *Neotropical Ichthyology*, **9**, 49–56.
- MELO, M. R. S., BUCKUP, P. A. & OYAKAWA, O. T. (2016). A new species of *Characidium* Reinhardt, 1867 (Characiformes, Crenuchidae) endemic to the Atlantic Forest in Paraná State, southern Brazil. *Neotropical Ichthyology*, **14**, 533–544.
- MENDONÇA, M. B., PEIXOTO, L. A. W., DUTRA, G. M. & NETTO-FERREIRA, A. L. (2016). A new miniature Xenurobryconini (Characiformes: Characidae) from the upper Rio Paraguay basin, Mato Grosso state, Brazil. *Neotropical Ichthyology*, **14**, 533–544.

- aciformes: Characidae) from the rio Tapajós basin, Brazil. *Neotropical Ichthyology*, **14**, 45–52.
- MENEZES, N. A. & WEITZMAN, S. H. (1990). Two new species of *Migmagoniates* (Teleostei: Characidae: Glandulocaudinae), their phylogeny and biogeography and a key to the glandulocaudin fishes of Brazil and Paraguay. *Proceedings of the Biological Society of Washington*, **103**, 380–426.
- MENEZES, N. A., WEITZMAN, S. H. & QUAGIO-GRASSIOTTO, I. (2013). Two new species and a review of the inseminating freshwater fish genus *Monotocheirodon* (Characiformes: Characidae) from Peru and Bolivia. *Papéis Avulsos de Zoologia*, **53**, 129–144.
- MIRANDE, J. M. (2010). Phylogeny of the family Characidae (Teleostei: Characiformes): from characters to taxonomy. *Neotropical Ichthyology*, **8**, 385–568.
- MIRANDE, J. M. (2018). Morphology, molecules and the phylogeny of Characidae (Teleostei, Characiformes). *Cladistics*, **2018**, 1–19.
- NETTO-FERREIRA, A. L. (2010). *Revisão taxonómica e relações interespecíficas de Lebiasininae (Ostariophysi: Characiformes: Lebiasinidae)*. Unpublished PhD Thesis. Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil.
- NETTO-FERREIRA, A. L. (2012). Three new species of *Lebiasina* (Characiformes: Lebiasinidae) from the Brazilian Shield border at Serra do Cachimbo, Pará, Brazil. *Neotropical Ichthyology*, **10**, 487–498.
- NETTO-FERREIRA, A. L. (2018). Family Lebiasinidae – pencilfishes, pp. 165–169 in: VAN DER SLEEN, P. & ALBERT, J. S. (eds) *Field guide to the fishes of the Amazon, Orinoco & Guianas*. Princeton, NJ: Princeton University Press.
- OLIVEIRA, C., AVELINO, G. S., ABE, K. T., MARIGUELA, T. C., BENINE, R. C., ORTÍ, G., VARI, R. P. & CASTRO, R. M. C. (2011). Phylogenetic relationships within the speciose family Characidae (Teleostei: Ostariophysi: Characiformes) based on multilocus analysis and extensive ingroup sampling. *BMC Evolutionary Biology*, **11**, 1–25.
- OYAKAWA, O. T. & MATTOX, G. M. T. (2009). Revision of the Neotropical trahiras of the *Hoplias lacerdae* species-group (Ostariophysi: Characiformes: Erythrinidae) with descriptions of two new species. *Neotropical Ichthyology*, **7**, 117–140.
- OYAKAWA, O. T. & MATTOX, G. M. T. (2018). Family Erythrinidae – wolf-fishes and yarrows, pp. 156–158 in: VAN DER SLEEN, P. & ALBERT, J. S. (eds) *Field guide to the fishes of the Amazon, Orinoco & Guianas*. Princeton, NJ: Princeton University Press.
- PEREIRA, L. H. G., MAIA, G. M. G., HANNER, R., FORESTI, F. & OLIVEIRA, C. (2011a). DNA barcodes discriminate freshwater fishes from the Paráíba do Sul River Basin, São Paulo, Brazil. *Mitochondrial DNA*, **22**, 71–79.
- PEREIRA, L. H. G., PAZIAN, M. F., HANNER, R., FORESTI, F. & OLIVEIRA, C. (2011b). DNA barcoding reveals hidden diversity in the Neotropical freshwater fish *Piabina argentea* (Characiformes: Characidae) from the Upper Paraná Basin of Brazil. *Mitochondrial DNA*, **22**, 87–96.
- PEREIRA, L. H. G., HANNER, R., FORESTI, F. & OLIVEIRA, C. (2013). Can DNA barcoding accurately discriminate megadiverse Neotropical freshwater fish fauna? *BMC Genetics*, **14**, 1–14.
- QUEIROZ, L. J., TORRENTE-VILARA, G., OHARA, W. M., PIRES, T. H. S., ZUANON, J. & DORIA, C. R. C. (2003). *Peixes do Rio Madeira, vol I*. São Paulo: Dialetô Latin American Documentary.
- RAMIREZ, J. L. & GALETTI JR, P. M. (2015). DNA barcode and evolutionary relationship within *Laemolyta* Cope 1872 (Characiformes: Anostomidae) through molecular analyses. *Molecular Phylogenetics and Evolution*, **93**, 77–82.
- ROBERTS, T. R. (1973). The glandulocaudine characid fishes of the Guayas Basin in western Ecuador. *Bulletin of the Museum of Comparative Zoology*, **144**, 489–514.
- ROBERTS, T. R. (1984). *Amazonsprattus scintilla*, new genus and species from the Rio Negro, Brazil, the smallest known clupeomorph fish. *Proceedings of the California Academy of Sciences*, **43**, 317–321.
- ROMAN-VALENCIA, C., PAEPKE, H.-J. & PANTOJA, F. (2003). Una especie nueva de *Grundulus* (Teleostei: Ostariophysi: Characidae) para Colombia y redescrición de *Grundulus bogotensis* (Humboldt y Valenciennes, 1833). *Memoria de la Fundación La Salle de Ciencias Naturales*, **155**, 51–72.
- ROMAN-VALENCIA, C., RUIZ-C., R. I. & BARRIGA, R. (2005). Una nueva especie ecuatoriana del género de peces andinos *Grundulus* (Characiformes: Characidae). *Revista de Biología Tropical*, **53**, 537–544.
- ROMAN-VALENCIA, C., GARCIA-ALZATE, C. A., RUIZ-C., R. I. & TAP-HORN, D. B. (2012). A new species of *Tyttocharax* (Characiformes: Characidae: Stevardiinae) from the Güejar River, Orinoco River Basin, Colombia. *Neotropical Ichthyology*, **10**, 519–525.
- ROSSINI, B. C., OLIVEIRA, C. A. M., MELO, F. A. G., BERTACO, V. A., ASTARLOA, J. M. D., ROSSO, J. J., FORESTI, F. & OLIVEIRA, C. (2016). Highlighting *Astyanax* species diversity through DNA barcoding. *PLoS ONE*, **11**, e0167203.
- SERRA, J. P. & LANGEANI, F. (2015). A new *Hasemania* Ellis from the upper rio Paraná basin, with the redescription of *Hasemania crenuchoides* Zarske & Géry (Characiformes: Characidae). *Neotropical Ichthyology*, **13**, 479–486.
- SILVA, G. S. C., MELO, B. F., OLIVEIRA, C. & BENINE, R. C. (2013). Morphological and molecular evidence for two new species of *Tetragonopterus* (Characiformes, Characidae) from central Brazil. *Journal of Fish Biology*, **82**, 1613–1631.
- TAMURA, K., STECHER, G., PETERSON, D., FILIPSKI, A. & KUMAR, S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*, **30**, 2725–2729.
- TAYLOR, W. R. & VAN DYKE, G. C. (1985). Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. *Cybium*, **9**, 107–109.
- TEUGELS, G. G. & ROBERTS, T. R. (1990). Description of a small distinctively coloured new species of the characoid genus *Neolebias* from the Niger delta, West Africa (Pisces; Distichodontidae). *Journal of African Zoology*, **104**, 61–67.
- TOLEDO-PIZA, M., MATTOX, G. M. T. & BRITZ, R. (2014). *Priocharax nanus*, a new miniature characid from the Rio Negro, Amazon basin (Ostariophysi: Characiformes), with an updated list of Neotropical freshwater miniature fishes. *Neotropical Ichthyology*, **12**, 229–246.
- TRAVASSOS, H. (1949). Notas ictiológicas I. *Characidium lauroi* n. sp. (Actinopterygii, Ostareophysi). *Revista Brasileira de Biologia*, **9**, 87–92.
- VIEIRA, F. G., MATSUZAKI, A. A., BARROS, B. S. F., OHARA, W. M., PAIXÃO, A. C., TORRENTE-VILARA, G., ZUANON, J. & DORIA, C. R. C. (2016). *Catálogo de Peixes da ESEC Cuniã*. Porto Velho: Edufro.
- VIEIRA, L. S. & NETTO-FERREIRA, A. L. (2019). New species of *Pyrrhulina* (Teleostei: Characiformes: Lebiasinidae) from the eastern Amazon, Pará, Brazil. *Neotropical Ichthyology*, **17**, e190013[1–8].
- WARD, R. D., ZEMLAK, T. S., INNES, B. H., LAST, P. R. & HEBERT, P. D. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**, 1847–1857.
- WEITZMAN, S. H. (1960). Further notes on the relationships and classification of the South American characid fishes of the subfamily Gasteropelecinae. *Stanford Ichthyological Bulletin*, **7**, 217–239.
- WEITZMAN, S. H. (1962). The osteology of *Brycon meeki*, a generalized characid fish, with an osteological definition of the family. *Stanford Ichthyological Bulletin*, **8**, 1–77.
- WEITZMAN, S. H. (1978). Three new species of fished of the genus *Nannostomus* from the Brazilian States of Pará and Amazonas (Teleostei: Lebiasinidae). *Smithsonian Contributions to Zoology*, **263**, 14p.
- WEITZMAN, S. H. & COBB, J. S., (1975). A revision of the South American fishes of the genus *Nannostomus* Günther (Family Lebiasinidae). *Smithsonian Contributions to Zoology*, **186**, 36p.

- WEITZMAN, S. H. & FINK, S. V. (1985). Xenobryconin phylogeny and putative pheromone pumps in Glandulocaudine fishes. *Smithsonian Contributions to Zoology*, **421**, 121p.
- WEITZMAN, S. H. & MALABARBA, L. R. (1999). Systematics of *Spintherobolus* (Teleostei: Characidae: Cheirodontinae) from eastern Brazil. *Ichthyological Exploration of Freshwaters*, **10**, 1–43.
- WEITZMAN, S. H. & ORTEGA, H. (1995). A new species of *Tyttocharax* (Teleostei: Characidae: Glandulocaudinae: Xenobryconini) from the Río Madre de Dios basin of Peru. *Ichthyological Exploration of Freshwaters*, **6**, 129–148.
- WEITZMAN, S. H. & VARI, R. P. (1987). Two new species and a new genus of miniature characid fishes (Teleostei: Characiformes) from Northern South America. *Proceedings of the Biological Society of Washington*, **100**, 640–652.
- WEITZMAN, S. H. & VARI, R. P. (1988). Miniaturization in South American freshwater fishes; an overview and discussion. *Proceedings of the Biological Society of Washington*, **101**, 444–465.
- WINEMILLER, K. O. & KELSO-WINEMILLER, L. C. (1993). Description of a new *Neolebias* (Characiformes: Distichodontidae) from the Upper Zambezi drainage of Zambia. *Copeia*, **1993**, 112–116.
- XIA, X. (2013). DAMBE5: a comprehensive software package for data analysis in molecular biology and evolution. *Molecular Biology and Evolution*, **30**, 1720–1728.
- XIA, X., XIE, Z., SALEM, M., CHEN, L. & WANG, Y. (2003). An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution*, **26**, 1–7.

Zoobank Registration

at <http://zoobank.org>

This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the International Commission on Zoological Nomenclature (ICZN). The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information can be viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org>. The LSID for this publication is as follows:

urn:lsid:zoobank.org:pub:6A49E4E4-429E-4923-B6E6-18A7087CBFF7

Comparative material examined

Priobarax ariel: Brazil: Amazonas, Santa Isabel do Rio Negro, rio Negro basin: MZUSP 39778, 4, 13.5–14.6 mm SL, rio Urubaxi; MZUSP 55099, 8, 12.4–14.2 mm SL, igarapé at São João, near Santa Isabel do Rio Negro; MZUSP 55097, 4 of 6, 12.2–12.7 mm SL, lagoon near Paricatuba; MZUSP 62230, 2 of 4, 15.1–15.2 mm SL, lagoon in island near Paricatuba. Rio Madeira basin: UFRO-ICT 5761, 1, 14.5 mm SL, Uirapurá, Nova Olinda do Norte. UFRO-ICT 19660, 1, 10.9 mm SL, Rio Marmelos upstream from Vila do Escondido, Manicoré. UFRO-ICT 20954, 8, 10.1–11.8 mm SL, rocks on the right margin of Rio Marmelos, Manicoré. Rondônia: UFRO-ICT 2151, 11, 13.7–16.1 mm SL (4 c&s, 13.7–16.1 mm SL), Lago Madalena, mouth of Rio Jaciparaná, Porto Velho. UFRO-ICT 5744, 3, 12.1–12.5 mm SL, unnamed tributary of Rio Machado, Porto Velho. UFRO-ICT 5745, 3, 11.6–12.9 mm SL, Lago Cuniã, Porto Velho. UFRO-ICT 8449, 1, 17.1 mm SL, Rio Jaciparaná, tributary of Rio Madeira, Porto Velho. UFRO-ICT 13825, 1, 14.2 mm SL, Igarapé C, ESEC Cuniã, Porto Velho. UFRO-ICT 17412, 2, 11.6–12.4 mm SL, Arrozal, Lago Cuniã, Porto Velho. UFRO-ICT 20075, 1, 17.4 mm SL, Igarapé A, ESEC Cuniã, Porto Velho. Venezuela: Territorio Federal Amazonas: MZUSP 36497, 50, 11.8–15.2 mm SL, MZUSP 55142, 12 paratypes, 12.0–14.7 mm SL (5 c&s, 12.0–14.0 mm SL), Caño Manu, tributary of río Casiquiare approximately 250 m upstream from Solano.

Priobarax nanus: MZUSP 114014, 13.8 mm SL, holotype, Brazil, Amazonas, Santa Isabel do Rio Negro, rio Negro basin, lake at right bank of rio Urubaxi, near igarapé Tapage, 0°33'44.2"S 64°49'40.8"W, 26 Oct 2011, M. Toledo-Piza, G. Mattox, M. Marinho & R. Britz. All paratypes: MZUSP 114015, 9,

12.1–15.3 mm SL (3 c&s, 14.1–15.3 mm SL), same data as holotype. MZUSP 114016, 5, 12.6–14.6 mm SL (2 c&s, 13.4–13.8 mm SL), Brazil, Amazonas, Santa Isabel do Rio Negro, rio Negro basin, igarapé Tapage at left bank of rio Urubaxi, 0°30'5.3"S 64°49'11.7"W, 26 Oct 2011, M. Toledo-Piza, G. Mattox, M. Marinho & R. Britz. MZUSP 114017, 3, 13.5–14.6 mm SL (1 c&s, 14.6 mm SL), Brazil, Amazonas, Santa Isabel do Rio Negro, rio Negro basin, first tributary of rio Negro above rio Daraá, 0°27'24.2"S 64°46'54.1"W, 27 Oct 2011, M. Toledo-Piza, G. Mattox, M. Marinho & R. Britz. INPA 39891, 4, 12.5–13.9 mm SL; MZUSP 114018, 11, 11.1–15.4 mm SL (5 c&s, 12.0–14.0 mm SL); USNM 427007, 4, 12.1–13.3 mm SL; Brazil, Amazonas, Santa Isabel do Rio Negro, rio Negro basin, rio Negro and tributaries near Santa Isabel do Rio Negro, 23–30 Oct 2011, M. Toledo-Piza, G. Mattox, M. Marinho & R. Britz.

Priobarax pygmaeus: Colombia: Departamento Amazonas: MZUSP 36498, 5 paratypes, 10.2–10.7 mm SL, Quebrada Parajito, tributary of Quebrada Bacada, tributary of Quebrada Matamata, tributary of río Amazonas, northwest of Leticia, about 04°41'S 69°57'W. Peru: Loreto, Requena, río Ucayali basin: MZUSP 85644, 1, 16.5 mm SL, small "quebrada", tributary of Quebrada Fierro Caño, ca. 4 km North of IIAP (2.7 km east of Jenaro Herrera).

Supplementary material




Fig. 1. Automatic partition of the dataset reporting the number of groups inside the initial and recursive partitions. Red symbols are overlapped by yellow symbols.




Fig. 2. Poisson Tree Processes (PTP) delimitation tests of species using the maximum likelihood tree.

