

## VEGETATION STRUCTURE AND SPECIES COMPOSITION AT THE NORTHERN PERIPHERY OF THE BOUMBA-BEK NATIONAL PARK, SOUTHEASTERN CAMEROON

Vice Clotèxe TAJEUKEM

*Department of Plant Biology, Faculty of Science, University of Yaoundé I*

Evariste FONGNZOSSIE FEDOUNG

*Higher Teacher's Training School for Technical Education (ENSET), University of Douala*

Victor Aimé KEMEUEZE

*Department of Plant Biology, University of Ngaoundéré*

Bernard-Aloys NKONGMENECK

*Department of Plant Biology, Faculty of Science, University of Yaoundé I*

**ABSTRACT** Forest conservation outside protected areas is taking center stage in global conservation discourse. This study was conducted to clarify the species composition, vegetation structure and plant diversity at the northern periphery of the Boumba-Bek National Park, whose timber and Non-Timber Forest Products (NTFPs) have been used by the local peoples and logging companies. A total of 16 transects measuring 5 km in length and 5 m in width were delineated. The survey recorded DBH of all tree individuals above 10 cm. The result shows a total number of 17,583 trees with a density of 439.6 stems/ha (total area = 40 ha). These trees belong to 51 families, 169 genera and 247 species. Shannon-Wiener diversity index ( $H'$ ) ranged from 5.94 to 6.51 and higher diversity was recorded in the Agroforest Zone. The family importance index (FIV) marked highest score for Euphorbiaceae and Combretaceae. The importance index for species (IVI) was higher for *Terminalia superba*, *Musanga cecropioides*, *Anonidium mannii* and *Celtis mildbraedii*. The height-class distribution of the species shows that the majority of trees belong to the height-class of 5–20 m, which accounts for the average of 87.4% of total stems. The diameter-class distribution of the trees shows an inverse J-shape curve. The study concludes that this forest, despite having undergone disturbance in past years driven by logging and agriculture, is relatively rich and diversified.

**Key Words:** Tropical moist forest; Transect survey; Species composition; Forest structure; Plant diversity.

### INTRODUCTION

During the 20th century, the establishment of protected areas became a key element of most tropical forest countries' efforts to achieve biodiversity conservation. This is because most conservationists assumed that biodiversity is best managed in protected areas and other areas where land has not been fragmented due to human activities and population pressure (Molnar et al., 2004; Brown et al., 2009; Watson et al., 2010).

However, the effectiveness of protected areas to reconcile biodiversity conservation and the development of forest dependent communities has been widely criticised in recent years (Hayes & Ostrom, 2005). Besides this, the projected

human population increase, 90% of which will occur in developing countries (Potts, 2007) which sustain the greatest proportion of the world's biodiversity, poses many challenges to ecological systems. This is due to the high global deforestation rate (FAO, 2005) and increasing demands for environmental products and services (Houghton, 1994). In recent decades, ecologists have come to realize that human impacts on surrounding lands may cross the boundaries into protected areas (Buechner, 1987; Dasmann, 1988; Schonewald-Cox, 1988). The creation of buffer zones around protected areas has been recommended to minimize negative boundary influences (Noss, 1983). Accordingly, UNESCO's Man and the Biosphere (MAB) program advocated managing the lands around protected areas along a gradient of decreasingly intense land use toward protected area boundaries (Ebregt & de Greve, 2000).

Indeed, biodiversity conservation outside protected areas has started to take centre stage in global conservation discourse (Anyonge-Bashir & Udoto, 2012). Many researchers agree that forests outside formally protected areas are necessary for the maintenance of biodiversity and ecosystems services. Such efforts to mitigate boundary influences on protected areas will be most effective if based on scientific understanding of the underlying ecological mechanisms. Knowledge of the ecological connections could help to answer several management oriented questions.

For example, in an effort to apply the Convention on Biological Diversity, Cameroon has delineated protected areas covering 7.4 million ha, for the moment, 16% of the national territory (WRI, 2011). Approximately 60% of this territory is located in the humid forest zone. The Boumba-Bek National Park is among the country's most important protected areas. The establishment of such a conservation area is associated with the limitation of customary resources use rights of the local peoples. Investigating resources used by the Baka in and around this national park, Njounan Tegomo et al. (2012) argued that to accommodate the Baka customary rights to access the forest resources, the management plan for Boumba-Bek National Park should be partly modified to sufficiently elaborate on the actuality of the Baka traditional use of land and resources.

Since 2012, efforts have been ongoing to combine forest conservation with sustainable use of Non-Timber Forest Products (NTFPs) in this area, within the framework of the Forest Savannah Sustainability project (FOSAS). The major project activities have been carried out in Grike village located in the northern periphery of the Boumba-Bek National Park. An ecological study aiming to clarify the state of natural vegetation has been conducted in the core zone of the park, with results revealing a rich biodiversity (Nkongmeneck, 1996). However, no study has focused on the periphery zone used by the villagers for NTFPs collection and agriculture. Understanding the forest status of this periphery zone from an ecological point of view is therefore essential for management purposes.

In light of this, the present study aims to clarify the species composition, vegetation structure and plant biodiversity of the forest used by Grike villagers. Moreover, the study provides a floristic list of the vascular plants as basic information.

## METHODS

## I. Study Area

The study was carried out in the forest expanding from the Gribé village, at the northern periphery of the Boumba-Bek National Park in the southeastern Cameroon (Fig. 1) from 2011 to 2012. The periphery stretches between 2°55' N and 3°20' N and 14°45' E and 15°00' E. Almost all the inhabitants belong to one of two ethnic groups: The Baka hunter-gatherers and the Konabembe; the latter are Bantu-speaking agriculturalists. The population of the village stands at 700 inhabitants, while the population ratio in terms of ethnic groups is almost equal.

The vegetation type is classified as semi-deciduous forest where the dominant families are Sterculiaceae and Ulmaceae (Letouzey, 1968). The annual precipitation ranges from 1200 to 1600 mm. The area's major water basin is the Boumba River, which measures 200 m in width, making it the biggest river of the region.

The studied forest around the Gribé village has been used for a long period of time. The village was established 100–150 years ago by a Kounabembe clan. Since this establishment, the inhabitants have moved their settlement many times, till 1960's during which a resettlement policy has been established by colonial government. Although they changed location, their moving range was often close to their original position (Toda, 2014, this issue). This indicates that the studied forest has been used by them in parallel with their moving history, and influenced by the shifting cultivation which is a major form of Konabembe livelihood.

The periphery zone of the park is classified into three categories according to the Cameroonian zoning plans established at the end of the 1990s; (1) Logging




Fig.1. Location of study area.

Zone and ZIC (Zone d'Intérêt Cynégétique) where farming and hunting by the local people is restricted, (2) Agroforest Zone allocated to the people for their agriculture and hunting under limited condition and (3) Community Forests managed by the local communities following a management plan approved by the government's forest authority (Table 1).

## II. Data Collections and Analysis

The sampling procedure was essentially based on the establishment of a baseline of approximately 16 km running from the village settlement to the park. From this baseline, 16 transects were delineated. Each transect measured 5 km in length, 5 m in width and 2.5 ha in area. The distance between two consecutive transects was 1 km. They were alternatively placed on both sides of the baseline. The baseline was oriented from NW to SE and all transects were oriented from SW to NE (Fig. 2).

Botanical data were collected along the 16 transects. All trees above 10 cm DBH were measured and species names were identified in the field according to available floras (Letouzey, 1985; Vivien & Faure, 1985; Wilks & Issembé, 2000).




Fig. 2. Transect sampling design in the periphery zone.

Voucher specimens were prepared for unidentified species and identified at the Millennium Ecologic Museum.

For the floristic analyses, overall species, genera and family level richness, tree density (per ha), species diversity and basal area were calculated. The Importance Value Index of species (IVI) was computed as the sum of relative frequency, relative abundance and relative dominance of species:

IVI = Relative frequency + Relative abundance + Relative dominance.

$$\text{Relative frequency (RFr)} = \frac{\text{Frequency of a species}}{\text{Total number of sampling unit}} \times 100$$

$$\text{Relative abundance} = \frac{\text{Individual numbers of each species}}{\text{Total number of individuals in the sample}} \times 100$$

$$\text{Relative dominance} = \frac{\text{Basal area of each species}}{\text{Total basal area}} \times 100$$

Where basal area =  $\pi \times (D^2/4)$  where D = DBH of the tree

The family Importance Index used by Campbell et al. (2006) in the Gamba Forest Complex in Gabon and by Fongzossie et al. (2008) in the Mengamé Gorilla Reserve in South Cameroon, is the sum of relative density, relative basal area and relative diversity of each family.

$$\text{FIV} = \frac{\text{Number of species of family X} \times 100}{\text{Total number of species identified}} + \frac{\text{Number of individuals of family X} \times 100}{\text{Total number of individuals}} + \frac{\text{Sum of basal areas of family X} \times 100}{\text{Total basal area}}$$

In order to demonstrate the plant diversity for all the plots, the Shannon-Wiener diversity index ( $H'$ ) was calculated. The equation is as follows:

$$H' = -\sum_{i=1}^S p_i \log_2(p_i) \text{ where } S \text{ is the number of species, } p_i \text{ is the proportion of the } i\text{th species.}$$

The 16 transects were categorized into three zones based on the land use classification established by the Cameroonian government and status of forest use (Table 1). The three zones are: (1) Agroforest Zone which includes transects 1 to 4, (2) Intermediate Zone which includes transects 5 to 8 and (3) Logging Zone

**Table 1.** Categorization of 16 transects according to the land classification

Land classification	Status of land or forest uses	Corresponding transects
Agroforest Zone	<ul style="list-style-type: none"> <li>- Shifting cultivation with a long fallow period, cacao farming and NTFPs collection have been practiced by both Konabembe and Baka.</li> <li>- The zone consists of many types of forest: Mature forest, old secondary forest, old and young forest fallows and grass stage fallow. These types of vegetation distribute mosaic-like in the zone.</li> <li>- The density of current agricultural plots with crops and fallows decreases from T1 to T4.</li> </ul>	T1 to T4
Intermediate Zone	<ul style="list-style-type: none"> <li>- Mature and old secondary forests represent the majority.</li> <li>- There are old fallows and old cacao plots, although this is rare.</li> <li>- The zone partly includes the Community Forest where the villagers forecast to cut down commercial trees. However, the logging has not yet started (2012).</li> </ul>	T5 to T8
Logging Zone	<ul style="list-style-type: none"> <li>- Mature and old secondary forests represent the majority.</li> <li>- A logging company has occupied this zone and has been cutting the commercial trees for a few decades.</li> <li>- Logging roads measuring 5–30 m in width run through the zone in a high density.</li> </ul>	T9 to T16

which includes transects 9 to 16. Table 1 shows the status of land and forest uses of each zone.

In order to examine floristic similarities in terms of species composition among the transects and categorized lands, the NNESS similarity index was calculated. The NNESS index was used because of size differences among samples. This index is used to compare, with minimum bias, the degree of similarity of the two samples (i and j) on the basis of an identical data size k, which is randomly selected from each sample. The similarity between the two samples (i and j) is expressed by the Morishita-Horn index and by its generalisation, the NNESS index, which is a variant of the NESS index (Grassle & Smith, 1976). The formula is given below and was computed using the software BiodivR 1.0 (Hardy, 2005).

$$\text{NNESS } ij/k = \frac{\text{ESS } ij/k}{(\text{ESS } ii/k + \text{ESS } jj/k)/2}$$

where  $\text{ESS}_{ij/k}$  is the expected number of species shared for random draws of k specimens from sample i and k specimens from sample j.

## RESULTS

### I. Dominant Families and Genera

A total of 17,583 individuals were recorded across all 16 transects. The average density was 439.6 stems/ha (Total area = 40 ha) while the average basal area was 43.6 m<sup>2</sup>/ha. These trees belong to 51 families, 169 genera and 247 species. The most dominant families in terms of density were: Annonaceae with density of 48.8 stems/ha, Euphorbiaceae (44.6 stems/ha), Meliaceae (36.7), Sterculiaceae (23.9), Ulmaceae (20.7), Myristicaceae (19.9), Cecropiaceae (19.8), Violaceae (18.1), Olacaceae (18.0) and Rubiaceae (17.0). These families accounted for 59.8% of all the families recorded in the study area.

The most diversified families were Caesalpiniaceae, with 21 species, followed by Sterculiaceae (17 species), Rubiaceae and Euphorbiaceae (15), Meliaceae (14), Sapotaceae (13), Annonaceae (10), Apocynaceae (10), Moraceae (8), Ulmaceae (8), Mimosaceae (8) and Clusiaceae (8).

Based on the FIV index, the 10 most leading dominant families were Euphorbiaceae, Combretaceae, Sterculiaceae, Mimosaceae, Annonaceae, Meliaceae, Ulmaceae, Caesalpiniaceae, Irvingiaceae and Cecropiaceae (Table 2).

The most diversified genera were *Celtis* (Ulmaceae), *Cola* (Sterculiaceae), *Garcinia* (Clusiaceae) with 6 species each; followed by *Drypetes* (Putranjivaceae), *Ficus* (Moraceae) *Irvingia* (Irvingiaceae), *Psychotria* (Rubiaceae) with 5 species, and *Beilschmiedia* (Lauraceae), *Homalium* (Salicaceae), *Diospyros* (Ebenaceae), *Khaya* (Meliaceae), *Rauvolfia* (Apocynaceae), *Rinorea* (Violaceae) with 4 species each. Some genera, including *Anthonotha* (Caesalpiniaceae), *Entandrophragma* (Meliaceae), *Gambeya* (Sapotaceae) and *Xylopia* (Annonaceae) with only 3 species each.

### II. Differences in Plant Diversity among Three Zones

Turkey-tests were performed to examine significant differences in the mean number of species, the mean number of individuals and the mean value of diversity index ( $H'$ ) among the Agroforest Zone (T1–T4, N = 4), Intermediate Zone (T5–T8, N = 4) and Logging Zone (T9–T16, N = 8). The results showed that there were significant differences in all three types of value. More specifically, a significant difference was found between the Agroforest Zone and the other two zones (Table 3).

The mean number of species in Agroforest Zone, Intermediate Zone and Logging Zone were 144 (Standard deviation = 9.9), 137 (SD = 5.5) and 124 (SD = 5.5), respectively. In addition, the numbers of species in the Agroforest Zone and the Intermediate Zone were significantly larger than that in the Logging Zone ( $P = 0.0011$  between Agroforest Zone and Logging Zone,  $P = 0.0262$  between Intermediate Zone and Logging Zone). This elucidated that the species richness were more abundant in Agroforest Zone and Intermediate Zone. The mean value of number of individuals in the Agroforest Zone, Intermediate Zone and Logging

**Table 2.** Family Importance value (FIV) index for all families recorded in the study site

Rank	Family	Number of species	Number of individuals	Basal area (m <sup>2</sup> /ha)	FIV
1	Euphorbiaceae	15	1,784	4.37	191.65
2	Combretaceae	2	285	3.97	161.29
3	Sterculiaceae	16	956	3.69	159.64
4	Mimosaceae	8	567	3.30	138.56
5	Annonaceae	10	1,950	2.78	126.85
6	Meliaceae	13	1,469	2.51	114.53
7	Ulmaceae	7	826	2.51	108.12
8	Caesalpiniaceae	21	444	1.97	89.84
9	Irvingiaceae	6	318	2.00	84.50
10	Cecropiaceae	2	792	1.92	82.32
11	Apocynaceae	10	329	1.58	69.14
12	Moraceae	8	570	1.50	66.62
13	Sapotaceae	13	379	1.42	64.16
14	Olacaceae	5	719	1.08	49.42
15	Papilionaceae	7	366	1.08	48.07
16	Myristicaceae	3	797	0.95	43.81
17	Tiliaceae	4	342	0.95	41.72
18	Rubiaceae	14	613	0.75	39.32
19	Violaceae	4	724	0.44	23.77
20	Flacourtiaceae	7	316	0.41	21.30
21	Ebenaceae	6	465	0.38	20.35
22	Lecythidaceae	3	128	0.42	18.65
23	Anacardiaceae	4	227	0.39	18.53
24	Putranjivaceae	5	339	0.28	15.43
25	Bombacaceae	3	16	0.34	14.87
26	Pandaceae	1	197	0.31	13.90
27	Bignoniaceae	6	132	0.26	13.78
28	Rhamnaceae	2	367	0.22	11.87
29	Phyllanthaceae	2	65	0.26	11.61
30	Lauraceae	4	117	0.19	9.92
31	Sapindaceae	3	141	0.19	9.82
32	Clusiaceae	8	134	0.13	9.41
33	Huaceae	1	120	0.16	7.69
34	Verbenaceae	1	105	0.15	7.21
35	Lepydobotryaceae	1	128	0.14	6.97
36	Burseraceae	2	135	0.12	6.28
37	Chrysobalanaceae	2	10	0.10	4.82
38	Myrtaceae	1	5	0.09	4.21
39	Salicaceae	2	54	0.07	3.77
40	Rutaceae	6	28	0.02	3.37
41	Passifloraceae	1	72	0.03	2.18
42	Melastomataceae	2	8	0.03	2.09
43	Rhizophoraceae	1	7	0.04	1.92
44	Boraginaceae	1	4	0.03	1.57
45	Ixonanthaceae	1	13	0.03	1.54
46	Loganiaceae	3	6	0.00	1.40
47	Anisophylleaceae	1	1	0.01	1.00
48	Thymelaeaceae	2	4	< 0.01	0.93
49	Thomandersiaceae	2	7	< 0.01	0.92
50	Capparaceae	1	1	< 0.01	0.49
51	Menispermaceae	1	1	< 0.01	0.42




**Table 3.** Difference in tree density, basal area and diversity among three zones

Forest category	Transect	Number of species	Number of individuals	Density (stems/ha)	Basal area (m <sup>2</sup> /ha)	H'
Agroforest Zone	T1	158	1,018	407.2	40.5	6.51
	T2	137	875	350.0	43.3	6.35
	T3	139	864	345.6	38.3	6.42
	T4	144	928	371.2	33.7	6.27
	Total	208	3,685			
	Mean	144 <sup>A</sup>	921.3 <sup>A</sup>	368.5 <sup>A</sup>	39.0 <sup>A</sup>	6.39 <sup>A</sup>
	SD	9.9	70.3	28.1	4.1	0.10
Intermediate Zone	T5	133	1,238	495.2	46.9	6.03
	T6	145	1,078	431.2	47.5	6.33
	T7	133	1,330	532	58.3	6.32
	T8	138	1,314	525.6	45.9	6.35
	Total	189	4,960			
	Mean	137 <sup>A</sup>	1,240 <sup>B**</sup>	496 <sup>B**</sup>	49.7 <sup>B*</sup>	6.25 <sup>AB</sup>
	SD	5.5	115.2	46.1	5.8	0.15
Logging Zone	T9	133	1,304	521.6	45.2	6.30
	T10	127	1,207	482.8	43.1	6.16
	T11	125	1,122	448.8	40.3	6.22
	T12	132	1,221	488.4	42.4	6.21
	T13	119	1,103	441.2	43.3	6.21
	T14	122	1,003	401.2	35.6	6.12
	T15	124	940	376	43.4	6.23
	T16	116	1,038	415.2	49.6	5.94
	Total	186	8,938			
	Mean	124 <sup>B** C*</sup>	1,117.3 <sup>B*</sup>	446.9 <sup>B*</sup>	42.9 <sup>AB</sup>	6.20 <sup>B*</sup>
SD	5.5	122.3	48.9	4.2	0.10	
Whole (16 transects)	Total	247	17,583			
	Mean		1,098.9	439.6	43.6	6.24
	SD		156.6	62.6	5.7	0.14

Turkey-tests (ANOVA) were performed to examine significant differences in the mean number of species, individuals or mean density, the mean basal area and the mean value of H' among three zones. Significant differences were examined in order of (1) from Agroforest Zone to Intermediate Zone, (2) from Agroforest Zone to Logging Zone and (3) from Intermediate Zone to Logging Zone. When significant differences were detected, \* (P < 0.05) or \*\* (P < 0.01) were marked on the latter zones. The mean values with the same letters (A, B and C) were not significantly different.

Zone were 921.3 (SD = 70.3), 1,240 (SD = 115.2) and 1,117.3 (SD = 122.3) respectively, while the Agroforest Zone showed a significantly lower number of individuals compared to the other two zones (P = 0.0011 and 0.0262, respectively). Since the areas of each transect were identical at 2.5 ha, the significant difference of the average density in the three zones corresponds to that of average number of species.

Diversity index (H') ranged from 5.94 (Transect 16) to 6.51 (T1) with a mean value of 6.24. There was a significant difference in the index among different three zones (Fig. 3); the diversity of the Agroforest Zone was higher than that of



**Fig. 3.** Difference in mean values of diversity index ( $H'$ ) among the three zones categorized according to the land use status.

See Table 1 for the categorization of the 16 transects into the three zones. The bars indicate standard deviations. A significant difference was found between Agroforest Zone and Logging Zone according to the Turkey-test. The P-values were 0.295, 0.030 and 0.525 between the Agroforest Zone–Intermediate Zone, Agroforest Zone–Logging Zone and Intermediate Zone–Logging Zone, respectively. The mean values with the same letter were not significantly different.

the Logging Zone, while the diversity of the intermediate Zone and Logging Zone was not significant.

### III. Differences in Species Composition among Three Zones

#### (1) Floristic similarity based on NNESS similarity index

The computation of the NNESS similarity index shows that the 16 transects did not differ considerably in terms of species composition (Table 4). This index uses values between zero and one. Where there is no species in common, the pair will be scored zero, and on the contrary, the pair will be scored one if all species match.

For the 16 transects, the number of pairs to be examined totaled 120. The results of the comparison of the 120 pairs yielded a mean, minimum and maximum score in the NNESS index of 0.79, 0.62 and 0.93, respectively. This mean score shall be regarded as relatively high. These findings indicated that the species composition in the 16 transects are generally similar, and thus clarified that many species are commonly found in all of the transects.

Similarities between the vegetation in the transects set in the same zone were much higher. In the Agroforest Zone (among which four transects and six pairs were examined), the mean, minimum and maximum score in the NNESS index were 0.83, 0.81 and 0.86, respectively. Those in the Intermediate Zone were 0.78,

**Table 4.** Comparison of the species compositions among the 16 transects based on NNESS index

	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10	T11	T12	T13	T14	T15	T16
T1	1															
T2	0.81	1														
T3	0.86	0.82	1													
T4	0.82	0.85	0.84	1												
T5	0.72	0.80	0.70	0.77	1											
T6	0.80	0.84	0.78	0.84	0.80	1										
T7	0.80	0.78	0.78	0.81	0.69	0.83	1									
T8	0.79	0.77	0.76	0.81	0.64	0.82	0.90	1								
T9	0.79	0.76	0.75	0.79	0.72	0.81	0.92	0.90	1							
T10	0.78	0.76	0.76	0.81	0.64	0.82	0.91	0.92	0.91	1						
T11	0.75	0.77	0.75	0.77	0.70	0.80	0.88	0.87	0.90	0.90	1					
T12	0.76	0.76	0.75	0.81	0.64	0.80	0.88	0.91	0.88	0.92	0.91	1				
T13	0.73	0.73	0.73	0.77	0.64	0.78	0.87	0.89	0.90	0.91	0.93	0.93	1			
T14	0.73	0.72	0.70	0.77	0.62	0.78	0.85	0.86	0.85	0.91	0.89	0.93	0.91	1		
T15	0.68	0.77	0.67	0.77	0.73	0.82	0.75	0.76	0.78	0.78	0.84	0.81	0.82	0.82	1	
T16	0.68	0.73	0.67	0.74	0.65	0.77	0.73	0.80	0.76	0.80	0.82	0.85	0.85	0.86	0.86	1


Pairs in low NNESS values are marked with solid lines.

0.64 and 0.9. Those in the Logging Zone were 0.89, 0.88 and 0.91.

Several pairs with rather low similarities were found across different zones. Among these were the pairs of T1 with T15, T1 with T16, T3 with T15 and T3 with T16. As Fig. 2 shows, these transects are distant from each other. As such, there is a possibility that the land use differences were reflected in the species composition as a reason for the relatively lower value shown in these pairs. Another group of pairs with relatively lower similarity were the pairs of T5 and T7 as well as nine consecutive transects up to T16. This may have been caused by a drastic change which took place in T5.

## (2) Identification of dominant species based on IVI values and zone-specific species

The number of common species found in the three zones was 145 (58.7%) out of the total 247 species. In addition, more than half of the 145 species marked high IVI values, which is an index used to comprehensively judge dominance of species, exceeding 100 (Fig. 4). These species were common species as well as dominant species in the three zones (hereinafter, co-dominant species). Unlike temperate forests, tropical rainforests, which have higher biodiversity, often encompass dominant species with the same degree of dominance. Based on this understanding, tropical rainforests are sometimes regarded as “forests without dominant species.” However, in this study, the following two conditions were set to identify species with even higher dominance. The conditions were; (1) species that marked over 100 in IVI value in all three zones, and (2) species with even IVI values in the three zones. Shannon-Wiener evenness index ( $J'$ ) value was used as the index to show evenness here. Co-dominant species whose two conditions were applicable totaled as many as 56 species.



**Fig. 4.** Species rank-Abundance (IVI) curve.

The vertical axis usually shows number of individuals, while the IVI value showing comprehensive dominance was employed in the axis for this case. The IVI value of the common species shows the mean of the IVI values of the species marked in the three zones.

Table 5 shows the top 20 species which marked the highest mean scores in IVI value among the 56 species in the three zones. They included typical pioneer species which have characteristics of being the first to establish themselves in the gap where the vegetation had been disturbed due to tree falling, land clearance and so on. The following are the species as such; *Terminalia superba*, *Musanga cecropioides*, *Ricinodendron heudelotii*, *Albizia glaberrima*, *Trilepisium madagascariense*, *Pentaclethra macrophylla* and *Alstonia boonei*. Generally speaking, pioneer species are rapidly increasing in terms of individual numbers, fast-growing, but light in wood density together with a short life span. However, co-dominant species included not only the species with strong pioneering characteristics, but also the tree species with high carrying capacity which are slow and low in growth rate and increasing rate of individual number. However, they have a long life with a firm trunk. Species such as *Polyalthia suaveolens*, *Entandrophragma cylindricum*, *Klainedoxa gabonensis*, *Rinorea elliotii* and *Duboscia macrocarpa* were included in the group.

According to the species rank-abundance (IVI) curve (Fig. 4) of these common species, the IVI values show a very gradual decline from *Terminalia superba*, which ranked first to *Baillonella toxisperma* in 145th. The gradual decline of the IVI value shows high diversity in tree species across all three zones, in addition to the significance in number of co-dominant species.

**Table 5.** Co-dominant species across and zone-specific species in three zones

	No. of Species	% to the total no. of species	Rank	Species	P*	IVI			J'
						Agroforest Zone	Intermediate Zone	Logging Zone	
Co-dominant species across three zones (Top 20 species)	145	58.7	1	<i>Terminalia superba</i>	pi	113.3	111.5	107.4	1
			2	<i>Musanga cecropioides</i>	pi	110.7	110.8	103.9	1
			3	<i>Celtis mildbraedii</i>	np	107.7	104.7	103.8	1
			4	<i>Anonidium mannii</i>	np	103.4	103.5	108.5	1
			5	<i>Ricinodendron heudelotii</i>	pi	103.7	104.2	105.5	1
			6	<i>Trichilia heudelotii</i>	np	104.5	104.2	104.4	1
			7	<i>Trilepisium madagascariense</i>	pi	104.4	106.3	101.9	1
			8	<i>Polyalthia suaveolens</i>	np	103.8	103.9	104.5	1
			9	<i>Albizia glaberrima</i>	pi	104.2	103.4	104.3	1
			10	<i>Entandrophragma cylindricum</i>	np	105.2	101.6	104.0	1
			11	<i>Staudtia kamerunensis</i>	np	103.3	103.8	103.1	1
			12	<i>Pentaclethra macrophylla</i>	pi	103.5	103.0	103.3	1
			13	<i>Alstonia boonei</i>	pi	102.0	103.0	104.4	1
			14	<i>Strombosia pustulata</i>	np	101.6	104.5	103.3	1
			15	<i>Rinorea elliotii</i>	np	103.4	102.9	102.6	1
			16	<i>Cola lateritia</i>	-	101.6	103.1	103.9	1
			17	<i>Duboscia macrocarpa</i>	np	102.7	102.2	103.3	1
			18	<i>Dichostemma glaucescens</i>	np	102.0	101.9	104.1	1
			19	<i>Klainedoxa gabonensis</i>	np	101.1	103.5	103.2	1
			20	<i>Celtis adolfi-friderici</i>	-	102.6	102.1	103.0	1
Zone-specific species in Agroforest Zone (Top 10 species)	32	13.0	1	<i>Irvingia</i> sp.	np	100.7			
			2	<i>Pavetta</i> sp.	np	76.0			
			3	<i>Copaifera mildbraedii</i>	np	75.1			
			4	<i>Ficus mucuso</i>	pi	50.6			
			5	<i>Samanea dinklagei</i>	pi	50.6			
			6	<i>Stemonocoleus micranthus</i>	-	50.5			
			7	<i>Calpocalyx</i> sp.	-	50.3			
			8	<i>Millettia</i> sp.	-	50.1			
			9	<i>Hallea ledermannii</i>	-	50.1			
			10	<i>Beilschmiedia</i> sp.	np	25.2			
Zone-specific species in Intermediate Zone (Top 10 species)	11	4.5	1	<i>Allanblachia gabonensis</i>	-		50.2		
			2	<i>Diospyros abyssinica</i>	-		25.1		
			3	<i>Zanthoxylon laurentii</i>	np		25.0		
			4	<i>Buchholzia</i> sp.	-		25.0		
			5	<i>Beilschmiedia</i> sp.2	np		25.0		
			6	<i>Stychnos</i> sp.2	-		25.0		
			7	<i>Macaranga spinosa</i>	pi		25.0		
			8	<i>Cyttaranthus congolensis</i>	-		25.0		
			9	<i>Parinari excelsa</i>	np		25.0		
			10	<i>Chytranthus atroviolaceus</i>	-		25.0		
Zone-specific species in Logging Zone (Top 10 species)	13	5.3	1	<i>Pterocarpus mildbraedii</i>	np			63.7	
			2	<i>Xylopia hypolampra</i>	-			62.9	
			3	<i>Dialium</i> sp.	np			62.6	
			4	<i>Fagara macrophylla</i>	-			25.1	
			5	<i>Entandrophragma utile</i>	np			12.9	
			6	<i>Poga oleosa</i>	-			12.6	
			7	<i>Brenania brieyi</i>	-			12.5	
			8	<i>Rauvolfia macrophylla</i>	-			12.5	
			9	<i>Celtis zenkeri</i>	-			12.5	
			10	<i>Anthocleista schweinfurthii</i>	-			12.5	

\*: Judgment of whether or not species were pioneering was made based on the vegetation observation in the study area. pi: Pioneer species, np: Non-pioneer, -: Unknown.

In comparison to the common species, the number of species specifically found in each zone was significantly low. Number of such zone-specific species (hereinafter, specific species) were 32, 11 and 13 species in the Agroforest Zone, Intermediate Zone and Logging Zone respectively (Table 5).

The specific species presumably included more species with high carrying capacity than tree species with pioneering characteristics. As far as we can tell at this point, such species were *Irvingia* sp. and *Pavetta* sp. (in Agroforest Zone), *Copaifera mildbraedii*, *Diospyros abyssinica* and *Beilschmiedia* sp. (in Intermediate Zone), and *Parinari excelsa*, *Entandrophragma utile*, *Dialium* sp. and *Pterocarpus mildbraedii* (in Logging Zone).


The species rank-abundance (IVI) curve of the specific species (Fig. 4) shows that only limited species marked high IVI values in all zones, in contrast with the common species. In addition, the curve shows a steep drop. Specific species may serve as indicators to show the states of the vegetation environment. However, we are not yet certain as to whether these specific species may become such indicators for each zone at this stage. This will need to be elucidated through further studies on the habitat preferred by the specific species and the characteristics of the species.

#### IV. Forest Structure: Diameter and Height-class Distribution

Distribution of density (or number of individuals) by DBH-class showed a common tendency for individuals with smaller diameter to be larger in number of individuals in the Agroforest Zone, Intermediate Zone and Logging Zone (Fig. 5 left). The ratio of the number of the individuals included in the smallest diameter-class (10–30 cm), in the total number of individuals was 71.2% in the Agroforest Zone ( $N = 3,685$ ), 74.2% in the Intermediate Zone ( $N = 4,960$ ) and 75.8% in the Logging Zone ( $N = 8,938$ ). The largest DBH-classes for the Agroforest Zone were 210–230 cm ( $N = 1$ , 0.03% of total individuals of the zone) and 250–310 cm for the Intermediate Zone and Logging Zone (one individual in each zone, the percentages were 0.02 in the former and 0.01 in the latter). It was clarified that the distribution of diameter-classes showed a reverse J shape pattern in each zone, which is often found in natural forests.

A Turkey-test was performed in order to examine significant differences in basal area between the three zones. The results (Table 3) showed that the basal area in the Agroforest Zone was significantly lower ( $P = 0.013$ ) than that in the Intermediate Zone, although it had no significant difference with the Logging Zone ( $P = 0.358$ ).

A similar pattern was found in distribution of the individual numbers by tree height-classes across three zones (Fig. 5 right). The individuals classified in the 5–20 m height-class were the most numerous in all the zones, and their ratio against total individuals in each zone was 92.7% in Agroforest Zone, 84.9% in the Intermediate Zone and 84.6% in the Logging Zone. The highest classes observed in each zone were 45–50 m for the Agroforest Zone and the Intermediate Zone ( $N = 12$  in each zone, with 0.3% for the former and 0.2 for the latter) and 50–55 m in the Logging Zone ( $N = 1$ , 0.01%).



**Fig. 5.** Size-class distribution by zone.

Left shows the tree density by DBH-class and right indicates tree density by height-class. All individuals with DBH larger than 10 cm were subject of the measurement, and their DBH and height were measured ( $N = 17,583$ ).

By examining Fig. 5 (right) in the view of the forest stratification, it was hypothesized that each zone was comprised of seven layers, namely (1) lower layer (0–5 m), (2) middle layer I (5–15 m), (3) middle layer II (15–20 m), (4) canopy layer I (20–25 m), (5) canopy layer II (25–35 m), (6) canopy layer III (35–40 m) and (7) emergent layer (40 m or higher). The fact that the forest has seven vertical layers of tree heights can be seen as evidence of the forest's development. It can be said that different species segregating themselves to grow in a vertical direction would mitigate the inter-species competition for light resources, and has contributed to the creation and maintenance of high diversity in plant species.

In order to examine significant differences in DBH and tree height among the three zones, an applicable one way ANOVA paired-test was performed. The result revealed no significant difference among the zones in terms of DBH. On the other hand, a significant difference was found between the Agroforest Zone and the Intermediate Zones in terms of tree height ( $P = 0.018$ ). From these findings, it was elucidated that the height of the trees grown in the Agroforest Zone tended to be lower than other zones in every tree height-class. However, the difference was slight, and one should observe the fact that the forest in this zone was also comprised of the seven layers in the same way as in other zones.

## DISCUSSION

### I. Global Floristic Composition

According to Letouzey (1968), the study area is classified as semi-deciduous forest dominated by Sterculiaceae and Ulmaceae. Our results showed that the studied forest was dominated by the families of Annonaceae, Euphorbiaceae, Meliaceae, Sterculiaceae, Ulmaceae, Myristicaceae and Cecropiaceae in terms of FIV values. This observation is in line with Letouzey's description and the findings of Nkongmeneck (1996) who reported the important occurrence of Euphorbiaceae, Sterculiaceae, Rubiaceae, Meliaceae, Caesalpiniaceae, Annonaceae, Fabaceae, Tiliaceae, Ulmaceae and Ebenaceae in the Boumba-Bek National Park, and Annonaceae, Euphorbiaceae, Caesalpiniaceae, Olacaceae, Fabaceae, Meliaceae, Rubiaceae, Sterculiaceae, Tiliaceae and Mimosaceae in the Nki National Park.

Studies conducted in neighboring forests reported the dominance of Euphorbiaceae, Rubiaceae, Caesalpiniaceae, Sapotaceae, Annonaceae, Meliaceae, Sterculiaceae, Mimosaceae, Flacourtiaceae, Apocynaceae, Olacaceae and Sapindaceae in the Dja Biosphere Reserve (Nguenang & Dupain, 2002; Mbolo, 2004; Sonké, 2005; Nzoo Dongmo, 2005), and Euphorbiaceae, Annonaceae, Sterculiaceae, Meliaceae, Caesalpiniaceae, Mimosaceae in the Kom-Mengamé Forest Complex (Fongzossie et al., 2010).

### II. Difference in Plant Biodiversity and Dominant Species among the Three Different Zones

The present study has considered vegetation differences in the three zones with different usage from three perspectives, namely diversity of the trees, species composition, and forest structure. At this point, we compare the three zones in each point of view.

The tree species observed totaled 247 with a density of 439.6 stems/ha across the three zones. Comparing to other regions, Lubini (1997) found 236 stems/ha at Luki. Lomba (2007) identified 183 species arranged in 37 families with a mean density of 506 stems/ha in the Yoko Reserve. Moreover, Nshimba (2008) reported a density of 506 stems/ha from Kisangani Mbiye's Island. Sunderland et al. (2004) conducted their study in the National Park of Mbe (Gabon), and found a density of 539 stems/ha, 97 species and a total basal area of 37.2 m<sup>2</sup>/ha. Whitmore (1990) and Doucet (2003) have all mentioned the specific richness of South America and tropical Asia, with approximately 300 species (dbh  $\geq$  10cm) per 1 ha and 711 species (dbh  $\geq$  10cm) per 6.6 ha, respectively. These values show a striking analogy with our results. Indeed, there are certain difficulties which arise when trying to conduct a comparative analysis of the diversity between the studied forest in this work and that on other sites. These difficulties have been cited by many authors, including Lejoly (2003), Senterre (2005), Ngok (2000) and Kanguuja (2009). These problems concern the diversity of methods used.

After comparing the diversity between the three divided zones in the study



area, it was clear that the diversity was higher in the Agroforest Zone than the Logging Zone and even in the Intermediate Zone. This is because both zones are subject to different land use intensities which affect the forest structure and species composition differently. In the Agroforest Zone, the villagers cut trees every year in order to clear new farming plots, although such cleared plots only constitutes approximately 1% of the whole Agroforest Zone area (Hirai, 2014, this issue). In addition, farming plots would rarely be maintained after three years of the clearance and would be gradually covered by pioneer species afterwards. The villagers only seek their new plots in forests which have not been cleared for 30 to 60 years. Agriculture in this region has been operated based on the premise of the long fallow period, and as such, patches of vegetation in various recovery stages from the clearance are distributed mosaic-like in this zone (Hirai, 2014, this issue). It is conceivable that this is a part of the mechanism which creates tree diversity in the zone. Another important point relating to diversity creation is that the villagers leave many trees after the forest clearance due to factors such as usability, growing location, and easiness to cut (firmness of trunks). Carrière et al. (2002) clarified that such trees left in the farming lands play important roles when abandoned plots are promptly recovered into forests.

In addition, tree regeneration composition in the gaps has been shown to be dependent upon the history of the forest community, seed availability and the biology of the species (Hubbell & Foster, 1992; Putul Bhuyan et al., 2003). This trend in the diversity pattern across three zones is consistent with the Intermediate disturbance hypothesis (Connell, 1979), according to which a forest stand reaches maximum diversity under an intermediate disturbance regime.

It is now fitting to compare the three zones in terms of species composition. The NNESS similarity coefficient between pairs of different transects were relatively high, thus indicating very low floristic difference among three zones. Moreover, the similarities between the transects in the same zone were higher. This indicates that many tree species are commonly found in the three zones. The results of the detailed investigation of common species and zone-specific species show greater dominance of *Terminalia superba*, *Musanga cecropioides*, *Trichilia heudelotii*, *Polyalthia suaveolens*, *Albizia glaberrima*, *Celtis mildbraedii* and *Entandrophragma cylindricum* across all three zones. Most of these leading dominant species are pioneer fast growing trees. Their dominance has been reported in several vegetation assessments in disturbed tropical rainforest areas. However, there were many co-dominant species and not all of them were pioneer species in the study area. For instance, *Duboscia macrocarpa* and *Entandrophragma cylindricum* are species grown often in mature forests. In addition to this, more species with carrying capacity than pioneer species were found among the specific species in the Agroforestry Zone.

### III. Vegetation Structure

Finally, we discuss the differences in the three zones from the point of view of forest structure. The tree distribution patterns of both diameter and height-class were common in the three zones. The distribution of trees in this survey into

different diameter-classes had an inverse J-shape. This pattern is similar to those obtained in the Dja Biosphere Reserve and the Mengame Gorilla Sanctuary in the southern Cameroon (Sonké, 2005; Fongnzossie et al., 2008). This distribution pattern, similar to the exponential model, characterizes a forest dominated by young and fast growing trees (Rollet, 1974). However, this exponential model does not always mean that the forest is dominated by sub-strata species but could prove that the considered species have a high regeneration level (Senterre & Nguema, 2001; Fongnzossie et al., 2008). Gentry (1990) has shown that this capacity of species to maintain the constant rhythm of installation of seedling can be favored by several factors including the faunal diversity, disturbance and the auto-ecology of each species. Compared with the other site (Fongnzossie et al., 2010), the average tree density (439.6 stems/ha) are relatively high and is contrasting with the disturbance history of the site. It is worth mentioning that the forest at the northern periphery of Boumba-Bek National Park still has many trees with high carrying capacity. A unimodal reverse J-shaped curve of diameter-class distribution in most zones is indicative of these forests' regenerating status.

The tree height-class distribution clarified that the forests were divided into seven vertical layers in every zone. Tree height was slightly lower in the Agroforestry Zone in general, although the same number of layers as other zones were found in the zone. The forests in the Agroforestry Zone are most frequently utilised by the villagers as a place for agriculture and collection of NTFPs. Since complicated layer division in forests may be employed as an indicator of forest development, the fact that the divisions were found in the forests under such circumstance suggests the villagers are not in existence only to destroy the forests, contrary to what was thought in the past. There is a need for further development when it comes to understanding the mutual relationship between people and forest through further research on forest vegetation.

## CONCLUSION

We conclude that the studied forest keeps a high degree of plant biodiversity despite the frequent use by the villagers. This higher biodiversity would be caused by small scale disturbances of the local peoples' livelihoods such as shifting cultivation with a long fallow period and NTFPs collection, whereas these human impacts have often been considered as major causes of forest degradation. The quantity of information treated through this study at the northern periphery of the Boumba-Bek National Park revealed the high variability of tree species in this habitat. This variation is seen at family and species level. A total of 17,583 trees belong to 51 families, 169 genera and 247 species have been inventoried. Despite the disturbance of the site driven by forest logging and agriculture, the analysis shows that this forest is relatively diverse and probably young. Though only tree species with DBH above 10 cm were inventoried, it is known that this layer contributes to nearly 60% of the total diversity of tropical ecosystems. However, investigations on other sub-strata might be interesting to complete the floral database of this forest. The diversity of tree species encountered in this study

is the basis of livelihoods of thousands of population living around this park. Investigation concerning the forest-people interaction will be crucial for establishing a multipurpose forestry system harnessing forest conservation, its uses and the improvement of the livelihood of adjacent communities.

**ACKNOWLEDGEMENTS** This study was financially supported by the FOSAS project. We express our gratitude to all participating institutions (IRAD, Millennium Ecologic Museum, Kyoto University, University of Yaoundé 1, University of Douala, and University of Dschang) and researchers for their support. We also acknowledge all our field guides and all residents of the Gribé village for their co-operation, assistance and warm hospitality during the survey. Our sincere gratitude also goes to Dr. Tsabang Nole for his assistance in identifying the numerous plant specimens collected during this study.

## REFERENCES

- Anyonge-Bashir, M. & P. Udoto 2012. Beyond philanthropy: Community nature-based enterprises as a basis for wildlife conservation. *The George Wright Forum*, 29(1): 67–73.
- Brown, S.C., C.A. Mason, J.L. Lombard, F. Martinez, E. Plater-Zyberk, A.R. Spokane, F.L. Newman, H. Pantin & J. Szapocznik 2009. The relationship of built environment to perceived social support and psychological distress in Hispanic Elders: The role of “eyes on the street”. *Journals of Gerontology: Psychological Sciences and Social Sciences*, 64B(2): 234–246.
- Buechner, M. 1987. Conservation in insular parks: Simulation models of factors affecting the movement of animals across park boundaries. *Biological Conservation*, 41(1): 57–76.
- Campbell, P., P. Rivera, D. Thomas, H. Bourobou-Bourobou, T. Nzabi, A. Alonso & F. Dallmeier 2006. Floristic structure, composition and diversity of an equatorial forest in Gabon. In (A. Alonso, M.E. Lee, P. Campbell, O.S.G. Pauwels, F. Dallmeier, eds.) *Gamba, Gabon: Biodiversity of an Equatorial African Rainforest*. Bulletin of the Biological Society of Washington, No. 12, pp. 29–52. Biological Society of Washington, Lawrence.
- Carrière, S.M., M. Andre, P. Letourmy, I. Olivier & D.B. Mckey 2002. Seed rain beneath remnant trees in a slash-and-burn agricultural system in southern Cameroon. *Journal of Tropical Ecology*, 18(3): 353–374.
- Connell, J.H. 1979. Intermediate-disturbance hypothesis. *Science*, 204(4399): 1345.
- Dasmann, R.F. 1988. Biosphere reserves, buffers, and boundaries. *BioScience*, 38(7) : 487–489.
- Doucet, J.-L. 2003. *L'Alliance Délicate de la Gestion Forestière et de la Biodiversité dans les Forêts du Centre du Gabon*. Thèse de Doctorat. Universitaire des sciences Agronomiques de Gembloux.
- Ebregt, A. & P. de Greve 2000. *Buffer Zones and Their Management: Policy and Best Practices for Terrestrial Ecosystems in Developing Countries*. National Reference Centre for Nature Management and International Agricultural Centre, Wageningen.
- Fongzossie, F.E., B.-A. Nkongmeneck, N. Tsabang & G.M. Nguenang 2010. The importance of habitat characteristics for tree diversity in the Mengamé Gorilla Reserve (South Cameroon). *Tropics*, 19(2): 53–66.
- Fongzossie, F.E., N. Tsabang, B.-A. Nkongmeneck, G.M. Nguenang, P. Auzel, E. Christina, E. Kamou, J.M. Balouma, P. Apalo, H. Mathieu, M. Valbuena & M. Valère 2008. Les peuplements d'arbres du sanctuaire à gorilles de Mengamé au sud Cameroun. *Tropical Conservation Science*, 1(3): 204–221.

- Food and Agriculture Organization (FAO) 2005. *Evaluation des Ressources Forestières Mondiales 2005 : Progrès vers la Gestion Forestière Durable*. Food and Agriculture Organization, Rome.
- Gentry, A.H. (ed.) 1990. *Four Neotropical Rainforests*. Yale University Press. New Haven.
- Grassle, J.F. & W. Smith 1976. A similarity measure sensitive to the contribution of rare species and its use in investigation of variation in marine benthic communities. *Oecologia*, 25: 13–22.
- Hardy, O.J. 2005. BiodivR 1.0: A program to compute statistically unbiased indices of species diversity within samples and species similarity between samples using rarefaction principles. Online. <http://ebe.ulb.ac.be/ebe/Software.html> (Accessed October 10, 2010).
- Hayes, T. & E. Ostrom 2005. Conserving the world's forests: Are protected areas the only way? *Indiana Law Review*, 38(3): 595–617.
- Hirai, M. 2014. Agricultural land use, collection and sales of non-timber forest products in the Agroforest Zone in southeastern Cameroon. *African Study Monographs Supplementary Issue*, 49: 169–202.
- Houghton, R.A. 1994. The worldwide extent of land-use change. *BioScience*, 44(5): 305–313.
- Hubbell, S.P. & R.B. Foster 1992. Short-term dynamics of a neotropical forest: why ecological research matters to tropical conservation and management. *Oikos*, 63: 48–61.
- Kanguéja, B.F. 2009. *Analyse de la Diversité des Ligneux Arborescents des Principaux Types Forestiers du Nord-est de la Réserve de Biosphère de Luki (Bas-Congo, RDC)*. Mémoire de Master, Université de Kisangani.
- Lejoly, J. 2003. *Gestion Durable des Forêts Tropicales*. Université Libre de Bruxelles, Brussel.
- Letouzey, R. 1968. Etude phytogéographique du Cameroun. In (P. Lechevalier, ed.) *Encyclopédie Biologique, LXIX*, p. 511. P. Lechevalier, Paris.
- 1985. *Notice de la Carte Phytogéographique du Cameroun au 1: 500000*. Institut de la Carte Internationale de la Végétation, Toulouse.
- Lomba, C. 2007. *Contribution à l'Étude de la Phytodiversité de la Réserve Forestière de Yoko*. Faculté des Sciences, Université de Kisangani, Kisangani.
- Lubini, A. 1997. *La Végétation de la Réserve de Biosphère de Luki au Mayombe (Zaire)*, *Opera Botanica Belgica, Series vol. 10*. Jardin Botanique National de Belgique, Meise.
- Mbolo, M. 2004. *Cartographie et Typologie de la Végétation de la Réserve du Dja*. Thèse de Doctorat. Département de Biologie et Physiologie Végétales, Université de Yaoundé I.
- Molnar, A., S.J. Scherr & A. Khare 2004. *Who Conserves the World's Forests? A New Assessment of Conservation and Investment Trends*. Forest Trends, Washington, D.C.
- Ngok, B.L. 2000. *Diversité Végétale des Inselbergs et des Dalles Rocheuses du Nord de Gabon*. Thèse de Doctorat. Laboratoire de Botanique Systématique et de Phytosociologie, Université Libre de Bruxelles.
- Nguenang, G.M. & J. Dupain 2002. *Typologie Morphostructurale de la Mosaïque Forestière du Dja : Cas du Site d'Étude de la Socioécologie des Grands Singes dans les Villages Malen V, Doumo Pierre et Mimpala (Est Cameroun)*. Rapport Société Royale Zoologique d'Anvers, Yaounde.
- Njounan Tegomo, O., L. Defo & L. Usongo 2012. Mapping of resource use area by the Baka pygmies inside and around Boumba-Bek National Park in southeast Cameroon, with special reference to Baka's customary rights. *African Study Monographs Supplementary Issue*, 43: 45–59.
- Nkongmeneck, B.-A. 1996. *The Boumba-Bek and Nki Forest Reserve: Botany and Ethnobotany*. World Wildlife Fund Central Africa Office, Cameroon, Yaoundé.

- Noss, R.F. 1983. A regional landscape approach to maintain diversity. *BioScience*, 33(11): 700–706.
- Nshimba, H. 2008. *Étude Floristique, Ecologique et Phytosociologique des Forêts Inondées de l'île Mbiye à Kisangani, R.D. Congo*. Thèse de Doctorat. Laboratoire de Botanique systématique et de Phytosociologie, Université Libre de Bruxelles.
- Nzoooh Dongmo, Z.L. 2005. *Biologie et Ecologie des Rotangs Dans la Réserve de Biosphère du Dja*. Thèse de Doctorat. Département de Biologie et Physiologie Végétales, Université de Yaoundé I.
- Potts, D. 2007. Natural resource access in Southern Africa: The influence of political change and development practice. In (C. Tornimbeni & M. Zamponi, eds.) *Afriche e Orienti, Special Issue 07, Terra e Risorse Naturali in Africa. Quali Diritti? Competing Rights. Land and Natural Resources in Africa*, pp. 67–80. Agenzia Internazionale Editoria Pubblicità Editore, Serravalle.
- Putul Bhuyan, M.L. Khan & R.S. Tripathi 2003. Tree diversity and population structure in undisturbed and human-impacted stands of tropical wet evergreen forest in Arunachal Pradesh, eastern Himalayas, India. *Biodiversity and Conservation*, 12: 1753–1773.
- Rollet, B. 1974. *L'Architecture des Forêts Denses Humides Sempervirentes de Plaines*. Centre Technique Forestier Tropical, Nogent-sur-Marne.
- Schonewald-Cox, C. M. 1988. Boundaries in the protection of nature reserves. *BioScience*, 38(7): 480–486.
- Senterre, B. 2005. *Recherches Méthodologiques pour la Typologie de la Végétation et la Phytogéographie des Forêts Denses d'Afrique Tropicale*. Thèse de Doctorat. Université Libre de Bruxelles.
- Senterre, B. & N. Nguema 2001. La diversité des ligneux dans la forêt de Nsork (Guinée Equatoriale). *Systematics and Geography of Plants*, 71(2): 837–846.
- Sonké, B. 2005. *Forêts de la Réserve du Dja (Cameroun)*. Thèse de Doctorat. Université libre de Bruxelles.
- Sunderland, T., G. Walters & Y. Issembe 2004. *Etude Préliminaire de la Végétation du Parc National de Mbe, Monts de Cristal, Gabon*. Central Africa Regional Program for the Environment, Kinshasa.
- Toda, M. 2014. People and social organizations in Gribé, southeastern Cameroon. *African Study Monographs Supplementary Issue*, 49: 139–168.
- Vivien J. & J.J. Faure 1985. *Arbres des Forêts Denses d'Afrique Centrale*. Agence de Coopération Culturelle et Technique, Paris.
- Watson, G.B., S.W. Chouinard, K.R. Cook, C. Geng, J.M. Gifford, G.D. Gustafson, J.M. Hasler, I.M. Larrinua, T.J. Letherer, J.C. Mitchell, W.L. Pak, V.L. Salgado, T.C. Sparks & G.E. Stilwell 2010. A spinosyn-sensitive *Drosophila melanogaster* nicotinic acetylcholine receptor identified through chemically induced target site resistance, resistance gene identification, and heterologous expression. *Insect Biochemistry and Molecular Biology*, 40(5): 376–384.
- Whitmore, T.C. 1990. *An Introduction to Tropical Rain Forest*. Oxford University Press, New York.
- Wilks C. & Y. Issembé 2000. *Guide Pratique d'Identification : Les Arbres de Guinée Equatoriale, Région Continentale*. Conservation et Utilisation Rationnelle des Ecosystèmes Forestiers, Bata.
- World Resources Institute 2011. *Atlas Forestiers Interactif du Cameroun, Version 3.0, Document de Synthèse*. World Resources Institute, Washington, D.C.

————— Accepted *March 10, 2014*

Corresponding Author's Name and Address: Vice Clotèxe TAJEUKEM, *Department of Plant Biology, Faculty of Science, University of Yaoundé 1, P.O. BOX 8038, Yaoundé, CAMEROON.*  
E-mail: tajeukem [at] yahoo.fr

**Appendix I.** All family and species recorded in the studied forest

No.	Family	Scientific name	Baka name	IVI
1	Rutaceae	<i>Aeglopsis chevalieri</i>	unknown 8	6.3
2	Huaceae	<i>Afrostryrax lepidophyllus</i>	guimba	101.1
3	Caesalpiniaceae	<i>Azelia bipindensis</i>	bimba	100.8
4	Mimosaceae	<i>Albizia ferruginea</i>	londa	12.6
5	Mimosaceae	<i>Albizia glaberrima</i>	bamba	104.0
6	Clusiaceae	<i>Allanblackia floribunda</i>	kouom 1	75.4
7	Clusiaceae	<i>Allanblackia gabonensis</i>	kouom 2	12.5
8	Apocynaceae	<i>Alstonia boonei</i>	gouga	103.5
9	Euphorbiaceae	<i>Amanoa bracteosa</i>	mbondo 1	102.0
10	Caesalpiniaceae	<i>Amphimas pterocarpoides</i>	kanga	63.5
11	Papilionaceae	<i>Angylocalyx pynaertii</i>	yonga 1	101.1
12	Papilionaceae	<i>Angylocalyx vermeulenii</i>	yonga 2	75.4
13	Sapotaceae	<i>Aningeria robusta</i>	moundonguè 3	37.7
14	Annonaceae	<i>Anonidium mannii</i>	ngwé	106.0
15	Rhizophoraceae	<i>Anopyxis klaineana</i>	boma	37.6
16	Loganiaceae	<i>Anthocleista schweinfurthii</i>	molondjo2	6.3
17	Caesalpiniaceae	<i>Anthonotha macrophylla</i>	koguimba	69.1
18	Caesalpiniaceae	<i>Anthonotha</i> sp.1	pfhouopfhouolo	6.3
19	Caesalpiniaceae	<i>Anthonotha</i> sp.2	pfouofoundo	6.3
20	Anacardiaceae	<i>Antrocaryon klaineinum</i>	gongó	81.6
21	Sapotaceae	<i>Baillonella toxisperma</i>	mabé	19.3
22	Passifloraceae	<i>Barteria fistulosa</i>	pambo	100.5
23	Lauraceae	<i>Beilschmiedia mannii</i>	békéssi	44.5
24	Lauraceae	<i>Beilschmiedia obscura</i>	mobakosso	69.0
25	Lauraceae	<i>Beilschmiedia</i> sp.1	bokondo 2	6.3
26	Lauraceae	<i>Beilschmiedia</i> sp.2	mbatama 2	6.3
27	Sapindaceae	<i>Blighia welwitschii</i>	toko	95.0
28	Bombacaceae	<i>Bombax buonopozense</i>	dombi 1	18.8
29	Rubiaceae	<i>Brenania brieiyi</i>	molondjo 1	6.3
30	Sapotaceae	<i>Brevice leptosperma</i>	koloka	94.7
31	Euphorbiaceae	<i>Bridelia grandis</i>	takou	81.8
32	Capparaceae	<i>Buchholzia</i> sp.	ngó'ndo	6.3
33	Flacourtiaceae	<i>Buchnerodendron speciosum</i>	gwagolo 1	37.6
34	Flacourtiaceae	<i>Caloncoba glauca</i>	gwagolo 2	62.6
35	Flacourtiaceae	<i>Caloncoba</i> sp.	gnangalé	6.3
36	Mimosaceae	<i>Calpocalyx</i> sp.	bambayoko	12.6
37	Meliaceae	<i>Carapa procera</i>	godjo 1	38.0
38	Meliaceae	<i>Carapa</i> sp.	unknown 19	6.3
39	Bombacaceae	<i>Ceiba pentandra</i>	koulou	32.0
40	Ulmaceae	<i>Celtis adolfi-friderici</i>	kakala	102.7
41	Ulmaceae	<i>Celtis mildbraedii</i>	ngombé	104.9
42	Ulmaceae	<i>Celtis philippensis</i>	gwèguè	107.8
43	Ulmaceae	<i>Celtis tessmannii</i>	kèkèlè 1	94.5
44	Ulmaceae	<i>Celtis zenkeri</i>	kongombé	6.3
45	Sapindaceae	<i>Chytranthus atroviolaceus</i>	tokoboli	6.3
46	Rutaceae	<i>Citropsis</i> sp.	adjomba	12.5
47	Annonaceae	<i>Cleistopholis patens</i>	kiyo	100.5
48	Myristicaceae	<i>Coelocaryon preussii</i>	bambayoko	32.1
49	Sterculiaceae	<i>Cola acuminata</i>	ligoh 1	101.1
50	Sterculiaceae	<i>Cola altissima</i>	banga	75.3
51	Sterculiaceae	<i>Cola ballayi</i>	ligoh (goro)	62.8
52	Sterculiaceae	<i>Cola lateritia</i>	p'hop'hoko	89.3
53	Sterculiaceae	<i>Cola</i> sp.	mbolé	63.8



## Appendix I. (continued)

No.	Family	Scientific name	Baka name	IVI
54	Sterculiaceae	<i>Cola pachycarpa</i>	<i>mboussoua</i> 2	6.3
55	Caesalpiniaceae	<i>Copaifera mildbraedii</i>	<i>modoumba</i>	18.8
56	Boraginaceae	<i>Cordia platythyrsa</i>	<i>gwabi</i>	25.1
57	Rubiaceae	<i>Corynanthe pachyceras</i>	<i>moka</i>	102.3
58	Rubiaceae	<i>Corynanthe</i> sp.	<i>ndôo</i>	12.5
59	Mimosaceae	<i>Cylicodiscus gabunensis</i>	<i>bolouma</i>	88.7
60	Euphorbiaceae	<i>Cyttaranthus congolensis</i>	<i>poungué</i>	6.3
61	Burseraceae	<i>Dacryodes klaineana</i>	<i>bo'o</i>	56.6
62	Tiliaceae	<i>Desplatsia dewevrei</i>	<i>liamba</i>	101.0
63	Tiliaceae	<i>Desplatsia</i> sp.	<i>libéa</i>	37.7
64	Caesalpiniaceae	<i>Detarium macrocarpum</i>	<i>mbili</i> 1	25.4
65	Caesalpiniaceae	<i>Dialium dinklagei</i>	<i>kombé</i> 2	25.1
66	Caesalpiniaceae	<i>Dialium</i> sp.	<i>mbéléngué</i> 3	31.3
67	Melastomataceae	<i>Dichaetanthera africana</i>	unknown 14	6.3
68	Melastomataceae	<i>Dichaetanthera</i> sp.	<i>mbondo</i> 2	25.1
69	Euphorbiaceae	<i>Dichostemma glaucescens</i>	<i>moungamba</i>	103.1
70	Thymelaeaceae	<i>Dicranolepis disticha</i>	<i>ngwi</i> 1	12.5
71	Thymelaeaceae	<i>Dicranolepis</i> sp.	<i>mapèbègnè</i>	6.3
72	Ebenaceae	<i>Diospyros abyssinica</i>	<i>djama</i>	6.3
73	Ebenaceae	<i>Diospyros iturensis</i>	<i>babango</i>	94.2
74	Ebenaceae	<i>Diospyros canaliculata</i>	<i>mbéla</i>	87.8
75	Ebenaceae	<i>Diospyros crassiflora</i>	<i>lembè</i>	101.3
76	Ebenaceae	<i>Diospyros</i> sp.1	<i>bandoguilè</i>	43.9
77	Ebenaceae	<i>Diospyros</i> sp.2	<i>mboloa</i>	101.4
78	Euphorbiaceae	<i>Discoglypemma caloneura</i>	<i>djila</i> 1	94.1
79	Caesalpiniaceae	<i>Distemonanthus benthamiana</i>	<i>sènè</i>	75.3
80	Sapotaceae	<i>Donella ubanguiensis</i>	<i>konya</i>	6.3
81	Putranjivaceae	<i>Drypetes gossweileri</i>	<i>bologa</i>	100.8
82	Putranjivaceae	<i>Drypetes ituriensis</i>	<i>gongo</i>	75.3
83	Putranjivaceae	<i>Drypetes principum</i>	<i>mototombo</i>	6.3
84	Putranjivaceae	<i>Drypetes</i> sp.1	<i>macép'ha</i>	101.6
85	Putranjivaceae	<i>Drypetes</i> sp.2	unknown 2	6.3
86	Tiliaceae	<i>Duboscia macrocarpa</i>	<i>goulouma</i>	102.9
87	Euphorbiaceae	<i>Euphorbia drupifera</i>	<i>songolibila</i>	18.8
88	Annonaceae	<i>Enantia chlorantha</i>	<i>ep'houé</i>	94.3
89	Meliaceae	<i>Entandrophragma candollei</i>	<i>kangá</i>	75.3
90	Meliaceae	<i>Entandrophragma cylindricum</i>	<i>boyo</i>	103.6
91	Meliaceae	<i>Entandrophragma utile</i>	<i>bokoulo</i>	6.5
92	Sterculiaceae	<i>Eribroma oblongum</i>	<i>egwoyo</i>	101.0
93	Caesalpiniaceae	<i>Erythrophleum suaveolens</i>	<i>ngwanda</i>	95.2
94	Rutaceae	<i>Fagara heitzii</i>	<i>bolongo</i> 1	6.3
95	Rutaceae	<i>Fagara macrophylla</i>	<i>bolongo</i> 3	12.5
96	Bignoniaceae	<i>Fernandoa adolfi-friderici</i>	<i>bongo</i>	37.8
97	Moraceae	<i>Ficus macrosperma</i>	<i>djolo</i>	6.3
98	Moraceae	<i>Ficus etrangulator</i>	<i>bongo'o</i>	82.1
99	Moraceae	<i>Ficus exasperata</i>	<i>soubéme</i>	43.9
100	Moraceae	<i>Ficus mucuso</i>	<i>nguéhi</i>	12.6
101	Moraceae	<i>Ficus</i> sp.	<i>bambamessambo</i>	37.6
102	Apocynaceae	<i>Funtumia africana</i>	<i>kondo</i>	94.6
103	Apocynaceae	<i>Funtumia elastica</i>	<i>kondo</i>	25.2
104	Sapotaceae	<i>Gambeya boukokoensis</i>	<i>madjédjé</i>	56.5
105	Sapotaceae	<i>Gambeya lacourtiana</i>	<i>bambou</i>	95.1
106	Sapotaceae	<i>Gambeya perpulchra</i>	<i>boundouguèl</i>	6.3



## Appendix I. (continued)

No.	Family	Scientific name	Baka name	IVI
107	Clusiaceae	<i>Garcinia afzelii</i>	ngambé 2	25.1
108	Clusiaceae	<i>Garcinia kola</i>	gwel	12.5
109	Clusiaceae	<i>Garcinia lucida</i>	bako 'o	6.3
110	Clusiaceae	<i>Garcinia punctata</i>	p'handaka	75.2
111	Clusiaceae	<i>Garcinia</i> sp.1	bongoli	25.1
112	Clusiaceae	<i>Garcinia</i> sp.2	kèkèlè 2	62.7
113	Tiliaceae	<i>Glyphaea brevis</i>	andaka	56.5
114	Meliaceae	<i>Guarea cedrata</i>	djombo	102.2
115	Meliaceae	<i>Guarea</i> sp.	linga	31.3
116	Rubiaceae	<i>Hallea ledermannii</i>	moissé	12.5
117	Rubiaceae	<i>Hallea stipulosa</i>	langago	56.6
118	Olacaceae	<i>Heisteria zimmereri</i>	molomba 2	75.3
119	Annonaceae	<i>Hexalobus crispiflorus</i>	pota	102.3
120	Ulmaceae	<i>Holoptelea grandis</i>	bèlè	44.0
121	Flacourtiaceae	<i>Homalium africanum</i>	ngongo	6.3
122	Salicaceae	<i>Homalium aylmeri</i>	bambi	31.3
123	Salicaceae	<i>Homalium letestui</i>	tembo	69.1
124	Flacourtiaceae	<i>Homalium</i> sp.	djèkè	101.2
125	Apocynaceae	<i>Hunteria umbellata</i>	moundanga	56.4
126	Irvingiaceae	<i>Irvingia excelsa</i>	payo	81.8
127	Irvingiaceae	<i>Irvingia gabonensis</i>	péké	101.6
128	Irvingiaceae	<i>Irvingia grandifolia</i>	solia	94.9
129	Irvingiaceae	<i>Irvingia robur</i>	kombélé	18.8
130	Irvingiaceae	<i>Irvingia</i> sp.	bondoulou	25.2
131	Phyllanthaceae	<i>Keayodendron bridelioides</i>	mbôndo	50.4
132	Meliaceae	<i>Khaya anthotheca</i>	p'houa	18.8
133	Meliaceae	<i>Khaya grandifolia</i>	djila 2	43.8
134	Meliaceae	<i>Khaya ivorensis</i>	ngolo	12.6
135	Meliaceae	<i>Khaya</i> sp.	mboussoua 1	6.3
136	Irvingiaceae	<i>Klainedoxa gabonensis</i>	bukoko	102.8
137	Anacardiaceae	<i>Lannea welwitschii</i>	kwa	94.5
138	Sapotaceae	<i>Lasersisia</i> sp.	touba 1	100.3
139	Rhamnaceae	<i>Lasiodiscus manni</i>	souma	102.4
140	Sapindaceae	<i>Lecaniodiscus</i> sp.	elinga	6.3
141	Lepidobotryaceae	<i>Lepidobotrys staudtii</i>	moussako asséko	101.1
142	Sterculiaceae	<i>Leptonychia</i> sp.	mboké	18.8
143	Flacourtiaceae	<i>Lindackeria dentata</i>	loagwiyé	81.5
144	Euphorbiaceae	<i>Macaranga saccifera</i>	moussassah 2	101.6
145	Euphorbiaceae	<i>Macaranga spinosa</i>	moussassah 3	6.3
146	Rhamnaceae	<i>Maesopsis eminii</i>	londô	69.0
147	Euphorbiaceae	<i>Mallotus oppositifolius</i>	ngoka 1	50.1
148	Sterculiaceae	<i>Mansonia altissima</i>	babandja	82.6
149	Chrysobalanaceae	<i>Maranthes glabra</i>	bokandja	19.0
150	Phyllanthaceae	<i>Margaritaria discoidea</i>	kango	81.8
151	Bignoniaceae	<i>Markhamia lutea</i>	gondja	101.1
152	Annonaceae	<i>Meiocarpidium lepidotum</i>	mabêlenguè	95.3
153	Caesalpiniaceae	<i>Milbraediodendron excelsum</i>	ekéla 1	31.4
154	Moraceae	<i>Milicia excelsa</i>	bangui	94.6
155	Papilionaceae	<i>Millettia</i> sp.	olinga	12.5
156	Papilionaceae	<i>Millettia sanagana</i>	nganda	18.8
157	Cecropiaceae	<i>Musanga cecropioides</i>	kombo	107.3
158	Rubiaceae	<i>Mussaenda</i> sp.	lip'huètè	12.5
159	Cecropiaceae	<i>Myrianthus arboreus</i>	ngata	101.6

## Appendix I. (continued)

No.	Family	Scientific name	Baka name	IVI
160	Euphorbiaceae	<i>Neoboutonia glabrescens</i>	toubou	75.5
161	Moraceae	<i>Neosloetiopsis kamerunensis</i>	doundou	94.6
162	Sterculiaceae	<i>Nesogordonia papaverifera</i>	tétéké	101.2
163	Ixonanthaceae	<i>Ochthocosmus africanus</i>	likoumbi	25.1
164	Sterculiaceae	<i>Octobolus</i> sp.	mbéléngué 2	43.9
165	Sterculiaceae	<i>Octobolus spectabilis</i>	gagoulou	43.9
166	Caesalpiniaceae	<i>Oddoniodendron micranthum</i>	bolongo 2	87.9
167	Sapotaceae	<i>Omphalocarpum lecomteanum</i>	mbaté	6.3
168	Olacaceae	<i>Ongokea gore</i>	bossolo	25.2
169	Flacourtiaceae	<i>Ophiobotrys zenkeri</i>	mogwala	101.1
170	Caesalpiniaceae	<i>Oxystigma oxyphyllum</i>	gondo	37.8
171	Caesalpiniaceae	<i>Pachyelasma tessmannii</i>	ngouo	25.4
172	Annonaceae	<i>Pachypodanthium staudtii</i>	molombo	100.9
173	Pandaceae	<i>Panda oleosa</i>	kana	101.8
174	Chrysobalanaceae	<i>Parinari excelsa</i>	nombokola	6.3
175	Mimosaceae	<i>Parkia bicolor</i>	ndembé	25.4
176	Rubiaceae	<i>Pausinystalia johimbe</i>	wassassah	95.6
177	Rubiaceae	<i>Pavetta</i> sp.	loapoula	19.0
178	Menispermaceae	<i>Penianthus longifolius</i>	sombolo 1	6.3
179	Mimosaceae	<i>Pentaclethra macrophylla</i>	mbalaka	103.3
180	Papilionaceae	<i>Pericopsis elata</i>	mobai	44.2
181	Lecythidaceae	<i>Petersianthus macrocarpus</i>	bossô	95.4
182	Apocynaceae	<i>Picralima nitida</i>	poussah	87.9
183	Mimosaceae	<i>Piptadeniastrum africanum</i>	koungou	95.0
184	Euphorbiaceae	<i>Plagiostyles africana</i>	ngolé 1	50.1
185	Anisophylleaceae	<i>Poga oleosa</i>	p'houo	6.3
186	Annonaceae	<i>Polyalthia suaveolens</i>	botounga	104.2
187	Anacardiaceae	<i>Pseudospondias microcarpa</i>	ewoungou	75.7
188	Rubiaceae	<i>Psychotria avakubensis</i>	nokomindo	6.3
189	Rubiaceae	<i>Psychotria peduncularis</i>	gwègwèlè 2	18.8
190	Rubiaceae	<i>Psychotria</i> sp.1	bitongo	6.3
191	Rubiaceae	<i>Psychotria</i> sp.2	wélaleko	56.4
192	Rubiaceae	<i>Psychotria vogeliana</i>	unknown 5	18.8
193	Combretaceae	<i>Pteleopsis hylo dendron</i>	mobito	75.9
194	Papilionaceae	<i>Pterocarpus mildbraedii</i>	nguèlè (b)	31.8
195	Papilionaceae	<i>Pterocarpus soyauxii</i>	nguèlè (r)	89.5
196	Sterculiaceae	<i>Pterygota bequaertii</i>	mawouya 2	18.9
197	Sterculiaceae	<i>Pterygota macrocarpa</i>	ngoaka	6.3
198	Myristicaceae	<i>Pycnanthus angolensis</i>	tengué	102.4
199	Apocynaceae	<i>Rauvolfia caffra</i>	mbonga 1	43.9
200	Apocynaceae	<i>Rauvolfia grandifolia</i>	mbonga 2	25.1
201	Apocynaceae	<i>Rauvolfia macrophylla</i>	mbonga 3	6.3
202	Apocynaceae	<i>Rauvolfia vomitoria</i>	loli	37.6
203	Bombacaceae	<i>Rhodognaphalon brevicuspe</i>	tenonou	6.3
204	Euphorbiaceae	<i>Ricinodendron heudelotii</i>	gobo	104.7
205	Violaceae	<i>Rinorea aylmeri</i>	nguindi 1	50.1
206	Violaceae	<i>Rinorea elliotii</i>	sandjabongo	102.8
207	Violaceae	<i>Rinorea lepidobotrys</i>	nguindi 2	18.8
208	Violaceae	<i>Rinorea</i> sp.	técendé	95.9
209	Caesalpiniaceae	<i>Samanea dinklagei</i>	bokondo 1	12.6
210	Burseraceae	<i>Santiria trimera</i>	libaba 2	63.2
211	Caesalpiniaceae	<i>Scorodophloeus zenkeri</i>	minguègnè	38.5
212	Euphorbiaceae	<i>Spondianthus preussii</i>	godjo 2	65.5

## Appendix I. (continued)

No.	Family	Scientific name	Baka name	IVI
213	Myristicaceae	<i>Staudtia kamerunensis</i>	malanga	103.4
214	Caesalpiniaceae	<i>Stemonocoleus micranthus</i>	gondou	12.6
215	Sterculiaceae	<i>Sterculia subviolacea</i>	yébolo	82.1
216	Sterculiaceae	<i>Sterculia tragacantha</i>	bototo	31.4
217	Olacaceae	<i>Strombosia pustulata</i>	bobongo	103.3
218	Olacaceae	<i>Strombosiopsis</i> sp.	p'huindo	50.4
219	Olacaceae	<i>Strombosiopsis tetrandra</i>	bossiko	102.4
220	Loganiaceae	<i>Strychnos ternata</i>	dingwa	18.8
221	Loganiaceae	<i>Strychnos</i> sp.2	boukou	6.3
222	Caesalpiniaceae	<i>Swartzia fistuloides</i>	eloukou	12.5
223	Caesalpiniaceae	<i>Swartzia</i> sp.	ganda	18.8
224	Sapotaceae	<i>Synsepalum aubrevillei</i>	djingou	75.2
225	Myrtaceae	<i>Syzygium rowlandii</i>	essossi	25.2
226	Apocynaceae	<i>Tabernaemontana crassa</i>	paadok	50.2
227	Combretaceae	<i>Terminalia superba</i>	ngolou	109.9
228	Caesalpiniaceae	<i>Tessmania africana</i>	paka 2	31.3
229	Mimosaceae	<i>Tetrapleura tetraptera</i>	djaga	100.5
230	Euphorbiaceae	<i>Tetrorchidium didymostemon</i>	djènè	62.7
231	Thomandersiaceae	<i>Thomandersia hensii</i>	ngoka 2	25.0
232	Thomandersiaceae	<i>Thomandersia laurifolia</i>	ngoka 3	6.3
233	Sapotaceae	<i>Tieghemella africana</i>	kolo	57.7
234	Ulmaceae	<i>Trema orientalis</i>	messiongo	25.2
235	Rubiaceae	<i>Trichalysia</i> sp.1	molomba 1	50.2
236	Rubiaceae	<i>Trichalysia</i> sp.2	molomba 2	6.3
237	Meliaceae	<i>Trichilia heudelottii</i>	mayimbo 1	104.3
238	Meliaceae	<i>Trichilia welwitschii</i>	mayimbo agwanga	102.6
239	Anacardiaceae	<i>Trichoscypha acuminata</i>	ngoyo	100.3
240	Sapotaceae	<i>Tridesmostemon omphalocarpoides</i>	touba 2	31.3
241	Moraceae	<i>Trilepisium madagascariense</i>	pongui	103.7
242	Sterculiaceae	<i>Triplochiton scleroxylon</i>	gwado	60.4
243	Euphorbiaceae	<i>Uapaca guineensis</i>	séngui	97.8
244	Rutaceae	<i>Vepriis louisii</i>	tanda	50.1
245	Verbenaceae	<i>Vitex grandifolia</i>	p'houlou 2	101.0
246	Annonaceae	<i>Xylophia hypolampra</i>	moundjiè	31.4
247	Annonaceae	<i>Xylophia phloidora</i>	sangué	100.9
248	Annonaceae	<i>Xylophia</i> sp.	loabano	87.9
249	Rutaceae	<i>Zanthoxylum laurentii</i>	djoumgwé	6.3

We did not include two species of No. 58 and No. 85 into our analysis due to a high ambiguity of identifications; the total number of species, therefore are 247 in the body text.



39	<i>Ceiba pentandra</i>	2	0.10	2	0.12	1	0.32	5	2.97	1	1.47	10	0.67	16	0.30	13	0.63	20	0.42	15	0.85	8	0.50	16	0.73	11	0.80	11	0.38	24	1.46			
40	<i>Celtis adolfi-friederici</i>	13	0.16	11	0.35	19	1.26	7	0.20	12	0.20	14	0.98	10	0.78	27	1.44	12	0.43	30	1.32	17	0.76	24	1.07	12	0.37	20	0.36	13	1.15	22	1.95	
41	<i>Celtis mildbraedii</i>	28	1.93	36	2.43	34	1.54	23	0.91	26	1.41	31	1.55	20	0.78	27	1.44	12	0.43	30	1.32	17	0.76	24	1.07	12	0.37	20	0.36	13	1.15	22	1.95	
42	<i>Celtis philippensis</i>	3	0.17	14	1.22	2	0.38	8	0.33	30	1.49	12	0.80	3	0.17	5	0.29	8	0.32	1	0.01	7	0.16	3	0.13	4	0.08	2	0.07	20	0.89	7	0.39	
43	<i>Celtis tessmannii</i>	5	0.31	7	0.29	9	0.30	5	0.09	1	0.04	6	0.10	4	0.03	3	0.05	8	0.21	4	0.33	5	0.09	3	0.02	4	0.06	2	0.48	3	0.09			
44	<i>Celtis zenkeri</i>																																	
45	<i>Chytranthus atroviolaceus</i>																																	
46	<i>Citropsis</i> sp.																																	
47	<i>Clastopholis patens</i>	2	0.02	5	0.13	1	0.03	5	0.10	4	0.03	3	0.16	5	0.17	3	0.06	3	0.13	1	0.04	3	0.09	3	0.03	5	0.05	7	0.16	3	0.04	2	0.10	
48	<i>Coelocaryon preussii</i>	5	0.20	18	0.43	21	0.43	38	0.75	19	0.40																							
49	<i>Cola acuminata</i>	7	0.05	7	0.05	3	0.05	10	0.09	8	0.12	21	0.23	15	0.13	8	0.10	15	0.24	6	0.07	11	0.08	7	0.05	13	0.09	7	0.05	12	0.15	8	0.06	
50	<i>Cola allissima</i>	4	0.19	3	0.08	1	<0.01			2	0.22			6	0.05	2	0.02	3	0.02	1	<0.01	2	<0.01	5	0.02	3	0.02	2	0.03			2	0.03	
51	<i>Cola ballayi</i>													5	0.07	12	0.12	9	0.18	2	0.01	2	<0.01	5	0.05	1	<0.01	1	<0.01	1	<0.01	1	<0.01	
52	<i>Cola laterita</i>	1	0.05	2	0.60	7	0.38	13	0.52	15	3.01	12	0.60	13	0.53	6	0.10	14	0.90	15	0.18	15	0.17	23	1.17	23	0.66	16	0.44	20	1.27	23	2.82	
53	<i>Cola pachycarpa</i>																																	
54	<i>Copaifera mildbraedii</i>																																	
55	<i>Cordia platyhyrsa</i>	1	0.05	1	0.07																													
56	<i>Corynanthe pachyceras</i>	11	0.12	9	0.11	18	0.35	11	0.25	24	0.50	15	0.20	10	0.13	9	0.19	14	0.23	22	0.37	24	0.49	13	0.18	26	0.53	12	0.18	30	0.58	28	0.58	
57	<i>Cylocodiscus gabunensis</i>	2	1.27	2	0.03									2	0.03	5	0.10	3	0.54	2	0.83	8	1.50	4	0.23	6	0.65	3	0.04	10	0.72	5	0.46	
58	<i>Cyrtanthus congolensis</i>													1	<0.01																			
59	<i>Daeryodes klainiana</i>	1	0.05	8	0.18	11	0.20	1	0.02	10	0.13	5	0.11	8	0.07	9	0.08	9	0.36	13	0.13	11	0.17	13	0.15	13	0.12	8	0.07	10	0.13	5	0.06	
60	<i>Desplatsia dewevrei</i>	4	0.11	4	0.07	1	0.01	9	0.19	10	0.13																							
61	<i>Desplatsia</i> sp.																																	
62	<i>Detarium macrocarpum</i>	1	<0.01	1	0.92									1	0.83																			
63	<i>Dialium dinklagei</i>																																	
64	<i>Dialium</i> sp.																																	
65	<i>Dichaetanthera africana</i>																																	
66	<i>Dichaetanthera</i> sp.	17	0.13	15	0.38	16	0.09	11	0.07	3	0.02	5	0.03	36	0.24	37	0.29	57	0.43	27	0.19	57	0.34	44	0.27	74	0.49	35	0.21	2	0.01	17	0.11	
67	<i>Dichostemma glaucescens</i>																																	
68	<i>Dieranalepis disticha</i>																																	
69	<i>Dieranalepis</i> sp.																																	
70	<i>Diospyros abyssinica</i>																																	
71	<i>Diospyros iturensis</i>	3	0.04	8	0.08	5	0.05	12	0.11					2	0.02	2	0.02	6	0.07	11	0.07	4	0.02	7	0.03	3	0.02	4	0.02	1	<0.01	1	<0.01	
72	<i>Diospyros canaliculata</i>	1	<0.01	2	0.02	3	0.02	2	0.05	5	0.05	1	<0.01																					
73	<i>Diospyros crassiflora</i>	2	0.02	7	0.40	1	<0.01	2	0.03	3	0.02	2	0.18	7	0.46	14	0.29	9	0.30	11	0.30	8	0.07	10	0.09	17	0.25	16	0.26	13	0.15	19	0.39	
74	<i>Diospyros</i> sp.1													2	0.05	2	0.04	3	0.04															
75	<i>Diospyros</i> sp.2	9	0.04	15	0.12	6	0.05	10	0.08	36	0.33	14	0.14	6	0.04	10	0.06	16	0.13	7	0.06	12	0.07	2	0.02	4	0.04	6	0.04	37	0.26	11	0.07	
76	<i>Discoglypemma caloneura</i>	3	0.05	1	0.01	1	0.05	2	0.25	4	0.10	1	0.06	4	0.02	4	0.07	1	0.01	3	0.11	1	0.11	2	0.03	2	0.02	2	0.03	3	0.04			
77	<i>Distemonanthus benthamiana</i>	1	<0.01	1	0.02	1	<0.01	1	<0.01	1	0.02			1	0.18	5	0.84																	
78	<i>Donella ubanguiensis</i>	1	<0.01																															
79	<i>Drypetes</i> sp.1	5	0.03	20	0.29	5	0.03	9	0.07	32	0.22	10	0.08	31	0.22	14	0.08	26	0.19	9	0.04	17	0.09	16	0.07	11	0.06	4	0.02	15	0.11	6	0.05	

Appendix II. (continued)

No.	Species	T1		T2		T3		T4		T5		T6		T7		T8		T9		T10		T11		T12		T13		T14		T15		T16				
		IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA	IN	BA			
80	<i>Drypetes gossweileri</i>	3	0.05	3	0.28	16	0.88	7	0.23	1	0.03	4	0.03	3	0.04	3	0.03	1	0.02	7	0.47	5	0.12	2	<0.01	2	0.04	4	0.12	5	0.11	5	0.03			
81	<i>Drypetes turinensis</i>	6	0.08	1	<0.01	9	0.09	1	0.02	4	0.04	1	<0.01	7	0.06	1	0.02	2	0.02	1	<0.01															
82	<i>Drypetes principum</i>	1	<0.01																																	
83	<i>Drypetes</i> sp.2	2	<0.01																																	
84	<i>Duboisia macrocarpa</i>	13	0.18	10	0.58	7	0.84	15	0.65	16	0.89	12	0.51	5	0.48	7	0.94	9	0.25	9	0.47	5	0.74	18	1.49	14	2.23	9	0.20	7	0.81	16	1.80			
85	<i>Euphorbia drupifera</i>	1	<0.01			1	<0.01			2	<0.01																									
86	<i>Enantia chlorantha</i>	2	0.01	6	0.13	3	0.05	5	0.04	4	0.08	5	0.19	7	0.31	4	0.06	5	0.16	2	0.04	4	0.11	7	0.14	3	0.04	3	0.04			2	0.02			
87	<i>Entandrophragma candollei</i>	2	0.01	1	<0.01	1	<0.01					1	0.04			1	0.04			2	0.02	2	0.02	3	0.03	3	0.02	3	0.46	4	0.07	7	0.15			
88	<i>Entandrophragma cylindricum</i>	9	2.62	11	2.24	8	0.77	3	1.08	2	0.15	6	0.69	8	0.85	9	0.56	7	0.73	15	2.38	5	1.07	12	2.84	8	1.23	6	1.69	6	1.22	3	0.02			
89	<i>Entandrophragma utile</i>																																			
90	<i>Eribronia oblongum</i>	1	0.02	2	0.04	3	0.68	3	0.04	3	0.40	2	0.08	3	0.29	7	0.08	4	0.06	7	0.26	9	0.62	8	0.72	7	0.19	3	0.06	7	0.31	1	1.47			
91	<i>Erythrophleum suaveolens</i>	3	0.34	2	0.32	4	1.18	1	0.03			3	0.51	2	0.22	4	0.62	4	0.50	3	0.52	3	0.85	4	0.39	3	0.31	3	0.65	3	1.27	2	0.59			
92	<i>Fagara heitzii</i>	3	0.01																																	
93	<i>Fagara macrophylla</i>																																			
94	<i>Fernandoa adolfi-friderici</i>	2	0.06	4	0.23							5	0.37	1	0.02																					
95	<i>Ficus macrosperma</i>	2	0.16																																	
96	<i>Ficus errangulata</i>	1	0.03							2	0.08	3	0.12	2	0.09	4	0.36	7	1.15	3	0.34	2	0.06	3	0.06	7	0.17	1	0.03	8	0.62			10	0.48	
97	<i>Ficus exasperata</i>			2	0.02							1	<0.01	15	0.20	1	0.01			1	<0.01	1	0.07													
98	<i>Ficus mucosa</i>	16	0.13			1	0.16																													
99	<i>Ficus</i> sp.			3	0.40	2	0.12	1	0.04	1	0.01																									
100	<i>Funtumia africana</i>	5	0.09	6	0.12	4	0.09	4	0.12			5	0.12	4	0.12	6	0.17	4	0.02	5	0.15	9	0.19	6	0.08	10	0.21	8	0.19	6	0.11	8	0.34			
101	<i>Funtumia elastica</i>	7	0.11							15	0.22																									
102	<i>Gambeya baukokoensis</i>									3	0.05	1	0.02	2	0.33	2	0.03			2	0.54			2	0.04	1	<0.01						2	<0.01	3	0.04
103	<i>Gambeya lacouriana</i>	2	0.02			1	<0.01	4	0.37	5	0.31	4	0.09	5	0.54	12	0.34	9	0.11	13	0.15	11	0.10	13	0.31	20	1.23	11	0.10	8	0.12	13	0.27			
104	<i>Gambeya perpulchra</i>									1	<0.01																									
105	<i>Garcinia afzeli</i>	1	<0.01			5	0.03	2	0.03	1	<0.01																									
106	<i>Garcinia kola</i>	2	0.02																																	
107	<i>Garcinia lucida</i>			1	0.01																															
108	<i>Garcinia punctata</i>	3	0.03	4	0.05	8	0.06	2	0.02			2	0.02	1	<0.01																					
109	<i>Garcinia</i> sp.1	1	<0.01	1	0.02			6	0.07	7	0.14																									
110	<i>Garcinia</i> sp.2			2	<0.01					1	<0.01	1	0.03	1	0.02																					
111	<i>Glyphaea brevis</i>			1	<0.01					1	<0.01																									
112	<i>Guarea cedrata</i>	12	0.22	9	0.28	14	0.66	8	0.06	2	0.01	12	0.16	25	0.38	20	0.38	17	0.20	32	0.39	19	0.45	24	0.63	19	0.45	15	0.17	19	0.36	11	0.28			
113	<i>Guarea</i> sp.	1	0.01	1	0.02	2	0.02	2	0.02	1	<0.01																									
114	<i>Hallea ledermannii</i>			1	<0.01																															
115	<i>Hallea stipulosa</i>	4	0.25			2	0.11	2	0.01			1	<0.01																							
116	<i>Heisteria zimmereri</i>	1	<0.01			3	0.03	1	0.06	1	<0.01	2	0.01	6	0.06			10	0.12	4	0.05	6	0.05	4	0.02											
117	<i>Hexalobus crispiflorus</i>	14	0.42	19	0.86	9	0.40	4	0.25	8	0.33	17	0.33	17	0.30	21	0.70	15	0.28	12	0.53	17	0.38	18	0.27	13	0.34	7	0.22	15	0.54	23	0.61			









