

## On Trees in Paddy Fields in Northeast Thailand

Hiroyuki WATANABE\*, Ken-ichi ABE\*\*, Tomoyuki HOSHIKAWA\*\*\*,  
Buared PRACHAIYO†, Pongsak SAHUNALU†† and Choob KHEMNARK††

### I Introduction

In contrast to the open spread of paddy fields in the Central Plain (Chao Praya Delta), northeast Thailand is strikingly characterized by the presence of trees. According to the land assessment using the LANDSAT imagery [Wacharakitti 1979], paddy and farm land where trees stand sparsely account for about 60% of the whole area of Khon Kaen and Mahasarakham provinces. The scenery of paddy field and farmland with trees is very common in northeast Thailand.

### II Research Sites and Methods

Research on the location and density of trees was done at Ban Sok Muang, Ban Tasala and Ban Non Tan, located

between Khon Kaen and Chumpae, along the national road the Route 12, and Ban Phone Phek, located at Phu Wiang district, along the Route 2038, in the vicinity of Khon Kaen (all sites are 200–240 meters above mean sea level).

Research was made in September 1986, and also in November 1986 after paddy fields dried up and the paddy was harvested.

At each research site, quadrats (50 m × 60 m–50 m × 100 m) were set up, location of trees was mapped; their diameter (DBH) and size of tree crown were estimated; and tree species were identified. In addition, locations of stump, snags and termite mound, etc., were also mapped when found. According to the information obtained at Ban Non Tan and Ban Phone Phek, about 20 years has passed since reclamation. At other two sites, a maximum of 20 years has also elapsed. Paddy fields are not irrigated.

### III Results and Discussion

#### *Tree Species and Density*

Table 1 shows the tree density, average diameter, basal area and tree species. In addition, Fig. 1 shows the size

---

\* 渡辺弘之, Faculty of Agriculture, Kyoto University, Kyoto 606, Japan

\*\* 阿部健一, The Center for Southeast Asian Studies, Kyoto University

\*\*\* 星川智之, Kasugai-nishi High School, Kasugai, Aichi 486, Japan

† Khon Kaen Regional Forest Office, Khon Kaen 40000, Thailand

†† Faculty of Forestry, Kasetsart University, Bangkok 10900, Thailand

**Table 1** Tree Density, Average Diameter and Basal Area

	Density/ha	Average diameter cm (Range)	Basal area m <sup>2</sup> /ha
Ban Phone Phek			
<i>Terminalia tomentosa</i>	20.0		2.35
<i>Shorea obtusa</i>	6.7		1.75
<i>Terminalia chebula</i>	3.3		0.26
Total	30.0	42.0 (31.7 - 61.6)	4.37
Ban Sok Muang			
<i>Dipterocarpus tuberculatus</i>	50.0		6.22
<i>Shorea obtusa</i>	13.3		1.07
<i>Dipterocarpus obtusifolius</i>	3.3		0.95
Total	66.6	38.7 (25.5 - 60.3)	8.24
Ban Non Tan			
<i>Diospyros rhodocalyx</i>	46.0		0.94
<i>Sindora siamensis</i>	12.0		0.19
<i>Lagerstroemia calyculata</i>	6.0		0.21
<i>Nauclea orientalis</i>	4.0		0.34
<i>Shorea siamensis</i>	2.0		0.18
<i>Dalbergia nigrescens</i>	2.0		0.17
<i>Terminalia chebula</i>	2.0		0.12
<i>Xylia xylocarpa</i>	2.0		0.10
<i>Pterocarpus macrocarpus</i>	2.0		0.02
<i>Diospyros mollis</i>	2.0		0.02
<i>Canarium kerrii</i>	2.0		0.03
<i>Cassia garrettiana</i>	2.0		0.02
Total	84.0	17.4 (8.9 - 36.4)	2.35
Ban Tasala			
<i>Shorea obtusa</i>	108.6		6.81
<i>Dipterocarpus tuberculatus</i>	34.3		3.33
<i>Xylia xylocarpa</i>	2.9		0.42
<i>Terminalia tomentosa</i>	2.9		0.33
Total	148.7	30.0 (20.7 - 43.5)	10.89

class distribution and Fig. 2 the location maps of trees in each sites. The least density (30 trees/ha, basal area 4.4 m<sup>2</sup>/ha) was recorded at Ban Phone Phek where average diameter (DBH 42 cm) was highest. It distinctly shows

that big trees were left sparsely. The maximum (148.7 trees/ha and 10.9 m<sup>2</sup>/ha) density was recorded at Ban Tasala. The size of *Dipterocarpus tuberculatus* is usually bigger than that of *Shorea obtusa*.




Fig. 1 Size Class Distribution of Trees at Each Site






Fig. 2 Maps of Tree Location at Each Site



**C**  
Ban Non Tan (100m×50m)

- C. g. *Cassia garrettiana*
- C. k. *Canarium kerrii*
- D. m. *Diospyros mollis*
- D. n. *Darbergia nigrescens*
- D. r. *Diospyros rhodocalyx*
- L. c. *Lagerstroemia calyculata*
- N. o. *Nauclea orientalis*
- P. m. *Pterocarpus macrocarpus*
  
- Sh. s. *Shorea siamensis*
- S. s. *Sindora siamensis*
- T. c. *Terminalia chebula*
- X. x. *Xylia xylocarpa*

- Termite mound
- Paddy dike
- Crown size

Fig. 2 Continued




Fig. 2 Continued

Tree species also vary greatly from site to site. At Ban Phone Phek with the lower density, *Terminalia tomentosa* was the major tree mixed with other two *Shorea obtusa* and *T. chebula*. At Ban Sok Muang, *Dipterocarpus tuberculatus* was dominant, and *S. obtusa* and *D. obtusifolius* were also found. These trees belong to the Dipterocarpaceae.

On the contrary, at Ban Tasala, *S. obtusa* was dominant and *D. tuberculatus* was the next dominant, and *Xylia xylocarpa* and *T. tomentosa* occurred instead of *D. obtusifolius*. At Ban Non Tan, twelve tree species, *Diospyros rhodocalyx*, *D. mollis*, *Sindora siamensis*, *Lagerstroemia calyculata*, *Nauclea orientalis*, *Dalbergia nigrescens*, *Canarium kerrii*, *Cassia garrettiana* and *Shorea siamensis* were found while total basal area was the lowest. The difference in the composition of tree species clearly reflects the differences in the original


vegetation and there was no tree recognized as planted or introduced after reclamation to paddy fields.

Wacharakitti [1979] described that the average density of trees in an undisturbed Dry Dipterocarp Forest in this area was 468 trees/ha and that of paddy field was 51/ha with a volume of 12.7 m<sup>3</sup>/ha (equivalent to 19.3 t/ha). In this study the maximum density in the paddy field was 150/ha but clearly much less than density in the surrounding forest.

Regarding trees found in the paddy fields, Grandstaff *et al.* [1986] listed 54 species at least 18 of which were evidently introduced, including lontar palm (*Borassus flabellifer*) and raintree (*Samanea saman*). They suggested that density and composition depended on ages of the paddy fields, among other factors.



A : Ban Phone Phek



B : Ban Sok Muang



C : Ban Non Tan



D : Ban Tasala

**Fig. 3** Scenery of Trees in the Paddy Fields

**Table 2** Uses of Trees in Paddy Fields

Local name	Scientific name	Major use	Specific use
Dipterocarpaceae			
Hiang/Saad	<i>Dipterocarpus obtusifolius</i>	Trunk : C	Resin : Torch
Kung/Pluang	<i>D. tuberculatus</i>	Trunk : C	Leaf : Wrapping, Resin : Torch
Chik/Teng	<i>Shorea obtusa</i>	Trunk : C, F	
Rang/Hang	<i>S. siamensis</i>	Trunk : C, F	
Combretaceae			
Rok Fah	<i>Terminalia chebula</i>	Trunk : T	Fruit : Tanning, Edible
Sa Mo Tai	<i>T. tomentosa</i>	Trunk : T	Bark : Dyeing, Tanning
Ebenaceae			
Muklua	<i>Diospyros mollis</i>	Trunk : T, F	Fruit : Dyeing, Medicine
Ta Ko Na	<i>D. rhodocalyx</i>	Trunk : T, F	Fruit : Edible, Dyeing
Leguminosae			
Khi Lek Pa	<i>Cassia garrettiana</i>	Trunk, Branch : F	
Pradu	<i>Pterocarpus macrocarpus</i>	Trunk : C, T	Leaf : Fodder, Bark : Dyeing
Tae	<i>Sindora siamensis</i>	Trunk : T, F	
Cha Nuan	<i>Dalbergia nigrescens</i>	Trunk : T, F	
Mai Daeng	<i>Xylocarpus xylocarpa</i>	Trunk : C, T, F	
Rubiaceae			
Kra Tum Nam	<i>Nauclea orientalis</i>	Trunk : C, T, F	
Lythraceae			
Tabaek	<i>Lagerstroemia calyculata</i>	Trunk : T	
Burseraceae			
Ma Kerm (Ma Kok Luem)	<i>Canarium kerrii</i>	Trunk : T	Fruit : Edible

C : Construction T : Tool and Furniture F : Fuel

*Use of Trees*

Table 2 shows the tree species found in the study sites and their uses indicated in interviews with farmers. Most important and daily uses of trees are 1) to gather fuel wood by collection of fallen branches and branches pruned to moderate the shade to paddy, and 2) shade for livestock and human in intervals of work. As described by Subhadhira *et al.* [1988], shortage of fuelwood due to the conversion of woodland to agricultural land in this region has become a seri-

ous problem.

In addition to the uses mentioned above, large leaves of *Dipterocarpus tuberculatus* are used for wrapping. Young leaves of Legume trees, such as *Pterocarpus macrocarpus* and *Cassia garrettiana*, etc., are fodder for livestock, and fruit of *Terminalia chebula*, *T. tomentosa*, *Diospyros mollis* and *D. rhodocalyx*, are for tanning and dyeing. Of course, big trunks of *Dipterocarpus tuberculatus*, *D. obtusifolius* and *Shorea obtusa* are used for construction and also making tool or furniture. Surely,





Fig. 4 Eucalyptus Planted on the Paddy Dikes (at Somdet)

those big trees are valuable property.

It can be said that trees in the paddy fields is usually left not for one specific purpose of use, but rather for multi-purpose or multi-uses. Farmers are still wedded to this traditional agro-forestry systems. For example, at Somdet near Mahasarakham, *Eucalyptus camaldulensis* is planted on the paddy dikes where paddy fields were newly opened and no native trees left (Fig. 4).

#### Acknowledgments

We wish to thank the late former Dean, Dr. Sathit Wacharakitti, and former Dean Dr. Somsak Sukwong, Faculty of Forestry, Kasetsart University, Bangkok for invaluable

advice throughout the present study. We also thank Ms. Michiko Takagaki, Faculty of Agriculture, Chiba University, Chiba, Japan for her assistance during our field study. Lastly, we are indebted to the National Research Council of Thailand (NRCT) and Khon Kaen Regional Forest Office for their support.

We would also like to thank Dr. Terry B. Grandstaff, KCU-Ford Rural System Research Project, Khon Kaen University, for his helpful comments on the manuscript and for improving the English text.

#### References

- Grandstaff, S. W. ; Grandstaff, T. B. ; Rathakette, P. ; Thomas, D. E. ; and Thomas, J.K. 1986. Trees in Paddy Fields in Northeast Thailand. In *Traditional Agriculture in Southeast Asia*, edited by G. G. Marten, pp. 273-292. Boulder & London : Westview Press.

- Phengkklai, C.; and Khamsai, S. 1985. Some Non-timber Species of Thailand. *Thai For. Bull.* 15: 108-148.
- Rathakette, P.; Somnasang, P.; Ratanapanya, S.; and Homchoen, S. 1985. Taboos and Traditions: Their Influence on the Conservation and Exploitation of Trees in Social Forestry Projects in Northeastern Thailand. In *Community Forestry: Socio-economic Aspects*, edited by Y.S. Rao; N.T. Vergara.; and G.W. Lovelace, pp.363-369. Bangkok: RAPA. FAO.
- Sahunalu, P. 1984. *Ecological Aspects of Agroforestry Systems in Thailand*. Technical Paper No.51. Bangkok: Fac. of Forestry, Kasetsart Univ. 28 p.
- Subhadhira, S.; Suphanchaimat, N.; Smutkupt, S.; Simaraks, S.; Pakuthai, W.; Subarnbhesaj, K.; and Petchsingha, P. 1988. Fuel Wood Situation and Farmer's Adjustment in Northeastern Thai Villages. In *Rapid rural Appraisal in Northeast Thailand* (Case studies), edited by W. George, *et al.*, pp.29-53. Khon Kean: KKU-Ford Rural System Research Project, Khon Kaen University.
- Takaya, Y.; and Tomosugi, T. 1972. Rice Lands in the Upland Hill Regions of Northeast Thailand-A Remark on "Rice Producing Forest." *Southeast Asian Studies* 10 (1): 77-85. (in Japanese with English summary)
- Vityakon, P.; Smutkupt, S.; and Prachaiyo, B. 1988. Trees in Paddy Fields: Their Contributions to Soil Fertility, and Sustainability of the Paddy Rice System. In *Sustainable Rural Development in Asia-Selected Papers from the 4th SUAN Regional Symposium on Agroecosystems Research Held at Khon Kaen University*, edited by T. Charoenwatana; and A.T. Rambo, pp.65-86. Khon Kaen.
- Wacharakitti, S. 1979. *Land Use*. Nam Pong Environmental Management Research Project Working Document No.3. Bangkok: Mekong Secretariat. 68 p.
- Wacharakitti, S.; Ruangpanit, N.; Sahunalu, P.; and Sanguantham, P. 1972. Forest Land Assessment of Khon Kaen and Mahasarakham Provinces. *Thai Jour. For.* 1(2): 1-22.