

## THE FLORA OF BOSSOU : ITS UTILIZATION BY CHIMPANZEES AND HUMANS

Yukimaru SUGIYAMA

*Primate Research Institute, Kyoto University*

Jeremy KOMAN

*Laboratoire Monts Nimba*

**ABSTRACT** Altogether 664 plant species from 392 genera have been identified and listed as the flora of Bossou, in the south-eastern corner of the Republic of Guinea. Previously, Bossou must have been covered with a primary and mature forest. However, due to destruction of forest, 60% of the 265 tree species identified were small trees of less than 10 m in height with 55% found mostly in secondary forest. Chimpanzees use 246 items (parts of plants) from 200 different plant species for food, but people use only 83 items from 76 species for food. A characteristic feature of human plant use is as traditional medicine, representing 113 items from 81 species. People also use particular species of plants as material for house construction, furniture and other purposes in order to make human life more convenient and comfortable.

**Key Words:** Flora; Bossou; Guinea; Traditional medicine; Deforestation.

### INTRODUCTION

A description of whole plant species and their traditional uses in a particular area or by a tribe is important basic information for the utilization of natural resources and the conservation of animals and their environment. Deforestation is rapidly expanding at present throughout the tropical world. However, in recent years there have been few listings of the flora in remaining forests. The effort required for identification of each plant is extraordinarily large and there are fewer plant taxonomists than before.

The present authors have been studying the behavior and ecology of chimpanzees (*Pan troglodytes verus*) at Bossou, south-eastern corner of the Republic of Guinea, west Africa, since 1976. An attempt has been made to describe all of the plants in this area as deforestation is destroying the original vegetation and the chimpanzee habitat. Deforestation combined with urbanization forces local people to forget their traditional habits concerning the use of plants and other natural resources. The following is a preliminary list of the flora of Bossou and its utilization by chimpanzees and local people as far as the authors have recognized until this point. We expect it will be revised in future as research continues.

## STUDY AREA AND METHODS

Bossou is located at the peripheral part of the tropical rain forest. It is 1,100 km from Conakry, the capital of Guinea, by road, and was far from civilization for many years (Fig. 1). Even when the present authors began their study in 1976, however, much of Bossou was already deforested due to shifting cultivation, and the lower part of the hills which surrounds the village was changing to secondary forest (Fig. 2, Fig. 3). The primary forest was found only in the higher areas and valleys of the small hills (Fig. 4), reaching to 710 m above sea level at the highest plateau. The village of Bossou is about 550 m above sea level. The lower parts of the hills contained patches of secondary forest and cultivated fields (Sugiyama & Koman, 1979). The study area is connected with Mt. Nimba by savanna which is burned by the people every dry season (Fig. 5).

Meteorological records are available only from the Nimba region of Liberia, which is within 10 km of Bossou (Fig. 6). Rain fall is about 2,500–3,000 mm a year, mostly between the middle of March and early November.

The identification of plants was done mainly by Koman through a comparison of specimens with the collection of the Institute of Medical Plant at Seredou, Macenta, and its confirmation was done by Sugiyama. "Flore descriptive des Monts Nimba" (Adam, 1971a; 1971b; 1975a; 1975b) and "Flora of west tropical Africa" (Hutchinson & Dalziel, 1954; 1958; 1963; 1968; 1972) were effectively used for these purposes.






Fig. 1. Map of Bossou and its adjacent area with distribution range of chimpanzees. Shadow: chimpanzee distribution is confirmed, double shadow: chimpanzees are recently extinct.




**Fig. 2.** Study area. Circles with a small triangle are the higher parts of hills which are surrounded by secondary open forests, cultivated fields and, then, savanna. Shadow is the village part. The circle enclosed by break line is the approximate range of the present study.



**Fig. 3.** Deforested slope of Guein. Only the plateau is covered with the closed canopy.




**Fig. 4.** Forests of Ban. The core-area of chimpanzees' territory which is covered with dense forests.



**Fig. 5.** South-eastern periphery of study area. Bossou and Nimba Mountains are connected with savanna of about 3 km.

The utilization of plants by chimpanzees as food was one of the primary targets of the study and was confirmed mainly by Sugiyama through direct observations (revised from Sugiyama & Koman, 1987). Information on the traditional use of plants by local people was collected through local assistants and older villagers. Human utilization was described only when people indicated a particular kind of plant. Non-selective use of plants was not included.



**Fig. 6.** Walter's climatograph at two adjacent points. Geologist's camp at Mt. Nimba (circles) is 1,340 m above sea level and Yekepa (triangles) is 500 m. The black marks show monthly rain fall between 1957 and 1964; 3,509 mm at Geologist's camp and 1,895 mm at Yekepa, average per year. Open marks show mean daily temperature per month between 1960 and 1964 (Data from Adam, 1971a).

## RESULTS AND DISCUSSION

Altogether, 664 species from 392 genera of plants were identified (See Appendix). Supposedly, Bossou was covered with primary and/or mature forests in the old days. However, much of this area is now covered with secondary forest, cultivated field and savanna, as a result of the heavy deforestation by shifting cultivation and field burning. Sixty percent of tree species, 159 out of 265, are small trees of less than 10 m in height, while more than half of them are found in the secondary forest (Table 1). The composition of plant species clearly shows the disappearing feature of primary and mature forest around Bossou.

Chimpanzees are basically frugivorous and forest dwellers. Everyday they travel from food patch to food patch, mainly searching for fruits and young leaves. They chose 246 different items (parts of plants) from 200 plant species as food, representing about 30% of the available species (Table 2). On the other hand, local people used only 83 different items from 76 plant species as food, 11% of all species. More than half of these, the people obtained from uncultivated areas. In spite of this, the people concentrate their interest on a far smaller number of plant species for food and get much of their volume through cultivation.

*Oryza* sp., *Manihot* spp. and *Musa* spp. are the most important foods for humans and are all cultivated. Fruits of *Citrus* spp., *Carica papaya*, as well as a few other trees are also important, but they represent seasonal foods and are planted. Fruits of *Mangifera indica*, that of *Persea gratissima*, and nuts and resin

**Table 1.** Number of plant species in each vegetation type.

Plant type	Vegetation type									
	Forest			Savanna	Cultivated field	Abandoned field	Marigo	Ruderal	Other	Total
dense	Transitional	secondary								
Tree										
big	19	4	1			3				27
middle	1	29	39	1	3	1	5			79
small	1	36	87	6	17	6	5	1		159
Bush/scrub		4			3	1				8
Grass/herb	36	40	70	39	38	35	18	2		278
Liana	15	75	2	6	15					113
Total	2	139	245	80	68	64	45	19	2	664

Plants which are found in two grades of forest are recorded as transitional forest in this table for convenience. Other plants which are distributed in more than one vegetation type are recorded in the one where they are found more frequently.

**Table 2.** Number of plant items used by chimpanzees and humans.

	Fruit	Leaf	Flower	Shoot	Tubercl	Kernel Seed	Bark	Twig Stem	Gum Resin	Tissue fibre	Wood, whole plant	Total
											Item	Sp
Chimpanzee												
food	128	44	11		6	16	4	33	3	1	246	200
Human	41	87	7	2	6	21	20	27	5	2	239	198
food/drink	37	18	4		3	16	1		4		83	76
(uncultivate)	(20)	(11)				(8)	(1)		(4)		(44)	(37)
medicine	1	62	3	2	2	5	19	18	1		113	81
other	3	7			1			9		2	21	41



**Fig. 7.** Abandoned field and trees of oil-palm, *Elaeis guineensis*. It is one of the most important plant species for humans and chimpanzees.

Table 3. Number of plant items used by humans for medicine.

Purpose	Plant part										Total
	Fr	Fe	Fl	Br	Tb	Gr	Ec	Tg	Gm	Lg	
Aphrodisiac	1	3			1	2	6	2			15
Anodyne		5	1		1	1	5	1			14
Anticough		8						4			12
Antiicterus		5					3	1			9
Febrifuge		8									8
Vermifuge		6					1		1		8
Antipox		4							4		8
Antidycentery		5						1	1		7
Trichomonacide		1						1	1		3
Antiatrepsia			1	2							3
Others	17	1				1	3	3	1		26
Total	1	62	3	2	2	5	19	18	1		113

Fr: fruit, Fe (feuille): leaf, Fl (fleur): flower, Br (bourgeon): bud or shoot, Tb: tubercle, Gr (graine): seed or kernel of nut, Ec (ecorce): bark, Tg (tige): twig or stem, Gm (gomme): gum or resin, Lg (ligneuse): woody tissue.

of *Elaeis guineensis* are also very important foods (or drink). However, they are rarely planted but left in the forest or field as they are (Fig. 7). The reason why some kinds of important trees are not artificially planted may be that they grow sufficiently well even in the forest or an abandoned field without special care.

The most characteristic feature of the human use of plants is for medicine (Table 3). People selectively use more items and species of plants as traditional medicine than as food. The largest variety of plant items used as medicine is as an aphrodisiac, though a scanty aphrodisia is not usually physical trouble. The second and third largest varieties are for minor but common troubles. From 4th to 8th are also minor and common troubles, or serious ones. Based on the ranking of varieties of plant species for each trouble, we can see how much public interest in aphrodisia exists among the people.

*Aspilia* spp. and *Vernonia* spp. are said to be used for medicine over a wide range of Africa by both humans and, perhaps, by chimpanzees in East Africa (Huffman & Seifu, 1989; Nishida, 1990). Altogether eight species of these two genera have been found in Bossou. However, none of them were recorded being us-

Table 4. Number of plant items used by humans.

Purpose	Plant part											Total
	Fr	Fe	Fl	Br	Tb	Gr	Ec	Tg	Gm	Lg	Wood, whole plant	
Medicine	1	62	3	2	2	5	19	18	1			113
House construct		1						1		1	3	6
Furniture, joinery		2						3			10	15
Dye	1	2			1			2				6
Ornament										6		6
Others	2	2						4		1	1	10
Total	4	69	3	2	3	5	19	28	1	2	20	156

Abbreviations as same as Table 3.

ed by people or chimpanzees as either medicine or food. Recently, a coughing chimpanzee mother with an extremely exhausted infant was observed to swallow some leaves of *Polycephalium capitatum* without chewing, as if for medicine (Matsuzawa, 1992). Leaves of this liana are also used by people as medicine, though as a hemostatic.

Additional human utilization of particular species of plants is mainly as material for house construction, furniture making and other tools (Table 4). Dye, soap and ornaments are also important for convenient and comfortable human life. Altogether, people use almost same number of items and species of plants, 239 and 198, as do chimpanzees, with 246 and 200 respectively. If we include nonselective uses of plants, e.g., timber for house construction, fences and fire-wood, far more plants are utilized by people than chimpanzees, but, of course, for many different kinds of purposes.

**ACKNOWLEDGMENTS** This field research was carried out with the collaboration of Kyoto University Primate Research Institute of Japan and Direction National de la Recherche Scientifique et Technique of Republique de Guinee. It was financed by grants under the Monbusho International Scientific Research Program. Manon names and other information on plants were primarily provided by Guano GUMI and Tino CAMARA. The preliminary draft was kindly commented and revised by Ellen INGMANSON. The long-term field research was shared with many colleagues. We would like to express our heartfelt thanks to these organizations and colleagues.

## REFERENCES

- Adam, J.-G. 1971a. *Flore Descriptive des Monts Nimba*. Editions du Museum, Paris.  
——— 1971b. *Flore Descriptive des Monts Nimba*, 2e partie. Editions du Museum, Paris.  
——— 1975a. *Flore Descriptive des Monts Nimba*, 3e partie. Editions du Museum, Paris.  
——— 1975b. *Flore Descriptive des Monts Nimba*, 4e partie. Editions du Museum, Paris.  
Huffman, M. & M. Seifu 1989. Observations on the illness and consumption of a possibly medicinal plant *Vernonia amygdalina* (Del.), by a wild chimpanzee in the Mahale Mountains National Park, Tanzania. *Primates*, 30(1):51–63.  
Hutchinson, J. & J. M. Dalziel 1954. *Flora of West Tropical Africa*, Vol.1, Part 1 (2nd edition, revised by R. W. J. Keay). Crown Agents for Overseas Governments and Administration (CAOGA), London.  
——— 1958. *Flora of West Tropical Africa*, Vol.1, Part 2 (2nd ed., rev. by R. W. J. Keay). CAOGA, London.  
——— 1963. *Flora of West Tropical Africa*, Vol.2 (2nd ed., rev. by F. N. Hepper). CAOGA, London.  
——— 1968. *Flora of West Tropical Africa*, Vol.3, Part 1 (2nd ed., rev. by F. N. Hepper). CAOGA, London.  
——— 1972. *Flora of West Tropical Africa*, Vol.3, Part 2 (2nd ed., rev. by F. N. Hepper). CAOGA, London.  
Matsuzawa, T. 1992. Death of a chimpanzee and care for it. *Hattatsu*, 49:95–104 (in Japanese).  
Nishida, T. 1990. A quarter century of research in the Mahale Mountains: An overview. In

- (T. Nishida, ed.) *The Chimpanzees of the Mahale Mountains*, pp.3-35, University of Tokyo Press, Tokyo.
- Sugiyama, Y. & J. Koman 1979. Social structure and dynamics of wild chimpanzees at Bossou, Guinea. *Primates*, 20(3): 323-339.
- 1987. A preliminary list of chimpanzees' alimentation at Bossou, Guinea. *Primates*, 28(1):133-147.

———Received June 8, 1992

Authors' Names and Addresses: Yukimaru SUGIYAMA, *Primate Research Institute, Kyoto University, Kanrin, Inuyama, Aichi 484, Japan*; Jeremy KOMAN, *Laboratoire Monts Nimba, Lola, Republique de Guinee*.

**Appendix. Flora of Bossou: Utilization by chimpanzees and humans.**

Brief explanations and abbreviations in Appendix are as follows:

(1) Vernacular name. Manon (m) is the local language of Bossou and Guerze (g) is that of an adjacent region. Only for some important plants are local names given in the list.

(2) Plant type. Large trees of more than approximately 20 m = gA (grande arbre), small tree of less than 10 m = pA (petite arbre), middle-sized tree = A, bush or scrub = Ar (arbuste), herb and grass = H, liana = L.

(3) Vegetation. Dense forest = Fd, secondary and deciduous forest = Fs, (transitional area is simply = F), scrub forest or savanna = S, cultivated field = C (champ), abandoned field = J (jachere recursus), swamp = M (marigot), ruderal = R.

(4) Part of the plant consumed and utilized (parties consommées). Fruit = Fr, leaf = Fe (feuille), flower = Fl (fleur), bud or shoot = Br (bourgeon), tubercle = Tb, seed or kernel of nut = Gr (graine), bark = Ec (ecorce), twig or stem = Tg (tige), gum or resin = Gm (gomme), woody tissue = Lg (ligneuse); young = j (jeune), dead = m (mort), base = b.

(5) Place of Bossou (see Fig. 2). Ban = 1, Guein = 2, Bouton = 3, Ueyaton = 4, Tonzuru = 5.

No.	Scientific name	Vernacular name	Plant type	Vegetation	Place	Consumed by		Medical use		Other use
						chimps	men	function	part	
1	<i>Abrus canescens</i> Welw ex Bak (Papilionaceae)		L	Fs, S	1 2 3					
2	<i>Abrus stictosperma</i> J.Berhant		L	Fs	1 2 3					
3	<i>Acacia ataxacantha</i> DC (Mimosaceae)	gban=g	pL	Fs	1 2 3					
4	<i>Acacia pennata</i> (L) Willd		pL	Fs	1 2 3	Gm				
5	<i>Acanthospermum hispidum</i> DC (Compositae)		H	R				antiicterus	Fe	
6	<i>Acacia tenuiflora</i> Dinkl & Engl (Rosaceae)		pA	F	1	Gr				
7	<i>Adathoda guineensis</i> H.Heine (Acanthaceae)		pA	F	1 2					
8	<i>Adathoda robusta</i> CB		pA	F						
9	<i>Adenia lobata</i> (Jacq) Engl (Passifloraceae)		L	Fs	1 2					
10	<i>Adenopus breviflorus</i> Benth (Cucurbitaceae)		L	Fs	1 2 3	Fr				
11	<i>Adenostemma perrottetii</i> DC (Compositae)		H	M						

12 <i>Adiantum philipense</i> L (Polypodiaceae)	H	Fs, S	1 2		
13 <i>Aedesia glabra</i> (Klatt) O.Hoffm(Compositae)	H	S			
14 <i>Neolanthus pubescens</i> Benth (Labiatae)	H	S			
15 <i>Aeschynomene pulchella</i> Planch ex Bak (Papilionaceae)	H	S			
16 <i>Aframomum exscapum</i> (Sims) daunti=m Hepper (Zingiberaceae)	H	Fs	1 2 3	Fr, Tg	
17 <i>Aframomum latifolium</i> Afz	H	S		Fr, Tg	
18 <i>Aframomum longiscapum</i> (Hook.f) K.Schum	H	Fs	1 2 3	Fr, Tg	
19 <i>Aframomum melegueta</i> K.Schum	H	C		Fr, Tg	Fr
20 <i>Aframomum sceptrum</i> (Oliv & Hanb)	H	Fs	1 2 3	Fr, Tg	
21 <i>Aframomum strobilaceum</i> (Sm) Hepper	H	Fs	1 2 3	Fr, Tg	
22 <i>Aframomum subsericeum</i> (Oliv & Hanb) K.Schum	H	Fs	1 2 3	Fr, Tg	
23 <i>Aframomum sulcatum</i> (Oliv & Hanb) K.Schum	H	Fs	1 2 3	Fr, Tg	
24 <i>Agelaea obliqua</i> (P.Beauv) Baill (Connaraceae)	L	F	1 2		
25 <i>Agelaea trifolia</i> (Lam) Gilg	L	F	1 2		
26 <i>Ageratum conyzoides</i> L (Compositae)	melapapa=g	H	J		vermifuge Fe antiseptic(vulnery)
27 <i>Albizia adianthifolia</i> (Sch) WF Wright (Mimosaceae)	pan=m gbaan=g	A	Fs	1 2 3	Fr
28 <i>Albizia ferruginea</i> (Guill & Perr) Benth	A	Fs	1 2 3		
29 <i>Albizia lebbeck</i> (L) Benth	A	Fs	1 2 3		
30 <i>Albizia zygia</i> (DC) JF Macbr	pmati=m gbaan, tei=g	A	Fs	1 2 3	Gm Fl
31 <i>Alchornea cordifolia</i> (Schum & Thonn Mull(Euphorbiaceae)	furome=m pelena=g	ps	Fs	1 2 3	Fr Tg(pith) antiseptic(vulnery) Fe anticough Tg(pith)

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
32	<i>Alchornea hirtella</i> Benth		pA	F	1 2				
33	<i>Allophylus africanus</i> P.Beaup (Sapindaceae)		pA	Fs	1 2 3 4			anti-rheumatism	Fe
34	<i>Alstonia congensis</i> Engl (Apocynaceae)		A	Fs	1 2 3				wood(music instrument)
35	<i>Alternanthera repens</i> (L) Link (Amaranthaceae)		H	R					
36	<i>Alternanthera sessilis</i> (L)R.Br.ex Roth		H	R					
37	<i>Amaranthus spinosus</i> L (Amaranthaceae)	gboo=g	H	C				Fe	
38	<i>Amaranthus viridis</i> L		H	C				Fe	
39	<i>Ammania senegalensis</i> Lam (Lythraceae)		H	S					
40	<i>Ampelocissus pentaphylla</i> (Ampelidaceae)		L	S, Fs	1 2 3				
41	<i>Anacardium occidentale</i> L (Anacardiaceae)		pA	C			Fr	Fr	
42	<i>Anadelphia</i> sp (Gramineae)		H	S					house roof
43	<i>Ananasa comosus</i> (L) Merrill (Bromeliaceae)	yerababi =m	Ar	C	1 2 3 4 5	Fr, Tg	Fr		
44	<i>Anchomanes difformis</i> Engl (Araceae)		H	Fs	1 2 3				
45	<i>Ancilobotrys scandens</i> (Schum & Thonn) Pichon (Apocynaceae)		L	Fs	1 2	Fr			
46	<i>Ancistrophyllum secundiflorum</i> lao=g Wendl (Palmace)		L	F	1		trichomonacide	Tg	
47	<i>Andropogon shirensis</i> Hochst (Gramineae)		H	S					
48	<i>Aneilema paludosum</i> A.Chev (Commelinaceae)		H	M					
49	<i>Anlingeria altissima</i> (A.Chev) Aubr & Pellegr (Sapotaceae)	iera=m	gA	Fsd	1 2	Fr			

50 <i>Annona muricata</i> L (Annonaceae)	susu=g	pA	C		Fr	Fr	
51 <i>Anthocleista nobilis</i> G.Don (Loganiaceae)		pA	Fs	1 2 3 4		febrifuge	Fe
52 <i>Anthocleista vogelii</i> Planch		A	M				
53 <i>Anthonotha fragrans</i> (Bak f) Exell & Hillcoat (Caesalpiniaceae)		pA	Fs	1 2 3 4			
54 <i>Anthonotha macrophylla</i> P.Beauv		A	Fs		jFe		
55 <i>Antiaris africana</i> Engl (Moraceae)	po=m	gA	Fsd	1 2 3	Fr		wood (furniture)
56 <i>Arachis hypogaea</i> L (Papilionaceae)	quo,tiga=m tian=g	H	C		Gr	Gr (oil)	
57 <i>Argemone mexicana</i> L (Papaveraceae)		H	village			antiicterus	Fe
58 <i>Artobotrys jollianus</i> (Annonaceae)		L	Fs	1			
59 <i>Asparagus africanus</i> Lam (Liliaceae)		H	Fs	1 2 3			
60 <i>Aspilia africana</i> (Pers) Adams (Compositae)		H	J				
61 <i>Aspilia helianthoides</i> (S.& Th) Oliv		H	J				
62 <i>Aspilia paludosa</i> Berh		H	J				
63 <i>Asplenium</i> sp (Polypodiaceae)		H	F	1 2			
64 <i>Asystacia calycina</i> Benth (Acanthaceae)		H	Fs	1 2			
65 <i>Asystacia scandens</i> Lindl		H	F	1 2			
66 <i>Asystacia vogeliana</i> Benth		Ar	F	1 2			
67 <i>Aubrevillea platycarpa</i> Pellegr (Mimosaceae)		A	F	1 2			
68 <i>Azadirachta indica</i> A.Juss (Meliaceae)		pA	C			antiicterus	Fe
69 <i>Azolla</i> sp (Lemnaceae)		H	S				

(cont.)

(Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
70	<i>Baissea breviloba</i> Stapf (Apocynaceae)		pA	Fs		1	2	3	
71	<i>Baissea lamepoolei</i> Stapf		L	Fs		1	2	3	
72	<i>Baissea</i> sp		L	Fs		1	2	3	
73	<i>Baissea zygodiooides</i> (K.Schum) Stapf		L	J					
74	<i>Balsamocitrus chevalieri</i> (Swingle) A.Chev (Rutaceae)		pA	Fs		1	2		
75	<i>Baphia nitida</i> Lodd (Papionaceae)	dolo=g	pA	F, Fs	1	2		regulation of Tg profuse menstruation	Tg(dye)
76	<i>Beilschmiedia Mannimeiss</i> Benth & Hook (Lauraceae)	gbaan=g	pA	Fs	1	2	Fr	Fr	
77	<i>Begonia emini</i> Warb (Begoniaceae)		H			1	2		
78	<i>Bequaertiadendron megalistemonatum</i> (Soud) Heine & JH (Sapotaceae)		A	F, M		1	2		
79	<i>Bequaertiadendron ob lanceolatum</i> (S.Moore) Heine & Hemsley (Sapotaceae)		A	F, M		1	2		
80	<i>Bertiera racemosa</i> (G.Don) K.Schum (Rubiaceae)		pA	M		1	2		
81	<i>Bertiera spicata</i> (Gaertn) Wernh		pA	F, M		1	2		
82	<i>Bidens bipinnata</i> L (Compositae)		H	J					
83	<i>Biophytum</i> sp (Oxalidaceae)		H	M					
84	<i>Blighia sapida</i> Konig (Sapindaceae)	kabaan=g	gA	F	1	2	Fr		
85	<i>Blighia unijugata</i> Bak	Kabaan=g	A	F	1	2	Fr		
86	<i>Blighia welwitschii</i> Hiern Radlk	gren=m	gA	F	1	2	Fr		
87	<i>Boerhaavia diffusa</i> L (Nyctagynaceae)		H	R					
88	<i>Boerhaavia erecta</i> L		H	R					

89 <i>Boerhaavia graminicola</i> Berh	H	R				
90 <i>Boerhaavia repens</i> L	H	R				
91 <i>Bolbitis heudelotii</i> (Bory) Alston (Polypodiaceae)	H	M				
92 <i>Bombax buonopozense</i> P.Beauv (Adansonia)	gA	Fs	1 2	Fr		
93 <i>Bonamia thunbergiana</i> (Roem & Schult) FN Williams (Convolvulaceae)	L	J				
94 <i>Borreria ocyoides</i> (Burm) DC (Rubiaceae)	H	J				
95 <i>Borreria scabra</i> (Schum & thonn) K.Schum	H	S				
96 <i>Borreria verticillata</i> (L) GFW Mey	H	R			antilichen	Fe
97 <i>Boscia angustifolia</i> A.Rich (Capparidaceae)	pH	R	1 2			
98 <i>Bosquiea angolensis</i> Ficahlo (Moraceae)	A	Fs	1 2 3	Ec		
99 <i>Brachypodium</i> sp (Gramineae)	H	Fs	1 2			
100 <i>Brachystegia leonensis</i> (Burrt) Davy et Hutch(Caesalpniaceae)	A	Fs	1 2			
101 <i>Brachysthephanus nimbae</i> H.Heine (Acanthaceae)	Ar	F	1 2 5			
102 <i>Bridelia ferruginea</i> Benth (Euphorbiaceae)	pA	S		Fr	Ec(alcohol fermentation)	
103 <i>Bridelia micrantha</i> (Hochst) Baill	pA	Fs	1 2			
104 <i>Brillantaisia lamium</i> (Nees) Benth (Acanthaceae)	H	J				
105 <i>Brillantaisia nitens</i> Lindan	H	Fs, J				
106 <i>Bryophyllum pinnatum</i> (Lam) Oken (Crassulaceae)	gbou-ouou =g	C			vermifuge anodyne	Fe
107 <i>Bulbophyllum barbigerum</i> (Orchidaceae)	lind=g	F	1			
108 <i>Bulbophyllum bifarium</i> Hook.f	H	F	1			

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use		Other use
						chimps	men	function	part	
109	<i>Bulbophyllum cochleatum</i> Lindl		H	F		1	2			
110	<i>Bulbophyllum cocoimum</i> Batem ex Lindl		H	F		1	2			
111	<i>Bulbophyllum falcatum</i> (Lindl) Rolfe		H	F		1				
112	<i>Bulbophyllum inflatum</i> Rolfe		H	F		1	2			
113	<i>Bulbophyllum intertextum</i> Lindl		H	F		1	2			
114	<i>Bulbophyllum linderi</i> Summerh		H	F		1	2			
115	<i>Bulbophyllum maximum</i> (Lindl) Rehd & Wils		H	F		1	2			
116	<i>Bulbophyllum phaeopogon</i> Schltr		H	F		1	2			
117	<i>Bulbophyllum recurvum</i> Lindl		H	F		1	2			
118	<i>Bulbophyllum scariosum</i> Summerh		H	F		1	2			
119	<i>Bulbophyllum velutinum</i> (Lindl) Rechb. f		H	F		1	2			
120	<i>Bussea occidentalis</i> Hutch (Caesalpiniaceae)	kpayeli	A	Fd		1	2	Gr	aphrodisiac	Ec, Gr Tg(pestle / mortar)
121	<i>Byrsocarpus coccineus</i> Schum & Thonn (Connaraceae)		pA	Fs		1	2			
122	<i>Cajanus cajan</i> (L) Millsp (Papilionaceae)		H	C				Gr	Gr	
123	<i>Caladium bicolor</i> (Ait) Vent (Araceae)		H	C						(ornament)
124	<i>Callichilia subsessilis</i> (Benth) Stapf (Apocynaceae)	gbakpawole	pA	Fs, S		1	2			
125	<i>Calopogonium mucunoides</i> Desv (Papilionaceae)		L	J						
126	<i>Calpocalyx aubrevillei</i> Pellegr (Mimosaceae)		pA	Fs		1	2	Fr		
127	<i>Canarium schweinfurthii</i> Engl (Burseraceae)		A	Fs		1		Fr		

128 <i>Canna bidentata</i> Bertol (Cannaceae)	mamanan=g	H	J				Fe (packing)
129 <i>Canthium horizontale</i> (Schum & Thonn) Hiern (Rubiaceae)		pA	Fsd	1 2			
130 <i>Canthium multiflorum</i> (Schum & Thonn) Hiern		pA	Fs	1 2			
131 <i>Canthium subcordatum</i> DC	wolowolokoi =g	pA	Fs	1 2 3			
132 <i>Canthium vulgare</i> K.Schum		pA	Fs	1 2			
133 <i>Carapa procera</i> DC (Meliaceae)	gbon=m	pA	Fs	1 2 3 4 5	jF1 Gr	anodyne	Gr (oil)
134 <i>Cardiospermum grandiflorum</i> Swartz (Sapindaceae)		L	Fs	1 2 3			
135 <i>Cardiospermum halicacabum</i> L		L	F	1 2 3 4			
136 <i>Carica papaya</i> L (Caricaceae)	iritike=m yeletiga=g	pA	C		Fr	vermifuge	Gr
137 <i>Carpodinus</i> sp (Apocynaceae)		pA	Fs		Fr		
138 <i>Cassia alata</i> L (Caesalpiniaceae)		pA	C, J			anti- dermatonosis	Fe
139 <i>Cassia mimosoides</i> L		H	S				
140 <i>Cassia siamea</i> Lam		pA	C			febrifuge	Fe
141 <i>Cassia sieberiana</i> DC		pA	Fs	1 2 3 4		aphrodisiac	Fe
142 <i>Cassia tora</i> L		pL	J			anodyne	Ec
143 <i>Cassytha filiformis</i> L (Lauraceae)		L	Fs	1 3		antidysentery	Tg
144 <i>Catharanthus roseus</i> (L) G.Don (Apocynaceae)		H	C				(ornament)
145 <i>Cathormion altissima</i> (Hook.f.) (Mimosaceae)		A	M	2			
146 <i>Ceiba pentandra</i> Gaerth (Bombacaceae)	gee=m	gA	Fsd	1 2	jFr F1		
147 <i>Celtis adulfi-friderici</i> Engl (Ulmaceae)	consinguan =m	A	Fsd	1 2	Fr		
148 <i>Celtis brownii</i> Rendle		A	Fs	1 2	Fr		
149 <i>Celtis mildbraedii</i> Engl		A	Fsd	1 2	Fr		

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Medical use part	Other use
						chimps	men			
150	<i>Centella asiatica</i> (L) Urb (Umbelliferae)		H	Fs, M	2			Fl	antiicterus anodyne antipaludian	Fe
151	<i>Cephaelis biaurita</i> (H & D) Hepper (Rubiaceae)		Ar	F	1 2					
152	<i>Cephaelis mangenotii</i> Ake-Assi		H	F	1 2					
153	<i>Cephaelis ombrophila</i> Schnell		H	F	1 2					
154	<i>Cephaelis peduncularis</i> salish		pA	F	1 2					
155	<i>Ceratopteris calomelanos</i> (L) Und (Polypodiaceae)		H	Fs, M	1 2					
156	<i>Cercestis afzelii</i> Schott (Araceae)		L	Fs, F	1 2					
157	<i>Cercestis stigmaticus</i> NE.Br		L	Fs, F	1 2					
158	<i>Chasmopodium caudatum</i> Stapf (Gramineae)		H	S						
159	<i>Chidlowia sanguinea</i> Hoyle (Caesalpiniaceae)		A	F	1 2					
160	<i>Chlorophora excelsa</i> Welw Benth (Moraceae)	guei=m gelieda=g	gA	Fsd	1 2	Fr, Fl		antiicterus	Ec	wood (furniture)
161	<i>Chlorophora regia</i> A.Chev	hea=g	gA	Fsd	1 2	Fr, Fl		antiicterus	Ec	wood (furniture)
162	<i>Chrysophyllum gigantium</i> A.Chev (Sapotaceae)	diera=m yalaulu=g	gA	F	1 2	Fr		antimyalgia	Ec	wood (furniture)
163	<i>Chrysophyllum perpulchrum</i> Mildbr ex H & D	winbenre =m	gA	F	1 2	Fr, jFe		antimyalgia	Ec	wood (furniture)
164	<i>Chrysophyllum welwitschii</i> Engl		L	Fs	1 2 3	Fe	Fe			
165	<i>Citrus aurantifolia</i> Swin (Rutaceae)	gein=m	pA	C	1 2 3 4	Fr	Fr			
166	<i>Citrus grandis</i> L		pA	C	1 2 3 4	Fr	Fr			
167	<i>Citrus limon</i> L	ganikpona =g	pA	C	1 2 3 4	Fr	Fr			
168	<i>Clematis grandiflora</i> DC (Ranunculaceae)		L	Fs	1 2 3					

169 <i>Clematis hirsuta</i> G.& P.		L	Fs	1 2 3			
170 <i>Clerodendrum buchholzii</i> Gurke (Verbenaceae)		L	J				
171 <i>Clerodendrum formicarum</i> Gurke		L	J				
172 <i>Clerodendrum splendens</i> G.Don		pgA, L	J				
173 <i>Clearodendrum thyrsoides</i> Gurke		L	J, Fs				
174 <i>Clerodendrum umbellatum</i> Poir		pgA, L	J				
175 <i>Clerodendrum volubile</i> P.Beauv		pgA, L	J				
176 <i>Clerodendrum poloycephalum</i> Bak		pgA, L	J				
177 <i>Cnestis corniculata</i> Lam (Connaraceae)		L	F	1 2			
178 <i>Cnestis ferruginea</i> DC		pA	Fs	1 2 3			
179 <i>Cnestis racemosa</i> G.Don		pA	Fs	1 2 3			
180 <i>Cocos nucifera</i> L (Palmae)		A	C		Fr	Fr	
181 <i>Coffea arabica</i> L (Rubiaceae)	Cafe	pA	C			Gr	
182 <i>Coix lacrima-jobi</i> L (Gramineae)		H	C				Fe (feed rabbit)
183 <i>Cola caricaefolia</i> G.Don (Sterculiaceae)	bumoguan =m	pA	Fs		Fr	aphrodisiac	Fe
184 <i>Cola cordifolia</i> (Cav)R.Br	buba=m	A	Fsd	1 2	Fr		
185 <i>Cola laurifolia</i> Mast		pA	Fsd	1 2			
186 <i>Cola nitida</i> (Vent) Sch & Endl	toole=g	A	C		jFe Fr	jFe Fr	antipapitation (cardiotonic) Fe
187 <i>Cola reticulata</i> A.Chev		pA	F	1 2	Fr		
188 <i>Colocassia esculenta</i> L Schott (Araceae)		H	C		Tb		
189 <i>Combretodendron macrocarpum</i> P.Beauv (Lecythidaceae)	keaye=g	A	F	1			
190 <i>Combretum hispidum</i> Laws (Combretaceae)		L	S				
191 <i>Combretum paniculatum</i> Vent		L	S				

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use		Other use
						chimps	men	function	part	
192	<i>Combretum platypterum</i> Welw		L	Fs		1	2	3		
193	<i>Combretum smethmannii</i> G.Don		L	Fs		1	3			
194	<i>Commelina africana</i> L (Commelinaceae)		H	S						
195	<i>Commelina erecta</i> L		H	S						
196	<i>Corynanthe pachyceras</i> K.Schum (Rubiaceae)		pA	Fs		1	2			
197	<i>Costus afer</i> Ker-Gawl (Zingiberaceae)	toha=g	H	Fs	1 4	2	3	Tg, jFe	F1	anticough antipox
198	<i>Costus deistelii</i> K.Schum	zen=m	H	F, Fs	1	2	3	Tg, jFe		anticough antipox
199	<i>Costus dubius</i> (Afz) K.Schum		H	J, Fs	1	2	3	Tg, jFe		anticough antipox
200	<i>Costus lucanusianus</i> J.Braun & K.Schum		H	Fs	1	2	3	Tg, jFe		anticough antipox
201	<i>Crassocephalum crepidioides</i> (Benth) Moore (Compositae)		H	J						
202	<i>Crassocephalum rubens</i> Juss. ex Jacq		H	J						
203	<i>Craterispermum laurinum</i> (Poir) Benth (Rubiaceae)	gbeke=g	pA	Fs	1 4	2	3	Ec	antiseptic(vulnery) antiicterous	Ec
208	<i>Cryptosepalum tetraphyllum</i> Hook.f Benth (Caesalpiniaceae)		pA	Fsd		1	2			
209	<i>Ctenitis protensa</i> (Afz) ex SW. Ching (Polypodiaceae)		H	M						
210	<i>Ctenium newtonii</i> Hack (Gramineae)		H	S						
211	<i>Cucumis melo</i> L (Cucurbitaceae)		L	C			Fr	Fr		
212	<i>Cucurbita pepo</i> L (Cucurbitaceae)		L	C			Gr	Gr		
213	<i>Culcasia angolensis</i> Welw ex Schott (Araceae)		L	Fs		1	2			

214 <i>Culcasia liberica</i> NE.Br		L	Fs	1 2		
215 <i>Culcasia scandens</i> P.Beauv		L	Fs	1 2		
216 <i>Culcasia sereti</i> DeWild		L	Fs	1 2		
217 <i>Culcasia striolata</i> Engl		L	Fs	1 2		
218 <i>Curcuma xanthoriza</i> Robs (Zingiberaceae)		H	C			Tb(dye)
219 <i>Cussonia barteri</i> Suman (Araliaceae)	melegbou- lou-g	pA	S			Tg(soap)
220 <i>Cyanotis longifolia</i> Benth (Commelinaceae)		H	S			
221 <i>Cyathea manniana</i> HK (Cyatheaceae)		pA	M		vermifuge	jFe
222 <i>Cyclosurus afer</i> {C.Chr} Ching (Polypodiaceae)		H	M			
223 <i>Cyclosurus goggilodus</i> (Schk) Link		H	M			
224 <i>Cyclosurus striatus</i> (Schum) Ching		H	M	2		
225 <i>Cynometra ananta</i> Hutch. & Dalz (Caesalpiniaceae)		A	F	1 2		
226 <i>Cyperus</i> sp L (Cyperaceae)		H	M			
227 <i>Cyrtosperma senegalense</i> (Schott) Engl (Araceae)		H	M			
228 <i>Dacryodes</i> sp (Burseraceae)		pA	Fs	1 2	Fr	
229 <i>Dalbergia afzeliana</i> G.Don Gen (Papilionaceae)		L	J			
230 <i>Dalbergia bignonae</i> J.Berhiant		L	Fs	1 2 3		
231 <i>Dalbergia saxatilis</i> Hook.f		L	Fs	1 2 3		
232 <i>Daniellia thurifera</i> Benn (Caesalpiniaceae)		A	Fs,J	1 2		
233 <i>Datura</i> sp (Solanaceae)		H	C		antifebrile	Fe
234 <i>Delonix regia</i> (Boj.ex Hook) Raf (Caesalpiniaceae)		pA	C			(ornament)

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
235	<i>Desmodium adscendens</i> (Sw) DC (Papilionaceae)		H	J				anti-dysentery	Fe
236	<i>Desmodium salicifolium</i> (Poir) DC		H	J					
237	<i>Detarium microcarpum</i> G & Perr (Caesalpiniaceae)		A	F	2	Fr	Fr		
238	<i>Dialium dinklagei</i> Harms (Caesalpiniaceae)		A	F	2				
239	<i>Dialium guineense</i> Willd peim		A	F	2	Fr	Fr		
240	<i>Dichrostachys glomerata</i> (Forsk) Chiov (Mimosaceae)	lana=g	pA	Fs	1 2 3 4	jFe		antivaginitis	pith Lg (rope)
241	<i>Dicranolepis laciniata</i> Gilg (Thymelaeaceae)		pA	F	1 2	Fr			
242	<i>Dicranopteris linearis</i> (Burns) Underwood (Gleicheniaceae)		H	M					
243	<i>Dinophora spenneroides</i> Benth (Melastomataceae)		pA	Fs	1 2 3				
244	<i>Diiodia scandens</i> Sw (Rubiaceae)		H	J, S					
245	<i>Dioscorea alata</i> L (Dioscoreaceae)		L	C					
246	<i>Dioscorea bulbifera</i> L		H	C		Tb			
247	<i>Dioscorea hirtiflora</i> Benth		L	Fs	1 2 3				
248	<i>Dioscoreophyllum</i> sp (Menispermaceae)		L	Fs	1 2 3				
249	<i>Diospyros sanza-minika</i> A.Chev (Ebenaceae)		pA	F	1 2				
250	<i>Diospyros thomasii</i> H. & D.		pA	Fs	1 2				
251	<i>Diospyros viridicans</i> Hiern		A	Fs	1 2				
252	<i>Dissotis grandiflora</i> (Sm) Benth (Melastomataceae)		H	S					
253	<i>Dissotis rotundifolia</i> (Sm) Triana		H	J, Fs	1 2 3 4	jFe	jFe anticough	Fe	

254 <i>Dissotis sylvestris</i> Jac-Fel		H	M			
255 <i>Distemonanthus benthamianus</i> Baill (Caesalpiniaceae)		A	Fs	1 2 3	Gr	
256 <i>Dolichos nimbaensis</i> Schnell (Papilionaceae)		H	S			
257 <i>Dracaena adamii</i> Hepper (Agavaceae)		H	F	1 2		
258 <i>Dracaena arborea</i> Willd	sirednen=m imanu=g	pA	Fs	1 2	bjFe	
259 <i>Dracaena draco</i> L		pA	Fsd	1 2	Gr	
260 <i>Dracaena humilis</i> Bak		H	Fs	1 2		
261 <i>Dracaena scoparia</i> A.Chev ex Hutch		pA	Fsd	1 2		
262 <i>Drepanocarpus lunatus</i> (L) GFW Mey (Papilionaceae)		L	Fs	1 2 3		
263 <i>Droogmansia scaettaiana</i> A.Chev & Sillans (Papilionaceae)		H	S			
264 <i>Drypetes chevalieri</i> Beille (Euphorbiaceae)		pA	F	1 2 3		
265 <i>Drypetes gulgiana</i> Pax & K.Hoffm		pA	F	1 2 3		
266 <i>Drypetes inaequalis</i> Hutch		pA	F	1 2		
267 <i>Elaeis guineensis</i> Jacq	tou=mg	A	Fs, C	1 2 3	bjFe, Gr Fr, Lg	Gr, Fr, Gm (oil) (alcohol)
268 <i>Elatostema paivaeanum</i> Wedd (Urticaceae)		H	M			
269 <i>Elephantopus scaber</i> A.Chev (Compositae)		H	R			
270 <i>Eleusine indica</i> (L) Gaertn (Gramineae)		H	R			
271 <i>Emilia coccinea</i> (Sims) G.Don (Compositae)		H	R			
272 <i>Emilia sonchifolia</i> (L) DC		H	J			
273 <i>Entada manni</i> (Oliv) Tisserant (Mimosaceae)		L	Fs	1 2 3		Lg(rope for roof)

(cont.)

(Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
274	<i>Entandrophragma angolense</i> (Welw) C.DC (Meliaceae)		A	Fs		1	2		
275	<i>Entandrophragma candollei</i> Harms		A	Fsd		1	2		
276	<i>Entandrophragma cylindricum</i> Sprague		A	Fsd		1	2		
277	<i>Entandrophragma utile</i> Dawe & Sprague		A	Fsd		1	2		
278	<i>Eremospatha hookeri</i> Mann & Wendl (Palmae)		L	Fsd		1	2		
279	<i>Eriocaulon pulchellum</i> Koern (Eriocaulaceae)		H	M					
280	<i>Eriosema glomeratum</i> G & Perr (Papilionaceae)		H	S					
281	<i>Eriosema parviflorum</i> E.Mey		H	S					
282	<i>Erythrina mildbraedii</i> (Harms) (Papilionaceae)		A	Fs		1	2	3	jFe
283	<i>Erythrina senegalensis</i> DC		pA	S					
284	<i>Ethulia conyzoides</i> L.f. (Compositae)		H	J				Fe	
285	<i>Euclinia longiflora</i> Salisb rugen-mm		pA	F		1	2	3	Fr
286	<i>Eupatorium</i> sp (Compositae)		H	S					
287	<i>Euphorbia hirta</i> L (Euphorbiaceae)		H	R				antidysentery aphrodisiac	Fe
288	<i>Euphorbia prostrata</i> Ait		H	R					
289	<i>Eurychone rothschildiana</i> (O.Brien) Schltr (Orchidaceae)		H	S					
290	<i>Fagara leprieurii</i> (G & Perr) Engl (Rutaceae)		pA	Fs		1	2		antidentalgia Ec
291	<i>Fagara macrophylla</i> Engl		A	Fs		1	2		antidentalgia Ec
292	<i>Fagara parvifoliola</i> A.Chev ex Keay		pA	Fs		1	2		antidentalgia Ec

293 <i>Ficus asperifolia</i> Miq (Moraceae)		pA	Fs	1 2	Fr
294 <i>Ficus capensis</i> Thunb		A	F	1 2 3	Fr
295 <i>Ficus congensis</i> Engl		A	M		
296 <i>Ficus exasperata</i> Vahl	nyanare=m	A	Fs	1 2 3	Fr, jFe
297 <i>Ficus glumosa</i> Del		pA	S		
298 <i>Ficus gnaphalocarpa</i> Miq	bre=m	A	Fs	1 2 3	Fr
299 <i>Ficus goliath</i> A.Chev		gA	Fs	2	
300 <i>Ficus leprieurii</i> Miq	pililapela =g	pA	F	1 2	Fr
301 <i>Ficus lyrata</i> Warb		pA	F	1 2	
302 <i>Ficus macroisperma</i> Warb ex Mildbr & Burret		gA	F	1 2	Fr
303 <i>Ficus mucoso</i> Welw ex Ficalho	serebro=m gnanale=g	gA	Fsd	1 2	jFe, Fr
304 <i>Ficus ovata</i> Vahl		pA	Fs	1 2	Fr
305 <i>Ficus polita</i> Vahl		pA	Fs	1 2	Fr
306 <i>Ficus sagittifolia</i> Warb ex Mildbr & Burret		pA	Fs	1 2	Fr
307 <i>Ficus thonningii</i> Blume	guru=m	A	F	1 2	Fr
308 <i>Ficus umbellata</i> Vahl	guruboa=m	A	Fs	1 2	jFe, Fr
309 <i>Floscopa africana</i> P.Beauv (Commelinaceae)		H	M		
310 <i>Fuirena umbellata</i> Rottb (Cyperaceae)	yalele=g	H	M		
311 <i>Funtumia africana</i> (Benth) Stapf (Apocynaceae)	tokurati =m	A	Fs	1 2 3	
312 <i>Funtumia elastica</i> (Preuss) Stapf		pA	F	1 2	jTg
313 <i>Gaertnera paniculata</i> Benth (Rubiaceae)		pA	Fs	1 2 3	
314 <i>Garcinia afzelii</i> Engl (Guttiferae)		pA	Fs	1 2 3	aphrodisiac Ec
315 <i>Garcinia kola</i> Heckel		pA	Fd	1 2	Gr Gr aphrodisiac Gr, Ec

(cont.)

(Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
316	<i>Garcinia polyantha</i> Oliv	zuruzei=m	pA	F	1 2				Fr(dye)
317	<i>Gardenia imperialis</i> K.Schum (Rubiaceae)		pA	F	1 2				
318	<i>Geophila afzeli</i> Hiern (Rubiaceae)		H	Fsd	1 2				
319	<i>Geophila obvallata</i> (Schumach) F.Didr		H	Fsd	1 2				
320	<i>Gilbertiodendron splendidum</i> A.Chev ex Hutch & Dalz(Caesalpiniaceae)		A	F	1 2				
321	<i>Gloriosa simplex</i> L (Liliaceae)		H	Fs	1 2 3				
322	<i>Gloriosa superba</i> L		H	Fs	1 2 3				
323	<i>Glyphaea brevis</i> Spreng (Tiliaceae)		pA	Fs	1 2 3				
324	<i>Gmelina arborea</i> Roxb (Verbenaceae)		pA	C					retimbering
325	<i>Gongronema latifolium</i> Benth (Asclepiadaceae)		L	Fs	1 2 3	Ec, Tg			
326	<i>Gossypium arboreum</i> L (Malvaceae)	yewulu=g	H	C			anti-blennorragia	Fe	
327	<i>Gouania longipetala</i> Hemsl (Rhamnaceae)		L	Fs	1 2 3		antivertigo	Fe	
328	<i>Grewia pubescens</i> P.Beauv (Tiliaceae)	uneparu=m	pA	Fs	1 2	Fr			
329	<i>Guibourtia copallifera</i> Benn (Caesalpiniaceae)		A	F	2				
330	<i>Gynandropsis gynandra</i> (L) Briq (Capparidaceae)		H	C			anodyne	Fe	
331	<i>Gynura micheliana</i> JC (Compositae)		H	S					
332	<i>Haemanthus multiflorus</i> Martyn (Amaryllidaceae)		H	Fs	1 2 3 4				
333	<i>Halopegia azurea</i> K.Schum (Marantaceae)	ma=g	H	F	1 2	Tg			Tg(mat)

334 <i>Hannoa klaineana</i> Pierre & Engl (Simaroubaceae)	vitri=m	pA	Fs	3	Fr		
335 <i>Harissonia abyssinica</i> Oliv (Simaroubaceae)		pA	S, J	1 2 3		aphrodisiac	Ec
336 <i>Harungana madagascariensis</i> Lam ex Poir (Hypericaceae)	loru=m lolo=g	pA	Fs	1 4	jFe, Fr	antiburn anodyne	Fe Tg(house construction)
337 <i>Heliotropium indicum</i> L (Boraginaceae)	boube=m gbegne=g	H	C, village			antialbuminuria	Fe
338 <i>Hibiscus esculentus</i> L (Malvaceae)		H	C		Fl, Fr	Fr, Fe	
339 <i>Hibiscus rostellatus</i> G & Perr		H	Fs	1 2 3			
340 <i>Hibiscus sabdariffa</i> L		H	C		Fl	Fl	
341 <i>Hippeastrum equestre</i> Ait (Amaryllidaceae)		H	C				(ornament)
342 <i>Hippocratea paniculata</i> Vahl (Celastraceae)		L	Fs	1 2 3	Fe		
343 <i>Holarrhena floribunda</i> (G.Don) Dur & Schinz (Apocynaceae)		A	Fs	1 2 3			
344 <i>Hugonia macrophylla</i> Oliv (Linaceae)		L	F	1 2			
345 <i>Hugonia planchonii</i> Hook.f		pA	Fs	1 2			
346 <i>Hugonia platysepala</i> Welw ex Oliv		pA	J			trichomonacide	Fe
347 <i>Hyparrhenia archaelymandra</i> J.Fel (Gramineae)		H	S				
348 <i>Hyparrhenia diplandra</i> (Hack) Stapf		H	S				(house roof)
349 <i>Hyparrhenia subplumosa</i> Stapf		H	S				
350 <i>Hypothymnum braunianum</i> K.Schum (Marantaceae)	gomoe=m	H	Fs	1 2 3	jFe, Gr	antiatrepsia	br
351 <i>Hypselodelphys poggeana</i> (K.Schum) Milne-Redh (Marantaceae)		H	Fs	1 2	Tg		
352 <i>Hypselodelphys violacea</i> Ridl		H	Fs	1 2	Tg		
353 <i>Hyptis lanceolata</i> Poir (Labiatae)		H	J				

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
354	<i>Impatiens nzoana</i> A.Chev (Balsaminaceae)		H	M				antiatrepsie	Fl,Fe
355	<i>Imperata cylindrica</i> P.Beaup (Gramineae)		H	S					
356	<i>Indigofera dalabaca</i> A.Chev (Papilionaceae)		H	S					
357	<i>Ipomoea dichroa</i> Choisy (Convolvulaceae)		H	S					
358	<i>Ipomoea involucrata</i> P.Beaup		L	S,Fs	1 2 3				
359	<i>Ipomoea batatas</i> (L) Lam	Cuission=m	L	C		Tb	Tb		
360	<i>Ipomoea quamoclit</i> L		L	Fs,R					
361	<i>Irvingia gabonensis</i> (Aubry-Lec ex O'Rorke) Baill(Irvingiaceae)		A	Fs	1 2				
362	<i>Ixora nimbania</i> Schnell (Rubiaceae)		H	F,Fs	1				
363	<i>Jatropha curcas</i> L (Euphorbiaceae)		Ar	C				laxative	Gr
364	<i>Jatropha gossypifolia</i> L		Ar	C					(ornament)
365	<i>Justicia tenella</i> (Nees) T.Anders (Acanthaceae)		H	Fs	1 2 3		F1		
366	<i>Justicia extensa</i> T.Andres		Ar	J					
367	<i>Justicia flava</i> (Forsk) Vahl		H	J					
368	<i>Khaya ivorensis</i> A.Chev (Meliaceae)		gA	Fsd	1 2				wood (furniture)
369	<i>Kigelia africana</i> (Lam) Benth (Bignoniaceae)		A	Fs	1				
370	<i>Kotschyia ochreata</i> (Taub) & Duvign (Papilionaceae)		H	S					
371	<i>Lagenaria siceraria</i> (Mollina) Standl (Cucurbitaceae)	koopilin =m koo=g	L	C					Fr(ladle)
372	<i>Laggera heudelotii</i> CD.Admas (Compositae)		H	J					

373	<i>Landolphia dulcis</i> (R.Br ex Sabine) Pichon (Apocynaceae)	bubwakara =m	L	Fsd	1 2	Fr
374	<i>Landolphia owariensis</i> P.Beaup	sean sean =m	L	Fs	2	Fr
375	<i>Lasiodiscus fasciculiflorus</i> Engl (Rhamnaceae)		pA	F	1 2	Fr
376	<i>Lasiodiscus marmoratus</i> Ch Wright		pA	F	1 2	Fr
377	<i>Lecanioidiscus cupanioides</i> Planch ex Benth (Sapindaceae)		pA	F	1 2	Fr
378	<i>Leea guineensis</i> G.Don (Ampelidaceae)		H	Fs	1 2 3	Tg
379	<i>Leptoderris reticulata</i> Dunn (Papilionaceae)		L	Fs	1 2 3	
380	<i>Lindackeria dentata</i> (Oliv) Gilg (Flacourtiaceae)		pA	Fs	1 3	
381	<i>Lindernia</i> sp (Scrophulariaceae)		H	M		
382	<i>Lippia</i> sp (Verbenaceae)		L	Fs	1 2 3	Fl
383	<i>Lomariopsis guineensis</i> (Underw) Alston (Polypodiaceae)		H	F	1 2	
384	<i>Lonchocarpus cyanescens</i> (Schum & Thonn) Benth (Papilionaceae)		L	Fs	2 3	Fe(dye)
385	<i>Lophira alata</i> Banks ex Gaertn (Ochnaceae)		gA	F	1 2	wood (furniture)
386	<i>Loranthus cupulatus</i> DC (Loranthaceae)	L (parasite)		Fs	1 2	
387	<i>Loudetia arundinacea</i> (Hochst ex A.Rich) Steud (Gramineae)		H	S		
388	<i>Luffa aegyptiaca</i> Mill (Cucurbitaceae)		L	J		Fr (sponge)
389	<i>Lycopodium cernuum</i> L (Lycopodiaceae)		H	M		
390	<i>Lycopodium microphyllum</i> (Cav) R.Br		H	M		
391	<i>Lycopodium scandens</i> SW		H	M		detoxication Fe, Tg
392	<i>Lycopodium smithianum</i> Pr.Z		H	M		

(cont.)

(Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
393	<i>Macaranga barteri</i> Mull (Euphorbiaceae)		pA	Fs	1 2				
394	<i>Macaranga heterophylla</i> Mull Arg		pA	Fs	1 2		Fr		
395	<i>Macaranga hurifolia</i> Beille		pA	Fs	1 2				
396	<i>Macaranga schweinfurthii</i> Pax		pA	M					
397	<i>Maesobotrya barteri</i> (Baill) Hutch (Euphorbiaceae)	kein=m tagna=g	pA	Fsd	1 2	Fr	Fr		
398	<i>Mangenotia eburnea</i> Pichon (Periplocaceae)		L	J					
399	<i>Mangifera indica</i> L (Anacardiaceae)	mangoro =mg	A	F			Fr	Fr	
400	<i>Manihot esculenta</i> Grantz (Euphorbiaceae)	npei=m māna=g	H	C			Tb, Fl	Tb	
401	<i>Manihot glaziovii</i> Mull Arg		H	C			Tb	Fe	
402	<i>Manihot utilissima</i> Pohl		H	C			Tb	Tb	
403	<i>Manniophytion africanum</i> (Euphorbiaceae)		pA	Fs	1 2 3			anticough	Fe
404	<i>Marantochloa congensis</i> (K.Schum) Leon & Mull (Marantaceae)		H	Fs	1 2		Tg		
405	<i>Marantochloa cuspidata</i> (Rosc) Milne-Redh		H	Fs	1 2		Tg		
406	<i>Marantochloa filipes</i> (Benth) birare=m Hutch		H	Fs	1 2		Tg		
407	<i>Marantochloa flexuosa</i> Hutch		H	Fs	1 2		Tg		
408	<i>Marantochloa leucantha</i> (K.Schum) Milne-Redh		H	Fs	1 2		jTg		
409	<i>Marantochloa purpurea</i> (Ridl) Milne-Redl		H	Fs	1 2		jTg		
410	<i>Marattia fraxinea</i> J.Sm (Marattiaceae)		H	M	2				
411	<i>Mareya micrantha</i> (Benth) Mull Arg (Euphorbiaceae)		pA	Fs, J	1 2 3			scabicide purgative	Fe Fe

412 <i>Massularia acuminata</i> (G.Don) Bullock ex Hoyle (Rubiaceae)	pA	Fs	1 2 3		
413 <i>Megaphrynum macrostachyum</i> yoro=m (Benth) Milne-Redh (Marantaceae)	H	F	1 2	Tg	Tg (mat)
414 <i>Melinis minutiflora</i> P.Beauv (Gramineae)	H	S, J			
415 <i>Merremia pterygocaulos</i> (Steud ex Choisy) Hallier (Convolvulaceae)	L	J			
416 <i>Merremia umbellata</i> (L) Hallier (Convolvulaceae)	H	J			
417 <i>Mesanthesum radicans</i> (Benth) Koern (Eriocaulaceae)	H	M			
418 <i>Mezoneuron benthamianum</i> Baill (Caesalpiniaceae)	konomini, gnale=g	A	J		aphrodisiac Tg
419 <i>Microdesmis puberula</i> Hook (Euphorbiaceae)	heli=g	pA	F, Fs	1 2 3	Fe, Fr Fe
420 <i>Microglossa afzeli</i> O.Hoffm (Compositae)	L	Fs	1 2 3	jFe	
421 <i>Microglossa caudata</i> O.Hoffm	L	J			
422 <i>Microglossa pyrifolia</i> (Lam) O.Ktze	H	J			
423 <i>Mikania cordata</i> (Burn) B.L.Rob (Compositae)	H	J			
424 <i>Millettia lanepoolei</i> Dunn (Papilionaceae)	pA	Fs	1 3		
425 <i>Millettia zechiana</i> Harms	pA	Fs	1 2 3 4		
426 <i>Mimosa pigra</i> L (Mimosaceae)	H	J			
427 <i>Mimosa pudica</i> L	H	J			
428 <i>Mitragyna stipulosa</i> (DC) Okzé (Rubiaceae)	boon=g	A	M		wood (furniture)
429 <i>Momordica cabraei</i> (Cogn) C.Jeffrey (Curcurbitaceae)	L	Fs	1 2 3		
430 <i>Momordica charantia</i> L	L	Fs	1 2 3		
431 <i>Mondia whitei</i> (Hook.f) Skeels (Periplocaceae)	H	S			

(cont.)

(Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
432	<i>Monodora tenuifolia</i> Benth (Annonaceae)	niyapelin, ulu=g	pA	Fs	1 2				
433	<i>Monopetalanthus compactus</i> Hutch & Dalz (Caesalpiniaceae)		A	F	1 2				
434	<i>Morinda geminata</i> DC (Rubiaceae)	zologba=g	pA	Fs	1 2 3	Fr		febrifuge	Fe
435	<i>Morinda longiflora</i> G.Don	botere=m	L	Fs	1 2 3	Fr		febrifuge	Fe
436	<i>Morinda lucida</i> Benth		pA	Fs	1 2 3			febrifuge	Fe
437	<i>Morinda morindoides</i> (Bak) Milne-Redh		pA	Fs	1 2 3	Fr		febrifuge	Fe
438	<i>Morus mesozugia</i> Stapf (Moraceae)	gangue=m Kagbe=g	gA	Fs	1 2	Fl, jFe Fr			wood (construction)
439	<i>Mucuna flagillipes</i> T.Vogel ex Hook.f (Papilionaceae)		L	Fs	2 3				
440	<i>Mucuna pruriens</i> (L) DC		L	Fs	1 2 3 4				Fe, Tg(dye)
441	<i>Mucuna sloanei</i> Fawc & Rendle		L	Fs	1 3 4 5				
442	<i>Musa sinensis</i> Sweer (Musaceae)	gvi=m	H	C		Tg, Fr	Fr		
443	<i>Musa paradisiaca</i> L		H	C		Tg, Fr	Fr		
444	<i>Musa sapientum</i> L	buro=m	H	C					
445	<i>Musaenda elegans</i> Schum & Thonn (Rubiaceae)		L	Fs	1 2 3				
446	<i>Musaenda erithrophylla</i> Schum & Thonn	teegbele =g	L	Fs	1 2 3			menstruation normalise	Fl
447	<i>Musanga cecropioides</i> R.Br (Moraceae)	uguru=m wei=g	A	Fs	1 2 3 4	jFe			
448	<i>Myrianthus arboreus</i> P.Beauv (Moraceae)	baru=m gbalo=g	pA	Fs	1 2 3	Fr, jFe (sauce)	jFe		
449	<i>Myrianthus libericus</i> Rendle		pA	Fs	1 2 3	jFe, Fr			
450	<i>Myrianthus serratus</i> (Trecul) Benth		pA	Fsd	1 2 3	Fr			

451 <i>Napoleona leonensis</i> Hutch & Daltz (Lecythidaceae)		pA	Fsd	1 2	Fr			
452 <i>Napoleona vogelii</i> Hook & Planch	dei=m	pA	Fsd	1 2	Fr			
453 <i>Nauclea latifolia</i> Sm (Rubiaceae)	bahila=g	pA	Fs, S	1 2 3	Fr	Fr	febrifuge	Fe
454 <i>Nelsonia canescens</i> (Lam) Spreng (Acanthaceae)		H	S					
455 <i>Nephrolepis biserrata</i> (Sw) (Polypodiaceae)		H	M					
456 <i>Nephrolepis undulata</i> Afz. J.Sm (Polypodiaceae)		H	S					
457 <i>Nesaea radicans</i> Guill & Perr (Lythraceae)		H	R					
458 <i>Neuroleptis</i> sp (Convolvulaceae)		L	Fs	1 2 3				
459 <i>Neurotheca</i> sp (Gentianaceae)		H	S					
460 <i>Newbouldia laevis</i> (P.Beaup) Seem ex Bureau (Bignoniaceae)		pA	Fs	1 2 3	jFe	trichomonacide	Ec	
461 <i>Nicotiana rustica</i> L (Solanaceae)		H	C					Fe(tabacco)
462 <i>Nymphaea lotus</i> L (Nymphaeaceae)		H	M					
463 <i>Ochna membranacea</i> Oliv (Ochnaceae)		pA	Fs, S	1 2 3				
464 <i>Ocimum basilicum</i> L (Labiatae)		H	C			jFe(Sauce)		
465 <i>Ocimum gratissimum</i> L		H	J			jFe(Sauce)		
466 <i>Olyra latifolia</i> L (Gramineae)		H	Fsd	1 2 3				
467 <i>Ophioglossum reticulatum</i> L (Ophioglossaceae)		H	M					
468 <i>Opisismenus burmannii</i> (Retz) P.Beaup (Gramineae)		H	Fs	1 2 3				
469 <i>Oryza</i> sp (Gramineae)	bu=m	H	C		Tg	Gr		Tg(mattress)
470 <i>Osbeckia porteresi</i> Jac.Feli (Melastomataceae)		H	S					
471 <i>Osbeckia tubulosa</i> Sm		H	J					

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
472	<i>Oxyanthus formosus</i> Hook ex Planch (Rubiaceae)		pA	Fs		1	2	3	
473	<i>Oxyanthus pallidus</i> Hiern		pA	Fs		1	2	3	
474	<i>Oxyanthus unilocularis</i> Hiern		pA	Fs	1	2	Fr	aphrodisiac	Fr, Fe
475	<i>Oxytenanthera abyssinica</i> (A.Rich) Munro (Gramineae)		pA	C					Tg(fence)
476	<i>Pachyopodanthium staudtii</i> Engl & Diels (Annonaceae)		pA	Fs		2			
477	<i>Pachystela pobeguiniana</i> Pierre ex Lecomte (Sapotaceae)	beke=m	A	Fs	1	2	Fr		
478	<i>Palisota bracteosa</i> C.B.CI (Commelinaceae)		H	Fs	1	2	3		
479	<i>Palisota hirsuta</i> (Thunb) K.Schum		H	Fs	1	2	3	Fr	
480	<i>Pandiaka</i> sp (Amarantaceae)		H	S					
481	<i>Panicum humilenees</i> (Gramineae)		H	M					
482	<i>Panicum maximum</i> Jacq		H	S					
483	<i>Panicum paludosum</i> Roxb		H	M					
484	<i>Panicum turgidum</i> Foresk		H	S					
485	<i>Parinari excelsa</i> Sabine (Rosaceae)	koen=m hagala=g	gA	S		Fr	Fr		
486	<i>Parkia bicolor</i> A.Chev (Mimosaceae)	komi=m koi=g	gA	Fsd	1	2	Fr	Fr	
487	<i>Paspalum scrobiculatum</i> L (Gramineae)		H	S					
488	<i>Passiflora foetida</i> L (Passifloraceae)		L	Fs	1	2	3		
489	<i>Paullinia pinnata</i> L (Sapindaceae)		L	Fs	1	2	3	antidysentery antihemorrhoids	Fe
490	<i>Pauridiantha afzelii</i> (Hiern) (Rubiaceae)		pA	J					Tg(tooth brush)
491	<i>Pauridiantha hirtella</i> (Benth) Bremek		pA	J					

492 <i>Pennisetum pedicellatum</i> Trin (Gramineae)	H	S					
493 <i>Pennisetum polystachyin</i> (L) Schult	H	S					
494 <i>Pennisetum purpureum</i> Schum ka=m	H	S, Fs, J	1 2	Tg		vermifuge	Fe, Tg
495 <i>Pennisetum subangustum</i> Stapf & C.E.Hubb	H	S					
496 <i>Pennisetum violaceum</i> (Lam) L.Rich	H	S					
497 <i>Pentaclethra macrophylla</i> Benth (Mimosaceae)	gA	Fsd	1 2 5	Gr			
498 <i>Pentadesma butyracea</i> Sabine (Guttiferae)	A	Fs	2				
499 <i>Peperomia molleri</i> C.DC (Piperaceae)	H	Fs	1 2				
500 <i>Persea gratissima</i> Gaertn (Lauraceae)	pA	C		Fr	Fr		
501 <i>Phaseolus lunatus</i> L (Papilionaceae)	L	C		Gr, jFe	Gr		
502 <i>Phyllanthus alpestris</i> Beille (Euphorbiaceae)	H	S					
503 <i>Phyllanthus discoideus</i> (Baill) Mull Arg	pA	S, Fs	2 3				
504 <i>Phyllanthus muellerianus</i> beibabure (O.Ktze) Exell zumure=m tolou, kalagbomon=g	L	Fs	1 2 3	Fr			
505 <i>Phyllanthus niruri</i> L	H	J					
506 <i>Physalis angulata</i> L (Solanaceae)	H	J					
507 <i>Physalis micrantha</i> Link	L	J					
508 <i>Physostigma venenosum</i> Balfour (Papilionaceae)	L	Fs	1 2				
509 <i>Piper guineense</i> Schum & Thonn (Piperaceae)				jFr, Fe	Fr, jFe (sauce)		
510 <i>Piper umbellatum</i> L	H	Fs	1 2	jFe	jFe	vermifuge	Fe
511 <i>Piptadeniastrum africanum</i> bela=m (Hook.f) Brenan(Mimosaceae)	gA	F	1 2	Fr			

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	part	Other use
						chimps	men			
512	<i>Platycerium angolense</i> Welw (Polypodiaceae)	gnamouta Koulata=g	H	Fsd	1 2 3	jFe				
513	<i>Polycephalum capitatum</i> (Baill) Keay (Icacinaceae)		L	Fs	1 2 3 4	Fe		hemostatic	Fe	
514	<i>Portulaca oleracea</i> L (Portulacaceae)		H	R						
515	<i>Premna hispida</i> Benth (Verbenace)		pA	Fs	1 2 3	Fr				
516	<i>Pseuderanthemum tunicatum</i> (Afzel)(Acanthaceae)		pA	F	1 2					
517	<i>Pseudonanthenum tunicatum</i> (Acanthaceae)		H	M	2					
518	<i>Pseudospondias microcarpa</i> (A.Rich) Engl(Anacardiaceae)	poni=m	A	Fs	1 2 3	Fr				
519	<i>Psidium guajava</i> (Ladd) L (Myrtaceae)		pA	C		Fr, jFe	Fr	antidysentery	Fe	
520	<i>Psychotria vogeliana</i> Benth (Rubiaceae)		pA	Fs	1 2					
521	<i>Pteridium aquilinum</i> (L) Kuhn (Polypodiaceae)		H	J		jFe	jFe	vermifuge	Fe	
522	<i>Pteris atrovirens</i> Willd (Polypodiaceae)		H	M						
523	<i>Pterocarpus santalinoides</i> L'Her ex DC(Papilionaceae)	gbano=g	pA	M		jFe, Gr	Gr	antidysentery	Fe, Ec	
524	<i>Pycnanthus angolensis</i> (Welw) Warb (Myristicaceae)	dini=m gboi=g	A	Fs	1 2 3 4	Fr		antidentalgia	Ec	wood (furniture)
525	<i>Pycreus</i> sp (Cyperaceae)		H	M						
526	<i>Pyrenacantha acuminata</i> Engl (Icacinaceae)		L	Fs	1 2 3	Fr				
527	<i>Raphia gracilis</i> Beck (Palmæ)	duo=m	A	M,C		Gr(pulp) Gm(sap)	Gr(pulp) Gm(sap)	antiburn	Tg	bFe(mat house roof) Fe(sack)
528	<i>Rauvolfia vomitoria</i> Afzel (Apocynaceae)		pA	Fs	1 2 3 4					

529 <i>Renealmia maculata</i> Stapf (Zingiberaceae)		H	F	1 2	Tg
530 <i>Rhaphiostylis beninensis</i> (Hook.f ex Planch) Planch	purpure m ex Benth (Icacinaceae)	pA	F	1 2 3	Fr
531 <i>Rhynchospora</i> sp (Cyperaceae)		H	M		
532 <i>Rhytachne</i> sp (Gramineae)		H	S		
533 <i>Ricinodendron heudelotii</i> (Baill) Pierr & Pax	koo=m gbolo=g (Euphorbiaceae)	gA	Fs	1 2 3	Fr,Gr Gr
534 <i>Ricinus communis</i> FL (Euphorbiaceae)		pA	C		
535 <i>Rinorea dentata</i> (P.Beauv) O.Ktze (Violaceae)		pA	Fs	1 2	
536 <i>Rinorea oblongifolia</i> (C.H.Wright) Marq ex Chip		pA	Fs	1 2	Fr
537 <i>Rinorea subintegritifolia</i> (P.Beauv) O.Ktze		pA	Fs	1 2	
538 <i>Rothmannia hispida</i> (K.Schum) Fagerlind (Rubiaceae)		pA	Fs	1 2 3	
539 <i>Rothmannia longiflora</i> Salisb		pA	Fs	1 2 3	
540 <i>Rubus</i> sp (Rosaceae)		pA	Fs	1 2 3	
541 <i>Ruthalicia eglandulosa</i> (Hook.f) C.Jefrey (Curcurbitaceae)	L	Fs	1 2 3		
542 <i>Rutidea parviflora</i> DC (Rubiaceae)		pA	Fs	1 4 2 3	Fr
543 <i>Sabicea discolor</i> Stapf (Rubiaceae)		L	Fs	1 2 3	
544 <i>Sabicea harleyae</i> Hepper		L	Fs	2 3	
545 <i>Sabicea venosa</i> Benth		L	Fs	1 2 3	
546 <i>Sabicea vogelii</i> Benth		L	Fs	1 2 3	
547 <i>Saccharum officinarum</i> L (Gramineae)	guilo=g	H	C	Tg	vermifuge Fe
548 <i>Sacciolepis</i> sp (Gramineae)		H	S		

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use function	Other use part
						chimps	men		
549	<i>Salacia columnna</i> H.Halle (Celastraceae)		L	F	2				
550	<i>Santiria trimera</i> (Oliv) Aubr (Burseraceae)		pA	F	1	Fr	Fr		
551	<i>Sarcophrynum brachystachys</i> (Benth) K.Schum (Märantaceae)		H	F	1	Tg			
552	<i>Sarcophrynum prionogonium</i> K.Schum		H	F	1	Tg			
553	<i>Schizachyrium platyphyllum</i> (Franch) Stapf (Gramineae)		H	S					
554	<i>Schenckia americana</i> L (Solanaceae)		H	S, R					
555	<i>Scirpus</i> sp (Cyperaceae)		H	S					
556	<i>Scleria barteri</i> (Cyperaceae)		H	J					
557	<i>Scoparia dulcis</i> L (Scrophulariaceae)		H	R					
558	<i>Securidaca welwitschii</i> Oliv (Polygalaceae)		L	F	1 2 3				
559	<i>Selaginella blepharophylla</i> Alston (Selaginellaceae)		H	F	1 2 3				
560	<i>Selaginella cathedrifolia</i> Spring		H	F	1 2 3				
561	<i>Selaginella molliceps</i> Spring		H	F	1 2 3				
562	<i>Sesamum indicum</i> L (Pedaliaceae)		H	C					
563	<i>Sesamum radiatum</i> Schum & Thonn		H	C					
564	<i>Sesbania sesban</i> L (Papilionaceae)		H	J					
565	<i>Setaria chevalieri</i> Stapf (Gramineae)		H	J					
566	<i>Setaria megaphylla</i> (Steud) Dur & Sching		H	J					

567 <i>Sherbournia bignoniflora</i> (Welw) Hua (Rubiaceae)		L	Fs	1 2 3	Fr	Fr	anticough	Fe
568 <i>Sherbournia calycina</i> (G.Don) Hua	kinegbo=g	L	Fs	1 2 3	Fr	Fr jFe	anticough	Fe
569 <i>Sida linifolia</i> Juss ex Cav (Malvaceae)		H	M					
570 <i>Sida rhombifolia</i> L		H	R					
571 <i>Sida stipulata</i> Cav		H	J					
572 <i>Sida urens</i> L		H	J					
573 <i>Smilax kraussiana</i> Meisn (Smilacaceae)	peango=m	L	Fs	1 2 3	jFe			
574 <i>Solanum aculeatissimum</i> Jacq (Solanaceae)		Ar	F, J	1 2 3				
575 <i>Solanum erianthum</i> D.Don		pA	J					
576 <i>Solanum lycopersicum</i> L		H	C		Fr	Fr		
577 <i>Solanum macrocarpon</i> L		H	C		Fr	Fr		
578 <i>Solanum nodiflorum</i> Jacq		H	C		Fr	Fe(sauce)		
579 <i>Solanum terminale</i> Forsk		L	J					
580 <i>Solenostemon</i> sp (Labiatae)		H	J					
581 <i>Sopubia ramosa</i> Hochst (Scrophulariaceae)		H	S					
582 <i>Sorghum exiguum</i> Forsk (Gramineae)		H	C		Gr			
583 <i>Spathodea campanulata</i> P.Beauv (Bignoniaceae)		A	Fs	1 2 3				
584 <i>Spigelia anthelmia</i> L (Loganiaceae)		H	R					
585 <i>Spilanthes uliginosa</i> SW (Compositae)		H	J,R			jFe (sauce)	antidentalgia	F1
586 <i>Spondias mombin</i> L (Anacardiaceae)	buna,dian =m	pA	Fs	1 2 3 4	Fr	Fr	antirheumatism	Fe
587 <i>Sporobolus festivus</i> Hochst ex A.Rich (Gramineae)		H	S					
588 <i>Stachytarpheta angustifolia</i> (Mill) Vahl (Verbenaceae)		H	S					

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use		Other use
						chimps	men	function	part	
589	<i>Sterculia tragacantha</i> Lind (Sterculiaceae)	tu=m ko=a=g	A	Fs	1 2 3	jFe,Fr		fontanel treatment		Gm
590	<i>Streptogyna crinita</i> P.Beauv (Gramineae)		H	F	1 2					
591	<i>Striga bilabiata</i> (Thunb) O.Ktze (Scrophulariaceae)		H	S						
592	<i>Striga rowlandi</i> Engl		H	S						
593	<i>Strophanthus hispidus</i> DC (Apocynaceae)		L	Fs	1 2 3					
594	<i>Strophanthus preussii</i> Engl & Pax		L	Fs	1 2 3					
595	<i>Strophanthus sarmentosus</i> DC		L	Fs	1 2 3					
596	<i>Strychnos aculeata</i> Solereder (Loganiaceae)		L	F	2					
597	<i>Strychnos floribunda</i> Gilg		L	Fs	1 2					
598	<i>Synedrella nodiflora</i> Gaertn (Compositae)		H	J						
599	<i>Syzygium</i> sp (Myrtaceae)		pA	Fs	1 2					
600	<i>Tabernaemontana glandulosa</i> (Stapf) Pichon (Apocynaceae)		pA	Fs	1 2	Fr				
601	<i>Tabernaemontana longiflora</i> Benth		pA	Fs	1 2	Fr				
602	<i>Tapinanthus bangwensis</i> (Engl & K.Krause) Danšer(Loranthaceae)	L (parasite)			1 2	Fr				
603	<i>Taraktogenus</i> sp (Flacourtiaceae)		A	F	2	Ec		aphrodisiac	Ec	
604	<i>Tarenna pavettoides</i> (Harv) Sim (Rubiaceae)		pA	S						
605	<i>Tarrietia utilis</i> Sprague (Sterculiaceae)		A	F	1 2					
606	<i>Tectona grandis</i> L.F. (Verbenaceae)		A	C						
607	<i>Telphairia</i> sp (Cucurbitaceae)		L	F	2		Gr			
608	<i>Tephrosia</i> sp (Papilionaceae)		H	S						

609 <i>Terminalia glaucescens</i> Planch ex Benth (Combretaceae)	A	Fs, S	1 2	jFe		
610 <i>Terminalia ivorensis</i> A.Chev	gA	F, Fs	1 2 3	jFe		
611 <i>Terminalia superba</i> Engl & Diels	A	F, Fs	1 2	jFe		
612 <i>Tetracera alnifolia</i> Willd (Dilleniaceae)	L	Fs	1 2 3	jFe	Gm (drink)	
613 <i>Tetracera potatoria</i> Afzel ex G.Don zupeibrebro	L	Fs	1 2 3	Fe	Gm (drink)	
614 <i>Tetrapleura tetraptera</i> Schum & Thonn (Mimosaceae)	A	Fs	1 2	jFe		
615 <i>Tetrorchidium didymostemon</i> (Baill) Pax & K.Hoffm (Euphorbiaceae)	pA	Fs	1 2	jFe		
616 <i>Thaumatococcus daniellii</i> sa=m (Benn) Benth (Marantaceae)	H	Fs	1 2	Fr	Fr	Fe(mat)
617 <i>Thevetia nerifolia</i> Vahl (Apocynaceae)	pA	C			antiicterus	Fe
618 <i>Theobroma cacao</i> Gaertn (Sterculiaceae)	pA	C		Fr	Gr	
619 <i>Thonningia sanguinea</i> Vahl loitou=g	H	Fs	1 2 3		antiicterus	Tg
620 <i>Thunbergia</i> sp (Acanthaceae)	L	J				
621 <i>Tithonia diversifolia</i> A.Gray (Compositae)	pA	C				
622 <i>Trema guineensis</i> (Schum & Thonn) Ficalho (Ulmaceae) waba=g	pA	Fs	1 2 3	Fr	filaricide	Fe
623 <i>Tricalysia bracteata</i> Hiern (Rubiaceae)	pA	Fs	1 2			
624 <i>Trichilia heudelotii</i> Planch ex Oliv (Meliaceae)	A	Fs	1 2 3	Fr		
625 <i>Trichilia lanata</i> A.Chev	A	Fs	1 2 3 5	Fr		
626 <i>Trichomanes erosum</i> Willd (Hymenophyllaceae)	H	F	1 2 3			
627 <i>Triplochiton scleroxylon</i> K.Schum (Sterculiaceae) jokuro=m zogoloulu=g	A	Fs, F	1 2 3	jFe		wood (furniture)
628 <i>Uapaca guineensis</i> Mull Arg (Euphorbiaceae)	sonan,doh =m	A	F	1 2 3	Fr	scabicide Fe

(cont.)

## (Appendix cont.)

No.	Scientific name	Vernacular name	Plant type	Vege-tation	Place	Consumed by		Medical use		Other use
						chimps	men	function	part	
629	<i>Vapaca paludosa</i> Aubr		pA	F		1	2			
630	<i>Urera cameroonensis</i> Wedd (Urticaceae)		L	Fs		1	2	3		
631	<i>Urera oblongifolia</i> Benth		L	Fs		1	2			
632	<i>Urera obovata</i> Benth		L	Fs		1	2			
633	<i>Urera rigida</i> (Benth) Keay		L	Fs		1	2			
634	<i>Utricularia subulata</i> L (Lentibulariaceae)		H	S						
635	<i>Vanquieriopsis discolor</i> (Benth) Rabyns (Rubiaceae)		L	J						
636	<i>Vernonia cinerea</i> (L) Less (Compositae)		H	S						
637	<i>Vernonia conferta</i> Benth		pA	Fs		1	2			
638	<i>Vernonia gerberiformis</i> Oliv & Hiern		H	S						
639	<i>Vernonia guineensis</i> Benth		H	S						
640	<i>Vernonia nimbaensis</i> CD.Adams		H	S						
641	<i>Vetiveria nigritana</i> (Benth) Stapf (Gramineae)		H	C						
642	<i>Vigna multiflora</i> Hook.f (Papilionaceae)		H	S						
643	<i>Virectaria major</i> (K.Schum) Verdc (Rubiaceae)		H	S						
644	<i>Virectaria multiflora</i> (Sm) Bremek		H	S						
645	<i>Vismia guineensis</i> (L) Choisy (Hypericaceae)		pA	Fs		1	2	3		
646	<i>Vitex cienkowskii</i> Kostachy & R (Verbenaceae)		A	Fs		1	2	3	Fr	
647	<i>Vitex Doniana</i> Sweet		A	Fs, S		1	2	Fr	Fr	
648	<i>Vitex ferruginea</i> Schum & Thonn		A	F		1	2	Fr		

649 <i>Vitex grandifolia</i> Gurke	A	Fs	1 2 3	Fr		
650 <i>Vitex madiensis</i> Oliv	A	S		Fr		
651 <i>Vitex micrantha</i> Gurke	A	F	1 2			
652 <i>Vitex oxycuspis</i> Bak	A	F	1 2	Fr		
653 <i>Voacanga africana</i> Stapf (Apocynaceae)	pA	Fs	1 2 3		febrifuge	Fe
654 <i>Voacanga thouarsii</i> Roem & Schult	pA	M				
655 <i>Voandzeia subterranea</i> (L) DC (Papilionaceae)	H	C		Gr		
656 <i>Waltheria americana</i> L (Sterculiaceae)	H	S			anodyne aphrodisiac	Tg, Tb
657 <i>Xylia evansii</i> Hutch (Mimosaceae)	A	Fs	1 2			
658 <i>Xylopia aethiopica</i> (Dunal) A.Rich (Annonaceae)	pA	Fs	1 2 3	Fr	Fr	
659 <i>Xylopia staudtii</i> Engl & Diels	pA	Fs	1 2 3	Fr	Fr	
660 <i>Xyris filiformis</i> Lam (Xyridaceae)	H	S				
661 <i>Xyris straminea</i> Nilss	H	S				
662 <i>Zea mays</i> L (Gramineae)	bae=m kpai=g	H	C	Fr	Fr	
663 <i>Zinnia elegans</i> Jacq (Compositae)		H	C			(ornament)
664 <i>Zornia</i> sp (Papilionaceae)	H	J				

---