


A History and Annotated Account of the Benthic Marine Algae of Taiwan



JANE E. LEWIS
and
JAMES N. NORRIS

SERIES PUBLICATIONS OF THE SMITHSONIAN INSTITUTION

Emphasis upon publication as a means of "diffusing knowledge" was expressed by the first Secretary of the Smithsonian. In his formal plan for the Institution, Joseph Henry outlined a program that included the following statement: "It is proposed to publish a series of reports, giving an account of the new discoveries in science, and of the changes made from year to year in all branches of knowledge." This theme of basic research has been adhered to through the years by thousands of titles issued in series publications under the Smithsonian imprint, commencing with *Smithsonian Contributions to Knowledge* in 1848 and continuing with the following active series:

Smithsonian Contributions to Anthropology
Smithsonian Contributions to Astrophysics
Smithsonian Contributions to Botany
Smithsonian Contributions to the Earth Sciences
Smithsonian Contributions to the Marine Sciences
Smithsonian Contributions to Paleobiology
Smithsonian Contributions to Zoology
Smithsonian Folklife Studies
Smithsonian Studies in Air and Space
Smithsonian Studies in History and Technology

In these series, the Institution publishes small papers and full-scale monographs that report the research and collections of its various museums and bureaux or of professional colleagues in the world of science and scholarship. The publications are distributed by mailing lists to libraries, universities, and similar institutions throughout the world.

Papers or monographs submitted for series publication are received by the Smithsonian Institution Press, subject to its own review for format and style, only through departments of the various Smithsonian museums or bureaux, where the manuscripts are given substantive review. Press requirements for manuscript and art preparation are outlined on the inside back cover.

Robert McC. Adams
Secretary
Smithsonian Institution

A History and Annotated Account
of the Benthic Marine Algae
of Taiwan

*Jane E. Lewis
and James N. Norris*



SMITHSONIAN INSTITUTION PRESS

Washington, D.C.

1987

A B S T R A C T

Lewis, Jane E., and James N. Norris. A History and Annotated Account of the Benthic Marine Algae of Taiwan. *Smithsonian Contributions to the Marine Sciences*, number 29, 38 pages, 1 figure, 1987.—Records of the benthic marine algae of the Island of Taiwan and neighboring islands have been organized in a floristic listing. All publications with citations of benthic marine green algae (Chlorophyta), brown algae (Phaeophyta), and red algae (Rhodophyta) in Taiwan are systematically arranged under the currently accepted nomenclature for each species. The annotated list includes names of almost 600 taxa, of which 476 are recognized today. In comparing the three major groups, the red algae predominate with 55% of the reported species, the green algae comprise 24%, and the browns 21%. *Laurencia bronniartii* J. Agardh is herein reported for Taiwan for the first time.

The history of modern marine phycology in the Taiwan region is reviewed. Three periods of phycological research are recognized: the western (1866–1905); Japanese (1895–1945); and Chinese (1950–present). Western phycologists have apparently overlooked the large body of Japanese studies, which included references and records of Taiwan algae.

By bringing together in one place all previous records of the Taiwanese marine flora, it is our expectation that this work will serve as a basis for further phycological investigations in the western Pacific region.

OFFICIAL PUBLICATION DATE is handstamped in a limited number of initial copies and is recorded in the Institution's annual report, *Smithsonian Year*. **SERIES COVER DESIGN:** Seascapes along the Atlantic Coast of eastern North America.

Library of Congress Cataloging in Publication Data

Lewis, Jane E.

A history and annotated account of the benthic marine algae of Taiwan.
(Smithsonian contributions to the marine sciences ; no. 29)

Bibliography: p.

Includes index.

Supt. of Docs. no.: SI 1.34 : 29

1. Marine algae—Taiwan—Classification—Bibliography. 2. Marine algae—Taiwan—Bibliography. 3. Marine algae—Research—Taiwan—History. I. Norris, James N. II. Title. III. Series

Z5356.A6L48 1987 016.5894'0951'249 86-600251

[Qk575.T28]

Contents

	<i>Page</i>
Introduction	1
Historical Review	1
Geographic and Oceanographic Features	4
Discussion	6
The Annotated List	6
Conclusion	7
Acknowledgments	7
Benthic Marine Algae from Taiwan	7
CHLOROPHYTA	7
ULOTRICHALES	7
ULOTRICHACEAE	7
CHAETOPHORALES	7
CHAETOPHORACEAE	7
ULVALES	7
ULVACEAE	7
CLADOPHORALES	8
CLADOPHORACEAE	8
CAULERPALES	8
CODIACEAE	8
UDOTEACEAE	9
CAULERPACEAE	9
BRYOPSIDACEAE	10
DERBESIACEAE	10
SIPHONOCLADALES	10
SIPHONOCLADACEAE	10
VALONIACEAE	10
ANADYOMENACEAE	11
DASYCLADALES	11
ACETABULARIACEAE	11
PHAEOPHYTA	11
ECTOCARPALES	11
ECTOCARPACEAE	11
CHORDARIALES	11
CORYNTHLAEACEAE	11
SPHACELARIALES	11
SPHACELARIACEAE	11
DICTYOTALES	11
DICTYOTACEAE	11
SCYTOSIPHONALES	12
ISHIGEACEAE	12
PUNCTARIACEAE	12
SCYTOSIPHONACEAE	13
FUCALES	13
CYSTOSEIRACEAE	13
SARGASSACEAE	13

RHODOPHYTA	14
GONIOTRICHALES	14
GONIOTRICHACEAE	14
BANGIALES	14
ERYTHROPELTIDACEAE	14
BANGIACEAE	14
NEMALIALES	15
ACHROCHAETIACEAE	15
DERMONEMATACEAE	15
LIAGORACEAE	15
CHAETANGIACEAE	15
BONNEMAISONIACEAE	16
GELIDIALES	17
GELIDIACEAE	17
GELIDIELLACEAE	17
CRYPTONEMIALES	17
DUMONTIACEAE	17
RHIZOPHYLLIDACEAE	17
PEYSSONNELIACEAE	18
CORALLINACEAE	18
ENDOCLADIACEAE	18
CRYPTONEMIACEAE	18
GIGARTINALES	19
SOLIERIACEAE	19
HYPNEACEAE	19
PLOCAMIACEAE	20
GRACILARIACEAE	20
SPHAEROCOCCACEAE	21
SARCODIACEAE	21
PHYLLOPHORACEAE	21
GIGARTINACEAE	21
RHODYMENIALES	21
RHODYMENIACEAE	21
CHAMPIACEAE	21
CERAMIALES	21
CERAMIACEAE	21
DELESSERIACEAE	22
DASYACEAE	22
RHODOMELACEAE	22
INCERTAE SEDIS	23
WURDEMANNIACEAE	23
Literature Cited	24
Index	34

A History and Annotated Account of the Benthic Marine Algae of Taiwan

Jane E. Lewis and James N. Norris

Introduction

Historical Review

The history of recognizing, naming, and using marine algae in China extends far back into the early Chinese literature. Although of uncertain antiquity, the specific mention of marine algae goes back at least to the publication of the *Er Ya** or *Literary Expositor* (Tseng and Chang, 1961), a dictionary that dates to the 3rd century B.C. (Needham, 1970). Application of Latin binomials to the Taiwan marine flora, however, began only in the late 19th century (Martens, 1866). After a period of European collections and reports, a period of extensive Japanese studies during the early to mid-20th century followed. The third, or current, period is one in which the activities of Chinese investigators predominate.

The first report of benthic Taiwan algae that followed Linnean taxonomy appeared in Georg von Martens' *Die Tange* (1866), based on the botanical explorations from the German expedition to East Asia, 1860 to 1862. During this voyage von Martens' son Eduard, chief zoologist for the expedition, collected marine and freshwater algae from Java, the Philippines, Singapore, Taiwan, Hong Kong, and other East Asian areas. In this work, the elder von Martens reported seven marine species from "Tamsui" (Danshui), a northeast Taiwan seaport, including three marine algal species, *Ulva lactuca* f. *lapathifolia*, *Gratelouphia filicina* f. *filiformis*, and *Caulacanthus ustulatus* var. *fastigiatus*, not again recorded from Taiwan.

During a second German expedition to East Asia (1886–1888) Dr. Warburg made phycological collections that were later published by Heydrich (1894) and De Toni (1895,

1905). This expedition covered the East Indian Ocean, the Mollucca Islands, China, Java, the southern Japanese islands, and a brief stop at Taiwan. Taiwan collection locations were noted as Jilong (a northeast seaport), south Taiwan, the east coast and "Long-kiau" on the south coast.

Among Warburg's Taiwanese algal collection, Heydrich identified 43 taxa, consisting of 24 red (Rhodophyta), 12 brown (Phaeophyta), and 7 green (Chlorophyta) species. In continued studies of the algae from this expedition, De Toni (1895) reported 36 taxa from Taiwan and in 1905 another two species. As a result of these early European investigators, over 60 marine benthic algae were reported for the Taiwan region, including many new to science.

The history of the exploration of Taiwan marine algae, as with other sciences and indeed all aspects of life, was greatly shaped during the 50-year period, beginning in 1895, of Japanese political dominance in Taiwan. Voluminous studies on the Japanese flora and fauna, then including Taiwan, were undertaken. It was during this time that exploration of the Taiwan marine flora was most intensive, dominated by the work of K. Okamura (e.g., 1900–1902, 1907–1942) and his student, Y. Yamada (1930–1944).

The first publication of this period, "New or Little Known Algae From Japan" (Okamura, 1895), included two new records of species from Taiwan. Many publications specifically addressed the Taiwan flora (Okamura, 1915b, 1931, 1935b; Oshima, 1915; Ariga, 1919, 1920, 1921; Horikawa, 1919; Yamada 1925a, 1925b, 1936a; Yamada and Tanaka, 1934; Tokida, 1941), but most references to the Taiwan flora were included as distributional notes or collection sites within general Japanese floristic works (Okamura, 1930, 1936; Yamada, 1928, 1934). Three important series con-

Jane E. Lewis, Department of Botany, University of Hawaii, and East-West Center, Honolulu, Hawaii 96822. James N. Norris, Department of Botany, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560.

* Throughout this text, Chinese place names and words not enclosed in quotation marks are spelled according to the Pin Yin romanization system.

taining numerous, but scattered, references to Taiwan algae are *Illustrations of the Marine Algae of Japan* (Okamura, 1900–1902) and *Icones of Japanese Algae* (Okamura, 1907, 1909, 1913, 1915a, 1921, 1923, 1932, 1934b, 1935a, 1937, 1942); Yamada's "Notes on Some Japanese Algae" (1930, 1931a, 1932b, 1933, 1935, 1936b, 1941, 1944b); and Yendo's "Notes on Algae New to Japan" (1909, 1914, 1915, 1916a, 1916b, 1918).

From an early time the Japanese were interested in floristic comparisons, sometimes speculating on the reasons for disjunct species distributions or factors that might cause adjacent areas to have divergent floras (Ariga, 1920, 1921; Okamura, 1915b, 1919). One of the first comparative phytogeographic studies of the western Pacific region (Yamada, 1926) included Taiwan as one of its eight floristic areas. Of the 42 taxa compared in the region, 16 were present in Taiwan.

Japanese monographs included new or additional records of taxa for Taiwan. Among these were studies of the Corallinaceae (Segawa, 1941; Yendo, 1902) and the Florideophyceae (Tanaka, 1944), and the genera *Gelidium* and *Pterocladia* (Okamura, 1934a), *Galaxaura* (Tanaka, 1935, 1936), *Hypnea* (Tanaka, 1941), *Laurencia* (Yamada, 1931b, 1936c), *Liagora* (Yamada, 1938a), *Porphyra* (Ueda, 1932), and *Sargassum* (Yamada, 1942).

Many of the papers written specifically on Taiwan's marine flora are in Japanese. Ariga (1919) describes a 14-day collecting trip and lists 43 taxa collected. Horikawa's (1919) "Marine Algae of Taiwan" reports 20 red, 10 brown, and 15 green species, based on determinations by Okamura of specimens that had been collected by Horikawa, Nagasawa, Maki, and their students at locales around Taiwan, including Danshui and Jilong in the north, Elanbi (Olanpi) in the south, Xiao Liuqiu (Shao Liou Chou) and Lan Yu, and the Penghu (= Pescadores) Islands. In another study a year later, Ariga (1920) reported 69 taxa for the Penghu Islands and compared them with his records from "Amoy," a small island off Fujian (= Fukien) Province, southeast China coast. Although only about 135 km of the shallow Taiwan Strait separate the two sites, the floras were found to be drastically different in composition. Even when the same species occurred in both places they often exhibited marked morphological or seasonal differences. It was speculated that tidal fluctuation or substrate type may have effected these differences. In determining species names, Ariga (1919, 1920) relied heavily on the papers of Yendo and Okamura.

The papers of Ariga (1919, 1920) and Horikawa (1919) were subsequently overlooked by later authors, perhaps due in part to having been written in Japanese. Both 1919 papers use the Latin scientific names and authors of the species followed by Japanese names. Ariga, in his subsequent paper (1920), uses these Japanese names almost exclusively. Few cases are known in other languages where common names apply solely to a single species. In the course of our research,

however, these Japanese names were found to be a special case because they are not common names but a Japanese version of the scientific name (Lai Chuen-fu, personal communication). When Latin binomials were introduced to Japan, the Japanese erected equivalent Japanese scientific names, one for each taxon, which were different from the Japanese common names. Because the Japanese names are listed with their Latin binomials in Okamura's flora (1936), it has been possible to translate Ariga's names into the binomial form for inclusion in the current listing.

Yamada's studies on the Chlorophyceae (1925a) and the Phaeophyceae (1925b) of Taiwan, representing his undergraduate thesis at Tokyo University under K. Okamura, are frequently cited in later publications. Most of the specimens of these papers were collected by Yamada during a short spring collecting trip to the north and south coasts of Taiwan and to the Penghu Islands, with a few additional specimens provided by T. Aoki, then a government official in Taiwan.

Economic considerations have also contributed to identification of Taiwan's marine flora. Some papers were devoted to the agarophytes in Taiwan at a time when the Japanese seaweed industry was flourishing (Okamura, 1915b, 1935b; Oshima, 1915). Taiwan's east coast was thought to possibly have a richer agarophyte flora than north Taiwan or any other coast in Japan. However, due to the presence of unfriendly aborigines (Okamura, 1915b), this hypothesis was not investigated until much later, and was found to be incorrect (Fan, 1951).

Several papers discuss Taiwan algae used as foods. The island of Lan Yu (also called Kōtōsho, Botel Tobago, or Orchid Island) has attracted much attention because of its aboriginal population, unique biology, and its close proximity (61 km) to the island of Taiwan. An issue of the *Bulletin of the Biogeographical Society of Japan* (1931, vol. 2, no. 2) was devoted to the anthropology and biology of Lan Yu, with Okamura (1931:95–122) reporting 92 taxa of benthic algae from collections made by S. Segawa during one summer month in 1930. Only about one-third of these algae (Okamura, 1931) were also known to be on both Taiwan and Lan Yu. The Lan Yu flora was considered to be Indo-Pacific in nature, whereas no specific comments were made on the affinities of the Taiwan Island marine flora. Of the many algae presumably eaten by the "Yami" tribe on Lan Yu, Okamura listed 12 (*Carpopeltis formosana*, *Laurencia* sp., *Acanthophora orientalis*, *Halymenia durvillaei* var. *formosa*, *Hypnea seticulosa*?, *Nemalion pulvinatum*, *Chondria armata*, and *Dermoneema dichotoma*), also giving a Japanese transliteration of their traditional Yami names. Subsequently, Tokida (1939) published an almost identical list of algae eaten by the "Ami" people on Taiwan with the same common names.

An identification book with photographs of Japanese seaweeds published during this period (Higashi, 1934) in-

cluded 12 records of Taiwan taxa. With the high scholarly level characteristic of the Japanese picturebook genre, specific names are generally considered accurate. Because these books contain records of Taiwan algae often not previously published, it is believed that distributional citations were based on Japanese herbarium material.

The Japanese flora *Nippon Kaiso-shi* (Okamura, 1936) contains numerous reports of algal species in Taiwan. This is an invaluable reference for the identification of benthic marine algae throughout the northwestern Pacific region.

In addition to the extensive Japanese publications of the period 1895–1945, some Chinese and western papers also included accounts of the Taiwan marine flora. A.D. Cotton (1915) reported nine taxa from Taiwan. Specimens of *Sargassum* were verified by K. Yendo, who visited the Royal Botanic Garden, Kew, while the paper was in preparation.

The lack of scientific communication between the East and West is perhaps characterized by J. Tilden's (1929) comments in "The Marine and Freshwater Algae of China." Although Professor Tilden claims to record "all species previously noted by investigators" for the algae of China, her account of Chinese phycological history only cites one non-western reference, a short paper by Okamura (1913) on "Chinese Edible *Nostoc*." She clearly agrees with Cotton (1915) that "Formosa [now known as Taiwan] . . . though belonging to Japan, must geographically be included with China." Though aware of Martens' (1866) account of algae, Tilden considered his determinations to be unreliable, and her account of the Taiwan marine flora was limited to only the nine taxa Cotton (1915) had reported.

Apparently unaware of the Japanese literature of 1895–1928, Tilden listed only 92 species of marine algae from the entire Chinese region, as "all that has been done in the study of Chinese algae." If the Japanese literature (e.g., Okamura, 1909–1926) and other accounts of European expeditions had been included, she would have found over 350 published records of more than 200 species for Taiwan alone by 1928. It is easy today to see the inaccuracy of Tilden's assessment; however, her conclusions were understandable considering the barriers of language, culture, and communication at that time.

Professor C.K. Tseng, as a young botany instructor from Fujian Province, made extensive collections in the 1930's throughout China. In studies of marine Chlorophyceae from Hainan Island (1936), Chinese Chaetangiaceae (1941b), Hong Kong *Polysiphonia* (1944), and "New and Unrecorded Marine Algae of Hong Kong" (1945), Tseng included several Taiwan collections, many of them new records.

The first publication by Chinese marine botanists on Taiwan after the Japanese occupation came from the Taiwan Fisheries Research Institute. Y.F. Shen and K.C. Fan (1950) compiled much of the earlier literature and studied their own collections as well as specimens from the herbar-

ium at the National Taiwan University made from Kōtōsho by Y. Yamamoto, which resulted in a list of 62 green, 50 brown, and 142 red taxa. Locations were given as "Taiwan," "Kōtōsho" (Lan Yu), "Kasyoto" (Lu Dao) and the "Pescadores" (Penghu Islands).

Fan (1951) described eight species and two forms of the economic genera *Gelidium* and *Pterocladia* from Taiwan, with English translations of Dr. Okamura's (1935) four new taxa. *Gelidium* and *Pterocladia* were found to be restricted to the north and northeast part of the island. The distributions of the various species were compared and found to belong to two distributional groups (a disjunct north-south distribution), but Fan concluded that "[the] chief factors that delimit the mutual exclusion of these species are at present uncertain."

A list of edible marine algae (Fan, 1952) from Taiwan included 30 taxa used by Taiwan aborigines as well as those with a long Chinese tradition, with brief descriptions and localities of the algae, preparation methods, and a table of their Chinese and aboriginal names. Many species were noted to be commonly occurring only in the spring, though no reason was suggested.

Since 1960, Y.M. Chiang (now a professor at Taiwan National University), has been a major investigator of the Taiwan marine algae. Studies on floristics (Chiang, 1960, 1962a,b, 1973a,b), taxonomy (Chiang, 1981; Chiang and Chen, 1982; Yang and Chiang, 1982), reproduction (Chen and Chiang, 1982; Chiang, 1969, 1970, 1971, 1972; Chiang and Chen, 1982), and aquaculture (Chiang, 1981, 1982; Chiang and Chou, 1980; Liaw and Chiang, 1979; Nelson et al., 1983) have been the focus of his phycological contributions.

Professor Chiang began phycological publications (as a student of Y.F. Shen, Botany Department, National Taiwan University) with his two-part "Marine Algae of Northern Taiwan," the blue-green, brown, and green algae (1960), and the red algae (1962a). Included in these works were information on collection locations, habitat notes and seasonality of the major algae, identification keys, and taxonomic summaries of each of the 96 species.

After a field trip of a few days to Penghu, Chiang (1962b) recorded 26 species and noted the fragmentary nature of earlier Japanese records in the Penghu Islands. Dr. Chiang began working on Taiwan's phytogeographic affinities with "Notes on Marine Algae of Taiwan" (1973a), which included north-south distributional observations and new records of six green, six brown, and nine red algae for Taiwan and its offshore islands. "Studies on the Marine Flora of Southern Taiwan" (1973b) reported on the algae at four southern localities and compared them with the north, finding that "some northern species do not occur in the southern waters. On the contrary, there are quite many species which occur in southern Taiwan but not in the northern regions." The southern flora was also considered

to be more depauperate than the northern and unique within Taiwan in having typically tropical elements, such as *Bornetella* and *Neomeris*. The water temperatures, warmer and less variable throughout the year in the south, were suggested to be partially responsible for this difference. Because Okinawa, located slightly to the north and east of Taiwan, shares tropical genera with southern Taiwan, Chiang also suggested that the Kuroshio (or Japanese) Current is likely to be a major factor in distribution.

A later series of papers focus on aspects of morphology and reproduction (Chiang and Chou, 1980; Chen, Chiang, and Chiang, 1981; Chen and Chiang, 1982; Chiang, 1982; Chiang and Chen, 1983) and vegetative reproduction in a brown alga (Chiang and Chou, 1980). These papers provide taxonomic information, some of which is new for the Taiwan flora, and others that substantiate previous records. Taxonomic studies on *Sargassum* (Chou and Chiang, 1981), *Liagora* (Chiang and Chen, 1982), and *Gracilaria* (Yang and Chiang, 1982) provide additional listings with distribution, morphology and seasonality information.

Aquaculture in Taiwan is a well-developed business principally of shellfish and fin fish, and in the last 10 years the culture of marine algae (Chueh and Chen, 1982), especially *Gracilaria* (Chen, 1976; Shang, 1976; Michanek, 1978; Ryther, 1979; Tseng, 1981a,b; Chiang, 1981; Doty, 1983), and *Porphyra* (Chiang, 1982), has received more attention from both growers and taxonomists. Of the numerous publications on seaweed culture, those providing records of native species in Taiwan were most appropriate for inclusion in this listing.

In a review of *Gracilaria* culture in Taiwan, Chiang (1981) and Hansen et al. (1981) provided taxonomic information on the three native species cultured (*G. edulis*, *G. gigas*, and *G. verrucosa*) and the techniques employed. The seaweed aquaculture program, centered in the southwest of Taiwan, began in 1961, and by 1979 about 12,000 tons of the dry seaweed were being harvested, predominately for the domestic agar industry, and about 120 tons of fresh seaweed went as feed to abalone farms.

Gracilaria is the major source of agar-agar, along with some *Gelidium*. Reports on seasonal variation of agar quality and quantity produced in Taiwan (Yang, 1982; Yang et al., 1981) provide records on the native agarophyte flora, and a comparison of agar from species in Taiwan and Micronesia (Nelson et al., 1983) provides information on Taiwan sites of natural populations.

In contrast to *Gracilaria*, *Porphyra* culture has been small-scale and irregular. Cultivation of local species in the Penghu Islands began in 1968, but studies on the life history of the local species were not initiated until 1975 (Chiang, 1982), when the "Conchocelis-stage" of *Porphyra* and monospore formation were investigated. This study and two on *Conchocelis* culture (Chiang and Chou, 1980; Liaw and Chiang, 1979) also provide taxonomic information on native Taiwan taxa.

Alginates, a family of chemicals used in food preparation and with a wide variety of industrial uses, which conventionally are extracted from cold-water kelps, may become an industry in Taiwan, using a tropical brown alga. Research has been conducted (Liu, 1982) on improving the quality of alginates from *Sargassum duplicatum*, a species growing naturally along the shoreline of the southern tip of Taiwan.

Green algae are also a part of Taiwan's aquaculture industry. Liu (1982) mentions use of *Chlorella*. Included in a report on chemical analysis and utilization of *Monostroma* (Wu, 1982) is information on the two native species and their distribution.

The study of biological activity and natural-product chemistry (e.g., Norris and Fenical, 1985) is a recent research interest in Taiwan. For example, in a recent paper, "Pharmacological Properties of Some Taiwan Seaweeds" (Su et al., 1982), 30 species from Taiwan were screened for antimicrobial activity.

Geographic and Oceanographic Features

The area encompassed in this study includes the main island of Taiwan and a few of its 14 associated islands as well as the 64 islands of the Pescadores Archipelago (Figure 1). The islands most commonly referred to in the phycological literature are those of Taiwan (= Formosa) ($21^{\circ}53' - 25^{\circ}18'N$, $120^{\circ}1' - 122^{\circ}0'E$), Orchid Island (= Kôtôsho or Lan Yu) ($22^{\circ}0' - 22^{\circ}5'N$, $121^{\circ}36' - 121^{\circ}30'E$), Green Island (= Kasyoto or Lu Dao) ($22^{\circ}38' - 22^{\circ}41'N$, $121^{\circ}28' - 121^{\circ}30'E$) and the Pescadores (= Penghu Islands) ($23^{\circ}11' - 23^{\circ}46'N$, $119^{\circ}18' - 119^{\circ}42'E$). The most frequented collecting sites on the Island of Taiwan have been Olanpi at the southern tip, "Tai Dung" on the east coast, and the general area of the northern tip. Bisected by the Tropic of Cancer, the Island of Taiwan is considered both tropical and subtropical. Geologically it is a continental island with a mostly sedimentary origin, and it has a coastline of roughly 1600 km.

Oceanographic conditions around the island vary in topography, temperature, and currents. The Taiwan Strait is shallow (60 m average depth) and turbid, extending westward from Taiwan some 140 to 200 nautical miles to the southeast coast of mainland China. Within the Strait, the Penghu Islands are some 40 km from Taiwan. Tidal currents around these islands reach almost 6 knots in places (USDD Nautical Map #94060). In the East China Sea northeast of Taiwan is the Ryukyu Island chain, beyond which are the main islands of Japan. To the south the Bashi Channel separates Taiwan and the Philippines. The east coast, often precipitous, marks the eastern edge of the continental shelf. The sharp drop-off continues some 4000 m below sea level, reaching the floor of the Philippine Basin. Along the southeast coast are areas of upwelling, with colder, nutrient-rich waters.

Many benthic marine algae are found growing in the




FIGURE 1.—Map of Taiwan, showing islands and locales most commonly referred to in the phycological literature.

nearshore coastal waters. Shoreline conditions in Taiwan vary from sandy to rocky shores with murky water in the north, to coral reefs with clear swift water in the south. To the east are precipitous cliffs and rough waters with deep-ocean upwellings along the coast and, in contrast, level, calm expanses of intertidal mud and sand flats along the western shoreline.

The two main currents affecting Taiwan are the Kuroshio (or Japanese) Current and the Taiwan coastal current. The Kuroshio Current brings water of high temperature and salinity up from the Philippines and the equatorial region toward southern Taiwan where it branches. The stable, main branch runs past the east coast of Taiwan at an average velocity of 30 to 40 nautical miles per day toward Japan (USDD Nautical Map #94010). A smaller branch, subject to seasonal variability, veers west through the Taiwan Strait.

Flowing south along the China mainland coastal region, the Taiwan coastal current carries colder water from north to south, and is strong in summer and weak in winter. Two smaller seasonal currents, the northeast monsoon current and the southwest seasonal current, result from an interaction between the two major currents and the prevailing winds.

Discussion

While compiling the records of Taiwan algae from the numerous publications, efforts were made to find the correct name as well as the valid date and place of publication for each of the taxa in the annotated list. Unfortunately, phycologists do not have a modern source for specific names such as *Index Kewensis* (Jackson et al., 1893 to date; see Stafleu and Cowan, 1979:397–398, for complete listing) for phanerogams, or *Index Filicum* (Christensen, 1905, 1906; see Stafleu and Cowan, 1976:501–501, for complete listing) for ferns. Our effort has shown how great the need is for such an index of algal species names, such as has recently been done for the generic names of fossil and living plants (Farr et al., 1979). In the absence of such a reference, we found the earlier works of De Toni (1895–1924) and Dawson (1962) helpful.

The Taiwan algae project was undertaken recognizing that a single bibliographic source on the Taiwan marine flora would facilitate identification of the algae and their distribution and would encourage further research within the region. It was evident that the existing literature was widely dispersed and not adequately referenced in most phycological bibliographies. It became necessary to carefully research the east Asiatic algal literature, particularly from Japan and China, and the whole Pacific region, in order to locate information on Taiwan algae. Work on this compilation commenced while the first author was employed at the Fisheries Biology Laboratory, Institute of Zoology, Academia Sinica, Taiwan. Additional extensive

searches were necessary to locate original publications to establish proper citation of the binomial and to subsequently determine the current taxonomic and systematic status of each taxon. For this work, the E. Yale Dawson Phycological Library and Department of Botany libraries of the National Museum of Natural History, Smithsonian Institution, were invaluable.

The annotated list stands as a compilation of reports of attached, benthic marine taxa occurring naturally in the region of Taiwan. Efforts were taken to locate all relevant literature, and although it is possible that additional reports can be found, the present work represents the most comprehensive compilation available for the region. It must be emphasized that records are taken from published papers and that the accuracy of these identifications has not been confirmed by herbarium investigations; reported names of taxa are simply noted and arranged systematically following current convention. Verification of the determinations awaits future investigation. Some indication of accuracy may, however, be gained from the frequency and dates of the reports for any given taxon. For example, the three taxa reported by Martens (1866) that have not again been recorded may be considered in need of verification. Nevertheless, it is felt that the present list reasonably reflects the general composition of the Taiwan marine flora.

The Annotated List

There are 476 taxonomic entries in this list, systematically arranged following Abbott and Hollenberg (1976) for the red algae, and Lobban and Wynne (1981) for the brown and green algae. Below family level, genera and species within each genus are listed alphabetically. Each entry consists of (1) the currently accepted taxon name, with its author(s), and date and page of valid publication, and (2) an alphabetical list of the investigators who have reported the taxon present in the Taiwan region.

Orthographic errors in publication of taxa were found to have occurred nine times in this list (M.J. Lai and D.H. Nicolson, personal communication). These were simply corrected, with the original spelling in single quotations following "Recommendation 50F.1" of the *International Code of Botanical Nomenclature* (Voss et al., 1983).

In some cases there are additional names within the entry. In reporting a taxon from Taiwan, authors sometimes used names that have since been considered taxonomic or nomenclatural synonyms, or have been re-determined by later investigators. In all cases, the actual names used by the authors for their algal records from Taiwan are retained in this list. However, when those names differ from the currently accepted ones, they are included in the alphabetical list of references to the taxon. It is hoped that by preserving the original names, the concepts held by the authors will be

indicated and later taxonomic and nomenclatural changes will be more easily integrated into this list.

Conclusion

The marine flora of this region appears to have tropical, subtropical, and temperate elements. Many authors have commented on the diversity of floral elements within this region (e.g., Ariga, Chiang, Horikawa, Okamura) and have speculated on the possible causes. As noted earlier, water quality and substrate vary greatly in the region, no doubt affecting floral elements. Southern Taiwan and especially the two southeast islands Lan Yu and Lu Dao are referred to as having a tropical/subtropical flora and the north and west as having a temperate flora, judged by the abundance of brown algae and rarity of red algal elements. These correlations are deserving of further study. Indications are that a number of physical and biotic factors interact to create this diversity of floristic types exhibited in the Taiwan region.

Based on written reports from the literature, this study makes available much previously inaccessible information basic to all phycological studies of the East Asiatic region. This comprehensive list of marine algae, treated in both historical and modern systematic contexts, constitutes a basis for further systematic and ecological investigations and biogeographical comparisons among the East Asian and

Pacific areas, and serves as a possible framework for a marine flora of Taiwan.

Acknowledgments

The first author is very grateful to Jang Kun-Hsiung, Director of the Academia Sinica Zoology Institute and its Fisheries Biology Laboratory, Taipei, Taiwan, for the time and support to initiate this project, Chiang Young-Meng, Oceanography Department, National Taiwan University, Taipei, for references and encouragement throughout the project, and the Department of Botany and the Office of Fellowships and Grants, Smithsonian Institution, for support and use of the libraries, laboratory, and computers. A special appreciation is due to Isabella A. Abbott for her time, guidance, and patience and for the use of her personal library. Japanese translations were kindly provided by Lai Chuen-fu, Rika Taga, and Mason E. Hale. We thank Lai Ming-Jou and Dan H. Nicholson for discussions on nomenclature, and for checking the unpublished algal-name file at the University of California, Berkeley, J. Olsen-Stojkovich and P.C. Silva. For comments on the text, thanks go to J. Berdach, M.I. Cannon, A. Medeiros, and K.E. Bucher. Our appreciation to I.A. Abbott, K.E. Bucher, Y.M. Chiang, M.S. Doty, M.M. Littler, H.E. Robinson, and C.K. Tseng for reviewing the paper.

Benthic Marine Algae from Taiwan

CHLOROPHYTA

ULOTRICHALES

ULOTRICHACEAE

Ulothrix flaccida (Dillwyn) Thuret in Le Jolis, 1863:56.
Taniguti, 1976.

CHAETOPHORALES

CHAETOPHORACEAE

Endophyton ramosum Gardner, 1909:372.
Chiang, 1973a.

ULVALES

ULVACEAE

Enteromorpha clathrata (Roth) Greville, 1830:181.
Chiang, 1960; Fan, 1953a; Okamura, 1931, 1936; Shen and Fan, 1950; Tokida, 1939.
Enteromorpha clathrata var. *crinita* (Roth) Hauck, 1884:429.

As "*E. crinita*": Ariga, 1920; Chiang, 1960; Chihara, 1970; De Toni, 1895; Fan, 1953a; Heydrich, 1894; Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Tokida, 1939.

Enteromorpha compressa (Linnaeus) Greville, 1830:180.
Chiang, 1960; Fan, 1953a; Okamura, 1931, 1936; Shen and Fan, 1950; Taniguti, 1976; Tokida, 1939; Yamada, 1925a, 1926, 1950; Yoshikawa and Yoshikawa, 1977.

Enteromorpha intestinalis (Linnaeus) Link ex Nees, 1820:5.
Ariga, 1919; Chiang, 1960, 1973b; Fan, 1953a; Horikawa, 1919; Shen and Fan, 1950.

Enteromorpha linza (Linnaeus) J. Agardh, 1883:134.
Chiang, 1960, 1973b; Chiang et al., 1974. As "*Phycoseris lanceolata* var. *angusta*": Martens, 1866.

Enteromorpha prolifera (O.F. Müller) J. Agardh, 1883:129.
Chiang, 1960; Fan, 1953a; Shen and Fan, 1950; Taniguti, 1976.

Enteromorpha sp.
Chiang, 1973b.
Monostroma latissimum (Kützing) Wittrock, 1866:33.
Chiang, 1973a, 1973b; Okamura, 1935b; Rho, 1958.

- Monostroma nitidum* Wittrock, 1866:41.
 Chiang, 1960; Fan, 1953a; Okamura, 1936; Shen and Fan, 1950; Taniguti, 1976; Yamada, 1925a, 1925b, 1934, 1950.
- Ulva angusta* Setchell et Gardner, 1920:283.
 Chiang, 1973a; Okamura, 1935b.
- Ulva conglobata* Kjellman, 1897b:10.
 Ariga, 1920, 1921; Chiang, 1960, 1962a; Shen and Fan, 1950; Taniguti, 1976; Tokida, 1939; Yamada, 1925a; Yoshikawa and Yoshikawa, 1977.
- Ulva fasciata* Delile, 1813:155.
 Chiang, 1960, 1973b; Fan, 1953a; Shen and Fan, 1950; Tokida, 1939; Tseng, 1983; Yamada, 1935; Yoshikawa and Yoshikawa, 1977.
- Ulva japonica* (Holmes) Papenfuss, 1960:309.
 As "Letterstedtia japonica": Chiang, 1973a; Okamura, 1935b; Taniguti, 1976.
- Ulva lactuca* Linnaeus, 1753:1163.
 Chiang, 1960, 1973b; Fan, 1953a; Heydrich, 1894; Okamura, 1930, 1931; Shen and Fan, 1950; Su et al., 1982; Yamada, 1950; Yoshikawa and Yoshikawa, 1977.
- Ulva lactuca* f. *lapathifolia* (Areschoug) Hauck, 1884:437.
 As "Phycoseris lapathifolia": Martens, 1866.
- Ulva pertusa* Kjellman, 1897b:4.
 Ariga, 1919, 1920; Chiang, 1960, 1973b; Fan, 1953a; Horikawa, 1919; Okamura, 1921; Rho, 1958; Shen and Fan, 1950; Taniguti, 1976; Tokida, 1939, 1954; Yamada, 1925a; Yoshikawa and Yoshikawa, 1977.
- Ulva reticulata* Forsskål, 1775:187.
 Arasaki, 1964; Ariga, 1920; Chiang, 1960, 1973b; Fan, 1953a; Okamura, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Tokida, 1939; Tseng, 1983; Yamada, 1925a.
- Ulva rigida* C. Agardh, 1823:410.
 Ariga, 1920; Yendo, 1916b.
- Ulva* sp.
 Chiang et al., 1974.
- CLADOPHORALES**
- CLADOPHORACEAE**
- Chaetomorpha aerea* (Dillwyn) Kützing, 1849:379.
 Ariga, 1921.
- Chaetomorpha aerea* f. *versata* Heydrich, 1894:273.
 De Toni, 1895; Heydrich, 1894; Okamura, 1936; Shen and Fan, 1950.
- Chaetomorpha basiretrorsa* Setchell, 1926:72.
 Chiang, 1960.
- Chaetomorpha brachygona* Harvey, 1858:87.
 Chiang, 1960; Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.
- Chaetomorpha crassa* (C. Agardh) Kützing, 1845:204.
 Ariga, 1920; Chiang, 1960, 1973b; Fan, 1953a; Okamura, 1931, 1936; Shen and Fan, 1950; Taniguti, 1976;
- Yamada, 1925a, 1950; Yoshikawa and Yoshikawa, 1977.
- Chaetomorpha linum* (O.F. Müller) Kützing, 1845:204.
 Ariga, 1919; Chiang, 1960; Fan, 1953a; Horikawa, 1919; Okamura, 1936; Shen and Fan, 1950; Tokida, 1954; Yamada, 1925a, 1926, 1950; Yendo, 1916a.
- Chaetomorpha spiralis* Okamura, 1912:162.
 Chiang, 1960, 1973b; Fan, 1953a.
- Cladophora aokii* Yamada, 1925a:85.
 Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.
- Cladophora fuliginosa* Kützing, 1849:415.
 Okamura, 1936; Tseng, 1983; Yamada, 1932b, 1934, 1950.
- Cladophora montagnei* var. *radicans* Yamada, 1925a:87.
 Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.
- Cladophora patentiramea* (Montagne) Kützing, 1849:416.
 Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925a, 1950.
- Cladophora pellucida* (Hudson) Kützing, 1845:208.
 Okamura, 1936; Shen and Fan, 1950. As "C. prolifera": Okamura, 1931.
- Cladophora rugulosa* Martens, 1866:112.
 Ariga, 1920.
- Cladophora scitula* (Suhr) Kützing, 1849:399.
 De Toni, 1895; Heydrich, 1894; Okamura, 1936; Shen and Fan, 1950.
- Cladophora sibogae* Reinbold, 1905:146.
 Okamura, 1931, 1936; Shen and Fan, 1950; Taniguti, 1976; Yamada, 1950.
- Cladophora* sp.
 Ariga, 1919, 1920; Fan, 1953a; Horikawa, 1919; Okamura, 1931; Shen and Fan, 1950; Tokida, 1939.

CAULERPALES

CODIACEAE

- Codium adhaerens* (Cabrera) C. Agardh, 1823:457.
 Ariga, 1919, 1920; Chiang, 1960, 1962b; Horikawa, 1919; Okamura, 1936.
- Codium arabicum* Kützing, 1856:35.
 Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925a.
- Codium contractum* Kjellman, 1897b:35.
 Ariga, 1920; Taniguti, 1976.
- Codium cylindricum* Holmes, 1896:250.
 Taniguti, 1976.
- Codium dichotomum* (Hudson) S.F. Gray, 1821:293.
 Fan, 1953a; Tokida, 1954. As "C. tomentosum": De Toni, 1895; Heydrich, 1894; Shen and Fan, 1950.
- Codium formosanum* Yamada, 1950:180.
 Fan, 1953a; Yamada, 1950; Yoshikawa and Yoshikawa, 1977.
- Codium fragile* (Suringar) Hariot, 1889:32.
 As "C. mucronatum": Ariga, 1919; Horikawa, 1919.

- Codium intricatum* Okamura, 1913:74.
 Ariga, 1919, 1920; Chiang, 1962b, 1973b; Fan, 1953a;
 Higashi, 1934; Horikawa, 1919; Shen and Fan, 1950;
 Yamada, 1950.
- Codium reediae* Silva in Egerod, 1952:389.
 Chiang, 1973a.
- Codium repens* (P. et H. Crouan) Vickers, 1905:56.
 Fan, 1953a; Okamura, 1931, 1936; Shen and Fan, 1950;
 Tseng, 1983; Yamada, 1925a.
- Codium tenue* Kützing, 1856:33.
 Fan, 1953a; Okamura, 1936; Rho, 1958; Shen and Fan,
 1950; Taniguti, 1976; Yamada, 1925a.

UDOTEACEAE

- Chlorodesmis caespitosa* J. Agardh, 1887:49.
 Chiang, 1973b; Chihara, 1975; Yoshikawa and Yoshi-
 kawa, 1977. As “*C. formosana*”: Arasaki, 1964; Chihara,
 1970; Okamura, 1930, 1936; Rho, 1958; Segawa, 1974;
 Shen and Fan, 1950; Yamada, 1925a.
- Remarks: Ducker et al. (1965) and Ducker (1967) pro-
 vide evidence for reducing *C. formosana* into synonymy
 with *C. caespitosa* (see also Fan, 1974).
- Chlorodesmis comosa* Bailey et Harvey in Harvey, 1858:29.
 Ariga, 1920; Higashi, 1934; Okamura, 1931, 1936; Rho,
 1958; Shen and Fan, 1950; Yamada, 1925a.
- Halimeda cuneata* Hering in Krauss, 1846:214.
 Higashi, 1934; Okamura, 1930, 1936; Shen and Fan,
 1950; Taniguti, 1976; Yamada, 1925a, 1950.
- Halimeda discoidea* Decaisne, 1842:102.
 Ariga, 1920.
- Halimeda incrassata* var. *ovata* (J. Agardh) Barton, 1901:27.
 Okamura, 1936; Yendo, 1909.
- Halimeda macroloba* Decaisne, 1841:118.
 Chiang, 1973a, 1973b; Okamura, 1915a, 1935b, 1936.
- Halimeda opuntia* (Linnaeus) Lamouroux, 1812:186.
 Ariga, 1920; Chiang, 1973b; Su et al., 1982; Taniguti,
 1976; Yamada, 1926.
- Halimeda renchii* Hauck, 1886:167.
 As “*H. opuntia* f. *renchii*”: Ariga, 1919, 1920; Chiang,
 1962b; Horikawa, 1919; Okamura, 1915a, 1931, 1936;
 Shen and Fan, 1950; Su et al., 1982; Yamada, 1925a,
 1950.
- Halimeda velasquezii* W.R. Taylor, 1962:176.
 As “*H. opuntia* f. *intermedia*” Chiang, 1962b.

CAULERPACEAE

- Caulerpa cupressoides* var. *lycopodium* f. *amicorum* (Harvey)
 Weber-van Bosse, 1898:337.
 Ariga, 1920.
- Caulerpa freycinetii* var. *freycinetii* f. *lata* Weber-van Bosse,
 1898:313.

- Ariga, 1920; Okamura, 1936; Shen and Fan, 1950; Yoshi-
 kawa and Yoshikawa, 1977. As “*C. freycinetii* var. *typica*”:
 Okamura, 1931.
- Caulerpa racemosa* (Forsskål) J. Agardh, 1873:35.
 Ariga, 1919; Fan, 1953a; Horikawa, 1919; Taniguti,
 1976; Tokida, 1939.
- Caulerpa racemosa* var. *clavifera* f. *macrophysa* (Kützing) We-
 ber-van Bosse, 1898:361.
 Ariga, 1920; Chiang, 1962b; Okamura, 1936; Rho,
 1958; Segawa, 1974; Yamada, 1925a, 1926, 1950. As
 “*C. racemosa* var. *clavifera*”: Okamura, 1913, 1931; Shen
 and Fan, 1950; Yamada, 1925a, 1926; Tseng, 1983.
- Caulerpa racemosa* var. *clavifera* f. *microphysa* (Kützing) We-
 ber-van Bosse, 1898:361.
 Okamura, 1931, 1936; Yamada, 1950.
- Caulerpa racemosa* var. *laetevirens* (Montagne) Weber-van
 Bosse, 1898:366.
 Chiang, 1960, 1973b; Okamura, 1930, 1931, 1936; Se-
 gawa, 1974; Shen and Fan, 1950; Taniguti, 1976; Ya-
 mada, 1925a, 1926, 1950; Yoshikawa and Yoshikawa,
 1977.
- Caulerpa racemosa* var. *occidentalis* (J. Agardh) Børgesen,
 1907:379.
 Yoshikawa and Yoshikawa, 1977.
- Caulerpa racemosa* var. *peltata* (Lamouroux) Eubank,
 1946:421.
 As “*C. peltata*”: Chiang, 1960, 1962b; Okamura, 1931,
 1932, 1936; Segawa, 1974; Shen and Fan, 1950; Tseng,
 1936; Yamada, 1950.
- Caulerpa racemosa* var. *turbinata* (J. Agardh) Eubank,
 1946:420.
 As “*C. racemosa* var. *chemnitizia*”: Chiang, 1960; Shen and
 Fan, 1950; Yamada, 1925a, 1926.
- Caulerpa serrulata* (Forsskål) C. Agardh, 1823:446.
 Eubank, 1946; Segawa, 1974; Tseng, 1936; Chihara,
 1975.
- Caulerpa serrulata* f. *lata* (Weber-van Bosse) Tseng
 1936:178.
 Tseng, 1983.
- Caulerpa sertularioides* (Gmelin) Howe, 1905:576.
 Ariga, 1919; Chihara, 1975; Horikawa, 1919; Yamada,
 1950.
- Caulerpa sertularioides* f. *longipes* (J. Agardh) Collins,
 1909:415.
 Ariga, 1920.
- Caulerpa taxifolia* (Vahl) C. Agardh, 1823:435.
 Yamada, 1926.
- Caulerpa webbiana* Montagne, 1838:18.
 Ariga, 1919; Horikawa, 1919.
- Caulerpa webbiana* f. *disticha* Weber-van Bosse, 1898:270.
 Okamura, 1931, 1936; Shen and Fan, 1950.
- Caulerpa webbiana* f. *tomentella* (Harvey) Weber-van Bosse,
 1898:270.
 Ariga, 1920; Chihara, 1975; Okamura, 1931, 1936; Se-
 gawa, 1974; Shen and Fan, 1950; Tseng, 1936.

BRYOPSIDACEAE

- Bryopsis harveyana* J. Agardh, 1887:22.
 Okamura, 1931, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925a, 1950.
- Bryopsis indica* A. et E.S. Gepp, 1908:169.
 Chiang, 1960, 1962b.
- Bryopsis mucosa* Lamouroux, 1809b:333.
 Chiang, 1960.
- Bryopsis plumosa* (Hudson) C. Agardh, 1823:448.
 Ariga, 1920; Chiang, 1960; De Toni, 1895; Heydrich, 1894; Okamura, 1936; Rho, 1958; Shen and Fan, 1950.

DERBESIACEAE

- Derbesia lamourouxii* (J. Agardh) Solier, 1847:162.
 Taniguti, 1976.

SIPHONOCLADALES**SIPHONOCLADACEAE**

- Boergesenia forbesii* (Harvey) Feldmann, 1938:584.
 Chiang, 1962b, 1973b; Tseng, 1983; Yamada, 1950. As "*Valonia forbesii*": Ariga, 1919; Horikawa, 1919; Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.

VALONIACEAE

- Boodlea coacta* (Dickie) Murray et De Toni in Murray, 1889:245.
 Yamada, 1926, 1950; Yoshikawa and Yoshikawa, 1977.
- Boodlea composita* (Harvey et J. Hooker) Brand, 1904:187.
 Chiang, 1960, 1973b; Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada 1925a, 1950.
- Boodlea montagnei* (Harvey ex J.E. Gray) Egerod, 1952:332.
 As "*B. paradoxa*": Chiang, 1960; Okamura, 1936; Shen and Fan, 1950.

- Boodlea siamensis* Reinbold, 1901:191.
 Chiang, 1962b; Okamura, 1931, 1936; Shen and Fan, 1950; Yamada, 1925a.

- Boodlea van-bosseae* Reinbold, 1905:148, 'van Bossei'.
 Okamura, 1919, 1931, 1936; Shen and Fan, 1950. As "*B. bosseae*": Ariga, 1919; Horikawa, 1919.
- Remarks: Following the examples given under Article 73.9 of the *International Code of Botanical Nomenclature* (Voss et al., 1983), "a hyphen is correctly used in an epithet after a word which could stand independently . . ." we spell the species name "van-bosseae."

- Chamaedoris orientalis* Okamura et Higashi in Okamura, 1931:98.

Okamura, 1931, 1932, 1936; Shen and Fan, 1950.

Cladophoropsis herpestica (Montagne) Howe, 1914:31.
 Shen and Fan, 1950.

Cladophoropsis sundanensis Reinbold, 1905:147.
 Chiang, 1960; Su et al., 1982.

Cladophoropsis zollingeri (Kützing) Reinbold, 1905:147.
 Chiang, 1960; Shen and Fan, 1950; Su et al., 1982; Taniguti, 1976; Tseng, 1983. As "*C. fasciculatus*": Okamura, 1930, 1931; Tseng, 1983; Yamada, 1925a.

Remarks: There has been some question as to when and where the combination *Cladophoropsis zollingeri* was validly published. Most authors (e.g., Papenfuss, 1950; Chiang, 1962a; Taniguti, 1976) have accepted *C. zollingeri* (Kützing) Børgesen (1905:288) as correct. According to Article 33.1 of the *International Code of Botanical Nomenclature* (Voss et al., 1983), "the combination is not validly published unless the author definitely indicates that the epithet or epithets concerned are to be used in that particular combination." Parallel to the examples, such as *Eulophus*, cited in the Code, Børgesen (1905) has merely stated that "*S. zollingeri*" belongs here but did not actually make the combination in *Cladophoropsis*. The earliest valid combination we are aware of is: *C. zollingeri* (Kützing) Reinbold 1905:147.

Dictyosphaeria bokotensis Yamada, 1925a:81.
 Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a, 1950.

Dictyosphaeria cavernosa (Forsskål) Børgesen, 1932:2.
 Chiang, 1960, 1973b; Shen and Fan, 1950; Taniguti, 1976; Tseng, 1983. As "*D. favulosa*": Ariga, 1920; Okamura, 1930, 1931, 1936; Yamada, 1925a, 1926.

Microdictyon japonicum Setchell, 1925:107.
 Chiang, 1962b; Tseng, 1983; Shen and Fan, 1950. As "*Rhipidiphyllum reticulatum*": Ariga, 1920; De Toni, 1895; Heydrich, 1894.

Remarks: Fan (1974:249–250, 253–254) reviews taxonomic problems with the genus *Rhipidiphyllum* Heydrich, 1894:281. The type-locality for this genus is in east Taiwan.

Microdictyon nigrescens (Yamada) Setchell, 1925:107.
 Okamura, 1936; Shen and Fan, 1950; Fan, 1974; Yamada, 1950. As "*Rhipidiphyllum nigrescens*": Yamada, 1925a.

Microdictyon okamurae Setchell, 1925:107, 'okamurae'.
 Okamura, 1931; Shen and Fan, 1950; Tseng, 1983.

Remarks: The correct spelling for the Latinization of Professor Okamura's name is "okamurae" and has been corrected throughout this study.

Struvea anastomosans (Harvey) Piccone et Grunow ex Piccone, 1884:20.

Ariga, 1920; Tseng, 1983.

Struvea delicatula Kützing, 1866:1.

Chiang, 1960; De Toni, 1895; Heydrich, 1894; Oka-

- mura, 1930, 1931, 1936; Shen and Fan, 1950; Yamada, 1925b, 1926.
- Valonia aegagropila* C. Agardh, 1823:429.
Chiang, 1973b; Chihara, 1975; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925a, 1950; Yoshikawa and Yoshikawa, 1977.
- Valonia fastigiata* Harvey in J. Agardh, 1887:101.
Okamura, 1931; Shen and Fan, 1950.
- Valonia utricularis* (Roth) C. Agardh, 1823:431.
Okamura, 1931, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1926; Yendo, 1914.
- Valonia verticillata* Kützing, 1849:508.
Okamura, 1930.
- Valoniopsis pachynema* (Martens) Børgesen, 1934:10.
Chiang, 1960, 1973b; Shen and Fan, 1950; Tseng, 1983; Yamada, 1950; Yoshikawa and Yoshikawa, 1977. As "*Valonia confervoides*": Ariga, 1920; Okamura, 1909, 1931, 1936; Yamada, 1925a.

ANADYOMENACEAE

- Anadyomene wrightii* Harvey in Gray, 1866:48.
Ariga, 1920; Chiang, 1973b; Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925a, 1926, 1950.

DASYCLADALES**ACETABULARIACEAE**

- Acetabularia gigas* Solms-Laubach, 1895:23.
Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.
- Acetabularia major* Martens, 1866:25.
Okamura, 1936; Shen and Fan, 1950; Yamada, 1925a.
- Bornetella sphaerica* (Zanardini) Solms-Laubach, 1893:92.
Chiang, 1973b.
- Neomeris annulata* Dickie, 1874:198.
Chiang, 1973b.

PHAEOPHYTA**ECTOCARPALES****ECTOCARPACEAE**

- Ectocarpus breviarticulatus* J. Agardh, 1847:7.
Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925b, 1950.
- Ectocarpus laurenciae* Yamada, 1931a:66.
Okamura, 1936; Shen and Fan, 1950; Yamada, 1931a.
- Ectocarpus spinosus* Kützing, 1843:288.
Heydrich, 1894; Shen and Fan, 1950.
- Ectocarpus van-bossea* Setchell et Gardner in Setchell, 1924:170.
Yamada, 1950.

Ectocarpus sp.

Chiang, 1976.

Feldmannia formosana (Yamada) Itono, 1973:162.
As "*Ectocarpus formosanus*": Yamada, 1950.*Giffordia mitchelliae* (Harvey) Hamel, 1939:xiv.As "*Ectocarpus mitchelliae*": Chiang, 1960; Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925b.**CHORDARIALES****CORYNOTHЛАEACEAE**

- Leathesia difformis* (Linnaeus) Areschoug, 1847:376.
Taniguti, 1976.

SPHACELARIALES**SPHACELARIACEAE**

- Sphacelaria furcigera* var. *tenuis* Yamada, 1941:196.
Shen and Fan, 1950; Yamada, 1941.
- Sphacelaria tribuloides* Meneghini, 1840:2.
Chihara, 1975; De Toni, 1895; Heydrich, 1894; Okamura, 1897, 1936; Segawa, 1974; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925b.

DICTYOTALES**DICTYOTACEAE**

- Dictyopteris delicatula* Lamouroux, 1809b:332.
Yamada, 1950.
- Dictyopteris repens* (Okamura) Børgesen, 1920a:265.
Tseng, 1983; Yoshikawa and Yoshikawa, 1977. As "*Neurocarpus repens*": Okamura, 1931, 1936; Shen and Fan, 1950.
- Dictyopteris undulata* Holmes, 1896:251.
Tseng, 1983. As "*Haliseris undulata*": Okamura, 1907.
- As "*Neurocarpus undulata*": Higashi, 1934; Okamura, 1930, 1936; Shen and Fan, 1950. As "*N. undulata* f. *plana*": Chiang, 1960; Okamura, 1930, 1936.
- Dictyota bartayresii* Lamouroux, 1809b:331.
Okamura, 1931, 1936; Shen and Fan, 1950.

- Dictyota ceylanica* var. *anastomosans* Yamada, 1950:186.
Yamada, 1950; Yoshikawa and Yoshikawa, 1977.
- Dictyota dichotoma* (Hudson) Lamouroux, 1809a:42.
Ariga, 1919; Higashi, 1934; Horikawa, 1919; Okamura, 1930; Rho, 1958; Shen and Fan, 1950; Tokida, 1954; Yamada, 1925b.
- Dictyota dilata* Yamada, 1925b:252.
Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Yamada, 1925b.
- Dictyota divaricata* Lamouroux, 1809a:43.
Chiang, 1960; Okamura, 1930; Shen and Fan, 1950; Yamada, 1925b.
- Dictyota hamifera* Setchell, 1926:92.
Yamada, 1950.
- Dictyota linearis* (C. Agardh) Greville, 1830:xlili.
Ariga, 1919, 1920; Horikawa, 1919.
- Dictyota patens* J. Agardh, 1882:93.
Taniguti, 1976.
Remarks: Howe (1920) and Allender and Kraft (1983) consider this taxon to be a synonym of *D. bartayresii* Lamouroux.
- Dictyota spinulosa* Hooker et Arnott, 1838:275.
Ariga, 1920.
- Dilophus okamurae* Dawson, 1950:83, 'okamurae'.
Chihara, 1970, 1975; Fan, 1953b; Segawa, 1974; Tseng, 1983. As "*D. marginatus*": Chiang, 1960; Okamura, 1936; Shen and Fan, 1950.
- Dilophus radicans* Okamura, 1916:7.
Chiang, 1960; Okamura, 1936; Shen and Fan, 1950.
Remarks: Fan (1953b) considers this name to be in need of synonymy since he believes *D. radicans* may be the prostrate form of *Padina commersonii* (see also Lewis et al., 1987).
- Lobophora variegata* (Lamouroux) Womersley, 1967:221.
Tseng, 1983. As "*Gymnosorus collaris*": Ariga, 1920. As "*Zonaria variegata*": Heydrich, 1894.
- Pachydictyon coriaceum* (Holmes) Okamura, 1899:39.
Ariga, 1920; Chiang, 1976; Chihara, 1970; Okamura, 1936; Rho, 1958; Shen and Fan, 1950.
- Padina arborescens* Holmes, 1896:251.
Higashi, 1934.
- Padina australis* Hauck, 1887:44.
Chiang, 1960, 1962b; Okamura, 1931, 1932, 1936; Rho, 1958; Shen and Fan, 1950; Su et al., 1982; Tseng, 1936, 1983; Yamada, 1925b, 1931a, 1950.
- Padina boryana* Thivy in Taylor, 1966:355.
Tseng, 1983.
Remarks: Papenfuss (1977) considered both *P. tenuis* (C. Agardh) Bory and *P. commersonii* Bory to be synonyms of *Lobophora variegata* (Lamouroux) Womersley, and recognized *P. boryana* Thivy as the name for this taxon. Tseng (1983) apparently follows this, listing *P. commersonii* as a synonym of *P. boryana*, whereas Womersley and Bailey (1970) and Allender and Kraft (1983) considered *P. commersonii* and *P. boryana* to be synonyms of *P. tenuis*.
- Padina crassa* Yamada, 1931a:67.
Ariga, 1920; Chiang, 1960; Yoshikawa and Yoshikawa, 1977.
- Padina durvillaei* Bory, 1827:591.
Ariga, 1920; Chiang, 1973a; De Toni, 1895; Heydrich, 1894; Okamura, 1935b; Shen and Fan, 1950.
- Padina japonica* Yamada, 1931a:69.
As "*P. pavonia*": Ariga, 1919; De Toni, 1895; Heydrich, 1894; Horikawa, 1919.
- Padina minor* Yamada, 1925b:251.
Chiang, 1960, 1962a, 1962b, 1973b, 1976; Chihara, 1975; Okamura, 1931, 1932, 1936; Rho, 1958; Shen and Fan, 1950; Taniguti, 1976; Tseng, 1983; Yamada, 1925b, 1931a, 1950; Yoshikawa and Yoshikawa, 1977.
- Padina tenuis* Bory, 1827:590.
As "*Padina commersonii*": Okamura, 1931, 1936; Shen and Fan, 1950; Yamada, 1925b, 1931a.
Remarks: See "Remarks" herein under *Padina boryana* Thivy in Taylor.
- Spatoglossum pacificum* Yendo, 1920:2.
Chiang, 1973a.
- Zonaria coriacea* Yamada, 1925b:249.
Chiang, 1960; Okamura, 1936; Shen and Fan, 1950; Yamada, 1925b.
- Zonaria diesingiana* J. Agardh, 1841:443.
Ariga, 1920; Chiang, 1960; Okamura, 1930; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925b.
- Zonaria flabellata* (Okamura) Papenfuss, 1944:341.
As "*Homoeostrichus flabellatus*": Taniguti, 1976.
- Zonaria harveyana* (Kützing) Areschoug, 1851:26.
As "*Homoeostrichus multifidus*": Chiang, 1960; Okamura, 1936; Shen and Fan, 1950; Yamada, 1925b.
- Zonaria nigrescens* Sonder, 1845:50.
Heydrich, 1894; Shen and Fan, 1950.
- Zonaria stipitata* Tanaka et Nozawa, 1962:183.
Chiang and Chou, 1980.
- Zonaria* sp.
Su et al., 1982.

SCYTOSIPHONALES

ISHIGEACEAE

- Ishige okamurae* Yendo, 1907:154, 'okamurae'.
Chiang, 1960, 1973b; Higashi, 1934; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Yamada, 1925b.
- Ishige sinicola* (Setchell et Gardner) Chihara, 1969:3.
As "*I. foliacea*": Rho, 1958; Tseng, 1936.

PUNCTARIACEAE

- Petalonia fascia* (O.F. Müller) Kuntze, 1898:419.
As "*Ilea fascia*": Chiang, 1960, 1976; De Toni, 1895; Rho, 1958; Shen and Fan, 1950; Tokida, 1954; Yamada, 1925b. As "*Phyllitis fascia*": Heydrich, 1894.

SCYTOSIPHONACEAE

- Chnoospora implexa* (Hering) J. Agardh, 1848:172.
 Chiang, 1960; Okamura, 1931, 1936; Shen and Fan, 1950; Taniguti, 1976; Tokida, 1939; Tseng, 1983; Yamada, 1925b, 1950.
- Chnoospora minima* (Hering) Papenfuss, 1956:69.
 Chiang, 1976; Segawa, 1974. As "*Chnoospora pacifica*": Okamura, 1931, 1936; Shen and Fan, 1950; Tokida, 1939; Yamada, 1950.
- Colpomenia sinuosa* (Mertens et Roth) Derbès et Solier in Castagne, 1851:95.
 Ariga, 1919, 1920; Chiang, 1960, 1976; De Toni, 1895; Heydrich, 1894; Horikawa, 1919; Okamura, 1930; Shen and Fan, 1950; Taniguti, 1976; Tokida, 1954; Yamada, 1925b, 1950; Yoshikawa and Yoshikawa, 1977.
- Endarachne binghamiae* J. Agardh, 1896:27.
 Ariga, 1919, 1920; Chiang, 1960, 1973b, 1976; Cotton, 1915; Fan, 1953a; Higashi, 1934; Horikawa, 1919; Okamura, 1930, 1936; Rho, 1958; Shen and Fan, 1950; Taniguti, 1976; Tilden, 1929; Tseng, 1983; Yamada, 1925b.
- Hydroclathrus clathratus* (Bory) Howe, 1920:590.
 Chiang, 1960, 1962a, 1973b; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Taniguti, 1976; Tseng, 1983; Yamada, 1950; Yoshikawa and Yoshikawa, 1977. As "*H. cancellatus*": De Toni, 1895; Heydrich, 1894; Higashi, 1934; Okamura, 1897, 1931; Yamada, 1925b.
- Rosenvingea orientalis* (J. Agardh) Børgesen, 1914:26.
 Okamura, 1936; Shen and Fan, 1950; Tseng, 1983. As "*Hydroclathrus orientalis*": De Toni, 1895; Heydrich, 1894.
- Scytoniphon lomentaria* (Lyngbye) J. Agardh, 1848:126, 'lomentarium'.
 Chiang, 1973a, 1976; Okamura, 1935b, 1936; Segawa, 1974; Tokida, 1954; Yamada, 1925b, 1928. As "*S. lomentarius*": Chihara, 1970, 1975.

FUCALES

CYSTOSEIRACEAE

- Cystoseira prolifera* J. Agardh, 1848:215.
 Chiang, 1976.
- Cystoseira* sp.
 Nizamuddin, 1970.
- Hormophysa triquetra* (Linnaeus) Kützing, 1843:359.
 Chiang, 1973a; Okamura, 1935b; Su et al., 1982.

SARGASSACEAE

- Sargassum amabile* Yamada, 1944a:1.
 Shen and Fan, 1950.
- Sargassum aquifolium* (Turner) C. Agardh, 1820:12.
 Okamura, 1931, 1936; Shen and Fan, 1950.
- Sargassum baccularia* (Mertens) C. Agardh, 1824:304.

- Chou and Chiang, 1981.
- Sargassum berberifolium* J. Agardh, 1848:337.
 Okamura, 1936; Shen and Fan, 1950; Yamada, 1925b.
- Sargassum binderi* Sonder in J. Agardh, 1848:328.
 Chiang, 1976; Chou and Chiang, 1981; Okamura, 1936; Shen and Fan, 1950; Yamada, 1925b.
- Sargassum carpophyllum* J. Agardh, 1848:304.
 Chou and Chiang, 1981; Yamada, 1942.
- Sargassum coriifolium* J. Agardh, 1889:86.
 Okamura, 1936; Shen and Fan, 1950; Yamada, 1925b.
- Sargassum coriifolium* f. *duplicatum* Yamada, 1950:192.
 Yamada, 1950; Yoshikawa and Yoshikawa, 1977.
- Sargassum coriifolium* f. *prolongatum* (Okamura) Yamada, 1950:192.
 Yamada, 1950. As "*S. prolongatum*": Okamura, 1931, 1932, 1936; Shen and Fan, 1950.
- Sargassum crassifolium* J. Agardh, 1848:326.
 Chihara, 1975; Chou and Chiang, 1981; Tseng, 1983; Yamada, 1942, 1950; Yoshikawa and Yoshikawa, 1977.
- Sargassum crispifolium* Yamada, 1931a:72.
 Chiang, 1960, 1962b.
- Sargassum cristaefolium* C. Agardh, 1820:13.
 Ariga, 1920; Chiang, 1976; Chou and Chiang, 1981; Cotton, 1915; Okamura, 1931, 1935b, 1936; Shen and Fan, 1950; Tilden, 1929. As "*S. biserrula*": De Toni, 1895; Heydrich, 1894.
- Sargassum duplicatum* J. Agardh, 1889:90.
 Ariga, 1921; Chiang, 1976; Chou and Chiang, 1981; Liu, 1982; Okamura, 1936; Shen and Fan, 1950; Taniguti, 1976; Yamada, 1925b.
- Sargassum echinocarpum* J. Agardh, 1848:327.
 Yamada, 1950.
- Sargassum fulvellum* (Turner) C. Agardh, 1820:34.
 Ariga, 1920.
- Sargassum glaucescens* J. Agardh, 1848:306.
 Chiang, 1973a; Chou and Chiang, 1981; Okamura, 1935b.
- Sargassum hemiphyllum* (Turner) C. Agardh, 1820:39.
 Ariga, 1920; Chiang, 1976; Chou and Chiang, 1981; Tilden, 1929; Tseng et al., 1985.
- Sargassum henslowianum* C. Agardh in J. Agardh, 1848:315.
 Chou and Chiang, 1981.
- Sargassum heterocystum* Montagne, 1842:250.
 Chou and Chiang, 1981.
- Sargassum horneri* (Turner) C. Agardh, 1820:38.
 Ariga, 1919; Chiang, 1960; Chou and Chiang, 1981; Horikawa, 1919; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Tokida, 1954; Tseng et al., 1962, 1985; Yamada, 1925b.
- Sargassum ilicifolium* (Turner) C. Agardh, 1820:11.
 Ariga, 1920; Chou and Chiang, 1981; Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1925b.
- Sargassum ilicifolium* var. *conduplicatum* Grunow in Weber-van Bosse, 1913a:160.
 Yamada, 1942.

- Sargassum kasyotense* Yamada, 1942:553.
Shen and Fan, 1950; Yamada, 1942.
- Sargassum kuetzingii* Setchell, 1931:249.
Chou and Chiang, 1981.
- Sargassum kushimotense* Yendo, 1905:157.
Chou and Chiang, 1981.
- Sargassum muriocystum* J. Agardh, 1848:314.
Ariga, 1920; De Toni, 1895; Okamura, 1895, 1936;
Shen and Fan, 1950.
- Sargassum plagiophyllum* C. Agardh, 1824:309.
Yamada, 1942.
- Sargassum polycystum* C. Agardh, 1824:304.
Ariga, 1920; Chou and Chiang, 1981; Okamura, 1931,
1936; Shen and Fan, 1950; Su et al., 1982; Tseng, 1983;
Yamada, 1942, 1950; Yoshikawa and Yoshikawa, 1977.
As "S. microphyllum": Chiang, 1976; Yamada, 1925b.
- Sargassum rostratum* J. Agardh, 1896:55.
Chou and Chiang, 1981; Okamura, 1936; Shen and Fan,
1950; Tseng et al., 1985.
- Sargassum sandei* Reinbold in Weber-van Bosse, 1913a:158.
Chiang, 1960, 1976; Chou and Chiang, 1981; Okamura,
1936; Segawa, 1974; Yamada, 1925b; Yoshikawa and
Yoshikawa, 1977.
- Sargassum sandei* f. *heterophyllum* Yamada, 1950:192.
Yamada, 1950.
- Sargassum serratifolium* C. Agardh, 1820:16.
Chiang, 1973a.
- Sargassum siliquosum* J. Agardh, 1848:316.
Chiang, 1976; Chou and Chiang, 1981; Okamura, 1931,
1936; Rho, 1958; Shen and Fan, 1950; Yamada, 1925b,
1942.
- Sargassum swartzii* (Turner) C. Agardh, 1820:11.
Yamada, 1942.
- Sargassum telephifolium* (Turner) C. Agardh in J. Agardh,
1889:107.
Okamura, 1931, 1936; Shen and Fan, 1950.
- Sargassum tenuifolium* Yamada, 1942:505.
Yamada, 1942.
- Sargassum vulgare* C. Agardh, 1820:3.
Ariga, 1919; Horikawa, 1919.
- Sargassum* sp.
Chiang, 1962b, 1976; Chiang et al., 1974; Chou and
Chiang, 1981; Su et al., 1982.
- Turbinaria conoides* (J. Agardh) Kützing, 1860:24.
Chiang, 1973b; Okamura, 1931, 1936; Shen and Fan,
1950; Tseng, 1983.
- Turbinaria filamentosa* Yamada, 1925b:243.
Chiang, 1960, 1976; Fan, 1974; Shen and Fan, 1950;
Yamada, 1925b. As "T. filiformis": Okamura, 1936; Yosh-
ikawa and Yoshikawa, 1977.
Remarks: Fan (1974:252, 254) suggests that the char-
acteristic of this species (i.e., elongate filiform receptacle)
may not reflect a species difference, but rather reflect an
ecological or physiological state.
- Turbinaria ornata* (Turner) J. Agardh, 1848:266.
Chiang, 1960, 1962b, 1976; De Toni, 1895; Okamura,
1895, 1931, 1936; Shen and Fan, 1950; Su et al., 1982;
Yamada, 1925b, 1950.
- Turbinaria trialata* (J. Agardh) Kützing, 1860:24.
Chiang, 1962b; Yamada, 1950.
- Turbinaria* sp.
Ariga, 1919; Horikawa, 1919.

RHODOPHYTA

GONIOTRICHALES

GONIOTRICHACEAE

- Chroodactylon ornatum* (C. Agardh) Basson, 1979:67.
As "Asteroctis ornata": Shen and Fan, 1950; Tanaka,
1944, 1952. As "Asteroctis ornata" f. *simplex*: Shen and
Fan, 1950; Tanaka, 1944, 1952. As "Asteroctis ramosa" f.
simplex: Tanaka, 1944.
- Stylonema alsidii* (Zanardini) Drew, 1956:72.
As "Goniotrichum alsidii": Tanaka, 1952.
Remarks: See Wynne (1985) for discussion of use of the
generic name *Stylonema* Reinsche, and Drew and Ross
(1965) on use of some generic names in the Bangiophy-
cidae.

BANGIALES

ERYTHROPELTIDACEAE

- Erythrotrichia biseriata* Tanaka, 1944:86.
Shen and Fan, 1950; Tanaka, 1944, 1951, 1952.

BANGIACEAE

- Bangia atropurpurea* (Roth) C. Agardh, 1824:76.
Ariga, 1920. As "B. fuscopurpurea": Chiang, 1962a,
1973b; Fan, 1953a; Okamura, 1936; Rho, 1958; Sheath
and Cole, 1984; Shen and Fan, 1950; Tanaka, 1950,
1952; Taniguti, 1976; Tokida, 1954.
- Bangia yamadae* Tanaka, 1944:84, 'yamadai'.
Shen and Fan, 1950; Tanaka, 1944, 1950, 1952.
- Bangia* sp.
Chiang, 1973b.
- Porphyra angusta* Okamura et Ueda in Ueda, 1932:28.
Liaw and Chiang, 1979; Wang and Chiang, 1977.
- Porphyra crispata* Kjellman, 1897a:15.
Ariga, 1920; Chiang, 1962a; Chihara, 1975; Dawson,
1954c; Fan, 1953a; Okamura, 1931, 1936; Shen and
Fan, 1950; Tanaka, 1952; Taniguti, 1976; Wang and
Chiang, 1977; Ueda, 1932.

- Porphyra dentata* Kjellman, 1897a:13.
Wang and Chiang, 1977.
Porphyra suborbiculata Kjellman, 1897a:10.
Okamura, 1930, 1936; Wang and Chiang, 1977.
Porphyra sp.
Wang and Chiang, 1977.

NEMALIALES

Remarks: For discussion on the correct spelling of the name of this order see Nicolson and Norris, 1983.

ACROCHAETIACEAE

- Liagorophila endophytica* Yamada, 1944b:16.
Fan, 1974; Shen and Fan, 1950; Yamada, 1944b. As "*Liagora orientalis*": Yamada, 1938a.

DERMONEMATACEAE

- Dermonema frappieri* (Montagne et Millardet) Børgesen, 1942:42.
Chiang, 1969, 1971, 1973b; Chihara, 1975; Segawa, 1974; Tseng, 1983.

- Dermonema gracile* Martens ex Weber-van Bosse, 1921:204.
Fan, 1953a; Okamura, 1931, 1936; Shen and Fan, 1950; Tokida, 1939; Tseng, 1945. As "*D. dichotomum*": Ariga, 1920; Chen and Chiang, 1982; De Toni, 1895; Fan, 1974; Heydrich, 1894.

- Dermonema pulvinata* (Grunow in Holmes) Fan, 1962:337.
Chen and Chiang, 1982; Chihara, 1975; Fan, 1953a; Tseng, 1983; Umezaki, 1972. As "*Nemalion pulvinatum*": Chiang, 1962a; Chihara, 1970; Okamura, 1931, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Tokida, 1939.

- Remarks: Papenfuss (1967:96–97) points out that *Dermonema* Heydrich had been a nomen nudum until Heydrich (1894:289) published *D. dichotomum*. The type-locality for the genus is therefore Taiwan (see also Fan 1974:249, 253).

- Dotyophycus yamadae* (Ohmi et Itono) Abbott et Yoshizaki, 1981:225.

- Chiang and Chen, 1983.
Liagoropsis schrammi (P. et H. Crouan in Mazè et Schramm) Doty and Abbott, 1964:443.

- Fan, 1974. As "*L. maxima*": Shen and Fan, 1950; Yamada, 1944c.

- Yamadaella cenomyce* (Decaisne) Abbott, 1970:117.
Chiang, 1973b. As "*L. annulata*": Okamura, 1931, 1936.
As "*L. caenomyce*": Chihara, 1975; Segawa, 1974; Shen and Fan, 1950; Tseng, 1941a; Yamada, 1938a.

LIAGORACEAE

- Helminthocladia australis* Harvey, 1863, pl. 272.
Chen et al., 1981.

- Liagora boergesenii* Yamada, 1938a:11.
Chiang, 1971, 1972; Chiang and Chen, 1982; Chihara, 1975; Shen and Fan, 1950; Yamada, 1938a.
Liagora ceranoides Lamouroux, 1816:239.
Chiang and Chen, 1982.
Liagora ceranoides var. *leprosa* (J. Agardh) Yamada, 1938a:21.
Segawa, 1974; Shen and Fan, 1950; Yamada, 1938a.
Liagora ceranoides var. *pulverulenta* (C. Agardh) Yamada, 1938a:21.
Shen and Fan, 1950; Yamada, 1938a.
Liagora decussata Montagne, 1849:64.
Chiang and Chen, 1982; Shen and Fan, 1950; Yamada, 1938a.
Liagora farinosa Lamouroux, 1816:240.
Chihara, 1975; Shen and Fan, 1950; Tseng, 1941a; Yamada, 1938a.
Remarks: Recently Fan and Wang (1974:492) proposed a new generic name, *Ganonema*, for this taxon. However, we follow the opinion of Abbott (1976:130), and for the time being continue to recognize this taxon as a *Liagora* (see also Norris and Bucher, 1982:190).
Liagora orientalis J. Agardh, 1896:99.
Ariga, 1920; Chiang and Chen, 1982; Fan, 1974; Okamura, 1936; Shen and Fan, 1950; Yamada, 1932b. As "*L. formosana*": Yamada, 1938a.
Remarks: Yamada (1944b:18–19) reduced his species *L. formosana* (1938a:32) into synonymy with *L. orientalis*.
Liagora rugosa Zanardini, 1851:36.
Ariga, 1920.
Liagora segawai Yamada, 1938a:1.
Chiang and Chen, 1982.
Liagora setchellii Yamada, 1938a:13.
Chiang and Chen, 1982; Shen and Fan, 1950; Tseng, 1941a; Yamada, 1938a; Yoshikawa and Yoshikawa, 1977.
Liagora valida Harvey, 1853:138.
Chiang and Chen, 1982; Okamura, 1931, 1936; Shen and Fan, 1950.

CHAETANGIACEAE

- Actinotrichia fragilis* (Forsskål) Børgesen, 1932:6.
Chihara, 1970, 1975; Shen and Fan, 1950; Tseng, 1941b. As "*Actinotrichia rigida*": Ariga, 1920.
Galaxaura arborea Kjellman, 1900:72.
Ariga, 1920; Chiang, 1962a; Yendo, 1918; Yoshikawa and Yoshikawa, 1977.
Remarks: The taxonomy and nomenclature of this genus have been confused by morphological plasticity and uncertain life histories. Since the hypothesis of an alternation of morphologically different sexual and tetrasporic forms of *Galaxaura* (Howe 1917, 1918), there has been much speculation as to which tetrasporic and sexual "species" represent stages within the same life history (e.g., Børgesen, 1920b; Chou, 1945, 1947; Dawson, 1953;

Taylor, 1960). We hold the opinion that until culture studies conclusively link the different reproductive stages into the life history, each reproductive form should be recognized by its previous species name.

Galaxaura canaliculata Kützing, 1849:530.

Heydrich, 1894; Shen and Fan, 1950.

Galaxaura clavigera Kjellman, 1900:76.

Chiang, 1962b; Itono, 1977a; Okamura, 1931, 1936; Shen and Fan, 1950; Tanaka, 1936.

Remarks: Papenfuss, Mshigeni, and Chiang (1982:411) consider this species to be a synonym of *Galaxaura marginata* (Ellis et Solander) Lamouroux.

Galaxaura distenta Harvey, 1859:331.

Ariga, 1920.

Galaxaura elegans Tanaka, 1935:52.

Okamura, 1936; Shen and Fan, 1950; Tanaka, 1935, 1936.

Galaxaura elongata J. Agardh, 1876:529.

Itono, 1977a; Okamura, 1936; Shen and Fan, 1950; Tanaka, 1936; Tseng, 1941b; Yendo, 1916b; Yoshikawa and Yoshikawa, 1977.

Remarks: Considered a synonym of *G. rugosa* (Ellis et Solander) Lamouroux (1816:263) by Papenfuss, Mshigeni, and Chiang (1982).

Galaxaura falcata Kjellman, 1900:73.

Ariga, 1919; Horikawa, 1919; Taniguti, 1976.

Galaxaura fasciculata Kjellman, 1900:53.

Itono, 1977a; Okamura, 1936; Shen and Fan, 1950; Tanaka, 1936; Tseng, 1941b.

Galaxaura fastigiata Decaisne 1842:116.

Ariga, 1920; Chiang, 1962a, 1962b; Chihara, 1970, 1975; Itono, 1977a; Okamura, 1931, 1936; Segawa, 1974; Shen and Fan, 1950; Tanaka, 1936; Tanaguti, 1976; Yoshikawa and Yoshikawa, 1977.

Remarks: Considered a synonym of *G. oblongata* (Ellis et Solander) Lamouroux (1816: 262) by Papenfuss, Mshigeni, and Chiang (1982).

Galaxaura filamentosa Chou in Taylor, 1945:139.

Itono, 1977a.

Galaxaura lapidescens (Ellis et Solander) Lamouroux, 1816:264.

De Toni, 1895.

Galaxaura lapidescens f. *villosa* J. Agardh, 1876:530.

Heydrich, 1894; Okamura, 1897; Shen and Fan, 1950.

Galaxaura latifolia Tanaka, 1935:54.

Okamura, 1935a, 1936; Shen and Fan, 1950; Tanaka, 1935, 1936.

Galaxaura oblongata (Ellis et Solander) Lamouroux, 1816:262.

Chiang, 1973a, 1973b; Okamura, 1935b; Su et al., 1982.

Galaxaura obtusata (Ellis et Solander) Lamouroux, 1816:262.

Itono, 1977a; Okamura, 1936; Papenfuss, Mshigeni, and

Chiang, 1982; Shen and Fan, 1950; Su et al., 1982; Tanaka, 1936; Tseng, 1941b.

Galaxaura pacifica Tanaka, 1935:55.

Itono, 1977a; Okamura, 1935a, 1936; Shen and Fan, 1950; Tanaka, 1935, 1936.

Remarks: Considered a synonym of *G. rugosa* (Ellis et Solander) Lamouroux (1816:262) by Papenfuss, Mshigeni, and Chiang (1982).

Galaxaura robusta Kjellman, 1900:85.

Chihara, 1970, 1975; Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Tanaka, 1936.

Remarks: Considered a synonym of *G. obtusata* (Ellis et Solander) Lamouroux by Papenfuss, Mshigeni, and Chiang (1982).

Galaxaura rudis Kjellman, 1900:43.

Okamura, 1931, 1936; Shen and Fan, 1950; Tanaka, 1936; Tseng, 1941b.

Remarks: Considered a synonym of *G. lapidescens* (Ellis et Solander) Lamouroux by Papenfuss, Mshigeni, and Chiang (1982).

Galaxaura marginata (Ellis et Solander) Lamouroux, 1816:264.

Papenfuss, Mshigeni, and Chiang, 1982. As "*Galaxaura tenera*": Chiang, 1973a, 1973b; Okamura, 1935a; Su et al., 1982.

Remarks: Material from Taiwan determined as *G. tenera* was re-identified by Papenfuss, Mshigeni, and Chiang (1982) to be this species.

Galaxaura veprecula Kjellman, 1900:80.

Ariga, 1920; Chihara, 1975; Itono, 1977a; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Tanaka, 1936; Tseng, 1983; Yendo, 1918.

Remarks: Papenfuss, Mshigeni, and Chiang (1982) consider this species to be a synonym of *G. marginata* (Ellis et Solander) Lamouroux.

Scinaia boergesenii Tseng, 1941:100.

Shen and Fan, 1950.

Scinaia cottonii Setchell, 1914:103.

Shen and Fan, 1950.

Scinaia moniliformis J. Agardh, 1885:72.

Chiang, 1962a; Yoshikawa and Yoshikawa, 1977.

Scinaia pseudojaponica Yamada et Tanaka in Yamada, 1938b:127.

Chiang, 1962a, 1970, 1973b; Shen and Fan, 1950.

BONNEMAISONIACEAE

Asparagopsis taxiformis (Delile) Trevisan, 1845:45.

As "*A. sandfordiana*": Ariga, 1919; Fan, 1953a; Horikawa, 1919; Okamura, 1931, 1936; Shen and Fan, 1950.

Delisea fimbriata (Lamouroux) Montagne, 1844:155.

Chihara, 1970, 1975. As "*D. japonica*": Cotton, 1915; Tilden, 1929.

GELIDIALES**GELIDIACEAE**

Gelidium amansii Lamouroux in Kützing 1868:16.

Cotton, 1915; Fan, 1953a; Okamura, 1913, 1915b, 1930; Oshima, 1915; Rho, 1958; Santelices and Stewart, 1985; Tilden, 1929; Tokida, 1954.

Gelidium amansii f. *elegans* Okamura, 1934a:56.

Chiang, 1962a; Fan, 1951.

Gelidium amansii f. *latioris* Okamura, 1935b:443.

Chiang, 1962a; Fan, 1951; Okamura, 1935b, 1936; Shen and Fan, 1950.

Gelidium "cartilagineum".

Ariga, 1919; Horikawa, 1919.

Remarks: As with all taxa in this paper, we are reporting the name used by the author in recording the taxon from Taiwan. We recognize that Dixon (1967) has shown the basionym *Fucus cartilagineus* Linnaeus (1753) to be correctly *Plocamium cartilagineum* (Linnaeus) Dixon. However, in this case, we suggest the material of Ariga (1919) and Horikawa (1919) probably belongs to *Gelidium*, rather than *Plocamium*, but cannot be sure what species it is.

Gelidium crinale (Turner) Lamouroux in Bory, 1825:191.

Ariga, 1920.

Gelidium divaricatum Martens, 1866:30.

Okamura, 1934a, 1936; Taniguti, 1976.

Gelidium japonicum (Harvey) Okamura, 1901:57.

Arasaki, 1964; Ariga, 1919, 1920; Chiang, 1962a; Fan, 1951, 1953a; Higashi, 1934; Horikawa, 1919; Okamura, 1901, 1915b, 1930, 1934a, 1936; Oshima, 1915; Santelices and Stewart, 1985; Shen and Fan, 1950.

Gelidium kintaroi (Okamura) Yamada, 1941:201.

Fan, 1951, 1953a; Shen and Fan, 1950. As "*G. clavatum*": Okamura, 1934a, 1935a, 1935b, 1936.

Gelidium latiusculum Okamura, 1935b:443.

Chiang, 1962a; Fan, 1951, 1953a; Okamura, 1935b, 1936; Shen and Fan, 1950.

Gelidium pacificum Okamura, 1914:99.

Okamura, 1915b; Oshima, 1915.

Gelidium planiusculum Okamura, 1935b:442.

Chiang, 1962a; Fan, 1951, 1953a; Okamura, 1935b, 1936; Santelices and Stewart, 1985; Shen and Fan, 1950.

Gelidium pusillum (Stackhouse) Le Jolis, 1863:139.

Chiang, 1962a; Fan, 1951; Santelices and Stewart, 1985; Taniguti, 1976. As "*Acrocarpus pusillus*": Martens, 1866.

Gelidium pusillum f. *foliaceum* Okamura, 1934a:51.

Ariga, 1920; Taniguti, 1976.

Gelidium subcostatum Okamura in Schmitz, 1894:1.

Ariga, 1919; Horikawa, 1919.

Gelidium yamadae (Okamura) Fan, 1951:10.

Fan, 1951, 1953a. As "*G. densum*": Okamura, 1935b; Shen and Fan, 1950.

Remarks: Fan (1951) considers *G. densum* invalid on the basis of an earlier homonym.

Gelidium sp.

Ariga, 1920.

Pterocladia nana Okamura, 1934a:64.

Chiang, 1962a, 1973b; Fan, 1951.

Pterocladia tenuis Okamura, 1934a:62.

Arasaki, 1964; Chiang, 1962a, 1962b, 1973b; Fan, 1951, 1953a; Okamura, 1935b, 1936; Segawa, 1974; Rho, 1958; Shen and Fan, 1950; Taniguti, 1976. As "*Gelidium corneum* var. *pinnatum*": Heydrich, 1894. As "*P. capillaceum*": Ariga, 1920; Chihara, 1970, 1975; Okamura, 1915b; Oshima, 1915.

GELIDIELLACEAE

Gelidiella acerosa (Forsskål) Feldmann et Hamel, 1934:533.

Ariga, 1920; Chiang, 1973b; Chihara, 1975; Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Yoshikawa and Yoshikawa, 1977. As "*Gelidiopsis rigida*": Okamura, 1931.

CRYPTONEMIALES**DUMONTIACEAE**

Dudresnaya japonica Okamura, 1908:209.

Shen and Fan, 1950.

RHIZOPHYLLIDACEAE

Remarks: Placed in the Cryptonemiales By Kylin (1956), this family was later transferred by Wiseman (1975) to the Gigartinales (see also Kraft (1981) and Wynne and Kraft (1981)). However, West and Hommersand (1981) noted "there is no major evidence to suggest that the [Rhizophyllidaceae] is closely related to either order," and retained it in the Cryptonemiales.

Chondrococcus hornemannii (Mertens) Schmitz, 1895:170, '*hornemannii*'.

Arasaki, 1964; Okamura, 1931, 1936; Segawa, 1974; Shen and Fan, 1950. As "*Desmia hornemannii*": Chiang, 1962a, 1973b; Chihara, 1970; Su et al., 1982; Yoshikawa and Yoshikawa, 1977.

Rhodopeltis borealis Yamada, 1931a:75.

Arasaki, 1964; Chihara, 1975; Nozawa, 1963, 1970; Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Yamada, 1931a.

Rhodopeltis gracilis Yamada et Tanaka in Yamada, 1935:30.

Arasaki, 1964; Nozawa, 1963, 1970; Okamura, 1936; Shen and Fan, 1950; Yamada, 1935.

Rhodopeltis setchelliae Yamada, 1935:33, '*setchellii*'.

Nozawa, 1963, 1970; Okamura, 1936; Shen and Fan, 1950; Yamada, 1935.

PEYSSONNELIACEAE

- Peyssonnelia caulifera* Okamura, 1899:8.
Ariga, 1920.
- Peyssonnelia distenta* (Harvey) Yamada, 1930:29.
Ariga, 1920; Chiang, 1962a, 1962b; Okamura, 1899, 1936; Shen and Fan, 1950; Yamada, 1930; Yoshikawa and Yoshikawa, 1977. As "*P. involvens*": Okamura, 1909.
- Peyssonnelia rubra* (Greville) J. Agardh, 1852:502.
De Toni, 1895; Heydrich, 1894; Okamura, 1897, 1931, 1936; Shen and Fan, 1950.

CORALLINACEAE

- Amphiroa beauvoisii* Lamouroux, 1816:299.
As "*A. pusilla*": Okamura, 1936; Segawa, 1974; Shen and Fan, 1950. As "*A. zonata*": Chiang, 1962a.
Remarks: Recently Norris and Johansen (1981:6) considered *A. zonata* to be a taxonomic synonym of *A. beauvoisii* Lamouroux.
- Amphiroa bowerbankii* Harvey, 1849b:97.
De Toni, 1895; Heydrich, 1894; Shen and Fan, 1950.
- Amphiroa ephedracea* (Lamarck) Decaisne, 1842:124.
Chihara, 1970; De Toni, 1895; Okamura, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950. As "*A. exilis*": Heydrich, 1894; Yendo, 1902.
- Amphiroa fragilissima* (Linnaeus) Lamouroux, 1816:298.
Yoshikawa and Yoshikawa, 1977.
- Amphiroa multifida* Kützing, 1858:27.
De Toni, 1895; Heydrich, 1894; Shen and Fan, 1950.
- Amphiroa pusilla* Yendo, 1902:13.
Rho, 1958; Su et al., 1982.
- Amphiroa* spp.
Ariga, 1920.
- Cheilosporum anceps* Yendo, 1902:18.
Shen and Fan, 1950.
- Cheilosporum jungermannioides* Ruprecht in J. Agardh, 1852:546.
Chihara, 1975; Okamura, 1931, 1936; Segawa, 1941, 1974; Shen and Fan, 1950; Tseng, 1983. As "*Amphiroa cultrata* var. *globulifera*": De Toni, 1895; Heydrich, 1894. As "*C. cultratum*": Yendo, 1902.
- Jania adhaerens* Lamouroux, 1816:270.
Chiang, 1962a, 1962b; Okamura, 1931, 1936; Shen and Fan, 1950; Su et al., 1982; Taniguti, 1976. As "*Corallina adhaerens*": De Toni, 1905; Heydrich, 1894; Yendo, 1902. As "*Corallina decussato-dichotoma*": Okamura, 1931, 1936; Segawa, 1974. As "*J. decussato-dichotoma*": Arasaki, 1964; Chiang, 1962a; Chihara, 1970, 1975; Shen and Fan, 1950; Taniguti, 1976.
- Jania radiata* Yendo, 1902:26.
Chiang, 1962a.
- Jania tenella* Kützing, 1858:41.
Chiang, 1962a.

JANIALEAE

- Jania undulata* Yendo, 1902:26.
Shen and Fan, 1950.
- Lithophyllum perulatum* Foslie, 1900:18.
De Toni, 1895; Okamura, 1936; Shen and Fan, 1950; Yendo, 1902. As "*Melobesia pustulata*": Heydrich, 1894.
- Lithophyllum* spp.
Ariga, 1920.
- Lithothamnium membranaceum* (Esper) Foslie, 1905:72.
De Toni, 1895; Okamura, 1936; Shen and Fan, 1950. As "*Melobesia membranaceae*": Heydrich, 1894; Yendo, 1902.
- Lithothamnium* spp.
Ariga, 1920.
- Mastophora pygmaea* Heydrich, 1894:300.
De Toni, 1895; Fan, 1974; Heydrich, 1894; Okamura, 1936; Shen and Fan, 1950; Yendo, 1902.
Remarks: Fan (1974:251, 254) considers the taxonomy of this species suspect and encourages study of the type specimen and type-locality specimens to confirm the identity of Heydrich's plant.
- Mastophora rosea* (C. Agardh) Setchell, 1943:129.
Chiang, 1973b; Chihara, 1975; Segawa, 1974; Su et al., 1982; Taniguti, 1976; Yoshikawa and Yoshikawa, 1977. As "*M. macrocarpa*": Chiang, 1962a; De Toni, 1905; Heydrich, 1894; Okamura, 1931, 1936; Shen and Fan, 1950; Yendo, 1902.
- Melobesia farinosa* Lamouroux, 1816:315.
De Toni, 1895; Heydrich, 1894; Okamura, 1936; Shen and Fan, 1950; Yendo, 1902.
- Melobesia* spp.
Ariga, 1920.
- Tenarea tumidulum* (Foslie) Adey, 1970:7.
As "*Dermatolithon tumidulum*": Chiang, 1973a; Okamura, 1935b.

ENDOCLADIACEAE

- Gloiopeltis complanata* (Harvey) Yamada, 1932a:117.
As "*Endocladia complanata*": Ariga, 1920. As "*Gloiopeltis cervicornis*": Ariga, 1921.
- Gloiopeltis furcata* (Postels et Ruprecht) J. Agardh, 1851:235.
Chiang, 1973b; Fan, 1953a; Shen and Fan, 1950.
- Gloiopeltis tenax* (Turner) J. Agardh, 1842:68.
Chiang, 1969, 1973a; Okamura, 1935b.

CRYPTONEMIACEAE

- Carpopeltis angusta* (Harvey) Okamura, 1910:66.
Ariga, 1920; Shen and Fan, 1950.
- Carpopeltis cornea* Okamura, 1936:553.
Arasaki, 1964; Chihara, 1975; Okamura, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Su et al., 1982.

Carpopeltis flabellata (Holmes) Okamura, 1935a:39.
Ariga, 1920; Chiang 1962a.

Carpopeltis formosana Okamura, 1931:110.
Fan, 1953a; Okamura 1931, 1936; Shen and Fan, 1950;
Tokida, 1939.

Carpopeltis rigida (Harvey) Schmitz, 1895:168.
Ariga, 1920; Chihara, 1975; Okamura, 1909, 1930,
1936; Rho, 1958; Segawa, 1974; Shen and Fan 1950;
Yoshikawa and Yoshikawa, 1977.

Grateloupia carnosa Yamada et Segawa in Yamada,
1938b:126.
Chiang, 1962a.

Grateloupia filicina (Wulfen) J. Agardh, 1851:180.
Ariga, 1919, 1920; Chiang, 1962a, 1973b; De Toni,
1895; Horikawa, 1919.

Grateloupia filicina f. *filiformis* (Kützing) Pilger, 1911:310.
As "G. filiformis": Martens, 1866.

Grateloupia okamurai Yamada, 1941:204, 'okamurai'.
Chiang, 1973a.

Grateloupia ramosissima Okamura, 1913:60.
Chiang, 1962a, 1973a; Chihara, 1970, 1975; Dawson,
1954c; Rho, 1958; Segawa, 1974; Shen and Fan, 1950.

Halymenia ceylanica (Harvey) Kützing, 1866:33.
As "H. formosa": Okamura, 1909. As "H. durvillaei" var.
formosa: Tokida, 1939. As "H. durvillaei" var. *ceylanica*":
Ariga, 1920; Chiang, 1962a; Okamura, 1936; Shen and
Fan, 1950.

Halymenia durvillaei Bory, 1826, pl. 15.
Fan, 1953a.

Polyopes polyideooides Okamura, 1895:447.
Chiang, 1973a, 1973b; Okamura, 1935b.

Polyopes sp.
Heydrich, 1894.

GIGARTINALES

SOLIERIACEAE

Eucheuma arnoldii Weber-van Bosse, 1928:421.
Kraft, 1972. As "E. cupressoideum": Shen and Fan, 1950.

Eucheuma "audiolis"
Tseng and Chen, 1977; Tseng, 1984.
Remarks: This species name is apparently a nomen nudum (fide M.S. Doty).

Eucheuma cottonii Weber-van Bosse, 1913b:115.
Tseng and Chen, 1977.

Eucheuma crassum Zanardini, 1878:36.
Shen and Fan, 1950; Yamada, 1936a.

Eucheuma crustaeforme Weber-van Bosse, 1928:415.
Okamura, 1931, 1936. As "E. cottonii": Okamura, 1931.

Eucheuma gelatinae (Esper) J. Agardh, 1852:628.
Chihara, 1975; Doty and Norris, 1985; Shen and Fan,
1950; Tseng, 1983; Tseng and Chen, 1977; Yamada,
1936a.

Eucheuma muricatum (Gmelin) Weber-van Bosse, 1928:413.
Higashi, 1934; Shen and Fan, 1950.

Eucheuma muricatum f. *depauperata* Weber-van Bosse,
1928:415.
Ariga, 1920; Okamura, 1936. As "E. muricatum": Oka-
mura, 1931. As "E. spinosum": Okamura, 1909, 1915b.

Eucheuma okamurai Yamada, 1936a:125, 'okamurai'.
Shen and Fan, 1950.

Eucheuma papulosa Cotton et Yendo in Cotton, 1914:220.
Cotton, 1915; Tilden, 1929.

Eucheuma serra J. Agardh, 1852:626.
Chiang, 1962a, 1973b; Chihara, 1975; Segawa, 1974;
Shen and Fan, 1950; Su et al., 1982; Tseng and Chen,
1977; Yamada, 1936a; Yoshikawa and Yoshikawa, 1977.

Eucheuma spp.
Ariga, 1920.

Meristotheca coacta Okamura, 1930:97.
Arasaki, 1964; Okamura, 1930, 1936; Segawa, 1974;
Yoshikawa and Yoshikawa, 1977.

Meristotheca papulosa (Montagne) J. Agardh, 1876:584.
Arasaki, 1964; Chiang, 1962a; Fan, 1953a; Okamura,
1936; Segawa, 1974; Tseng, 1983.

HYPNEACEAE

Hypnea boergesenii Tanaka, 1941:233.
Chiang, 1962a; Fan, 1953a; Shen and Fan, 1950; Tan-
aka, 1941; Tseng, 1983.

Hypnea crenomycé J. Agardh, 1852:452.
Shen and Fan, 1950; Tanaka, 1941; Yoshikawa and
Yoshikawa, 1977.

Hypnea cervicornis J. Agardh, 1852:451.
Yoshikawa and Yoshikawa, 1977.

Hypnea charoides Lamouroux, 1813:131.
Chiang, 1973b; Fan, 1953a; Shen and Fan, 1950; Tseng,
1983.

Hypnea chordacea Kützing, 1847:776.
Arasaki, 1964; Shen and Fan, 1950; Tanaka, 1941.

Hypnea chordacea f. *simpliciuscula* (Okamura) Tanaka,
1941:232.
Chiang, 1962a; Chihara, 1975; Shen and Fan 1950;
Tanaka, 1941.

Hypnea cornuta (Lamouroux) J. Agardh, 1852:449.
Shen and Fan, 1950; Tanaka, 1941.

Hypnea esperi Bory, 1828:157.
Shen and Fan, 1950; Tanaka, 1941.

Hypnea hamulosa (Turner) Lamouroux, 1813:44.
Okamura, 1931, 1936; Shen and Fan, 1950; Tanaka,
1941.
Remarks: Several authors (e.g., J. Agardh, 1852:447;
De Toni 1924:477; Okamura, 1936:611; Tanaka,
1941:245; Shen and Fan, 1950:339; Weber-van Bosse,
1928:453; and Zanardini, 1858:270) have incorrectly

cited "Montagne 1850:n.16 (page 9)" as the original place of publication.

Hypnea japonica Tanaka, 1941:236.

Chiang, 1962a; Chihara, 1970; Segawa, 1974; Shen and Fan 1950; Tanaka, 1941; Tseng, 1983.

Hypnea nidulans Setchell, 1924:161.

Chihara, 1975; Okamura, 1931, 1936; Shen and Fan, 1950; Su et al., 1982; Tanaka, 1941; Yoshikawa and Yoshikawa, 1977.

Hypnea pannosa J. Agardh, 1847:14.

Okamura, 1931, 1936; Shen and Fan, 1950; Tanaka, 1941; Yoshikawa and Yoshikawa, 1977.

Hypnea saidana Holmes, 1896:256.

Chiang, 1973b.

Hypnea seticulosa J. Agardh, 1852:446.

Ariga, 1920; Okamura, 1931, 1936; Tokida, 1939.

Hypnea spinella (Greville) Kützing, 1849:759.

Chiang, 1962a.

Hypnea sp.

Su et al., 1982.

PLOCAMIACEAE

Plocamium oviforme Okamura, 1896:23.

Ariga, 1920.

Plocamium serratulum Okamura, 1932:100.

Okamura, 1936; Shen and Fan, 1950. As "*P. costatum*":

Okamura, 1923, 1931; Yendo, 1918.

Plocamium telfairiae (J. Agardh) Harvey in Kützing, 1849:885.

Shen and Fan, 1950; Yendo, 1915. As "*P. abnorme*":

Ariga, 1919, 1920; Horikawa, 1919; Okamura, 1931.

Plocamium telfairiae f. *uncinatum* Okamura, 1936:615.

As "*P. abnorme* f. *uncinatum*": Okamura, 1913.

GRACILARIACEAE

Ceratodictyon spongiosum Zanardini, 1878:37.

Arasaki, 1964; Ariga, 1920; Okamura, 1936. As "*C. spongoides*": Okamura, 1931; Shen and Fan, 1950.

Gelidiopsis hachijoensis Yamada et Segawa, 1953:112.

Yoshikawa and Yoshikawa, 1977.

Gelidiopsis repens (Kützing) Schmitz, 1895:148.

Chiang, 1962a, 1973b; Chihara, 1975; Okamura, 1931, 1936; Shen and Fan, 1950; Taniguti, 1976; Yamada 1932b; Yoshikawa and Yoshikawa, 1977.

Gelidiopsis variabilis (Greville) Schmitz, 1895:148.

Yamada, 1932b.

Gracilaria arcuata Zanardini, 1858:265.

Chang and Xia, 1976; Chiang, 1973a, 1973b, 1985; Okamura, 1935b; Su et al., 1982.

Gracilaria blodgettii Harvey, 1853:111.

Chiang, 1985; Ohmi, 1958; Yang and Chiang, 1982.

Remarks: Although "*G. blodgettii*" has been reported

from Taiwan (Chiang, 1985), the South China Sea (Xia, 1985), and southern Japan (Yamamoto, 1985), the relationship between them and Caribbean type specimens of *G. blodgettii* Harvey is still to be resolved.

Gracilaria bursapastoris (Gmelin) Silva, 1952:265.

As "*G. compressa*": Chen, 1976; Shang, 1976.

Gracilaria canaliculata (Kützing) Sonder, 1871:56.

As "*Corallopsis opuntia*": Arasaki, 1964; Segawa, 1974.

Gracilaria chorda Holmes, 1896:253.

Ariga, 1920; Chen, 1976.

Gracilaria coronopifolia J. Agardh, 1852:592.

Chang and Xia, 1976; Chiang, 1985; Ohmi, 1958; Tseng, 1983; Yamada, 1941; Yang and Chiang, 1982.

As "*G. lichenoides* f. *coronopifolia*": Fan, 1953a.

Gracilaria crassa Harvey ex J. Agardh, 1876:417.

Chang and Xia, 1976; Chihara, 1975; Chiang, 1985; Ohmi, 1958; Okamura, 1936; Shen and Fan, 1950; Tseng, 1983; Yamada, 1933; Yang and Chiang, 1982.

Gracilaria denticulata (Kützing) Schmitz in Mazza, 1907:138.

Chiang, 1962a, 1985; Chihara, 1975; Okamura, 1931, 1936; Yang and Chiang, 1982.

Gracilaria edulis (Gmelin) Silva, 1952:293.

Nelson et al., 1983. As "*G. lichenoides*": Chen, 1976; Chiang, 1981; Shang, 1976.

Gracilaria eucheumoides Harvey, 1859:331.

Chang and Xia, 1976; Chiang, 1973a, 1985; Okamura, 1935b; Yang and Chiang, 1982.

Gracilaria gigas Harvey, 1859:330.

Chiang, 1981; Chen, 1976; Shang, 1976.

Gracilaria incurvata Okamura, 1931:41.

Chiang, 1973a; Okamura, 1935b.

Gracilaria punctata (Okamura) Yamada, 1941:203.

Ohmi, 1958; Shen and Fan, 1950; Yamada, 1941.

Gracilaria purpurascens Harvey in J. Agardh, 1885:63.

Ohmi, 1958; Shen and Fan, 1950; Yamada, 1938b; Yoshikawa and Yoshikawa, 1977.

Gracilaria spinulosa (Okamura) Chang et Xia, 1976:148.

Tseng, 1983. As "*G. purpurascens* f. *spinulosa*": Ohmi, 1958; Shen and Fan, 1950; Yamada, 1941; Yoshikwa and Yoshikwa, 1977.

Gracilaria salicornia (C. Agardh) Dawson, 1954b:4.

Chiang, 1985; Yang and Chiang, 1982.

Gracilaria "verrucosa".

G. verrucosa sensu Chiang, 1981, 1985; Nelson et al., 1983; Rho, 1958; Segawa, 1974; Yang et al., 1981; Yang and Chiang, 1982 [non *G. verrucosa* (Hudson) Papenfuss, 1950:195]. As "*G. confervoides*": De Toni, 1895; Fan, 1953a; Martens, 1866; Shang, 1976; Shen and Fan, 1950; Tokida, 1954.

Remarks: A comparison with type-locality (England) specimens of *G. verrucosa* showed the Taiwan specimens are a different species (Chiang, 1985; Abbott et al., 1985), and that the Chinese and Japanese specimens identified

as "*G. verrucosa*" are the same (Xia and Yamamoto, 1985; Yamamoto, 1985; Xia, 1985) and that they too are different from the British *G. verrucosa*. Recently, Zhang and Xia (1985:177) described *G. asiatica* for the specimens from Japan and China.

Gracilaria sp.

Ariga, 1920.

SPHAEROCOCCACEAE

Caulacanthus okamurae Yamada, 1933:278, 'okamurae'.

Rho, 1958; Taniguti, 1976; Yamada, 1933.

Caulacanthus spinellus (Hooker et Harvey) Kützing, 1849:753.

Chiang, 1973b; Shen and Fan, 1950; Yamada, 1933.

Caulacanthus ustulatus var. *fastigiatus* (Kützing) Pilger, 1920:5.

As "C. fastigiatus": Martens, 1866.

Phacelocarpus japonicus Okamura, 1902:79.

Shen and Fan, 1950.

SARCODIACEAE

Sarcodia ceylanica Harvey ex Kützing, 1869, pl. 33.

Chiang, 1962a; Shen and Fan, 1950; Yoshikawa and Yoshikawa, 1977.

PHYLLOPHORACEAE

Ahnfeltia paradoxa (Suringar) Okamura, 1934b:13.

As "Gymnogongrus paradoxus": Ariga, 1920.

Gymnogongrus flabelliformis Harvey, 1856:332.

Shen and Fan, 1950; Taniguti, 1976.

GIGARTINACEAE

Chondrus crispus Stackhouse, 1797:xxiv.

Ariga, 1920; Mikami, 1965.

Chondrus ocellatus Holmes, 1896:252.

Chiang, 1973b; Fan, 1953a; Tseng, 1983. As "C. ocellatus f. typicus": Chiang, 1962a.

Chondrus ocellatus f. *canaliculatus* Okamura, 1932:84.

Shen and Fan, 1950.

Gigartina intermedia Suringar, 1870:30.

Ariga, 1921; Chiang, 1962a; Fan, 1953a; Shen and Fan, 1950; Taniguti, 1976.

Gigartina ochotensis Ruprecht in Kjellman, 1889:31.

Ariga, 1920.

Gigartina tenella Harvey, 1859:331.

Arasaki, 1964; Okamura, 1930, 1936; Rho, 1958; Shen and Fan, 1950.

Rhodoglossum affine (Harvey) Kylin, 1928:49.

As "Chondrus affinis": Heydrich, 1894.

RHODYMENIALES

RHODYMENIACEAE

Chrysomenia procumbens Weber-van Bosse, 1928:470.

Okamura, 1931, 1936; Shen and Fan, 1950.

Erythrocolon podagraria (Harvey ex J. Agardh in Grunow) J. Agardh ex Kylin, 1931:14.

Shen and Fan, 1950.

Remarks: For discussion on the complicated nomenclature of this taxon see Abbott and Littler (1969:168).

Rhodymenia spinulosa Okamura, 1934b:33.

Okamura, 1934b, 1936; Yoshikawa and Yoshikawa, 1977.

Weberella micans Hauptfleisch in Schmitz and Hauptfleisch, 1897:402.

Arasaki, 1964; Okamura, 1936; Segawa, 1974; Shen and Fan, 1950; Yamada, 1932b.

CHAMPIACEAE

Champia parvula (C. Agardh) Harvey, 1853:76.

Chiang, 1962a; Higashi, 1934; Okamura, 1931, 1936; Shen and Fan, 1950.

CERAMIACEALES

CERAMIACEAE

Carpoblepharis schmitziana var. *erecta* Yamada, 1932b:273.

Okamura, 1936; Shen and Fan, 1950.

Carpoblepharis warburgii Heydrich, 1894:297.

De Toni, 1895; Heydrich, 1894; Okamura, 1936; Shen and Fan, 1950.

Remarks: Hommersand (1963:196) and Fan (1974) noted "C. warburgii" has never been re-examined, but . . . it appears unlikely that this taxon belongs in *Carpoblepharis*."

Centroceras clavulatum (C. Agardh) Montagne, 1846:140.

Ariga, 1920; Chiang, 1962a, 1962b, 1973b; Chihara, 1970, 1975; De Toni, 1895; Okamura, 1931, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Taniguti, 1976; Yamada, 1928; Yoshikawa and Yoshikawa, 1977.

As "Ceramium clavulatum": Heydrich, 1894.

Centroceras minutum Yamada, 1944c:42.

Chiang, 1962a.

Ceramium aduncum Nakamura, 1950:159.

Nakamura, 1950, 1965; Itono, 1972, 1977b. As "Ceramium clarionense": Dawson, 1954c.

Ceramium ciliatum (Ellis) Ducluzeau, 1805:64, var. *robustum* (J. Agardh) Mazoyer, 1938:322.

Nakamura, 1965; Itono, 1972, 1977b.

Ceramium flaccidium (Kützing) Ardisson, 1971:40.

As "C. gracillimum" var. *byssoideum*: Itono, 1972, 1977b.

Remarks: See Womersley (1978:234–238) for discussion on the nomenclature and taxonomy of this taxon.

Ceremium gracillimum (Kützing) Griffiths et Harvey in Harvey, 1848b, pl. 206.

Nakamura, 1965; Yoshikawa and Yoshikawa, 1977.

Remarks: Womersley (1979:234) considers *C. gracillimum* to be a synonym of *C. flaccidum* (Kützing) Ardisson.

Ceremium nakamurai Dawson, 1954a:6.

Nakamura, 1965; Itono, 1972, 1977b. As “*C. equisetoides*”: Nakamura, 1950.

Ceremium paniculatum Okamura, 1896:36.

Nakamura, 1965.

Ceramium tenerimum (Martius) Okamura, 1921:112.

Chiang, 1962a; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Itono, 1972, 1977b.

Ceramium tenuissimum (Lyngbye) J. Agardh, 1851:120.

Ariga, 1921; De Toni, 1895; Heydrich, 1894; Okamura, 1936; Rho, 1958; Shen and Fan, 1950; Tseng, 1983; Yamada, 1928.

Dasyphila plumariooides Yendo, 1920:7.

Arasaki, 1964; Itono, 1977b; Okamura, 1923, 1931, 1936; Segawa, 1974; Yendo, 1920.

Gymnothamnion elegans (C. Agardh) J. Agardh, 1892:27.

Itono, 1977b. As “*Plumaria ramosa*”: Okamura, 1936; Shen and Fan, 1950; Yamada and Tanaka, 1934.

Microcladia elegans Okamura 1907:1.

Chiang, 1962a.

Ptilothamnion cladophorae (Yamada et Tanaka) Feldmann-Mazoyer, 1941:375.

Fan, 1974; Itono, 1977b. As “*Spermothamnion cladophorae*”: Okamura 1936, 1937; Shen and Fan, 1950; Yamada and Tanaka, 1934.

Reinboldiella schmitziana (Reinbold) De Toni, 1895:35.

Ariga, 1920.

Spyridia filamentosa (Wulfen) Harvey ex Hooker, 1833:337.

Ariga, 1920; Chihara, 1975; De Toni, 1895; Heydrich, 1894; Okamura, 1913, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950.

Tiffaniella codicola (Yamada et Tanaka) Doty et Meñez, 1960:137.

Fan, 1974; Itono, 1977b. As “*Spermothamnion codicola*”: Okamura, 1936, 1937; Shen and Fan, 1950; Yamada and Tanaka, 1934.

Wrangelia velutina Harvey, 1854:546.

Okamura, 1931; Shen and Fan, 1950.

DELESSERIACEAE

Caloglossa bombayensis Børgesen, 1933:127.

Fan, 1952.

Claudea batanensis Tanaka, 1967:18.

Tanaka, 1967; Yoshikawa and Yoshikawa, 1977.

Haloplegma duperreyi Montagne, 1842:258.

Yamada, 1936b; Tseng, 1983.

Holmesia neurymenioides Okamura, 1932:98.

Okamura, 1936; Shen and Fan, 1950.

Martensia denticulata Harvey, 1854:537.

Ariga, 1919, 1920; Horikawa, 1919.

Martensia flabelliformis Harvey ex J. Agardh, 1863:826.

Chiang, 1962b; Chihara, 1975; Segawa, 1974; Shen and Fan, 1950; Yamada, 1936b.

Nitophyllum sp.

Ariga, 1920.

DASYACEAE

Dasya sp.

Ariga, 1919, 1920; Horikawa, 1919.

RHODOMELACEAE

Acanthophora aokii Okamura, 1934b:35.

Okamura, 1934b, 1936; Shen and Fan, 1950.

Acanthophora muscoidea (Linnaeus) Bory, 1828:156.

Shen and Fan, 1950.

Acanthophora orientalis (Sonder) J. Agardh, 1863:820.

Ariga, 1920; Chiang, 1962a, 1973b; Chihara, 1975; Fan, 1953a; Okamura, 1931, 1936; Segawa, 1974; Shen and Fan, 1950; Tokida, 1939; Yoshikawa and Yoshikawa, 1977.

Acrocystis nana Zanardini, 1872:145.

Chiang, 1973b; Chihara, 1975; Okamura, 1931, 1936; Segawa, 1974; Shen and Fan, 1950; Tseng, 1983.

Amansia glomerata C. Agardh, 1822:194.

Ariga, 1920; Chiang, 1973b; Chihara, 1975; Okamura, 1930, 1931, 1936; Segawa, 1974; Shen and Fan, 1950; Yoshikawa and Yoshikawa, 1977.

Bostrychia tenella (Vahl) J. Agardh, 1863:869.

Ariga, 1920; Chiang, 1962a; Taniguti, 1976; Yoshikawa and Yoshikawa, 1977.

Chondria armata (Kützing) Okamura, 1907:69.

Ariga, 1919, 1920; Chiang, 1962a, 1973b; Chihara, 1970, 1975; Fan, 1953a; Horikawa, 1919; Okamura, 1930, 1931, 1936; Segawa, 1974; Shen and Fan, 1950; Tokida, 1939; Tseng, 1983.

Chondria dasypylla (Woodward) C. Agardh, 1822:350.

Ariga, 1920; De Toni, 1895; Yoshikawa and Yoshikawa, 1977. As “*Laurencia dasypylla*”: Martens, 1866.

Digenia simplex (Wulfen) C. Agardh, 1822:389.

Ariga, 1920; Chihara, 1970, 1975; Fan, 1953a; Higashi, 1934; Okamura, 1931, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Tseng, 1983.

Herposiphonia subdisticha Okamura, 1899:37.

Chiang, 1962a; Okamura, 1931, 1936; Shen and Fan, 1950.

Herposiphonia sp.

Chiang, 1962b.

Laurencia bronniartii J. Agardh, 1841:20.

Remarks: Specimens from Nanwan, Taiwan (collected

- by Ger Dzeng-Joung; 21 August 1979; #US-071851), were identified by J. Norris as *L. bronniartii* and establish the presence of this taxon in southern Taiwan.
- Laurencia flexilis* var. *tropica* (Yamada) Xia et Zhang, 1982:538.
Tseng, 1983.
Laurencia forsteri (Mertens ex Turner) Greville, 1830:lii.
Ariga, 1920.
Laurencia glandulifera Kützing, 1849:855.
Ariga, 1920.
Laurencia grevilleana Harvey, 1854:545.
Ariga, 1920, Su et al., 1982.
Remarks: Saito and Womersley (1974:839) considered *L. grevilleana* to be a taxonomic synonym of *L. bronniartii* J. Agardh.
- Laurencia obtusa* var. *densa* Yamada, 1931b:226.
Dawson, 1954c; Shen and Fan, 1950; Yamada, 1931b, 1936c.
- Laurencia palizada* Yamada, 1931b:196.
Chiang, 1962a; Shen and Fan, 1950; Su et al., 1982; Yamada, 1931b, 1936c.
- Laurencia papillosa* (Forsskål) Greville, 1830:lii.
Ariga, 1920; Chihara, 1975; Cotton, 1915; Okamura, 1931; Shen and Fan, 1950; Tilden, 1929; Yamada, 1936c; Yoshikawa and Yoshikawa, 1977.
- Laurencia perforata* (Montagne) J. Agardh, 1876:648.
Ariga, 1920; De Toni, 1895; Heydrich, 1894.
- Laurencia pinnatifida* (Gmelin) Lamouroux, 1813:42.
As "*L. pinnatifida* var. *simplex*": De Toni, 1895; Heydrich, 1894.
- Laurencia tropica* Yamada, 1931b:223.
Okamura, 1931; Shen and Fan, 1950; Taniguti, 1976; Yamada, 1931b, 1936c.
- Laurencia venusta* Yamada, 1931b:203.
Okamura, 1931; Shen and Fan, 1950; Yamada, 1936c.
- Laurencia* sp.
Ariga, 1919; Fan, 1953a; Horikawa, 1919; Taniguti, 1976; Tokida, 1939.
- Leveillea jungermannioides* (Martius et Hering) Harvey, 1854:539.
- Ariga, 1920; De Toni, 1895; Chiang, 1962b; Heydrich, 1894; Okamura, 1931, 1936; Rho, 1958; Shen and Fan, 1950; Tseng, 1983; Yoshikawa and Yoshikawa, 1977. As "*Polyzonia jungermannioides*": Okamura, 1897.
- Murrayella periclados* (C. Agardh) Schmitz, 1893:227.
Tokida, 1941.
- Murrayella squarrosa* (Harvey) Schmitz, 1893:228.
Shen and Fan, 1950; Yamada, 1936b.
- Neurymenia fraxinifolia* (Mertens ex Turner) J. Agardh, 1863:1135.
Chiang, 1973b; Okamura, 1931; Shen and Fan, 1950; Tanaka and Itono, 1969.
- Polysiphonia harlandii* Harvey, 1859:330.
Segi, 1951; Tseng, 1944, 1983; Yamada, 1933; Yoshikawa and Yoshikawa, 1977.
- Polysiphonia kampsaxii* Børgesen, 1939:122.
Segi, 1951.
- Polysiphonia pulvinata* J. Agardh, 1842:124.
Segi, 1951.
- Sympyocladia marchantioides* (Harvey) Falkenberg, 1901: 277.
Ariga, 1920; Chiang 1962a; Chihara, 1970, 1975; Okamura 1930, 1931, 1936; Rho, 1958; Segawa, 1974; Shen and Fan, 1950; Tseng, 1983. As "*Placophora marchantioides*": Heydrich, 1894.
- Vidalia obtusiloba* (Mertens) J. Agardh, 1863:1123.
Ariga, 1920; Chihara, 1975; Okamura, 1931, 1936; Shen and Fan, 1950.

INCERTAE SEDIS

WURDEMANNIACEAE

Remarks: Taylor (1960:348, 361, 633) assigned the Wurdemaniaceae to the Gelidiales; however, Farr et al. (1979) have referred *Wurdemannia* to incertae sedis.

Wurdemannia setacea Harvey, 1853:246.

Okamura, 1931, 1936; Shen and Fan, 1950.

Wurdemannia sp.

Okamura, 1931.

Literature Cited

- Abbott, I.A.
- 1970. *Yamadaella*, a New Genus in the Nemaliales (Rhodophyta). *Phycologia*, 9:115–123.
 - 1976. *Dotyophycus pacificum* gen. et sp. nov., with a Discussion of Some Families of Nemaliales (Rhodophyta). *Phycologia*, 15:125–132.
 - 1984. Two New Species of *Liagora* (Nemaliales, Rhodophyta) and Notes on *Liagora farinosa* Lamouroux. *American Journal of Botany*, 71: 1015–1022.
- Abbott, I.A., and G.J. Hollenberg
- 1976. *Marine Algae of California*. xii + 827 pages. Stanford, California: Stanford University Press.
- Abbott, I.A., and M.M. Littler
- 1969. Some Rhodymeniales from Hawaii. *Phycologia*, 8:165–169.
- Abbott, I.A., and M. Yoshizaki
- 1981. A Second Species of *Dotyophycus* (Nemaliales, Rhodophyta) That Emphasizes the Distinctness of a Diffuse Gonimoblast. *Phycologia*, 20:222–227.
- Abbott, I.A., Y.M. Chiang, S. Fredericq, J.N. Norris, R.T. Tsuda, B.M. Xia, and H. Yamamoto
- 1985. The Red Alga *Gracilaria* Greville (Gracilariaeae, Gigartinales): Introduction. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds* . . . , pages 67–68. La Jolla, California: California Sea Grant College Program, University of California, La Jolla.
- Adey, W.H.
- 1970. A Revision of the Foslie Crustose Coralline Herbarium. *Det Kongelige Norske Videnskabers Selskab Skrifter*, 1:1–46.
- Agardh, C.A.
- 1820. *Species algarum* Volume 1, part 1, . . . 168 pages. Lund: Berling.
 - 1822. *Species algarum* . . . Volume 1, part 2, [v-vi] + 169–398 pages. Lund: Berling.
 - 1823. *Species algarum* . . . Volume 1, part 2, [vii–viii] + 399–531 pages. Greifswald: Ernst Mauritus.
 - 1824. *Systema algarum*. xxxviii + 312 pages. Lund: Berling.
- Agardh, J.G.
- 1841. In historiam algarum symbolae. *Linnaea*, 15:1–50, 443–457.
 - 1842. *Algae maris Mediterranei et Adriatici* . . . x + 164 pages. Paris: Fortin, Masson.
 - 1847. Nya alger från Mexico. *Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar*, 4(1):5–17.
 - 1848. *Species genera et ordines algarum, seu descriptiones succinctae specierum, generum et ordinum, quibus algarum regnum constituitur*. Volume 1, viii + 363 pages. Lund: C.W.K. Gleerup.
 - 1851. *Species genera et ordines algarum* . . . Volume 2, part 1, xii + 336 + [14] pages. Lund: C.W.K. Gleerup.
 - 1852. *Species genera et ordines algarum* . . . Volume 2, part 2, [2] + 337–700 + [20] + 701–786 pages. Lund: C.W.K. Gleerup.
 - 1863. *Species genera et ordines algarum* . . . Volume 2, part 3, fascicle 2, [2] + 787–1291 pages. Lund: C.W.K. Gleerup.
 - 1873. Till algernes systematik . . . Part 1, numbers i–iii. *Lunds Universitets Årsskrift*, 9:1–76.
 - 1876. *Species genera et ordines algarum* . . . Volume 3, part 1, *Epicrisis systematis floridearum*. viii + 724 pages. Leipzig: Malmström.
 - 1882. Till algernes systematik . . . Part 2, numbers iv–v. *Lunds Universitets Årsskrift*, 17:1–134 + [2], 3 plates.
 - 1883. Till algernes systematik . . . Part 3, number vi. *Lunds Universitets Årsskrift*, 19(2):1–177 + [4], 4 plates.
 - 1885. Till algernes systematik . . . Part 4, number vii. *Lunds Universitets Årsskrift*, 21(8):1–117 + [3], 1 plate.
 - 1887. Till algernes systematik . . . Part 5, number viii. *Lunds Universitets Årsskrift*, 23:1–174, 5 plates.
 - 1889. *Species Sargassorum Australiae descriptae et dispositae*. 133 pages, 31 plates. Stockholm: P.A. Norstedt & Söner.
 - 1892. Analecta algologica. *Lunds Universitets Årsskrift*, 28(6):1–182, 3 plates.
 - 1896. Analecta algologica, continuatio III. *Lunds Universitets Årsskrift*, 33(9):1–106, 1 plate.
- Allender, B.M., and G.T. Kraft
- 1983. The Marine Algae of Lord Howe Island (New South Wales): The Dictyotales and Cutleriales (Phaeophyta). *Brunonia*, 6:73–130.
- Arasaki, S.
- 1964. *How to Know the Seaweeds of Japan and Its Vicinity Fully Illustrated in Colours*. x + 217 pages, 491 + 7 + 24 figures. Tokyo: Hokuryukan. [In Japanese.]
- Ardisson, F.
- 1871. Rivista dei Ceramii della flore Italiana. *Nuovo Giornale Botanico Italiano*, 3:32–50.
- Areschoug, J.E.
- 1847. Phycarum, quae in maribus Scandinaviae crescunt, enumeratio, sectio prior. *Societas Scientiarum Upsaliensis, Nova Acta*, 13:223–382.
 - 1851. *Phyceae capenses* . . . 32 pages. Uppsala: Reg. Acad. Typographus.
- Ariga, K.
- 1919. Taiwan kaiso saishu nisshi. [An Account of Taiwan Marine Algae Collections.] *Transactions of the Natural History Society of Taiwan*, 9:10–13.
 - 1920. Taiwan oyobi Amonsan kaiso hikaku. [A Comparison of the Marine Algae of Taiwan and Amoy.] *Transactions of the Natural History Society of Taiwan*, 10:54–63.
 - 1921. Kamayamashima no kaiso. [The Marine Algae of Turtle Island.] *Transactions of the Natural History Society of Taiwan*, 11(53):21.
- Barton, E.S.
- 1901. The Genus *Halimeda*. In M. Weber, editor, *Siboga-Expedition, Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebied* . . . , monograph LX:1–32, plates 1–4. Leiden: E.J. Brill.
- Basson, P.W.
- 1979. Marine Algae of the Arabian Gulf Coast of Saudi Arabia (Second Half). *Botanica Marina*, 22(2):56–82.
- Børgesen, F. [C.E.]
- 1905. Contributions à la connaissance du genre *Siphonocladus* Schmitz. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger*, 3:259–291.
 - 1907. An Ecological and Systematic Account of the Caulerpas of the Danish West Indies. *Det Kongelige Danske Videnskabernes Selskabs Skrifter Naturvidenskabelige og Matematiske Afhandlinger*, 7th series, 4(5):337–392, 31 figures.
 - 1914. The Marine Algae of the Danish West Indies, Part 2: Phaeophyceae. *Dansk Botanisk Arkiv*, 2(2):1–66 + [2].
 - 1920a. Marine Algae from Easter Island. In C. Skottsberg, editor, *The Natural History of Juan Fernandez and Easter Island*, 2:247–309. Uppsala.
 - 1920b. The Marine Algae of the Danish West Indies, III: Rhodophyceae, Part 6, with Addenda to the Chlorophyceae, Phaeophyceae and Rhodophyceae. *Dansk Botanisk Arkiv*, 3:369–498 + [6].
 - 1932. A Revision of Forsskål's Algae Mentioned in Flora Aegyptiaco-

- Arabica and Found in His Herbarium in the Botanical Museum of the University of Copenhagen. *Dansk Botanisk Arkiv*, 8(2): 15 pages, 4 figures, 1 plate.
1933. Some Indian Rhodophyceae Especially from the Shores of the Presidency of Bombay. *Bulletin of Miscellaneous Information* (Royal Botanic Gardens, Kew), 3:113–142, 20 figures, 9 plates.
1934. Some Marine Algae from the Northern Part of the Arabian Sea with Remarks on Their Geographical Distribution. *Det Kongelige Danske Videnskabernes Selskabs, Biologiske Meddelelser*, 11(6):1–72, 8 figures, 2 plates.
1939. Marine Algae from the Iranian Gulf Especially from the Innermost Part Near Bushire and the Island Kharg. In K. Jessen and R. Spärck, editors, *Danish Scientific Investigations in Iran*, part I, pages 47–141. Copenhagen: E. Munksgaard.
1942. Some Marine Algae from Mauritius, III: Rhodophyceae, Part 1: Porphyridiales, Bangiales, Nenialionales. *Det Kongelige Danske Videnskabernes Selskabs, Biologiske Meddelelser*, 17(5):[1–3] + 4–63 + [64], 27 figures, 2 plates.
- Bory de Saint-Vincent, J. [B.G.M.]
1825. *Dictionnaire classique d'Histoire naturelle*. Volume 7, iii + 626 pages. Paris: Rey et Gravier.
1826. *Dictionnaire classique d'Histoire naturelle*. Volume 10, iii + 642 pages. Paris: Rey et Gravier.
1827. *Dictionnaire classique d'Histoire Naturelle*. Volume 12, iii + 634 pages. Paris: Rey et Gravier.
1828. Cryptogamie. In M.M. D'Urville, Botanique. In L.I. Duperrey, *Voyage autour du monde, exécuté par ordre du roi, sur la corvette de Sa Majesté, la Coquille, pendant les années 1822, 1823, 1824 et 1825, sous le Ministère et conformément avec instructions de S.G.M. le marquis Clermont Tonnerre, ministre de la marine . . .*, 2(fascicle 6):249–301. Paris: Arthus Bertrand.
- Brand, F.
1904. Über die Anheftung der Cladophoraceen und über verschiedene polynesische Formen dieser Familie. *Beihete zum Botanischen Zentralblatt*, 18:165–193.
- Castagne, J.L.M.
1851. *Catalogue des plantes qui croissent naturellement aux environs de Marseille, supplément*. i + 125 pages, plates 8–11. Aix: Nicot et Pardigon.
- Chang, C.F., and Xia Bangmei
1976. Studies on Chinese Species of *Gracilaria*. *Studia Marina Sinica*, 11(12):91–163, plates 1, 2.
- Chen, C., and Y.M. Chiang
1982. Morphological Studies on the Family Helminthocladiaeae (Nemaliales, Rhodophycophyta) of Taiwan, II: *Dermonema*. *Biological Bulletin, National Taiwan University*, 17:95–101.
- Chen, C., Y.M. Chiang, and S.H. Tsai Chiang
1981. Morphological Studies on the Family Helminthocladiaeae (Nemaliales, Rhodophycophyta) of Taiwan, I: *Helminthocladia*. *Biological Bulletin, National Taiwan University*, 16:27–32.
- Chen, T.P.
1976. Culture of *Gracilaria*. In *Aquaculture Practices in Taiwan*, pages 145–149. Taipei, Taiwan: Fishing News Books, Ltd.
- Chiang, Y.M.
1960. Marine Algae of Northern Taiwan (Cyanophyta, Chlorophyta, Phaeophyta). *Taiwania*, 7:51–75.
- 1962a. Marine Algae of Northern Taiwan (Rhodophyta). *Taiwania*, 8:143–165.
- 1962b. Marine Algae Collected from Penghu (Pescadores). *Taiwania*, 8:167–180.
1970. Observations on the Development of the Carposporophyte of *Scinaia pseudajaponica* Yamada et Tanaka (Nemaliales, Chaetangiaceae). *Journal of Phycology*, 6(3):289–292, 10 figures.
1971. Observations on the Development of Spermatangia in Some Genera of the Helminthocladiaeae. *Phycologia*, 10:163–167.
1972. Observations on the Development of the Carposporophyte of *Liagora boergesenii* Yamada (Nemaliales, Helminthocladiaeae). In K. Nisizawa, editor, *Proceedings of the 7th International Seaweed Symposium, Sapporo, Japan*, pages 111–113. New York: John Wiley & Sons.
- 1973a. Notes on Marine Algae of Taiwan. *Taiwania*, 18:13–17.
- 1973b. Studies on the Marine Flora of Southern Taiwan. *Bulletin of the Japanese Society of Phycology*, 21:97–102.
1981. Cultivation of *Gracilaria* (Rhodophycophyta, Gigartinales) in Taiwan. In T. Levring, editor, *Proceedings of the Xth International Seaweed Symposium*, pages 569–574. New York: Walter de Gruyter & Co.
1982. Cultivation of *Porphyta* in Taiwan. In R.T. Tsuda and Y.M. Chiang, editors, *Proceedings of Republic of China–United States Cooperative Science Seminar on Cultivation and Utilization of Economic Algae*, pages 81–86. Mangilao: University of Guam Marine Laboratory.
1985. *Gracilaria* from Taiwan: Key, List and Distribution of the Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds . . .*, pages 81–83. La Jolla, California: California Sea Grant College Program, University of California.
- Chiang, Y.M., and C. Chen
1982. The Genus *Liagora* of Taiwan. *Acta Oceanographie Taiwanica*, 13:181–196.
1983. Studies on *Datyophycus yamadae* (Ohmi et Itono) Abbott et Yoshizaki (Nemaliales, Rhodophycophyta) from Southern Taiwan. *Japanese Journal of Phycology*, 31:10–15.
- Chiang, Y.M., and H.N. Chou
1980. The Occurrence of *Zonaria stipitata* on the Southern Coasts of Taiwan. *Japanese Journal of Phycology*, 28:165–167.
- Chiang, Y.M., R. Huang, and H.M. Su
1974. The Uptake of ^{65}Zn by Marine Benthic Algae. *Acta Oceanographie Taiwanica*, 4:241–248.
- Chihara, M.
1969. *Pseudogliophloea okamurae* (Setchell) comb. nov. and *Ishige sinicola* (Setchell and Gardner) comb. nov. *Bulletin of the Japanese Society of Phycology*, 17:1–4.
1970. *Common Seaweeds of Japan in Color*. 173 pages, 64 plates. Osaka: Hoikusha Publishing Co. Ltd.
1975. *Marine Algae*. 290 pages. Tokyo: Gakken Co., Ltd. [In Japanese.]
- Chou, H.N., and Y.M. Chiang
1976. Studies on Algin from Brown Algae of Taiwan, I: Estimation of Yield and Quality of Algin. *Acta Oceanographie Taiwanica*, 6:135–139.
1981. The *Sargassum* of Taiwan. *Acta Oceanographie Taiwanica*, 12:132–149.
- Chou, R.C.Y.
1945. Pacific Species of *Galaxaura*, I: Asexual Types. *Papers of the Michigan Academy of Science, Arts, and Letters*, 30:33–55, plates 1–11.
1947. Pacific Species of *Galaxaura*, II; Sexual Types. *Papers of the Michigan Academy of Science, Arts, and Letters*, 31:3–24, plates 1–13.
- Christensen, C.F.A.
1905. *Index Filicum . . .* 384 pages. Copenhagen: H. Hagerup.
1906. *Index Filicum . . .* pages i–ix + 385–744 + xiii–lix. Copenhagen: H. Hagerup.
- Chueh, C.T., and C.C. Chen
1982. Seaweed Economics in Taiwan. In R.T. Tsuda and Y.M. Chiang, editors, *Cultivation and Utilization of Economic Algae*, pages 9–16. Mangilao, Guam: University of Guam Marine Laboratory.
- Collins, F.S.
1909. The Green Algae of North America. *Tufts College Studies, Scientific Series*, 2:79–480, 18 plates.
- Cotton, A.D.
1914. The Japanese Seaweed, *Tosaka nori*. *Bulletin of Miscellaneous Information* (Royal Gardens, Kew), 1914:219–222.

1915. Some Chinese Marine Algae. *Bulletin of Miscellaneous Information* (Royal Gardens, Kew), 1915:107–113.
- Dawson, E.Y.
1950. Notes on Some Pacific Mexican Dictyotaceae. *Bulletin of the Torrey Botanical Club*, 77:83–93.
1953. Marine Red Algae of Pacific Mexico, Part 1: Bangiales to Corallinaceae subf. Corallinoideae. *Allan Hancock Pacific Expeditions*, 17(1):1–239.
- 1954a. The Marine Flora of Isla San Benedicto Following the Volcanic Eruption of 1952–53. *Allan Hancock Foundation Publications, Occasional Papers*, 16:1–13.
- 1954b. Notes on Tropical Pacific Marine Algae. *Bulletin of the Southern California Academy of Sciences*, 53:1–7.
- 1954c. Marine Plants in the Vicinity of Nha Trang, Việt Nam. *Pacific Science*, 8:373–481.
1962. *New Taxa of Benthic Green, Brown and Red Algae*. 105 pages. Santa Ynez, California: Beaudette Foundation for Biological Research.
- Decaisne, J.
1841. Plantes de l'Arabie heureuse, recueillies par M.P.E. Botta. *Archives du Muséum National d'Histoire Naturelle* (Paris), 2:89–199.
1842. Mémoire sur les Corallines ou Polypiers Calcifères. *Annales des Sciences Naturelles*, Botanical series, II, 18:96–128.
- Delile, A.R.
1813. Flore de l'Egypte: Explication des planches. In France (Commission d'Egypte). Description de l'Egypte ou recueil des observations et des recherches qui ont été faites en Egypte pendant l'expédition de l'armée Francaise (1798–1801). *Histoire Naturelle de Paris*, 2(4):i + 145–320, plates 1–62. [Plates published in 1826.]
- De Toni, G.B.
1895. Phycæa japonicae novæ addita enumeratio algarum in ditione maritima japoniae hucusque collectarum: Alge marine del Giapponè ed isole ad esso appartenenti con illustrazione di alcune specie nuove (con 2 tavole). *Memorie del Real Istituto Veneto di Scienze, Letere ed Arti*, 25(5):1–78, plates 1, 2.
1905. *Sylloge algarum omnium hucusque cognitarum*. Volume 4, section 4: *Sylloge Floridearum* Pages [v] + 1523–1973. Padova: Typis Seminarii.
1924. *Sylloge algarum omnium hucusque cognitarum*. Volume 6: *Sylloge Floridearum* . . . sectio 5: additamenta. xi + 767 pages. Padova: Typis Seminarii.
- Dickie, G.
1874. On the Algae of Mauritius. *The Journal of the Linnean Society (Botany)*, 14:190–202.
- Dixon, P.S.
1967. The Typification of *Fucus cartilagineus* L. and *F. corneus* Huds. *Blumea*, 15:55–62.
- Doty, M.S.
1983. Coral Reef Diversified Farming. In C.K. Tseng, editor, *Proceedings of the Joint China-U.S. Phycology Symposium*, pages 437–478. Beijing, China: Science Press.
- Doty, M.S., and I.A. Abbott
1964. Studies in the Helminthocladiaeae, III: *Liagoropsis*. *Pacific Science*, 18:441–452.
- Doty, M.S., and E.G. Meñez
1960. *Tiffaniella*, a New Genus in the Ceramiales. *Transactions of the American Microscopical Society*, 79:135–144.
- Doty, M.S., and J.N. Norris
1985. *Eucheuma* Species (Solieriaceae, Rhodophyta) That Are Major Sources of Carrageenan. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds* . . . , pages 47–61. La Jolla, California: California Sea Grant College Program, University of California, La Jolla.
- Drew, K.M.
1956. *Confervula ceramicola* Lyngbye. *Botanisk Tidsskrift*, 53:67–74.
- Drew, K.M., and R. Ross
1965. Some Generic Names in the Bangiophyceae. *Taxon*, 14:93–99.
- Ducker, S.C.
1967. The Genus *Chlorodesmis* (Chlorophyta) in the Indo-Pacific Region. *Nova Hedwigia*, 13:145–182, plates 26–43.
- Ducker, S.C., W.T. Williams, and G.N. Lance
1965. Numerical Classification of the Pacific Forms of *Chlorodesmis* (Chlorophyta). *Australian Journal of Botany*, 13:489–499.
- Ducluzeau, J.A.P.
1805. *Essai sur l'histoire naturelle des conserves des environs de Montpellier*. i + 89 [90–92 + présentation] pages. Montpellier: Auguste Ricard.
- Egerod, L.E.
1952. An Analysis of the Siphonous Chlorophycophyta with Special Reference to the Siphonocladales, Siphonales, and Dasycladales of Hawaii. *University of California Publications in Botany*, 25(5):325–454, 13 plates.
- Eubank, L.L.
1946. Hawaiian Representatives of the Genus *Caulerpa*. *University of California Publications in Botany*, 18(18):409–432, 1 plate.
- Falkenberg, P.
1901. Die Rhodomelaceen des Golfes Neapel und der angrenzenden Meeres-Abschnitte. *Fauna und Flora des Golfs von Neapel und der angrenzenden Meeresteile*. Volume 26, xvi + 754 pages, 10 figures, 24 plates. Berlin: R. Friedlander & Sohns.
- Fan, K.C.
1951. The Genera *Gelidium* and *Pterocladia* of Taiwan. *Taiwan Fisheries Research Institute, Laboratory of Biology Report*, 2:1–22, 5 plates.
1952. The Structure, Methods of Branching and Tetrasporangia Formation of *Caloglossa*. *Taiwan Fisheries Research Institute, Hydrobiology Report*, 4:1–16.
- 1953a. A List of Edible Sea-weeds in Taiwan. *Taiwan Fisheries Research Institute, Hydrobiology Report*, 5/6:1–11.
- 1953b. The Identity of *Dilophus radicans* Okamura. *Taiwan Fisheries Research Institute, Hydrobiology Report*, 5/6:12–13.
1962. Studies on the Reproductive Organs of Red Algae, II: The Genus *Dermonema*. *Acta Botanica Sinica*, 10:336–340.
1974. Notes on Algal Taxonomy, I: A Review of Certain New Taxa on the Marine Algae of Taiwan Province, China. *Acta Phytotaxonomica Sinica*, 12:249–255.
- Fan, K.C., and Y.C. Wang
1974. Studies on the Marine Algae of the Hsisha Islands, China, I: New Species and New Records of the Family Nemalionaceae (Rhodophyta). *Acta Phytotaxonomica Sinica*, 12:489–495.
- Farr, E.R., J.A. Leussink, and F.A. Stafleu
1979. *Index nominum genericorum (plantarum)*. Volume I: Aa-Epochnum, xxvi + 630 pages; volume II: Eprolithus-Peersia, pages 631–1276; volume III: Pegaeophyton-Zzygium, pages 1277–1896. Utrecht: Bohn, Scheltema & Holkema.
- Feldmann, J.
1938. Sur la classification de l'ordre des Siphonocladals. *La Revue Générale de Botanique*, 50:571–597.
- Feldmann, J., and G. Hamel
1934. Observations sur quelques Gélidiacées. *Revue Générale de Botanique*, 46:528–549.
- Feldmann-Mazoyer, G.
1941. *Recherches sur les Céramiacées de la Méditerranée occidentale*. ii + 510 pages, 4 plates. Alger: Imprimerie Minerva.
- Forsskål, P.
1775. *Flora Aegyptiaco-Arabica* . . . [1]–32 + i–xxvi + [1] + [1]–219 + [220] pages + frontispiece + map. Copenhagen: Möller.
- Foslie, M.H.
1900. Revised Systematical Survey of the Melobesiaeae. *Det Kongelige Norske Videnskabers Selskabs Skrifter*, 1900(5):1–22.

1905. Remarks on Northern Lithothamnia. *Det Kongelige Norske Videnskabers Selskabs Skrifter*, 1905(3):1–138.
- Gaillon, F.B.
1828. Résumé méthodique des classifications des thalassophytes In F. Cuvier, *Dictionnaire des sciences naturelles* . . . , second edition, 53:1–59, plates 1, 2. Strasbourg: F.G. Levraut.
- Gardner, N.L.
1909. New Chlorophyceae from California. *University of California Publications in Botany*, 3:371–375, 1 plate.
- Gepp, A., and E.S. Gepp
1908. Marine Algae (Chlorophyceae and Phaeophyceae) and Marine Phanerogams of the "Sealark" Expedition, Collected by J. Stanley Gardiner *Transactions of the Linnean Society*, 7:163–188.
- Gray, J.E.
1866. On *Anadyomene* and *Microdictyon*, with the Description of Three New Allied Genera, Discovered by Menzies in the Gulf of Mexico. *Journal of Botany, British and Foreign*, 4:41–51, 65–72.
- Gray, S.F.
1821. *A Natural Arrangement of British Plants* Volume 1, xxviii + 824 pages, 21 plates. London.
- Greville, R.K.
1830. *Algae Britannicae, or Descriptions of the Marine and Other Articulated Plants of the British Islands, Belonging to the Order Algae*. lxxxviii + 218 pages, 19 plates. Edinburgh: MacLachlan and Stewart.
- Hamel, G.
1939. *Pheophycées de France*. Part 5, pages i–lvii, 337–432, 10 plates. Paris.
- Hansen, J.E., J.E. Packard, and W.T. Doyle
1981. *Mariculture of Red Seaweeds*. x + 42 pages. La Jolla, California: California Sea Grant College Program, University of California, La Jolla.
- Hariot, P.A.
1889. Algues. *Mission Scientifique du Cap Horn, 1882–1883*. Volume 5(Botanique), pages 3–119, plates 1–9. Paris: Gauthier-Villars et fils.
- Harvey, W.H.
- 1847–1849a [1849]. *Phycologia britannica* Volume 2, vi + plates 121–240 with text [1847, fascicles 21–24 comprising plates 121–144 with text; 1848a, fascicles 25–36 comprising plates 145–216 with text; 1849a, fascicles 37–40 comprising plates 217–240 with text]. London: Reeve, Benham & Reeve.
- 1848b [1847]. *Nereis australis* Part 1, [iii] + viii + 64 pages, plates 1–25. London: Reeve Brothers.
- 1849b. *Nereis australis* Part II, pages 65–124, plates 26–50. London: Reeve Brothers.
1853. *Nereis boreali-americana*, Part II: *Rhodospirmeae*. First issue, ii + 258 pages, plates 13–26. Washington, D.C.: Smithsonian Institution and London: John Van Voorst. [Third issue, 1858 (dated 1853), published as *Smithsonian Contributions to Knowledge*, 5(5):ii + 258 pages, plates 13–36.]
1854. Some Account of the Marine Botany of the Colony of Western Australia. *Transactions of the Royal Irish Academy*, 22:525–566.
1856. Algae of Japan. In A.S. Gray, Account of the Botanical Specimens. In *Narrative of the Expedition of an American Squadron to the China Seas and Japan*, 2:331–332. Washington: Beverley Tucker. [Listed under "Commodore M.C. Perry" in most libraries.]
1858. *Nereis boreali-americana*, Part III: *Chlorospermeae*. First issue, [ii] + 140 pages, plates 37–50. Washington, D.C.: Smithsonian Institution and London: John van Voorst. [Also issued as *Smithsonian Contributions to Knowledge*, 10:1–140, plates 1–14.]
1859. Characters of New Algae, Chiefly from Japan and Adjacent Regions, Collected by Charles Wright in the North Pacific Ex-
- ploring Expedition under Captain John Rodgers. *Proceedings of the American Academy*, 4:327–335.
1863. *Phycologia Australica* Volume 5, vi + plates 241–300 with text, index [vii]–x, and synoptic catalogue [v]–lxiii. London: Lovell Reeve.
- Hauck, F.
1884. Die Meeresalgen Deutschlands und Österreichs. In L. Rabenhorst, *Kryptogamen-Flora von Deutschland, Österreich und der Schweiz*, edition 2, 2:321–512. Leipzig: Edward Kummer.
1886. Über einige von J.M. Hildenbrandt in Rothen Meere und in Indischen Ocean gesammelte Algen, I. *Hedwigia*, 25(5):165–168.
1887. Über einige von J.M. Hildebrandt in Rothen Meere und in Indischen Ocean gesammelte Algen, IV. *Hedwigia*, 26:41–45.
- Heydrich, F.
1894. Beitrag zur Kenntnis der Algenflora von Ost-Asien besonders der Insel Formosa, Mollukken und Liu-kiu-Inseln. *Hedwigia*, 33:267–306, 2 plates.
- Higashi, M.
1934. *Natural Coloured Figures and Descriptions of the Japanese Seaweeds*. Pages i, 1–4, 1–5, 1–8, 1–80, 80 plates. Tokyo.
- Holmes, E.M.
- 1896 [1895]. New Marine Algae from Japan. *Journal of the Linnaean Society of Botany*, 31:248–269, 12 plates.
- Hommersand, M.H.
1963. The Morphology and Classification of Some Ceramiaceae and Rhodomelaceae. *University of California Publications in Botany*, 35(2):165–366.
- Hooker, W.J.
1833. Class XXIV: Cryptogamia. *The English Flora of Sir James Edward Smith*. Volume 5, pages 322–385. London.
- Hooker, W.J., and G.A. Arnott
1838. *The Botany of Captain Beechey's Voyage*. Part 6, pages 241–288, plates 50–59. London: Henry G. Bohn.
- Horikawa, Y.
1919. Taiwan kaiso. [Marine Algae of Taiwan.] *Transactions of the Natural History Society Formosa*, 9:13–16.
- Howe, M.A.
1905. Phylogenetic Studies, II: New Chlorophyceae, New Rhodophyceae, and Miscellaneous Notes. *Bulletin of the Torrey Botanical Club*, 32:563–586, 6 plates.
1914. The Marine Algae of Peru. *Memoirs of the Torrey Botanical Club*, 15:[i] + 185 pages, 66 plates.
1917. A Note on the Structural Dimorphism of Sexual and Tetrasporic Plants of *Galaxaura obtusata*. *Bulletin of the Torrey Botanical Club*, 43:621–624.
1918. Further Notes on the Structural Dimorphism of Sexual and Tetrasporic Plants in the Genus *Galaxaura*. *Brooklyn Botanical Garden Memoir*, 1:191–197.
1920. Algae. In N.L. Britton and C.F. Millspaugh, *The Bahama Flora*, pages 553–618. New York: Published by the authors.
- Itono, H.
1972. The Genus *Ceramium* (Ceramiaceae, Rhodophyta) in Southern Japan. *Botanica Marina*, 15:74–86.
1973. Notes on Marine Algae from Hateruma Island, Ryukyu. *Botanical Magazine (Tokyo)*, 86:155–168.
- 1977a. Studies on the Southern Japanese Species of *Galaxaura* (Rhodophyta). *Micronesica*, 13:1–26.
- 1977b. Studies on the Ceramiaceous Algae (Rhodophyta) from Southern Parts of Japan. *Bibliotheca Phycologica*, 35: 499 pages. Vaduz: J. Cramer.
- Jackson, B.D.
1893. *Index Kewensis* xvi + 728 + vii + 729–1268 pages. Oxford: Clarendon Press.

- Kjellman, F.R.
- 1889. Om Beringhavets algflora. *Det Kongelige Vetenskaps Akademiens Handlingar*, 23(8): 58 pages.
 - 1897a. Japanska arter af slägten *Porphyra*. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar*, 23(4):1–34.
 - 1897b. Marine Chlorophyceer från Japan. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar*, 23(11):1–44, 4 plates.
 - 1900. Om Floridé-Slägten *Galaxaura* dess organografi och systematik. *Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar*, 33(1):1–107, 20 plates.
- Kraft, G.T.
- 1972. Preliminary Studies of Philippine *Eucheuma* Species (Rhodophyta), Part 1: Taxonomy and Ecology of *Eucheuma arnoldii* Weber-van Bosse. *Pacific Science*, 26:318–334.
 - 1981. Rhodophyta: Morphology and Classification. In C.S. Loban and M.J. Wynne, editors, *The Biology of Seaweeds*, pages 6–51. Berkeley: University of California Press.
- Krauss, C.F.F. von
- 1846. Algae. In *Beiträge zur Flora des Cap- und Natallandes . . .*, 29:192–215.
- Kuntze, C.E.O.
- 1898. *Revisio generum plantarum . . .* Volume 3, part 3, pages 1–576. Leipzig: Arthur Felix.
- Kützing, F.T.
- 1843. *Phycologia generalis oder Anatomie, Physiologie und Systemkunde der Tange . . .* xxxii + 458 pages, 80 plates. Leipzig: F.A. Brockhaus.
 - 1845. *Phycologia Germanica: Deutschlands Algen in bündigen Beschreibungen*. vii + 240 pages. Nordhausen: Wilh. Köhne.
 - 1847. Diagnosen einiger neuen auständischen Algenspecies . . . *Flora oder Allgemeine Botanische Zeitung*, 30(48):773–776.
 - 1849. *Species algarum . . .* vi + 922 pages. Leipzig: F.A. Brockhaus.
 - 1856. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 6, vi + 35 pages, 100 plates. Nordhausen: Kosten des Verfassers.
 - 1858. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 8, parts 1–2, ii + 48 pages, 100 plates. Nordhausen: Kosten des Verfassers.
 - 1860. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 10, parts 1–2, iv + 39 pages, 100 plates. Nordhausen: Kosten des Verfassers.
 - 1866. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 16, parts 1–2, i + 35 pages, 100 plates. Nordhausen: Kosten des Verfassers.
 - 1868. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 18, parts 1–2, i + 35 pages, 100 plates. Nordhausen: Kosten des Verfassers.
 - 1869. *Tabulae phycologicae oder Abbildungen der Tange*. Volume 19, parts 1–2, 36 pages, 100 plates. Nordhausen: Kosten des Verfassers.
- Kylin, H.
- 1928. Entwicklungsgeschichtliche Florideenstudien. *Lunds Universitets Årsskrift*, 24(4):1–127.
 - 1931. Die Florideenordnung Rhodymeniales. *Lunds Universitets Årsskrift*, 27(11):1–48, 20 plates.
 - 1956. *Die Gattungen der Rhodophyceen*. xv + 673 pages. Lund: C.W.K. Gleerups.
- Lamouroux, J.V.F.
- 1809a. Exposition des caractères du genre *Dictyota* (1), et tableau des espèces qu'il renferme. *Journal de Botanique* (Paris), 2:38–44.
 - 1809b. Observations sur la physiologie des algues marines, et description de cinq nouveaux genres de cette famille. *Nouveau Bulletin des Sciences de la Société Philomatique de Paris*, 1:330–333.
 - 1812. Extrait d'un mémoire sur la classification des polypiers coralligènes non entièrement pierreux. *Nouveau Bulletin des Sciences de la Société Philomatique de Paris*, 3:181–188.
 - 1813. Essai sur les genres de la famille des thalassiphyses non articulés.
- Le Jolis, A.
- 1863. *Liste des algues marines de Cherbourg*. 168 pages, 6 plates. Paris: J.B. Baillière et Fils. [Preprinted from *Mémoires de la Société Impériale des Sciences Naturelles et Mathématiques de Cherbourg*, 1864, 10:5–168.]
- Lewis, S.M., J.N. Norris, and R.B. Searles
- 1987. The Regulation of Morphological Plasticity in Tropical Reef Algae by Herbivory. *Ecology*, 68.
- Liaw, J.P., and Y.M. Chiang
- 1979. Culture Studies on the *Concoocelis* of *Porphyra angusta* Udea. *Journal of the Fisheries Society of Taiwan*, 6:59–65.
- Linnaeus, C.
- 1753. *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum sistema sexuale digestas*. Volume 2, pages 561–1200. Stockholm: Impensis Laurentii Salvii.
- Liu, T.Y.
- 1982. Quality Improvement of Algin from *Sargassum*. In R.T. Tsuda and Y.M. Chiang, editors, *Proceedings of Republic of China–United States Cooperative Science Seminar on Cultivation and Utilization of Economic Algae*, pages 65–80. Mangilao: University of Guam Marine Laboratory.
- Lobban, C.S., and M.J. Wynne
- 1981. *The Biology of Seaweeds*. xi + 786 pages. Berkeley: University of California Press.
- Martens, G. von
- 1866. *Die preussische Expedition nach Ost-Asiens, Botanischer Theil. Die Tange*. 152 pages, plates 1–8. Berlin: Ober-hofbuchdruckerei, R.V. Decker.
- Mazoyer, G.
- 1938. Les Céramiacées de l'Afrique du Nord. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 29: 317–331.
- Mazza, A.
- 1907. Saggio di algologia oceanica . . . *Nuova Notarisia, Rassegna Consacrata allo Studio della Alghe*, 18:126–152.
- Meneghini, G.
- 1840. *Lettera del Prof. Giuseppe Meneghini al Dottore Jacob Corinaldi a Pisa*. [4] pages. Padova: Tipografia Prospieri.
- Michanek, G.
- 1978. Trends in Applied Phycology, with a Literature Review: Seaweed Farming on an Industrial Scale. *Botanica Marina*, 21:469–475.
- Mikami, H.
- 1965. A Systematic Study of the Phyllophoraceae and Gigartinaceae from Japan and Its Vicinity. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido University*, 5:181–285, plates I–XI.
- Montagne, J.P.F.C.
- 1838. De l'organisation et du mode de reproduction des Caulerpées et en particulier du *Caulerpa webbiana*, espèce nouvelle des îles Canaries. *Annales des Sciences Naturelles*, 2nd series, 9:129–150. Paris.
 - 1842. Troisième centurie de cellulaires exotiques nouvelles . . . *Annales des Sciences Naturelles, Botanique*, 18:241–282.
 - 1844. Quelques observations touchant la structure et la fructification des genres *Ctenodus*, Kütz., *Delisea*, Lamx., et *Lenormandia*, Montag., de la famille des Floridées. *Annales des Sciences Naturelles, Botanique*, series 3, 1:126–129, 151–161, plates 10, 11.
 - 1846. Phyceae. In M.C. Durieu de Maisonneuve, Flore d'Algérie, Botanique, 1: Cryptogamie. In *Exploration scientifique de l'Algérie . . .*, pages 1–197, plates 1–16. Paris: Imprimerie Royal.

1849. Sicerne Centurie de Plantes cellulaires nouvelles, tant indigenes qu'exotiques, Decades III. a VI. *Annales des Sciences Naturelles*, 11:33–66.
1850. Pugillus Algarum Yemenium *Annales des Sciences Naturelles*, 13:236–248.
- Murray, G.
1889. On *Boodlea*, a New Genus of Siphonocladaceae. *Journal of the Linnean Society, Botany*, 25:243–245.
- Nakamura, Y.
1950. New *Ceramiums* and *Campylaephoras* from Japan. *Science Papers, Institute of Algal Research, Hokkaido University Faculty of Science*, 3:155–172.
1965. Species of the Genera *Ceramium* and *Campylaephora*, Especially Those of Northern Japan. *Science Papers, Institute of Algal Research, Hokkaido University Faculty of Science*, 5:119–180.
- Needham, J.
1970. *Clerks and Craftsmen in China and the West*. 470 pages. Cambridge: Cambridge University Press.
- Nees von Esenbeck, C.
1820. *Horae physicae berolinenses Part 1*, pages 1–7, plate 1. Bonn: Adolphii Marcus.
- Nelson, S.G., S.S. Yang, C.Y. Wang, and Y.M. Chiang
1983. Yield and Quality of Agar from Species of *Gracilaria* (Rhodophyta) Collected from Taiwan and Micronesia. *Botanica Marina*, 26:361–366.
- Nicholson, D.H., and J.N. Norris
1983. Ordinal Orthography: Nemalionales vs. Nemaliales (Rhodophyta). *Taxon*, 32(2):288–292.
- Nizamuddin, M.
1970. Phytogeography of the Fucales and Their Seasonal Growth. *Botanica Marina*, 12:131–139.
- Norris, J.N., and K.E. Bucher
1982. Marine Algae and Seagrasses from Carrie Bow Cay, Belize. In K. Rützler and I.G. Macintyre, editors, *The Atlantic Barrier Reef Ecosystem at Carrie Bow Cay, Belize, I: Structure and Communities*. *Smithsonian Contributions to the Marine Sciences*, 12:167–223.
- Norris, J.N., and W.H. Fenical
1985. Natural Products Chemistry: Uses in Ecology and Systematics. In M.M. Littler and D.S. Littler, editors, *Handbook of Phycological Methods, V: Ecological Field Methods for Macroalgae*, pages 121–145. New York/Cambridge: Cambridge University Press.
- Norris, J.N., and H.W. Johansen
1981. The Articulated Coralline Algae of the Gulf of California, Mexico, I: *Amphiroa* Lamouroux. *Smithsonian Contributions to the Marine Sciences*, 9:1–29.
- Nozawa, Y.
1963. Systematic Anatomy of the Genus *Rhodopeltis*. *Memoirs of the Janshin Junior College*, 5:1–48, 2 plates.
1970. Systematic Anatomy of the Red Algal Genus *Rhodopeltis*. *Pacific Science*, 24:99–133.
- Ohmi, H.
1958. The Species of *Gracilaria* and *Gracilariaopsis* from Japan and Adjacent Waters. *Memoirs of the Faculty of Fisheries, Hokkaido University*, 6(1):1–66.
- Okamura, K.
1895. New or Little Known Algae from Japan. *The Botanical Magazine* (Tokyo), 9:445–455, 472–482, 1 plate.
1896. Contributions to the Knowledge of the Marine Algae of Japan, II. *The Botanical Magazine* (Tokyo), 10:23–36.
1897. On the Algae from Ogasawarajima (Bonin Islands). *The Botanical Magazine* (Tokyo), 11:1–17, 1 plate, 4 figures.
1899. Contributions to the Knowledge of the Marine Algae of Japan, III. *The Botanical Magazine* (Tokyo), 13:2–10, 35–43, 11 plates.
1901. *Illustrations of the Marine Algae of Japan*. Volume 1, part 5, pages 57–74 [Japanese text, pages 79–102], 5 plates. Tokyo: Keigyosha & Co.
1902. *Illustrations of the Marine Algae of Japan*. Volume 1, part 6, pages 75–93 [Japanese text, pages 103–128], 5 plates. Tokyo: Keigyosha & Co.
1907. *Icones of Japanese Algae*. Volume 1, pages 1–119, plates 1–25. Tokyo: Kazamashobo.
1908. *Icones of Japanese Algae*. Volume 1, pages 121–233, plates 26–45. Tokyo: Kazamashobo.
1909. *Icones of Japanese Algae*. Volume 2, pages 1–62, plates 51–65. Tokyo: Kazamashobo.
1910. *Icones of Japanese Algae*. Volume 2, pages 63–126, plates 66–85. Tokyo: Kazamashobo.
1912. *Illustrations of the Marine Algae of Japan*. Volume 2, pages 127–191, plates 86–100. Tokyo: Kazamashobo.
1913. *Icones of Japanese Algae*. Volume 3, pages 1–77, plates 101–120. Tokyo: Kazamashobo.
1914. *Icones of Japanese Algae*. Volume 3, pages 79–121, plates 121–130. Tokyo: Kazamashobo.
- 1915a. *Icones of Japanese Algae*. Volume 3, pages 123–218, plates 131–150. Tokyo: Kazamashobo.
- 1915b. Taiwan no ten gusa. [*Gelidium* of Taiwan.] *Shokubutsugaku Zasshi*, 29:57–58. [In Japanese.]
1916. List of Marine Algae Collected in the Caroline and Mariana Islands. *The Botanical Magazine* (Tokyo), 30:1–14.
1919. Notes on Some Marine Algae from Amoy, I. *The Botanical Magazine* (Tokyo), 33:12–14.
1921. *Icones of Japanese Algae*. Volume 4, pages 63–149, plates 166–185. Tokyo: Kazamashobo.
1923. *Icones of Japanese Algae*. Volume 4, pages 185–205, plates 196–200. Tokyo: Kazamashobo.
1930. On the Algae from the Island Hatidyo. *Records of Oceanographic Works in Japan*, 2(2):92–110, 2 plates.
1931. On the Marine Algae from Kōtōsho (Botel Tobago). *Bulletin of the Biogeographical Society of Japan*, 2(2):95–122, 3 plates.
1932. *Icones of Japanese Algae*. Volume 6, pages 49–101 [Japanese text, 47–96], plates 276–300. Tokyo: Kazamashobo.
- 1934a. On *Gelidium* and *Pterocladia* of Japan. *Journal of the Imperial Fisheries Institute*, 29(2):47–67, 18 plates.
- 1934b. *Icones of Japanese Algae*. Volume 7, pages 9–37 [Japanese text, 9–34], plates 306–320. Tokyo: Kazamashobo.
- 1935a. *Icones of Japanese Algae*. Volume 7, pages 39–72 [Japanese text 35–63], plates 321–340. Tokyo: Kazamashobo.
- 1935b. Taiwan-san Tengusa ni Tsuite. *Japanese Association for the Advancement of Science*, 10:441–443.
1936. Nippon Kaiso-shi. [Marine Algae of Japan.] Frontispiece + 9 + 6 + 964 + 11 pages. Tokyo: Published by the author.
1937. *Icones of Japanese Algae*. Volume 7, pages 73–79, plates 341–345. Tokyo: Kazamashobo.
- Oshima, M.
1915. Taiwan san Tengusa. [*Gelidium* in Taiwan.] *Transactions of the Natural History Society of Taiwan*, 5:55.
- Papenfuss, G.F.
1944. Notes on Algal Nomenclature, III: Miscellaneous Species of Chlorophyceae, Phaeophyceae and Rhodophyceae. *Farlowia*, 1(3):337–346.
1950. Review of the Genera of Algae Described by Stackhouse. *Hydrobiologia*, 2:181–208.
1956. Notes on South African Marine Algae, IV. *The Journal of South African Botany*, 22(2):65–77.
1960. On the Genera of the Ulvales and the Status of the Order. *Journal of the Linnean Society, Botany*, 56:303–318.
1967. Notes on Algal Nomenclature, V: Various Chlorophyceae and

- Rhodophyceae. *Phykos*, 5:95–105.
1977. Review of the Genera of Dictyotales (Phaeophycophyta). *Bulletin of the Japanese Society of Phycology*, 25(supplement):271–287.
- Papenfuss, G.F., K.E. Mshigeni, and Y.M. Chiang
1982. Revision of the Red Algal Genus *Galaxaura* with Special Reference to the Species Occurring in the Western Indian Ocean. *Botanica Marina*, 25:401–444.
- Piccone, A.
1884. Alge. In *Crociera del Corsaro alle Isole Madera e Canarie del Capitano Enrico d'Albertis*, 60 pages. Genova: R. Instituto Sordomuti.
- Pilger, R.
1911. Die Meeresalgen von Kamerun, nach der Sammlung von C. Ledermann. *Sonderabdruck aus Engler's Botanischen Jahrbüchern*, 46(3):294–323, 26 figures.
1920. Algae Mildbraediana Annobonenses. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie Herausgegeben von A. Engler*, 57(1):1–14, 34 figures.
- Reinbold, T.
1901. The Marine Algae. In J. Schmidt, Flora of Koh Chang: Contributions to the Knowledge of the Vegetation in the Gulf of Siam, Part IV. *Botanisk Tidsskrift*, 24(2):187–201.
1905. Einige neue Chlorophyceen aus dem Ind. Ocean (Niederl. Indien), gesammelt von A. Weber van Bosse. *La Nuova Notarisia*, 16:145–149.
- Rho, J.H.
1958. A Preliminary Survey of the Marine Algae of Korea. *Collectio Theseon, Scientia Naturalis, Universitas Sung Kyun Kwan*, 3:41–143.
- Ryther, J.H.
1979. Aquaculture in China. *Oceanus*, 22:21–28.
- Saito, Y., and H.B.S. Womersley
1974. The Southern Australian Species of *Laurencia* (Ceramiales: Rhodophyta). *Australian Journal of Botany*, 22:815–874.
- Santelices, B., and J.G. Stewart
1985. Pacific Species of *Gelidium* Lamouroux and Other Gelidiales (Rhodophyta), with Keys and Descriptions to the Common or Economically Important Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds* . . . , pages 17–31. La Jolla, California: California Sea Grant College Program, University of California, La Jolla.
- Schmitz, F.
1893. Die Gattung *Lophothalia* J. Ag. *Sonderabdruck aus den Berichten der Deutschen Botanischen Gesellschaft*, 11(3):212–232.
1894. Neue japanische Florideen von K. Okamura. *Hedwigia*, 33:190–201, 1 plate.
1895. Marine Florideen von Deutsch-Ostafrika. In A. Engler, Beiträge zur Flora von Afrika. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*, 22:137–177.
- Schmitz, F., and P. Hauptfleisch
1897. Rhodymeniaceae. In H.G.A. Engler and K.A.E. Prantl, *Die natürlichen Pflanzenfamilien* . . . , 1, 2(149–150):396–405, 4 figures. Leipzig: Wilhelm Engelmann.
- Segawa, S.
1941. Systematic Anatomy of the Articulated Corallines (VI). *The Journal of Japanese Botany*, 17(8):450–456, 4 figures.
1974. *Coloured Illustrations of the Seaweeds of Japan* [Gen shoku Nihon Kaijou Zukan]. 175 pages, 72 figures + 12 plates. Tokyo: Hoikusha Publ. Co.
- Segi, T.
1951. Systematic Study of the Genus *Polysiphonia* from Japan and Its Vicinity. *Journal of the Faculty of Fisheries, Prefectural University, Mie*, 1(2):169–272.
- Setchell, W.A.
1914. The *Scinaia* Assemblage. *University of California Publications in Botany*, 6:79–152, 6 plates.
1924. American Samoa, Part I: Vegetation of Tutuila Island. *Carnegie Institution of Washington Publication*, 20:i–vi + 1–275 [Algae, 28–46; 55–57], 37 plates.
1925. Notes on *Microdictyon*. *University of California Publications in Botany*, 13:101–107.
1926. Tahitian Algae Collected by W.A. Setchell, C.B. Setchell, and H.E. Parks. *University of California Publications in Botany*, 12(5):61–142, 16 plates.
1931. Hong Kong Seaweeds, II. *Hong Kong Naturalist*, 2(4):237–253.
1943. Mastophora and the Mastophoreae: Genus and Subfamily of the Corallinaceae. *Proceedings of the National Academy of Sciences*, 29(5):127–135.
- Setchell, W.A., and N.L. Gardner
1920. Phycological Contributions, I. *University of California Publications in Botany*, 7:279–324, 11 plates.
- Shang, Y.C.
1976. Economic Aspects of *Gracilaria* Culture in Taiwan. *Aquaculture*, 8:1–7.
- Sheath, R.G., and K.M. Cole
1984. Systematics of *Bangia* (Rhodophyta) in North America, I: Biogeographic Trends in Morphology. *Phycologia*, 23:383–396.
- Shen, Y.F., and K.C. Fan
1950. Marine Algae of Formosa. *Taiwania*, 1(2–4):317–345.
- Silva, P.C.
1952. A Review of Nomenclatural Conservation in the Algae from the Point of View of the Type Method. *University of California Publications in Botany*, 25(4):241–324.
- Solier, A.J.J.
1847. Sur deux algues zoosporées formant le nouveau genre *Derbesia*. *Annales des Sciences Naturelles* (Botany), 7:157–166.
- Solms-Laubach, H.
1893. Über die Algengenera *Cymopolia*, *Neomeris* und *Bornetella*. *Annales du Jardin Botanique de Buitenzorg*, 11:61–97, 2 plates.
1895. Monograph of the Acetabulariae. *Transactions of the Linnean Society of London*, 2nd series, Botany, 2:1–39.
- Sonder, O.G.
1845. Nova algarum genera et species, quas in itinere ad oras occidentales Novae Hollandiae, collegit L. Preiss. *Botanische Zeitung*, 3:49–57.
1871. Die Algen des tropischen Australiens. *Abhandlungen aus dem Gebiete der Naturwissenschaften* (Hamburg), 5(2):35–74, 6 plates.
- Stackhouse, J.
1797. *Nereis Britannica* . . . Fascicle 2, ix–xxiv + pages 31–70, plates 9–13.
- Stafleu, F.A., and R.S. Cowan
1976. *Taxonomic Literature* . . . Volume I: A–G, xl + 1136 pages. Utrecht: Bohn, Scheltema & Holkema.
1979. *Taxonomic Literature* . . . Volume II: H–Le, xviii + 991 pages. Utrecht: Bohn, Scheltema & Holkema.
- Su, Y.C., J.H. Huang, Y.C. Lin, C.Y. Chen, and F.C. Chen
1982. Pharmacological Properties of Some Taiwan Seaweeds. In R.T. Tsuda and Y.M. Chiang, editors, *Proceedings of Republic of China–United States Seminar on Cultivation and Utilization of Economic Algae*, pages 51–53. Mangilao: University of Guam Marine Laboratory.
- Suringar, W.F.R.
1870. *Algae Japonicae musei botanici Lugduno-Batavia*. Pages i–viii + 1–39, 25 plates. Haarlem: Typis Heredum Loosjes.
- Tanaka, T.
1935. Four New Species of *Galaxaura* from Japan. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1(1):52–57.
1936. The Genus *Galaxaura* from Japan. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1(2):141–174, 41 figures, 12 plates.
1941. The Genus *Hypnea* from Japan. *Scientific Papers of the Institute of*

- Algological Research, Faculty of Science, Hokkaido Imperial University*, 2(2):227–250.
1944. The Japanese Species of Protoflorideae. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 3:79–97, 16 figures.
1950. On the Species of *Bangia* from Japan. *The Botanical Magazine (Tokyo)*, 63:163–169, 3 figures.
1951. The Species of *Erythrotrichia* from Japan (I). *Acta Phytotaxonomica et Geobotanica*, 14:96–100, 1 figure.
1952. The Systematic Study of the Japanese Protoflorideae. *Memoirs of the Faculty of Fisheries, Kagoshima University*, 2:1–92, plates 1–23.
1967. Some Marine Algae from Bataan and Lamiguru Islands, N. Philippines, I. *Memoires of the Faculty of Fisheries, Kagoshima University*, 16:13–27, 10 figures, 2 plates.
- Tanaka, T., and H. Itono
1969. Studies on the Genus *Neurymenia* (Rhodonelaceae) from Southern Japan and Vicinities. *Memoirs of the Faculty of Fisheries, Kagoshima University*, 18:7–27.
- Tanaka, T., and K. Nozawa
1962. Some Notes on the Genera *Padina* and *Zonaria* in the Southwestern Islands of Japan. *Memoirs of the Faculty of Fisheries, Kagoshima University*, 11:179–187.
- Taniguti, M.
1976. Phytosociological Study of Marine Algae in Taiwan. *Bulletin of the Mie University*, 27:51–57.
- Taylor, W.R.
1945. Pacific Marine Algae of the Allan Hancock Expeditions to the Galapagos Islands. *Allan Hancock Pacific Expeditions*, 12:iv + 528 pages.
1960. *Marine Algae of Tropical and Subtropical Coasts of the Americas*. ix + 870 pages. Ann Arbor: University of Michigan Press.
1962. Two Undescribed Species of *Halimeda*. *Bulletin of the Torrey Botanical Club*, 89(3):172–177.
1966. Records of Asian and Western Pacific Marine Algae, Particularly Algae from Indonesia and the Philippines. *Pacific Science*, 20:342–359.
- Tilden, J.E.
1929. The Marine and Freshwater Algae of China. *Lignan Science Journal*, 7:349–399.
- Tokida, J.
1939. On Some Edible Seaweeds Utilized among a Native Race "Ami" of Formosa, Collected by Mr. N. Nakanome. *Botany and Zoology (Tokyo)*, 7:1547–1552.
1941. On Some Little Known Marine Algae of Japan, 2. *Botany and Zoology (Tokyo)*, 9:49–56.
1954. The Marine Algae of Southern Saghalien. *Memoirs of the Faculty of Fisheries, Hokkaido University*, 2(1):1–264, 15 plates.
- Trevisan, V.B.A.
1845. *Nomenclator algarum, ou collection des noms imposés aux plantes de la famille des algues*. Volume 1, 80 pages. Padua.
- Tseng, C.K.
1936. Studies of the Marine Chlorophyceae from Hainan. *Amoy Marine Biology Bulletin*, 1(5):129–200.
- 1941a. Studies on the Chinese Species of *Liagora*. *Bulletin of the Fan Institute of Biology*, Botanical series, 10:265–281.
- 1941b. Studies on the Chaetangiaceae of China. *Bulletin of the Fan Institute of Biology*, Botanical series, 11:83–118.
1944. Marine Algae of Hong Kong, VI: Genus *Polysiphonia*. *Papers of the Michigan Academy of Science, Arts and Letters*, 29:67–82.
1945. New and Unrecorded Marine Algae of Hong Kong. *Papers of the Michigan Academy of Science, Arts and Letters*, 30:157–172.
- 1981a. Marine Phycoculture in China. In T. Levring, editor, *Proceedings of the Xth International Seaweed Symposium*, pages 123–152. New York: Walter de Gruyter & Co.
- 1981b. Commercial Cultivation. In C.S. Lobban and M.J. Wynne, editors, *The Biology of Seaweeds*, pages 680–725. Berkeley: University of California Press.
- Tseng, C.K., editor
1983. *Common Seaweeds of China*. x + 316 pages. Beijing: Science Press.
- Tseng, C.K., and C.F. Chang
1961. On the Botanical Names of Economic Marine Algae in Old Chinese Literature. *Acta Botanica Sinica*, 9:316–336.
- Tseng, C.K., and T.J. Chang
1962. *Economic Seaweed Flora of China*. vii + 193 pages, 10 plates. Beijing: Science Press.
- Tseng, C.K., Y. Yoshida, and Y.M. Chiang
1985. East Asiatic Species of *Sargassum* subgenus *Bactrophycus* J. Agardh (Sargassaceae, Fucales), with Keys to the Sections and Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds . . .*, pages 1–14. La Jolla, California: California Sea Grant College Program, University of California, La Jolla.
- Tseng, W.Y.
1984. *Eucheuma* of Taiwan—Emphasizing Its Mariculture Potential. *Hydrobiologia*, 116/117:237–242.
- Tseng, W.Y., and C.S. Chen
1977. *Eucheuma* of Taiwan. *Bulletin of the Taiwan Fisheries Research Institute*, 28:103–112.
- Ueda, S.
1932. A Systematic Study of Genus *Porphyra* Found on the Japanese Coasts. *Journal of the Imperial Fisheries Institute (Tokyo)*, 28:1–45.
- Umezaki, I.
1972. The Life Histories of Some Nemaliales Whose Tetrasporophytes Were Unknown. In I.A. Abbott and M. Kurogi, editors, *Contributions to the Systematics of Benthic Algae of the North Pacific*, pages 231–242. Kobe, Japan: Japanese Society of Phycology.
- Vickers, A.
1905. Liste des algues marines du Barbade. *Annales des Sciences Naturelles*, series 9, 1:45–66.
- Voss, E.G., et al.
1983. *International Code of Botanical Nomenclature*. [Adopted by the Thirteenth International Botanical Congress, Sydney, August 1981.] xv + 472 pages. Utrecht: Bohn, Scheltema & Holkema.
- Wang, J.C., and Y.M. Chiang
1977. Notes on Marine Algae of Taiwan, II: The Genus *Porphyra*. *Taiwania*, 22:105–112.
- Weber-van Bosse, A.
1898. Monographie des Caulerpes. *Annales du Jardin Botanique de Buitenzorg*, 15(2):243–401, 14 plates.
- 1913a. Liste des algues du Siboga, I: Myxophyceae, Chlorophyceae, Phaeophyceae. In M. Weber, editor, *Siboga-Expedition: Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebied . . .*, 59a:1–186, plates 1–52. Leiden: E.J. Brill.
- 1913b. Marine Algae, *Rhodophyceae* of the "Sealark" Expedition. *Transactions of the Linnean Society, London*, series 2 (Botany), 8:105–142, plates 12–14.
1921. Liste des algues du Siboga, II: Rhodophyceae, Part 1: Protoflorideae, Nemalionales, Cryptonemiales. In M. Weber, editor, *Siboga-Expedition: Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebied . . .*, 59b:187–310, plates 6–8. Leiden: E.J. Brill.
1928. Liste des algues du Siboga, Part IV: Rhodophyceae, Part 3: Gigartinales et Rhodymeniales. In M. Weber, editor, *Siboga-Expedition: Uitkomsten op Zoologisch, Botanisch, Oceanographisch en Geologisch Gebied . . .*, 59d:393–533, plates 11–16. Leiden: E.J. Brill.
- West, J.A., and M.H. Hommersand
1981. Rhodophyta: Life Histories. In C.S. Lobband and M.J. Wynne, editors, *The Biology of Seaweeds*, pages 133–193. Berkeley: University of California Press.

- Wiseman, D.R.
- 1975. On the Status of the Red Algal Family, the Rhizophyllidaceae (Gigartinales). *Taxon*, 24:489-490.
- Wittrock, V.B.
- 1866. *Försök till en monographie öfver alsgläget Monostroma*. 66 pages, 4 plates. Stockholm.
- Womersley, H.B.S.
- 1967. A Critical Survey of the Marine Algae of Southern Australia, II: Phaeophyta. *Australian Journal of Botany*, 15:189-270.
 - 1979. Southern Australian Species of *Ceramium* Roth (Rhodophyta). *Australian Journal of Marine and Freshwater Research*, 29:205-257.
- Womersley, H.B.S., and A. Bailey
- 1970. Marine Algae of the Solomon Islands. *Philosophical Transactions of the Royal Society of London*, series B (Biological Sciences), 259(830):257-352.
- Wu, C.S.
- 1982. Chemical Analysis and Utilization of *Monostroma latissimum*. In R.T. Tsuda and Y.M. Chiang, editors, *Proceedings of the Republic of China—United States Seminar on Cultivation and Utilization of Economic Algae*, pages 55-60. Mangilao: University of Guam Marine Laboratory.
- Wynne, M.J.
- 1985. Nomenclature Assessment of *Goniotrichum* Kützing, *Erythrotricha* Areschoug, *Diconia* Harvey, and *Stylonema* Reinsch (Rhodophyta). *Taxon*, 34:502-505.
- Wynne, M.J., and G.T. Kraft
- 1981. Appendix: Classification Summary. In C.S. Lobban and M.J. Wynne, editors, *The Biology of Seaweeds*, pages 743-750. Berkeley: University of California Press.
- Xia, B.M.
- 1985. *Gracilaria* from China: Key, List and Distribution of the Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds*, pages 71-76. La Jolla: California Sea Grant College Program, University of California, La Jolla.
- Xia, B.M., and H. Yamamoto
- 1985. *Gracilaria* Species from Both China and Japan: Key, List, and Distribution of Common and Economically Important Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds*, pages 69-70. La Jolla: California Sea Grant College Program, University of California, La Jolla.
- Xia, B.M., and J.F. Zhang
- 1982. A New Raw Material for the Manufacture of Agar-agar from Hainan Island, China. *Oceanologica et Limnologia Sinica*, 13(6):538-543, plate 1.
- Yamada, Y.
- 1925a. Studien über die Meeresalgen von der Insel Formosa, I: Chlorophyceae. *The Botanical Magazine (Tokyo)*, 39:77-95, 5 figures.
 - 1925b. Studien über die Meeresalgen von der Insel Formosa, II: Phaeophyceae. *The Botanical Magazine (Tokyo)*, 39:239-254, 6 figures.
 - 1926. The Phylogeographical Relation between the Chlorophyceae of the Mariannes, Carolines and Marshall Islands and Those of the Malay Archipelago, Australia and Japan. *Proceedings of the Third Pan-Pacific Science Congress*. Pages 964-966. Tokyo.
 - 1928. Report of the Biological Survey of Mutsu Bay, 9: Marine Algae of Mutsu Bay and Adjacent Waters, II. *Science Report, Tohoku Imperial University*, 4th series (Biology), 3:497-534.
 - 1930. Notes on Some Japanese Algae, I. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1:27-36.
 - 1931a. Notes on Some Japanese Algae, II. *Journal of the Faculty of Science, Hokkaido Imperial University*, series V, 1(2):65-76, 3 figures, 5 plates.
 - 1931b. Notes on *Laurencia*, with Special Reference to the Japanese Species. *University of California Publications in Botany*, 16:185-310, 20 figures, 30 plates.
 - 1932a. Notes on Some Japanese Algae, III. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1:109-123.
 - 1932b. Notes on Some Japanese Algae, IV. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1:267-276.
 - 1933. Notes on Some Japanese Algae, V. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 2:277-285.
 - 1934. The Marine Chlorophyceae from Ryukyu, Especially from the Vicinity of Nawa. *Journal of the Faculty of Science, Hokkaido Imperial University*, series V, 3(2):33-88, 55 figures.
 - 1935. Notes on Some Japanese Algae, VI. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1(1):27-35.
 - 1936a. The Species of *Eucheuma* from Ryukyu and Formosa. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1(2):119-134, 9 plates.
 - 1936b. Notes on Some Japanese Algae, VII. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 1(2):135-140, 3 figures, 4 plates.
 - 1936c. [Laurencia.] In K. Okamura, *Nippon Kaiyō-shi* [Marine Algae of Japan], pages 851-861. Tokyo: Published by the author.
 - 1938a. The Species of *Liagora* from Japan. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 2:1-34, 22 figures, 15 plates.
 - 1938b. Notes on Some Japanese Algae, VIII. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 2:119-130.
 - 1941. Notes on Some Japanese Algae, IX. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 2:195-216, 9 plates.
 - 1942. Notes on *Sargassum* from the Southern Parts of Japan, I, II, III. *Japanese Journal of Botany*, 18:369-381, 504-519, 553-562.
 - 1944a. Diagnoses of New Sargassums from Japan. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 3:1-10.
 - 1944b. Notes on Some Japanese Algae, X. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 3:11-25, 18 figures.
 - 1944c. A List of the Marine Algae from the Atoll of Ant. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 3:31-45.
 - 1950. A List of Marine Algae from Ryukyusho, Formosa, I: Chlorophyceae and Phaeophyceae. *Scientific Papers of the Institute of Algological Research, Faculty of Science, Hokkaido Imperial University*, 3:173-194.
 - Yamada, Y., and S. Segawa
 - 1953. On Some New or Noteworthy Algae from Hachijo Island. *Records of Oceanographic Works in Japan*, new series, 1:109-114.
 - Yamada, Y., and T. Tanaka
 - 1934. Three New Red Algae from Formosa. *Transactions of the Natural History Society of Formosa*, 24:342-349, 5 figures.
 - Yamamoto, H.
 - 1985. *Gracilaria* from Japan: Vegetative and Reproductive Keys and List of the Species. In I.A. Abbott and J.N. Norris, editors, *Taxonomy of Economic Seaweeds*, pages 77-80. La Jolla: California Sea Grant College Program, University of California, La Jolla.
 - Yang, H.N., and Y.M. Chiang
 - 1982. Taxonomical Study on the *Gracilaria* of Taiwan. *Journal of the Fisheries Society of Taiwan*, 9:55-71.

- Yang, S.S., C.Y. Wang, and H.H. Wang
1981. Seasonal Variation of Agar-agar Produced in Taiwan Area. In T. Levring, editor, *Proceedings of the Xth International Seaweed Symposium*, pages 737–742. New York: Walter de Gruyter & Co.
- Yendo, K.
1902. Corallinae verae Japonicae. *Journal of the College of Science, Imperial University*, 16:1–36.
1905. Preliminary List of the Fucaceae of Japan. *The Botanical Magazine (Tokyo)*, 19(222):149–161.
1907. The Fucaceae of Japan. *Journal of the College of Science, Imperial University*, 21:1–174, 8 plates.
1909. Notes on Algae New to Japan, I. *The Botanical Magazine (Tokyo)*, 23:117–133.
1914. Notes on Algae New to Japan, II. *The Botanical Magazine (Tokyo)*, 28:263–281, 1 figure.
1915. Notes on Algae New to Japan, III. *The Botanical Magazine (Tokyo)*, 29:99–117.
1916a. Notes on Algae New to Japan, IV. *The Botanical Magazine (Tokyo)*, 30:47–65.
1916b. Notes on Algae New to Japan, V. *The Botanical Magazine (Tokyo)*, 30:243–263.
1918. Notes on Algae New to Japan, VIII. *The Botanical Magazine (Tokyo)*, 32:65–81.
1920. Novae algae Japoniae, Decas I–III. *The Botanical Magazine (Tokyo)*, 34:1–12.
- Yoshikawa, H., and K. Yoshikawa
1977. A List of Marine Algae Collected at Maopit'ou, Taiwan. *Joshigaku yo Daigaku Kiyo* [Records of the Women's Home Economics College], 8:127–130.
- Zanardini, J.
1851. Algae novae vel minus cognitae in mari rubro a Portiero collectae. In *Flora oder Allgemeine Botanische Zeitung*, Jena, 34(3):33–35.
1858. Plantarum in mari rubro hucusque collectarum enumeratio. *Memorie della Istituto Veneto di Scienze, Lettere ed Arti*, 7:209–309.
1872. Phycearum indicarum pugillus. *Memorie della Istituto Veneto di Scienze, Lettere ed Arti*, 17:129–170.
1878. Phyceae Papuanae Novae vel minus cognitae a Cl. O. Beccari in itinere ad Novam Guineam annis 1872–75 Collectae. *Nuovo Giornale Botanico Italiano*, 10(1):34–40.
- Zhang, J.F. [Chang, C.F.] and B.M. Xia
1985. On *Gracilaria asiatica* sp. nov. and *G. verrucosa* (Hudson) Papenfuss. *Oceanologia et Limnologica Sinica*, 16(3):175–180.

Index

- Acanthophora
aokii, 22
muscoides, 22
orientalis, 2, 22
- Acetabularia
gigas, 11
major, 11
- Acetabulariaceae, 11
- "*Acrocarpus pusillus*," 17
- Acrochaetiaceae, 15
- Acrocystis nana*, 22
- Actinotrichia *fragilis*, 15
 "rigida," 15
- Ahnfeltia *paradoxa*, 21
- Amansia *glomerata*, 22
- Amphiroa
beauvoisii, 18
bowerbankii, 18
"cultrata var. globulifera", 18
ephedraceae, 18
"exilis", 18
fragilissima, 18
multifida, 18
pusilla, 18
"pusilla", 18
"zonata", 18
 spp., 18
- Anadyomenaceae, 11
- Anadyomene *wrightii*, 11
- Asparagopsis
"sanfordiana", 16
taxiformis, 16
- "Asteroecytis"
"ornata", 14
"ornata f. simplex", 14
"ramosa f. simplex", 14
- Bangia
atropurpurea, 14
"fuscopurpurea", 14
yamadae, 14
'yamadai', 14
 sp., 14
- Bangiaceae, 14
- Bangiales, 14
- Boergesenia *forbesii*, 10
- Bonnemaisoniaceae, 16
- Boodlea
"bosseae", 10
coacta, 10
composita, 10
montagnei, 10
"paradoxa", 10
siamensis, 10
van-bosseae, 10
'van-Bossei', 10
- Bornetella, 4
sphaerica, 11
- Bostrychia *tenella*, 11
- Bryopsidaceae, 10
- Bryopsis
harveyana, 10
indica, 10
mucosa, 10
plumosa, 10
- Caloglossa *bombayensis*, 22
- Carpolepharis
schmitziana var. *erecta*, 21
warburgii, 21
- Carpopeltis
angusta, 18
cornea, 18
flabellata, 19
formosana, 2, 19
rigida, 19
- Caulacanthus
"fastigiatus", 21
okamurae, 21
'okamurai', 21
spinellus, 21
ustulatus var. *fastigiatus*, 1, 21
- Caulerpa
cupressoides var. *lycopodium f. amicorum*, 9
freycinetti var. *freycinetti f. lata*, 9
"freycinetti var. typica", 9
"peltata", 9
racemosa, 9
"racemosa var. chenmunitzia", 9
"racemosa var. clavifera", 9
racemosa var. *clavifera f. macrophysa*, 9
racemosa var. *clavifera f. microphysa*, 9
racemosa var. *laetevirens*, 9
racemosa var. *occidentalis*, 9
racemosa var. *peltata*, 9
racemosa var. *turbanata*, 9
serrulata, 9
serrulata f. lata, 9
sertularioides, 9
sertularioides f. longipes, 9
taxifolia, 9
webbiana, 9
webbiana f. disticha, 9
webbiana f. tomentella, 9
- Caulerpaceae, 9
- Caulerpales, 8
- Centroceras
clavulatum, 21
minutum, 21
- Ceramiaceae, 21
- Ceramiales, 21
- Ceramium
aduncum, 21
ciliatum var. *robustum*, 21
"clarionense", 21
- "clavulatum," 21
 "equisetoides," 22
flaccidium, 21
gracillimum, 22
"gracillimum var. byssoideum", 21
nakamurai, 22
paniculatum, 22
tenerimum, 22
tenuissimum, 22
- Ceratodictyon
spongiosum, 20
 "spongioides," 20
- Chaetangiaceae, 15
- Chaetomorpha
aerea, 8
aerea f. versata, 8
basiretrorsa, 8
brachygona, 8
crassa, 8
linum, 8
spiralis, 8
- Chaetophoraceae, 7
- Chaetophorales, 7
- Chamaedoris *orientalis*, 10
- Champia *parvula*, 21
- Champiaceae, 21
- Cheilosporum
anceps, 18
 "cultratum," 18
jungermanniooides, 18
- Chlorella, 4
- Chlorodesmis
caespitosa, 9
comosa, 9
 "formosana," 9
- Chlorophyta, 1, 7
- Chnoospora
implexa, 13
minima, 13
 "pacifica," 13
- Chondria
armata, 2, 22
dasyphylla, 22
- Chondrococcus
hornemannii, 17
'hornemannii', 17
- Chondrus
"affinis", 21
crispus, 21
ocellatus, 21
ocellatus f. canaliculatus, 21
 "ocellatus f. typicus," 21
- Chordariales, 11
- Chroodactylon *ornatum*, 14
- Chrysomenia *procumbens*, 21
- Cladophora
aokii, 8
fuliginosa, 8

montagnei var. *radicans*, 8
patentiramea, 8
pellucida, 8
 "prolifera," 8
rugulosa, 8
scitula, 8
sibogae, 8
 sp., 8
 Cladophoraceae, 8
 Cladophorales, 8
 Cladophoropsis
 "fasciculatus," 10
herpestica, 10
sundanensis, 10
zollingeri, 10
Claudea bataensis, 22
 Codiaceae, 8
Codium
 adhaerens, 8
arabicum, 8
contractum, 8
cylindricum, 8
dichotomum, 8
formosanum, 8
fragile, 8
intricatum, 9
 "mucronatum," 8
reediae, 9
repens, 9
tenue, 9
 "tomentosum," 9
Colpomenia sinuosa, 13
 "Conchocelis," 4
 "Corallina"
 "adhaerens," 18
 "decussato-dichotoma," 18
 Corallinaceae, 2, 18
 "Corallopsis opuntia," 20
 Corynothlaeacea, 11
 Cryptonemiaceae, 18
 Cryptonemiales, 17
Cystoseira
 prolifera, 13
 sp., 13
 Cystoseiraceae, 13

Dasya sp., 22
 Dasyaceae, 22
Dasycladales, 11
 Delesseriaceae, 22
Dasyphila plumarioides, 22
Delisea
 fimbriata, 16
 "japonica," 16
Derbesia lamourouxii, 10
 Derbesiaceae, 10
 "Dermatolithon tumidulum," 18
Dermonema
 frappieri, 15
 gracile, 15
dichotoma, 2
 "dichotomum," 15
pulvinata, 15
 Dermonemataceae, 15
 "Desmia hornemannii," 17

Dictyopteris
delicatula, 11
repens, 11
undulata, 11
Dictyosphaeria
bokokensis, 10
cavernosa, 10
 "favulosa," 10
Dictyota
bartayresii, 11
ceylanica var. *anastomosans*, 12
dichotoma, 12
dilata, 12
divaricata, 12
hamifera, 12
linearis, 12
patens, 12
spinulosa, 12
Dictyotaceae, 11
Dictyotales, 11
Digenia simplex, 22
Dilophus
okamurae, 12
 'okamurai,' 12
 "marginatus," 12
radicans, 12
Dotyophycus yamadae, 15
Dudresnaya japonica, 17
Dumontiaceae, 17

Ectocarpaceae, 11
Ectocarpales, 11
Ectocarpus
breviarticulatus, 11
 "formosanus," 11
laurenciae, 11
 "mitchelliae," 11
spinosus, 11
vanbossea, 11
van-bossea, 11
 sp., 11
Endarachne binghamiae, 13
 "Endocladia complanata," 18
Endocladiaeae, 18
Endophyton ramosum, 7
Enteromorpha
clathrata, 7
clathrata var. *crinita*, 7
 "crinita," 7
compressa, 7
intestinalis, 7
linza, 7
prolifera, 7
 sp., 7
Erythrocolon podagraria, 21
Erythropeltidaceae, 14
Erythrotrichia
biseriata, 14
carnea f. *tenuis*, 14
Eucheuma
arnoldii, 19
 "audiolis," 19
cottonii, 19
 "cottonii," 19
crassum, 19

crustaeforme, 19
 "cupressoideum," 19
gelatinae, 19
muricatum, 19
 "muricatum," 19
muricatum f. *depauperata*, 19
okamurae, 19
 'okamurai,' 19
papulosa, 19
serra, 19
 "spinosum," 19
 spp., 19

Feldmannia formosana, 11
Fucales, 13

Galaxaura, 2
arborea, 15
canaliculata, 16
clavigera, 16
distenta, 16
elegans, 16
elongata, 16
falcata, 16
fasciculata, 16
fastigiata, 16
filamentosa, 16
lapidescens, 16
lapidescens f. *villosa*, 16
latifolia, 16
marginata, 16
oblongata, 16
obtusata, 16
pacifica, 16
robusta, 16
rudis, 16
rugosa, 16
 "tenera," 16
veprecula, 16
Ganonema, 15
Gelidiaceae, 17
Gelidiella
acerosa, 17
Gelidiellaceae, 17
Gelidiopsis
hachijoensis, 20
repens, 20
 "rigida," 17
variabilis, 20
Gelidium, 2, 3, 4
amansii, 17
amansii f. *elegans*, 17
amansii f. *latioris*, 17
 "cartilagineum," 17
 "clavatum," 17
 "corneum" var. *pinnatum*, 17
crinale, 17
 "densum," 17
divaricatum, 17
japonicum, 17
kintaroi, 17
latiusculum, 17
pacificum, 17
planiusculum, 17
pusillum, 17

pusillum f. foliaceum, 17
subcostatum, 17
yamadae, 17
 sp., 17

Giffordia mitchelliae, 11

Gigartina
intermedia, 21
ochotensis, 21
tenella, 21

Gigartinaceae, 21

Gigartinales, 19

Gloiopektis

cervicornis, 18
complanata, 18
furcata, 18
tenax, 18

Coniotrichaceae, 14

Coniotrichales, 14

"*Goniotrichum alsidii*", 14

Gracilaria, 4

arcuata, 20
blodgettii, 20
bursapastoris, 20
canaliculata, 20
chorda, 20
 "compressa", 20
 "confervoides", 20
coronopifolia, 20
crassa, 20
denticulata, 20
edulis, 4, 20
eucheumoides, 20
gigas, 4, 20
incurvata, 20
 "lichenoides", 20
 "lichenoides f. coronopifolia", 20
punctata, 20
purpurascens, 20
 "purpurascens f. spinulosa", 20
salicornia, 20
spinulosa, 20
 "verrucosa", 4, 20
 sp., 21

Graciliariaceae, 20

Gratelouphia
carnosa, 19
filicina, 19
filicina f. filiformis, 1, 19
 "filiformis", 19
okamurae, 19
 "okamurae", 19
ramosissima, 19

Gymnogongrus
flabelliformis, 21
 "paradoxus", 21

"*Gymnosorus collaris*", 12

Gymnothamnion elegans, 22

Halimeda

cuneata, 9
discoidea, 9
incrassata var. *ovata*, 9
macroloba, 9
opuntia, 9
 "opuntia f. intermedia", 9

"*opuntia f. renschii*", 9
renschii, 9
velasquezii, 9
 "Halisseris undulata", 11
Haloplegma duperreyi, 22
Halymenia
ceylanica, 19
durvillei, 19
 "durvillei var. *formosa*", 2, 19
 "durvillei var. *ceylanica*", 19
 "formosa", 19

Helminthocladia australis, 15

Herposiphonia
subdisticha, 22
 sp., 22

Holmesia neurymenioides, 22

"*Homoeostrichus*"
flabellatus, 12
multifidus, 12

Hormophysa triquetra, 13

Hydroclathrus
cancellatus, 13
clathratus, 13
orientalis, 13

Hypnea, 2

boergesenii, 19
cenomyce, 19
cervicornis, 19
charoides, 19
chordacea, 19
chordacea f. simpliciuscula, 19
cornuta, 19
esperi, 19
hamulosa, 19
japonica, 20
nidulans, 20
pannosa, 20
saidana, 20
seticulosa, 2, 20
spinella, 20
 sp., 20

Hypnaceae, 19

"*Ilea fascia*", 12

Ishige
okamurae, 12
 'okamurae', 12
simicola, 12
 "foliacea", 12

Ishigeaceae, 12

Jania

adhaerens, 18
 "decussato-dichotoma", 18
radiata, 18
tenella, 18
undulata, 18

Laurencia, 2

brongniartii, ii, 22
 "dasyphylla", 22
flexilis var. *tropica*, 23
forsteri, 23
glandulifera, 23

grevilleana, 23
obtusa var. *densa*, 23
palizada, 23
papillosa, 23
perforata, 23
pinnatifida, 23
 "pinnatifida" var. *simplex*, 23
tropica, 23
venusta, 23
 sp., 2, 23

"*Letterstedtia japonica*", 8

Leveillea jungermannioides, 23

Liagora, 2, 4

"*annulata*", 15
boergesenii, 15
 "caenomyce", 15
ceranoides, 15
ceranoides var. *leprosa*, 15
ceranoides var. *pulverulenta*, 15
decussata, 15
farinosa, 15
 "formosana", 15
 "maxima", 15
orientalis, 15
 "orientalis", 15
rugosa, 15
segawai, 15
setchellii, 15
valida, 15

Liagoraceae, 15

Liagorophila endophytica, 15

Liagoropsis schrammi, 15

Lithophyllum
perulatum, 18
 spp., 18

Lithothamnium

membranaceum, 18
 spp., 18

Lobophora variegata, 12

Martlesia

denticulata, 22
flabelliformis, 22

Mastophora

pygmaea, 18
rosea, 18
 "macrocarpa", 18

Melobesia

farinosa, 18
 "membranaceae", 18
 "pustulata", 18
 spp., 18

Meristotheca

coacta, 19
papulosa, 19

Microcladia elegans, 22

Microdictyon

japonicum, 10
nigrescens, 10
okamurae, 10

Monostroma, 4
latisimum, 7
nitidum, 8

- Murrayella
periclados, 23
squarrosa, 23
- Nemaliales, 15
"Nemalion pulvinatum," 2, 15
- Neomeris, 4
annulata, 11
- "Neurocarpus"
"repens," 11
"undulata," 11
"undulata f. plana," 11
- Neurymenia fraxinifolia, 23
- Nitophyllum sp., 22
- Nostoc, 3
- Pachydictyon coriaceum, 12
- Padina
arborescens, 12
australis, 12
boryana, 12
commersonii, 12
"commersonii," 12
crassa, 12
durvillaei, 12
japonica, 12
minor, 12
"pavonia," 12
tenuis, 12
- Petalonia fascia, 12
- Peyssonnelia
caulifera, 18
distenta, 18
"involvens," 18
rubra, 18
- Peyssonneliaceae, 18
- Phacelocarpus japonicus, 21
- Phaeophyta, 1, 11
"Phycoseris"
"lanceolata var. angusta," 7
"lapathifolia," 8
- "Phyllitis fascia," 12
- Phyllophoraceae, 21
- "Placophora marchantioides," 23
- Plocamiaceae, 20
- Plocamium
"abnorme," 20
"abnorme f. uncinatum," 20
"costatum," 20
oviforme, 20
serratum, 20
telfairiae, 20
telfairiae f. uncinatum, 20
- "Plumaria ramosa," 22
- Polyopales
polyideoides, 19
sp., 19
- Polysiphonia
harlandii, 23
kampsaxii, 23
pulvinata, 23
- "Polyzonaria jungermannioides," 23
- Porphyra, 2, 4
angusta, 14
crispata, 14
- dentata, 15
suborbiculata, 15
sp., 15
- Pterocladia, 2, 3
"capillaceum," 17
nana, 17
tenuis, 17
- Ptilothamnion cladophorae, 22
- Punctariaceae, 12
- Reinboldiella schmitziana, 22
- "Rhipidiphyllon"
"reticulatum," 10
"nigrescens," 10
- Rhizophyllidaceae, 17
- Rhodoglossum affine, 21
- Rhodomelaceae, 22
- Rhodopeltis
borealis, 17
gracilis, 17
setchelliae, 17
'setchelii,' 17
- Rhodophyta, 1, 14
- Rhodymenia spinulosa, 21
- Rhodymeniaceae, 21
- Rhodymeniales, 21
- Rosenvingea orientalis, 13
- Sarcodiaceae, 21
- Sarcodia ceylanica, 21
- Sargassaceae, 13
- Sargassum, 2, 3, 4
amabile, 13
aquifolium, 13
baccularia, 13
berberifolium, 13
binderi, 13
"biserrula," 13
carpophyllum, 13
coriifolium, 13
coriifolium f. dupicum, 13
coriifolium f. prolongatum, 13
crassifolium, 13
crispifolium, 13
cristae folium, 13
duplicatum, 4, 13
echinocarpum, 13
fulvellum, 13
glaucescens, 13
hemiphyllum, 13
henslowianum, 13
heterocystum, 13
horneri, 13
ilicifolium, 13
ilicifolium var. conduplicatum, 13
kasyotense, 14
kuetzingii, 14
kushimotense, 14
"microphyllum," 14
muriocystum, 14
plagiophyllum, 14
polycystum, 14
"prolongatum," 13
rostratum, 14
- sandeii, 14
sandeii f. heterophyllum, 14
- serratifolium, 14
- siliquosum, 14
- swartzii, 14
- telephifolium, 14
- tenuifolium, 14
- vulgare, 14
- sp., 14
- Scinaia
boergesenii, 16
cottonii, 16
moniliformis, 16
pseudojaponica, 16
- Scytoniphon
lomentaria, 13
'lomentarium,' 13
"lomentarius," 13
- Scytoniphonaceae, 13
- Scytoniphonales, 12
- Siphonocladaceae, 10
- Siphonocladales, 10
- Solieriaceae, 19
- Spatoglossum pacificum, 12
- "Spermothamnion"
"cladophorae," 22
"codicola," 22
- Sphaelaria
furcigera var. tenuis, 11
tribuloides, 11
- Sphaelariaceae, 11
- Sphaelariales, 11
- Sphaerococcaceae, 21
- Spyridia filamentosa, 22
- Struvea
anastomosans, 10
delicatula, 10
- Symphyocladia marchantioides, 23
- Tenarea tumidulum, 18
- Tiffaniella codicola, 22
- Turbinaria
conoides, 14
filamentosa, 14
"filiformis," 14
ornata, 14
trialata, 14
sp., 14
- Udoteaceae, 9
- Ulothrix flaccida, 7
- Ulrichaceae, 7
- Ulotrichales, 7
- Ulva
angusta, 8
conglobata, 8
fasciata, 8
japonica, 8
lactuca, 8
lactuca f. lapathifolia, 1, 8
pertusa, 8
reticulata, 8
rigida, 8
sp., 8
- Ulvaceae, 7
- Ulvales, 7

Valonia
 , 11
 "confervoides," 11
 fastigiata, 11
 "forbesii," 10
 utricularis, 11
 verticillata, 11
Valoniaceae, 10
Valoniopsis pachynema, 11

Vidalia obtusiloba, 23
Weberella micans, 21
Wrangelia velutina, 22
Wurdemannia
 setacea, 23
 sp., 23
Wurdemaniaceae, 23
Yamadaella cconomyce, 15

Zonaria
 coriacea, 12
 diesingiana, 12
 flabellata, 12
 harveyana, 12
 nigrescens, 12
 stipitata, 12
 "variegata," 12
 sp., 12

REQUIREMENTS FOR SMITHSONIAN SERIES PUBLICATION

Manuscripts intended for series publication receive substantive review (conducted by their originating Smithsonian museums or offices) and are submitted to the Smithsonian Institution Press with Form SI-36, which must show the approval of the appropriate authority designated by the sponsoring organizational unit. Requests for special treatment—use of color, foldouts, case-bound covers, etc.—require, on the same form, the added approval of the sponsoring authority.

Review of manuscripts and art by the Press for requirements of series format and style, completeness and clarity of copy, and arrangement of all material, as outlined below, will govern, within the judgment of the Press, acceptance or rejection of manuscripts and art.

Copy must be prepared on typewriter or word processor, double-spaced, on one side of standard white bond paper (not erasable), with 1 1/4" margins, submitted as ribbon copy (not carbon or xerox), in loose sheets (not stapled or bound), and accompanied by original art. Minimum acceptable length is 30 pages.

Front matter (preceding the text) should include: title page with only title and author and no other information, abstract page with author, title, series, etc., following the established format; table of contents with indents reflecting the hierarchy of heads in the paper; also, foreword and/or preface, if appropriate.

First page of text should carry the title and author at the top of the page; **second page** should have only the author's name and professional mailing address, to be used as an unnumbered footnote on the first page of printed text.

Center heads of whatever level should be typed with initial caps of major words, with extra space above and below the head, but no other preparation (such as all caps or underline, except for the underline necessary for generic and specific epithets). Run-in paragraph heads should use period/dashes or colons as necessary.

Tabulations within text (lists of data, often in parallel columns) can be typed on the text page where they occur, but they should not contain rules or numbered table captions.

Formal tables (numbered, with captions, boxheads, stubs, rules) should be submitted as carefully typed, double-spaced copy separate from the text; they will be typeset unless otherwise requested. If camera-copy use is anticipated, do not draw rules on manuscript copy.

Taxonomic keys in natural history papers should use the aligned-couplet form for zoology and may use the multi-level indent form for botany. If cross referencing is required between key and text, do not include page references within the key, but number the keyed-out taxa, using the same numbers with their corresponding heads in the text.

Synonymy in zoology must use the short form (taxon, author, year:page), with full reference at the end of the paper under "Literature Cited." For botany, the long form (taxon, author, abbreviated journal or book title, volume, page, year, with no reference in "Literature Cited") is optional.

Text-reference system (author, year:page used within the text, with full citation in "Literature Cited" at the end of the text) must be used in place of bibliographic footnotes in all Contributions Series and is strongly recommended in the Studies Series: "(Jones, 1910:122)" or "... Jones (1910:122)." If bibliographic

footnotes are required, use the short form (author, brief title, page) with the full citation in the bibliography.

Footnotes, when few in number, whether annotative or bibliographic, should be typed on separate sheets and inserted immediately after the text pages on which the references occur. Extensive notes must be gathered together and placed at the end of the text in a notes section.

Bibliography, depending upon use, is termed "Literature Cited," "References," or "Bibliography." Spell out titles of books, articles, journals, and monographic series. For book and article titles use sentence-style capitalization according to the rules of the language employed (exception: capitalize all major words in English). For journal and series titles, capitalize the initial word and all subsequent words except articles, conjunctions, and prepositions. Transliterate languages that use a non-Roman alphabet according to the Library of Congress system. Underline (for italics) titles of journals and series and titles of books that are not part of a series. Use the parentheses/colon system for volume (number); pagination: "10(2):5-9." For alignment and arrangement of elements, follow the format of recent publications in the series for which the manuscript is intended. Guidelines for preparing bibliography may be secured from Series Section, SI Press.

Legends for illustrations must be submitted at the end of the manuscript, with as many legends typed, double-spaced, to a page as convenient.

Illustrations must be submitted as original art (not copies) accompanying, but separate from, the manuscript. Guidelines for preparing art may be secured from Series Section, SI Press. All types of illustrations (photographs, line drawings, maps, etc.) may be intermixed throughout the printed text. They should be termed **Figures** and should be numbered consecutively as they will appear in the monograph. If several illustrations are treated as components of a single composite figure, they should be designated by lowercase italic letters on the illustration; also, in the legend and in text references the italic letters (underlined in copy) should be used: "Figure 9b." Illustrations that are intended to follow the printed text may be termed **Plates**, and any components should be similarly lettered and referenced: "Plate 9b." Keys to any symbols within an illustration should appear on the art rather than in the legend.

Some points of style: Do not use periods after such abbreviations as "mm, ft, USNM, NNE." Spell out numbers "one" through "nine" in expository text, but use digits in all other cases if possible. Use of the metric system of measurement is preferable; where use of the English system is unavoidable, supply metric equivalents in parentheses. Use the decimal system for precise measurements and relationships, common fractions for approximations. Use day/month/year sequence for dates: "9 April 1976." For months in tabular listings or data sections, use three-letter abbreviations with no periods: "Jan, Mar, Jun," etc. Omit space between initials of a personal name: "J.B. Jones."

Arrange and paginate sequentially every sheet of manuscript in the following order: (1) title page, (2) abstract, (3) contents, (4) foreword and/or preface, (5) text, (6) appendixes, (7) notes section, (8) glossary, (9) bibliography, (10) legends, (11) tables. Index copy may be submitted at page proof stage, but plans for an index should be indicated when manuscript is submitted.

