

## TWO INNOVATIONS IN MEXICAN MALVACEAE

Laurence J. Dorr

Department of Botany

National Museum of Natural History, MRC-166

Smithsonian Institution

P.O. Box 37012

Washington, D.C. 20013-7012, U.S.A.

dorr@si.edu

### ABSTRACT

Two innovations in the nomenclature of Mexican Malvaceae are proposed. First, *Ayenia sidifolia* is reestablished as the correct name for the species currently known as *A. mexicana*: this latter name was incorrectly proposed as a replacement name in *Ayenia* for *Cybiostigma sidifolium*. Second, the monotypic genus *Veeseria* endemic to Mexico is placed in synonymy under the Asian genus *Reevesia*, and the new combination *Reevesia clarkii* is proposed.

KEY WORDS: Malvaceae, Mexico, *Ayenia*, *Cybiostigma*, *Reevesia*, *Veeseria*

### RESUMEN

Se proponen dos cambios en la nomenclatura de las Malvaceae mexicanas. En primer lugar, *Ayenia sidifolia* se restablece como el nombre correcto para la especie actualmente conocida como *A. mexicana*: este último nombre fue propuesto incorrectamente como un nombre de reemplazo en *Ayenia* para *Cybiostigma sidifolium*. En segundo lugar, el género monotípico *Veeseria* endémico de México se pone en sinonimia bajo el género asiático *Reevesia* y se propone la nueva combinación *Reevesia clarkii*.

PALABRAS CLAVES: Malvaceae, Mexico, *Ayenia*, *Cybiostigma*, *Reevesia*, *Veeseria*

The recent publication of a checklist of the native vascular plants of Mexico (Villaseñor 2016) has prompted me to propose two changes in the nomenclature of Mexican Malvaceae. The first innovation emanates from a review of the American taxa of vascular plants described by Loefling (Dorr & Wiersema 2010a, b) and especially several of his species of *Ayenia* L. The second innovation is one that came to my attention in the early 1980s, but was set aside then out of deference to a student who undertook a project to revise the genus *Reevesia* Lindl. (Solheim 1991).

### Reestablishment of *Ayenia sidifolia* (Malvaceae: Byttnerioideae)

When Turczaninow (1863) synonymized *Cybiostigma* Turcz. with *Ayenia* he proposed the replacement name *A. mexicana* Turcz. because he thought that *A. sidifolia* Loefl. ex DC. prevented him from making a new combination based on *C. sidifolium* Turcz. However, *A. sidifolia* is a nomen nudum; an unintentional orthographic corruption of the earlier *A. sidiformis* Loefl., nom. rej. There was therefore no impediment to making a combination based on *C. sidifolium* and the correct name for this species in the genus *Ayenia* as a consequence is *A. sidifolia* (Turcz.) Hemsl.

Standley (1923) treated *Ayenia mexicana* as a doubtful species, but it is not clear why he did this. Cristóbal (1960, 1961), who revised the genus *Ayenia*, adopted *A. mexicana* and her choice, albeit incorrect, has been followed in the few floras that treat this species (Fryxell 2001; Machuca-Machuca 2015) as well as in the checklist of the vascular flora of Mexico (Villaseñor 2016).

***Ayenia sidifolia* (Turcz.) Hemsl.**, Biol. Cent.-Amer. Bot. 1(2):135. 1879 (as “*sidæfolia*”), non *Ayenia sidiformis* Loefl., Iter. Hispan. 257. 1758 (as “*Sidaformis*”), nom. rej. *Cybiostigma sidifolium* Turcz., Bull. Soc. Imp. Naturalistes Moscou 25(3):155. 1852. *Ayenia mexicana* Turcz., Bull. Soc. Imp. Naturalistes Moscou 36(2):569. 1863, nom. superfl. TYPE: MEXICO. Oaxaca: savannah near the Pacific, 3000 ft, 1840 (fl), H. Galeotti 326 (LECTOTYPE, as “tipo” designated by Cristóbal 1960: G [G00358304 as image!]; ISOLECTOTYPES: BR [BR0000005423101 as image!], K [K000543778!], K [K000543779!], KW [KW001000156 as image!]P [P02286136 as image!]).

*Ayenia nelsonii* Rose, Contr. U.S. Natl. Herb. 8(4):321. 1905 (as “nelsonii”). TYPE: MEXICO. CHIAPAS: roadside between San Ricardo and Ocozontla, 1600–3300 ft, 18 Aug 1895 (fl), E.W. Nelson 2982 (HOLOTYPE: US [US000102169]!).

**Distribution.**—This species occurs in central, western, and southern Mexico and adjacent Guatemala; 400–1850 m. Its precise range needs clarification; in particular the state records from the Yucatán Peninsula cited by Villaseñor (2016) are suspect and need to be verified.

Representative specimens examined: **MEXICO. Chiapas:** Mpio. Osumacinta: steep canon between Soyalo and La Bombana on road to Chicoasen, 10 km WNW of Soyalo, 7 Aug 1981 (fl), D.E. Breedlove 51986 (US); Hacienda Monserrate, Sep 1923 (fl), C.A. Purpus 9228 (MO, US). **Guerrero:** Mpio. Apango: Tlamamacan, 17°54'48" N, 099°31'50" W, 1 Sep 2005 (fl), J. Amith et al. 669 (US); Mpio. Tepecoaculco de Trujano: Cerca de Iguala, en Valerio Trujano, 30 Aug 1960 (fl), C. Cristóbal 601 (MO); Coyuco, Jaripo, 5 Nov 1934 (fr), G.B. Hinton 6928 (K). Near Iguala, 3500 ft, 11 Oct 1900 (fl, fr), C.G. Pringle 9254 (MO, US); bluffs of Balsas River, 2000 ft, 27 Sep 1905 (fl), C.G. Pringle 10070 (K, MO, P [P06697391 as image], US(2)). **México:** District of Temascaltepec, Chorrera, 15 Oct 1932 (fr), G.B. Hinton 2193 (ASU [ASU0049915 as image], DES [DES00005210 as image], GBH [as image], US).

### A new combination in *Reevesia* (Malvaceae: Helicteroideae)

When the monotypic American genus *Veeseria* Monach. & Moldenke was described, its authors (Monachino & Moldenke in Monachino 1940) recognized that it was closely related to the Asian genus *Reevesia*. They thought their new genus differed from *Reevesia* principally in the presence of staminodes. Solheim (1987, 1991) synonymized the two genera because this principal differentiating character is not supported; both genera have minute tooth-like staminodes at the summit of the anther column. Subsequently, Terada & Suzuki (1998), Pire & Cristóbal (2001), Bayer (2003), and Tang et al. (2007) have considered these two genera to be synonymous. The failure to formally make this synonymy explicit by transferring *Veeseria clarkii* Monach. & Moldenke to *Reevesia* muddles discussions about relationships between the floras of eastern Asia and North America (see e.g., Cristóbal 1987; Manchester et al. 2009) and about the phylogenetic relationships of fossil woods (Terada & Suzuki 1998; Kvaček 2006; Jeong et al. 2009; Manchester et al. 2006; Lim et al. 2010; Li et al. 2015).

***Reevesia clarkii*** (Monach. & Moldenke) Solheim ex Dorr, comb. nov. (**Fig. 1**). *Veeseria clarkii* Monach. & Moldenke, Bull. Torrey Bot. Club 67:621. 1940 (as “Clarkii”). TYPE: MEXICO. HIDALGO: mountainsides N of Chapahuacan, 8000 ft, 2 Jul 1935 (fl), O.M. Clark 7401 (HOLOTYPE: NY [NY00222433]!; ISOTYPES: MO [MO-194628]!, NY [NY00023376]!).

**Distribution.**—Endemic to Mexico where it is found in the mountains of Hidalgo and Querétaro; 800–1300 m. A Guatemalan record (Parker, 2008, as “*Veeseria clarkii*”) is not vouchered and cannot be verified. Several Nicaraguan records (Cristóbal, 2001, as “*Veeseria clarkii*”) probably represent a distinct species of *Reevesia* (Solheim, 1991).

**Common names.**—Aguiche, conote [sic], jonote, and jonote amargoso.

Representative specimens examined: **MEXICO. Hidalgo.** Mpio. Chapulhuacan: N of Chapahuacan, 2 Jul 1935, O.M. Clark 2212 (UNM [UNM00035 as image]); 2 km N of Chapulhuacan along trail to transmission tower W of road from Chapulhuacan to Hacienda de Cahuazas, ca. 1.5 km E of Hwy 85, ca. 17 km SW of Tamazunchale, S.L.P., 21°10'N, 098°53'W, 22 Jun 1986, G.E. Schatz 1200 (MO); 2 km N of Chapulhuacan around trail to transmission tower W of road from Chapulhuacan to Hacienda de Cahuazas, ca. 1.5 km E of Mexico Highway 85, ca. 14 km SW of Tamazunchale, S.L.P., 21°10'N, 098°53'W, 7 Jul 1984 (fl), S.L. Solheim 1765 (BR [BR0000005422852 as image], K, MO, NY, P [P02286062 as image], US); 2 km N of Chapulhuacan, near transmission tower W of road to Cahuazas, ca. 1.5 km E of Mex. Hwy 85, 14 km SW of Tamazunchale, 21°10'N, 098°53'W, 25 Oct 1986 (fr), S.L. Solheim 2067 (MO, NY(2), US(2)). **Querétaro:** Mpio. Jalpan: 3–4 km al Norte de La Parada, 28 Jun 1991 (fl), B. Servín 1144 (MO, US); 5–6 km al Norte de La Parada, 19 Sep 1991 (fr), B. Servín 1337 (US); Mpio. Landa: Al W de El Humo, 23 Jul 1991 (fl), E. Carranza 3295 (US); 2 km al Norte de Neblinas, 29 Sep 1988 (fr), H. Rubio 183 (US); Rancho Nuevo, 2 km al Sureste de San Onofre, 7 Jul 1989 (fl), H. Rubio 875 (US); 1 km al Sureste de San Onofre, 17 Jun 1991 (fl), H. Rubio 2435 (US); 1.5 km al Noroeste de San Onofre, 30 Jun 1991 (fl), H. Rubio 2444 (US); 1 km al Noroeste de San Onofre, 3 Aug 1991 (im fr), H. Rubio 2522 (US); 1 km al Poniente de El Humo, 19 Aug 1991 (fr), H. Rubio 2557 (US); 10 km al NE de Agua Zarga, sobre el camino a Neblinas, 23 Jun 1988 (fl), Rzedowski 46826 (MO, US).



FIG. 1. *Reevesia clarkii*. A. Habit. B. Flower at anthesis and floral buds. C. Anthers clustered at apex of staminal column; 5-lobed stigma visible in center. D. Fruit. E. Fruit with one locule open to show the apical attachment of a seed. F, G. Pair of seeds showing development of the wings. (A, C, Servín 1144; B, Rubio 2444; D–G, Solheim 2067. Illustration by C. Pasquale).

## ACKNOWLEDGMENTS

Erika Pagaza-Calderón (HJBC) and an anonymous reviewer provided helpful comments on the manuscript.

## REFERENCES

- BAYER, C. 2003. Malvaceae. In: K. Kubitzki & C. Bayer, eds. The families and genera of vascular plants. Volume 5. Springer-Verlag, Berlin, Germany. Pp. 225–311.
- CRISTÓBAL, C.L. 1960. Revisión del género *Ayenia* (Sterculiaceae). Opera Lilloana 4:1–230.
- CRISTÓBAL, C.L. 1961 [1962]. Nueva contribución al estudio del género *Ayenia* (Sterculiaceae). Anales Inst. Biol. Univ. Nac. Mexico 32:191–200.
- CRISTÓBAL, C.L. 1987. Un *Helicteres* nuevo de México, nexo entre las especies Asiáticas y Americanas. Bonplandia 6:71–81.
- CRISTÓBAL, C.L. 2001. Sterculiaceae. In: W.D. Stevens, C. Ulloa Ulloa, A. Pool, & O.M. Montiel, eds. Flora de Nicaragua. Tomo 3. Monogr. Syst. Bot. Missouri Bot. Gard. 85:2428–2437.
- DORR, L.J. & J.H. WIERSEMA. 2010a. Names of American vascular plants published in Loefling's *Iter Hispanicum* (1758) and its German translation (1766). Taxon 59:1245–1262.
- DORR, L.J. & J.H. WIERSEMA. 2010b. (1947–1958) Proposals to reject twelve names emanating from Loefling's *Iter Hispanicum* (1758), *Ayenia sidiformis* (Malvaceae), *Cofer* (Symplocaceae), *Cruzeta* and *C. hispanica* (Amaranthaceae), *Edechia inermis* and *E. spinosa* (Rubiaceae), *Justicia putata* (Acanthaceae), *Menais* and *M. topiaria* (? Boraginaceae), *Muco* (Capparaceae), *Samyda parviflora* (Salicaceae), and *Spermacoce suffruticosa* (Rubiaceae). Taxon 59:1280–1282.
- FRYXELL, P.A. 2001. Sterculiaceae. In: R. McVaugh, ed. Flora Novo-Galiciano. Volume 3. The University of Michigan Herbarium, Ann Arbor, U.S.A. Pp. 110–147.
- JEONG, E.K., K. KYUNGSIK, M. SUZUKI, & J.W. KIM. 2009. Fossil woods from the Lower Coal-bearing Formation of the Janggi Group (Early Miocene) in the Pohang Basin, Korea. Rev. Palaeobot. Palynol. 153:124–138. doi:10.1016/j.revpalbo.2008.07.006
- KVAČEK, Z. 2006. Fossil fruits of *Reevesia* (Malvaceae, subfam. Helicteroideae) and associated plant organs (seeds, foliage) from the Lower Miocene of North Bohemia (Czech Republic). Neues Jahrb. Geol. Paläontol., Monatsh. 2006 (7):431–448.
- LI, Y.-J., A.A. OSKOLSKI, F.M.B. JACQUES, & Z.-K. ZHOU. 2015. New middle Miocene fossil wood of *Wataria* (Malvaceae) from Southwest China. IAWA J. 36:345–357.
- LIM, J.-D., E.-K. JEONG, K. KIM, M. SUZUKI, I.S. PAIK, & H.-M. KIM. 2010. Miocene woods of the Janggi Basin in Korea: Implications for paleofloral changes. Geosciences J. 14:11–22. doi:10.1007/s12303-010-0002-8
- MACHUCA-MACHUCA, K. 2015. Sterculiaceae. In: R. Medina Lemos, ed. Flora del valle de Tehuacán-Cuicatlán. Volume 128. Instituto de Biología, Universidad Nacional Autónoma de México, México, D.F. Pp. 1–43.
- MANCHESTER, S.R., Z.-D. CHEN, A.-M. LU, & K. UEMURA. 2009. Eastern Asian seed plant endemic genera and their paleogeographic history through the northern hemisphere. J. Syst. Evol. 47:1–42. doi:10.1111/j.1759-6831.2009.00001.x
- MANCHESTER, S.R., Z. CHEN, & Z. ZHOU. 2006. Wood anatomy of *Craigia* (Malvales) from southeastern Yunnan, China. IAWA J. 27:29–136. doi:10.1163/22941932-90000142
- MONACHINO, J. 1940. A new genus and species of Sterculiaceae. Bull. Torrey Bot. Club 67:621–622. doi:10.2307/2481584
- PARKER, T. 2008. Trees of Guatemala. The Tree Press, Austin, Texas, U.S.A. Pp. i–vi, 1–1033.
- PIRE, S.M. & C.L. CRISTÓBAL. 2001. El polen de *Helicteres* (Sterculiaceae) y su comparación con géneros vecinos. Bonplandia 11:207–230.
- SOLHEIM, S.L. 1987. *Reevesia* (Sterculiaceae, Helicteroideae): Systematics and biogeography of an Asian-Eastern American disjunction. Amer. J. Bot. 74:754. [Abstract].
- SOLHEIM, S.L. 1991. *Reevesia* and *Ungeria* (Sterculiaceae): A taxonomic and biogeographic study. Ph.D. Dissertation, University of Wisconsin, Madison, Wisconsin, U.S.A. Pp. i–xv, 1–383.
- STANLEY, P.C. 1923. Trees and shrubs of Mexico. Contr. U.S. Natl. Herb. 23(3):i–iii, 517–848, v–xxviii.
- TANG, Y., M.G. GILBERT, & L.J. DORR. 2007. Sterculiaceae. In: X. Wu, P.H. Raven & D. Hong, eds., Flora of China. Volume 12. Science Press, Beijing/Missouri Botanical Garden Press, St. Louis, Missouri, U.S.A. Pp. 302–330.
- TERADA, K. & M. SUZUKI. 1998. Revision of the so-called 'Reevesia' fossil woods from the Tertiary in Japan—a proposal of the new genus *Wataria* (Sterculiaceae). Rev. Palaeobot. Palynol. 103:235–251.
- TURCZANINOW, N. 1863. Animadversiones ad catalogum primum et secundum herbaria universitatis charkoviensis. Bull. Soc. Imp. Naturalistes Moscou 36(2):545–615.
- VILLASEÑOR, J.L. 2016. Checklist of the native vascular plants of Mexico. Revista Mex. Biol. 87:559–902. doi:10.1016/j.rmb.2016.06.017



**Barney Lipscomb**  
**Botanical Research Institute of Texas Press**  
**1700 University Dr., Fort Worth TX 76107-3400**  
**Phone: 817-332-4441; Fax: 817-332-4112**  
**E-Mail [barney@brit.org](mailto:barney@brit.org)**

RE: [JBotResInstTexas-04]---Page proof of your Mexican Malvaceae paper

21 Jun 2017

Dear Larry,

I have page proof of your **Mexican Malvaceae** paper and I'm attaching it here in PDF format. Please acknowledge receipt of the pdf file of your proofs and that you can print out hard copies for proofing purposes (see instructions below). IF you cannot open the pdf file, PLEASE send an email or fax (817-332-4112) immediately indicating you need to receive page proofs either in hard copy via the USPS mail or via fax. IF fax, please send a fax number.

The **Spanish Resumen** will be added/checked by *J. Bot. Res. Inst. Texas'* Spanish editor before publication. IF no Spanish Resumen was provided the English abstract is duplicated as the Spanish Resumen place holder.

### **Instructions:**

1) OPEN the pdf file in Adobe Acrobat and edit with Tool box but do not use Touchup Text Tool in Advanced Editing. Feel free to **PRINT out a hard copy, MARK corrections CLEARLY** on the hard copy, and **RETURN** it to me preferably as an email attachment or mail if need.

2) \*\*PLEASE NOTE: Figures are low resolution (FPO, for position only) and they will be much sharper and crisper in the final printing. Please be sure to check the running heads for errors. Also, check for mathematical operators such as multiplication signs, greater than/less than/equal to; equals symbol, identical symbol, Greek characters; fractions, etc. to see that they have translated OK. Also check for temperature degree symbol and longitude and latitude marks and any other special symbols used.

3) → **Please proof and return by 30 Jun 2017 or sooner!!**

**Reprint QTY:**  None  50 [\$16]  100 [\$28]  low res PDF [complimentary] **[\$100]** hi-res Open Access PDF

♣ **Page costs for this paper are \$160 [4pp@\$40/p=\$160].** If you have support for publication costs and need an invoice before actual publication please let me know. **Accepted Payment Methods:** Visa, MasterCard, Check, Purchase Order, Money Order, Invoice, Bank Draft, Bank/Wire Transfer, Installments, Split Invoices. Please let me know if there is anything else needed in order to help with publication costs for this paper. Partial support is appreciated. Subscribers to the journal receive 15% discount on publication costs. An invoice will be sent after official publication unless requested earlier. We appreciate your support in keeping *J. Bot. Res. Inst. Texas* strong. Thank you!—*Barney Lipscomb*—817-332-7432  
VOICE; 817-332-4112 FAX; e-mail [barney@brit.org](mailto:barney@brit.org) ♣

4) **Please print this page, mark the number of reprints desired if any, and return with your hard copy or digital copy of page proofs.**

5) Expected publication date for *J. Bot. Res. Inst. Texas* 11(1) is July 2017.

I look forward to hearing from you. The text of this email may also be found as the last page in the attached PDF.

Best regards,—Barney