



K.P. 22
CHECKLIST
OF
GEKKONOID LIZARDS



ARNOLD G. KLUGE
Museum of Zoology
&
Department of Biology
University of Michigan



SMITHSONIAN
HERPETOLOGICAL INFORMATION
SERVICE
NO. 85

1991

SMITHSONIAN
HERPETOLOGICAL
INFORMATION
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.



INTRODUCTION

The following list includes all genus and species group names of extant gekkonoid lizards, their authors and years of publication. Most of the taxa I recognize are based on current common usage, and no attempt has been made to assess the monophyly of any genus. Junior objective and subjective synonyms are placed in parentheses. According to Article 80c of the ICZN, I follow existing usage in those cases under consideration by the Commission (e.g., Case 2527; BZN 46:38-40, 46-48, regarding *Nactus arnouxii* A. Duméril [1851] and *Nactus pelagicus* Girard [1858]). An asterisk is applied to all recognized subspecies, taxa followed by a question mark require further study, and a nomen nudum is followed by n.n. Not all unjustified emendations are noted in this abbreviated compilation. The list was derived from my manuscript of gekkonoid synonymies, wherein complete literature references, type localities and primary types are given. I would greatly appreciate notification of additions and corrections to the following list so that the complete version can be published with minimal errors. I wish to thank Aaron Bauer, Ross Sadlier and Glenn Shea for their assistance.

CHECKLIST

Aclys Kluge, 1974

concinna Kluge, 1974 (*major** Storr, 1987)

Aeluroscalabotes Boulenger, 1887 (*Aelurosaurus* Boulenger, 1885)

felinus Günther, 1864 (*borneensis* Günther, 1864;

dorsalis W. Peters, 1871; *longicaudatus*

Andersson, 1924; *multituberculatus** [?]

Kopstein, 1927)

Afroedura Loveridge, 1944

africana Boulenger, 1888 (*namaquensis** FitzSimons, 1938; *tirasensis** Haacke, 1965)

amatolica Hewitt, 1925

bogerti Loveridge, 1944

hawequensis Mouton and Mostert, 1985

karroica Hewitt, 1925 (*wilmoti* Hewitt, 1926; *halli** Hewitt, 1935)

nivaria Boulenger, 1894

pondolia Hewitt, 1925 (*multiporis** Hewitt, 1925; *langi** FitzSimons, 1930; *marleyi** FitzSimons, 1930; *haackei** Onderstall, 1984; *major** Onderstall, 1984)

tembulica Hewitt, 1926

transvaalica Hewitt, 1925 (*platyceps* Hewitt, 1925; *loveridgei** Broadly, 1963)

Agamura Blanford, 1874 (*Rhinogekko* de Witte, 1973)

femoralis M. A. Smith, 1933

gastropholis Werner, 1917
misonnei de Witte, 1973
persica A. Duméril, 1856 (*cruralis* Blanford, 1874)

Ailuronyx Fitzinger, 1843 (*Theconyx* Gray, 1845;
Aeluronyx Agassiz, 1846)
seychellensis Duméril and Bibron, 1836
trachygaster A. Duméril, 1851

Alsophylax Fitzinger, 1843 (*Altiphylax* Jeriomtschenko
and Shcherbak, 1984)
laevis Nikolsky, 1907 (*kashkarovi* Andrushko, 1968)
loricatus Strauch, 1887 (*szczerbaki** Golubev and
Sattarov, 1979)
pipliens Pallas, 1811
przewalskii Strauch, 1887 (*microtis* Blanford,
1875)
tadjikiensis Golubev, 1979
tokobajevi Jeriomtschenko and Shcherbak, 1984

Aprasia Gray, 1839 (*Ophioseps* Barboza du Bocage, 1873;
Ophiopsiseps Boulenger, 1887)
aurita Kluge, 1974
fusca Storr, 1979
haroldi Storr, 1978
inaurita Kluge, 1974
parapulchella Kluge, 1974
pseudopulchella Kluge, 1974
pulchella Gray, 1839 (*nasutus* Barboza du Bocage,
1873; *brevirostris* Werner, 1909)
repens Fry, 1914
rostrata Parker, 1956
smithi Storr, 1970
striolata Lütken, 1863 (*lineolata* Lütken, 1863;
octolineata W. Peters, 1864; *glauerti* Parker,
1956)

Aristelliger Cope, 1862 (*Idiodactylus* Bocourt, 1870;
Aristelligella Noble and Klingel, 1932)
barbouri Noble and Klingel, 1932
cochranae Grant, 1931 (*expectatus** Cochran, 1933)
georgeensis Bocourt, 1873 (*irregularis* Cope, 1885)
hechti Schwartz and Crombie, 1975
lar Cope, 1862
praesignis Hallowell, 1857 (*nelsoni** Barbour,
1914)

Asaccus Dixon and S. Anderson, 1973
elisae Werner, 1895 (*eugeniae* Nikolskij, 1907;
ingae Eiselt, 1973)
gallagheri Arnold, 1972

griseonotus Dixon and S. Anderson, 1973

Asiocolotes Golubev, 1984

depressus Minton and J. A. Anderson, 1965
levitoni Golubev and Shcherbak, 1979

Bavayia Roux, 1913

crassicollis Roux, 1913)
cyclura Günther, 1872 (*neocaledonicus* Barboza du
 Bocage, 1873; *bavayi* Sauvage, 1878)
montana Roux, 1913
ornata Roux, 1913
sauvagii Boulenger, 1883
septuiclavis Sadlier, 1988
validiclavis Sadlier, 1988

Bogertia Loveridge, 1941

lutzae Loveridge, 1941

Briba Amaral, 1935

brasiliiana Amaral, 1935

Bunopus Blanford, 1874 (*Trachydactylus* Haas and Battersby, 1959)

crassicauda Nikolsky, 1907
spatalurus J. Anderson, 1901 (*jolensis* Haas and
 Battersby, 1959; *hajarensis** Arnold, 1980)
tuberculatus Blanford, 1874 (*blanfordii* Strauch,
 1887; *gabrielis* Werner, 1936; *biporus* Werner,
 1938; *abudhabi* Leviton and S. Anderson, 1967)

Calodactylodes Strand, 1870 (*Calodactylus* Beddome, 1870)

aureus Beddome, 1870
illingworthi Deraniyagala, 1953

Carinatogecko Golubev and Shcherbak, 1981

aspratilis S. Anderson, 1973
heteropholis Minton, S. Anderson, and J. A.
 Anderson, 1970

Carphodactylus Günther, 1897

laevis Günther, 1897

Chondrodactylus W. Peters, 1870

angulifer W. Peters, 1870 (*weirei* Boulenger, 1887;
 *namibensis** Haacke, 1975)

Christinus Wells and Wellington, 1984 (*Ridgeekko* Wells
 and Wellington, 1985)

guentheri Boulenger, 1885
marmoratus Gray, 1845 (*peronii* Fitzinger, 1843;
 australis Gray, 1845; *macrodactylus*
 Boulenger, 1885; *affinis* Boulenger, 1885;

biggsae Wells and Wellington, 1985;
*alexanderi** Storr, 1987)

- Cnemaspis* Strauch, 1887 (*Ancylodactylus* L. Müller,
1907; *Paragonatodes* Noble, 1921)
affinis Stoliczka, 1870 (*penangensis* Flower, 1896)
africana Werner, 1895
argus Dring, 1978
barbouri Perret, 1986
beddomei Theobald, 1870 (*marmoratus* Beddome, 1870)
boiei Gray, 1842
boulengerii Strauch, 1887 (*glaucus* M. A. Smith,
1921)
dickersoni K. P. Schmidt, 1919
dilepis Perret, 1963
elgonensis Loveridge, 1935
lavolineata Nicholls, 1949
gigas Perret, 1986
goaensis Sharma, 1976
indica Gray, 1846
jerdonii Theobald, 1868 (*scalpensis** Ferguson,
1877)
kandiana Kelaart, 1852 (*malabaricus* Jerdon, 1853;
gracilis Beddome, 1870; *wicksii* Stoliczka,
1873; *humei* Theobald, 1876; *andersonii*
Annandale, 1904)
kendalli Gray, 1845
koehleri Mertens, 1937
kumpoli Taylor, 1963
littoralis Jerdon, 1853 (*planiceps* Beddome, 1871)
mysoriensis Jerdon, 1853
nairi Inger, Marx and Koshy, 1984
nigridius M. A. Smith, 1925
occidentalis Angel, 1943
ornata Beddome, 1870
petrodroma Perret, 1986
podihuna Deraniyagala, 1944
quattuorseriata Sternfeld, 1912 (*bohmanni* L.
Müller and Uthmoller, 1950)
siamensis M. A. Smith, 1925
sisparensis Theobald, 1876 (*maculatus* Beddome,
1870; *bireticulatus* Annandale, 1915)
spinicollis L. Müller, 1907
timoriensis Duméril and Bibron, 1836
tropidogaster Boulenger, 1885
uzungwae Perret, 1986
wynadensis Beddome, 1870

Coleodactylus Parker, 1926
amazonicus Andersson, 1918 (*zernyi* Wettstein,
1928)
brachystoma Amaral, 1935 (*pfrimeri*

Miranda-Ribeiro, 1939)
guimaraesi Vanzolini, 1957
meridionalis Boulenger, 1888
septentrionalis Vanzolini, 1980

Coleonyx Gray, 1845 (*Brachydactylus* W. Peters, 1863;
 Anarbylus Murphy, 1974)
brevis Stejneger, 1893
elegans Gray, 1845 (*scapularis* A. Duméril, 1851;
 coleonyx A. Duméril, 1856; *nemoralis**
 Klauber, 1945)
fasciatus Boulenger, 1885
mitratus W. Peters, 1863 (*dovii* Boulenger, 1885)
reticulatus Davis and Dixon, 1958
switaki Murphy, 1974 (*gypsicolus** Grismer and
 Ottley, 1988)
variegatus Baird, 1859 (*abbotti** Klauber, 1945;
 *bogerti** Klauber, 1945; *peninsularis**
 Klauber, 1945; *slevini** Klauber, 1945;
 *sonoriensis** Klauber, 1945; *utahensis**
 Klauber, 1945)

Colopus W. Peters, 1869
wahlbergii W. Peters, 1869 (*kalaharicus*
 FitzSimons, 1932; *furcifer** Haacke, 1976)

Cosymbotus Fitzinger, 1843 (*Platyurus* Oken, 1836;
 Nycteridium Günther, 1864; *Mimetozoon*
 Boulenger, 1896)
craspedotus Mocquard, 1890 (*floweri* Boulenger,
 1896)
platyurus Schneider, 1792 (*schneideriana* Shaw,
 1802; *marginatus* Cuvier, 1829; *schneideri*
 Günther, 1864; *himalayanum* J. Anderson, 1871;
nepalensis Annandale, 1907)

Crenadactylus Dixon and Kluge, 1964
ocellatus Gray, 1845 (*bilineatus* Gray, 1845;
 *horni** Lucas and Frost, 1895; *naso** Storr,
 1978; *rostralis** Storr, 1978)

Crossobamon Boettger, 1888 (*Ptenodactylus* Strauch, 1887)
eversmanni Wiegmann, 1834 (*atropunctatus*
 Lichtenstein and von Martens, 1856; *lumsdenii*
 Boulenger, 1887; *maynardi** M. A. Smith, 1933)
orientalis Blanford, 1876 (*dunstervillei* Murray,
 1884)

Cyrtopodion Fitzinger, 1843 (*Dinosastra* Gistel, 1868;
 Mesodactylus Shcherbak and Golubev, 1984)
agamurooides Nikolsky, 1900
elongatus Blanford, 1875

kachhensis Stoliczka, 1872 (*petrensis* Murray,
1884)
montiumsalsorum Annandale, 1913
scaber Heyden, 1827 (*basoglu* Baran and Gruber,
1982)
watsoni Murray, 1892 (*ingoldbyi* Procter, 1923)

Delma Gray, 1831 (*Nisara* Gray, 1867; *Pseudodelma* Fischer, 1882)
australis Kluge, 1974
borea Kluge, 1974
butleri Storr, 1987 (*haroldi* Storr, 1987)
elegans Kluge, 1974
fraseri Gray, 1831
grayii A. Smith, 1849
impar Fischer, 1882 (*lineata* Rosen, 1905)
inornata Kluge, 1974
labialis Shea, 1987
mitella Shea, 1987
mollerii Lütken, 1863
nasuta Kluge, 1974
pax Kluge, 1974
plebeia De Vis, 1888 (*wollemi* [n.n.] Wells and
Wellington, 1985)
tincta De Vis, 1888 (*reticulata* Garman, 1901)
torquata Kluge, 1974

Diplodactylus Gray, 1832 (*Stenodactylopsis*
Steindachner, 1870; *Lucasius* Kinghorn, 1929;
Lucasium Wermuth, 1965; *Turnerdactylus* Wells
and Wellington, 1989; *Ozziedactylus* Wells and
Wellington, 1989; *Manwellisaurus* Wells and
Wellington, 1989)
alboguttatus Werner, 1910
byrnei Lucas and Frost, 1896 (*dorotheae* Wells and
Wellington, 1985)
conspicillatus Lucas and Frost, 1897 (*hillii*
Longman, 1915; *laevis* Sternfeld, 1924;
platyurus Parker, 1926)
damaeus Lucas and Frost, 1896
fulleri Storr, 1978
galeatus Kluge, 1963
granariensis Storr, 1979 (*rex** Storr, 1988)
immaculatus Storr, 1988
kenneallyi Storr, 1988
maini Kluge, 1962
mitchelli Kluge, 1963
occultus King, Braithwaite and Wombey, 1982
ornatus Gray, 1845
polyophthalmus Günther, 1867
pulcher Steindachner, 1870 (*bilineatus* Lucas and
Frost, 1903; *dorsotaeniata* Pellegrin, 1909;
dorsalis Werner, 1910; *lucasi* Fry, 1914)

savagei Kluge, 1963
squarrosum Kluge, 1962
steindachneri Boulenger, 1885 (*stenurus* Werner,
 1909; *jonathoni* Wells and Wellington, 1985)
stenodactylus Boulenger, 1896 (*woodwardi* Fry,
 1914)
taeniocauda De Vis, 1886 (*taeniocauda* Boulenger,
 1887)
tessellatus Günther, 1875 (*pachyurus* Werner, 1909)
vittatus Gray, 1832 (*furcosus* W. Peters, 1864;
barbouri Angel, 1936)
wombeysi Storr, 1978

Dravidogecko M. A. Smith, 1933
anamallensis Günther, 1875

Ebenavia Boettger, 1878
inunguis Boettger, 1878 (*boettgeri* Boulenger,
 1885)

Eublepharis Gray, 1827
angramainyu S. Anderson and Leviton, 1966 (*ensafi*
 Baloutch and Thireau, 1986)
hardwickii Gray, 1827 (*lunatus* Blyth, 1847)
macularius Blyth, 1854 (*fasciolatus** Günther,
 1864; *afghanicus** Borner, 1974; *gracilis*
 Borner, 1974; *montanus** Borner, 1976; *fuscus**
 Borner, 1981; *smithi** Borner, 1981)
turcmenicus Darevsky, 1978

Eurydactylodes Wermuth, 1965 (*Eurydactylus* Sauvage, 1878)
symmetricus Andersson, 1908
vieillardi Bavay, 1869

Geckoella Gray, 1867
collegalensis Beddome, 1870 (*speciosus* Beddome,
 1870)
deccanensis Günther, 1864 (*albofasciatus*
 Boulenger, 1885; *dekkanensis* M. A. Smith,
 1935)
jeyporensis Beddome, 1877
madarensis Sharma, 1980
nebulosus Beddome, 1870
triedrus Günther, 1864 (*punctata* Gray, 1867)
yakhuna Deraniyagala, 1945 (*zonatus* Deraniyagala,
 1945)

Geckolepis Grandidier, 1867
anomala Mocquard, 1902
maculata W. Peters, 1880 (*humbloti* Vaillant, 1887)
modesta Methuen and Hewitt, 1913
petiti Angel, 1942

polylepis Boettger, 1893
typica Grandidier, 1867

Geckonia Mocquard, 1895
chazaliae Mocquard, 1895 (*malazodes* Ahl, 1930)

Gehyra Gray, 1834 (*Peropus* Wiegmann, 1835; *Phyria* Gray, 1842; *Dactyloperus* Fitzinger, 1843;
Perodactylus Fitzinger, 1843; *Peripia* Gray, 1845; *Chalinocnemis* Duges, 1883; *Spasmocnemis* Duges, 1883; *Phyriadora* Wells and Wellington, 1985)
angusticaudata Taylor, 1963
australis Gray, 1845 (*punctulata* Gray, 1842)
baliola A. Duméril, 1851 (*marmorata* Macleay, 1877;
 brevicaudis Macleay, 1877)
barea Kopstein, 1926
borroloola King, 1983
brevipalmata W. Peters, 1874
butleri Boulenger, 1900
catenata Low, 1979
dubia Macleay, 1877 (*longicaudis* Macleay, 1877;
 torresiana Günther, 1877)
fehlmanni Taylor, 1962
intermedia Brown, 1902
interstitialis Oudemans, 1894
kimberleyi Börner and Schuttler, 1983
lacerata Taylor, 1962
lampei Andersson, 1913
leopoldi Brongersma, 1930
marginata Boulenger, 1887 (*fischeri* Strauch, 1887;
 marginata Boettger, 1895)
membranacralis King and Horner, 1989
minuta King, 1982
montium Storr, 1982
mutilata Wiegmann, 1835 (*peronii* Duméril and Bibron, 1836; *platurus* Bleeker, 1858;
 insulensis Girard, 1858; *pardus* Tytler, 1865;
 perdicolor Cope, 1868; *packardii* Cope, 1869;
 navarri Duges, 1883; *beebei* Annandale, 1913)
nana Storr, 1978
occidentalis King, 1984
oceanica Lesson, 1830 (*pacifica* Gray, 1834;
 oualensis Duméril and Bibron, 1836; *gularis* Gray, 1842)
pamela King, 1982
papuana Meyer, 1874
pilbara Mitchell, 1965 (*cognata* Börner and Schuttler, 1982)
punctata Fry, 1914 (*fenestra* Mitchell, 1965)
purpurascens Storr, 1982
robusta King, 1983

variegata Duméril and Bibron, 1836 (*grayi*
Steindachner, 1867; *pusillus* [?] Cope, 1868;
mestoni de Vis, 1890; *annetteae* Wells and
Wellington, 1985; *kingi* Wells and Wellington,
1985; *lazelli* Wells and Wellington, 1985)
vorax Girard, 1858
xenopus Storr, 1978

Gekko Laurenti, 1768 (*Gecko* Brongniart, 1800; *Gecus*
Rafinesque, 1810; *Platydactylus* Goldfuss,
1820; *Gecco* Ritgen, 1828; *Lomatodactylus* van
der Hoeven, 1833; *Scelotretus* Fitzinger,
1843)
athymus Brown and Alcala, 1962
auriverrucosus Zhou and Liu, 1982
chinensis Gray, 1842 (*similignum* M. A. Smith,
1923)
gecko Linnaeus, 1758 (*verticillatus* Laurenti,
1768; *teres* Laurenti, 1768; *guttatus* Daudin,
1802; *verus* Merrem, 1820; *annulatus* Kuhl,
1820; *reevesii* Gray, 1831; *tenuis* Hallowell,
1857; *indicus* Girard, 1858; *azhari** Mertens,
1955)
gigante Brown and Alcala, 1978
hokouensis Pope, 1928 (*amissus* Taylor, 1962)
japonicus Schlegel (*in* Duméril and Bibron), 1836
(jamori Temminck and Schlegel, 1838; *nanus*
Cantor, 1842; *yamori* Fritze, 1891)
kikuchii Oshima, 1912
liboensis Zhou and Li, 1982
mindorensis Taylor, 1919
palawanensis Taylor, 1925
palmatus Boulenger, 1907
petricolus Taylor, 1962
porosus Taylor, 1922
romblon Brown and Alcala, 1978
scabridus Liu and Zhou, 1982
smithii Gray, 1842 (*stentor* Cantor, 1847;
albomaculatus Giebel, 1861; *albofasciolatus*
Günther, 1867)
subpalmatus Günther, 1864 (*melli* Vogt, 1922)
swinhonis Günther, 1864
taibaiensis Mingtao, 1985
tawaensis Okada, 1956
tuberculatus Daudin, 1802 (*monarchus* Schlegel, *in*
Duméril and Bibron, 1836; *burmeisteri* Giebel,
1861; *deissneri* Giebel, 1861)
verreauxii Tytler, 1865
vittatus Houttuyn, 1782 (*unistriata* Shaw, 1792;
bivittatus Duméril and Bibron, 1836;
trachyloemus W. Peters, 1872)
yakuensis Matsui and Okada, 1968

Gonatodes Fitzinger, 1843

albogularis Duméril and Bibron, 1836 (*fuscus**
 Hallowell, 1855; *varius* A. Duméril, 1856;
*notatus** Reinhardt and Lütken, 1862;
maculatus Steindachner, 1867; *braconnieri*
 O'Shaughnessy, 1875; *bodinii** Rivero Blanco,
 1964)
annularis Boulenger, 1887 (*boonii* Lidth de Jeude,
 1904; *beebei* Noble, 1923)
antillensis Lidth de Jeude, 1887
atricucullaris Noble, 1921
caudiscutatus Günther, 1859 (*collaris* Garman,
 1892)
ceciliae Donoso-Barros, 1965
concinnatus O'Shaughnessy, 1881 (*buckleyi*
 O'Shaughnessy, 1881; *ligiae* Donoso-Barros,
 1967)
eladioi Nascimento, Avila-Pires and Cunha, 1987
falconensis Shreve, 1947
hasemani Griffen, 1917 (*spinulosus* Amaral, 1932)
humeralis Guichenot, 1855 (*ferriugineus* 24Cope,
 1864; *incertus* W. Peters, 1871; *sulcatus*
 O'Shaughnessy, 1875)
ocellatus Gray, 1831 (*confidentialis* Marcuzzi,
 1950)
petersi Donoso-Barros, 1967
seigliei Donoso-Barros, 1965
taniae Roze, 1963
tapajonicus Rodrigues, 1980
vittatus Wiegmann, 1856 (*gilli* Cope, 1864;
*roquensis** Roze, 1956)

Goniurosaurus Barbour, 1908 (*Amamisaurus* Borner, 1981)

kuroiwae Namiye, 1912 (*orientalis** Maki, 1931;
yamashinae Okada, 1936; *splendens** Nakamura
 and Ueno, 1959)
lichtenfelderi Mocquard, 1897 (*hainanensis**
 Barbour, 1908)

Gonydactylus Kuhl and van Hasselt, 1822 (*Goniodactylus*

Schlegel, 1826; *Cyrtodactylus* Gray, 1827;
Gonyodactylus Wagler, 1830; *Goniodactylus*
 Gray, 1842; *Gonyodactylus* Fitzinger, 1843;
Puellula Blyth, 1860; *Quantasia* Wells and
 Wellington, 1985)
agusianensis Taylor, 1915
angularis M. A. Smith, 1921
annulatus Taylor, 1915
baluensis Mocquard, 1890
biordinis Brown and McCoy, 1980
brevipalmatus M. A. Smith, 1923

- cavernicolus* Inger and King, 1961 (*kinabalensis*
Malkmus, 1989)
chitralensis M. A. Smith, 1935
condorensis M. A. Smith, 1921
consobrinooides Annandale, 1905
consobrinus W. Peters, 1871
darmandvillei Weber, 1890 (*defossei* Dunn, 1927)
dattanensis Khan, 1980
derongo Brown and Parker, 1973
deveti Brongersma, 1948
elok Dring, 1978
fasciolatus Blyth, 1860
feae Boulenger, 1893
fraenatus Günther, 1864
fumosus F. Müller, 1895 (*halmahericus** Mertens,
1929)
gubernatoris Annandale, 1913
himalayensis Duda and Sahi, 1978
indusoani Khan, 1988
ingeri Hikida, 1990
intermedius M. A. Smith, 1917
irregularis M. A. Smith, 1921
jellesmae Boulenger, 1897
khasiensis Jerdon, 1870 (*himalayicus* Annandale,
1906; *tamaiensis** M. A. Smith, 1940)
kirmanensis Nikolsky, 1900
laevigatus Darevsky, 1964 (*uniformis** Auffenberg,
1980)
lateralis Werner, 1896
lawderanus Stoliczka, 1871 (*himalayensis*
Annandale, 1913)
loriae Boulenger, 1897
louisiadensis de Vis, 1892 (*tuberculatus* Lucas and
Frost, 1900; *olivii* Garman, 1901)
malcolmsmithi Constable, 1949
malayanus de Rooij, 1915
mansarulus Duda and Sahi, 1978
marmoratus Gray, 1831 (*agamensis* Bleeker, 1860)
matsuii Hikida, 1990
mimikanus Boulenger, 1914
mintoni Golubev and Shcherbak, 1981
novaeguineae Schlegel, 1844
oldhami Theobald, 1876
papuensis Brongersma, 1928 (*novaeguineae*
Brongersma, 1928)
peguensis Boulenger, 1893 (*zebraicus** Taylor,
1962)
philippinus Steindachner, 1867
pubisulcus Inger, 1957
pulchella Gray, 1827
quadrivirgatus Taylor, 1962
rohtasfortai Khan and Tasnim, 1990

redimiculus King, 1962
rubidus Blyth, 1860 (*tigris* Tytler, 1864)
sadleiri Wells and Wellington, 1985
sermowaiensis de Rooij, 1915
stoliczkai Steindachner, 1867 (*yarkandensis* J. Anderson, 1872; *walli* Ingoldby, 1922)
sworderi M. A. Smith, 1925
tibetinus Boulenger, 1905
variegatus Blyth, 1859
wetariensis Dunn, 1927
yoshii Hikida, 1990

Gymnodactylus Spix, 1825 (*Dasyderma* Fitzinger, 1843)
geckoides Spix, 1825 (*spixii* Gray, 1831;
gymnodactylus Schinz, 1833; *spinulosus* Fitzinger, 1843; *darwini** Gray, 1845;
girardi Steindachner, 1867; *amarali** Barbour, 1925; *conspicuus* Amaral, 1932; *helgae* Amaral, 1950)
guttulatus Vanzolini, 1982

Hemidactylus Gray, 1825 (*Gecus* Rafinesque, 1810;
Hemidact Oken, 1817; *Boltalia* Gray, 1842;
Hoplopodion Fitzinger, 1843; *Microdactylus* Fitzinger, 1843; *Onychopus* Fitzinger, 1843;
Tachybates Fitzinger, 1843; *Pnoepus* Fitzinger, 1843; *Doryura* Gray, 1845;
Velernesia Gray, 1845; *Leiurus* Gray, 1845;
Nubilia Gray, 1845; *Eurhous* Fitzinger, 1861;
Liurus Cope, 1862; *Emydactylus* Bocourt, 1870;
Bunocnemis Günther, 1894; *Lophopholis* M. A. Smith and Deraniyagala, 1934; *Aliurus* Dunn and Dunn, 1940)
agrius Vanzolini, 1978
albopunctatus Loveridge, 1942
aporus Boulenger, 1906
arnoldi Lanza, 1978
barodanus Boulenger, 1901
bavazzanoi Lanza, 1978
bayonii Barboza du Bocage, 1893
bouvieri Bocourt, 1870 (*cessacii* Barboza du Bocage, 1873; *boavistensis** Boulenger, 1906; *chevalieri* Angel, 1935; *razoensis** Gruber and Schleich, 1982)
bowringii Gray, 1845 (*berdmorei* Blyth, 1853)
brookii Gray, 1845 (*angulatus** Hallowell, 1854; *tytleri* Tytler, 1865; *guineensis* W. Peters, 1868; *affinis* Steindachner, 1870; *kushmorensis* Murray, 1884; *gleadowi* Murray, 1884; *stellatus* Boulenger, 1885; *murrayi* Gleadow, 1887; *tenkatei* Lidth de Jeude, 1895; *togoensis* Werner, 1897; *haitianus** Meerwarth,

- 1901; *subtriedroides** Annandale, 1905;
*leightoni** Boulenger, 1911; *luzonensis*
Taylor, 1915; *neotropicalis* Shreve, 1936;
*parvimaculatus** Deraniyagala, 1953)
citernii Boulenger, 1912
curlei Parker, 1942
depressus Gray, 1842 (*argentii* Gray, 1845;
pieresii Kelaart, 1853)
echinus O'Shaughnessy, 1875
fasciatus Gray, 1842 (*ornatus* Gray, 1845; *formosus*
Hallowell, 1857; *ituriensis** K. P. Schmidt,
1919)
flaviviridis Rüppell, 1835 (*coctaei* Duméril and
Bibron, 1836; *sublevis* Gray, 1842; *cocteau*
Fitzinger, 1843; *rupPELLi* Fitzinger, 1843;
bengaliensis J. Anderson, 1871; *sublaevis*
Boulenger, 1885; *bengalensis* Boulenger, 1885;
zolii Scortecci, 1929)
forbesii Boulenger, 1899
frenatus Duméril and Bibron, 1836 (*bojeri*
Fitzinger, 1843; *javanicus* Fitzinger, 1843;
vittatus Gray, 1845; *punctatus* Jerdon, 1853;
inornatus Hallowell, 1861; *pumilus* Hallowell,
1861; *caracal* Tytler, 1865; *chaus* Tytler,
1865; *hexaspis* Cope, 1869; *longiceps* Cope,
1869; *papuensis* Macleay, 1877; *tristis*
Sauvage, 1878; *fragilis* Calabresi, 1915;
vandermeermohri Brongersma, 1928; *okinawensis*
Okada, 1936)
funaiolii Lanza, 1978
garnotii Duméril and Bibron, 1836 (*peruvianus*
Wiegmann, 1835; *vulpecula* Girard, 1858;
ludekingii Bleeker, 1858; *guadama* Theobald,
1868; *mortonii* Theobald, 1868; *mandelianus*
Stoliczka, 1871; *blanfordii* Boulenger, 1885)
giganteus Stoliczka, 1871
gracilis Blanford, 1870 (*platyceps* Annandale,
1912)
granchii Lanza, 1978
granti Boulenger, 1899
greefi Barboza du Bocage, 1886
homoeolepis Blanford, 1881
intestinalis Werner, 1897 (*ansorgii* Boulenger,
1901)
isolepis Boulenger, 1895
jubensis Boulenger, 1895
karenorum Theobald, 1868
klauberi Scortecci, 1948
laevis Boulenger, 1901
laticaudatus Andersson, 1910 (*fossatii** Scortecci,
1928)
lemurinus Arnold, 1980

- leschenaultii* Duméril and Bibron, 1836 (*bellii*
 Gray, 1845; *pustulosus* Lichtenstein and von
 Martens, 1856; *kelaarti* Theobald, 1868;
marmoratus Blanford, 1870)
- longicephalus* Barboza du Bocage, 1873 (*longiceps*
 O'Shaughnessy, 1873; *bocagii* Boulenger, 1885;
mollerii Bedriaga, 1892; *hecqui* Boulenger,
 1901; *steindachneri* Tornier, 1902)
- mabouia* Moreau de Jonnès, 1818 (*tuberculosus*
 Raddi, 1823; *incanescens* Wied-Neuwied, 1824;
armatus Wied-Neuwied, 1824; *aculeatus* Spix,
 1825; *cruciger* Spix, 1825; *mabuia* Cuvier,
 1829; *mabuya* Fitzinger, 1843; *calabaricus*
 Boettger, 1878; *benguellensis* Barboza du
 Bocage, 1893)
- macropholis* Boulenger, 1896
- maculatus* Duméril and Bibron, 1836 (*sykesii*
 Günther, 1864; *hunae** Deraniyagala, 1937)
- marmoratus* Hallowell, 1861
- matschiei* Toriner, 1901
- megalops* Parker, 1932
- mercatorius* Gray, 1842 (*gardineri* Boulenger, 1909;
persimilis Barbour, 1928; *mandanus* Loveridge,
 1936)
- modestus* Günther, 1894
- muriceus* W. Peters, 1870
- newtoni* Ferreira, 1897
- nigriventris* Lidth de Jeude, 1905
- ophiolepis* Boulenger, 1903
- ophiolepoïdes* Lanza, 1978
- oxyrhinus* Boulenger, 1899
- palaichthus* Kluge, 1969
- persicus* J. Anderson, 1872 (*bornmuelleri* Werner,
 1895)
- platycephalus* W. Peters, 1854
- prashadi* M. A. Smith, 1935
- puccionii* Calabresi, 1927
- pumilio* Boulenger, 1899 (*pumilus* Boulenger, 1899)
- reticulatus* Beddome, 1870
- richardsonii* Gray, 1845
- ruspolii* Boulenger, 1896 (*erlangeri* Steindachner,
 1907)
- scabriceps* Annandale, 1906
- sinaitus* Boulenger, 1885 (*shugraensis* Haas and
 Battersby, 1959)
- smithi* Boulenger, 1895
- somalicus* Parker, 1932
- squamulatus* Tornier, 1896 (*bocagei* Toriner, 1896;
wernerii Tornier, 1897; *tornieri* Mocquard,
 1902; *floweri** Werner, 1908; *alluandi* Angel,
 1923; *barbouri** Loveridge, 1942)
- stejnegeri* Ota and Hikida, 1989

subtriedrus Jerdon, 1853
tanganicus Loveridge, 1929
tasmani Hewitt, 1932
taylori Parker, 1932
triedrus Daudin, 1802 (*lankae** Deraniyagala, 1953)
tropidolepis Mocquard, 1888
turcicus Linnaeus, 1758 (*cyanodactylus* Rafinesque,
 1810; *meridionalis* Risso, 1826; *granosus*
 Heyden, 1827; *robustus* Heyden, 1827;
verruculatus Cuvier, 1829; *verrucosus* Gray,
 1831; *karachiensis* Murray, 1884; *exsul*
 Barbour and Cole, 1906; *parkeri** Loveridge,
 1936; *spinalis** Buchholz, 1954)
vietnamensis Darevsky and Kupriyanova, 1984
yerburii J. Anderson, 1895 (*pauciporusus** Lanza,
 1978)

Hemiphyllodactylus Bleeker, 1860 (*Spathodactylus*
 Günther, 1872; *Spathoscalabotes* Boulenger,
 1885; *Cainodactylus* Barbour, 1924)
harterti Werner, 1900 (*larutensis* Boulenger, 1900)
insularis Taylor, 1918
typus Bleeker, 1860 (*gracilis* Bleeker, 1860;
crepuscularis Bavay, 1869; *aurantiacus**
 Beddome, 1870; *mutilatus* Günther, 1872;
ceylonensis Boulenger, 1885; *leucostictus*
 Stejneger, 1899; *margarethae* Brongersma,
 1931; *chapaensis** Bourret, 1937; *pallidus**
 Auffenberg, 1980)
yunnanensis Boulenger, 1903 (*dushanensis** Zhou and
 Liu, 1981; *jinpingsis** Zhou and Liu, 1981;
*longlingensis** Zhou and Liu, 1981)

Hemitheconyx Stejneger, 1893 (*Psilodactylus* Gray, 1864)
caudicinctus A. Duméril, 1851
taylori Parker, 1930

Heteronotia Wermuth, 1965 (*Heteronota* Gray, 1845)
binoei Gray, 1845 (*derbianus* Gray, 1845; *australis*
 Steindachner, 1867; *anomalus* W. Peters, 1868;
annulatus Macleay, 1877; *horneri* Wells and
 Wellington, 1985; *wadei* Wells and Wellington,
 1985)
planiceps Storr, 1989
spelea Kluge, 1963

Holodactylus Boettger, 1893
africanus Boettger, 1893 (*aculeatus* Calabresi,
 1915)
cornii Scortecci, 1931

Homonota Gray, 1845 (*Cubina* Gray, 1845; *Cubinia* Gray, 1845; *Wallsaurus* Underwood, 1954; *Garthia* Donoso-Barros and Vanzolini, 1965; *Garthia* Vanzolini and Donoso-Barros, 1966)
andicola Cei, 1978
borellii Peracca, 1897
darwinii Boulenger, 1885 (*macrocephala** Cei, 1978)
fasciatus Duméril and Bibron, 1836 (*pasteuri* Wermuth, 1965)
gaudichaudii Duméril and Bibron, 1836 (*dorbignii** Duméril and Bibron, 1836; *klugei** Donoso-Barros, 1970)
horrida Burmeister, 1861 (*mattogrossensis* Berg, 1895)
penai Donoso-Barros, 1965
underwoodi Kluge, 1964
uruguayensis Vaz-Ferreira and Soriano, 1961
whitii Boulenger, 1885

Homopholis Boulenger, 1885 (*Platypholis* Boulenger, 1890; *Blaesodactylus* Boettger, 1893)
antongilensis Bohme and Meier, 1980
boivini A. Duméril, 1856
fasciata Boulenger, 1890 (*erlangeri** Steindachner, 1906)
mulleri Visser, 1987
sakalava Grandidier, 1867 (*heterolepis* Boulenger, 1896)
walbergii A. Smith, 1849 (*macrolepis* Boulenger, 1885; *arnoldi* Loveridge, 1944)

Hoplodactylus Fitzinger, 1843 (*Pentadactylus* Gray, 1845; *Dactylocnemis* Steindachner, 1867; *Woodworthia* Garman, 1901)
chrysosireticus Robb, 1980
delcourtii Bauer and Russell, 1986
duvaucelii Duméril and Bibron, 1836
granulatus Gray, 1845 (*brevidactylus* Gray, 1845; *greyii* Knox, 1870; *sylvestris* Buller, 1881; *versicolor* Colenso, 1885; *nebulosus* McCann, 1955)
kahutarae Whitaker, 1985
maculatus Boulenger, 1885
pacificus Gray, 1842 (*pomarii* Girard, 1858; *wullerstorfi* Fitzinger, 1861; *brunneus* Cope, 1868; *digitatus* Garman, 1901)
rakiurae Thomas, 1981
stephensi Robb, 1980

Kaokogecko Steyn and Haacke, 1966
vanzylis Steyn and Haacke, 1966

Lepidoblepharis Peracca, 1897 (*Lathrogecko* Ruthven, 1916)

- buchwaldi* Werner, 1910
- colombianus* Mechler, 1968
- duolepis* Ayala and Castro, 1983
- festae* Peracca, 1897
- grandis* Miyata, 1985
- heyerorum* Vanzolini, 1978
- intermedius* Boulenger, 1914
- miyata* Lamar, 1985
- microlepis* Noble, 1923
- oxycephalus* Werner, 1894
- peraccae* Boulenger, 1908
- ruthveni* Parker, 1926
- sanctaemartae* Ruthven, 1916 (*fugax* Ruthven, 1928)
- williamsi* Ayala and Serna, 1986
- xanthostigma* Noble, 1916

Lepidodactylus Fitzinger, 1843 (*Amydosaurus* Gray, 1845)

- aureolineatus* Taylor, 1915
- balioburius* Ota and Crombie, 1989
- browni* Pernetta and Black, 1983
- christiani* Taylor, 1917
- euaensis* Gibbons and Brown, 1988
- gardineri* Boulenger, 1897
- guppyi* Boulenger, 1884
- herrei* Taylor, 1923 (*medianus** Brown and Alcala, 1978)
- listeri* Boulenger, 1889
- lugubris* Duméril and Bibron, 1836 (*neglectus* Girard, 1858; *meijeri* Bleeker, 1859; *pomareae* Fitzinger, 1861; *cantoris* Günther, 1864; *harrieti* Tytler, 1865; *moestus* W. Peters, 1867; *roseus* Cope, 1869; *caudeloti* Bavay, 1869; *mysorensis* Meyer, 1874; *ornata* Macleay, 1877; *divergens* Taylor, 1918; *lombocensis* Mertens, 1929; *ogasawarasimae* Okada, 1930; *intermedius** Darevsky, 1964)
- magnus* Brown and Parker, 1977
- manni* K. P. Schmidt, 1923
- mutahi* Brown and Parker, 1977
- novaeguineae* Brown and Parker, 1977
- oorti* Kopstein, 1926
- orientalis* Brown and Parker, 1977
- planicaudus* Stejneger, 1905 (*naujanensis* Taylor, 1919)
- pulcher* Boulenger, 1885
- pumilus* Boulenger, 1885
- pusillus* [?] Cope, 1868
- ranauensis* Ota and Hikida, 1988
- shebae* Brown and Tanner, 1949
- woodfordi* Boulenger, 1887

yami Ota, 1987

Lialis Gray, 1835 (*Ophiophthalmus* Fitzinger, 1843;
Alopekosaurus Lindholm, 1905)
burtonis Gray, 1835 (*bicatenata* Gray, 1842;
punctulata Gray, 1842; *leptorhyncha* W.
 Peters, 1874; *concolor* W. Peters, 1874)
jicari Boulenger, 1903 (*cuneirostris* Lindholm,
 1905; *inornata* Lindholm, 1905)

Luperosaurus Gray, 1845
brooksii Boulenger, 1920
browni Russell, 1979
cumingii Gray, 1845
joloensis Taylor, 1918
macgregori Stejneger, 1907
palawanensis Brown and Alcala, 1974

Lygodactylus Gray, 1864 (*Scalabotes* W. Peters, 1880;
Vanzoia H. M. Smith, Martin and Swain, 1977)
angolensis Barboza du Bocage, 1896 [1897] (*laurae*
 K. P. Schmidt, 1933)
angularis Günther, 1893 (*heeneni** DeWitte, 1933;
*grzimeki** Bannikov and Darevsky, 1969)
arnoulti Pasteur, 1964
bernardi FitzSimons, 1958 (*bonsi** Pasteur, 1962)
blanci Pasteur, 1967
bradfieldi Hewitt, 1932
capensis A. Smith, 1849 (*strigatus* Gray, 1864;
*grotei** Sternfeld, 1911; *ngamiensis*
 FitzSimons, 1932; *pakenhami** Loveridge, 1941)
chobiensis FitzSimons, 1932
conradti Matschie, 1892
conraui Tornier, 1902 (*strongi* Barbour and
 Loveridge, 1927)
decaryi Angel, 1930
depressus K. P. Schmidt, 1919
expectatus Pasteur and Blanc, 1967
fischeri Boulenger, 1890
grandisonae Pasteur, 1962
gravis Pasteur, 1964
guibei Pasteur, 1964
gutturalis Barboza du Bocage, 1873 (*paurospilus**
 Laurent, 1952; *dysmicus** Perret, 1963)
heterurus Boettger, 1913
inexpectatus Pasteur, 1964
insularis Boettger, 1913
keniensis Parker, 1936
klemmeri Pasteur, 1964
klugei H. M. Smith, Martin and Swain, 1977
lawrencei Hewitt, 1926
luteopicturatus Pasteur, 1964 (*zanzibaritis**

- Pasteur, 1964)
madagascariensis Boettger, 1881 (*pictus* W. Peters,
 1883; *petteri** Pasteur and Blanc, 1967)
manni Loveridge, 1928
methueni FitzSimons, 1937
miops Günther, 1891 (*spinulifer* Boettger, 1913)
montanus Pasteur, 1964
ocellatus Roux, 1907
ornatus Pasteur, 1964
picturatus W. Peters, 1868 (*variegatus* W. Peters,
 1868; *griseus* Toriner, 1896; *septemlineatus*
 Toriner, 1896; *quinquelineatus* Tornier, 1896;
*mombasicus** Loveridge, 1935; *sudanensis**
 Loveridge, 1935; *ukerewensis* Loveridge, 1935)
pictus W. Peters, 1883 (*robustus* Boettger, 1913)
rarus Pasteur and Blanc, 1973
rex Broadly, 1963
scheffleri Sternfeld, 1912 (*laterimaculatus**
 Pasteur, 1964; *compositus** Pasteur, 1964;
*ulugurensis** Pasteur, 1964)
scorteccii Pasteur, 1959
septemtuberculatus Angel, 1942
somalicus Loveridge, 1931 (*fisheri* Scortecci,
 1931; *annectans* Loveridge, 1935; *battersbyi**
 Pasteur, 1962; *howelli** Pasteur and Broadly,
 1988)
stevensonii Hewitt, 1926
thomensis W. Peters, 1880 (*rolasi* Greef, 1885;
*delicatus** Pasteur, 1962; *wermuthi** Pasteur,
 1962)
tolampyae Grandidier, 1872 (*tuberifer* Boettger,
 1913)
tuberosus Mertens, 1965
verticillatus Mocquard, 1895
wetzeli H. M. Smith, Martin and Swain, 1977
williamsi Loveridge, 1952
- Mediodactylus* Shcherbak and Golubev, 1977
amictopholis Hoofien, 1967
brevipes Blanford, 1874
heterocercus Blanford, 1874 (*mardinensis** Mertens,
 1924)
kotschi Steindachner, 1870 (*cyprius* Fitzinger,
 1843; *concolor** Bedriaga, 1882; *maculatus**
 Bedriaga, 1882; *danilewskii** Strauch, 1887;
*oertzeni** Boettger, 1888; *colchicus**
 Nikolsky, 1902; *bartoni** Stepanek, 1934;
*fitzingeri** Stepanek, 1937; *orientalis**
 Stepanek, 1937; *solerii** Wettstein, 1937;
*steindachneri** Stepanek, 1937; *stepaneki**
 Wettstein, 1937; *syriacus** Stepanek, 1937;
unicolor Wettstein, 1937; *wettsteini**

Stepanek, 1937; *bureschi* Stepanek, 1937;
*rumelicus** L. Müller, 1939; *saronicus**
Werner, 1939; *kalypsae** Stepanek, 1939;
*lycaonicus** Mertens, 1952; *rarus* Wettstein,
1952; *stubbei* Wettstein, 1952; *christianae*
Buchholz, 1955; *skopjensis** Karaman, 1965;
*buchholzi** Beutler and Gruber, 1977;
*schultzewestrumi** Beutler and Gruber, 1977;
gruberi Beutler and Gruber, 1977; *fuchsii**
Beutler and Gruber, 1977; *bibroni** Beutler
and Gruber, 1977; *adelphiensis** Beutler and
Gruber, 1978; *bileki* Tiedemann and Haupl,
1980; *tinensis** Beutler and Frör, 1980;
*beutleri** Baran and Gruber, 1981; *karabagi**
Baran and Gruber, 1982; *ciliciensis** Baran
and Gruber, 1982; *ponticus** Baran and Gruber,
1982)
russowii Strauch, 1887 (*zarudnyi** Nikolsky, 1899;
copalensis Schnitnikow, 1928)
sagittifer Nikolsky, 1899
spinicauda Nikolsky, 1887

Microgecko Nikolsky, 1907
helenae Nikolsky, 1907 (*fasciatus** Schmidtler and
Schmidtler, 1972)
latifi Leviton and S. Anderson, 1972
persicus Nikolsky, 1903 (*bakhtiari** Minton, S.
Anderson, J. A. Anderson, 1970;
*euphorbiacola** Minton, S. Anderson, and J. A.
Anderson, 1970)

Microscalabotes Boulenger, 1883
bivittis W. Peters, 1883 (*hildebrandti* W. Peters,
1883; *cowanni* Boulenger, 1883)

Millotisaurus Pasteur, 1962
mirabilis Pasteur, 1962

Nactus Kluge, 1983 (*Mascarenogecko* Ulber and Gericke, 1988)
pelagicus Girard, 1858 (*arnouxii* A. Duméril, 1851;
multicarinatus Günther, 1872; *arfakianus*
Meyer, 1874; *fasciata* Macleay, 1877;
marmorata Macleay, 1877; *eboracensis* Macleay,
1877; *cheverti* Boulenger, 1885; *heteronotus*
Boulenger, 1885; *undulatus** Kopstein, 1926)
coindermirensis Bullock, Arnold and Bloxam, 1985
galgajuga Ingram, 1978
serpensinsula Loveridge, 1951
vankampeni Brongersma, 1933

Narudasia Methuen and Hewitt, 1914
festiva Methuen and Hewitt, 1914

Naultinus Gray, 1842 (*Heteropholis* Fischer, 1882;
Naultinulus Chrapliwy, H. M. Smith, and
Grant, 1961)
elegans Gray, 1842 (*punctatus** Gray, 1843;
lineatus Gray, 1869; *sulphurus* Buller, 1871;
pentagonalis Colenso, 1880; *ocellatus* McCann,
1955)
gemmeus McCann, 1955
grayii Bell, 1843 (*simpsoni* Robb, 1980)
manukanus McCann, 1955
poecilochlorus Robb, 1980
rudis Fischer, 1881 [1882]
stellatus Hutton, 1872 (*pulcherrimus* Buller, 1877)
tuberculatus McCann, 1955

Nephrurus Günther 1876 (*Underwoodisaurus* Wermuth, 1965)
asper Günther, 1876
deleani Harvey, 1983
laevissimus Mertens, 1958
laevis De Vis, 1886 (*platyurus* Boulenger, 1886;
*occidentalis** Storr, 1963; *pilbarensis**
Storr, 1963)
miliusii Bory de Saint-Vincent, 1825 (*miliusii*
Duméril and Bibron, 1836; *blavieri* de
Rochebrune, 1884; *asper* Boulenger, 1913;
husbandi Wells and Wellington, 1984)
sphyrurus Ogilby, 1892 (*walshi* Kinghorn, 1931)
stellatus Storr, 1968
vertebralis Storr, 1963
wheeleri Loveridge, 1932 (*cinctus** Storr, 1963)

Oedura Gray, 1842 (*Pachyurus* Fitzinger, 1843; *Amalosia*
Wells and Wellington, 1984)
castelnauui Thominot, 1889 (*mayeri* Garman, 1901)
coggeri Bustard, 1966
filicipoda King, 1984
gemmaata King and Gow, 1983
gracilis King, 1984
lesueurii Duméril and Bibron, 1836 (*phillipsi*
Wells and Wellington, 1985)
marmorata Gray, 1842 (*fracticolor* De Vis, 1884;
cincta De Vis, 1888; *attenboroughi* Wells and
Wellington, 1985; *derelicta* Wells and
Wellington, 1985; *greeri* [n.n.] Wells and
Wellington, 1985)
monilis De Vis, 1888
obscura King, 1984
reticulata Bustard, 1969
rhombifer Gray, 1845
robusta Boulenger, 1885
tryoni De Vis, 1884 (*ocellata* Boulenger, 1885)

verrilli [?] Cope, 1868

Ophidiocephalus Lucas and Frost, 1897
taeniatus Lucas and Frost, 1897

Pachydactylus Wiegmann, 1834 (*Colobopus* Fitzinger,
 1843; *Cantinia* Gray, 1845; *Homodactylus* Gray,
 1864; *Elasmodactylus* Boulenger, 1894)
amoenus Werner, 1910
austeni Hewitt, 1923
bibronii A. Smith, 1845 [1846] (*turneri* Gray,
 1864; *stellatus* Werner, 1910; *pulitzerae** K.
 P. Schmidt, 1933)
bicolor Hewitt, 1926
capensis A. Smith, 1846 (*elegans* Gray, 1845;
mentomarginatus A. Smith, 1849;
*oshaughnessyi** Boulenger, 1885; *obscurus*
 Thominot, 1889; *affinis** Boulenger, 1896;
tessellatus Werner, 1910; *leopardinus*
 Sternfeld, 1911; *levyi* FitzSimons, 1933;
*vansonii** FitzSimons, 1933; *katanganus**
 De Witte, 1953)
caraculicus FitzSimons, 1959
fasciatus Boulenger, 1888
formosus [?] A. Smith, 1849
gaiasensis Steyn and Mitchell, 1967
geitje Sparrman, 1778 (*ocellatus* Cuvier, 1817;
inunguis Cuvier, 1817; *bergii* Wiegmann, 1834)
kobosensis FitzSimons, 1938
kochii FitzSimons, 1959
labialis FitzSimons, 1938
laevigatus Fischer, 1888 (*tessellatus* FitzSimons,
 1938; *fitzsimonsi** Loveridge, 1947)
maculatus Gray, 1845 (*albomarginatus* Hewitt, 1932;
microlepis Hewitt, 1935)
mariquensis A. Smith, 1849 (*latirostris** Hewitt,
 1923; *macrolepis* FitzSimons, 1939)
mentalis Hewitt, 1926
monticolus [?] FitzSimons, 1943
namaquensis Sclater, 1898
oculatus Hewitt, 1927
oreophilus McLachlan and Spence, 1967
punctatus W. Peters, 1854 (*brunnthaleri* Werner,
 1913; *langi* FitzSimons, 1932; *amoenoides**
 Hewitt, 1935; *scherzi** Mertens, 1954)
rugosus A. Smith, 1849 (*frater* Hewitt, 1935;
*barnardi** FitzSimons, 1941)
sansteyni Steyn and Mitchell, 1967
scutatus Hewitt, 1927 (*robertsi** FitzSimons, 1938;
*angolensis** Loveridge, 1944)
serval Werner, 1910 (*purcelli** Boulenger, 1910;
pardus Sternfeld, 1911; *montanus* Methuen and

Hewitt, 1914; *onscepensis** Hewitt, 1935)
tetensis Loveridge, 1953
tigrinus Van Dam, 1921 (*rhodesianus* Loveridge,
1947)
tsodiloensis Haacke, 1966
tuberculosus Boulenger, 1894 (*triedrus* Boulenger,
1913; *boulengeri* Toriner, 1896)
weberi Roux, 1907 (*garieensis* Hewitt, 1932;
*wernerii** Hewitt, 1935; *acuminatus**
FitzSimons, 1941)

Palmatogecko Andersson, 1908 (*Syndactylosaura* Werner, 1910)
rangei Andersson, 1908 (*schultzei* Werner, 1910)

Paradelma Kinghorn, 1926
orientalis Günther, 1876

Paragehyra Angel, 1929
petiti Angel, 1929

Paroedura Günther, 1879
androyensis Grandidier, 1867 (*porogaster*
Boulenger, 1896)
bastardi Mocquard, 1900
gracilis Boulenger, 1896
guibae Dixon and Kroll, 1974
homalorhinus Angel, 1936
oviceps Boettger, 1881
pictus W. Peters, 1854 (*madagascariensis* Mocquard,
1895; *robustus* Boulenger, 1896)
sanctijohannis Günther, 1879
stumpffi Boettger, 1879

Perochirus Boulenger, 1885
ateles A. Duméril, 1856 (*articulatus* Fischer,
1882; *depressus* Fischer, 1882; *klugei*
Wermuth, 1965)
guentheri Boulenger, 1885
scutellatus Fischer, 1882

Phelsuma Gray, 1825 (*Anoplopus* Wagler, 1830; *Chiroporus*
Wiegmann, 1834; *Rhoptropella* Hewitt, 1937;
Macrophelsuma Hoffstetter, 1946;
Archaeophelsuma Borner, 1972; *Neophelsuma*
Borner, 1972)
abbotti Stejneger, 1893 (*sumptio** Cheke, 1982)
andamanensis Blyth, 1860 (*chamaelon* Tytler, 1864)
astriata Tornier, 1901 (*carinatum* [?] Rendahl,
1939; *maheense* Rendahl, 1939; *astovei**
FitzSimons, 1948; *semicarinata** Cheke, 1982)
barbouri Loveridge, 1942
befotakensis Borner and Minuth, 1982

borbonica Mertens, 1966 (*agalegae** Cheke, 1975)
cepediana Merrem, 1820
chekei Borner and Minuth, 1982
comorensis Boettger, 1913
dubia Boettger, 1881
flavigularis Mertens, 1962
gigas Lienard, 1842 (*newtonii* Günther, 1877;
 newtonii Boulenger, 1884; *edwardnewtonii*
 Vinson and Vinson, 1969)
guentheri Boulenger, 1885
guimbeaui Mertens, 1963 (*rosangularis** Vinson and
 Vinson, 1969)
guttata Kaudern, 1922
laticauda Boettger, 1880 (*angularis** Mertens,
 1964)
lineata Gray, 1842 (*bifasciata** Boettger, 1913;
 *chloroscelis** Mertens, 1962; *bombetokensis**
 Mertens, 1964; *dorsivittata** Mertens, 1964;
 *pusilla** Mertens, 1964; *punctulata** Mertens,
 1970; *leiogaster* Mertens, 1973)
madagascariensis Gray, 1831 (*tetradactylus* [?]
 Schneider, 1797; *sarroubea* [?] Daudin, 1802;
 *grandis** Gray, 1870; *kochi** Mertens, 1954;
 martensi Mertens, 1962; *venusta** Mertens,
 1964; *notissima* Mertens, 1970; *boehmei**
 Meier, 1982)
minuthi Borner, 1980
modesta Mertens, 1970
mutabilis Grandidier, 1869 (*breviceps** Boettger,
 1894; *androyense* Mocquard, 1901; *micropholis*
 Boettger, 1913)
nigristriata Meier, 1984
ocellata Boulenger, 1885
ornata Gray, 1825 (*vinsoni* Mertens, 1963;
 *inexpectata** Mertens, 1966)
parkeri Loveridge, 1941
quadriocellata W. Peters, 1883 (*bimaculata**
 Kaudern, 1922; *leiura** Meier, 1983; *parva**
 Meier, 1983)
robertmertensi Meier, 1980
serraticauda Mertens, 1963
standingi Methuen and Hewitt, 1913
sundbergi Rendahl, 1939 (*longinsulae** Rendahl,
 1939; *pulchra* Rendahl, 1939; *cousinense*
 Rendahl 1939; *menaiensis* Mertens, 1966;
 *ladiguensis** Bohme and Meier, 1981; *rubra*
 Borner and Minuth, 1982; *umbrae* Borner and
 Minuth, 1982)
trilineata Gray, 1842
v-nigra Boettger, 1913 (*pasteuri** Meier, 1984)

Phyllodactylus Gray, 1828 (*Euleptes* Fitzinger, 1843;
Discodactylus Fitzinger, 1843; *Gerrhopygus*
 Gray, 1845)
angelensis Dixon, 1966
angustidigitus Dixon and Huey, 1970
ansorgii Boulenger, 1907
apricus Dixon, 1966
barringtonensis Van Denburgh, 1912
baurii Garman, 1892 (*gorii** Lanza, 1973)
bordai Taylor, 1942
bugastrolepis Dixon, 1966
clinatus Dixon and Huey, 1970
darwini Taylor, 1942
davisi Dixon, 1964
delcampi Mosauer, 1936
dixoni Rivero-Blanco and Lancini, 1968
duellmani Dixon, 1960
europaeus Gene, 1839 (*wagleri* Fitzinger, 1843;
doriae Lataste, 1877)
galapagensis W. Peters, 1869 (*daphnensis** Van
 Denburgh, 1912; *duncanensis* Van Denburgh,
 1912; *maresi** Lanza, 1973; *olschkii** Lanza,
 1973)
gerrhopygus Wiegmann, 1835 (*gymnopygus* Duméril and
 Bibron, 1836)
gilberti Heller, 1903
heterurus Werner, 1907
homolepidurus H. M. Smith, 1935 (*nolascoensis**
 Dixon, 1964)
inaequalis Cope, 1875
insularis Dixon, 1960
interandinus Dixon and Huey, 1970
johnwrighti Dixon and Huey, 1970
juliensi Cope, 1885
kofordi Dixon and Huey, 1970
lanei H. M. Smith, 1935 (*magnatuberculatus* Taylor,
 1940; *rupinus5** Dixon, 1964)
leei Cope, 1889
lepidopygus Tschudi, 1845 (*phacophorus* Tschudi,
 1845; *nigrofasciatus* Cope, 1878; *variegatus*
 Werner, 1900)
lineatus Gray, 1838 (*essexi** Hewitt, 1925;
*rupicolus** FitzSimons, 1938)
martini Lidth de Jeude, 1887
melanostictus Taylor, 1962
microlepidotus FitzSimons, 1939
microphyllus Cope, 1876
muralis Taylor, 1939 (*isthmus** Dixon, 1964)
palmeus Dixon, 1968
partidus Dixon, 1966
paucituberculatus Dixon, 1960

- peringueyi* [?] Boulenger, 1910
porphyreus Daudin, 1802 (*brevipes* Mocquard, 1900;
namaquensis Hewitt, 1935; *cronwrighti* Hewitt,
 1937)
pulcher Gray, 1828 (*spatulatus* Cope, 1863)
pumilis Dixon and Huey, 1970
reissii W. Peters, 1862 (*baessleri* Werner, 1900;
guayaquilensis Werner, 1910; *abrupteseriatus*
 Werner, 1913; *magister* Noble, 1924)
riebeckii W. Peters, 1882
rutteni Hummelinck, 1947
santacruzensis Dixon, 1966
sentosus Dixon and Huey, 1970
siamensis Boulenger, 1898 (*paviei* Mocquard, 1904;
burmanicus Annandale, 1905)
tinklei Dixon, 1966
trachyrhinus Boulenger, 1899
transversalis Huey, 1975
tuberculatus Wiegmann, 1835 (*magnus** Taylor, 1940;
edwardofischeri Mertens, 1952; *ingeri** Dixon,
 1964; *saxatilis** Dixon, 1964)
unctus Cope, 1864
ventralis O'Shaughnessy, 1875 (*mulleri* Parker,
 1935; *underwoodi* Dixon, 1962)
wirshingi Kerster and H. M. Smith, 1955
 (*hispaniolae** Schwartz, 1979; *sommeri**
 Schwartz, 1979)
xanti Cope, 1864 (*mentalis* Werner, 1910;
*nocticolus** Dixon, 1964; *zweifeli** Dixon,
 1964; *acorius** Dixon, 1966; *angulus** Dixon,
 1966; *circus** Dixon, 1966; *estabanensis**
 Dixon, 1966; *coronatus** Dixon, 1966; *sloani*
 Bostic, 1971)

Phyllopezus W. Peters, 1877
periosus Rodrigues, 1986
pollicaris Spix, 1825 (*goyazensis* W. Peters, 1877;
*przewalskii** Koslowsky, 1894; *spixii*
 Schlegel, 1933)

Phyllurus Goldfuss, 1820 (*Geckoides* Peron, 1807;
Phyllurus [n.n.] Oken, 1817; *Phyllura* Kaup,
 1827; *Anomalurus* Fitzinger, 1843)
caudiannulatus Covacevich, 1975
cornutus Ogilby, 1892 (*lichenosus* Günther, 1897;
swaini Wells and Wellington, 1985)
platurus Shaw, 1790 (*phyllurus* Schneider, 1797;
discosura Merrem, 1820; *platyura* Merrem,
 1820; *novae-hollandiae* Cuvier, 1822; *whitii*
 Gray, 1825; *cuvieri* Bory de Saint-Vincent,
 1828; *platycaudus* Schinz, 1835; *australis*
 Swainson, 1839; *inermis* Gray, 1845)

salebrosus Covacevich, 1975

Pletholax Cope, 1864

gracilis Cope, 1864 (*edelensis** Storr, 1978)

Pristurus Rüppell, 1835 (*Spatialura* Gray, 1863;

Geisopristurus Lanza and Sassi, 1968)

abdelkuri Arnold, 1986

carteri Gray, 1863 (*tuberculatus** Parker, 1931)

celerrimus Arnold, 1977

collaris Steindachner, 1867

crucifer Valenciennes, 1861 (*longipes* W. Peters,

1871; *stefaninii* Calabresi, 1927;

laticephalus Scortecci, 1933)

flavipunctatus Rüppell, 1835 (*percristatus*

Boulenger, 1896; *pseudoflavipunctatus*

Scortecci, 1933)

gasperetti Arnold, 1986 (*gallagheri** Arnold, 1986)

guichardi Arnold, 1986

insignis Blanford, 1881

insignoides Arnold, 1986

minimus Arnold, 1977

ornithocephalus Arnold, 1986

phillipsii Boulenger, 1895 (*somalicus* Parker,
1932)

popovi Arnold, 1982

rupestris Blanford, 1874 (*migiurtinicus* Scortecci,
1933; *guweirensis** Haas, 1951; *iranicus** K.

P. Schmidt, 1952)

saada Arnold, 1986

simonettai Lanza and Sassi, 1968

sokotranus Parker, 1938

Pseudogekko Taylor, 1922

brevipes Boettger, 1897

compressicorpus Taylor, 1915

labialis W. Peters, 1867

smaragdinus Taylor, 1922

Pseudogonatodes Ruthven, 1915

barbouri Noble, 1921

furvus Ruthven, 1915

guianensis Parker, 1935 (*amazonicus** Vanzolini,
1967)

lunulatus Roux, 1927

peruvianus Huey and Dixon, 1970

Ptenopus Gray, 1865

carpi Brain, 1962

garrulus A. Smith, 1849 (*maculatus** Gray, 1865)

kochi Haacke, 1964

Ptychozoon Kuhl, 1822 (*Ptyxozoon* Kuhl and Hasselt,
 1822; *Pteropleura* Gray, 1827)
horsfieldii Gray, 1827
intermedium Taylor, 1915
kuhlai Stejneger, 1902 (*homalocephala* Creveldt,
 1809)
lionotum Annandale, 1905
rhacophorus Boulenger, 1899

Ptyodactylus Goldfuss, 1820
guttatus Heyden, 1827 (*maculatus* Schinz, 1833;
lacazzi Boutan, 1892; *siphonorrhina* J.
 Anderson, 1896)
hasselquistii Donndorff, 1798 (*ascalabotes* Merrem,
 1820; *lobatus* Geoffroy, 1823; *ragazzi** J.
 Anderson, 1898; *togoensis* Tornier, 1901)
homolepis Blanford, 1876 (*socotranus**
 Steindachner, 1902)
oudrii Lataste, 1880
puiseuxi Bouton, 1893 (*bischoffsheimi* Boutan,
 1893; *montmahoui* Bouton, 1893; *barroisi*
 Bouton, 1893; *syriacus* Peracca, 1894;
sancti-montis Barbour, 1914)

Pygopus Merrem, 1820 (*Hysteropus* A. Duméril, in Duméril
 and Bibron, 1839; *Cryptodelma* Fischer, 1882)
lepidopodus Lacepede, 1804 (*novaehollandiae* Oppel,
 1811; *lepidopus* Merrem, 1820; *squamiceps*
 Gray, 1845; *longicaudatus* Tepper, 1882)
nigriceps Fischer, 1882 (*baileyi* Günther, 1897;
*schraderi** Boulenger, 1913; *klugei* Wells and
 Wellington, 1985; *territorianus* [n.n.] Wells
 and Wellington, 1985)

Quedenfeldtia Boettger, 1883
trachyblepharus Boettger, 1874 (*moerens* Chabanaud,
 1916)

Rhacodactylus Fitzinger, 1843 (*Correlophus* Guichenot,
 1866; *Ceratolophus* Barboza du Bocage, 1873;
Chameleonurus Boulenger, 1878; *Chamaeleonurus*
Carus, 1889; *Pseudothecadactylus* Brongersma,
 1936; *Torresia* Brongersma, 1934)
auriculatus Bavay, 1869 (*hexaceros* Barboza du
 Bocage, 1873)
australis Günther, 1877
cavaticus Cogger, 1975
chahoua Bavay, 1869
ciliatus Guichenot, 1866
leachianus Cuvier, 1829 (*aubryanus* Barboza du
 Bocage, 1873)
lindneri Cogger, 1975

sarasinorum Roux, 1913

trachyrhynchus Barboza du Bocage, 1873
 (*trachycephalus* Boulenger, 1878)

Rhoptropus W. Peters, 1869 (*Dactychylion* Thominot, 1878)

afer W. Peters, 1869

barnardi Hewitt, 1926

biporusus FitzSimons, 1957

boultoni K. P. Schmidt, 1933 (*benguellensis**
 Mertens, 1938; *montanus** Laurent, 1964)

braconnieri Thominot, 1878

bradfieldi Hewitt, 1935 (*diporus** Haacke, 1965)

taeniostictus Laurent, 1964

Rhynchoedura Günther, 1867

ornata Günther, 1867 (*paraornatus* Wells and
 Wellington, 1984; *ormsbyi* Wells and
 Wellington, 1985)

Sauromactylus Fitzinger, 1843

fasciatus Werner, 1931

mauritanicus Duméril and Bibron, 1836 (*desertorum*
 Fitzinger, 1843; *brosseti** Bons and Pasteur,
 1957)

Sphaerodactylus Wagler, 1830 (*Sphaeriodactylus* Gray, 1831)

alayoi Grant, 1959

altavelensis Noble and Hassler, 1933

(*brevirostratus** Shreve, 1968;

*enriguilloensis** Shreve, 1968; *lucioi* Thomas
 and Schwartz, 1983)

argivus Garman, 1888 (*bartschi** Cochran, 1934;
*lewisi** Grant, 1941)

argus Gosse, 1850 (*henriquesi* Grant, 1940;
*andresensis** Dunn and Saxe, 1950)

armasi Schwartz and Garrido, 1974

armstrongi Noble and Hassler, 1933 (*hypnephes**
 Thomas and Schwartz, 1983)

asterulus Schwartz and Graham, 1980

beattyi Grant, 1937 (*seamani** Thomas and Schwartz,
 1966)

becki K. P. Schmidt, 1919

bromeliarum G. Peters and Schwartz, 1972

caicosensis Cochran, 1934

calloricrus Schwartz, 1976

cinereus Wagler, 1830

clenchi Shreve, 1968 (*apocoptus** Schwartz, 1983)

cochranae Ruibal, 1946

copei Steindachner, 1867 (*anthracinus* Cope, 1862;

*picturatus** Garman, 1887; *asper* Garman, 1888;

*cataplexis** Schwartz and Thomas, 1965;

*enochrus** Schwartz and Thomas, 1965;

*polyommatus** Thomas, 1968; *astreptus**
 Schwartz, 1975; *pelates** Schwartz, 1975;
*websteri** Schwartz, 1975; *deuterus** Schwartz,
 1975)
corticola Garman, 1888 (*aporrox** Schwartz, 1968;
*campter** Schwartz, 1968; *soter** Schwartz,
 1968)
cryphius Thomas and Schwartz, 1977
darlingtoni Shreve, 1968 (*noblei** Shreve, 1968;
*bobilini** Thomas and Schwartz, 1983;
*mekistus** Thomas and Schwartz, 1983)
difficilis Barbour, 1914 (*lycauges** Schwartz,
 1983; *euopter** Schwartz, 1983; *typhlops**
 Schwartz, 1983; *peratus** Schwartz, 1983;
*diolenius** Schwartz, 1983; *anthracomus**
 Schwartz, 1983)
dunni K. P. Schmidt, 1936
elasmorhynchus Thomas, 1966
elegans Macleay, 1834 (*punctatissimus* Duméril and
 Bibron, 1836; *alopex* Cope, 1862)
elegantulus Barbour, 1917
fantasticus Duméril and Bibron, 1836
 (*ligniservulus** King, 1962; *anidrotus**
 Thomas, 1965; *fuga** Thomas, 1965; *hippomanes**
 Thomas, 1965; *karukera** Thomas, 1965;
*orescius** Thomas, 1965; *phyzacinus** Thomas,
 1965; *tartaropylorus** Thomas, 1965)
gaigeae Grant, 1932
gilvitorques Cope, 1862
glaucus Cope, 1865 (*inornatus* W. Peters, 1873;
torquatus Strauch, 1887)
goniorhynchus Cope, 1895
graptolaemus Harris and Kluge, 1983
heliconiae Harris, 1982
homolepis Cope, 1886 (*carinatus* Andersson, 1916;
imbricatus Andersson, 1916; *mertensi* Wermuth,
 1965)
inaguae Noble and Klingel, 1932
intermedius Barbour and Ramsden, 1919 (*drapetiscus*
 Schwartz, 1958)
klauberi Grant, 1931
ladae Thomas and Hedges, 1988
lazelli Shreve, 1968
leucaster Schwartz, 1973
levinsi Heatwole, 1968
lineolatus Lichtenstein and von Martens, 1856
 (*casicolus* Cope, 1862)
macrolepis Günther, 1859 (*grandisquamis* Stejneger,
 1904; *danforthi* Grant, 1931; *parvus** King,
 1962; *ateles** Thomas and Schwartz, 1966;
*guarionex** Thomas and Schwartz, 1966; *inigoi**
 Thomas and Schwartz, 1966; *mimetus** Thomas

- and Schwartz, 1966; *phoberus** Thomas and Schwartz, 1966; *spanius** Thomas and Schwartz, 1966; *stibarus** Thomas and Schwartz, 1966)
mariguanae Cochran, 1934
microlepis Reinhardt and Lütken, 1863
 (*melanospilos* Bocourt, 1873; *thomasi**
 Schwartz, 1965)
micropithecus Schwartz, 1977
millepunctatus Hallowell, 1861 (*continentalis*
 Werner, 1896)
molei Boettger, 1894 (*buergeri* Werner, 1900;
 venezuelanus Roux, 1927; *boettgeri* [n.n.]
 Donoso Barros, 1968)
monensis Meerwarth, 1901
nicholsi Grant, 1931
nigropunctatus Gray, 1845 (*decoratus** Garman,
 1888; *flavicauda** Barbour, 1904; *gibbus**
 Barbour, 1921; *atessares* Thomas and Schwartz,
 1966; *granti** Thomas and Schwartz, 1966;
*lissodesmus** Thomas and Schwartz, 1966;
*strategus** Thomas and Schwartz, 1966;
*porrasi** Schwartz, 1972)
notatus Baird, 1858 (*exsul** Barbour, 1914;
 *amaurus** Schwartz, 1966; *atactus** Schwartz,
 1966; *peltastes** Schwartz, 1966)
nycteropus Thomas and Schwartz, 1977
ocoae Schwartz and Thomas, 1977
oliveri Grant, 1944 (*storeyae** Grant, 1944)
omoglaux Thomas, 1982
oxyrhinus Gosse, 1850 (*dacnicolor** Barbour, 1910)
pacificus Stejneger, 1903
parkeri Grant, 1939
parthenopion Thomas, 1965
perissodactylus Thomas and Hedges, 1988
ramsdeni Ruibal, 1959
randi Shreve, 1968 (*methorius** Schwartz, 1977;
 *strahmi** Schwartz, 1977)
rhabdotus Schwartz, 1970
richardsoni Gray, 1845 (*gossei** Grant, 1939)
roosevelti Grant, 1931
rosaurae Parker, 1940
ruibali Grant, 1959
sabanus Cochran, 1938
samanensis Cochran, 1932
savagei Shreve, 1968 (*juanilloensis** Shreve, 1968)
scaber Barbour and Ramsden, 1919
scapularis Boulenger, 1902
semasiops Thomas, 1975
shrevei Lazell, 1961
sommeri Graham, 1981
sputator Sparrman, 1784 (*pictus* Garman, 1888)
streptophorus Thomas and Schwartz, 1977

- (*sphenophanes** Thomas and Schwartz, 1983)
stejnegeri Cochran, 1931
thompsoni Schwartz and Franz, 1976
torrei Barbour, 1914 (*spielmani** Grant, 1958;
 *ocujal** Thomas and Schwartz, 1966)
townsendi Grant, 1931
underwoodi Schwartz, 1968
vincenti Boulenger, 1891 (*festus** Barbour, 1915;
 *monilifer** Barbour, 1921; *adamas** Schwartz,
 1965; *diamesus** Schwartz, 1965; *josephinae**
 Schwartz, 1965; *pheristus** Schwartz, 1965;
 *psammius** Schwartz, 1965; *ronaldi** Schwartz,
 1965)
williamsi Thomas and Schwartz, 1983
zygaena Schwartz and Thomas, 1977
- Stenodactylus* Fitzinger, 1826 (*Goniodactylus* Schlegel,
 1826; *Tolarenta* Gray, 1842; *Ceramodactylus*
 Blanford, 1872; *Pseudoceramodactylus* Haas,
 1957; *Trigonodactylus* Haas, 1957; *Garzoniella*
 Perret, 1976)
affinis Murray, 1884
arabicus Haas, 1957
doriae Blanford, 1874 (*major* Parker, 1930)
grandiceps Haas, 1952
khobarensis Haas, 1957
leptocosymbotes Leviton and S. Anderson, 1967
petrii J. Anderson, 1896 (*stenurus* Werner, 1899;
 elimensis Barbour, 1914)
pulcher J. Anderson, 1896
slevini Haas, 1957 (*arabicus* Haas, 1957; *hassi*
 Kluge, 1967)
sthenodactylus Lichtenstein, 1823 (*elegans*
 Fitzinger, 1826; *savignyi* Audouin, 1827;
 guttatus Cuvier, 1829; *wilkinsonii* Gray,
 1842; *mauritanicus* Guichenot, 1850; *hirouxii*
 Doumergue, 1899; *longipes* Perret, 1976)
yemenensis Arnold, 1980
- Strophurus* Fitzinger, 1843 (*Strophura* Gray, 1845;
 Oedurella Lönnberg and Andersson, 1913;
 Eremiastrophurus Wells and Wellington, 1985)
assimilis Storr, 1988
ciliaris Boulenger, 1885 (*aberrans** Glauert, 1952)
elderi Stirling and Zietz, 1893 (*mahoodi* Wells and
 Wellington, 1985)
intermedius Ogilby, 1892
jeanae Storr, 1988
mcmillani Storr, 1978
michaelseni Werner, 1910
rankini Storr, 1979
spinigerus Gray, 1842 (*inornatus** Storr, 1988)

- strophurus* Duméril and Bibron, 1836 (*dumerili*
Fitzinger, 1843)
taeniatus Lönnberg and Andersson, 1913
wellingtonae Storr, 1988
williamsi Kluge, 1963
wilsoni Storr, 1983
- Tarentola* Gray, 1825 (*Makariogecko* Joger, 1984;
Sahelogecko Joger, 1984; *Saharogecko* Joger,
1984; *Neotarentola* Joger 1984)
americana Gray, 1831 (*milberti* Duméril and Bibron,
1836; *cubana* Gundlach and W. Peters, 1865;
*warreni** Schwartz, 1968)
angustimentalis Steindachner, 1891
annularis I. Geoffroy, 1827 (*savignyi* Audouin,
1827; *aegyptiacus* Cuvier, 1829; *senegalensis*
Boulenger, 1885; *quadraticauda* Toriner, 1905;
*relicta** Joger, 1984)
boehmei Joger, 1984
boettgeri Steindachner, 1891 (*hierrensis** Joger
and Bischoff, 1983; *bischoffi** Joger, 1984)
borneensis Gray, 1845 (*gigas** Barboza du Bocage,
1875; *rudis** Boulenger, 1906; *protogigas**
Joger, 1984; *maioensis** Schleich, 1984;
brancoensis Schleich, 1984)
caboverdianus Schleich, 1984 (*substituta**
Schleich, 1984; *antoensis* Joger, 1984;
*razianus** Schleich, 1984; *nicolauensis**
Schleich, 1984)
clypeata [?] Gray, 1842
darwini Joger, 1984
delalandii Duméril and Bibron, 1836
deserti Boulenger, 1891
ephippiata O'Shaughnessy, 1875 (*hoggarensis**
Werner, 1937; *panousei* Pasteur, 1959;
*senegambiae** Joger, 1984; *nikolausi** Joger,
1984)
gomerensis Joger and Bischoff, 1983
mauritanica Linnaeus, 1758 (*muricatus* Laurenti,
1768; *fascicularis** Daudin, 1802; *stellio*
Merrem, 1820; *muralis* Duméril and Bibron,
1836; *facetanus* Strauch, 1862; *atlantica*
Doumergue, 1899; *gracilis* Doumergue, 1899;
lissoide Doumergue, 1899; *saharae* Doumergue,
1899; *tuberculata* Rosen, 1905; *juliae** Joger,
1984)
neglecta Strauch, 1887 (*angusticeps* Strauch, 1887;
*greyi** Joger, 1984)
parvicarinata Joger, 1980
- Tenuidactylus* Shcherbak and Golubev, 1984
caspius Eichwald, 1831 (*fasciatus* Menetries, 1832;

insularis Akhmedow and Shcherbak, 1978)
fedtschenkoi Strauch, 1887
longipes Nikolsky, 1896 (*microlepis** Lantz, 1918)
turcmenicus Shcherbak, 1978
voraginosus Leviton and S. Anderson, 1984

Teratolepis Günther, 1870
albofasciatus Grandison and Soman, 1963
fasciata Blyth, 1853

Teratoscincus Strauch, 1863
bedriagai Nikolsky, 1899
microlepis Nikolsky, 1899
przewalskii Strauch, 1887
scincus Schlegel, 1858 (*keyserlingii** Strauch,
1863; *zarudnyi* Nikolsky, 1896; *roborowskii*
Bedriaga, 1905; *rustamowi** Shcherbak, 1979)

Thecadactylus Oken, 1817 (*Thecodactylus* Wagler, 1830)
rapicauda Houttuyn, 1782 (*perfoliatus* Schneider,
1793; *laevis* Daudin, 1802; *surinamensis*
Daudin, 1802; *theconyx* Duméril and Bibron,
1836; *tristis* Hallowell, 1854)

Tropiocolotes W. Peters, 1880
persicus Nikolsky, 1903
scorteccii Cherchi and Spano, 1963
steudneri W. Peters, 1869 (*petersii* Boulenger,
1891; *natterei* Steindachner, 1900)
tripolitanus W. Peters, 1880 (*occidentalis**
Parker, 1942; *somalicus** Parker, 1942;
*algericus** Loveridge, 1947; *apoklomax*
Papenfuss, 1970)

Urocotyledon Kluge, 1983
inexpectata Stejneger, 1893
palmata Mocquard, 1902
weileri L. Müller, 1909
wolterstorffi Tornier, 1900

Uroplates A. Duméril, 1805 [1806] (*Uroplates* Gray,
1825; *Rhacoessa* Wagler, 1830; *Oiacurus*
Fitzinger, 1843; *Lonchurus* Fitzinger, 1843)
alluaudi Mocquard, 1894
ebenaui Boettger, 1879 (*boettgeri* Fischer, 1883
[1884]; *phantasticus* Boulenger, 1888;
schniederi Lamberton, 1913)
fimbriatus Schneider, 1797 (*hypoxantha* Wagler,
1833)
guentheri Mocquard, 1908
lineatus Duméril and Bibron, 1836
sikorae Boettger, 1913