

A new havanensin-type limonoid from *Chisocheton* *macrophyllus*

by Mulyadi Tanjung

Submission date: 05-Aug-2021 01:06PM (UTC+0800)

Submission ID: 1627931511

File name: A_new_havanensin-type_limonoid_from_Chisocheton_macrophyllus.pdf (992.86K)

Word count: 4016

Character count: 19665

ARTICLE

Open Access

A new havanensin-type limonoid from *Chisocheton macrophyllus*



Nurlelarsari¹, Intan Rahmayanti¹, Supriatno Salam¹, Agus Safari¹, Desi Harneti¹, Rani Maharani^{1,2}, Ace Tatang Hidayat^{1,2}, Mulyadi Tanjung³, Rurini Retnowati⁴, Yoshihito Shiono⁵ and Unang Supratman^{1,2*}

Abstract

A new havanensin-type limonoid, 16 β -hydroxydysobinin (**1**), along with four known limonoids (**2–5**), have been isolated from the seeds of *Chisocheton macrophyllus*. The chemical structure of the new compound was determined by referencing spectroscopic data, and by comparison to those related spectra previously reported. Each compound was evaluated for their cytotoxic effects against Michigan Cancer Foundation-7 (MCF-7) breast cancer cells and display no significant activity.

Keywords: 16 β -hydroxydysobinin, *Chisocheton macrophyllus*, MCF-7, Limonoid, Meliaceae

Introduction

Limonoids, known as degraded triterpenes, are derived from a precursor with a 4,4,8-trimethyl-17-furanylsteroid four-ring skeleton labelled as A, B, C and D rings [1]. Limonoids are a class of secondary metabolites found in the order Rutales and the Meliaceae and Rutaceae family [2]. Meliaceae is a family of timber trees that are a rich source for limonoids and are widely distributed in tropical and subtropical regions with 50 genera and more than 1400 species [3, 4]. Limonoids isolated from species of the family Meliaceae have been reported to have biological activity as antifeedant, antimicrobial, anti-malarial, and cytotoxins [5–8]. Genus *Chisocheton* is the second largest in the family Meliaceae, consisting of more than 50 species distributed across Nepal, India, Bhutan, Myanmar, South China, Thailand, Indonesia, Malaysia, and Papua New Guinea [7, 9].

Chisocheton macrophyllus is a species distributed in the Nicobar Islands, peninsular Thailand, peninsular Malaysia, Singapore, Sumatera, Anambas Islands, Java and Borneo Islands [10]. Its seeds have been

reported to yield bioactive limonoids such as dysobinol, 7 α -hydroxyneotricilenone, dysobinin and nimonol with cytotoxic activity against P-388 murine leukemia cells [11], whereas the leaves to yield Epstein-Barr virus activation of Triterpenoids [12]. After further investigations for cytotoxic limonoids from the seeds of *C. macrophyllus*, we found and structural elucidation of a new havanensin-type limonoids (**1**) and four known limonoids (**2–5**), along with their cytotoxic activity against MCF-7 breast cancer cells. Herein, the isolation, structural elucidation and cytotoxic activity against MCF-7 breast cancer cells are discussed.

Materials and methods

Plant materials

Seeds of *C. macrophyllus* were collected from Bogor Botanical Garden, Bogor, West Java Province, Indonesia. The plant was identified by Mr. Harto, the staff of Bogoriense Herbarium, Research Center for Biology, Indonesia Science Institute, Bogor, Indonesia and a voucher specimen (No. Bo-1295453) was deposited at the Herbarium.

Instruments and reagents

Optical rotations were measured on a Perkin Elmer 341 Polarimeter (Waltham, MA, USA). UV spectra

*Correspondence: unang.supratman@unpad.ac.id

¹ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor 45363, Indonesia

Full list of author information is available at the end of the article

was measured using a TECAN Infinite M200 pro with MeOH. Furthermore, the IR spectra and mass spectra were recorded on a One PerkinElmer spectrum-100 FT-IR in KBr and Waters Xevo QTOF MS, respectively. NMR spectra were obtained with Bruker Topspin at 500 MHz for ^1H and 125 MHz for ^{13}C (compound 1) and for compounds 2–5 using JEOL JNM-ECZ500R/S1 at 500 MHz for ^1H and 125 MHz for ^{13}C , using tetramethylsilane (TMS) as the internal standard. Chromatographic separations were carried out on the silica gel 60 (70–230 and 230–400 mesh, Merck). Thin layer chromatography (TLC) analysis was carried out on 60 GF₂₅₄ (Merck, 0.25 mm) using various solvent systems, and measured by irradiation under ultraviolet–visible light Vilber Lourmat (λ 254 nm dan 365 nm) followed by heating of silica gel plates, sprayed with 10% H₂SO₄ in ethanol and Ehrlich's reagent (*p*-Dimethylaminobenzaldehyde in ethanol).

Extraction and isolation of *C. macrophyllus*

The dried and powdered seeds of *C. macrophyllus* (2.5 kg) were extracted with methanol at room temperature for 3 days (3 × 5 L). After removal of the solvent under a vacuum, a total of 360 g of methanol extract was obtained and partitioned with *n*-hexane (3 × 3 L), ethyl acetate (3 × 2 L) and *n*-butanol (3 × 2 L). Evaporation resulted in crude extracts of *n*-hexane (146.6 g), ethyl acetate (60.8 g) and *n*-butanol (14.6 g) respectively. The *n*-hexane soluble fraction (140 g) was subjected to vacuum-liquid chromatography (VLC) column packed with silica gel 60 using a gradient of *n*-hexane, ethyl acetate and methanol (10% stepwise) to afford thirteen fractions (A–M). Fraction D (5.4 g) was subjected to silica gel column chromatography using a gradient of *n*-hexane and ethyl acetate (5% stepwise) as eluting solvent to afford five subfractions (D1–D5). Subfraction D2 (165.7 mg) was chromatographed on a column of silica gel eluted with *n*-hexane: dichloromethane: ethyl acetate (2:7.5:0.5) to give 1 (15.3 mg). Fraction F (4.4 g) was subjected to a silica gel column chromatography using a gradient of *n*-hexane and ethyl acetate (5% stepwise) as eluting solvent to afford twelve subfractions (F1–F12). Subfraction F5 (1.2 g) was chromatographed on a column of silica gel eluted with *n*-hexane: dichloromethane: ethyl acetate (2:7.5:0.5) to give 3 (19.7 mg) and four subfractions (F5A–F5D). Furthermore, subfraction F5D (308.3 mg) was chromatographed on a column of silica gel eluted with *n*-hexane: dichloromethane: ethyl acetate (1:8.5:0.5) to give 2 (12.8 mg). Fraction H (1.8 g) was subjected to a silica gel column chromatography using a gradient of *n*-hexane and ethyl acetate (5% stepwise) as eluting solvent to give 5 (12.0 mg). Fraction J (1.5 g) was subjected to

a silica gel column chromatography using a gradient of *n*-hexane and ethyl acetate (5% stepwise) to afford nineteen subfractions (J1–J19). Subfraction J9 (50.3 mg) was chromatographed on a column of silica gel eluted with *n*-hexane: dichloromethane: ethyl acetate (4:5.5:0.5) to give 4 (3.0 mg).

16 β -hydroxydysobinin (1): Colorless needle crystals; mp: 205–207 °C; $[\alpha]_D^{27} +122.5^\circ$ (*c* 0.2, MeOH); UV (MeOH) λ_{max} 284 nm; IR (KBr) ν_{max} 3509, 2929, 1744, 1670, 1502, 1366, 1386, 1248 cm^{-1} ; HR-TOFMS m/z 511.2634 $[\text{M} + \text{H}]^+$, (calcd. for C₃₀H₃₉O₇ m/z 511.2696); $^1\text{H-NMR}$ (CDCl₃, 500 MHz) and $^{13}\text{C-NMR}$ (CDCl₃, 125 MHz) see Table 1.

Table 1 NMR spectral data for 1 (500 MHz for ^1H and 125 MHz for ^{13}C in CDCl₃)

Position	δ_{H} (ΣH , mult, <i>J</i> =Hz)	δ_{C} (mult.)
1	8.32 (1H, d, 10.5)	158.1 (d)
2	5.88 (1H, d, 10.5)	124.5 (d)
3	–	204.4 (s)
4	–	45.0 (s)
5	2.60 (1H, d, 12.5)	47.5 (d)
6	5.46 (1H, m)	69.7 (d)
7	5.41 (1H, d, 2.6)	74.0 (d)
8	–	41.6 (s)
9	2.12 (1H, dd, 6.5, 14.5)	45.4 (d)
10	–	42.7 (s)
11	1.86 (1H, d, 14.5)	34.3 (t)
	2.51 (1H, m)	
12	2.37 (1H, m)	46.4 (t)
	1.67 (1H, m)	
13	–	46.8 (s)
14	–	160.6 (s)
15	5.48 (1H, d, 7.7)	119.3 (d)
16	4.49 (1H, t, 7.7)	67.2 (d)
17	2.82 (1H, dd, 7.7, 11.0)	51.2 (d)
18	1.33 (3H, s)	28.7 (q)
19	1.28 (3H, s)	31.6 (q)
20	–	124.0 (s)
21	7.28 (1H, s)	139.7 (d)
22	6.30 (1H, d, 1.4)	110.9 (d)
23	7.40 (1H, d, 1.4)	142.7 (d)
28	0.95 (3H, s)	20.5 (q)
29	1.20 (3H, s)	20.6 (q)
30	1.33 (3H, s)	20.9 (q)
1'	2.04 (3H, s)	21.3 (q)
2'	–	170.1 (s)
1''	2.07 (3H, s)	22.0 (q)
2''	–	170.3 (s)

Cytotoxic activity test

The cytotoxicity of compounds 1–5 was determined with a cell viability test using PrestoBlue[®] assay. The cells were maintained in a Roswell Park Memorial Institute (RPMI) medium with 10% (v/v) Fetal Bovine Serum (FBS) and 1 μ L/1 mL antibiotics (1% Penicillin–Streptomycin). Cultures were incubated at 37 °C in a humidified atmosphere of 5% CO₂. MCF-7 cells plated in 96 multiwell culture plates at a density of 1.7×10^4 cells/well. After twenty-four hours, the medium was discarded and fresh medium containing sample with different concentrations 7.81, 15.63, 31.25, 62.50, 125.00, 250.00, 500.00, 1000.00 μ g/mL and control was added. After incubation with the sample for 24 h, PrestoBlue[®] reagent (resazurin dye) was added. The PrestoBlue[®] assay results were read using a multimode reader at 570 nm. The IC₅₀ values were determined by linier regression method using Microsoft Excel software. The IC₅₀ value corresponds to the concentration of compounds that decreases by 50% the number of viable cells and the absorbance in control corresponds to 100% viability.

Results and discussion

The *n*-hexane fraction from the seeds of *C. macrophyllus* was subjected to vacuum-liquid chromatography (VLC) column packed with silica gel 60 by gradient elution. The VLC fractions were repeatedly subjected to normal phase column chromatography on silica gel to yield compounds 1–5 (Fig. 1).

Compound 1 was isolated as colorless needle crystals. The molecular formula was determined to be C₃₀H₃₉O₇ based on the high resolution time-of-flight mass spectrometry (HR-TOFMS) spectra (Additional file 1; Fig. S8) at *m/z* 511.2634 [M+H]⁺ (calcd. for C₃₀H₃₉O₇ *m/z* 511.2696) and nuclear magnetic resonance (NMR) data (Table 1), indicating the presence of twelve degrees of unsaturation. The ultraviolet (UV) spectrum showed maximum absorption at 284 nm, indicating the presence of an α, β -unsaturated ketone [13, 14]. Infrared (IR) absorptions spectra suggested the presence of hydroxyl (3509 cm⁻¹), aliphatic (2929 cm⁻¹), carbonyl ester (1744 cm⁻¹), α, β -unsaturated carbonyl (1670 cm⁻¹), olefinic (1502 cm⁻¹), *gem*-dimethyl (1366 and 1386 cm⁻¹) and ether groups (1248 cm⁻¹).






Fig. 1 Structures of compounds 1–5

The $^1\text{H-NMR}$ spectrum (Additional file 1; Fig. S1) showed five tertiary methyls at δ_{H} 0.95 (3H, s, Me-28), 1.20 (3H, s, Me-29), 1.28 (3H, s, Me-19), and 1.33 (6H, s, Me-18 and Me-30) as well as two acetoxy groups at δ_{H} 2.04 (3H, s, Me-1') and 2.07 (3H, s, Me-1''). In addition, three oxygenated protons δ_{H} 5.46 (1H, m, H-6), 5.41 (1H, d, $J=2.6$ Hz, H-7) and 4.49 (1H, t, $J=7.7$ Hz, H-16), a β -furan moiety δ_{H} 6.30 (1H, d, $J=1.45$ Hz, H-22), 7.28 (1H, s, H-21), and 7.40 (1H, d, $J=1.45$ Hz, H-23) and three olefinic protons at δ_{H} 5.48 (1H, d, $J=7.7$ Hz, H-15), 5.88 (1H, d, $J=10.5$ Hz, H-2) and 8.32 (1H, d, $J=10.5$ Hz, H-1) were also observed in the $^1\text{H-NMR}$ spectrum. The $^{13}\text{C-NMR}$ (Additional file 1; Fig. S2) along with distortions enhancement by polarization transfer (DEPT) (Additional file 1; Fig. S3) and heteronuclear single quantum coherence (HSQC) spectra (Additional file 1; Fig. S4) showed thirty carbons consisting of an α,β -unsaturated carbonyl at δ_{C} 204.4 (C-3), two acetoxy groups at δ_{C} 21.3 (C-1'), 170.1 (C-2'), 22.0 (C-1'') and 170.3 (C-2'') and five methyls at δ_{C} 20.5 (Me-28), 20.6 (Me-29), 20.9 (Me-30), 28.7 (Me-18) and 31.6 (Me-19). The spectra also showed two methylene carbons at δ_{C} 34.3 (C-11) and 46.4 (C-12), three sp^3 methine carbons at δ_{C} 45.4 (C-9), 47.5 (C-5) and 51.2 (C-17), four sp^2 methine carbons at δ_{C} 110.9 (C-22), 119.3 (C-15), 124.5 (C-2) and 158.1 (C-1), three oxygenated sp^3 methine carbons at δ_{C} 67.2 (C-16), 69.7 (C-6) and 74.0 (C-7), two oxygenated sp^2 methine carbons at δ_{C} 139.7 (C-21) and 142.7 (C-23), four sp^3 quaternary carbons at δ_{C} 41.6 (C-8), 42.7 (C-10), 45.0 (C-4), and 46.8 (C-13) and two sp^2 quaternary carbons at δ_{C} 124.0 (C-20) and 160.6 (C-14). These functionalities accounted for seven out of the twelve degrees of unsaturation, while the remaining five degrees of unsaturation corresponded to the pentacyclic limonoid structure [6, 11, 15, 16]. The NMR spectra data of **1** resembled those of previously reported dysobinin [16, 17], except for the appearance of oxygenated signals [δ_{H} 4.49 (1H, t, $J=7.7$ Hz), δ_{C} 67.2], thus suggesting that **1** was a hydroxyl analog of dysobinin. Position of the hydroxyl group at C-16 was determined through the $^1\text{H-}^1\text{H}$ correlated spectroscopy ($^1\text{H-}^1\text{H}$ COSY) and proton multiple bond connectivity (HMBC) experiments (Fig. 2, Additional file 1; Fig. S5 and Fig. S6). Correlations from methyl protons at δ_{H} 1.33 (CH_3 -18) to δ_{C} 51.2 (C-17), oxygenated sp^3 methine at δ_{H} 4.49 (H-16) to δ_{C} 124.0 (C-20) and methyne proton δ_{H} 2.82 (H-17) to δ_{C} 139.7 (C-21) and δ_{C} 110.9 (C-22) were used to assign the hydroxyl group and a furan ring attached at C-16 and C-17, respectively.

Based on the $^1\text{H-}^1\text{H}$ COSY spectrum of **1**, correlation in $\text{H}_1\text{-H}_2$, $\text{H}_5\text{-H}_6\text{-H}_7$, $\text{H}_9\text{-H}_{11}\text{-H}_{12}$, $\text{H}_{16}\text{-H}_{17}$ and $\text{H}_{22}\text{-H}_{23}$ supported the presence of a havanensin-type limonoid structure in **1** [15, 17, 18]. The HMBC spectrum showed 3J correlations between sp^2 methine proton signal δ_{H}



8.32 (H-1) to δ_{C} 47.5 (C-5) and carbonyl at δ_{C} 204.4 (C-3) and δ_{H} 5.88 (H-2) to δ_{C} 42.7 (C-10), indicating the presence of α, β -unsaturated ketone located at C-1, C-2 and C-3, respectively. Correlations from oxygenated sp^3 methine protons at δ_{H} 5.46 (H-6) to δ_{C} 45.0 (C-4) and δ_{C} 170.1 (C-2') and δ_{H} 5.41 (H-7) to δ_{C} 45.4 (C-9) as well as δ_{H} 2.04 (H-1') to δ_{C} 170.1 (C-2') and δ_{H} 2.07 (H-1'') to δ_{C} 170.3 (C-2''), indicate that an acetyl group was attached at C-6 and C-7, respectively.

The relative stereochemistry of hydroxyl group at C-16 of **1** was determined by a nuclear overhauser and exchange spectroscopy (NOESY) experiment (Fig. 3 and Additional file 1; Fig. S7). Comparison of oxygenated sp³ methine protons at δ_{H} 4.49 (H-16) and CH₃-18 (δ_{H} 1.33) with α -oriented, indicated that H-16 was α -oriented and hydroxyl group at C-16 is β -oriented. Correlations between δ_{H} 5.41 (H-7) and CH₃-30 (δ_{H} 1.33) with β -oriented, indicated that H-7 was β -oriented and acetyl group at C-7 is α -oriented. Correlations between δ_{H} 5.46 (H-6) and CH₃-19 (δ_{H} 1.28) with β -oriented, indicated that H-6 was β -oriented and acetyl group at C-6 is α -oriented. Furthermore, the optical rotation of **1**, $[\alpha]_{\text{D}}^{27} + 122.5^{\circ}$ (*c* 0.2, MeOH) is the same sign to those of previously reported dysobinin (**6**) ($[\alpha]_{\text{D}}^{20} + 150^{\circ}$) [17, 18]. Therefore, the structure of **1** was elucidated as the new havanensin-type of limonoid derivative and named 16 β -hydroxydysobinin.

Four known compounds, 7-deacetylepoxызadiradione (**2**), were previously synthesized as a derivative of epoxyzadiradione [19, 20], but isolated from a natural source for the first time. In addition, 6 α -acetoxyepoxyzadiradione (**3**) and 6 α -acetoxygedunin (**4**) [21] as well as dysobinin (**5**) [11, 17, 18] were identified by comparison of their spectroscopic data with previously reported values.

Cytotoxic activity

All isolated compounds were evaluated for the cytotoxic activity against MCF-7 breast cancer cell line and cisplatin is used as a positive control according to the method previously described [16, 22] and the results are shown in Table 2. Compound **1** showed the strongest activity against MCF-7 breast cancer cell with IC₅₀ (inhibitory concentration, 50%) values of 45.91 μM , suggesting that the presence of hydroxyl at C-16 can increase the cytotoxic activity. In addition, the presence of epoxy ring and ketone group, like in compound **2–4**, showed weak activity, indicated the presence of epoxy ring and ketone group can decrease activity.

Table 2 Cytotoxicity activity of compounds **1–5** against MCF-7 breast cancer cell line

Compounds	IC ₅₀ (μM)
16 β -hydroxydysobinin (1)	45.91
7-deacetylepoxызadiradione (2)	94.62
6 α -acetoxyepoxyzadiradione (3)	105.16
6 α -acetoxygedunin (4)	121.82
Dysobinin (5)	68.15
Cisplatin*	38.06

* Positive control

18

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13765-021-00606-5>.

Additional file 1: Figure S1. ¹H-NMR Spectrum of **1** (500 MHz in CDCl₃). **Figure S2.** ¹³C-NMR Spectrum of **1** (125 MHz in CDCl₃). **Figure S3.** DEPT-135° Spectrum of **1** (135 MHz in CDCl₃). **Figure S4.** HSQC Spectrum of **1**. **Figure S5.** ¹H-¹H COSY Spectrum of **1**. **Figure S6.** HMBC Spectrum of **1**. **Figure S7.** ¹H-¹H NOESY Spectrum of **1**. **Figure S8.** HRTOFMS spectrum of **1**. **Figure S9.** ¹H-NMR Spectrum of **2** (500 MHz in CDCl₃). **Figure S10.** ¹³C-NMR and DEPT-135° Spectrum of **2**. **Figure S11.** HMQC Spectrum of **2**. **Figure S12.** ¹H-¹H COSY Spectrum of **2**. **Figure S13.** HMBC Spectrum of **2**. **Figure S14.** ¹H-¹H NOESY Spectra of **2**. **Figure S15.** HRTOFMS spectrum of **2**.

Acknowledgements

This investigation was financially supported by Universitas Padjadjaran, Indonesia (PPKI Grant, No: 1427/UN6.3.1/LT/2020 by US) by Airlangga University, Indonesia (PPKI Grant, No: 305/UN3.14/PT/2020 by MT) and Brawijaya University, Indonesia (PPKI Grant, No: 455.5/UN10.C10/PN/2020 by RR).

Authors' contributions

NS, IR and SS: Isolation of limonoid from the *n*-hexane extract; AS, DH, RM, ATH, MT: Structural elucidation of isolated compounds; RR, DH: cytotoxic assay; NS, IR, US, YS: preparing and completing manuscript. All authors read and approved the final manuscript.

Declarations

Competing interests

There is no conflict of interests.

Author details

¹ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor 45363, Indonesia. ² Central Laboratory, Universitas Padjadjaran, Jatinangor 45363, Indonesia. ³ Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60286, Indonesia. ⁴ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Malang 65145, Indonesia. ⁵ Department of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata University, Tsuruoka, Yamagata 997-8555, Japan.

Received: 25 January 2021 Accepted: 22 March 2021

Published online: 07 April 2021

References

- Fang X, Di YT, Hao XJ (2011) The advances in the limonoid chemistry of the Meliaceae Family. *Curr Org Chem* 15:1363–1391
- Li H, Peng Y, Zheng J (2016) (2016) Metabolic activation and toxicities of furanoterpenoids. *Adv Mol Toxicol* 10:55–97
- Paritala V, Chiruvella KK, Thammineni C, Ghanta RG, Mohammed A (2015) Phytochemicals and antimicrobial potentials of mahogany family. *Rev Bras Farmacogn* 25:61–83
- Tan QD, Luo XD (2011) Meliaceae Limonoids: chemistry and biological activities. *Chem Rev* 111:7437–7522
- Yang MH, Wang JS, Luo JG, Wang XB, Kong LY (2009) Tetranortriterpenoids from *Chisocheton paniculatus*. *J Nat Prod* 72:2014–2018
- Wong CP, Shimada M, Nagakura Y, Nugroho AE, Hirasawa Y, Taneda T, Awang K, Hadi AHA, Mohamad K, Shiro M, Morita H (2011) Ceramicines E–I, new limonoids from *Chisocheton ceramicus*. *Chem Pharm Bull* 59:407–411
- Shilpi JA, Sahab S, Chong SL, Nahard L, Sarker SD, Awang K (2016) Advances in chemistry and bioactivity of the genus *Chisocheton* BLUME. *Chem Biodiv* 13:483–503

8. Chong SL, Hematpoor A, Hazni H, Azirun MS, Litaudon M, Supratman U, Murata M, Awang K (2019) Mosquito larvicidal limonoids from the fruits of *Chisocheton erythrocarpus* Hiern. *Phytochem Lett* 30:69–73
9. Katja DG, Farabi K, Nurlelasari HD, Mayanti T, Supratman U, Awang K, Hayashi H (2017) Cytotoxic constituents from the bark of *Chisocheton cumingianus* (Meliaceae). *J Asian Nat Prod Res* 6:1–5
10. Vossen VD and Umali BE (2002) Plant Resources of South East Asia, No. 14, Vegetable oil and fats, Prosea Foundation, Bogor, Indonesia.
11. Nurlelasari KDG, Harneti D, Wardayo MM, Supratman U, Awang K (2017) Limonoids from the seeds of *Chisocheton macrophyllus*. *Chem Nat Compd* 53:83–87
12. Inada A, Somekawa M, Murata H, Nakanishi T, Tokuda H, Nishino H, Iwashima A, Darnaedi D, Murata J (1993) Phytochemical studies on meliaceous plants. VIII structures and inhibitory effects on Epstein-Barr Virus activation of triterpenoids from leaves of *Chisocheton macrophyllus* KING. *Chem Pharm Bull* 41(3), 617–619.
13. Shiono Y, Miyazani N, Murayama T, Koseki T, Harizon KDG, Supratman U, Nakata J, Kakihara Y, Saeki M, Yoshida J, Uesugi S, Kimura K (2016) GSK-3 β inhibitory activities of novel dichloresorcinol derivatives from *Cosmospora villiar* isolated from a mangrove plant. *Phytochem Lett* 18:122–127
14. Aisyah LS, Yun YF, Herlina T, Julaha E, Zainuddin A, Nurfarida I, Hidayat AT, Supratman U, Shiono Y (2017) Flavonoid compounds from the leaves of *Kalanchoe prolifera* and their cytotoxic activity against P-388 murine leukemia cells. *Nat Prod Sci* 23(2):139–145
15. Najmuldeen IA, Hadi AHA, Mohamad K, Awang K, Ketuly KA, Mukhtar MR, Taha H, Nordin M, Litaudon M, Gueritte F, Nugroho AE, Morita H (2012) Hisomicians D and E, two new limonoids from *Chisocheton ceramicus*. *Heterocycles* 84:1265–1270
16. Supriatno S, Nurlelasari HT, Harneti D, Maharani R, Hidayat AT, Mayanti T, Supratman U, Azmi MN, Shiono Y (2018) A new limonoid from stem bark of *Chisocheton pentandrus* (Meliaceae). *Nat Prod Res* 25:1–7
17. Singh S, Garg HS, Khanna NM (1976) Dysobinin, a new tetranortriterpene from *Dysoxylum binectariferum*. *Phytochemistry* 15:2001–2002
18. Maneerat WS, Laphookhieo S, Koysoomboon K, Chantrapromma K (2008) Antimalarial, antimycobacterial and cytotoxic limonoids from *Chisocheton siamensis*. *Phytomedicine* 15:1130–1134
19. Haldar S, Kolet SP, Thulasiram HV (2013) Biocatalysis: fungi mediated novel and selective 12 β - or 17 β -hydroxylation on the basic limonoid skeleton. *Green Chem* 15:1311–1317
20. Yadav PA, Kumar CP, Siva B, Babu KS, Allanki AD, Sijwali PS, Jain N, Rao AV (2017) Synthesis and evaluation of anti-plasmodial and cytotoxic activities of epoxyazadiradione derivatives. *Euro J of Med Chem* 134:242–257
21. Pereira TB, Silva L, Amorim R, Melo M, Souza R, Eberlin M, Lima ES, Vasconcellos MC, Pohlit AM (2014) In vitro and in vivo anti-malarial activity of limonoids isolated from the residual seed biomass from *Carapa guianensis* (andiroba) oil production. *Malar J* 13:317
22. Examinati RRIN, Wulandari AP, Harneti D, Poniah A (2018) Cytotoxicity of aromatic compound from an endophytic fungus, *Cladosporium* sp EN-S01. *Int J of Curr Pharm Res*, 10, 10–12.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

A new havanensin-type limonoid from *Chisocheton macrophyllus*

ORIGINALITY REPORT

20%

SIMILARITY INDEX

11%

INTERNET SOURCES

19%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 "Natural Antimicrobial Agents", Springer Science and Business Media LLC, 2018
Publication 1%
- 2 Mc Jesus Kinyok, Susan Bonnet, Olivier Placide Noté, Joséphine Ngo Mbing et al. " A new flavanolignan and a new alkane from the Stem bark of ", Natural Product Research, 2017
Publication 1%
- 3 Celia Bustos-Brito, Antonio Nieto-Camacho, Simón Hernandez-Ortega, José Rivera-Chávez et al. "Structural Elucidation of Malonylcommunol and 6 β -Hydroxy-trans-communic Acid, Two Undescribed Diterpenes from *Salvia cinnabarina*. First Examples of Labdane Diterpenoids from a Mexican *Salvia* Species", *Molecules*, 2020
Publication 1%
- 4 S. A. M. Abdelgaleil. "Antifeeding activity of limonoids from *Khaya senegalensis* 1%

(Meliaceae)", Journal of Applied Entomology,
5/2003

Publication

5 Bin Wu, Wen Hui Lin, Hui Yuan Gao, Lu Zheng,
Li Jun Wu, Chul Sa Kim. " Four New
Antibacterial Constituents from . ",
Pharmaceutical Biology, 2008

Publication

6 Yuan Li. "A New Stilbene from Cercis
chinensis Bunge", Journal of Integrative Plant
Biology, 8/2005

Publication

7 journal.waocp.org

Internet Source

8 Hong-Ying Wang, Jun-Song Wang, Si-Ming
Shan, Xiao-Bing Wang, Jun Luo, Ming-Hua
Yang, Ling-Yi Kong. "Chemical Constituents
from Trichilia connaroides and Their Nitric
Oxide Production and α -Glucosidase
Inhibitory Activities", Planta Medica, 2013

Publication

9 Kai-Long Ji, Shang-Gao Liao, Xiao-Ling Zheng,
Zhi Na, Hua-Bin Hu, Ping Zhang, You-Kai Xu.
"Limonoids from the Fruits of Khaya
ivorensis", Molecules, 2014

Publication

10 innovareacademics.in

Internet Source

11	thieme-connect.de Internet Source	1 %
12	Claudio Madeddu, Maria Cinta Roda-Serrat, Knud Villy Christensen, Rime B. El-Houri, Massimiliano Errico. "A Biocascade Approach Towards the Recovery of High-Value Natural Products from Biowaste: State-of-Art and Future Trends", Waste and Biomass Valorization, 2020 Publication	<1 %
13	M. Mostafa, Nilufar Nahar, M. Mosihuzzaman, Selestin D. Sokeng, Naheed Fatima, Atta-ur-Rahman, M. Iqbal Choudhary. " Phosphodiesterase-I inhibitor quinovic acid glycosides from ", Natural Product Research, 2006 Publication	<1 %
14	Dan Yuan. "Inhibitory activity of isoflavones of Pueraria flowers on nitric oxide production from lipopolysaccharide-activated primary rat microglia", Journal of Asian Natural Products Research, 06/2009 Publication	<1 %
15	mbio.asm.org Internet Source	<1 %
16	vjs.ac.vn Internet Source	<1 %

17	www.thieme-connect.de Internet Source	<1 %
18	Zhao-Qing Zeng, Wen-Ying Zhuang. "Our current understanding of the genus <i>Pseudocosmospora</i> (Hypocreales, Nectriaceae) in China", <i>Mycological Progress</i> , 2021 Publication	<1 %
19	Poitras, Marie-Eve, Martin Fortin, Catherine Hudon, Jeannie Haggerty, and JosÃ© Almirall. "Validation of the disease burden morbidity assessment by self-report in a French-speaking population", <i>BMC Health Services Research</i> , 2012. Publication	<1 %
20	www.journaltoocs.ac.uk Internet Source	<1 %
21	www.tandfonline.com Internet Source	<1 %
22	Di, Ying-Tong, Hong-Ping He, Chun-Shun Li, Jun-Mian Tian, Shu-Zhen Mu, Shun-Lin Li, Suo Gao, and Xiao-Jiang Hao. "Yuzurimine-Type Alkaloids from <i>Daphniphyllum yunnanense</i> ", <i>Journal of Natural Products</i> , 2006. Publication	<1 %
23	Qing-Wen Shi, Zuo-Ping Li, Ding Zhao, Ji-Shun Gu, Hiromasa Kiyota. " Isolation and structure	<1 %

revision of 10-deacetyltaxinine from the seeds of the Chinese yew, ", Natural Product Research, 2006

Publication

24

Ahmed I. Khodair. " SYNTHESIS OF 2-THIOHYDANTOINS AND THEIR - GLUCOSYLATED DERIVATIVES AS POTENTIAL ANTIVIRAL AND ANTITUMOR AGENTS ", Nucleosides, Nucleotides and Nucleic Acids, 2001

Publication

25

Marc Litaudon, Hadjira Bousserouel, Khalijah Awang, Olivier Nosjean et al. " A Dimeric Sesquiterpenoid from a Malaysian as a New Inhibitor of Bcl-xL/BakBH3 Domain Peptide Interaction ", Journal of Natural Products, 2009

Publication

26

Wen-Chen Chen, Jyh-Horng Sheu, Lee-Shing Fang, Wan-Ping Hu, Ping-Jyun Sung. " 3 α ,7 α ,12 α -Triacetoxo-5 β -cholanic acid, a steroid from the Formosan soft coral sp. (Alcyoniidae) ", Natural Product Research, 2006

Publication

27

hal-univ-rennes1.archives-ouvertes.fr

Internet Source

<1 %

<1 %

<1 %

<1 %

28

Internet Source

<1 %

29

repository.um.edu.my

Internet Source

<1 %

30

www.degruyter.com

Internet Source

<1 %

31

edepot.wur.nl

Internet Source

<1 %

32

jurnal.unpad.ac.id

Internet Source

<1 %

33

www.omicsonline.org

Internet Source

<1 %

34

www.rroij.com

Internet Source

<1 %

35

yamagata.repo.nii.ac.jp

Internet Source

<1 %

36

www.infona.pl

Internet Source

<1 %

37

Hosoya, T.. "Antioxidant phenylpropanoid glycosides from the leaves of Wasabia japonica", *Phytochemistry*, 200802

Publication

<1 %

38

Shadid, K.A.. "Cytotoxic caged-polyprenylated xanthonoids and a xanthone from *Garcinia cantleyana*", *Phytochemistry*, 200710

<1 %

39

Swati P. Kolet, Saikat Haldar, Siddiqui Niloferjahan, Hirekodathakallu V. Thulasiram. "Mucor hiemalis mediated 14 α -hydroxylation on steroids: In vivo and in vitro investigations of 14 α -hydroxylase activity", Steroids, 2014

Publication

40

Y. Venkateswarlu, P. Ramesh, N. Srinivasa Reddy, T. Prabhakara Rao. "A Novel Polyhydroxylated Epoxy Steriod From the Soft Coral Sinularia Dissecta", Natural Product Letters, 1999

Publication

41

downloads.hindawi.com

Internet Source

42

spandidos-publications.com

Internet Source

43

ugspace.ug.edu.gh

Internet Source

44

univmed.org

Internet Source

45

J. G. Urones, D. Díez, P. M. Gómez, I. S. Marcos, P. Basabe, R. F. Moro. " Drimane Homochiral Semisynthesis: Pereniporin a, - Warburganal and C-9 Nitrogenated Drimanes ", Natural Product Letters, 1998

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

46

Mireku, Evelyn Afua, Souvik Kusari, Dennis Eckelmann, Abraham Yeboah Mensah, Ferdinand M. Talontsi, and Michael Spiteller. "Anti-inflammatory tirucallane triterpenoids from *Anopyxis klaineana* Pierre (Engl.), (Rhizophoraceae)", *Fitoterapia*, 2015.

Publication

<1 %

47

P. Ashok Yadav, Ch. Pavan Kumar, Bandi Siva, K. Suresh Babu et al. "Synthesis and evaluation of anti-plasmodial and cytotoxic activities of epoxyazadiradione derivatives", *European Journal of Medicinal Chemistry*, 2017

Publication

<1 %

48

Qiang-Qiang Shi, Jing Lu, Xing-Rong Peng, Da-Shan Li, Lin Zhou, Ming-Hua Qiu. "Cimitriteromone A–G, Macromolecular Triterpenoid–Chromone Hybrids from the Rhizomes of ", *The Journal of Organic Chemistry*, 2018

Publication

<1 %

49

Sasaki, Hisako, Hirofumi Shibata, Kiyoshi Imabayashi, Yoshihisa Takaishi, and Yoshiki Kashiwada. "Prenylated Flavonoids from the Stems and Leaves of *Desmodium caudatum* and Evaluation of Their Inhibitory Activity against the Film-Forming Growth of

<1 %

Zygosaccharomyces rouxii F51", Journal of Agricultural and Food Chemistry

Publication

50

Sato, Kimihiko, Yasunori Inaba, Hyun-Sun Park, Toshiyuki Akiyama, Tetsuo Koyama, Haruhiko Fukaya, Yutaka Aoyagi, and Koichi Takeya. "Cytotoxic Bisnor- and Norditerpene Dilactones Having 7 $\hat{\pm}$,8 $\hat{\pm}$ -Epoxy-9,11-enolide Substructure from Podocarpus macrophyllus D. DON", CHEMICAL & PHARMACEUTICAL BULLETIN, 2009.

Publication

51

Uppuluri V. Mallavadhani, Anita Mahapatra, Satyabrata Mohapatra. "An efficient chemical method for separation of corsolic acid from its isomeric maslinic acid", Natural Product Research, 2006

Publication

52

escholarship.org

Internet Source

53

repository.unpad.ac.id

Internet Source

54

thno.org

Internet Source

55

www.pubmedcentral.nih.gov

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

56

Judit Hohmann, Ferenc Evanics, György Dombi, Pál Szabó. "Salicifoline and salicinolide, new diterpene polyesters from *Euphorbia salicifolia*", *Tetrahedron Letters*, 2001

Publication

<1 %

57

Michel Souza Passos, Thalya Soares Ribeiro Nogueira, Otoniel de Aquino Azevedo, Milena Gonçalves Curcino Vieira et al. "Limonoids from the genus *Trichilia* and biological activities: review", *Phytochemistry Reviews*, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

A new havanensin-type limonoid from *Chisocheton macrophyllus*

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
