



Short Communication

Otolith Dimensions versus Fish Lengths Estimated for Five Carangids (Pisces) in Pakistan

Nazia Qamar* and Sher Khan Panhwar

Center of Excellence in Marine Biology, University of Karachi, Karachi 75270

ABSTRACT

Sagitta otoliths of five species *Scomberoides commersonnianus* (414), *S. lysan* (15), *S. tala* (8), *S. tol* (344) and *Megalaspis cordyla* (277) were collected from the commercial catches from July 2013 to March 2015. Fish length and otolith length for five species recorded as *S. commersonnianus* TL_{cm} = 16.3–88.4, OL_{cm} = 0.3–0.9; *S. lysan* TL=23.2–73, OL = 0.2–0.7; *S. tala* TL = 48.5–55.5, OL = 0.6–0.9; *S. tol* TL = 18.9–68, OL = 0.3–0.9; *M. cordyla* TL= 18.4–47, OL = 0.3–0.9. The otolith dimensions such as length, height, weight were plotted against fish body parameters length and weights. The regression models estimate for otolith dimensions and fish body parameters estimates can be useful for studies on population structure of five important carangids species.

Article Information

Received 07 September 2016

Revised 13 November 2017

Accepted 22 December 2017

Available online 17 June 2019

Authors' Contributions

NQ prepared manuscript and performed statistical data analysis. SKP guided in the research and helped in manuscript writing.

Key words

Sagitta otolith, Linear model, Queen fishes, Torpedo.

Four queen fishes (*Scomberoides commersonnianus*, *S. tala*, *S. lysan* and *S. tol*) and a horse mackerel (*Megalaspis cordyla*) are co-occurring species in the waters off northern Arabian Sea coast of Pakistan are economically and ecologically important to the fisheries in the area. The recorded contribution of *M. cordyla* was 25% among overall carangid species caught in Pakistan. Although, no separate data is available for queen fishes, hence genus *Scomberoides* combined catch had been reduced from 17779mt in 1999 to 9073 mt in 2009 (Handbook, 2012). The FAO (2012) recorded increased landing of *S. commersonnianus* in the Western Indian Ocean from 4,994 mt in 2001 to 11,374 mt in 2010. Fish otoliths functions as balancing, hearing and an indicator of variability of fish habitat. In comparison to the fast swimming fishes broader otoliths found in slow swimming fishes (Parafkande-Haghighi, 2008). These hard sensory structures in fishes help in understanding feeding, fish stock monitoring and management (Harvey *et al.*, 2000) and its size can be affected by the process of digestion in the stomach. The aim of this paper was to compare variability between otolith dimensions and fish body parameters and find inter species shape variations in five species of carangid fishes.

Material and methods

A total of 773 individuals of five species of

Scomberoides and *Megalaspis* was collected from commercial catches from the Karachi fish harbor from July 2013 to March 2015. Species were identified using FAO species catalogs (Bianchi, 1985). Standard morphometric parameters such as total length, standard length, fork length, girth measured in (cm) and body weight in (grams). The sagitta otolith (pair) were extracted out and measured to (mm) and weighed at 0.0001-gram precision using digital balance (Diamond MCT500, Ohaus, USA). No significant difference in right and left otolith were tested (t test, P>0.05). However, right otoliths were used for each calculation and damaged otoliths were separated and did not use. Regression model were applied on total length-verses otolith length, otolith weight, and otolith length (OL) verses otolith weight (OW). Otolith length and width description presented in Supplementary Figure S1.

Results

The otolith shape parameters for five carangids showed considerable variations within the inter-intra species. Of all five species otolith have sulcus with heterosulcoid in shape and ostium is funnel-like while cauda tubular (Supplementary Table I). The parameters of fish (length and weight) and otolith (length and weight) for four queen fishes and one horse mackerel are summarized in Table I.

Regression model on TL~OL, TL~OW, OL~OW was applied separately for all five species (Fig. 1). We found strong correlation between otolith weight (OW)/otolith length and total length (TL) in *S. tala* (R²=1).

* Corresponding author: dr.naziaedu@yahoo.com
0030-9923/2019/0005-0001 \$ 9.00/0

Copyright 2019 Zoological Society of Pakistan




Fig. 1. Regression estimation of *S. commersonianus*, *S. lysan*, *S. tala*, *S. tol* and *M. cordyla* collected from northern Arabian Sea.

Table I.- Fish and otolith lengths and weights (total fish length TL, Otolith length OL, Otolith height OH), Otolith weight OW for five carangid species.

Species name	n	TL (cm) Min-Max	OL (cm) Min-Max	OH (cm) Min-Max	OW (g) Min-Max
<i>Scomberoides commersonianus</i> (Lacepède, 1801)	414	16.3–88.4	0.3–0.9	0.2–0.5	0.0006–0.00162
<i>S. lysan</i> (Forsskål, 1775)	15	23.2–73	0.2–0.7	0.15–0.35	0.0012–0.0078
<i>S. tala</i> (Cuvier, 1832)	8	48.5–55.5	0.6–0.9	0.3–0.35	0.00098–0.0105
<i>S. tol</i> (Cuvier, 1832)	344	18.9–68	0.3–1.1	0.15–0.5	0.0007–0.0155
<i>M. cordyla</i> (Linnaeus, 1758)	277	18.4–47	0.3–0.9	0.15–0.3	0.0008–0.024

Supplementary Figure S2 shows a very weak relation between otolith length and otolith weight of *S. commersonianus* (OL~OW, $R^2= 0.53$) (Fig. 1). The regression between TL~OL and OL~OW was estimated to be $R^2=0.74$ and 0.53 , respectively for *S. commersonianus*. For *S. lysan*, relationship between OL~OW and OL~TL was $R^2= 0.92$ and 0.91 , respectively. R^2 in *S. tala* for TL~OL was 1 and for OL~OW was 0.82 . For *S. tol* this relationship between TL~OW and OL~TL was 0.81 and 0.77 , respectively. In *M. cordyla* R^2 for TL~OL was $R^2=0.74$ and for OL~OW it was 0.53 (Fig. 1).

Discussion

The morphometry is widely used to evaluate fish population variations, monitor fish stocks and paleontological studies. Use of sulcus inscription structure in otoliths is rational method for distinguishing species in most of carangid species. The fast and slow swimming, nature can also be detected from the otolith shape in carangids, slow moving fishes have a larger otolith as compared to the fast swimming fishes. The sulcus shape correlates with the special nature of diet of *S. commersonianus*, *S. tol* and *M. cordyla* (Qamar *et al.*, 2015, 2016b). Somatic growth depends on availability of food, whereas temperature influences both metabolism and somatic growth. Larger otolith can be a result of slower somatic growth, whereas fast growth may produce elongated otolith. The rounded otolith may be the result of reduced growth. This study showed that the elongated otolith is because of slow growth rate. Qamar *et al.* (2016a) and Panhwar *et al.* (2014) have reported slow growth rate of *M. cordyla* ($K=0.73$), *S. commersonianus* ($K=0.25$) and *S. tol* ($K= 0.37$). Previous studies revealed that all these species are slow growing. Regression estimates showed positive correlation between otolith growth and fish length, whereas weak relationship was estimated in the otolith dimensions versus total length of *S. tala*. Otolith length increases till the fish attains its maximum length, the otolith continue to grow in thickness (Blacker, 1974). However, in three species (*S. commersonianus*, *S. tala* and *M. cordyla*) otolith length is the best predictor of fish length and in *S. tol* and *S. lysan*, otolith weight was supposed to be forecaster of fish length.

Supplementary material

There is supplementary material associated with this article. Access the material online at: <http://dx.doi.org/10.17582/journal.pjz/2017.x.x.xxx.xxxxxxxx>

Statement of conflict of interest

Authors have declared no conflict of interest.

References

- Blacker, R.W., 1974. In: *Sea fisheries research* (ed. F.R. Harden-Jones). John Wiley and Sons, pp. 67-90.
- Bianchi, G., 1985. *Field guide commercial marine and brackish water: FAO identification sheets for fishery purpose*. UNDP/ FAO/PAK/77/033
- FAO, 2012. *World fisheries production*, FAO official publication, Rome, Italy
- Harvey, J., Loughlin, T., Perez, M. and Oxman, D., 2000. *Relationship between fish size and otolith length for 63 species of fishes from Eastern North Pacific Ocean*. NOAA Technical Report, 150.
- Handbook, 2012. *Fisheries statistics of Pakistan*, Marine Fisheries Department, Karachi, Pakistan.
- Panhwar, S.K., Qamar, N. and Jahangir, S., 2014. *Pakistan J. Agric. Sci.*, **51**: 1111-1116.
- Parafkandeh-Haghighi, F., 2008. *Age determination in aquatic animals*. Fisheries Research Organization Publications, Tehran, pp. 139.
- Qamar, N., 2015a. *Population characteristics, reproductive seasonality, spawning, feeding habits and stock assessment of selected species of the family Carangidae (Pisces) in Pakistan*. Ph.D thesis, Center of Excellence in Marine Biology, University of Karachi, pp. 250.
- Qamar, N., Panhwar, S.K. and Jahangir, S., 2015b. *Pakistan J. Zool.*, **47**: 1171-1179.
- Qamar, N., Panhwar, S.K. and Brower, S., 2016a. *Pakistan J. Zool.*, **48**: 869-874.
- Qamar, N., Panhwar, S.K. and Siddiqui, G., 2016b. In: *Fisheries and aquaculture in the modern world* (ed. H. Mikkola). InTech Open, ISBN: 978-953-51-4732-9, pp. 169-201. <https://doi.org/10.5772/62627>




Supplementary Material

Short Communication: Otolith Dimensions versus Fish Lengths Estimated for Five Carangids (Pisces) in Pakistan

Nazia Qamar* and Sher Khan Panhwar

Center of Excellence in Marine Biology, University of Karachi, Karachi 75270

* Corresponding author: dr.naziaedu@yahoo.com
0030-9923/2019/0005-0001 \$ 9.00/0
Copyright 2019 Zoological Society of Pakistan



Supplementary Fig. S1. A general description of otolith length, height, anterior and poster positions of *Scomberoides tol* in this study.

Supplementary Table I.- Otolith shape description for five carangid species with common heterosulcoid, sulcus acoustic and funnel like ostium with tubular cauda.

Species name	Shape	Dorsal margin	Ventral margin	Rostrum	Anti-rostrum	Posterior position
<i>Scomberoides commersonianus</i>	lanceolate-triangular	Sinuate	Crenate	Narrow, Pointed, elongated	Broad, short	Round-irregular
<i>S. lysan</i>	Triangular	Sinuate	Crenate	Narrow, Pointed, elongated	Short, narrow, broad	Oblique
<i>S. tala</i>	lanceolate-triangular	Sinuate	Serrate	Narrow, elongated, pointed	Pointed, short, broad	Oblique-irregular
<i>S. tol</i>	Lanceolate, curved	Sinuate	Crenate	Narrow, long and pointed	Short, broad	Oblique
<i>M. cordyla</i>	lanceolate	Sinuate	dentate	Elongated	No	Oblique

Supplementary Fig. S2. Five species of fish and otolith collected from the Arabian Sea.



Scomberoides commersonianus



S. tol



S. lysan



S. tala



Megalaspis cordyla

