

Review

Lichen algae: the photosynthetic partners in lichen symbioses

William B. Sanders¹  and Hiroshi Masumoto²

¹Department of Biological Sciences, Florida Gulf Coast University, Ft. Myers, FL 33965-6565, USA and ²Laboratory of Terrestrial Microbiology and Systematics, Graduate School of Global Environmental Studies, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto 606-8501, Japan

Abstract

A review of algal (including cyanobacterial) symbionts associated with lichen-forming fungi is presented. General aspects of their biology relevant to lichen symbioses are summarized. The genera of algae currently believed to include lichen symbionts are outlined; approximately 50 can be recognized at present. References reporting algal taxa in lichen symbiosis are tabulated, with emphasis on those published since the 1988 review by Tschermak-Woess, and particularly those providing molecular evidence for their identifications. This review is dedicated in honour of Austrian phycologist Elisabeth Tschermak-Woess (1917–2001), for her numerous and significant contributions to our knowledge of lichen algae (some published under the names Elisabeth Tschermak and Liesl Tschermak).

Key words: chlorobiont, cyanobacteria, cyanobiont, lichenized stramenopiles, phaeobiont, photobiont, phycobiont, xanthobiont

(Accepted 17 June 2021)

Introduction

The principal components of the lichen symbiosis are fungus and alga. Their intimate trophic relationship remains central to the lichen concept, despite our growing appreciation that other microorganisms harboured within the thallus might also play significant roles (Lakatos *et al.* 2004; Grube & Berg 2009; Bates *et al.* 2011; Grube *et al.* 2015; Spribille *et al.* 2016; Muggia & Grube 2018; Mark *et al.* 2020; Smith *et al.* 2020; Tzovaras *et al.* 2020). The lichen-forming fungi typically build distinctive vegetative tissues and characteristic sexual structures, providing numerous biological features for study and significant clues about phylogenetic relationships, which are now relatively well delimited at broader taxonomic levels (Jaklitsch *et al.* 2016; Lücking *et al.* 2017a). Lichen algae, by contrast, have proved much more elusive. Most are unicells or simple filaments, with sexual structures unknown or seldom reported. The paucity of phenotypic characters is often aggravated by their plasticity. Lichen algae may look and behave quite differently in symbiosis with different lichen-forming fungi, in the free-living condition in nature and in aposymbiotic laboratory culture (Fig. 1; Ahmadjian 1967; Bubrick 1988). All this has hindered progress in clarifying their identities, phylogenies and life histories. Schwendener (1869) was the first to survey lichen ‘gonidia’ in a phycological context, recognizing them as organisms distinct from the surrounding fungus that correspond to known taxa of free-living algae. In the last half-century, the diversity of lichen-forming algae has been reviewed by various authors (Ahmadjian 1967; Létrouit-Galinou 1968; Henssen & Jahns 1974; Friedl & Büdel 2008), with a particularly thorough literature summary compiled and annotated by Tschermak-Woess (1988a).

In recent decades, our understanding of algal diversity and biosystematics has advanced substantially with the accumulation, analysis and integration of DNA sequence data. Systematic schemes for the eukaryotic algae have changed considerably, as the broad contours of consensus emerge concerning phylogenies and their reconstruction. Recent works have reviewed the current status of some principal algal groups with lichen-forming taxa, such as the genus *Trebouxia* (Muggia *et al.* 2017), the class *Trebouxiophyceae* (Muggia *et al.* 2018), the *Coccomyxa-Elliptochloris* clade (Gustavs *et al.* 2017), the *Trentepohliaceae* (Grube *et al.* 2017a), and the cyanobacteria (Rikkinen 2017). Yet most taxa remain insufficiently understood. Even the most intensively studied genera, such as *Trebouxia*, are still unresolved with respect to species delimitation, and much new diversity continues to be uncovered (Muggia *et al.* 2020). A great many algal symbionts, identified phenotypically (often without isolation into culture) or recorded merely as ‘trebouxioid’ or ‘chlorococcalean’, have yet to be revisited with DNA sequence analyses. Identities and relationships remain especially problematic among the cyanobacteria (blue-green algae), where sexual reproduction is absent, diversification is ancient (Garcia-Pichel 2009) and horizontal gene transfer events may obscure the vertical components of phylogenies (Zhaxybayeva *et al.* 2006). The aposymbiotic lives of lichen algae also remain largely unknown, despite their potential importance in active genetic mixing. Here an attempt is made to focus more attention on the algal side of the lichen partnership, still relatively neglected compared to that of the fungus. We include a synopsis of the relevant genera and list citations of algal taxa in lichen symbiosis (Table 1), emphasizing those published since Tschermak-Woess’s (1988a) landmark review, and particularly those accompanied by genetic sequence data.

The Major Algal Groups Involved

Lichen algae are diverse. This may contribute to the distinct distributions and climatic preferences of the symbiotic thalli that

Author for correspondence: William B. Sanders. E-mail: wsanders@fgcu.edu

Cite this article: Sanders WB and Masumoto H (2021) Lichen algae: the photosynthetic partners in lichen symbioses. *Lichenologist* 53, 347–393. <https://doi.org/10.1017/S0024282921000335>

© The Author(s), 2021. Published by Cambridge University Press on behalf of the British Lichen Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. Three filamentous lichen photobiont genera in aposymbiotic and symbiotic states. A–C, *Trentepohlia*. A, branching filament free-living on bark. B, lichenized by *Coenogonium* hyphae (arrows) growing over morphologically unchanged algal filament and its new branches (horizontal arrow). C, lichenized by *Arthonia rubricincta*; the alga is largely broken up into individual cells or short segments. D–F, *Rhizonema*. D, cultured isolate from *Dictyonema*; note false branching (arrowhead). E, trichome ensheathed by cells of mycobiont *Dictyonema*. F, contorted or broken filaments (arrow) within thallus of *Coccocarpia palmicola*. G–J, *Nostoc*. G, free-living thallus-like macrocolony on soil. H, cultured strain. I, more or less intact filaments (arrows) within thallus of *Collema furfuraceum*. J, contorted or broken up into cell groups (arrows) within cyanomorph of *Sticta canariensis*. Scales: A–F, H–J = 10 µm; G = 1 cm.

enclose them (Marini *et al.* 2011). Most are green algae, a paraphyletic grouping of two major clades: the charophytes (*Streptophyta*), from which embryophytes descend, and the *Chlorophyta* s. str. (Leliaert *et al.* 2012). The latter includes nearly all green algae reported as lichen symbionts. Within the *Chlorophyta*, lichen symbionts are found principally in the classes *Trebouxiophyceae* and *Ulvophyceae*. A third class, the *Chlorophyceae*, is known or suspected to include the partners of several lichens. The prokaryotic blue-green algae (cyanobacteria) encompass most of the remainder, occurring in *c.* 10% of the nearly 20 000 known lichen associations (Rikkinen 2017). Additionally, two stramenopile algae (a xanthophyte and a phaeophyte) are known to enter into lichen symbioses. The full range of phylogenetic disparity among lichen-forming algae is therefore much wider than that found among the lichen-forming fungi, which all fall within the kingdom's *Dikarya* crown group (mostly *Ascomycota*, with several genera of *Basidiomycota*). Just what common features might permit those disparate algal lineages to form comparable symbioses with lichen-forming fungi remain enigmatic. As colonizers of exposed, subaerial substrata, potentially suitable algae may be pre-adapted to coping with hydric stresses and high radiation loads (Lange *et al.* 1990; Gustavs *et al.* 2010; Candotto Carniel *et al.* 2015). It is striking that most lineages of basidiomycete fungi that independently adopted the lichen lifestyle did not domesticate novel algal genera; instead they chose taxa that associate with ascolichens, such as *Coccomyxa*, *Elliptochloris* and *Rhizonema* (Oberwinkler 2012; Dal Forno *et al.* 2020; Masumoto 2020; but see Hodgkinson *et al.* (2014) concerning *Lepidostromatales*). It is also noteworthy

that quite a number of lichen algae belong to genera (e.g. *Chlorella* s. str., *Coccomyxa*, *Elliptochloris* and *Nostoc*) that include species occurring in symbiosis (often endosymbioses) with diverse protists, plants and animals (Adams *et al.* 2012; Grube *et al.* 2017b).

Algal partners in lichen symbioses were termed phycobionts by Scott (1957). Subsequently, Ahmadjian (1982) proposed that *photobiont* replace *phycobiont* where cyanobacteria are meant to be included, because they 'are not algae per se but actually bacteria'. No further argumentation was provided; it was presumed self-evident that *algae* and *bacteria* must denote mutually exclusive concepts. Some contemporary treatments distinguish cyanobacteria from algae (e.g. Friedl & Büdel 2008; Grube *et al.* 2017b), while others consider them as algae (e.g. Graham *et al.* 2009; Büdel & Kauff 2012; Lee 2018). Clearly, there are significant differences between prokaryotes and eukaryotes. At issue, however, is whether those differences are relevant to the concept of algae. This term has no biosystematic status and cannot attain any by exclusion of the blue-greens. The emblematic algal trait, oxygen-generating photosynthesis, is ultimately derived from cyanobacteria. It was subsequently acquired in multiple events involving primary, secondary and tertiary endosymbioses (Keeling 2004, 2013), and now characterizes diverse lineages included within most of the major eukaryote clades (*Archaeplastida*, *Alveolata*, *Excavata*, *Rhizaria*, *Stramenopila*, *Cryptista* and *Haptista*). The one and only unifying thread in this polyphyletic algal tapestry (Delwiche 1999) is the common photosynthetic apparatus, originating in cyanobacteria and passed on vertically as well as horizontally. The present work therefore uses the term algae to encompass all non-embryophyte lineages that inherited oxygenic photosynthesis. Phycobiont and photobiont are considered synonymous terms.

The Algal Role in Lichen Symbiosis

The algal partner is the primary producer, sustaining the lichen association by supplying the fungal partner with carbohydrate products of photosynthesis (Smith 1974). Those with pyrenoids (Fig. 2) possess CO₂-concentrating mechanisms that improve the efficiency of carbon fixation (Smith & Griffiths 1996). Green algal symbionts (chlorobionts) transfer their photosynthate as polyol sugar alcohols such as ribitol (Richardson *et al.* 1968). Significantly, these compounds also confer desiccation tolerance by providing osmolarity and protecting cell membranes from damage as water is lost (Smith 2019). Polyols are likewise produced by non-symbiotic, aeroterrestrial green algae, particularly under osmotic stress conditions (Darienko *et al.* 2010; Gustavs *et al.* 2010, 2011). Blue-green symbionts (cyanobionts) transfer glucose, or glucan, which their fungal partners take up and immediately convert into the sugar alcohol mannitol (Smith & Drew 1965; Hill 1972). When lichenized, the algal symbionts are somehow induced to leak large amounts of carbohydrate to the surrounding fungal cells, a process that quickly ceases when the algae are isolated into culture (Drew & Smith 1967). Fungal penetration of photobionts may occur to varying degrees (Geitler 1934; Tschermak 1941a; Plessl 1963; Galun *et al.* 1970, 1971; Honegger 1986; Matthews *et al.* 1989), but these so-called haustoria do not appear to be principal conduits of carbohydrate transfer in ascolichens (Jacobs & Ahmadjian 1971; Collins & Farrar 1978; Hessler & Peveling 1978). The intrusive hyphae of certain basidiolichens that deeply penetrate longitudinally through the centre of their cyanobiont trichomes (Roskin 1970; Oberwinkler 1980, 2012) have not yet been examined with respect to substance transfer. In most foliose and fruticose lichens examined, haustorial penetrations are either absent altogether or do not fully traverse the

algal cell wall. To facilitate transfer, the mycobiont secretes a hydrophobic sealant that envelops the cell surfaces of both symbionts at their contact zones, thereby funnelling carbohydrate released by the alga to the fungus (Honegger 1991; Trembley *et al.* 2002a). At least that is the case in the selection of taxa examined so far. Where cyanobacterial symbionts are involved, they provide the lichen fungus with fixed nitrogen as well as carbon (Millbank & Kershaw 1974). In those lichens (chiefly *Peltigerales*) where a chlorobiont constitutes the main algal layer and cyanobionts are localized within nodules known as cephalodia, the cyanobacteria become highly specialized for nitrogen fixation, with an elevated percentage of cells differentiating as heterocysts (Hitch & Millbank 1975). In lichens with only cyanobacterial photobionts, heterocyst frequency can be much lower at the growing margins of the thallus (Bergman & Hällbom 1981), where photosynthate may be in higher demand.

Whether any substance is transferred from fungus to alga in exchange has yet to be demonstrated. At least some genes relevant to such metabolic transfers appear to be differentially expressed in symbiosis (Kono *et al.* 2020). Certainly, there has been speculation that the fungal partner might apportion carbohydrate, nitrogen, or other substances back to the algal symbiont to regulate its growth (Ahmadjian 1995) in coordination with that of the mycobiont (Greenhalgh & Anglesea 1979; Hill 1985, 1989; Honegger 1987). The heterotrophic tendencies shown by many lichen algae (*Trebouxia*, *Asterochloris*, *Elliptochloris*, *Coccomyxa*, *Apatococcus*) when cultured in the laboratory (Ahmadjian 1993; Gustavs *et al.* 2016, 2017) suggest the possibility that they could be susceptible to such control. Indeed, Ahmadjian (2001) proposed that *Trebouxia* is fully dependent upon its mycobiont for nutrition and is therefore unable to survive in the free-living state (Ahmadjian 1988).




Fig. 2. TEM micrographs of some photobiont pyrenoids, with plastoglobuli (round black dots) and penetrating membranes in various positions and orientations. A, *Trebouxia*, within thallus of *Lasallia pustulata*. Note pyrenoid structure here more closely resembles that of distantly related *Heveochlorella* (B) than that of another species (C) of *Trebouxia*. B, *Heveochlorella*, within thallus of *Calopadia*. C, *Trebouxia*, within thallus of *Ramalina usnea*. D, bulging exserted pyrenoid of *Petroderma maculiforme*. E, *Diplosphaera*, within thallus of *Endocarpon pusillum*. S = starch grain or plates. Scales: A = 1 μ m; B = 200 nm; C–E = 500 nm.

However, he also promoted the seemingly contradictory viewpoint that *Trebouxia* is a victim of fungal parasitism rather than a mutualist partner (Ahmadjian 1993, 1995, 2002). This would make *Trebouxia* a host that cannot survive without its parasite.

In any event, proof of fungus-to-alga nutrient transfer is not required to make the case that lichen symbiosis offers advantages to the algal partner. There is considerable evidence that the surrounding fungal tissues and their secondary metabolites may help protect the lichenized alga from desiccation, photoinhibition, temperature extremes and herbivory (e.g. Solhaug & Gauslaa 1996; Kranner *et al.* 2008; Kosugi *et al.* 2009; Asplund & Wardle 2013; Gauslaa *et al.* 2017; Míguez *et al.* 2017; Sadowsky & Ott 2016; Beckett *et al.* 2019; Fernández-Marín *et al.* 2019). Symbiosis may significantly improve the alga's ability to avoid cellular damage caused by highly reactive forms of oxygen (ROS) generated under stress conditions (Kranner *et al.* 2005). With these protections, and the facilitated display for light capture afforded by a supportive mycobiont structure, lichen algae may greatly expand their ecological range and abundance via symbiosis (Honegger 2012). On the other hand, lichen symbioses are diverse and it is likely that the parameters of the relationship vary among taxa, along environmental gradients, and perhaps also during the course of a single lichen's development. The long history of attempts to maintain or resynthesize lichens in the laboratory has provided a key insight into the nature of this seemingly well-integrated association: it is very much a relationship of contingency. That the partners can often be cultured separately on appropriate media in the laboratory (Ahmadjian 1993; Crittenden *et al.* 1995; Stocker-Wörgötter & Hager 2008) shows there is no strict physiological impediment to growth without symbiosis. To initiate and support lichen formation, a fluctuating balance of conditions suboptimal for separate fungal or algal growth appears to be necessary. Any combination of culture conditions (light, moisture, nutrient availability) that continuously favours either fungal or algal growth results in the breakdown of symbiotic structures, and the dissociated proliferation of the micro-organisms separately (Thomas 1939; Scott 1960; Ahmadjian 1962; Stocker-Wörgötter 2001; but see Marton & Galun 1976). It therefore seems logical to view the lichen symbiosis as a more or less mutualistic response to conditions that permit neither partner to thrive independently.

Although both partners may derive benefits, the lichen symbiosis is clearly not symmetrical (Hill 2009). The heterotrophic mycobionts, with their elaborate structural adaptations for algal cultivation, are more fully committed to symbiosis than their trophically autonomous photobionts. The mycobiont frees itself of symbiosis only in spore dispersal, seeking algal partners again immediately upon germination. To carry out sexual reproduction, it must be in symbiosis, whereas its photobiont appears to need aposymbiotic freedom to do so. From the alga's point of view, whenever unfavourable conditions reduce its possibilities of aposymbiotic success, the benefits of lichenization may begin to outweigh any disadvantages. Photobionts may rely on lichen symbioses for long-term persistence in habitats periodically subject to adverse conditions, while needing intervals of independence under favourable conditions to complete their life cycles. Thus, mycobiont and photobiont life histories do not fully coincide, but produce a lichen where they intersect compatibly. To varying degrees, natural selection has optimized the mycobiont principally for symbiosis, the photobiont for autonomy as well as symbiosis. The trade-off is that greater adaptation to symbiotic compatibility is likely to

constrain the possibilities for competitive success in the aposymbiotic state. However, the lingering notion that certain photobionts may not ever occur free-living is probably attributable to insufficient sampling, and the conflation of invisibility with absence. Unsurprisingly, those photobionts that are macroscopically visible (*Nostoc*, *Cephaleuros*, *Phycopeltis*, *Trentepohlia*, *Prasiola*, *Petroderma*) have not had their aposymbiotic occurrence disputed.

Both fungus and alga must adapt, at least to some extent, to be compatible symbionts. For some authors, such mutual adaptation constitutes coevolution (Ahmadjian 1987; Saini *et al.* 2019); for others, coevolution supposes parallel cladogenesis in partners' phylogenies, a criterion not generally met by lichen symbioses analyzed in this regard (Piercey-Normore & DePriest 2001; Stenroos *et al.* 2006). However, it has been argued that focusing exclusively on this fine scale ignores broader patterns of co-adaptation, whereby 'guilds' of different mycobionts share common pools of photobionts to mutual advantage (Rikkinen 2003, 2013). According to Hill (2009), photobionts cannot coevolve with their mycobionts because they lack sexual reproduction in the thallus, are not subject to natural selection from one lichen to the next, and are not perpetuated when a lichen thallus dies. Yet photobionts are continually escaping from lichen thalli by means of soredia, isidia, thallus fragments, co-dispersed hymenial, epithelial or conidiomatal algae (Fig. 3), and the excreta of lichenivorous invertebrates (Frøberg *et al.* 2001; Meier *et al.* 2002; Boch *et al.* 2011). Such diaspores afford many chances of finding microconditions where independent algal growth is favoured; aposymbiotic, potentially sexual populations may then develop, be they brief or enduring. Selection among genotypes for compatibility (or resistance) will occur when the opportunity for relichenization next presents itself. Compatible genotypes incorporated into a developing lichen may then be subject to further winnowing selection in the course of thallus growth.

Patterns of Symbiont Pairing

The asymmetrical needs of the lichen symbionts are reflected in the non-reciprocal patterns of pairing that have evolved between mycobionts and photobionts. Photobiont genera frequently associate with multiple, phylogenetically disparate lineages of lichen-forming fungi. The converse, however, is much less common; mycobiont genera, and often families and even orders, generally tend to lichenize a single algal genus (Rambold *et al.* 1998; Peršoh *et al.* 2004). There are a number of notable exceptions. Lichen-forming fungi of the family *Verrucariaceae* partner with an extremely diverse array of eukaryotic algae, including the only reported cases of stramenopile phycobionts (Thüs *et al.* 2011). The pin-lichen genus *Chaenotheca* (*Coniocybomyces*) includes species associating with *Trebouxia*, *Trentepohlia*, *Symbiochloris* or *Tritostichococcus* (Tibell 2001; Škaloud *et al.* 2016; Pröschold & Darienko 2020). The fruticose lichen genus *Stereocaulon* may harbour thallus photobionts of either *Asterochloris*, *Vulcanochloris* or *Chloroidium* (Vančurová *et al.* 2018). Species of *Sticta* may partner with chlorobionts of *Symbiochloris*, *Coccomyxa*, *Elliptochloris*, *Heveochlorella* or *Chloroidium* (Lindgren *et al.* 2020). Squamulose *Psora decipiens* is reported to partner with either *Asterochloris*, *Trebouxia*, *Chloroidium* (Ruprecht *et al.* 2014) or *Myrmecia* photobionts (Williams *et al.* 2017; Moya *et al.* 2018). In addition, it is well known that many individual mycobionts, particularly in the *Peltigerales*, may associate with both green and blue-green algae simultaneously, giving rise to cyanobacterial cephalodia within or upon a chlorophyte-




Fig. 3. Liberation and potential co-dispersal of photobionts from the spore-producing structures of certain mycobionts. A, *Diplosphaera* photobiont (arrows) within perithecium of *Endocarpon pusillum*; note much smaller size compared to photobiont cells within thalline tissue (t); s = ascospore. B, apothecial surface of foliicolous lichen colonizing plastic cover slip; note epithelial algal cells (arrows) among emerging ascospores (s). C, *Heveochlorella* photobionts (vertical arrow) within conidiogenous tissue of campylidia and intermixed among filiform macroconidia (oblique arrow). D, hyphophore of *Gyalectidium paolae* showing diahyphal propagules (bundles of conidial chains dispersed as a unit) with adhering or intermixed *Heveochlorella* photobionts (arrows). E, campylidial macroconidia, with co-dispersed *Heveochlorella* photobionts loosely encircled, germinating (arrowheads) on a plastic cover slip. F, diahyphal propagules of *Gyalectidium* germinating (arrowheads) on a plastic cover slip, with co-dispersed *Heveochlorella* photobionts. Scales: A, C & D = 20 μm ; B = 50 μm ; E & F = 10 μm .




Fig. 4. Dichotomously lobed chloromorphs of *Sticta canariensis* emerging from lower surfaces of cyanomorph thalli (arrows). Scale = 5 mm.

containing thallus, or distinct cyanomorph and chloromorph thalli separately or conjoined (Fig. 4) via a common fungal individual (e.g. James & Henssen 1976). Association with both a chlorobiont and a cyanobiont in separate thallus components has also been reported for certain basidiolichen species in *Cyphellostereum* (Oberwinkler 2012) and *Lichenomphalia* (Gasulla *et al.* 2020). In a small number of lichens, green and blue-green photobionts are known to occur intermixed within the same thallus structure (Büdel & Henssen 1987; Henskens *et al.* 2012). There are distinct physiological advantages to each of these two kinds of photobionts. Cyanobionts can fix nitrogen as well as carbon but require liquid water to rehydrate and resume physiological activity, whereas chlorobionts can rehydrate from vapour, although their CO₂ fixation rates may be more adversely affected by high thallus water contents (Lange *et al.* 1986, 1993; Green *et al.* 1993, 2002). Less obvious are the implications of choosing *Trentepohlia* (*Ulvophyceae*) versus *Trebouxia* (*Trebouxiophyceae*) photobionts; neither fix nitrogen, although they may differ in their tolerance of freezing temperatures (Nash *et al.* 1987). Interestingly, mycobiont genera *Ionaspis* and *Hymenelia* (*Lecanoromycetes*) include trentepohliophilic and trebouxiophilic taxa, and the single species *H. epulotica* can apparently associate with photobionts of either of these two very different genera (Lutzoni & Brodo 1995; McCune *et al.* 2018). Recently, Ertz *et al.* (2018) demonstrated that the lichen fungus *Lecanographa amylicata* can form morphologically distinct sexual and asexual thalli with *Trentepohlia* and *Trebouxia* photobionts, respectively. While the above examples show that significant divergences in photobiont selection have arisen in a number of mycobiont lineages, far more conservative tendencies appear to predominate in the majority of lichen-forming fungal groups.

Photobiont choice and the range of compatible pairings for a given mycobiont were first explored experimentally in classic laboratory resynthesis studies using *Cladonia cristatella* and *Lecanora chrysoleuca* (Ahmadjian *et al.* 1980; Ahmadjian & Jacobs 1981). Varying degrees of compatibility were observed, with thallus formation reaching different developmental stages depending on the photobiont strain introduced. Nonetheless, overall results generally reflected patterns observable in natural lichens:

Cladonia successfully lichenized strains of *Asterochloris* but not those of *Trebouxia* (as currently defined), while *Lecanora* did just the opposite. In the last two decades, genetic markers have been used to characterize the range of photobiont diversity chosen by individual lichen-forming fungal species in nature, and to assess the parameters that might affect their choices. This complex topic has attracted much attention and merits a review of its own, but some general findings can be summarized here. Most mycobiont species appear to be fairly selective; they tend to partner with a limited range of strains or species within a single photobiont genus, but to differing degrees. Some mycobionts accept a substantially broader range of taxa within the photobiont partner genus; this relative liberality is often characteristic of lichen-forming fungi that have attained wider, more cosmopolitan distributions (Blaha *et al.* 2006; Guzew-Krzemińska 2006; Leavitt *et al.* 2013; Muggia *et al.* 2014; Magain *et al.* 2017; Vančurová *et al.* 2018), or those capable of colonizing extreme environments with probably fewer photobiont options available (Romeike *et al.* 2002; Wirtz *et al.* 2003; Engelen *et al.* 2010; Pérez-Ortega *et al.* 2012; Osyczka *et al.* 2021; Rola *et al.* 2021). Such mycobionts may be closely related to species that accept a much narrower range of photobiont partners (Piercey-Normore 2004; Yahr *et al.* 2004; Otálora *et al.* 2010; Onuț-Brännström *et al.* 2017). Some studies have correlated symbiont selection patterns with environmental parameters, such as latitude (Singh *et al.* 2017), climate (Řídká *et al.* 2014) and ecological conditions that influence the distribution and availability of photobionts (Yahr *et al.* 2006; Fernández-Mendoza *et al.* 2011; Peksa & Škaloud 2011; Vargas Castillo & Beck 2012; Werth & Sork 2014). Photobiont tolerance of heavy metals appears to influence their selection by mycobionts in some lichen communities colonizing metal-rich substrata (Vančurová *et al.* 2018; Rola *et al.* 2021) but not others (Beck 2002; Hauck *et al.* 2007; Bačkor *et al.* 2010). Many studies stress the intrinsic compatibility requirements of individual fungal taxa as primary determinants of pairing patterns (Yahr *et al.* 2004; Stenroos *et al.* 2006; Myllys *et al.* 2007; Leavitt *et al.* 2015; Joneson & O'Brien 2017), often in conjunction with ecological factors (Elvebakk *et al.* 2008; O'Brien *et al.* 2013; Dal Grande *et al.* 2018; Jüriado *et al.* 2019; Pino-Bodas & Stenroos 2020). In some communities, mycobionts may have adapted to utilize a common pool or pools of photobionts, whose local availability might thereby be sustained for all users (Beck *et al.* 2002; Rikkinen *et al.* 2002; Rikkinen 2003; Sanders *et al.* 2016; Onuț-Brännström *et al.* 2018; Cardós *et al.* 2019). Thallus growth form may also affect photobiont selection patterns. Some authors have suggested that crustose lichens may associate with a broader range of photobionts than do related foliose and fruticose taxa (Helms *et al.* 2001), perhaps because their more extensive and intimate contact with the substratum offers more opportunity to take up additional algae in the course of development. Lichen reproductive mode can also be superimposed upon these factors. Some studies have found that lichens reproducing primarily by vegetative propagules, such as soredia or isidia, associate with a narrower range of photobiont genotypes, presumably due to chiefly vertical transmission of both symbionts together (Dal Grande *et al.* 2012; Werth & Scheidegger 2012; Otálora *et al.* 2013; Cao *et al.* 2015; Hestmark *et al.* 2016; Steinová *et al.* 2019). However, other vegetatively reproducing lichens accept a much broader range of photobionts, suggesting that the fungus does not necessarily maintain partnership with its co-dispersed photobiont throughout development (Ohmura *et al.* 2006, 2019; Nelsen & Gargas 2008, 2009; Wornik & Grube 2010).

Acquisition of New Algal Symbionts

Acquisition of new and different photobionts, 'photobiont switching', has clearly been significant in the evolution of lichen relationships. However, this phrase may refer variably to events occurring at different levels of organization. A single mycobiont individual might acquire new photobionts at different times in the course of its development (Friedl 1987; Wedin *et al.* 2016), or at separate places along its somatic extension (Létrouit-Galinou & Asta 1994). The degree to which the newly lichenized alga may differ genetically from algal strain(s) already in possession will be limited by the innate compatibility range of that mycobiont individual. In contrast, a new fungal individual developing from a meiospore may encounter and select a photobiont strain different from the one its parental genotypes associated with. In this case, a generational change in photobiont partner could be enabled by a generational change in mycobiont genotype. At a phylogenetic level, a cladogram may provide evidence that a fungal lineage has changed its association from one photobiont to another in the course of evolution. But at a finer scale, a great many photobiont switches, perhaps back and forth, might have taken place over many generations; comparing taxa will indicate only the overall result.

New photobionts may be acquired in multiple ways. Contact and capture of free-living photobionts in nature by hyphae emerging from germinated spores (Fig. 5), once thought to be unlikely (Lamb 1959), has been documented in a number of studies (Ward 1884; Werner 1931; Bublick *et al.* 1984; Garty & Delarea 1988; Scheidegger 1995; Sanders & Lücking 2002; Sanders 2014). In theory, a single compatible algal individual might be sufficient to generate the entire population within a developing thallus. However, there appear to be many opportunities for additional photobionts to be incorporated from exterior sources. Particularly in early developmental stages, prothallial hyphae extending outward along the substratum from the

lichenized portions of the organizing thallus can incorporate additional algal cells (Sanders & Lücking 2002; Sanders 2014). Vegetative propagules, such as soredia or isidia, also begin development with the emergence and proliferation of such hyphae (Jahns *et al.* 1979; Schuster *et al.* 1985), anchoring the structure and greatly expanding the available surfaces for potential contact with other compatible photobionts as the thallus is organized. In many crustose lichens, a prothallus remains active at the growing margins of the lichen and may continue to incorporate compatible photobionts falling upon it or encountered on the substratum (Fig. 6; see also Galløe 1927: p. 40, 1932: p. 78; Letrouit-Galinou & Asta 1994). The multitude of discrete, lichenized units that comprise the thallus of squamulose lichens probably also arise from repeated algal capture by a network of prothallial hyphae interconnecting the squamules. Certain soil- and rock-colonizing squamulose lichens produce hyphal aggregates (cords or rhizomorphs) of indeterminate growth that penetrate the substratum extensively (Poelt & Baumgärtner 1964; Sanders *et al.* 1994), giving rise to new thallus squamules where compatible algal symbionts are encountered and lichenized (Wagner & Letrouit-Galinou 1988; Sanders & Rico 1992; Sanders 1994). The structurally similar rhizomorphs of certain umbilicate lichens also appear to have this capability (Schuster 1992). In some foliose and fruticose lichens, organized thallus surfaces may themselves be capable of incorporating compatible algal cells that make external contact (Bitter 1904). Lichens that form cephalodia and/or joined chloromorph and cyanomorph thalli clearly retain this ability (see discussion under *Nostoc* below). Additionally, certain lichen-forming fungi appear capable of obtaining photobionts from other lichens, upon which their spores may germinate (Hawksworth *et al.* 1979). The host thallus is eventually destroyed as its photobionts are taken over by the invading hyphae of the aggressor, giving rise to a new lichen (Poelt 1958; Friedl 1987; Feige *et al.* 1993; Lücking & Grube 2002; Wedin *et al.* 2016). Thus, capture of free-living algae by spore germlings is clearly not the only opportunity for a mycobiont to acquire new photobionts. On the other hand, some interesting transplant experiments with *Psora decipiens* suggest that lichens may not always be able to switch to more favourable photobionts when needed (Williams *et al.* 2017).

If acquisition of additional photobionts is indeed a common occurrence in the course of lichen development, lichen thalli may be expected to contain a heterogeneous photobiont population, at least at certain stages. Some authors have observed and illustrated quite different chlorobionts occurring together within single thalli (Voytsekhovich *et al.* 2011). Data from molecular markers have also addressed this question. Some authors found no evidence of multiple photobiont genotypes in single thalli examined (Paulsrud & Lindblad 1998; Beck & Koop 2001; Singh *et al.* 2017; Škaloud *et al.* 2018); others found occasional occurrences (Guzow-Krzemińska 2006; Bačkor *et al.* 2010; Muggia *et al.* 2013; Nyati *et al.* 2013; Řídka *et al.* 2014; Onuț-Brännström *et al.* 2018; Vančurová *et al.* 2018; Molins *et al.* 2020), or frequent presence (Piercey-Normore 2006; Muggia *et al.* 2014; Park *et al.* 2015; Dal Grande *et al.* 2018; Osyczka *et al.* 2021). Intrathalline populations of *Trebouxia* can also vary in simple sequence DNA regions, which may result from clonal replication errors (Mansournia *et al.* 2012; Dal Grande *et al.* 2014a). Individual thalli of *Parmotrema pseudotincturum* from the Canary Islands were reported to encompass distinct lineages of *Trebouxia* as well as *Asterochloris* (Molins *et al.* 2013). According to Casano *et al.* (2011), two genetically distinct strains of *Trebouxia* are always present together in thalli of



Fig. 5. Muriform ascospore (a), probably of *Calopadia*, germinating on a plastic cover slip placed in a south-west Florida oak hammock, and lichenizing a group of algal cells (arrow), most likely *Heveochlorella*. Scale = 20 μ m.




Fig. 6. *Phycopeltis* free-living and in stages of lichenization. A, free-living. B, edge of developed *Phycopeltis* thallus (left) lichenized by a network of hyphae (probably foliicolous *Porina* sp.) that extend over substratum and capture additional young *Phycopeltis* germlings (arrows). Scales: A = 20 μ m; B = 10 μ m.

Ramalina farinacea, and high-throughput sequencing results suggest that a number of other, minority algae might also be present in this lichen (Moya *et al.* 2017). One constant challenge in assessing photobiont identities is that lichen thallus surfaces are colonized by epibiontic algae (including possible photobionts of other lichens) that are not intimate symbionts of the lichen in question, yet may figure prominently in cultures established or samples obtained from thallus fragments (Warén 1920; Muggia *et al.* 2013). Confidence that sampled algae are indeed the thallus photobionts can be improved by establishing cultures from single algal cells extracted from within the thallus using a micromanipulator (Beck & Koop 2001), although the procedure is time-consuming. Additional evidence may be sought in TEM micrographs of photobionts within the same thallus (e.g. Catalá *et al.* 2016; Molins *et al.* 2018), particularly where more than one pyrenoid type (Friedl 1989) is present. However, variability should first be assessed among individuals of the same genetic strain because chloroplast structure may vary from cell to cell and often looks substantially different according to the plane of ultrathin section examined. In sequencing, conventional dideoxy chain termination (Sanger) technology will reliably identify a predominant photobiont and ignore any others present in low abundance, while the procedure fails if there are secondary photobionts in sufficient abundance (c. 30%; Paul *et al.* 2018). High-throughput sequencing will detect minority photobionts but will also be more sensitive to epibiontic algae. A recent comparison of the two sequencing approaches concluded that in most lichens there is a single dominant photobiont genotype, representative of most of the thallus population (Paul *et al.* 2018).

The Genera of Lichen Algae

Approximately 50 algal genera are currently said to include lichen photobionts. Some may represent identifications that are erroneous or based on outdated circumscriptions of taxa. Others may spin off new genera as their cryptic genetic diversity is further elucidated. It is evident that a small number of very prominent photobiont genera (*Asterochloris*, *Nostoc*, *Rhizonema*, *Trebouxia*, *Trentepohlia*) each partner with many hundreds or thousands of lichen-forming fungal species; a number of others (e.g. *Coccomyxa*, *Elliptochloris*, *Heveochlorella*, *Symbiochloris*) are lichenized by many dozens or hundreds of different mycobiont species, while much of the remainder participate in only a small number of known lichen associations. It seems probable that further surveys will uncover more photobiont genera in the latter category. While it is widely agreed that the diversity of

lichen-forming algae remains considerably less well known than that of lichen-forming fungi, this fact alone is unlikely to account for the enormous disparity between the currently recognized number of photobiont genera (c. 50) and that of mycobiont genera (c. 1000; Lücking *et al.* 2017a). The number of photobiont species described, estimated at c. 100 not long ago (Škaloud & Peksa 2010), shows a similar disparity with the number of lichen-forming fungal species (20 000). Indeed, both the generic and species estimates differ between mycobiont and phycobiont by the same factor of 20. Thus, the imbalance is not likely due to differences in genus/species concepts between algae and fungi. Of course, much of the genetic diversity discovered within photobiont genera in the last few years has been reported as clades that still lack taxonomic recognition; species numbers will surely increase substantially in the near future as such diversity becomes formalized biosystematically. However, this still seems unlikely to close the enormous gap with mycobiont species numbers. Rather, the disparities probably indicate a real ecological asymmetry: the large number of lichen-forming fungal taxa may be partnering with a substantially smaller pool of photobiont taxa, many of which are shared among mycobionts. Such was the conclusion reached recently by Dal Forno *et al.* (2020) in their detailed comparison of genetic diversity in *Dictyonema* and its *Rhizonema* photobionts.

A synopsis of algal genera to which lichen photobionts are currently attributed is given below.

Cyanobacteria

Anabaena Bory ex É. Bornet & C. Flahault — See *Nostoc*. Strains of *Anabaena* versus *Nostoc* are resolved in some analyses (Henson *et al.* 2002; Rajaniemi *et al.* 2005; Liu *et al.* 2013; Elshobary *et al.* 2015) but formal distinction of the two genera remains controversial (Makra *et al.* 2019). Tschermak-Woess (1988a) recommended re-examination of earlier reports that *Anabaena* occurs as cephalodial photobiont of *Stereocaulon*.

Anacystis Meneghini — According to Bold & Wynne (1985), this generic name has been applied to ellipsoid to cylindrical cyanobacteria that often accumulate in a common gelatinous matrix, with some authors also including spheroidal-celled taxa such as *Gloeocapsa* and *Chroococcus*. The much-studied '*Anacystis nidulans*' is usually treated now under *Synecococcus*; other taxa are currently placed in *Microcystis*. Photobionts attributed to *Anacystis* in the past include the partners of a small number of *Peltula* species and the cephalodial symbionts of a

Stereocaulon (see Tschermak-Woess 1988a); determining their identities with confidence will require further study.

Brasilonema Fiore *et al.* — This cyanobacterial genus, forming a distinct clade in molecular analyses (Fiore *et al.* 2007), has aggregated filaments morphologically similar to *Scytonema* but only rarely showing false branching. A recent paper reported new species of both *Brasilonema* and *Chroococcidiopsis* as co-occurring photobionts of an unidentified lichen growing on gravestones in a northern Florida cemetery (Villanueva *et al.* 2018). However, as no description or evidence of this association has yet been published, the status of *Brasilonema* as lichen photobiont awaits corroboration.

Calothrix C. Agardh ex É. Bornet & C. Flahault and *Dichothrix* G. Zanardini ex É. Bornet & C. Flahault — These filamentous cyanobacteria are members of the *Rivulariaceae*; their trichomes have a basal heterocyte and gradually narrow towards the apex. The two genera are morphologically similar and both have been reported as lichen photobionts, particularly in association with certain species of *Lichina* (see Tschermak-Woess 1988a). However, DNA sequences obtained from two such examples instead placed the algae in question in the genus *Rivularia* (Ortiz-Álvarez *et al.* 2015). The photobiont of *Placynthium nigrum* isolated into culture also shows the distinctive *Rivulariaceae* morphology (apically tapering filaments with basal heterocytes) while the lichenized filaments rather resemble those now placed in *Rhizonema* (see Geitler 1934). The circumscription of *Calothrix* and *Dichothrix* with respect to lichen photobionts currently remains unresolved.

Chroococcidiopsis Geitler (and *Myxosarcina* H. Printz) — These unicellular cyanobacteria are found in a great diversity of habitats and include extremophiles. Cells divide in sequence by binary fission, often in alternating planes to produce more or less cubical packages of cells. Cells can also undergo multiple fission to produce four or more autospore-like products known as baeocytes, initially contained within the sheath-like, fibrous outer wall layer of the mother cell (Waterbury & Stanier 1978). The baeocytes of *Myxosarcina*, unlike those of *Chroococcidiopsis*, have a brief stage of gliding motility; the genera are said to be otherwise indistinguishable morphologically. The baeocyte-forming cyanobacteria were formerly grouped together in the order *Pleurocapsales* (Waterbury & Stanier 1978), but SSU sequence analysis has shown this trait to be a convergence shared by a number of lineages of quite different origin (Fewer *et al.* 2002). In that study, several photobionts isolated from *Lichinaceae* appear within the same clade as *Chroococcidiopsis thermalis*, sister to the heterocyte-forming *Stigonematales* and *Nostocales*, and distant from *Myxosarcina* as well as other morphologically similar taxa formerly attributed to *Chroococcidiopsis* (Fewer *et al.* 2002). Sequences obtained from photobionts of several *Peltula* species collected in Vietnam also suggested affinities within a broad ‘*Chroococcidiopsidales*’ clade (Võ 2016). Other algal partners of *Lichinaceae* have been attributed to *Chroococcidiopsis* based on morphology and the production of baeocytes observed in cultured isolates (Büdel & Henssen 1983). Tschermak-Woess (1988a) suggested that some taxa identified as *Chroococcidiopsis* might actually belong to *Gloeocapsa* and require study in culture. Most photobiont isolates attributed to *Chroococcidiopsis* and *Myxosarcina* await more detailed molecular scrutiny.

Chroococcus Nägeli — A morphologically distinctive cyanobacterial genus, *Chroococcus* has relatively large, spherical cells that divide at consecutive right angles to produce small packets of cells, often within concentric, gelatinous sheath layers. A number of reports, compiled by Tschermak-Woess (1988a), attribute thallus and cephalodial photobionts of various lichens to this genus or merely to *Chroococcaceae*, or *Chroococcales*. Many are anecdotal and most await reinvestigation with molecular sequence comparisons. The photobionts of certain *Dictyonema* species, once attributed to *Chroococcus*, have been shown to belong instead to *Rhizonema*, a usually filamentous taxon that may be greatly altered morphologically in certain lichen associations (Lücking *et al.* 2009). The circumscription of *Chroococcus* and its status as a lichen photobiont genus remain uncertain at present.

Gloeocapsa Kützing — This colonial cyanobacterium has roundish to oblong cells surrounded individually and communally by successive layers of dense mucilage, reflecting the sequence of cell divisions. Morphologically defined at present, *Gloeocapsa* commonly occurs free-living in moist terrestrial habitats and is also reported as thallus photobiont in several genera of *Lichinaceae*, and as cephalodial symbiont in certain species of *Stereocaulon* and *Amygdalaria* (Tschermak-Woess 1988a). In the lichen *Gonohymenia*, contacting mycobiont hyphae broadly invaginate the cells of its photobiont, identified as *Gloeocapsa* (Paran *et al.* 1971). Geitler (1933) described appressorial hyphae in the lichen *Synalissa* that branch in synchrony with the binary fission of its *Gloeocapsa* photobiont.

Molecular sequence data are much needed to understand the relationship among taxa currently assigned to *Gloeocapsa*.

Hyella É. Bornet & C. Flahault — The filamentous cyanobacterium *Hyella* is a widespread inhabitant of the marine intertidal zone, where it colonizes calcareous substrata such as mollusc shells. The substratum is penetrated by threads arising from a basal system at the surface; endospore-like baeocytes may be formed (Fritsch 1945). Genomic analysis shows *Hyella* phylogenetically nearest to the genus *Chroococcidiopsis* (Brito *et al.* 2020). *Hyella* is reported to be the photobiont of some species of fungi now assigned to *Collemopsisidium* (Mohr *et al.* 2004). However, details of the symbiotic interaction are few; other genera of cyanobacteria, such as *Gloeocapsa* and *Nostoc*, are also said to be photobionts for *Collemopsisidium* [= *Pyrenocollema*] (Purvis *et al.* 1992).

Hyphomorpha A. Borzi — These seldom encountered cyanobacteria occur as epiphytes upon tropical liverworts and tree bark, where they form a prostrate filament system. The filaments have an apical cell producing derivatives that may later divide periclinally to become pluriseriate, as do structurally similar species of *Stigonema*. Cells of these older portions tend to fall out of alignment and become jumbled into a ‘chroococcoid stage’ (Fritsch 1945). *Hyphomorpha* was first identified as photobiont in two species of *Spilonema* lichens by Henssen (1981), who reported confirmation of the alga’s identity by eminent phycologist Lothar Geitler. One of these mycobiont species has been recently reclassified as *Erinacellus dendroides* (Spribille *et al.* 2014). At present, the algal genus *Hyphomorpha* is phenotypically defined; it is currently placed in *Fischerellaceae* (Büdel & Kauff 2012) or included under *Hapalosiphonaceae* (Komárek *et al.* 2014) within the *Nostocales*.

Nostoc Vaucher ex É. Bornet & C. Flahault — This genus accommodates cyanobacteria occurring worldwide in fresh water and upon soil, bark and low-growing plants, with some strains highly desiccation-tolerant (Dodds *et al.* 1995). Phenotypically defined at present, taxa attributed to *Nostoc* fall within several distinct clades of the *Nostocales*, making the genus polyphyletic (Rajaniemi *et al.* 2005; Gagunashvili & Andrésson 2018). These algae typically form darkly pigmented, mucilaginous macrocolonies of highly variable size and shape, ranging from spheres to irregularly pustulose mats to tangles of cord-like axes. Embedded within the gelatinous matrix are uniseriate trichomes markedly constricted at the cross walls, giving individual cells an almost spherical to barrel-shaped form and the filaments a characteristic string-of-beads appearance. Cell division is diffuse, without apical cells or directional polarity. At intervals along the chain of vegetative cells are slightly larger, thicker-walled, lighter-coloured heterocytes (heterocysts) that specialize as centres of nitrogen fixation. Since the enzyme involved in this process is inhibited by the presence of oxygen, heterocytes lack oxygen-generating Photosystem II (Wolk *et al.* 1994); electron donors are imported and fixed nitrogen is exported via microplasmodesmatal connections with neighbouring vegetative cells (Giddings & Staehelin 1981; Kumar *et al.* 2010). Thus, prokaryotic *Nostoc* and its heterocytic relatives show degrees of cell specialization and intercellular transport characteristic of true multicellular organization (Garcia-Pichel 2009).

Nostoc, like many filamentous cyanobacteria, has a motile phase. Short filament segments known as hormogonia are produced by multiple divisions of the vegetative cells between two heterocytes, then break free (Boissière *et al.* 1987; Paulsrud 2001). The segments disperse or migrate directionally by a gliding motion that involves secretion of polysaccharide, against which proteinaceous pili appear to push or pull the trichome (Khayatan *et al.* 2015). Under favourable conditions, the hormogonia lose motility and differentiate heterocytes as they transition to vegetative filaments (Paulsrud 2001). It is conceivable that motile hormogonia might facilitate symbiont encounters in the formation of cyanolichens, as also suspected of flagellate stages in eukaryotic photobionts, but direct evidence is lacking. In the establishment of plant-*Nostoc* symbioses, the role of hormogonia as infective agents is well known (Adams *et al.* 2012), and genes related to hormogonial function have been identified in lichen-symbiotic strains (Gagunashvili & Andrésson 2018). *Nostoc* may also disperse temporally by forming akinetes, a kind of resistant spore that develops from a vegetative cell and endures adverse conditions.

Nostoc is photobiont in the majority of cyanophilic lichens. In the *Peltigerales*, *Nostoc* serves as principal thallus photobiont, or as secondary photobiont specialized for nitrogen fixation within discrete structures known as cephalodia; these are formed upon or within a thallus that has a green alga as principal photobiont. In a number of cases, *Nostoc* may serve as both principal and secondary photobiont of a single mycobiont species or individual; this results in cyanomorph and cephalodiate chloromorph thalli that may be either separate or conjoined (James & Henssen 1976; Brodo & Richardson 1978; Tønsberg & Holtan-Hartwig 1983; Armaleo & Clerc 1991; Stenroos *et al.* 2003; Moncada *et al.* 2013; Simon *et al.* 2018). The same strain of *Nostoc* may occur in both morphs (Paulsrud *et al.* 1998, 2001). In many such instances, chloromorph and cyanomorph are both foliose, but in some species

of *Lobaria* and *Sticta*, the *Nostoc*-containing cyanomorph is a branching, fruticose growth that bears no resemblance to the foliose chloromorph (Jordan 1972; James & Henssen 1976; Tønsberg & Goward 2001; Magain *et al.* 2012); when growing separately, the two morphs were long presumed to represent very different taxa. Thallus morphology would appear to be influenced by the distinct photobionts in such cases. In certain species of *Pseudocyphellaria* on the other hand, the independently growing 'cyanomorphs' include numerous clusters of the green algal symbiont (probably *Symbiochloris*) spread among the *Nostoc* within the algal layer (Henskens *et al.* 2012), with no visible alterations to thallus morphology. Even when *Nostoc* serves as a secondary (cephalodial) photobiont in a mature lichen, it may be acquired at a very early stage of lichen formation through contact and capture by the developing mycobiont prothallus (Ott 1988; de los Ríos *et al.* 2011). Once organized, thallus lobes containing green algae may secondarily encounter and incorporate compatible *Nostoc* on the lower surface (Jordan 1970; Jordan & Rickson 1971), or either the upper or lower surface (Cornejo & Scheidegger 2013). Mycobiont selectivity for particular strains of *Nostoc* can be very high (Paulsrud *et al.* 2001). The *Nostoc*-containing cyanomorph may in turn capture compatible green algal symbionts that contact the tomentum hyphae of the lower cortex, from which chloromorph lobes arise (Sanders 2001).

In most lichens where it is primary photobiont, *Nostoc* is confined to a discrete algal layer; its filaments are often broken up or contorted into cell clusters with little secretion of mucilaginous sheath material (Fig. 1J). When isolated into culture, it reverts to the morphology and growth pattern typical of its free-living state (Kardish *et al.* 1989). However, in many of the so-called gelatinous lichens, the form of the *Nostoc* is not fundamentally altered in lichenization; it maintains the necklace-like filaments and extensive surrounding gelatinous sheath, through which the mycobiont hyphae penetrate (Fig. 1I). In such cases, the photobiont constitutes the main structural component of the lichen, which may maintain an appearance and texture rather similar to that of free-living *Nostoc* macrocolonies. A recent study suggests that these differences in phenotypic expression, leading to stratified versus gelatinous lichens, may be associated with different genetic strains of *Nostoc* (Magain & Sérusiaux 2014). This would appear to be another example where major differences in thallus structure may be correlated with photobiont identity.

Cyanophilic mycobionts can be highly selective of their *Nostoc* partner strains, often overriding geographical factors (Paulsrud *et al.* 1998, 2000; Stenroos *et al.* 2006; Myllys *et al.* 2007), although a considerably lower selectivity was observed in lichen communities in maritime Antarctica (Wirtz *et al.* 2003). Within a single clade of *Peltigera*, both highly selective and less discriminating generalist species can be recognized (Magain *et al.* 2017, 2018). A study of temperate and boreal communities reported genetically distinct terricolous and epiphytic pools of *Nostoc*, from which *Peltigera* and *Nephroma* spp. colonizing those respective substrata select their photobionts (Rikkinen *et al.* 2002). Using a larger data set, Stenroos *et al.* (2006) found *Nostoc* photobiont strains to be correlated with mycobiont identity rather than ecological guild. However, fungal preference for the *Nostoc* photobiont strains of other community members over those sampled from the substratum has been reported in other lichen communities (Cardós *et al.* 2019). In other studies, involving

Pannaria and other cyanophilic lichens, both corticolous and saxicolous species sometimes chose closely related strains of *Nostoc*, and more complex combinations of variable mycobiont selectivity and ecological factors were observed (Elvebakk *et al.* 2008).

Nostoc participates in a range of symbioses besides those it forms with lichen-forming fungi (Adams *et al.* 2012). It is taken up by the locally emergent protoplast of the coenocytic, glomeromycete fungus *Geosiphon pyriformis*, which then produces a swollen bladder within which the endosymbiotic (endocytobiotic) *Nostoc* is housed. The intracellular location of the algal symbiont and the close affinities of the fungal component to arbuscular mycorrhizal fungi make the *Geosiphon-Nostoc* symbiosis quite distinct from fungal-algal symbioses treated under the lichen concept (Kluge *et al.* 2002; Schüssler 2012). *Nostoc* also includes obligatory partners of plants representing several major clades of embryophytes; motile hormogonia are the usual infective agent, and fixed nitrogen, usually in the form of ammonium, is supplied to the host from the numerous heterocytes that differentiate in the symbiotic state (Meeks 1998). In hornworts and the liverwort *Blasia*, hormogonia enter and inhabit specialized, mucilage-secreting chambers within the gametophytes (Adams & Duggan 2002). Branched filamentous outgrowths from the inner surfaces of these chambers then develop and increase surface contact between the host and the cyanobacterial colonies (Rodgers & Stewart 1977). In cycad gymnosperms, *Nostoc* colonizes radial cavities in the cortex of specialized, upward-growing coralloid roots (Costa & Lindblad 2002). Symbiosis with the floating aquatic fern *Azolla* is unique in that the *Nostoc* (or *Anabaena*; Svenning *et al.* 2005) is vertically inherited through plant generations, obviating the need for new symbiont capture; the principal cyanobacterium involved cannot be cultivated separately, since its genome shows considerable gene degradation (Ran *et al.* 2010). In the angiosperm *Gunnera*, symbiotic *Nostoc* occurs intracellularly in leaf base tissue (Bergman *et al.* 1992). Some of these symbiotic strains, as well as free-living isolates, appear to be similar or closely related to those occurring within lichen thalli or cephalodia, whereas certain other *Nostoc* strains might be more specialized as lichen photobionts (O'Brien *et al.* 2005; Stenroos *et al.* 2006). Recent genomic comparisons identified certain genes of potential relevance to symbiosis in *Nostoc*, suggesting also that symbiotic strains may have larger genomes than non-symbiotic ones (Gagunashvili & Andr sson 2018).

Rhizonema L cking & Barrie — This cyanobacterial genus was resurrected recently to accommodate filamentous, heterocyte-producing photobionts previously assumed to belong to *Scytonema*, but distinct from that lineage in their 16S rRNA sequences (L cking *et al.* 2009). *Rhizonema* species may be boreal as well as tropical; they are at present known mainly from lichen symbioses but free-living or liverwort-associated populations have also been reported (Cornejo *et al.* 2016). The filaments may be broken up into cell clusters or remain as discrete trichomes (Fig. 1E & F), with sporadic lateral proliferation that has been interpreted as true branching based on the appearance of a mature branch junction (L cking *et al.* 2014). This would presumably distinguish *Rhizonema* from *Scytonema*, which shows false branching. Thus, when V  (2016) observed paired false branching in photobionts of Vietnamese *Cyphellostereum* and *Dictyonema*, she concluded

that the algae were *Scytonema* rather than *Rhizonema*, apparently without corroborating molecular data. However, recent observations of *Rhizonema*, isolated into culture from *Dictyonema* and identified with genetic sequence comparisons, show branching that appears distinctly false (Fig. 1D). Interestingly, a 19th century illustration of a *Dictyonema sericeum* thallus (Bornet 1873: plate 12) depicted the photobiont with both double-false branching and seemingly true branching with a junction similar to that shown in L cking *et al.* (2014). The range of branch development modes possible in *Rhizonema* strains clearly requires further study in both lichenized and aposymbiotic material.

Major genera of lichen-forming fungal partners known so far include *Coccocarpia*, *Erioderma* (Peltigerales), and the basidiomycetes *Acantholichen*, *Dictyonema*, *Cora*, *Corella* and *Cyphellostereum* (all *Hygrophoraceae*). In those basidiolichens, the *Rhizonema* trichome is usually penetrated longitudinally by a single, central mycobiont haustorium quite unlike anything reported in other lichen groups (Roskin 1970; Oberwinkler 1980, 1984, 2012; Slocum 1980; Tschermak-Woess 1983). Such elaborate intrusive structures differ dramatically from the very limited penetrations known in other lichenized algae and might represent specialized absorptive structures. Carbon transfer has not yet been studied in basidiolichens.

Rivularia C. Agardh ex  . Bornet & C. Flahault — The trichomes of this cyanobacterial genus occur in clusters, often on submerged rocks; each filament has a heterocyte at the base and tends to taper gradually towards the apex. The genus includes the photobionts of a couple of maritime species of *Lichina*, whose algal symbionts were previously attributed to the morphologically similar genus *Calothrix* (Ortiz- lvarez *et al.* 2015).

Scytonema C. Agardh ex  . Bornet & C. Flahault — This aquatic or aerophilic genus of cyanobacteria has trichome walls unconstricted at the septa, with vegetative cells usually wider than long, prominent heterocytes, and thick sheaths that are often darkly pigmented. *Scytonema* is traditionally recognized by the frequently paired ('double') false branches, where segments created by a break in the trichome continue linear growth by simply reorienting laterally and emerging from their formerly common sheath. Trichome breaks may arise where intercellular material is deposited as a separation disc, or one or more cells degenerate, or at intercalary heterocyte positions (Bh radw ja 1933). Once considered a significant photobiont genus, including both principal and secondary (cephalodial) lichen symbionts, *Scytonema* in its current sense encompasses an uncertain but much reduced number of lichen algae. Photobionts previously ascribed to *Scytonema* have been shown by DNA sequence analyses to belong to a quite distinct clade, now designated *Rhizonema* (L cking *et al.* 2009). Nevertheless, at least one recent photobiont sequence (16s rRNA), from a *Heppia* thallus, appears to fall within *Scytonema* in the strict sense (V  2016). This may provide some corroboration for previous attributions of *Heppia* photobionts to *Scytonema* based on morphology of cultured isolates (Wetmore 1970). The cell shape and division planes of the *Heppia* photobionts are radically transformed to produce cell clusters in the lichenized state, reverting quickly to typical filamentous growth when cultured aposymbiotically (Marton & Galun 1976). In

Pyrenothrix nigra, the lichenized filamentous cyanobiont shows the double-false branching typical of *Scytonema* (Tschermak-Woess *et al.* 1983), although Lücking *et al.* (2009) suggested that its photobiont might be *Rhizonema*. This is quite plausible, since there is some doubt as to whether the two cyanobacterial genera can be reliably distinguished by their mode of branching (see comments under *Rhizonema*). More sequence data are clearly needed to clarify the extent to which lichen symbioses may involve the genus *Scytonema* in its current, more restricted sense.

Stigonema C. Agardh ex É. Bornet & C. Flahault — This cyanobacterial genus is recognized by its complex, branching axes with cells dividing in perpendicular planes as in true parenchyma. Filaments are uniseriate at the apex but become locally multiseriate proximally by periclinal divisions, often but not necessarily associated with the formation of true branches laterally. After division, cells retain continuity at the central portion of the septum, where micropores traverse the septal wall (Butler & Allsopp 1972). *Stigonema* has been reported as thallus photobiont in *Ephebe* and *Spilonema*, and also as cephalodial partner in numerous species of *Stereocaulon* (Tschermak-Woess 1988a). The genus awaits molecular treatment, remaining morphologically defined for the time being.

Tolypothrix Kützing ex É. Bornet & C. Flahault — These are filamentous cyanobacteria resembling *Scytonema* but with usually single- rather than double-false branches emerging from filament breaks; one side of the break grows out as the false branch, the other usually differentiates as a heterocyte. *Tolypothrix* has been reported as photobiont of the ‘primitively lichenized’ *Thermutopsis jamesii* based on morphology in collected material (Henssen 1990). Molecular sequences obtained from cephalodia of *Placopsis* placed the cyanobionts in or near *Tolypothrix* (Raggio *et al.* 2012).

Green algae (Viridiplantae – Archaeplastida)

Apatococcus F. Brand — Abundant and widely distributed as a free-living organism, *Apatococcus* has long been known as an omnipresent subaerial unicellular alga, inevitably encountered but not chosen by discriminating germling hyphae of lichen-forming fungi. Now it appears that *Apatococcus* includes lichen symbionts as well. Light microscopic observations of algal symbionts cultured from several maritime lichen species first implicated *Apatococcus* as a photobiont (Watanabe *et al.* 1997); molecular sequence comparisons later identified *Apatococcus* strains as partners of *Scoliosporum* (Beck 2002) and *Fuscidea* species (Zahradníková *et al.* 2017). Cells are spherical with alternating perpendicular planes of division, producing cuboidal packets of transiently adherent daughter cells. Autospores and biflagellate zoospores are also formed (Ettl & Gärtner 2014). Autospores may be of unequal size within a sporangium (Gärtner & Ingolić 1989), as also occurs in Watanabean genera such as *Chloroidium* and *Jaagichlorella*. As with *Elliptochloris* and *Trebouxia*, *Apatococcus* is facultatively heterotrophic: it is very slow growing in culture unless carbohydrate is supplied (Gustavs *et al.* 2016). This observation is particularly interesting because the similarly heterotrophic behaviour of *Trebouxia* in culture was central to Ahmadjian’s (1988, 2002) argument that *Trebouxia* cannot exist free-living. The seemingly ubiquitous *Apatococcus* shows quite clearly that

a photobiont exhibiting strongly heterotrophic tendencies in culture may nonetheless abound free-living in nature.

Asterochloris Tschermak-Woess — First described to accommodate the trebouxoid photobiont of a single lichen in the *Pertusariaceae* (Tschermak-Woess 1980a), this major photobiont clade now encompasses the former *Trebouxia* subgenus *Eleutherococcus* (Tschermak-Woess 1989; Škaloud & Peksa 2010). It corresponds roughly to Archibald’s (1975) restricted concept of genus *Trebouxia*, a source of continual confusion. *Asterochloris* species produce aplanospores, or zoospores in culture, but most strains do not form the appressed, low-number autospores characteristic of *Trebouxia* in the current sense. Its deeply-lobed chloroplast becomes flattened and parietal during cell division, while that of *Trebouxia* remains more or less central (Tschermak-Woess 1989). Pyrenoids are present; in TEM they may be distinguished as the *irregularis*-, *erici*-, or *magna*-types of Friedl (1989). Chloroplast morphology is highly variable and its utility as a marker in species delimitation was emphasized by Škaloud *et al.* (2015). As with *Trebouxia*, a considerable amount of genetic diversity is revealed at the molecular level in *Asterochloris* (Škaloud & Peksa 2010; Peksa & Škaloud 2011).

Sexual fusion of biflagellate isogametes to form a quadriflagellate zygote has been documented in cultures of *A. woessiae* (Škaloud *et al.* 2015). The detection of genes specific to meiosis in *A. glomerata* (Armaleo *et al.* 2019) provides further support for a functioning sexual cycle in *Asterochloris*.

Asterochloris is associated principally with mycobionts of the *Cladoniaceae*, *Stereocaulaceae*, and the genus *Lepraria*. These fungal partners appear to range from moderately to rather highly selective of their *Asterochloris* symbionts; there is also some indication that mycobionts of different clades are choosing particular *Asterochloris* lineages, showing distinct climatic preferences related to rainfall regime (Peksa & Škaloud 2011).

Auxenochlorella (I. Shihira & R. W. Krauss) T. Kalina & M. Puncová — Within the *Chlorellaceae*, *Auxenochlorella* is related to the fully heterotrophic genus *Prototheca*, and its type species, *A. protothecoides*, is also known for its heterotrophic tendencies in culture (Darienko & Pröschold 2015). *Auxenochlorella* has been implicated in regard to the identity of the photobiont associated with *Psoroglaena stigonemoides* in the *Verrucariaceae* (Nyati *et al.* 2007; Thüs *et al.* 2011). Unlike *Chlorella*, *Auxenochlorella* lacks a pyrenoid. The genus also includes ‘zoochlorellae’ symbionts of the cnidarian *Hydra* that are now considered a new species, *A. symbiontica* (Darienko & Pröschold 2015).

Bracteacoccus Tereg — *Bracteacoccus* are small, spherical unicells that have a multinucleate stage as they mature, and reproduce by zoospores or aplanospores; chloroplasts lack pyrenoids (Kouwets 1996). Currently included in the *Sphaeropleales* (Fučíková *et al.* 2014), *Bracteacoccus* appears at present to be the only genus of the class *Chlorophyceae* into which lichen photobionts have been placed with supporting DNA sequence data. The corresponding mycobionts are two species of the basidiomycete *Sulzbacheromyces* in the *Lepidostromatales* (Hodkinson *et al.* 2014; Masumoto 2020).

Cephaleuros Kunze ex E. M. Fries — These foliicolous relatives of *Trentepohlia* form macroscopic, multicellular thalli visible as

small, fuzzy yellow-orange patches on leaves and fruit in tropical and subtropical climates. *Cephaleuros* species typically grow beneath the cuticle of the leaf substratum, forming rounded to lobed thalli of more or less integrated horizontal filaments. These give rise to the erect setae and sporangiophores that emerge through the overlying cuticle. Usually, *Cephaleuros* develops within the space it excavates between the host cuticle and epidermis; in some cases, filaments penetrate deeper among the epidermal or mesophyll cells of the leaf, provoking a localized phellogen wound response. The alga can therefore be mildly pathogenic, but it is more often described as 'parasitic', despite an absence of information concerning any nutritional exchange with the plant host. Occasionally, the alga may develop upon the leaf cuticle, like other epiphylls. The behaviour may vary according to the species of *Cephaleuros* or that of the host plant (Ward 1884; Suto & Ohtani 2009; Brooks *et al.* 2015). Lichenization by foliicolous *Strigula* fungi is said to curb the alga's invasion of host tissue and its localized pathogenic effects (Joubert & Rijkenberg 1971).

Morphologically, *Cephaleuros* can somewhat resemble the related foliicolous genus *Phycopeltis*, which is not subcuticular and generally lacks vertical hairs and complex, long-stalked sporangiophores. According to molecular sequence analyses, however, the nearest relatives of *Cephaleuros* lie not within *Phycopeltis* but rather *Stomatochroon* (Zhu *et al.* 2017), a microscopic colonizer of the leaf's substomatal cavities. While trentepohliaceous taxa currently ascribed to *Phycopeltis* and *Trentepohlia* are phylogenetically intertwined, *Cephaleuros* appears to be essentially monophyletic (López-Bautista *et al.* 2006; Rindi *et al.* 2009; Nelsen *et al.* 2011; Zhu *et al.* 2017).

Cephaleuros is one of the few lichen photobiont genera for which life cycle events have been observed in some detail. Thompson & Wujek (1997) describe a haplodiplontic life cycle with heteromorphic multicellular phases. The familiar thallus corresponds to the gametophyte; fusion of gametes produces a zygote that germinates into a short-stalked, dwarf sporophyte bearing a putative meiosporangium. Flagellate meiospores presumably develop into new gametophytic thalli. The *Cephaleuros* gametophyte is the phase known to serve as phycobiont for the fungus *Strigula*. Whether or not the sporophytes can also be lichenized is unknown. Perhaps they are too highly reduced or short-lived, but the question does not seem to have been explored. On gametophyte thalli, two distinct structures produce flagellate zoospores or gametes (often called zooids, or swarmers, when their function is uncertain or polyvalent). Zoosporangia are elevated in groups upon vertical stalks; they produce quadri-flagellate zoospores that have been observed to round off, germinate and reproduce the gametophyte thallus asexually (Ward 1884; Thompson & Wujek 1997). The mature sporangia detach readily as units of dispersal, for which both wind and insects act as vectors. On the horizontal filament system, single, usually terminal cells may enlarge to become what are referred to as gametangia; these produce biflagellate cells that may fuse sexually (Thompson & Wujek 1997). However, other authors have been unable to observe any instances of sexual fusions in the taxa they studied, instead reporting that the biflagellate zooids germinate directly as zoospores to form new gametophyte thalli (Suto & Ohtani 2013).

Ward (1884) described in detail the course of lichenization of *Cephaleuros* by *Strigula*. Young germlings of *Cephaleuros* are often quickly overrun by the mycelium of *Strigula*, suppressing algal reproduction, while individuals contacted at more

advanced stages of development may produce abundant sporangia from portions of its thalli remaining relatively free of mycobiont domination. Interestingly, both symbionts grow and sporulate independently upon the leaf substratum, although the fungus *Strigula* will produce pycnidia and perithecia only after successful lichenization. These observations highlight the flexibility of the symbionts in this particular association.

Chlamydomonas Ehrenberg — This well-known unicellular green algal genus chiefly encompasses aquatic taxa that are flagellate in the vegetative state and unlikely candidates for lichen symbiosis. However, a number of aeroterrestrial species are also known (Ettl & Gärtner 2014). One species of *Chlamydomonas* (*C. augustae*) was described in association with the ascomycete *Pyronema laetissimum*, growing on leaf litter in Latvia (Skuja 1943). It was included in Tschermak-Woess's (1988a) review of phycobionts as 'facultatively lichenized'. However, Skuja (1943) distinguished this association from lichen and lichenoid symbioses, making comparisons instead with green algae known to grow abundantly on the surfaces of perennial basidiocarps. The *Chlamydomonas* was abundantly present among the dense hyphae below the *Pyronema* apothecium, but no tissue layer was differentiated, nor were any distinctive contact interfaces noted between symbionts. Skuja also mentioned that other apothecia of the same fungus were fruiting nearby without the alga present. The operculate discomycetes (*Pezizales*), to which *Pyronema* belongs, are not otherwise known to include lichen-forming members. The *Pyronema-Chlamydomonas* association is worthy of further investigation but seems unlikely to fit the criteria usually ascribed to lichen symbioses.

Chlorella Beijerinck — A once-notorious miscellany of indistinguishable 'little round green things', this trebouxiphycean genus has been radically deconstructed, particularly with the help of DNA sequence comparisons (Huss *et al.* 1999). Many formerly included species have been moved to different genera, orders, even classes, while taxa surrounding the type species *C. vulgaris*, and *C. sorokiniana*, are retained as true *Chlorella*. In TEM, they show a distinctive pyrenoid surrounded by a thick sheath of starch and bisected centrally by a single thylakoid (Ikeda & Takeda 1995; Němcová & Kalina 2000; Hoshina *et al.* 2010). Flagellate cells and sexual reproduction are unknown. True *Chlorella* also includes a number of mucilaginous, colonial forms in its current circumscription (Luo *et al.* 2010; Bock *et al.* 2011). Many of the lichen photobionts previously attributed to *Chlorella* s. lat. (e.g. Tschermak-Woess 1988b) are among those taxa moved to other genera, especially *Chloroidium*; others await re-examination. At present, only a couple of lichen-forming fungal species have photobionts of corroborated placement within *Chlorella* (*Porpidia crustulata*; Li *et al.* 2013) or *Chlorellales*. The genus has also been long identified with endosymbiotic algal symbionts of diverse protists and invertebrates. Molecular sequences confirm that true *Chlorella* occur as endosymbionts of the ciliate *Paramecium bursaria* (Hoshina *et al.* 2004; Summerer *et al.* 2008) and the cnidarian *Hydra* (Kovačević *et al.* 2010), which may also utilize *Auxenochlorella* as its algal symbiont (Pröschold *et al.* 2011). Chloroplast ultrastructure likewise suggests that the green endosymbiont of the colonial ciliate *Ophrydium versatile* is a true *Chlorella* (Forsberg & Lindblad 1996). However, the phylogenetic affinities of other 'zoochlorellae' symbionts appear to fall elsewhere in the

Trebouxiophyceae (Lewis & Muller-Parker 2004; Kovačević *et al.* 2010; Pröschold *et al.* 2011), while many have yet to be explored with molecular sequence comparisons.

Chloroidium Nadson — Resurrected to accommodate segregates from *Chlorella* s. lat. (Darienko *et al.* 2010), *Chloroidium* falls within the trebouxiophycean assemblage now formalized as *Watanabeales* (Li *et al.* 2021). Cells have a parietal chloroplast with or without a pyrenoid; in *C. saccharophilum*, a prominent pyrenoid with surrounding plastoglobuli and traversing membranes has been observed (González *et al.* 2013). Reproduction is by autospores, often of variable number and different sizes within a single sporangium. The genus encompasses diverse taxa found in a wide variety of habitats (Darienko *et al.* 2018), including extremophiles capable of using a variety of carbon sources (Nelson *et al.* 2017). Since its recent emendation, *Chloroidium* includes photobiont partners of a growing number of lichen-forming fungi, including some species of *Gomphillaceae*, *Verrucariaceae*, *Psora*, *Stereocaulon* and *Sticta*.

Chlorosarcinopsis Herndon — In the course of her studies on lichen haustoria, Plessl (1963) identified as *Chlorosarcina* [= *Chlorosarcinopsis*] *minor* the photobionts she isolated from two species of *Lecidea*, *L. plana* and *L. lapicida*. *Chlorosarcinopsis* has traditionally accommodated spherical unicellular green algae dividing to form cuboidal packets. According to Neustupa (2015), the genus is polyphyletic, with members scattered among the *Chlamydomonadales* (*Chlorophyceae*). As this clade is not otherwise known for lichen symbionts (but see Skuja 1943), the photobionts of the *Lecidea* species in question need further study.

Coccobotrys Chodat (now *Uvulifera* Molinari-Novoa) — This green alga forms irregular cuboidal cell packages or branched multiseriate filaments in culture (Neustupa 2015). The genus *Coccobotrys* was described by Chodat (1913) and emended by Vischer (1960) to accommodate the putative photobiont *C. verrucariae* isolated from a thallus of *Verrucaria nigrescens*. Thüs *et al.* (2011), on the other hand, reported *Diplosphaera* as photobiont of the *V. nigrescens* thallus they sampled. *Coccobotrys verrucariae* was also cited among algae isolated from soil crusts (Flechtner *et al.* 2009), and a photobiont identified with microscopy as ‘probably *Coccobotrys*’ (Canals *et al.* 1997) was isolated from *Botrylepraria lesdainii*, another member of the *Verrucariales* (Kukwa & Pérez-Ortega 2010). A second species of *Coccobotrys* was described by Warén (1920) as the photobiont of *Lecidea fuliginosa*, but Tschermak-Woess (1988a) expressed doubt that the alga he described belongs in *Coccobotrys*. The photobiont status of species in this genus should be corroborated. Genetic sequence analyses place *Coccobotrys* in the *Trebouxiophyceae* (e.g. Thüs *et al.* 2011; Mikhailyuk *et al.* 2020), but its affinities among the defined clades within this class remain uncertain. Molinari-Novoa (2016) recently found *Coccobotrys* Chodat to be a later homonym of a name applied to an anamorphic basidiomycete and renamed the algal genus *Uvulifera*.

Coccomyxa Schmidle — This trebouxiophycean algal genus is notable for the diversity of habitats and ecological circumstances in which its species are known to occur. Environmental surveys have found *Coccomyxa* sequences to be among the most widely

distributed OTUs, and notably well represented in cold high-latitude climates (Metz *et al.* 2019). It is commonly reported free-living on terrestrial substrata and in aquatic environments, including those highly polluted with heavy metals and radioactive materials (see Gustavs *et al.* 2017). *Coccomyxa* species are subspherical to ovoid-ellipsoidal unicells, often embedded colonially in thick gelatinous sheath material with concentric layering that reflects the cell division pattern. The chloroplast is parietal, not markedly lobed, and lacks a pyrenoid. In TEM, thylakoid bands often show a distinctly longitudinal orientation over the length of the chloroplast, with interspersed starch grains (Peveling & Galun 1976; Palmqvist *et al.* 1997). Flagellate cells and sexual reproduction are unknown; cells subdivide into packages of 2–8 autospores (Tschermak-Woess 1988a). Recent assessments of species number within the genus range from seven (Darienko *et al.* 2015) to as many as 27 (Malvasi *et al.* 2016). The genus is thought to include the photobionts of diverse lichen-forming fungi, such as species of *Icmadophila*, *Micarea*, *Nephroma*, *Peltigera*, *Solorina*, the stalked-apotheciate genera *Baeomyces*, *Dibaeis* and *Phyllobaeis*, and the basidiomycete *Lichenomphalia* (Table 1). Some of these reports await confirmation with genetic sequence data. The photobionts do not form a single clade but instead represent several distinct lineages within *Coccomyxa*, intermixed among free-living isolates (Darienko *et al.* 2015). In lichen symbiosis, the cells are often more spheroidal, and extensive gelatinous sheath material is not usually produced (Tschermak-Woess 1988a). Interestingly, while the cells of *Coccomyxa* and *Elliptochloris* photobionts are tightly enveloped by mycobiont hyphae, their walls are usually not penetrated (Tschermak 1941a; Geitler 1955; Plessl 1963; but see Coppins 1983: figs 2 & 55). This has been attributed to degradation-resistant polymers resembling sporopollenin in the multi-layered cell wall (Honegger & Brunner 1981; Brunner & Honegger 1985). However, *Coccomyxa* cells are fully penetrated by *Aphelidium collabens*, a parasitoid basal within, or sister to, the kingdom *Fungi* (Seto *et al.* 2020).

Species of the genus *Coccomyxa* also live in poorly understood symbioses within molluscs (Stevenson & South 1974; Syasina *et al.* 2012) and echinoderms, and endocytotically within ovules and other tissues of the gymnosperm *Ginkgo biloba* (Trémouillaux-Guiller *et al.* 2002; Trémouillaux-Guiller & Huss 2007). Molecular sequence comparisons have shown that some zoochlorellae isolated from certain strains of *Paramecium bursaria* correspond to *Coccomyxa*, while most others are true *Chlorella* (Hoshina & Imamura 2008).

Deuterostichococcus Pröschold & Darienko — A recent segregate of *Stichococcus* s. lat. (Pröschold & Darienko 2020), this trebouxiophycean genus currently includes, in addition to free-living isolates, the photobiont of two *Placopsis* species (Beck *et al.* 2019) and *Staurothele clopima* (Hodač *et al.* 2016); the latter is also known to partner with *Diplosphaera* algae (Thüs *et al.* 2011).

Dictyochloropsis Geitler — See *Symbiochloris*.

Dilabifilum Tschermak-Woess — Polymorphic, unicellular to sarcinoid to filamentous algae with pyrenoids and quadriflagellate zoospores have been included in this ulvacean genus. They occur free-living, as photobionts, or both. Recently, Darienko & Pröschold (2017) deconstructed *Dilabifilum*, recognizing at generic level several distinct clades resolved in their gene-based

phylogenies. A number of photobionts previously contained therein are now distributed in *Halofilum*, *Lithotrichon*, *Paulbroadya* and *Pseudendoconium*, while others await reassessment.

Diplosphaera Bialosuknia — This prasiolalean genus appears to contain the majority of the unicellular photobiont strains attributed until recently to the related *Stichococcus*. Apparently, the two morphologically plastic genera are often not distinguishable microscopically, although *Diplosphaera* may produce distinctive, adherent two-celled clusters in division. Pyrenoids may be absent (Pröschold & Darienko 2020) or weakly visible (Ettl & Gärtner 2014) but some taxa falling within the *Diplosphaera* clade, including lichen photobionts, show prominent pyrenoids in TEM (Fig. 2E). The main fungal partners of *Diplosphaera* are members of the *Verrucariaceae*. Photobiont strains compared so far appear to represent the same species and are similar to free-living collections (Pröschold & Darienko 2020).

In association with certain lichen genera, such as *Endocarpon* and *Staurothele*, *Diplosphaera* photobionts ‘escape’ vegetative hyphal contacts and penetrate into the hymenial layer of developing perithecia, where they freely intermix among the asci (Fig. 3A). These algal cells are typically much smaller than those within the algal layer of the vegetative thallus; they scatter everywhere when a hand-cut section is water-mounted, indicating that unlike the photobionts in the vegetative thallus, those entering the perithecia are not bound in place by lichenizing contacts with the mycobiont. The unassociated photobionts may adhere to the large ascospores as they are ejected, and can be dispersed with them. Many readily detach and divide aposymbiotically; they are available to the mycobiont if the spore germinates successfully (Stahl 1877; Bertsch & Butin 1967; Ahmadjian & Heikkilä 1970) or might otherwise divide to form free-living populations. Potentially co-dispersible photobionts also occur in the conidiomata and ascomal epithelia of many foliicolous lichens of the *Gomphillaceae* and *Pilocarpaceae* (see *Heveochlorella*). Dispersal of liberated photobionts can thereby provide a direct connection between lichenized and free-living populations of the alga.

Elliptochloris Tschermak-Woess — Like its sister genus *Coccomyxa*, *Elliptochloris* has subspherical to ellipsoidal unicells with a parietal chloroplast, bearing two opposed indentations in the type species *E. bilobata* (Tschermak-Woess 1980b). Sexual or flagellate stages are unknown; reproduction occurs by autospores, of which there are usually two morphologically distinct types. Autosporangia may contain a low number (usually four in cultured *E. bilobata*) of spherical spores appressed together at flattened junctions, or more numerous (16–32) cylindrical-ellipsoidal spores (Tschermak-Woess 1980b; Darienko *et al.* 2016). The multilayered cell wall, as in *Coccomyxa*, is impregnated with degradation-resistant polymers, which are thought to explain the lack of haustorial penetration by their lichen-forming partners (Brunner & Honegger 1985). However, haustoria have been noted in certain species of *Micarea* (Coppins 1983: figs 2 & 55), a lichen-forming genus known to partner with *Elliptochloris* and *Coccomyxa* photobionts. Unlike *Coccomyxa*, at least some species of *Elliptochloris* possess pyrenoids, and layered mucilaginous sheaths are typically lacking. However, gelatinous extracellular material may be copious in free-living populations, and was

observed in association with *Protothelenella* thalli where the photobiont population grew beyond the reach of mycobiont hyphae (Tschermak-Woess 1985).

Elliptochloris is somewhat less often reported than *Coccomyxa* but is known from a similarly diverse array of habitats. It is said to be quite strongly heterotrophic in culture, where it depends heavily on organic materials to thrive; this might in part account for its less frequent recovery in isolation procedures (Gustavs *et al.* 2017). Species of *Elliptochloris* are known to partner with mycobionts of diverse genera including *Catillaria*, *Catolechia*, *Fuscidea*, *Micarea*, *Sticta*, *Stictis*, *Verrucaria*, and the basidiolichen-forming *Bryoclavula* and *Multiclavula* (see Table 1). They also occur as endosymbionts of the marine anenome *Anthopleura* (Letsch *et al.* 2009).

Gloeocystis Nägeli — Taxa treated under this genus are unicellular green algae that form occasionally macroscopic colonies of ellipsoidal cells with a parietal chloroplast possessing a pyrenoid. Thick, colourless, often layered gelatinous sheaths surround the cells. Reports of *Gloeocystis* as photobiont of *Cryptodiscus* [*Bryophagus*] *gloeocapsa* and *Epigloea bactrospora* were cited by Ahmadjian (1967) and Tschermak-Woess (1988a). There is doubt as to whether *Epigloea* is lichenized (Kirk *et al.* 2001; not included in Lücking *et al.* (2017a)), although distinctly symbiotic contacts with living, unicellular green algae were illustrated by Jaag & Thomas (1934) and Döbbler (1984).

According to Neustupa (2015), the algal genus *Gloeocystis* is highly polyphyletic, encompassing members of both *Chlorophyceae* and *Trebouxiophyceae*. The identities of the photobionts associated with the mycobionts mentioned above will therefore require further study.

Halofilum Darienko & Pröschold — This is another genus that now accommodates taxa previously treated under *Dilabifilum* (Darienko & Pröschold 2017). These algae consist of branched filaments with parietal chloroplasts containing pyrenoids; flagellated stages are unknown. The species *H. ramosum* occurs as photobiont of *Hydropunctaria maura* and *Wahlenbergiella striatula* (*Verrucariaceae*), as well as free-living (Darienko & Pröschold 2017).

Heveochlorella J. Zhang *et al.* — Unicellular algae attributed to *Heveochlorella* have a prominent, somewhat lobed chloroplast with a central pyrenoid that is readily visible with light microscopy. TEM shows the pyrenoid surrounded by several irregular starch plates and penetrated centripetally by thylakoid-derived tubules that are lined with pyrenoglobuli (Fig. 2B). Cells reproduce by autospores, usually in low number (2–8) and not infrequently of unequal size within sporangia, at least in culture (Zhang *et al.* 2008; Ma *et al.* 2013; Sanders *et al.* 2016). Darienko & Pröschold (2019) recently subsumed both *Heveochlorella* (Zhang *et al.* 2008) and the related *Heterochlorella* (Neustupa *et al.* 2009), which has not been reported from lichen symbioses, into the resurrected genus *Jaagichlorella*. These algae belong to the trebouxiophycean clade recently formalized as *Watanabeales* (Li *et al.* 2021).

The first indication that lichen symbionts belonged in this group was the report of *Heveochlorella* isolated as photobiont from one specimen of *Sticta* and two of *Pseudocyphellaria* from Taiwan (Dal Grande *et al.* 2014b). Soon thereafter, the ‘trebouxioid’ photobionts associated with foliicolous

Gomphillaceae and *Pilocarpaceae* were also attributed to this genus (Sanders *et al.* 2016). More recently, a study of *Sticta* lichens worldwide reported *Heveochlorella* to be the photobiont of numerous specimens from New Zealand and Indian Ocean islands, including six identified species and many undetermined collections (Lindgren *et al.* 2020). In the opinion of Darienko & Pröschold (2019), the algae encompassed by *Jaagichlorella*, though distributed worldwide, are rare taxa. While more surveys will be necessary to evaluate this view, a number of observations suggest that these algae might be in fact quite common and merely overlooked. We know, for example, that foliicolous lichens of the *Gomphillaceae* and *Pilocarpaceae* occur in abundance throughout much of the humid tropics (Santesson 1952; Lücking 2008), although it is not yet clear how consistently they harbour *Heveochlorella* (*Jaagichlorella*) photobionts. Recent sampling of the phyllosphere community in Asian tropical forests has revealed a major representation of *Heveochlorella* genotypes (Zhu *et al.* 2018), as well as several new species in related genera (Li *et al.* 2020, 2021). Some of the most frequently detected OTUs in environmental surveys of marine habitats (Metz *et al.* 2019) were also attributed to *Heveochlorella*.

In many foliicolous lichens, dividing *Heveochlorella* photobionts may escape the lichenizing vegetative hyphae and proliferate among the spore-generating fungal structures, upon apothecia and within specialized conidiomata such as campylidia and hyphophores (Fig. 3B & C). They can be dispersed from these structures, as are the fungal spores or diahyphae to which the algal cells may adhere (Fig. 3D–F). Once dispersed, they may become lichenized by the germinating fungal propagules, or divide to produce independent populations on the substratum (Sanders 2014; Sanders & de los Ríos 2015). Co-dispersal and relichenization thereby provide *Heveochlorella* with abundant opportunities for exchange between lichenized and free-living populations.

Interfilum Chodat — This genus of aeroterrestrial charophytes (*Streptophyta*) includes taxa that form single, paired or sarcinoid packets of cells or grow filamentously, often closely resembling unrelated *Chlorophyta*, such as *Desmococcus* (Mikhailyuk *et al.* 2008). It is sister to clades of the widely distributed *Klebsormidium* (Rindi *et al.* 2011). *Interfilum* was reported by Voytsekhovich *et al.* (2011) as a secondary photobiont within the algal layer of *Micarea* and *Placynthiella* thalli collected in Ukraine, based on light microscopic examination of thalli and cultured isolates. The principal photobionts in those lichens were reported to be *Elliptochloris* and *Radiococcus*, respectively. As the charophytes are not otherwise known as lichen symbionts, and other algal genera were cited as the main photobionts within the thalli in question, further study of the reported associations is warranted.

Jaagichlorella Reisinger — See *Heveochlorella*.

Leptosira A. Borzi — This photobiont grows as unicells tightly wrapped by mycobiont hyphae or separated by copious sheath material free of the mycobiont; in agar culture, it produces short filaments (Tscheramak-Woess 1953). *Leptosira* is a trebouxiophycean of uncertain placement, appearing in the vicinity of the *Microthamniales* clade in recent gene-based phylogenies (Lemieux *et al.* 2014; Neustupa 2015; Hallmann *et al.* 2016). According to Mattox & Stewart (1984),

'*Pleurastrum terrestre*' (a synonym of *Leptosira obovata*, now *L. terrestre*; Friedl 1996) is so similar ultrastructurally to the genus *Trebouxia* that they could be combined in the same genus. Ahmadjian (1988) went one step further, opining that *Trebouxia* was merely the lichenized form of this taxon. However, the aforementioned gene-based cladograms do not show a close relationship between *Leptosira* and the *Trebouxiales*.

Leptosira terrestris, in lichen symbiosis with *Vezeadaea aestivalis*, grows subcuticularly (Tscheramak-Woess & Poelt 1976), a distinction shared with the photobiont *Cephaleuros*. *Leptosira* is also among the very few photobiont genera (*Phycopeltis* and *Cephaleuros* are others) reported to produce zoospores in the lichenized state (Tscheramak-Woess & Poelt 1976).

Lithotrichon Darienko & Pröschold — Another genus separated from the *Dilabifilum* (*Ulvales*) complex, *Lithotrichon* forms clustered cell packets as well as branching filaments and is distinguished from similar genera by SSU and ITS sequence data. The species *L. pulchrum* occurs as photobiont of the freshwater lichen *Hydropunctaria rheitrophila* (Dariencko & Pröschold 2017).

Myrmecia Printz — These spherical to pyriform unicells have a parietal chloroplast, without a pyrenoid, extending around most of the cell, with 2–4 broad lobes defined by deep notches. Cells proliferate via zoospores, aplanospores, or autospores (Ettl & Gärtner 2014). Gene-based phylogenies consistently place *Myrmecia* in the *Trebouxiales*, sister to *Trebouxia* (Muggia *et al.* 2020) or to the *Asterochloris* + *Vulcanochloris* clade (Vančurová *et al.* 2015). *Myrmecia* occurs free-living as well as in lichen symbiosis. An aerophilic alga, originally described as *Friedmannia* from Negev Desert rocks, is now recognized as *Myrmecia israelensis* (Friedl 1995) and was recently reported as lichen photobiont (Thüs *et al.* 2011; Moya *et al.* 2018). *Psora decipiens* and a number of species in the *Verrucariaceae* are among the lichen-forming fungi known to partner with *Myrmecia*.

Nannochloris Naumann — The genus *Nannochloris* has encompassed simple, extremely tiny (1.5–2 µm) chlorophyte algae that reproduce by binary division or autospores. Circumscription of the genus has been controversial, but molecular data indicate that most of the species belong in *Chlorellales* (Henley *et al.* 2004). Tscheramak-Woess (1981) recognized *Nannochloris normandinae* as the photobiont partner of lichen-forming *Normandina pulchella*; in other works, *Nannochloris* has been mentioned more indirectly in the context of photobionts (e.g. Lohtander *et al.* 2003). However, Thüs *et al.* (2011) found only *Diplosphaera* as photobiont in the 10 *Normandina* thalli they examined and, more recently, Pröschold & Darienko (2020) reduced *Nannochloris normandinae* to synonymy with *Diplosphaera chodatii* (*Prasiolales*). Thus, clear evidence of lichen photobionts belonging in *Nannochloris* appears to be lacking at present.

Neocystis F. Hindák — Members of this trebouxiophycean genus produce mucilaginous colonies of spherical to ellipsoidal or crescent-shaped cells that reproduce by autospores (Neustupa 2015). Cultures assigned to *Neocystis* as well as other genera were recently reviewed with molecular sequence analyses, revealing considerable taxonomic redundancy assigned to only two closely related,

genetically distinct but morphologically plastic species (Elišá *et al.* 2013). An alga identified as *Neocystis* sp. was cited as ‘additional photobiont’ of *Micarea misella*, in thalli having *Elliptochloris bilobata* as principal photobiont (Voytsekhovich *et al.* 2011).

Paulbroadya Darienko & Pröschold — This recently recognized clade in the *Ulvales* is distinguished by SSU-ITS sequences from other taxa previously treated under *Dilabifilum* (Darienko & Pröschold 2017). The species *Paulbroadya petersii* occurs as photobiont of the marine intertidal lichen *Wahlenbergiella mucosa* (Darienko & Pröschold 2017).

Phycopeltis Millardet — Members of this trentepohliaceous genus are most often seen as coppery orange discs a few mm in diameter on leaf surfaces in humid subtropical and tropical regions, with one or two species extending to cooler regions such as oceanic Europe (Rindi *et al.* 2004). Thallus discs consist of a monostromatic layer of closely appressed, bifurcating filaments (Fig. 6A). Unlike those of *Cephaleuros*, *Phycopeltis* thalli are supracuticular, non-pathogenic, and at least some species readily colonize other favourably displayed plant substrata besides leaves. Sporangia are usually borne erect on a very short stalk, and release quadriflagellate zoospores through a pore at the end opposite to the point of attachment. Gametangia are sessile and develop from intercalary compartments of the horizontal filament system in most species; gametes are biflagellate and isomorphic, and their fusion has been observed (Thompson & Wujek 1997). The life cycle of *Phycopeltis* is believed to be haplodiplontic, with alternation of gametophytes and sporophytes that are isomorphic, rather than heteromorphic as in *Cephaleuros* and *Stomatochroon* (Thompson & Wujek 1997). If this is the case, recognizing meiosporangia by the presence of tetrads might be the only means of distinguishing the phases phenotypically, but there do not appear to be such reports. Whether the gametophytes and sporophytes are equally susceptible to lichenization would be an interesting question to examine.

Although distinguishing *Phycopeltis* from *Trentepohlia* under current morphological concepts appears fairly straightforward, DNA sequence data show species of the two genera to be intertwined phylogenetically (Zhu *et al.* 2015, 2017; Grube *et al.* 2017a). *Phycopeltis* is particularly under-sampled at present. The morphological distinction between the two genera may also break down in the lichenized condition. Although *Phycopeltis* species can retain their placoid thallus characteristics when partnering with certain foliicolous mycobionts (Grube & Lücking 2002), in other lichens the algal filaments may be broken up into individual cells indistinguishable from those of *Trentepohlia* (see fig. 9; Lücking 2008). Using TEM, Matthews *et al.* (1989) believed they could differentiate the two genera in such cases by features of the septal wall near plasmodesmata. It would be useful to test how well such traits correlate with molecular markers.

As widespread colonizers of the warm-temperate and tropical phyllosphere, species of *Phycopeltis* are important photobionts in foliicolous lichen communities, where they partner with diverse leaf-dwelling mycobionts including species of *Arthonia*, *Chroodiscus*, *Mazosia*, *Opegrapha*, *Porina*, *Trichothelium*, and supracuticular taxa of *Strigulaceae*, among others (Santesson 1952; Lücking 2008). Multiple *Phycopeltis* thalli may occur edge-to-edge within a single foliicolous lichen, as additional individuals are incorporated by the mycobiont’s expanding

prothallus (Sanders 2002). There are some reports of algal gametangia or sporangia being produced in the lichenized state, particularly in those taxa where the *Phycopeltis* thalli are sparsely covered by the mycobiont (Santesson 1952; Lücking 1994; Sanders 2002).

In at least one species of *Phycopeltis* (*P. epiphyton*), the highly degradation-resistant biopolymer sporopollenin was detected in the cell wall (Good & Chapman 1978). Its presence in the walls of other photobionts (*Coccomyxa* and *Elliptochloris*) has been correlated with the absence of haustorial penetration by mycobionts (Honegger & Brunner 1981; Brunner & Honegger 1985). However, at least some strains of lichenized *Phycopeltis* may be deeply penetrated by mycobiont haustoria, such as those of *Porina* (Matthews *et al.* 1989).

Prasiola Meneghini — This trebouxioephycean seaweed of high-latitude supratidal zones is exceptional for its class in having a multicellular, macroscopic blade-like thallus. It is often abundant and readily visible both free-living and in symbiosis with mycobiont *Mastodia tessellata* (*Verrucariaceae*). Two or three distinct species of *Prasiola* appear to serve as photobiont to the bipolarly distributed *Mastodia* (Garrido-Benavent *et al.* 2017, 2018). There has been some discussion, on structural grounds, as to whether this fungal-algal partnership ought to be considered a true lichen (Lud *et al.* 2001; Kohlmeyer *et al.* 2004; Pérez-Ortega *et al.* 2010). There is no fungal cortex, nor does symbiosis substantially change algal thallus morphology; its anatomy, however, is significantly altered, as algal cells become well separated by a proliferation of encircling mycobiont hyphae (Kovačik & Batista Pereira 2001; Lud *et al.* 2001). The symbiosis therefore entails considerably more structural transformation than that produced by the marine fungus *Mycophycias* upon its seaweed host *Ascophyllum* (Xu *et al.* 2008), or *Turgidosculum* upon *Blidingia* (Pérez-Ortega *et al.* 2018). From a phylogenetic perspective, it is worth noting that close relatives of both the mycobiont (*Verrucariaceae*) and the alga (*Prasiolales*) participate in symbioses that are unambiguously lichenic.

Pseudendoconium Wille — These ulvalean algae have variably packet-forming to filamentous morphologies and may be differentiated into prostrate and erect filament systems. Darienko & Pröschold (2017) moved into this genus a couple of photobionts previously treated under *Dilabifilum*, recognizing the photobiont of *Arthopyrenia kelpii* as *Pseudendoconium arthopyreniae*, and the photobiont of *Hydropunctaria maura* as *P. commune*, which is also widespread as a free-living alga on intertidal rocks. *Pseudendoconium arthopyreniae* has a pyrenoid surrounded by thick plates of starch and traversed by several narrow, thylakoid-derived membranes lacking pyrenoglobuli (Namba & Nakayama 2021).

Pseudochlorella J. W. G. Lund — This unicellular genus of *Chlorella*-like algae is now placed in the *Prasiolales*. Molecular data support inclusion of the photobiont of at least one lichen-forming fungus, *Trapelia coarctata* (Darienko *et al.* 2016). Other reports attribute to *Pseudochlorella* the photobionts of certain *Micarea*, *Placynthiella* and *Stereocaulon* species (see Tschermak-Woess 1988a; Voytsekhovich *et al.* 2011). However, molecular sequence studies have so far identified photobionts from *Micarea* as *Coccomyxa* and *Elliptochloris*, and those from the green algal layer of *Stereocaulon* as *Asterochloris*, *Chloroidium* and *Vulcanochloris*.

Pseudococcomyxa Korshikov — Isolates identified as *Pseudococcomyxa simplex* have been reported as photobionts of a maritime *Leproloma* sp. (Watanabe *et al.* 1997) and also *Micarea prasina* (Voytsekhovich *et al.* 2011), based on light microscopy and culture studies. The genus *Pseudococcomyxa* has been distinguished morphologically from *Coccomyxa* by the polarized secretion of mucilage to form a cap at one end of the cell. However, Darienko *et al.* (2015) found this character to be culture-dependent, and the *Pseudococcomyxa* strains they analyzed phylogenetically appeared intermixed with those of *Coccomyxa* (see also Yahr *et al.* 2015). Isolates attributed to *P. simplex* in particular occurred in several distinct clades. Darienko *et al.* (2015) reassigned all these strains to *Coccomyxa*. While lichen photobiont isolates attributed to *Pseudococcomyxa* remain to be examined, support for distinction of the genus from *Coccomyxa* now appears to be lacking.

Pseudostichococcus L. Moewus — Morphologically similar to *Stichococcus*, this genus was recently revised with molecular data (Pröschold & Darienko 2020). It currently includes the photobiont partner of *Neocatapyrenium rhizinosum* (Hodač *et al.* 2016) in the *Verrucariaceae*.

Pseudotreboxia P. A. Archibald — See *Treboxia*.

Radiococcus Schmidle — Species of this genus have been reported to occur as principal photobiont in thalli of two species of *Placynthiella* (*P. icmalea* and *P. uliginosa*) from the Ukraine (Voytsekhovich *et al.* 2011). Corroboration with DNA sequence data is needed, particularly since diverse, unrelated taxa have been repeatedly ascribed to this genus in the past (Wolf *et al.* 2003). According to a recent taxonomic treatment, *Radiococcaceae* and *Radiococcus* belong in the order *Sphaeropleales* of the *Chlorophyceae* (Neustupa 2015), although these names are still being applied to taxa falling in other groups, such as the *Treboxiophyceae* (e.g. Metz *et al.* 2019).

Stichococcus Nägeli s. lat. — In its broad sense, *Stichococcus* (*Prasiolales*) has encompassed smallish unicellular to filamentous algae of notably labile morphology, the most commonly recognized form represented by short-cylindrical cells. The chloroplast is parietal, often extending to no more than half of the cell circumference, not markedly lobed, with or without a pyrenoid. Culture conditions appear to have a significant effect on cell form. The straight or slightly curved, rod-shaped cells may separate or remain together after division to form very short filaments or swell to more spherical shapes, and may or may not produce a surrounding gelatinous sheath (Ettl & Gärtner 2014). Quite a number of lichen photobionts have been ascribed to *Stichococcus*, but as their diversity is studied at the molecular level, these taxa are being placed in segregate genera or other prasiolalean clades. Some seven to nine clades have now been recognized within *Stichococcus* s. lat. (Hodač *et al.* 2016; Pröschold & Darienko 2020). All *Stichococcus*-like photobionts examined in a study of the *Verrucariaceae* by Thüs *et al.* (2011), were shown to belong in *Diplosphaera*. Others now appear to fall within *Pseudostichococcus*, *Deuterostichococcus* or *Tritostichococcus* (Pröschold & Darienko 2020). It is not yet clear whether *Stichococcus* in the restricted sense (near to type species *S. bacillaris*) includes lichen photobionts.

Symbiochloris Škaloud *et al.* — Formally described by Škaloud *et al.* (2016), the genus corresponds to a distinct clade of *Dictyochloropsis* s. lat. previously recognized by Dal Grande *et al.* (2014b). *Symbiochloris* is currently thought to include all lichen photobionts previously included in *Dictyochloropsis*, as well as some free-living taxa. Principal mycobiont partner genera are *Lobaria*, *Pseudocyphellaria*, *Sticta* and their recent segregates *Crocodia*, *Dendrosticta* and *Ricasolia*, all members of the *Lobariaceae*. Other lichens reported harbouring *Symbiochloris* photobionts include species of *Biatora*, *Brigantiaea*, *Chaenotheca*, *Megalospora* and *Phlyctis*.

The net-like chloroplast of *Symbiochloris*, similar to that of *Dictyochloropsis*, has reticulations that vary in form, thickness and orientation according to species and developmental stage (Škaloud *et al.* 2005, 2016). Lichenized populations reproduce by aplanospores, but zoospore production may be observed in isolated culture. Cells of free-living populations often attain much larger sizes and their surfaces may be covered with scales (Tschermak-Woess 1995).

Treboxia Puymaly — The principal crop of the alga-farming fungi, unicellular *Treboxia* is thought to include the photobionts chosen by the largest proportion (nearly half) of known mycobiont species. Together with the closely related *Asterochloris*, *Treboxia* is chlorobiont of most *Lecanorales* and *Teloschistales*, as well as many other taxa of the other species-rich lecanoromycetid orders (Miadlikowska *et al.* 2014). Species of *Treboxia* are spherical or occasionally ellipsoidal unicells with a variously lobed, axial chloroplast that fills much of the cell and bears a prominent pyrenoid (Fig. 2A & C). There is considerable diversity of pyrenoid ultrastructure within the genus, involving differences in the morphology of penetrating membranes and the distribution of starch deposits and pyrenoglobuli, when present (Peveling 1968, 1969; Fisher & Lang 1971; Friedl 1989). CO₂-fixing Rubisco is concentrated in the pyrenoids, which in some instances also comprise additional, smaller, satellite substructures within the chloroplast (Ascaso *et al.* 1995). Although pyrenoid types do not correspond precisely to the *Treboxia* clades supported in molecular sequence analyses, and several are strikingly convergent with pyrenoids of distantly related algae, they can nonetheless be useful in distinguishing certain groupings of taxa at close range (Friedl 1989; but see also Muggia *et al.* (2010)). Some 30 species of *Treboxia*, distributed among four major clades, are currently recognized. However, this figure is believed to grossly underestimate the true genetic diversity present in the genus (Leavitt *et al.* 2015; Muggia *et al.* 2020). The boundaries among the formally described species remain largely unresolved, since much of the genetic diversity uncovered in recent studies is not fully congruent with the phenotypically defined taxa. Muggia *et al.* (2017) postulated that the application of a phylogenetic species concept would at least triple the number of species currently recognized in *Treboxia*.

Two different groups were long distinguished within *Treboxia* s. lat. (Ahmadjian 1960), which was previously known as *Cystococcus*. Archibald (1975) recognized two genera, *Treboxia* and *Pseudotreboxia*, based on differences in cell division which were judged sufficient to separate them into two distinct orders. However, Gärtner (1985) and Tschermak-Woess (1989) found Archibald's subdivision untenable and reunited the genus, while acknowledging that differences in cell division were present. Tschermak-

Woess (1989) distinguished two subgenera: *Trebouxia* (corresponding roughly to *Pseudotreboxia*), which forms aplanospores (or zoospores in culture) and also autospores, and *Eleutherococcus* (later *Asterochloris*), which produces aplanospores/zoospores but not autospores. Autospores are distinguishable from aplanospores in that they are produced in lower numbers and are tightly appressed together within the sporangium such that their walls form angular junctions between them (Tschermak-Woess 1989). Another difference is the position of the chloroplast during cell division, which remains more or less central in *Trebouxia* but becomes parietal and flattened in *Asterochloris* (Ahmadjian 1960; Tschermak-Woess 1989). Molecular data firmly distinguish the two clades, which have been formally recognized as distinct genera for the past decade (Škaloud & Peksa 2010).

As a lichen symbiont, *Trebouxia* is abundant in a great diversity of habitats worldwide. It is said to be infrequently reported in the free-living state, although researchers who sample substrata with microscopy have often found it (Tschermak-Woess 1978; Bubrick *et al.* 1984; Mukhtar *et al.* 1994; Sanders 2005; Handa *et al.* 2007; Uher 2008; Neustupa & Štifterová 2013), with the notable exception of Degelius (1964). Clearly, the germinating spores of trebouxiphilic mycobionts manage to obtain it, often without needing to produce an extensive mycelium (Werner 1931; Clayden 1998). Recent environmental sequencing studies have found *Trebouxia* on a variety of surfaces (Darienko *et al.* 2013; Hallmann *et al.* 2013, 2016; Yung *et al.* 2014) and well represented in soil, fresh water and even marine environments (Metz *et al.* 2019), although one cannot be certain that the detected sequences represent free-living individuals. By contrast, two other principal lichen photobiont genera, *Trentepohlia* and *Nostoc*, are uncontroversially well known in the free-living state. The comparison may not be fair, however, because *Trentepohlia* and *Nostoc* both form easily recognized macrocolonies (bright orange tufts and distinctive gelatinous globs, respectively) whereas *Trebouxia* cannot be distinguished without a microscope and some degree of effort. In any case, a shadow of doubt still seems to haunt the status of free-living *Trebouxia* populations, to judge from the cautious wording in even quite recent literature (e.g. Friedl & Büdel 2008). Although he never claimed to have searched for it in nature, Ahmadjian (1988, 1993, 2001, 2002) repeatedly affirmed that *Trebouxia* existed only in highly coevolved symbiosis with lichen fungi and did not occur free-living. Yet he acknowledged that aposymbiotic populations of *Trebouxia* could appear in nature. He even proposed, as have others, that they arose from the breakdown of lichenized propagules, such as soredia and isidia, that reach microhabitats unsuitable for the partners to develop symbiotically (Ahmadjian 1988). Ahmadjian asserted, however, that such populations were not truly free-living, except in a 'secondary sense'. Apparently, he meant that they were ephemeral rather than stably established, but stable or not, aposymbiotic populations of *Trebouxia* are likely to be significant. Like other micro-organisms, many algae take advantage of ephemeral resources and transiently favourable microenvironments, then complete their life cycles with sexual reproduction when conditions deteriorate. Some then survive as resistant spores; others may escape adversity by entering into lichen symbioses. Within a lichen thallus, an algal population may be perpetuated for many years, yet continually disperse via soredia, isidia, lichenized

fragments and other propagules that can seed new free-living populations. This has been characterized as photobiont 'escape' from the lichen fungus (Werth 2010). It may be equally valid to view relichenization as photobiont escape from conditions that aposymbiotic populations might not long endure.

Although stages of flagellar development within a lichen thallus were reported (Slocum *et al.* 1980), authors have expressed scepticism that *Trebouxia* could produce motile or sexual cells in the symbiotic state (Tschermak-Woess 1989), where all algal cells are held by one or more appressorial hyphae (Honegger 1990). In aposymbiotic culture, by contrast, the production and release of *Trebouxia* zooids are well documented (Ahmadjian 1960, 1967; Tschermak-Woess 1989; Takeshita 2001). The huge genetic diversity present (Muggia *et al.* 2020) and its structure within populations (Kroken & Taylor 2000) suggest that *Trebouxia* is reproducing sexually, but virtually nothing is known about how or when the sexual cycle proceeds in nature. Although it is often said that sexual reproduction has not been observed in this genus, both Warén (1920) and Ahmadjian (1960) reported and illustrated the fusion of flagellate isogametes in *Trebouxia* cultures. However, Ahmadjian (1988, 2001) believed that these features were vestiges of the alga's free-living ancestry that no longer play any role in their present life histories. Further investigation of aposymbiotic populations is needed, since considerable indirect evidence suggests that they may reveal key events in the *Trebouxia* life cycle.

Trentepohlia C. Martius — The filamentous taxa currently treated under this cosmopolitan genus are among the most familiar of subaerial algae, often forming readily visible yellowish orange tufts on bark, rocks and other substrata in a wide variety of environments. They are also among the phycobionts chosen by the most diverse lichen-forming ascomycetes, including members of the *Arthoniomycetes*, *Coniocybomyces*, *Dothidiomycetes*, *Eurotiomycetes* (*Pyrenulales*) and ostopalean *Lecanoromycetes* such as the species-rich *Graphidaceae*. Members of the order *Trentepohliales* and its sole family *Trentepohliaceae* present a distinctive combination of features: phragmoplastic cell division with plasmodesmata (otherwise characteristic of charophycean algae), a uniquely structured flagellar apparatus, peculiar sporangiophores, and distinctive orange pigmentation. Consequently, widely divergent interpretations of their phylogenetic affinities have been proposed, with some authors even placing the group in a separate class of its own (van den Hoek *et al.* 1995). However, rDNA sequence data firmly place the subaerial *Trentepohliales* among orders of mainly marine taxa within the *Ulvophyceae* (López-Bautista & Chapman 2003; Leliaert *et al.* 2012).

Among the taxa currently treated under *Trentepohlia*, a number of genera were described to accommodate the morphological diversity represented, most recently *Printzina* (Thompson & Wujek 1992). However, DNA sequence analyses have so far shown that the phenotypic similarities recognized are unreliable indicators of phylogenetic affinity (López-Bautista *et al.* 2006; Rindi *et al.* 2009). This also applies to some of the morphological traits currently used to distinguish *Trentepohlia* species from those of *Phycopeltis*. Free-living and lichenized isolates of *Trentepohlia* occur intermixed in gene-based phylogenies (Nelsen *et al.* 2011; Hametner *et al.*

2014a, b; Kosecka *et al.* 2020). Due to its visible and widespread presence in the free-living state, *Trentepohlia* is an excellent subject for studying the relationship between lichenized and aposymbiotic populations in nature (Fig. 1A–C). So far, however, the genus has been the focus of relatively few modern phylogenetic studies, despite its visibility and primary importance in lichen symbioses.

The pigments characteristic of the *Trentepohliaceae*, called ‘haematochrome’ in the older literature, are carotenoids that occur abundantly as lipidic globules in the cytoplasm. Some authors have attributed the colour to astaxanthin (Thompson & Wujek 1997; Grube *et al.* 2017a), a deep red carotenoid known from other chlorophytes such as *Haematococcus* and the snow alga *Chlamydomonas nivalis*. However, analyses of *Trentepohlia* haematochrome show principally beta-carotenes (Czeczuga & Maximov 1996; Mukherjee *et al.* 2010; Kharkongor & Ramanujam 2015; Chen *et al.* 2016). Located outside the plastids, these secondary carotenoids do not participate in photosynthetic light-harvesting but are hypothesized to filter excess light and suppress any damaging reactive oxygen species thereby generated (Solovchenko 2013). This may contribute to the visible success of *Trentepohlia* in colonizing exposed substrata in diverse environments.

The sporangia of *Trentepohlia* are themselves units of dispersal, easily detached when mature and vectored by wind, rain or insects. They then initiate a second round of shorter-distance dispersal by releasing quadriflagellate zoospores (Thompson & Wujek 1997). *Trentepohlia* also produces putative gametangia that are morphologically distinct from sporangia. However, the biflagellate zooids released have most often been observed germinating as spores rather than fusing as gametes (Rindi & Guiry 2002).

Cellular contacts between mycobionts and their trentepohliaceous photobionts often appear to be superficial. However, most of these lichens, when carefully examined microscopically, reveal haustorial penetration, often deeply into the algal cells (Tschermak 1941a; Withrow & Ahmadjian 1983; Matthews *et al.* 1989; but see Meier & Chapman 1983).

Trentepohlia photobionts have been occasionally reported to grow out from the algal layer and emerge as free filaments projecting from the lichen thallus or thalline apothecial margin (Zahlbruckner 1907: p. 126; McGee 2002). In one case, such a filament was seen bearing a sporangium (Tschermak 1941a: p. 289). However, some authors have expressed scepticism about this interpretation, suggesting that epiphytic *Trentepohlia* might instead develop upon, and then grow into, an already formed thallus (Henssen & Jahns 1974: p. 196). More detailed observations are clearly required, but either explanation could represent another potentially significant mechanism by which exchange may occur between lichenized and free-living populations.

Tritostichococcus Pröschold & Darienko — This recent segregate of *Stichococcus* s. lat. (Pröschold & Darienko 2020) includes the *Stichococcus*-like photobionts that associate with *Chaenotheca*, a genus of lichen-forming fungi that partner with a remarkably broad spectrum of photobionts (Tibell 2001).

Trochiscia Kützing — This genus of unicellular algae is characterized by an often-thick cell wall with spine- or wart-like projections, an irregularly stellate chloroplast, and two forms of endogenous spore production, resulting in hundreds of

small cylindrical autospores or just two rounded endospores (Tschermak 1941b). *Trochiscia* currently appears to be placed among the *Chlorophyceae*, in or near *Sphaeropleales* (Fučíková *et al.* 2019). It was identified as photobiont of *Polyblastia amota* and *P. hyperborea* (Tschermak 1941b; Ahmadjian 1967) in the *Verrucariaceae*, but those reports appear to be in doubt (Ettl & Gärtner 2014) and further studies are needed. *Trochiscia* was not among the photobionts detected in the survey by Thüs *et al.* (2011) of the algal partners of *Verrucariaceae*.

Vulcanochloris Vancurová *et al.* — This newest addition to the *Trebouxia* family encompasses three recently described species with a distinctive, highly dissected chloroplast structure, and molecular sequences that place them as sister to *Asterochloris* (Vančurová *et al.* 2015). They are known mainly as principal photobionts from some thalli of *Stereocaulon vesuvianum* in the Canary Islands, although there is also a very recent report of *Vulcanochloris* from a *Stereocaulon* thallus collected in highland Bolivia (Kosecka *et al.* 2021). Most other *Stereocaulon* lineages investigated to date appear to associate with *Asterochloris* or *Chloroidium* (Vančurová *et al.* 2018). *Vulcanochloris* has also been recently reported as a minority photobiont in thalli of *Ramalina farinacea* (Moya *et al.* 2017).

Stramenopila (Heterokontae)

Heterococcus Chodat — The yellow-green (xanthophyte) algae lack fucoxanthin, the golden brownish plastidial carotenoid otherwise characteristic of the photosynthetic stramenopiles. The absence of fucoxanthin makes them rather easy to confuse with green algae. Their zoospores, however, will have the characteristic stramenopilous flagellum bearing stiff, hollow, tripartite appendages (mastigonemes). *Heterococcus* forms irregular filaments and/or cell clusters when isolated into culture (Zeitler 1954). Molecular sequences supported the light microscope identification of *Heterococcus* as photobiont in thalli of three species of *Verrucariaceae* (*Hydropunctaria rheitrophila*, *Verrucaria funckii* and *V. hydrela*) that are each in separate clades and not closely related to one another (Thüs *et al.* 2011). Another xanthophyte, *Heterothrix* (now *Xanthonema*; Silva 1979) was identified via light microscopy as photobiont of *Staurothele clopimoides* (Pereira Riquelme 1992) but that interesting report requires corroboration.

Petroderma Kuckuck — *Petroderma maculiforme* is a small crustose brown alga (*Phaeophyceae*) found on rocks in the lower intertidal zone of western North America and Europe. In San Francisco Bay, it is particularly common on discarded plastic (Sanders *et al.* 2004). The alga is a disc of tightly branched, radiating horizontal filaments, rather similar in morphology to the chlorophyte *Phycopeltis* but with a dense carpet of short, erect filaments arising proximally from the horizontal system. In the free-living state, these erect filaments may bear unilocular and/or plurilocular sporangia (zoidangia) terminally (Fritsch 1945). Chloroplasts typically possess one or several large pyrenoids that are prominent in electron micrographs (Fig. 2D) but not readily visible with light microscopy. The pyrenoids are traversed by branching tubules arising from invagination of the plastidial boundary membranes (rather than

thylakoids, as in *Trebouxia* and *Heveochlorella*); the position of the pyrenoid may be laminar, protruding to exerted, or enfolded by chloroplast lobes (Sanders *et al.* 2005). The alga was first brought to the attention of lichenologists by a footnote in a phycology dissertation (Wynne 1969) that reported it in symbiosis with a *Verrucaria* species on intertidal rocks in northern California. The lichen was not studied further until Moe (1997) re-collected it and formally described the fungal symbiont as *Verrucaria tavaresiae* (now *Wahlenbergiella tavaresiae*; Gueidan *et al.* 2011). When lichenized, the *Petroderma*

filaments are separated by fungal tissue, into which they grow and branch downwards rather than upwards as in the free-living condition (Sanders *et al.* 2004). *Petroderma* is the only member of the *Phaeophyceae* known to enter into lichen symbiosis. However, certain larger brown seaweeds, such as *Ascophyllum*, have intimate, mutualistic associations with verrucariacean fungi (Garbary & London 1995; Garbary & MacDonald 1995) that are generally not considered to be lichens on structural grounds, since the fungus grows within algal tissues as a conventional mycelium (Hawksworth 1988).

Table 1. Taxonomically grouped list of photobiont genera and mycobionts reported in association with them. The family names of the mycobionts are included in places where emphasis might be useful. id = procedures used in the study to identify the photobiont. LM = light microscopy, TEM = transmission electron microscopy. See table 1 in Tschermak-Woess (1988a) for a comprehensive list of photobiont reports prior to 1988. Taxon names follow those used in the original articles.

Alga	Fungal symbionts	References & comments
Domain Bacteria, Phylum Cyanobacteria		
[<i>Brasilonema</i>] sp.	?	Villanueva <i>et al.</i> (2018). Lichen not identified; need evidence that algal isolate is lichen photobiont.
<i>Calothrix</i> and <i>Dichothrix</i>	<i>Lichina</i> spp., <i>Placynthium nigrum</i>	see Tschermak-Woess (1988a)
<i>Chroococidiopsis</i> sp.	<i>Anema nummularium</i> , <i>Peccania cerebriiformis</i> , <i>Psorotichia columnaris</i> , <i>Gonohymenia</i> sp.	Büdel & Henssen (1983); Büdel (1985). id: LM, culture.
<i>Chroococidiopsis</i> sp.	<i>Peltula</i> spp. and 'other cyanolichen genera'	Büdel (1999). Unpublished 16S rDNA data of B. Büdel & T. Friedl. cited.
<i>Chroococidiopsis</i> sp.	<i>Anema nummularium</i> , <i>Peltula euploca</i> , <i>Thyrea pulvinata</i>	Fewer <i>et al.</i> (2002)
[<i>Chroococidiopsis</i>] sp.	?	Villanueva <i>et al.</i> (2018). Lichen not identified; need evidence that algal isolate is a lichen photobiont.
<i>Chroococidiopsis</i> / <i>Chroococidiopsidales</i>	<i>Peltula bolanderi</i> , <i>P. clavata</i> , <i>P. euploca</i> , <i>P. impressa</i> , <i>P. obscurans</i> , <i>P. placodizans</i>	V6 (2016)
<i>Gloeocapsa</i> sp.	Species of <i>Anema</i> , <i>Edwardiella</i> , <i>Gonohymenia</i> , <i>Heppia</i> , <i>Jenmania</i> , <i>Peccania</i> , <i>Phylliscum</i> , <i>Psorotichia</i> , <i>Pyrenopsis</i> , <i>Synalissa</i> and <i>Thyrea</i> ; cephalodia of <i>Amygdalaria</i> and <i>Stereocaulon</i> spp.	Numerous LM reports summarized by Tschermak-Woess (1988a). Friedl & Büdel (2008); Brodo <i>et al.</i> (2001). id: LM.
<i>Hyella</i> sp.	<i>Collemopsisidium</i> sp.	see Ahmadjian (1967). id: LM.
<i>Hyphomorpha</i> sp.	<i>Spilonema dendroides</i> , <i>S. schmidtii</i>	Henssen (1981). id: LM.
<i>Myxosarcina</i> sp.	<i>Peltula euploca</i>	Friedl & Büdel (2008)
<i>Nostoc</i> sp.	<i>Nephroma resupinatum</i> , <i>Peltigera britannica</i> , <i>P. canina</i> , <i>P. membranacea</i> ; cephalodia of <i>Peltigera aphthosa</i> , <i>P. venosa</i>	Paulsrud <i>et al.</i> (2000, 2001). High genetic specificity, also for cephalodia formation, overrules geography.
<i>Nostoc</i> sp.	<i>Collema multipartitum</i>	Oksanen <i>et al.</i> (2002)
<i>Nostoc</i> sp.	<i>Nephroma</i> , numerous spp.	Lohtander <i>et al.</i> (2003)
<i>Nostoc</i> sp.	<i>Leptogium puberulum</i> , <i>Massalongia carnosa</i> , <i>Placopsis contortuplicata</i> , <i>P. parellina</i> , <i>Psoroma cinnamomeum</i>	Wirtz <i>et al.</i> (2003). Maritime Antarctica; two main clades that also include free-living <i>Nostoc</i> .
<i>Nostoc</i> sp.	<i>Collema crispum</i> , <i>Leptogium gelatinosum</i> , <i>Lobaria amplissima</i> , <i>L. hallii</i> , <i>Massalongia carnosa</i> , <i>Nephroma bellum</i> , <i>N. helveticum</i> , <i>Pannaria conoplea</i> , <i>Peltigera canina</i> , <i>P. degenii</i> , <i>P. didactyla</i> , <i>P. horizontalis</i> , <i>P. lepidophora</i> , <i>P. membranacea</i> , <i>P. neopolydactyla</i> , <i>P. rufescens</i> , <i>Sticta beauvoisii</i> , <i>S. fuliginosa</i> ; cephalodia of <i>Peltigera aphthosa</i>	O'Brien <i>et al.</i> (2005)
<i>Nostoc</i> sp.	<i>Pseudocyphellaria crocata</i> , (=) <i>P. neglecta</i> , (=) <i>P. perpetua</i>	Summerfield & Eaton-Rye (2006)
<i>Nostoc</i> sp.	<i>Pannaria andina</i> , <i>P. araneosa</i> , <i>P. athroophila</i> , <i>P. conoplea</i> , <i>P. durietzii</i> , <i>P. elixii</i> , <i>P. euphylla</i> , <i>P. farinosa</i> , <i>P. fulvescens</i> , <i>P. isabellina</i> , <i>P. lobulifera</i> , <i>P. mosenii</i> , <i>P. obscura</i> , <i>P. pallida</i> , <i>P. patagonica</i> , <i>P. rubiginella</i> , <i>P. rubiginosa</i> , <i>P. sphinctrina</i> , <i>P. tavaresii</i>	Elvebakk <i>et al.</i> (2008)
<i>Nostoc</i> sp.	<i>Degeelia atlantica</i> , <i>D. plumbea</i>	Otálora <i>et al.</i> (2013)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Nostoc</i> sp.	<i>Polychidium muscicola</i>	Jayalal <i>et al.</i> (2012)
<i>Nostoc</i> sp.	Species of <i>Fuscopannaria</i> , <i>Kroswia</i> , <i>Pannaria</i> , <i>Parmeliella</i> and <i>Physma</i>	Magain & Sérusiaux (2014). Homoiomerous (gelatinous) vs stratified anatomy correlated with <i>Nostoc</i> strain, not fungal family, across <i>Pannariaceae</i> and <i>Collemaaceae</i> .
<i>Nostoc</i> sp.	<i>Leptogium lichenoides</i> , <i>Peltigera islandica</i> , <i>P. ponojensis</i>	Joneson & O'Brien (2017)
<i>Nostoc</i> sp.	<i>Peltigera</i> sections <i>Peltigera</i> and <i>Retifoviatae</i>	Magain <i>et al.</i> (2018)
<i>Nostoc</i> sp.	<i>Pectenium plumbea</i>	Cardós <i>et al.</i> (2019)
<i>Rhizonema andinum</i>	<i>Cora arachnodavidea</i> , <i>C. barbulate</i> , <i>C. bovei</i> , <i>C. dewisanti</i> , <i>C. dewisanti-mexicana</i> , <i>C. elephas</i> , <i>C. hawksworthiana</i> , <i>C. hochesuordensis</i> , <i>C. pseudobovei</i> , <i>Cora</i> sp.; cephalodia of <i>Stereocaulon fronduliferum</i>	Dal Forno <i>et al.</i> (2020)
<i>R. interruptum</i> lineage	<i>Coccocarpia filiformis</i> , <i>C. palmicola</i> , <i>C. stellata</i> , <i>Cora applanata</i> , <i>C. aspera</i> , <i>C. auriculeslia</i> , <i>C. canari</i> , <i>C. ciferrii</i> , <i>C. crispoleslia</i> , <i>C. galapagoensis</i> , <i>C. hymenocarpa</i> , <i>C. melvinii</i> , <i>C. paraciferrii</i> , <i>C. reticulifera</i> , <i>C. smaragdina</i> , <i>C. soledavidia</i> , <i>C. squamiformis</i> , <i>C. strigosa</i> , <i>C. suturifera</i> , <i>C. terrestris</i> , <i>C. viliewoa</i> , <i>Cora</i> sp. <i>Cyphellostereum</i> sp., <i>Dictyonema aeruginosum</i> , <i>D. barbatum</i> , <i>D. darwinianum</i> , <i>D. hernandezii</i> , <i>D. interruptum</i> [= <i>D. coppinsii</i>], <i>D. phyllogenum</i> , <i>D. sericeum</i> , <i>D. schenkianum</i> , <i>Dictyonema</i> sp.; cephalodia of <i>Stereocaulon ramulosum</i>	Dal Forno <i>et al.</i> (2020)
<i>R. neotropicum</i>	<i>Acantholichen pannarioides</i> , <i>Coccocarpia palmicola</i> , <i>Cora gigantea</i> , <i>C. leslactuca</i> , <i>C. rubrosanguinea</i> , <i>Corella</i> sp., <i>Dictyonema</i> sp.	Dal Forno <i>et al.</i> (2020)
<i>Rhizonema</i> sp.	<i>Acantholichen pannarioides</i> , <i>Coccocarpia filiformis</i> , <i>C. palmicola</i> , <i>C. stellata</i> , <i>Dictyonema aeruginosum</i> , <i>D. coppinsii</i> , <i>D. glabratum</i> , <i>D. hernandezii</i> , <i>D. phyllogenum</i> , <i>D. schenkianum</i> , <i>D. sericeum</i> , <i>Coccocarpia</i> ; cephalodia of <i>Stereocaulon fronduliferum</i> and <i>S. ramulosum</i>	Lücking <i>et al.</i> (2009). Shows photobiont more closely related to <i>Nostoc</i> clade than to <i>Scytonema</i> s. str. Lücking <i>et al.</i> (2014): formal description of genus <i>Rhizonema</i> .
<i>Rhizonema</i> sp.	<i>Athelia andina</i> , <i>Cyphellostereum pusiolum</i>	Oberwinkler (2012). Mycobiont also associates with green photobiont.
<i>Rhizonema</i> sp.	<i>Athelia phycophila</i>	Jülich (1972). Reported as <i>Scytonema</i> . <i>Athelia poeltii</i> also said to associate with filamentous cyanobacteria; Jülich (1978).
<i>Rhizonema</i> sp.	<i>Coccocarpia palmicola</i> , <i>Erioderma pedicellatum</i> , <i>E. solediatum</i> , <i>Leptogidium</i> sp., <i>Lichinodium</i> sp., <i>Moelleropsis nebulosa</i> , <i>Parmeliella parvula</i>	Cornejo & Scheidegger (2016); Cornejo <i>et al.</i> (2016). Boreal env. (Newfoundland). Also symbiotic among liverwort <i>Frullania asagrayana</i> .
<i>Rhizonema</i> sp.	<i>Lichinodium ahlneri</i> , <i>L. sirosiphoideum</i> (Leotiomycetes!)	Prieto <i>et al.</i> (2019)
<i>Rhizonema</i> sp.	<i>Dictyonema moorei</i>	Masumoto (2020)
<i>Rivularia</i> sp.	<i>Lichina confinis</i> , <i>L. pygmaea</i>	Ortiz-Álvarez <i>et al.</i> (2015)
<i>Scytonema</i> sp.	<i>Pyrenothrix nigra</i>	Tschermak-Woess <i>et al.</i> (1983). id: LM & TEM.
<i>Scytonema</i> sp.	Species of <i>Heppia</i> , <i>Lichinodium</i> , <i>Thermutis</i> and <i>Zahlbrucknerella</i>	Henssen (1994). DNA sequence data needed. Might be <i>Rhizonema</i> , but see Vö (2016) concerning <i>Heppia</i> .
<i>Scytonema</i> sp.	<i>Heppia lutosa</i>	Vö (2016). A 16s rRNA sequence suggests true <i>Scytonema</i> , not <i>Rhizonema</i> .
<i>Stigonema</i> sp.	Species of <i>Ephebe</i> and <i>Spiloneima</i>	Henssen & Jahns (1974). id: LM.
<i>Stigonema</i> sp.	Cephalodia of <i>Stereocaulon</i>	Lavoie <i>et al.</i> (2020)
<i>Tolypothrix</i> sp.	<i>Thermutopsis</i> sp.	Henssen (1990)
cf. <i>Tolypothrix</i>	Cephalodia of <i>Placopsis perrugosa</i> and <i>P. stenophylla</i>	Raggio <i>et al.</i> (2012)
Domain Eukarya, Eukaryote supergroup Archaeplastida, Kingdom Plantae, Division Chlorophyta Class Trebouxiophyceae, Order Chlorellales		
<i>Auxenochlorella</i>	<i>Psoroglaena stigonemoides</i>	Thüs <i>et al.</i> (2011)
Near <i>A. protothecoides</i>	<i>Psoroglaena stigonemoides</i>	Nyati <i>et al.</i> (2007). Strain P-1015.

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
Near <i>Chlorella sorokiniana</i>	<i>Porpidia crustulata</i>	Li <i>et al.</i> (2013)
<i>Nannochloris normandinae</i>	<i>Normandina pulchella</i>	Tschermak-Woess (1988a). Pröschold & Darienko (2020) synonymize <i>N. normandinae</i> with <i>Diplosphaera chodatii</i> .
Class Trebouxiphyceae, Order Prasiolales		
<i>Deuterostichococcus allas</i> and <i>D. antarcticus</i>	<i>Placopsis antarctica</i> , <i>P. contortuplicata</i>	Beck <i>et al.</i> (2019). Reclassified by Pröschold & Darienko (2020).
<i>D. deasonii</i>	<i>Staurothele clopima</i>	Hodač <i>et al.</i> (2016); Pröschold & Darienko (2020).
<i>Diplosphaera chodatii</i>	<i>Dermatocarpon luridum</i> var. <i>luridum</i>	Fontaine <i>et al.</i> (2012); Doering <i>et al.</i> (2020).
<i>Diplosphaera</i> sp.	<i>Agonimia koreana</i> , <i>A. opuntiella</i> , <i>A. repleta</i> , <i>A. tristicula</i> , <i>Bagliettoa parmigera</i> , <i>Catapyrenium cinereum</i> , <i>C. daedaleum</i> , <i>Dermatocarpon luridum</i> , <i>D. miniatum</i> , <i>Endocarpon adscendens</i> , <i>E. diffractellum</i> , <i>E. pallidulum</i> , <i>E. pusillum</i> , <i>Flakea papillata</i> , <i>Neocatapyrenium rhizinosum</i> , <i>Normandina acroglypta</i> , <i>N. pulchella</i> , <i>Placidopsis cartilaginea</i> , <i>Placopyrenium bucekii</i> , <i>Polyblastia cupularis</i> , <i>P. viridescens</i> , <i>Staurothele areolata</i> , <i>S. clopima</i> , <i>S. clopimoides</i> , <i>S. drummondii</i> , <i>S. fissa</i> , <i>S. frustulenta</i> , <i>Verrucaria elaeina</i> , <i>V. hochstetteri</i> , <i>V. nigrescens</i> , <i>V. praetermissa</i> , <i>V. rupestris</i> (Verrucariaceae)	Thüs <i>et al.</i> (2011). Morphologically similar to <i>Stichococcus</i> , but not close to type sp. <i>S. bacillaris</i> .
<i>Diplosphaera</i> sp.	<i>Ramalina farinacea</i>	Moya <i>et al.</i> (2017). Minority photobiont.
<i>Prasiola borealis</i>	<i>Mastodia tessellata</i>	Moniz <i>et al.</i> (2012). Authors refer to symbiotic form as lichenized. Stipe section shows photobiont well-embedded in fungal tissue.
<i>P. borealis</i> and one undescribed sp.	<i>Mastodia tessellata</i>	Pérez-Ortega <i>et al.</i> (2012); Garrido-Benavent <i>et al.</i> (2017).
<i>P. delicata</i>	<i>Mastodia tessellata</i>	Moniz <i>et al.</i> (2014); Garrido-Benavent <i>et al.</i> (2018).
<i>Prasiola</i> sp.	<i>Mastodia tessellata</i>	Thüs <i>et al.</i> (2011)
<i>Pseudochlorella pyrenoidosa</i>	<i>Micarea assimilata</i>	Zeitler (1954); Tschermak-Woess (1988a).
<i>P. signiensis</i>	<i>Trapelia coarctata</i>	Darienko <i>et al.</i> (2016)
<i>Pseudochlorella</i> sp.	<i>Stereocaulon strictum</i>	Brunner & Honegger (1985). Pyrenoid ultrastructure suggests this strain might be <i>Chloroidium</i> , as indicated by DNA sequences from other <i>Stereocaulon</i> photobionts.
<i>Pseudochlorella</i> sp.	<i>Placynthiella dasaea</i>	Voytsekhovich <i>et al.</i> (2011). id: LM, culture.
<i>Pseudochlorella</i> sp.	<i>Umbilicaria antarctica</i>	Park <i>et al.</i> (2015). In some thallus samples, along with <i>Trebouxia</i> . Status as photobiont needs corroboration.
<i>Pseudostichococcus</i> clade	<i>Neocatapyrenium rhizinosum</i>	Hodač <i>et al.</i> (2016); Pröschold & Darienko (2020).
<i>Tritostichococcus conicybes</i>	<i>Chaenotheca</i> sp.	Pröschold & Darienko (2020)
Class Trebouxiphyceae, Order Trebouxiales		
<i>Asterochloris antarctica</i>	<i>Cladonia</i> sp.	Kim <i>et al.</i> (2020)
<i>Asterochloris</i> clade I1	<i>Cladonia corymbescens</i> , <i>C. furcata</i> , <i>C. pyxidata</i> , <i>C. rangiferina</i>	Řídká <i>et al.</i> (2014)
<i>Asterochloris</i> clade I2	<i>Cladonia furcata</i>	Řídká <i>et al.</i> (2014)
<i>Asterochloris</i> clade II	<i>Cladonia subtenuis</i>	Yahr <i>et al.</i> (2006)
<i>Asterochloris</i> clade 9	<i>Cladonia cariosa</i> , <i>C. coniocraea</i> , <i>C. delavayi</i> , <i>C. fruticulosa</i> , <i>C. praetermissa</i> , <i>C. scabriuscula</i> , <i>C. verticillata</i>	Řídká <i>et al.</i> (2014)
<i>Asterochloris</i> clade 12	<i>Cladonia pyxidata</i>	Řídká <i>et al.</i> (2014)
<i>Asterochloris</i> clade 16	<i>Cladonia fruticulosa</i>	Řídká <i>et al.</i> (2014)
<i>A. echinata</i>	<i>Lepraria caesioalba</i> , <i>L. rigidula</i>	Škaloud <i>et al.</i> (2015)
<i>A. erici</i>	<i>Cladonia calycanta</i> , <i>C. chlorophaea</i> , <i>C. crispata</i> , <i>C. floerkeana</i> , <i>C. gracilis</i> ssp. <i>turbinata</i> , <i>C. ramulosa</i> , <i>C. squamosissima</i>	Nakano & Iguchi (1994). id: LM, culture.

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Asterochloris excentrica</i>	<i>Lepraria caesioalba</i> , <i>L. neglecta</i> , <i>L. rigidula</i>	Škaloud & Peksa (2010)
<i>A. friedlii</i>	<i>Lepraria caesioalba</i> , <i>L. lobificans</i>	Škaloud et al. (2015)
<i>A. friedlii</i>	<i>Lepraria achariana</i> , <i>L. finkii</i>	Kosecka et al. (2021)
<i>A. friedlii</i>	<i>Cladonia fimbriata</i>	Pino-Bodas & Stenroos (2020)
<i>A. gaertneri</i>	<i>Lepraria incana</i> , <i>L. rigidula</i>	Škaloud et al. (2015)
<i>A. glomerata</i>	<i>Cladonia humilis</i>	Nakano & Iguchi (1994). id: LM, culture.
<i>A. glomerata</i>	<i>Diploschistes muscorum</i>	Škaloud & Peksa (2010)
<i>A. glomerata</i>	<i>Stereocaulon pileatum</i> , <i>S. saxatile</i> , <i>Stereocaulon</i> sp.	Peksa & Škaloud (2011)
<i>A. glomerata</i>	<i>Cladonia coniocraea</i> , <i>C. squamosa</i> , <i>Stereocaulon evolutoides</i> , <i>S. pileatum</i>	Řídká et al. (2014)
<i>A. glomerata</i>	<i>Cladonia deformis</i> , <i>C. pleurota</i>	Steinová et al. (2019)
<i>A. glomerata</i>	<i>Cladonia grayi</i>	Armaleo et al. (2019)
<i>A. glomerata</i>	<i>C. caroliniana</i> , <i>C. cornuta</i> , <i>C. deformis</i> , <i>C. hondoensis</i> , <i>C. krogiana</i> , <i>C. mitis</i> , <i>C. monomorpha</i> , <i>C. oricola</i> , <i>C. phyllophora</i> , <i>C. piedmontensis</i> , <i>C. pleurota</i> , <i>C. submitis</i> , <i>C. subulata</i> , <i>C. uncialis</i>	Pino-Bodas & Stenroos (2020)
<i>A. glomerata/irregularis</i> clade	<i>Cladonia cariosa</i> , <i>C. cervicornis</i> ssp. <i>verticillata</i> , <i>C. phyllophora</i>	Rola et al. (2021)
<i>A. irregularis</i>	<i>Squamarina lentigera</i>	Beck et al. (2002)
<i>A. irregularis</i>	<i>Cladonia furcata</i> , <i>Lecidea fuscoatra</i> , <i>L. plana</i> , <i>Lepraria caesioalba</i> , <i>Porpidia crustulata</i> , <i>P. flavocaerulescens</i> , <i>P. macrocarpa</i> , <i>Porpidia</i> spp., <i>Stereocaulon vesuvianum</i>	Beck (2002)
<i>A. irregularis</i>	<i>Cladonia arbuscula</i> , <i>C. mitis</i> , <i>Stereocaulon botryosum</i> , <i>S. pileatum</i> , <i>S. subcoralloides</i>	Škaloud & Peksa (2010); Škaloud et al. (2015).
<i>A. irregularis</i>	<i>Stereocaulon alpinum</i> , <i>S. apocalypticum</i> , <i>S. intermedium</i> , <i>S. paschale</i> , <i>S. symphycheilum</i> , <i>S. tomentosum</i> , <i>S. vesuvianum</i>	Vančurová (2012)
<i>A. irregularis</i>	<i>Stereocaulon botryosum</i> , <i>S. pileatum</i> , <i>S. subcoralloides</i> , <i>S. vesuvianum</i> , <i>Stereocaulon</i> sp.	Peksa & Škaloud (2011)
<i>A. irregularis</i>	<i>Cladonia deformis</i> , <i>C. pleurota</i>	Steinová et al. (2019)
<i>A. irregularis</i>	<i>Cladonia albonigra</i> , <i>C. alinii</i> , <i>C. amaurocraea</i> , <i>C. botrytes</i> , <i>C. crispata</i> , <i>C. ecmocyna</i> , <i>C. gracilis</i> , <i>C. granulans</i> , <i>C. kanewskii</i> , <i>C. labradorica</i> , <i>C. macrophylla</i> , <i>C. uliginosa</i> , <i>C. uncialis</i>	Pino-Bodas & Stenroos (2020)
<i>A. italiana</i>	<i>Cladonia capitellata</i> , <i>C. scabriuscula</i>	Škaloud et al. (2015)
<i>A. italiana</i>	<i>Cladonia coccifera</i> , <i>C. diversa</i>	Steinová et al. (2019)
<i>A. italiana</i>	<i>Stereocaulon condensatum</i>	Vančurová (2012)
<i>A. italiana</i>	<i>Cladonia bellidiflora</i> , <i>C. borbonica</i> , <i>C. brevis</i> , <i>C. callosa</i> , <i>C. capitellata</i> , <i>C. ciliata</i> , <i>C. foliacea</i> , <i>C. groenlandica</i> , <i>C. islandica</i> , <i>C. lepidophora</i> , <i>C. mawsonii</i> , <i>C. merochlorophaea</i> , <i>C. neozelandica</i> , <i>C. pleurota</i> , <i>C. portentosa</i> , <i>C. prolifica</i> , <i>C. pulvinata</i> , <i>C. rappi</i> , <i>C. rigida</i> , <i>C. subcervicornis</i> , <i>C. subsubulata</i> , <i>C. ustulata</i> , <i>C. weymouthii</i>	Pino-Bodas & Stenroos (2020)
<i>A. italiana</i> clade	<i>Diploschistes muscorum</i>	Rola et al. (2021)
<i>A. leprarii</i>	<i>Lepraria caesioalba</i> , <i>L. neglecta</i>	Škaloud et al. (2015)
<i>A. lobophora</i>	<i>Cladonia</i> cf. <i>bacillaris</i> , <i>Diploschistes muscorum</i> , <i>Lepraria alpina</i> , <i>L. borealis</i> , <i>L. caesioalba</i>	Škaloud et al. (2015)
<i>A. lobophora</i>	<i>Cladonia coccifera</i>	Steinová et al. (2019)
<i>A. lobophora</i>	<i>Cladonia rei</i>	Pino-Bodas & Stenroos (2020)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Asterochloris lobophora</i> / <i>phycobiontica</i> clade	<i>Cladonia cariosa</i> , <i>C. rei</i> , <i>Diploschistes muscorum</i>	Rola <i>et al.</i> (2021)
<i>A. magna</i>	<i>Porpidia contraponenda</i>	Beck (2002)
<i>A. magna</i>	<i>Cladonia chlorophaea</i> , <i>C. decorticata</i>	Pino-Bodas & Stenroos (2020)
<i>A. mediterranea</i>	<i>Cladonia cervicornis</i> , <i>C. concoluta</i> , <i>C. foliacea</i> , <i>C. rangiformis</i>	Moya <i>et al.</i> (2015)
<i>A. mediterranea</i>	<i>Cladonia calycantha</i> , <i>Stereocaulon pachycephalum</i>	Kosecka <i>et al.</i> (2021)
<i>A. mediterranea</i>	<i>Cladonia corsicana</i> , <i>C. crispata</i> , <i>C. cryptochlorophaea</i> , <i>C. diversa</i> , <i>C. glauca</i> , <i>C. rei</i>	Pino-Bodas & Stenroos (2020)
<i>A. phycobiontica</i>	<i>Anzina</i> [Varicellaria] <i>carneonivea</i>	Tschermak-Woess (1980a); Škaloud <i>et al.</i> (2015).
<i>A. phycobiontica</i>	<i>Stereocaulon alpinum</i>	Vančurová (2012)
<i>A. phycobiontica</i>	<i>Lepraria neglecta</i>	Škaloud & Peksa (2010)
<i>A. phycobiontica</i>	<i>Lepraria alpina</i> , <i>L. caesioalba</i> , <i>L. neglecta</i> , <i>Lepraria</i> sp.	Peksa & Škaloud (2011)
<i>A. pseudoirregularis</i>	<i>Cladonia gracilis</i> , <i>Cladonia</i> sp.	Kim <i>et al.</i> (2020)
<i>A. pseudoirregularis</i>	<i>Cladonia amaurocraea</i> , <i>C. gracilis</i> , <i>C. granulans</i> , <i>C. vulcanii</i> , <i>Pycnothelia papillaria</i>	Pino-Bodas & Stenroos (2020)
<i>A. sejongensis</i>	<i>Cladonia pyxidata</i> , <i>Sphaerophorus globosus</i>	Kim <i>et al.</i> (2017)
<i>A. stereocaulonica</i>	<i>Stereocaulon alpinum</i>	Kim <i>et al.</i> (2020)
<i>A. woessiae</i>	<i>Cladonia foliacea</i> , <i>Lepraria borealis</i> , <i>L. caesioalba</i> , <i>L. crassissima</i> , <i>L. nigrocincta</i> , <i>L. nylanderiana</i> , <i>Lepraria</i> sp., <i>Stereocaulon saxatile</i>	Škaloud <i>et al.</i> (2015)
<i>A. woessiae</i>	<i>Cladonia coccifera</i>	Steinová <i>et al.</i> (2019)
<i>A. woessiae</i>	<i>Cladonia conista</i> , <i>C. foliacea</i>	Pino-Bodas & Stenroos (2020)
<i>A. woessiae</i> clade	<i>Diploschistes muscorum</i>	Rola <i>et al.</i> (2021)
<i>Asterochloris</i> sp.	<i>Bagliettoa cazzae</i>	Thüs <i>et al.</i> (2011)
<i>Asterochloris</i> sp.	<i>Psora decipiens</i>	Ruprecht <i>et al.</i> (2014). Also <i>Trebouxias</i> and a <i>Chloroidium</i> in this lichen sp., but see Williams <i>et al.</i> (2017).
<i>Asterochloris</i> sp.	<i>Lepraria borealis</i> , <i>Ochrolechia frigida</i>	Engelen <i>et al.</i> (2010, 2016). Maritime Antarctica.
<i>Asterochloris</i> sp.	<i>Lecidea lurida</i> , <i>Psora decipiens</i> , <i>Squamarina cartilaginea</i>	Schaper & Ott (2003). Gotland, Sweden.
<i>Asterochloris</i> sp.	<i>Ramalina farinacea</i>	Moya <i>et al.</i> (2017). Minority photobiont.
<i>Asterochloris</i> sp.	<i>Porpidia</i> from southern South America	Ruprecht <i>et al.</i> (2020). All mycobiont species also partnered with <i>Trebouxia</i> .
<i>Asterochloris</i> sp.	<i>Cladonia acuata</i> , <i>C. aggregata</i> , <i>C. andesita</i> , <i>C. arbuscula</i> ssp. <i>boliviana</i> , <i>C. calycantha</i> , <i>C. chlorophaea</i> , <i>C. confusa</i> , <i>C. dactylota</i> , <i>C. didyma</i> , <i>C. furcata</i> , <i>C. granulosa</i> , <i>C. isabellina</i> , <i>C. melanopoda</i> , <i>C. pocillum</i> , <i>C. ramulosa</i> , <i>C.</i> cf. <i>subradiosa</i> , <i>C. vouauxii</i> , <i>Lepraria cryptovouauxii</i> , <i>L. hodkinsoniana</i> , <i>Stereocaulon myriocarpum</i> , <i>S. tomentosum</i>	Kosecka <i>et al.</i> (2021)
<i>Asterochloris</i> sp.	<i>Cladonia arbuscula</i> , <i>C. fimbriata</i> , <i>C. foliacea</i> , <i>C. rei</i> , <i>Diploschistes muscorum</i> , <i>Lepraria alpina</i> , <i>L. borealis</i> , <i>L. caesioalba</i> , <i>L. lobificans</i> , <i>L. neglecta</i> , <i>L. nigrocincta</i> , <i>L. nylanderiana</i> , <i>L. rigidula</i> , <i>Pilophorus</i> sp., <i>Stereocaulon</i> <i>botryosum</i> , <i>S. paschale</i> , <i>S. pileatum</i> , <i>S. saxatile</i> , <i>S. subcoralloides</i> , <i>S. tomentosum</i> , <i>S. vesuvianum</i>	Nelsen & Gargas (2008). Formal emendation of genus by Škaloud & Peksa (2010) to include <i>Trebouxia</i> subgen. <i>Eleutherococcus</i> Tschermak-Woess.
<i>Asterochloris</i> sp.	<i>Cladia aggregata</i> , <i>Cladonia</i> cf. <i>bacillaris</i> , <i>C. evansii</i> , <i>C.</i> <i>fimbriata</i> , <i>C. perforata</i> , <i>C. rangiferina</i> , <i>Lepraria</i> <i>atromentosa</i> , <i>L. caesiella</i> , <i>L. caesioalba</i> , <i>L. incana</i> , <i>L. lobificans</i> , <i>L. nigrocincta</i> , <i>L. nivalis</i> , <i>L. nylanderiana</i> , <i>Lepraria</i> spp., <i>Pilophorus acicularis</i> , <i>P.</i> cf. <i>cereolus</i> , <i>Stereocaulon dactylophyllum</i> , <i>S. paschale</i> , <i>S. saxatile</i> , <i>S. tomentosum</i> , <i>S. vesuvianum</i> , <i>Stereocaulon</i> spp.	Nelsen & Gargas (2008)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Asterochloris</i> spp.	Species of <i>Lepraria</i> and <i>Stereocaulon</i>	Peksa & Škaloud (2011)
<i>Asterochloris</i> spp.	<i>Cladonia furcata</i>	Řídká et al. (2014)
<i>Myrmecia</i> biatorellae	<i>Catapyrenium rufescens</i> , <i>Dermatocarpon hepaticum</i> , <i>D. tuckermanni</i> , <i>D. velebiticum</i> , <i>Lecidea berengeriana</i> , <i>Lobaria linita</i> , <i>Polysporina simplex</i> , <i>Psora decipiens</i> , <i>P. globifera</i> , <i>Psoroma hypnorum</i> , <i>Sarcogyne privigna</i> , <i>Verrucaria submersella</i>	Tschermak-Woess (1988a)
<i>M. biatorellae</i>	<i>Fuscidea cyathoides</i> var. <i>japonica</i> , <i>Leproloma</i> sp.	Watanabe et al. (1997). Maritime lichens. id: LM.
<i>M. biatorellae</i>	<i>Placidium squamulosum</i>	Voytsekhovich & Beck (2016)
<i>M. biatorellae</i>	<i>Heteroplacidium contumescens</i> , <i>H. imbricatum</i> , <i>Placidium acarosporoides</i> , <i>P. arboreum</i> , <i>P. chilense</i> , <i>P. lachneum</i> , <i>P. lacinulatum</i> , <i>P. squamulosum</i> , <i>P. umbrinum</i> (Verrucariaceae)	Thüs et al. (2011)
<i>M. biatorellae</i>	<i>Psora decipiens</i>	Williams et al. (2017). Austrian samples.
<i>M. israelensis</i>	<i>Psora decipiens</i> , <i>P. saviczii</i> , <i>Placidium pilosellum</i> , <i>Placidium</i> spp., <i>Clavascidium</i> spp.	Moya et al. (2018)
<i>Myrmecia</i> sp.	<i>Psora decipiens</i>	Williams et al. (2017). Transplant study; algal layer degenerates, no photobiont switching.
<i>Trebouxia aggregata</i>	<i>Xanthoria parietina</i>	Beck & Mayr (2012)
<i>T. angustilobata</i>	<i>Acarospora sinopica</i> , <i>Porpidia tuberculosa</i> , <i>Tremolecia atrata</i>	Beck (2002)
<i>T. angustilobata</i>	<i>Protoparmelia badia</i>	Singh et al. (2017)
<i>T. angustilobata</i>	<i>Pseudephebe</i> sp.	Garrido-Benavent et al. (2020)
<i>T. arboricola</i>	<i>Lecania cyrtella</i> , <i>Lecanora</i> sp., <i>Lecidella elaeochroma</i> , <i>Xanthoria parietina</i>	Beck et al. (1998)
<i>T. arboricola</i>	<i>Anaptychia ciliaris</i>	Helms et al. (2001); Dahlkild et al. (2001).
<i>T. arboricola</i>	<i>Pleurosticta acetabulum</i>	Beck & Koop (2001)
<i>T. arboricola</i>	<i>Chaenotheca phaeocephala</i>	Tibell & Beck (2002)
<i>T. arboricola</i>	<i>Ramalina farinacea</i> , <i>R. pollinaria</i> , <i>R. siliquosa</i>	Tschaikner et al. (2007). id: LM, cultures.
<i>T. arboricola</i>	<i>Caloplaca fernandeziana</i> , <i>C. orthoclada</i>	Vargas Castillo & Beck (2012)
<i>T. arboricola</i>	<i>Xanthoria aureola</i> , <i>X. calcicola</i> , <i>X. capensis</i> , <i>X. ectaneoides</i> , <i>X. flammea</i> , <i>X. hirsuta</i> , <i>X. karrooensis</i> , <i>X. ligulata</i> , <i>X. parietina</i> , <i>X. polycarpa</i> , <i>X. turbinata</i>	Nyati et al. (2013, 2014)
<i>T. arboricola</i>	<i>Ochrolechia</i> sp., <i>Xanthoria calcicola</i> , <i>X. parietina</i>	Voytsekhovich & Beck (2016)
<i>T. arboricola</i> clade	<i>Acarospora conafii</i> , <i>Ramalina thrausta</i>	Jung et al. (2019)
<i>T. arboricola</i> / <i>T. gigantea</i> clade	<i>Xanthoparmelia</i> spp.	Leavitt et al. (2013)
<i>T. arboricola</i> / <i>T. gigantea</i> clade	<i>Cladonia cariosa</i> , <i>C. rei</i> , <i>Diploschistes muscorum</i>	Osyczka et al. (2021)
Clade A (= <i>arboricola</i> / <i>gigantea</i> group)	Species of <i>Melanelixia</i> , <i>Melanohalea</i> , <i>Montanelia</i> , <i>Oropogon</i> , <i>Parmotrema</i> , <i>Protoparmeliopsis</i> , <i>Rhizoplaca</i> and <i>Xanthoparmelia</i>	Leavitt et al. (2015)
<i>T. 'arnoldoi'</i>	<i>Buellia zoharyi</i>	Molins et al. (2020)
<i>T. asymmetrica</i>	<i>Buellia zoharyi</i>	Helms et al. (2001); Molins et al. (2020).
<i>T. asymmetrica</i>	<i>Fulgensia fulgida</i> , <i>Toninia sedifolia</i>	Beck et al. (2002)
<i>T. asymmetrica</i>	<i>Protoparmeliopsis muralis</i>	Guzow-Krzemińska (2006)
<i>T. asymmetrica</i>	<i>Caloplaca teicholyta</i> , <i>Circinaria contorta</i>	Voytsekhovich & Beck (2016)
<i>T. asymmetrica</i>	<i>Circinaria</i> spp.	Molins et al. (2018)
<i>T. asymmetrica</i> clade	<i>Placidium</i> sp.	Jung et al. (2019)
<i>T. australis</i>	<i>Lecanora farinacea</i> , <i>L. polytropha</i> , <i>Rhizocarpon geographicum</i>	Beck (2002)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Trebouxia brindabellae</i>	<i>Protoparmelia badia</i>	Singh <i>et al.</i> (2017)
<i>T. corticola</i>	<i>Parmotrema tinctorum</i>	Ohmura <i>et al.</i> (2006, 2019)
<i>T. corticola</i>	<i>Umbilicaria muehlenbergii</i>	Cao <i>et al.</i> (2015)
<i>T. crenulata</i>	<i>Ramalina capitata</i>	Tschaikner <i>et al.</i> (2007)
<i>T. crenulata</i>	<i>Xanthoria parietina</i>	Beck & Mayr (2012)
<i>T. crenulata</i>	<i>Caloplaca aurantia</i> , <i>C. xerica</i> , <i>Candelariella medians</i> , <i>Lecanora swartzii</i> , <i>Leproplaca xantholyta</i> , <i>Tephromela atra</i> , <i>Xanthoria calcicola</i>	Voytsekhovich & Beck (2016)
<i>T. crespoana</i>	<i>Parmotrema pseudotinctorum</i>	Škaloud <i>et al.</i> (2018)
<i>T. cretacea</i>	<i>Aspicilia desertorum</i> , <i>Rusavskia papillifera</i>	Voytsekhovich & Beck (2016)
<i>T. cretacea</i>	<i>Buellia zoharyi</i>	Molins <i>et al.</i> (2020)
<i>T. decolorans</i>	<i>Lecanora rupicola</i>	Blaha <i>et al.</i> (2006)
<i>T. decolorans</i>	<i>Xanthoria parietina</i>	Beck & Mayr (2012)
<i>T. decolorans</i>	<i>Xanthomendoza borealis</i> , <i>Xm. fulva</i> , <i>Xm. hasseana</i> , <i>Xanthoria candelaria</i> , <i>X. elegans</i> , <i>X. parietina</i> , <i>X. polycarpa</i>	Nyati <i>et al.</i> (2013, 2014)
<i>T. decolorans</i>	<i>Ramalina menziesii</i>	Werth & Sork (2014)
<i>T. decolorans</i>	<i>Caloplaca orthoclada</i>	Vargas Castillo & Beck (2012)
<i>T. decolorans</i>	<i>Ramalina fraxinea</i>	Catalá <i>et al.</i> (2016)
<i>T. decolorans</i>	<i>Anaptychia ciliaris</i> , <i>Xanthoria parietina</i>	Dal Grande <i>et al.</i> (2014a)
<i>T. decolorans</i>	<i>Anaptychia setifera</i> , <i>Candelariella medians</i>	Voytsekhovich & Beck (2016)
<i>Trebouxia</i> clade G (<i>galapagensis</i> / <i>usneae</i> group)	Species of <i>Parmotrema</i> and <i>Xanthoparmelia</i>	Leavitt <i>et al.</i> (2015)
<i>T. gelatinosa</i>	<i>Caloplaca</i> spp., <i>Hymenelia</i> sp., <i>Lecanora subimergens</i> , <i>Lecidella enteroleucella</i> , <i>Ochrolechia parellula</i> , <i>Rinodina</i> sp., <i>Verrucaria</i> sp.	Watanabe <i>et al.</i> (1997). Maritime lichens. id: LM.
<i>T. gelatinosa</i>	<i>Physcia semipinnata</i>	Helms <i>et al.</i> (2001)
<i>T. gelatinosa</i>	<i>Flavoparmelia caperata</i> , <i>Punctelia subrudecta</i> , <i>Teloschistes</i> sp., <i>Xanthoria</i> sp.	Doering & Piercey-Normore (2009)
<i>T. gelatinosa</i>	<i>Josefpoeltia parva</i> , <i>Teloschistes chrysophthalmus</i> , <i>T. hosseusianus</i> , <i>Xanthomendoza novozelandica</i> , <i>Xm. weberi</i>	Nyati <i>et al.</i> (2013, 2014)
<i>T. gelatinosa</i>	<i>Rhizocarpon geographicum</i>	Voytsekhovich & Beck (2016)
<i>T. gigantea</i>	<i>Parmelia subramigera</i>	Watanabe <i>et al.</i> (1997). Maritime. id: LM.
<i>T. gigantea</i>	<i>Rinodinella controversa</i>	Helms <i>et al.</i> (2001)
<i>T. gigantea</i>	<i>Protoparmeliopsis muralis</i>	Guzow-Krzemińska (2006)
<i>T. gigantea</i>	<i>Caloplaca orthoclada</i>	Vargas Castillo & Beck (2012)
<i>T. gigantea</i>	<i>Caloplaca erythrocarpia</i> , <i>Candelariella medians</i>	Voytsekhovich & Beck (2016)
<i>T. glomerata</i>	Species of <i>Polyblastia</i> , <i>Rinodina</i> and <i>Verrucaria</i>	Watanabe <i>et al.</i> (1997). Maritime lichens. id: LM.
<i>T. higginsiae</i>	<i>Aspicilia</i> sp., <i>Buellia stellulata</i> , <i>Caloplaca kobeana</i> , <i>Hymenelia</i> sp., <i>Lecidella enteroleucella</i> , <i>Parmelia congensis</i> , <i>Rhizocarpon</i> sp., <i>Rinodina</i> sp., <i>Rinodinella</i> sp.	Watanabe <i>et al.</i> (1997). Maritime lichens. id: LM.
' <i>T. hypogymniae</i> '	<i>Hypogymnia physodes</i>	Hauck <i>et al.</i> (2007)
' <i>T. hypogymniae</i> '	<i>Evernia divaricata</i> , <i>E. mesomorpha</i>	Piercey-Normore (2009)
<i>T. impressa</i>	<i>Parmelia mexicana</i>	Watanabe <i>et al.</i> (1997). Maritime lichens. id: LM.
<i>T. impressa</i>	<i>Phaeophyscia orbicularis</i> , <i>Physcia adscendens</i> , <i>P. tenella</i>	Beck <i>et al.</i> (1998); Helms <i>et al.</i> (2001).
<i>T. impressa</i>	<i>Dimelaena oreina</i> , <i>Physcia aipola</i> , <i>Physconia perisidiosa</i> , <i>Rinodina capensis</i> , <i>R. milvina</i>	Helms <i>et al.</i> (2001)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Trebouxia impressa</i>	<i>Phaeophyscia orbicularis</i> , <i>Physcia caesia</i> , <i>P. tenella</i> , <i>Physconia distorta</i>	Dahlkild <i>et al.</i> (2001)
<i>T. impressa</i>	<i>Umbilicaria kappenii</i>	Romeike <i>et al.</i> (2002)
<i>T. impressa</i>	<i>Melanelia glabra</i> , <i>Parmelina carporrhizans</i> , <i>P. tiliacea</i> , <i>Phaeophyscia orbicularis</i> , <i>Physcia caesia</i> , <i>P. stellaris</i> , <i>P. tenella</i>	Doering & Piercey-Normore (2009)
<i>T. impressa</i>	<i>Xanthomendoza fallax</i> , <i>X. ulophyllodes</i>	Nyati <i>et al.</i> (2014)
<i>T. impressa</i>	<i>Lecanora rupicola</i>	Blaha <i>et al.</i> (2006)
<i>T. impressa</i>	<i>Lecanora fuscobrunnea</i> , <i>Lecidea andersonii</i> , <i>L. atrobrunnea</i> , <i>L. cancriformis</i> , <i>L. tessellata</i> , <i>Lecidella carpathica</i>	Ruprecht <i>et al.</i> (2012)
<i>T. impressa</i>	<i>Thamnolia vermicularis</i>	Onuț-Brännström <i>et al.</i> (2017)
<i>T. impressa</i> / <i>T. gelatinosa</i>	<i>Cladonia cariosa</i> , <i>C. rei</i> , <i>Diploschistes muscorum</i>	Oszczka <i>et al.</i> (2021)
Clade I (<i>impressa</i> / <i>gelatinosa</i> group)	Species of <i>Melanelixia</i> , <i>Melanohalea</i> , <i>Montanelia</i> , <i>Oropogon</i> , <i>Parmelia</i> , <i>Protoparmeliopsis</i> , <i>Rhizoplaca</i> and <i>Xanthoparmelia</i>	Leavitt <i>et al.</i> (2015)
<i>T. impressa</i> / <i>T. potteri</i>	<i>Ramalina americana</i> , <i>R. sinensis</i>	Francisco de Oliveira <i>et al.</i> (2012)
<i>T. impressa</i> / <i>T. potteri</i>	<i>Pleurosticta koflerae</i>	Voytsekhovich & Beck (2016)
<i>Trebouxia</i> clade including <i>T. impressa</i> , <i>T. flava</i> and <i>T. potteri</i>	<i>Physconia distorta</i> , <i>P. grisea</i>	Wornik & Grube (2010)
<i>T. incrustata</i>	<i>Xanthoria candelaria</i>	Aoki <i>et al.</i> (1998). Antarctica. id: LM, culture.
<i>T. incrustata</i>	<i>Rinodina atrocineria</i>	Helms <i>et al.</i> (2001)
<i>T. incrustata</i>	<i>Acarospora bullata</i> , <i>Lecanora muralis</i> , <i>L. stenotropa</i> , <i>Neofuscelia stygiodes</i> , <i>Protoparmelia badia</i> , <i>Rhizocarpon cf. cyanescens</i> , <i>R. cf. viridiatrum</i>	Beck (2002)
<i>T. incrustata</i>	<i>Lecanora rupicola</i>	Blaha <i>et al.</i> (2006)
<i>T. incrustata</i>	<i>Protoparmelia incrustans</i>	Muggia <i>et al.</i> (2013)
<i>T. incrustata</i>	<i>Protoparmeliopsis muralis</i>	Guzow-Krzemińska (2006)
<i>T. incrustata</i>	<i>Caloplaca crenulatella</i> , <i>C. squamulosa</i> , <i>Circinaria contorta</i> , <i>Lecanora muralis</i> , <i>Staurothele sp.</i> , <i>Xanthoparmelia pulla</i> , <i>X. stenophylla</i>	Voytsekhovich & Beck (2016)
<i>T. irregularis</i>	<i>Diploschistes muscorum</i>	Friedl (1989). Photobiont taken from parasitized <i>Cladonia</i> ; later substituted by <i>T. showmanii</i> . id: LM.
<i>T. jamesii</i>	<i>Candelaria concolor</i> , <i>Candelariella sp.</i>	Beck <i>et al.</i> (1998)
<i>T. jamesii</i>	<i>Tremolecia atrata</i>	Beck & Koop (2001)
<i>T. jamesii</i>	<i>Chaenotheca subroscida</i>	Tibell & Beck (2002)
<i>T. jamesii</i>	<i>Umbilicaria antarctica</i> , <i>U. decussata</i> , <i>U. kappenii</i> , <i>U. umbilicarioides</i>	Romeike <i>et al.</i> (2002)
<i>T. jamesii</i>	<i>Evernia mesomorpha</i> , <i>Melanelia exasperatula</i>	Piercey-Normore (2006, 2009). Multiple clades within.
<i>T. jamesii</i>	<i>Evernia divaricata</i> , <i>E. prunastri</i> , <i>Ramalina fraxinea</i> , <i>R. siliquosa</i>	Tschaikner <i>et al.</i> (2007)
<i>T. jamesii</i>	<i>Ramalina farinacea</i>	Casano <i>et al.</i> (2011); Moya <i>et al.</i> (2017).
<i>T. jamesii</i>	<i>Ramalina fraxinea</i>	Catalá <i>et al.</i> (2016)
<i>T. jamesii</i>	<i>Carbonea vorticosa</i> , <i>Lecanora fuscobrunnea</i> , <i>L. physciella</i> , <i>Lecidea auriculata</i> , <i>L. cancriformis</i> , <i>L. lapicida</i> , <i>L. obluridata</i> , <i>Lecidella greenii</i> , <i>L. siplei</i>	Ruprecht <i>et al.</i> (2012)
<i>T. jamesii</i>	<i>Lecidea cancriformis</i>	Pérez-Ortega <i>et al.</i> (2012). Hap4, well-supported clade.
<i>T. jamesii</i>	<i>Lepraria borealis</i>	Engelen <i>et al.</i> (2010). Maritime Antarctica.

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Trebouxia jamesii</i>	<i>Ramalina menziesii</i>	Werth & Sork (2014). Only 6% of sampled thalli; majority had <i>T. decolorans</i> .
<i>T. jamesii</i>	<i>Candelariella vitellina</i> , <i>Lecanora frustulosa</i> , <i>Lecidea</i> sp., <i>Protoparmelia psarophana</i> , <i>Ramalina calicaris</i> , <i>R. capitata</i> , <i>R. pollinaria</i> , <i>R. pontica</i> , <i>Ramalina</i> sp., <i>Rhizocarpon geographicum</i> , <i>Rhizoplaca</i> sp., <i>Umbilicaria grisea</i>	Voytsekhovich & Beck (2016)
<i>T. jamesii</i>	<i>Protoparmelia badia</i> , <i>P. montagnei</i>	Singh et al. (2017)
<i>T. jamesii</i> clade 'T. lethariae'	<i>Letharia barbata</i> , <i>L. gracilis</i> , <i>L. lucida</i> , <i>L. lupina</i> , <i>L. rugosa</i>	Kroken & Taylor (2000)
<i>T. jamesii</i> clade 'T. vulpinae'	<i>Letharia vulpina</i>	Kroken & Taylor (2000)
<i>T. jamesii</i> clade 'T. vulpinae'	<i>Cetraria aculeata</i>	Fernández-Mendoza et al. (2011); Onuț-Brännström et al. (2018).
<i>T. jamesii</i> and <i>Trebouxia</i> clades A, I, S, G	<i>Thamnia vermicularis</i>	Nelsen & Gargas (2009)
<i>T. jamesii</i> 'group A'	<i>Boreoplaca ultrafrigida</i> , <i>Umbilicaria esculenta</i> , <i>U. muehlenbergii</i>	Cao et al. (2015)
<i>T. jamesii</i> 'group B'	<i>Boreoplaca ultrafrigida</i> , <i>U. muehlenbergii</i>	Cao et al. (2015)
<i>T. jamesii</i> ssp. <i>angustilobata</i>	<i>Acarospora sinopica</i> , <i>Lecidea lapicida</i>	Beck (1999)
<i>T. jamesii</i> ssp. <i>jamesii</i>	<i>Acarospora rugulosa</i> , <i>Bellemeria diamartha</i> , <i>Lecanora polytropica</i> , <i>L. subaurea</i> , <i>Lecidea silacea</i> , <i>Rhizocarpon geographicum</i> , <i>Umbilicaria cylindrica</i>	Beck (1999)
<i>T. potteri</i>	<i>Pertusaria</i> spp.	Takeshita et al. (1989). id: LM, culture.
<i>T. potteri/flava</i>	<i>Lecanora rupicola</i>	Blaha et al. (2006)
<i>T. showmanii</i>	<i>Caloplaca</i> spp.	Watanabe et al. (1997). Maritime lichens. id: LM, culture.
<i>T. showmanii</i>	<i>Diploschistes muscorum</i>	Friedl (1989). id: LM.
<i>T. simplex</i>	<i>Acarospora rugulosa</i> , <i>A. sinopica</i> , <i>A. smaragdula</i> , <i>Bellemeria alpina</i> , <i>B. cinereorufescens</i> , <i>B. subsorediza</i> , <i>Carbonea vorticosa</i> , <i>Lecidea confluens</i> , <i>L. epanora</i> , <i>L. handelii</i> , <i>L. lapicida</i> , <i>L. paupercula</i> , <i>L. polytropica</i> , <i>L. silacea</i> , <i>L. subaurea</i> , <i>L. tessellata</i> , <i>Miriquidica atrofulva</i> , <i>M. garovaglii</i> , <i>Pleopsidium chlorophanum</i> , <i>Porpidia tuberculosa</i> , <i>Rhizocarpon geographicum</i> , <i>R. lecanorinum</i> , <i>R. norvegicum</i> , <i>R. polycarpum</i> , <i>R. ridescens</i> , <i>R. simillimum</i> , <i>Sporastatia testudinea</i> , <i>Tremolecia atrata</i> , <i>Umbilicaria crustulosa</i> , <i>U. cylindrica</i> , <i>U. torrefacta</i>	Beck (2002)
<i>T. simplex</i>	<i>Lecanora rupicola</i>	Blaha et al. (2006)
<i>T. simplex</i>	<i>Lecanora conizaeoides</i>	Hauck et al. (2007)
<i>T. simplex</i>	<i>Evernia mesomorpha</i> , <i>Everniastrum catawbiense</i> , <i>Imshaugia aleurites</i> , <i>Pseudevernia consocians</i>	Piercey-Normore (2009)
<i>T. simplex</i>	<i>Evernia mesomorpha</i> , <i>Imshaugia placorodia</i> , <i>Lecanora conizaeoides</i> , <i>Pseudevernia cladoniae</i> , <i>P. consocians</i>	Doering & Piercey-Normore (2009)
<i>T. simplex</i>	<i>Bryoria americana</i> , <i>B. bicolor</i> , <i>B. capillaris</i> , <i>B. fremontii</i> , <i>B. furcellata</i> , <i>B. glabra</i> , <i>B. implexa</i> , <i>B. lanestris</i> , <i>B. nadvornikiana</i> , <i>B. nitidula</i> , <i>B. simplicior</i> , <i>B. smithii</i> , <i>B. subcana</i> , <i>B. tenuis</i> , <i>Bryoria</i> sp.	Lindgren et al. (2014)
<i>T. simplex</i>	<i>Protoparmelia hypotremella</i> , <i>P. ochrococca</i> , <i>P. oleagina</i>	Singh et al. (2017)
<i>T. simplex</i>	<i>Thamnia vermicularis</i>	Onuț-Brännström et al. (2017)
<i>T. simplex/jamesii</i>	<i>Lasallia pustulata</i> , <i>L. hispanica</i> , <i>Umbilicaria spodochoa</i>	Hestmark et al. (2016); Dal Grande et al. (2018).
Clade S (<i>simplex/letharii/jamesii</i> group)	Species of <i>Cetraria</i> , <i>Letharia</i> , <i>Melanohalea</i> , <i>Motanelia</i> , <i>Rhizoplaca</i> and <i>Xanthoparmelia</i>	Leavitt et al. (2015)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
Clade S (<i>Trebouxia simplex/letharii/jamesii</i> group)	<i>Pseudephebe minuscula</i> , <i>P. pubescens</i>	Garrido-Benavent <i>et al.</i> (2020)
<i>T. simplex/suecica</i> (S clade)	<i>Cetraria aculeata</i> , <i>C. ericetorum</i> , <i>C. islandica</i> , <i>C. muricata</i> , <i>C. sepincola</i> , <i>Flavocetraria cucullata</i> , <i>F. nivalis</i> , <i>Melanelia agnata</i> , <i>M. hepatizon</i> , <i>M. stygia</i> , <i>Tuckermannopsis chlorophylla</i> , <i>Vulpicida pinastris</i>	Xu <i>et al.</i> (2020)
<i>T. simplex</i> s. lat.	<i>Tephromela atra</i> s. str.	Muggia <i>et al.</i> (2008). <i>Tephromela</i> photobionts also found in two other <i>Trebouxia</i> clades.
<i>T. simplex</i> subclades 1 and 2	<i>Cetraria aculeata</i> , <i>C. islandica</i> , <i>Thamnolia subuliformis</i>	Onuț-Brännström <i>et al.</i> (2018)
<i>Trebouxia solaris</i>	<i>Aspicilia cinerea</i> , <i>Caloplaca aractina</i> , <i>Candelariella vittellina</i> , <i>Circinaria contorta</i> , <i>Ramalina capitata</i> , <i>Rhizocarpon geographicum</i>	Voytsekhovich & Beck (2016)
<i>T. aff. solaris</i>	<i>Chrysothrix candelaris</i>	Ertz <i>et al.</i> (2018)
<i>T. aff. solaris</i>	<i>Lecanographa amylacea</i> (' <i>Buellia violaceofusca</i> ')	Ertz <i>et al.</i> (2018). Same mycobiont produces morphologically distinct, sexual thallus with <i>Trentepohlia</i> sp.!
<i>T. suecica</i>	<i>Acarospora sinopica</i> , <i>Candelariella vitellina</i> , <i>Immersaria athrocarpa</i> , <i>Lecanora polytropha</i> , <i>Mirioidica atrofulva</i> , <i>Protoparmelia atriseda</i> , <i>P. badia</i> , <i>Rhizocarpon geographicum</i> , <i>Tremolecia atrata</i> , <i>Umbilicaria torrefacta</i>	Beck (2002)
<i>T. suecica</i>	<i>Protoparmelia badia</i> , <i>P. memnonia</i>	Singh <i>et al.</i> (2017)
<i>T. suecica</i>	<i>Pseudephebe</i> sp.	Garrido-Benavent <i>et al.</i> (2020)
<i>T. vaga</i>	<i>Candelariella vitellina</i> , <i>Circinaria contorta</i> , <i>Diploschistes diacapsis</i> , <i>Porpidia crustulata</i>	Voytsekhovich & Beck (2016)
<i>T. vaga</i>	<i>Thamnolia vermicularis</i>	Onuț-Brännström <i>et al.</i> (2017)
Near <i>T. vaga</i>	<i>Caloplaca tubelliana</i> , <i>Dirina massiliensis</i> [Arthoniomycetes!], <i>Lecanora albescens</i>	Voytsekhovich & Beck (2016)
<i>Trebouxia</i> clade IV (including <i>T. corticola</i> , <i>T. galapagensis</i> , <i>T. higginsiae</i> and <i>T. usneae</i>)	Tropical <i>Ramalina anceps</i> , <i>R. complanata</i> , <i>R. dendroides</i> , <i>R. gracilis</i> , <i>R. peruviana</i> , <i>R. sorediosa</i> , <i>R. sprengelii</i>	Cordeiro <i>et al.</i> (2005)
<i>Trebouxia</i> A 02	<i>Austrolechia</i> sp., <i>Buellia frigida</i> , <i>Carbonea vorticosa</i> , <i>Huea</i> sp., <i>Lecidea cancriformis</i> , <i>L. polypycnidophora</i> , <i>Lecidella greenii</i> , <i>Rhizoplaca macleanii</i>	Wagner <i>et al.</i> (2020). Antarctic Dry Valleys.
<i>Trebouxia</i> sp. clade 1	<i>Fulgensia bracteata</i> , <i>F. fulgens</i> , <i>Toninia sedifolia</i>	Schaper & Ott (2003). Götland, Sweden.
<i>Trebouxia</i> sp. 'D11'	<i>Buellia frigida</i> , endolithic <i>Lecidea</i> sp., <i>Umbilicaria aprina</i>	de los Ríos <i>et al.</i> (2005). Continental Antarctic.
<i>Trebouxia</i> TR9	<i>Ramalina farinacea</i>	Casano <i>et al.</i> (2011); Moya <i>et al.</i> (2017).
<i>Trebouxia</i> sp. 'URa1'	<i>Carbonea vorticosa</i> , <i>Lecidea cancriformis</i>	Ruprecht <i>et al.</i> (2012)
<i>Trebouxia</i> sp. 'URa2'	<i>Carbonea vorticosa</i> , <i>Lecidea andersonii</i> , <i>L. atrobrunnea</i> , <i>L. cancriformis</i> , <i>L. fuscoatra</i> , <i>L. lapicida</i> , <i>Lecidea</i> sp., <i>Lecidella greenii</i> , <i>L. siplei</i> , <i>Rhizoplaca macleanii</i>	Ruprecht <i>et al.</i> (2012)
<i>Trebouxia</i> sp. 'URa3'	<i>Carbonea vorticosa</i> , <i>Lecanora physciella</i> , <i>Lecidea andersonii</i> , <i>L. cancriformis</i> , <i>L. lapicida</i> , <i>L. polypycnidophora</i> , <i>Lecidella greenii</i> , <i>L. patavina</i> , <i>L. siplei</i> , <i>Rhizoplaca macleanii</i>	Ruprecht <i>et al.</i> (2012)
<i>Trebouxia</i> sp. 'URa4'	<i>Lecidea andersonii</i> , <i>Lecidella stigmata</i>	Ruprecht <i>et al.</i> (2012)
<i>Trebouxia</i> sp.	<i>Buellia papillata</i> , <i>Caloplaca johnstonii</i> , <i>C. lewis-smithii</i> , <i>Candelariella flava</i> , <i>Lecidella pataviana</i> , <i>Lepraria cacuminum</i> , <i>Pseudephebe minuscula</i> , <i>Psoroma</i> sp., <i>Tephromela atra</i> , <i>T. disciformis</i> , <i>Usnea lambii</i> , <i>Xanthoria elegans</i>	Engelen <i>et al.</i> (2016). Maritime Antarctic.
<i>Trebouxia</i> sp.	<i>Bagliettoa marmorea</i>	Thüs <i>et al.</i> (2011)
<i>Trebouxia</i> sp.	<i>Lasallia pustulata</i>	Sadowksa-Deś <i>et al.</i> (2014)
<i>Trebouxia</i> sp.	<i>Cetraria aculeata</i>	Lutsak <i>et al.</i> (2016)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Trebouxia</i> sp.	<i>Austrolechia</i> spp., <i>Buellia frigida</i> , <i>Caloplaca</i> cf. <i>citrina</i> , <i>C. sublobata</i> , <i>Caloplaca</i> sp., <i>Carbonea vorticosa</i> , <i>Huea</i> sp., <i>Lecanora cancriformis</i> , <i>L. flotowiana</i> , <i>L. fuscobrunnea</i> , <i>Lecidella greenii</i> , <i>Rhizoplaca macleanii</i> , <i>Rhizoplaca</i> sp., <i>Umbilicaria aprina</i>	Pérez-Ortega <i>et al.</i> (2012). Crustose lichens of McMurdo Dry Valleys, probably representing at least two undescribed clades.
<i>Trebouxia</i> sp.	<i>Circinaria</i> spp.	Molins <i>et al.</i> (2018)
<i>Trebouxia</i> sp. B	<i>Boreoplaca ultrafrigida</i> , <i>Parmelia omphalodes</i>	Cao <i>et al.</i> (2015)
<i>Trebouxia</i> lineages lacking formal recognition	<i>Rhizoplaca</i> spp., <i>Protoparmeliopsis</i> spp.	Leavitt <i>et al.</i> (2016). Some 15 spp., collected worldwide.
<i>Vulcanochloris</i> sp.	<i>Stereocaulon vesuvianum</i>	Vančurová <i>et al.</i> (2015). From La Palma.
<i>Vulcanochloris</i> sp.	<i>Ramalina farinacea</i>	Moya <i>et al.</i> (2017). Minority photobiont.
<i>Vulcanochloris</i> sp.	<i>Stereocaulon pityrizans</i>	Kosecka <i>et al.</i> (2021)
Class <i>Trebouxiophyceae</i>, Order <i>Prasiolales</i>, <i>Elliptochloris</i> clade		
(<i>Coccomyxa greatwallensis</i>)	Isolated from <i>Psoroma hypnorum</i>	Cao <i>et al.</i> (2018). Symbiotic status not established.
<i>C. icmadophilae</i>	<i>Dibaeis baeomyces</i>	Beck (2002)
<i>C. solorinae</i> var. <i>croceae</i> , var. <i>saccatae</i> and <i>C. peltigerae</i>	<i>Peltigera variolosa</i> , <i>Solorina bispora</i> , <i>S. crocea</i> , <i>S. saccata</i>	Malavasi <i>et al.</i> (2016)
<i>C. subellipsoidea</i>	<i>Lichenomphalia ericetorum</i> , <i>L. grisella</i> , <i>L. hudsoniana</i> , <i>L. luteovitellina</i> , <i>L. velutina</i> , <i>Lichenomphalia</i> sp.	Zoller & Lutzoni (2003)
<i>C. subellipsoidea</i>	<i>Lichenomphalia hudsoniana</i> , <i>L. meridionalis</i> , <i>L. umbellifera</i>	Masumoto <i>et al.</i> (2019); Gasulla <i>et al.</i> (2020); Masumoto (2020).
<i>Coccomyxa</i> sp.	<i>Dacampia hookeri</i>	Henssen (1995). id: LM.
<i>Coccomyxa</i> sp.	<i>Nephroma arcticum</i> , <i>N. expallidum</i> , <i>Peltigera aphthosa</i> , <i>P. britannica</i> , <i>P. leucophebia</i>	Lohtander <i>et al.</i> (2003)
<i>Coccomyxa</i> sp.	<i>Peltigera variolosa</i> , <i>Solorina bispora</i> , <i>S. crocea</i> , <i>S. saccata</i>	Darienko <i>et al.</i> (2015). Photobionts not monophyletic, related to free-living <i>C. subellipsoidea</i> .
<i>Coccomyxa</i> sp.	<i>Micarea byssacea</i> , <i>M. denigrata</i> , <i>M. leprosula</i> , <i>M. lignaria</i> , <i>M. micrococca</i> , <i>M. nitschkeana</i> , <i>M. peliocarpa</i> , <i>M. prasina</i> (<i>Pilocarpaceae</i>)	Yahr <i>et al.</i> (2015)
<i>Coccomyxa</i> sp.	<i>Icmadophila aversa</i> , <i>I. ericetorum</i> , <i>I. japonica</i> , <i>I. splachnirima</i>	Ludwig (2015)
<i>Coccomyxa</i> sp.	<i>Lichenomphalia oreades</i>	Lücking <i>et al.</i> (2017b)
<i>Coccomyxa</i> sp.	<i>Sticta laciniosa</i> from Cuba	Lindgren <i>et al.</i> (2020)
Related to <i>Coccomyxa</i> and <i>Pseudococcomyxa</i>	<i>Schizoxylon albescens</i> (<i>Stictidaceae</i>)	Muggia <i>et al.</i> (2011)
<i>Elliptochloris bilobata</i>	<i>Fuscidea cyathea</i> var. <i>japonica</i>	Watanabe <i>et al.</i> (1997). Maritime. id: LM, culture.
<i>E. bilobata</i>	Species of <i>Buellia</i> and <i>Leproloma</i>	Aoki <i>et al.</i> (1998). Antarctic. id: LM, culture.
<i>E. bilobata</i>	<i>Baeomyces rufus</i>	Trembley <i>et al.</i> (2002b)
<i>E. bilobata</i>	<i>Verrucaria sublobulata</i>	Thüs <i>et al.</i> (2011)
<i>E. bilobata</i>	<i>Micarea misella</i> , <i>M. prasina</i>	Voytsekhovich <i>et al.</i> (2011). id: LM, culture.
<i>E. bilobata</i>	<i>Micarea prasina</i>	Yahr <i>et al.</i> (2015)
<i>E. bilobata</i>	<i>Catolechia wahlenbergii</i>	Darienko <i>et al.</i> (2016)
<i>E. parasphaerica</i> ad int.	<i>Multiclavula petricola</i> , <i>M. vernalis</i>	Reported by H. Masumoto & S. Handa in Masumoto (2020).
<i>E. perforata</i>	<i>Micarea prasina</i>	Darienko <i>et al.</i> (2016)
<i>E. reniformis</i>	<i>Fuscidea cyathoides</i> var. <i>japonica</i> , <i>Leproloma</i> sp., <i>Ochrolechia parellula</i>	Watanabe <i>et al.</i> (1997); Darienko <i>et al.</i> (2016). Maritime. id: LM, culture.
<i>E. reniformis</i>	<i>Micarea peliocarpa</i>	Voytsekhovich <i>et al.</i> (2011). id: LM, culture.
<i>E. subsphaerica</i>	<i>Aspicilia</i> sp., <i>Fuscidea cyathoides</i> var. <i>japonica</i>	Watanabe <i>et al.</i> (1997). Maritime. id: LM, culture. (= <i>Chlorella reisiigii</i> Watanabe; Darienko <i>et al.</i> 2016)
<i>E. subsphaerica</i>	<i>Micarea melanobola</i> , <i>M. prasina</i> , <i>M. subnigrata</i>	Voytsekhovich <i>et al.</i> (2011). id: LM, culture.

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Elliptochloris subsphaerica</i>	<i>Multiclavula mucida</i>	Masumoto (2020)
<i>Elliptochloris</i> sp.	<i>Stictis urceolatum</i>	Thüs et al. (2011)
<i>Elliptochloris</i> sp.	<i>Catillaria chalybeia</i>	Dal Grande et al. (2014b)
<i>Elliptochloris</i> sp.	<i>Ramalina farinacea</i>	Moya et al. (2017). Minority photobiont.
<i>Elliptochloris</i> sp.	<i>Bryoclavula phycophila</i>	Masumoto (2020)
<i>Elliptochloris</i> sp.	<i>Sticta filix</i> , <i>S. menziesii</i> , <i>S. stipitata</i> , <i>S. subcaperata</i>	Lindgren et al. (2020)
<i>Pseudococcomyxa simplex</i>	<i>Leproloma</i> sp.	Watanabe et al. (1997). Maritime. id: LM, culture.
<i>Pseudococcomyxa</i> sp.	<i>Micarea prasina</i>	Voytsekhovich et al. (2011). id: LM, culture.
Class Trebouxiophyceae, Order Watanabeales		
<i>Chloridium angustoellipsoideum</i>	<i>Stereocaulon nanodes</i> , <i>S. spathuliferum</i>	Vančurová (2012); Vančurová et al. (2018). Darienko et al. (2018): <i>C. angustoellipsoideum</i> now <i>C. lichenum</i> .
<i>C. ellipsoideum</i>	<i>Trapelia obtegens</i>	Beck (2002)
<i>C. ellipsoideum</i>	<i>Stereocaulon nanodes</i> , <i>S. pileatum</i> , <i>S. vesuvianum</i> , <i>S. spathuliferum</i> , <i>Stereocaulon</i> sp.	Vančurová (2012); Vančurová et al. (2018). <i>Stereocaulon vesuvianum</i> also hosts <i>Asterochloris</i> and <i>Vulcanochloris</i> .
Near <i>C. ellipsoideum</i>	<i>Trapeliopsis flexuosa</i> , <i>T. granulosa</i>	Schmitt & Lumbsch (2001). Amplified unintentionally with non-specific primers. 'Chlorella' <i>ellipsoidea</i> sequence X63520. Known photobionts of other <i>Trapeliaceae</i> are in <i>Prasiolales</i> .
<i>C. saccharophilum</i>	<i>Trapelia coarctata</i>	Tschermak-Woess (1988b)
<i>C. saccharophilum</i>	<i>Lecidea inops</i> , <i>Psilolechia leprosa</i> , <i>Stereocaulon nanodes</i>	Beck (2002)
<i>C. saccharophilum</i>	<i>Psora decipiens</i>	Ruprecht et al. (2014). One record; all other thalli had <i>Asterochloris</i> or <i>Trebouxia</i> . Contrast Williams et al. (2017).
<i>C. viscosum</i>	<i>Bacidina</i> [Woessia] <i>fusarioides</i>	Tschermak-Woess (1988b); Darienko et al. (2018).
<i>Chloridium</i> sp.	<i>Verrucaria praetermissa</i>	Thüs et al. (2011). Possibly an accessory alga rather than main photobiont.
<i>Chloridium</i> sp.	<i>Verrucaria nigrescens</i> , <i>Verrucaria</i> sp.	Voytsekhovich & Beck (2016). id: LM.
<i>Chloridium</i> sp.	Species of <i>Gomphillus</i> and <i>Gyalidea</i>	Sanders et al. (2016)
<i>Chloridium</i> sp.	<i>Sticta latifrons</i> , <i>S. subcaperata</i>	Lindgren et al. (2020)
Near <i>Chloridium</i> and <i>Parachloridium</i>	<i>Bapalmua lineata</i>	Sanders et al. (2016)
<i>Heveochlorella</i> sp. (<i>Jaagichlorella</i> Darienko & Pröschold 2019)	<i>Pseudocyphellaria</i> sp., <i>Sticta</i> sp.	Dal Grande et al. (2014b). Three specimens; authors confident alga is photobiont.
<i>Heveochlorella</i> sp.	<i>Aspidothelium cinerascens</i> , <i>A. fugiens</i> , <i>A. geminiparum</i> (<i>Thelenellaceae</i>); <i>Aderkomyces heterellus</i> , <i>Echinoplaca</i> spp., <i>Gyalectidium appendiculatum</i> , <i>G. floridense</i> , <i>G. imperfectum</i> , <i>G. paolae</i> , <i>G. ulloae</i> , <i>Gyalideopsis sessilis</i> , <i>Gyalideopsis</i> sp. (<i>Gomphillaceae</i>); <i>Byssoloma discordans</i> , <i>Calopadia fusca</i> , <i>C. perpallida</i> , <i>C. puiggarii</i> (<i>Pilocarpaceae</i>)	Sanders et al. (2016). Mainly foliicolous.
<i>Heveochlorella</i> sp.	<i>Sticta caperata</i> , <i>S. dichotoma</i> , <i>S. latifrons</i> , <i>S. squamata</i> , <i>S. subcaperata</i> , <i>S. variabilis</i> , <i>Sticta</i> spp.	Lindgren et al. (2020)
[unidentified, falling 'near' <i>Heterochlorella</i>]	<i>Psoroglaena epiphylla</i>	Nyati et al. (2007); Thüs et al. (2011).
Class Trebouxiophyceae, incertae sedis		
<i>Apatococcus lobatus</i>	<i>Caloplaca</i> sp., <i>Fuscidea cyathoides</i> var. <i>japonica</i> , <i>Lecanora subimergens</i> , <i>Pyrenopsis</i> sp., <i>Verrucaria</i> sp.	Watanabe et al. (1997). Maritime. id: LM.
<i>A. lobatus</i>	<i>Scoliosporum umbrinum</i>	Beck (2002)
<i>A. fuscidae</i>	<i>Fuscidea arboricola</i> , <i>F. cyathoides</i> , <i>F. kochiana</i> , <i>F. pusilla</i>	Zahradníková et al. (2017)
<i>Apatococcus</i> sp.	<i>Fuscidea lightfootii</i>	Zahradníková et al. (2017)

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
<i>Coccobotrys verrucariae</i>	<i>Verrucaria nigrescens</i>	Chodat (1913); Vischer (1960).
<i>Coccobotrys</i> sp. (?)	<i>Botryolepraria lesdainii</i>	Canals et al. (1997). id: LM, culture.
<i>Dictyochloropsis splendida</i>	<i>Phlyctis argena</i>	Tschermak-Woess (1995). id: LM.
<i>D. symbiontica</i>	<i>Sphaerophorus diplotypus</i> , <i>S. fragilis</i> , <i>S. melanocarpus</i> , <i>S. melophorus</i>	Ihda et al. (1997). id: LM, culture.
<i>D. symbiontica</i>	Species of <i>Caloplaca</i> and <i>Leproloma</i>	Watanabe et al. (1997). Maritime. id: LM.
<i>D. symbiontica</i>	<i>Lecania naegelii</i>	Beck et al. (1998)
<i>Dictyochloropsis</i> clade 2	<i>Brigantiaea ferruginea</i> , <i>B. leucoxantha</i> (<i>Brigantiaeeaceae</i>); <i>Crocodia aurata</i> , <i>Lobaria oregana</i> , <i>L. patinifera</i> , <i>L. pulmonaria</i> , <i>Lobariella crenulata</i> , <i>L. pallidocrenulata</i> , <i>Lobariella</i> sp., <i>Pseudocyphellaria fimbriata</i> , <i>P. homeophylla</i> , <i>P. lindsayi</i> , <i>P. lividofusca</i> , <i>P. multifida</i> , <i>Sticta canariensis</i> , <i>S. latifrons</i> , <i>S. pulmonariodes</i> , <i>S. subcaperata</i> , <i>Sticta</i> spp. (<i>Lobariaceae</i>); <i>Megalospora sulphurata</i> (<i>Megalosporaceae</i>); <i>Chaenothecopsis consociata</i> (<i>Mycocaliciaceae</i>); <i>Phlyctis argena</i> (<i>Phlyctidaceae</i>); <i>Biatora</i> sp. (<i>Ramalinaceae</i>); <i>Chaenotheca brunneola</i> (<i>Coniocybomycetes</i>)	Dal Grande et al. (2014b). <i>Dictyochloropsis</i> . clade 2 = <i>Symbiochloris</i> ; Škaloud et al. (2016).
<i>Leptosira terrestris</i>	<i>Vezeada aestivalis</i>	Tschermak-Woess & Poelt (1976). id: LM, culture.
<i>L. thombii</i>	<i>Thrombium epigaeum</i>	Tschermak-Woess (1953). id: LM, culture.
<i>Neocystis</i> sp.	<i>Micarea misella</i>	Voytsekhovich et al. (2011). Reported as an 'additional photobiont' to <i>Elliptochloris</i> . id: LM, culture.
<i>Symbiochloris</i> sp.	<i>Brigantiaea ferruginea</i> , <i>B. leucoxantha</i> (<i>Brigantiaeeaceae</i>); <i>Crocodia aurata</i> , <i>Dendriscosticta platyphylla</i> , <i>D. wrightii</i> ; <i>Lobaria macaronesica</i> , <i>L. oregana</i> , <i>L. patinifera</i> , <i>L. pulmonaria</i> ; <i>Lobariella crenulata</i> , <i>L. pallidocrenulata</i> , <i>Lobariella</i> sp., <i>Pseudocyphellaria</i> sp., <i>Ricasolia amplissima</i> , <i>Sticta canariensis</i> , <i>S. neopulmonaria</i> (<i>Lobariaceae</i>); <i>Megalospora sulphurata</i> (<i>Megalosporaceae</i>); <i>Chaenothecopsis consociata</i> (<i>Mycocaliciaceae</i>); <i>Phlyctis argena</i> (<i>Phlyctidaceae</i>); <i>Chaenotheca brunneola</i> (<i>Coniocybomycetes</i>).	Škaloud et al. (2016). All known photobionts previously included in <i>Dictyochloropsis</i> s. lat.
<i>Symbiochloris</i> sp.	<i>Sticta ainoae</i> , <i>S. cinereoglauca</i> , <i>S. granatensis</i> , <i>S. laciniata</i> , <i>S. laciniosa</i> , <i>S. puracensis</i> , <i>Sticta</i> spp.	Lindgren et al. (2020)
Class <i>Ulvophyceae</i>, Order <i>Trentepohliales</i>		
<i>Cephaleuros</i> sp.	<i>Strigula smaragdula</i>	Green (2012)
<i>Cephaleuros</i> sp.	<i>Strigula smaragdula</i>	Jiang et al. (2020)
<i>Phycopeltis</i> sp.	<i>Tenuitholiascus porinoides</i>	Jiang et al. (2020)
<i>Trentepohlia lagenifera</i>	<i>Graphina cleistoblephara</i> , <i>G. inabensis</i> , <i>G. intortura</i> , <i>G. mendax</i> , <i>G. undulata</i> , <i>Graphis aperiens</i> , <i>G. aphanes</i> , <i>G. batanensis</i> , <i>G. cognata</i> , <i>G. connectans</i> , <i>G. dupaxana</i> , <i>G. handelii</i> , <i>G. intricata</i> , <i>G. proserpens</i> , <i>G. rikuzensis</i> , <i>G. rufia</i> , <i>G. scripta</i> , <i>G. subdura</i> , <i>G. subtropica</i> , <i>G. subvirginea</i> , <i>G. tenella</i> , <i>Phaeographina chlorocarpoides</i> , <i>P. endofusca</i> , <i>Phaeographis asteriformis</i> , <i>P. exaltata</i> , <i>P. pruinosa</i>	Nakano (1988). id: LM, culture.
<i>T. lagenifera</i>	<i>Pyrenula japonica</i>	Nakano & Ihda (1996). id: LM, culture.
<i>Trentepohlia</i>/'Printzina'	<i>Acanthothecis</i> sp., <i>Anthracotheccium</i> sp., <i>Astrothelium cinnamomeum</i> , <i>A. galbineum</i> , <i>A. leucoconicum</i> , <i>A. versicolor</i> , <i>Astrothelium</i> sp., <i>Coenogonium linkii</i> , <i>Cryptothecia assimilis</i> , <i>Cryptothelium pulchrum</i> , <i>Cryptothelium</i> sp., <i>Dendrographa alectoroides</i> , <i>Dichosporidium boschianum</i> , <i>Graphis scripta</i> , <i>Graphis</i> sp., <i>Laurera megasperma</i> , <i>Myriotrema peninsulae</i> , <i>Porina distans</i> , <i>P. dolichophora</i> , <i>P. imitatrix</i> , <i>P. nucula</i> , <i>P. tetracerae</i> , <i>Porina</i> spp., <i>Racodium rupestre</i> , <i>Thalloloma hypoleptum</i> , <i>Thelotrema pachysporum</i> , <i>Trypethelium aenum</i> , <i>T. nitidiusculum</i> , <i>Trypethelium</i> spp.	Nelsen et al. (2011). 'Printzina' clades occur scattered throughout <i>Trentepohlia</i> . Mycobionts represent <i>Arthoniomycetes</i> , <i>Dothideomycetes</i> and <i>Lecanoromycetes</i> .

(Continued)

Table 1. (Continued)


Alga	Fungal symbionts	References & comments
<i>Trentepohlia</i> /'Printzina'	<i>Astrothelium variolosum</i> , <i>Graphis endoxantha</i> , <i>G. pulverulenta</i> , <i>G. scripta</i> , <i>Herpothallon rubrocinctum</i> , <i>Phaeographina</i> sp., <i>Phaeographis inusta</i> , <i>P. punctiformis</i> , <i>Pyrenula confinis</i> , <i>P. cruenta</i> , <i>P. pseudobufonia</i> , <i>P. sexlocularis</i> , <i>Strigula subtilissima</i> , <i>Trypethelium virens</i>	Green (2012). Photobiont of one <i>Strigula</i> sp. included.
<i>Trentepohlia</i> 'Printzina'	<i>Acrocordia gemmata</i> , <i>Arthonia cinnabarina</i> , <i>A. radiata</i> , <i>Arthothelium ruanum</i> , <i>Cystocoleus ebeneus</i> , <i>Dimerella pineti</i> , <i>Graphis propinqua</i> , <i>G. scripta</i> , <i>G. submarginata</i> , <i>Gyalecta jenensis</i> , <i>Mycoporum sparsellum</i> , <i>Opegrapha atra</i> , <i>Pyrenula laevigata</i> , <i>Roccella decipiens</i> , <i>R. galapagoensis</i> , <i>R. linearis</i> , <i>R. lirellina</i> , <i>R. maderensis</i> , <i>R. cf. montagnei</i> , <i>R. phycopsis</i> , <i>R. tinctoria</i> , <i>R. tuberculata</i> ; <i>Thelotrema lepadinum</i>	Hametner et al. (2014a, b)
<i>Trentepohlia</i> sp.	<i>Lecanographa amylicea</i>	Ertz et al. (2018). Mycobiont also forms heteromorphic, asexual thalli with <i>Trebouxia</i> sp.!
<i>Trentepohlia</i> sp.	<i>Dichosporidium</i> sp., <i>Diorygma antillarum</i> , <i>D. confluens</i> , <i>D. nigrocinctum</i> , <i>Diorygma</i> sp., <i>Herpothallon echinatum</i> , <i>H. rubroechinatum</i> , <i>Herpothallon</i> sp., <i>Ocellularia soreddiigera</i> , <i>Ocellularia</i> spp., <i>Porina</i> sp., <i>Sagenidiopsis isidiata</i> , <i>Sagenidiopsis</i> sp., <i>Synccesia farinacea</i>	Kosecka et al. (2020)
Class Ulvophyceae, Order Ulvales		
<i>Blidingia minima</i>	<i>Turgidosculum ulvae</i>	Pérez-Ortega et al. (2018). Form 'borderline lichen' but mycobiont belongs to lichen-forming clade.
(<i>Dilabifilum</i>)	<i>Hydropunctaria adriatica</i>	Tschermak-Woess (1976). id: LM, culture.
(<i>Dilabifilum</i>)	<i>Hydropunctaria maura</i> , <i>H. rheitrophila</i> , <i>Wahlenbergiella striatula</i> , <i>Verrucaria aquatilis</i>	Thüs et al. (2011)
(<i>Dilabifilum</i>)	<i>Verrucaria glaucina</i>	Voytsekhovich & Beck (2016). id: LM.
<i>Halofilum ramosum</i>	<i>Hydropunctaria amphibia</i> , <i>H. maura</i> , <i>Wahlenbergiella striatula</i>	Darienko & Pröschold (2017); Gasulla et al. (2019).
<i>Lithotrichon pulchrum</i>	<i>Hydropunctaria rheitrophila</i>	Darienko & Pröschold (2017). Submerged lichen.
<i>Paulbroadya petersii</i>	<i>Wahlenbergiella mucosa</i>	Darienko & Pröschold (2017)
<i>Pseudendozonium arthopyreniae</i>	Species of <i>Caloplaca</i> , <i>Rinodina</i> , <i>Thelidium</i> and <i>Verrucaria</i>	Watanabe et al. (1997). Maritime lichens. According to Thüs et al. (2011), maritime specimens of <i>Caloplaca</i> etc. may overgrow <i>Verrucariaceae</i> containing <i>Dilabifilum</i> (<i>Pseudendozonium</i>). id: LM, culture.
<i>P. incrustans</i>	<i>Verrucaria aquatilis</i>	Tschermak-Woess (1970). <i>Dilabifilum</i> , reclassified in Darienko & Pröschold (2017).
<i>P. commune</i>	<i>Hydropunctaria maura</i>	<i>Dilabifilum</i> , reclassified in Darienko & Pröschold (2017).
Class Chlorophyceae		
<i>Bracteacoccus</i> sp.	' <i>Multiclavula clara</i> ' [= <i>Sulzbacheromyces sinensis</i>]	Takeshita et al. (2010). id: LM, culture.
<i>Bracteacoccus</i> sp.	<i>Sulzbacheromyces caatingae</i>	Hodkinson et al. (2014). <i>rbcL</i> sequences.
<i>Bracteacoccus</i> sp.	<i>Sulzbacheromyces sinensis</i>	Masumoto (2020). ITS and <i>rbcL</i> sequences.
<i>Chlamydomonas</i> sp.	<i>Pyronema</i>	Skuja (1943). Association doubtfully lichenic.
<i>Chlorosarcinopsis minor</i>	<i>Lecidea lapicida</i> , <i>L. plana</i>	Plesl (1963). Genus polyphyletic (Neustupa 2015).
<i>Gloeocystis</i> sp.	<i>Bryophagus gloeocapsa</i> , <i>Epigloea bactrospora</i>	Tschermak-Woess (1988a). <i>Gloeocystis</i> s. lat. highly polyphyletic (Neustupa 2015).
<i>Radiococcus signiensis</i>	<i>Placynthiella icmalea</i> , <i>P. uliginosa</i>	Voytsekhovich et al. (2011). Other photobionts found in lesser abundance within same algal layer. id: LM, culture.
<i>Trochiscia</i> sp.	<i>Polyblastia amota</i> , <i>P. hyperborea</i>	Tschermak (1941b); Ahmadjian (1967); but Ettl & Gärtner (2014) consider ID questionable.
Domain Eukarya, Eukaryote supergroup Archaeplastida, Kingdom Plantae, Division Streptophyta (Charophytes)		
<i>Interfilum</i> sp.	<i>Micarea prasina</i> ; <i>Placynthiella icmalea</i> , <i>P. uliginosa</i>	Voytsekhovich et al. (2011). Reported as secondary (less abundant) photobiont within algal layer. id: LM, culture.

(Continued)

Table 1. (Continued)

Alga	Fungal symbionts	References & comments
Domain Eukarya, Eukaryote supergroup: SAR (Stramenopila Alveolata Rhizaria), Kingdom Stramenopila (Heterokontae)		
Class Phaeophyceae		
<i>Petroderma maculiforme</i>	<i>Wahlenbergiella tavaresiae</i>	Moe (1997); Peters & Moe (2001); Gueidan <i>et al.</i> (2011).
Class Xanthophyceae		
<i>Heterococcus</i> sp.	<i>Verrucaria funckii</i> , <i>V. laevata</i>	Zeitler (1954); Tschermak-Woess (1988a).
<i>Heterococcus</i> sp.	<i>Hydropunctaria maura</i>	Parra & Redón (1977). id: LM.
<i>Heterococcus</i> sp.	<i>Hydropunctaria rheitrophila</i> ; <i>Verrucaria funckii</i> , <i>V. hydrela</i>	Thüs <i>et al.</i> (2011)
<i>Xanthonema</i> sp.	<i>Staurothele clopimoides</i>	Pereira Riquelme (1992). id: LM.

Acknowledgements. The manuscript benefited from critical review by Dr Richard L. Moe and two anonymous referees, and the expert scrutiny of the journal production staff. Funding for colour in print was kindly provided by the College of Arts and Sciences and the Department of Biological Sciences, Florida Gulf Coast University.

Author ORCID.  William B. Sanders, 0000-0001-9572-4244.

References

- Adams DG and Duggan PS (2002) Cyanobacteria-bryophyte symbioses. *Journal of Experimental Botany* **59**, 1047–1058.
- Adams DG, Duggan PS and Jackson O (2012) Cyanobacterial symbiosis. In Whitton BA (ed.), *Ecology of Cyanobacteria II: Their Diversity in Space and Time*. Heidelberg: Springer, pp. 593–647.
- Ahmadjian V (1960) Some new and interesting species of *Trebouxia*, a lichenized alga. *American Journal of Botany* **47**, 677–683.
- Ahmadjian V (1962) Investigations on lichen synthesis. *American Journal of Botany* **49**, 277–283.
- Ahmadjian V (1967) A guide to the algae occurring as lichen symbionts: isolation, culture, cultural physiology, and identification. *Phycologia* **6**, 127–160.
- Ahmadjian V (1982) Holobionts have more parts. *International Association for Lichenology Newsletter* **15**, 19.
- Ahmadjian V (1987) Coevolution in lichens. *Annals of the New York Academy of Sciences* **503**, 307–315.
- Ahmadjian V (1988) The lichen alga *Trebouxia*: does it occur free-living? *Plant Systematics and Evolution* **158**, 243–247.
- Ahmadjian V (1993) *The Lichen Symbiosis*. New York: John Wiley and Sons.
- Ahmadjian V (1995) Lichens are more important than you think. *BioScience* **45**, 124.
- Ahmadjian V (2001) *Trebouxia*: reflections on a perplexing and controversial lichen photobiont. In Seckbach J (ed.), *Symbiosis: Mechanisms and Model Systems*. Dordrecht: Kluwer Academic Publishers, pp. 373–383.
- Ahmadjian V (2002) Lingering lichen myths are hard to dispel. *ISS Symbiosis International* **2**, 1–2.
- Ahmadjian V and Heikkilä H (1970) The culture and synthesis of *Endocarpon pusillum* and *Staurothele clopima*. *Lichenologist* **4**, 259–267.
- Ahmadjian A and Jacobs JB (1981) Relationship between fungus and alga in the lichen *Cladonia cristatella* Tuck. *Nature* **289**, 169–172.
- Ahmadjian A, Russell LA and Hildreth KC (1980) Artificial reestablishment of lichens. I. Morphological interactions between the phycobionts of different lichens and the mycobionts *Cladonia cristatella* and *Lecanora chrysoleuca*. *Mycologia* **72**, 73–89.
- Aoki M, Nakano T, Kanda H and Deguchi H (1998) Photobionts isolated from Antarctic lichens. *Journal of Marine Biotechnology* **6**, 39–43.
- Archibald PA (1975) *Trebouxia* de Puymaly (*Chlorophyceae*, *Chlorococcales*) and *Pseudotrebouxia* gen. nov. (*Chlorophyceae*, *Chlorosarcinales*). *Phycologia* **14**, 125–137.
- Armaleo D and Clerc P (1991) Lichen chimeras: DNA analysis suggests that one fungus forms two morphotypes. *Experimental Mycology* **15**, 1–10.
- Armaleo D, Müller O, Lutzoni F, Andrésón ÓS, Blanc G, Bode HB, Collart FR, Dal Grande F, Dietrich F, Grigoriev IV, *et al.* (2019) The lichen symbiosis re-viewed through the genomes of *Cladonia grayi* and its algal partner *Asterochloris glomerata*. *BMC Genomics* **20**, 605.
- Ascaso C, Valladares F and de los Rios A (1995) New ultrastructural aspect of pyrenoids of the lichen photobiont *Trebouxia* (*Microthamniales*, *Chlorophyta*). *Journal of Phycology* **31**, 114–119.
- Asplund J and Wardle DA (2013) The impact of secondary compounds and functional characteristics on lichen palatability and decomposition. *Journal of Ecology* **101**, 689–700.
- Báčkor M, Peksa O, Škaloud P and Báčkorová M (2010) Photobiont diversity in lichens from metal-rich substrata based on ITS rDNA sequences. *Ecotoxicology and Environmental Safety* **73**, 603–612.
- Bates ST, Cropsey GWG, Caporaso JG, Knight R and Fierer N (2011) Bacterial communities associated with the lichen symbiosis. *Applied and Environmental Microbiology* **77**, 1309–1314.
- Beck A (1999) Photobiont inventory of a lichen community growing on heavy-metal-rich rock. *Lichenologist* **31**, 501–510.
- Beck A (2002) *Selektivität der symbionten schwermetalltoleranter flechten*. Ph.D. thesis, Ludwig Maximilian University of Munich.
- Beck A and Koop H-U (2001) Analysis of the photobiont population in lichens using a single-cell manipulator. *Symbiosis* **31**, 57–67.
- Beck A and Mayr C (2012) Nitrogen and carbon isotope variability in the green-algal lichen *Xanthoria parietina* and their implications on mycobiont–photobiont interactions. *Ecology and Evolution* **2**, 3132–3144.
- Beck A, Friedl T and Rambold G (1998) Selectivity of photobiont choice in a defined lichen community: inferences from cultural and molecular studies. *New Phytologist* **139**, 709–720.
- Beck A, Kasalický T and Rambold G (2002) Myco-photobiont selection in a Mediterranean cryptogam community with *Fulgensia fulgida*. *New Phytologist* **153**, 317–326.
- Beck A, Bechteler J, Casanova-Katny A and Dzhilyanova I (2019) The pioneer lichen *Placopsis* in maritime Antarctica: genetic diversity of their mycobionts and green algal symbionts, and their correlation with deglaciation time. *Symbiosis* **79**, 1–24.
- Beckett RP, Solhaug KA, Gauslaa Y and Minibayeva F (2019) Improved photoprotection in melanized lichens is a result of fungal solar radiation screening rather than photobiont acclimation. *Lichenologist* **51**, 483–491.
- Bergman B and Hällbom L (1981) *Nostoc* of *Peltigera canina* when lichenized and isolated. *Canadian Journal of Botany* **60**, 2092–2098.
- Bergman B, Johanssen C and Söderbäck E (1992) The *Nostoc-Gunnerya* symbiosis (Tansley Review No. 42). *New Phytologist* **122**, 379–400.
- Bertsch A and Butin H (1967) Die Kultur der Erdflechte *Endocarpon pusillum* im Labor. *Planta* **72**, 29–42.
- Bhâradwâja Y (1933) False branching and sheath structure in the *Myxophyceae*, with special reference to the *Scytonemataceae*. *Archiv für Protistenkunde* **81**, 243–283.
- Bitter G (1904) *Peltigera*-Studien II. Das Verhalten der oberseitigen Thallusschuppen der *Peltigera lepidophora* (Nyl). *Berichte der Deutschen Botanischen Gesellschaft* **22**, 251–254.

- Blaž J, Baloch E and Grube M** (2006) High photobiont diversity associated with the euryoecious lichen-forming ascomycete *Lecanora rupicola* (Lecanoraceae, Ascomycota). *Biological Journal of the Linnean Society* **88**, 283–293.
- Boch S, Prati D, Werth S, Rüetschi J and Fischer M** (2011) Lichen endozoochory by snails. *PLoS ONE* **6**, e18770.
- Bock C, Krienitz L and Pröschold T** (2011) Taxonomic reassessment of the genus *Chlorella* (Trebouxiophyceae) using molecular signatures (barcodes), including description of seven new species. *Fottea* **11**, 293–312.
- Boissière J-C, Boissière M-C, Champion-Arnaud P, Lallmant R and Wagner J** (1987) Le cycle des *Nostoc* des genres *Peltigera* et *Collema* en cultures *in vitro* et dans le thalle liquéfactive. *Canadian Journal of Botany* **65**, 1468–1477.
- Bold HC and Wynne M** (1985) *Introduction to the Algae*, 2nd Edn. Englewood Cliffs, New Jersey: Prentice Hall.
- Bornet E** (1873) Recherches sur les gonidies des algues. *Annales des Sciences Naturelles (5me Série, Botanique)* **17**, 45–110.
- Brito A, Vieira J, Vieira CP, Zhu T, Leão PN, Ramos V, Lu X, Vasconcelos VM, Gugger M and Tamagnini P** (2020) Comparative genomics discloses the uniqueness and the biosynthetic potential of the marine cyanobacterium *Hyella patelloides*. *Frontiers in Microbiology* **11**, 1527.
- Brodo IM and Richardson DHS** (1978) Chimeroid association in the genus *Peltigera*. *Lichenologist* **10**, 157–170.
- Brodo IM, Sharnoff SD and Sharnoff S** (2001) *Lichens of North America*. New Haven and London: Yale University Press.
- Brooks F, Rindi F, Suto S, Ohtani S and Green M** (2015) The *Trentepohliales* (Ulvophyceae, Chlorophyta): an unusual algal order and its novel plant pathogen, *Cephaleuros*. *Plant Disease* **99**, 740–753.
- Brunner U and Honegger R** (1985) Chemical and ultrastructural studies on the distribution of sporopollenin-like biopolymers in six genera of lichen photobionts. *Canadian Journal of Botany* **63**, 2221–2230.
- Bubrick P** (1988) Effects of symbiosis on the photobiont. In Galun M (ed.), *CRC Handbook of Lichenology*. Boca Raton, Florida: CRC Press, pp. 133–144.
- Bubrick P, Galun M and Frensdorff A** (1984) Observations on free-living *Trebouxia* de Puymaly and *Pseudotrebouxia* Archibald, and evidence that both symbionts from *Xanthoria parietina* (L.) Th. Fr. can be found free-living in nature. *New Phytologist* **97**, 455–462.
- Büdel B** (1985) Blue-green phycobionts in the lichen family *Lichinaceae*. *Archiv für Hydrobiologie, Supplement* **71** [Algalogical Studies **38/39**], 355–357.
- Büdel B** (1999) Ecology and diversity of rock-inhabiting cyanobacteria in tropical regions. *European Journal of Phycology* **34**, 361–370.
- Büdel B and Henssen A** (1983) *Chroococcidiopsis* (Cyanophyceae), a phycobiont in the lichen family *Lichinaceae*. *Phycologia* **22**, 367–375.
- Büdel B and Henssen A** (1987) *Trebouxia aggregata* und *Gloeocapsa sanguinea*, phycobionten in *Euopsis granatina* (*Lichinaceae*). *Plant Systematics and Evolution* **158**, 235–241.
- Büdel B and Kauff F** (2012) Blue-green algae. In Frey G (series ed.), *Syllabus of Plant Families: Adolf Engler's Syllabus der Pflanzenfamilien, Part VI*. Stuttgart: Borntraeger Verlagbuchhandlung, pp. 5–39.
- Butler RD and Allsopp A** (1972) Ultrastructural investigations in the *Stigonemataceae* (Cyanophyta). *Archiv für Mikrobiologie* **82**, 283–299.
- Canals A, Hernández-Mariné M, Gómez-Bolea A and Llimona X** (1997) *Botryolepraria*, a new monotypic genus segregated from *Lepraria*. *Lichenologist* **29**, 339–345.
- Candotto Carniel F, Zanelli D, Bertuzzi S and Tretiach M** (2015) Desiccation tolerance and lichenization: a case study with the aeroterrestrial microalga *Trebouxia* sp. (Chlorophyta). *Planta* **242**, 493–505.
- Cao S, Zhang F, Liu C, Hao Z, Tian Y, Zhu L and Zhou Q** (2015) Distribution patterns of haplotypes for symbionts from *Umbilicaria esculenta* and *U. muehlenbergii* reflect the importance of reproductive strategy in shaping population genetic structure. *BMC Microbiology* **15**, 212.
- Cao S, Zhang F, Zheng H, Peng F, Liu C and Zhou Q** (2018) *Coccomyxa greatwallensis* sp. nov. (Trebouxiophyceae, Chlorophyta), a lichen epiphytic alga from Fildes Peninsula, Antarctica. *PhytoKeys* **110**, 39–50.
- Cardós JLH, Prieto M, Jylhä M, Aragón G, Molina MC, Martínez I and Rikkinen J** (2019) A case study on the re-establishment of the cyanolichen symbiosis: where do the compatible photobionts come from? *Annals of Botany* **124**, 379–388.
- Casano LM, del Campo E, García-Breijo FJ, Reig-Armiñana J, Gasulla F, del Hoyo A, Guéra A and Barreno E** (2011) Two *Trebouxia* algae with different physiological performances are ever-present in lichen thalli of *Ramalina farinacea*. Coexistence versus competition? *Environmental Microbiology* **13**, 806–818.
- Catalá S, del Campo EM, Barreno E, García-Breijo FJ, Reig-Armiñana J and Casano LM** (2016) Coordinated ultrastructural and phylogenomic analyses shed light on the hidden phycobiont diversity of *Trebouxia* microalgae in *Ramalina fraxinea*. *Molecular Phylogenetics and Evolution* **94**, 765–777.
- Chen L, Zhang L and Liu T** (2016) Concurrent production of carotenoids and lipid by a filamentous microalga *Trentepohlia arborum*. *Bioresource Technology* **214**, 567–573.
- Chodat R** (1913) Monographie d'algues en culture pure. *Matériaux par la Flore Cryptogamique Suisse [Beiträge zur Kryptogamenflora der Schweiz]* **4**, 1–226.
- Clayden SR** (1998) Thallus initiation and development in the lichen *Rhizocarpon lecanorinum*. *New Phytologist* **139**, 685–695.
- Collins CR and Farrar JF** (1978) Structural resistances to mass transfer in the lichen *Xanthoria parietina*. *New Phytologist* **81**, 71–83.
- Coppins BJ** (1983) A taxonomic study of the lichen genus *Micarea* in Europe. *Bulletin of the British Museum (Natural History), Botany Series* **11**, 17–214.
- Cordeiro LM, Reis RA, Cruz LM, Stocker-Wörgötter E, Grube M and Iacomini M** (2005) Molecular studies of photobionts of selected lichens from the coastal vegetation of Brazil. *FEMS Microbiology Ecology* **54**, 381–390.
- Cornejo C and Scheidegger C** (2013) New morphological aspects of cephalodium formation in the lichen *Lobaria pulmonaria* (Lecanorales Ascomycota). *Lichenologist* **45**, 77–87.
- Cornejo C and Scheidegger C** (2016) Cyanobacterial gardens: the liverwort *Frullania asagrayana* acts as a reservoir of lichen photobionts. *Environmental Microbiology Reports* **8**, 352–357.
- Cornejo C, Nelson PR, Stepanchikova I, Himelbrant D, Jørgensen P-M and Scheidegger C** (2016) Contrasting pattern of photobiont diversity in the Atlantic and Pacific populations of *Erioderma pedicellatum* (Pannariaceae). *Lichenologist* **48**, 275–291.
- Costa J-L and Lindblad P** (2002) Cyanobacteria in symbiosis with cycads. In Rai AN, Bergman B and Rasmussen U (eds), *Cyanobacteria in Symbiosis*. Dordrecht: Kluwer Academic Publishers, pp. 195–205.
- Crittenden P, David JC, Hawksworth DL and Campbell FS** (1995) Attempted isolation and success in the culturing of a broad spectrum of lichen-forming and lichenicolous fungi. *New Phytologist* **130**, 267–297.
- Czeczuga B and Maximov OM** (1996) Carotenoids in the cells of the alga *Trentepohlia gobii* Meyer. *Acta Societatis Botanicorum Poloniae* **65**, 273–276.
- Dahlkild A, Kallersjö M, Lohtander K and Tehler A** (2001) Photobiont diversity in the *Physciaceae* (Lecanorales). *Bryologist* **104**, 527–536.
- Dal Forno M, Lawrey JD, Sikaroodi M, Gillevet PM, Schuettelpelz E and Lücking R** (2020) Extensive photobiont sharing in a rapidly radiating cyanolichen clade. *Molecular Ecology* **30**, 1755–1776.
- Dal Grande F, Widmer I, Wagner HH and Scheidegger C** (2012) Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. *Molecular Ecology* **21**, 3159–3172.
- Dal Grande F, Alors D, Divakar PK, Bálint M, Crespo A and Schmitt I** (2014a) Insights into intrathalline genetic diversity of the cosmopolitan lichen symbiotic green alga *Trebouxia decolorans* Ahmadjian using microsatellite markers. *Molecular Phylogenetics and Evolution* **72**, 54–60.
- Dal Grande F, Beck A, Cornejo C, Singh G, Cheenacharoen S, Nelsen MP and Scheidegger C** (2014b) Molecular phylogeny and symbiotic selectivity of the green algal genus *Dictyochloropsis* s. l. (Trebouxiophyceae): a polyphyletic and widespread group forming photobiont-mediated guilds in the lichen family *Lobariaceae*. *New Phytologist* **202**, 455–470.
- Dal Grande F, Rolshausen G, Divakar PK, Crespo A, Otte J, Schleuning M and Schmitt I** (2018) Environment and host identity structure communities of green algal symbionts in lichens. *New Phytologist* **217**, 277–289.
- Darienko T and Pröschold T** (2015) Genetic variability and taxonomic revision of the genus *Auxenochlorella* (Shihira et Krauss) Kalina et Puncocharova (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* **51**, 394–400.
- Darienko T and Pröschold T** (2017) Toward a monograph of non-marine *Ulvophyceae* using an integrative approach (Molecular phylogeny and systematics of terrestrial *Ulvophyceae* II.). *Phytotaxa* **324**, 1–41.

- Darienko T and Pröschold T** (2019) The genus *Jaagichlorella* Reisigl (*Trebouxiophyceae*, *Chlorophyta*) and its close relatives: an evolutionary puzzle. *Phytotaxa* **388**, 47–68.
- Darienko T, Gustavs L, Mudimu O, Menendez CR, Schumann R, Karsten U, Friedl T and Pröschold T** (2010) *Chloroidium*, a common terrestrial coccoid green alga previously assigned to *Chlorella* (*Trebouxiophyceae*, *Chlorophyta*). *European Journal of Phycology* **45**, 79–95.
- Darienko T, Gruber M, Pröschold T and Schagerl M** (2013) *Terrestrial microalgae on Viennese buildings*. Final report of project H-2081/2010. Vienna: Universität Wien.
- Darienko T, Gustavs L, Eggert A, Wolf W and Pröschold T** (2015) Evaluating the species boundaries of green microalgae (*Coccomyxa*, *Trebouxiophyceae*, *Chlorophyta*) using integrative taxonomy and DNA bar-coding with further implications for the species identification in environmental samples. *PLoS ONE* **10**, e0127838.
- Darienko T, Gustavs L and Pröschold T** (2016) Species concept and nomenclatural changes within the genera *Elliptochloris* and *Pseudochlorella* (*Trebouxiophyceae*) based on an integrative approach. *Journal of Phycology* **52**, 1125–1145.
- Darienko T, Lukešová A and Pröschold T** (2018) The polyphasic approach revealed new species of *Chloroidium* (*Trebouxiophyceae*, *Chlorophyta*). *Phytotaxa* **372**, 51–66.
- de los Ríos A, Sancho LG, Grube M, Wierzos J and Ascaso C** (2005) Endolithic growth of two *Lecidea* lichens in granite from continental Antarctica detected by molecular and microscopy techniques. *New Phytologist* **165**, 181–190.
- de los Ríos A, Raggio J, Pérez-Ortega S, Vivas M, Pintado A, Green TGA, Ascaso C and Sancho LG** (2011) Anatomical, morphological and ecophysiological strategies in *Placopsis pycnotheca* (lichenized fungi, *Ascomycota*) allowing rapid colonization of recently deglaciated soils. *Flora* **206**, 857–864.
- Degelius G** (1964) Biological studies of the epiphytic vegetation on twigs of *Fraxinus excelsior*. *Acta Horti Gotoburgensis* **27**, 11–55.
- Delwiche CF** (1999) Tracing the thread of plastid diversity through the tapestry of time. *American Naturalist* **154**, S164–S177.
- Döbbler P** (1984) Symbiosen zwischen Gallertalgen und Gallertpilzen der Gattung *Epigloea* (*Ascomycetes*). *Beiheft zur Nova Hedwigia* **79**, 203–239.
- Dodds WK, Gudder DA and Mollenhauer D** (1995) The ecology of *Nostoc*. *Journal of Phycology* **31**, 2–18.
- Doering JA, Booth T, Wiersma YF and Piercey-Normore MD** (2020) How do genes flow? Identifying potential dispersal mode for the semi-aquatic lichen *Dermatocarpon luridum* using spatial modelling and photobiont markers. *BMC Ecology* **20**, 56.
- Doering M and Piercey-Normore MD** (2009) Genetically divergent algae shape an epiphytic lichen community on Jack Pine in Manitoba. *Lichenologist* **41**, 69–80.
- Drew EA and Smith DC** (1967) Studies in the physiology of lichens VII. The physiology of the *Nostoc* symbiont of *Peltigera polydactyla* compared with cultured and free-living forms. *New Phytologist* **66**, 379–388.
- Eliš M, Neustupa J, Pažoutová M and Škaloud P** (2013) A case of taxonomic inflation in coccoid algae: *Ellipsoidium parvum* and *Neocystis vischeri* are conspecific with *Neocystis* (= *Nephrodiella*) *brevis* (*Chlorophyta*, *Trebouxiophyceae*). *Phytotaxa* **76**, 15–27.
- Elsbary ME, Osman MEH, Abushady AM and Piercey-Normore MD** (2015) Comparison of lichen-forming cyanobacterial and green algal photobionts with free-living algae. *Cryptogamie, Algologie* **36**, 81–100.
- Elvebakk A, Papaefthimiou D, Robertsen EH and Liaimer A** (2008) Phylogenetic patterns among *Nostoc* cyanobionts within bi- and tripartite lichens of the genus *Pannaria*. *Journal of Phycology* **44**, 1049–1059.
- Engelen A, Convey P and Ott S** (2010) Life history strategy of *Lepraria borealis* at an Antarctic inland site, Coal Nunatak. *Lichenologist* **42**, 339–346.
- Engelen A, Convey P, Popa O and Ott S** (2016) Lichen photobiont diversity and selectivity at the southern limit of the maritime Antarctic region (Coal Nunatak, Alexander Island). *Polar Biology* **39**, 2403–2410.
- Ertz D, Guzow-Krzemińska B, Thor G, Łubek A and Kukwa M** (2018) Photobiont switching causes changes in the reproduction strategy and phenotypic dimorphism in the *Arthoniomycetes*. *Scientific Reports* **8**, 4952.
- Ettl H and Gärtner G** (2014) *Syllabus der Boden-, Luft-, und Flechtentalgen*, 2nd Edn. Berlin and Heidelberg: Springer Spektrum.
- Feige GB, Lumbsch HT and Mies B** (1993) Morphological and chemical changes in *Roccella* thalli infected by *Lecanactis grumulosa*. *Cryptogamic Botany* **3**, 101–107.
- Fernández-Marín B, López-Pozo M, Perera-Castro AV, Irati Arzac M, Sáenz-Ceniceros A, Colesie C, de los Ríos A, Sancho LG, Pintado A, Laza JM, et al.** (2019) Symbiosis at its limits: ecophysiological consequences of lichenization in the genus *Prasiola* in Antarctica. *Annals of Botany* **124**, 1211–1226.
- Fernández-Mendoza F, Domaschke S, García MA, Jordan P, Martín MP and Printzen C** (2011) Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*. *Molecular Ecology* **20**, 1208–1232.
- Fewer D, Friedl T and Büdel B** (2002) *Chroococcidiopsis* and heterocyst-differentiating cyanobacteria are each other's closest living relatives. *Molecular Phylogenetics and Evolution* **23**, 82–90.
- Fiore MF, Sant'Anna CL, Azevedo MTDP, Komárek J, Kaštovský J, Sulek J and Lorenzi AAS** (2007) The cyanobacterial genus *Brasilonema*, gen. nov., a molecular and phenotypic evaluation. *Journal of Phycology* **43**, 789–798.
- Fisher KA and Lang NJ** (1971) Comparative ultrastructure of cultured species of *Trebouxia*. *Journal of Phycology* **7**, 155–165.
- Flechtner VR, Johansen JR and Belpaj J** (2009) The biological soil crusts of the San Nicolas Island: enigmatic algae from a geographically isolated ecosystem. *Western North American Naturalist* **68**, 405–436.
- Fontaine KM, Beck A, Stocker-Wörgötter E and Piercey-Normore MD** (2012) Photobiont relationships and phylogenetic history of *Dermatocarpon luridum* var. *luridum* and related *Dermatocarpon* species. *Plants* **1**, 39–60.
- Forsberg A and Lindblad P** (1996) The *Ophrydium-Chlorella* (*Chlorophyceae*, *Chlorophyta*) symbiosis: an ultrastructural characterization. *Phycologia* **35**, 44–47.
- Francisco de Oliveira PM, Timsina B and Piercey-Normore MD** (2012) Diversity of *Ramalina sinensis* and its photobiont in local populations. *Lichenologist* **44**, 649–660.
- Friedl T** (1987) Thallus development and phycobionts of the parasitic lichen *Diploschistes muscorum*. *Lichenologist* **19**, 183–191.
- Friedl T** (1989) Comparative ultrastructure of pyrenoids in *Trebouxia* (*Microthamniales*, *Chlorophyta*). *Plant Systematics and Evolution* **164**, 145–159.
- Friedl T** (1995) Inferring taxonomic positions and testing genus level assignments in coccoid green lichen algae: a phylogenetic analysis of 18S ribosomal RNA sequences from *Dictyochloropsis reticulata* and from members of the genus *Myrmecia* (*Chlorophyta*, *Trebouxiophyceae* cl. nov.). *Journal of Phycology* **31**, 632–639.
- Friedl T** (1996) Evolution of the polyphyletic genus *Pleurastrum* (*Chlorophyta*): inferences from nuclear-encoded ribosomal DNA sequences and motile cell ultrastructure. *Phycologia* **35**, 456–469.
- Friedl T and Büdel B** (2008) Photobionts. In Nash, TH III (ed.), *Lichen Biology*. New York: Cambridge University Press, pp. 9–26.
- Fritsch FE** (1945) *The Structure and Reproduction of the Algae*, Vol. II. Cambridge: Cambridge University Press.
- Fröberg L, Björn LO, Baur A and Baur B** (2001) Viability of lichen photobionts after passing through the digestive tract of a land snail. *Lichenologist* **33**, 543–545.
- Fučíková K, Lewis PO and Lewis LA** (2014) Putting *incertae sedis* taxa in their place: a proposal for ten new families and three new genera in *Sphaeropleales* (*Chlorophyceae*, *Chlorophyta*). *Journal of Phycology* **50**, 14–25.
- Fučíková K, Lewis PO, Neupane S, Karol KG and Lewis LA** (2019) Order, please! Uncertainty in the ordinal-level classification of *Chlorophyceae*. *PeerJ* **7**, e6899.
- Gagunashvili AN and Andrésón ÓS** (2018) Distinctive characters of *Nostoc* genomes in cyanolichens. *BMC Genomics* **19**, 434.
- Galløe O** (1927, 1932) *Natural History of the Danish Lichens*, Parts I and V. Copenhagen: H. Aschehoug and Co, Levin & Munksgaard.
- Gallun [sic] M, Ben-Shaul Y and Paran N** (1971) The fungus-alga association in the *Lecideaceae*: an ultrastructural study. *New Phytologist* **70**, 483–485.
- Galun M, Paran N and Ben-Shaul Y** (1970) The fungus-alga association in the *Lecanoraceae*: an ultrastructural study. *New Phytologist* **69**, 599–603.

- Garbary DJ and London JF (1995) The *Ascophyllum/Polysiphonia/Mycosphaerella* symbiosis. V. Fungal infection protects *A. nodosum* from desiccation. *Botanica Marina* **38**, 529–533.
- Garbary DJ and MacDonald KA (1995) The *Ascophyllum/Polysiphonia/Mycosphaerella* symbiosis. IV. Mutualism in the *Ascophyllum/Mycosphaerella* interaction. *Botanica Marina* **38**, 221–225.
- García-Pichel F (2009) Cyanobacteria. In Schaechter M (ed.), *Encyclopedia of Microbiology*, Vol. 2, 3rd Edn. Oxford and San Diego: Elsevier [Academic Press], pp. 107–124.
- Garrido-Benavent I, Pérez-Ortega S and de los Ríos A (2017) From Alaska to Antarctica: species boundaries and genetic diversity of *Prasiola* (*Trebouxiophyceae*), a foliose chlorophyte associated with the bipolar lichen-forming fungus *Mastodia tessellata*. *Molecular Phylogenetics and Evolution* **107**, 117–131.
- Garrido-Benavent I, de los Ríos A, Fernández-Mendoza F and Pérez-Ortega S (2018) No need for stepping stones: direct, joint dispersal of the lichen-forming fungus *Mastodia tessellata* (*Ascomycota*) and its photobiont explains their bipolar distribution. *Journal of Biogeography* **45**, 213–224.
- Garrido-Benavent I, Pérez-Ortega S, de los Ríos A and Fernández-Mendoza F (2020) Amphitropical variation of the algal partners of *Pseudophebe* (*Parmeliaceae*, lichenized fungi). *Symbiosis* **82**, 35–48.
- Gärtner G (1985) Die Gattung *Trebouxia* Puymaly (*Chlorellales*, *Chlorophyceae*). *Archiv für Hydrobiologie, Supplement* **71** [Algalological Studies **41**], 495–548.
- Gärtner G and Ingolić E (1989) Ein Beitrag zur Kenntnis von *Apatococcus lobatus* (*Chlorophyta*, *Chaetophorales*, *Leptosiroideae*). *Plant Systematics and Evolution* **164**, 133–143.
- Garty J and Delarea J (1988) Evidence of liberation of lichen ascospores in clusters and reports on contact between free-living algal cells and germinating lichen ascospores under natural conditions. *Canadian Journal of Botany* **66**, 2171–2177.
- Gasulla F, Guéra A, de los Ríos A and Pérez-Ortega S (2019) Differential responses to salt concentrations of lichen photobiont strains isolated from lichens occurring in different littoral zones. *Plant and Fungal Systematics* **64**, 149–162.
- Gasulla F, Barrasa JM, Casano LM and del Campo EM (2020) Symbiont composition of the basidiolichen *Lichenomphalia meridionalis* varies with altitude in the Iberian Peninsula. *Lichenologist* **52**, 17–26.
- Gauslaa Y, Alam MA, Lucas P-L, Chowdhury DP and Solhaug KA (2017) Fungal tissue *per se* is stronger as a UV-B screen than secondary fungal extroliths in *Lobaria pulmonaria*. *Fungal Ecology* **26**, 109–113.
- Geitler L (1933) Beiträge zur Kenntnis de Flechtensymbiose. I–III. *Archiv für Protistenkunde* **80**, 378–409.
- Geitler L (1934) Beiträge zur Kenntnis de Flechtensymbiose. IV, V. *Archiv für Protistenkunde* **82**, 51–85.
- Geitler L (1955) *Clavaria mucida*, eine extratropische Basidiolichene. *Biologisches Zentralblatt* **74**, 145–159.
- Giddings TH, Jr, and Staehelin LA (1981) Observations of microplasmodesmata in both heterocyst-forming and non-heterocyst forming filamentous cyanobacteria by freeze-fracture electron microscopy. *Archives of Microbiology* **129**, 295–298.
- González MA, Pröschold T, Palacios Y, Aguayo P, Inostroza I and Gómez PI (2013) Taxonomic identification and lipid production of two Chilean *Chlorella*-like strains isolated from a marine and an estuarine coastal environment. *AoB PLANTS* **5**, plt020.
- Good BH and Chapman RL (1978) The ultrastructure of *Phycopeltis* (*Chroolepidaceae*: *Chlorophyta*). I. Sporopollenin in the cell walls. *American Journal of Botany* **65**, 27–33.
- Graham LE, Graham JM and Wilcox LW (2009) *Algae*, 2nd Edn. San Francisco: Pearson Education Inc.
- Green MS (2012) *The Trentepohliales* (*Ulvophyceae*, *Chlorophyta*) from coastal South Carolina. Ph.D. thesis, Tennessee Technological University.
- Green TGA, Büdel B, Heber U, Meyer A, Zellner H and Lange OL (1993) Differences in photosynthetic performance between cyanobacterial and green algal components of lichen photosymbiodemes measured in the field. *New Phytologist* **125**, 723–731.
- Green TGA, Schlenos M, Sancho LG, Winkler JB, Broom FD and Schroeter B (2002) The photobiont determines the pattern of photosynthetic activity within a single lichen thallus containing cyanobacterial and green algal sectors (photosymbiodeme). *Oecologia* **130**, 191–198.
- Greenhalgh GN and Anglesea D (1979) The distribution of algal cells in lichen thalli. *Lichenologist* **11**, 283–292.
- Grube M and Berg B (2009) Microbial consortia of bacteria and fungi with focus on the lichen symbiosis. *Fungal Biology Reviews* **23**, 72–85.
- Grube M and Lücking R (2002) Fine structure of foliicolous lichens and their lichenicolous fungi studied by epifluorescence. *Symbiosis* **32**, 229–246.
- Grube M, Cernava T, Soh J, Fuchs S, Aschenbrenner I, Lassek C, Wegner U, Becher D, Riedel K, Sensen CW, et al. (2015) Exploring functional contexts of symbiotic sustain within lichen-associated bacteria by comparative omics. *ISME Journal* **9**, 412–424.
- Grube M, Muggia L, Baloch E, Hametner C and Stocker-Wörgötter E (2017a) Symbiosis of lichen-forming fungi with trentepohlialean algae. In Grube M, Seckbach J and Muggia L (eds), *Algal and Cyanobacterial Symbioses*. London: World Scientific Publishing Europe Ltd, pp. 85–110.
- Grube M, Seckbach J and Muggia L (2017b) *Algal and Cyanobacterial Symbioses*. London: World Scientific Publishing Europe Ltd.
- Gueidan C, Thüs H and Pérez-Ortega S (2011) Phylogenetic position of the brown algae-associated lichenized fungus *Verrucaria tavaresiae* (*Verrucariaceae*). *Bryologist* **114**, 563–569.
- Gustavs L, Eggert A, Michalik D and Karsten U (2010) Physiological and biochemical responses of green microalgae from different habitats to osmotic and matrix stress. *Protoplasma* **243**, 3–14.
- Gustavs L, Görs M and Karsten U (2011) Polyol patterns in biofilm-forming aeroterrestrial green algae (*Trebouxiophyceae*, *Chlorophyta*). *Journal of Phycology* **47**, 533–537.
- Gustavs L, Schumann R and Karsten U (2016) Mixotrophy in the terrestrial green alga *Apatococcus lobatus* (*Trebouxiophyceae*, *Chlorophyta*). *Journal of Phycology* **52**, 311–314.
- Gustavs L, Schiefelbein U and Darienko T (2017) Symbioses of the green algal genera *Coccomyxa* and *Elliptochloris* (*Trebouxiophyceae*, *Chlorophyta*). In Grube M, Seckbach J and Muggia L (eds), *Algal and Cyanobacterial Symbioses*. London: World Scientific Publishing Europe Ltd, pp. 169–208.
- Guzow-Krzemińska B (2006) Photobiont flexibility in the lichen *Protarmeliopsis muralis* as revealed by ITS rDNA analyses. *Lichenologist* **38**, 469–476.
- Hallmann C, Stannek L, Fritzl D, Hause-Reitner D, Friedl T and Hoppert M (2013) Molecular diversity of phototrophic biofilms on building stone. *FEMS Microbiology Ecology* **84**, 355–372.
- Hallmann C, Hoppert M, Mudimu O and Friedl T (2016) Biodiversity of green algae covering artificial hard substrate surfaces in a suburban environment: a case study using molecular approaches. *Journal of Phycology* **52**, 732–744.
- Hametner C, Stocker-Wörgötter E and Grube M (2014a) New insights into diversity and selectivity of trentepohlialean lichen photobionts from the extratropics. *Symbiosis* **63**, 31–40.
- Hametner C, Stocker-Wörgötter E, Rindi F and Grube M (2014b) Phylogenetic position and morphology of lichenized *Trentepohliales* (*Ulvophyceae*, *Chlorophyta*) from selected species of *Graphidaceae*. *Phycological Research* **62**, 170–186.
- Handa S, Ohmura Y, Nakano T and Nakahara-Tsubota M (2007) Airborne green microalgae (*Chlorophyta*) in snowfall. *Hikobia* **15**, 109–120 [In Japanese, with English abstract and figure legends].
- Hauck M, Helms G and Friedl T (2007) Photobiont selectivity in the epiphytic lichens *Hypogymnia physodes* and *Lecanora conizaeoides*. *Lichenologist* **39**, 195–204.
- Hawksworth DL (1988) The variety of fungal-algal symbioses, their evolutionary significance, and the nature of lichens. *Botanical Journal of the Linnean Society* **96**, 3–20.
- Hawksworth DL, Coppins BJ and James PW (1979) *Blarneya*, a lichenized hyphomycete from southern Ireland. *Botanical Journal of the Linnean Society* **79**, 357–367.
- Helms G, Friedl T, Rambold G and Mayrhofer H (2001) Identification of photobionts from the lichen family *Physciaceae* using algal-specific ITS rDNA sequencing. *Lichenologist* **33**, 73–86.
- Henley WJ, Hironaka JL, Gillou L, Buchheim MA, Buchheim JA, Fawley MW and Fawley KP (2004) Phylogenetic analysis of the *Nannochloris-*

- like' algae and diagnoses of *Picochlorum oklahomensis* gen. et sp. nov. (*Trebouxiophyceae*, *Chlorophyta*). *Phycologia* **43**, 641–652.
- Henskens FL, Green TGA and Wilkins A** (2012) Cyanolichens can have both cyanobacteria and green algae in a common layer as major contributors to photosynthesis. *Annals of Botany* **110**, 555–563.
- Henson BJ, Watson LE and Barnum SR** (2002) Molecular differentiation of the heterocystous cyanobacteria, *Nostoc* and *Anabaena*, based on complete *NifD* sequences. *Current Microbiology* **45**, 161–164.
- Henssen A** (1981) *Hyphomorpha* als Phycobiont in Flechten. *Plant Systematics and Evolution* **137**, 139–143.
- Henssen A** (1990) *Thermutopsis jamesii*, a new member of the *Lichinaceae* from Antigua. *Lichenologist* **22**, 253–259.
- Henssen A** (1994) Contribution to the morphology and species delimitation in *Heppia* sensu stricto (lichenized *Ascomycotina*). *Acta Botanica Fennica* **150**, 57–73.
- Henssen A** (1995) Studies on the biology and structure of *Dacampia* (*Dothideales*), a genus with lichenized and lichenicolous species. *Cryptogamic Botany* **5**, 149–158.
- Henssen A and Jahns HM** (1974) *Lichenes: Eine Einführung in die Flechtenkunde*. Stuttgart: Georg Thieme.
- Hessler R and Peveling E** (1978) Die Lokalisation von ¹⁴C-Assimilaten in Flechtenthalli von *Cladonia incrassata* Floerke und *Hypogymnia physodes* (L.) Ach. *Zeitschrift für Pflanzenphysiologie* **86**, 287–302.
- Hestmark G, Lutzoni F and Miallikowska J** (2016) Photobiont associations in co-occurring umbilicate lichens with contrasting modes of reproduction in coastal Norway. *Lichenologist* **48**, 545–557.
- Hill DJ** (1972) The movement of carbohydrate from the alga to the fungus in the lichen *Peltigera polydactyla*. *New Phytologist* **71**, 31–39.
- Hill DJ** (1985) Changes in photobiont dimensions and numbers during co-development of lichen symbionts. In Brown DH (ed.), *Lichen Physiology and Cell Biology*. New York and London: Plenum Press, pp. 303–317.
- Hill DJ** (1989) The control of the cell cycle in microbial symbionts. *New Phytologist* **112**, 175–184.
- Hill DJ** (2009) Asymmetrical co-evolution in the lichen symbiosis caused by a limited capacity for adaptation in the photobiont. *Botanical Review* **75**, 326–338.
- Hitch CBJ and Millbank JW** (1975) Nitrogen metabolism in lichens VII. Nitrogenase activity and heterocyst frequency in lichens with blue-green phycobionts. *New Phytologist* **75**, 239–244.
- Hodač L, Hallmann C, Spitzer K, Elster J, Faßhauer F, Brinkmann N, Lepka D, Diwan V and Fried T** (2016) Widespread green algae *Chlorella* and *Stichococcus* exhibit polar-temperate and tropical-temperate biogeography. *FEMS Microbiology Ecology* **92**, fiw122.
- Hodkinson BP, Moncada B and Lücking R** (2014) *Lepidostromatales*, a new order of lichenized fungi (*Basidiomycota*, *Agaricomycetes*), with two new genera, *Ertzia* and *Sulzbacheromyces*, and one new species, *Lepidostroma winklerianum*. *Fungal Diversity* **64**, 165–179.
- Honegger R** (1986) Ultrastructural studies in lichens I. Haustorial types and their frequencies in a range of lichens with trebouxioid photobionts. *New Phytologist* **103**, 785–795.
- Honegger R** (1987) Questions about pattern formation in the algal layer of lichens with stratified (heteromorous) thalli. *Bibliotheca Lichenologica* **25**, 59–71.
- Honegger R** (1990) Haustoria-like structures and cell wall surface layers in lichens. In Mendgen K and Leseman DE (eds), *Electron Microscopy of Plant Pathogens*. Berlin: Springer-Verlag, pp. 277–290.
- Honegger R** (1991) Functional aspects of the lichen symbiosis. *Annual Review of Plant Physiology and Plant Molecular Biology* **42**, 553–578.
- Honegger R** (2012) The symbiotic phenotype of lichen-forming ascomycetes and their endo- and epibionts. In Hock B (ed.), *The Mycota IX: Fungal Associations*. Berlin and Heidelberg: Springer-Verlag, pp. 287–339.
- Honegger R and Brunner U** (1981) Sporopollenin in the cell wall of *Coccomyxa* and *Myrmecia* phycobionts of various lichens: an ultrastructural and chemical investigation. *Canadian Journal of Botany* **59**, 2713–2734.
- Hoshina R and Imamura N** (2008) Multiple origins of the symbioses in *Paramecium bursaria*. *Protist* **159**, 53–63.
- Hoshina R, Kamako S-i and Imamura N** (2004) Phylogenetic position of endosymbiotic green algae in *Paramecium bursaria* Ehrenberg from Japan. *Plant Biology* **6**, 447–453.
- Hoshina R, Iwataki M and Imamura N** (2010) *Chlorella variabilis* and *Micractinium reisseri* sp. nov. (*Chlorellaceae*, *Trebouxiophyceae*): redescription of the endosymbiotic green algae of *Paramecium bursaria* (*Peniculia*, *Oligohymenophorea*) in the 120th year. *Phycological Research* **58**, 188–201.
- Huss VAR, Frank C, Hartmann EC, Hirmer M, Kloboucek A, Seidel B, Wenzeler P and Kessler E** (1999) Biochemical taxonomy and molecular phylogeny of the genus *Chlorella sensu lato* (*Chlorophyta*). *Journal of Phycology* **35**, 587–598.
- Ihda T, Nakano T and Deguchi H** (1997) Photobionts of Japanese *Sphaerophorus*. *Symbiosis* **23**, 1–8.
- Ikeda T and Takeda H** (1995) Species-specific differences of pyrenoids in *Chlorella* (*Chlorophyta*). *Journal of Phycology* **31**, 813–818.
- Jaag O and Thomas E** (1934) Neue Untersuchungen über die Flechte *Epigloea bactrospora* Zukul. *Berichte der Schweizerischen Botanischen Gesellschaft* **43**, 77–89.
- Jacobs JB and Ahmadjian V** (1971) The ultrastructure of lichens. IV. Movement of carbon products from alga to fungus as demonstrated by high resolution autoradiography. *New Phytologist* **70**, 47–50.
- Jahns HM, Mollenhauer D, Jenniger M and Schönborn D** (1979) Die Neubesiedlung von Baumrinde durch Flechten I. *Natur und Museum* **109**, 40–51.
- Jaklitsch W, Baral H-O, Lücking R and Lumbsch HT** (2016) *Syllabus of Plant Families: A. Engler's Syllabus der Pflanzenfamilien. Part 1/2, Ascomycota*. Stuttgart: Borntraeger Verlagsbuchhandlung.
- James PW and Henssen A** (1976) The morphological and taxonomic significance of cephalodia. In Brown DH, Hawksworth DL and Bailey RH (eds), *Lichenology: Progress and Problems*. London: Academic Press, pp. 27–77.
- Jayalal U, Aptroot A and Hur J-S** (2012) The lichen genus *Polychidium* new to South Korea. *Mycobiology* **40**, 252–254.
- Jiang SH, Hawksworth DL, Lücking R and Wei JC** (2020) A new genus and species of foliicolous lichen in a new family of *Strigulales* (*Ascomycota: Dothideomycetes*) reveals remarkable class-level homoplasy. *IMA Fungus* **11**, 1.
- Joneson S and O'Brien H** (2017) A molecular investigation of free-living and lichenized *Nostoc* sp. and symbiotic lifestyle determination. *Bryologist* **120**, 371–381.
- Jordan WP** (1970) The internal cephalodia of the genus *Lobaria*. *Bryologist* **73**, 669–681.
- Jordan WP** (1972) Erumpent cephalodia, an apparent case of phycobial influence on lichen morphology. *Journal of Phycology* **8**, 112–117.
- Jordan WP and Rickson F** (1971) Cyanophyte cephalodia in the lichen genus *Nephroma*. *American Journal of Botany* **58**, 562–568.
- Joubert JJ and Rijkenberg FJH** (1971) Parasitic green algae. *Annual Review of Phytopathology* **9**, 45–64.
- Jülich W** (1972) Monographie der *Atheliae* (*Corticaceae*, *Basidiomycetes*). *Willdenowia, Beiheft* **7**, 1–283.
- Jülich W** (1978) A new lichenized *Athelia* from Florida. *Persoonia* **10**, 149–151.
- Jung P, Emrich D, Briegel-Williams L, Schermer M, Weber L, Baumann K, Colesie C, Clerc P, Lehnert LW, Achilles S, et al.** (2019) Ecophysiology and phylogeny of new terricolous and epiphytic chlorolichens in a fog oasis of the Atacama Desert. *MicrobiologyOpen* **8**, e894.
- Jürjado I, Kaasalainen U, Jylhä and Rikkinen J** (2019) Relationships between mycobiont identity, photobiont specificity and ecological preferences in the lichen genus *Peltigera* (*Ascomycota*) in Estonia (northeastern Europe). *Fungal Ecology* **39**, 45–54.
- Kardish N, Kessel M and Galun M** (1989) Characterization of symbiotic and cultured *Nostoc* of the lichen *Nephroma laevigatum* Ach. *Symbiosis* **7**, 257–266.
- Keeling PJ** (2004) Diversity and evolutionary history of plastids and their hosts. *American Journal of Botany* **91**, 1481–1493.
- Keeling PJ** (2013) The number, speed, and impact of plastid endosymbioses in eukaryotic evolution. *Annual Review of Plant Biology* **64**, 583–607.
- Kharkongor D and Ramanujam P** (2015) Spatial and temporal variation of carotenoids in four species of *Trentepohlia* (*Trentepohliales*, *Chlorophyta*). *Journal of Botany* **2015**, 201641.
- Khayatan B, Meeks J and Risser DD** (2015) Evidence that a modified type IV pilus-like system powers gliding motility and polysaccharide secretion in filamentous cyanobacteria. *Molecular Microbiology* **98**, 1021–1036.

- Kim JI, Nam SW, So JE, Hong SG, Choi HG and Shin W (2017) *Asterochloris sejongensis* sp. nov. (Trebouxiophyceae, Chlorophyta) from King George Island, Antarctica. *Phytotaxa* **295**, 60–70.
- Kim JI, Kim YJ, Nam SW, So JE, Hong SG, Choi H-G and Shin W (2020) Taxonomic study of three new Antarctic *Asterochloris* (Trebouxiophyceae) based on morphological and molecular data. *Algae* **35**, 17–32.
- Kirk PM, Cannon PF, David JC and Stalpers JA (2001) *Ainsworth and Bisby's Dictionary of the Fungi*, 9th Edn. Wallingford: CAB International.
- Kluge M, Mollenhauer D, Wolf E and Schüßler A (2002) The *Nostoc-Geosiphon* endocytobiosis. In Rai AN, Bergman B and Rasmussen U (eds), *Cyanobacteria in Symbiosis*. Dordrecht: Kluwer Academic Publishers, pp. 19–30.
- Kohlmeyer J, Hawksworth DL and Volkmann-Kohlmeyer B (2004) Observations on two marine and maritime 'borderline' lichens: *Mastodia tessellata* and *Collemopsidium pelvetiae*. *Mycological Progress* **3**, 51–56.
- Komárek J, Kaštovský J, Mareš J and Johansen JR (2014) Taxonomic classification of cyanoprokaryotes (cyanobacterial genera) 2014, using a polyphasic approach. *Preslia* **86**, 295–335.
- Kono M, Kon Y, Ohmura Y, Satta Y and Terai Y (2020) In vitro resynthesis of lichenization reveals the genetic background of symbiosis-specific fungal-algal interaction in *Usnea hakonensis*. *BMC Genomics* **21**, 1–16.
- Kosecka M, Jabłońska A, Flakus A, Rodriguez-Flakus P, Kukwa M and Guzow-Krzemińska B (2020) Trentepohlialean algae (Trentepohliales, Ulvophyceae) show preference to selected mycobiont lineages in lichen symbioses. *Journal of Phycology* **56**, 979–993.
- Kosecka M, Guzow-Krzemińska B, Čemajová I, Škaloud P, Jabłońska A and Kukwa M (2021) New lineages of photobionts in Bolivian lichens expand our knowledge on habitat preferences and distribution of *Asterochloris* algae. *Scientific Reports* **11**, 8701.
- Kosugi M, Arita M, Shizuma R, Moryama Y, Kashino Y, Koike H and Satoh K (2009) Responses to desiccation stress in lichens are different from those in their photobionts. *Plant and Cell Physiology* **50**, 879–888.
- Kouwets FAC (1996) Ultrastructural studies of the cell cycle in a multicentriolar form of *Bracteacoccus minor* (Chlorophyceae, Chlorellales). *Protoplasma* **191**, 191–204.
- Kovačević G, Franjević D, Jelenčić B and Kalafatić M (2010) Isolation and cultivation of endosymbiotic algae from green hydra and phylogenetic analysis of 18S rDNA sequences. *Folia Biologica (Kraków)* **58**, 135–143.
- Kovačik L and Batista Pereira A (2001) Green alga *Prasiola crispa* and its lichenized form *Mastodia tessellata* in Antarctic environment: general aspects. *Nova Hedwigia* **123**, 465–478.
- Kranner I, Cram WJ, Zorn M, Wornik S, Yoshimura I, Stabentheiner E and Pfeifhofer HW (2005) Antioxidants and photoprotection in a lichen as compared with its isolated symbiotic partners. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 3141–3146.
- Kranner I, Beckett R, Hochman A and Nash, TH III (2008) Desiccation-tolerance in lichens: a review. *Bryologist* **111**, 576–593.
- Kroken S and Taylor JW (2000) Phylogenetic species, reproductive mode, and specificity of the green alga *Trebouxia* forming lichens with the fungal genus *Letharia*. *Bryologist* **103**, 645–660.
- Kukwa M and Pérez-Ortega S (2010) A second species of *Botryolepraria* from the Neotropics and the phylogenetic placement of the genus within *Ascomycota*. *Mycological Progress* **9**, 345–351.
- Kumar K, Mella-Herrera RA and Golden JW (2010) Cyanobacterial heterocysts. *Cold Spring Harbor Perspectives in Biology* **2**, a000315.
- Lakatos M, Lange-Bertalot H and Büdel B (2004) Diatoms living inside the thallus of the green algal lichen *Coenogonium linkii* in neotropical lowland rain forests. *Journal of Phycology* **40**, 70–73.
- Lamb IM (1959) Lichens. *Scientific American* **201**, 144–159.
- Lange OL, Kilian E and Ziegler H (1986) Water vapor uptake and photosynthesis of lichens: performance differences in species with green and blue-green algae as phycobionts. *Oecologia* **71**, 104–110.
- Lange OL, Pfanz H, Kilian E and Meyer A (1990) Effect of low water potential on photosynthesis in intact lichens and their liberated algal components. *Planta* **182**, 467–472.
- Lange OL, Büdel B, Meyer A and Kilian E (1993) Further evidence that activation of net photosynthesis by dry cyanobacterial lichens requires liquid water. *Lichenologist* **25**, 175–189.
- Lavoie C, Renaudin M, McMullin RT, Ganon J, Roy C, Beaulieu M-E, Bellenger JP and Villareal JC (2020) Extremely low genetic diversity of *Stigonema* associated with *Stereocaulon* in eastern Canada. *Bryologist* **123**, 188–203.
- Leavitt SD, Nelsen MP, Lumbsch HT, Johnson LA and St Clair LL (2013) Symbiont flexibility in subalpine rock shield lichen communities in the Southwestern USA. *Bryologist* **116**, 149–161.
- Leavitt SD, Kraichak E, Vondrak J, Nelsen MP, Altermann S, Divakar PK, Alors D, Esslinger TL, Crespo A and Lumbsch HT (2015) Fungal specificity and selectivity for algae play a major role in determining lichen partnerships across diverse ecogeographic regions in the lichen-forming family *Parmeliaceae* (Ascomycota). *Molecular Ecology* **24**, 3779–3797.
- Leavitt SD, Kraichak E, Vondrak J, Nelsen MP, Sohrabi M, Pérez-Ortega S, St Clair LL and Lumbsch HT (2016) Cryptic diversity and symbiont interactions in rock-posed lichens. *Molecular Phylogenetics and Evolution* **99**, 261–274.
- Lee RE (2018) *Phycology*, 5th Edn. Cambridge: Cambridge University Press.
- Leliaert F, Smith DR, Moreau H, Herron MD, Verbruggen H, Delwiche CF and De Clerck O (2012) Phylogeny and molecular evolution of the green algae. *Critical Reviews in Plant Sciences* **31**, 1–46.
- Lemieux C, Otis C and Turmel M (2014) Chloroplast phylogenomic analysis resolves deep-level relationships within the green algal class *Trebouxiophyceae*. *BMC Evolutionary Biology* **14**, 211.
- Letrouit-Galinou M-A (1968) Les algues des lichens. *Bulletin de la Société Botanique de France* **115** (Supp. 2), 35–77.
- Letrouit-Galinou M-A and Asta J (1994) Thallus morphogenesis in some lichens. *Cryptogamic Botany* **4**, 274–282.
- Letsch MR, Muller-Parker G, Friedl T and Lewis LA (2009) *Elliptochloris marina* sp. nov. (Trebouxiophyceae, Chlorophyta), symbiotic green alga of the temperate Pacific sea anemones *Anthopleura xanthogrammica* and *A. elegantissima* (Anthozoa, Cnidaria). *Journal of Phycology* **45**, 1127–1135.
- Lewis L and Muller-Parker G (2004) Phylogenetic placement of 'Zoochlorella' (Chlorophyta), algal symbiont of the temperate sea anemone *Anthopleura elegantissima*. *Biological Bulletin* **207**, 87–92.
- Li B, Feng J and Xie S (2013) Morphological and phylogenetic study of algal partners associated with the lichen-forming fungus *Porpidia crustulata* from the Guancen Mountains, northern China. *Symbiosis* **61**, 37–46.
- Li S, Sun H, Hu Y, Liu B, Zhu H, Hu Z and Liu G (2020) Four new members of foliicolous green algae within the *Watanabea* clade (Trebouxiophyceae, Chlorophyta) from China. *Journal of Eukaryotic Microbiology* **67**, 369–382.
- Li S, Tan H, Liu B, Zhu H, Hu Z and Liu G (2021) *Watanabeales* ord. nov. and twelve novel species of *Trebouxiophyceae* (Chlorophyta). *Journal of Phycology*, doi: 10.1111/JPY.13165
- Lindgren H, Velmala S, Högnabba F, Goward T, Holien H and Myllys L (2014) High fungal selectivity for algal symbionts in the genus *Bryoria*. *Lichenologist* **46**, 681–695.
- Lindgren H, Moncada B, Lücking R, Magain N, Simon A, Goffinet B, Sérusiaux E, Nelsen M, Mercado-Díaz J, Widhelm T, et al. (2020) Cophylogenetic patterns in algal symbionts correlate with repeated symbiont switches during diversification and geographic expansion of lichen-forming fungi in the genus *Sticta* (Ascomycota: Peltigeraceae). *Molecular Phylogenetics and Evolution* **150**, 106860.
- Liu A, Zhu T, Lu X and Song L (2013) Hydrocarbon profiles and phylogenetic analyses of diversified cyanobacterial species. *Applied Energy* **111**, 383–393.
- Lohtander K, Oksanen I and Rikkinen J (2003) Genetic diversity of green algal and cyanobacterial photobionts in *Nephroma* (Peltigerales). *Lichenologist* **35**, 325–339.
- López-Bautista JM and Chapman RL (2003) Phylogenetic affinities of the *Trentepohliales* inferred from small-subunit rDNA. *International Journal of Systematic and Evolutionary Microbiology* **53**, 2099–2106.
- López-Bautista JM, Rindi F and Guiry MD (2006) Molecular systematics of the subaerial green algal order *Trentepohliales*: an assessment based on morphological and molecular data. *International Journal of Systematic and Evolutionary Microbiology* **56**, 1709–1715.
- Lücking R (1994) A new foliicolous species of *Microtheliopsis* (Lichens, *Microtheliopsidaceae*) from Costa Rica. *Mycotaxon* **51**, 69–73.
- Lücking R (2008) Follicolous lichenized fungi. *Flora Neotropica Monograph* **103**, 1–866.

- Lücking R and Grube M (2002) Facultative parasitism and reproductive strategies in *Chroodiscus* (Ascomycota, Ostropales). *Stappia* **80**, 267–292.
- Lücking R, Lawrey JD, Sikaroodi M, Gillevé PM, Chaves JL, Sipman HJM and Bungartz F (2009) Do lichens domesticate photobionts like farmers domesticate crops? Evidence from a previously unrecognized lineage of filamentous cyanobacteria. *American Journal of Botany* **96**, 1409–1418.
- Lücking R, Barrie FR and Genney D (2014) *Dictyonema coppinsii*, a new name for the European species known as *Dictyonema interruptum* (Basidiomycota: Agaricales: Hygrophoraceae), with a validation of its photobiont *Rhizonema* (Cyanoprokaryota: Nostocales: Rhizonemataceae). *Lichenologist* **46**, 261–267.
- Lücking R, Hodgkinson BP and Leavitt SD (2017a) The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota – approaching one thousand genera. *Bryologist* **119**, 361–416.
- Lücking R, Thorn RG, Saar I, Piercey-Normore MD, Moncada B, Doering J, Mann H, Lebeuf R, Voitk M and Voitk A (2017b) A hidden basidiolichen rediscovered: *Omphalina oreades* is a separate species in the genus *Lichenomphalia* (Basidiomycota: Agaricales: Hygrophoraceae). *Lichenologist* **49**, 467–481.
- Lud D, Huiskes AHL and Ott S (2001) Morphological evidence for the symbiotic character of *Turgidosculum complicatulum* Kohlm. & Kohlm. (= *Mastodia tessellata* Hook. f. & Harvey). *Symbiosis* **31**, 141–151.
- Ludwig LR (2015) *The reproductive ecology of Icmadophila splashnirima, including aspects of the reproduction in additional members of Icmadophilaceae*. Ph.D. thesis, University of Otago.
- Luo W, Pröschold T, Bock C and Krienitz L (2010) Generic concept in *Chlorella*-related coccoid green algae (Chlorophyta, Trebouxiophyceae). *Plant Biology* **12**, 545–553.
- Lutsak T, Fernández-Mendoza F, Kirika P, Wondafra M and Printzen C (2016) Mycobiont-photobiont interactions of the lichen *Cetraria aculeata* in high alpine regions of East Africa and South America. *Symbiosis* **68**, 25–37.
- Lutzoni FM and Brodo IM (1995) A generic redelimitation of the *Ionaspis-Hymenelia* complex (lichenized Ascomycotina). *Systematic Botany* **20**, 224–258.
- Ma S, Huss VAR, Tan D, Sun X, Chun J, Xie Y and Zhang J (2013) A novel species in the genus *Heveochlorella* (Trebouxiophyceae, Chlorophyta) witnesses the evolution from an epiphytic into an endophytic lifestyle in tree-dwelling green algae. *European Journal of Phycology* **48**, 200–209.
- Magain N and Sérusiaux E (2014) Do photobiont switch and cephalodia emancipation act as evolutionary drivers in the lichen symbiosis? A case study in the *Pannariaceae* (Peltigerales). *PLoS ONE* **9**, e89876.
- Magain N, Goffinet B and Sérusiaux E (2012) Further photomorphs in the lichen family *Lobariaceae* from Réunion (Mascarene archipelago) with notes on the phylogeny of *Dendriscoaulon* cyanomorphs. *Bryologist* **115**, 243–254.
- Magain N, Miadlikowska J, Goffinet B, Sérusiaux E and Lutzoni F (2017) Macroevolution of specificity in cyanolichens of the genus *Peltigera* section *Polydactylon* (Lecanoromycetes, Ascomycota). *Systematic Botany* **66**, 74–99.
- Magain N, Truong C, Goward T, Niu D, Goffinet B, Sérusiaux E, Vitikainen O, Lutzoni F and Miadlikowska J (2018) Species delimitation at a global scale reveals high species richness with complex biogeography and patterns of symbiont association in *Peltigera* section *Peltigera* (lichenized Ascomycota: Lecanoromycetes). *Taxon* **67**, 836–870.
- Makra N, Gell G, Juhász A, Soós V, Kiss T, Molnár Z, Ördög V, Vörös L and Balázs E (2019) Molecular taxonomic evaluation of *Anabaena* and *Nostoc* strains from the Mosonmagyaróvár Algal Culture Collection. *South African Journal of Botany* **124**, 80–86.
- Malavasi V, Škaloud P, Rindi F, Tempesta S, Paoletti M and Pasqualetti M (2016) DNA-based taxonomy in ecologically versatile microalgae: a re-evaluation of the species concept within the coccoid green algal genus *Coccomyxa* (Trebouxiophyceae, Chlorophyta). *PLoS ONE* **11**, e0151137.
- Mansournia MR, Wu B, Matsushita N and Hogetsu T (2012) Genotypic analysis of the foliose lichen *Parmotrema tinctorum* using microsatellite markers: association of mycobiont and photobiont, and their reproductive modes. *Lichenologist* **44**, 419–440.
- Marini L, Nascimbene J and Nimis PL (2011) Large-scale patterns of epiphytic lichen species richness: photobiont-dependent response to climate and forest structure. *Science of the Total Environment* **409**, 4381–4386.
- Mark K, Laanisto L, Bueno CG, Niinemets Ü, Keller C and Scheidegger C (2020) Contrasting co-occurrence patterns of photobiont and cystobasidiomycete yeast associated with common epiphytic lichen species. *New Phytologist* **227**, 1362–1375.
- Marton K and Galun M (1976) *In vitro* dissociation and reassociation of the symbionts of the lichen *Heppia echinulata*. *Protoplasma* **87**, 135–143.
- Masumoto H (2020) *Taxonomic studies on lichenized basidiomycetes and their photobionts in Japan: towards the establishment of a model co-culture system of lichen symbiosis*. Ph.D. thesis, University of Tsukuba.
- Masumoto H, Ohmura Y and Degawa Y (2019) *Lichenomphalia meridionalis* (Hygrophoraceae, lichenized Basidiomycota) new to Asia. *Opuscula Philolichenum* **18**, 379–389.
- Matthews SW, Tucker SC and Chapman RL (1989) Ultrastructural features of mycobionts and trentepohliaceous phycobionts in selected tropical crustose lichens. *Botanical Gazette* **150**, 417–438.
- Mattox KR and Stewart KD (1984) Classification of the green algae: a concept based on comparative cytology. In Irvine DEG and John DM (eds), *Systematics of the Green Algae*. London: Academic Press, pp. 29–72.
- McCune B, Arup U, Breuss O, Di Meglio E, Di Meglio J, Esslinger TL, Magain N, Miadlikowska J, Miller AE, Muggia L, et al. (2018) Biodiversity and ecology of lichens of Katmai and Lake Clark National Parks and Preserves, Alaska. *Mycosphere* **9**, 859–930.
- McGee M (2002) Back cover. *Bulletin of the California Lichen Society* **9**, 25–26.
- Meeks JC (1998) Symbiosis between nitrogen-fixing cyanobacteria and plants. *BioScience* **48**, 266–276.
- Meier FA, Scherrer S and Honegger R (2002) Faecal pellets of lichenivorous mites contain viable cells of the lichen-forming ascomycete *Xanthoria parietina* and its green algal photobiont, *Trebouxia arboricola*. *Biological Journal of the Linnean Society* **76**, 259–268.
- Meier JL and Chapman RL (1983) Ultrastructure of the lichen *Coenogonium interplexum* Nyl. *American Journal of Botany* **70**, 400–407.
- Metz S, Singer D, Domaizon I, Unrein F and Lara E (2019) Global distribution of *Trebouxiophyceae* diversity explored by high-throughput sequencing and phylogenetic approaches. *Environmental Microbiology* **21**, 3885–3895.
- Miadlikowska J, Kauff F, Högnabba F, Oliver JC, Molnár K, Fraker E, Gaya E, Hafellner J, Hofstetter V, Gueidan C, et al. (2014) A multigene phylogenetic synthesis for the class *Lecanoromycetes* (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. *Molecular Phylogenetics and Evolution* **79**, 132–168.
- Míguez F, Schiefelbein U, Karsten U, García-Plazaola JI and Gustavs L (2017) Unraveling the photoprotective response of lichenized and free-living green algae (Trebouxiophyceae, Chlorophyta) to photochilling stress. *Frontiers in Plant Science* **8**, 1144.
- Mikhailyuk TI, Sluiman HJ, Massalski A, Mudimu O, Demchenko EM, Kondratyuk SY and Friedl T (2008) New streptophyte green algae from terrestrial habitats and an assessment of the genus *Interfilum* (Klebsormidiophyceae, Streptophyta). *Journal of Phycology* **44**, 1586–1603.
- Mikhailyuk T, Holzinger A, Tsarenko P, Glaser K, Demchenko E and Karsten U (2020) *Dictyosphaerium*-like morphotype in terrestrial algae: what is *Xerochlorella* (Trebouxiophyceae, Chlorophyta)? *Journal of Phycology* **56**, 671–686.
- Millbank JW and Kershaw KA (1974 [1973]) Nitrogen metabolism. In Ahmadjian V and Hale ME (eds), *The Lichens*. New York: Academic Press, pp. 289–307.
- Moe RL (1997) *Verrucaria tavaresiae* sp. nov., a marine lichen with a brown algal photobiont. *Bulletin of the California Lichen Society* **4**, 7–11.
- Mohr F, Ekman S and Heegaard E (2004) Evolution and taxonomy of the marine *Collembosporium* species (lichenized Ascomycota) in north-west Europe. *Mycological Research* **108**, 515–532.
- Molinari-Novoqa EA (2016) *Uvulifera*, a new generic name for *Coccombortry* (Chaetophoraceae). *Notulae Algarum* **5**, 1–2.
- Molins A, García-Breijo F, Reig-Armiñana J, del Campo E, Casano L and Barreno E (2013) Coexistence of different intrathalline symbiotic algae and bacterial biofilms in the foliose Canarian lichen *Parmotrema pseudotinctorum*. *Vieraea* **41**, 249–270.
- Molins A, Moya P, García-Breijo F, Reig-Armiñana J and Barreno E (2018) Molecular and morphological diversity of *Trebouxia* microalgae in sphaerothalloid *Circinaria* spp. lichens. *Journal of Phycology* **54**, 494–504.

- Molins A, Chiva S, Calatayud A, Marco F, García-Breijo F, Reig-Armiñana J, Carrasco P and Moya P (2020) Multidisciplinary approach to describe *Trebouxia* diversity within lichenized fungi *Buellia zoharyi* from the Canary Islands. *Symbiosis* **82**, 19–34.
- Moncada B, Coca LF and Lücking R (2013) Neotropical members of *Sticta* (lichenized Ascomycota: Lobariaceae) forming photosymbiodemes, with the description of seven new species. *Bryologist* **116**, 169–200.
- Moniz MBJ, Rindi F and Guiry MD (2012) Phylogeny and taxonomy of *Prasiolales* (Trebouxiophyceae, Chlorophyta) from Tasmania, including *Rosenvingiella tasmanica* sp. nov. *Phycologia* **51**, 86–97.
- Moniz MBJ, Guiry MD and Rindi F (2014) TufA phylogeny and species boundaries in the green algal order *Prasiolales* (Trebouxiophyceae, Chlorophyta). *Phycologia* **53**, 396–406.
- Moya P, Škaloud P, Chiva S, García-Breijo FJ, Reig-Armiñana J, Vančurová L and Barreno E (2015) Molecular phylogeny and ultrastructure of the lichen microalga *Asterochloris mediterranea* sp. nov. from Mediterranean and Canary Islands ecosystems. *International Journal of Systematic and Evolutionary Microbiology* **65**, 1838–1854.
- Moya P, Molins A, Martínez-Alberola F, Muggia L and Barreno E (2017) Unexpected associated microalgal diversity in the lichen *Ramalina farinacea* is uncovered by pyrosequencing analyses. *PLoS ONE* **12**, e0175091.
- Moya P, Chiva S, Molins A, Jadrná I, Škaloud P, Peksa O and Barreno E (2018) *Myrmecia israelensis* as the primary symbiotic microalga in squamulose lichens growing in European and Canary Island terricolous communities. *Fottea* **18**, 72–85.
- Muggia L and Grube M (2018) Fungal diversity in lichens: from extremotolerance to interactions with algae. *Life* **8**, 15.
- Muggia L, Grube M and Tretiach M (2008) Genetic diversity and photobiont association in selected taxa of the *Tephromela atra* group (Lecanorales, lichenized Ascomycota). *Mycological Progress* **7**, 147–160.
- Muggia L, Zellnig G, Rabensteiner J and Grube M (2010) Morphological and phylogenetic study of algal partners associated with the lichen-forming fungus *Tephromela atra* from the Mediterranean region. *Symbiosis* **51**, 149–160.
- Muggia L, Baloch E, Stabentheiner E, Grube M and Wedin M (2011) Photobiont association and genetic diversity of the optionally lichenized fungus *Schizoxylon albescens*. *FEMS Microbiology Ecology* **75**, 255–272.
- Muggia L, Vancurova L, Škaloud P, Peksa O, Wedin M and Grube M (2013) The symbiotic playground of lichen thalli – a highly flexible photobiont association in rock-inhabiting lichens. *FEMS Microbiology Ecology* **85**, 313–323.
- Muggia L, Pérez-Ortega S, Kopun T, Zellnig G and Grube M (2014) Photobiont selectivity leads to ecological tolerance and evolutionary divergence in a polymorphic complex of lichenized fungi. *Annals of Botany* **114**, 463–475.
- Muggia L, Candotto-Carniel F and Grube M (2017) The lichen photobiont *Trebouxia*: towards and appreciation of species diversity and molecular studies. In Grube M, Seckbach J and Muggia L (eds), *Algal and Cyanobacterial Symbioses*. London: World Scientific Publishing Europe Ltd, pp. 111–146.
- Muggia L, Leavitt S and Barreno E (2018) The hidden diversity of lichenized *Trebouxiophyceae* (Chlorophyta). *Phycologia* **57**, 503–524.
- Muggia L, Nelsen MP, Kirika PM, Barreno E, Beck A, Lindgren H, Lumbsch HT, Leavitt SD and *Trebouxia* working group (2020) Formally described species woefully underrepresent phylogenetic diversity in the common lichen photobiont genus *Trebouxia* (Trebouxiophyceae, Chlorophyta): an impetus for developing an integrated taxonomy. *Molecular Phylogenetics and Evolution* **149**, 106821.
- Mukherjee R, Borah SP and Goswami BC (2010) Biochemical characterization of carotenoids in two species of *Trentepohlia* (Trentepohliales, Chlorophyta). *Journal of Applied Phycology* **22**, 569–571.
- Mukhtar A, Garty J and Galun M (1994) Does the lichen alga *Trebouxia* occur free-living in nature: further immunological evidence. *Symbiosis* **17**, 247–253.
- Myllys L, Stenroos S, Thell A and Kuusinen M (2007) High cyanobiont selectivity of epiphytic lichens in old growth boreal forest of Finland. *New Phytologist* **173**, 621–629.
- Nakano T (1988) Phycobionts of some Japanese species of the *Graphidaceae*. *Lichenologist* **20**, 353–360.
- Nakano T and Iguchi K (1994) Photobionts isolated from some Japanese species of *Cladonia* (lichens). *Symbiosis* **17**, 65–73.
- Nakano T and Ihda T (1996) The identity of photobionts from the lichen *Pyrenula japonica*. *Lichenologist* **28**, 437–442.
- Namba N and Nakayama T (2021) Taxonomic study of a new green alga, *Annulotesta cochlephila* gen. et sp. nov. (Kornmanniaceae, Ulvales, Ulvophyceae), growing on the shells of door snails. *Journal of Plant Research* **134**, 77–89.
- Nash, TH III, Kappen L, Lösch R, Larson DW and Matthes-Sears U (1987) Cold resistance of lichens with *Trentepohlia*- or *Trebouxia*-photobionts from the North American west coast. *Flora* **170**, 241–251.
- Nelsen MP and Gargas A (2008) Dissociation and horizontal transmission of codispersing lichen symbionts in the genus *Lepraria* (Lecanorales: Stereocaulaceae). *New Phytologist* **177**, 264–275.
- Nelsen MP and Gargas A (2009) Symbiont flexibility in *Thamnolia vermicularis* (Pertusariales: Icmadophilaceae). *Bryologist* **104**, 404–417.
- Nelsen MP, Rivas Plata E, Andrew CJ, Lücking R and Lumbsch HT (2011) Phylogenetic diversity of trentepohlialean algae associated with lichen-forming fungi. *Journal of Phycology* **47**, 282–290.
- Nelson DR, Khraiweh B, Fu W, Alseekh S, Jaiswal A, Chaiboonchoe A, Hazzouri KM, O'Connor MJ, Butterfoss GL, Dro N, et al. (2017) The genome and phenome of the green alga *Chloroidium* sp. UTEX 3007 reveal adaptive traits for desert acclimatization. *eLife* **6**, e25783.
- Němcová Y and Kalina T (2000) Cell wall development, microfibril and pyrenoid structure in type strains of *Chlorella vulgaris*, *C. kessleri*, *C. sorokiniana* compared with *C. luteoviridis* (Trebouxiophyceae, Chlorophyta). *Algological Studies* **100**, 95–105.
- Neustupa J (2015) Division *Chlorophyta*. In Frey W (series ed.), *Syllabus of Plant Families: 2/1 Photoautotrophic Eukaryotic Algae*. Stuttgart: Borntraeger Verlagsbuchhandlung, pp. 191–247.
- Neustupa J and Štíferová S (2013) Distribution patterns of subaerial corticolous microalgae in two European regions. *Plant Ecology and Evolution* **146**, 279–289.
- Neustupa J, Němcová Y, Eliáš M and Škaloud P (2009) *Kalinella bambusicola* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel coccoid *Chlorella*-like subaerial alga from Southeast Asia. *Phycological Research* **57**, 159–169.
- Nyati S, Beck A and Honegger R (2007) Fine structure and phylogeny of green algal photobionts in the microfilamentous genus *Psoroglaena* (Verrucariaceae, lichen-forming ascomycetes). *Plant Biology* **9**, 390–399.
- Nyati S, Bhattacharya D, Werth S and Honegger R (2013) Phylogenetic analysis of LSU and SSU rDNA group I introns of lichen photobionts associated with the genera *Xanthoria* and *Xanthomendoza* (Teloschistaceae, lichenized Ascomycetes). *Journal of Phycology* **49**, 1154–1166.
- Nyati S, Scherrer S, Werth S and Honegger R (2014) Green-algal photobiont diversity (*Trebouxia* spp.) in representatives of *Teloschistaceae* (Lecanoromycetes, lichen-forming ascomycetes). *Lichenologist* **46**, 189–212.
- O'Brien H, Miadlikowska J and Lutzoni F (2005) Assessing host specialization in symbiotic cyanobacteria associated with four closely related species of the lichen fungus *Peltigera*. *European Journal of Phycology* **40**, 363–378.
- O'Brien H, Miadlikowska J and Lutzoni F (2013) Assessing population structure and host specialization in lichenized cyanobacteria. *New Phytologist* **198**, 557–566.
- Oberwinkler F (1980) Symbiotic relationship between fungus and alga in basidiolichens. In Schwemmler W and Schenk HEA (eds), *Endocytobiology: Endosymbiosis and Cell Biology, Vol. I*. Berlin: Walter de Gruyter, pp. 305–315.
- Oberwinkler F (1984) Fungus-alga interactions in basidiolichens. In Hertel H and Oberwinkler F (eds), *Festschrift Josef Poelt: Beiheft Beiträge zur Lichenologie*. Vaduz: J. Cramer, pp. 739–774.
- Oberwinkler F (2012) Basidiolichens. In Hock B (ed.), *The Mycota IX. Fungal Associations*, 2nd Edn. Berlin: Springer-Verlag, pp. 341–362.
- Ohmura Y, Kawachi M, Kasai F, Watanabe MM and Takeshita S (2006) Genetic combinations of symbionts in a vegetatively reproducing lichen, *Parmotrema tinctorum*, based on ITS rDNA sequences. *Bryologist* **109**, 43–59.
- Ohmura Y, Takeshita S and Kawachi M (2019) Photobiont diversity within populations of a vegetatively reproducing lichen, *Parmotrema tinctorum*, can be generated by photobiont switching. *Symbiosis* **77**, 59–72.
- Oksanen I, Lohtander K, Paulsrud P and Rikkinen J (2002) A molecular approach to cyanobacterial diversity in a rock-pool community involving

- gelatinous lichens and free-living *Nostoc* colonies. *Annales Botanici Fennici* **39**, 93–99.
- Onuț-Brännström I, Tibell L and Johannesson H** (2017) A worldwide phylogeography of the whiteworm lichens *Thamnolia* reveals three lineages with distinct habitats and evolutionary histories. *Ecology and Evolution* **7**, 3602–3615.
- Onuț-Brännström I, Benjamin M, Scofield DG, Heidmarsson S, Andersson MGI, Lindström ES and Johannesson H** (2018) Sharing of photobionts in sympatric populations of *Thamnolia* and *Cetraria* lichens: evidence from high-throughput sequencing. *Scientific Reports* **8**, 4406.
- Ortiz-Álvarez R, de los Ríos A, Fernández-Mendoza F, Torralba-Burrial A and Pérez-Ortega S** (2015) Ecological specialization of two photobiont-specific maritime cyanolichen species of the genus *Lichina*. *PLoS ONE* **10**, e0132718.
- Oszycza P, Lenart-Borón A, Borón P and Rola K** (2021) Lichen-forming fungi in postindustrial habitats involve alternative photobionts. *Mycologia* **113**, 43–55.
- Otálora MAG, Martínez I, O'Brien H, Molina MC, Aragón G and Lutzoni F** (2010) Multiple origins of high reciprocal symbiotic specificity at an inter-continental spatial scale among gelatinous lichens (*Collema* spp., *Lecanoromycetes*). *Molecular Phylogenetics and Evolution* **56**, 1089–1095.
- Otálora MAG, Salvador C, Martínez I and Aragón G** (2013) Does the reproductive strategy affect the transmission and genetic diversity of bionts in cyanolichens? A case study using two closely related species. *Microbial Ecology* **65**, 517–530.
- Ott S** (1988) Photosymbiodemes and their development in *Peltigera venosa*. *Lichenologist* **20**, 361–368.
- Palmqvist K, de los Ríos A, Ascaso C and Samuelsson G** (1997) Photosynthetic carbon acquisition in the lichen photobionts *Coccomyxa* and *Trebouxia* (*Chlorophyta*). *Physiologia Plantarum* **101**, 67–76.
- Paran N, Ben-Shaul Y and Galun M** (1971) Fine structure of the blue-green phycobiont and its relation to the mycobiont in two *Gonohymenia* lichens. *Archiv für Mikrobiologie* **76**, 103–113.
- Park CH, Kim KM, Elvebakk A, Kim O-S, Jeong G and Hong SG** (2015) Algal and fungal diversity in Antarctic lichens. *Journal of Eukaryotic Microbiology* **62**, 196–205.
- Parra OO and Redón J** (1977) Aislamiento de *Heterococcus caespitosus* Vischer ficobionte de *Verrucaria maura*. *Boletín de la Sociedad de Biología de Concepción* **51**, 219–224.
- Paul F, Otte J, Schmitt I and Dal Grande F** (2018) Comparing Sanger sequencing and high-throughput metabarcoding for inferring photobiont diversity in lichens. *Scientific Reports* **8**, 8624.
- Paulsrud P** (2001) The *Nostoc* symbiosis in lichens: diversity, specificity and cellular modifications. *Acta Universitatis Upsaliensis: Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science and Technology* **662**, 1–55.
- Paulsrud P and Lindblad P** (1998) Sequence variation of the tRNA^{Leu} intron as a marker for genetic diversity and specificity of symbiotic cyanobacteria in some lichens. *Applied and Environmental Microbiology* **64**, 310–315.
- Paulsrud P, Rikkinen J and Lindblad P** (1998) Cyanobiont specificity in some *Nostoc*-containing lichens and in a *Peltigera aphthosa* photosymbiodeme. *New Phytologist* **139**, 517–524.
- Paulsrud P, Rikkinen J and Lindblad P** (2000) Spatial patterns of photobiont diversity in some *Nostoc*-containing lichens. *New Phytologist* **146**, 291–299.
- Paulsrud P, Rikkinen J and Lindblad P** (2001) Field investigations on cyanobacterial specificity in *Peltigera aphthosa*. *New Phytologist* **152**, 117–123.
- Peksa O and Škaloud P** (2011) Do photobionts influence the ecology of lichens? A case study of environmental preferences in symbiotic green alga *Asterochloris* (*Trebouxiophyceae*). *Molecular Ecology* **20**, 3936–3948.
- Pereira Riquelme I** (1992) *Flora, vegetación y ecología de los líquenes acuáticos de España*. Ph.D. thesis, University of Barcelona.
- Pérez-Ortega S, de los Ríos A, Crespo A and Sancho LG** (2010) Symbiotic lifestyle and phylogenetic relationships of the bionts of *Mastodia tessellata* (*Ascomycota, incertae sedis*). *American Journal of Botany* **97**, 738–752.
- Pérez-Ortega S, Ortiz-Álvarez R, Green TGA and de los Ríos A** (2012) Lichen myco- and photobiont diversity and their relationships at the edge of life (McMurdo Dry Valleys, Antarctica). *FEMS Microbiology Ecology* **82**, 429–448.
- Pérez-Ortega S, Miller KA and de los Ríos A** (2018) Challenging the lichen concept: *Turgidisculum ulvae* (*Verrucariaceae*) represents an independent photobiont shift to a multicellular blade-like alga. *Lichenologist* **50**, 341–356.
- Peršoh D, Beck A and Rambold G** (2004) The distribution of ascus types and photobiontal selection in *Lecanoromycetes* (*Ascomycota*) against the background of a revised SSU nrDNA phylogeny. *Mycological Progress* **3**, 103–121.
- Peters AF and Moe RL** (2001) DNA sequences confirm that *Petroderma maculiforme* (*Phaeophyceae*) is the brown algal phycobiont of the marine lichen *Verrucaria tavaresiae* (*Verrucariales, Ascomycota*) from central California. *Bulletin of the California Lichen Society* **8**, 41–43.
- Peveling E** (1968) Pyrenoidstrukturen in symbiotisch lebenden *Trebouxia*-Arten. *Zeitschrift für Pflanzenphysiologie* **59**, 393–396.
- Peveling E** (1969) Elektronenoptische Untersuchungen an Flechten. II. Die Feinstruktur von *Trebouxia*-Phycobionten. *Planta* **87**, 69–85.
- Peveling E and Galun M** (1976) Electron-microscopical studies on the phycobiont *Coccomyxa* Schmidle. *New Phytologist* **77**, 713–718.
- Piercey-Normore MD** (2004) Selection of algal genotypes by three species of lichen fungi in the genus *Cladonia*. *Canadian Journal of Botany* **82**, 947–961.
- Piercey-Normore MD** (2006) The lichen-forming ascomycete *Evernia mesomorpha* associates with multiple genotypes of *Trebouxia jamesii*. *New Phytologist* **169**, 331–344.
- Piercey-Normore MD** (2009) Vegetatively reproducing fungi in three genera of the *Parmeliaceae* share divergent algal partners. *Bryologist* **112**, 773–785.
- Piercey-Normore MD and DePriest PT** (2001) Algal switching among lichen symbioses. *American Journal of Botany* **88**, 1490–1498.
- Pino-Bodas R and Stenroos A** (2020) Global biodiversity patterns of the photobionts associated with the genus *Cladonia* (*Lecanorales, Ascomycota*). *Microbial Ecology*, doi:10.1007/s00248-020-01633-3
- Plessl A** (1963) Über die Beziehung von Haustorientypus und Organisationshöhe bei Flechten. *Österreichische Botanische Zeitschrift* **110**, 194–269.
- Poelt J** (1958) Über parasitische Flechten. II. *Planta* **51**, 288–307.
- Poelt J and Baumgärtner H** (1964) Über Rhizinenstränge bei placodialen Flechten. *Österreichische Botanische Zeitschrift* **111**, 1–18.
- Prieto M, Schultz M, Olariaga I and Wedin M** (2019) *Lichinodium* is a new lichenized lineage in the *Leotiomyces*. *Fungal Diversity* **94**, 23–39.
- Pröschold T and Darienko T** (2020) The green puzzle *Stichococcus* (*Trebouxiophyceae, Chlorophyta*): new generic and species concept among this widely distributed genus. *Phytotaxa* **441**, 113–142.
- Pröschold T, Darienko T, Silva P, Reisser W and Krienitz L** (2011) The systematics of *Zoochlorella* revisited employing a molecular approach. *Environmental Microbiology* **13**, 350–364.
- Purvis OW, Coppins BJ, Hawksworth DL, James PW and Moore DM** (1992) *The Lichen Flora of Great Britain and Ireland*. London: British Lichen Society.
- Raggio J, Green TGA, Crittenden PD, Pintado A, Vivas M, Pérez-Ortega S, de los Ríos A and Sancho LG** (2012) Comparative ecology of three *Placopsis* species, pioneer lichens in recently exposed Chilean glacial forelands. *Symbiosis* **56**, 55–66.
- Rajaniemi P, Hrouzek P, Kaštovská K, Willame R, Rantala A, Hoffmann L, Komárek J and Sivonen K** (2005) Phylogenetic and morphological evaluation of the genera *Anabaena*, *Aphanizomenon*, *Trichormus* and *Nostoc* (*Nostocales, Cyanobacteria*). *International Journal of Systematic and Evolutionary Microbiology* **55**, 11–26.
- Rambold G, Friedl T and Beck A** (1998) Photobionts in lichens: possible indicators of phylogenetic relationships? *Bryologist* **101**, 392–397.
- Ran L, Larsson J, Vigil-Stenman T, Nylander JAA, Ininbergs K, Zheng W-W, Lapidus A, Lowry S, Haselkorn R and Bergman B** (2010) Genome erosion in a nitrogen-fixing vertically transmitted endosymbiotic multicellular cyanobacterium. *PLoS ONE* **5**, e11486.
- Richardson DHS, Hill DJ and Smith DC** (1968) Lichen physiology. XI. The role of the alga in determining the pattern of carbohydrate movement between lichen symbionts. *New Phytologist* **67**, 469–486.
- Řídká T, Peksa O, Rai H, Upreti DK and Škaloud P** (2014) Photobiont diversity in Indian *Cladonia* lichens, with special emphasis on the geographical patterns. In Rai H and Upreti DK (eds), *Terricolous Lichens in India*. New York: Springer, pp. 53–71.
- Rikkinen J** (2003) Ecological and evolutionary role of photobiont-mediated guilds in lichens. *Symbiosis* **34**, 99–110.

- Rikkinen J** (2013) Molecular studies on cyanobacterial diversity in lichen symbiosis. *MycKeys* **6**, 3–32.
- Rikkinen J** (2017) Symbiotic cyanobacteria in lichens. In Grube M, Seckbach J and Muggia L (eds), *Algal and Cyanobacterial Symbioses*. London: World Scientific Publishing Europe Ltd, pp. 147–167.
- Rikkinen J, Oksanen I and Lohtander K** (2002) Lichen guilds share related cyanobacterial symbionts. *Science* **297**, 357.
- Rindi F and Guiry MD** (2002) Diversity, life history, and ecology of *Trentepohlia* and *Printzina* (Trentepohliales, Chlorophyta) in urban habitats in western Ireland. *Journal of Phycology* **38**, 39–54.
- Rindi F, Menéndez JL, Guiry MD and Rico JM** (2004) The taxonomy and distribution of *Phycopeltis* (Trentepohliaceae, Chlorophyta) in Europe. *Cryptogamie, Algologie* **25**, 3–17.
- Rindi F, Lam DW and López-Bautista JM** (2009) Phylogenetic relationships and species circumscription in *Trentepohlia* and *Printzina* (Trentepohliales, Chlorophyta). *Molecular Phylogenetics and Evolution* **52**, 329–339.
- Rindi F, Mikhailyuk TI, Sluiman HJ, Friedl T and López-Bautista JM** (2011) Phylogenetic relationships in *Interfilum* and *Klebsormidium* (Klebsormidiophyceae, Streptophyta). *Molecular Phylogenetics and Evolution* **58**, 218–231.
- Rodgers GA and Stewart WDP** (1977) The cyanophyte-hepatic symbiosis. *New Phytologist* **78**, 441–458.
- Rola K, Lenart-Borón A, Borón P and Osyczka P** (2021) Heavy-metal pollution induces changes in the genetic composition and anatomical properties of photobionts in pioneer lichens colonizing post-industrial habitats. *Science of the Total Environment* **750**, 141439.
- Romeike J, Friedl T, Helm G and Ott S** (2002) Genetic diversity of algal and fungal partners in four species of *Umbilicaria* (lichenized ascomycetes) along a transect of the Antarctic Peninsula. *Molecular Biology and Evolution* **19**, 1209–1217.
- Roskin PA** (1970) Ultrastructure of the host-parasite interaction in the basidiolichen *Cora pavonia* (Web.) E. Fries. *Archiv für Mikrobiologie* **70**, 176–182.
- Ruprecht U, Brunauer G and Printzen C** (2012) Genetic diversity of photobionts in Antarctic lecodeoid lichens from an ecological viewpoint. *Lichenologist* **44**, 661–678.
- Ruprecht U, Brunauer G and Türk R** (2014) High photobiont diversity in the common European soil crust lichen *Psora decipiens*. *Biodiversity and Conservation* **23**, 1771–1785.
- Ruprecht U, Fernández-Mendoza F, Türk R and Fryday AM** (2020) High levels of endemism and local differentiation in the fungal and algal symbionts of saxicolous lecodeoid lichens along a latitudinal gradient in southern South America. *Lichenologist* **52**, 287–303.
- Sadowska-Deś AD, Dal Grande F, Lumsch HT, Beck A, Otte J, Hur J-S, Kim JA and Schmitt I** (2014) Integrating coalescent and phylogenetic approaches to delimit species in the lichen photobiont *Trebouxia*. *Molecular Phylogenetics and Evolution* **76**, 202–210.
- Sadowsky A and Ott S** (2016) Symbiosis as a successful strategy in continental Antarctica: performance and protection of *Trebouxia* photosystem II in relation to lichen pigmentation. *Polar Biology* **39**, 139–151.
- Saini KC, Nayaka S and Bast F** (2019) Diversity of lichen photobionts: their coevolution and bioprospecting potential. In Satyanarayana T, Johri BN and Das SK (eds), *Microbial Diversity in Ecosystem Sustainability and Biotechnological Applications*. Singapore: Springer Singapore, pp. 307–323.
- Sanders WB** (1994) Role of lichen rhizomorphs in thallus propagation and substrate colonization. *Cryptogamic Botany* **4**, 283–289.
- Sanders WB** (2001) Composite thalli of *Sticta* sp. from Brazil with morphologically similar lobes containing either a chlorobiont or a cyanobiont layer. *Symbiosis* **31**, 47–55.
- Sanders WB** (2002) In situ development of the foliicolous lichen *Phyllophiale* (Trichotheliaceae) from propagule germination to propagule production. *American Journal of Botany* **89**, 1741–1746.
- Sanders WB** (2005) Observing microscopic phases of lichen life cycles on transparent substrata placed *in situ*. *Lichenologist* **37**, 373–382.
- Sanders WB** (2014) Complete life cycle of the lichen fungus *Calopadia puigarii* (Pilocarpaceae, Ascomycetes) documented *in situ*: propagule dispersal, establishment of symbiosis, thallus development, and formation of sexual and asexual reproductive structures. *American Journal of Botany* **101**, 1836–1848.
- Sanders WB and de los Ríos A** (2015) Structure and *in situ* development of the microlichen *Gyalectidium paolae* (Gomphillaceae, Ascomycota), an overlooked colonist on palm leaves in southwest Florida. *American Journal of Botany* **102**, 1403–1412.
- Sanders WB and Lücking R** (2002) Reproductive strategies, relichenization and thallus development observed *in situ* in leaf-dwelling lichen communities. *New Phytologist* **155**, 425–435.
- Sanders WB and Rico VJ** (1992) Lichenizing rhizomorphs and thallus development in the squamulose lichen *Aspicilia crespiana* Rico ined. (Lecanorales, Ascomycetes). *Botanica Acta* **105**, 449–456.
- Sanders WB, Wierzychos J and Ascaso C** (1994) Physical interactions of two rhizomorph-forming lichens with their rock substrate. *Acta Botanica* **107**, 432–439.
- Sanders WB, Moe RL and Ascaso C** (2004) The intertidal marine lichen formed by the pyrenomycete fungus *Verrucaria tavaresiae* and the brown alga *Petroderma maculiforme* (Phaeophyceae): thallus organization and symbiont interaction. *American Journal of Botany* **91**, 511–522.
- Sanders WB, Moe RL and Ascaso C** (2005) Ultrastructural study of the brown alga *Petroderma maculiforme* (Phaeophyceae) in the free-living state and in lichen symbiosis with the intertidal marine fungus *Verrucaria tavaresiae* (Ascomycotina). *European Journal of Phycology* **40**, 353–361.
- Sanders WB, Pérez-Ortega S, Nelsen MP, Lücking R and de los Ríos A** (2016) *Heveochlorella* (Trebouxiophyceae): a little-known genus of unicellular green algae outside the *Trebouxiiales* emerges unexpectedly as a major clade of lichen photobionts in foliicolous communities. *Journal of Phycology* **52**, 840–853.
- Santesson R** (1952) Follicolous lichens I. A revision of the taxonomy of the obligately foliicolous, lichenized fungi. *Symbolae Botanicae Upsaliensis* **12**, 1–590.
- Schaper T and Ott S** (2003) Photobiont selectivity and interspecific interactions in lichen communities. I. Culture experiments with the mycobiont *Fulgensia bracteata*. *Plant Biology* **5**, 441–450.
- Scheidegger C** (1995) Reproductive strategies in *Vezeadaea* (Lecanorales, lichenized Ascomycetes): a low-temperature scanning electron microscope study of a ruderal species. *Cryptogamic Botany* **5**, 163–171.
- Schmitt I and Lumsch HT** (2001) Identification of the photobionts in *Trapeliopsis* and *Pertusaria* using SSU ribosomal DNA sequences obtained from PCR amplification with a non-green-algal primer. *Mycotaxon* **78**, 407–411.
- Schüßler A** (2012) The *Geosiphon-Nostoc* endosymbiosis and its role as a model for arbuscular mycorrhiza research. In Hock B (ed.), *The Mycota IX. Fungal Associations*, 2nd Edn. Berlin: Springer-Verlag, pp. 77–91.
- Schuster G** (1992) Development of adventive thalli in *Umbilicaria* Hoffm. *Flora* **187**, 201–207.
- Schuster G, Ott S and Jahns HM** (1985) Artificial culture of lichens in the natural environment. *Lichenologist* **17**, 147–153.
- Schwendener S** (1869) *Die Algentypen der Flechtengonidien*. Basel: Universitätsbuchdruckerei von C. Schultze.
- Scott GD** (1957) Lichen terminology. *Nature* **179**, 486–487.
- Scott GD** (1960) Studies of the lichen symbiosis I. The relationship between nutrition and moisture content in the maintenance of the symbiotic state. *New Phytologist* **59**, 374–381.
- Seto K, Matsuzawa T, Kuno H and Kagami M** (2020) Morphology, ultrastructure, and molecular phylogeny of *Aphelidium collabens* sp. nov. (*Aphelida*), a parasitoid of a green alga *Coccomyxa* sp. *Protist* **171**, 125728.
- Silva PC** (1979) Review of the taxonomic history and nomenclature of the yellow-green algae. *Archiv für Protistenkunde* **121**, 20–63.
- Simon A, Goffinet B, Magain N and Sérusiaux E** (2018) High diversity, high insular endemism and recent origin in the lichen genus *Sticta* (lichenized Ascomycota, Peltigerales) in Madagascar and the Mascarenes. *Molecular Phylogenetics and Evolution* **122**, 15–28.
- Singh G, Dal Grande F, Divakar PK, Otte J, Crespo A and Schmitt I** (2017) Fungal-algal association patterns in lichen symbiosis linked to macroclimate. *New Phytologist* **214**, 317–329.
- Škaloud P and Peksa O** (2010) Evolutionary inferences based on ITS rDNA and actin sequences reveal extensive diversity of the common lichen alga *Asterochloris* (Trebouxiophyceae, Chlorophyta). *Molecular Phylogenetics and Evolution* **54**, 36–46.

- Škaloud P, Neustupa J, Radochová B and Kubínová L (2005) Confocal microscopy of chloroplast morphology and ontogeny in three strains of *Dictyochloropsis* (Trebouxiophyceae, Chlorophyta). *Phycologia* **44**, 261–269.
- Škaloud P, Steinová J, Řídká T and Vančurová L (2015) Assembling the challenging puzzle of algal biodiversity: species delimitation within the genus *Asterochloris* (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* **51**, 507–527.
- Škaloud P, Friedl T, Hallmann C, Beck A and Dal Grande F (2016) Taxonomic revision and species delimitation of coccoid green algae currently assigned to the genus *Dictyochloropsis* (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* **52**, 599–617.
- Škaloud P, Moya P, Molins A, Peksa O, Santos-Guerra A and Barreno E (2018) Untangling the hidden intrathalline microalgal diversity in *Parmotrema pseudotinctorum*: *Trebouxia crespoana* sp. nov. *Lichenologist* **50**, 357–369.
- Skuja H (1943) Ein Fall von fakultativer Symbiose zwischen operculatem Discomycet und einer Chlamydomonade. *Archiv für Protistenkunde* **96**, 365–376.
- Slocum RD (1980) Light and electron microscopic investigations in the *Dictyonemataceae* (basidiolichens) II. *Dictyonema irpicinum*. *Canadian Journal of Botany* **58**, 1005–1015.
- Slocum RD, Ahmadjian V and Hildreth KC (1980) Zoosporogenesis in *Trebouxia gelatinosa*: ultrastructure potential for zoospore release and implications for the lichen association. *Lichenologist* **12**, 173–187.
- Smith DC (1974) Transport from symbiotic algae and symbiotic chloroplasts to host cells. *Symposia of the Society for Experimental Biology* **28**, 485–520.
- Smith DC (2019) What can lichens tell us about 'real fungi'? *Symbiosis* **77**, 93–98.
- Smith DC and Drew EA (1965) Studies in the physiology of lichens. V. Translocation from the algal layer to the medulla in *Peltigera polydactyla*. *New Phytologist* **64**, 195–200.
- Smith EC and Griffiths H (1996) The occurrence of the chloroplast pyrenoid is correlated with the activity of a CO₂-concentrating mechanism and carbon isotope discrimination in lichens and bryophytes. *Planta* **198**, 6–16.
- Smith H, Dal Grande F, Muggia L, Keuler R, Divakar PK, Grewe F, Schmitt I, Lumbsch HT and Leavitt SD (2020) Metagenomic data reveal diverse fungal and algal communities associated with the lichen symbiosis. *Symbiosis* **82**, 133–147.
- Solhaug KA and Gauslaa Y (1996) Parietin, a photoprotective secondary product of the lichen *Xanthoria parietina*. *Oecologia* **108**, 412–418.
- Solovchenko AE (2013) Physiology and adaptive significance of secondary carotenogenesis in green microalgae. *Russian Journal of Plant Physiology* **60**, 1–13.
- Spribille T, Tønsberg T, Stabentheiner E and Muggia L (2014) Reassessing evolutionary relationships in the filamentous cyanolichen genus *Spilonema* (Peltigerales, Lecanoromycetes). *Lichenologist* **46**, 373–388.
- Spribille T, Tuovinen V, Resl P, Vanderpool D, Wolinski H, Aime MC, Schneider K, Stabentheiner E, Toome-Heller M, Thor G, et al. (2016) Basidiomycete yeasts in the cortex of ascomycete macrolichens. *Science* **353**, 488–492.
- Stahl E (1877) *Beiträge zur Entwicklungsgeschichte der Flechten. II. Ueber die Bedeutung der Hymenialgonidien*. Leipzig: Arthur Felix.
- Steinová J, Škaloud P, Yahr R, Bestová H and Muggia L (2019) Reproductive and dispersal strategies shape the diversity of mycobiont-photobiont association in *Cladonia* lichens. *Molecular Phylogenetics and Evolution* **134**, 226–237.
- Stenroos S, Stocker-Wörgötter E, Yoshimura I, Myllys L, Thell A and Hyvönen J (2003) Culture experiments and DNA sequence data confirm the identity of *Lobaria* photomorphs. *Canadian Journal of Botany* **81**, 232–247.
- Stenroos S, Högnabba F, Myllys L, Hyvönen J and Thell A (2006) High selectivity in symbiotic associations of lichenized ascomycetes and cyanobacteria. *Cladistics* **22**, 230–238.
- Stevenson RN and South GR (1974) *Coccomyxa parasitica* sp. nov. (Coccomyxaceae, Chlorococcales), a parasite of giant scallops in Newfoundland. *British Phycological Journal* **9**, 319–329.
- Stocker-Wörgötter E (2001) Experimental lichenology and microbiology of lichens: culture experiments, secondary chemistry of cultured mycobionts, resynthesis, and thallus morphogenesis. *Bryologist* **104**, 576–581.
- Stocker-Wörgötter E and Hager A (2008) Culture methods for lichens and lichen symbionts. In Nash, TH III (ed.), *Lichen Biology*. New York: Cambridge University Press, pp. 353–363.
- Summerer M, Sonntag B and Sommaruga R (2008) Ciliate-symbiont specificity of freshwater endosymbiotic *Chlorella* (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* **44**, 77–84.
- Summerfield TC and Eaton-Rye JJ (2006) *Pseudocypbellaria crocata*, *P. neglecta* and *P. perpetua* from the Northern and Southern Hemispheres are a phylogenetic species and share cyanobionts. *New Phytologist* **170**, 597–607.
- Suto Y and Ohtani S (2009) Morphology and taxonomy of five *Cephaleuros* species (Trentepohliaceae, Chlorophyta) from Japan, including three new species. *Phycologia* **48**, 213–236.
- Suto Y and Ohtani S (2013) Seasonal development of five *Cephaleuros* species (Trentepohliaceae, Chlorophyta) on the leaves of woody plants and the behaviors of their gametes and zoospores. *Phycological Research* **61**, 105–115.
- Svenning MM, Eriksson T and Rasmussen U (2005) Phylogeny of symbiotic cyanobacteria within the genus *Nostoc* based on 16S rDNA sequence analyses. *Archives of Microbiology* **183**, 19–26.
- Syasina IG, Kukhlevsky AD, Kovaleva AL and Vaschenko MA (2012) Phylogenetic and morphological characterization of the green alga infesting the horse mussel *Modiolus* from Vityaz Bay (Peter the Great Bay, Sea of Japan). *Journal of Invertebrate Pathology* **111**, 175–181.
- Takeshita S (2001) A taxonomic revision of the genus *Trebouxia*. *Hikobia* **13**, 425–455.
- Takeshita S, Nakano T and Iwatsuki Z (1989) Phycobionts of some Japanese species of *Pertusaria* (Pertusariaceae). *Plant Systematics and Evolution* **165**, 49–54.
- Takeshita S, Tokizawa M, Handa S and Okamoto T (2010) A report of the photobiont isolated from *Multiclavula clara* (Berk. & Curt.) R. H. Peterson (Clavariaceae, lichenized Basidiomycetes). *Hikobia* **15**, 493–497. [In Japanese with English abstract and figure legends]
- Thomas EA (1939) Über die Biologie von Flechtenbildnern. *Beiträge zur Kryptogamenflora der Schweiz* **9**, 1–208 (+ 6 tab.).
- Thompson RH and Wujek DE (1992) *Printzina* gen. nov. (Trentepohliaceae), including a description of a new species. *Journal of Phycology* **28**, 232–237.
- Thompson RH and Wujek DE (1997) Trentepohliaceae: Cephaleuros, Phycopeltis and Stomatochroon. *Morphology Taxonomy and Ecology*. Enfield, New Hampshire: Science Publishers.
- Thüs H, Muggia L, Pérez-Ortega S, Favero-Longo SE, Joneson S, O'Brien H, Nelsen MP, Duque-Thüs R, Grube M, Friedl T, et al. (2011) Revisiting photobiont diversity in the lichen family Verrucariaceae (Ascomycota). *European Journal of Phycology* **46**, 399–415.
- Tibell L (2001) Photobiont association and molecular phylogeny of the lichen genus *Chaenotheca*. *Bryologist* **104**, 191–198.
- Tibell L and Beck A (2002) Morphological variation, photobiont association and ITS phylogeny of *Chaenotheca phaeocephala* and *C. subrosida* (Coniocybaceae, lichenized ascomycetes). *Nordic Journal of Botany* **21**, 651–660.
- Tønsberg T and Goward T (2001) *Sticta oroborealis* sp. nov., and other Pacific North American lichens forming dendriscoauloid cyanotypes. *Bryologist* **104**, 12–23.
- Tønsberg T and Holtan-Hartwig J (1983) Phycotype pairs in *Nephroma*, *Peltigera* and *Lobaria* in Norway. *Nordic Journal of Botany* **3**, 681–688.
- Trembley ML, Ringli C and Honegger R (2002a) Differential expression of hydrophobins *DGH1*, *DGH2* and *DGH3* and immunolocalization of *DGH1* in strata of the lichenized basidiocarp of *Dictyonema glabratum*. *New Phytologist* **154**, 185–195.
- Trembley ML, Ringli C and Honegger R (2002b) Morphological and molecular analysis of early stages in the resynthesis of the lichen *Baeomyces rufus*. *Mycological Research* **106**, 768–776.
- Trémouillaux-Guiller J and Huss VAR (2007) A cryptic intracellular green alga in *Ginkgo biloba*: ribosomal DNA markers reveal worldwide distribution. *Planta* **226**, 553–557.
- Trémouillaux-Guiller J, Rohr T, Rohr R and Huss VAR (2002) Discovery of an endophytic alga in *Ginkgo biloba*. *American Journal of Botany* **89**, 727–733.
- Tschaikner A, Ingolić E, Holzinger A and Gärtner G (2007) Phycobionts of some species of *Evernia* and *Ramalina*. *Herzogia* **20**, 53–60.
- Tschermak E (1941a) Untersuchungen über die Beziehung von Pilz und Alge im Flechtenthallus. *Österreichische Botanische Zeitschrift* **90**, 233–307.

- Tschermak L** (1941b) Beitrag zur Entwicklungsgeschichte und Morphologie der Protococcale *Trochiscia granulata*. *Österreichische Botanische Zeitschrift* **90**, 67–73.
- Tschermak-Woess E** (1953) Über wenig bekannte und neue Flechtengonidien III. Die Entwicklungsgeschichte von *Leptosira thrombii* nov. spec., der Gonidie von *Thrombium epigaeum*. *Österreichische Botanische Zeitschrift* **100**, 203–216.
- Tschermak-Woess E** (1970) Über wenig bekannte und neue Flechtengonidien V. Der Phycobiont von *Verrucaria aquatilis* und die Fortpflanzung von *Pseudopleurococcus arthopyreniae*. *Österreichische Botanische Zeitschrift* **118**, 443–455.
- Tschermak-Woess E** (1976) Algal taxonomy and the taxonomy of lichens: the phycobiont of *Verrucaria adriatica*. In Brown DH, Hawksworth DL and Bailey RL (eds), *Lichenology: Progress and Problems*. London: Academic Press, pp. 79–88.
- Tschermak-Woess E** (1978) *Myrmecia reticulata* as a phycobiont and free-living – free-living *Trebouxia* – the problem of *Stenocybe septata*. *Lichenologist* **10**, 69–79.
- Tschermak-Woess E** (1980a) *Asterochloris phycobiontica*, gen. et spec. nov., der Phycobiont der Flechte *Varicellaria carneonivea*. *Plant Systematics and Evolution* **135**, 279–294.
- Tschermak-Woess E** (1980b) *Elliptochloris bilobata*, gen. et spec. nov., der Phycobiont von *Catolechia wahlenbergii*. *Plant Systematics and Evolution* **136**, 63–72.
- Tschermak-Woess E** (1981) Zur Kenntnis der Phycobionten von *Lobaria limita* und *Normandina pulchella*. *Nova Hedwigia* **35**, 63–73.
- Tschermak-Woess E** (1983) Das Haustorialsystem von *Dictyonema* kennzeichnend für die Gattung. *Plant Systematics and Evolution* **143**, 109–115.
- Tschermak-Woess E** (1985) *Elliptochloris bilobata* kein ganz seltener Photobiont. *Herzogia* **7**, 105–116.
- Tschermak-Woess E** (1988a) The algal partner. In Galun M (ed.), *CRC Handbook of Lichenology*. Boca Raton, Florida: CRC Press, pp. 39–92.
- Tschermak-Woess E** (1988b) New and known taxa of *Chlorella* (*Chlorophyceae*): occurrence as lichen phycobionts and observations on living dictyosomes. *Plant Systematics and Evolution* **159**, 123–139.
- Tschermak-Woess E** (1989) Developmental studies in trebouxoid algae and taxonomical consequences. *Plant Systematics and Evolution* **164**, 161–195.
- Tschermak-Woess E** (1995) *Dictyochloropsis splendida* (*Chlorophyta*), the correct phycobiont of *Phlyctis argena* and the high degree of selectivity involved. *Lichenologist* **27**, 169–187.
- Tschermak-Woess E and Poelt J** (1976) *Vezdaea*, a peculiar lichen genus, and its phycobiont. In Brown DH, Hawksworth DL and Bailey RL (eds), *Lichenology: Progress and Problems*. London: Academic Press, pp. 89–105.
- Tschermak-Woess E, Bartlett J and Peveling E** (1983) *Lichenothrix riddlei* is an ascolichen and also occurs in New Zealand – light and electron microscopical investigation. *Plant Systematics and Evolution* **143**, 293–309.
- Tzovaras BG, Segers FHID, Bicker A, Dal Grande F, Otte J, Anvar SY, Hankeln T, Schmitt I and Ebersberger I** (2020) What is in *Umbilicaria pustulata*? A metagenomic approach to reconstruct the holo-genome of a lichen. *Genome Biology and Evolution* **12**, 309–324.
- Uher B** (2008) Spatial distribution of cyanobacteria and algae from the tombstone in a historic cemetery in Bratislava, Slovakia. *Fottea* **9**, 81–92.
- van den Hoek C, Mann DG and Jahns HM** (1995) *Algae: An Introduction to Phycology*. Cambridge: Cambridge University Press.
- Vančurová L** (2012) *Diverzita fotobiontů ve stélkách lišejníků rodu Stereocaulon [Photobiont diversity in the lichen genus Stereocaulon (Lecanoromycetes, Ascomycota)]*. Master's thesis, Charles University, Prague. [In Czech, with English summary].
- Vančurová L, Peksa O, Němcová Y and Škaloud P** (2015) *Vulcanochloris* (*Trebouxiales, Trebouxiophyceae*), a new genus of lichen photobiont from La Palma, Canary Islands, Spain. *Phytotaxa* **219**, 118–132.
- Vančurová L, Muggia L, Peksa O, Řídká T and Škaloud P** (2018) The complexity of symbiotic interactions influences the ecological amplitude of the host: a case study in *Stereocaulon* (lichenized *Ascomycota*). *Molecular Ecology* **27**, 3016–3033.
- Vargas Castillo R and Beck A** (2012) Photobiont selectivity and specificity in *Caloplaca* species in a fog-induced community in the Atacama Desert, northern Chile. *Fungal Biology* **116**, 665–676.
- Villanueva CD, Hašler P, Dvořák P, Pouličková A and Casamatta DA** (2018) *Brasilonema lichenoides* sp. nov. and *Chroococcidiopsis lichenoides* sp. nov. (*Cyanobacteria*): two novel cyanobacterial constituents isolated from a tripartite lichen of headstones. *Journal of Phycology* **54**, 224–233.
- Vischer W** (1960) Reproduktion und systematische Stellung einiger Rinden- und Bodenalgae. *Schweizerische Zeitschrift für Hydrologie* **22**, 330–349.
- Võ GTP** (2016) *Cyanobacterial lichenized fungi and their photobionts in Vietnam*. Ph.D. thesis, Technische Universität Kaiserslautern.
- Voytsekhovich A and Beck A** (2016) Lichen photobionts of the rocky outcrops of Karadag massif (Crimean Peninsula). *Symbiosis* **68**, 9–24.
- Voytsekhovich A, Dymytrova L and Nadyeina O** (2011) Photobiont composition of some taxa of the genera *Micarea* and *Placynthiella* (*Lecanoromycetes*, lichenized *Ascomycota*) from Ukraine. *Folia Cryptogamica Estonica* **48**, 135–148.
- Wagner J and Létrouit-Galinou M-A** (1988) Structure et ontogenèse du thalle squamuleux du lichen *Endocarpon pusillum*. *Canadian Journal of Botany* **66**, 2118–2127.
- Wagner M, Bathke AC, Cary CS, Green TGA, Junker RR, Trutschnig W and Ruprecht U** (2020) Myco- and photobiont associations in crustose lichens in the McMurdo Dry Valleys (Antarctica) reveal high differentiation along an elevational gradient. *Polar Biology* **43**, 1967–1983.
- Ward HM** (1884) On the structure, development and life history of a tropical epiphyllous lichen (*Strigula complanata* Fée, *vide* Rev. J. M. Crombie). *Transactions of the Linnean Society of London* **2**, 87–119.
- Warén H** (1920) Reinkultur von Flechtengonidien. *Finska Vetenskaps-Societetens Förhandlingar* **61**, 1–79.
- Watanabe S, Nakano T and Deguchi H** (1997) Photobionts isolated from maritime lichens. *Journal of Marine Biotechnology* **5**, 103–112.
- Waterbury JB and Stanier RY** (1978) Pattern of growth and development in pleurocapsalean cyanobacteria. *Microbiological Reviews* **42**, 2–44.
- Wedin M, Maier S, Fernández-Brime S, Cronholm B, Westberg M and Grube M** (2016) Microbiome change by symbiotic invasion in lichens. *Environmental Microbiology* **18**, 1428–1439.
- Werner R-G** (1931) Histoire de la synthèse liquéfique. *Mémoires de la Société des Science Naturelles du Maroc* **27**, 7–45.
- Werth S** (2010) Biogeography and phylogeography of lichen fungi and their photobionts. In Fontaneto D (ed.), *Biogeography of Microscopic Organisms: Is Everything Small Everywhere?* Cambridge: Cambridge University Press, pp. 191–208.
- Werth S and Scheidegger C** (2012) Congruent genetic structure in the lichen-forming fungus *Lobaria pulmonaria* and its green-algal photobiont. *Molecular Plant-Microbe Interactions* **25**, 220–230.
- Werth S and Sork VL** (2014) Ecological specialization in *Trebouxia* (*Trebouxiophyceae*) photobionts of *Ramalina menziesii* (*Ramalinaceae*) across six range-covering ecoregions of western North America. *American Journal of Botany* **101**, 1127–1140.
- Wetmore C** (1970) The lichen family *Heppiaceae* in North America. *Annals of the Missouri Botanical Gardens* **57**, 158–209.
- Williams L, Colesie C, Ullmann A, Westberg M, Wedin M and Büdel B** (2017) Lichen acclimation to changing environments: photobiont switching vs. climate-specific uniqueness in *Psora decipiens*. *Ecology and Evolution* **7**, 2560–2574.
- Wirtz N, Lumbsch HT, Green TGA, Türk R, Pintado A, Sancho LG and Schroeter B** (2003) Lichen fungi have low cyanobiont selectivity in maritime Antarctica. *New Phytologist* **160**, 177–183.
- Withrow K and Ahmadjian V** (1983) The ultrastructure of *Chiodecton sanguineum*. *Mycologia* **75**, 337–339.
- Wolf M, Hepperle D and Krienitz L** (2003) On the phylogeny of *Radiococcus*, *Planktosphaeria* and *Schizochlamydes* (*Radiococcaceae*, *Chlorophyta*). *Biologia, Bratislava* **58**, 759–765.
- Wolk CP, Ernst A and Elhai J** (1994) Heterocyst metabolism and development. In Bryant DA (ed.), *The Molecular Biology of Cyanobacteria*. Dordrecht: Kluwer Academic Publishers, pp. 769–823.
- Wornik S and Grube M** (2010) Joint dispersal does not imply maintenance of partnerships in lichen symbioses. *Microbial Ecology* **59**, 150–157.

- Wynne MJ (1969) Life history and systematic studies of some Pacific North American *Phaeophyceae* (brown algae). *University of California Publications in Botany* **50**, 1–88.
- Xu H, Deckert RJ and Garbary DJ (2008) *Ascophyllum* and its symbionts. X. Ultrastructure of the interaction between *A. nodosum* (*Phaeophyceae*) and *Mycophycias ascophylli* (*Ascomycetes*). *Botany* **86**, 185–193.
- Xu M, De Boer H, Olafsdottir ES, Omarsdottir S and Heidmarsson S (2020) Phylogenetic diversity of the lichenized algal genus *Trebouxia* (*Trebouxiophyceae*, *Chlorophyta*): a new lineage and novel insights from fungal-algal association patterns of Icelandic cetrarioid lichens (*Parmeliaceae*, *Ascomycota*). *Botanical Journal of the Linnean Society* **194**, 460–468.
- Yahr R, Vilgalys R and DePriest PT (2004) Strong fungal specificity and selectivity for algal symbionts in Florida scrub *Cladonia* lichens. *Molecular Ecology* **13**, 3367–3378.
- Yahr R, Vilgalys R and DePriest PT (2006) Geographic variation in algal partners of *Cladonia tenuis* (*Cladoniaceae*) highlights the dynamic nature of a lichen symbiosis. *New Phytologist* **171**, 847–860.
- Yahr R, Florence A, Škaloud P and Voytsekhovich A (2015) Molecular and morphological diversity in photobionts associated with *Micarea* s. str. (*Lecanorales*, *Ascomycota*). *Lichenologist* **47**, 403–414.
- Yung CCM, Chan Y, Lacap DC, Pérez-Ortega S, de los Ríos A, Lee CK, Cary SC and Pointing S (2014) Characterization of chasmoendolithic community in Miers Valley, McMurdo Dry Valleys, Antarctica. *Microbial Ecology* **68**, 351–359.
- Zahlbruckner A (1907) Spezieller Teil: Ascolichenes (Schlauchflechten); Hymenolichenes (Basidiomycetenflechten). In Engler A and Prantl K (eds), *Die Naturlichen Pflanzenfamilien*. I. Teil, 1. Abteilung*. Leipzig: Wilhelm Engelmann, pp. 49–243.
- Zahradníková M, Andersen HL, Tonsberg T and Beck A (2017) Molecular evidence of *Apatococcus*, including *A. fuscideae* sp. nov., as photobiont in the genus *Fuscidea*. *Protist* **168**, 425–438.
- Zeitler I (1954) Untersuchungen über die Morphologie, Entwicklungsgeschichte und Systematik von Flechtengonidien. *Österreichische Botanische Zeitschrift* **101**, 453–487.
- Zhang J, Huss VAR, Sun X, Chang K and Pang D (2008) Morphology and phylogenetic position of a trebouxiophycean green alga (*Chlorophyta*) growing on the rubber tree, *Hevea brasiliensis*, with the description of a new genus and species. *European Journal of Phycology*, **43**, 185–193.
- Zhaxybayeva O, Gogrtzen JP, Charlebois RL, Doolittle WF and Papke RT (2006) Phylogenetic analyses of cyanobacterial genomes: quantification of horizontal gene transfer events. *Genome Research* **16**, 1099–1108.
- Zhu H, Zhao Z, Xia S, Hu Z and Liu G (2015) Morphological examination and phylogenetic analyses of *Phycopeltis* spp. (*Trentepohliales*, *Ulvophyceae*) from tropical China. *PLoS ONE* **10**, e0114936.
- Zhu H, Hu Z and Liu G (2017) Morphology and molecular phylogeny of *Trentepohliales* (*Chlorophyta*) from China. *European Journal of Phycology* **52**, 330–341.
- Zhu H, Li S, Hu Z and Liu G (2018) Molecular characterization of eukaryotic algal communities in the tropical phyllosphere based on real-time sequencing of the 18S rDNA gene. *BMC Plant Biology* **18**, 365.
- Zoller S and Lutzoni F (2003) Slow algae, fast fungi: exceptionally high nucleotide substitution rate differences between lichenized fungi *Omphalina* and their symbiotic green algae *Coccomyxa*. *Molecular Phylogenetics and Evolution* **29**, 629–640.