

MYCOTAXON

<http://dx.doi.org/10.5248/130.721>

Volume 130, pp. 721–729

July–September 2015

A new species and a new record of *Anthostomella* on *Alnus* leaf-litter from Argentina

ESTEBAN B. SIR^{1,2*}, ANDREA I. ROMERO³, & ADRIANA I. HLADKI¹

¹*Laboratorio Criptogámico, Fundación Miguel Lillo,
Miguel Lillo 251, San Miguel de Tucumán (CP 4000), Tucumán, Argentina*

²*Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)*

³*PRHIDEB-CONICET, Dep. de Biodiversidad y Biología Experimental,
Facultad de Ciencias Exactas y Naturales (UBA),
Ciudad Universitaria, Pabellón II, 4to. Piso, CP1428EHA Buenos Aires, Argentina*

* CORRESPONDENCE TO: sirestebanbenjamin@gmail.com

ABSTRACT — A study on saprobic fungi occurring on litter of *Alnus acuminata* from northwestern Argentina yielded an unknown *Anthostomella* species (described here as *Anthostomella saltensis*) and the first record of *A. scotina* for Argentina. A table is provided comparing *A. saltensis* with similar species.

KEY WORDS — *Ascomycota*, *Betulaceae*, *Xylariaceae*, Yungas

Introduction

There have been few investigations on the genus *Anthostomella* in Argentina. The main publications are those by Spegazzini (1880, 1881, 1887, 1899, 1910, 1915) recording eight species, of which only four are currently accepted: *A. achira* (Speg.) Speg., *A. fuegiana* Speg., *A. sphaeroidea* Speg., and *A. sepelibilis* (Berk. & M.A. Curtis) Sacc. Lu & Hyde (2000) synonymized or excluded the remaining species, treating *A. nummularioides* Speg. and *A. phoenicicola* Speg. as synonyms of *A. tenacis* (Cooke) Sacc. and *A. platensis* Speg. as a synonym of *A. tumulosa* (Roberge ex Desm.) Sacc. and excluding *A. lonchosperma* Speg. from the genus. Additionally, they synonymised *Phaeophomatospora argentinensis* Speg. with *Anthostomella limitata* Sacc., *Anthostoma yatay* Speg. with *Anthostomella dilatata* (Berk. & Broome) Petch, and *Entosordaria fuegiana* Speg. (nom. illeg.) with *Anthostomella phaeosticta* (Berk.) Sacc. Lu & Hyde (2000) also cited a collection from Argentina on a *Bromelia* sp.

when they proposed the new species *Anthostomella raphidophylli* B.S. Lu & K.D. Hyde. An erroneous record of *A. tomicoides* Sacc. from Argentina (Lu & Hyde 2000: 188, as *A. italicica* Sacc. & Speg.) is based on a Spegazzini collection from Bellano, Italy.

Recently, Capdet (2012) has reported *Anthostomella nitidissima* (Durieu & Mont.) Sacc., *A. palmaria* B.S. Lu & K.D. Hyde, *A. puiggarii* Speg., and *A. spiralis* K.D. Hyde & B.S. Lu on fallen woody parts of native palms from Argentina.

Overall, fourteen species growing on herbaceous plants (*Cannaceae*, *Juncaceae*, *Agavaceae*, *Apiaceae*, *Smilacaceae*, *Asparagaceae*) and woody parts of palms (*Arecaceae*) (Lu & Hyde 2000, Capdet 2012) have been reported so far from Argentina.

Alnus acuminata Kunth (“aliso del cerro”; *Betulaceae*) is a tree species present from northern Mexico to northwest Argentina (www.theplantlist.org). It is the dominant plant of the Argentina Yungas at the highest altitudes between 700 and 1500 m.s.l. (Zuloaga & Morrone 1999). Little is known about *Xylariaceae* on *A. acuminata*. Sir et al. (2012) recently reported *Hypoxylon rubiginosum* var. *microsporum*, *Xylaria luxurians*, and *X. mellissii* from Tucumán province (Argentina), but no *Anthostomella* species have been recorded on this host.

The present work deals with *Anthostomella* species collected in leaf-litter of *Alnus acuminata* during a survey of the *Xylariaceae* within the Argentine Yungas. We propose a new species and record *A. scotina* for the first time for the country.

Materials & methods

The sampling area (23–29°S 64–68°W) is part of the ‘Yunga’ phytogeographic region (Cabrera 1971) in northwest Argentina, involving the Salta, Jujuy, Catamarca, and Tucumán provinces (Sir et al. 2012: fig. 1). Microscopic preparations and observations, terminology, and the criteria used for descriptions follow Lu & Hyde (2000) and Francis (1975).

Taxonomy

Anthostomella saltensis Sir, Hladki & A.I. Romero, sp. nov.

PLATE 1

INDEX FUNGORUM IF550513

Differs from *Anthostomella clypeata* by its slightly smaller, inequilaterally ellipsoidal ascospores with a germ slit extending the full length of the spore.

TYPE: Argentina, Salta, Department Santa Victoria, National Reserve El Nogalar de los Toldos, 27-XII-2011, Sir & Hladki 118 (Holotype, LIL).

ETYMOLOGY: “saltensis” refers to the name of the Salta province.

ASCOMATA peritheciate, immersed, clustered or solitary, visible as black and conical areas, dome-shaped in vertical sections (183–)186–250(–274) × (188–)




Plate 1 (FIGS 1–8). *Anthostomella saltensis* (Holotype). 1: Clypeus on surface of *Alnus acuminata* leaf; 2: Longitudinal section of immersed perithecioid; 3: Peridium (arrow); 4: Ascii; 5: Ascal apical apparatus; 6, 8: Ascospores; 7: Ascospores showing germ slit. Bars: 1 = 200 µm; 2, 3 = 100 µm; 4–8 = 10 µm.

193–250(–273) µm ($x = 225 \times 222.8$ µm, $n = 11$). CLYPEUS black, (160–)162.5–250 (–314) µm diam. ($x = 208.6$ µm, $n = 11$); ostiolar papilla (30–)37–60(–61) µm diam. ($x = 47.7$ µm, $n = 11$), periphysate ostiolar canal 27 µm broad. PERIDIUM

10–16.5 μm thick ($x = 14 \mu\text{m}$). Ascii cylindrical, unitunicate, 8-spored, short-pedicellate (64–)68–97(–98) \times (5–)5.5–7.5(–8) μm ($x = 83.2 \times 6.1 \mu\text{m}$, $n = 50$), apically rounded with IK+ apical ring inverted hat-shaped, 2–3 \times 1–2 μm ($x = 2.1 \times 1.7 \mu\text{m}$, $n = 50$). PARAPHYSES septate, hyaline, 2–5(–5.5) μm diam. ($x = 3.5 \mu\text{m}$, $n = 50$). ASCOSPORES (9–)9.5–13(–14) \times 4–5.5(–6) μm ($x = 11.2 \times 4.9 \mu\text{m}$, $n = 100$), two-celled with a hyaline globose basal cell, 1–2 \times 1.5–2(–2.5) μm ($x = 1.4 \times 2 \mu\text{m}$), and a larger brown cell, 8–11(–12) \times 4–5.5(–6) μm ($x = 9.5 \times 4.8 \mu\text{m}$), inequilaterally ellipsoidal, usually with narrowly rounded ends, with one end narrowly rounded and the other end truncate; germ slit straight extending over the full length, ventral. Mucilaginous sheath lacking.

HOST — leaf-litter of *Alnus acuminata*.

DISTRIBUTION — Argentina (Salta province).

ADDITIONAL MATERIAL EXAMINED — ARGENTINA, SALTA, Department Santa Victoria National Reserve El Nogalar de los Toldos: near the river, 26-VI-2013, Sir & Hladki 499 (LIL); road to the National Park Baritú, 28-XII-2011, Sir & Hladki 119 (LIL); 27-VI-2013 Sir & Hladki 500 (LIL). CHILE, Los Ríos, Valdivia, on leaves and stems of *Uncinia erinacea*, Jan. 1909, C. Spegazzini (LPS 6537, holotype of *Paranthostomella unciniicola*).

NOTES — TABLE 1 summarizes the morphological differences between *Anthostomella saltensis* and the three most similar *Anthostomella* species. *Anthostomella saltensis* is close to *Anthostomella clypeata* (De Not.) Sacc. as described by Lu & Hyde (2000), but our examination of the holotype of *Paranthostomella unciniicola* Speg. (considered a synonym of *A. clypeata* by Lu & Hyde 2000) revealed differences between the Argentine collection and the accepted concept of *A. clypeata*. *Anthostomella clypeata* differs from *A. saltensis* by its ascospores being equilaterally ellipsoidal, without a germ slit, and slightly larger ($x = 13.3 \times 5.3 \mu\text{m}$; Lu & Hyde 2000). Another allied species, *A. clypeoides* Rehm, has ascospores of a similar size and shape, but they lack a germ slit, have brown cells with one conic end, and are surrounded by a thin mucilaginous sheath (Lu & Hyde 2000). *Anthostomella sabinianae* S.M. Francis is also similar but has larger ascospores with one end widely rounded.

As *Anthostomella* spores do not easily germinate, cultures are not available and few sequences are registered at GenBank (<http://www.ncbi.nlm.nih.gov/>), making molecular comparisons impractical. There is no phylogenetic study on the genus available at present. Most new taxa have been based on morphological characters (Lu & Hyde 2000; Lee & Crous 2003). There are a few general studies on some *Anthostomella* sequences such as Zhang et al (2006) and Pelaez et al. (2008), and sequences have been generated for five species by Lu & Hyde (2000) and for two other species by Crous et al. (2006) and Crous & Groenewald (2010).

TABLE 1. Comparison of ascospores and the ascal apical ring in *Anthostomella saltensis* and similar species.

| SPECIES | ASCOSPORES (all two-celled) | | | | | | APICAL RING (μm) | SOURCE |
|--------------------------|--------------------------------|--|------------------------|---------------------------------|----|----|------------------------|---------------------------------------|
| | TOTAL LENGTH (μm) | BROWN CELL SHAPE | SIZE (μm) | HYALINE CELL SIZE (μm) | G* | M* | | |
| <i>A. sabiniana</i> | 14–15 | ± ellipsoid, one end broadly rounded | 12–13 × 6–7 | 2 × 3 | + | + | 4 × 3 | Francis 1975 |
| | 13.5–16 | ± ellipsoid, wider at base | 11.5– 14 × 5–7 | 2–2.5 × 2.5–3 | + | – | 2–2.5 × 2.5–3 | Lu & Hyde 2000 |
| <i>A. clypeata</i> | 12–15 | ellipsoid | 10.5– 13 × 5–5.5 | 1.5–2 × 1.5–2 | – | – | 1.5–2 × 1.3–1.5 | Lu & Hyde 2000; Francis 1975 |
| <i>A. clypeoides</i> | 10–14.5 | oval- ellipsoid, one end conic | 9–13 × 3–5 | 1–1.5 × 2 | – | – | 2 × 2 | Francis, 1975 |
| | 11.5–14 | ± ellipsoid, one side flattened | 10– 12.5 × 5–6.5 | 1.5–2 × 2–2.5 | – | + | 1.3–1.5 × 2–2.5 | Lu & Hyde 2000 |
| <i>A. saltensis</i> | 9–14 | ± ellipsoid, one end narrowly rounded | 8–12 × 4–6 | 1–2 × 1.5–2.5 | + | – | 2–3 × 1–2.5 | This paper |

* G = germ slit; M = mucilaginous sheath; +/– = present/absent.

Anthostomella scotina (Durieu & Mont.) Sacc., Syll. Fung. 1: 288. 1882. PLATE 2

ASCOMATA immersed, clustered or solitary, visible as black and conical areas, dome-shaped, in adaxial and abaxial leaf, globose or subglobose in vertical sections, (220–)235–300(–310) × (180–)200–295(–300) μm ($x = 258.1 \times 248.3$ μm, $n = 13$). CLYPEUS black (100–)200–255(–280) μm diam. ($x = 225.1$ μm, $n = 13$); conical ostiolar papilla 50–98 μm diam. ($x = 63.7$, $n = 13$), periphysate ostiolar canal 14–37.5 μm diam. ($x = 25.7$ μm). PERIDIUM (8–)9–16(–17) μm diam. ($x = 13.1$ μm). ASCI cylindrical, unitunicate, short-pedicellate, 8-spored, (85–)87.5–117.5 × 4.5–7 μm ($x = 101.7 \times 5.4$ μm, $n = 50$), apically rounded, with a J+ cylindrical-shaped apical ring (1.5–)2–3–x 1–2 μm ($x = 2.4 \times 1.3$ μm, $n = 50$). PARAPHYSES septate, hyaline (3–)3.5–6(–7) μm diam. ($x = 4.7$ μm, $n = 50$). ASCOSPORES (20–)21–29(–30) μm total length ($x = 25.7$ μm, $n = 100$), fusiform, three cells, with inequilaterally ellipsoidal or cylindrical brown central cell,




Plate 2 (FIGS 9-14). *Anthostomella scotina* (Sir & Hladki 120). 9: Clypeus on surface of *Alnus acuminata* leaf; 10: Longitudinal section of immersed perithecium; 11: Peridium (arrow); 12: Ascus apical apparatus; 13: Ascospores; 14: Ascospores showing germ slit (arrows). Bars: 9 = 450 µm; 10, 11 = 100 µm; 12-14: 10 µm.

with one broadly rounded basal end and one narrowly rounded distal end (9-) 9.5-14(-15) × (3-)3.5-4.5(-5) µm ($x = 11.3 \times 4.1$ µm), straight and faint germ slit. Hyaline globose basal cell on the widest part of the brown cell (6-)6.5-8 (-9) × 2-3 µm ($x = 7.4 \times 2.5$ µm) and hyaline curved distal cell (5-)6-9(-10) × 1-2.5 ($x = 7.1 \times 1.9$ µm), both becoming narrower towards the end, lacking mucilaginous sheath.

MATERIAL EXAMINED — ARGENTINA: CATAMARCA: Department Aconquija, near road, 13-V-2013, leg. Sir & Hladki 405-406 (LIL). JUJUY: Department Ledesma, entering the National Park Calilegua, 12-V-2012, leg. Sir & Hladki 120 (LIL). TUCUMÁN: Department Yerba Buena, Cerro San Javier, Memorial Park, Aliso forests, 27-X-2011, leg. Sir & Hladki 366 (LIL); Villa Nougués, near road, 25-XI-2011, leg. Sir & Hladki 362 (LIL); Department Monteros, Las Azucenas, 27-X-2011, leg. Sir & Hladki 364 (LIL); La Heladera, 27-X-2011, leg Sir & Hladki 365 (LIL); Department Chicligasta, Provincial Park El Cochuna, near road, 26-XI-2011, leg. Sir & Hladki 363 (LIL); La Banderita, Aliso

forests, 13-V-2013, leg. Sir & Hladki 404 (LIL); Las Lenguas, Aliso forests, 13-V-2013, leg. Sir & Hladki 403 (LIL); Department Tafi del Valle, near the los Sosa river, 7-VI-2012, leg. Sir & Hladki 361 (LIL). CHILE: Bío Bío, Concepción, Caracol, on *Persea lingue*, Jan. 1909, C. Spegazzini (LPS 7098, holotype of *Entosordaria perseicola*).

HOSTS — *Alnus* (*Betulaceae*), *Betula* (*Betulaceae*), *Cladium* (*Cyperaceae*), *Lithocarpus* (*Fagaceae*); *Persea* (*Lauraceae*).

DISTRIBUTION — Algeria, Argentina (Catamarca, Jujuy, and Tucumán provinces), Belgium, Chile, UK, USA (California).

NOTES — There are some discrepancies between our ascospores and those described by Francis (1975) and Lu & Hyde (2000). We follow Lu & Hyde (2000) in interpreting the ascospores as 3-celled, while Francis (1975) interpreted the distal hyaline cell as a gelatinous appendage, consequently citing a much smaller “total” spore length; the total ascospore length measured by Lu & Hyde (2000) is similar to ours. The length of the brown cell measured by Francis (1975) is similar to ours, but Lu & Hyde (2000) cite a shorter length. Lu & Hyde (2000) described a very thin gelatinous sheath and absence of a germ slit; whereas both Francis (1975) and we observed a faint germ slit but no mucilaginous sheath.

Provisional key to species of *Anthostomella* from northwestern Argentina

1. Ascospores with a single hyaline basal dwarf cell *A. saltensis*
1. Ascospores with two hyaline cells 2
2. Brown cell $9\text{--}15 \times 3\text{--}5 \mu\text{m}$, inequilaterally ellipsoidal,
hyaline cells $5\text{--}10 \times 1\text{--}3 \mu\text{m}$, apical ring cylindrical shaped *A. scotina*
2. Brown cell $12\text{--}16.5 \times 5\text{--}7 \mu\text{m}$, inequilaterally ellipsoid,
dwarf cells $1.5\text{--}2.5 \times 2\text{--}3 \mu\text{m}$ apical ring wedge shaped *A. sepelibilis*

Discussion

Eleven xylariaceous genera have been reported in the protected areas of the Argentine Yungas (Sir et al. 2012) but the only report of *Anthostomella* species from this region is *A. sepelibilis*, cited from Tucumán (Lu & Hyde 2000). Our survey in the Argentine Yungas yielded two additional *Anthostomella* species. The characters of one did not match those of any previously described *Anthostomella*. Its morphology differs enough that the specimens can be described as a new species, *A. saltensis*. Future molecular research will test our hypothesis and contribute more information about the affinities among *Anthostomella* species. The other species found, *A. scotina*, is incorporated into the mycobiota of Argentina.

Anthostomella species are found on the stems and leaves of 82 plant families (Lu & Hyde 2000), but only *A. scotina* was cited on *Betulaceae* (*Betula alba* L., *B. pendula* Roth). Our study reports for the first time an *Anthostomella* species on another genus of this family, *Alnus*.

Alnus acuminata is a tree of great ecological importance for Argentina's Yungas. It is a pioneer species of forest succession, appearing mainly after landslides or large fires (Arturi et al. 1998) and is associated with an unusual and diverse mycobiota. Walker et al. (2012) reported a new member of *Gnomoniaceae* on *Alnus acuminata* leaves from Tucumán (Argentina) and our research added two species of a xylariaceous genus, one new to science and one new to the country.

Acknowledgements

The authors express their appreciation to the authorities of Fundation Miguel Lillo, Tucumán, for financial support. Dr. Andrea I. Romero thanks the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET-Argentina), PROPLAME-PRHIDEB (publication N° 201). The authors also thank the pre-submission reviewers Dr. Liliane Petrini and Dr. Yu-Ming Ju for valuable comments and suggestions.

Literature cited

- Arturi MF, Grau HR, Aceñolaza PG, Brown AD. 1998. Estructura y sucesión en bosques montanos del Noroeste de Argentina. Revista Biología Tropical 46: 525–532.
- Cabrera A. 1971. Fitogeografía de la de la República Argentina. Boletín de la Sociedad Argentina de Botánica 14(1–2): 147–159.
- Capdet M. 2012. Biodiversidad de Ascomycetes sexuales y asexuales xilófilos sobre palmeras nativas en la República Argentina. Tesis Doctoral. Universidad de Buenos Aires en el área Biología. Buenos Aires, Argentina. 214 p.
- Crous PW, Groenewald JZ. 2010. *Anthostomella pinea*. Fungal Planet 53. Persoonia 25: 126–127.
- Crous PW, Groenewald JZ, Wingfield MJ. 2006. *Anthostomella eucalyptorum*. Fungal Planet 1.
- Francis SM. 1975. *Anthostomella* Sacc. (Part I). Mycological Papers 139. 197 p.
- Lee S, Crous PW. 2003. New species of *Anthostomella* on fynbos, with a key to the genus in South Africa. Mycological Research 107(3): 360–370. <http://dx.doi.org/10.1017/S0953756203007184>.
- Lu B, Hyde KD. 2000. A world monograph of *Anthostomella*. Fungal Diversity Research Series 4. 376 p.
- Peláez F, Gonzalez V, Platas G, Sanchez-Ballesteros J. 2008. Molecular phylogenetic studies within the family Xylariaceae based on ribosomal DNA sequences. Fungal Diversity 31: 111–134.
- Sir EB, Parrado MF, Romero AI, Hladki AI. 2012. Biodiversity of Xylariaceae (Ascomycota) and their hosts in protected areas from Tucumán (Argentina). Kurtziana 37(2): 35–48.
- Spegazzini CL. 1880. Fungi argentini. Pugillus primus. Annales de la Sociedad Científica Argentina. 9: 158–192.
- Spegazzini CL. 1881. Fungi argentini. Pugillus quartus. Annales de la Sociedad Científica Argentina. 12: 97–117, 174–189.
- Spegazzini CL. 1887. Fungi fuegiana. Boletín de la Academia Nacional de Ciencias de Córdoba 11: 135–308.
- Spegazzini CL. 1899. Fungi argentini novi vel critici. Annales Museo Nacional Buenos Aires 6: 81–365.
- Spegazzini CL. 1910. Fungi chilenses. Revista de la Facultad de Agronomía y Veterinaria de La Plata 6: 1–205.
- Spegazzini CL. 1915. Fungi nonnulli senegalenses et canarienses. Annales Museo Nacional Buenos Aires 26: 117–134.

- Walker DC, Castelebury LA, Rossman AY, Mejia L C, White JF. 2012. Phylogeny and taxonomy of *Ophiognomonia* (*Gnomoniaceae, Diaphorales*); including twenty-five new species in this highly diverse genus. *Fungal Diversity* 57: 85–147. <http://dx.doi.org/10.1007/s13225-012-0200-y>
- Zhang N, Castlebury LA, Miller AN, Huhndorf SM, Schoch CL, Seifert KA, Rossman AY, Rogers JD, Kohlmeyer J, Volkmann-Kohlmeyer B, Sung GH. 2006. An overview of the systematics of the *Sordariomycetes* based on a four-gene phylogeny. *Mycologia* 98(6): 1076–1087. <http://dx.doi.org/10.3852/mycologia.98.6.1076>
- Zuloaga FO, Morrone O (eds). 1999. Catálogo de las plantas vasculares de la República Argentina. II. *Acanthaceae – Euphorbiaceae (Dicotyledoneae)*. Monographs in Systematic Botany from the Missouri Botanical Garden 74: 1– 621.