

Dendritic Ferroselite (FeSe_2) with 2D Carbon Based Nanosheets of rGO and g- C_3N_4 as Efficient Catalysts for Electrochemical Hydrogen Evolution

R. Shwetharani^a, Samadhan Kapse^c, Ranjit Thapa^c, D. H. Nagaraju^{ab*}, R. Geetha Balakrishna^{a*}

^aCentre for Nano and Material Sciences, Jain University, Bangalore 562112, Karnataka, India

^bDepartment of Chemistry, School of Applied Sciences, REVA University, Bangalore 560064, Karnataka, India

^cDepartment of Physics, SRM University – AP, Amaravati 522502, Andhra Pradesh, India

*Mailing Address: br.geetha@jainuniversity.ac.in, dhnagu@gmail.com




Fig. S1. FESEM image of FeSe_2/rGO




Fig. S2. XPS spectra of FeSe₂




Fig S3. TEM image of FeSe₂/rGO




Fig S4. TGA curve for 2:8 rGO/FeSe₂ hybrid electro-catalyst




Fig S5. Mass loading of FeSe₂ on g-C₃N₄ in hybrid catalyst




Fig S6. LSV polarization plot of FeSe₂/g-C₃N₄, FeSe₂/GO and FeSe₂/rGO




Fig S7. Impedance spectra of rGO/FeSe₂, g-C₃N₄/FeSe₂ and GO/FeSe₂ composite




Fig S8. FESEM images of (a) GO and (b) GO/FeSe₂




Fig S9. TEM image of GO/FeSe₂