



Report on the **Subtidal Habitats**
and Associated Biological Taxa **in San Francisco Bay**

Editors and Contributors

Editors

Korie Schaeffer, *National Marine Fisheries Service (NMFS), Southwest Region*

Katie McGourty, *NMFS Southwest Region*

Natalie Cosentino-Manning, *NMFS Restoration Center*

Contributing Authors

Sarah Allen, *National Park Service*

Andrew Cohen, *San Francisco Estuary Institute*

Meredith Elliot, *Point Reyes Bird Observatory*

Natalie Cosentino-Manning, *NMFS Restoration Center*

Kathryn Hieb, *California Department of Fish and Game (CDFG)*

Katie McGourty, *NMFS Southwest Region*

Korie Schaeffer, *NMFS Southwest Region*

Janet Thompson, *US Geological Survey (USGS)*

Susan Wainwright-De La Cruz, *USGS*

Maps and Figures

Charleen Gavette, *NMFS Southwest Region*

Michele Jacobi, *National Ocean Service*

Jill Bodner, *National Ocean Service*

Reviewers

Chris Brown, *Smithsonian Institute*

Brenda Goeden, *San Francisco Bay Conservation and Development Commission*

Kathryn Hieb, *CDFG*

April Hennessy, *CDFG*

Laura Hoberecht, *NMFS Southwest Region*

Kathy Ann Miller, *University of California Berkeley*

David Schoellhamer, *USGS*

Janet Thompson, *USGS*

Robert Tasto, *NMFS Southwest Region*

Layout and Design

Milena Viljoen, *NMFS Restoration Center*

Cover Photo

Dennis Anderson, *Dennis Anderson Photography*

LIST OF TABLES *i*

LIST OF FIGURES *ii*

Introduction *1*

PURPOSE AND NEED	<i>1</i>
PHYSICAL ENVIRONMENT	<i>1</i>
<i>Topography and Substrates</i>	<i>1</i>
<i>Tides and Tidal Currents</i>	<i>7</i>
<i>Freshwater Inflows</i>	<i>8</i>
<i>Coastal Oceanographic Conditions</i>	<i>8</i>
<i>Salinity, Stratification and Gravitational Circulation</i>	<i>8</i>
<i>Temperature</i>	<i>13</i>
<i>Sediment Dynamics</i>	<i>13</i>
<i>Oxygen</i>	<i>15</i>
<i>Contaminant Inflows and Legacy Pollutants</i>	<i>15</i>
DELINEATION OF HABITATS	<i>16</i>
BIOTIC ASSEMBLAGES	<i>16</i>

Habitat Types and Associated Biological Taxa *36*

SOFT BOTTOM	<i>36</i>
<i>General Description</i>	<i>36</i>
<i>Taxonomic Groups</i>	<i>38</i>
<i>Plants</i>	<i>38</i>
<i>Invertebrates</i>	<i>38</i>
<i>Fishes</i>	<i>43</i>
<i>Birds</i>	<i>44</i>
<i>Marine Mammals</i>	<i>45</i>

HARD BOTTOM	45
General Description	45
Taxonomic Groups	46
<i>Plants</i>	46
<i>Invertebrates</i>	46
<i>Fishes</i>	47
<i>Birds</i>	48
<i>Marine Mammals</i>	48
SHELLFISH BEDS	49
General Description	49
California Mussel (<i>Mytilus californianus</i>) Beds	50
Bay Mussel (<i>Mytilus trossulus/galloprovincialis</i>) Beds	50
Ribbed Horsemussel (<i>Geukensia demissa</i>) Beds	51
Green Bagmussel (<i>Musculista senhousia</i>) Beds	51
Olympia Oyster (<i>Ostrea conchaphila</i>) Beds	51
PLANT BEDS	56
General Description	56
Macroalgae Beds	56
<i>Ulva Beds</i>	56
<i>Gracilaria Beds</i>	56
<i>Fucus Beds</i>	57
<i>Sargassum Beds</i>	57
Submerged Aquatic Vegetation	58
<i>Sago Pondweed (Potamogeton pectinatus) Beds</i>	58
<i>Widgeon Grass (Ruppia maritima) Beds</i>	58
<i>Eelgrass (Zostera marina) Beds</i>	60
<i>Surfgrass (Phyllospadix scouleri and torreyi) Beds</i>	63
WATER COLUMN	64
General Description	64
Taxonomic Groups	64
<i>Plants (phytoplankton)</i>	64
<i>Invertebrates (zooplankton)</i>	67
<i>Fishes</i>	71
<i>Birds</i>	72
<i>Marine Mammals</i>	73

Literature Cited 74

Index 83

- Table 1** Area, depth and volume of San Francisco Bay after (a) Conomos (1979), (b) Jassby (1992), (c) Monroe and Kelly (1992), and (d) Goals Project (1999).
- Table 2** Common subtidal and intertidal plants of San Francisco Bay.
- Table 3** Common sessile and small, motile invertebrates of San Francisco Bay for (a) soft bottom substrate (sand/silt/clay), (b) soft bottom substrate (shell mix), (c) hard bottom substrate (natural and artificial) and (d) eelgrass beds.
- Table 4** Common fish and large, motile invertebrates of San Francisco Bay after K. Hieb (CDFG, pers. comm.), Merkel & Associates (2005), and Entrix (1998).
- Table 5** Common birds that use subtidal habitats in San Francisco Bay.
- Table 6** Common marine mammals of San Francisco Bay.
- Table 7** Sessile invertebrates sampled at native oyster restoration sites at Marin Rod and Gun Club and Redwood City in San Francisco Bay after MACTEC (2006).
- Table 8** Common phytoplankton taxa in San Francisco Bay after Cloern and Dufford (2005).
- Table 9** Common zooplankton taxa in San Francisco Bay after Kimmerer et al. (1999).

- Figure 1** Area map of the San Francisco Bay showing landforms, major roads and bridges, and bay regions.
- Figure 2** Sea level rise from 15,000 years ago to present time from Cohen (2000).
- Figure 3** Bathymetry (shown in meters) of San Pablo, Central, and South Bays within San Francisco Bay.
- Figure 4** Map of generalized sediment types in San Francisco Bay from Monroe and Kelly (1992).
- Figure 5** Schematic of tidal cycles near San Francisco Bay from Cohen (2000).
- Figure 6** Average salinity during winter and summer months in San Francisco Bay.
- Figure 7** Average salinity for flood years 1996 and 1997 and drought years 1987 to 2002 by Venice Salinity categories for San Francisco Bay.
- Figure 8** Schematic of estuarine circulation as occurs in San Francisco Bay from Cohen (2000).
- Figure 9** Average winter and summer water temperature profiles for San Francisco Bay.
- Figure 10** Soft bottom and hard bottom substrates in San Francisco Bay interpreted from multibeam bathymetric images and side-scan sonar mosaics after Greene and Bizarro (2003).
- Figure 11** Historic native oyster distribution, sites of known oyster occurrence at present date, and native oyster restoration sites in San Francisco Bay.
- Figure 12** Distribution of eelgrass (*Zostera marina*), the marine alga, *Gracilaria*, and widgeon grass (*Ruppia maritima*) in San Francisco Bay after Merkel & Associates (2003) and N. Cosentino-Manning (NMFS, pers. comm.).
- Figure 13** Eelgrass (*Zostera marina*) density in San Francisco Bay in 2003 after Merkel & Associates (2004) and Wyllie Echeverria and Rutten (1989).

Introduction

(by Andrew Cohen, Korie Schaeffer, Katie McGourty, Natalie Cosentino-Manning, Susan Wainwright-De La Cruz, Meredith Elliot, and Sarah Allen)

PURPOSE AND NEED

This report describes the current status of knowledge of the primary subtidal habitat types in San Francisco Bay and their associated biota. For the most part, we did not attempt to describe trends over time, nor do we predict future conditions. Rather, information sources dating from the last 30 years are used to describe what we expect to find in San Francisco Bay in 2007. This report is meant to provide a common understanding of subtidal habitats in San Francisco Bay in support of the San Francisco Bay Subtidal Habitat Goals project. In compiling this report, we also provide a basis for future comparisons of habitat conditions and biotic assemblages.

The geographic scope of the report includes San Francisco Bay from Sherman Island at the base of the Sacramento-San Joaquin Delta (Delta) seaward to the outer Golden Gate Channel (i.e., Point Bonita and Point Lobos) and south to the southern extent of San Francisco Bay (Figure 1). The scope includes only tidally influenced areas within tributaries to San Francisco Bay, and does not include the Delta. Within this geographic area, we focus on subtidal habitats, which are below mean low water tidal height. We recognize, however, that subtidal habitats continue into intertidal areas, and that while there are certainly differences, the biota of subtidal and intertidal habitats overlap. Therefore, when appropriate and when information is available, we also discuss intertidal habitats.

Every effort was made to compile the best available information, including sources from peer-reviewed journals, agency reports, unpublished data, and expert knowledge of both scientists and resource managers working in San Francisco Bay.

PHYSICAL ENVIRONMENT

Topography and Substrates

San Francisco Bay contains four main basins (Figure 1). Suisun Bay and the diked wetlands of Suisun Marsh form the most upstream basin of the northern reach of the Bay. San Pablo Bay is the next downstream basin, and below that the Central Bay basin connects with the ocean through the Golden Gate. The large, shallow South Bay basin extends off the Central Bay. Various authorities have drawn the boundaries between basins in different places (Gunther 1987). Carquinez Strait has been included in either the Suisun or San Pablo Bay basins, or treated as a separate reach. The boundary between the San Pablo Bay and Central Bay basins has generally been set between San Pablo Strait (on a line connecting Point San Pedro to Point San Pablo) and the Richmond-San Rafael Bridge. Placement of the Central Bay-South Bay boundary has ranged from the Oakland Bay Bridge to the San Bruno shoal. The South Bay is sometimes divided into upper and lower basins, or subdivided into 2 or 3 sub-basins, with boundaries variously placed at San Bruno shoal, San Mateo Bridge or Dumbarton Bridge. Table 1 shows the basic measurements of these basins based on different segmentation schemes.



Figure 1. Area map of the San Francisco Bay showing landforms, major roads and bridges, and bay regions.

Table 1a. The area, depth and volume of San Francisco Bay after Conomos (1979).

Section	Area at MLLW (mi ²)	Area including mudflats (mi ²)	Average depth at MLLW (feet)	Average depth including mudflats ^a (feet)	Volume at MLLW (acre-feet)
Total	402	479	20	6.5	5,400,000

^a Estimated graphically from hypsometric curve.

Table 1b. The area, depth and volume of San Francisco Bay after Jassby (1992).

Section	Area at MLLW (mi ²)	Area at MHHW (mi ²)	Mean Depth at MLLW (feet)	Volume at MLLW (acre-feet)
Suisun Bay / Chipps Island to Benicia Bridge	39	66	10	251,000
San Pablo Bay / Benicia Bridge to San Pablo Strait	100	170	11	697,000
Central Bay / San Pablo Strait to Golden Gate and Oakland Bay bridges	85	97	36	2,027,000
South Bay / South of Oakland Bay Bridge	181	236	13	1,540,000
Total	405	568	17	4,515,000

Table 1c. The area, depth and volume of San Francisco Bay after Monroe & Kelly (1992).

Section	Area at MLLW ^a (mi ²)	Mean Depth at MLLW (feet)	Volume at MLLW (acre-feet)
Suisun Bay / Chipps Island to Benicia Bridge	36	14	323,000
Carquinez Strait	12	29	223,000
San Pablo Bay / Mouth of Carquinez Strait to Richmond-San Rafael Bridge	105	9	605,000
Central Bay / Between Richmond-San Rafael, Golden Gate and Oakland Bay bridges	103	35	2,307,000
South Bay / South of Oakland Bay Bridge	214	11	1,507,000
Total	470	17	4,965,000

^a Including saturated mudflats.

Table 1d. The area of San Francisco Bay after Goals Project (1999).

Section ^a	Subtidal Area ^a (mi ²)	Intertidal Area ^b (mi ²)	Total Area (mi ²)
Suisun Bay Subregion / Antioch to Carquinez Bridge	53	23	76
North Bay Subregion / Carquinez Bridge to San Pablo Strait	100	40	140
Central Bay Subregion / San Pablo Strait to Golden Gate Bridge and Coyote Point-San Leandro Marina	168	8	176
South Bay Subregion / South of Coyote Point-San Leandro Marina	76	38	114
Total	397	109	506

^a Deep Bay/Channel plus Shallow Bay/Channel habitats.

^b Tidal Flat plus Tidal Marsh habitats; boundary between Tidal Flat and Shallow Bay/Channel habitats appear to be between MLW and MLLW (Goals Project, Fig. 2.3).

INTRODUCTION – Physical Environment

Fifteen to eighteen thousand years ago during the last Ice Age, sea level was 120 meters (m) lower, the ocean shore lay west of the Farallon Islands, and the San Francisco Bay basins were a chain of broad river valleys with narrow canyons at Carquinez Strait, Raccoon Strait (between Tiburon and Angel Island), and the Golden Gate. Around 10,000 years ago, rising sea level brought seawater in through the Golden Gate, and the Bay began to fill at a rate of nearly 2 centimeters (cm) per year (Atwater 1979) (Figure 2). By about 5,000 years ago, sea level was only about 8 m lower than today, the rate of rise slowed to 0.1 to 0.2 cm per year, sediments accreted, and mudflats and marshes formed around the San Francisco Bay (Atwater 1979).

In the past 160 years, human activities have dramatically reshaped the Bay by changing sediment supply, dredging channels and diking off large portions of the Bay (Herbold *et al.* 1992). Since 1990, mean sea level increased by 22 cm per century, and mean high water rose about 19 percent faster than mean sea level, resulting in an increase in tide range of about 60 millimeters (mm) per century (Flick *et al.* 2003). Global warming studies suggest that relative to 1990 global average sea level will rise by 8.9 to 87.9 cm by 2100 (IPCC 2001).

Today, the Bay at high tide covers approximately 500 mi² (Table 1), having been reduced by about 40 percent since 1850 (Conomos 1979, Monroe and Kelly 1992). A strong low tide exposes about one-sixth of this area (Conomos 1979), with the largest intertidal mudflats extending across the eastern and southern parts of the South Bay and the northern parts of San Pablo and Suisun Bays. Another third of the Bay is less than 2 m

South Bay and the northern parts of San Pablo and Suisun Bays. Another third of the Bay is less than 2 m deep on a good low tide (Conomos 1979) (Figure 3). Narrow channels 10 to 20 m deep lead across the main basins into harbors and shipping terminals, maintained partly by dredging and partly by the scouring action of river and tidal currents. The deepest spot in San Francisco Bay, under the Golden Gate Bridge, is more than 110 m deep at low tide (Conomos 1979). Just outside the bay's mouth, a curving sandbar rises to within 9 m of the surface, built of sediments transported from the bay.

Most of the Bay's bottom is covered with sand, silt or clay, along with significant quantities of oyster shell fragments (of the native Olympia oyster *Ostrea conchaphila* and the exotic Virginia oyster *Crassostrea virginica* and Pacific oyster *C. gigas*). The channel floors in the northern reach are mostly sand, including much of the western portion of the Central Bay. Shell fragments occur over a good part of the southeastern and southwestern shallows in the South Bay. Otherwise, the Bay bottom is mostly mud, often quite soft, consisting of 80 percent or more of silt and clay (Figure 4). A few areas of bedrock rim the western part of the Central Bay and crop out at islands and a few shoreline locations. Artificial hard substrate is scattered across the Bay, including rip-rapped banks, jetties, breakwaters, seawalls, pilings, docks and piers, bridge and powerline supports, and debris.



Figure 2. Sea level rise from 15,000 years ago to the present from Cohen (2000).



California Department of Fish & Game San Francisco Bay Bathymetry (Feet)



Figure 3. Bathymetry (shown in feet) of San Pablo, Central, and South Bays within San Francisco Bay.




Figure 4. Map of generalized sediment types in San Francisco Bay from Monroe and Kelly (1992).

Tides and Tidal Currents

Tides are complex waves hundreds of miles long that are caused by the gravitational pull of the moon and the sun. The tidal range (the difference in height between high and low water) changes as the moon circles the earth every 28 days. The tides with the greatest range, called spring tides, occur during full and new moons, when the moon, sun, and earth are nearly aligned and the gravitational pulls of the moon and sun reinforce each other. Neap tides, with the least tidal range, occur during the moon's quarters, when the gravitational pulls from the moon and sun tend to cancel each other (Figure 5). Tidal ranges also vary over the year, with the highest highs and the lowest lows in the bay typically occurring around June and December (Cohen 2000).

Mixed, semi-diurnal tides, with two unequal high tides and two unequal low tides occurring about every 24 hours and 50 minutes, are typical of the west coast (Conomos 1979, Smith 1987). Twice each day, with each tidal cycle, a huge volume of salt water moves in and out of the Bay. This tidal prism, as it is called, averages about 1,300,000 acre-feet¹, or nearly a quarter of the Bay's total volume (Conomos 1979, Conomos *et al.* 1985, Largier 1996), though much of it consists of water that had flowed out of the Bay during recent tidal cycles (Monroe and Kelly 1992). In comparison, the average daily inflow of freshwater to the bay is only about 50,000 acre-feet.

The rate at which tides move inward through the Bay is determined primarily by water depth and bottom friction. The tides progress quickly up channels (where tidal currents may be >1 m/second) and spread from the channels into the shallows (where tidal currents are <0.2 m/second). The northern reach's deep channel allows a greater volume of water to be moved by tidal action through San Pablo and Suisun Bays than through the South Bay (Herbold *et al.* 1992), and the tidal patterns in the two reaches are different. In the northern reach, the tidal range shrinks with distance from the ocean, from a mean range of about 1.7 m at the Golden Gate to about

¹ One acre-foot of water will cover an acre to a foot deep, and is equal to about 326,000 gallons.




Figure 5. Schematic of tidal cycles near San Francisco Bay from Cohen (2000).

1.2 m at Chipps Island, to 0.9 m in the Sacramento River at Sacramento. The high and low tides at Chipps Island occur about 3 to 3.5 hours later than the corresponding tides at the Golden Gate, and at Sacramento about 8 hours later than at the Golden Gate. In contrast, the tidal range increases as one proceeds south through the South Bay, due to reflection and reinforcement of the tide wave within the semi-enclosed basin. The tide range reaches 2.6 m at the southern end of the South Bay, where the tides lag the Golden Gate tides by 1 to 2 hours (Conomos 1979, Smith 1987, Monroe and Kelly 1992).

Tidal currents dominate the current patterns in the Bay. They are stronger in channels than in shallows, and usually parallel the depth contours. In the South Bay,

maximum currents occur at mid-tide and slack water occurs around high and low tide, but in the northern reach maximum currents occur later, just before high and low tide, with slack water at 1 to 2 hours after high and low tide. As a result, the South Bay begins to flood while San Pablo Bay is still ebbing, and some northern reach water thus enters the South Bay (Smith 1987).

Freshwater Inflows

The bay's watershed covers about 60,000 mi², or about 40 percent of California (Conomos 1979, Conomos *et al.* 1985). Roughly half of California's surface water supply falls as rain or snow within this region, and about half of that is diverted for human use. The remainder flows downstream into San Francisco Bay, with about 14 percent coming from local rivers (primarily Denverton Creek draining into Suisun Bay, Petaluma and Napa Rivers into San Pablo Bay, and Alameda and Coyote Creeks into South Bay (Herbold *et al.* 1992), and the rest entering through the Sacramento-San Joaquin Delta.

Located immediately upstream of the Bay's northern reach, the Delta is a thousand-square-mile triangle of diked and drained swampland. The remains of once-extensive tule marshes, fringe sloughs and channels that wind between "islands" of flat, levee-rimmed farmlands. The Delta is the central component of a vast plumbing system that extends through nearly the entire state. Water released from reservoirs in northern California flows down the Sacramento Valley and into the northern Delta, while large pumping stations pull water from the southern Delta and deliver it to southern cities and farms. Other systems divert water from tributaries in the Sierra Nevada; some is used locally, and some is piped to Bay Area cities. The long-term average outflow from the Delta into the bay is estimated at 17 million acre-feet per year, with another 2.7 million acre-feet entering from the local watersheds (Monroe and Kelly 1992).

Coastal Oceanographic Conditions

Conditions in the coastal ocean outside of the Golden Gate typically progress through three seasons over the course of the year. During the upwelling season, from March or April to July or August, strong and persistent winds from the northwest induce a southward-running surface current called the California Current. The Coriolis effect and Ekman transport produce a net offshore movement of water, which in turn, cause upwelling of bottom water along the coast. This cold, nutrient-rich water brought to the surface in the Gulf of the Farallones is then available for exchange with Bay water over the shallow bar outside the mouth of the Bay. During the relaxation, or oceanic, season from August to October or November, the winds and currents weaken, upwelling stops and surface waters warm, dissipating the coastal fog. In the winter, or Davidson Current season, winds are often from the south or southeast although winter storms bring intermittent strong winds from the west and southwest. The Davidson Current runs northward, producing a downwelling of surface water (Smith 1987, Herbold *et al.* 1992, Largier 1996).

This pattern is interrupted in El Niño years. In those years, warm tropical surface waters move northward, creating vertical density differences that can prevent upwelling (Herbold *et al.* 1992).

Salinity Stratification and Gravitational Circulation

An estuary is a partially enclosed body of water where fresh water meets and mixes with salt water from the ocean. The mixing zone in San Francisco Bay, where fresh and salt water meet, can move tens of miles upstream and downstream as river flows decrease and increase, and even further under extreme conditions. The 1862 flood pushed the mixing zone out beyond the Golden Gate for several weeks and freshened the ocean's surface as far as 40 miles from shore. At the opposite extreme, the 1931 drought pulled the mixing zone inland as far as Courtland on the Sacramento River and Stockton on the San Joaquin River (Smith 1987, Cohen 2000).

Because freshwater is lighter than water that contains a significant amount of dissolved salts, inflowing river water tends to float on top of, and only gradually mixes with, sea water in the bay. This stratification is usually stronger in winter and in wet years when river flows are greater (Conomos 1979), and occurs primarily in channels. Tidal currents during spring tides are typically 2 to 3 times stronger than during neap tides, and tend to increase vertical mixing and reduce stratification (Smith 1987). In the shallows, currents generated by wind and tides usually keep water mixed throughout the water column.

Freshwater inflows also create horizontal salinity gradients. Water from the Sacramento River and the eastside streams entering the Delta averages less than 0.1 part by weight of salt per thousand parts of water (parts per thousand or ppt), while San Joaquin River water entering the Delta has about 0.4 ppt of salt, primarily because of farm drainage (Monroe and Kelly 1992). Outflow from the Delta to the Bay is typically under 0.5 ppt, with the salinity increasing as one proceeds downstream to about 30 ppt near the mouth of the Bay, and about 33 ppt in the coastal ocean (Conomos 1979). At any location in the Bay, the salinity is generally lower when river flows are high.

Within San Francisco Bay, salinity also depends on water year type, and season (Cloern *et al.* 1999). In this report, we discuss salinity categories consistent with the Venice System (Anonymous 1959): limnetic (0 to 0.5 ppt), oligohaline (0.5 to 5.0 ppt), mesohaline (5.0 to 18.0 ppt), polyhaline (18.0 to 30.0 ppt), and euhaline (30.0 to 35.0 ppt). Based on recent work by Thompson *et al.* (2000) and Lee *et al.* (2003), Suisun Bay is typically oligohaline or mesohaline, San Pablo Bay and South Bay are typically polyhaline, and Central Bay is typically euhaline. Within each Bay region, salinities are typically lower in winter and spring than in summer and fall (Figure 6). As expected, salinities throughout San Francisco Bay tend to be lower in flood years and higher in drought years (Figure 7). Average winter salinity ranges from fresh to euhaline in flood years and from oligohaline to euhaline in drought years. Average summer salinity ranges from oligohaline to euhaline in flood years and mesohaline to euhaline in drought years.

The combination of horizontal and vertical salinity gradients can cause tidally-averaged currents of saltier water to flow landward along the bottom of the channels while fresher water flows seaward at the surface, a pattern known as gravitational (or estuarine) circulation (Figure 8). The place where the upstream and downstream currents meet and cancel out along the bottom is the null zone (Herbold *et al.* 1992, Cohen 2000). An entrapment zone can form downstream of a null zone, where suspended sediments carried by river flows clump up and start to settle on contact with saltier water, and suspended sediments, phytoplankton, zooplankton, and the drifting eggs and larvae of fish are concentrated by the downstream and upstream mix of currents (Arthur and Ball 1979, Herbold *et al.* 1992, Cohen 2000).

Bottom topography and other factors complicate this picture in the northern reach of San Francisco Bay. Research conducted during the 1990s has increased our understanding of gravitational circulation. In Suisun Bay, gravitational circulation increases with water depth, is suppressed by vertical mixing during spring tides, and is driven by the horizontal salinity gradient rather than absolute salinity (Schoellhamer and Burau 1998). Gravitational circulation is often well-developed in Carquinez Strait, with a null zone typically found near the upper end of the Strait (SFEP 1997). A zone of concentrated sediments, nutrients and small organisms often develops in Suisun Bay during spring and summer where the salinity is about 2 ppt, but not always close to a null zone (SFEP 1997).

Salinities are generally more uniform and higher in the South Bay than in the northern regions of San Francisco Bay (i.e., San Pablo and Suisun Bays), with high evaporation contributing to salinity (Conomos 1979). Freshwater discharge from the South Bay's tributaries is usually too little to stratify the water and cause gravitational circulation (Conomos 1979). Thus, residence time (the average length of time it takes for a water molecule or a dissolved contaminant to leave the system) is somewhat longer in the South Bay than in the northern reach (Conomos 1979, Smith 1987, Davis *et al.* 1991). In wet winters, however, flood flows from the northern reach enter the South Bay, causing two-layered

INTRODUCTION – Physical Environment

gravitational flow with the fresher surface water flowing southward, and the saltier bottom water flowing northward towards the mouth of the Bay, the opposite pattern to that of gravitational circulation in the northern part of the Bay. This gravitational flow can substantially reduce residence times, and flush contaminants out of the South Bay (Cohen 2000). After the high discharges subside, the intrusion of ocean water raises the salinity in

the Central Bay over that in the South Bay, and the direction of gravitational circulation reverses (Smith 1987). San Bruno Shoal may often restrict gravitational circulation to the northern part of the South Bay, though flows from Coyote Creek and Guadalupe River may also cause gravitational circulation at the southern end of the South Bay in extremely wet winters (Smith 1987).




Figure 6. San Francisco Bay average salinity (winter and summer).

The salinity profiles for the months of February (winter) and September (summer) were derived from data found on USGS Water Quality of San Francisco Bay Website (<http://sfbay.wr.usgs.gov/access/wqdata>). Data for 39 monitoring stations were retrieved, plotted and averaged. Using GIS, an interpolated grid was produced of the salinity averages (Inverse Distance Weighting, fixed radius). Dates ranged from 1969 to 2006. Number of records per station ranged from 200-400.




Figure 7. San Francisco Bay average salinity (winter and summer) for flood years 1996 and 1997 (top) and drought years 1987 to 1992 (bottom).

The salinity profiles for the months of February (winter) and September (summer) were derived from data found on USGS Water Quality of San Francisco Bay Website (<http://sfbay.wr.usgs.gov/access/wqdata>). Data for monitoring stations were retrieved, plotted and averaged. Using GIS, an interpolated grid was produced of the salinity averages (Inverse Distance Weighting, fixed radius). Number of records per station ranged from 0-70. Summer records omitted 6 stations located in southern San Francisco Bay.




Figure 8. Schematic of estuarine circulation as occurs in San Francisco Bay from Cohen (2000).

Temperature

Water temperatures in San Francisco Bay range from about 10 to 20° C (Monroe and Kelly 1992), and are highest in shallow areas in the summer (Figure 9). Since the Bay is much shallower than the nearby coastal ocean, its seasonal temperature swings are greater, so that the Bay tends to be colder than the ocean during the winter, and warmer than the ocean the rest of the year (Largier 1996, Conomos 1979). Water in the Bay is warmer than inflow from tributaries, which also contributes to seasonal fluctuations in temperature. Baywide average winter temperatures are lower than summer temperatures. Average winter temperature in Suisun Bay, which receives flow directly from the Sacramento-San Joaquin Delta tends to be colder than the rest of San Francisco Bay. In summer when freshwater inflow is low, the more marine influenced Central Bay has the lowest average temperature, with average temperature increasing as you move northeast into San Pablo and Suisun Bays and as you move south through the South Bay.

Sediment Dynamics

The primary source of sediment in San Francisco Bay is erosion from the watershed and transport into the Bay with freshwater flow from tributaries. Large plumes of sediment are carried into the Bay during the first large storm of the year; less erodible sediments are then carried in during subsequent storms (Oltmann *et al.* 1999, Goodwin and Denton 1991, Ruhl *et al.* 2001). Most suspended sediment is from the Sacramento and San Joaquin Rivers, but sediment also comes from the Yolo Bypass, Mokelumne River, Calaveras River, Cosumnes River, and several other smaller streams (Oltmann 1996). From 1856 to 1887, more than 230 million cubic meters of sediment were deposited in San Pablo Bay (Jaffe *et al.* 1998), and approximately 115 million cubic meters of sediment were deposited in Suisun Bay (Cappiella *et al.* 1999). The majority of this sediment was debris from hydraulic gold mining in the Sierra Nevada (Cappiella *et al.* 1999, Jaffe *et al.* 1998). During this period, deposition occurred in nearly the entire Bay.

In contrast, from 1951 to 2001 the watershed sediment yield decreased by about one-half, and much of San Francisco Bay became erosional (Jaffe *et al.* 1998, Wright and Schoellhamer 2003). This decline is a result of the cessation of hydraulic mining, water distribution projects decreasing sediment supply by reducing frequency and duration of peak flow conditions, trapping of sediment in reservoirs, and riverbank protection (Wright and Schoellhamer 2003, Cappiella *et al.* 1999, Jaffe *et al.* 1998). The decrease in suspended sediment resulted in a decrease in the area of tidal flats, particularly in San Pablo Bay and Grizzly and Honker Bays within the Suisun Bay region (Cappiella *et al.* 1999, Jaffe *et al.* 1998).

As sediment is carried into San Francisco Bay with freshwater inflow, it deposits on the Bay bottom and begins a cycle of resuspension, transport, and deposition (Krone 1979). In San Francisco Bay, suspended sediment is driven by resuspension of sediments by currents and wind waves, and not from riverine input (Schoellhamer 2002). Physical processes that affect suspension and deposition, and thus, suspended sediment concentration, include diurnal, semidiurnal, monthly and semiannual tidal cycles, freshwater inflow, and wind-wave stress (Schoellhamer 2002, 2001). Suspended sediments accumulate in the water column as a spring tide approaches and slowly deposit as a neap tide approaches. This dynamic occurs because the duration of deposition and consolidation of newly deposited bed sediments is limited during a relatively short duration of slack water (Schoellhamer 1996). In San Francisco Bay, suspended sediment concentrations typically are greatest in spring when wind waves resuspend sediment recently delivered during high winter flows. Stronger winds during summer months cause wind shear that resuspends surface sediments in shallow water (Krone 1979, Schoellhamer 1996). Suspended sediment concentration decreases in late summer and autumn as supply of erodible sediment decreases (Schoellhamer 2002).




Figure 9. San Francisco Bay average temperature (winter and summer).

The temperature profiles for the months of February (winter) and September (summer) were derived from data found on USGS Water Quality of San Francisco Bay Website (<http://sfbay.wr.usgs.gov/access/wqdata>). Data for 39 monitoring stations were retrieved, plotted and averaged. Using GIS, an interpolated grid was produced of the temperature averages (Inverse Distance Weighting, fixed radius). Dates ranged from 1969 to 2006. Number of records per station ranged from 200 to 400.

Oxygen

Unlike many estuaries, mixing due to strong winds and shallow depths keeps San Francisco Bay waters well oxygenated, with oxygen saturation typically found all the way through the water column down to the sediment surface (Herbold *et al.* 1992). Prior to the 1960s, high concentrations of nutrients (nitrates and phosphates) in municipal wastewater discharges frequently caused rampant algal growth (known as eutrophication), whose subsequent decomposition depleted oxygen levels and sometimes produced noxious, rotten-egg odors along the shore (Monroe and Kelly 1992). However, the construction of primary and secondary treatment plants in the 1950s and 1960s (see below) largely eliminated these problems. On rare occasions, wastewater inputs, limited water circulation, and high summer temperatures at the southern end of the South Bay still combine to reduce oxygen levels (Monroe and Kelly 1992).

Contaminant Inflows and Legacy Pollutants

Serious contamination of San Francisco Bay waters started in the 1850s, with the mining and use of an estimated 3,500 tons of toxic mercury to separate gold from ore. By the late 1800s, oil spills and discharges and untreated domestic sewage were also taking a toll. During the first half of the 20th century, increasing amounts of industrial, agricultural, and automotive wastes entered the Bay (Monroe and Kelly 1992). Visual and nasal evidence of the fouling of the Bay eventually led to efforts to control this pollution. Municipal wastewater plants began the primary treatment of sewage (removal of solids and disinfection) in the 1950s, followed by secondary

treatment (biological breakdown of organic material) in the 1960s, with tertiary treatment (targeting persistent pollutants or treating the effluent for reclamation) and pretreatment programs (reducing the industrial wastes discharged into municipal systems) starting at some plants in the 1970s. During the same period, outfalls were also moved to deeper water to dilute and disperse discharges with larger water volumes and stronger currents.

Currently, 47 municipal and 15 major industrial plants discharge trace metals and other contaminants into the Bay and Delta (SFWQCB 2007). Urban runoff carries metals, PAHs, PCBs and pesticides, along with floatable debris (predominantly plastic and polystyrene pellets) washed from streets, lawns, and industrial properties. Urban and agricultural runoff contains pesticides and herbicides, nitrates, phosphates applied as fertilizer, and selenium leached from the soil, which reaches the Bay in Delta outflow. Runoff from historic mining districts carries mercury and other toxic metals. Oil and petroleum products enter the Bay from accidental spills, leaks from boat and ship engines, and from storm drains when leaking or improperly disposed oil, grease, and antifreeze are washed off the streets. Chemical spills, leachate from landfills, herbicides used to control aquatic weeds, and contaminants that settle from the air can also end up in the Bay (Monroe and Kelly 1992, Cohen 2000).

Past discharges have also left significant quantities of mercury and other "legacy pollutants" buried in the Bay's sediments, which under some circumstances may become available for uptake by organisms. At some "hot spots" of former industrial activity or waste dumping, sediments show greatly elevated contaminant concentrations.

DELINEATION OF HABITATS

There are many ways to define and categorize habitat types. For the purpose of this report, habitat types were delineated by physical substrate and the presence of three-dimensional structure. The main habitat categories are: soft bottom (mud, sand, cobble/pebble/gravel, shell mix), natural and artificial hard bottom, shellfish beds (native oysters, clams, mussels), plant beds (marine algae beds and submerged aquatic vegetation beds), and the water column. It is important to note that within this list, two categories (plant beds and shellfish beds) are defined by the presence of biological organisms that are found on soft bottom or hard bottom substrates. These habitat categories only include those areas where individual organisms are at a high enough concentration to be considered a bed rather than any place an individual or a few individuals of that organism are found. We also have to recognize that presently the native Olympia oyster (*Ostrea conchaphila*) is not found in concentrations normally considered a “bed.” In contrast to our method of habitat delineation, we include Olympia oyster bed as a habitat type in this report because of the historic presence of Olympia oyster beds in San Francisco Bay, and because scientists, restoration groups, and resource agencies have recently shown increasing interest in the status, function, and restoration of Olympia oysters in San Francisco Bay.

BIOTIC ASSEMBLAGES

For each habitat type, we describe existing taxa of plants, including both algae and angiosperms, invertebrates, fishes, birds, and marine mammals. In many cases, little information exists to describe these assemblages, and in those cases the lack of available information is noted. We do not attempt to describe organisms beyond these taxonomic groups (e.g., bacteria, protozoa). For many

taxa, salinity is a driving factor in determining distribution and abundance by habitat type and location at a given point in time. Where appropriate and when information is available to do so, biotic assemblages are described by salinity ranges. If salinity association is not available, taxa are described by region of the Bay or location of biological sampling.

There are over 162 species of macroalgae (Josselyn and West 1985) and four species of submerged aquatic vegetation (SAV) in San Francisco Bay (Table 2). Both groups occur throughout the Bay, and provide habitat for a variety of fish, invertebrates, and birds. Some of the more common attached macroalgae include *Ulva* (*Enteromorpha*) *clathrata*², *U. (E.) intestinalis*, *U. (E.) linza*, *U. californica* (formally *angusta*) and *U. lactuca*, *Bryopsis hypnoides*, *Cladophora sericea*, *Fucus gardneri*, *Polysiphonia denudata*, *Cryptopleura violacea*, *Gelidium coulteri*, *Gracilaria pacifica* (formally *verrucosa*), *Halkymenia schizymenioides*, *Polyneura latissima* (Kathy Ann Miller, University of California at Berkeley, pers. comm.), and *Chondracanthus* (formally *Gigtina*) *exaspertata*. Most of these species can be found attached to hard substrates, soft bottoms (mudflats) or epiphytic on SAV.

Data sources for invertebrate assemblages vary by habitat type and are location-based. For each species within an assemblage we indicate the general taxonomic group, the feeding mode, and the growth position or structure orientation (Table 3a-d). Feeding modes include: filter-feeders (filter out pelagic food particles), suspension feeders (trap particles from the water column), surface deposit feeders (feed on particles that have settled onto or grown on a substrate), sub-surface deposit feeders (depend on buried carbon sources), and carnivores (consume epifaunal and infaunal organisms). Growth position or structures built by an animal give them access to specialized food sources (e.g., surface tube dwelling, surface dwelling, deep tube dwelling, free living, etc).

² Hayden et al. 2003 merged species in the genus *Enteromorpha* into the genus *Ulva*, the current correct name.

Table 2. Common subtidal and intertidal plants within San Francisco Bay after Josselyn and West (1985).

Taxon	Division	Habitat	Native/non-native
<i>Ulva</i> (includes <i>Enteromorpha</i> *)	Chlorophyta	Mud/rocks	Native
<i>Bryopsis hypnoides</i>	Chlorophyta	Rocks/sand	Native
<i>Cladophora sericea</i>	Chlorophyta	Rocks	Native
<i>Codium fragile</i> subspecies <i>tomentosoides</i>	Chlorophyta	Rocks/sand	Non-native
<i>Fucus gardneri</i>	Phaeophyta	Rocks	Native
<i>Laminaria sinclairii</i>	Phaeophyta	Rocks	Native
<i>Egregia menziesii</i>	Phaeophyta	Rocks	Native
<i>Sargassum muticum</i>	Phaeophyta	Rocks	Non-native
<i>Polyneura latissima</i>	Rhodophyta	Rocks	Native
<i>Halkymenia schizymenioides</i>	Rhodophyta	Rocks	Native
<i>Polysiphonia denudata</i>	Rhodophyta	Drift/epiphytic	Non-native
<i>Cryptopleura violacea</i>	Rhodophyta	Rocks	Native
<i>Gelidium coulteri</i>	Rhodophyta	Rocks	Native
<i>Gracilaria verrucosa</i> (formerly <i>pacifica</i>)	Rhodophyta	Sand/eelgrass/drift	Native
<i>Chondracanthus</i> (formerly <i>Gigartina</i>) <i>exasperata</i>	Rhodophyta	Rocks/sand	Native
<i>Ahnfeltiopsis</i> (formerly <i>Gymnogongrus</i>) <i>leptophyllus</i>	Rhodophyta	Rocks/sand	Native
<i>Phyllospadix scoleri/torreyi</i>	Magnoliophyta	Rocks	Native
<i>Zostera marina</i>	Magnoliophyta	Mud/sand	Native
<i>Ruppia maritima</i>	Magnoliophyta	Mud/sand	Native
<i>Potamogeton pectinatus</i>	Magnoliophyta	Mud/sand	Native

* Hayden et al. (2003) merged species in the genus *Enteromorpha* into the genus *Ulva*, the current correct name.

Table 3a. Common invertebrate taxa on soft bottom substrates (sand/silt/clay) in San Francisco Bay by salinity category and topography.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations	
OLIGOHALINE						
Channel						
<i>Americorophium spinicorne</i>	amphipod	tube, free living	interannually/seasonally ephemeral		DWR-D4C	
<i>Americorophium stimpsoni</i>	amphipod	tube, free living	interannually/seasonally ephemeral			
<i>Gammarus daberi</i>	amphipod	free living/surface	interannually/seasonally ephemeral			
<i>Corbicula fluminea</i>	bivalve	surface-subsurface	common/persistent			
<i>Corbula amurensis</i>	bivalve	surface-subsurface	dry years or dry months only			
<i>Limnodrilus hoffmeisteri</i>	oligochaete	free living subsurface	common/persistent	spring & fall peak		
<i>Varichaetadrilus angustipennis</i>	oligochaete	free living subsurface	common/persistent			
<i>Marenzelleria viridis</i>	polychaete	tube to surface	common/persistent			
Channel Edge						
<i>Corbicula fluminea</i>	bivalve	surface-subsurface	common/persistent	spring peak common		DWR D4L & D4R
<i>Varichaetadrilus angustipennis</i>	oligochaete	free living subsurface	common/persistent			
<i>Limnodrilus hoffmeisteri</i>	oligochaete	free living subsurface	common/persistent			
<i>Marenzelleria viridis</i>	polychaete	tube to surface	common/persistent	spring & fall peak		
<i>Laonome sp.</i>	polychaete	tube to surface	low number/persistent			
<i>Gammarus daberi</i>	amphipod	free living/surface	interannually/seasonally ephemeral	late spring/summer peak		
<i>Americorophium spinicorne</i>	amphipod	tube, free living	interannually/seasonally ephemeral	late spring/summer peak		
<i>Americorophium stimpsoni</i>	amphipod	tube, free living	interannually/seasonally ephemeral	late spring/summer peak		
<i>Corbula amurensis</i>	bivalve	surface-subsurface	dry years or dry months only	spring and/or fall peaks		
<i>Rhithropanopeus harrissi</i>	decapod	surface	low numbers/persistent			
<i>Palaemon macrodactylus</i>	decapod	surface	common/persistent			
MESOHALINE						
Channel						
<i>Corbula amurensis</i>	bivalve	surface-subsurface	common/persistent	spring and/or fall peaks	EMAP 2001 1.1.1	
<i>Corbicula fluminea</i>	bivalve	surface-subsurface	wet years			
<i>Nippoleucon hinumensis</i>	cumacean	free living/surface-subsurface	low number/persistent			
<i>Heteromastus filiformis</i>	polychaete	free living subsurface	low number/persistent			
<i>Marenzelleria viridis</i>	polychaete	tube to surface	low number/persistent			
Channel Edge						
<i>Ampelisca abdita</i>	amphipod	tube to surface	dry years - common			DWR D6, EMAP 2001 10.1, 8.1
<i>Grandidierella japonica</i>	amphipod	tube, free living	low number/persistent	late fall/winter peak		
<i>Corbula amurensis</i>	bivalve	surface-subsurface	common/persistent	spring/summer peak		
<i>Nippoleucon hinumensis</i>	cumacean	free living/surface-subsurface	common/persistent			
<i>Synidotea laevidorsalis</i>	isopod	free living surface	dry years - common			
<i>Marenzelleria viridis</i>	polychaete	tube to surface	common/persistent			
<i>Heteromastus filiformis</i>	polychaete	free living subsurface	low number/persistent			
	filter & deposit feeder					
	filter & deposit feeder					
	deposit & scraper					
	filter & deposit feeder					
	filter feeder					
	subsurface deposit					
	subsurface deposit					
	surface deposit					
	filter & deposit feeder					
	subsurface deposit					
	subsurface deposit					
	surface deposit					
	filter feeder					
	filter & deposit feeder					
	filter feeder					
	surface deposit					
	subsurface deposit					
	surface deposit					
	filter feeder					
	filter & deposit feeder					
	filter feeder					
	surface deposit					
	subsurface deposit					
	surface deposit					
	filter feeder					
	filter & deposit feeder					
	filter feeder					
	surface deposit					
	subsurface deposit					
	surface deposit					

Table 3a. Common invertebrate taxa on soft bottom substrates (sand/silt/clay) in San Francisco Bay by salinity category and topography.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
MESOHALINE (continued)					
Slough Channels					
<i>Ampelisca abdita</i>	filter feeder	tube to surface	common*	most common bayward	
<i>Grandidierella japonica</i>	filter & deposit feeder	tube, free living	common*		
<i>Monocorophium alienense</i>	filter & deposit feeder	tube, free living	common*		EMAP 2001 1.3-1.6
<i>Corbula amurensis</i>	filter feeder	surface-subsurface	common*		EMAP 2000 7.1, 7.4, 7.6, 6.4
<i>Nippoleucon hinumensis</i>	surface deposit	free living/surface-subsurface	common*		
Tubificidae - unidentified	free living deposit	free living subsurface	common*		
<i>Heteromastus filiformis</i>	subsurface deposit	free living subsurface	common*		
<i>Marenzelleria viridis</i>	surface deposit	tube to surface	common*		
Shallow Subtidal					
<i>Corbula amurensis</i>	filter feeder	surface-subsurface	dry years – common/persistent	summer and fall peak	
<i>Marenzelleria viridis</i>	surface deposit	tube to surface	common/persistent	late spring/summer peak	
<i>Nippoleucon hinumensis</i>	surface deposit	free living/surface-subsurface	common/persistent	dry years winter/spring peak; wet years spring peak	
<i>Monocorophium alienense</i>	filter & deposit feeder	tube, free living	common/persistent	late fall/early winter peak	DWR D7, EMAP 2001 5.1; 2000 4.2, 3.3
<i>Limnodrilus hoffmeisteri</i>	subsurface deposit	free living subsurface	low number/persistent	summer peak	
<i>Americorophium stimpsoni</i>	filter & deposit feeder	tube, free living	wet years	late fall/early winter peak	
<i>Gammarus daberi</i>	deposit & scraper	free living/surface	wet years		
POLYHALINE					
Channel					
<i>Ampelisca abdita</i>	filter feeder	tube to surface	common/persistent		
<i>Corophium acherusicum</i>	filter & deposit feeder	tube, free living	common/persistent		
<i>Corophium heteroceratum</i>	filter & deposit feeder	tube, free living	common/persistent		
<i>Molgula marhattensis</i>	filter feeder	surface	low numbers/persistent**		
<i>Corbula amurensis</i>	filter feeder	surface-subsurface	common/persistent	spring and/or fall peaks	
<i>Theora lubrica</i>	surface deposit	subsurface	common/persistent		
<i>Macoma petalum</i>	filter & deposit feeder	subsurface	low numbers/persistent		
<i>Musculista senhousia</i>	filter feeder	surface	low numbers/persistent**		
<i>Mya arenaria</i>	filter feeder	subsurface	low numbers/persistent**		
<i>Venerupis philippinarum</i>	filter feeder	subsurface	low numbers/persistent**		
<i>Nippoleucon hinumensis</i>	surface deposit	free living/surface-subsurface	common/persistent		
<i>Pyrosoma tuberculata</i>	subsurface deposit	surface	common/persistent		
<i>Tubificoides</i> spp	subsurface deposit	free living subsurface	common/persistent		
<i>Cirriformia spirabranchia</i>	surface deposit	surface-subsurface	common/persistent		
<i>Exogone lourei</i>	carnivore	subsurface	low numbers/persistent	more common in south bay	
<i>Glycera armigera</i>	carnivore	subsurface	low numbers/persistent		
<i>Heteromastus filiformis</i>	subsurface deposit	free living subsurface	low numbers/persistent		
<i>Marenzelleria viridis</i>	surface deposit	tube to surface	low numbers/persistent		
<i>Neanthes succinea</i>	surface deposit & scavenger	free living/surface-subsurface	low numbers/persistent		
<i>Polydora cornuta</i>	filter & deposit feeder	tube to surface	low numbers/persistent		
<i>Sabaco elongatus</i>	subsurface deposit	tubed, surface to .8m	low numbers/persistent		
<i>Streblospio benedicti</i>	surface deposit	tube to surface	low numbers/persistent		

REM SPDeep, SBDeep, DWR D41, USGS SM31, SM49, D6, RMP BD41

Table 3a. Common invertebrate taxa on soft bottom substrates (sand/silt/clay) in San Francisco Bay by salinity category and topography.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
POLYHALINE (continued)					
Slough Channels					
<i>Grandidierella japonica</i>	amphipod	filter & deposit feeder	common in south bay		
<i>Monocorophium alienense</i>	amphipod	filter & deposit feeder	common in south bay	most common bayward	
<i>Ampelisca abdita</i>	amphipod	filter feeder	common*		
<i>Monocorophium acherusicum</i>	amphipod	filter & deposit feeder	common*		
<i>Macoma petalum</i>	bivalve	filter & deposit feeder	common in south bay		
<i>Mya arenaria</i>	bivalve	filter feeder	common in south bay		
<i>Corbula amurensis</i>	bivalve	filter feeder	common*		
<i>Nippoleucon hinumensis</i>	cumacean	surface deposit	common in south bay		
Tubificidae - unidentified	oligochaete	surface deposit	common*		
<i>Neanthes succinea</i>	polychaete	surface deposit & scavenger	common in south bay		
<i>Sabaco elongatus</i>	polychaete	subsurface deposit	common in south bay		
<i>Heteromastus filiformis</i>	polychaete	subsurface deposit	common*		
<i>Streblospio benedict</i>	polychaete	surface deposit	common*		
Shallow Subtidal					
<i>Ampelisca abdita</i>	amphipod	filter feeder	common/persistent	summer, early winter peak	
<i>Corophium heteroceratum</i>	amphipod	filter & deposit feeder	common/persistent	spring and/or fall peak	
<i>Monocorophium acherusicum</i>	amphipod	filter & deposit feeder	sporadic	summer peak	
<i>Corbula amurensis</i>	bivalve	filter feeder	common/persistent	spring peak plus fall peak	
<i>Gemma gemma</i>	bivalve	filter & deposit feeder	common/persistent	some years	
<i>Macoma petalum</i>	bivalve	filter & deposit feeder	common/persistent	higher number in South Bay - peak in summer	
<i>Musculista senhousia</i>	bivalve	filter feeder	low numbers/persistent	higher number in South Bay - peak in summer	
<i>Mya arenaria</i>	bivalve	filter feeder	low numbers/persistent	higher number in South Bay - peak in summer	
<i>Venerupis philippinarum</i>	bivalve	filter feeder	low numbers/persistent	higher number in South Bay - peak in summer	
<i>Anguinella palmata</i>	bryozoa	filter feeder	common/persistent	spring peak	
<i>Nippoleucon hinumensis</i>	cumacean	surface deposit	common/persistent		
<i>Pyromma tuberculata</i>	decapod	debris & omnivore	common/persistent		
<i>Ilyanassa obsoleta</i>	gastropod	algae-scraiper	common/persistent		
<i>Sakuraeolis erosimensis</i>	gastropod	carnivore-hydroids	common/persistent		
<i>Euchone limicola</i>	polychaete	filter feeder	low number/persistent	higher number in South Bay - may be spreading north	
<i>Heteromastus filiformis</i>	polychaete	subsurface deposit	low numbers/persistent		
<i>Sabaco elongatus</i>	polychaete	subsurface deposit	low numbers/persistent	more common in south bay	
		tubed, surface to .8m			DWR 41a, REM Coyote Pt., REM PA, USGS PA

Table 3a. Common invertebrate taxa on soft bottom substrates (sand/silt/clay) in San Francisco Bay by salinity category and topography.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
EUHALINE					
Channel (Deep Water)					
<i>Ampelisca abilita</i>	amphipod	filter feeder	common/persistent		
<i>Monocorophium acherusicum</i>	filter & deposit feeder	tube to surface	common/persistent		
<i>Molgula marhattensis</i>	filter feeder	tube, free living	low numbers/persistent**	more common in South Bay	
<i>Tellina nucuoides</i>	subsurface deposit	surface	common/persistent		
<i>Mya arenaria</i>	filter feeder	subsurface	low numbers/persistent	common in Central Bay	RMP, EMAP, USGSEPA, REM SB Deep
<i>Styātula elongata</i>	filter feeder	surface-subsurface	common/persistent		
<i>Anneana</i> spp.	detritus & omnivore	tube to surface	common/persistent		
<i>Euchone limicola</i>	filter feeder	tube to surface	common/persistent		
<i>Exogone lourei</i>	carnivore	subsurface	common/persistent		
<i>Heteropodarke heteromorpha</i>	carnivore	free living subsurface	common/persistent		
<i>Mediomastus</i> spp	subsurface deposit	free living subsurface	common/persistent		
<i>Sabaco elongatus</i>	subsurface deposit	tubed, surface to .8m	low numbers/persistent	more common in South Bay	
Slough Channels					
<i>Tharyx</i> sp.	surface deposit	free living surface	common*		
<i>Ampelisca abilita</i>	filter feeder	tube to surface	common*		
<i>Grandidierella japonica</i>	filter & deposit feeder	tube, free living	common*		
<i>Monocorophium acherusicum</i>	filter & deposit feeder	tube, free living	common*		
<i>Macoma petalum</i>	filter & deposit feeder	subsurface	common*		
<i>Mya arenaria</i>	filter feeder	subsurface	common*		
Tubificidae - unidentified	subsurface deposit	free living subsurface	common*		EMAP 2000 46.4; EMAP 2001 63-65,96
<i>Cossura</i> spp.	polychaete		common*		
<i>Euchone limicola</i>	filter feeder	tube to surface	common*		
<i>Mediomastus californiensis</i>	subsurface deposit	free living subsurface	common*		
<i>Polydora cornuta</i>	filter & deposit feeder	tube to surface	common*		
<i>Pseudopolydora diopatra</i>	surface deposit	tube to surface	common*		
<i>Streblospio benedicti</i>	surface deposit	tube to surface	common*		
Harbors					
<i>Ampelisca abilita</i>	filter feeder	tube to surface	common*		
<i>Grandidierella japonica</i>	filter & deposit feeder	tube, free living	common*		
<i>Monocorophium acherusicum</i>	filter & deposit feeder	tube, free living	common*		
<i>Monocorophium alienense</i>	filter & deposit feeder	tube, free living	common*		
<i>Eudrella pacifica</i>	surface deposit	free living/surface-subsurface	common*		
<i>Nippoleucon hinumensis</i>	surface deposit	free living/surface-subsurface	common*		
Tubificidae - unidentified	subsurface deposit	free living subsurface	common*		
<i>Cossura</i> spp.	polychaete		common*		EMAP 2000 19.2, 19.3, 47.3, 47.4, 32.2, 32.3, 32.6, 25.3, 26.1, 26.2; EMAP 2001 57,62, 66-67, 45-49, 39, 94, 35, 89, 90
<i>Euchone limicola</i>	filter feeder	tube to surface	common*		
<i>Exogone lourei</i>	carnivore	free living	common*		
<i>Glycinde armigera</i>	carnivore	subsurface	common*		
<i>Heteromastus filiformis</i>	subsurface deposit	free living subsurface	common*		
<i>Mediomastus</i> spp.	subsurface deposit	free living subsurface	common*		
<i>Pseudopolydora diopatra</i>	surface deposit	tube to surface	common*		
<i>Sabaco elongatus</i>	subsurface deposit	tubed, surface to .8m	common*		
<i>Streblospio benedicti</i>	surface deposit	tube to surface	common*		
<i>Harmothoe imbricata</i>	carnivore	free living	common*		

Table 3a. Common invertebrate taxa on soft bottom substrates (sand/silt/clay) in San Francisco Bay by salinity category and topography.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
EUHALINE (continued)					
Shallow Subtidal					
<i>Ampelisca abdita</i>	filter feeder	surface tube	common/persistent		
<i>Corophium acherusicum</i>	filter & deposit feeder	tube, free living	common/persistent		
<i>Corophium heteroceratum</i>	filter & deposit feeder	tube, free living	common/persistent		
<i>Corophium insidiosum</i>	filter & deposit feeder	tube, free living	common/persistent		
<i>Musculista senhousia</i>	filter feeder	surface	low numbers/persistent**		
Tubificidae - unidentified	subsurface deposit	free living subsurface	low numbers/persistent		
<i>Phoronis</i> spp.	suspension feeder	tube to surface	common/persistent		
<i>Euchone limicola</i>	filter feeder	tube to surface	common/persistent		
<i>Exogone laurei</i>	carnivore	subsurface	common/persistent		
<i>Mediomastus</i> spp.	subsurface deposit	free living subsurface	common/persistent		
<i>Sphaerosyllis californiensis</i>	surface deposit	surface	common/persistent		
<i>Glycinde polygnatha</i>	carnivore	subsurface	low numbers/persistent		
<i>Leptochelia dubia</i>		surface	common/persistent		REM Berkeley, San Leandro, RMP

Table 3b. Common invertebrate taxa on soft bottom substrate (shell mix) in San Francisco Bay.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
<i>Diadumene lineata</i>	suspension feeder	surface-attached	present		
<i>Molgula manhattensis</i>	filter feeder	surface-attached	low numbers/persistent**		
<i>Amphibalanus improvisus</i>	filter feeder	surface-attached	common/persistent		
<i>Balanus crenatus</i>	filter feeder	surface-attached	common/persistent		
<i>Corbula amurensis</i>	filter feeder	surface-subsurface	low numbers/persistent**		
<i>Musculista senhousia</i>	filter feeder	surface-attached	low numbers/persistent**		
<i>Venerupis philippinarum</i>	filter feeder	subsurface	low numbers/persistent**		
<i>Amphipholis squamata</i>	filter feeder	subsurface	low numbers/persistent**		
<i>Watersipora subtorquata</i>	surface deposit	surface-subsurface	present		
<i>Eriocher sinensis</i>	omnivores	surface-motile	present		
<i>Hemigrapsus oregonensis</i>	omnivores	surface-motile	present		
<i>Crepidula convexa</i>	scraper, algae	surface-motile	common/persistent		
<i>Crepidula plana</i>	scraper, algae	surface-motile	common/persistent		
<i>Clytia aff. hemisphaerica</i>	suspension feeder	surface-attached	common/persistent		
<i>Obelia longissima</i>	suspension feeder	surface-attached	present		
<i>Gnoringosphaeroma oregonense</i>	surface deposit	surface-motile	present		
<i>Sphaeroma quonianum</i>	algae	burrows, wood, mud	present		
<i>Mytilus trossulus/gallopvinctialis</i>	filter feeder	surface-attached	present		
<i>Haminoea japonica</i>	opistobranch	surface-motile	present		
<i>Ammonothea hilgendorfi</i>	carnivore	surface-motile	present		
<i>Assiminea californica</i>	surface deposit	surface-motile	low numbers/persistent**		
<i>Urosalpinx cinerea</i>	carnivore	surface-motile	present		
<i>Diadumene lineata</i>	suspension feeder	surface-attached	present		

REM San Leandro (some samples) USGS SM35, SM42; Rapid Assessment Intertidal Benthos - epifauna at two or more sites

Table 3c. Common invertebrate taxa on hard bottom substrate (natural and artificial) in San Francisco Bay.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
MESOHALINE					
<i>Amphitoe valida</i>	algae	surface tubes among algae	present		
<i>Eogammarus confervicolus</i>	algae	surface-motile	present		
<i>Monocorophium acherusicum</i>	filter & deposit feeder	tube, free living	present		
<i>Diadumene</i> sp.	suspension feeder	surface-attached	present		
<i>Amphibalanus improvisus</i>	filter feeder	surface-attached	present		
<i>Conopeum</i> cf. <i>tenuissimum</i>	suspension feeder	surface-attached	present		
<i>Gamella franciscana</i>	suspension feeder	surface-attached	present		
<i>Gnirimosphaeroma oregonensis</i>	surface deposit	surface-motile	present		
<i>Pseudosphaeroma campbellensis</i>		surface-motile	present		
<i>Sphaeroma caoianum</i>	algae	surface-motile	present		
<i>Symidotea laevidorsalis</i>	surface deposit	surface-motile	present		
<i>Mytilus trossulus/galloprovincialis</i>	filter feeder	surface-attached	present		
<i>Okenia plana</i>	carnivore - bryozoans	surface-motile	present		
<i>Ficopomatus enigmaticus</i>	filter feeder	surface-attached	common/persistent		
<i>Clathria prolifera</i>	suspension feeder	surface-attached	common/persistent		
<i>Halichondria bowerbanki</i>	suspension feeder	surface-attached	common/persistent		
<i>Haliciona</i> sp.	suspension feeder	surface-attached	common/persistent		
POLYHALINE					
<i>Caprella mutica</i>	filter feeder	surface-motile	common/persistent		
<i>Iniscocallope derzhavini</i>		surface-motile	present		
<i>Jassa marmorata</i>		surface-motile	present		
<i>Monocorophium acherusicum</i>	filter & deposit feeder	tube, free living	present		
<i>Stenothoe valida</i>		surface-motile	present		
<i>Diadumene lineata</i>	suspension feeder	surface-attached	present		
<i>Diadumene</i> sp.	suspension feeder	surface-attached	present		
<i>Molgula manhattensis</i>	filter feeder	surface-attached	present		
<i>Amphibalanus "amphitrite"</i>	filter feeder	surface-attached	present		
<i>Amphibalanus improvisus</i>	filter feeder	surface-attached	present		
<i>Balanus glandula</i>	filter feeder	surface-attached	present		
<i>Musculista senhousia</i>	filter feeder	surface-attached	present		
<i>Conopeum</i> cf. <i>tenuissimum</i>	suspension feeder	surface-attached	present		
<i>Schizoporella unicornis</i>	suspension feeder	surface-attached	present		
<i>Polycera hedgepethi</i>	suspension feeder	surface-motile	present		
<i>Ectopleura crocea</i>	carnivore	surface-attached	present		
<i>Gamella franciscana</i>	suspension feeder	surface-attached	present		
<i>Paranthura japonica</i>		surface-motile	present		
<i>Synidotea laevidorsalis</i>	isopod	surface-motile	present		
<i>Mytilus trossulus/galloprovincialis</i>	filter feeder	surface-attached	present		
<i>Ostrea conchaphila</i>	oyster	surface-attached	present		
<i>Neanthes succinea</i>	polychaete	surface	present		
<i>Clathria prolifera</i>	suspension feeder	surface-attached	common/persistent		
<i>Haliciona</i> sp.	suspension feeder	surface-attached	common/persistent		
<i>Leucilla nuttingi</i>	suspension feeder	surface-attached	common/persistent		
<i>Halichondria bowerbanki</i>	suspension feeder	surface-attached	present		
<i>Mycale macginitiei</i>	suspension feeder	surface-attached	present		
<i>Sycon</i> sp.	suspension feeder	surface-attached	present		

Rapid Assessment Channel substrate and pilings habitat - seen at half or more of sites

Rapid Assessment Channel substrate and pilings habitat - seen at half or more of sites

Table 3c. Common invertebrate taxa on hard bottom substrate (natural and artificial) in San Francisco Bay.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
POLYHALINE (continued)					
<i>Ascidia zara</i>	filter feeder	surface-attached	common/persistent		
<i>Botrylloides</i> spp.	filter feeder	surface-attached	common/persistent		
<i>Botryllus</i> spp.	filter feeder	surface-attached	common/persistent		
<i>Ciona intestinalis</i>	filter feeder	surface-attached	common/persistent		
<i>Ciona savignyi</i>	filter feeder	surface-attached	common/persistent		
<i>Styela clava</i>	filter feeder	surface-attached	common/persistent		
<i>Bugula neritina</i>	filter feeder	surface-attached	common/persistent		
<i>Bugula stolonifera</i>	filter feeder	surface-attached	common/persistent		
<i>Cryptosula pallasiana</i>	filter feeder	surface-attached	common/persistent		
<i>Schizoporella unicornis</i>	filter feeder	surface-attached	common/persistent		
<i>Waterspora subtorquata</i>	filter feeder	surface-attached	common/persistent		
Slough Channels					
<i>Amphibalanus improvisus</i>	filter feeder	surface-attached	present		Rapid Assessment Channel substrate and pilings habitat - seen at half or more of sites
<i>Victorella pavidia</i>	suspension feeder	surface-attached	present		Rapid Assessment Channel substrate and pilings habitat - seen at half or more of sites
EUHALINE					
<i>Iniscocallope derzhavini</i>			present		
<i>Jassa marmorata</i>		tube	present		
<i>Monocorophium achenusicum</i>	filter & deposit feeder	free, among hydroids	present		
<i>Podocerus brazilensis</i>			present		
<i>Stenothoe valida</i>			present		
<i>Metridium senile</i>	suspension feeder	surface-attached	present		
<i>Amphibalanus improvisus</i>	filter feeder	surface-attached	present		
<i>Balanus crenatus</i>	filter feeder	surface-attached	present		
<i>Balanus glandula</i>	filter feeder	surface-attached	present		
<i>Hiatella artica</i>	filter feeder	surface-attached	present		
<i>Modiolus rectus</i>	filter feeder	surface-attached	present		
<i>Musculista senhousia</i>	filter feeder	surface-attached	present		
<i>Alicyoniidum parasiticum</i>	suspension feeder	surface-attached	present		
<i>Farella elongata</i>	suspension feeder	surface-attached	present		
<i>Hippothoe hyalina</i>	suspension feeder	surface-attached	present		
<i>Caprella equibrata</i>	detritivores, carnivores, deposit feeders	surface-motile	present		
<i>Caprella penantis</i>	detritivores, carnivores, deposit feeders	surface-motile	present		
<i>Caprella verrucosa</i>	detritivores, carnivores, deposit feeders	surface-motile	present		
<i>Ecioptleura crocea</i>	suspension feeder	surface-attached	present		
<i>Mytilus trossulus/galloprovincialis</i>	filter feeder	surface-attached	present		
<i>Nereis</i> sp (not succinea)	omnivore	surface-motile	present		
Phylodocidae	carnivore, scavengers	surface, free living	present		
Terebellidae	surface deposit feeder		present		
<i>Typosyllis nipponica</i>	carnivore	surface-motile	present		
<i>Harmothoe imbricata</i>	carnivore	surface-motile	present		
Polyoidae	carnivore	surface-motile	present		
<i>Haliclioniidae bowerbanki</i>	suspension feeder	surface-attached	present		

Table 3d. Common invertebrate taxa on soft bottom substrate (shell mix) in San Francisco Bay.

Species	Feeding Mode	Growth Position	Relative Frequency	Seasonal Notes	Reference Locations
<i>Ampithoe valida</i>	herbivore	on leaves			
<i>Caprella mutica</i>	filter feeder	on leaves			
<i>Corophium</i> sp.	filter feeder	on leaves			
<i>Musculista senhousia</i>	filter feeder	among roots			
<i>Conopeum tenuissimum</i>	suspension feeder	encrusting on leaves			

We summarize the fishes found in San Francisco Bay based on review of ongoing surveys and existing reports by the California Department of Fish and Game (CDFG) (CDFG 2001; Table 4). CDFG began regular monitoring of fish and macroinvertebrate populations throughout the Bay in 1980, including the San Francisco Bay Study midwater and otter trawl surveys conducted from South San Francisco Bay to the Western Delta and beach seine data collected in South Bay from 1980 to 1987. CDFG currently samples at 52 sites throughout the Bay and Delta with nine sites within the South Bay (Orsi *et al.*, 1999). Results of this sampling is summarized for 1980 to 1995 by Orsi *et al.* (1999) and the San Francisco Bay Fish Index (The Bay Institute 2003).

Fish taxa by habitat are described by habitat category and salinity range, with a slight variation from the Venice Symposium categories. Table 4 includes species by salinity category within each habitat type. The table also includes information on whether a species is resident, using the San Francisco Bay during a part of its life history (e.g., juvenile nursery, adult), or whether the species are migratory, moving through San Francisco Bay between freshwater tributaries and the Pacific Ocean. Species that reproduce in freshwater and migrate to the ocean after birth are called anadromous. Species that reproduce in the ocean and migrate to freshwater after birth are called catadromous.

Table 4. Common fish and mobile invertebrate species of San Francisco Bay after K. Hieb (pers. comm.), Merkel & Associates (2005), and Entrix (1998).

Salinity	Scientific Name	Common Name	Status	Subtidal Habitat Type							Water Column				
				Mud	Sand	Pebble / Cobble	Hard Bottom Substrate	Shellfish Beds	Algae	Eelgrass		Plant Beds	Surf Grass	Widgeon Grass	
Euhaline (25-35 ppt)	<i>Crangon nigromaculata</i>	blackspotted bay shrimp	N	J, A										R	
	<i>Lissocrangon stylirostris</i>	smooth bay shrimp	N	J, A										R	
	<i>Cancer gracilis</i>	graceful rock crab	N	J, A										R	
	<i>Cancer antennarius</i>	Pacific rock crab	N		J, A	J, A								R	
	<i>Cancer productus</i>	red rock crab	N		J, A	J, A								R	
	<i>Mustelus henlei</i>	brown smoothhound	N	J, A											
	<i>Triakis semifasciata</i>	leopard shark	N	J, A											
	<i>Raja binoculata</i>	big skate	N	J, A											
	<i>Myliobatis californica</i>	bat ray	N	J, A											
	<i>Sardinops sagax</i>	Pacific sardine	N							J, A					
	<i>Engraulis mordax</i>	northern anchovy	N												
	<i>Porichthys notatus</i>	plainfin midshipman	N	J, A											
	<i>Leuresthes tenuis</i>	California grunion	N												
	<i>Atherinopsis californiensis</i>	jacksmelt	N	R											
	<i>Atherinops affinis</i>	topsmelt	N	R											
	<i>Syngnathus leptorhynchus</i>	bay pipefish	N												
	<i>Sebastes auriculatus</i>	brown rockfish	N			J, A									
	<i>Ophiodon elongatus</i>	lingcod	N			J									
	<i>Geyonemus lineatus</i>	white croaker	N	J, A											
	<i>Rhacochilus vacca</i>	pile perch	N	J, A											
	<i>Cymatogaster aggregata</i>	shiner perch	N												
	<i>Hyperprosopon argenteum</i>	walleye surfperch	N												
	<i>Phanerodon furcatus</i>	white seaperch	N												
	<i>Lepidogobius lepidus</i>	bay goby	N	J, A											
	<i>Paralichthys californicus</i>	California halibut	N	J, A	J, A										
	<i>Oithichthys sigmaeus</i>	speckled sanddab	N	J	J										
	Polyhaline (20-28 ppt)	<i>Crangon nigricauda</i>	blacktail bay shrimp	N	J, A										
		<i>Cancer magister</i>	Dungeness crab	N	J										R
<i>Clupea pallasii</i>		Pacific herring	N											R, J, A	
<i>Leptocottus armatus</i>		Pacific staghorn sculpin	N	J, A										R	
<i>Parophris vetulus</i>		English sole	N	J, A											

R Reproduction, includes eggs on substrate and pelagic eggs and larvae, but not pupping areas

J Juvenile nursery area

A Adult residence or forage area

N Native species

I Introduced species

☐ Catadromous: rears in freshwater, reproduces in brackish salinities or higher

☐ Anadromous: reproduces in freshwater, rears in brackish salinities or higher, or migrates through San Francisco Bay as juveniles

Table 4. Common fish and mobile invertebrate species of San Francisco Bay after K. Hieb (pers. comm.), Merkel & Associates (2005), and Entrix (1998).

Salinity	Scientific Name	Common Name	Status	Subtidal Habitat Type														
				Mud	Sand	Pebble/ Cobble	Hard Bottom Substrate	Shellfish Beds	Algae	Eelgrass	Plant Beds	Surf Grass	Widgeon Grass	Water Column				
Oligohaline / Mesohaline (0.5-20 ppt)	<i>Palaemon macrodactylus</i>	oriental shrimp	I	J, A												R		
	<i>Crangon franciscorum</i>	California bay shrimp	N	J, A			J, A									R		
	<i>Acipenser medirostris</i>	green sturgeon	N	J, A														
	<i>Acipenser transmontanus</i>	white sturgeon	N	J, A														
	<i>Pogonichthys macrolepidotus</i>	spittail	N	J, A														
	<i>Hypomesus transpacificus</i>	delta smelt	N														R, J, A	
	<i>Sprinichus thaleichthys</i>	longfin smelt	N														R, J, A	
	<i>Morone saxatilis</i>	striped bass	I	J, A													R, J, A	
	<i>Acanthogobius flavimanus</i>	yellowfin goby	I	J, A													R	
	<i>Platichthys stellatus</i>	starry flounder	N	J, A		J, A												
	Fresh (<0.5 ppt)	<i>Exopalaemon modestus</i>	Siberian prawn	I	J, A													R
		<i>Eriocheir sinensis</i>	Chinese mitten crab	I	J, A			J, A										R
		<i>Lampetra tridentata</i>	Pacific lamprey	N														J, A
		<i>Lampetra ayresii</i>	river lamprey	N														J, A
<i>Alosa sapidissima</i>		American shad	I														R, J, A	
<i>Dorosoma petenense</i>		threadfin shad	I														R, J, A	
<i>Ictalurus punctatus</i>		channel catfish	I	J, A														
<i>Ameiurus catus</i>		white catfish	I	J, A														
<i>Oncorhynchus tshawytscha</i>		Chinook salmon	N														J, A	
<i>Oncorhynchus mykiss</i>		steelhead	N														J, A	
<i>Cottus asper</i>		prickly sculpin	N	J, A				J, A									R	

R Reproduction, includes eggs on substrate and pelagic eggs and larvae, but not pupping areas

J Juvenile nursery area

A Adult residence or forage area

N Native species

I Introduced species

■ Catastrophic: rears in freshwater, reproduces in brackish salinities or higher

■ Anadromous: reproduces in freshwater, rears in brackish salinities or higher, or migrates through San Francisco Bay as juveniles

Bird and marine mammal taxa can not be characterized by salinity category, as individuals move throughout the Bay independent of salinity in search of prey items. Surveys of bird and marine mammal abundance and distribution in San Francisco Bay focus on seasonal counts or abundances at shore-based nests or haul-out sites. A general discussion of bird and marine mammal use of San Francisco Bay is included here because of the importance of subtidal habitats for their breeding and feeding. Information on specific habitat use is provided below in the habitat sections.

The bird species that utilize subtidal habitats in San Francisco Bay are listed in Table 5. The bird communities discussed in this chapter are divided into foraging guilds, including diving benthivores (feed in deeper water on benthic invertebrates), dabblers (feed in the upper water column of the shallow subtidal), piscivores (fish consumers), and opportunistic predators (Takekawa *et al.* 2001). San Francisco Bay is a critical Pacific Flyway wintering and stop-over site for migratory birds that depend on subtidal habitats. More than 300,000 wintering waterfowl use San Francisco Bay and associated ponds each year (Accurso 1992). Nearly 75 percent of these birds are bay and sea ducks that forage on benthic invertebrates in a variety of subtidal habitats.

During the spring and summer months, various seabird species use the Bay for breeding and foraging. Alameda Point hosts the largest least tern (*Sterna antillarum*) colony in the Bay, comprising an estimated 424 breeding pairs in 2005 (Hurt 2006). In 2005, approximately 450 nesting attempts for double-crested cormorant (*Phalacrocorax auritus*) were recorded on the San Francisco-Oakland Bay Bridge, and the Richmond-San Rafael Bridge had over 350 attempts (PRBO, unpub. data). Brandt's cormorants (*Phalacrocorax penicillatus*), pelagic cormorants (*P. pelagicus*), pigeon guillemots (*Cepphus columba*), western gulls (*Larus occidentalis*), and since 2004, California gulls (*L. californicus*), breed on Alcatraz Island in central San Francisco Bay. Recent Alcatraz population estimates are 1010 breeding attempts for Brandt's cormorants, 7 breeding attempts for pelagic cormorants, 30 to 40 breeding sites for pigeon guillemots, 1033 breeding pairs for western gulls, and 21 nests for

California gulls (Acosta and Thayer 2006). While the Brandt's cormorant, western gull, and California gull colonies on Alcatraz have been expanding, pelagic cormorant and pigeon guillemot populations have been in decline (Acosta and Thayer 2006).

The most numerous diving duck species wintering on San Francisco Bay are greater scaup (*Aythya marila*), lesser scaup (*A. affinis*), canvasback (*A. valisineria*), and surf scoter (*Melanitta perspicillata*). Based on USFWS mid-winter waterfowl surveys, an average of 44 percent of greater and lesser scaup, 42 percent of scoters (three species combined), and 35 percent of canvasback in the lower Pacific Flyway winter in the San Francisco Bay estuary (Trost 2002). The greater scaup, lesser scaup, and surf scoter populations are all undergoing continental decline (USFWS 2006, Sea Duck Joint Venture Management Board 2001, Afton and Anderson 2001, Austin *et al.* 2000). Nationwide, total canvasback numbers are currently holding at their management goal of approximately 500,000 birds (USFWS 2006), but canvasback in San Francisco Bay are declining (Trost 2002). This species is closely monitored and hunting restrictions are frequently imposed (USFWS 2006). Causes of decline for these species are difficult to discern; however, San Francisco Bay remains a crucial over-wintering location for all four of them.

Two endangered bird species, the least tern and the brown pelican (*Pelecanus occidentalis*) are also migratory species that utilize San Francisco Bay. The largest least tern breeding colony in California north of San Luis Obispo County is located in San Francisco Bay. Due to their breeding habitat preferences of open gravel areas with sparse vegetation, human development has significantly decreased available breeding habitat in the San Francisco Bay area (Goals Report 2000). While the least tern population is increasing, breeding success has shown signs of decline since the mid-1990s (Elliott *et al.* 2006). Historic abundance of pelicans within San Francisco Bay is unknown. In 2006, over 2000 brown pelicans were observed on the Alameda breakwater island in July and August (Euing 2006).

Table 5. Common bird species that use subtidal habitats in San Francisco Bay.

Foraging Guild	Scientific Name	Common Name	Time of Year	Subtidal Habitat Type												
				Soft Bottom Substrate	Hard Bottom Substrate	Shellfish Beds	Algal Beds	Eelgrass	Plant Beds	Surf Grass	Widgeon Grass	Water Column				
				Mud	Sand	Cobble / Pebble										
Dabblers	<i>Anas clypeata</i>	northern shoveler	winter, more intertidal													
	<i>Anas strepera</i>	gadwall	winter, year-round, more intertidal												R, F	
	<i>Anas acuta</i>	northern pintail	winter, more intertidal												R	
	<i>Anas platyrhynchos</i>	mallard	year-round, more intertidal												R, F	
Diving Benthivores	<i>Podiceps nigricollis</i>	eared grebe*	winter												R, F	
	<i>Podiceps auritus</i>	horned grebe*	winter, uncommon												R, F	
	<i>Aythya marila</i>	greater scaup	winter		F	F									R	
	<i>Aythya affinis</i>	lesser scaup	winter	F	F	F									R	
	<i>Aythya valisineria</i>	canvasback	winter	F	F	F									R	
	<i>Aythya americana</i>	redhead	winter, uncommon												R	
	<i>Oxyura jamaicensis</i>	ruddy duck	winter	F	F	F									R	
	<i>Bucephala clangula</i>	common goldeneye	winter	F	F	F									R	
	<i>Bucephala islandica</i>	Barrow's goldeneye	winter	F	F	F									R	
	<i>Bucephala albeola</i>	bufflehead	winter	F	F	F									R	
	<i>Melanitta fusca</i>	white-winged scoter	winter, uncommon	F	F	F									R	
	<i>Melanitta perspicillata</i>	surf scoter	winter	F	F	F									R	
	Piscivores	<i>Podilymbus podiceps</i>	pie-billed grebe**	year-round												R, F
		<i>Podiceps griseogena</i>	red-necked grebe**	winter, uncommon												R, F
<i>Gavia pacifica</i>		Pacific loon	winter												R, F	
<i>Aechmophorus occidentalis</i>		Western grebe	winter												R, F	
<i>Aechmophorus clarkii</i>		Clark's grebe	winter												R, F	
<i>Pelecanus erythrorhynchos</i>		American white pelican	winter, post-breeding												R, F	
<i>Pelecanus occidentalis</i>		brown pelican	winter, post-breeding, juveniles year-round												R, F	
<i>Phalacrocorax auritus</i>		double-crested cormorant	year-round	F	F	F									R, F	
<i>Phalacrocorax penicillatus</i>		Brandt's cormorant	year-round	F	F	F									R, F	
<i>Phalacrocorax pelagicus</i>		pelagic cormorant	year-round	F	F	F									R, F	
<i>Uria aalge</i>		common murre	year-round, but concentrated more in late summer w/chicks												R, F	
<i>Cephus columba</i>		pigeon guillemot	summer	F	F	F									R, F	
<i>Mergus merganser</i>		common merganser	year-round in upper estuary												R, F	
<i>Mergus serrator</i>		red-breasted merganser	winter												R, F	
<i>Pandion haliaetus</i>	osprey	year-round												F		
<i>Sterna caspia</i>	Caspian tern	summer												F		
<i>Sterna forsteri</i>	Forster's tern	year-round	F											F		
<i>Sterna elegans</i>	elegant tern	migration												F		
<i>Sterna anthillanum</i>	least tern	summer	F	F	F									F		
<i>Rynchops niger</i>	black skimmer	summer												F		

* Also feeds on invertebrates and fish on surface or in water column
 ** Also feeds on benthic invertebrates
 ... Members of this guild may also feed in the intertidal or in terrestrial habitats, but are surface feeders on the open water

F Forage
 R Roost or Raft

Table 5. Common bird species that use subtidal habitats in San Francisco Bay.

Foraging Guild	Scientific Name	Common Name	Time of Year	Subtidal Habitat Type																			
				Soft Bottom Substrate	Hard Bottom Substrate	Shellfish Beds	Algal Beds	Plant Beds	Surf Grass	Widgeon Grass													
				Mud	Sand	Pebble / Cobble																	
Foraging Guild ***	<i>Larus occidentalis</i>	Western gull	year-round																				
	<i>Larus californicus</i>	California gull	year-round																				
	<i>Larus delawarensis</i>	ring-billed gull	winter	F	F																		
	<i>Larus philadelphia</i>	Bonaparte's gull	winter	F	F																		
	<i>Larus argentatus</i>	herring gull	winter	F	F			F															
	<i>Larus glaucescens</i>	glaucous-winged gull	winter	F	F																		
	<i>Larus canus</i>	mew gull	winter	F	F																		
	<i>Larus heermanni</i>	Heermann's gull	winter, summer juveniles & post-breeding adults																				
		<i>Phalaropus lobatus</i>	red-necked phalarope	winter, uncommon																			
Other	<i>Branta canadensis</i>	Canada goose	year-round, more intertidal																				
	<i>Fulica americana</i>	American coot	year-round, more intertidal																				

* Also feeds on invertebrates and fish on surface or in water column

** Also feeds on benthic invertebrates

*** Members of this guild may also feed in the intertidal or in terrestrial habitats, but are surface feeders on the open water

F Forage
R Roost or Raft

Seven species of marine mammals occur in San Francisco Bay (Table 6). The harbor seal (*Phoca vitulina richardii*), California sea lion (*Zalophus californianus*), harbor porpoise (*Phocoena phocoena*), and the Eastern Pacific stock of the gray whale (*Eschrichtius robustus*) are the most common. In general, habitat association of marine mammals in San Francisco Bay is related to distribution of prey species. For example, gray whales might occur over mud habitat where they suck up invertebrates, and seals occur in habitats where fish prey concentrate, such as eelgrass beds. Additionally, harbor seals and sea lions use various intertidal substrates that are exposed at low to medium tide levels for resting and breeding.

Harbor seals are the only year-round resident of San Francisco Bay, using the area for breeding, pupping, foraging, and refugia. Harbor seals have been observed in waters as far inland as Sacramento, though use of habitat north of Suisun Bay is irregular (Goals Report 2000). Harbor seals haul out onshore at specific locations within San Francisco Bay, utilizing mostly rocks and mud flats exposed at low tides, sloughs, islands, and beaches, likely in proximity to food resources and distant from human activities (Allen 1991). Haul-out use varies by day, season, and year (Green *et al.* 2006). The primary colonies within San Francisco Bay are at Castro Rocks in the San Pablo Bay, Yerba Buena Island in Central Bay and Mowry Slough in the South Bay. Some colonies are only accessible at medium to high tides, such as the tidal mudflats and pickleweed marshes of Mowry and Newark Sloughs. Seals have abandoned several locations used in the past (e.g., Strawberry Spit) within San Francisco Bay due to human activities (Allen 1991).

Current population estimates are derived from aerial and land-based surveys conducted by the CDFG, the National Marine Fisheries Service, the U.S. Fish and Wildlife Service, San Francisco State University, the Marine Mammal Center of Sausalito, and the National Park Service (Lowry and Forney 2005, Vanderhoof and Allen 2005, Manna *et al.* 2006, Green *et al.* 2006). Surveys estimate the resident San Francisco Bay population to be between 500 and 700 individuals (Richmond-Bridge Harbor Seal Survey 2001, Grigg *et al.* 2004). Between 1965 and 2002, harbor seals are thought to have

increased slightly in abundance at two sites in San Francisco Bay and to have abandoned one site completely (Allen 1991, Grigg *et al.* 2004). The average number of seals counted during the 2002 breeding/molt seasons was 117.5 at Castro Rocks, 96.6 at Yerba Buena Island and 147.6 at Mowry Slough. Because of their year-round residency within the Bay, harbor seals are exposed to elevated pollutant loads compared to outer coastal seals and have been a subject of research on marine mammal health (Risebrough *et al.* 1981, Kopec and Harvey 1995). There is a high incidence of red colored harbor seals in San Francisco Bay (~40%), a condition related to iron oxide deposition on the seal fur, and the condition may be related to elevated pollutant levels (Allen *et al.* 1993, Kopec and Harvey 1995). Seals forage throughout the Bay and feed on what is seasonally abundant such as Pacific herring during the winter spawn. They eat small schooling fish such as smelt, anchovies and herring, rockfish, sculpin, perch, and midshipman, and invertebrates such as squid and mysid shrimp. Surprisingly, they also forage on non-native species such as yellowfin goby. During the breeding and molt seasons, harbor seals likely do not forage more than a couple of miles from their colonies because of the requirement of attending pups, mating, or molting.

California sea lions use San Francisco Bay for refugia and foraging, but do not breed or pup within the Bay. California sea lions are most abundant within the Bay while migrating to and from their primary breeding areas on the Farallon and California Channel Islands, and when Pacific herring and salmon are spawning in the bay. Sea lions can travel far up into the Delta, but most concentrate feeding in Central Bay and where herring spawn. Similar to harbor seals, sea lions haul-out onshore, often utilizing anthropogenic structures such as boat docks and navigational buoys, although individuals may also haul out also on islands within San Francisco Bay, such as Alcatraz and Angel Islands. Historically, sea lions hauled out on Seal Rock near the Cliff House, but they abandoned the site in the 1980s. The largest California sea lion haul-out in San Francisco Bay is at the Port of San Francisco Pier 39, where up to 800 sea lions have been counted. Sea lions often float on the surface in large groups of 10 to 20 after feeding.

The harbor porpoise is a near-shore species, commonly observed near the Golden Gate Bridge and areas of Central Bay. A distinct genetic stock, the San Francisco-Russian River Stock, ranges from Point Arena to Monterey Bay. There are no good harbor porpoise population estimates. In the early 1980s, hundreds were killed incidentally in gill net fisheries, but since those fisheries were shifted offshore, few deaths have been reported. Harbor porpoise eat mostly small schooling fish and invertebrates, and along with seals and sea lions, will feed on herring and anchovies.

The Eastern Pacific gray whale migrates between calving grounds in Baja, Mexico to primary feeding grounds in Alaska and Canada on an annual basis. Gray whales are commonly sighted near the Golden Gate during peak migration periods (northward migration in spring and southward migration in winter), and annually a few individuals are observed within the Bay (Green *et al.* 2006). Occasionally, gray whales on their migration will forage in nearshore waters such as San Francisco Bay, Drakes Bay, Tomales Bay, and Monterey Bay. They prey mostly on invertebrates that live on or in soft sediments.

Table 6. Common marine mammal species of San Francisco Bay.

Scientific Name	Common Name	Subtidal Habitat Type									
		Soft Bottom Substrate		Hard Bottom Substrate	Shellfish Beds	Plant Beds		Water Column			
		Mud	Sand	Pebble / Cobble			Algal Beds	Eelgrass	Surf Grass	Widgeon Grass	
<i>Eumetopias jubatus</i>	Steller sea lion	x	x	x			x	x	x	x	
<i>Zalophus californianus</i>	California sea lion	x	x	x			x	x	x	x	
<i>Phoca vitulina</i>	Harbor seal	x	x	x					x	x	
<i>Eschrichtius robustus</i>	Gray whale	x	x			x			x	x	
<i>Megaptera novaeangliae</i>	Humpback whale	x	x						x	x	
<i>Phocoena phocoena</i>	Harbor porpoise	x	x	x					x	x	
<i>Enhydra lutra</i>	California sea otter	x	x	x			x	x	x	x	

Note: habitat association is based on location of prey items

Habitat Type and Associated Biological Assemblages

SOFT BOTTOM SUBSTRATE

(by Janet Thompson, Kathryn Hieb, Katie McGourty, Natalie Cosentino-Manning, Susan Wainwright-De La Cruz, Meredith Elliot and Sarah Allen)

General Description

Soft bottom substrate habitats include mud/silt/clay (particles 0.001 to 0.062 mm in diameter), sand (particles 0.062 to 2.0 mm in diameter), pebble/cobble (particles 2 to 256 mm in diameter), and shell mix (a mix of mud/silt/clay or sand and shell fragments). Soft bottom substrates are characterized by a lack of large stable surfaces for plant and animal attachment. Exposure to wave and current action, temperature, salinity, and light penetration determine the composition and distribution of organisms within the sediments (USGS 1998). While comprehensive bottom surveys are unavailable for San Francisco Bay, survey data (Greene and Bizzaro 2003, San Francisco Bay Regional Monitoring Program (RMP), San Francisco Bay Regional Effects Monitoring Program (REM), National Oceanic and Atmospheric Association's Regional Monitoring Program (EMAP)) and anecdotal information from scientific sampling and commercial and recreational fishing, and from commercial uses of the Bay (e.g., dredging for vessel traffic, sediment extraction) indicate unconsolidated sediments are present throughout and are the most common substrate type in the San Francisco Bay (Figures 4, 10).

Unconsolidated bottoms provide an array of functions to San Francisco Bay flora and fauna. Mud/silt/clay substrates provide a source and sink of suspended sediment; they store, transport, transform and make available trace elements and organic compounds (e.g., carbon, nitrogen, phosphorous, sulfur); they provide a substrate for invertebrate epifauna (organisms on the surface) and infauna (sessile organisms which live within sediments); and they provide a substrate for fish, birds, and marine mammals to reproduce, rear, feed, and grow. Sand habitats provide similar functions for invertebrate epifauna and infauna, fish, birds, and marine mammals, and provide for bed load transport. Shell mix and pebble-cobble habitats provide similar functions for invertebrate epifauna and infauna, fish, birds, and marine mammals, but the substrate is more three-dimensional and more stable, and so provides a higher level of refuge from predators, and a substrate for attached organisms.



Figure 10. Soft bottom and hard bottom substrates in San Francisco Bay interpreted from multibeam bathymetric images and side-scan sonar mosaics after Greene and Bizarro (2003).

Taxonomic Groups

Plants

Plants associated with soft bottom habitats include algae and submerged aquatic vegetation (SAV; Table 2). In most cases, these plants need coarse grain particles such as pebbles or shells to become established in this habitat. There are very few species that exclusively inhabit soft bottoms and when they do, they are mainly found in sheltered embayments such as Richardson Bay (Josselyn and West 1985). Such genera include the macroalgae *Ulva/Enteromorpha*, which are found in the euhaline to the mesohaline parts of the Bay.

Eelgrass (*Zostera marina*) is found in both fine and coarse grain sediments. Eelgrass can grow in thick, muddy sediments like those off Point Molate and Point San Pablo, or in sandy sediments like those off Crown Beach. Very little is known about relationships between grain size and distribution of eelgrass around San Francisco Bay.

Invertebrates

In this report, we aggregate the sessile invertebrate infauna into assemblages by location as a proxy for salinity (Table 3). Because of their usefulness as lifetime integrators of *in situ* water and sediment quality, benthic communities are usually monitored at fixed locations over time. This monitoring scheme has been used in San Francisco Bay, thus, our discussion reflects data from these locations. Data from monitoring stations have been grouped based on location that is representative of a given salinity category during normal hydrologic years. As discussed in the Physical Description above, we can anticipate the general salinity range within a specific area of San Francisco Bay during different seasons in normal water years. Soft bottom invertebrate taxa at the fresh and euhaline ends of San Francisco Bay traditionally have the most stable species composition. Communities between these end points show the most seasonal and inter-annual variability, reflecting changes in freshwater flow at tidal and seasonal time scales. However, all communities change with flood and drought events.

When possible, seasonal and inter-annual patterns are noted for dominant species in each assemblage within each habitat. Using existing information, we further divide these assemblages into main channel, channel edge, slough channel, and shallow subtidal topographical regions. Data is not always available for all topographic regions at all locations. These sub-assemblages allow us to consider benthic communities by feeding-type and availability to predators whose feeding capabilities are depth and habitat specific (e.g., diving ducks that feed only on surface dwellers), and to identify assemblages most likely to be affected by depth-specific anthropogenic alterations of the system. For the most part, information on invertebrate taxa on soft bottom substrate does not allow us to separate taxa found on specific sediment types (i.e., sand, silt, cobble). There is information, however, on taxa found on shell mix, which is discussed separately at the end of this section.

There is no recent summary of the soft bottom benthic community, and it is likely that all summaries, including this one, will be outdated soon after they are written due to the high rate of non-native species introductions in the system (Cohen and Carlton 1995, Cohen and Carlton 1998). The only comprehensive summary of any compartment of the soft bottom benthos was written in 1986 (Nichols and Pamamat 1988), just prior to the invasion of *Corbula* (formerly known as *Potamocorbula*) *amurensis*, a bivalve that altered the soft-bottom benthos community structure (Nichols *et al.* 1990, Peterson 2002) and changed the trophic state of northern San Francisco Bay (Alpine and Cloern 1992, Thompson 2005). Two long term studies demonstrate that the soft bottom infaunal community has been substantially changed by exotic species: (1) Only 2 of the 58 species seen in Grizzly Bay are confirmed native species with 25 of the species being confirmed as exotic and 3 as cryptogenic species (Peterson 2002); (2) Of the 37 species seen on a mudflat in South Bay, only one is a confirmed native species with 27 species confirmed as exotic and 3 as cryptogenic (Nichols and Thompson 1985 updated with new findings in Cohen and Carlton 1995). Lee *et al.* (2003) included data from throughout the bay, including the less-invaded euhaline section, and reported that 19 to 41 percent of

the species during a four-year period were non-indigenous or cryptogenic.

The data used to describe infaunal benthos in this system are limited to the post-*Corbula* invasion period due to the dominance of the bivalve in the communities where it resides. The major data sources and their abbreviations include: (1) long-term monitoring data collected monthly from the freshwater Delta to the Richmond San Rafael Bridge by the California Department of Water Resources (DWR); (2) 2 to 3 year bimonthly data collected by the Regional Effects Monitoring Program in 1986 to 1989 (REM); (3) long-term near-monthly data collected by the USGS south of Dumbarton Bridge in Palo Alto (USGS-PA); (4) the summary of semi-annual data collected by various agencies as listed in Thompson *et al.* 2000 (Regional Monitoring Program [RMP], Long-term monitoring program [LMP], and Bay Protection and Toxic Clean-Up Program [BPTCP]); (5) samples taken as part of NOAA's National Benthic Investigation in August 2000 and 2001 (Environmental Monitoring Program [EMAP]); (6) a monthly study of the bivalves south of San Mateo Bridge collected by the USGS in 1990 to 1996 (Thompson 2005, Thompson 1999); and (7) unpublished rapid assessment survey data from the California Academy of Sciences (C. Brown, Smithsonian Institute, pers. comm.). The focus here is on the most common species found within samples and does not include all species found at every sampling location during each sampling event. Because only a few studies include biomass data, the discussion will be limited to abundances unless otherwise noted.

Lower Estuarine River

The Lower Estuarine River region incorporates the summer extent of X2¹ during most non-drought years and includes the estuarine river upstream of Honker Bay to Colinsville on the Sacramento River. During normal hydrologic years the benthic community in this location would be considered oligohaline. Soft bottom, sessile invertebrates found in this region are divided into channel and channel edge locations. The channel assemblage is

found mostly in sand habitats and is characterized by organisms living in high velocity areas among and in sand waves. The community is dominated by two invasive filter-feeding bivalves, *Corbicula fluminea* and *Corbula amurensis*, with *C. amurensis* appearing during dry months (summer and fall) and dry years. Both bivalves filter-feed, but *C. fluminea* is also capable of pedal deposit-feeding. *C. amurensis* lives at or just below the surface and the presence of barnacles on their anterior shell is an indication that extending this part of the body into the flow is common. *C. fluminea* may be slightly less available to some surface predators as they live slightly below the surface, and their shell is considerably thicker. Small but interannually persistent populations of subsurface, deposit feeding oligochaetes (*Varichaetadrilus angustipenis* and *Limnodilus hoffmeisteri*) and tube-dwelling, surface deposit-feeding spionid polychaetes (*Marenzelleria viridis*) co-occur with the bivalves. Amphipods are seasonally and interannually ephemeral with the surface-tube dwelling *Americorophium spinicorne*, *Americorophium stimpsoni*, and free living *Gammarus daiberi* commonly seen in small numbers.

The channel edge community is also dominated by *C. fluminea* and *C. amurensis*. These two species are much more numerous in the channel edge than in the channel, with an order of magnitude increase in *C. fluminea* abundance and at least twice as many individuals of *C. amurensis* as seen in the channel. *C. fluminea* abundance tends to peak in spring, whereas *C. amurensis* can have either or both a spring and fall abundance peak. Unlike the channel, *C. amurensis* becomes the dominant bivalve in dry years on the channel edge. In the channel edge community, oligochaetes are 10 times more plentiful and include more species than in the channel; *V. angustipenis* and *L. hoffmeisteri* are still the dominant oligochaetes. Surface feeding polychaetes are at least twice, and up to 10 times, as abundant as in the channel. This niche is still dominated by *M. viridis* but also includes a sabellid polychaete, *Laonome* sp. *M. viridis* shows a seasonal cycle with peak abundances in spring that is not seen in the sabellid. Amphipods are an order of magnitude more abundant in the channel and consistently peak in spring and summer with the same *Americorophium* species dominating the channel edge community as in the

¹ X2 is defined as the distance upstream from the Golden Gate Bridge to the point where daily average salinity at 1 meter from the bottom is 2 ppt (Jassby *et al.* 1995).

channel community. The free-living *Gammarus daiberi* is inter-annually persistent with slightly smaller abundances on the channel edge versus in the channel.

Common mobile invertebrates found in oligohaline salinity zones in San Francisco Bay soft bottom habitat include the oriental shrimp (*Palaemon macrodactylus*) and the mud crab (*Rhithropanopeus harrisi*), both of which are non-native introduced species.

Suisun Bay Region

The Suisun Bay reach includes two shallow bays (Honker and Grizzly Bays), a deeper bay (Suisun Bay), and a deep water channel that connects these bays and San Pablo Bay (Figure 1). During normal hydrologic years, the benthic community at these locations would be considered a mesohaline community. Invertebrate communities in soft bottom sediments have been further classified by channel, channel edge, slough channel, and shallow subtidal topography. *C. amurensis* is the dominant bivalve in the channel with *C. fluminea* arriving during wet years, and once settled, remaining as adults for 2 to 3 years beyond the low salinity years. In addition to the bivalves, the spinoid polychaete *M. viridis*, the subsurface deposit feeding polychaete *Heteromastus filiformis*, and the surface dwelling cumacean *Nippoleucon hinumensis* occur in small numbers in the channel (EMAP).

The channel edges tend to have muddier sediments with higher organic carbon content (Hymanson *et al.* 1994) and the benthic community has a higher number of species and individuals than seen in the channel areas. The channel edge community is dominated by *C. amurensis*, which tends to peak in fall at this location. The oligochaetes are much less numerous than in the up-bay Estuarine River region and are joined by *Heteromastus filiformis* in the subsurface deposit feeding niche. The tube dwelling polychaetes (mostly *M. Viridis*) are also less abundant than in up-bay regions. The most abundant crustacean is the cumacean *N. hinumensis*, with the deposit feeding isopod *Synidotea laevidorsalis* and filter feeding barnacle *Balanus improvisus* being consistent members of the community. The surface dwelling crustaceans and polychaetes show peak abundances in

spring whereas the deposit feeding species and *C. amurensis* tend to peak in fall.

The benthic communities in the two shallow bays are similar, with variation reflecting their relative position to freshwater inflow. Honker Bay, the more upstream bay, has a high percentage of coarse plant material and gravel/sand (EMAP 2000). *C. amurensis* occurs throughout Honker Bay, but in higher numbers in the western than in the eastern bay (EMAP). A similar abundance gradient occurs for *M. viridis*. Grizzly Bay has been described by Hymanson *et al.* (1994) and Peterson (2002). The species richness and abundance is lower than at estuarine river and San Pablo Bay locations. The community is dominated by *C. amurensis*, which peaks in abundance in fall during the wet and normal water years, and peaks in summer and fall in dry and below normal years. Minimum abundance for this bivalve is seen in spring or early summer in most years. Abundance patterns for the tube-dwelling, surface-deposit feeding spinoid polychaete *M. viridis* (peak abundance late spring-early summer, followed by a several month minimum) covaries with *Monocorophium alienense*, an amphipod with similar habits and food but with which peaks in abundance in late fall/early winter.

Montezuma and Suisun Slough Channels occur in the Suisun Marsh and have similar sediment (clay and silty clay) and benthic communities as seen in the nearby shallow Grizzly Bay areas (EMAP). The abundances of all species, including *Corbula amurensis*, *Marenzelleria viridis*, and *Nippoleucon hinumensis* are much smaller in the Suisun Marsh sloughs than in Grizzly Bay. *Ampelisca abdita* is found in the marsh sloughs but occurs in highest numbers near the interface with the bay.

Similar to estuarine river regions, the exotic oriental shrimp (*Palaemon macrodactylus*) is common in mesohaline regions of San Francisco Bay. Also, *Crangon franciscorum* is the most common shrimp in the Bay during most years and is targeted by shrimp trawlers.

San Pablo and South Bays

The San Pablo Bay and South Bay regions are separated by the connection to the Pacific Ocean (Figure 1). Sessile invertebrate communities in both bays are considered mesohaline during normal hydrologic years, although the proximity of San Pablo Bay to freshwater flow from the Sacramento and San Joaquin Rivers can lead to seasonal and interannual patterns in the benthic community that differ from those in the South Bay.

The soft bottom invertebrate communities in these bays have been further classified by main channel, slough channel, and shallow subtidal topography. The main channel sediment is mostly muddy sand with low organic carbon content (RMP, REM, EMAP). The bivalves *C. amurensis*, *Mya arenaria*, *Venerupis japonica*, *Macoma petalum*, and *Musculista senhousia* are common inhabitants that as a group can dominate the biomass but not necessarily the abundance of individuals in this community. All of these bivalves are capable of filter-feeding although *M. petalum* can also surface deposit feed. *M. arenaria* and *M. petalum* live deep in the sediment with the rest residing in the near surface. The most accessible bivalve species to predators is the thin-shelled *M. senhousia* that lives on the surface in a mat of byssal threads. *C. amurensis* has a strong seasonal pattern with abundance peaking in summer or fall. The surface tube-dwelling amphipod, *Ampelisca abdita*, can reach extremely high numbers in the estuarine channels (>40,000/m²); maximum abundance commonly occurs in summer and fall. Other tube-dwelling amphipods, including several species of *Corophium* and spionid worms (mostly *M. viridis* and *Streblospio benedicti*) can also reach moderate abundances. Surface and subsurface deposit feeders are present with several species such as the cumacean *Nippoleucon hinumensis*, the polychaete *Cirriformia spirabranca*, oligochaete *Tubificoides* species, and the bivalve *Theora lubrica* being common.

San Pablo Bay and South Bay include many small slough channels, including those in the Napa and Petaluma Rivers that flow into San Pablo Bay. The sediment in these sloughs is mostly mud with some localized areas of sandy mud. Species richness and abundance is higher in

downstream Napa River benthic communities with the community being quite similar to that seen at the channel edge. The lower Napa River benthic community can include large populations of *Ampelisca abdita* (EMAP).

South Bay sloughs differ from those in the Petaluma and Napa Rivers due to larger populations of the amphipods *Monocorophium alienense* and *Grandidierella japonica*, the cumacean *Nippoleucon hinumensis*, the bivalve *Macoma petalum*, and two polychaetes (*Sabaco elongates* and *Neanthes succinea*). Large populations of the maldanid polychaete *Sabaco elongatus*, a head down deposit feeder with tubes that can be close to a meter in length can structure the rest of the community. Surprisingly, *A. abdita* and the bivalves mentioned above are capable of living among *S. elongatus* tubes which extend 1 to 2 cm above the substrate surface. Many of the species in this assemblage are patchy in space and time with some, like *M. senhousia* and *A. abdita*, having very high abundance one year and low abundance the next year (REM, USGS).

The species composition of the benthic community in the shallow, subtidal areas is similar to that seen in the main channel. The bivalves *C. amurensis*, *Mya arenaria*, *Venerupis japonica*, *Macoma petalum*, and *Musculista senhousia* are common, but show very strong seasonal patterns with declines in bivalve abundances to near zero each winter/early spring. The bivalves are therefore mostly annual species in this habitat with peaks in abundance occurring in late spring/early summer. The amphipods (*Corophium heteroceratum* and *Ampelisca abdita*) can show similar annual patterns except during dry years when *A. abdita* in particular seems to persist through the winter. As seen in the channel, the cumacean *Nippoleucon hinumensis* is common and peaks in spring of most years.

Five large, motile invertebrates are common in San Pablo Bay and South Bay. Dungeness crab reproduces in the ocean and rears in nearshore and estuarine areas. Juvenile Dungeness crabs immigrate to San Francisco Bay from May to early June, using bottom currents, and rear in the Bay for 8 to 15 months (Tasto 1983, Hieb 2006). Juveniles that rear in the Bay grow faster than and reach adult size one to two years before ocean-reared individuals (Hieb 2005). Dungeness crab abundance is typically

lowest when ocean temperatures are warm (i.e., El Niño years) and highest when ocean temperatures are cool. Abundance in 2005 was low, likely due to above-average sea surface temperatures in the Gulf of the Farallones in winter 2004 to 2005 (Hieb 2006). Dungeness crab support a valuable sport and commercial fishery along the coast. The blackspotted shrimp (*Crangon nigromaculata*) is the second most common shrimp in the Bay overall and the most common shrimp in some years. *Ilyanassa obsoleta* is the most abundant intertidal snail on San Francisco Bay mudflats and in the lower reaches of marsh channels, where it is often found in large groups. It has been collected in the Bay in salinities of 10 to 32 ppt and water temperatures of 13 to 22° C (Cohen 2005). The American spider crab, *Pyromaia tuberculata*, and the nudibranch *Sakuraeolis enosimensis* are also common.

Central Bay

The Central Bay region is at the nexus of the South Bay, San Pablo Bay and Pacific Ocean, and includes the northern extreme area of South Bay (Figure 1). Sessile invertebrates in Central Bay are representative of euhaline communities during most years, with extreme freshwater flood events affecting only the shallow sections. The soft bottom invertebrate taxa in Central Bay have been further classified into deep water channel, slough channel, harbor, and shallow subtidal topography. Within these topographic areas, there are typically moving sand and sand waves in Central Bay channels, muddy-sand and sandy-mud in the lee of islands, and sand and sandy-mud areas south of the Bay Bridge (EMAP).

The invertebrate taxa in the sand substrate is characterized by the free living, predatory polychaete *Heteropodarke heteromorpha*, several low abundance species of amphipods (*A. abdita*, *Monocorophium acherusicum*) and cumaceans, and a few bivalves including the filter feeding *Mya arenaria* and the surface deposit feeder *Tellina nuculoides* (EMAP, RMP). The muddy-sand benthic community in Central Bay and northern South Bay have a diverse polychaete community represented by several subsurface deposit feeding capitellid species, a tube dwelling filter feeding species (*Euchone limnicola*), a carnivorous species (*Exogone lourei*), and the maldanid

polychaete *Sabaco elongatus*. There are also several surface deposit feeding *Ameana* spp. persisting throughout the year (Thompson and Peterson 2004). The ascidean *Molgula manhattensis* is found throughout euhaline areas of northern South Bay, adhering to any solid object and developing “bloom” abundances during some years.

The slough channels in Central Bay that have been sampled are near harbors in China Basin and Corte Madera Creek. Communities in these sloughs are more similar to those in San Pablo and South Bay than the benthic community seen in the euhaline shallow subtidal zone in Central Bay. In most cases, the same species occur as in San Pablo Bay, with a few additional species such as the amphipod *Monocorophium acherusicum*, the bivalve *Mya arenaria*, the surface deposit-feeding polychaetes *Pseudopolydora diopatra* and *Tharyx* sp., and the tube dwelling filter-feeding polychaetes *Euchone limnicole* and *Polydora cornuta*. Major harbors and ship channels in the estuary are located in Central Bay and are a mix of the benthic community from surrounding areas (deep and shallow-water and slough marine communities). The only abundant obligate filter-feeders in the harbors are the tube dwelling amphipod *Ampelisca abdita* and tube dwelling polychaete *Euchone limnicola*. Probably indicative of the increased water column residence time and increased sedimentation in harbors, the majority of the species are deposit feeders, or in the case of the amphipods *Grandidierella japonica*, *Monocorophium acherusicum*, and *Monocorophium alienense*, are species that both filter and deposit feed. The two surface deposit feeding polychaetes *Streblospio benedicti* and *Pseudopolydora diopatra* are tube dwellers. There is a relatively high number of subsurface deposit feeding polychaetes and oligochaetes in these areas; *Tubificidae* spp., *Mediomastus* spp., *Heteromastus filiformis*, and *Sabaco elongatus* dominate the areas studied. There is also sufficient community complexity and abundance to support relatively high abundances of three carnivorous polychaete species: *Exogone lourei*, *Harmothoe imbricata*, and *Glycinde armigera*.

Mobile invertebrates of the euhaline salinity zone are characterized by crustaceans such as blackspotted shrimp (*Crangon nigromaculata*), California bay shrimp (*Crangon franciscorum*), and the slender rock crab (*Cancer gracilis*),

which may crawl on or burrow in bottom sediments. They provide an important food source for carnivorous fishes, marine mammals, and birds. In San Francisco Bay, the abundance of blackspotted bay shrimp peaks from May through August and again from December to February (Hieb 1999b). The California bay shrimp is the most common *Crangon* spp. in San Francisco Bay (CDFG 1992). It is found throughout the Bay, but the center of juvenile distribution is normally between San Pablo Bay and the western Delta (Hieb 1999b). Juveniles rear in shallow, low salinity areas and migrate to deeper, higher salinities areas as they grow (Hieb 1999b). There is a strong positive relationship between California bay shrimp annual abundance and freshwater outflow in spring (CDFG 1987, CDFG 1992). The slender rock crab is smaller than other common Cancer crabs, rarely exceeding 85 mm carapace width (Hieb 2005). Slender rock crabs are most common in Central Bay, and probably utilize San Francisco Bay as an extension of nearshore coastal habitat (Hieb 1999a). Slender rock crab abundance in San Francisco Bay decreased in both 2004 and 2005 following a decade of relatively high abundances (Hieb 2006).

Within the soft bottom substrate habitat, taxa on shell mix habitat differ from taxa on other soft bottom substrate habitat (Table 3b). Two clams, a mussel and an ascidian (*Corbula amurensis*, *Molgula manhattensis*, *Musculista senhousia*, and *Venerupis philippinarum*, respectively) are found on shell mix habitat, as well as on other soft bottom sediment types in San Francisco Bay. An anemone, a mussel, two barnacles, and an isopod (*Diadumene lineate*, *Mytilus tossulus/galloprovincialis*, *Balanus crenatus*, *Amphibalanus improvisus*, and *Gnorimosphaeroma oregonense*, respectively) are found on both shell mix habitat and hard bottom substrate. All other species are common only on shell mix substrate.

Fishes

Mesohaline/Oligohaline

In the mesohaline/oligohaline community of San Francisco Bay, common fishes over unconsolidated

sediments include sturgeon (*Acipenser* spp.), Sacramento splittail (*Pogonichthys macrolepidotus*), longfin smelt (*Spirinchus thaleichthys*), and the starry flounder (*Platichthys stellatus*) (Table 4). Sturgeon are anadromous, spending most of their lives in marine or estuarine waters and moving to the Sacramento River to spawn. Sturgeon are associated with turbid water, and use their specialized tube-like mouth to prey on benthic organisms. There are two species of sturgeon in San Francisco Bay: the white sturgeon (*Acipenser transmontanus*) and the federally threatened green sturgeon (*Acipenser medirostris*). In San Francisco Bay, the white sturgeon is more abundant than green sturgeon and spawn in the Sacramento River system during the winter months (Moyle 2002). White sturgeon are an important recreational species, and good management practices (size limits) have helped bring this species from near extinction due to commercial fishing in the late 19th and early 20th century back to stable population levels (Moyle 2002). Fishing pressure has had less of an impact on green sturgeon, as their poor taste makes them less desirable to consumers. Green sturgeon spawning grounds are limited due to habitat alteration, however, making it vulnerable to decline (Moyle 2002).

The Sacramento splittail was federally listed as threatened from 1999 to 2003. Young-of-the-year and yearlings are most abundant in the summer months in shallow water areas such as Suisun Marsh (Moyle 2002). Adults move upstream during the winter and spring to forage and spawn in flooded areas (Moyle 2002). The longfin smelt is an anadromous species that spawns in the Delta and rears in the brackish areas of the San Francisco Bay and Delta. It is common in Suisun Bay and upper San Pablo Bay during winter and spring, and disperses throughout San Francisco Bay in late spring, summer and fall (Messersmith 1966, Hieb and Baxter 1993). Longfin smelt abundance in San Francisco Bay is a function of freshwater outflow (Stevens and Miller 1983), although the relationship changed to substantially lower longfin smelt abundance with outflow after the introduction of the invasive clam *Corbula amurensis* in the late 1980s. This corresponded with a decline in phytoplankton and zooplankton abundance due to grazing by *C. amurensis* (Kimmerer 2002). Adult starry flounder primarily inhabit and spawn in marine waters, while juveniles seek shallow

fresh to brackish water in bays and estuaries to rear (Orcutt 1950, Haertel and Osterberg 1967, Bottom *et al.* 1984, Hieb and Baxter 1993). Juveniles enter San Francisco Bay via strong gravitational currents present during wet years (Moyle 2002) and appear to remain in the Bay through at least their second year (Baxter 1999). Juvenile abundance declined through the mid-1980s and early 1990s, increased with the high outflow years of 1995 to 1998, and has been cyclic since, with a recent increase from 2002 to 2005. Juveniles are caught from San Pablo Bay upstream to the Delta, with the younger fish further upstream. (Greiner *et al.* 2006).

Polyhaline

In the polyhaline community of San Francisco Bay, fish assemblages over unconsolidated sediments include Pacific staghorn sculpin (*Leptocottus armatus*) and English sole (*Parophrys vetulus*). English sole adults enter San Francisco Bay to spawn, and juveniles use intertidal and subtidal sand and mudflats for refuge and feeding (Pearson and Owen 2001). Age-0 English sole are found primarily in areas where temperatures are between 12.8 and 14.5 °C and salinities are between 12 and 24 ppt (Baxter 1999). As temperatures increase in spring and summer, fish move to deeper and cooler areas.

Euhaline

Within the euhaline community, the fish community is primarily dominated by elasmobranchs, such as the brown smoothhound (*Mustelus henlei*) and leopard shark (*Triakis semifasciata*), brown rockfish (*Sebastes auriculatus*), plainfin midshipman (*Porichthys notatus*), and flatfishes such as the California halibut (*Paralichthys californicus*) and the speckled sanddab (*Citharichthys stigmaeus*). Leopard shark individuals may remain in San Francisco Bay year-round or utilize the Bay seasonally, emigrating to the ocean in fall and winter (Smith and Abramson 1990). Adults migrate to shallow water in San Francisco Bay in late winter and spring to pup and forage. Although found from South Bay to San Pablo Bay, leopard shark abundance is highest in South Bay. Reproduction occurs between April and May, with each female producing four to 29 pups. Catch in San Francisco

Bay has exhibited a downward trend since 1984 (Greiner *et al.* 2006). Leopard shark is fished commercially and recreationally in the Bay, although the recreational fishery accounts for the majority of the catch (Smith and Abramson 1990). Juvenile brown rockfish emigrate to the Bay from nearshore coastal areas in spring to rear (Baxter 1999). Juveniles rear in the Bay for 3 to 4 years and have high site fidelity (Kendall and Lenarz 1986). Brown rockfish is found from South Bay to San Pablo Bay, but abundance is highest in Central Bay (Baxter 1999). It is found in rocky areas and near piers and other structured habitats. California halibut became common in San Francisco Bay during the 1980s and 1990s when abundances increased, apparently a result of a succession of warm water and El Niño years (Baxter 1999). Adult California halibut enter San Francisco Bay to forage and spawn, and juveniles use intertidal sand and mud flats for refuge and feeding (Pearson and Owen 2001). California halibut spawning and survivorship and growth of juveniles increase when temperatures are between 15 and 16.5 oC (Gadomski and Caddell 1991, Caddell *et al.* 1990). California halibut is found from South Bay upstream to the Carquinez Straight, but highest juvenile catches are in South Bay (Greiner *et al.* 2005). The California halibut supports an important offshore commercial fishery and is targeted in the Bay by sport anglers. The plainfin midshipman is a demersal, marine fish that buries into soft sediments during the day and moves into the water column to feed at night (Fitch and Lavenberg 1971). Plainfin midshipman adults migrate from coastal areas to bays and estuaries in late spring and summer to spawn (Wang 1986). Juveniles rear in estuaries through at least December (Baxter 1999). In San Francisco Bay, plainfin midshipmen are found primarily in Central Bay, with record high catches from 2001 to 2004 (Greiner *et al.* 2006).

Birds

The diving benthivore feeding guild uses unconsolidated sediment habitat for foraging on benthic infauna (Table 5). Among this guild, canvasback (*Aythya valisineria*) within San Francisco Bay are declining (Trost 2002). Members of the genus *Aythya* including greater and lesser

scaup and canvasback are diving ducks which use San Francisco Bay for overwintering, and are associated with together because they are difficult to tell apart during aerial surveys. Scaup feed on benthic infauna including mussels and clams. Canvasbacks forage in depths up to 9 m on submerged aquatic vegetation and small invertebrates (Fix and Bezener 2000). Canvasbacks are most abundant from October to January in Suisun Bay, San Pablo Bay, and South Bay (Goals Project 2000). The San Pablo National Wildlife Refuge was established, in part, for the protection of canvasbacks (Goals Project 2000). The surf scoter is associated with euhaline and polyhaline salinities, where they dive for bivalves, molluscs, crustaceans, and other marine and estuarine invertebrates. Scoters are most abundant in Central Bay and South Bay (Goals Project 2000).

Marine Mammals

Soft bottom habitats are used for feeding by several marine mammal species (Table 6). Harbor seals (*Phoca vitulina*) will feed throughout the year on fish and invertebrates such as California halibut, speckled sanddabs, and mysid shrimp that occur in and on top of soft bottom substrates. Harbor porpoise (*Phocoena phocoena*) also often feed on species associated with soft bottom substrates such as rockfish, flatfish, and speckled sanddabs. California and lions (*Zalophus californianus*) also occasionally feed on species associated with soft bottom habitat. The Eastern Pacific Gray whale (*Eschrichtius robustus*) uses soft bottom substrates in San Francisco Bay during stopovers on bi-annual migrations between calving grounds in Baja, Mexico and feeding grounds in Alaska and Canada. Gray whales feed on small crustaceans, polychaete worms, mollusks, amphipods, and other sessile invertebrates by actively sucking up sediments, entangling food in baleen, and using the tongue to push out sediment, leaving food items behind.

Harbor seals also use intertidal soft bottom habitats to rest, molt, and breed. Most of the haul-out sites within San Francisco Bay are on soft bottom habitat in the South Bay including the primary breeding site within the Bay, Mowry Slough. Nearly 100 pups have been documented

shallow, polyhaline to oligohaline salinities. Greater (*A. marila*) and lesser scaup (*A. affinis*) are often grouped at this site compared to 50 at Castro Rocks, the next largest colony site (Risebrough *et al.* 1981, Green *et al.* 2006). Haul-out sites usually have a small slope and immediate access to deep water, and are remote from human activities and close to foraging areas.

HARD BOTTOM SUBSTRATE

(by Natalie Cosentino-Manning, Janet Thompson, Katie McGourty, Susan Wainwright-De La Cruz and Sarah Allen)

General Description

Hard bottom substrate in San Francisco Bay consists of natural and artificial surfaces. Natural substrates include boulders, rock face outcrops, and low relief rock. These habitats are found primarily in Central San Francisco Bay, where soft sediment has been scoured by tidal flow (Figure 10). Boulders occur around islands including Yerba Buena, Angel, and Alcatraz Islands. Boulders are also found around peninsulas and in channels such as Marin and Tiburon Peninsulas, Racoon Strait, Belvedere Dumbarton Narrows, San Pablo Point, and the north and south sides of the Golden Gate channel. Submerged rocks are found in west Central Bay including Arch Rock, Harding Rock, Shag Rock, and Blossom Rock. The tops of these rocks were flattened by the U.S. Army Corps of Engineers to minimize navigation hazards; they are approximately 50 meters high and 11 to 12 meters deep. Rock face outcrops can be found around Yellow Bluff and in northern Golden Gate channel. Artificial hard substrate includes vessel structures, pilings, rip rap, and pipelines. Pilings, rip rap, and pipelines can be found in every San Francisco Bay region.

Hard substrate provides habitat for an assemblage of marine algae, invertebrates and fishes. All hard substrate in San Francisco Bay provides substrate for invertebrate attachment and refugia and foraging for fishes and invertebrates. Natural hard substrate provides substrate

for algae and diatoms and foraging areas for birds and marine mammals. Boulders and rock face outcrops provide substrate for fish rearing, spawning, and growth.

Unfortunately, natural hard substrate within the bay has been poorly studied; whereas artificial hard structures have been examined to describe invasion of exotic species over time. The existing knowledge of benthic invertebrate communities in natural hard substrate of San Francisco Bay consists of only one 4-day ROV survey of Harding, Shag, Arch, and Blossom rocks conducted September 11 to 15, 2001 (Garcia & Associates 2001). A complete investigation of the community was not attempted. The most common species noted were “turf” organisms (hydroids, bryozoans, and anenomes) and sea stars.

Since 1993, the San Francisco Estuary Institute has been performing Rapid Assessment Surveys (RAS) during summer months to describe exotic species distribution throughout San Francisco Bay on artificial hard substrates including pilings, docks, and boat ramps. Currently, there are 896 records of 294 distinct taxa (Cohen *et al.* 2005). The majority of these records (62%) are from floating docks, followed by intertidal benthos (20%) and benthic grabs (13%). The Smithsonian Institute also completes periodic RAS; information from these surveys are cited here as C. Brown (Smithsonian Institute, pers. comm.). Table 3c represents the most commonly observed species on artificial hard substrate. The majority of species in this habitat must be filter- or suspension-feeders with a much smaller number of surface deposit feeders that live within these communities benefiting by the left-over food, pseudo-feces and feces produced by the filter and suspension feeders (J. Thompson, USGS, pers. comm.). Some scrapers also survive in this habitat by eating algae that has adhered to the substrate or by eating encrusting bryozoan colonies (J. Thompson, USGS, pers. comm.).

Taxonomic Groups

Plants

Natural hard bottom substrates within the intertidal and subtidal are more common along the shores closest to the

opening of the Bay and are almost absent in Suisun, San Pablo and South San Francisco Bays. The lack of hard substratum for attachment is one of the major limiting factors for the occurrence of macroalgae in San Francisco Bay. Silva (1979) found that hard substrata such as outcroppings, rocks, pebbles, seawalls and shells supported the greatest diversity of macroalgae compared to soft substrates. In a long-term study conducted by San Francisco State and University of California Berkeley, nine sites around San Francisco Bay were surveyed for macroalgae from 1978 to 1983 (Josselyn and West 1985). Of the 162 species found, 129 species resembled assemblages found on the rocky, euhaline shores outside of the Bay and only 33 were determined to be polyhaline. Five of the species found were non-native. Two of these, *Codium fragile* subsp. *Tomentosoides* and *Sargassum muticum* were introduced in the 1970s and are known to be nuisance species in other estuaries (Silva 1979).

The majority of the species found during the surveys were located in the middle to upper intertidal zones and 21 species were found to occur in the subtidal zone. The Twin Sisters, a rocky outcropping in San Pablo Bay, was found to have the highest diversity of species due to the strong reversing currents which scour the lower intertidal and subtidal during each tidal cycle. The kelps, *Laminaria* and *Egregia* are found here. Other species such as *Ahnfeltiopsis* (formerly *Gymnogongrus*) and *Halymenia* that are common on the outer exposed rocky coast, were also known to be present.

The only angiosperm, of the genus *Phyllospadix*, is found at the entrance to San Francisco Bay and is not found any further east than Fort Baker and Fort Point, which flank the Golden Gate Bridge. *Phyllospadix* requires a euhaline environment with heavy wave exposure.

Invertebrates

Mesohaline

The large filter-feeding mussel *Mytilus trossulus*/*galloprovincialis* and filter-feeding barnacle *Amphibalanus improvisus* are the most visible members of the attached

invertebrate taxa in mesohaline regions (RAS). Other attached, non-mobile species include an anemone, *Diadumene* sp., a hydrozoan, *Garveia franciscana*, the sponges, *Halichondria bowerbanki*, *Haliclona* sp., and *Clanthea prolifera*, and a bryozoan, *Conopeium cf. tenuissimum*; all species that suspension feed (RAS, C. Brown, Smithsonian Institute, pers. comm.).

In mesohaline/oligohaline regions, several species of isopods and amphipods can be found crawling among sessile invertebrate species. These species include surface deposit-feeders (i.e., *Gnorimosphaeroma* sp., *Synidotea laevidorsalis*), grazers on attached algae (i.e., *Ampithoe valida*, *Sphaeroma quoianum*, *Eogammarus confervicolus*), and carnivores on the bryozoan colonies (i.e., the opisthobranch, *Hopkinsia plana*) (RAS).

Polyhaline

Invertebrate taxa in polyhaline regions are much more diverse than that seen in mesohaline regions. The attached filter-feeding species include three bivalves, *M. trossulus/gallopogonialis*, *Musculista senhousia*, and *Ostrea conchalpila*, and seven ascidians (RAS, C. Brown, Smithsonian Institute, pers. comm.). The barnacles *Balanus glandula* and *A. amphitrite*, in addition to *A. improvisus*, are seen throughout this zone. Attached suspension feeders include at least six sponges (e.g., *Halichondria bowerbanki*, *Clathra prolifera*, *Mycale macginitiei*, *Leucilla nuttingi*, *Sycon* sp., *Haliclona* sp.), the hydrozoans *Ectopleura crocea* and *G. franciscana*, seven bryozoans, and at least two species of the anemone genus *Diadumene* (RAS, C. Brown, Smithsonian Institute, pers. comm.). Hard bottom substrate in polyhaline sloughs is poorly represented in available studies, but the small amount completed shows the barnacle *A. improvisus* and the bryozoan *Victorella pavidia* as present in the community (RAS). The commercially important Dungeness crab, discussed above, is also found in polyhaline hard bottom substrate.

In polyhaline regions, surface deposit feeding isopods are similar to those seen in mesohaline/oligohaline regions, with the addition of *Paranthura japonica* (RAS). Several additional amphipods are also present including

Incisocalliope derzhavini, *Jassa marmorata* and *Stenothoe valida* (RAS). Dungeness crabs (*Cancer magister*) occur in hard bottom polyhaline regions, as discussed above in the soft bottom substrate section.

Euhaline

The species richness of sessile invertebrates on hard bottom euhaline regions increases relative to polyhaline regions with the addition of filter-feeding bivalves *Hiatella artica* and *Modiolus rectus*, several bryozoans, the barnacle *Balanus crenatus*, the anemone *Metridium senile*, and the echinoderm *Pisaster brevispinus* (RAS, C. Brown, Smithsonian Institute, pers. comm.).

In euhaline regions, amphipods are still a major component of the community, but the isopods are less prevalent (RAS). Three species of multi-functioning caprellids (i.e., they are detritivores, carnivores, and deposit feeders) are present (RAS, C. Brown, Smithsonian Institute, pers. comm.). Omnivorous and carnivorous polychaetes, including two species of scale worms join the caprellids moving among the bases of the many attached epifauna and epiflora (RAS). Pacific rock crab (*Cancer antennarius*) and the red rock crab (*C. productus*) inhabit rocky, intertidal and subtidal areas in the Pacific Ocean, and likely use San Francisco Bay as an extension of their coastal habitats (Hieb 1999a). Adult (age 1+) Pacific rock crabs are most commonly found in Central Bay in both the fall and spring months. Juveniles are most common in Central Bay from January to May and in South Bay from July to December (Hieb 1999a). Pacific rock crabs move seasonally from channels (January to April) to shoals (June to December) (Hieb 1999a). The Pacific and red rock crabs are targeted by sport anglers from piers and jetties (Hieb 2006). Rock crab abundance increased from the mid-1990s to 2004, but in 2005, decreased to levels typical of the 1980s (Hieb 2006).

Fishes

Oligohaline/Mesohaline

Within oligohaline/mesohaline hard bottom taxa, the prickly sculpin (*Cottus asper*) is found in every freshwater tributary of San Francisco Bay. As a demersal species, prickly sculpins camouflage themselves in areas of hard substrate, and spawn in areas with flowing water and loose rocks, where males will locate nests (Moyle 2002). In San Francisco Bay, prickly sculpins are most abundant during periods of high freshwater outflow (Baxter 1999). Larvae are commonly found January through May, with a peak occurring in March; adults are most common from May through July (Baxter 1999).

Polyhaline

Fish species occurring over hard substrate in polyhaline salinity zones have not been documented.

Euhaline

In the euhaline community, brown rockfishes (*Sebastes auriculatus*) and surfperches (family *Embiotocidae*) are the most common fishes associated with natural hard substrates. All surfperches are livebearers, with the young born fully developed after a three to six month gestation period (DeLeon 1999). Most species are transient, immigrating to bays and estuaries to give birth in spring and summer. Both species of surfperch are distributed from South to San Pablo Bays. Shiner perch move from South and San Pablo Bays to Central Bay as they mature (DeLeon 1999). There are both recreational and commercial fisheries for surfperch in San Francisco Bay for use as food and live bait. Recreational anglers, mostly fishing from piers, docks, and along the shore, catch surfperch primarily in Central Bay. All surfperch common to San Francisco Bay underwent a precipitous abundance decline in the mid-1980s (DeLeon 1999). In response to the decline, CDFG changed their sportfishing regulations to be more protective, and abundance of four surfperch species (walleye surfperch, white seaperch, shiner perch, and dwarf perch) increased between 2000 and 2004 in the Bay (Greiner *et al.* 2006). However, shiner perch

abundance decreased again to 10% of the 2004 abundance in 2005 (Greiner *et al.* 2006).

During winter months, Pacific herring (*Clupea pallasii*) enter euhaline areas of San Francisco Bay to spawn during periods of low salinity. Schools of adult herring enter the Bay during fall and winter, depositing adhesive eggs onto submerged aquatic vegetation and hard bottom substrate (O'Farrell and Larson 2005). CDFG conducts herring spawning surveys November through March throughout Central San Francisco Bay, the area of highest spawning concentration (Watters *et al.* 2004).

Birds

Piscivorous birds such as cormorants (*Phalacrocorax* spp., including the double-crested cormorant, Brandt's cormorant, and the pelagic cormorant), the pigeon guillemot (*Cepphus columba*), the herring gull (*Larus argentatus*), and the mew gull (*L. canus*) forage over hard bottom substrate in San Francisco Bay. For more in depth discussion on cormorants, please refer to the Water Column Habitat discussion below. The pigeon guillemot, a member of the family *Alcidae*, which includes the common murre, murrelets, auklets, and puffins, is commonly seen in summer months. Pigeon guillemots nest and forage in nearshore rocky habitats from March to September including sea cliffs, offshore rocks, and periodically docks and bridges. Pigeon guillemots forage by diving for fishes, mollusks, crustaceans, and worms (Fix and Bezener 2000). Both the mew and herring gull are observed during winter months and occupy similar niches, feeding opportunistically on fishes, crustaceans, and other small prey items. The mew gull remains in near-shore areas, however, avoiding anthropogenic food source areas such as garbage dumps and in-shore agricultural areas where herring gull are commonly observed (Fix and Bezener 2000).

Marine Mammals

Hard bottom substrates attract fish assemblages, providing prey for harbor seals and sea lions, including

sculpins, rockfish, perch and herring. Hard substrates are an important feeding habitat for harbor seals, including from the Golden Gate east to Treasure Island, northwest to Tiburon Peninsula, and southward to Yerba Buena Island (Goals Project 2000, Green *et al.* 2006). Also, many harbor seal haul-out sites are located in close proximity to subtidal hard substrate including Point Bonita, Castro Rocks, Yerba Buena Island, and Angel Island (Goals Project 2000). Sea lions also feed in deep, marine waters of San Francisco Bay; however, they feed mostly on seasonally abundant herring associated with hard structures and eel grass beds in the Central Bay, and on spawning salmonids that are migrating up the Bay across multiple habitats to freshwater tributaries. Pacific herring is a major prey item for many marine mammal species that are drawn to the Bay, including harbor porpoise and Steller sea lions.

SHELLFISH BEDS

(by Andrew Cohen, Natalie Cosentino-Manning and Korie Schaeffer)

General Description

Past studies in San Francisco Bay have used the term "shellfish bed" to refer to locations where commonly harvested mollusks or crustaceans were found in sufficient concentrations to warrant their harvest. Earlier studies (e.g., Bonnot 1932 and 1935, Skinner 1962, Barrett 1963) primarily referred to locations of commercial interest, including planted or cultured beds of exotic bivalves. The species involved included the native Olympia oyster (*Ostrea conchaphila*), the exotic Virginia oyster (*Crassostrea virginica*) and Pacific oyster (*C. gigas*), the exotic soft-shell clam (*Mya arenaria*) and Japanese littleneck or Manila clam (*Venerupis philippinarum*), and the native bay shrimp (*Crangon* spp.), all of which were commercially harvested for food. More recent studies (e.g., Wooster 1968, USEPA 1972, Dahlstrom 1977, Jones & Stokes 1977, Sutton 1978, 1981) have focused on concentrations that are or could be sport-harvested, primarily beds of the clams, *V. philippinarum* and *M.*

arenaria. Sutton (1978) also delineated beds of the rough piddock (*Zirfaea pilsbryi*), a native clam that bores into soft rock or clays, as a potential species for sport harvest, and suggested that there may be significant subtidal beds of the native bentnose clam (*Macoma nasuta*), which was commonly harvested by Native Americans. Some studies refer in passing to beds of the exotic freshwater Asiatic clam (*Corbicula fluminea*; harvested for food or bait), native California mussel (*Mytilus californianus*), bay mussel (consisting of the native *M. trossulus*, the exotic *M. galloprovincialis*, and their hybrids), exotic ribbed horsemussel (*Geukensia demissa*), native ghost shrimp (*Neotrypaea californica*), and the blue mud shrimp (*Upogebia pugettensis*). Jones & Stokes (1977) and Sutton (1978) implied that the ribbed horsemussel might be a sport-harvested species in the San Francisco Bay. The ghost shrimp and blue mud shrimp are used for fish bait, though it's not clear that there are or ever were beds of blue mud shrimp in San Francisco Bay.

In this report, "shellfish bed" is defined structurally rather than in terms of human usage. As a working definition, we propose that in order to constitute a bed, living specimens of the nominal bivalve must cover at least 50 percent of the surface over at least several square meters and, in concentration, must provide a distinct, three-dimensional substrate. We discuss five types of shellfish beds that occur or may occur in San Francisco Bay: beds of California mussels and bay mussels, which occur on hard surfaces to which they attach by byssal threads; beds of the ribbed horsemussel, which typically occur either partially buried in the sediment in salt marshes, or on hard surfaces similar to California and bay mussels; beds of the green bagmussel, which in dense concentrations occur in a mat of interwoven byssal threads on the sediment surface, and beds of the Olympia oyster, which cement to hard substrates including other oysters, and which in the past built up in extensive congregations on bottom sediments in the Bay.

Little to no information is available on biotic assemblages associated with specific types of shellfish beds. Where specific information is available, it is discussed below. Some information on species use of shellfish beds in general exists based on observations within San Francisco

Bay and elsewhere. For example, three marine mammal species, the harbor seal (*Phoca vitulina*), the California sea lion (*Zalophus californianus*) and the California sea otter (*Enhydra lutris*) are thought to be commonly associated with shellfish beds based on location of prey items (S. Allen, National Park Service, pers. comm.). Harbor seals have been observed foraging at two oyster restoration sites, Bair Island and the Marin Rod and Gun Club. Sea otters, if feeding within the Bay, would be attracted to shellfish beds for food, as they do within other estuaries such as Elkhorn Slough in Monterey Bay.

California Mussel (*Mytilus californianus*) Beds

Mytilus californianus is a native, primarily outer coastal species that ranges from Baja California to the Aleutian Islands. It forms large beds on rocks and pilings exposed to the surf. It feeds mainly on suspended organic detritus and plankton, especially dinoflagellates (one species of which, *Gonyaulax catenella*, produces a neurotoxin that accumulates in *M. californianus* and may render the species unsafe to eat during the summer months). *M. californianus* attains a maximum length of 25 cm, but usually grows no bigger than 13 cm in intertidal populations in California (Haderlie and Abbott 1980).

In San Francisco Bay, *M. californianus* is found only in the western part of the Central Bay, with records inside the Bay as far as the east side of the Tiburon Peninsula, Angel Island, Alcatraz Island and northwestern Yerba Buena Island (A.N. Cohen unpublished data). Here it is primarily found on rocks and seawalls, and occasionally on floating docks or buoys. No studies have been made of the distribution or species composition of its beds in the Bay. There are, however, many studies of *M. californianus* beds on the outer coast, where it typically forms beds down to around MLLW and up to the surge zone in exposed sites. It is usually associated with the gooseneck barnacle (*Pollicipes polymerus*), either intermixed or in alternating beds. Other organisms typically found on or among them on the outer coast include sea cucumbers, *Cucumaria* spp., the pile worm, *Nereis vexillosa*, the isopods, *Cirolana harfordi*, *Idotea wosnesenskii* and *Idotea stenops*, the barnacles, *Semibalanus cariosus*, *Balanus*

glandula and *Tetraclita rubescens*, and feeding on the mussels, the sea star, *Pisaster ochraceus* (Kozloff 1973). Some of these may be associated with *M. californianus* near the mouth of San Francisco Bay.

Bay Mussel (*Mytilus trossulus/galloprovincialis*) Beds

In California, the bay mussel, identified in older literature as *Mytilus edulis*, is in fact a mix of two species, the native *Mytilus trossulus* and the Mediterranean mussel (*Mytilus galloprovincialis*), which are difficult to differentiate without molecular analysis. *M. trossulus* is the dominant mussel from northern California to Alaska, and *M. galloprovincialis* is dominant in southern California, where it was introduced sometime prior to 1947 (Cohen and Carlton 1995). San Francisco Bay and nearby bays in central California are in the boundary zone between the two populations, where both of the species and their hybrids occur in substantial numbers (McDonald and Koehn 1988, Sarver and Foltz 1993, Geller *et al.* 1993, 1994). It is not known whether the location of this boundary in central California is determined by the two species' environmental requirements, and thus is relatively stable, or is an invasion front, and thus likely to shift further north as *M. galloprovincialis* continues to expand its range². In the latter case, we would expect the bay mussels in San Francisco Bay to become increasingly dominated by *M. galloprovincialis*.

The larvae of bay mussels can settle on the outer coast, and adults are occasionally found there. In wave-exposed situations, however, they are out-competed by *M. californianus*, which attach more strongly and grow larger (Haderlie and Abbott 1980). In San Francisco Bay, bay mussels are found mainly from the northern South Bay to southern San Pablo Bay, with a few records ranging as far as the Dumbarton Bridge, Port Sonoma, and Martinez (Hopkins 1986, Cohen and Chapman 2005, A.N. Cohen unpublished data). Roughly the same distribution was reported by Packard (1918) in dredge samples from the 1911 to 1912 Albatross survey, suggesting that the arrival

² Global warming could shift an environmental boundary northward also, though probably not quickly as an invasion front.

of *M. galloprovincialis* has not markedly affected the range in the San Francisco Bay.

In the Bay, bay mussels are commonly observed to form beds on seawalls, dock sides and pilings, and possibly also on bedrock or riprap, though there are no studies of the distribution of these beds. Similarly, although there are published studies of the bay mussel bed community in other Pacific Coast bays, there are none in San Francisco Bay, and studies from these other bays may not apply to the distinct, invasion dominated biota of San Francisco Bay. In general, however, we expect a wide variety of organisms in bay mussel beds in the Bay, on and among the shells and among their byssal threads, (e.g., sponges, hydroids, snails (limpets and *Urosalpinx cinerea*), barnacles, tanaids, sphaeromatid and cirrolanid isopods, gammarid and caprellid amphipods, bryozoans and tunicates) with the communities varying with salinity, tidal exposure, and other variables. One mussel bed observed over the past decade, on pilings and rocks near the Fruitvale Bridge in Oakland, has consistently yielded numerous large specimens of the yellow sponge, *Halichondria bowerbanki*, large colonies of the hydroid, *Obelia cf. longissima*, large numbers of the barnacles, *Balanus glandula* and *Amphibalanus improvisus* along with a few *Amphibalanus amphitrite*, the tanaid, *Sinelobus* sp., the isopod, *Cirolana cf. harfordi*, the amphipod, *Ampithoe valida* and the tunicates, *Molgula manhattensis* and *Styela clava* (Cohen *et al.* 2005, A.N. Cohen unpublished data).

Ribbed Horsemussel (*Geukensia demissa*) Beds

The ribbed horsemussel (*Geukensia demissa*) is a non-indigenous, invasive species that was first collected in South San Francisco Bay in 1894 (Stearns 1899). It is now one of the most abundant bivalves in San Francisco Bay (Cohen and Carlton 1995). It commonly forms beds in salt marshes and along the edge of steep salt marsh channels from the South Bay to San Pablo Bay, where it frequently lies embedded with its posterior margin protruding above the mud (Cohen and Carlton 1995, A. Cohen, San Francisco Estuary Institute, pers. comm.). The ribbed horsemussel is also found on rocks and along seawalls in Lake Merritt, a shallow, brackish lagoon on

San Francisco Bay. It may also occur in beds on soft bottom and natural and artificial hard bottom subtidal areas from the South Bay to San Pablo Bay, but no surveys have been completed to identify or document such beds. There is no documented information on taxonomic groups associated with ribbed horsemussel beds, but native barnacles, *Balanus glandula*, have been observed attached to individual mussels within the beds (A. Cohen, San Francisco Estuary Institute, pers. comm.).

Green Bagmussel (*Musculista senhousia*) Beds

The green bagmussel (*Musculista senhousia*) is native to Japan and China and was introduced to central California with Japanese oysters (*Crassostrea gigas*) for harvest (Kincaid 1949). It was first collected in San Francisco Bay in 1946 (Carlton 1979). Green bagmussels tend to occur in very high densities along the eastern shore of Central San Francisco Bay (A. Cohen, San Francisco Estuary Institute, pers. comm.). It occurs in the bottom of Lake Merritt in dense byssal mats that can be pulled from the bottom in sheets and in the Oakland estuary mixed with bay mussels and *Diadumene* sp. anemones (A. Cohen, San Francisco Estuary Institute, pers. comm.). It is frequently the most common benthic organism from South Bay to San Pablo Bay, where it has been collected at densities of up to 1,000 to 2,000 clams/m² and is occasionally collected in Suisun Bay but not necessarily at densities high enough to form beds (Nichols and Thompson 1985, Hopkins 1986, Markmann 1986).

Olympia Oyster (*Ostrea conchaphila*) Beds

Shellfish bed habitat includes areas of living Olympia oysters (*Ostrea conchaphila*, formerly *O. lurida*), and remnant beds composed of dead shell material. Olympia oysters can survive in a broad range of habitats but are most abundant in estuaries, small rivers, and streams. Along the Pacific coast, Olympia oyster beds are formed in the subtidal zone and are bordered by mud flats at high tidal elevations and by eelgrass beds at low tidal elevations. They are found at depths of 0 to 71 meters (Hertlein 1959). Oysters may attach to the underside of

rocks higher in the intertidal zone, where the bottom is gravel or rock (Kozloff 1973).

Native oyster bed habitat is undoubtedly the most poorly understood of any San Francisco Bay subtidal habitat types. While anecdotal information indicates that historically Olympia oysters were a large component of the Bay ecosystem, to date no live subtidal Olympia oyster beds have been documented in San Francisco Bay. Intertidal populations, however, have been found throughout the Bay (Figure 11) and are currently receiving much attention by researchers and restoration practitioners. Elsewhere along the Pacific coast, Olympia oyster shell serves as a substrate for epifauna such as mussels, *Mytilus galloprovincialis* and *M. trossulus*, barnacles, and boring sponges (Baker 1995). Information on invertebrate taxa associated with oyster beds in the Bay is limited to on-going monitoring at three restoration sites (Richardson Bay, Bair Island, and Marin Rod and Gun Club), which use Pacific oyster shell as substrate for Olympia oyster recruitment (MACTEC 2006; Table 7).

The Richardson Bay oyster restoration site included a fish-monitoring component. Long-line and minnow trap techniques were used on a monthly basis for a period of 8 months. The most abundant species included the bat ray (*Myliobatis californica*), leopard shark (*Triakis*

semifasciata), shiner surf perch (*Cymatogaster aggregata*), diamond turbot (*Hypsopsetta guttulata*), thornback (*Ptyrrhinoidis trisderiata*), and bay pipefish (*Syngnathus leptorhynchus*) (McGowan 2005).

Diving benthivore birds such as the eared grebe (*Podiceps nigricollis*), the ruddy duck (*Oxyura jamaicensis*), and the common goldeneye (*Bucephala clangula*) have been observed foraging in shellfish beds (Table 5). Eared grebes may be observed throughout the year in San Francisco Bay (Goals Project 2000). Eared grebes make shallow dives for prey items such as small crustaceans, fishes, and mollusks (Fix and Bezener 2000). An estimated 85 percent of the Northern American ruddy duck population primarily uses the San Francisco Bay as over-wintering habitat (Goals Project 2000). The greatest ruddy duck abundance has been observed in the South Bay (Goals Project 2000), which is a historic area for native oysters (Figure 11). Ruddy ducks feed on submerged aquatic vegetation and small crustaceans (Fix and Bezener 2000). Similar to other diving benthivores, the common goldeneye is most abundant during the winter months, with San Francisco Bay representing a major over-wintering area. Goldeneyes feed on a variety of prey including crustaceans, mollusks, small fishes, and plant material (Fix and Bezener 2000).

HABITAT TYPE AND ASSOCIATED BIOLOGICAL ASSEMBLAGES – Shellfish Beds

Table 7. Sessile invertebrates associated with native oyster restoration sites at Marin Rod and Gun Club and Redwood City sites in San Francisco Bay from MACTEC (2006).

	Species	Native Oyster Restoration		
		RC Pallets	MRGC Necklace	MRGC Pallets
Sponges	<i>Halichondria bowerbanki</i>	X	X	X
	<i>Haliclona</i> sp.	X		
	<i>Clathria prolifera</i>	X		
Hydroids	<i>Tubularia</i> sp.	X	X	X
	<i>Obelia</i> sp.	X	X	X
Anenomes	<i>Diadumene</i> sp.	X	X	X
	<i>Haliplanelia lineata</i>	X	X	
Flatworms	<i>Platyhelminthes</i> sp.	X	X	
Scale Worms	<i>Halosydna brevisetosa</i>		X	
	<i>Harmothoe "imbricata"</i>	X		
	<i>Lepidontus squamatus</i>			
Family Syllidae	<i>Typosyllis nipponica</i>	X		X
Family Neridae	<i>Neanthes succinea</i>	X	X	X
	<i>Nereis vexillosa</i>		X	
	<i>Nereis latescens</i>		X	
Family Eunicidae	<i>Marphysa</i> sp.	X		
Family Terebellidae		X		
Barnacle	<i>Balanus</i> sp.	X	X	
Isopods	<i>Synidotea laevidorsalis</i>	X	X	X
	<i>Paranthura elegans</i>	X		
	Sphaeromatid type	X	X	X
Amphipods	Unknown gammaridian sp.	X	X	X
	<i>Ampelisca</i> sp.	?	X	
	<i>Corophium</i> sp.	X	X	X
Caprellid shrimp	<i>Metacaprella kenerlyi</i>		X	
	<i>Caprella</i> sp.			
Caridea shrimp	<i>Palamon macrodactylus</i>	X		X
	<i>Hemigrapsus nudus</i>	X		
	<i>Hemigrapsus oregonensis</i>	X	X	X
	<i>Rhithropanopeus harrisi</i>	X		
Pycnogonida		X		
Gastropods	<i>Urosalpinx cinera</i>	X		
	<i>Crepidula plana</i>	X		
	<i>Crepidula convexa</i>	X		
	<i>Philine</i> sp.	X		
Opisthobranchs	<i>Sakuraeolis enosimensis</i>	X	X	
	<i>Dirona picta</i>		X	
	<i>Elysia hedgpethi</i>		X	
	<i>Haminoea</i> sp.	X		
RC Pallets	Oyster shell pallets at Redwood City site.			
MRGC Necklace	Oyster shell necklace at Marin Rod and Gun Club			
MRGC Pallets	Oyster shell pallets at Marin Rod and Gun Club site.			

Shellfish Beds – HABITAT TYPE AND ASSOCIATED BIOLOGICAL ASSEMBLAGES

Table 7. Sessile invertebrates associated with native oyster restoration sites at Marin Rod and Gun Club and Redwood City sites in San Francisco Bay from MACTEC (2006).

	Species	Native Oyster Restoration		
		RC Pallets	MRGC Necklace	MRGC Pallets
Bivalves	<i>Venerupis philipparum</i>	X		X
	<i>Musculista senhousia</i>	X	X	X
	<i>Ostrea conchaphila</i>	X	X	X
	<i>Mytilus edulis</i> (mussel)		X	
Bryozoans	<i>Schizoporella</i> sp.	X	X	X
	<i>Bugula</i> sp.	X	X	X
	<i>Watersipora subtorquata</i>	X		
	<i>Conopeum</i> sp.			X
	Other unknown species	X	X	X
Tunicates	<i>Molgula manhattensis</i>	X	X	X
	<i>Styela clava</i>	X	X	X
	<i>Ciona</i> sp.	X		
	<i>Botrylloides</i> sp.	X		

RC Pallets Oyster shell pallets at Redwood City site.
MRGC Necklace Oyster shell necklace at Marin Rod and Gun Club
MRGC Pallets Oyster shell pallets at Marin Rod and Gun Club site.




Figure 11. Historic native oyster distribution from Barrett (1963), sites of known oyster occurrence at present from Harris (2004), and native oyster restoration sites in San Francisco Bay. Historic oyster distribution is for the period prior to 1915 and is an approximation.

PLANT BEDS

(by Natalie Cosentino-Manning, Katie McGourty and Susan Wainwright-De La Cruz)

General Description

Subtidal plant beds in San Francisco Bay include two major groups: algal beds (both macro and micro) and angiosperm beds, more commonly referred to as submerged aquatic vegetation (SAV). Although there is very little historical data (pre-gold rush) for the two groups, surveys in the past 20 years described the Bay as having the same taxonomic diversity as other estuaries on the west coast (Josselyn and West 1985).

Macroalgae Beds

Macroalgae occur in San Francisco Bay both as floating drift and as attached plants. Most of the macroalgae seen as drift in the Bay has been ripped off the outer coast by strong currents or storms and carried into the Bay with the tides (N. Cosentino-Manning, NMFS, pers. comm.). This generally occurs during winter storms or during the upwelling months (April to June), when offshore winds are strong. The macroalgae (benthic and drift) belong to the taxonomic groups Chlorophyta (green), Rhodophyta (red), and Phaeophyta (brown) and are mostly found within Central Bay, where salinity levels are mainly euhaline. However, the green algal distribution extends further into the polyhaline and mesohaline parts of the Bay, although exact distribution limits for a particular species are difficult to determine (Wilkinson 1980).

Evidence from other estuaries within the United States show that estuarine macroalgae contribute significantly to a number of estuarine processes. Productivity of macroalgae can at times match or exceed that of phytoplankton (Dillion 1971) and can act as a nutrient sink. Algae also deteriorate faster than terrestrial plants and therefore can contribute significantly to the particulate organic matter in estuaries (Josselyn and Mathieson 1980).

In this report we focus on four species of bed forming macroalgae, *Ulva*, *Gracilaria*, *Fucus*, and *Sargassum*.

Ulva Beds

Probably the most conspicuous alga along the shores of San Francisco Bay, especially at low tide, is *Ulva*. *Ulva* species live primarily in euhaline to polyhaline environments. It lives attached to rocks or on mudflats in the middle to low intertidal zone, and as deep as 10 meters in calm, protected harbors. *Ulva* plants are usually seen in dense groups. Commonly known as the sea lettuce, the genus includes species that look like bright, green sheets. Shapes of *Ulva* species are quite varied from circular to oval to long and narrow and tubular, ranging in size from microscopic to 65 cm.

Ulva species have high nitrogen requirements, a reduced ability to take up nitrogen in low nitrogen conditions, and a limited ability to store nitrogen. These features can limit the distribution of *Ulva* to nitrogen-rich environments. When nitrogen is available in particularly high concentrations, however, *Ulva* species are able to take up nitrogen quickly and use it to grow rapidly. Thus, nuisance growth of *Ulva* species can occur in areas with nitrogen-rich sewage pollution, especially if they are within enclosed or semi-enclosed sub-embayments and experience little mixing (Valiela *et al.* 1997). In these areas, *Ulva* species comprise a large proportion of drift plants, and can smother the benthic communities below. Species associated with *Ulva* include the isopods *Idotea* sp. and *Sphaeroma* sp. and the amphipod *Gammarus* sp. Small littorine snails are also associated with *Ulva*. (N. Cosentino-Manning, NMFS, pers. comm.).

Gracilaria Beds

Gracilaria pacifica (formerly *verrucosa*), also known as the red spaghetti alga or agar, is a resident species to San Francisco Bay and can be found within quiet embayments such as Richardson Bay or on the leeward side of islands such as Treasure Island and Angel Island. *Gracilaria* requires a euhaline to polyhaline environment and can be found

attached or floating (free-living). Many times it is found in areas where eelgrass (*Zostera marina*) is growing. It is unknown if there is a link between the two species or if similar habitats types are preferred.

Gracilaria appearance ranges from dark brown to bright red in coloration. *Gracilaria* is considered a “bloom” species because it has the ability to grow and dominate a community under increased nitrogen loads. In some estuaries, *Gracilaria* has even replaced *Zostera* beds (Valiela *et al.* 1997). It is unknown whether the current abundance levels for *Gracilaria* in the San Francisco Bay are at normal or bloom levels.

Species associated with *Gracilaria* include the isopods, *Idotea* sp. and *Sphaeroma* sp., and the amphipod, *Gammarus* sp. (N. Cosentino-Manning, NMFS, pers. comm.). The red algal parasite, *Gracilariophila oryzoides*, was found on specimens collected from 1970 to 1980 (Silva 1979). The Pacific herring is the only documented fish associated with *Gracilaria*. The CDFG has documented herring spawning on *Gracilaria* (Tom Moore, CDFG, pers. comm.). Herring may benefit from high abundance of *Gracilaria* because of its increased surface area for spawning.

Fucus Beds

Another conspicuous alga on the intertidal shores of San Francisco Bay is the brown furoid algae *Fucus gardneri*, also known as rockweed. *Fucus* is flat and dichotomously branched and ranges in color from golden brown to almost black if dessicated. When reproductive, the tips of the plant become inflated and a mucus discharge is evident. *Fucus* is found throughout the euhaline Central Bay in sheltered and exposed intertidal habitats. It is most apparent on hard bottom substrate such as rip-rap, boulders, and concrete harbor pilings.

Fucus provides habitat for many sessile and motile invertebrates. Small acorn barnacles such as *Balanus glandula* and *Chthamalus* sp. reside under the wet blades of *Fucus* and are protected from desiccation during the low tide cycle. Gammaridian amphipods are also found in

large congregations between and under the blades. (N. Cosentino-Manning, NMFS, pers. comm.).

Sargassum Beds

Sargassum muticum is a brown algae (Phaeophyta) originating from Japan and introduced to the West coast of the United States in 1945 (Abbott and Hollenberg 1976). The alga is yellowish- or olive-brown and can be distinguished from most other Pacific coast seaweeds by its small, spherical air bladders (pneumatocysts) that allow the alga to rest on the water at high tide. In San Francisco Bay, *Sargassum* can be found throughout the euhaline and polyhaline portions of the Bay and is mostly restricted to calm embayments or within harbors and marinas. It grows attached to rocks, shells or other hard objects, in the lower intertidal zone to the shallow subtidal (3 m to 5 m). It can grow up to 4 m in length.

There are many reports of *Sargassum muticum* competing with and displacing native species of seaweed and eelgrass, at least in part by shading and reduction of light levels. At Santa Catalina Island in southern California, it appears to have taken over large shallow beds of the seaweed *Scytosiphon* (Nicholson *et al.* 1981). It may also exclude native species by occupying areas when they become temporarily vacant, as was reported following a die-back of the kelp *Macrocystis pyrifera* at Santa Catalina Island, and for eelgrass die-backs in Atlantic France (Den Hartog 1997). Silva (1979) reported that "there is no evidence that *Sargassum muticum* is displacing the native biota of San Francisco Bay." There is no data on the distribution or abundance of *Sargassum* in San Francisco Bay and how it affects other species in the Bay.

Sargassum muticum beds do provide habitat to several invertebrate species such as caprellidae amphipods, isopods, littorine snails and juveniles of both the red rock crab and the Dungeness crab.

Submerged Aquatic Vegetation

Submerged aquatic vegetation (SAV) are rooted, vascular, flowering plants that, except for some of the flowering structures, live and grow below the water surface. SAV can be found in euhaline to oligohaline environments within the Bay. Although SAV sometimes occurs intertidally or extends to the water's surface, these plants are generally submerged and cannot survive if removed from the water for any length of time (Hurley 1990). The leaves and stems of SAV have specialized thin-walled cells with large intracellular air spaces to provide buoyancy and support. Leaves and stems are generally thin and lack the waxy cuticle found in terrestrial plants. The lack of a waxy cuticle increases the exchange of water, nutrients, and gases between the plant and the water (Hurley 1990). The extensive root and rhizome system anchors the plants, and also absorbs nutrients (Thayer *et al.* 1984). Reproduction occurs both sexually and asexually.

SAV in San Francisco Bay has been poorly studied and very little is known about its distribution and abundance. Four types of SAV communities are known to occur within San Francisco Bay; surfgrass (*Phyllospadix scouleri*), eelgrass (*Zostera marina*), widgeon grass (*Ruppia maritima*) and sago pondweed (*Potamogeton pectinatus*). Each species of SAV has varying salinity requirements and is found in distinct parts of the Bay. Based on limited surveys of the Bay, the most widely distributed SAV habitat is eelgrass (Merkel & Associates, Inc. 2003), followed by widgeon grass, sago pondweed and surf grass (Figure 12). All four SAV species provide primary productivity and decrease erosion by dampening wave action, preventing sediment resuspension, increasing sedimentation, providing attachment for sessile organisms, and providing a resource area for invertebrates, fishes, birds, and marine mammals.

Sago Pondweed (*Potamogeton pectinatus*) Beds

Sago pondweed (*Potamogeton pectinatus*) is found in fresh non-tidal to moderately oligohaline waters; sago pondweed can die back when water salinity exceeds 15 ppt (Kantrud 1990). It can tolerate high alkalinity and grows on silty-muddy sediments. It tolerates strong

currents and wave action better than most bay grasses because of its long rhizomes and runners. The precise abundance and distribution of sago pondweed in San Francisco Bay have not been well documented, but anecdotal evidence suggests that beds are mainly located in the shallow areas of Suisun Bay surrounding Simmons, Ryer, and Roe Islands (N. Cosentino-Manning, NMFS, pers. comm.) and in Little Honker Bay (S. Siegel, Wetlands and Water Resources, Inc., unpub. data). Feminella and Resh (1989) conducted a 3-year study on sago pondweed density in Coyote Hills Marsh, a man-made marsh in Alameda County. The study cover area contained 71 percent sago pondweed in 1984. By 1986, sago pondweed had been eliminated by the invasive crawfish *Procambarus clarkii*.

Sago pondweed is considered one of the most valuable food sources for waterfowl in North America. Its highly nutrient seeds and tubers, as well as leaves, stems, and roots, are consumed by numerous species of ducks, geese, and marsh and shorebirds. Martin *et al.* (1951) found widgeongrass composed 10 to 25 percent of the diet of redheads (*Aythya americana*) and scaup (*Aythya* spp.); 5 to 10 percent of the diet of pintails (*Anas acuta*) and ruddy ducks (*Oxyura jamaicensis*); and 2 to 5 percent of the diet of mallards (*Anas platyrhynchos*), canvasback (*Aythya valisineria*), and green-winged teal (*Anas crecca*). Sago pondweed has been found to compose 50 percent or more of the diet of canvasbacks; 25 to 50 percent of the diet of mallards and redheads; and 10 to 25 percent of the diet of pintails, green-winged teal, and scaup (Martin *et al.* 1951). The importance of sago pondweed as a waterfowl food resource in San Francisco Bay is unknown.

Widgeon Grass (*Ruppia maritima*) Beds

Like eelgrass, widgeongrass (*Ruppia maritima*) is found in polyhaline to euhaline regions, however, widgeon grass can tolerate and grow in both fresh water and high-salinity environments. Widgeon grass has been documented within Suisun Bay, Lake Merritt (MSI 2005), and within one South Bay salt pond (Anderson 1970). Widgeon grass has become a management issue for Lake Merritt and is routinely harvested for removal.



Figure 12. Distribution of eelgrass (*Zostera marina*), the marine alga, *Gracilaria*, and widgeon grass (*Ruppia maritima*) in San Francisco Bay after Merkel & Associates (2003) and N. Cosentino-Manning (NMFS, pers. comm.).

Eelgrass (*Zostera marina*) Beds

Eelgrass, *Zostera marina*, is a subtidal marine flowering vascular plant, with all stages of the life cycle occurring underwater, including flowering, pollination, and seed germination. Eelgrass is widespread through the Atlantic and Pacific and has a restricted distribution in the Mediterranean. It is the only SAV species that extends into the Arctic Circle.

Intermittent eelgrass surveys suggest eelgrass abundance has varied greatly in San Francisco Bay in the last several decades. In the late 1920s, eelgrass was reported as an abundant species along the shores of San Francisco Bay (Setchell 1929). In 1987, 60 years later, a survey of the Bay revealed only 128 hectares (only 0.1 percent total Bay bottom coverage) of eelgrass throughout the San Francisco Bay, with much of the existing habitat exhibiting conditions of environmental stress (Wyllie-Echeverria and Rutten 1989, Wyllie-Echeverria 1990). A decade later, surveys of the San Pablo Peninsula documented over 162 hectares of eelgrass (SAIC and Merkel & Associates, Inc. 1999a, 1999b). However, in a series of acoustic surveys between June 4 and October 12, 2003, 1165.7 hectares of eelgrass were documented (Merkel & Associates, Inc. 2004, Figure 13).

Currently, eelgrass beds within San Francisco Bay are found in euhaline to polyhaline salinities and are found mainly along the eastern shores of San Pablo and Central Bays. Existing eelgrass beds cover approximately 1 percent of San Francisco Bay, which is at least an order of magnitude less than seen in other large estuarine systems (Merkel & Associates, Inc. 2003) (Figure 12). Due to the large size of the San Francisco Bay, however, even a minor proportional representation amounts to substantial eelgrass resource on a state wide basis.

The majority of the eelgrass in San Francisco Bay is located on the east shoreline between Point Pinole and Bayfarm Island (Figure 13). A few solitary plants are recorded north of Point Pinole near Wilson Point and several individual plants and two small patches are observed south of the San Mateo Bridge along the Alameda shoreline. The range of eelgrass distribution

during the Merkel 2003 survey and the Wyllie-Echeverria and Rutten 1989 survey were similar.



The largest eelgrass bed in the Bay is the Point San Pablo bed, which is located between Point Pinole and Point San Pablo north of the Richmond-San Rafael Bridge (Figure 13). This bed was approximately 608.9 hectares (1,504.5 acres) during 2003, and comprised 52.2 percent of the total eelgrass coverage of the Bay (Merkel & Associates, Inc. 2003). The bed lies on a shallow depositional shoal approximately -1.5 to -0.5 m MLLW.

The second largest eelgrass bed is found in Richardson Bay near Sausalito in Marin County (Figure 13). This bed is approximately 176.7 hectares (436.7 acres) during the 2003 survey, the densest part of which was located among the boat moorings and near the marinas on the western side of the Bay. Eelgrass in Richardson Bay occurred between -3.0 and -0.5 m MLLW.

Eelgrass beds are important habitats in euhaline and polyhaline waters because they are home to many small organisms that are food for larger species and they provide protective cover for migrating salmon, provide spawning substrate for Pacific herring, and act as a nursery for many other smaller fish such as gobies. Eelgrass stabilizes and binds substrates and absorbs nutrients from sediments. They reduce water currents by frictional forces, dampen wave energy, and slow erosional processes. They are primary producers removing inorganic nutrients from the sediments and the water column and through photosynthesis convert them into organic material.

Plants

Epiphytic plants, such as diatoms and microscopic green algae, can be found on eelgrass blades. *Gracilaria* sp. often co-occurs with SAV for reasons not understood. Other macroalgal genera found in eelgrass include salt/brackish (*Ulva*, *Codium*, *Gracilaria*) species and the macroalgae, *Ectocarpus*, *Cladomorpha*, and *Chaetomorpha* (Mallin *et al.* 2000, Thayer *et al.* 1984).



2003 Eelgrass (*Zostera marina*) Density (Percent Cover)



Figure 13. Eelgrass (*Zostera marina*) density in San Francisco Bay in 2003 after Merkel & Associates (2004) and Wyllie Echeverria et al. (1989).

Invertebrates

Epiphytic invertebrates such as bryozoans and hydroids are commonly observed on eelgrass blades (Kozloff 1973). Herbivorous snails such as *Littorina* sp. and *Tegula* sp. graze on epiphytic diatoms and algae on eelgrass blades, while carnivorous whelks such as *Nassarius* sp. and hermit crabs are commonly observed (Ricketts *et al.* 1985). Along the Pacific Coast, several species of nudibranchs (sea slugs) can be found among eelgrass blades, including *Mebile leonina*, which feeds on small crustaceans, *Aeolidia papillosa*, feeding on anemones, and *Phyllaplysia taylora*, a photosynthetic nudibranch endemic to eelgrass beds (Kozloff 1973). Crustaceans inhabiting Pacific Coast eelgrass beds include skeleton shrimp (*Caprella californica*), the Dungeness crab, the red crab, the graceful crab (*Cancer gracilis*), and amphipods (small crustaceans). Similar species may inhabit eelgrass beds in San Francisco Bay. Merkel & Associates, Inc. (2003) conducted six benthic infaunal samples from the San Francisco Bay from August 5 to 8 and September 22 to 25. Arthropods (composed primarily of amphipods), mollusks, and annelid worms were the most abundant phyla found (Merkel & Associates, Inc. 2005).

Fishes

Eelgrass beds throughout California serve as a nursery area for several fish species including rockfish (*Sebastes* sp.) and surfperch (*Embiotocidae*), and a refuge and foraging area for year round resident fish species such as the bat ray (*Myliobatis californica*) and the bay pipefish (*Syngnathus leptorhynchus*) (Table 4). Two studies have been undertaken in San Francisco Bay to determine fish species utilization of eelgrass beds. ENTRIX, Inc. (1998) conducted a survey of fish communities in eelgrass in relation to the Port of Oakland's Harbor Navigation Improvement Project in the spring, summer, and fall of 1997. Twenty-two fish species were observed; the most abundant were Pacific herring (*Clupea pallasii*), shiner surfperch (*Cymatogaster aggregata*), speckled sanddab (*Citharichthys stigmaeus*), bat ray, the bay pipefish, and black perch (*Embiotoca jacksoni*). Merkel & Associates, Inc. (2005) and NOAA Fisheries conducted fish surveys from June 14 to 17 in 2004 using four different gear

types (purse seine, large seine, otter trawl, and minnow traps). Thirty-six fish species were observed, with topsmelt (*Atherinops affinis*), shiner surfperch, dwarf surfperch (*Micrometrus minimus*), and northern anchovy (*Engraulis mordax*) most abundant.

Eelgrass beds provide refuge for a diverse assemblage of fishes uncommon in other areas of San Francisco Bay. Black perch (*Embiotoca jacksoni*) are relatively rare in San Francisco Bay; CDFG found only 77 individuals over 15 years of monthly monitoring (DeLeon 1999). Black perch mate from April to June, and after a six-month gestation period give birth to an average brood of 14 offspring from September to November (DeLeon 1999, Baltz 1984). The bay pipefish is closely related to the Pacific seahorse. Members of this family are unique because males brood and care for young in a specialized brooding pouch.

Birds

Similar to the unconsolidated sediment and shellfish bed habitats, diving benthivores commonly forage in SAV (Table 5). In addition to diving benthivores, piscivorous terns including the Caspian tern (*Sterna caspia*), the Forester's tern (*S. forsteri*), and the elegant tern (*S. elegans*) forage in eelgrass beds. Forester's tern can be observed year-round in San Francisco Bay. During spring and fall, Forester's terns migrate from breeding grounds to feeding grounds and become more abundant within San Francisco Bay. During the winter months, a comparatively smaller number of Forester's terns (28 breeding colonies) breed in South Bay (Goals Project 2000). Caspian terns, the largest of tern species found in San Francisco Bay, migrate great distances between breeding and foraging grounds and can be found on every continent except Antarctica. Caspian terns arrive to San Francisco Bay in late spring to breed and are present until August (Goals Project 2000). Caspian tern breeding sites are concentrated in Central and San Pablo Bays (Goals Project 2000). Elegant terns use San Francisco Bay as a stop-over point during their migration from late spring through early fall. Annual abundance varies in accordance with oceanographic conditions (Fix and Bezener 2000). All three tern species feed by hovering over the water and making quick dives for small, schooling fishes.

Marine Mammals

Marine mammals often feed in submerged aquatic vegetation where high abundances of fish and invertebrates occur. Eelgrass, in particular, provides an important foraging habitat for all seven species of marine mammals found within San Francisco Bay (Table 6) due to high concentrations of prey items such as schooling fishes. Eelgrass is the most productive foraging area for California sea lions and harbor seals during winter months when Pacific herring are in high abundance (S. Allen, National Park Service, pers. comm.). Harbor seals in this habitat will also forage on perch, top smelt, anchovies, California halibut, and speckled sanddabs. Gray whales periodically feed on epiphytic growth on eelgrass, with the ability to feed in depths as shallow as 1.5 to 2.5 m (S. Allen, National Park Service, pers. comm.). *Gracillaria* beds are similarly important foraging habitat for marine mammals because herring and other fish congregate there. Harbor seals will also rest in eelgrass beds, laying on the bottom and surfacing to breathe.

Surfgrass (*Phyllospadix scouleri* and *torreyi*) Beds

The surfgrasses, *Phyllospadix scouleri* and *torreyi* are similar to eelgrass in that they are angiosperms with true leaves, stems, rootstocks, and flowers. Surfgrass can be found from Alaska to Baja California, Mexico. Surfgrass grows attached to rocks, many of which are exposed at low tide. Surfgrass occurs below the mean lower low water level (Ricketts *et al.* 1985) and requires euhaline salinity with exposure to wave action. Surfgrass is found only at the entrance to San Francisco Bay and is not found any further east than the two forts (Fort Baker and Fort Point) that flank the Golden Gate Bridge (N. Cosentino-Manning, NMFS, pers. comm.).

The ecology and importance of *Phyllospadix* is not known nearly as well as that of eelgrass. *Phyllospadix* vegetation

can protect rocky substrate from erosion and transform it into sandy beaches or sublittoral sand flats by accumulating sand in and between the tussock (Phillips 1979). Surfgrass provides habitat for many species of algae (Stewart and Myers 1980) and provides shelter for many invertebrates. The red algae, *Smithora naiadum* and *Melobesia mediocris*, and the green algae, *Kormmannia zostericola*, (Kozloff 1973) are exclusively epiphytic on sea grasses, such as surfgrass (Abbott and Hollenberg 1976). Small herbivorous snails including the chink snail (*Lacuna* sp.) and limpets such as *Notoacmea paleacea* graze on epiphytic algae found on surfgrass (Kozloff 1973).

Colonial tunicates, sponges, various species of amphipods, the limpet, *Notoacmea insessa*, the sea anemone, *Anthopleura xanthogrammica*, and crabs such as the kelp crab (*Pugettia producta*), and Cancer crabs (*Cancer antennarius* and *productus*) inhabit surfgrass beds. Isopods (*Isopodea* sp.), a type of crustacean, cling to blades and feed on surfgrass leaves (Kozloff 1973). Several species of nudibranchs can also be found in protected areas of surfgrass, including *Hermisenda crassicornis*, the Spanish shawl (*Flabellinopsis iodinea*), and the sea lemon (*Anisodris nobilis*). Surfgrass also provides nursery habitat for fishes such as gobies, gunnels, and the monkey-faced eel (*Cebidichthys violaceus*) (Engle 1979). Marine mammals such as the harbor seal (*Phoca vitulina*) can be seen resting at low tide on *Phyllospadix* covered rocks. (N. Cosentino-Manning, NMFS, pers. comm.).

Phyllospadix is susceptible to desiccation and heat stress during low midday tides (Raimondi *et al.* 1999). It is also sensitive to sewage (Litler and Murray 1975) and oiling (Foster *et al.* 1998). If the rhizome systems remain viable, then recovery following disturbance can be fairly rapid; however, recovery is long if the entire bed is lost because recruitment is irregular (Turner 1983, 1985), and restoration projects have been unsuccessful.

WATER COLUMN

(by Korie Schaeffer, Kathryn Hieb, Katie McGourty, Susan Wainwright-De La Cruz, and Sarah Allen)

General Description

The water column consists of the area between the benthos and the water surface. Temperature, salinity, dissolved oxygen, and turbidity vary within the water column depending on depth, location, and season. Species assemblages occur in the lower, middle, and upper portion of the water column depending on available cover, light, and predator-prey interactions. The water column may be classified as a channel (deeper area with “u” or “v” shape geomorphology with stronger currents) or shoals (shallow, flat areas with weaker currents).

Water column habitat allows for horizontal and vertical water transport and mixing, tidal propagation, and suspension or deposition of sediments, and provides a medium for primary and secondary production, foraging areas for invertebrates, fishes, birds, and marine mammals, and nursery and spawning areas for fishes and invertebrates. Shoals function as recipients and dispersers of non-local sources of detritus. Channels provide a connection from marine to freshwater ecosystems.

Taxonomic Groups

Plants (phytoplankton)

Phytoplankton of San Francisco Bay includes those species produced within the water column, those that are exchanged between the water column and the sediments, those transported into the Bay with freshwater flow, and those transported into the Bay with tidal exchange from the Pacific Ocean. Cloern and Dufford (2005) characterized the phytoplankton community based on seasonal samples taken along the salinity gradient from the Sacramento River to Central Bay to South San Francisco Bay between 1992 and 2001. The phytoplankton assemblage was dominated by a small

number of species; diatoms were the dominant phytoplankton type, followed by dinoflagellates and cryptophytes. Each of these three groups of phytoplankton contributed 81, 11, and 5 percent, respectively, to cumulative biomass. A species list from this study is provided in Table 8. Within this list, the top 17 ranked species include 12 diatoms, 2 dinoflagellates, 2 cryptophytes, and *Mesodinium rubrum*. During the period of sampling, some phytoplankton taxa apparently disappeared after 1996 while others first appeared and have occurred yearly since 1997 to 1998. The general timing of these changes mirror large-scale changes in ocean temperatures, regional wind patterns, and biological communities across the Pacific basin (Chavez *et al.* 2003). (Cloern and Dufford 2005)

Many phytoplankton species occur across broad salinity and temperature ranges, and are not easily classified into specific salinity categories. For example, picocyanobacteria and some small eukaryotes (e.g., *Nannochloropsis* sp., *Teleaulax amphioxeia*, *Plagioselmis prolonga*) are persistent and ubiquitous across large habitat gradients in San Francisco Bay. Exceptions to this general rule include more commonly marine species (e.g., *Thalassiosira frauenfeldii*, *Ceratium furca*, *Pyramimonas parkeae*, *Ceratium* spp., *Alexandrium catenella*, *Prorocentrum micans*, *P. gracile*, *Dinophysis acuminata*, *Heterosigma akashiwo*), which are confined to areas of higher salinity; and freshwater taxa, such as *Skeletonema potamos*, which are transported to San Francisco Bay with river flow. (Cloern and Dufford 2005)

Historically, median phytoplankton primary production in San Francisco Bay has been relatively low (120 gCm⁻²yr⁻¹) compared to other estuaries world wide (average 200 gCm⁻²yr⁻¹) (Jassby *et al.* 2002). Eutrophication and hypoxia, which are of concern in many estuaries, including Chesapeake Bay, are not high-priority concerns for San Francisco Bay. Despite the relatively low production levels, phytoplankton photosynthesis is the primary energy supply to food webs of San Francisco Bay (Jassby *et al.* 1993, Sobczak *et al.* 2002, 2005).

Table 8. Common phytoplankton taxa in San Francisco Bay from Cloern and Dufford (2005).

Only those species that occurred in >10 (of 599) samples and contributed >0.1% of cumulative biomass as biovolume contained in all samples; n=number of occurrences.

Species	Division	Biomass (%)	n
<i>Thalassiosira rotula</i>	Bacillariophyta	20.97	140
<i>Chaetoceros socialis</i>	Bacillariophyta	11.98	48
<i>Skeletonema costatum</i>	Bacillariophyta	9.51	277
<i>Ditylum brightwellii</i>	Bacillariophyta	7.44	103
<i>Gymnodinium sanguineum</i>	Pyrrophyta	7.40	41
<i>Coscinodiscus oculus-iridis</i>	Bacillariophyta	6.30	66
<i>Thalassiosira hendeyi</i>	Bacillariophyta	4.85	203
<i>Thalassiosira punctigera</i>	Bacillariophyta	3.31	27
<i>Plagioselmis prolonga</i> var. <i>nordica</i>	Cryptophyta	2.65	495
<i>Coscinodiscus curvatulus</i>	Bacillariophyta	2.15	49
<i>Mesodinium rubrum</i>	Holotrich ciliate	2.03	190
<i>Teleaulax amphioxeia</i>	Cryptophyta	2.02	375
<i>Chaetoceros debilis</i>	Bacillariophyta	1.84	31
<i>Eucampia zodiacus</i>	Bacillariophyta	1.80	29
<i>Coscinodiscus radiatus</i>	Bacillariophyta	1.77	66
<i>Thalassiosira eccentrica</i>	Bacillariophyta	1.48	136
<i>Protoperdinium</i> sp.	Pyrrophyta	1.45	21
<i>Thalassiosira decipiens</i>	Bacillariophyta	0.90	136
<i>Coscinodiscus centralis</i> var. <i>pacifica</i>	Bacillariophyta	0.72	23
<i>Rhizosolenia setigera</i>	Bacillariophyta	0.49	102
<i>Noctiluca scintillans</i>	Pyrrophyta	0.46	36
<i>Nitzschia bilobata</i>	Bacillariophyta	0.44	33
<i>Cyclotella atomus</i>	Bacillariophyta	0.43	304
<i>Coscinodiscus jonesianus</i>	Bacillariophyta	0.33	17
<i>Pyramimonas orientalis</i>	Chlorophyta	0.31	175
<i>Rhodomonas marina</i>	Cryptophyta	0.26	116
<i>Protoperdinium depressum</i>	Pyrrophyta	0.23	41
<i>Heterocapsa triquetra</i>	Pyrrophyta	0.23	138
<i>Protoperdinium claudicans</i>	Pyrrophyta	0.22	52
<i>Cyclotella choctawhatcheeana</i>	Bacillariophyta	0.22	74
<i>Alexandrium tamarense</i>	Pyrrophyta	0.21	109
<i>Nannochloropsis</i> sp.	Chrysophyta	0.20	424
<i>Thalassiosira nodulolineata</i>	Bacillariophyta	0.18	40
<i>Chlorella salina</i>	Chlorophyta	0.17	97
<i>Chaetoceros wighamii</i>	Bacillariophyta	0.17	43
<i>Eutreptia lanowii</i>	Euglenophyta	0.16	167
<i>Prorocentrum minimum</i>	Pyrrophyta	0.16	116
<i>Aulacoseira lirata</i>	Bacillariophyta	0.17	43
<i>Rhizosolenia styliformis</i>	Bacillariophyta	0.13	22
<i>Entomoneis paludosa</i>	Bacillariophyta	0.12	30
<i>Odontella mobiliensis</i>	Bacillariophyta	0.11	17
<i>Gyrosigma balticum</i>	Bacillariophyta	0.11	36

Phytoplankton biomass and growth in San Francisco Bay is controlled by a number of factors, including light availability (Alpine and Cloern 1988, Cloern 1999), residence time driven by horizontal transport by freshwater inflow, and predation by benthic suspension feeders (Jassby *et al.* 2002). Nutrient limitation is not usually a controlling factor for phytoplankton in San Francisco Bay. (Jassby *et al.* 2002)

Jassby *et al.* (2002) reconstructed annual primary productivity from historical water quality data for 1975 to 1995 for the Delta, including Suisun Bay. These reconstructed values do not apply to all of San Francisco Bay, but provide an idea of levels in northern San Francisco Bay. Annual primary production averaged 70 gCm⁻² but varied by over a factor of five among years, likely due to: invasion of the clam, *Corbula amurensis*; long-term decline in total suspended solids and associated increase in water transparency; fluctuation in residence time with river inflow; and unexplained winter biomass declines (Jassby *et al.* 2002).

The most notable of these factors is the invasion of *Corbula amurensis*, which became widely established in 1987. Following invasion, chlorophyll biomass became persistently low and primary production was reduced 5-fold in the northern part of San Francisco Bay (Alpine and Cloern 1992). The concurrent decrease in total suspended sediments represented a 25 percent increase in the photic zone depth. This should have translated to a higher growth rate and higher primary productivity for light-limited phytoplankton in the Bay. It appears, however, the increased predation by *Corbula* more than compensated for this potential increase in production due to increased light availability (Jassby *et al.* 2002).

Since the late 1990s, the “baseline” or annual minimum chlorophyll in San Francisco Bay downstream of Suisun Bay has increased 5 to 10 percent per year, blooms are observed in winter-fall and not just in the spring and spring blooms are larger between San Pablo Bay and San Mateo Bridge. As a result average primary productivity has increased from 120 gCm⁻²yr⁻¹ in 1993 to 1996 to 215 gCm⁻²yr⁻¹ in 2001 to 2004 (Jassby *et al.* 2002), bringing productivity levels in these regions of San Francisco Bay

within the range often measured in temperate latitude estuaries. The specific reasons for these changes are unknown, but Cloern *et al.* (2006) proposed relaxed light limitation, reduced metal loading from progress in wastewater treatment, increased ocean sources of phytoplankton, and increased predation of bivalves by an opisthobranch, fishes, or birds and of zooplankton by an invasive predatory copepod. In contrast to southern San Francisco Bay, productivity in Suisun Bay where *Corbula* persists remains low. (Cloern *et al.* 2006)

In addition to inter-annual trends in production, phytoplankton blooms can occur in both the northern and southern parts of San Francisco Bay causing high seasonal variability. The timing, location, and magnitude of blooms change from year to year, depending on tides, river flow, wind dynamics, and nutrients.

South San Francisco Bay has been described as a lagoon-type estuary, with a longer residence time, lower turbidity levels, and higher phytoplankton population growth than northern San Francisco Bay (i.e., San Pablo Bay and Suisun Bay) (Cloern *et al.* 1996). Historically, there was slow phytoplankton growth in South San Francisco Bay through the winter, with increasing growth during spring that culminated in a spring bloom. *Mesodinium* blooms occur during years when the water column stratifies and dissolved inorganic nitrogen and silicon in the water column are depleted (Cloern 1996, Cloern *et al.* 1994). As noted above, smaller blooms have recently been observed during fall and winter, with the bloom intensity increasing towards the southern extreme end of the San Francisco Bay. Spring blooms are the source of much of the total annual primary production in South San Francisco Bay (Cole *et al.* 1986).

Annual blooms in South San Francisco Bay occur when physical processes, such as freshwater inflow and weak tidal mixing, result in reduced vertical mixing. The resulting decrease in turbidity releases phytoplankton populations from strong light limitation (Cloern 1996, Lucas *et al.* 1999a, 1999b, May *et al.* 2003). During years with high freshwater inflow, salinity stratification can isolate some of the phytoplankton cells in the euphotic zone resulting in large magnitude, persistent blooms.

Stratification aids phytoplankton growth by increasing light availability for the surface-dwelling phytoplankton and by reducing the loss of phytoplankton to the bottom-dwelling, benthic grazers (Cloern *et al.* 1985, Lucas *et al.* 1998). Smaller, localized blooms can also occur during periods with weak tides and low wind mixing.

The general absence of blooms in summer and early fall in South San Francisco Bay is caused by high vertical mixing rates due to the strong diurnal winds during this period, as well as from an increase in predation by benthic predators, whose biomass and grazing rates are highest in the summer (Cloern 1996).

The northern San Francisco Bay is more river-influenced than South San Francisco Bay. As such, phytoplankton residence time is lower and turbidity is higher. Historically, there was a prolonged summer bloom of netplanktonic diatoms in Suisun Bay, coinciding with the accumulation of suspended particulates at the convergence of freshwater outflow and tidal inflow (i.e., convergence zone). This convergence only occurred when river discharge fell within a narrow flow range (100 to 400 m³/s; Cloern *et al.* 1985). This freshwater flow range results in higher water column residence times and places the convergence zone adjacent to shallow embayments in the northern Bay. Thus, shallow water-generated phytoplankton was retained in the convergence zone, resulting in a bloom. At higher discharge, the convergence zone moves to the west and phytoplankton residence times are short relative to phytoplankton population growth rates, preventing populations from increasing to bloom densities. At lower discharge, the convergence zone moves east to the Delta. During more recent years, blooms have disappeared, likely due to the high rate of benthic grazing (Alpine and Cloern 1992, Cloern 1996) and the high ambient ammonia concentrations, which inhibit nitrate uptake by phytoplankton for growth (Wilkerson *et al.* 2006).

At smaller geographic scales, mechanisms of spatial variability in phytoplankton production throughout San Francisco Bay include seasonal blooms, exchanges between sediment and water column, and horizontal dispersion from regions of high productivity in the

shallows and landward areas to areas of low productivity in the deep channels and seaward areas (May *et al.* 2003, Cloern *et al.* 1985). May *et al.* (2003) modeled effects of turbidity and horizontal and vertical transport of shoal and channel areas of San Francisco Bay and determined that, given constant benthic grazing losses, different rates of vertical and horizontal clearing determine if blooms will be channel or shoal supported and whether they will be local or systemwide. Also, while nutrients are not generally limiting in the Bay, localized concentrations of nutrients can be depleted by phytoplankton blooms, causing a bloom to die off within a given area.

Invertebrates (zooplankton)

Zooplankton includes small organisms which spend their entire life cycles within the water column such as copepods, as well as life history stages of organisms with a pelagic life history phase, such as fishes and large invertebrates. The primary taxa in San Francisco Bay are comprised of small forms (microzooplankton) including tintinnids, rotifers, and copepod nauplii, larger zooplankton including copepods, meroplankton (larvae of benthic animals and fish), and cladocerans (Table 9).

Zooplankton are often divided into freshwater, brackish, and marine species. Because zooplankton are incapable of swimming against tidal or freshwater currents, they move with a water body, and thus, are more likely to be oriented by salinity than geography. In this section, zooplankton are often discussed as northern Bay (Suisun and sometimes San Pablo Bay) and southern bay (South Bay, Central Bay and sometimes San Pablo Bay). The demarcation between northern and southern Bay for zooplankton is driven primarily by salinity.

Some zooplankton are present in San Francisco Bay throughout the year, while others have distinct seasonal patterns. Abundance patterns for microplankton have not been well studied in the Bay, so the focus here is on copepods, mysid shrimp, juveniles of fish and other invertebrates, and hydromedusae.

Table 9. Zooplankton taxa from San Francisco Bay after Kimmerer et al. 1999.

Taxonomic Group		Species	Status
Copepods	Calanoid	<i>Acartia californiensis</i>	native
		<i>Acartia tonsa</i>	native
		<i>Acartia (Acartiura)</i>	native
		<i>Acartia danae</i>	native
		<i>Acartiella sinensis</i>	introduced
		<i>Epilabidocera longipedata</i>	neritic
		<i>Labidocera trispinosa</i>	
		<i>Eurytemora affinis</i>	cryptogenic
		<i>Pseudodiaptomus forbesi</i>	introduced
		<i>Pseudodiaptomus marinus</i>	introduced
		<i>Diaptomus</i> sp.	
		<i>Paracalanus quasimodo</i>	native
		<i>Sinocalanus doerrii</i>	introduced
		<i>Tortanus dextrilobatus</i>	
	<i>Calanus pacificus</i>	introduced	
	Cyclopoids	<i>Oithona davisae</i>	introduced
		<i>Oithona similis</i>	neritic
		<i>Limnoithona tetraspina</i>	introduced
		<i>Acanthocyclops vernalis</i>	
	Siphonostomes	<i>Corycaeus anglicus</i>	neritic
Harpacticoids	<i>Euterpina acutifrons</i>	cryptogenic	
	<i>Coullana canadensis</i>		
Nauplius			
Other Crustaceans	Cladocera	<i>Podon intermedius</i>	
		<i>Evadne</i> sp.	
		<i>Bosmina longirostris</i>	
		<i>Daphnia</i> sp.	
		<i>Ceriodaphnia</i> sp.	
		<i>Diaphanosoma</i> sp.	
	Mysida	<i>Neomysis mercedis</i> <i>Acanthomysis bowmani</i>	introduced
	Crangon sp.		
	Isopoda		
	Amphipoda	<i>Nippoleucon hinumensis</i> <i>Ampelisca abdita</i> <i>Eogammarus confervicolus</i>	
Ostracod			
Meroplankton	Barnacle nauplius		
	Barnacle cyprid		
	Crab zoea		
	Shrimp larvae		
	Snail veliger		
	Bivalve larvae		
	Ctenophore		
	Polychaete larvae		
	Trochophore larvae		
	Echinopluteus larvae		
	Asteroid larvae		
	Medusa		
	Fish eggs		
Fish larvae			
Other taxa	Rotifer		
	Chaetognath	<i>Sagitta</i> sp.	
	Larvacean	<i>Oikopleura dioica</i>	
	Ascidian larva		

Other taxa found in the zooplankton include the larvacean, *Oikopleura dioica*, barnacle nauplii, polychaete worm larvae (Kimmerer 1999), ghost shrimp (*Callinassa californiensis*) larvae, and krill (i.e., *Nematoscelis difficilis*, *Thysanoessa gregaria*, and *Nyctiphanes simplex*) (CDFG 1987).

Ambler *et al.* (1985) describes zooplankton collections in San Francisco Bay from 1978 to 1981. During these years, the most frequently collected zooplankton in the South, Central and San Pablo Bays included the copepods, *Acartia clausi*, *Acartia californiensis*, *Oithona davisae*, harpacticoid copepods, tintinnids, and larvae of gastropods, bivalves, barnacles, and polychaetes. Most frequently collected zooplankton in Suisun Bay were *Eurytemora affinis*, *Sinocalanus doerrii*, cyclopoid copepods, *Bosmina* sp., the cladoceran, *Daphnia pulex*, *Brachionus* sp., and bivalve veligers. Microplankton was also present in all embayments, including the rotifers, *Synchaeta*, and tintinnids *Tintinnopsis* spp. Tintinnids were the most numerous taxa, while copepods and meroplankton dominated by biomass. Mean zooplankton biomass ranged from 10 to 50 mg Cm⁻³, and was highest in southern San Francisco Bay during winter and spring (due to an increase in *A. clausi* and meroplankton). In the northern Bay, biomass peaked during summer and fall, due to an increase of *A. californiensis*, *E. affinis*, *S. doerrii*, and meroplankton. In general, zooplankton biomass was highest where primary productivity was highest, suggesting a strong influence of food availability on zooplankton dynamics. (Ambler *et al.* 1985).

The zooplankton assemblage in San Pablo, Central, and South Bays was described again by Kimmerer (1999), who noted a number of changes from the early 1980s. Overall, there was a decrease in zooplankton abundance in San Pablo and South Bay, particularly during summer months. While the assemblage was still dominated by copepods and meroplankton, dominant species changed drastically, with four new species introduced after the earlier study. The most dominant species in the Kimmerer (1999) study was *Oithona davisae*, which was only moderately abundant in earlier sampling. *Pseudodiaptomus marinus* was introduced into the Bay in the late 1980s (Orsi and Walter 1991), and was the third

most abundant copepod in the lower estuary. *Paracalanus quasimodo* and *Euterpina acutifrons* were quite abundant in higher salinity regions throughout the year, in contrast to earlier sampling. *Tortanus dextrilobatus*, which likely was introduced since the earlier sampling, was most abundant in mid- salinity (15 ppt) areas. Central Bay showed little change between the two collection periods.

Gewant and Bollens (2005) describe the meroplankton (including macrozooplankton and micronekton) of San Francisco Bay based on samples collected from 1997 through 2000. Eighty-two taxa were collected, but 11 taxa accounted for 98 percent of the total abundance. Dominant taxa included four species of fish and seven species of invertebrates: northern anchovy (*E. mordax*), longfin smelt (*Spirinchus thaleichthys*), Pacific herring (*Clupea pallasii*), plainfin midshipman (*Porichthys notatus*), the ctenophore, *Pleurobrachia bachei*, the isopod, *Syndotea laticauda*, the shrimps, *Palaemon macrodactylus*, *Crangon franciscorum*, and *C. nigricauda*, the mysid, *Neomysis kadiakensis*, and the medusa, *Polyorchis* spp. Variation in abundances was driven mostly by distance from the Golden Gate Bridge, seasonality, and life history. Highest abundances occurred in summer and lowest abundances occurred in spring. The authors characterized the meroplankton into four assemblages: taxa spawned from neritic adults that used mostly northern San Francisco Bay (e.g., herring, longfin smelt and plainfin midshipman), estuary-dependent taxa with broad distribution throughout San Francisco Bay year-round (e.g., *C. franciscorum*, *C. nigricauda*, and northern anchovy), resident species that remain in the estuary but occur mostly in the South Bay during the wet season (e.g., *P. macrodactylus*, *S. laticauda*, and *N. kadiakensis*), and gelatinous taxa that occur throughout San Francisco Bay with a single peak in North Bay and South Bay in December and January. Gewant and Bollens (2005).

Further details of abundance and changes since the 1980s are described below by individual taxa.

Copepods

Copepods are small crustaceans evolutionarily derived from oceanic animals. Copepoda is the most well-studied

zooplankton taxon of San Francisco Bay. Despite this, it is difficult to summarize the assemblage because of the number of factors that affect population dynamics (location, season, freshwater inflow, coastal hydrography, and primary productivity) (Ambler *et al.* 1985) and the numerous introductions of non-indigenous copepods since the 1980s (Cohen and Carlton 1995). In fact, *E. affinis*, one of the more abundant copepods in the Bay, is a non-native species from the Atlantic (Lee 2000).

In northern San Francisco Bay (Suisun Bay and northern San Pablo Bay) from 1978 to 1981, copepod dominance followed a pattern with decreasing salinity from east to west: *S. doerrii* was most dominant at the easternmost boundary; *E. affinis* was dominant in the oligohaline mixing zone; *Acartia* spp. were dominant in polyhaline waters; and *Paracalanus parvus* was dominant at the seaward boundary. *S. doerrii*, *E. affinis*, and *A. clausi* were present year-round, while *A. californiensis* was present from August to October. This pattern is absent in southern San Francisco Bay where *Acartia* spp. had increased biomass and showed a distinct seasonal switch from higher biomass of *A. clausi* during the cold, wet season to higher biomass of *A. californiensis* in the warm, dry season. (Ambler *et al.* 1985)

Major changes have occurred in the zooplankton assemblage of San Francisco Bay since 1981. *Sinocalanus doerrii* was first recorded in 1979. Within five years of its introduction, abundance began to decline and has continued to do so through 2005 (Baxter 2006). In 1987-88, the native copepods, *Acartia* spp. and *E. affinis* declined 53 to 91 percent coincidental with the establishment of the exotic clam, *Corbula amurensis* (Kimmerer *et al.* 1994). *E. affinis* abundance has remained at low levels through 2005 (Baxter 2006). In 1989, two calanoid copepods, *Pseudodiaptomus forbesi* and *P. marinus* became established (Kimmerer 2004). Abundance of *P. forbesi* has declined since its introduction, but remains relatively abundant in summer and fall months (Baxter 2006). In 1993, three more exotic copepods, *Limnithona tetraspina*, *Tortanus dextrilobatus* and *Acartiella sinensis*, became established (Orsi and Ohtsuka 1999, Kimmerer 2004, Bouley and Kimmerer 2006). *Limnithona sinensis* is reported to have

disappeared about the time *L. tetraspina* became abundant (Orsi and Ohtsuka 1999). In fact, this species has not disappeared, but is caught at low numbers and is not separated from other *Limnithona* species in collections (A. Hennessy, CDFG, pers. comm.).

Rotifers

Rotifers are microscopic multicellular invertebrates most common in freshwater. Most species are sessile, but about 100 species are planktonic. *Synchaeta*, *Keratella*, and *Brachionus* are the most abundant rotifer genera in San Francisco Bay (Herbold 1992). *Synchaeta* is common at salinities greater than 5 ppt, and is most abundant in the South Bay (Ambler *et al.* 1985). In the Central Bay and upstream, rotifer populations undergo seasonal cycles possibly due to changes in salinity. *Keratella* is usually only found in Suisun Bay and then only in spring when salinities are minimal (Turner and Chadwick 1972). Rotifer abundance in Suisun Bay decreased from 1972 to 1979, apparently associated with declining chlorophyll concentrations (Herbold 1992). *Synchaeta* was the only taxa that did not exhibit this trend, but abundance fell to a record low in 1988 coincident with establishment of *Corbula amurensis* (Herbold 1992). No exotic rotifers have been found in San Francisco Bay (Orsi 1999b).

Cladocera

Cladocera, or water fleas, are very abundant in freshwater systems worldwide, seldom occurring in salinities higher than 1 ppt. They are much more abundant in the Delta than they are in San Francisco Bay, but a few species, including *Bosmina longirostris*, *Daphnia pulex*, and *Diaphanosoma leuchtenbergianum* are found in Suisun Bay and Carquinez Strait during periods of high Delta outflow (Herbold 1992). No exotic cladocera have been found in San Francisco Bay (Orsi 1999b).

Mysid Shrimp

Nine species of mysid shrimp have been reported in San Francisco Bay and the Delta, including six native species and three non-native species. The native species are *Neomysis mercedis*, *N. Kadiakensis*, *N. costata* (now *Holmsimysis costata*), *N. macropsis* (now *Alienacanthomysis*

macropsis), *Deltamysis holmquistae*, and *N. rayi*. *N. rayi* is a coastal species that may only occasionally enter San Francisco Bay. *D. holmquistae* is extremely rare; only a few specimens are collected from the Delta each year. The non-native mysids are *Acanthomysis bowmani* and *A. aspera*, two Chinese species introduced in 1992, and *A. hwanhaiensis*, a Korean species found in 1998. Recent information indicates *Neomysis kadiakensis* collected in and upstream from San Pablo Bay may actually be the introduced *N. japonica*, which would add a tenth mysid species and a fourth introduced species. (Mecum 2006).

Mysid shrimp are most abundant in low salinity (0 to 7.2 ppt) regions, and therefore within the San Francisco Bay are most abundant in Suisun Bay (Heubach 1969, Herbold 1992). Orsi and Knutson (1979) collected six species of mysid shrimp throughout the Delta and Suisun Bay, but only *Neomysis mercedis*, the opossum shrimp, was abundant. *Acanthomysis bowmani* was introduced in 1993 and quickly increased to higher abundances than *Neomysis* (Orsi 1999b). Mysid shrimp have not shown the consistent declines shown by other zooplankton; while there are increasing frequencies of low population levels, the population occasionally rebounds (Herbold 1992).

Large seasonal fluctuations in mysid shrimp abundance from mean winter densities of less than 10 m⁻³ to mean spring densities of almost 1,000 m⁻³ have been attributed to temperature (Heubach 1969, Orsi and Knutson 1979, Siegfried *et al.* 1979), low DO (Turner and Heubach 1966, Orsi and Knutson 1979), predation (Heubach 1969), reproduction (Orsi and Knutson 1979), tidal currents and estuarine circulation (Orsi and Knutson 1979), and seasonal declines in phytoplankton (Orsi and Knutson 1979). Mysid shrimp distribution is also limited to areas with water velocity <0.12 ms⁻¹ and light intensity <10-5 lux (Heubach 1969). (Herbold 1992).

Scyphozoan and Hydrozoan Jellyfish

No hydrozoan jellyfish are known to be native to San Francisco Bay. Three introduced species of hydroid jellyfish, thought to be native to the Black Sea, have been reported since the 1950s. CDFG field notes and USFWS collections indicate *M. marginata* has been present in San

Francisco Bay since at least 1959 (Mills and Rees 2000). *Blackfordia virginica* was collected in the Napa and Petaluma Rivers in 1970 and 1974, respectively. These two species were collected in the Petaluma River and Napa River in 1992, 1993, and 1995 (Mills and Sommer 1995). *Moerisia* sp. was collected in the Petaluma River in 1993 (Mills and Rees 2000). In 1997 and 1998, *M. marginata* and *Moerisia* sp. were collected in Suisun Bay.

All three species are thought to be present in the plankton from at least May through November, with peak abundances from August through October (Rees 1999). Salinity ranges at collection sites ranged from 0 to 20 ppt (Mills and Sommer 1995, Mills and Rees 2000, Rees 1999). Temperature at collection sites ranged from 16.5 to 25 C (Rees 1999, Mills and Rees 2000). Most observations have been made in accessible rivers and sloughs of northern San Francisco Bay. It is unknown if these species are also present over shoals and channels elsewhere in the estuary. Rees and Gershwin (2000) theorize that both species have the ability to invade all low-salinity areas of San Francisco Bay through attachment to the underside of boats.

One scyphozoan jellyfish, *Aurelia* sp., has been observed in San Francisco Bay near Foster City since 1988. Morphology and genetic examinations suggest that the specimens seen in San Francisco Bay are not the native species frequently seen in Monterey Bay and Vancouver Island, but a second, introduced species from Tokyo Bay. (Greenberg *et al.* 1996).

Fishes

Two species of *Salmonidae* occur in fresh-brackish salinity gradients: steelhead (*Oncorhynchus mykiss*), which is listed as threatened pursuant to the Federal Endangered Species Act (ESA), and Chinook salmon (*O. tshawytscha*), of which the winter-run is listed as endangered and the spring-run is listed as threatened pursuant to the ESA. Salmonids are anadromous; adults emigrate from the ocean to spawn in freshwater rivers and streams. Offspring hatch and rear in freshwater and then immigrate to the ocean to forage until maturity.

Abundance of both species has declined in part because of human-induced degradation of their spawning and rearing habitats and loss of smolts in water diversions (Leidy *et al.* 2005). Highly productive estuaries are important feeding and acclimation areas for juvenile salmonids preparing to enter marine waters.

Oligohaline/mesohaline

Common fishes in oligohaline/mesohaline water column areas include the federally threatened delta smelt (*Hypomesus transpacificus*) (58 FR 12854 12864) and striped bass (*Monore saxatilis*). Delta smelt, which are endemic, planktonic feeders in San Francisco Bay and the Delta, prefer habitat within shallow, open water areas in close proximity to transition zones between fresh and oligohaline areas (Moyle 2002). Larvae are most abundant during years of high riverine outflow from April to July, peaking in May (Baxter 1999). Juvenile and adult delta smelt distribution concentrate in Suisun Marsh (Baxter 1999). Striped bass were introduced to San Francisco Bay in 1879 from New Jersey (Moyle 2002). A commercial fishery for striped bass existed in San Francisco Bay from 1889 to 1935 (Goals Project 2000). A recreational fishery still exists today. Striped bass concentrate in San Pablo and Suisun Bay in fall months and migrate to freshwater areas to spawn in winter through early spring (Moyle 2002, Goals Project 2000).

Polyhaline

Pacific herring and larval Pacific staghorn sculpin (*Leptocottus armatus*) are common to polyhaline water column environments (Table 4). The Pacific staghorn sculpin is a common marine and estuarine fish (Miller and Lea 1972). Spawning occurs from October to April. Eggs are demersal and deposited in clusters on a variety of substrates (Wang 1986). Larvae are planktonic and then develop into juveniles at about 10 to 15 mm in length, at which point they settle out and become demersal for the remainder of their life history (Wang 1986).

Eubaline

The Pacific sardine (*Sardinops sagax*), northern anchovy (*Engraulis mordax*), and topsmelt (*Atherinops affinis*) are

among the most abundant species present in euhaline, water column habitat (Table 4). The Pacific sardine is primarily an offshore pelagic species but enters San Francisco Bay in the summer months to feed. A commercial sardine fishery in San Francisco Bay existed from the early 1900s to the 1950s (Skinner 1962) until the fishery collapsed due to over-fishing and changes in off-shore ocean conditions (sardines prefer warmer ocean conditions). Since 1993, there have been seasonal populations of Pacific sardines in South Bay (MSI 2005), and Pacific sardines have been observed year round in Central Bay (Fleming 1999). The northern anchovy is the most common fish in San Francisco Bay and is an important prey species for many other animals (Greiner *et al.* 2005). Northern anchovy abundance in San Francisco Bay was below average from 1999 to 2005 (Greiner *et al.* 2006). San Francisco Bay falls between the northern and central subpopulations of northern anchovy in the Pacific Ocean. The decreased abundance in San Francisco Bay may have been due to a southward migration of the central population in response to a cool ocean regime. Abundance also varies seasonally, with highest abundance in from April to October (Fleming 1999). A small live bait fishery is supported by the northern anchovy. The topsmelt uses San Francisco Bay as a nursery area and spawning ground. Topsmelt spawn in shallow water in April-September and migrate to deeper water areas during the winter (Flemming 1999). Highest abundance of topsmelt occurs in South Bay (Flemming 1999).

Birds

Birds from all San Francisco Bay feeding guilds use the water column habitat for foraging and resting (Table 5). Some of the more abundant species include the western grebe (*Aechmophorus occidentalis* grouped with Clark's grebe, *A. clarkia*, during annual bird counts), the double-crested cormorant (*Phalacrocorax auritus*), the Brandt's cormorant (*Phalacrocorax penicillatus*), and the western gull (*Larus occidentalis*). Species of concern include the osprey (*Pandion haliaetus*), the least tern (*Sterna antillarum*), and the brown pelican (*Pelecanus occidentalis*). Grebes use San Francisco Bay primarily for foraging, peaking in abundance from October to April

(DeSante and Ainley 1980, Briggs *et al.* 1987, Shuford *et al.* 1989). Grebes prefer sheltered coves, rarely found in the open bay except along tidal currents in Raccoon Straits and around Angel Island (Goals Project 2000). Grebes feed on small planktivorous fishes such as sardines and anchovies and small crustaceans (Fix and Bezener 2000).

Double-crested cormorants can be found throughout San Francisco Bay year-round, and use the estuary for both breeding and foraging (Goals Project 2000). As colonial breeders, breeding birds concentrate in San Pablo Bay, Central Bay, and South Bay (Goals Project 2000). Cormorants dive up to depths of 9 m for planktivorous fish prey (Fix and Bezener 2000). The three different species of cormorants that utilize the Bay consume prey with different characteristics. The less common pelagic cormorant (*Phalacrocorax pelagicus*) consumes solitary, cryptic prey on rocky or flat habitats; double-crested cormorants eat mostly schooling fishes that occur above the benthos; and Brandt's cormorants take fish from both the benthic habitat and the water column (Ainley *et al.* 1981). Rocky substrates appeared important for pelagic cormorant prey, while rock, sand, and mud substrates were important habitats for Brandt's cormorant prey (Ainley *et al.* 1981). Brandt's cormorants nesting on Alcatraz Island in San Francisco Bay have been shown to consume northern anchovies as well as benthic fishes such as flatfishes, sculpins, and gobies (Yakich 2005).

The western gull achieves high numbers around San Francisco Bay due to its success at acclimating to urban areas. Western gull feeding preference has shifted from oceanic and intertidal feeding grounds to industrial waterfronts, parks, and garbage dumps where food scraps are readily available (Fix and Bezener 2000). Western gull are year-round residents of San Francisco Bay.

The endangered brown pelican is a seasonal migrant to San Francisco Bay, arriving during the summer months and persisting through late fall (Goals Project 2000). However, during El Niño years, brown pelicans persist year round in San Francisco Bay (Goals Project 2000). Pelicans feed on planktivorous fishes, plunging into the water bill-first. No bay-wide survey exists to count brown pelican abundance, though numbers are estimated to be

several hundred each summer and fall (Goals Project 2000). Osprey, a California state species of special concern, can be viewed year-round in San Francisco Bay, using the area for both rearing and foraging. Osprey are piscivorous, and fish averaging two pounds in weight make up 98 percent of their diet (Fix and Bezener 2000). The endangered least tern (*Sterna antillarum*) migrates to San Francisco Bay during the summer months, using San Francisco Bay as a critical breeding location (Goals Report 2000). Least terns hover over shallow to deep water areas, feeding on planktivorous fishes such as northern anchovy and silversides (*Atherinopsidae*).

Marine Mammals

Marine mammals use water column habitat in San Francisco Bay for migrating, foraging, and resting. Seasonally migrating cetaceans, such as the gray whale (*Eschrichtius robustus*) and humpback whale (*Megaptera novaengliae*), enter Central Bay during their migrations to feed. The harbor porpoise (*Phocoena phocoena*), which preys on pelagic fishes, is commonly sighted in areas of Central Bay and near the Golden Gate Bridge. The harbor seal (*Phoca vitulina*) and California sea lion (*Zalophus californianus*) feed primarily on fishes within San Francisco Bay, including schooling northern anchovy and Pacific herring, but also will feed on migratory Pacific eels, lamprey, salmonids and mysid shrimp and other invertebrates within the water column (Sarah Allen, National Park Service, pers. comm.). Weaned harbor seal pups, in particular, have been documented feeding on mysid shrimp. Both sea lions and harbor seals will sleep on the surface of the water column between foraging bouts. Additionally, harbor seals mate in the water column adjacent to colony haul out sites (Hayes *et al.* 2006, Boness *et al.* 2006).

LITERATURE CITED

- Abbott, I.A. and G.J. Hollenberg. 1976. Marine algae of California. Stanford Univ Press, Stanford, CA
- Accurso, L. M. 1992. Distribution and abundance of wintering waterfowl on San Francisco Bay, 1988-1990. Unpubl. Masters Thesis. Humboldt State University. Arcata, CA. 252 pp.
- Acosta, S. and J. Thayer. 2006. Ecological studies of seabirds on Alcatraz Island, 2006. PRBO Conservation Science Report to the Golden Gate National Recreation Area (GGNRA), National Park Service (NPS).
- Afton, A. D. and M. G. Anderson. 2001. Declining scaup populations: A retrospective analysis of long-term population and harvest survey data. *Journal of Wildlife Management* 65:781-796.
- Ainley, D. G., D. W. Anderson, and P. R. Kelly. 1981. Feeding ecology of marine cormorants in southwestern North America. *Condor* 83: 120-131.
- Allen, S.G. 1991. Harbor seal habitat restoration at Strawberry Spit, San Francisco Bay. Marine Mammal Comm., Contract MM29108990-9. Natl Tech Infor Serv, Alexandria, VA.
- Allen, S.G., M. Stephenson, R. W. Risebrough, L. Fancher, A. Shiller, and D. Smith. 1993. Red-pelaged harbor seals of the San Francisco Bay Region. *J. Mamm.* 74(3):588-593.
- Alpine, A. E., and J. E. Cloern. 1992. Trophic interactions and direct physical effects control phytoplankton biomass and production in an estuary. *Limnology and Oceanography* 37: 946-955.
- Alpine, A.E. and J.E. Cloern. 1988. Phytoplankton growth rate in a light-limited environment, San Francisco Bay. *Mar Ecol Prog Ser* 44:167-173.
- Ambler, J.W., J.E. Cloern and A Hutchinson. 1985. Seasonal cycles of zooplankton from San Francisco Bay. *Hydrobiologia* 129:177-197.
- Anderson, W. 1970. A preliminary study of the relationship of saltponds and wildlife - South San Francisco Bay. *Calif Fish and Game* 56(4): 240-252.
- Anonymous. 1959. Symposium on the classification of brackish waters. Venice 8-14th April 1958. *Archivio di Oceanografia Limnologia* Volume 11. Supplemento (Simposio sulla Classificazione della Acque Salmastre. Venezia 8-14 Aprile, 1958).
- Aplin, J.A. 1967. biological survey of San Francisco Bay, 1963-1966. California Department of Fish and Game Marine Resources Operations. MRO Ref. 67-4.
- Arthur, J.F. and M.D. Ball. 1979. Factors influencing the entrapment of suspended material in the San Francisco Bay estuarine system. Pages 143-174 in: *San Francisco Bay: The Urbanized Estuary*, Conomos, T.J. (ed.) Pacific Division, American Association for the Advancement of Science, San Francisco, CA.
- Atwater, B.F. 1979. Ancient processes at the site of Southern San Francisco Bay: movement of the crust and changes in sea level. Pages 31-45 in: *San Francisco Bay: The Urbanized Estuary*, Conomos, T.J. (ed.) Pacific Division, American Association for the Advancement of Science, San Francisco, CA.
- Austin, J. E., A. D. Afton, M. G. Anderson, R.G. Clark, C. M. Custer, J. S. Lawrence, J. B. Pollard, and J. K. Ringelman. 2000. Declining scaup populations: Issues, hypotheses, and research needs. *Wildlife Society Bulletin* 28:254-263.
- Baker, P. 1995. Review of ecology and fishery of the Olympia oyster, *Ostrea lurida* with annotated bibliography. *Journal of Shellfish Research* 14(2): 501-518.
- Barrett, E.M. 1963. *The California Oyster Industry*. Fish Bulletin 123, California Department of Fish and Game, Sacramento, CA.
- Baxter, R. 2006. Zooplankton monitoring 2005. IEP Newsletter Spring 2006.
- Baxter, R. 1999. Pleuronectiformes. In: Orsi, James J., editor. Report on the 1980-1995 Fish, Shrimp, and Crab Sampling in the San Francisco Estuary, California. http://www.estuaryarchive.org/archive/orsi_1999
- Bay Protection and Toxic Clean-up Program (BPTCP). State Water Resources Control Board. <http://www.swrcb.ca.gov/bptcp/>
- Bonnot, P. 1935. The California oyster industry. *California Fish and Game* 21(1): 65-80.
- Bonnot, P. 1932. Soft shell clam beds in the vicinity of San Francisco Bay. *California Fish and Game* 18: 64-66.
- Bottom, D.L., K.K. Jones, and M.J. Herring. 1984. Fishes of the Columbia River Estuary. Astoria, Oregon, USA Columbia River Estuary Data Development Program.
- Bouley, P. and W.J. Kimmerer. 2006. Ecology of a highly abundant, introduced cyclopoid copepod in a temperate estuary. *Mar Ecol Prog Ser* 324:219-228.
- Briggs, K.T., Tyler, W.B., Lewis, D.B., and Carlson, D.R. 1987. Bird communities at sea off California: 1975-1983. *Studies Avian Biology* No. 11.
- Caddell 1990. Development of California halibut, *Paralichthys californicus*, culture. *J Applied Aquaculture* 14(3/4):143-154. California Department of Fish and Game (CDFG). 2001. California's living marine resources: a status report. University of California, Library of congress control number: 2001098707
- CDFG. 1992. Estuary dependent species. Entered by the CDFG for the State Water Resources Control Board 1992 Water Quality/Water Rights Proceedings on the San Francisco Bay/ Sacramento-San Joaquin Delta. WRINT-DFG Exhibit 6.
- CDFG. 1987. Delta outflow effects on the abundance and distribution of San Francisco Bay fish and invertebrates, 1980-1985. Entered by the CDFG for the State Water Resources Control Board 1987 Water Quality/Water Rights Proceedings on the San Francisco Bay/ Sacramento-San Joaquin Delta. DFG Exhibit 60.

LITERATURE CITED

- Caddell, S.M., D.M. Gadowski, and L.R. Abbot. 1990. Induced spawning of the California halibut, *Paralichthys californicus*, (Pisces: Paralichthyidae) under artificial and natural conditions. Pages 175-198 In C.W. Haugen editor. The California halibut, *Paralichthys californicus*, resource and fisheries. California Department of Fish and Game, Fish Bulletin 174.
- Cappiella, K., C. Malzone, R. Smith and B. Jaffe. 1999. Sedimentation and bathymetric changes in Suisun Bay: 1867-1990. USGS Open File Report 99-563.
- Carlton, J.T. 1979. *History, Biogeography, and Ecology of the Introduced Marine and Estuarine Invertebrates of the Pacific Coast of North America*. Ph.D. dissertation, Univ California, Davis, 904 pp.
- Chavez, F.P., J. Ryan, S.E. Lluch-Cota and C.M. Niquen. 2003. From anchovies to sardines and back: multidecadal change in the Pacific Ocean. *Science* 299: 217-221.
- Cloern, J.E. 1999. The relative importance of light and nutrient limitation of phytoplankton growth: a simple index of coastal ecosystem sensitivity to nutrient enrichment. *Aquat Ecol* 33:3-15.
- Cloern, J.E. 1996. Phytoplankton bloom dynamics in coastal ecosystems: A review with some general lessons from sustained investigation of San Francisco Bay, California. *Reviews of Geophysics* 34(2):127-168.
- Cloern, J.E. and R. Dufford. 2005. Phytoplankton community ecology: principles applied in San Francisco Bay. *Mar Ecol Prog Ser* 285:11-28.
- Cloern, J.E., A.D. Jassby, T.S. Schraga and K.L. Dallas. 2006. The Pulse of the Estuary: Monitoring and Managing Water Quality in the San Francisco Estuary. SFEI Contribution 517. San Francisco Estuary Institute, Oakland, CA.
- Cloern, J.E., B.E. Cole, J.L. Edmunds, T.S. Schraga and A. Arnsberg. 1999. Patterns of water-quality variability in San Francisco Bay during the first six years of the RMP, 1993-1998. Prepared for: Regional Monitoring Program for Trace Substances, San Francisco Estuary Institute.
- Cloern, J.E., B.E. Cole, R.L.J. Wong and A.E. Alpine. 1985. Temporal dynamics of estuarine phytoplankton: a case study of San Francisco Bay. *Hydrobiologia* 129:153-176.
- Cohen, A. N. 2005. *Guide to the Exotic Species of San Francisco Bay*. San Francisco Estuary Institute, Oakland, CA, www.exoticguide.org.
- Cohen, A.N. 2000. *An Introduction to the San Francisco Estuary*. Third Edition. San Francisco Estuary Project, Oakland, CA, Save The Bay, Oakland, CA, and San Francisco Estuary Institute, Oakland, CA.
- Cohen, A.N. and J.T. Carlton. 1998. *Accelerating invasion rate in a highly invaded estuary*. San Francisco Estuary Project, Oakland, CA.
- Cohen, A.N. and J.T. Carlton. 1995. *Nonindigenous Aquatic Species in a United States Estuary: A Case Study of the Biological Invasions of the San Francisco Bay and Delta*. U. S. Fish and Wildlife Service, Washington DC.
- Cohen, A.N. and J.W. Chapman. 2005. *Rapid Assessment Channel Survey for Exotic Species in San Francisco Bay - November 2005*. Final Report for the California State Coastal Conservancy. San Francisco Estuary Institute, Oakland, CA.
- Cohen, A.N., D.R. Calder, J.T. Carlton, J.W. Chapman, L.H. Harris, T. Kitayama, C.C. Lambert, G. Lambert, C. Piotrowski, M. Shouse and L.A. Solórzano. 2005. *Rapid Assessment Shore Survey for Exotic Species in San Francisco Bay - May 2004*. Final Report for the California State Coastal Conservancy, Association of Bay Area Governments/San Francisco Bay-Delta Science Consortium, National Geographic Society and Rose Foundation. San Francisco Estuary Institute, Oakland, CA.
- Cole, B. E., J.E. Cloern and A.E. Alpine. 1986. Biomass and productivity of three phytoplankton size classes in San Francisco Bay. *Estuaries* 9: 117-126.
- Conomos, T.J. 1979. Properties and circulation of San Francisco Bay waters. Pages 47-84 in: *San Francisco Bay: The Urbanized Estuary*, Conomos, T.J. (ed.) Pacific Division, American Association for the Advancement of Science, San Francisco, CA.
- Conomos, T.J., R.E. Smith and J.W. Gartner. 1985. Environmental setting of San Francisco Bay. In: *Temporal Dynamics of an Estuary: San Francisco Bay*, Cloern, J.E. and F.H. Nichols (eds.). *Hydrobiologia* 129: 1-12.
- Dahlstrom, W. 1977. Unpublished study for California Department of Fish and Game, Marine Resources Operations, Menlo Park, CA, cited by Jones & Stokes 1977.
- Davis, J.A., A.J. Gunther, B.J. Richardson, J.M. O'Connor, R.B. Spies, E. Wyatt, E. Larson and E.C. Meiorin. 1991. *Status and Trends Report on Pollutants in the San Francisco Estuary*. San Francisco Estuary Project, Oakland, CA.
- DeLeón, S. 1999. Embiotocidae. In: Orsi, James J., editor. Report on the 1980-1995 Fish, Shrimp, and Crab Sampling in the San Francisco Estuary, California. http://www.estuaryarchive.org/archive/orsi_1999.
- Den Hartog, C. 1997. Is *Sargassum muticum* a threat to eelgrass beds? *Aquatic Botany* 58: 37-41.
- DeSante, D.F. and D.G. Ainley. 1980. *The Avifauna of the South Farallon Islands, California*. *Studies in Avian Biology* 4.
- Dillion, C.R., 1971 A comparative study of the primary productivity of estuarine phytoplankton and macrobenthic plants. Ph.D. Thesis. University. North Carolina, Chapel Hill 119 pp.
- Elliott, M. L., R. Hurt, and W. J. Sydeman. 2006. Breeding biology and status of the California Least Tern *Sterna antillarum browni* at Alameda Point, San Francisco Bay, California. Submitted for publication.
- Engle, J.M. 1979. Ecology and growth of juvenile California spiny lobster, *Panulirus interruptus* (Randall). Ph.D. Dissertation, University of Southern California.
- Environmental Monitoring and Assessment Program (EMAP). U.S. Environmental Protection Agency (EPA). <http://www.epa.gov/emfjulte/index.html>.
- ENTRIX, Inc. 1998. Biological assessment for the Berths 55-58 and Oakland Harbor Navigation Improvement (-50') Projects. Report for Port of Oakland. Project No. 377301.
- Euing, S. 2006. Breeding Status of the California Least Tern at Alameda Point, Alameda, California, 2006. Unpublished Report Prepared for the U.S. Navy, U.S. Fish and Wildlife Service, Fremont, California.

- Feminella, J.W and V.H. Resh. 1989. Submersed macrophytes and grazing crayfish: an experimental study of herbivory in a California freshwater marsh. *Holarctic Ecology* 12: 1-8.
- Fitch, J.E. and R.J. Lavenberg. 1971. Marine food and game fishes of California. University of California Press, Berkeley, California.
- Fix, D. and A. Bezener. 2000. Birds of Northern California. Lone Pine Publishing: Auburn, Washington. 384 pp.
- Fleming, K. 1999. Northern Anchovy. *In*: Orsi, James J., editor. Report on the 1980-1995 Fish, Shrimp, and Crab Sampling in the San Francisco Estuary, California. http://www.estuaryarchive.org/archive/orsi_1999
- Flick, R.E., J.F. Murray and L.C. Ewing. 2003. Trends in United States Tidal Datum Statistics and Tide Range, *J. Waterway, Port, Coastal and Ocean Eng.*, Amer. Soc. Civil Eng. 129(4):155-164.
- Foster, M.S., A.P. DeVogelaere, C. Harrold, J.S. Pearse and A.B. Thum. 1988. Causes of spatial and temporal patterns in rocky intertidal communities of Central and Northern California. *Memoirs of the California Academy of Sciences* Number 9, San Francisco, California
- Gadomski, D.M. and S.M. Caddell. 1991. Effects of temperature on early-life-history stages of California halibut *Paralichthys californicus*. *Fish Bull* 89:567-576.
- Garcia & Associates. 2001. Final Survey Report: San Francisco rock removal benthic survey off Alcatraz Island, CA. *for* San Francisco District, U.S. Army Corps of Engineers. DACW07-01-002, Task Order No. 0001.
- Geller, J.B., J.T. Carlton and D.A. Powers. 1994. PCR-based detection of mtDNA haplotypes of native and invading mussels on the northeastern Pacific coast: latitudinal pattern of invasion. *Mar Biol* 119: 243-249.
- Geller, J.B., J.T. Carlton and D.A. Powers. 1993. Interspecific and intrapopulation variation in mitochondrial ribosomal DNA sequences of *Mytilus* spp. (Bivalvia: Mollusca). *Molecular Mar Biol Biotechnol* 2(1): 44-50.
- Gewant, D.S. and S.M. Bollens. 2005. Macrozooplankton and micronekton of the lower San Francisco Estuary: seasonal, interannual, and regional variation in relation to environmental conditions. *Estuaries* 28(3):473-485.
- Goals Project. 1999. Baylands Ecosystem Habitat Goals. A Report by the San Francisco Bay Area Wetlands Ecosystem Goals Project. First Reprint. U.S. Environmental Protection Agency, San Francisco, Calif./S.F. Bay Regional Water Quality Control Board, Oakland, Calif.
- Goals Project. 2000. Baylands ecosystems species and community profiles: life histories and environmental requirements of key plants, fish and wildlife. Prepared by the San Francisco Bay Area Wetlands Ecosystem Goals Project. P.R. Olofson, editor. San Francisco Bay Regional Water Quality Control Board, Oakland, California.
- Goodwin, P. and R.A. Denton. 1991. Seasonal influences on the sediment transport characteristics of the Sacramento River. *In*: Proceedings of the Institute of Civil Engineers, part 2, volume 91, p. 163-172.
- Green, D.E., E. Grigg, S. Allen, and H. Markowitz. 2006. Monitoring the potential impact of the seismic retrofit construction activities at the Richmond-San Rafael Bridge on harbor seals (*Phoca vitulina richardsi*).
- Greene, H.G. and J.J. Bizarro. 2003. Benthic habitat maps of San Francisco Bay interpreted from multibeam bathymetric images and side-scan sonar mosaics. Prepared for NOAA National Ocean Service. <http://response.restoration.noaa.gov/cpr/cpr.html>.
- Greenberg, N., R.L. Garthwaite and D.C. Potts. 1996. Allozyme and morphological evidence for a newly introduced species of *Aurelia* in San Francisco Bay, California. *Mar Bio* 125(2):401-410.
- Greiner, T., K. Hieb, S. Slater, and M. Sandford. 2006. 2005 Fishes annual status and trends report for the San Francisco Estuary. IEP Newsletter, Spring 2006.
- Grigg, E.K., S.G. Allen, D.E. Green, and H. Markowitz. 2004. Harbor seal, *Phoca vitulina richardsi*, population trends in the San Francisco Bay Estuary, 1970-2002. *California Fish and Game* 90(2):51-70.
- Gunther, A.J. 1987. *Segmentation of the San Francisco Bay-Delta*. Aquatic Habitat Institute, Richmond, CA.
- Haderlie, E.C. and D.P. Abbott. 1980. Bivalvia: the clams and allies. Pages 355-411 in: *Intertidal Invertebrates of California*, R.H. Morris, D.P. Abbott and E.C. Haderlie (eds.), Stanford University Press, Stanford, CA.
- Haertel, L. and C. Osterberg. 1967. Ecology of zooplankton, benthos and fishes in the Columbia River Estuary. *Ecology* 48:459-472.
- Hayden, H.S., J. Blomster, C. Maggs, P. Silva, M. Stanhope, and R. Waaland. 2003. Linnaeus was right all along: *Ulva* and *Enteromorpha* are not distinct genera. *European J Phyc* 38:277-294.
- Hayes, S.A., D.P. Costa, J.T. Harvey, and B.J. le Boeuf. (2004). Aquatic mating strategies of the male Pacific Harbor Seal (*Phoca vitulina richardsi*): Are males defending the hotspot? *Marine Mammal Science* 20(3):639-656.
- Herbold, B., A.D. Jassby and P.B. Moyle. 1992. *Status and Trends report on Pollutants in the San Francisco Estuary*. San Francisco Estuary Project, Oakland, CA.
- Hertlein, L.G. 1959. Notes on California oysters. *Veliger* 2(1): 5-10.
- Heubach, W. 1969. *Neomysis awatschensis* in the Sacramento-San Joaquin River Estuary. *Limnol Ocean* 14(4):533-546.
- Hieb, K. 2006. 2005 annual status and trends report – Common crabs of the San Francisco Estuary. IEP Newsletter, Spring 2006.
- Hieb, K. 2005. Common crabs of the San Francisco Estuary. IEP Newsletter, Spring 2005.
- Hieb, K. 1999a. *Cancer Crabs*. *In*: Orsi, James J., editor. Report on the 1980-1995 Fish, Shrimp, and Crab Sampling in the San Francisco Estuary, California. http://estuaryarchive.org/archive/orsi_1999.
- Hieb, K. 1999b. *Caridean Shrimp*. *In*: Orsi, James J., editor. Report on the 1980-1995 Fish, Shrimp, and Crab Sampling in the San Francisco Estuary, California. http://estuaryarchive.org/archive/orsi_1999.
- Hieb, K. and R. Baxter. 1993. Delta outflow/San Francisco Bay. Pages 101-116 In P.L. Hergesell, editor. 1991 Annual Report-Interagency Ecological Studies Program for the Sacramento-San Joaquin Estuary.

LITERATURE CITED

- Hieb, K., M. Bryant, M. Dege, T. Greiner, K. Souza, and S. Slater. 2005. Fishes in the San Francisco Estuary, 2004 status and trends. IEP Newsletter, Spring 2005.
- Hopkins, D. R. 1986. Atlas of the Distributions and Abundances of Common Benthic Species in San Francisco Bay, California. Water Resources Investigations Report 86-4003, U. S. Geological Survey.
- Hurley, L.M. 1990. Field guide to the submerged aquatic vegetation of Chesapeake Bay. Annapolis (MD): US Fish and Wildlife Service, Chesapeake Bay Estuary Program. 51 p.
- Hurt, R. 2006. Breeding status of the California Least Tern at Alameda Point, Alameda, California, 2005. Unpublished report prepared for the U.S. Navy, U.S. Fish and Wildlife Service, Fremont, California.
- Hymanson, Z., D. Mayer and J. Steinbeck. 1994. Long-term trends in benthos abundance and persistence in the upper Sacramento-San Joaquin Estuary: Summary report 1980-1990. Report No. 38, Interagency Ecological Program for the San Francisco Bay/Delta Estuary.
- Intergovernmental Panel on Climate Change. 2001. Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881 pp.
- Jaffe, B.E., R.E. Smith and L.Z. Torresan. 1998. Sedimentation and bathymetric change in San Pablo Bay: 1856-1983. USGS Open File Report 98-759.
- Jassby, A.D. 1992. Organic carbon sources for the food web of San Francisco Bay. Appendix A in: *Status and Trends Report on Pollutants in the San Francisco Estuary*, Herbold, B., A.D. Jassby and P.B. Moyle, San Francisco Estuary Project, Oakland, CA.
- Jassby, A.D., J.E. Cloern and B.E.Cole. 2002. Annual primary production: patterns and mechanisms of change in a nutrient-rich tidal ecosystem. *Limnol and Ocean* 47(3): 698-712.
- Jassby, A. D., W. J. Kimmerer, S. G. Monismith, C. Armor, J. E. Cloern, T. M. Powell, J. R. Schubel, and T. J. Vendlinski. 1995. Isohaline position as a habitat indicator for estuarine populations. *Ecol App* 5:272-289.
- Jassby, A. D., J. E. Cloern and T. M. Powell. 1993. Organic carbon sources and sinks in San Francisco Bay: variability induced by river flow. *Mar Ecol Prog Ser* 95:39-54.
- Jones & Stokes Associates. 1977. *San Francisco Bay Shellfish: An Assessment of the Potential for Commercial & Recreational Harvesting*. Association of Bay Area Governments, Berkeley, CA.
- Josselyn, M.N. and A.C. Mathieson. 1980. Seasonal influx and decomposition of autochthonous macrophyte litter in a north temperate estuary. *Hydrobiologica* 71:197-208.
- Josselyn, M. and J. West, 1985. The distribution and temporal dynamics of the estuarine macroalgal community of San Francisco Bay. *Hydrobiologica* 129:139-152
- Kantrud, HA. 1990. *Sago pondweed (Potamogeton pectinatus): A literature review*. U.S. Fish and Wildlife Service, Fish and Wildlife Resource Publication 176. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. <http://www.chesapeake.org/SAV/literature/Ruppialitrev.html>
- Kendall Jr., A.W. and W.H. Lenarz. 1986. Status of early life history studies of northeast Pacific rockfishes; October 1986; Anchorage, Alaska. pp 99-128.
- Kimmerer, W.J. 2004. Open-Water Processes of the San Francisco Estuary: from physical forcing to biological responses. *San Francisco Estuary and Watershed Science* 2(1): Article 1.
- Kimmerer, W. J. 2002. Effects of freshwater flow on abundance of estuarine organisms: Physical effects or trophic linkages? *Mar Ecol Prog Series* 243:39-55.
- Kimmerer, W. 1999. Zooplankton in the lower San Francisco Estuary. IEP Newsletter Spring 1999.
- Kimmerer, W.J., E. Gartside and J.J. Orsi. 1994. Predation by an introduced clam as the likely cause of substantial declines in zooplankton of San Francisco Bay. *Mar Ecol Prog Ser* 113:81-93.
- Kincaid, T.B. 1949. Japanese mollusks in seed oyster boxes from Japan shipped to Willapa Bay. *Min Conchol Club So Calif* 88:1.
- Kitting, C.K. 1998. Bathymetry and Underwater Vegetation Mapping in Horseshoe Bay, Golden Gate National Recreation Area. Report to Golden Gate National Parks Ass. 13 pp. plus figures.
- Kopec, D. and J. Harvey. 1995. Toxic pollutants, health indices, and population dynamics of harbor seals in San Francisco Bay, 1989-91: a final report. Technical publication. Moss Landing, CA: Moss Landing Marine Labs.
- Kozloff, E.N. 1983. Seashore life of the Northern Pacific Coast, an illustrated guide to Northern California, Oregon, Washington, and British Columbia. University of Washington Press; Seattle, Washington. 370 pp.
- Krone, R.B. 1979. Sedimentation in the San Francisco Bay system. In: Conomos, T.J. (Ed.) *San Francisco Bay: The urbanized estuary*. Pacific Division of the American Association for the Advancement of Science, San Francisco. Pp. 85-96.
- Largier, J.L. 1996. Hydrodynamic exchange between San Francisco Bay and the ocean: the role of ocean circulation and stratification. Pages 69-104 in: *San Francisco Bay: The Ecosystem*, Hollibaugh, J.T. (ed.) Pacific Division, American Association for the Advancement of Science, San Francisco, CA.
- Lee, C.E. 2000. Global phylogeography of a cryptic copepod species complex and reproductive isolation between genetically proximate "populations." *Evol* 54(6):2014-2027.
- Lee, H., II, B. Thompson, and S. Lowe. 2003. Estuarine and scalar patterns of invasion in the soft-bottom benthic communities of the San Francisco Estuary. *Biological Invasions* 5:1-2.
- Leidy, R.A., S.B. Becker and B.N. Harvey. 2005. Historical distribution and current status of Steelhead/rainbow trout (*Oncorhynchus mykiss*) in streams of the San Francisco Estuary, California. Lowry, M.S. and K.A. Forney. 2005. Abundance of California sea lions (*Zalophus californianus*) in central and northern California during 1998 and summer 1999. *Fish Bull* 103(2):331-343.
- Litler, M.M. and D. Murray. 1975. Impact of sewage on the distribution, abundance and community structure of rocky intertidal macro-organisms. *J Mar Biol* 30: 277-291.
- Lucas, L.V., J.E. Cloern, J.R. Koseff, S.G. Monismith, and J.K. Thompson. 1998. Does the Sverdrup critical depth model

- explain bloom dynamics in estuaries?: *Journal of Marine Research* 56 (2):375-415.
- Lucas, L.V., J.R. Koseff, J.E. Cloern, S.G. Monismith and J.K. Thompson. 1999a. Processes governing phytoplankton blooms in estuaries, I, the local production-loss balance: *Marine Ecology Progress Series* 187:1-15.
- Lucas, L.V., J.R. Koseff, S.G. Monismith, J.E. Cloern and J.K. Thompson. 1999b. Processes governing phytoplankton blooms in estuaries, II, the role of horizontal transport: *Marine Ecology Progress Series* 187:17-30.
- MACTEC. 2006. Marin Rod and Gun Club Native Olympia Oyster Habitat Restoration Project (2005- 2006). A final report submitted to the NOAA Fisheries Restoration Center
- Mallin, M.A., J.M. Burkholder, L.B. Cahoon and M.H. Posey. 2000. The North and South Carolina coasts. *Mar Poll Bull* 41:56-75.
- Manna, J., D. Roberts, D. Press and S. Allen. 2006. Harbor seal monitoring, San Francisco Bay Area. Annual Report for the National Park Service. 21 pp.
- Marine Science Institute (MSI). 2005. Trends in south San Francisco Bay fish populations from 1972-2002. <http://www.cmi-registration.com/user/org/category.jsp?id=6538&org=261>
- Markmann, C. 1986. Benthic monitoring in the Sacramento-San Joaquin Delta: Results from 1975 through 1981. Interagency Ecological Studies Program for the Sacramento-San Joaquin Estuary, Tech Rep 12 (WQ/BIO-rATR/87-12), CA Dept Water Resources, Sacramento, 51 pp.
- Martin, A.C., H.S. Zim and A.L. Nelson. 1951. *American wildlife and plants*. New York: McGraw-Hill. 500 pp.
- May, C.L., J.R. Koseff, L.V. Lucas, J.E. Cloern, and D.H. Schoellhamer. 2003. Effects of spatial and temporal variability of turbidity on phytoplankton blooms. *Mar Ecol Prog Ser* 254:111-128.
- McDonald, J.H. and R.K. Koehn. 1988. The mussels *Mytilus galloprovincialis* and *M. trossulus* on the Pacific coast of North America. *Mar Bio* 99: 111-118.
- McGowan, M.F. 2005. Tiburon Audubon Center native oyster (*Ostrea conchaphila*) restoration project monitoring report (2004-2005) prepared for the NOAA Fisheries Restoration Center.
- Mecum, W.L. 2006. A key to the Mysidacea of the upper San Francisco Estuary. California Department of Fish and Game. November 20, 2006. 13 pp.
- Merkel & Associates, Inc. 2005. Eelgrass community pilot study for San Francisco Bay: techniques for examining invertebrate and fish assemblages within multiple eelgrass beds. Report for :San Francisco-Oakland Bay Bridge East Span Seismic Safety Project.
- Merkel & Associates, Inc. 2004. Baywide eelgrass inventory of San Francisco Bay. Report for San Francisco-Oakland Bay Bridge East Span Seismic Safety Project.
- Merkel & Associates, Inc. 2003. San Francisco Bay eelgrass inventory. A report for the Bay Bridge eelgrass mitigation project. Prepared for Caltrans and NOAA Fisheries.
- Merkel & Associates, Inc. 1999a. Middle harbor habitat design: Governing design and engineering criteria for target habitat elements. Prepared for Winzler and Kelly, San Francisco, CA and the U.S. Army Corps of Engineers, San Francisco District. 19 pp.
- Merkel & Associates, Inc. 1999b. Richmond Harbor Navigation Improvement Project. Post-dredging eelgrass survey. Prepared for Tetra Tech, Inc. San Francisco, CA. 22 pp.
- Messersmith, J. 1966. Fishes collected in the Carquinez Strait in 1961-1962. Pages 57-63 In D.W. Kelley, editor. *Ecological Studies of the Sacramento-San Joaquin Estuary: Part II. Fishes of the Delta*. California Department of Fish and Game, Fish Bull 136.
- Miller, D.J. and R.N. Lea. 1972. Guide to the coastal marine fishes of California. California Department of Fish and Game, Fish Bull 157.
- Mills, C.E. and J.T. Rees. 2000. New observations and corrections concerning the trio of invasive hydromedusae *Maeotias marginata* (= *M. inexpectata*), *Blackfordia virginica*, and *Moerisia* sp. in the San Francisco Estuary. *Mar Sci* 64 (Supl 1):151-155.
- Mills, C.E. and F. Sommer. 1995. Invertebrate introductions in marine habitats: two species of hydromedusae (Cnidaria) native to the Black Sea, *Maeotias inexpectata* and *Blackfordia virginica*, invade San Francisco Bay. *Mar Bio* 122: 279-288.
- Monroe, M.W. and J. Kelly. 1992. *State of the Estuary: A Report on Conditions and Problems in the San Francisco Bay/Sacramento-San Joaquin Delta Estuary*. Association of Bay Area Governments, Oakland, CA.
- Moyle, P. B. 2002. Inland fishes of California. University of California Press; Berkeley, CA. 501 pp.
- Nichols, F. H. and M. M. Pamatmat. 1988. The ecology of the soft-bottom benthos of SF Bay: a community profile. U.S. Fish and Wildlife Biological Report 85(7.19): 73 pp.
- Nichols, F.H. and J.K. Thompson. 1985. Time scales of change in the San Francisco Bay benthos. *Hydrobiologia* 129:121-138.
- Nichols, F. H., J. K. Thompson, and L. E. Schemel. 1990. Remarkable invasion of SF Bay (California, USA) by the Asian clam *Potamocorbula amurensis*. 2. Displacement of a former community. *Marine Ecol Prog Ser*. Oldendorf 66:95-101.
- Nicholson, N., H. Hosmer, K. Bird, L. Hart, W. Sandlin, C. Shoemaker and C. Sloan. 1981. The biology of *Sargassum muticum* (Yendo) Fensholt at Santa Catalina Island, California. Proceedings of the 8th International Seaweed Symposium, pp. 416-424.
- O'Farrell, M. and R.J. Larson. 2005. Year-class formation in Pacific herring (*Clupea pallasii*) estimated from spawning-date distributions of juveniles in San Francisco Bay, California. *Fish Bulletin* 103:130-141. California Department of Fish and Game.
- Oltmann, R.N. 1996. Sediment inflow to the Delta from the Sacramento and San Joaquin rivers. *IEP Newsletter* 9(2):22-26.
- Oltmann, R.N., D.H. Schoellhamer and R.L. Dinehart. 1999. Sediment inflow to the Sacramento-San Joaquin Delta and the San Francisco Bay. *IEP Newsletter* 12(1):30-33.
- Orcutt, H.G. 1950. The life history of the starry flounder, *Platichthys stellatus* (Pallas). California Department of Fish and Game, Fish Bull 78.
- Orsi, J (ed.). 1999. Report on the 1980-1995 fish, shrimp, and crab sampling in the San Francisco Estuary, California. California Department of Fish and Game Technical Report 63.
- Orsi, J. 1999. Long-term trends in mysid shrimp and zooplankton. *IEP Newsletter* Spring 1999.

LITERATURE CITED

- Orsi, J.J. and A.C. Knutson, Jr. 1979. The role of mysid shrimp in the Sacramento-San Joaquin Estuary and factors affecting their abundance and distribution. Pages 401-408 *In*: T.J. Conomos (ed.) San Francisco Bay: The Urbanized Estuary.
- Orsi, J. and S. Ohtsuka. 1999. Introduction of the Asian copepods *Acartiella sinensis*, *Tortanus dextrilobatus* (Copepoda: Calanoida), and *Limnoithona tetraspina* (Copepoda: Cyclopoida) to the San Francisco Estuary, California, USA. *Plankton Bio and Ecol* 46(2):128-131.
- Orsi, J. J. and T. C. Walter. 1991. *Pseudodiaptomus forbesi* and *P. marinus* (Copepoda: Calanoida), the latest copepod immigrants to California's Sacramento-San-Joaquin estuary. *In*: S. -i. Uye, S. Nishida, and J. -S. Ho, eds., Proceedings of the Fourth International Conference on Copepoda. Pp. 553-562. Bull Plank Soc Japan, special volume: i-xi, 1-645.
- Packard, E.L. 1918. Molluscan fauna from San Francisco Bay. *Univ Calif Pub Zoo* 14(2): 199-452.
- Pearson, D. E. and S. L. Owen. 2001. *In* California's Living Marine Resources: a Status Report. W. S. Leet, C. M. Dewees, R. Klingbeil, and E. J. Larson, eds. University of California Press, Berkeley, CA: 384-385.
- Peterson, H.A. 2002. Long-term benthic community change in a highly invaded estuary. Master's Thesis, Marine Biology, San Francisco State University. San Francisco, CA. 108 pp
- Phillips, R. C. 1979. Ecological notes on. Phyllospadix (Potamogetonaceae) in the Northeast. *Pacific. Aquatic Bot* 6:1-170.
- Raimondi, P.T., R.F. Ambrose, J.M. Engle, S.N. Murray and M. Wilson. 1999. Monitoring of rocky intertidal resources along the central and southern California mainland. 3-Year Report for San Luis Obispo, Santa Barbara, and Orange Counties (Fall 1995-Spring 1998). OCS Study, MMS 99-0032, U.S. Minerals Management Service, Pacific OCS Region
- Rees, J. 1999. Non-indigenous jellyfish in the upper San Francisco Estuary: potential impacts on zooplankton and fish. IEP Newsletter Summer 1999.
- Rees, J.T. and L. Gershwin. 2000. Non-indigenous hydromedusa in California's upper San Francisco Estuary: life cycles, distribution, and potential environmental impacts. *Sci Mar* 64 (supl 1):73-86.
- Regional Monitoring Program (RMP). San Francisco Estuary Institute. <http://www.sfei.org/rmp>.
- Ricketts, E.F., J. Calvin, J., and J.W. Hedgpeth. *Between Pacific Tides*, 5th edition. Stanford University Press; Stanford, California. 652 pp.
- Ruhl, C.A., D.H. Schoellhamer, R.P. Stumpf and C.L. Lindsay. 2001. Combined use of remote sensing and continuous monitoring to analyze the variability of suspended-sediment concentrations in San Francisco Bay, California. *Estuarine, Coastal, and Shelf Sci* 53:801-812.
- San Francisco Estuary Project (SFEP). 1997. *State of the Estuary 1992-1997*. San Francisco Estuary Project, Oakland, CA.
- San Francisco Bay Regional Water Quality Control Board. Water Quality Control Plan (Basin Plan). <http://www.swrcb.ca.gov/rwqcb2/basinplan.htm>
- Sarver, S.K. and D.W. Foltz. 1993. Genetic population structure of a species' complex of blue mussels (*Mytilus* spp.). *Mar. Biol.* 117: 105-112.
- Schoellhamer, D.H. 2002. Variability of suspended-sediment concentration at tidal to annual time scales in San Francisco Bay. *Cont Shelf Research* 22:1857-1866.
- Schoellhamer, D.H. 2001. Influence of salinity, bottom topography, and tides on estuarine turbidity maxima in northern San Francisco Bay. *In*: McAnally WH, Mehta AJ, editors. Coastal and estuarine fine sediment processes. Amsterdam: Elsevier. P 343-356.
- Schoellhamer, D.H. 1996. Factors affecting suspended-solids concentrations in south San Francisco Bay, California. *J Geophysical Research* 101:12087-12095.
- Schoellhamer, D.H. and J.R. Burau. 1998. Summary of findings about circulation and the estuarine turbidity maximum in Suisun Bay, California, U.S. Geological Survey Fact Sheet FS-047-98, 6 p. Sea Duck Joint Venture Management Board. 2001. Sea Duck Joint Venture Strategic Plan: 2001 – 2006. SDJV Continental Technical Team. Unpubl. Rept. [c/o USFWS, Anchorage, Alaska; CWS, Sackville, New Brunswick]. 14pp. + appendices.
- Setchell, W.A. 1929. Morphological and phonological notes on *Zostera marina* L. *Univ Calif Publ Bot* 14: 389-452.
- Shuford, W.D., G.W. Page, J.G. Evans and L.E. Stenzel. 1989. Seasonal abundance of waterbirds at Point Reyes: a coastal California perspective. *Western Birds* 20:137-265.
- Silva, P.C. 1979. The benthic algal flora of central San Francisco Bay. *In* T.J. Conomos (ed.), San Francisco Bay: the urbanized estuary. Pacific Div. Am. Ass. Adv. Sci, San Francisco, California.: 287-346.
- Skinner, J.E. 1962. *An Historical Review of the Fish and Wildlife Resources of the San Francisco Bay Area*. Water Projects Branch Report No. 7, California Department of Fish and Game, Sacramento, CA.
- Smith, L.W. 1987. *A Review of Circulation and Mixing Studies of San Francisco Bay, California*. USGS Circular 1015.
- Smith, S.E. and N.J. Abramson. 1990. Leopard shark (*Triakis semifasciata*) distribution, mortality rate, yield, and stock replenishment estimates based on a tagging study in San Francisco Bay. *Fish Bull* 88:371-381.
- Sobczak, W.V., J.E. Cloern, A.D. Jassby, B.E. Cole, T.S. Schraga, and A. Arnsberg. 2005. Detritus fuels ecosystem metabolism but not metazoan food webs in San Francisco Estuary's freshwater delta. *Estuaries* 28(1):122-135.
- Sobczak, W.V., J.E. Cloern, A.D. Jassby and A.B. Muller-Solger. 2002. Bioavailability of organic matter in a highly disturbed estuary: The role of detrital and algal resources. *PNAS* 99(12):8101-8105.
- Stearns, R.E.C. 1899. *Modiola plicatula* Lamarck, in San Francisco Bay. *Nautilus* 13:86.
- Stevens, D. E. and L. W. Miller. 1983. Effects of river flow on abundance of young Chinook salmon, American shad, longfin smelt, and delta smelt in the Sacramento-San Joaquin River System. *N Amer J Fish Manage* 3(4):425-437.
- Sutton, J.E. 1981. *Shellfish Resources of Eastern San Francisco Bay: Distribution, Abundance, Public Access and Use*. East Bay Municipal Utility District, Oakland CA.

- Sutton, J.E. 1978. *Survey of Sport Shellfishing Potential in San Francisco Bay, in Southern San Francisco and Northern San Mateo Counties*. San Francisco Wastewater Program, San Francisco, CA.
- Takekawa, J.Y., T.L. Corinna and R.T. Pratt. 2001. Avian communities in baylands and artificial salt evaporation ponds of the San Francisco Bay estuary. *Hydrobiologica* 466: 317-328.
- Tasto, R.N. 1983. Juvenile Dungeness crab, *Cancer magister*, studies in the San Francisco Bay area. Pages 135-154 in *Life history, environment, and mariculture studies of the Dungeness crab, Cancer magister*, with emphasis on the central California fishery resource, Wild, P.W. and R.N. (eds). Calif Dep Fish Game Fish Bull 172 pp.
- Thayer, G.W., W.J. Kenworthy, and M.S. Fonseca. 1984. The Ecology of Eelgrass Meadows of the Atlantic Coast: A Community Profile. U.S. Fish and Wildlife Service. FWS/OBS-84/02. 147p.
- The Bay Institute. 2003. San Francisco Bay Fish Index Indicator Analysis and Evaluation. The Bay Institute Ecological Scorecard. October 17, 2003. <http://www.bay.org/Scorecard/Fish.pdf>
- Thompson, J.K. 2005. One estuary, one invasion, two responses: phytoplankton and benthic community dynamics determine the effects of an estuarine invasive suspension feeder. Pages 221-316 in *The comparative roles of suspension feeders in ecosystems*, S. Olenin and R. Dame, Eds.
- Thompson, J.K. 1999. The effect of infaunal bivalve grazing on phytoplankton bloom development in south San Francisco Bay. Stanford University, Stanford, Calif., Ph.D. dissertation, 419 p.
- Thompson, B., R. Hoenicke, J. A. Davis, and A. Gunther. 2000. An overview of contaminant-related issues identified by monitoring in SF Bay. *Environmental Monitoring and Assessment*. 64:409-419.
- Trost, RE. 2002. Pacific Flyway 2001-2002 Fall and Winter Waterfowl Survey Report.
- Turner, T. 1983. Complexity of early and middle successional stages in a rocky intertidal surfgrass community. *Oecologia* 60:56-65.
- Turner, T. 1985. Stability of rocky intertidal surfgrass beds: persistence, preemption, and recovery. *Ecology* 66:83-92.
- Turner, J.L., and H.K. Chadwick. 1972. Distribution and abundance of young-of-the-year striped bass, *Morone saxatilis*, in relation to river flow in the Sacramento – San Joaquin estuary. *Trans Am Fish Soc* 101: 442-452.
- Turner, J.L. and W. Heubach. 1966. Distribution and concentration of *Neomysis awatschensis* in the Sacramento-San Joaquin Delta, p. 105-112, In D. W. Kelley, (ed), *Ecological studies of the Sacramento-San Joaquin Estuary*. Calif Fish and Game, Fish Bull 133:1-133.
- U.S. Environmental Protection Agency. 1972. Shellfish Study of San Francisco Bay. Technical Report EPA-909/9-74-003.
- U.S. Fish and Wildlife Service (USFWS). 2006. Waterfowl population status, 2006. U.S. Department of the Interior, Washington, D.C. U.S.A.
- U.S. Geological Survey (USGS). 1998. Classification of Wetlands and Deepwater Habitats of the United States Unconsolidated Bottom.
- Valiela, I, J. McClelland, J. Hauxwell, P. Behr, D. Hersh and K. Forman. 1997. Macroalgal blooms in shallow estuaries: Controls and ecophysical and ecosystem consequences. *Limnology and Oceanography*, 42:1105-1118.
- Vanderhoof, M. and S.G. Allen. 2005. Harbor Seal monitoring at Pt. Reyes National Seashore and Golden Gate National Recreational Area. Annual Report.
- Wang, J.C.S. 1986. Fishes of the Sacramento-San Joaquin Estuary and adjacent waters, California: A Guide to the early life histories. Interagency Ecological Study Program for the Sacramento-San Joaquin Estuary, Technical Report 9, Sacramento, California.
- Watters, D.L.H.M. Brown, F.J. Griffin, E.J. Larson and G.N. Cherr. 2004. Pacific herring *Clupea pallasii* spawning grounds in San Francisco Bay: 1973 to 2000. Pages 3-14 in *Early life history of fishes in the San Francisco Estuary and watershed*, F. Feyrer, L.R. Brown, R.L. Brown, and J.J. Orsi, eds. American Fisheries Society Symposium 39, Bethesda, MD.
- Wilkerson, F.P., R.C. Dugdale, V.E. Hogue and A. Marchi. 2006. Phytoplankton blooms and nitrogen productivity in San Francisco Bay. *Estuaries and Coasts* 29(3): 401-416.
- Wilkinson. M. 1980. Estuarine benthic algae and their environment: a review. Pages 425-486 in *The Shore Environment*, J. H. Price, D. E. G. Irvine and W. F. Farnham (eds).. Ecosystems. Academic Press, London, Vol. 2.
- Wooster, T.W. 1968. Clams, native oysters and mussels. In *Fish and Wildlife Resources of San Francisco Bay and the Delta: Description, Environmental Requirements, Problems, Opportunities and the Future*. California Department of Fish and Game, Sacramento, CA.
- Wright, S.A. and D.H. Schoellhamer. 2003. Trends in sediment yield of the Sacramento River, 1957-2001. Proceedings of 2003 CALFED Science Conference, Sacramento, California, January 14-16, 2003. p. 177.
- Wyllie-Echeverria, S. 1990. Geographic range and distribution of *Zostera marina*, eelgrass in San Francisco Bay. In: K. Merkel and R. Hoffman (eds.), Proceedings of the California Eelgrass Symposium, pp. 65-69. Sweetwater River Press, National City, CA.
- Wyllie-Echeverria, S. and P. J Ruttan. 1989. Inventory of eelgrass (*Zostera marina* L.) in San Francisco/San Pablo Bay. Southwest Region, NOAA Administrative Report SWR-89-05. 18 pp.
- Yakich, J. D. 2005. A dietary analysis of Brandt's cormorants (*Phalacrocorax penicillatus*) breeding in central San Francisco Bay. M.S. Thesis, San Francisco State University.

Index

- Acanthomysis*, 77, 81
Acartia, 77, 79, 80
Acipenser, 41, 45
Aeolidia, 68
Abnfeltiopsis, 49
Alcidae, 51
Alexandrium, 71, 73
Algae, 39, 60
Ameana, 27, 44
Americorophium, 19, 20, 22, 40, 41
Ampelisca, 21, 23, 24, 25, 26, 27, 28, 29, 42, 43, 44, 77
Amphibalanus, 31, 33, 34, 35, 36, 45, 49, 54
amphipods, 42, 43, 44, 47, 49, 50, 54, 61, 62, 68, 69
Ampithoe, 33, 37, 49, 54
amurensis, 19, 20, 21, 22, 23, 24, 25, 31, 39, 40, 41, 42, 45, 46, 74, 80, 81
Anas, 45, 63
anchovy, 39, 68, 79, 83, 84
anemone, 31, 32, 33, 34, 36, 45, 49, 50, 70
anemones, 55, 68
Anisodris, 70
Anthopleura, 70
ascidean, 44
ascidians, 49
Atherinops, 39, 68, 83
Atherinopsidae, 84
Aurelia, 82
Aythya, 43, 45, 47, 63
- Balanus*, 31, 34, 36, 41, 45, 50, 53, 54, 55, 57, 61
barnacle, 31, 33, 34, 35, 36, 41, 49, 50, 53, 79
bass, 41, 82
bat ray, 39, 55, 68
bivalve, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 31, 34, 36, 37, 39, 40, 41, 42, 43, 44, 52, 79
Bosmina, 77, 79, 81
Brachionus, 79, 81
Bryopsis, 16, 18
bryozoan, 31, 48, 49, 50
bryozoans, 33, 48, 50, 54, 68
- Bucephala*, 45, 56
- C. amurensis*, 40, 41, 42, 43, 46
C. fluminea, 40, 41
Callianassa, 79
Cancer, 39, 40, 45, 50, 68, 70
canvasback, 43, 45, 47, 63
Caprella, 34, 36, 37, 57, 68
Cebidichthys, 70
Cepphus, 43, 46, 51
Ceratium, 71
Chondracanthus, 16, 18
Chthamalus, 61
Cirolana, 53, 54
Cirriformia, 23, 43
Citharichthys, 40, 46, 68
cladocerans, 75
Cladophora, 16, 18
clams, 16, 45, 47, 52, 55
Clanthria, 49
Clathra, 50
Clupea, 40, 51, 68, 79
Codium, 18, 49, 66
Conopeium, 49
copepod, 74, 75, 79, 80
copepods, 75, 76, 79, 80
Corbula, 19, 20, 21, 22, 23, 24, 25, 31, 39, 40, 42, 45, 46, 74, 80, 81
Corophium, 23, 25, 29, 37, 42, 43, 57
Cottus, 42, 51
crab, 31, 39, 40, 42, 41, 43, 45, 50, 62, 68, 70
Crangon, 39, 40, 41, 42, 43, 45, 52, 77, 80
Crassostrea, 4, 52, 55
Cryptopleura, 16, 18
ctenophore, 79
Cucumaria, 53
cumacean, 20, 21, 22, 23, 25, 26, 28, 41, 42, 43
Cymatogaster, 40, 56, 68

- Daphnia*, 77, 79, 81
Diadumene, 31, 32, 33, 34, 45, 49, 50, 55, 57
Diaphanosoma, 77, 81
Dinophysis, 71
 duck, 43, 45, 56

E. affinis, 79, 80
Ectopleura, 34, 36, 50
 eelgrass, 18, 49, 39, 55, 61, 62, 63, 65, 68, 69
Egretta, 18, 49
 elasmobranchs, 46
Embiotocidae, 51, 68
Engraulis, 39, 68, 83
Enteromorpha, 16, 18, 39
Eogammarus, 33, 49, 77
Eschrichtius, 49, 51, 47, 84
Euchone, 26, 27, 28, 29, 44
Eurytemora, 77, 79
Exogone, 24, 27, 28, 30, 44

Flabellinopsis, 70
 flounder, 41, 45, 46
Fucus, 16, 18, 60, 61

Gammarus, 19, 20, 22, 40, 41, 60, 61
Garveia, 33, 34, 49
Gelidium, 16, 18
Geukensia, 52, 54
Glycinder, 28, 30, 44
Gnorimosphaeroma, 31, 33, 45, 49
 goldeneye, 45, 56
Gonyaulax, 53
Gracilaria, 16, 18, 60, 61, 65
Gracilariophila, 61
Grandidierella, 21, 24, 27, 28, 43, 44
 gray whale, 49, 50, 84
 grebe, 45, 46, 56, 83
 gull, 43, 47, 48, 51, 83, 84

 halibut, 40, 46, 47, 69
Halichondria, 33, 35, 37, 50, 54, 57
Haliclona, 33, 35, 49, 50, 57
Halkymenia, 16, 18
Harmothoe, 29, 37, 44, 57
Hermisenda, 70

 herring, 40, 47, 49, 50, 51, 52, 61, 65, 68, 69, 79, 83, 84
Heteromastus, 20, 21, 22, 24, 25, 26, 29, 41, 44
Heteropodarke, 27, 44
Heterosigma, 71
Hiarella, 36, 50
Hopkinsia, 49
 hydroid, 54, 82
 hydromedusae, 76
 hydrozoan, 31, 49, 82
Hypomesus, 41, 82
Hypsopsetta, 56

Idotea, 53, 60, 61
 Ilyanassa, 26, 43
Incisocalliope, 34, 36, 50
 isopod, 21, 31, 33, 34, 35, 41, 45, 54, 79

Jassa, 34, 36, 50
 jellyfish, 82

 kelp, 62, 70
Keratella, 81
Kornmannia, 69
 krill, 79

L. hoffmeisteri, 41
Lacuna, 69
Laminaria, 18, 49
Larus, 43, 47, 48, 51, 83
 larvacean, 79
Leucilla, 35, 50
Limnodilus, 40
Limnoithona, 77, 80
 limpet, 69

M. viridis, 41, 42
Macoma, 23, 24, 25, 27, 42, 43, 52
 Macroalgae, 60
 Macrocystis, 62
macrodactylus, 20, 41, 42, 57, 80
 mallards, 63
Marenzelleria, 19, 20, 21, 22, 24, 40, 42
Mebile, 68
Mediomastus, 27, 28, 29, 30, 44
 medusa, 80

INDEX

- Megaptera*, 51, 84
Melanitta, 43, 46
Melobesia, 69
meroplankton, 75, 79
Mesodinium, 71, 73, 74
Metridium, 36, 50
midshipman, 39, 49, 46, 79
Modiolus, 36, 50
Moerisia, 82
Molgula, 23, 26, 31, 34, 44, 45, 54, 58
monkey-faced eel, 70
Monocophium, 44
Monocorophium, 21, 22, 24, 25, 26, 27, 28, 33, 34, 36, 42, 44
Morone, 41
Musculista, 23, 25, 29, 31, 34, 36, 37, 42, 43, 45, 49, 55, 58
mussel, 32, 33, 35, 37, 45, 49, 52, 53, 54, 58
Mustelus, 39, 46
Mya, 23, 24, 25, 27, 42, 43, 44, 52
Mycale, 35, 50
Myliobatis, 39, 55, 68
mysid, 49, 47, 76, 80, 81, 84
Mytilus, 32, 33, 35, 37, 45, 49, 52, 53, 55, 58
- Nannochloropsis*, 71, 73
Nassarius, 68
Neanthes, 24, 25, 35, 43, 57
Nematoscelis, 79
Neomysis, 77, 80, 81
Neotrypaea, 52
Nereis, 37, 53, 57
Nippoleucon, 20, 21, 22, 23, 25, 26, 28, 41, 42, 43, 77
Notoacmea, 69, 70
nudibranch, 43, 68
Nyctiphanes, 79
- Obelia*, 31, 54, 57
Oikopleura, 78, 79
Oithona, 77, 79
oligochaete, 19, 22, 23, 25, 27, 28, 29, 43
Oncorhynchus, 42, 82
opisthobranch, 32, 33, 49, 74
osprey, 47, 83
Ostrea, 4, 16, 35, 49, 52, 55, 58
- Oxyura*, 45, 56, 63
- Palaemon*, 20, 41, 42, 80
Paracalanus, 77, 79, 80
Paralichthys, 40, 46
Paranthura, 34, 50, 57
Parophrys, 46
Pelecanus, 43, 46, 83
pelican, 43, 46, 83, 84
perch, 40, 49, 51, 52, 56, 68, 69
Phalacrocorax, 43, 46, 51, 83
Phocoena, 49, 51, 47, 84
Phyllaplysia, 68
Phyllospadix, 18, 49, 62, 69, 70
phytoplankton, 9, 46, 60, 71, 73, 74, 75, 81
pigeon guillemot, 43, 46, 51
pintail, 45
pipefish, 39, 56, 68
Pisaster, 50, 53
Plagioselmis, 71, 73
Platichthys, 41, 45
Pleurobrachia, 79
Plyrhinoidis, 56
Podiceps, 45, 46, 56
Pogonichthys, 45
Pollicipes, 53
polychaetes, 40, 41, 43, 44, 50, 79
Polydora, 24, 28, 44
Polyorchis, 80
Polysiphonia, 16, 18
Porichthys, 39, 46, 79
porpoise, 49, 50, 51, 47, 52, 84
Potamogeton, 18, 62, 63
Prorocentrum, 71, 73
Pseudodiaptomus, 77, 79, 80
Psuedopolycora, 44
Psuedopolydora, 28, 29, 44
Pugettia, 70
Pyramimonas, 71, 73
- Rhithropanopeus*, 20, 41, 57
rockfish, 40, 49, 46, 47, 52, 68
rotifer, 81
Ruppia, 18, 62, 63

- S. doerrii*, 79, 80
Sabaco, 24, 25, 26, 27, 29, 43, 44
 sago pondweed, 62, 63
Sakuraeolis, 26, 43, 58
 salmon, 42, 49, 65, 82
 sanddab, 40, 46, 68
 sardine, 39, 83
Sardinops, 39, 83
Sargassum, 18, 49, 60, 61, 62
 SAV, 16, 39, 60, 62, 65, 66, 68
 scaup, 43, 45, 47, 63
 sculpin, 40, 42, 49, 46, 51, 83
 Scytosiphon, 62
 sea cucumbers, 53
 sea lion, 49, 50, 51, 52, 53, 69, 84
 sea star, 48, 53
 seal, 49, 51, 52, 53, 70, 84
 seals, 49, 50, 47, 52, 53, 69, 84
Sebastes, 40, 46, 51, 68
Semibalanus, 53
 shark, 39, 46, 55
 shrimp, 39, 40, 41, 49, 41, 42, 43, 44, 47, 52, 57, 68, 76,
 79, 81, 84
 silversides, 84
Sinelobus, 54
Sinocalanus, 77, 79, 80
Skeletonema, 71, 73
 smelt, 41, 49, 45, 46, 69, 79, 82
Smithora, 69
 smoothound, 46
 snail, 32, 43, 69
 sole, 40, 46
Sphaeroma, 31, 33, 49, 60, 61
Spirinchus, 41, 45, 79
 splittail, 41, 45
 sponges, 49, 50, 54, 55, 69
 steelhead, 42, 82
Streblospio, 24, 25, 28, 29, 42, 44
 sturgeon, 41, 45
Styela, 35, 54, 58
 surf scoter, 43, 46, 47
 surfgrass, 62, 69, 70
 surfperch,, 51, 68
Sycon, 35, 50
Synchaeta, 79, 81
Syndotea, 79
Syngnathus, 39, 56, 68
Synidotea, 21, 33, 35, 41, 49, 57


 tanaids, 54
Teleaulax, 71, 73
 tern, 43, 47, 68, 83, 84
Tetraclita, 53
Thalassiosira, 71, 73
Tharyx, 27, 44
Theora, 23, 43
Thysanoessa, 79
Tomentosoides, 49
 topsmelt, 39, 68, 83
Tortanus, 77, 79, 80
Triakis, 39, 46, 55
Tubificidae, 22, 25, 27, 28, 29, 44
Tubificoides, 23, 43
 turbot, 56

Ulva, 16, 18, 39, 60, 66
Upogebia, 52

Varichaetadrilus, 19, 40
Venerupis, 23, 26, 31, 42, 43, 45, 52, 58
Victorella, 36, 50

 whale, 49, 50, 51, 47, 69, 84

Zalophus, 49, 51, 47, 53, 84
 zooplankton, 9, 46, 74, 75, 76, 79, 80, 81
Zostera, 18, 39, 61, 62, 65



August 2007

National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Santa Rosa Office

777 Sonoma Avenue
Santa Rosa, CA 95404

An online version of this publication is available at:
www.noaa.gov/

AR 0025882