

AN ANNOTATED CHECKLIST OF PHILIPPINE FLATFISHES: ECOLOGICAL IMPLICATIONS¹

Annadel Cabanban

*IUCN Commission on Ecosystem Management, Southeast Asia
Dumaguete, Philippines; Email: annadel_cabanban@yahoo.com.sg*

Emily Capuli

*SeaLifeBase Project, Aquatic Biodiversity Informatics Office
Khush Hall, IRRI, Los Baños, Laguna, Philippines; Email: e.capuli@cgiar.org*

Rainer Froese

*IFM-GEOMAR, University of Kiel
Duesternbrooker Weg 20, 24105 Kiel, Germany; Email: rfroese@ifm-geomar.de*

Daniel Pauly

*The Sea Around Us Project, Fisheries Centre, University of British Columbia,
2202 Main Mall, Vancouver, British Columbia, Canada, V6T 1Z4; Email: d.pauly@fisheries.ubc.ca*

ABSTRACT

An annotated list of the flatfishes of the Philippines was assembled, covering 108 species (vs. 74 in the entire North Atlantic), and thus highlighting this country's feature of being at the center of the world's marine biodiversity. More than 80 recent references relating to Philippine flatfish are assembled. Various biological inferences are drawn from the small sizes typical of Philippine (and tropical) flatfish, and pertinent to the "systems dynamics of flatfish". This was facilitated by FishBase, which documents all data presented here, and which was used to generate the graphs supporting these biological inferences.

INTRODUCTION

Taxonomy, in its widest sense, is at the root of every scientific discipline, which must first define the objects it studies. Then, the attributes of these objects can be used for various classificatory and/or interpretive schemes; for example, the table of elements in chemistry or evolutionary trees in biology. Fisheries science is no different; here the object of study is a fishery, the interaction between species and certain gears, deployed at certain times in certain places. This interaction determines some of the characteristics of the resource (e.g., recruitment to the exploited stock), and generates catches.

For conventional fisheries research to work, however, the underlying taxonomy must have been done: the species caught must be known, and catch statistics must be available, at least at species level. Without these, state-of-the art methods of fisheries research cannot be used, and emphasis must then be given, to various indirect methods and to inferences by analogy. This, indeed, is the reason for the renaissance of comparative methods in fishery research (Bakun, 1985).

Flatfish (Order Pleuronectiformes) support substantial single-species fisheries in the North Atlantic and North Pacific, besides forming a sizeable by-catch in various medium-latitude trawl fisheries. On the other hand, the many species of flatfish occurring in the inter-tropical belt do not support directed fishery, nor

¹ Cite as: Cabanban, A., Capuli, E., Froese, R., Pauly, D. 2010. An annotated checklist of Philippine flatfishes: ecological implications. In: Palomares, M.L.D., Pauly, D. (eds.), *Marine Biodiversity of Southeast Asian and Adjacent Seas*. Fisheries Centre Research Reports 18(3), p15-31. Fisheries Centre, University of British Columbia [ISSN 1198-6727].

do they contribute much to the by-catch of the multispecies (trawl) fisheries common in tropical shelves (Pauly, 1994). Thus, studying the fishery biology of tropical flatfish cannot proceed as does the study of flatfish resource species in temperate waters, and comparative approaches must make use of the facts that are known about the distribution and occurrence, the morphology and other features of the fish under study, in an attempt to compensate as far as possible for the unavailability of abundance data and of catch time series.

Relational databases are ideal for assembling, recombining and analyzing such facts, and this report relied heavily on the FishBase 96 CD-ROM (Froese and Pauly, 1996), and subsequent updates² which anticipates the release of FishBase 97. The usefulness of FishBase for the comparative study of flatfish in general (and by extension, of any other fish group) was highlighted in Froese and Pauly (1994). Hence this contribution focuses on the narrower issue of its use for generating inferences on the ecology of flatfishes (here taken as representing any other group of teleosts) in data-sparse, but species-rich tropical areas, here represented by the waters within the Philippine EEZ.

MATERIALS AND METHODS

The first task was to complete the FishBase coverage of Philippine flatfish; this was achieved by (1) scanning the Philippines (Evermann and Scale, 1907; Fowler, 1934; Herre, 1953) and regional taxonomic literature (e.g., Weber and de Beaufort, 1929; Menon and Monkolprasit, 1974; Amaoka and Hensley, 2001; Hensley and Amaoka, 2001), and (2) interacting with taxonomists, notably at the FAO/ICL ARM/MS I workshop held on 1 - 10 October 1995 for the production of an FAO Identification Guide to Living Marine Resources of the Western Central Pacific, and at the Smithsonian Institution, Washington, D.C. The pleuronectids in Herre (1953) were checked against Menon's (1977) revision of the Cynoglossidae and revisions of *Engyprosopon* (Amaoka et al., 1993) and *Paraplagiisia* (Chapleau and Renaud, 1993), while Eschmeyer (1990) was consulted for the validity of the generic names. Distribution records were taken from Herre (1953), from revisions, redescriptions (e.g., *Pseudorhombus megalops*; Hensley and Amaoka 1989), museum records, and the general scientific literature on Philippine demersal fish and fisheries.

Biological and ecological information on Philippine flatfish were gleaned mainly from the *Philippine Journal of Fisheries*, the *Philippine Journal of Science*, and the *Philippine Scientist*. Also, various bibliographies were examined for entries on flatfish (Blanco and Montalban, 1951; Gomez, 1980; Aprieto et al., 1986; Pauly et al., 1986), complemented by a search of the Aquatic Sciences and Fisheries Abstracts CD-ROM, and of the personal reprint collections of colleagues both at ICLARM³, Manila and the Smithsonian Institution, Washington, D.C.

The second task was to create, for each species of flatfish reported from the Philippines at least one georeferenced occurrence record with sampling depth and environmental temperature. The plot of




Figure 1. Relationship between mean annual sea temperature (in °C) and depth (in m) for various locations in the Philippines. Source: Dalzell and Ganaden (1987) based on Selga (1931) and Labao (1980).

² The original version of this, now slightly updated, paper was presented at the Symposium on System Dynamics of Flatfish, held 2-8 November 1996 at the Netherlands Institute for Sea Research, Texel, The Netherlands and was previously available from <http://filaman.uni-kiel.de/geomar/rfroese/Philippines%20Flatfish.pdf>. The coverage of flatfishes by FishBase now includes the data therein and additional information.

³ Now the WorldFish Center, Penang, Malaysia.

temperature *vs.* depth in Figure 1 was used to infer temperature from position and depth in cases where the temperatures had been missing from an original record. Our major source of occurrence records was a printout from the Smithsonian Institution listing all Philippine flatfish in their collection (courtesy of Dr. Leslie W. Knapp), the results of the MUSORSTOM Expedition to the Philippines (Fourmanoir, 1976 in Fourmanoir, 1981), and the definitions of the type locality for the species described (mainly by Fowler, 1934).

Biological characteristics (catch data and derived features do not exist for Philippine flatfish) were entered into the appropriate fields of FishBase, which also documents their sources. Also, the FishBase coverage of non-Philippine flatfish was boosted such as to provide sufficient contrast to Philippine species. The various graphing and reporting routines of FishBase were then evoked, and used to generate the exhibits presented below.

RESULTS AND DISCUSSION

There are at least 108 species of flatfish in the Philippines, distributed in 8 families and 36 genera (Appendix 1). The type locality of 22 nominal flatfish species is in the Philippines (W.N. Eschmeyer, pers. comm.). As predicted by Pauly (1994) for tropical species in general, Philippine flatfish tend to remain small, ranging from 6 to 80 cm in standard length (SL) with most species reaching 15 cm (SL) or less.

During the October 1995 FAO-ICLARM workshop for the testing of the FAO Western Central Pacific Field Guide, the fish markets of Cebu, Manila, and Bolinao were sampled by groups of taxonomists, and specimens were bought for identification and collection purposes. The relatively few flatfish found by that survey consisted of 19 flatfish species with an average maximum size of about 21 cm SL (Table 1), thus confirming the low abundance, high diversity, small size, and low economic importance of Philippine flatfish.

Figure 2 compares the maximum size distribution of Philippine flatfish with that of North Atlantic species (FAO areas 21 and 27). Two ecological implications of this are that Philippine flatfish are limited to smaller prey than their North Atlantic counterpart, while simultaneously being susceptible to (numerous) smaller predators. The implications of reduced size and increased temperature for population dynamics are faster turnover rates, i.e., the asymptotic size is approached rapidly due to high values of the parameter *K* of the von Bertalanffy growth function (Pauly, 1980, 2010). This leads to reduced longevity (Figure 3), and high natural mortality (Figure 4).

Table 1. List of flatfishes surveyed during the October 1995 FAO-ICLARM workshop.

Family	Species	Length (cm)
Bothidae	<i>Arnoglossus aspidos</i>	–
	<i>Arnoglossus taenio</i>	–
	<i>Bothus pantherinus</i>	15.4 SL
	<i>Chascanopsetta micrognathus</i>	–
	<i>Engyprosopon grandisquama</i>	–
Citharidae	<i>Citharoides macrolepidotus</i>	–
Cynoglossidae	<i>Cynoglossus cynoglossus</i>	10.4 SL
	<i>Cynoglossus kopsii</i>	–
	<i>Pseudorhombus arsius</i>	24.5 SL
	<i>Pseudorhombus duplucicellatus</i>	29.0 SL
Psettodidae	<i>Psettodes erumei</i>	25.5 SL
	<i>Psettodes</i> sp.	–
Soleidae	<i>Aseraggodes</i> sp.	–
	<i>Dexillichthys muelleri</i>	21.0 SL
	<i>Euryglossa</i> sp.	23.8 TL
	<i>Pardachirus pavoninus</i>	13.2 SL
	<i>Synaptura orientalis</i>	–
	<i>Synaptura sorsogonensis</i>	20.5 SL




Figure 2. Frequency distribution of maximum reported lengths in Philippine and North Atlantic flatfish, highlighting small sizes of Philippine species (data from FishBase, August 1996).




Figure 3. Longevity is in most organisms related to size, and neither the fish nor the Pleuronectiformes are an exception (data from FishBase, August 1996).




Figure 5. Within groups of similar fishes (here in the Pleuronectiformes), the maximum size reached by different species decreases with environmental temperature, although this effect is not seen when data for all orders of fish are pooled.





Figure 4. In Pleuronectiformes, as in other fishes, natural mortality (M) is strongly related to the parameters of the von Bertalanffy growth equation, K and L_{∞} . The plot in the right panel also shows the effect of temperature.

The maximum size that can be reached by fish of various taxa is largely independent of temperature; there are small and large fish at almost all temperatures. However, within groups, the size reduction of maximum size imposed by environmental temperature (for which Pauly, 1994 suggests a mechanism) does show, and this is confirmed by Figure 5 for the Pleuronectiformes.

Tropical demersal environments are usually characterized by high fish diversity (Aprieto and Viloso, 1979; Gloerfelt-Tarp and Kailola, 1984; Sainsbury *et al.*, 1985; Dredge, 1989a, 1989b; Kulbicki and Wantiez, 1990; Cabanban, 1991). Several surveys of demersal fishes were conducted in the Philippines (Warfel and Manacop, 1950; Ronquillo *et al.*, 1960; Viloso and Hermosa, 1982) which provided checklists of fishes and their relative abundances (Aprieto and Viloso, 1979; Viloso and Aprieto, 1983). Furthermore, catch rate data are available for several decades but have tended to remain underutilized (Silvestre *et al.* 1986b). These data allow rough assessments of the status of the demersal stocks (Silvestre *et al.* 1986a, 1986b) and inference on growth, mortality and recruitment patterns based on analysis of lengthfrequency data (Ingles and Pauly, 1984), though inferences on Pleuronectiformes are few, due to their scarcity.

The flatfish of the Philippines are diverse but compose a small percentage of the total catch of demersal fisheries. To date, there is a lack of scientific investigation on the systematics, biology, population ecology, and fisheries of Philippine flatfish. The high diversity and low abundance of flatfish in the tropics [e.g., Sunda Shelf (see contributions in Pauly and Martosubroto, 1996); North Western Australia (Sainsbury *et al.*, 1985); northern part of Australia (Rainer and Munro, 1982; Rainer, 1984); Cleveland Bay, Australia (Cabanban, 1991)] has been highlighted by Pauly (1994), who argued that the low biomass and recruitment rates of flatfish in the tropics are primarily based on environmental physiology (temperature-mediated difference of metabolic rate) and diet. He also suggests that flatfish are overadapted to feeding on zoobenthic epi- and infauna such that low availability of food limits the production of biomass and recruitment.

Flatfish are considered 'trashfish' (Saila, 1983; Dredge, 1989a, 1989b) in most warm water developed countries, e.g., in Australia (Rainer, 1984), but enter markets in the Philippines, often as dried packs of juveniles of various species used for snacks. As for the adults, their small sizes reduce their value, substantially, except for *Psettodes erumei*, a high "quality" fish (Aprieto and Villosa, 1979). Flatfish in Southeast Asia generally feed on benthic invertebrates (Chan and Liew, 1986). In turn, these fish form part of the prey items of medium-sized (*Saurida* spp.; Cabanban, 1991) and large-sized carnivores. As such they may form a significant link in those demersal ecosystems where terrigenous input of nutrients leads to high benthos biomasses (Belperio, 1983).

We conclude by pointing out that there is a need to revise the systematics of the Philippine Pleuronectiformes, many species of which have not been reported since they were originally described. Also there is a need to study their spatial and temporal distribution and abundances in various habitats. Furthermore, studies on the diet, growth, reproduction, and recruitment of these fish are required if understanding of their population dynamics is to improve. Except for taxonomic studies, dedicated work on flatfish may not be of high priority in the Philippines. However, it is hoped that Philippine Pleuronectiformes will be studied further, at least in the context of their relationships in multispecies assemblages.

REFERENCES

- Amaoka K., Hensley, D.A., 2001. Paralichthyidae. Sand flounders. In: Carpenter, K.E., Niem, V. (eds.), *The Living Marine Resources of the Western Central Pacific. Volume 6: Bony Fishes Part 4 (Labridae to Latimeriidae), Estuarine Crocodiles, Sea Turtles, Sea Snakes and Marine Mammals*, p. 3842-3862. FAO identification guide for fishery purposes. FAO, Rome.
- Amaoka, K., Yamamoto, E., 1984. Review of the genus *Chascanopsetta*, with the description of a new species. *Bulletin of the Faculty of Fisheries, Hokkaido University* 35(4), 201-224.
- Amaoka, K., Mihara, E., Rivaton, J., 1993. Pisces, Pleuronectiformes: Flatfishes from the waters around New Caledonia. A revision of the genus *Engyproson*. In: Crosnier, A. (ed.), *Resultats des Campagnes MUSORSTOM* 11(158), p. 377-426. Memoire du Muséum national d'Histoire naturelle, Paris.
- Anon., 1994. Printout of all Philippine flatfish in the collection of the Smithsonian, courtesy of Dr. Leslie Knapp (personal communication).
- Aprieto, V.L., Villosa, E.P., 1979. Catch composition and relative abundance of trawl-caught fishes in the Visayan Sea. *Fisheries Research Journal of the Philippines* 4(1), 9-18.
- Aprieto, V., Saeger, J., Pauly, D., Editors, 1986. Selected Papers on Philippine Marine Fisheries Resources (1947-1986). Department of Marine Fisheries, University of the Philippines, Visayas, Tech. Rep. No. 9, 435 p.
- Bakun, A., 1985. Comparative studies and the recruitment problem: searching for generalization. *CalCOFI Report* 26, 30-40.
- Bawazeer, A.S., 1987. Stock assessment of the large toothed flounder (khoffah, *Pseudorhombus arsius*) in Kuwait waters. *Kuwait Bulletin of Marine Science* 9, 207-214.
- Belperio, A.P., 1983. Terrigenous sedimentation in the central Great Barrier Reef Lagoon: a model from the Burdekin region. *BMR Journal of Australian Geology and Geophysics* 8, 179-190.
- Blaber, S.J.M., 1980. Fish of the Trinity Inlet System of North Queensland with notes on the ecology of fish faunas of tropical Indo-Pacific estuaries. *Australian Journal of Marine and Freshwater Research* 31, 137-46.
- Blanco, G.J., Montalban, H.R., 1951. A bibliography of Philippine fishes and fisheries. *Philippine Journal of Fisheries* 1(2), 107-130.
- Cabanban, A.S., 1991. The dynamics of the Leiognathidae in a tropical demersal ichthyofaunal community. James Cook University of North Queensland, Australia, 262 p. Ph. D. dissertation.
- Chan, E.H., Liew, H.C., 1986. A study on tropical demersal species (Malaysia). International Development Research Centre, IDRC/3-A-83-1905, Singapore, 64 p.
- Chapleau, F., Renaud, C.B., 1993. *Paraplagusia sinerama* (Pleuronectiformes: Cynoglossidae), a new Indo-Pacific tongue sole with a revised key to species of the genus. *Copeia* 3, 798-807.
- Conlu, P.V., 1979. *Guide to Philippine Flora and Fauna*. Volume III, Fishes. Natural Science Research Centre, Quezon City.
- Dalzell, P., Ganaden, R., 1987. A review of the fisheries for small pelagic fishes in Philippine waters. Bureau of Fisheries and Aquatic Resources Technical Paper Seriea 10(1), 58 p.
- Devadoss, P., Pillai, P.K.M., Natarajan, P., Muniyandi, K., 1977. Observations on some aspects of the biology and fishery of *Psettodes erumei* (Bloch) at Porto Novo. *Indian Journal of Fisheries* 24(1/2), 62-68.
- Dou, S., 1992. Feeding habit and seasonal variation of food constituents of left-eyed flounder, *Paralichthys olivaceus*, of the Bohai Sea. *Marine Science* 4(4), 277-281.
- Dredge, M.C.L., 1989a. By-catch from the Central Queensland prawn fisheries. The prawn fisheries, species composition, site associations from the by-catch. Fisheries Research Branch, Queensland Primary Industries. Technical Report FRB 88/04.

- Dredge, M.C.L., 1989b. By-catch from the Central Queensland prawn fisheries. Part 2. Spatial and temporal changes in by-catch composition and community assemblages. Fisheries Branch, Queensland Department of Primary Industries, Brisbane. 37 p.
- Edwards, R.R.C., Shaher, S., 1991. The biometrics of marine fishes from the Gulf of Aden. *Fishbyte* 9(2), 27-29.
- Erzini, K., 1991. A compilation of data on variability in length-age in marine fishes. Working Paper 77. Fisheries Stock Assessment, Title XII, Collaborative Research Support Program, University of Rhode Island.
- Eschmeyer, W.N., 1990. *Catalog of the Genera of Recent Fishes*. California Academy of Sciences, U.S.A. 697 p. 10
- Evermann, B.W., Scale, A., 1907. Fishes of the Philippine Islands. Bulletin of the United States Bureau of Fisheries 26(1906), 49-110.
- Fischer, W., Whitehead, P.J.P., Editors, 1974. FAO species identification sheets: eastern Indian Ocean (Fishing Area 57), Western Central Pacific (Fishing Area 71). Volumes 1- 4. FAO, Rome, pag. var.
- Fourmanoir, P., 1981. Poissons (première liste). In: Forest, J. (ed.), *Results of the MUSORSTOM Expeditions to the Philippine Islands (18-28 mars 1976)*, p. 85-102. Éditions de l'Office de la Recherche Scientifique et Technique Outre-Mer avec le concours du Muséum National d'Histoire Naturelle, Paris.
- Fowler, H.W., 1934. Descriptions of new species obtained 1907 to 1910, chiefly in the Philippine Islands and adjacent seas. *Proceedings of the Academy of Natural Sciences, Philadelphia* 85, 233-367.
- Froese, R., Pauly, D., 1994. FishBase as tool for comparing the life history patterns of flatfish. *Netherlands Journal of Sea Research* 32(3/4), 235-239.
- Froese, R., Pauly, D., Editors, 1996. *FishBase 96: Concepts, Design and Data Sources*. ICLARM, Manila. 179 p.
- Gloerfelt-Tarp, T., Kailola, P.J., 1984. *Trawled fishes of Southern Indonesia and Northwestern Australia*. Australian International Development Assistance Bureau, Australia, The Directorate General of Fisheries, Indonesia, and The German Agency for Technical Cooperation. 406 p.
- Gomez, E.D., 1980. *Bibliography of Philippine Marine Science 1978*. Filipinas Foundation, Inc., Makati, Metro Manila. 178 p.
- Heemstra, P.C., 1986a. Cynoglossidae. In: Smith, M.M., Heemstra, P.C. (eds.), *Smiths' Sea Fishes*, p. 865-868. Springer-Verlag, Berlin.
- Heemstra, P.C. 1986b. Pleuronectidae. In: Smith, M.M., Heemstra, P.C. (eds.), *Smiths' Sea Fishes*, p. 863-865. Springer-Verlag, Berlin.
- Hensley, D.A., 2001. Citharidae. Largescale flounders. In: Carpenter, K.E., Niem, V. (eds.), *The Living Marine Resources of the Western Central Pacific. Volume 6: Bony Fishes Part 4 (Labridae to Latimeriidae), Estuarine Crocodiles, Sea Turtles, Sea Snakes and Marine Mammals*, p. 3794-3798. FAO identification guide for fishery purposes. FAO, Rome.
- Hensley D.A., Amaoka, K., 2001. Bothidae. Lefteye flounders. In: Carpenter, K.E., Niem, V. (eds.), *The Living Marine Resources of the Western Central Pacific. Volume 6: Bony Fishes Part 4 (Labridae to Latimeriidae), Estuarine Crocodiles, Sea Turtles, Sea Snakes and Marine Mammals*, p. 3799-3814. FAO identification guide for fishery purposes. FAO, Rome.
- Hensley, D.A., Amaoka, K., 1989. A redescription of *Pseudorhombus megalops*, with comments on *Cephalopsetta ventrocellota* (Osteichthyes: Pleuronectiformes: Paralichthyidae). *Proceedings of the Biological Society of Washington* 102(3), 577-585. 11
- Hensley, D.A., Randall, J.E., 1990. A redescription of *Engyprosopon macrolepis* (Teleostei: Bothidae). *Copeia* (3), 674-680.
- Herre, A.W., 1953. Checklist of Philippine fishes. Research Report 20, 977 p. Fish and Wildlife Service and United State Department of the Interior, U.S.A.
- Ingles, J., Pauly, D., 1984. An atlas of the growth, mortality and recruitment of Philippines fishes. ICLARM Technical Report 13, 127 p.
- Kimura, S., 1995. A checklist of the marine fishes collected around northern Palawan and Calait islands, Philippines. In: Pawikan Conservation Project-PAWB, DENR, Philippines and Toba Aquarium, Japan, *Dugongs Dugong dugong (Miller 1776) of the Philippines*, p. 158-167. A report of the Joint Dugong Research and Conservation Program. PANB, DENR, Philippines and Toba Aquarium, Japan.
- Kottelat, M., 1993. Technical report on the fishes from fresh and brackish waters of Leyte, Philippines. Technical Report prepared for the Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH and ViSCA-GTZ Ecology Program, Visayan State College of Agriculture, Philippines. 54 p.
- Kulbicki, M., Wantiez, L., 1990. Variations in the fish catch composition in the Bay of St. Vincent, New Caledonia, as determined by experimental trawling. *Journal of Marine and Freshwater Research* 41, 121-144.
- Kurouma, K., Abe, Y., 1986. *Fishes of the Arabian Gulf*. Kuwait Institute for Scientific Research, State of Kuwait, 356 p.
- Labao, E., 1980. Oceanographic survey of Samar Sea. Marine Demersal Fisheries Resources and Management Project and Mid-water Trawl Exploration. University of the Philippines, College of Fisheries, Dept. of Marine Fisheries, Technical Report No. 3. Mimeo, pag. var.
- Livingston, P.A., 1993. Importance of predation by groundfish, marine mammals and birds on walleye pollock *Theragra chalcogramma* and Pacific herring *Clupea pallasii* in the eastern Bering Sea. *Marine Ecology (Progress Series)* 102, 205-215.
- Masuda, H., Amaoka, K., Araga, C., Uyeno, T., Yoshino, T. 1984a. *The Fishes of the Japanese Archipelago*. Vol. 1 (text). Tokai University Press, Tokyo, Japan. 437 p.

- Masuda, H., Amaoka, K., Araga, C., Uyeno, T., Yoshino, T., 1984b. *The Fishes of the Japanese Archipelago*, (plates) Tokai University Press, Shinjuku Tokai Building, Tokyo, Japan. 437 p.
- Matsuura, S., 1961. Age and growth of flatfish Ganzobriame, *Pseudorhombus cinamoneus* (Temminck & Schlegel). *Records of Oceanographic Works in Japan Sp.* (5), 103-110.
- McManus, J.W., Nanola, Jr., C.L., Reyes, Jr., R.B., Kesner, K.N., 1992. Resource ecology of the Bolinao coral reef system. *ICLARM Studies and Reviews* 22, 117 p.
- Menon, A.O.K., 1977. A systematic monograph of the tongue soles of the genus *Cynoglossus* Hamilton-Buchanan (Pisces: Cynoglossidae). *Smithsonian Contributions to Zoology* 238, 129 p.
- Menon, A.O.K., 1984. Soleidae, In: Fischer, W., Bianchi, G. (eds.), *FAO Species Identification Sheets for Fishery Purposes. Western Indian Ocean (Fishing Area 51)*. Volume 4. FAO, Rome. pag. var.
- Menon, A.O.K., Monkolprasit, S., 1974. Cynoglossidae. In: Fischer, W., Whitehead, P.J.P. (eds.), *FAO Species Identification Sheets for Fishery Purposes. Eastern Indian Ocean (fishing area 57) and Western Central Pacific (fishing area 71)*. Volume II. FAO, Rome, pag. var.
- Munroe, T., 2001a. Cynoglossidae. Tongue soles. In: Carpenter, K.E., Niem, V. (eds.), *The Living Marine Resources of the Western Central Pacific. Volume 6: Bony Fishes Part 4 (Labridae to Latimeriidae), Estuarine Crocodiles, Sea Turtles, Sea Snakes and Marine Mammals*, p. 3890-3901. FAO identification guide for fishery purposes. FAO, Rome.
- Munroe, T., 2001b. Soleidae. Soles. In: Carpenter, K.E., Niem, V. (eds.), *The Living Marine Resources of the Western Central Pacific. Volume 6: Bony Fishes Part 4 (Labridae to Latimeriidae), Estuarine Crocodiles, Sea Turtles, Sea Snakes and Marine Mammals*, p. 3878-3889. FAO identification guide for fishery purposes. FAO, Rome.
- Myers, R.F., 1991. *Micronesian Reef Fishes*. 2nd ed. Coral Graphics, Barrigada, Guam. 298 p.
- Nielsen, J., 1984a. Bothidae. In: Fischer, W., Bianchi, G. (eds.), *FAO Species Identification Sheets for Fishery Purposes. Western Indian Ocean Fishing Area 51*. Volume 1. FAO, Rome. pag. var.
- Nielsen, J., 1984b. Psettodidae. In: Fischer, W., Bianchi, G. (eds.), *FAO Species Identification Sheets for Fishery Purposes. Western Indian Ocean Fishing Area 51*. Volume 3. FAO, Rome.
- Pauly, D., 1978. A preliminary compilation of fish length growth parameters. *Ber. Inst. Meereskd. Christian-Albrechts Univ. Kiel*, 55, 200 p.
- Pauly, D., 1980. On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal du Conseil International sur l'Exploration de la Mer* 39(3), 175-192.
- Pauly, D., 1994. A framework for latitudinal comparisons of flatfish recruitment. *Netherlands Journal of Sea Research* 32(2), 107-118.
- Pauly, D., 2010. *Gasping Fish and Panting Squids: Oxygen, Temperature and the Growth of Water-Breathing Animals*. Excellence in Ecology (22), International Ecology Institute, Oldendorf/Luhe, Germany, xxviii + 216 p.
- Pauly, D., Martosubroto, P., Editors, 1996. Baseline studies of biodiversity: the fish resources of Western Indonesia. *ICLARM Studies and Reviews* 23, 312 p.
- Pauly, D., Saeger, J., Silvestre, G., Editors, 1986. Resources management and socioeconomics of Philippine marine fisheries. University of the Philippines in the Visayas, College of Fisheries, Technical Report of the Department of Marine Fisheries 10, 217 p.
- Pradhan, M.J., 1969. Fishery biology of *Psettodes erumei* (Schneider) - an Indian Ocean flatfish. III. The fishery of *Psettodes erumei*. *Bulletin of the National Institute of Science, India* (38), 906-926.
- Rainer, S.F., 1984. Temporal changes in a demersal fish and cephalopod communities of an unexploited coastal environment in northern Australia. *Australian Journal of Marine and Freshwater Research* 35, 747-768.
- Rainer, S.F., Munro, I.S.R., 1982. Demersal fish and cephalopod communities of an unexploited coastal environment in northern Australia. *Australian Journal of Marine and Freshwater Research* 33, 1039-1055.
- Rajaguru, A., 1992. Biology of two co-occurring tonguefishes, *Cynoglossus arel* and *C. lida* (Pleuronectiformes: Cynoglossidae), from Indian waters. *United States Fishery Bulletin* 90, 328-367.
- Ramanathan, N., Natarajan, R., 1979. Breeding biology of *Psettodes erumei* (Bloch & Schn.) and *Pseudorhombus arsius* (Hamilton-Buchanan). Pisces: Pleuronectiformes along Porto Novo coast (S. India). *Aquaculture* 18(3), 269-282.
- Randall, J.E., 1985. *Guide to Hawaiian Reef Fishes*. Harrowood Books, Pennsylvania.
- Randall, J.E., 1995. *Coastal Fishes of Oman*. University of Hawaii Press, Honolulu, Hawaii. 439 p.
- Randall, J.E., Allen, G.R., Steene, R.C., 1990. *Fishes of the Great Barrier Reef and Coral Sea*. University of Hawaii Press, Honolulu, Hawaii. 506 p.
- Ronquillo, I.A., Caces-Borja, P., Mines, A.N., 1960. Preliminary observations on the otter trawl fishery of Manila Bay. *Philippine Journal of Fisheries* 8(1), 47-56.
- Saila, S.B., 1983. The importance and assessment of discards in commercial fisheries. FAO Fisheries Circular No. 765, 62 p.
- Sainsbury, K.J., Kailola, P.J., Leyland, G.G., 1985. *Continental Shelf Fishes of Northern and North-western Australia*. Clouston and Hall and Peter Pownall Fisheries Information Service, Australia. 375 p.

- Sano, M., Shimizu, M., Nose, Y., 1984. *Food Habits of Teleostean Reef Fishes in Okinawa Island, Southern Japan*. University of Tokyo Press, Tokyo, Japan. 128 p.
- Seigel, J., Adamson, T.A., 1985. First record of the genus *Zebrias* (Pisces: Pleuronectiformes: Soleidae) from the Philippine Islands, with the description of a new species. *Proceedings of the Biological Society of Washington* 98(1), 13-16.
- Selga, M., 1931. *Sea Surface Temperature in the Philippines*. Publications of the Manila Observatory, Manila, Philippines.
- Silvestre, G., Regalado, R., Pauly, D., 1986a. Status of Philippine demersal stocks inferences from underutilized catch rate data. In: Pauly, D., Saeger, J., Silvestre, G. (eds.), *Resources, Management and Socio-economics of Philippine Marine Fisheries*, p. 47-96. Department of Marine Fisheries Technical Report 10.
- Silvestre, G., Hammer, C., Sambalay, Jr., V., Torres, Jr., F., 1986b. Size selection and related morphometrics of trawl-caught fish species from the Samar Sea. In: Pauly, D., Saeger, J., Silvestre, G. (eds.), *Resources, Management and Socio-economics of Philippine Marine Fisheries*, p. 107-138. Department of Marine Fisheries Technical Report 10.
- Villoso, E.P., Hermosa, Jr., G.V., 1982. Demersal trawl fish resources of Samar Sea and Carigara Bay, Philippines. *Fisheries Research Journal of the Philippines* 7(2), 59-68.
- Villoso, E.P., Aprieto, V.L., 1983. On the relative abundance and distribution of slipmouths (Pisces: Leiognathidae) in Lingayen Gulf, Philippines. *Fisheries Research Journal of the Philippines* 8(1), 26-43.
- Warfel, E., Manacop, P.R., 1950. Otter trawl explorations in Philippine waters. Research Report 25, Fish and Wildlife Service, U.S. Department of the Interior, Washington, D.C.
- Weber, M., de Beaufort, L.F., 1929. *The Fishes of the Indo-Australian Archipelago. V. Anacanthini, Allotriognathi, Heterostomata, Berycomorphi, Percomorphi: Kuhliidae, Apogonidae, Plesiopidae, Pseudopleusiopidae, Priacanthidae, Centropomidae*. E.J. Brill Ltd., Leiden. 458 p.
- Winterbottom, R., 1993. *Philippine Fishes*. Computerized catalog of the fish collection in the Royal Ontario Museum, Toronto, Canada.

APPENDIX 1. ANNOTATED CHECKLIST OF THE FLATFISHES OF THE PHILIPPINES

Bothidae

Arnoglossus aspilos (Bleeker 1851) Max. length: 19 cm TL; Museum: Eastern Luzon, 4.9 miles off Caringo I., in 11 fathoms (20 m), R/V *Albatross* collection, Stn. 5461, USNM 137659 (Anon. 1994). Sold in dried form called *palad*. See also Kuronuma and Abe (1996).

Arnoglossus brunneus (Fowler 1934) Max. length: 18.3 cm TL; Museum: R/V *Albatross* collections, as *Bothus bnmneus*. east coast of Luzon, in 146 fathoms (267 m), Stn. D. 5453, USNM 93074 (holotype, 18.3 cm) (Fowler, 1934); Sombrero I., Batangas, 118 fathoms (216 m), USNM 93543, and Uanivan I., Batanes, USNM 93544 (paratypes) (Anon., 1994). See also Herre (1953) and Hensley and Amaoka (2001).

Arnoglossus elongatus Weber 1913 Max. length: 11 cm TL; Inhabits coral-sand bottoms from depths of 100-224 m (Hensley and Amaoka, 2001).

Arnoglossus polyspilus (Günther 1880) Max. length: 24 cm TL; Museum: East coast of Luzon, in 195 fathoms (357 m), R/V *Albatross* collection, Stn. 5475, USNM 93076 (as *Bothus tchangi*, 21 cm) (Fowler, 1934). See also Hensley and Amaoka (2001). Additional reference: Morphology in Masuda *et al.* (1984a).

Arnoglossus tapeinosoma (Bleeker 1865) Max. length: 13 cm TL; Museum: R/V *Albatross* collections: western coast of Luzon, off San Fernando Pt., 45 fathoms (82.4 m), USNM 138709; Sulu Sea, off western Mindanao I., off Panabutan Pt., USNM 138712 (Anon., 1994).

Asterorhombus fijiensis (Norman 1931) Max. length: 15 cm TL; Museum: Palawan, Putic I., 0-15 ft (0-4.6 m), USNM 260364. Ajong, Negros I., 0-8 ft (0-2.4 m), USNM 260365. Balicasag I., 0-80 ft (0-24.4 m), USNM 260366. Siquijor I., 0-35 ft (0-11 m), USNM 260367 (Anon., 1994). See also Hensley and Amaoka (2001).

Asterorhombus intermedius (Bleeker 1865) Max. length: 15 cm TL; Museum: Bais Bay, Negros I., 0-120 ft (0-36.6 m), USNM 260363 (Anon., 1994). Additional reference: Morphology in Myers (1991).

Bothus mancus (Broussonet 1782) Max. length: 42 cm SL; Museum: Tagburos, Puerto Princesa, USNM 227085. West of Engano Point, Barrio Anqib., Santa Ana, Cagayan Prov., USNM 309422. Fuga I. (Babuyan Is.) USNM 318329. Maybag I., (Babuyan Is.), USNM 318330 (Anon., 1994). See also Herre (1953) and Randall *et al.* (1990). Additional references: Morphology in Myers (1991). Diet in Randall (1985).

Bothus myriaster (Temminck & Schlegel 1846) Max. length: 27 cm TL; A rare species found in sand and mudd bottoms of continental shelves (Hensley and Amaoka, 2001). See also Conlu (1979). Additional reference: Morphology in Masuda *et al.* (1984a).

Bothus pantherinus (Rüppell 1830) Max. length: 30 cm TL; Reported from southern to western Luzon to Cagayan Prov., Palawan, the Visayas (Panay, Negros, Cebu, Bohol), and northern Mindanao. Museum: ANSP 63543, 63483; LACM 347416, 42485-7; USNM 260373, 260471. Two specimens, 5.2 and 6 in (13 and 15 cm), were collected from Bacon, Sorsogon (Evermann and Scale, 1907). See also Herre (1953), Randall *et al.* (1990), Myers (1991), and Anon. (1994). Additional reference: Morphology in Myers (1991).

Chascanopsetta lugubris Alcock 1894 Max. length: 38 cm SL; Museum: Balayan Bay, Luzon, USNM 138016. Gulf of Davao, Dumalag I. USNM 138017. Northern Mindanao, USNM 138018. Luzon coast, USNM 138019-20 (Anon. 1994). See also Masuda *et al.* (1984a, 1984b). Additional reference: Morphology in Masuda *et al.* (1984a).

Chascanopsetta micrognatha Amaoka & Yamamoto 1984 Max. length: 27.4 cm. Reported by Kunio Amaoka (pers. comm.) using samples collected by him during the WCP Workshop 1995.

Crossorhombus valderostratus (Alcock 1890) Max. length: 14 cm TL; Museum: China Sea, vicinity s. Luzon, Malavatuan I., 80 fathoms (146 m), R/V *Albatross* collection, Stn. 5277, USNM 137391 (Anon., 1994).

Engyprosopon grandisquama (Temminck & Schlegel 1846) Max. length: 15 cm TL; Reported from Sulu archipelago to Corregidor I., Manila Bay. Museum: USNM 137924-41 (Anon., 1994). Sold in the market in dried form called *palad*. See also Herre (1953). Additional reference: Morphology in Masuda *et al.* (1984a).

Engyprosopon latifrons (Regan 1908) Max. length: 8 cm SL. Inhabits sandy bottoms at depths of 37-68 m (Hensley and Amaoka, 2001).

Engyprosopon macrolepis (Regan 1908) Max. length: 5.9 cm SL. Inhabits sandy and muddy bottoms (Hensley and Amaoka, 2001). Museum: BPBM 26860; USNM 260378; CAS-SU 33678. Species redescribed by Hensley and Randall (1990). Additional reference: Morphology in Hensley and Randall (1990).

Engyprosopon maldivensis (Regan 1908) Max. length: 12.7 cm SL. Museum: R/V *Albatross* collections, as *Arnoglossus maculipinnis*: vicinity of Jolo, in 20 to 76 fathoms (37-139 m), Stn. D. 5140, USNM 93098 (10.1 cm) (Fowler, 1934); between Samar and Leyte, vicinity of Surigao Strait, Tabuc Pt., (Leyte), 62 fathoms (113.5 m), Stn. 5480, USNM 93570 (Anon., 1994). See also Amaoka *et al.* (1993) and Hensley and Amaoka (2001). Additional reference: Morphology in Masuda *et al.* (1984a).

Engyprosopon mogkii (Bleeker 1854) Max. length: 11 cm SL. Known from Mindanao, southern Negros, Palawan, to southern Luzon. Museum: USNM 137960-81, 260468 (Anon., 1994). Based on records, this species occurs in estuarines, reef sand flats, and embayments.

Engyprosopon obliquiocolatum (Fowler 1934) Max. length: 7.6 cm. Museum: collected most likely from deep water, R/V *Albatross* collection, as *Bothus obliquiocolatits*, USNM 93077 (holotype, 7.6 cm); USNM 93078 (4 paratypes) (Anon., 1994).

Grammatobothus polyophthalmus (Bleeker 1865) Max. length: 21 cm TL. Reported from southern Negros to Masbate Is. and off entrance to Manila Bay (Herre, 1953). Museum: USNM 260448, 160480-1 (Anon., 1994). See also Weber and de Beaufort (1929).

Kamoharaia megastoma (Kamohara 1936) Max. length: 22.5 cm TL. Inhabits sandy and muddy bottoms (Hensley and Amaoka, 2001). Additional reference: Morphology in Masuda *et al.* (1984a).

Laeops clarus Fowler 1934 Max. length: 15.5 cm TL. Museum: R/V *Albatross* collections: between Cebu and Bohol, in 162 fathoms (296.5 m), Stn. D. 5412, USNM 93083 (holotype, 15.5 cm) (Fowler, 1934); east coast of Luzon, San Bernadino Strait to San Miguel Bay; Atulayan I., 0-560 fathoms (1025 m), USNM 93560 (paratype) (Anon., 1994). See also Herre (1953) and Hensley and Amaoka (2001).

Laeops cypho Fowler 1934 Max. length: 14.9 cm. Museum: R/V *Albatross* collections: off northern Mindanao, in 182 fathoms (333 m), Stn. D. 5519, USNM 93085 (holotype, 14.9 cm) (Fowler, 1934); Sombrero I., Batangas, 118 fathoms (216 m), USNM 93567 (paratype) (Anon., 1994). Type locality identified as off Point Tagolo, Zamboanga (Herre, 1953).

Laeops gracilis Fowler 1934 Max. length: 16.5 cm TL. Museum: East of Masbate, in 108 fathoms (197 m), R/V *Albatross* collection, Stn. D. 5212, USNM 93084 (holotype, 16.5 cm) (Fowler, 1934). See also Herre (1953) and Hensley and Amaoka (2001).

Laeops guentheri Alcock 1890 Max. length: 14 cm TL. Museum: West coast of Luzon, from Manila Bay to Lingayen Gulf, S. Fernando Pt., in 45 fathoms (82.4 m), R/V *Albatross* collection, Stn. 5442, USNM 137394 (Anon., 1994).

Laeops parviceps Günther 1880 Max. length: 14 cm TL. Museum: R/V *Albatross* collections: east coast of Luzon, San Bernardino Strait to San Miguel Bay, Legaspi, 146 fathoms (267 m), Stn. 5453, USNM 137395; west coast of Luzon, Manila Bay to Lingayen Gulf, San Fernando Pt., 45 fathoms (82.4 m), Stn. 5442, USNM 137396; Visayan Sea between northern Negros and Masbate Is., se. Tanguingui I., 0-69.5 m, USNM 260451 (Anon., 1994).

Neolaeops microphthalmus (von Bonde 1922) Max. length: 21 cm SL. Inhabits sandy and muddy bottoms (Hensley and Amaoka, 2001). Additional reference: Morphology in Masuda *et al.* (1984a).

Psettina brevirectis (Alcock 1890) Max. length: 8 cm SL. Museum: R/V *Albatross* collections: western Samar, Taratara I., 20 fathoms (37 m), Stn. D.5209, USNM 137389; off eastern Panay, Antonia I., 24 fathoms (44 m), Stn. 5182, USNM 137390 (Anon., 1994).

Psettina gigantea Amaoka 1963 Max. length: 13 cm SL. Museum: Visayan Sea between northern Negros and Masbate Is. northwest Guintacan I., 0-80.5 m, USNM 260446; southwest of Caduruan Point, 0-78.7 m, USNM 260482 (Anon., 1994). Additional reference: Morphology in Masuda *et al.* (1984a).

Psettina variegata (Fowler 1934) Max. length: 9.2 cm SL. Museum: between Samar and Leyte Islands in 61 fathoms (112 m), R/V *Albatross* collection Stn. D. 5481, USNM 93091 (as *Bothus variegatus*, holotype, 9.2 cm) (Fowler, 1934; Herre, 1953). See also Hensley and Amaoka (2001).

Taeniopsetta ocellata (Günther 1880) Max. length: 11.4 cm. Specimens 6.9 to 11.4 cm were trawled from Stn. 16, at depth of 150 to 164 m during the R/V *Vauban* expedition (Fourmanoir, 1976 in Fourmanoir, 1981). Additional reference: Morphology in Masuda *et al.* (1984a).

Citharidae

Brachypleura novaezeelandiae Günther 1862 Max. length: 14 cm TL. Museum: Marinduque and vicinity, USNM 137708. Off Luzon, Sueste Pt., USNM 137709-10. Manila Bay, Corregidor Lt., USNM 137711. S. Mindanao, eastern Illana Bay, USNM 137712. E. Mindanao, Nagubat I., USNM 137714. Visayan Sea, between northern Negros and Masbate I., USNM 261361, 261363-4, 261526. Carigara Bay, Samar Sea, USNM 228536-9 (Anon., 1994). See also Herre (1953) and Kuronuma and Abe (1986).

Citharoides axillaris (Fowler 1934) Max. length: 19.5 cm. Museum: *Albatross* collections, as *Erachyphurops axillaris*: Balayan Bay and Verde Island Passage, in 118 fathoms (216 m), R/V, Stn. D. 5117, USNM 93080 (holotype, 19.5 cm) (Fowler, 1934; Herre, 1953); China Sea, vicinity of southern Luzon, Malavatuan I., USNM 93545; Balabac Strait, Cape Melville, USNM 93547 (Anon., 1994).

Citharoides macrolepidotus Hubbs 1915 Max. length: 29 cm TL. A rare species found at depths of 121-240 m (Hensley, 2001).

Lepidoblepharon ophthalmolepis Weber 1913 Max. length: 36 cm TL. Museum: Balanja Pt., Mindoro Is., in 234 fathoms (428 m), R/V *Albatross* collection, Stn. 5260, USNM 137408 (Anon., 1994).

Cynoglossidae

Cynoglossus arel (Bloch & Schneider 1801) Max. length: 40 cm TL. Inhabits muddy and sandy bottoms of the continental shelf down to 125 m (Munroe, 2001). Additional references: Growth in Pauly (1980). Food, Diet, Reproduction, and Spawning in Rajaguru (1992).

Cynoglossus bilineatus (Lacepède, 1802) Max. length: 44 cm SL. Museum: R/V *Albatross* collections: Cavite Mkt., USNM 137616; Manila Mkt., USNM 137617, 137652; Palawan, Verde del Sur, reef sand flat, USNM 137618; Manila Bay, USNM 137620; Limbones Cove, USNM 286919 (removed from 113179 and recatalogued) (Anon., 1994). See also Herre (1953) and Menon (1977). Additional reference: Food in Blaber (1980).

Cynoglossus cynoglossus (Hamilton 1822) Max. length: 20 cm TL. Museum: ANSP 49038-9; NHV 43826. See also Herre (1953) and Menon (1977).

Cynoglossus kopsii (Bleeker 1851) Max. length: 17.7 cm SL. Museum: Iloilo, U.S.N. Eclipse Expedition USNM 112872-4. collections: Pt. Tagalo, 102 fathoms (187 m), Stn. 5520, USNM 113186; Lingayen Gulf, e. of Pt. Guacet, Stn. 5442, USNM 113187; Marinduque and vicinity, Tayabas, Stn. 5371, USNM 113188; Tawi Tawi, 34 fathoms (62.6 m), Stn. D.5152, USNM 137653; Cotabato, USNM 137656; Off San Fernando, 45 fathoms (82 m), Stn. D.5442, USNM 137657; Corregidor, 12 fathoms (22 m), Stn. 5360, USNM 137658; Panay, Iloilo, USNM 148586 (Anon., 1994). See also Herre (1953) and Menon (1977).

Cynoglossus lida (Bleeker 1851) Max. length: 21.3 cm SL. Museum: R/V *Albatross* collections: Davao, USNM 137952; Abuyog, Leyte, USNM 137953, 137957; Hinunangan B., USNM 137954; Iloilo Mkt., USNM 137955; offcast coast of Leyte I., Tacloban Anchorage, USNM 137956; Palawan, Mantaquin B., USNM 137958; Cotabato, below river mouth, USNM 137959 (Anon., 1994). BMNH 1872.4.6.96 (Menon, 1977). See also Herre (1953) and Heemstra (1986a). Additional references: Growth, Food, Diet, Reproduction, and Spawning in Rajaguru (1992).

Cynoglossus lingua Hamilton 1822 Max. length: 45 cm TL. Museum: China Sea, off s. Luzon, 17.5 miles from Malavatuan I., 525 fathoms (961 m), Stn. D.5274, USNM 137410 (Anon., 1994).

Cynoglossus monopus (Bleeker 1849) Max. length: 18.8 cm SL. Found on muddy substrates from 13-18.3 m (Menon, 1977). Museum: AMNH 19645.

Cynoglossus puncticeps (Richardson 1846) Max. length: 18 cm TL. Specimens were collected during the U.S.N. Eclipse Expedition and R/V *Albatross* from Cotabato, Mindanao, central and eastern Visayas to southern Philippines (Anon., 1994). Museum: ANSP 63524, 82548; LACM 42475-47. See also Herre (1953) and Menon and Monkolprasit (1974). Additional references: Morphology in Fischer and Whitehead (1974). Growth in Pauly (1994).

Cynoglossus suyeni Fowler 1934 Max. length: 27.5 cm SL. Museum: R/V *Albatross* collections: off southern Luzon, China Sea (Verde I. Passage, off Escarceo Light, Mindoro), in 173 fathoms (317 m), Stn. D. 5291, USNM 93086 (holotype, 15.5 cm) (Fowler, 1934); USNM 113189-113194, 137941-8,- 137950 (Anon., 1994). See also Herre (1953) and Menon (1977).

Paraplagusia bilineata (Bloch 1787) Max. length: 60 cm TL. Museum: R/V *Albatross* collections: Iloilo Mkt., USNM 138070, 138071; Manila Mkt, USNM 138072; Chase Head Endeavor St., Palawan, USNM 138073; Paluan Bay, Mindoro, USNM 138074; Mansalay Bay, southeastern Mindoro, USNM 138075; Lingayen Gulf, USNM 138076; Siquijor I., Santa Maria, USNM 138077; Abuyog, Leyte, USNM 138079; Subig Bay, USNM 138080; Port San Pio Quinto, Camiguin I., 1-6 m, USNM 138082; Panabutan Bay, Mindanao, USNM 138083; Cotabato, USNM 138084; Davao, USNM 138085; Balayan Bay, Luzon, Taal Anchorage, USNM 138086; Bolinao lagoon, Pangasinan, USNM 228535; northeastern side of Siquijor, tidal lagoon, USNM 273773; USNM 138081 (Anon., 1994). See also Herre (1953) and Heemstra (1986a). Additional references: Growth in Pauly (1978) and Erzini (1991). Food in Livingston (1993).

Paraplagusia blochii (Bleeker 1851) Max. length: 20 cm SL. Museum: R/V *Albatross* collections: Philippine Sea, off Daet, Luzon, 15 June 1909, USNM 138087 (7, 11.6-22 cm) (Chapleau and Renaud, 1993); Limbones Cove, USNM 113179. Iloilo, U.S.N. Eclipse Expedition, USNM 112870 (Anon., 1994). ANSP 77427. Also known from Dumaguete, Negros Oriental (Herre, 1953). See also Winterbottom (1993) and Randall (1995).

Symphurus gilesii (Alcock 1889) Max. length: 14 cm. Two specimens collected between 70 to 215 m during the R/V *Vauban* expedition (Fourmanoir, 1976 in Fourmanoir, 1981).

Symphurus marmoratus Fowler 1934 Max. length: 9.8 cm. Museum: Jolo I. and vicinity, in 10 fathoms (18.3 m), R/V *Albatross* collection, Stn. D. 5561, USNM 93092 (holotype, 9.8 cm) (Fowler, 1934).

Symphurus regani Weber & Beaufort 1929 Max. length: 12 cm. Museum: Between Siquijor and Bohol Is., Balicasag I., 805 fathoms (1473 m), R/V *Albatross* collection, Stn. 5526, USNM 138045 (Anon., 1994).

Symphurus septemstriatus (Alcock 1891) Max. length: 10 cm TL. Museum: R/V *Albatross* collections: Verde I. Passage and Batangas Bay, Matocot Pt., 135 fathoms (247 m), Stn. 5265, USNM 138023 and in 170 fathoms (311 m), Stn. 5268, USNM 163654; between Burias and Luzon, Anima Sola I., 215 fathoms (393 m), Stn. 5216, USNM 138026; China Sea, vicinity s. Luzon, Matocot Pt., 140 fathoms (256 m), Stn. 5298, USNM 138028; between Samar and Masbate, Tubig Pt., Destacado I., 118 fathoms (216 m), Stn. 5391 and in 135 fathoms (247 m), Stn. 5392, USNM 138032; between Cebu and Bohol, Luis Pt., 145 fathoms (265 m), Stn. 5411, USNM 138037; off n. Luzon, Hermanos I., 230 fathoms (421 m), Stn. 5326, USNM 138040; between Burias and Luzon, Bagatao I., 226 fathoms (414 m), Stn. 5388, USNM 138041 and in 209 fathoms (382 m), Stn. 5387, USNM 138042; Camp Overton Lt, Iligan Bay, Stn. 5508, USNM 163655; Dupon Bay (Leyte) and vicinity, Ponson I., 262 fathoms (479 m), Stn. 5405, USNM 163657 (Anon., 1994).

Symphurus strictus Gilbert 1905 Max. length: 14 cm. Museum: R/V *Albatross* collections: Verde I. Passage and Batangas Bay, Matocot Pt., 220 fathoms (402 m), Stn. 5269, USNM 138024; China Sea, vicinity s. Luzon, Matocot, 214 fathoms (392 m), Stn. 5290, USNM 138027 and, Escarceo, 244 fathoms (446 m), Stn. 5294, USNM 138030 (Anon., 1994).

Symphurus woodmasoni (Alcock 1889). Known in the Visayan and Mindanao area, R/V *Albatross* collections (Anon., 1994).

Paralichthyidae

Paralichthys olivaceus (Temminck & Schlegel 1846) Max. length: 80 cm SL. Inhabits muddy and sandy bottoms of shallow waters (Amaoka and Hensley, 2001). Additional reference: Diet in Dou (1992).

Pseudorhombus argus Weber 1913 Max. length: 25 cm SL. Museum: Buton Strait, Kalono Pt., in 39 fathoms (71.4 m), R/V *Albatross* collection, Stn. 5641, USNM 137393 (Anon., 1994). Additional reference: Morphology Amaoka and Hensley (2001).

Pseudorhombus arsius (Hamilton 1822) Max. length: 45 cm TL. Known from northwestern Mindanao to southern and western Luzon, R/V *Albatross* collections. Museum: Davao, USNM 137985. Malabang, USNM 137986. Cavite Mkt, USNM 137987. Manila Mkt., USNM 137988, 137993, 137996. North of Malampaya R., USNM 137989. Mantaquin B., Palawan, USNM 137990. Endeavor Pt., in 14-25 fathoms (26-46 m), Stn. 5342, USNM 137991. Abuyog, Leyte, USNM 137992] Outside Harbor of Manila Bay, USNM 137994. Iloilo Mkt, USNM 137995, 138000. Ragay R., tidewater, USNM 137998. Samar I., Catbalogan, USNM 137999. Cuyo Is., USNM 138001 (Anon., 1994). LACM 42475-33. See also Weber and de Beaufort (1929) and Herre (1953). Additional references: Morphology in Amaoka and Hensley (2001). Growth in Bawazeer (1987). Food in Blaber (1980).

Pseudorhombus cinnamoneus (Temminck & Schlegel 1846) Max. length: 35 cm SL. A 17.7 cm specimen was caught between 150 to 164 m during the 1976 R/V *Vauban* expedition (Fourmanoir, 1976 in Fourmanoir, 1981). Museum: Puerto Princesa Market, USNM 227078 (Anon., 1994). See also Herre (1953) and Masuda *et al.* (1984a). Additional references: Morphology in Amaoka and Hensley (2001). Growth in Matsuura (1961).

Pseudorhombus diplospilus Norman 1926 Max. length: 40 cm SL. Museum: Visayan Sea between northern Negros and Masbate Is., southwest of Caduruan Point, in 75 m, USNM 260477 (Anon., 1994). Additional reference: Morphology in Amaoka and Hensley (2001).

Pseudorhombus dupliocellatus Regan 1905 Max. length: 40 cm SL. Museum: Visayan Sea, between northern Negros and Masbate I., southwest of Caduruan Pt., 0-75 m, USNM 260478; north of Tanguingui I., USNM 260479; northwest Guintacan I., USNM 260687 (Anon., 1994). One large sample collected from Bulan, Sorsogon, USNM 55898 (as *Platophrys palad*, holotype, 15.5 in (39 cm)) (Evermann and Scale, 1907). Additional reference: Morphology in Amaoka and Hensley (2001).

Pseudorhombus javanicus (Bleeker 1853) Max. length: 35 cm SL. Museum: Bulan, USNM 55967. Panabutan Bay, Mindanao, USNM 138714. Buena Vista, Guimaras I. (Iloilo Strait), USNM 138715. Manila Bay, Corregidor Lt., USNM 138716. Visayan Sea between northern Negros and Masbate Is., southeast south Gigante, USNM 260447 (Anon., 1994). ANSP 49030, 49272. One specimen, 8.25 in. (21 cm), collected from Bulan, Sorsogon (Evermann and Seale, 1907). See also Herre (1953) and Nielsen (1984a). Additional references: Morphology in Amaoka and Hensley (2001). Growth in Chan and Liew (1986).

Pseudorhombus malayanus Bleeker 1865 Max. length: 35 cm SL. Museum: R/V *Albatross* collections: off east coast of Leyte I., Mariquitdaquit I., 15 fathoms (27 m), Stn. 5204, USNM 137420; Manila Bay, Corregidor Lt., 12 fathoms (22 m), Stn. 5361, USNM 137421; Bacoor Beach, USNM 137422; Manila Mkt., USNM 137423; Western Samar, Taratara I., 20 fathoms (37 m), Stn. D.5209, USNM 137424 (Anon., 1994). LACM 35964-9, 35957-15. Additional reference: Morphology in Amaoka and Hensley (2001).

Pseudorhombus megalops Fowler 1934. Max. length: 22 cm SL. Museum: Between Samar and Masbate in 135 fathoms (247 m), R/V *Albatross* collection, Stn D.5392, USNM 93082 (holotype, 22 cm) (Fowler, 1934). USNM 93548-51 (paratypes). Morphological information found also in Hensley and Amaoka (1989).

Pseudorhombus micrognathus Norman 1927. Museum: R/V *Albatross* collections: Balayan Bay, Luzon, C. Santiago Lt, 214 fathoms (392 m), Stn. 5365, USNM 137654; Sulu Archipelago, Tawi-tawi group, Tinakta I., 18 fathoms (33 m), Stn. 5157, USNM 137655 (Anon., 1994).

Pseudorhombus neglectus Bleeker 1865 Max. length: 25 cm SL. Museum: Bulan I., USNM 55968. Panay I., Iloilo, Naval Eclipse Expedition, USNM 102648 (Anon., 1994). Three specimens collected from San Fabian, Pangasinan, 3.5-6.75 in (9-17 cm) (Evermann and Seale, 1907). Also known from Dumaguete, Negros Oriental. Additional reference: Morphology in Amaoka and Hensley (2001).

Pseudorhombus oligodon (Bleeker 1854) Max. length: 30 cm SL. Inhabits muddy and sandy bottoms of continental shelves. Morphological information found also in Amaoka and Hensley (2001). See also Weber and de Beaufort (1929).

Pseudorhombus pentopthalmus Günther 1862 Max. length: 18 cm SL. Museum: Samar I., Catbalogan, USNM 137923. Visayan Sea between northern Negros and Masbate, southeast south Gigante, USNM 260384. Visayan Sea, east of Sicogon I., USNM 260385 (Anon., 1994). See also Masuda *et al.* (1984a, 1984b). Additional reference: Morphology in Amaoka and Hensley (2001).

Pseudorhombus polyspilos (Bleeker 1853) Max. length: 27 cm. Inhabits muddy and sandy bottoms of shallow waters (Amaoka and Hensley, 2001). See also Weber and de Beaufort (1929).

Pseudorhombus russellii (Gray 1834) Max. length: 23 cm. Museum: ANSP 63710, 63544 (12.81 and 16.34 cm). One specimen, 23 cm, was also collected from Bulan, Sorsogon (Evermann and Seale, 1907).

Pleuronectidae

Nematops macrochirus Norman 1931 Max. length: 8.2 cm TL. Museum: China Sea, off southern Luzon at 135 fathoms (247 m), R/V *Albatross* collection, D. 5110, USNM 93087 (holotype, 8.2 cm) (Fowler, 1934). Type locality described as near Corregidor I. See also Herre (1953) and Hensley (2001).

Poecilopsetta colorata Günther 1880 Max. length: 17 cm TL. Museum: Vicinity of southern Luzon, Malavatuan I., 117 fathoms (214 m), R/V *Albatross* collection, Stn. 5275, USNM 137392 (Anon., 1994).

Poecilopsetta megalepis Fowler 1934 Max. length: 18 cm TL. Museum: R/V *Albatross* collections: Balayan Bay and Verde I. Passage, in 118 fathoms (216 m), Stn. D. 5117, USNM 93094 (holotype, 12.8 cm) (Fowler, 1934; Herre, 1953). Balabac Strait: Cape Melville, 148 fathoms (271 m), USNM 93576 (Anon., 1994).

Poecilopsetta plinthus (Jordan & Starks 1904) Max. length: 19 cm TL. Fourmanoir (1976 in Fourmanoir, 1981) reported two specimens (9.6 and 9.9 cm) caught between 185 and 200 m during the R/V *Vauban* expedition. See also Herre (1953).

Poecilopsetta praelonga Alcock 1894 Max. length: 17.5 cm TL. Reported from Davao, Mindanao, central Visayas to the west coast of Luzon; specimens caught between 247-511 m, USNM 138004-138015 (Anon., 1994).

Psettodidae

Psettodes erumei (Bloch & Schneider 1801) Max. length: 64 cm. Known from Iloilo, west to Palawan, and north to western Luzon (Herre, 1953). Occurs from shallow waters to over 300 m deep; most abundant between 22 to 40 m (Warfel and Manacop, 1950). Museum: LACM 35957-12. Off El Nido, gill net, FRLM 11761 (Kimura, 1995). Additional references: Morphology in Nielsen (1984b). Growth in Pradhan (1969), Pauly (1978), and Edwards and Shaher (1991). Food in Devadoss *et al.* (1977) and Cabanban (1991). Diet and Reproduction in Devadoss *et al.* (1977). Spawning in Devadoss *et al.* (1977) and Ramanathan and Natarajan (1979).

Samaridae

Plagiopsetta glossa Franz 1910 Max. length: 19 cm TL. Specimens were collected between 150 and 164 m (Fourmanoir, 1976 in Fourmanoir, 1981).

Samaris cristatus Gray 1831 Max. length: 22 cm TL. Museum: R/V *Albatross* collections: Between Samar and Leyte, vicinity of Surigao Strait, Tabuc Ft., 62 fathoms (114 m), Stn. 5480, USNM 00137649; Buton Strait, Kalono Ft., 39 fathoms (71 m), Stn. 5641, USNM 137650. Samar Sea collection, Carigara Bay, USNM 228532 (Anon., 1994). A specimen, 12 cm, was caught between 70 and 76 m. See also Herre (1953) and Heemstra (1986b). Additional reference: Morphology in Hensley (2001).

Samariscus huysmani Weber 1913 Max. length: 11.5 cm TL. Museum: Samar Sea, Carigara Bay, 0-65 m, USNM 27534 (Anon., 1994).

Samariscus longimanus Norman 1927 Max. length: 12 cm TL. Museum: R/V *Albatross* collection: Between Cebu and Bohol, Luis Ft., 145-162 fathoms (265-297 m), Stns. 5411, 5412, 5418, USNM 137384-6; Ft. Tagolo, 182 fathoms (333 m), Stn. 5519, USNM 137387; Balayan Bay and Verde I. Passage, Sombrero I., 118 fathoms (216 m), Stn. 5117, USNM 137388 (Anon., 1994).

Samariscus luzonensis Fowler 1934 Max. length: 7.6 cm TL. Museum: West coast of Luzon, in 45 fathoms (82.4 m), R/V *Albatross* collection, Stn. D. 5442, USNM 93089 (holotype, 7.6 cm) (Fowler, 1934). Type locality identified as off San Fernando, La Union, Luzon. See also Herre (1953), Anon. (1994), and Hensley (2001).

Samariscus macrognathus Fowler 1934 Max. length: 5.5 cm TL. Museum: West coast of Luzon, in 45 fathoms (82.4 m), R/V *Albatross* collection, Stn. D. 5442, USNM 93088 (holotype, 5.4 cm) (Fowler, 1934). Type locality identified as San Fernando, La Union, Luzon (Anon., 1994). See also (Hensley, 2001).

Samariscus triocellatus Woods 1960 Max. length: 9 cm TL. Museum: Siquijor I., 80-100 ft (24-30 m), USNM 273792. White Beach, past Mahatae, Batan I., Batanes, 50-70 ft (15-21 m), USNM 298212 (Anon., 1994). Additional reference: Morphology in Myers (1991).

Soleidae

Aesopia cornuta Kaup 1858 Max. length: 20 cm SL. Caught by trawl in the seagrass beds of Bolinao (McManus *et al.*, 1992).

Aesopia heterorhinos (Bleeker 1856) Max. length: 11 cm SL. Museum: As *Soleichthys heterorhinos*: Bacon, USNM 55963; R/V *Albatross* collections-Cebu Mkt. USNM 137412 and Batan I., Caracaran Bay, USNM 137413; Sombrero I., Batangas, USNM 28550; west side of Solino (Selinog) I., Zamboanga del Norte, Mindanao, 0-15 ft (4.6 m), USNM 273795; near Tonga Pt., Siquijor I., 0-1.2 m, USNM 273796; tidal lagoon, northeastern side of Siquijor, 0-1 m, USNM 273800 (Anon., 1994). A 4.2 in (11 cm) specimen was collected from Bacon, Sorsogon (Evermann and Scale, 1907). See also Weber and de Beaufort (1929). Additional reference: Morphology in Myers (1991).

Aseraggodes cyaneus (Alcock 1890) Max. length: 8.3 cm SL. Museum: R/V *Albatross* collections: Balayan Bay and Verde I. Passage, Sombrero I., 340 fathoms (622.2 m), Stn. 5114, USNM 137674; China Sea, vicinity of southern Luzon, Corregidor, in 114 fathoms (208.6 m), USNM 137675 and in 118 fathoms (216 m), USNM 137676; east coast of Luzon, Legaspi, USNM 137678 (Anon., 1994). LACM 42475-47.

Aseraggodes dubius Weber 1913 Max. length: 8.5 cm. Museum: R/V *Albatross* collections: Davao, USNM 137667; China Sea, off s. Luzon, Sueste Pt., 25 fathoms (46 m), Stn. 5105, USNM 137668; Verde I. Passage and Batangas Bay, Matocot Pt., 100 fathoms (183 m), Stn. 5266, USNM 137669; Marinduque I. and vicinity, Tayabas, 90 fathoms (165 m), Stn. 5376, USNM 137671 and in 83 fathoms (152 m), Stn. 5371, USNM 137672; Batangas River, Luzon, USNM 137673 (Anon., 1994).

Aseraggodes filiger Weber 1913 Max. length: 11 cm. Collected from Manila Bay, 8 miles from Corregidor Is. in 15-25 fathoms (27-46 m) (Herre, 1953).

Aseraggodes kaianus (Günther 1880) Max. length: 11.3 cm. Forty specimens, ranging from 7.2-9 cm, were collected between 150-164 m, during the R/V *Vauban* expedition (Fourmanoir, 1976 in Fourmanoir, 1981).

Brachirus aspilos (Bleeker 1852) Max. length: 38 cm. Museum: Ulugan Bay near mouth of Baheli River, USNM 137679; Cebu Mkt., USNM 137680-1; Nasugbu Bay, Luzon, USNM 137682 (Anon., 1994).

Brachirus muelleri (Steindachner 1879) Max. length: 18 cm. Museum: Carigara Bay, Samar Sea, 50-70 m, USNM 228530. Sorsogon Mkt., USNM 286939 and 291084 (Anon., 1994).

Brachirus orientalis (Bloch & Schneider 1801) Max. length: 30 cm SL. Inhabits shallow sand and muddy bottoms of coastal waters (Menon and Monkolprasit, 1974).

Brachirus sorsogonensis Evermann & Seale 1907. Max. length: 23 cm. Museum: Bacon, Sorsogon, USNM 55916 (holotype, 9 in. (23 cm)) (Evermann and Seale, 1907). Cuyo Is., USNM 72194 (Anon., 1994).

Heteromycteris hartzfeldii (Bleeker, 1853) Max. length: 11.4 cm; Museum: R/V *Albatross* collections: Leyte, Hinunangan B., USNM 137718; Mindanao, Davao, USNM 137719; Cotabato, USNM 137720; Palawan, Verde del Sur, USNM 137721; Port Bais, eastern Negros, USNM 137722; Mantaquin Bay, Palawan, USNM 137723; Subic Bay, Olongapo, USNM 137724 (Anon. 1994). A 4.5 in (11.4 cm) specimen was collected from the country (Evermann and Seale 1907). See also Herre (1953).

Liachirus melanospilus (Bleeker 1854) Max. length: 7.5 cm SL. Reported from Manila Bay (Herre, 1953).

Pardachirus pavoninus (Lacepède 1802) Max. length: 25 cm TL. Museum: Cebu Mkt., USNM 137624-29. Bacon, USNM 55966. Zamboanga, USNM 84258. Jolo Mkt. USNM 137622. Bolinao Bay, USNM 137623. Pagapas Bay, Santiago R., USNM 137630. Senora Ascion, n. of Dumaguete, Negros O., USNM 273799. Tagbueros, Puerto Princesa City Mkt., USNM 286974 (Anon., 1994). LACM 37398-9, 37397-2, 37398-9, 42471-4. Marketable in Jolo, Sulu, and Cebu. A specimen, 5.5 in (14 cm) in length, was collected from Bacon, Sorsogon (Evermann and Scale, 1907). See also Herre (1953) and Randall *et al.* (1990). Additional references: Morphology in Myers (1991). Food in Sano *et al.* (1984).

Pardachirus poropterus (Bleeker 1851) Max. length: 6.6 cm TL. Museum: Rio Grande, Mindanao, USNM 56164 (Anon., 1994). Three specimens were caught at depths between 122 and 205 m during the 1976 R/V *Vauban* expedition (Fourmanoir, 1976 in Fourmanoir, 1981). See also Herre (1953) and Kottelat (1993).

Solea humilis Cantor 1849 Max. length: 8.9 cm. Considered a commercial fish in the country (Warfel and Manacop, 1950). See also Weber and de Beaufort (1929).

Solea ovata Richardson 1846 Max. length: 10 cm TL. Museum: R/V *Albatross* collections: Manila Mkt. USNM 137397, 137399-404; Sorsogon Mkt., USNM 137405 (Anon., 1994). See also Munroe (2001).

Synaptura marginata Boulenger 1900 Max. length: 50 cm TL. Caught in seagrass beds. Museum: Tagbueros, Puerto Princesa City Mkt., USNM 226832 (Anon., 1994).

Synaptura megalepidoura (Fowler 1934) Max. length: 24.3 cm. Museum: R/V *Albatross* collections, as *Brachirus megalepidoura*: offeast coast of Leyte, 15 fathoms (27 m), Stn. D. 5204, USNM 93081 (holotype, 24.3 cm) (Fowler, 1934); western Samar, Taratara I., 20 fathoms (37 m), Stn. D.5209, USNM 93554 (Anon., 1994). See also Herre (1953).

Zebrias lucapensis Seigel & Adamson 1985 Max. length: 8.4 cm SL. Museum: Lingayen Gulf, LACM 37436-6 (holotype); LACM 37436-8 (paratype). Morphological information found also in Seigel and Adamson (1985).

Zebrias quagga (Kaup 1858) Max. length: 15 cm TL. Inhabits shallow coastal waters (Menon 1984).

Zebrias zebra (Bloch 1787) Max. length: 19 cm TL. Museum: Tigbauan, Panay, USNM 106828 (Anon., 1994).