

H53-340

State or Country: Pacific Ocean Field No. TP-40

County: Map: 1

Locality: Leaving Station 55

Trawl # 6

Water: Salt Lat. $41^{\circ} 18.0'$ N. Long. $168^{\circ} 21.1' E$ Vegetation: $41^{\circ} 05.5'$ To $168^{\circ} 12.0' E$ $41^{\circ} 11.7' N$ $168^{\circ} 16.5' E$

Bottom:

Cover: Temp:

Shore: Current:

Dist. offshore: Stream width:

Depth of capture: 2000-3000(?) fms Depth of water: 2950 fms

Collected by R. K. Wisnet et al

Tide: Date: 10: IX: 53

Method of capture: 10' Mid water trawl

Orig. preserv.: 10% formalin Time: 0849-1935

9. Gadylagus pacificus Gilbert 1HG(53)10. Argyroleucus ~~hemigymnus~~ ~~Gilbert~~ ~~intermedia~~ 1HG(18)Cyclathene atraria Gilbert 25Yq-Ad(19-55)C. pallida Brauer 1Ad(50)C. signata Garman 22HG(23-27)10 Cyclathene, ~~sp~~ pseudopallida 26(27-43)10A Ganostoma gracile Günther 58Yq-Ad(73-100)

10 specimens R to UBC, exchange, I: 31:62

Ichthyococcus elongatus Imai 2Yq-HG(17-37)

Ident. by R. K. Wisnet

10B ~~Photichthys~~ Tactostoma macropus 1Ad(290)Sternoptyx obscura Garman 1Ad(42)

2 samples. Cod end closed off before ascent.

OVER

- 10. ~~Physanactis, n. sp.~~
Opostomias mitsuui 1(58)
- 10¹ Vinciguerria nimbaria (Jordan & Williams) 1Ad(27)
GR (total) = nimbaria Upper lateral photophore = lucetia.
- 25. Ceratospelus warmingi
~~townsendi~~ (Eig. and Eig.) 7Yg-H6 (28-46)
Ident. by R. L. Wisner.
- 25A Ditropichthys storeri (Goode and Bean) 1(104)
Ident. Huxley, 1952; Par, 1934; Goode & Bean, 1895.
Agrees well with above descriptions except meristic ~~count~~
Sometime 1 higher (D. 21 (vs. 20); A 18 (vs. 17); GR 4+1+10-3+1+10
(v. 7+1+8)). LL. pinn = 13-green.
Reident. as new species by R. Rosen. II: 2: 1959.
- 25D Cetomimus
~~Cyprinomimus~~, sp. evermanni - 1(34) → 1Yg (56)
25 Symbolophorus ~~sp.~~ Finnesqui
25 Diaphus ~~stagens~~ Brauer 1 ~~stagens~~ (34-)
Ident. by R. L. Wisner
25 Diaphus theta 1(51)
25 Diogenichthys atlanticus (Taning) 1H6(16.5)
- 25 Lampanyctus jordani Gilbert 11Yg-Ad (24-71)
7Yg-Ad (24-70) Rept. by R. L. Wisner III: 19: 1958
- 25 Stenobrachius
~~leucopsarus~~ (Eig. and Eig.) 1Ad(105)
Ident. by R. L. Wisner.
- 25 Stenobrachius
~~nanochir~~ (Gilbert) 4H6-Ad (67-90)
- 25B Benthalbella
Neoscopelarchoides dentata (Chapman) 1H6(57)
Ident. Marshall, 1955; Chapman, 1939
- 25C Benthalbella linguoides
~~diaphanus~~ (Rass) 2Yg (50)⁵⁴
Ident. Rass, 1955.
- 31 Poyomitra crassiceps 2Yg (19-28)
~~Metamphac~~ tuposus Chapman
- 31 Metamphac ~~sp.~~ Scopeloberyx robustus 2Ad (69-70)
Aff. nucterinus
- 65. Gigantactis ~~mesonema~~ savagei 1(39)
- add 61A Bassozetus zenkevitchi Rass 1H6(100)
D. 110; A 191; P 25; P. 1; GR = 4+2+1+1+2. Agrees in most counts with type desc. (Rosen, 1955). GR count not high. Ident. Huxley, 1952; Rosen, 1959; Chapman, 1939.
- 6R. Melanonus, sp. 1(17)

Alfred W. Chelung
II: 4: 1939.