# PATIENT INFORMATION FOR CATARACT SURGERY

# G. RICHARD COHEN, MD

## JOSHUA D. COHEN, MD



BAPTIST HEALTH SURGERY CENTER

1905 CLINT MOORE ROAD, SUITE 115 BOCA RATON, FL 33496 P 561-807-8090



BOCA RATON OUTPATIENT SURGERY AND LASER CENTER

501 GLADES ROAD BOCA RATON, FL 33432 P 561-362-4440

Pre	-Operative Regimen SURGERY DATE: / /							
	🗆 RIGHT EYE 🗆 LEFT EYE							
ONE	<ul> <li>PROLENSA / ILEVRO: 1 drop 1x/day, start 2 days before surgery</li> <li>BROMSITE: 1 drop 2x/day, start 2 days before surgery</li> </ul>							
DWD	<ul> <li>PRN Dry Eye Omega Benefits / Hydroeyes <sup>®</sup></li> <li>Liquid: 1 teaspoon per day</li> <li>Softgels: 4 per day</li> <li>Start Omega 3s as soon as possible</li> </ul>							
<ul> <li>HYPOCHLOR/ AVENOVA *: Use lid cleaner daily or until finished. Start 3 days prior to surgery in both eyes</li> <li>FRESHKOTE* PF: 1 drop in both eyes 3 or 4 times daily for 1 month</li> </ul>								
Day of Surgery Regimen								

- Use one drop of **PROLENSA** or **BROMSITE** when you wake up the morning of surgery.
- Do not eat or drink 6 HOURS before surgery.
- After surgery, take your daily dose of PRN Dry Omega Benefits®

## **General Instructions**

- Follow all of your primary care physician's instructions regarding all medications.
- Notify Dr. Cohen if you are taking Flomax or prostate medications.
- Take any medically necessary medicines with a small sip of water.
- Wash hands before using drops.
- Wait **5 MINUTES** between different drops.
- Store PRN Dry Eye Omega Benefits <sup>®</sup> Liquid in a refrigerator after first use.
- If you currently take other eye drops, please continue to take them as you normally do unless instructed otherwise.

**\*\*VISION MAY BE BLURRY FOR THE FIRST WEEK AFTER SURGERY\*\*** 

Pos	t-Operative Regimen <b>**</b> PLEASE BRING ALL DROPS TO OFFICE VISITS**
ANTIBIOTIC	<ul> <li>PREVENTS INFECTIONS</li> <li>BESIVANCE: One drop 3x/day for 1-2 weeks (as directed)</li> <li>POLYMYXIN: One drop 4x/day for 1-2 weeks (as directed)</li> <li>MOXI/CIPROFLOX: One drop 4x/day for 1-2 weeks</li> </ul>
NSAID	<ul> <li>PREVENTS INFLAMMATION AND PAIN</li> <li>BROMSITE: One drop 2x/day for 1 week, then 1x/day for 3 weeks</li> <li>PROLENSA: One drop 1x/day for 4-6 weeks (as directed)</li> <li>ILEVRO: One drop 1x/day for 4-6 weeks (as directed)</li> </ul>
STEROID	<ul> <li>REGULATES HEALING AND INFLAMMATION</li> <li>FLAREX 0.1%/PREDNISOLONE: One drop 4x/day for 4-6 week taper (as directed)</li> <li>LOTEMEX SM: One drop 3x/day for 4-6 week taper (as directed)</li> <li>DUREZOL: One drop 2-4x/day for 4-6 weeks (as directed)</li> </ul>

DROP START	PRE-OP	DAY OF	WEEK 1	WEEK 2	WEEK 3	WEEK 4
DATE ->						
ANTIBIOTIC		2x	4x	stop		
NSAID	1-2x	2x	1-2x	1x	1x	1x
STEROID		2x	2-4x	2-3x	2x	1x

**\*\*VISION MAY BE BLURRY FOR THE FIRST WEEK AFTER SURGERY\*\*** 

#### Post-Operative Reminders

- DO NOT RUB THE OPERATED EYE.
- Wear eye shield at night for 1 week.
- Avoid heavy lifting, bending, straining, and exercising for 1 week.
- Wear your old glasses if more comfortable. Wear sunglasses or tinted shades while outside during the day if more conformable.
- You may take a full-body shower and shampoo 24 hrs after surgery. Close eyes to avoid getting water in operated eye.
- Avoid swimming and gardening for 2 weeks.
- Do not wear eye makeup for 1 week.
- Most people can resume driving several days after surgery. Use your best judgment.
- Bring all medications to our office on each post-operative visit.
- Continue taking **PRN Dry Eye Omega Benefits** <sup>•</sup> (liquid or softgels) and **FRESHKOTE PF** <sup>°</sup> post-surgery to make sure that eye is producing a healthy tear and to improve quality of vision.
- **Tylenol** may be taken for discomfort. No Advil or aspirin unless approved.

### Normal Post-Operative Symptoms

**Blurred** vision, halos around light, eye tearing, eye redness, and small amount of **mucous** that collects on the eyelids is normal after surgery. A slight **burning** sensation from the drops is also expected. Symptoms should **improve gradually**.

#### THE FOLLOWING SYMPTOMS REQUIRE URGENT ATTENTION:

- Extreme pain or swelling
- Sudden vision loss
- Significant or thick discharge
   Flashing lights or floaters
- COHEN LASER AND VISION CENTER

**Cohen Laser and Vision Center** 3020 N Military Trail, Suite 150

> Boca Raton, FL 33431 P 561-981-8400 www.cohenlaser.com