

Isolation and cross-species characterization of polymorphic microsatellites for the orchid bee *Eulaema meriana* (Hymenoptera: Apidae: Euglossini)

Margarita M. López-Uribe · Amy N. Green ·
Santiago R. Ramírez · Steve M. Bogdanowicz ·
Bryan N. Danforth

Received: 1 April 2010/Accepted: 16 April 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract We describe and characterize eight polymorphic microsatellite loci for the orchid bee species *Eulaema meriana*, an abundant species and important pollinator in wet lowland forests in tropical America. We also tested the cross-species amplification of these microsatellite loci in seven other species of the genus *Eulaema*. For *E. meriana*, number of alleles per locus ranged from four to nine and expected heterozygosity ranged from 0.377 to 0.854. Seven out of the eight loci described amplified in all seven other *Eulaema* species. These microsatellite loci will be of practical use for population structure, mating system and inbreeding studies in euglossine bees.

Keywords *Eulaema* · Euglossine bees · Microsatellites · Diploid males

The tribe Euglossini comprises 218 species in 5 genera, including the genus *Eulaema* with 29 described species (Oliveira 2006; Nemésio 2009). Euglossine bees (Hymenoptera: Apidae), commonly known as orchid bees, are charismatic insects characterized by extremely long tongues and shiny iridescent colors (Roubik and Hanson

2004). Orchid bees are abundant in the Neotropics (López-Uribe et al. 2008) and are considered keystone species in lowland forests because of the ecological role that they play as pollinators of orchids (Dressler 1982) and many other flowering plants (Ramírez et al. 2002).

Orchid bees have recently been targets of conservation concern (Zayed 2004). There is evidence demonstrating that the species diversity of euglossine bees is negatively affected by habitat fragmentation (Brosi 2009). In addition, genetic studies using allozyme markers have shown that some orchid bee populations exhibit high frequencies of diploid males indicating high levels of inbreeding and/or low effective population size (Roubik et al. 1996; Zayed 2004; López-Uribe et al. 2007; but see Takahashi et al. 2001). However, a recent study looking for diploid males using microsatellite markers (Souza et al. 2010) found diploid males to be rare in euglossine bee populations suggesting that the high frequency of diploid males previously reported may be the result of technical flaws in the allozyme-based studies. Therefore, the development of microsatellite markers is essential for the study of population structure and conservation genetics of this group of bees. Here, we describe and characterize eight polymorphic microsatellite loci in *Eulaema meriana*, and tested these loci across seven other *Eulaema* species.

A genomic DNA library enriched for 12 microsatellite repeat motifs was created from one individual of *E. meriana* using a universal linker and ligation procedure (Hamilton et al. 1999; Grant and Bogdanowicz 2006). Transformed bacterial colonies were then screened for microsatellites through hybridization to ^{33}P -radiolabeled oligonucleotides. More than 800 positive clones were obtained from this method and ~200 of them were sequenced with universal M13 primers that flank the cloned insert for microsatellite primer design. PCR primer

M. M. López-Uribe (✉) · A. N. Green · B. N. Danforth
Department of Entomology, Cornell University, Ithaca,
NY 14853, USA
e-mail: mml82@cornell.edu

S. R. Ramírez
Department of Environmental Sciences, University of California
Berkeley, Berkeley, CA 94720, USA

S. M. Bogdanowicz
Evolutionary Genetics Core Facilities, Department of Ecology
and Evolutionary Biology, Cornell University, Ithaca,
NY 14853, USA

Table 1 Primer sequences, repeat motif and annealing temperature (T_a) of eight microsatellite loci isolated from *Eulaema meriana*

Locus	Primer sequence (5' → 3')	Repeat motif	T_a	Allele size (bp)	N_A	H_E	GenBank accession no.
EM8	F: CAG CGT CGC GAT TGG TTC TAC A R: TCA GCT TTG TCA CCG GCA CTG T	(GA) ₁₄	55	301–317	6	0.723	GU997087
EM13	F: GGC GCA ATG ACT AAG GGA ACG R: CCC ACG GGC TAA CGA TGT ATC TT	(TGC) ₇	55	173–185	4	0.637	GU997088
EM16	F: AGC GCA ATT ACA TAT GCA AAA ACA R: TCC GGT GGT ATC TGA GCA TTA TTC	(CAG) ₆ (CCACAG) ₂	55	190–211	4	0.377	GU997089
EM17	F: GGG CGA CGG CGA AGA TTT R: CGT TGC GCC CGA CTT TAC A	(CTT) ₁₀	55	157–187	6	0.731	GU997090
EM40	F: CGA CGC AGA CGC AGC AAC AG R: CCC GCG GAC TAA ACG ACA ACA CT	(CAA) ₁₀	57	143–164	7	0.722	GU997091
EM70	F: GTA CCA CTG CGA GAG CGA AGA AAA R: CCA GTG GCC CGA AGT AGA AAC A	(AG) ₂ G(AG) ₉ A(AG) ₇ A(AG) ₆	55	280–288	5	0.596	GU997092
EM106	F: GAC GTG GAT GAG CCG CAG AAG AC R: TCC GAC GAT GTA CGA GCA CGA A	(AAG) ₉ GAG(AAG) ₉	55	261–300	9	0.854	GU997093
EM107	F: CGA GCC CCG ACG ACG AAC R: GAC CGG AAC GAG CTG GAT GAA T	(TCT) ₂ TCC(TCT) ₁₂	57	200–221	8	0.841	GU997094

Allele size range, number of alleles (N_A) and expected heterozygosity (H_E) were calculated for the population from La Selva, Costa Rica ($N = 40$ haploid males)

Table 2 Characterization of microsatellite loci isolated from *Eulaema meriana* from four localities and cross-species amplification for seven other species of the genus *Eulaema*

Locus	<i>Eulaema meriana</i> ($N = 55$)	<i>Eulaema cingulata</i> ($N = 15$)	<i>Eulaema bombifloris</i> ($N = 10$)	<i>Eulaema chocoana</i> ($N = 3$)	<i>Eulaema luteola</i> ($N = 3$)	<i>Eulaema mocsaryi</i> ($N = 2$)	<i>Eulaema nigrifacies</i> ($N = 1$)	<i>Eulaema nigrita</i> ($N = 1$)
EM8	8 (301–323)	3 (301–311)	3 (307–313)	3 (309–315)	2 (290–292)	1 (301)	1 (317)	1 (309)
EM13	4 (173–185)	6 (161–185)	5 (173–188)	2 (197–203)	2 (192–198)	1 (173)	1 (164)	1 (182)
EM16	4 (190–211)	2 (199–202)	5 (201–216)	1 (202)	1 (202)	1 (199)	1 (210)	1 (190)
EM17	7 (157–187)	2 (169–184)	5 (178–196)	1 (172)	1 (157)	1 (184)	1 (178)	1 (220)
EM40	9 (140–167)	4 (143–158)	5 (152–173)	1 (147)	2 (158–164)	2 (161–164)	1 (164)	1 (167)
EM70	5 (280–288)	5 (284–294)	2 (261–294)	1 (288)	3 (225–273)	1 (287)	1 (297)	1 (297)
EM106	11 (261–300)	5 (268–295)	7 (261–294)	1 (264)	2 (225–273)	2 (271–283)	1 (282)	1 (305)
EM107	8 (200–221)	4 (196–221)	3 (203–212)	1 (203)	1 (199)	2 (199–202)	1 (202)	—

Above: number of alleles per locus; below: allele range size per locus ($N =$ number of haploid males)

pairs were designed for 29 microsatellite loci using the software PrimerSelect (DNASTAR). Nine of these loci were tested for PCR amplification quality and variability.

For microsatellite PCR amplifications, a universal tag method with three primers was employed (Schuelke 2000).

This approach allows fluorescent labeling of PCR fragments with a single dye-labeled tag used simultaneously with the unlabeled locus-specific (ULS) forward primer containing 20 additional bases at the 5'-end and the ULS reverse primer. The reverse primer was modified by adding

a six base pair ‘pigtail’ (GTTTCT) to the 5'-end (Brownstein et al. 1996) to facilitate genotyping by reducing stutter. PCR amplifications contained 5× GoTaq buffer pH 8.5, 2 mM MgCl₂, 0.2 mM dNTPs, 0.1 μM ULS forward primer, 0.2 μM dye-labeled tag, 0.2 μM ULS reverse primer, 1U GoTaq DNA polymerase (Promega) and 10–50 ng DNA in 20 μl total volume. PCR cycling conditions consisted of one cycle at 94°C for 30 s, 35 cycles at 94°C for 30 s, 45 s at the locus-specific annealing temperature (Table 1) and 45 s at 72°C, followed by one step of 7 min at 72°C. Cycling was carried out using a Biometra TGradient thermal cycler. Labeled PCR products were analyzed on an Applied BioSystems 3730 1× DNA Analyzer using the allele size standard GeneScan-500 LIZ and called using the software PeakScanner (Applied BioSystems).

Genomic DNA was extracted from males of *E. meriana* ($N = 55$), *Eulaema cingulata* ($N = 15$), *Eulaema bombiformis* ($N = 10$), *Eulaema choocoana* ($N = 3$), *Eulaema luteola* ($N = 2$), *Eulaema mocsaryi* ($N = 2$), *Eulaema nigrifacies* ($N = 1$) and *Eulaema nigrita* ($N = 1$) (Table 1) using the QIAGEN DNeasy Tissue kit. Characterization of each locus was based on one *E. meriana* population ($N = 40$) from La Selva, Costa Rica (Table 1). All loci were checked for amplification variability in four *E. meriana* populations and across the other seven *Eulaema* species (Table 2). Due to the haploid nature of the data, tests for Hardy–Weinberg equilibrium and linkage disequilibrium were not performed. Number of alleles per locus (N_A) and expected heterozygosity (H_E) were calculated using Microsatellite Analyser (MSA) (Dieringer and Schlotterer 2003).

The number of alleles per locus for *E. meriana* ranged from 4 to 9 in the population from La Selva (Costa Rica) (Table 1) and from 4 to 11 when including individuals from the other 3 populations analyzed (Table 2). Null alleles were only detected for the locus EM40 in one *E. meriana* individual. All microsatellite loci were easily genotyped in all species except for locus EM107 in *E. luteola* and *E. nigrita*. Stutter was only evident in locus EM70 for *E. luteola*. None of the 90 individuals analyzed showed a diploid genotype. Successful cross-species amplification of these loci shows that the microsatellite markers here described will be useful tools for future population and conservation genetic studies in *E. meriana* and several species of the genus *Eulaema*.

Acknowledgments We would like to thank R. Paxton, A. Soro and S. Cardinal for their comments on the manuscript. This work was supported by the Griswold Endowment, the Organization of Tropical Studies and the Lewis and Clark Fund for Exploration.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Brosi B (2009) The effects of forest fragmentation on euglossine bee communities (Hymenoptera: Apidae: Euglossini). Biol Conserv 142:414–423
- Brownstein MJ, Carpten JD, Smith JR (1996) Modulation of non-templated nucleotide addition by Taq DNA Polymerase: primer modifications that facilitate genotyping. BioTechniques 20:1004–1010
- Dieringer D, Schlotterer C (2003) MICROSATELLITE ANALYSER (MSA): a platform independent analysis tool for large microsatellite data sets. Mol Ecol Notes 3:167–169
- Dressler RL (1982) Biology of the orchid bees (Euglossini). Annu Rev Ecol Syst 13:373–392
- Grant J, Bogdanowicz S (2006) Isolation and characterization of microsatellite markers from the panic moth, *Saucrobotys futilalis* L. (Lepidoptera: Pyralidae: Pyraustinae). Mol Ecol Notes 6:353–355
- Hamilton MB, Pincus EL, Di Fiore A, Fleischer RC (1999) Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites. BioTechniques 27:500–506
- López-Uribe MM, Almanza MT, Ordóñez M (2007) Diploid male frequencies in Colombian populations of euglossine bees (Hymenoptera: Apidae: Euglossini). Biotropica 39:660–662
- López-Uribe MM, Oi CA, Del Lama MA (2008) Nectar-foraging behavior of euglossine bees (Hymenoptera: Apidae) in urban areas. Apidologie 39:410–418
- Nemésio A (2009) Orchid bees (Hymenoptera: Apidae) of the Brazilian Atlantic forest. Zootaxa 2041:242
- Oliveira ML (2006) Nova hipótese de relacionamento filogenético entre os gêneros de Euglossini e entre as espécies de *Eulaema* Lepeléter, 1841 (Hymenoptera: Apidae: Euglossini). Acta Amazonica 36:273–286
- Ramírez S, Dressler RL, Ospina M (2002) Abejas euglossinas (Hymenoptera: Apidae) de la región neotropical: listado de especies con notas sobre su biología. Biota Colombiana 3:7–118
- Roubik DW, Hanson PE (2004) Orchid Bees of Tropical America: Biology and Field Guide. Instituto Nacional de Biodiversidade (INBio), Santo Domingo de Heredia
- Roubik DW, Weight LA, Bonilla MA (1996) Population genetics, diploid males, and limits to social evolution of euglossine bees. Evolution 50:931–935
- Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. Nat Biotechnol 18:233–234
- Souza RO, Del Lama MA, Cervini C, Mortari N, Eltz T, Zimmermann Y, Bach C, Brosi BJ, Suni S, Quezada-Euán JJG, and Paxton RJ (2010) Conservation genetics of Neotropical pollinators revisited: microsatellite analysis demonstrates that diploid males are rare in orchid bees. Evolution. doi:10.1111/j.1558-5646.2010.01052.x
- Takahashi NC, Peruquetti RC, Del Lama MA, Campos LAO (2001) A reanalysis of diploid male frequencies in euglossine bees (Hymenoptera: Apidae). Evolution 55:1897–1899
- Zayed A (2004) Effective population size in Hymenoptera with complementary sex determination. Heredity 93:627–630