

## *Ozognathus cornutus* (LeConte) – new record for the Palaearctic Region (Coleoptera: Anobiidae)

Petr ZAHRADNÍK <sup>1)</sup> & David MIFSUD <sup>2)</sup>

- <sup>1)</sup> Forestry and Game Management Research Institute Jiloviště-Strnady,  
CZ-156 04 Praha 5 – Zbraslav, Czech Republic; e-mail: zahradnik@vulhm.cz  
<sup>2)</sup> Ministry for Rural Affairs and the Environment, Department of Plant Health,  
Agricultural Research & Development Centre, Ghammieri, Marsa CMR 01, Malta;  
e-mail: david.a.mifsud@gov.mt

**Distribution, Coleoptera, Anobiidae, *Ozognathus*, Palaearctic Region**

**Abstract.** The occurrence of *Ozognathus cornutus* (LeConte, 1859) in the Mediterranean Region (Madeira Is., Malta and Tunisia) is documented for the first time. The genus is new for the Palaearctic Region, where it was only previously recorded from the Nearctic and Neotropical Regions.

### INTRODUCTION

The genus *Ozognathus* (Anobiidae, Ernobiinae, Ozognathini) was described by LeConte (1861) for *Anobium cornutum* LeConte, 1859 from California (USA). At that time, the newly erected genus was included in the subfamily Dryophilinae (which subfamily formed part of the family Ptinidae, subfamily Anobiinae, tribe Anobiini, group Dryophili). The genus *Ozognathus* was placed besides the genera *Dryophilus* Chevrolat, 1832, *Xestobium* Motschulsky, 1845 and *Philoxylon* Le Conte, 1861 (= *Ernobius* Thomson, 1859) and all these mentioned genera were included in the subfamily Ernobiinae. Fall (1905) included three North American *Ozognathus* species and later, Pic (1912) grouped together seven species, namely from USA, Mexico, Guatemala and Chile. Later, Pic (1923) described two other species from Chile, and eight year later (Pic, 1931) other species were described from Colombia. Recently, a further species was described from Mexico (White, 1975).

*Ozognathus cornutus* (LeConte, 1859) was described from California (USA) and White (1982) confirmed its occurrence in the mentioned area only. For the first time, the species was found in the Palaearctic Region, first in Madeira, then Malta and later in Tunisia. It is clear, that these findings are not only adventitious, but that this species is adventive, and is able to acclimatize perfectly in the hot parts of the Euro-Mediterranean Region. According to our opinion, it is possible that this species will be found in other Mediterranean territories in a relatively short time. In the literature, there is no information about the biology and early stages of this species. Specimens from Tunisia were reared from non specified dry fruit.

## MATERIAL

### *Ozognathus cornutus* (LeConte, 1859)

(Figs 1-2)

**Material examined.** Madeira, Funchal, 23.-30.xii.1996, 1 ♂, 1 ♀, W. Kronblad lgt., G. Gillerfors coll.; Malta, Marsa, Ghammieri, 20.vi.2004, 1 ♂; 15.x.2004, 2 ♂♂; Marsa, Ghammieri, 23.x.2004, 4 ♂♂, 2 ♀♀; Marsa, Ghammieri, 10.x.2004, 5 ♂♂; Marsa, Ghammieri, 10-30.iii.2005, 7 ♂♂ (all specimens were collected by the window sill in an office building); Tal-Munxar (St. Thomas Bay), 5.i.2003, 1 ♂ (coastal, in decaying seaweed); Zejtun, 17.x.2004, 1 ♂, all D. Mifsud lgt., D. Mifsud and P. Zahradnik coll.; Tunisia, Djerba I., Aghir env., 20.xi.-2.xii.2004, 40 ♂♂, 50 ♀♀, J. Borowski lgt., J. Borowski and P. Zahradnik coll..

**Short description.** *Ozognathus cornutus* (LeConte, 1859) is a small species, 1.5-2.8 mm long. Dorsal surface is blackish brown to black, with short dense adjacent pubescence. It is a species with distinct sexual dimorphism, with males having a long horn arising from each mandible whereas this is completely absent in females. Angles between both horns, between horn and head and pronotum, and form and curvature of horns show extreme variability in the material examined.

## CATALOGUE OF THE WORLD SPECIES GENUS *OZOGNATHUS*

*Ozognathus* LeConte, 1861

*Durangoum* Pic, 1903

*Micranobium* Gorham, 1883



Fig. 1: Lateral view a) – male, b) – female.

Fig. 2: Aedeagus (scale 0.1 mm).

*cornutus* (LeConte, 1859) – Madeira Is., Malta, Tunisia (all three introduced), USA (California)  
*missellus* LeConte, 1865  
*dubius* Fall, 1905 – USA (Arizona, Colorado)  
*elongatus* Pic, 1923 – Chile  
*exiguus* (Gorham, 1883) – Grenada, Guatemala, St. Vincent I.  
*floridanus* LeConte, 1878 – USA (Florida, Kentucky, New Jersey, Texas, Virginia)  
*grossus* White, 1975 – Mexico  
*hirsutus* Pic, 1923 – Chile  
*inarmatus* Pic, 1904 – Chile  
*mexicanus* (Pic, 1903) – Mexico  
*rufescens* Pic, 1904 – Chile  
*seileri* Pic, 1931 – Colombia

ACKNOWLEDGEMENTS. We wish to thank Mr. Gösta Gillerfors (Varberg, Sweden) and Mr. Jerzy Borowski (Rogow, Poland) for the kind loan of the material.

#### REFERENCES

- FALL H. C. 1905: Revision of the Ptinidae of Boreal America. *Transaction of the American Entomological Society* 31: 97-296.  
LECONTE J. L. 1861: *Classification of the Coleoptera of North America. Part I.* Washington: Smithsonian Institute, 214 pp.  
PIC M. 1912: *Coleopterorum Catalogus. Pars 48 - Anobiidae.* In: Junk W. & Schenkling S. (eds.): *Coleopterorum Catalogus.* Berlin: W. Junk, 92 pp.  
PIC M. 1923: Nouveautés diverses. *Mélanges Exotico-entomologiques* 38: 1-32.  
PIC M. 1931: Nouveaux Coléoptères de la Colombie. *Bulletin de la Société Entomologique de France* 1931: 192-194.  
WHITE R. E. 1975: Sixteen new neotropical Anobiidae with a new genus and keys (Coleoptera). *Proceedings of the Entomological Society of Washington* 77: 169-188.  
WHITE R. E. 1982: *A catalog of the Coleoptera of America north of Mexico: Family Anobiidae.* United States Department of Agriculture, Agriculture Handbook Number 529-70: i-xi + 1-58.