

Index to Volume VIII

- Abies bracteata*, 349
religiosa, 348, 349
Abralia astrosticta, 365
Acanthaster, 259
planci, 259
Acanthomysis alaskensis, 131–134, 136
borealis, 133–136
costata, 133–135
davisi, 133–134, 138
mitsukurii, 133
nephrophtalma, 138
pseudomacropsis, 131
sculpta, 129, 131, 133, 136
sculpta var. *nuda*, 138
stelleri, 133–134
Acanthophora, 375
spicifera, 456
Acetabularia, 375
caliculus, 396
moebii, 397
Achras sapota, 181
Acrocarpus capitatus, 424
intricatus, 423
Acrochaetium gracile, 412–414
lauterbachii, 414
occidentale, 412–414
robustum, 412–414
sancti-thomae, 412
sargassicola, 414
subseriatum, 412–414
Acrocystis nana, 376, 461
Actinometra gracilis, 248
maculata, 249
robustispinna, 249
Actinotrichia, 375
fragilis, 416
rigida, 416
ADACHI, MARIAN S. and D. ELMO HARDY:
 Fruit Flies of the Philippines, Indonesia and
 Malaya. Part I. Dacini, 147–204
Adocia cinerea, 337
gellinara, 337
aholehole, 3
 food of, 3–10
Alecto bennetti, 248
palmata, 249
parvicirra, 249
Alphitonia excelsa, 143
ponderosa, 143
Amansia jungermannioides, 461
Amblyceraeus luteus, 219
Americanus giganteus, 344, 345
Americus gigantea, 343, 345, 351
Amphiroa foliacea, 374, 430
fragilissima, 430
Anadyomene, 374, 375, 376
wrightii, 390
Anous minutus marcusi, 89, 92
stolidus, 85, 88
stolidus pileatus, 92
Antecaridina, 368
Antedon protectus, 249
spicata, 249
Antithamnion antillanum, 444
lherminieri, 444
sp., 443
Apodemus agrarius, 12, 14, 16, 17, 18, 22
speciosus, 14, 17, 22
Araeosoma, 243
Archaster typicus, 254
Archeomyysis grebnitzkii, 125, 126
maculata, 126
Ardea purpurea, 89, 93
Areca
catechu, 162, 166
sapida, 37
Arenaria interpres interpres, 93
Argemone alba var. *glauca*, 144
mexicana, 144
Artocarpus dadah, 181
integra, 165, 166, 185, 191
integrifolia, 185
polyphemus, 181, 185
Asparagopsis taxiformis, 414
Asperococcus orientalis, 404
Asterias carinifera, 258
cephea, 258
exigua, 258
laevigata, 258
milleporella, 255
nodosa, 254
planci, 259
Asterina, 258
cepaea, 258
Asterocytis ornata, 367, 411
Astrope, 258
carinifera, 258
Astropyga, 244
Australian Species of *Poecilocryptus* Cameron
(Ichneumonidae: Hymenoptera), 239–242
Averrhoa bilimbi, 166
carambola, 165, 166, 191
Avicennia marina var. *resinifera*, 37
officinalis, 37
resinifera, 37
Avrainvillea, 375, 376
erecta, 395
lacerata, 367
Baccaurea angulata, 165, 166, 191
motleyana, 172
Bacillaria, 6
BANNER, ALBERT H.:
 New Records of Mysidacea and Euphausiacea from
 the Northeastern Pacific and Adjacent Areas,
 125–139
Bassogigas, 79
Benicasa cerifera, 165, 191
BERNER, LEO D.:
 On the Previously Undescribed Aggregate Form of
 the Pelagic Tunicate *Ritteriella picteti* (Apstein)
 (1904), 121–124
Bibliographic Notes on G. Forster's "De plantis
esculentis insularum oceanii australis" (1786),
 35–40
Bidens coimoides, 143
Biphora africana, 300
cofoederata, 317
democratica, 323
depressa, 297
maxima, 300

- mucronata*, 323
pinnata, 283
punctata, 295
tricuspidata, 297
zonaria, 326
- Boergesenia forbesii*, 376, 388
- BOLIN, ROLF L.:
 Report on a Fatal Attack by a Shark, 105-108
- Boedea*, 374
composita, 390
- Boreomysis kincaidi*, 138
nobilis, 125
- Bornetella*, 375
capitata, 396
nitida, 396
oligospora, 396
sphaerica, 396
- Bostrychia radicans*, 452
- Brachytrichia maculans*, 379, 380
quoyi, 379, 380
- Brissus brissus*, 245
latecarinatus, 245, 254
- BROCK, VERNON E.:
 Some Aspects of the Biology of the Aku, *Katsuwonus pelamis*, in the Hawaiian Islands, 94-104
- BROCK, V. E. et al.:
 Fishes Killed by the 1950 Eruption of Mauna Loa
 I. The Origin and Nature of the Collections, 23-27
- Brookzia*, 276, 293
rostrata, 280, 281, 293-295
- Brotula*, 70, 81
multibarbata, 82
- Broussaisia pellucida*, 143
- Bryopsis furcellata*, 395
pennata, 367, 393
- Bulweria bulweri bulweri*, 90
- Caesalpinia Bonduc*, 144
Bonduccella, 144
- Caesaramysis vanclevei*, 138
- Calappa hebatica*, 209-217
- Calidris acuminata*, 93
ferruginea, 90
- Callithamnion*, 375
 sp., 444
- Caloglossa*, 376
adnata, 451
- Calophyllum inophyllum*, 154
- Calothrix aeruginea*, 379
contareni, 379, 380
pilosa, 379
- Canangium odoratum*, 162, 166
- Cancer oculatus*, 126
- Canthium odoratum*, 145
- Capparis sandwichiana*, 145
- Capsicum annuum*, 166
- Carcharodon carcharias*, 108
- Carnavalia ensiformis*, 142
galeata, 142
- Carpopeltis formosana*, 436
- Cataetyx*, 69, 81
hawaiiensis, 69, 71-73, 76-78, 82
laticeps, 76, 78
messieri, 76, 78
rubiostris, 76, 78
- Catenella*, 376
nipae, 443
- Catharacta skua*, 89, 93
- Caulerpa ambigua*, 392
fastigiata, 392
racemosa var. *macrophysa*, 392, 393
serrulata, 367, 374, 392, 393
verticillata, 392
- Caulinia ovalis*, 376
- Centroceras clavatum*, 446
- Ceramiella buysmansii*, 446
- Ceramium*, 375
aduncum, 448
byssoidium, 448
clarionense, 446, 448
clavulatum, 446
cruciatum, 448
equisetoides, 450
fimbriatum, 446
gracillimum var. *byssoidium*, 446, 448
buysmansii, 375, 446
maryae, 446, 448
masonii, 448
mauritianum, 450
mazatlanense, 446, 448
procumbens, 446, 448
taylorii, 375, 446
- Ceratodictyon*, 374
spongiosum, 438
- Cetorhinus maximus*, 108
- Chaetomorpha*, 375
aerea, 386
antennina, 375, 384, 386, 412
crassa, 384, 386
indica, 384, 386
javanica, 384
torulosa, 386
- Chamaecyparis nootkatensis*, 350
- Champia compressa*, 443
parvula, 443, 450
vieillardii, 443
- Charpentiera ovata*, 143
- Chauvinia macrophysa*, 393
- Chondria*, 375
baileyana, 460
dangeardii, 460
obtusa var. *paniculata*, 458
parvula, 443
platyclada, 460
repens, 460
- Chondroclonium cornutum*, 436
- Chnoospora implexa*, 404
obtusangula, 405
pacifica, 375, 398, 405
- Chrysophyllum cainito*, 166
- Cidarites imperialis*, 249
metularia, 249
- Citrus aurantiifolia*, 201
- Cladophora*, 375
albida, 386
anastomosans, 390
composita, 390
inserta var. *ungulata*, 386, 388
mauritiana var. *ungulata*, 388
patentiramea f. *longiarticulata*, 386, 388
perpusilla, 386, 387
- Cladophoropsis herpestica*, 388
membranacea, 390
- Cladostiphon frappieri*, 414
- CLARK, AUSTIN H.:
 Records of Indo-Pacific Echinoderms, 243-263

- Clethrionomys rufocanus*, 12, 17
Clorodesmis hildebrandii, 394
Clunio littoralis, 6, 8
Clypeaster, 252
 latissimus, 252
 reticulatus, 252
Cocos nucifera, 37, 86
 rubra, 37
Codium, 374
 geppii, 395
Colobocentrotus, 244, 252
 atratus, 244, 252
 mertensi, 244, 252
 pedifer, 244, 252
Colpomenia, 374, 375
 sinuosa, 402
Comantheria polycnemis, 248
Comanthus, 248
 bennetti, 248
 parvicirra, 249
 samoana, 249
Comaster, 248
 gracilis, 248
Comatella, 249
 maculata, 249
Compsopogon, 5
Conferva albida, 386
 antennina, 386
 clathrata, 384
 crassa, 386
 herpestica, 390
 intertexta, 452
 membranacea, 390
 ornata, 411
Convolvulus chrysorrhizus, 37
Corallina decussato-dichotoma, 430
 fragillissima, 430
 opuntia, 395
 ungulata f. brevior, 430
Corallopsis opuntia, 438
Cordia subcordata, 142
Cordyline indivisa, 38
Coriaria sarmentosa, 38
Coriopsis cosmoides, 143
Cox, DOAK C.:
 Research in Ground-Water Hydrology in Hawaii, 230-233
Crataeva religiosa, 38
Cucurbita maxima, 165, 191, 194
Culcita, 244, 254
 novae-guineae, 254
Cypressus nootkatensis, 350
Cyclopalpa, 276, 282
 affinis, 280, 281, 284
 bakeri, 277, 280, 281, 283, 286
 dolicosoma-virgula, 286
 floridana, 277, 280, 281, 286, 287
 komaii, 276, 290
 pinnata, 280, 281, 283, 287
 pinnata var. polae, 283
 polae, 283
 retracta, 121, 307, 308
 virgula, 276, 289, 290
Dacini, index to generic and infrageneric categories, 203
Dagysa notata, 314
 strumosa, 314
Dardanus asper, 30, 33
Dasya pedicellata, 451
Dasyopsis pilosa, 451
DAWSON, E. YALE:
 Marine Plants of Nha Trang, 373-469
Decapodous Cephalopod
 Mollusks from the Marshall Islands, 363-366
DELAUBENFELS, M. W.:
 Occurrence of Sponges in an Aquarium, 377-340
Delesseria adnata, 451
Derbesia attenuata, 390
 neglecta, 392
Dermonema frappieri, 375, 414
 gracilis, 414
Development of the Spermatophoric Mass of the Rock Lobster, Parribacus antarcticus (Lund), 28-34
Diadema, 250
 paucispinum, 250
 setosum, 250
Dianella ensifolia, 143
 sandwicensis, 143
Diasterina, 258
 spinulifera, 258
Dichonema erecta, 395
Dictyosphaeria, 374, 376
 australis, 388
 cavernosa, 388
 favulosa, 388
 setchellii, 388
 vanbosseae, 388
 versluyssii, 388
Dictyota dichotoma, 401
 friabilis, 401
 patens, 401
 variegata, 400
Dinemictichthys, 69, 81
Diomedea immutabilis, 87, 90
 nigripes, 85, 87, 90
Dioscorea bulbifera var. *sativa*, 142
 sativa, 142
Diplacanthopoma, 69, 70, 81
 alcocki, 75, 82
 brachysoma, 75, 76, 82
 brunnea, 70, 75, 82
 jordani, 70, 75, 82
 nigripinnis, 75, 82
 raniceps, 75, 82
 rivers-andersoni, 69, 71-76
 sp., 71-73, 75-77
Diplanthera uninervis, 376
*Distribution of the Algal Genera *Rhipilia* and *Sargassum* in the Central Pacific*, 367-368
Dodonaea eriocarpa var. *obtusior*, 145
 viscosa, 145
DOTY, MAXWELL S.:
 Distribution of *Rhipilia* and *Sargassum* in the Pacific, 367-368
Dracaena indivisa, 38
Dracontomelum dao, 165, 174, 191
Dubreuillia cirrhosa, 324
Echinocaulon acerosum, 422
 myriocladium, 422
Echinocyamus, 253
 megapetalus, 253
Echinometra lucunter, 245
 mathaei, 251
 mathaei var. *oblonga*, 251
 oblongus, 251
 setosa, 250

- Echinoneus*, 253
cyclostomus, 253
Echinostrephus, 251
aciculatus, 251
Echinothrix, 250
calamaris, 250
diadema, 250
Echinus atratus, 252
calamaris, 250
diadema, 250
globulus, 250
gratilla, 250
mammillatus, 252
mathaei, 251
ovulum, 253
pedifer, 252
reticulatus, 252
spatangus, 254
trigonarius, 252
Ectocarpus breviarticulatus, 375, 398
indicus, 400
irregularis, 398
mitchellae, 398, 400
Egretta sp., 93
Encoelium clathratum, 403
Enoplomenopus occidentalis, 115-119
Enteromorpha, 375
clathrata, 383, 384
intestinalis, 383
intestinalis var. *tubulosa*, 384
kylinii, 383, 384
prolifera, 384
tubulosa, 383, 384
Entocladia viridis, 388
Entophysalis conferta, 377, 379
Erythrina monosperma, 144
sandwicensis, 144
Erythrotrichia parietalis, 412
Euchaetomeropsis pacifica, 138
Eucidaris, 249
metularia, 249
Eudistyla polymorpha, 59
vancouveri, 59
Eugenia jambos, 166
javanica, 165, 191
malaccensis, 142, 166
Euplana clippertonii, 333
concolor, 333
tropicalis, 331-333
Euprymnea morsei, 364
scolopes, 364
stenodactyla, 363, 364
Eurypleta affinis, 219
Euschöngassia ikaoensis, 12, 14
koreensis, 12, 14-15
Falkenbergia billebrandii, 414
Felisacus amboinæ, 42, 45, 49
auritulus, 49
carpenterae, 49
crassicornis, 48
elegantulus, 41, 42, 45, 47, 49
elegantulus var. *filicicola*, 44
filicicola, 42, 44, 46, 49
glabratus, 49
jacobsoni, 47, 49
longiceps, 48, 49
madagascarensis, 49
magnificus, 42, 48, 49
nigricornis, 47, 49
oehraceus, 42, 46, 49
philippinensis, 42, 48, 49
pulchellus, 48, 49
usingeri, 42, 47, 49
Ferdina, 255
offreti, 255
Fibularia, 253
australis, 253
ovulum, 253
Ficus aspera, 38
granatum, 38
Fishes Killed by the 1950 Eruption of Mauna Loa
I. The Origin and Nature of the Collections,
23-27
Fishes Killed by the 1950 Eruption of Mauna Loa
II. Brotulidae, 68-83
Food of the Aholehole, *Kuhlia sandvicensis*
(Steindachner), in Hawaiian Waters, 3-10
Fosliella farinosa, 425
paschalisi, 425
Fregata aquila, 89
minor minor, 89
Freycinetia arborea, 142
Fromia, 255
balansae, 255
hemipla, 255
millepora, 255
monilis, 255
pacifica, 255
Fucus acerosa, 422
crinalis, 420
filamentosus, 444
filicinus, 432
fragilis, 416
miniatus, 424
papillosum, 458
pusillus, 420
serrulatus, 393
spiciferus, 456
turbanatus var. *ornatus*, 405
valentiae, 436
verrucosus, 438
Gadus macrocephalus, 128, 131, 137
Gabrieliepa brennani var. *ventralis*, 11, 12, 13
saduski, 14
Garcinia dulcis, 158
laeana, 183
Galaxaura, 374
fasciculata, 416, 419
fastigiata, 416, 419
filamentosa, 416, 419
magna, 417
vietnamensis, 374, 416-418
Gelidiella, 375
acerosa, 375, 422
adnata, 422
myrioclada, 422
pannosa, 422
tenuissima, 422
Gelidiopsis intricata, 423
Gelidium crinale, 419, 420
crinale var. *perpusillum*, 421
divaricatum, 420, 421
pannosum, 422
pseudo intricatum, 421
pulchellum, 420, 421
pusillum, 376, 419, 420, 450

- Geranium cuneatum*, 143
bumile var. *kauaiense*, 143
- Gigantabies taxifolia*, 343, 346
Wellingtonia, 344, 346, 347, 356
- Gigartina intermedia*, 443
- Gnetum gnemon*, 177
- Gomophia*, 255
egyptica, 255
- Goniolithon frutescens*, 367
myriocarpum, 428
- Goniotrichum bumphreyi*, 412
- GOSLINE, WILLIAM A.:**
 Fishes Killed by the 1950 Eruption of Mauna Loa
 II. Brotulidae, 68-83
- GOSLINE, W. A. et al.:**
 Fishes Killed by the 1950 Eruption of Mauna Loa
 I. The Origin and Nature of the Collections,
 23-27
- Gossypium tomentosum*, 144
- Gracilaria*, 375
confervoidea, 438
corticata, 439
crassa, 374, 438
eucheumoides, 375, 438
lingula, 439
verrucosa, 438, 439
- Gracilariaopsis rhodotricha*, 439
 sp., 440
- Gratelouphia*, 375
filicina, 432
ramosissima, 432
- Griffithsia argus*, 444
metcalfii, 450
ovalis, 450
tenuis, 450
- Gygis alba*, 88
alba candida, 92
- Gymnogongrus*, 375
flabelliformis, 440
japonicus, 440, 441
pygmaeus, 440
serenei, 440, 441
tenuis, 443
- Gymnophloea gracilis*, 414
- Haliclona permollis*, 337
- Halicystis*, 375
pyriformis, 388
- Halimeda*, 375
gracilis, 396
opuntia, 395
- Halobates*, 8
- Halophila beccarii*, 376
ovalis, 376
ovata, 376
- Halymenia*, 374
dilatata, 433
maculata, 433
- HAND, CADET:**
 Three Pacific Species of "Lar" (Including a New Species), Their Hosts, Medusae, and Relationships (Coelenterata, Hydrozoa), 51-67
- HARDY, D. ELMO and MARIAN S. ADACHI:**
 Fruit Flies of the Philippines, Indonesia and Malaia. Part I. Dacini, 147-204
- Helicosalpa*, 276, 288
komaii, 277, 280, 281, 288, 290-292
virgula, 280, 281, 288-290, 292
- Herposiphonia insidiosa*, 452
tenella, 452
- Heterocentrotus*, 245, 252
mammillatus, 245, 252
trigonarius, 245, 252
- Heteromysis intermedia*, 128
- Hibiscus Arnotianus*, 143
Kokio, 143
tiliaeus, 142
waimeae, 143
Youngianus, 143
- Hildenbrandia*, 376
prototypus, 424
- Hillebrandia sandwicensis*, 143
- Himerometra*, 249
robustispinna, 249
- Homarus americanus*, 115
- Hormothamnion enteromorphoides*, 378, 379
solutum, 378, 379
- Hutchinsia glomerulata*, 452
obscura, 451
tenella, 452
- Hyaloscytus elegantus*, 42
- Hyaloteuthis pelagicus*, 365
- Hydrichthys*, 63
- Hydroclathrus*, 374
clathratus, 403
- Hydrocoleum lyngbyaceum*, 379, 380
- Hydrolithon reinboldii*, 425
- HYMAN, LIBBIE H.:**
 The Polyclad Genus *Pseudoceros*, 219-225
 Polyclad Flatworms from Hawaiian Islands, 331-340
- Hypnea*, 374, 375
boergesenii, 434, 436
cervicornis, 434, 437
charoides, 436
cornuta, 434, 436
esperi, 412, 434, 436
nidulans, 436, 438
valentiae, 434, 436
- Hypoglossum*, 375
attenuatum, 451
spatulatum, 451
- Hypolepis muelleri*, 44
- Iasis*, 276, 326
cylindrica, 305
zonaria, 280, 281, 326-328
- Ichthyocodium*, 63
- Ibla*, 276, 295
asymmetrica, 295-297
magalhanica, 280, 282, 295
punctata, 281, 295-297
- Inocarpus edulis*, 171
- Investigation of Hematochrome Accumulation in the Alga *Phycopeltis hawaiiensis* n. sp., 205-208
- Ipomea*, 86
acetosaefolia, 141
batatas, 37, 142
cairica var. *bederacea*, 143
congesta, 143
insularis, 143
palmata, 143
pes-caprae, 86, 144
stolonifera, 144
- Isurus glaucus*, 108
- JAMESON, E. W. JR. AND SEIICHI TOSHIOKA:**
 Notes on Some Chiggers (Acarina: Trombiculidae) from Southern Korea, 11-27
- Jania capillacea*, 430, 432

- decussato-dichotoma*, 430
longiarthra, 430, 431
ungulata f. *brevior*, 430
- Jasis hexagona*, 313
polycratica, 326
tilesi, 315
zonaria, 326
- Jasus lalandii*, 115
- Katsuwonus pelamis*, 94, 234
- KING, JOHN W.:
 Hematochrome Accumulation in *Phycopeltis hawaiiensis* n. sp., 205-208
- Kotimaea moorei*, 339
tethya, 338
- Kublia sandvicensis*
 food of, 3-10
- KURODA, NAGAHISA:
 Report on a Trip to Marcus Island with Notes on the Birds, 84-93
- Kylinia dubosquii*, 412
pulchellum, 412
- Laganum*, 253
lesueuri, 253
depressum, 253
- Lagenaria leucantha*, 165, 191
vulgaris, 191
- Lamna ditropis*, 108
- Lamprometra*, 249
palmata palmata, 249
- Lar sabellarum*, 51, 64
- Larus argentatus vegae*, 93
- Laurencia*, 375
articulata, 456, 458
baileyana, 460
brachyclados, 458
corymbosa, 456, 458
obtusa var. *densa*, 456, 458
paniculata, 456, 458
papillosa, 456, 458
parvipapillata, 456, 458
pygmaea, 456, 458
tenera, 458
- Lepidium oleraceum*, 38
piscidium, 38
- Leucosolenia eleanor*, 339
- Leveillea jungermannioides*, 461
- Liagora*, 374
ceranoides, 415, 416
divaricata, 415
farinosa, 415
formosana, 415
orientalis, 415
- Linckia guildingii*, 245
laevigata, 258
multifora, 257
pauciforis, 255
pustulata, 256
- Liocoris*, 42
- Lipochaeta australis*, 144
connata, 144
profusa, 144
- Lithoporella*, 375
pacifica, 428
- Lithophyllum affine*, 427
andrussowii, 427
fasciculatum, 427
kaiserii, 427
okamurae, 427
- pallescens*, 427
racemus, 427
reinboldii, 425
samoense, 427
trichotomum, 426
- Lithothamnion myriocarpum*, 428
trichotomum, 426
- Lithothamnum erubescens* var. *subflabellatum*, 426
- Loligo oualaniensis*, 365
- Lophosiphonia obscura*, 451
subadunca, 452
villum, 451
- Luciobrotula*, 79
- Luffa acutangula*, 152, 165, 191
cylindrica, 165, 191
- Lycium sandwicense*, 144
- Lyngbya*, 375
aestuarii, 379, 380
confervoides, 379, 380
majuscula, 379, 380
- Lysimachia daphnooides*, 143
Hillebrandi, 143
- Maba major*, 38
- Macrozanonia macrocarpa*, 187
- Mangifera foetida*, 181
pinnata, 39
- Marcus Island, Report on a Trip to, 84-93
- Mareta*, 254
ovata, 254
- Marine Plants in the Vicinity of the Institut Océanographique de Nha Trang, Viêt Nam, 373-469
- MATTHEWS, DONALD C.:
 The Development of the Spermatophoric Mass of the Rock Lobster, *Parribacus antarcticus* (Lund), 28-34
- The Origin and Development of the Spermatophoric Mass of a Nephropsid Lobster, *Enoplometopus occidentalis* (Randall), 115-120
- Melobesia farinosa*, 425
pacifica, 428
- MERRILL, E. D.:
 Bibliographic Notes on G. Forster's "De plantis esculentis insularum oceanii australis" (1786), 35-40
- Mesocaris*, 368
- Mesospora*, 374
schmidti, 400
- Mesothamnion caribaeum*, 444
- Mespilia*, 250
globulus, 250
- Messerschmidia argentea*, 86
- Metalia*, 253
dicrania, 254
spatagus, 254
sternalis, 254
- Metcalfina*, 276, 312
hexagona, 281, 282, 312-314
- Metopograpsus messor*, 209-217
- Metrosideros Collina* var. *glaberrima*, 142
polymorpha, 142
- Micranous marcusii*, 87
- Microbrotula*, 69, 81
rubra, 69
- Microtus montebelloi*, 16, 17, 20, 22
 sp., 14
- Misakia typica*, 63, 65
- Momordica charantia*, 165, 185, 191, 198
cochininchinensis, 162, 191, 198

- Monobrachium*, 63
Monospora indica, 450
 MOORE, H. L., et al.:
 Fishes Killed by the 1950 Eruption of Mauna Loa
 I. The Origin and Nature of the Collections,
 23-27
Morinda citrifolia, 142
Musa acuminata, 39
 acuminata, 39, 226
 balfisiana, 39, 226
 coarctata, 39
 coriacea, 39
 corniculata, 40
 dorsata, 40
 exsucca, 40
 fatua, 40
 febi, 39, 226
 granulosa, 40
 lunaris, 40
 mensaria, 40
 papillosa, 40
 paradisiaca, 39, 226
 pumila, 40
 punctata, 40
 purpurascens, 40
 regia, 40
 sapientum, 39, 226
 tetragona, 40
 troglodytarum, 39, 226
Myotis lucifugus, 20
Mysidella americana, 138
Mysis awatschensis, 128
 oculata, 126-128
 rayii, 128
 relicta, 127, 128
Myxicola, 59
Myxocephalus, 70
Nardoa, 255
 frianti, 256
 mollis, 255
 pauciforis, 255
Neobythites, 79
Neogoniolithon myriocarpum, 428
Neomeris, 374
 annulata, 396
 sphaerica, 396
Neomonospora indica, 451
 pedicellata var. *tenuis*, 450
Neomysis awatschensis, 128-130, 135
 franciscorum, 131
 intermedia, 128-130
 isaza, 128
 kadiakensis, 128
 mercedis, 128-130
 nigra, 128
 pseudomacropsis, 131
 rayi, 128, 131, 133, 135
 sculpta, 131
Neothunnus macropterus, 115, 234
Nephelium lappaceum, 181
 New Records of Mysidacea and Euphausiacea from
 the Northeastern Pacific and Adjacent Areas,
 125-139
 Notes on Banana Varieties in Hawaii, 226-229
 Notes on Some Chiggers (Acarina: Trombiculidae)
 from Southern Korea, 11-22
Nudiclava, 63
Numenius phaeopus variegatus, 89, 93
 Objective Estimates of Hawaiian Rainfall, 264-275
- Observations on the Oxygen Consumption of
 Certain Marine Crustacea, 209-218
 Occurrence of Sponges in an Aquarium, 337-340
Oceanodroma castro (or *leucorhoa*), 84
 furcata furcata, 90
 tristami (or *matsudairae*), 84
Ochotona hyperborea, 19
Oculospinus brevis, 76
Ommastrephes oualaniensis, 365
 pelagicus, 365
 On the Distribution of the Big-Eyed Tuna,
 Parathunnus sibi, in the Tropical Eastern
 Pacific Ocean, 234-235
 On the Genus *Felisacus* Distant (Heteroptera;
 Miridae; Bryocorinae), 41-50
 On the Previously Undescribed Aggregate Form of
 the Pelagic Tunicate *Ritteriella picteti* (Apstein)
 (1904), 121-124
Oocystis, 6
Ophiactis, 259
 savignyi, 259
Ophiarachna infernalis, 261
 gorgonia, 261
 spinosa, 261
Ophiarachnella, 261
 infernalis, 261
 gorgonia, 261
Ophiarthrum, 261
 elegans, 261
 pictum, 261
Ophidiaster, 256
 granifer, 256
 perplexus, 256
 pustulatus, 256
 squameus, 257
Ophiocoma, 260
 anaglyptica, 260
 brevipes, 260
 erinaceus, 260
 pica, 260
 picta, 261
 schoenleinii, 260
 scolopendrina, 260
Ophiocomella, 260
 clippertoni, 260
Ophiolepis, 262
 cincta, 262
 imbricata, 262
 savignyi, 259
Ophiomastix, 261
 annulosa, 261
 bispinosa, 261
 mixta, 261
 notabilis, 261
Ophionereis, 259
 porrecta, 259
Ophiopezella, 261
 spinosa, 262
Ophiopterus, 262
 imbricatus, 262
Ophiotrix, 259
 demessa, 259
 longipedata, 259
 picteti, 259
 propinqua, 259
 trilineata, 259
Ophiura, 262
 annulosa, 261
 kinbergi, 262

- longipeda*, 259
scolopendrina, 260
Opius angaleti, 152, 165, 185, 191
fletcheri, 165, 174
incisi, 166, 172, 181
longicaudatus, 147, 165, 166, 174
longicaudatus malayensis, 166, 181
makii, 166, 191
manii, 181
oophilus, 147, 166
skinneri, 174, 181
vandenboschi, 147, 165, 166
- Origin and Development of the Spermatophoric Mass of a Nephropsid Lobster, *Enaplonotopus occidentalis* (Randall), 115-120
- Orthocoea pinnata*, 283
Oscarella tenuis, 339
Oscillatoria margaritifera, 379, 380
nigro-viridis, 379, 380
Ostreaeum reineckei, 396
Othilia, 258
luzonica, 258
Pachycrepoides dubius, 165, 166, 174
Padina, 374
commersonii, 401
Palmella conferta, 379
Panulirus argus, 28, 115
interruptus, 28, 115
penicillatus, 28, 33, 115, 118
Paraplanocera oligoglena, 333
Parasalenia, 251
pöhlii, 251
Parabunus mebachi, 234
sibi, 234, 235
Parribacus antarcticus, 28-34, 115
PARROTT, ARTHUR W.:
Australian Species of *Poecilocryptus*, 239-242
Patiriella, 258
exigua, 258
Pegea, 276, 317
confederata, 278, 280, 282, 317-319
octofora, 317
Pentaceraster, 244
Peronella, 253
lesueuri, 253
- PETTERSSON, HANS:
Results from the Swedish "Albatross" Cruise, 359-362
Peyssonnelia calcea, 425
gunniana, 424
rubra var. *orientalis*, 424
Phaethon lepturus dorothea, 90
rubricaudus rothschildi, 90
Phormidium corium, 379, 380
Photocrypus, 239
Phycopeltis hawaiensis, 205-208
Phyllacanthus, 249
imperialis, 249
Phymodius unguilatus, 209-217
Pinus religiosa, 349
Pisonia grandis, 86
Pittosporum acuminatum, 145
coniferiflorum, 145
glabrum, 145
Planocera crosslandi, 336
hawaiensis, 331
pacifica, 333-336
tridentata, 336
Platypodia granulosa, 209-217
- Plectronia odorata*, 145
Pluvialis dominicus fulvus, 90
Poecilla polynema, 64
Pocockiella variegata, 400
Podophthalmus vigil, 209-217
Poecilocryptus, 239
inflexa, 239, 240
nigripectus, 239, 240
nigromaculatus, 239-241
nigromaculata var. *straminea*, 240, 241
straminea, 240, 241
Poecilopimpla, 239
nigromaculata, 240
Polyclad Genus *Pseudoceros*, with Special Reference to the Indo-Pacific Region, 219-225
Polysiphonia, 375
coacta, 456
forcipata, 453
fragilis, 452
billebrandii, 414
insidiosa, 452
obscura, 451
perpusilla, 451
reptabunda, 452
snyderae, 454
subtilissima, 454
tongatensis, 454
villum, 451
Porolithon aequinoctiale, 368
gardineri, 368
onkodes, 368
Porphyra, 375
crispata, 412
Potamobius fluviatilis, 215
trowbridgii, 115
Prionace glauca, 108
Proboscidacyla circumcincta, 52, 58, 64-66
flavicirrata, 51, 52, 64-66
mutabilis, 66
occidentalis, 51, 52, 64-66
ornata, 65, 66
ornata var. *geminifera*, 65, 66
ornata var. *stolonifera*, 63, 66
polynema, 63
stellata, 59, 64
Proceros concinnus, 220, 221
Procersterna cerulea saxatilis, 92
Protohydra, 62
Protoreaster, 244
nodosus, 254
Pseudobryopsis, 375
bainanensis, 393
mauritiana, 393
mucronata, 393
myura, 393
oahuensis, 393
papillata, 393
parva, 393
Pseudoceros bedfordi, 220
canadensis, 219
concinnus, 220-223
corallophilus, 223
maculosus, 219
splendidus, 219
superbus, 219
Pseudochlorodesmis furcellata, 395
Pseudo potamilla, 52
intermedia, 59, 62
myriops, 56

- neglecta*, 56, 62
ocellata, 56, 59
- Pseudosquilla ciliata*, 209-217
- Pseudoxius caystrus*, 209-217
- Psidium guajava*, 169, 181
- Psychocidaris*, 245
- Pteridium allenii*, 76
- Pteris tremula*, 44
- Pterocladia*, 375
 — *parva*, 421
- Pterodroma brevipes hypoleuca*, 85, 90
- Pugetomyia litoralis*, 126, 127
- Puffinus carneipes*, 84
 — *leucomelas*, 84
 — *nativitatis*, 84, 85, 88, 90
 — *pacificus*, 88
 — *pacificus chlororhynchos*, 90
 — *tenuirostris*, 84
- Pycnocraspedum*, 80, 81
 — *armatum*, 71-73, 77, 80, 82
 — *squamipinne*, 80, 81
- Ralfsia* sp., 400
- Rainfall, Objective Estimates of, 264
- RANDALL, JOHN E., M. TAKATA and P. B. VAN WEEL:
 On the Oxygen Consumption of Marine Crustacea, 209-218
- Rattus bouverii*, 14
 — *norvegicus*, 14, 16, 17, 20, 22
 — *rattus*, 14, 16, 17, 20
 — *rattus alexandrinus*, 86
- Rauvolfia sandwicensis*, 145
- Records of Indo-Pacific Echinoderms, 243-263
- Report on a Fatal Attack by a Shark, 105-108
- Report on a Trip to Marcus Island with Notes on the Birds, 84-93
- Research in Ground-Water Hydrology in Hawaii, 230-233
- Review of Mrs. Sinclair's "Indigenous Flowers of the Hawaiian Islands." Hawaiian Plant Studies 23, 140-146
- Rhipidosiphon javensis*, 395
- Rhipilia geppii*, 367-368
- Rhizoclonium kernerii*, 386
- Rhodomela radicans*, 452
- Rhopalostylis sapida*, 37
- Ritterella*, 276, 306
 — *amboinensis*, 121-124, 278, 281, 282, 306-308
 — *picteti*, 121-124, 281, 282, 306-312
- Rosenvingea*, 375
 — *fastigiata*, 404
 — *nbatrangensis*, 403
 — *orientalis*, 404
- Rubus hawaiiensis*, 143
- Sabella media*, 59
- Saccogaster*, 70
- ST. JOHN, HAROLD:
 Review of "Indigenous Flowers of the Hawaiian Islands," 140-146
- ST. JOHN, HAROLD and ROBERT W. KRAUSS:
 Taxonomy of the Big Tree known as *Sequoia gigantea*, 341-358
- Salpa*, 276, 297
 — *affinis*, 284
 — *africana*, 300
 — *africana-maxima*, 300
 — *amboinensis*, 121, 307, 308
 — *antarctica*, 300
 — *aspera*, 297
- asymmetrica*, 295
bakeri, 287
bicaudata, 317
biensis, 312
bigibbosa, 315
birostratus, 300
caboti, 324
caudata, 283
chamissonis, 285, 300
clostra, 297
coerulea, 305
coerulescens, 305
confoederata, 315
confoederata-bicaudata, 317
confoederata-scutigera, 317
cordiformis, 326
cordiformis-quadratica, 326
cordiformis-zonaria, 326
costata, 314
costata-tilesii, 315
cristata, 283
cyanea, 283
cyanogaster, 323
cylindrica, 276, 304
democratica, 323
democratica-mucronata, 324
democratica var. *orientalis*, 324
dolicosoma, 289
dolicosoma-virgula, 289
dolium, 317
echinata, 298
elongata, 305
femoralis, 317
ferruginea, 317
floridana, 286
forskali, 300
fusiformis, 121, 123, 280, 282, 297-300
fusiformis-runcinata, 298
garnotti, 305
gibba, 317
gibbosa, 315
hensenii, 319
hexagona, 276, 312
informis, 317
infundibuliformis, 315
laevis, 317
lineata, 312
maxima, 121, 281, 282, 297, 300-304
microstoma, 326
monotoma, 312
mucosa, 283
mucronata, 323
multitentaculata, 319
multitentaculata-bicristata, 319
musculosa, 295
musculosa-punctata, 295
naepolitana, 300
nephodea, 317
nitida, 326
octophora, 317
pelasgica, 283
picteti, 121, 308
pinnata, 283
pinnata var. *polae*, 283
polycratica, 326
proboscidalis, 283
punctata, 295
pyramidalis, 324
quadrangleularis, 326

- quadrata*, 317
radiata, 319
retracta, 121, 308
rhomboidea, 324
rhomboides, 324
rostrata, 293
runcinata, 297
runcinata-fusiformis, 298
scutigera, 317
scutigera-confoederata, 317
socia, 317
spinosa, 323
thalia, 283
tilesii, 314
tilesii-costata, 315
triangularis, 312
tricuspidata, 326
unicuspidata, 326
vagina, 315
verrucosa, 319
virgula, 288-290
vivipara, 317
zonaria, 326
zonaria-cordiformis, 326
Santalum ellipticum, 143
pyrularium, 143
Sarcocara, 70
Sargassum, 374, 375, 376
crassifolium, 406
mcclurei, 406
polycystum, 406
sandeii, 406
spp., 406-408
Scaevola frutescens var. *sericea*, 144
glabra, 143
kauaiensis, 143
Koenigii, 144
Schizobranchia, 52
insignis, 56
Schubertia? *semperfiriens*, 343
Scutella latissima, 252
Sepia bandensis, 363
papillata, 363
pelagica, 365
tuberculata, 363
Sepiola stenodactyla, 364
Sepioteuthis arctipinnis, 365
lessoniana, 364-365
Sequoia gigantea, 341-357
religiosa, 343
semperfiriens, 341-356
taxifolia, 343
washingtoniana, 344, 353, 356
Wellingtonia, 344, 351, 354-356
Sequoiadendron giganteum, 341, 343, 357
Sesbania grandiflora, 145
tomentosa, 144
SHIMADA, BELL M.:
 Distribution of the Big-Eyed Tuna in the Tropical
 Eastern Pacific, 234-235
Shunennia tarsalis, 12
Sida fallax, 145
Sideroxylon macranthum, 168
SIMMONDS, N. W.:
 Notes on Banana Varieties in Hawaii, 226-229
Smilax sandwicensis, 142
Solanum anthropophagorum, 39
aviculare, 38
incanum, 172
indicum, 172
sarmentosum, 172
uporo, 38
verbascifolium, 166, 172
viride, 38
 Some Aspects of the Biology of the Aku,
Katsuwonous pelamis, in the Hawaiian Islands,
 94-104
 Some Polyclad Flatworms from the Hawaiian
 Islands, 331-340
 Some Results from the Swedish "Albatross" Cruise,
 359-362
Spatangus brissus var. *latecarinatus*, 254
ovatus, 254
sternalis, 253
Sphaerelaria furcigera, 400
novaehollandiae, 400
tribuloides, 400
Sphaerococcus intricatus, 424
pedicellatus, 451
Spondias cytherea, 39
dulcis, 39
pinnata, 39
Spyridia, 375
filamentosa, 444
Steinbaueria gigantea, 344, 345
semperfiriens, 343
Stephanometra, 249
indica protectus, 249
spicata, 249
Stercorarius longicaudus, 85
pomarinus, 85
Sterna anaethetus anaethetus, 91
fuscata, 85, 87, 88
fuscata oahuensis, 91
birundo longipennis, 90
lunata, 90
Stictocardia campanulata, 142
STIDD, CHARLES K.:
 Objective Estimates of Hawaiian Rainfall, 264-275
Strongylodon lucidum, 142
Struvea, 375
anastomosans, 390
 Studies in the Fruit Flies of the Philippine Islands,
 Indonesia, and Malaya. Part I. Dacini
 (Tephritidae-Diptera), 147-204
Stylochoplana inquilina, 331
Sula cyanops, 89
dactylatra personata, 89
leucogaster plotus, 89
piscator, 89
sula rubripes, 89
Symplectoteuthis oualaniensis, 365
Symploca hydnoides, 379, 380
Syntomosphyrum indicum, 165, 166
Taenioma perpusillum, 451
Taenioplana teredini, 331
**TAKATA, M., P. B. VAN WEEL, and JOHN E.
 RANDALL:**
 On the Oxygen Consumption of Marine
 Crustacea, 209-218
Taxodium giganteum, 344, 354
nootkatensis, 349, 350
semperfiriens, 343, 349
Washingtonium, 344, 351, 353-356
 Taxonomic Position and the Scientific Name of the
 Big Tree known as *Sequoia gigantea*, 341-358
 Taxonomy of the Salpidae (Tunicata) of the Central
 Pacific Ocean, 276-330

- Terminalia catappa*, 39
glabrata, 39
Terpios zeteki, 339
- TESTER, ALBERT L., AND SHIRLEY M. TREFZ:
 The Food of the Aholehole, *Kuhlia sandvicensis* (Steindachner), in Hawaiian Waters, 3-10
- Tethya diploderma*, 339
- Tetragonia expansa*, 39
balmifolia, 39
- Tetraplatia*, 62
- Thalasseus bergi cristatus*, 90
- Thalassia hemprichii*, 376, 377
- Thalia*, 276, 323
democratica, 278, 279, 281, 295, 323-326
lingulata, 323
longicauda, 281, 282
mucronata, 324
- Theragra chalcogramma*, 131, 133, 137
- Thespesia populnea*, 142
- Thetys*, 276, 314
vagina, 277, 280, 281, 314-317
- Three Pacific Species of "Lar" (Including a New Species), Their Hosts, Medusae, and Relationships (Coelenterata, Hydrozoa), 51-67
- Thysanoessa inermis*, 137
longipes, 137
raschi, 136-137
- Thysanopoda dubia*, 138
raschi, 136
- Timea parasitica*, 338
squamata, 338
tetractis, 338
xena, 338
- Titanophora*, 374
calcareum, 435
incrustans, 435
mauritianum, 435
pikeana, 435
pulchra, 434
weberi, 435
- Tolypiocladia glomerulata*, 452
- TOSHIOKA, SEIICHI AND E. W. JAMESON, JR.: Notes on Some Chiggers (Acarina: Trombiculidae) from Southern Korea, 11-27
- Toxadocia violacea*, 338
- Traustedtia*, 276, 319
multotentaculata, 280, 282, 319-323
- TREFZ, SHIRLEY M., AND ALBERT L. TESTER: The Food of the Aholehole, *Kuhlia sandvicensis* (Steindachner), in Hawaiian Waters, 3-10
- Tribulus cistoides*, 144
- Trichosanthes anguina*, 165
- Tringa incana incana*, 90
- Tripheustes*, 250
gratilla, 250
- Trombicula* (*Leptotrombidium*)
akamushi, 19
burnsi, 17
hiranumai, 12, 18
insolii, 20
intermedia, 17
lanceolata, 17
murotoensis, 17
myotis, 12, 19
- orientalis*, 12, 16
pallida, 12, 17
palpalis, 12, 14
russica, 20
subakamushi, 12, 20, 21
subintermedia, 12, 17, 18
(Neotrombicula)
japonica, 12, 22
microti, 22
nagayoi, 12, 22
pomeranzevi, 22
sp., 12, 22
tamijai, 12, 20
- Tryblionographa daci*, 165, 166
- Turbinaria*, 374
ornata, 405
- Tylocidaris*, 245
- Udotea*, 375
javensis, 376, 395
- Ulva*, 374
dichotoma, 401
intestinalis, 383
lactuca, 383
sinuosa, 402
- Valonia*, 375
aegagropila, 376, 382, 388
forbesii, 388
ventricosa, 382, 388
- VAN WEEL, P. B., JOHN E. RANDALL, AND M. TAKATA: On the Oxygen Consumption of Marine Crustacea, 209-218
- Vespertilio superans*, 20
- Vigna lutea*, 144
marina, 144
- Vitex trifolia* var. *simplicifolia*, 144
trifolia var. *unifoliata*, 144
- Volcanus*, 78, 81
lineatus, 71-73, 77-80
- VOSS, GILBERT L.: Decapodous Cephalopod Mollusks from the Marshall Islands, 363-366
- Waltheria americana*, 144
- Washingtonia Americana*, 344
Californica, 344, 351, 353-356
- Weelia*, 276, 304
cylindrica, 281, 282, 304-306
- Wellingtonia gigantea*, 343-356
- Willia occidentalis*, 60
pacifica, 65
stellata, 51, 65
- WOODWARD, T. E.: On the Genus *Felisacus* Distant (Heteroptera; Miridae; Bryocorinae), 41-50
- Wrangelia argus*, 444
- Wurdemannia miniata*, 424
- YAMAGUCHI, Y. et al.: Fishes Killed by the 1950 Eruption of Mauna Loa I. The Origin and Nature of the Collections, 23-27
- YOUNT, JAMES L.: The Taxonomy of the Salpidae (Tunicata) of the Central Pacific Ocean, 276-330
- Zenocentrotus*, 244
- Zonaria variegata*, 400
- Zostera uninervis*, 376

