

adansonia

2022 • 44 • 26



An annotated checklist of the
tree species of French Guiana,
including vernacular nomenclature

Jean-François MOLINO, Daniel SABATIER, Pierre GRENAND,
Julien ENGEL, Dawn FRAME, Piero G. DELPRETE, Marie FLEURY,
Guillaume ODONNE, Damien DAVY, Eve J. LUCAS & Claire A. MARTIN

DIRECTEUR DE LA PUBLICATION / *PUBLICATION DIRECTOR*: Bruno David
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Thierry Deroïn

RÉDACTEURS / *EDITORS*: Porter P. Lowry II; Zachary S. Rogers

ASSISTANT DE RÉDACTION / *ASSISTANT EDITOR*: Emmanuel Côtez (adanson@mnhn.fr)

MISE EN PAGE / *PAGE LAYOUT*: Emmanuel Côtez

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

P. Baas (National Herbarium Nederland, Wageningen)
F. Blasco (CNRS, Toulouse)
M. W. Callmander (Conservatoire et Jardin botaniques de la Ville de Genève)
J. A. Doyle (University of California, Davis)
P. K. Endress (Institute of Systematic Botany, Zürich)
P. Feldmann (Cirad, Montpellier)
L. Gautier (Conservatoire et Jardins botaniques de la Ville de Genève)
F. Ghahremaninejad (Kharazmi University, Téhéran)
K. Iwatsuki (Museum of Nature and Human Activities, Hyogo)
A. A. Khapugin (Tyumen State University, Russia)
K. Kubitzki (Institut für Allgemeine Botanik, Hamburg)
J.-Y. Lesouef (Conservatoire botanique de Brest)
P. Morat (Muséum national d'Histoire naturelle, Paris)
J. Munzinger (Institut de Recherche pour le Développement, Montpellier)
S. E. Rakotoarisoa (Millennium Seed Bank, Royal Botanic Gardens Kew, Madagascar Conservation Centre, Antananarivo)
P. H. Raven (Missouri Botanical Garden, St. Louis)
G. Tohmé (Conseil national de la Recherche scientifique Liban, Beyrouth)
J. G. West (Australian National Herbarium, Canberra)
J. R. Wood (Oxford)

COUVERTURE / *COVER*:

Réalisée à partir des Figures de l'article/*Made from the Figures of the article*.

Adansonia est indexé dans / *Adansonia* is indexed in:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Agriculture, Biology, and Environmental Sciences®
- Scopus®

Adansonia est distribué en version électronique par / *Adansonia* is distributed electronically by:

- BioOne® (<http://www.bioone.org>)

Adansonia est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Adansonia is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / The Museum Science Press also publish: *Geodiversitas*, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Cryptogamie sous-sections Algologie*, *Bryologie*, *Mycologie*, *Comptes Rendus Palevol*

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2022
ISSN (imprimé / print): 1280-8571/ ISSN (électronique / electronic): 1639-4798

An annotated checklist of the tree species of French Guiana, including vernacular nomenclature

Jean-François MOLINO
Daniel SABATIER

AMAP, IRD, Université de Montpellier, CIRAD, CNRS, INRAE,
Boulevard de la Lironde, TA A-51/PS2, F-34398 Montpellier cedex 5 (France)
and AMAP, IRD, Herbier de Guyane, B.P. 90165, 97323 Cayenne (French Guiana)
jean-francois.molino@ird.fr (corresponding author)

Pierre GRENAND

IRD, OHM Oyapock, F-97300 Cayenne (French Guiana)

Julien ENGEL
Dawn FRAME
Piero G. DELPRETE

AMAP, IRD, Université de Montpellier, CIRAD, CNRS, INRAE,
Boulevard de la Lironde, TA A-51/PS2, F-34398 Montpellier cedex 5 (France)
and AMAP, IRD, Herbier de Guyane, B.P. 90165, 97323 Cayenne (French Guiana)

Marie FLEURY

Patrimoines locaux, Environnement et Globalisation (PALOC),
MNHN, IRD, B.P. 90165, F-97323 Cayenne (French Guiana)

Guillaume ODONNE
Damien DAVY

LEEISA, CNRS, Université de Guyane, IFREMER, B.P. 70620, F-97334 Cayenne (French Guiana)

Eve J. LUCAS

Royal Botanic Gardens, Kew, Richmond, Surrey (United Kingdom)

Claire A. MARTIN

AMAP, IRD, Université de Montpellier, CIRAD, CNRS, INRAE,
Boulevard de la Lironde, TA A-51/PS2, F-34398 Montpellier cedex 5 (France)

Submitted on 1 September 2021 | accepted on 9 May 2022 | published on 13 December 2022

Molino J.-F., Sabatier D., Grenand P., Engel J., Frame D., Delprete P. G., Fleury M., Odonne G., Davy D., Lucas E. J. & Martin C. A. 2022. — An annotated checklist of the tree species of French Guiana, including vernacular nomenclature. *Adansonia*, sér. 3, 44 (26): 345–903. <https://doi.org/10.5252/adansonia2022v44a26>. <http://adansonia.com/44/26>

ABSTRACT

We present a list of tree species of French Guiana, based on data collected from herbarium specimens and tree inventories, and verified against all available taxonomic literature to date. The list contains 1811 taxa representing 87 families and 421 genera, including 143 unnamed species which either may be species not previously recorded in French Guiana, or taxa new to Science. Original demographic data such as global abundance, local maximum frequency and geographic dispersion, are given for the 1293 species present in tree inventories. Our work has led to the recognition of certain taxonomic and nomenclatural novelties, including a new combination in *Lecythis* Loefl. (Lecythidaceae),

KEY WORDS
 French Guiana,
 trees,
 scientific names,
 vernacular names,
 abundance data,
 lectotypifications,
 new combination,
 new synonyms.

MOTS CLÉS
 Guyane française,
 arbres,
 noms scientifiques,
 noms vernaculaires,
 données d'abondance,
 lectotypifications,
 combinaison nouvelle,
 synonymes nouveaux.

several lectotypifications and new synonyms, as well as reinstatement of *Sloanea acutiflora* Uittien (Elaeocarpaceae), *Eugenia sinemariensis* Aubl. (Myrtaceae), and four species of *Inga* (Leguminosae). In addition, we provide the most comprehensive list available to date of corresponding vernacular names (4354 for 1157 species) in nine languages spoken in French Guiana: Palikur, Kali'na, Teko, Wayápi, Wayana, Nengee tongo (Aluku/Ndjuka/Paramaka), Creole, French, and Brasilian Portuguese. The checklist is complemented by indexes of exsiccata, vernacular names and scientific names. The introductory part is available in French and English.

RÉSUMÉ

Catalogue annoté des espèces d'arbres de Guyane française, avec la nomenclature vernaculaire.

Nous présentons une liste des espèces d'arbres de Guyane française, basée sur des données issues de spécimens d'herbiers et d'inventaires d'arbres. Les noms ont été vérifiés au regard de toute la littérature taxonomique disponible à ce jour. La liste contient 1811 taxons représentant 87 familles et 421 genres, dont 143 espèces non nommées qui peuvent être soit des espèces non répertoriées précédemment en Guyane française, soit des taxons nouveaux pour la Science. Des données démographiques originales telles que l'abondance globale, la fréquence maximale locale et la dispersion géographique, sont données pour les 1293 espèces présentes dans les inventaires d'arbres. Nos travaux ont conduit à la reconnaissance de quelques nouveautés taxonomiques et nomenclaturales, dont une nouvelle combinaison dans le genre *Lecythis* Loefl. (Lecythidaceae), plusieurs lectotypifications et nouveaux synonymes, ainsi que la réinstallation de *Sloanea acutiflora* Uittien (Elaeocarpaceae), d'*Eugenia sinemariensis* Aubl. (Myrtaceae), et de quatre espèces du genre *Inga* (Leguminosae). En outre, nous fournissons la liste la plus complète disponible à ce jour des noms vernaculaires correspondants (4354 pour 1157 espèces) dans neuf langues parlées en Guyane française : Palikur, Kali'na, Teko, Wayápi, Wayana, Nengee tongo (Aluku/Ndjuka/Paramaka), Créole, Français, et Portugais du Brésil. Le catalogue est complété par des index des exsiccata, des noms vernaculaires et des noms scientifiques. La partie introductive est proposée en français et en anglais.

PRESENTATION

Amazonian forests are considered hyper-diverse (ter Steege *et al.* 2020) and perhaps the most visible expression of this is their extreme tree species richness. A first correlate of this great diversity is that most species are very rare, and therefore difficult to observe, collect and describe; a second correlate is the very high variability of the species mix, at all scales of observation, which means that the species composition of a forest area, whatever its size, can only be very imperfectly predicted from that of its surroundings (ter Steege *et al.* 2020). It is thus very likely that vast still unexplored areas of tropical forest are home to many species still unknown to science. So too, it is hardly surprising that the estimation of Amazonian tree species has recently been the subject of heated debate: on the lower end of the range, Cardoso *et al.* (2017) project 6727 species whereas ter Steege *et al.* (2020) estimate up to 15 874 species (trees are defined as having a trunk of 10 cm or more in diameter at 1.3 m from the ground). At the root of these numerical discrepancies lies a lack of basic data, rather than methodological differences used in estimation, the later explanation suggested by Cardoso *et al.* (2017).

The case of French Guiana illustrates the variation in completeness of floristic inventory in general, and that of tree species in particular. French Guiana is one of the most botanically explored regions in Amazonia. Yet the

PRÉSENTATION

L'expression la plus visible de l'hyperdiversité des forêts amazoniennes est probablement leur extrême richesse en espèces d'arbres (ter Steege *et al.* 2020). Un premier corollaire de cette grande diversité est que la plupart des espèces sont très rares et donc difficiles à observer, à collecter et à décrire; le second corollaire est la très forte variabilité du mélange d'espèces, à toutes les échelles d'observation, ce qui signifie que la composition en espèces d'une surface de forêt, quelle que soit sa taille, ne peut être que très imparfaitement prédite à partir de la composition des forêts environnantes (ter Steege *et al.* 2020). Il est donc très probable que les vastes zones forestières encore inexplorées abritent de nombreuses espèces inconnues de la science. Dans ce contexte, il n'est guère surprenant que l'estimation du nombre d'espèces d'arbres en Amazonie ait récemment fait l'objet d'un débat animé: d'un côté Cardoso *et al.* (2017) annoncent 6 727 espèces, tandis qu'à l'autre extrême ter Steege *et al.* (2020) prédisent jusqu'à 15 874 espèces (les arbres étant définis comme ayant un tronc d'au moins 10 cm de diamètre à 1,3 m du sol). Ces divergences de chiffres tiennent surtout au manque de données de base, bien plus qu'à des différences dans les méthodes d'estimation, une explication pourtant suggérée par Cardoso *et al.* (2017).

Le cas de la Guyane française illustre bien les lacunes de l'inventaire floristique en général, et de celui des espèces d'arbres en particulier. La Guyane française a beau être l'un des territoires les mieux inventoriés d'Amazonie, les forêts de l'intérieur sont

TABLE 1. — Taxonomic and nomenclatural novelties. The numbers between brackets refer to accepted taxa, as numbered in the checklist/Nouveautés taxonomiques et nomenclaturales. Les numéros entre parenthèses font référence aux taxons acceptés, tels que numérotés dans le catalogue.

LECTOTYPIFICATIONS

- Blondea latifolia* Rich., new lectotype, *J.B. Leblond* 209 (lecto-, P[P02440487]; isolecto-, G[G00104361], P[P02440486, P02440488]) [463]
Byrsonima aerugo Sagot, new lectotype, *P.A. Sagot* 102 (lecto-, P[P05529869]; isolecto-, BM[BM000796094], BR[BR000008497468], GOET[GOET007083, GOET007084], P[P05529871], S[S08-15040]) [989]
Clusia palmicida Rich. ex Planch. & Triana, new lectotype, *L.C. Richard s.n.* (lecto-, P[P01901181]; isolecto-, P01901182) [371]
Elizabetha princeps R.H.Schomb. ex Benth., new lectotype, *R.H. Schomburgk s.n.* (lecto-, K[K000264886]; isolecto-, K[K000264887, K000264888]) [907]
Endlicheria grandis Mez, comment and amendment [666]
Gardenia integra A.Rich., new lectotype, *L.C. Richard s.n.* (lecto-, P[P00836504]) [1481]
Guarea megantha A.Juss., new lectotype, *J. Martin s.n.* (lecto-, P[P02288028]; isolecto-, P[P02288029, P02288030, P02288031, P02288032]) [1123]
Inga cayennensis Sagot ex Benth., comment and amendment [821]
Licania canescens Benoit, new lectotype, *Mélinon* 13 (lecto-, P[P00745971]) [322]
Moquilea licaniflora Sagot, new lectotype, *J. Martin s.n.* (lecto-, P[P00746011]; isolecto-, P[P00746012, P00746013], K[K000220643]) [360]
Moronobea coccinea Aubl., new lectotype, *J.B. Aublet s.n.* (lecto-, BM[BM000611835]) [378]
Ocotea rubra Mez, new lectotype, *Mélinon s.n.* (lecto-, P[P00711131]; isolecto-, B[B100244360, B100244361], P[P00711132, P00711133]) [673]

- Pourouma minor* Benoit, new lectotype, *R. Benoit* 960 (lecto-, P[P00756800]; isolecto-, P[P00756799]) [1753]

- Tariri guianensis* Aubl., new lectotype, *J.B. Aublet s.n.* (lecto-, P-JU[P00678733]; isolecto-, fide Howard [1983: 287]: BM, P[not seen]) [1395]

NEW COMBINATION

- Lecythis praeclera* (Sandwith) S.A.Mori ex Molino & Sabatier, comb. nov. [727]

REINSTATEMENTS

- Eugenia sinemariensis* Aubl. [1252]
Guarea megantha A.Juss. [1123]
Inga jenmanii Sandwith [835]
Inga longipedunculata Ducke [840]
Inga mitaraka Poncy [845]
Inga sarmentosa Glaz. ex Harms [857]
Sloanea acutiflora Uittien [449]

NEW SYNONYMIES

- Aydendron salicifolium* (Sw.) Nees, syn. nov. [575]
Caraipa latifolia Aubl., syn. nov. [338]
Eugenia coffeifolia DC., syn. nov. [1252]
Guarea grandifolia DC., syn. nov. [1123]
Icica acuminata Poir., syn. nov. [192]
Miconia punctata var. *latifolia* Cogn., syn. nov. [1062]
Moronobea grandiflora Choisy, syn. nov. [378]
Oreodaphne cayennensis Meisn., syn. nov. [625]
Tovomita bahiensis Engl., syn. nov. [389]
Tovomita choisyana Planch. & Triana, syn. nov. [389]
Tovomita excelsa Andrade-Lima & G.Mariz, syn. nov. [389]
Tovomita melinonii Vesque, syn. nov. [389]

forests of the interior are still little explored, and their flora poorly known. Although a first work on the trees of French Guiana was produced as early as 1933 (Benoist 1933), the last complete flora of the territory dates from the 1950's (Lemée 1953) and is obsolete. The *Guide to the Vascular Plants of Central French Guiana* (Mori *et al.* 2002), as well as regional or continental monographs published by the *Flora Neotropica* and *Flora of the Guianas* consortia are irreplaceable sources of information, but until now, there has been no reliable checklist of tree species for French Guiana.

After years of effort, we fill this debilitating scientific lacuna by producing the most comprehensive autochthonous tree species list for French Guiana; not merely a list, this work contains exhaustive synonymies and all known vernacular names, as well as demographic data derived from inventories of over 143 000 trees in 317 plots of varying shapes and size.

encore très peu explorées, et leur flore est toujours mal connue. Bien qu'un premier travail sur les arbres de la Guyane ait été publié dès 1933 (Benoist 1933), la dernière flore complète du territoire remonte aux années 1950 (Lemée 1953) et elle est obsolète. Le *Guide to the Vascular Plants of Central French Guiana* (Mori *et al.* 2002), ainsi que les monographies régionales ou continentales publiées par les consortiums *Flora Neotropica* et *Flora of the Guianas* sont des sources d'informations irremplaçables, mais il n'existe toujours aucune liste fiable des espèces d'arbres de Guyane.

Après des années d'efforts, nous comblons aujourd'hui cette lacune scientifique avec la liste la plus exhaustive possible des espèces d'arbres autochtones de Guyane française. Loin d'être seulement une liste, ce travail présente une synonymie exhaustive de chaque espèce, ainsi que d'autres informations telles que les listes de tous les noms vernaculaires connus et des données démographiques issues de l'inventaire de plus de 143 000 arbres sur 317 relevés botaniques de tailles et de formes variées.

MATERIAL AND METHODS

STUDY AREA

French Guiana is a single territorial collectivity of France located on the north-east Atlantic coast of South America. Together with Guyana, Suriname, and neighbouring parts of Venezuela, Brazil and Colombia, it lies on the Guiana Shield. This geological formation, one of three Precambrian cratons of the South American Plate, is characterised by very eroded reliefs culminating in the west with the tepuis of Venezuela, Guyana and Roraima, and by very poor, acidic soils. Among Amazonian forests, those of the Guiana Shield are characterised by a relatively low tree species diversity and by a dominance of families rich in large-seeded and hardwood tree species (notably Leguminosae, Lecythidaceae and Chrysobalanaceae) (ter Steege *et al.* 2000).

Due to its eastern location on the northern slope of the Guiana Shield, French Guiana has both a low relief (mean altitude ca. 140 m above sea level), with rare summits reaching 800 m above sea level, and a hydrographic network made up of coastal rivers; the most important of these rivers, the Oyapock and the Maroni, mark the borders with Brazil (State of Amapá) and Suriname, respectively. French Guiana's climate is equatorial, having annual rainfall ranging from 2000 mm in the south and extreme northwest to more than 4000 mm in the northeast (Héritier 2011). Equatorial lowland rainforests still cover about 90% of its 83 000 km² area.

Although environmental conditions lack strong gradients and are fairly uniform throughout the territory, French Guayan forest cover is variable, and forests vary both in structure and floristic composition. In particular, there is a marked difference between the forests of the north, which belong to the "Guianan dense forest arch" (Gond *et al.* 2011), and forests of the south. The former have a more closed canopy, are less dynamic and are dominated by Chrysobalanaceae, Lecythidaceae and slow-growing, hard-wooded Leguminosae, whereas the latter are more open and dynamic, comparatively less species-rich and are dominated by Burseraceae, Urticaceae and fast-growing Leguminosae species of the mimosoid clade (Guitet *et al.* 2018). In the extreme south, the forests of the Tumuc-Humac range are just as diverse as those of the north and center, but their composition exhibits a greater affinity to the forests of central Amazonia.

The attractiveness of French Guayan forests for scientific study by botanists and ecologists lies largely in their relatively high level of conservation compared to that found in neighbouring countries. These well-preserved forests are not the result of a political will to protect them, which is real today but relatively recent, but rather by neglect because anthropic pressure on these forests has, at least since the colonization and until recently, always been limited. This relatively low human impact on forests is attributable to overall moderate population density, primarily concentrated in the northern coastal zone. This said, it is also a consequence of French Guiana being a French territorial collectivity and as such benefits by having a higher standard of living than neighbouring countries and imports from overseas a large part of

MATÉRIEL ET MÉTHODES

ZONE D'ÉTUDE

La Guyane est une collectivité territoriale française située sur la côte nord-est de l'Amérique du Sud. De même que le Guyana, le Surinam, et les parties voisines du Venezuela, du Brésil et de la Colombie, elle repose sur un vaste plateau appelé Bouclier Guyanais. Cette formation géologique, qui est un des trois cratons précambriens de la plaque Sud-Américaine, est caractérisée par des reliefs très érodés qui culminent à l'ouest avec les tepuis du Venezuela, du Guyana et du Roraima, et par des sols acides et très pauvres. Parmi les forêts amazoniennes, celles du Bouclier Guyanais se caractérisent par une relativement faible diversité d'espèces d'arbres et par une dominance des familles riches en espèces à grosses graines et à bois dur (en particulier les Leguminosae, les Lecythidaceae et les Chrysobalanaceae) (ter Steege *et al.* 2000).

En raison de sa localisation sur le versant nord-est du Bouclier Guyanais, la Guyane présente à la fois un faible relief (l'altitude moyenne est d'environ 140 m, avec de rares sommets atteignant 800 m), et un réseau hydrographique constitué de fleuves côtiers. Les plus importants de ces fleuves, l'Oyapock et le Maroni, constituent des frontières naturelles avec, respectivement, le Brésil (Etat de l'Amapá) et le Surinam. Le climat de la Guyane est équatorial, avec une pluviométrie annuelle moyenne qui varie de 2000 mm dans le sud et à l'extrême nord-ouest, à plus de 4000 mm dans le nord-est (Héritier 2011). La forêt équatoriale de plaine couvre encore environ 90% des 83 000 km² du territoire.

Malgré l'absence de gradients environnementaux marqués d'un bout à l'autre du territoire, le couvert forestier est hétérogène, et la structure autant que la composition floristique des forêts sont variables. En particulier, on constate des différences marquées entre les forêts du nord, qui appartiennent à "l'arc des forêts denses guyanaises" (Gond *et al.* 2011), et celles du sud. Les premières ont une canopée plus fermée, sont moins dynamiques et sont dominées par les Chrysobalanaceae, les Lecythidaceae et les Leguminosae à croissance lente et à bois dur, tandis que les secondes sont plus ouvertes et plus dynamiques, comparativement moins riches en espèces et sont dominées par les Burseraceae, les Urticaceae et les Leguminosae à croissance rapide du clade mimosoïde (Guitet *et al.* 2018). À l'extrême sud, les forêts de la chaîne des Tumuc-Humac sont tout aussi diversifiées que celles du nord et du centre, mais leur composition montre une plus grande affinité avec les forêts de l'Amazonie centrale.

L'avantage des forêts guyanaises en tant que cadre et/ou objet d'études en botanique et en écologie réside en grande partie dans un niveau de conservation relativement élevé par rapport aux forêts des pays voisins. Cet état de préservation n'est pas le résultat d'une volonté politique (bien que réelle aujourd'hui, celle-ci est assez récente), mais plutôt d'un désintérêt puisque la pression anthropique sur ces forêts a, au moins depuis la colonisation et jusqu'à récemment, toujours été limitée. Ce faible impact humain résulte avant tout d'une densité de population globalement basse, et d'une concentration du peuplement dans la zone côtière nord. Mais c'est aussi



FIG. 1. — French Guiana map, showing the Guiana Shield (source Wikipedia, modified for the article)/Carte de la Guyane française, montrant le Bouclier Guyanais (source Wikipédia, modifiée pour l'article).

its primary needs; this translates into less need to clear forest as a means of subsistence.

Possibly the earliest botanical and ethnobotanical account of French Guiana is Pierre Barrère's little known *Es-*

une conséquence du fait que la Guyane est une collectivité territoriale française, puisqu'à ce titre elle bénéficie d'un niveau de vie plus élevé que les pays voisins et importe une grande partie de ses besoins primaires, ce qui se traduit par un faible

sai sur l'*histoire naturelle de la France Equinoxiale* (Barrère 1741). This French natural historian's book predates that of Aublet's *Histoire des plantes de la Guyane françoise* (Aublet 1775), but has been unjustly overlooked because his plant classification was based on that of Tournefort and was pre-Linnaean. Notwithstanding, in many ways the ethnobotanical information contained therein is much richer than that found in Aublet's work. Following in Barrère's footsteps, Jean Baptiste Fusée-Aublet continued botanical exploration mostly along the coast of French Guiana, culminating in the publication of his flora of the region (Aublet 1775), and as his treatise was post-Linnaean, it has been the source of many generic names.

Exploration continued sporadically until the middle of the twentieth century. From 1950 onwards, collection of material for herbarium specimens accelerated rapidly, continuing until today (Haripersaud *et al.* 2010). Most botanical collections were the product of classic botanical exploration missions, but beginning in the 1970s plot-based tree inventories conducted for community ecology research also contributed substantial amounts of material.

Created in 1965 in Cayenne, French Guiana, the *Herbier IRD de Guyane* (CAY) (Gonzalez *et al.* 2021) houses at least one duplicate of almost all collections made in French Guiana since the 1970s. Older collections are principally held in P, and any duplicates in European (mainly U, K, BM, B, MPU, G) or US-American (mainly NY, MO, US) herbaria. Only a few collections by Aublet in BM, as well as by botanists from the late 18th and early 19th centuries (e.g. Martin, Patris, von Rohr) in BM and G, do not have duplicates in P.

BUILDING THE CHECKLIST

For a species to be included in the list, three criteria had to be met: first, that it is a tree species, or at least that some of its representatives are trees; second, that at least one of its representatives reaches 10 cm dbh or 10 m in height; and third, that it is spontaneous, or even naturalized, in French Guiana.

As there is no consensus on what an individual tree is, nor on what a tree species is, we adopted the most widespread standard in tropical forest tree inventories: a tree is a self-supporting plant having a trunk, which has a diameter at breast height (dbh, i.e. 1.3 m from the ground) ≥ 10 cm. To account for species whose dbh has never been measured, in particular those that have never been found in tree inventories, we set a threshold of 10 m for the total height of the tree. We have therefore accounted for all species of which at least one representative is a tree with dbh ≥ 10 cm and/or with height ≥ 10 m. Therefore, throughout this checklist, dimensions in cm correspond to dbh, those in m to tree heights. When both dimensions are given for the same tree, it is in the form *height m × dbh cm*.

Given this, our list includes some small statured and/or slender species that only occasionally reach 10 cm dbh, in addition to some species that are more often hemi-epiphytes or lianas than real trees rooted in the ground.

The evidence for species size came primarily from actual tree measurements during plot inventories. For each species

recours aux ressources locales (donc à l'espace forestier) pour assurer la subsistance des populations.

Le plus ancien ouvrage de botanique et d'ethnobotanique sur la Guyane française est probablement le méconnu *Essai sur l'histoire naturelle de la France Équinoxiale* de Pierre Barrère (1741). Le travail de ce naturaliste français est antérieur à l'*Histoire des plantes de la Guyane françoise* d'Aublet (1775), mais il a été injustement négligé parce que sa classification des plantes était basée sur celle de Tournefort, et donc pré-linnéenne. Néanmoins, les informations ethnobotaniques qu'il contient sont à bien des égards beaucoup plus riches que celles que l'on trouve dans l'ouvrage d'Aublet. Suivant les traces de Barrère, Jean Baptiste Fusée-Aublet a poursuivi l'exploration botanique, principalement le long de la côte de la Guyane française, aboutissant à la publication de sa flore de la région (Aublet 1775). Son traité étant post-linnéen, il a notamment été la source de nombreux noms de genres botaniques.

L'exploration s'est poursuivie de manière sporadique jusqu'au milieu du vingtième siècle. À partir de 1950, la collecte de matériel d'herbier s'est accélérée rapidement, et s'est poursuivie à un rythme soutenu jusqu'à aujourd'hui (Haripersaud *et al.* 2010). La plupart des collectes botaniques sont le produit de missions classiques d'exploration, mais à partir des années 1970, les inventaires d'arbres sur parcelles menés dans le cadre de recherches en écologie des communautés ont également fourni des quantités substantielles de matériel.

Créé à Cayenne en 1965, l'*Herbier IRD de Guyane* (CAY) héberge au moins un double de la quasi-totalité des collectes réalisées en Guyane depuis les années 1970 (Gonzalez *et al.* 2021). Les collections plus anciennes sont principalement à P, avec des doubles dans d'autres institutions européennes (surtout U, K, BM, B, MPU, G) ou états-unies (notamment NY, MO, US). Seules quelques collectes d'Aublet à BM, ainsi que d'autres de botanistes de la fin du 18^e siècle et du début du 19^e siècle (e.g. Martin, Patris, von Rohr) à BM et G, n'ont pas de double à P.

CONSTRUCTION DE LA LISTE

Nous avons défini trois critères pour qu'une espèce soit incluse dans la liste: premièrement, ce doit être une espèce arborescente, ou a minima qu'une partie de ses représentants soient des arbres; deuxièmement, il faut qu'au moins un de ses représentants ait un tronc de diamètre à hauteur de poitrine (ou dbh, pour *diameter at breast height*, c'est-à-dire à 1,3 m du sol) de 10 cm ou plus, ou atteigne au moins 10 m de hauteur; et troisièmement, qu'elle soit spontanée (ou éventuellement naturalisée) en Guyane française.

En l'absence de consensus sur ce qu'est un arbre ou une espèce d'arbre, nous avons adopté le standard le plus répandu pour les inventaires d'arbres dans les forêts tropicales: un arbre est une plante autoportante avec un tronc, lequel a un dbh ≥ 10 cm. Pour prendre en compte les espèces pour lesquelle le dbh n'a jamais été mesuré, en particulier celles qui n'ont pas été rencontrées dans les inventaires d'arbres, nous avons fixé un minimum de 10 m pour la hauteur totale de l'arbre. Nous avons donc pris en compte toutes les espèces dont au moins un représentant est un arbre de dbh ≥ 10 cm et/ou de



Fig. 2. — Ethnobotanical research: **A**, survey in the Wayapi village of Zidock (Trois-Sauts) in 1980, from right to left: Raymond Lassouka, Gaétan Lassouka, Gérard Kwanu, Pierre Grenand and Françoise Grenand; **B**, drying of herbarium specimens in the Palikur village of Payuyu, Crique Gabaret, in 1979; the fruits on the right are those of *Manicaria saccifera* Gaertn.; **C**, P. Grenand preparing herbariums in the village of Zidock in 1980; **D**, drying of fruits on a traditional home barbacot ("boucan") for the carpotheque of CAY; village Zidock, 1978/Recherches ethnobotaniques: **A**, enquête dans le village Wayapi de Zidock (Trois-Sauts) en 1980, de droite à gauche: Raymond Lassouka, Gaétan Lassouka, Gérard Kwanu, Pierre Grenand et Françoise Grenand; **B**, séchage des spécimens d'herbier au village Palikur de Payuyu, Crique Gabaret, en 1979; les fruits à droite sont ceux de *Manicaria saccifera* Gaertn.; **C**, P. Grenand préparant des herbiers au village Zidock en 1980; **D**, séchage de fruits sur un boucan domestique pour la carpothèque de CAY; village Zidock, 1978. A, C, © M.-F. Prévost/IRD; B, D, © J.-M. Beaudet/Univ. Paris Nanterre.

encountered in those censuses, we indicate the maximum dbh measured, as dbh_{inv} . It must be emphasised that dbh_{inv} is by no means the actual maximum dbh of the species. Indeed, for many species the representatives present in our plots are far from being the largest known. For species not found in

hauteur ≥ 10 m. Ainsi, les dimensions en cm qui apparaissent dans le catalogue correspondent à des dbh, tandis que celles en m concernent des hauteurs. Quand les deux dimensions sont indiquées pour le même arbre, c'est sous la forme hauteur m \times dbh cm.

our inventories, evidence for species size is provided either in the “Herbarium data (FG)” section, or in the “Size” section when it was recorded from field notes on herbarium specimens from outside FG, or reliable published sources.

The best evidence for presence in French Guiana is that original material of the species or one of its synonyms has been collected there. Failing that, a French Guianan collection reliably identified is cited as an example (preceded by “Sel. exs.:”), in the “Herbarium data (FG)” section. The *Checklist of the Plants of the Guiana Shield* often served as a starting point for clues to likely presence in FG, although this essential volume does not designate plant habit (Funk *et al.* 2007; Feuillet 2009); notwithstanding, we independently verified all species by the aforementioned means.

Given that approximately 90% of French Guiana’s land area is covered by forests (including *terra firme*, secondary, swamp, riverine and mangrove forests), it is no surprise that forest species make up the overwhelming majority of taxa in this checklist. Notwithstanding, there are also tree species found exclusively in non-forest environments (savannas, sandy shorelines, rocky outcrops), and a few others that have been introduced and are now naturalised and sometimes even invasive.

SYSTEMATICS, TAXONOMY AND NOMENCLATURE

In most cases, we have adopted the species delimitations of the specialists of the group concerned, as they appear in recent literature, but also as they are reflected in the determinants of the specimens, especially for genera and families under revision. With respect to groups for which there was no recent or ongoing revision (i.e. no active specialist), we complemented published taxonomic works by informed use of the following online databases: IPNI (2022), Tropicos (2022), the World Checklist of Selected Plant Families (WCSP 2022), The Plant List (2013), Plants Of the World Online (POWO 2022) and for Leguminosae the International Legume Database & Information Service (ILDIS 2018). The familial classification of genera follows APG IV (Angiosperm Phylogeny Group 2016) except for the recognition of Cordiaceae (Stevens 2019), and for the choice of Leguminosae against the alternative name Fabaceae (LPWG *et al.* 2017).

In a few cases, our own studies of herbarium specimens and field knowledge have led us to conclusions that differ from those of previous published works. Such discrepancies in nomenclature or systematics are explained in the notes.

Lastly, we found 143 entities that we were unable to associate with any known French Guianan species, or to any species described for the relevant genus or family. These entities are characterised by character combinations that make them unambiguously distinguishable from all known French Guianan species. In our opinion, they are all distinct at the species level, and we treat them as unnamed species. For each, we cite a representative specimen.

To avoid inflation of published binomials, we excluded all wrong interpretations of formerly published names, i.e. those cited in previous publications or databases with an authorship beginning with “*auct. non*”.

En conséquence, notre liste comprend quelques espèces de petite taille qui n’atteignent qu’occasionnellement 10 cm de dbh, ainsi que quelques autres qui sont plus souvent des hémépiphytes ou des lianes que de véritables arbres toujours enracinés dans le sol.

Les preuves que ces espèces ont bien la taille minimale proviennent principalement de véritables mesures de tronc acquises lors de relevés botaniques. Pour chaque espèce rencontrée dans ces inventaires, nous indiquons le dbh maximum mesuré, appelé dbh_{inv} . Il faut souligner que ce dbh_{inv} n’est que rarement le vrai dbh maximum de l’espèce. De fait, pour beaucoup d’espèces, les individus recensés dans nos parcelles sont loin d’être les plus grands que l’on connaisse. Dans le cas des espèces absentes de nos relevés, la preuve qu’elles atteignent la taille limite est fournie soit dans la rubrique “Herbarium data (FG)” (données d’herbier de Guyane), soit dans la rubrique “Size” (taille) quand l’information provient de notes sur des spécimens d’herbier collectés hors de Guyane, ou bien de sources bibliographiques.

La meilleure preuve qu’une espèce est présente sur le territoire est que le matériel original à partir duquel elle (ou un de ses synonymes) a été décrite en provient. À défaut, nous citons en exemple (précédé de “Sel. exs.:”) une collecte de Guyane identifiée de manière sûre, dans la rubrique “Herbarium data (FG)”. La *Checklist of the Plants of the Guiana Shield* a souvent servi de point de départ pour obtenir des indications sur la présence probable en Guyane, bien que cet ouvrage essentiel ne donne pas d’indications sur le port des plantes (Funk *et al.* 2007; Feuillet 2009); néanmoins, nous avons vérifié indépendamment toutes les espèces par les moyens mentionnés ci-dessus.

Étant donné qu’environ 90% de la surface de la Guyane est couverte de forêts (qu’elles soient de terre ferme, ripicoles ou marécageuses, de mangrove, anciennes ou secondaires), il n’est pas surprenant que ce catalogue ne contienne quasiment que des espèces forestières. On y trouve tout de même des espèces qui poussent exclusivement dans des milieux non forestiers (savannes, cordons sableux littoraux, inselbergs), et quelques autres qui ont été introduites et qui sont aujourd’hui naturalisées, voire même parfois invasives.

SYSTÉMATIQUE, TAXONOMIE ET NOMENCLATURE

Dans la plupart des cas, nous avons suivi les spécialistes des groupes concernés dans leur délimitations des espèces, telles qu’elles apparaissent dans les publications récentes mais aussi telles qu’on peut les percevoir dans les *determinavit* qu’ils ont apposés sur des spécimens, notamment pour les genres et les familles en cours de révision. Dans le cas des groupes pour lesquels il n’y a pas de révision récente ni en cours (autrement dit pas de spécialiste en activité), nous avons complété les publications taxonomiques existantes par un usage réfléchi des grandes bases de données en ligne : IPNI (2022), Tropicos (2022), World Checklist of Selected Plant Families (WCSP 2022), The Plant List (2013), Plants Of the World Online (POWO 2022), et pour les Legu-



Fig. 3. — **A, B**, The fruits of yanipa (*Genipa americana* L.) are used by the Wayäpi of Camopi for temporary body paintings; **C**, the wood of singapeetu (*Dicorynia guianensis* Amshoff) is one of most used by the Aluku to build their large monoxylean dugout canoes, here in the village of Sparouine; **D**, traditional Teko carbet roof, village of Kayodé; the frame is made of various species of round wood, the roofing is made of small palms of *Geonoma baculifera* Kunth/A, **B**, les fruits de yanipa (*Genipa americana* L.) sont utilisés par les Wayäpi de Camopi pour réaliser des peintures corporelles temporaires; **C**, le bois du singapeetu (*Dicorynia guianensis* Amshoff) est l'un des plus utilisés par les Aluku pour construire leurs grandes pirogues monoxyles, ici au village Sparouine; **D**, toit de carbet traditionnel Teko, village de Kayodé; la charpente est constituée de diverses essences de bois ronds, la couverture est en palmes naines de *Geonoma baculifera* Kunth. © G. Odonne/CNRS.

Authors of plant names are abbreviated according to Brummitt & Powell (1992) and IPNI (2022), while Herbarium acronyms follow Index Herbariorum (2020). Protoglosses are cited in abridged form following Taxonomic Literature 2 (Stafleu & Cowan 1976-1988; Stafleu & Mennega 1992-2000; Dorr & Nicolson 2008-2009).

minosae l'International Legume Database & Information Service (ILDIS 2018). La classification familiale des genres suit l'APG IV (Angiosperm Phylogeny Group 2016), sauf pour la reconnaissance des Cordiaceae (Stevens 2019), et le choix du nom Leguminosae à la place du nom alternatif Fabaceae (LPWG *et al.* 2017).

SOURCES OF INFORMATION

Our work is based on three complementary and occasionally overlapping sources of information:

– Herbarium data. This information was used in three ways. First, to derive an indication on species geographic distribution. To this end, we counted the number of collection events in French Guiana from material held at CAY. These values correspond closely to the number of collections made in French Guiana since 1970. Second, we selected one (occasionally more) herbarium collections bearing recent and reliable identifications as evidence that the species is indeed present in French Guiana. Third, for species for which we did not have inventory data (see below), we selected one herbarium collection, not necessarily from French Guiana, having reliable information on tree size, as evidence that the species actually can attain the minimum dbh and/or height thresholds. When specimens were not available for physical examination, digital images of specimens available through herbarium websites or aggregation portals, i.e., JSTOR's Global Plants (JSTOR 2022), e-ReColNat (2022), and iDigBio (2022), were studied. In almost all cases, we have provided hypertext links that allows direct access to the specimen's record and/or scans on such websites.

– Inventory data. The GUYADIV and part of the GUYAFOR plot networks databases (Engel 2015) were the source for our dataset, which includes the identification and trunk measurement of more than 143 000 trees having a dbh \geq 10 cm, censused on 297 old-growth forest plots of varying type and size and in various forest types. Among these plots, 138 are in the range 0.88–1.25 ha, and called hereafter “1-ha plots”. Thousands of herbarium samples were collected during or after these inventories and some of the information from these inventories can also be found in the herbarium data. For each species present in this dataset, we computed four values: the total number of trees censused in our 297 plots; the number of plots where the species is present; F_{\max} , its highest frequency (or relative abundance, in %) per plot, among 1-ha plots; and dbh_{inv} , its maximum dbh in our dataset. Again, dbh_{inv} should not be interpreted as the largest known dbh of the species. It is only relative to our inventory data set. In addition, we pointed out the species that are considered hyperdominant, i.e. they are among the 218 species that together represent more than 50% of all trees present in Amazonia, according to ter Steege *et al.* 2000).

– Taxonomic literature. The Biodiversity Heritage Library (BHL 2022) and the Biblioteca Digital del Real Jardín Botánico (RJB 2022) were invaluable for consulting taxonomic literature (in particular protologues) to which we did not have physical access. Some references were found thanks to more generalist bibliographic sites, in particular Gallica (BNF 2022) and Europeana (2021). Wherever possible, we provide a hyperlink on the protologue citation to an online version of the protologue itself.

Pour quelques taxa, nos propres études des spécimens d'herbier et notre expertise (de terrain et en herbier) nous ont amenés à des conclusions différentes de celles parues dans des publications antérieures. Le cas échéant, ces divergences dans la nomenclature ou la systématique sont expliquées dans la rubrique “Notes”.

Enfin, nous avons identifié 143 entités que nous n'avons pu associer à aucune espèce connue de Guyane, ni à aucune espèce décrite dans le genre ou la famille correspondants. Ces entités présentent des combinaisons de caractères qui permettent de les distinguer sans ambiguïté de toutes les espèces connues de Guyane. De notre point de vue, elles s'en distinguent au niveau spécifique, et nous les considérons donc comme des espèces non nommées. Pour chacune d'elles nous citons un spécimen représentatif.

Afin d'éviter une inflation inutile de binômes, nous avons exclu de la liste des synonymes tous les noms qui ne font que refléter une mauvaise interprétation d'un nom existant, c'est-à-dire ceux dont la paternité commence par “*auct. non*” dans certaines publications ou bases de données.

Les auteurs de noms de taxa sont abrégés selon Brummitt & Powell (1992) et l'IPNI (2022), et les acronymes d'herbiers correspondent à l'Index Herbariorum (2020). Les protologues sont cités sous forme abrégée, conformément à *Taxonomic Literature 2* (Stafleu & Cowan 1976–1988 ; Stafleu & Mennega 1992–2000 ; Dorr & Nicolson 2008–2009).

SOURCES D'INFORMATION

Ce travail repose sur trois sources d'information distinctes et complémentaires, qui se recoupent parfois :

– Les données d'herbier. Elles sont utilisées de trois manières différentes. Premièrement, pour en déduire des indications sur la répartition géographique des espèces. À cette fin, nous avons compté le nombre d'évenements de collecte en Guyane sur la base des spécimens présents à CAY. Les valeurs obtenues sont très proches du nombre réel de collectes en Guyane depuis 1970. Deuxièmement, nous avons sélectionné pour chaque espèce un (parfois plus) spécimen(s) d'herbier portant une identification récente et fiable, en tant que preuve que l'espèce est bien présente en Guyane. Troisièmement, pour les espèces pour lesquelles nous ne disposons pas de données d'inventaire (voir plus loin), nous avons choisi une collecte d'herbier ne provenant pas nécessairement de Guyane mais portant des informations crédibles sur la taille de l'arbre, comme preuve que l'espèce atteint bien le seuil limite de dbh et/ou de hauteur. Lorsque nous n'avons pas eu la possibilité d'examiner les spécimens cités, nous nous sommes basés sur l'étude des images digitales (scans) visibles sur les sites web des herbiers eux-mêmes, ou sur les portails web d'agrégation de données naturalistes, c'est-à-dire JSTOR's Global Plants (JSTOR 2022), e-ReColNat (2022), et iDigBio (2022). Dans presque tous les cas, nous avons fourni des liens hypertextes qui permettent d'accéder directement à la fiche du spécimen et/ou à ses scans sur ces sites web.



FIG. 4. — Some species, whose fruits are highly valued by traditional forest societies, are considered semi-domesticated since pre-Columbian times (Clement 1999; Levis et al. 2017)/Certaines espèces, dont les fruits sont très appréciés des sociétés traditionnelles vivant en milieu forestier, sont considérées comme semi-domestiquées depuis l'époque précolombienne (Clement 1999; Levis et al. 2017). **A**, *Oenocarpus bacaba* Mart.; **B**, *Theobroma cacao* L.; **C**, *Spondias mombin* L.; **D**, *Caryocar villosum* (Aubl.) Pers. A, C, © D. Sabatier/IRD; B, © J.-F. Molino/IRD; D, © G. Odonne/CNRS.

VERNACULAR NAMES

Ethnobotanical lexicons have been recorded and even published for almost all languages spoken in French Guiana. The main Amerindian language missing is Arawak-Lokono, which is rarely spoken today in French Guiana. There are, however, good ethnobotanical nomenclatures of this language in works

— Les données d'inventaires. Les bases de données du réseau de parcelles GUYADIV et une partie de celles du réseau GUYAFOR (Engel 2015) ont été agrégées en un jeu de données unique contenant les identifications et les mesures de dbh de plus de 143 000 arbres de dbh \geq 10 cm, recensés sur 297 relevés de nature et de taille

devoted to Suriname and Guyana (Forte 1996; van Andel 2000; van't Klooster *et al.* 2003). The Maroon languages, Aluku, Ndjuka and Paamaka are very close linguistically and the names tend to pass from one language to another, for these reasons, we chose here to consider them as a homogeneous linguistic whole, and to group them together under the name Nengee tongo. Another Maroon language, Saamaka, is missing from our nomenclature because it is insufficiently studied in French Guiana. However, a good coverage of the botanical lexicon of this language in Suriname appears in van't Klooster *et al.* (2003).

Tree names in Amerindian and Nengee tongo languages presented here have mostly been collected by co-authors PG, MF, GO, DD and their collaborators and represent the fruit of ethnobotanical surveys and botanical inventories involving local knowledge holders. Some of these names have already been published (Fleury 1991 for Nengee tongo; Ogeron *et al.* 2018 for Palikur; Grenand 1989 for Wayápi; Grenand *et al.* 2004 for Wayápi, Palikur and Creole), but many are recorded here for the first time. Moreover, our lists were complemented by other published sources: Roosmalen (1985), Guitet *et al.* (2014) and ONF (2004) for Nengee tongo; Aublet (1775), Ahlbrinck (1931), Kloos (1971), van Andel (2000) and Courtz (2008) for Kali'na; Hurault (1965) and Camargo & Tapinkili (2009) for Wayana. Most of the Creole names were compiled by J.-J. de Granville from CAY specimens labels; this list was subsequently updated by C. Moretti, and later by O. Bruneau. Names in Brazilian Portuguese were compiled on the basis of collective experience and the abundant existing literature, notably Lorenzi (2002), Pio Corrêa (1926-1975), Ribeiro *et al.* (1979, 1999) and Silva *et al.* (1977).

For each language, the number of named species primarily depends on the duration of the ethnobotanical investigation, which ranges from a few years to several decades of survey and varies significantly according to species presence in the ecological zones crossed by the different ethnic groups under study. The tendency of all languages, but particularly of the vehicular languages (Creole, French, Portuguese) to assimilate names from other languages sometimes makes linguistic delimitations complex. Vernacular names also evolve through time, which is usual in languages of oral tradition. They may also vary according to the context and/or the knowledge holder. Identifying the most accurate/consensual name is part of the ethnobiologist's work; we chose to retain only the most agreed upon names among the different locutors and to minimize as much as possible intracultural lexical variations. In this way, we offer readers an accessible panorama of French Guiana's biocultural diversity and to bring the most common names to their attention.

We did not list all orthographical variants that appear on specimen labels because most of them are written according to the orthography of the collectors' native language. Although they are often recognizable to ethnobiologists, they are almost always misspelled and we considered it useless to list them indiscriminately especially considering that collective efforts are being made by Indigenous and local knowledge holders and researchers to properly record these names.

variées, dans des forêts anciennes de différents types. Parmi ces relevés, 138 ont entre 0,88 et 1,25 ha, et sont appelés ici "parcilles de 1 ha" ("1-ha plots"). Des milliers de spécimens d'herbier ont été collectés sur ces relevés, pendant ou après les inventaires, si bien qu'une partie des informations provenant de ces inventaires est aussi présente dans les données d'herbier. Pour chaque espèce présente dans le jeu de données d'inventaires, nous avons calculé quatre indices : le nombre total d'arbres recensés dans nos 297 relevés ; le nombre de relevés dans lesquels l'espèce a été observée ; F_{\max} , la fréquence (ou abondance relative, en %) maximale par parcelle de 1 ha ; et dbh_{inv} , le dbh maximum dans notre jeu de données. Encore une fois, le dbh_{inv} ne doit pas, sauf exception, être interprété comme le dbh maximum connu pour l'espèce. Ce n'est qu'un maximum au sein de notre jeu de données. Par ailleurs, nous indiquons les espèces considérées comme hyperdominantes, c'est-à-dire celles qui font partie des 218 espèces dont les effectifs cumulés représentent plus de 50% de tous les arbres présents en Amazonie selon ter Steege *et al.* (2020).

— La littérature taxonomique. La Biodiversity Heritage Library (BHL 2022) et la Biblioteca Digital del Real Jardín Botánico (RJB 2022) ont été des sources inestimables pour accéder à la littérature taxonomique (en particulier les protologues) à laquelle nous n'avons pas pu avoir accès physiquement. Quelques références ont été trouvées grâce à des sites bibliographiques plus généralistes, en particulier Gallica (BNF 2022) et Europeana (2021). Dans la mesure du possible, nous avons fourni un lien hypertexte sur la citation du protologue vers une version en ligne du protologue lui-même.

NOMS VERNACULAIRES

Des lexiques ethnobotaniques ont été enregistrés et même publiés pour presque toutes les langues parlées en Guyane. La principale langue amérindienne qui n'en dispose pas est l'arawak-lokono, mais elle est aujourd'hui peu parlée en Guyane. Il existe cependant de bonnes nomenclatures ethnobotaniques de cette langue dans les ouvrages consacrés au Suriname et au Guyana (Forte 1996 ; van Andel 2000 ; van't Klooster *et al.* 2003). Les langues noir-marron, l'aluku, le ndjuka et le paamaka, sont très proches linguistiquement et les mots ont tendance à passer d'une langue à l'autre. C'est pourquoi nous avons choisi ici de les considérer comme un ensemble linguistique unique, et de les regrouper sous le nom de nengee tongo. Une autre langue noir-marron, le saamaka, ne fait pas partie de notre nomenclature car elle est trop peu étudiée en Guyane. Un bon recensement du vocabulaire botanique dans cette langue est cependant fourni par van't Klooster *et al.* (2003).

Les noms d'arbres en langues amérindiennes et noir-marron présentés ici ont pour l'essentiel été collectés par les co-auteurs PG, MF, GO, DD et leurs collaborateurs au cours d'enquêtes ethnobotaniques et d'inventaires botaniques auxquels participaient des détenteurs de savoirs locaux.




FIG. 5. — Evolution of the census of tree species in French Guiana. For each work, we counted the number of species among the 1783 of the present list that have been included, regardless of the name used. Species that have been listed under two or more different names were counted once. Aublet (1775): 215 spp.; Lemée (1953-1956): 872 spp.; Sabatier (1993): 1200 spp.; Molino *et al.* (2009): 1592 spp./Évolution du recensement des espèces d’arbres en Guyane française. Pour chaque ouvrage, nous avons compté le nombre d’espèces parmi les 1783 de la liste actuelle qui ont été incluses, quel que soit le nom utilisé. Les espèces qui ont été répertoriées sous deux ou plusieurs noms différents ont été comptées une seule fois. Aublet (1775): 215 spp.; Lemée (1953-1956): 872 espèces; Sabatier (1993): 1200 spp.; Molino *et al.* (2009): 1592 spp.

Spellings used by contemporary ethnobiologists are based on the International Phonetic Alphabet, adapted by input from the people concerned and linguists. Our final spellings were the product of careful deliberation and are valid for French Guiana. Degree and extent of spelling adaptation often depends on the colonial experience of the various peoples, this is particularly true for Creole and Nengee tongo. Faithful language transcription is a form of respect for the speakers of the language and recognises the thoughtful choices they have made. We have also taken into account pronunciation variants, as is the case with Kali’na, which is spoken from Venezuela to Amapá.

The spelling of Palikur names is adapted from Green & Green (1998). That of Kali’na names is the one adopted by the Kali’na community of French Guiana (Renault-Lescure 2009), where we kept the archi-phoneme /k/ (today most often pronounced /g/). Wayana names are spelled according to Camargo & Tapinkili (2009), and Wayápi names according to Grenand (1989). The spelling of Teko names is the one adopted by Maurel *et al.* (2020). Nengee tongo names are spelled according to SIL (2003) and were revised by J. Ateni and C. Ansoe-Tareau. French Guianese Creole names are according to Barthélémi (2007) and Renault-Lescure & Goury (2009), and were revised by F. Grenand and M.-A. Tareau. A few names given by Aublet (1775), mostly Galibi (i.e., Kali’na) and “French” (i.e., Creole in its old meaning) were conserved in their original spelling.

Une partie de ces noms ont déjà été publiés (Fleury 1991 pour le nengee tongo ; Ogeron *et al.* 2018 pour le palikur ; Grenand 1989 pour le wayápi ; Grenand *et al.* 2004 pour le wayápi, le palikur et le créole), mais beaucoup sont publiés ici pour la première fois. Nous avons complété ces listes grâce à d’autres ressources bibliographiques : Roosmalen (1985), Guitet *et al.* (2014) et ONF (2004) pour le nengee tongo ; Aublet (1775), Ahlbrinck (1931), Kloos (1971), van Andel (2000) et Courtz (2008) pour le kali’na ; Hurault (1965) et Camargo & Tapinkili (2009) pour le wayana. La plupart des noms créoles ont été compilés par J.-J. de Granville à partir des étiquettes de spécimens de CAY, et cette liste a été ensuite mise à jour par C. Moretti, puis par O. Bruneau. Les noms en portugais du Brésil ont été compilés à partir de notre expérience collective, ainsi que de l’abondante littérature existante, notamment Lorenzi (2002), Pio Corrêa (1926-1975), Ribeiro *et al.* (1979, 1999) et Silva *et al.* (1977).

Le nombre d’espèces nommées dans chaque langue dépend avant tout de la durée des recherches ethnobotaniques, laquelle varie de quelques années à plusieurs décennies, mais est aussi fonction de la présence des espèces dans les zones écologiques traversées par les groupes ethniques étudiés. La tendance, dans toutes les langues mais particulièrement dans les langues véhiculaires (créole, français, portugais), à assimiler des noms d’autres langues complique parfois les délimitations linguistiques. Les noms vernaculaires peuvent évoluer au cours du temps,

TABLE 2. — Families with 20 or more tree species in French Guiana. For Leguminosae, the number of tree species and genera is given for each subfamily/ Familles comptant 20 espèces d’arbres ou plus en Guyane française. Pour les Leguminosae, le nombre d’espèces et de genres d’arbres est indiqué pour chaque sous-famille.

Family	Nb of genera	Nb of species
Leguminosae, including:		
Caesalpinioideae (non-mimosoid)	9	28
Caesalpinioideae (mimosoid)	18	106
Cercidoideae	1	2
Detarioideae	8	37
Dialioideae	3	3
Papilionoideae	26	75
Lauraceae	14	116
Myrtaceae	9	112
Sapotaceae	10	106
Chrysobalanaceae	12	96
Annonaceae	16	75
Rubiaceae	30	69
Melastomataceae	5	69
Moraceae	13	52
Burseraceae	3	47
Lecythidaceae	7	44
Malvaceae	16	41
Euphorbiaceae	18	40
Sapindaceae	10	38
Clusiaceae	7	37
Apocynaceae	12	35
Elaeocarpaceae	1	32
Meliaceae	5	30
Salicaceae	7	29
Ochnaceae	5	27
Urticaceae	4	27
Vochysiaceae	4	26
Violaceae	6	21
Rutaceae	9	20

Some phonemes used here deserve explanation. In most languages, the phoneme /u/ is pronounced as the “ou” (in “genou”) in French and Creole. The phoneme /i/, specific to most Amerindian languages of French Guiana, is a vowel pronounced between the French “i” (in “liste”) and “u” (in “rue”); it is written “í” in Wayápi, Teko and Kali’na, and “í” in Wayana. In most languages, the phoneme /e/ is always pronounced “é” as in the French “été”; in French and Creole, its pronunciation varies according to the context. In Wayana, /é/ corresponds to “eu” as in the French “beurre”. In Palikur, /h/ is close to the Spanish /j/ in “navaja”; but in Wayana, /h/ is exhaled. Nasal vowels of Amerindian languages are: /é/ pronounced as in the French “peint”, /ó/ as in “bon”, /á/ as in “banc”; /í/ and /ú/ are pronounced from the nose, as in Portuguese “mirim” and “atum”; /í/, an uncommon phoneme of the Wayápi, is the /i/ pronounced from the nose. In contrast the Creole “en” is pronounced “ein” as in the French “sein”. The /y/, as in the German “ya” is mainly written this way, except in Wayana where it is replaced by “j”. The Teko’s /b/ is pronounced “mb”. The /s/ of the Kali’na is pronounced as in the English “short”. Final consonants, like g, k, m, n, p, t, s are pronounced in Teko, Wayana, Palikur, Crémole and Nengee tongo.

Most names listed here are associated with herbarium vouchers, the majority of which being held at CAY. For each

ce qui est habituel dans les langues de tradition orale. Il arrive aussi qu’ils changent en fonction du contexte et/ou du détenteur de savoirs. C’est une partie du travail de l’ethnobiologiste que d’identifier le nom le plus approprié/consensuel. Nous avons choisi de ne retenir que les noms les plus partagés par les différents locuteurs et de minimiser autant que possible les variations lexicales au sein d’une même langue. Nous espérons ainsi offrir aux lecteurs un panorama accessible de la diversité bioculturelle de la Guyane, et attirer leur attention sur les noms communs les plus répandus.

Nous n’avons pas listé tous les variants orthographiques qui apparaissent sur les étiquettes des spécimens d’herbier, car la plupart sont écrits selon l’orthographe de la langue maternelle du collecteur. Bien que ces variants soient souvent identifiables par les ethnobiologistes, il sont presque toujours mal orthographiés, et nous pensons qu’il est inutile de les lister de manière indiscriminée, d’autant plus que de gros efforts collectifs sont faits par les détenteurs de savoirs locaux et les scientifiques pour que ces noms soient enregistrés de manière standardisée.

Les graphies utilisées par les ethnobiologistes contemporains sont basées sur l’alphabet phonétique international, avec des adaptations provenant des populations concernées et des linguistes. Les orthographies présentées ici sont le résultat de choix mûrement réfléchis, et sont valides pour la Guyane française. Le degré et l’étendue des adaptations orthographiques dépend souvent de l’expérience de la colonisation vécue par chaque peuple, notamment dans le cas du créole et du nengee tongo. Transcrire fidèlement une langue est une forme de respect pour ses locuteurs, et une reconnaissance des choix éclairés qu’ils ont faits. Nous avons parfois pris en compte des variations de prononciation, en particulier pour le kali’na qui est parlé du Venezuela à l’Amapá.

La graphie des noms palikur est adaptée de Green & Green (1998). Pour le kali’na, c’est celle qui a été adoptée par la communauté Kali’na de Guyane (Renault-Lescure 2009), dans laquelle nous avons gardé l’archi-phonème /k/ (aujourd’hui le plus souvent prononcé /g/). Les noms wayana sont orthographiés d’après Camargo & Tapinkili (2009), et ceux en wayápi selon Grenand (1989). La graphie des noms tekó est celle adoptée par Maurel *et al.* (2020). Les noms nengee tongo sont orthographiés d’après SIL (2003) et ont été vérifiés par J. Ateni et C. Ansoe-Tareau. Les noms en créole suivent Barthélemy (2007) et Renault-Lescure & Goury (2009), et ont été revus par F. Grenand and M.-A. Tareau. Les quelques noms donnés par Aublet (1775), surtout en galibi (c’est-à-dire en kali’na) et en “français” (ce qui, à l’époque, signifiait en créole) ont été conservés dans leur orthographe d’origine.

Certains phonèmes utilisés ici nécessitent une explication. Dans la plupart des langues, le phonème /u/ est prononcé “ou” comme dans “genou” en français et en créole. Le phonème /i/, présent dans la plupart des langues amérindiennes de Guyane, est une voyelle dont la prononciation est entre le “i” français (dans “liste”) et

species, all vernacular names are grouped in a single paragraph, the languages being separated by a bullet point and designated by the following abbreviations: **Pa**, Palikur • **Ka**, Kali'na • **Te**, Teko • **Wp**, Wayápi • **Wn**, Wayana • **Nt**, Nengee tongo (Aluku/Ndjuka/Paramaka) • **Cr**, Creole • **Fr**, French • **Br**, Brasilian Portuguese. Within each language, names are listed in alphabetical order.

RESULTS AND DISCUSSION

Our list consists of 1811 taxa (i.e. 1783 species, some of which represented by two or more infraspecific taxa) belonging to 422 genera and 87 families, and 7720 synonyms (see taxonomic novelties in Table 1). It includes 11 introduced, naturalised species. Eight families account for nearly 50 % of all species: Leguminosae, Lauraceae, Myrtaceae, Sapotaceae, Chrysobalanaceae, Annonaceae, Melastomataceae and Rubiaceae (Table 2). Among the Leguminosae, the subfamily Caesalpinoideae is by far the richest having 134 species representing 27 genera, which alone is more than any other family in French Guiana. Within Caesalpinoideae, 106 species representing 18 genera belong to the mimosoid clade as defined by LPWG *et al.* (2017). Sixteen families are represented by a single species, and 40 families by five or less (Table 2).

Of the 46 genera that together represent more than 50 % of all tree species, the richest are *Eugenia*, *Inga*, *Pouteria*, *Ocotea*, *Miconia*, *Myrcia*, *Protium*, *Sloanea* and *Licania* (Table 3). Conversely, 178 genera are represented by a single tree species, and 85 by two.

Comparison of our list with previous floristic references for the whole of French Guiana (Fig. 5) illustrates both an acceleration during the last few decades of the rate at which knowledge of the flora has accumulated, and the distance still to be covered to complete the inventory of the flora, of which trees are only one part. The incompleteness of the tree flora inventory is further illustrated by the evolution of the number of unnamed species: this figure has risen from 117 in Molino *et al.* (2009) to 143 today, despite the identification of 52 taxa of the 2009 list and the removal of 10 others based on insufficiency of data (Appendix 6).

Our list provides for the first time demographic data on many species, most of which are very poorly known in this respect. More specifically, total abundance, local maximum frequency, geographic dispersion, are given for the 1293 species that are present in our inventory plots. This information, along with the number of herbarium specimens collected in French Guiana, may be useful for future ecological studies, as well as for assessing the IUCN Red List status of these species (IUCN 2021).

Of the 218 species considered hyperdominant by ter Steege *et al.* (2020), 143 (66%) appear in our list. However, these hyperdominant species in Amazonia are not all abundant in French Guiana: 41 of them are represented by less than 20 individuals in our inventory network. In contrast, only 9 hyperdominant species are among the 20 most abundant species (> 1000 trees) in our dataset.

le "u" (dans "rue") ; il s'écrit "í" en wayápi, en teko et en kali'na, et "í" en wayana. Dans presque toutes les langues, le phonème /e/ est toujours prononcé "é" comme dans le mot français "été" ; en français et en créole, sa prononciation dépend du contexte. En wayana, /ë/ correspond au "eu" du mot français "beurre". En palikur, /h/ est proche de l'espagnol /j/ dans "navaja", mais en wayana, /h/ est un h expiré. Les voyelles nasales des langues amérindiennes sont : /ë/ prononcé comme dans le français "peint", /ó/ comme dans "bon", et /á/ comme dans "banc" ; /í/ et /ú/ sont nasalisés comme dans les mots portugais "mirim" et "atum" ; /í/, un phonème wayápi peu courant, est le /i/ nasalisé. Par ailleurs, le créole "en" est prononcé "ein" comme dans le français "sein". Le /y/, comme dans l'allemand "ya" est presque toujours écrit "y", sauf en wayana où il est remplacé par "j". Le phonème /b/ en teko est prononcé "mb". Le /š/ du kali'na est prononcé comme dans l'anglais "short". Les consonnes finales comme g, k, m, n, p, t, s sont prononcées en teko, wayana, palikur, créole et nengee tongo.

Beaucoup des noms cités ici sont associés à des spécimens d'herbier, pour la plupart déposés à CAY. Pour chaque espèce, tous les noms vernaculaires sont regroupés en un seul paragraphe, les langues étant séparées par une puce et abrégées comme suit: **Pa**, palikur • **Ka**, kali'na • **Te**, teko • **Wp**, wayápi • **Wn**, wayana • **Nt**, nengee tongo (aluku/ndjuka/paramaka) • **Cr**, créole • **Fr**, français • **Br**, portugais du Brésil. Dans chaque langue, les noms sont présentés dans l'ordre alphabétique.

RÉSULTATS ET DISCUSSION

Notre catalogue comprend 1811 taxa (c'est-à-dire 1783 espèces, dont certaines représentées par deux ou plusieurs taxa infraspécifiques) appartenant à 422 genres et 87 familles, et 7720 synonymes (voir nouveautés taxonomiques, Tableau 1). Il inclut 11 espèces introduites et naturalisées. Huit familles représentent ensemble près de 50 % des espèces : les Leguminosae, Lauraceae, Myrtaceae, Sapotaceae, Chrysobalanaceae, Annonaceae, Melastomataceae et Rubiaceae (Tableau 2). Au sein des Leguminosae, la sous-famille des Caesalpinoideae est de loin la plus riche, avec 134 spp. pour 27 genres, c'est-à-dire plus que n'importe quelle autre famille présente en Guyane. Au sein des Caesalpinoideae, 106 espèces représentant 18 genres font partie du clade mimosoïde tel que défini par LPWG (2017). Seize familles ne comptent qu'une seule espèce, et 40 familles cinq ou moins (Tableau 2).

Parmi les 46 genres qui ensemble regroupent plus de 50 % de toutes les espèces d'arbres, les plus diversifiés sont *Eugenia*, *Inga*, *Pouteria*, *Ocotea*, *Miconia*, *Myrcia*, *Protium*, *Sloanea* et *Licania* (Tableau 3). À l'opposé, 178 genres sont représentés par une seule espèce, et 85 par seulement deux.

La comparaison de notre liste avec les références floristiques antérieures pour l'ensemble de la Guyane fran-

TABLE 3. — Genera with ten or more tree species in French Guiana/Genres comportant dix espèces d'arbres ou plus en Guyane française.

Genus (Family)	Nb of species
<i>Eugenia</i> (Myrtaceae)	62
<i>Inga</i> (Leguminosae-Caesalpinoideae, mimosoid clade)	59
<i>Pouteria</i> (Sapotaceae)	59
<i>Ocotea</i> (Lauraceae)	40
<i>Miconia</i> (Melastomataceae)	38
<i>Myrcia</i> (Myrtaceae)	35
<i>Protium</i> (Burseraceae)	35
<i>Sloanea</i> (Elaeocarpaceae)	32
<i>Licania</i> (Chrysobalanaceae)	31
<i>Ficus</i> (Moraceae)	26
<i>Casearia</i> (Salicaceae)	18
<i>Cordia</i> (Cordiaceae)	18
<i>Guatteria</i> (Annonaceae)	18
<i>Eschweilera</i> (Lecythidaceae)	18
<i>Aniba</i> (Lauraceae)	17
<i>Chrysophyllum</i> (Sapotaceae)	17
<i>Hirtella</i> (Chrysobalanaceae)	17
<i>Mouriri</i> (Melastomataceae)	17
<i>Terminalia</i> (Combretaceae)	16
<i>Tovomita</i> (Clusiaceae)	16
<i>Ormosia</i> (Leguminosae-Papilionoideae)	15
<i>Swartzia</i> (Leguminosae-Papilionoideae)	15
<i>Talisia</i> (Sapindaceae)	15
<i>Annona</i> (Annonaceae)	14
<i>Guarea</i> (Meliaceae)	13
<i>Micromelis</i> (Sapotaceae)	13
<i>Vochysia</i> (Vochysiaceae)	13
<i>Clusia</i> (Clusiaceae)	12
<i>Couepia</i> (Chrysobalanaceae)	12
<i>Erythroxylum</i> (Erythroxylaceae)	12
<i>Hymenopus</i> (Chrysobalanaceae)	12
<i>Lecythis</i> (Lecythidaceae)	12
<i>Rinorea</i> (Violaceae)	12
<i>Trichilia</i> (Meliaceae)	12
<i>Xylopia</i> (Annonaceae)	12
<i>Licaria</i> (Lauraceae)	11
<i>Pourourma</i> (Urticaceae)	11
<i>Quiina</i> (Ochnaceae)	11
<i>Aspidosperma</i> (Apocynaceae)	10
<i>Byrsinima</i> (Malpighiaceae)	10
<i>Cybianthus</i> (Primulaceae)	10
<i>Diospyros</i> (Ebenaceae)	10
<i>Duguetia</i> (Annonaceae)	10
<i>Ouratea</i> (Ochnaceae)	10
<i>Tachigali</i> (Leguminosae-Caesalpinoideae)	10

Lastly, the 4354 vernacular names in nine languages that are listed here for 1157 species, make up the first vernacular nomenclature published so far for French Guiana.

Acknowledgements

We dedicate this work to our dear friends and colleagues Marie-Françoise “Fanchon” Prévost (1941-2013) and Scott A. Mori (1941-2020), who devoted so much passion and effort to our understanding of the flora of French Guiana. This work owes a lot to the richness and quality of the Herbier IRD de Guyane (CAY) (Gonzalez et al. 2021). We are indebted to its Curator Sophie Gonzalez and to the technical staff, as well as the many specialists who checked our tentative determinations and identified the collections held in CAY. We thank Georges Elfort, Jean-Louis Smock, Chantal Geniez, Michel Tarcy, Desmo and Wemo Betian, pour leur assistance technique sur le terrain lors des inventaires des arbres, et Pascal Petronelli pour sa contribution aux identifications botaniques. Nous remercions également Olivier Brunaux, Joseph Ateni et Hélène Richard pour leur contribution au recensement des noms vernaculaires utilisés par les forestiers et leurs informateurs en nengee tongo, français, créole et portugais du Brésil. Nous remercions chaleureusement Clarisse Ansoe-Tareau, Marc-Alexandre Tareau et Jammes Panapuy

çaise (Fig. 5) illustre à la fois une accélération du rythme d’accumulation des connaissances au cours des dernières décennies, et le chemin qui reste à parcourir pour compléter l’inventaire de la flore, dont les arbres ne sont qu’une partie. L’incomplétude de l’inventaire de la flore arborescente est illustrée par l’évolution du nombre d’espèces non nommées : ce chiffre est passé de 117 dans Molino et al. (2009) à 143 aujourd’hui, malgré l’identification de 52 taxons de la liste de 2009 et la suppression de 10 autres en raison de l’insuffisance des données (Annexe 6).

Ce catalogue fournit pour la première fois des données démographiques sur de nombreuses espèces, pour la plupart très mal connues à cet égard. Plus précisément, l’abondance totale, la fréquence maximale locale et la distribution géographique sont données pour les 1293 espèces qui sont présentes dans nos relevés. Ces informations, avec le nombre de spécimens d’herbier collectés en Guyane française, peuvent servir à de futures études en écologie, ainsi qu’à l’évaluation du statut de conservation de ces espèces selon les critères de la liste rouge de l’IUCN (2021).

Sur les 218 espèces considérées comme hyperdominantes par ter Steege et al. (2020), 143 (66%) sont présentes dans notre liste. Bien qu’hyperdominantes en Amazonie, elles ne sont cependant pas toutes abondantes en Guyane : 41 d’entre elles ne sont représentées que par 20 individus ou moins dans nos inventaires d’arbres. À l’opposé, seules neuf espèces hyperdominantes sont parmi les 20 espèces les plus abondantes (> 1000 arbres) dans notre jeu de données.

Enfin, les 4354 noms vernaculaires dans neuf langues recensés ici pour 1157 espèces constituent la première nomenclature vernaculaire des arbres de Guyane publiée à ce jour.

Remerciements

Nous dédions ce travail à nos regrettés amis et collègues Marie-Françoise “Fanchon” Prévost (1941-2013) et Scott A. Mori (1941-2020), qui ont consacré tant de passion et d’efforts à l’étude de la flore de la Guyane. Ce travail doit beaucoup à la richesse et à la qualité de l’Herbier IRD de Guyane (CAY) (Gonzalez et al. 2021). Nous sommes redevables à sa conservatrice Sophie Gonzalez et au personnel technique, ainsi qu’aux nombreux spécialistes qui ont vérifié nos déterminations et identifié les collections de CAY. Nous remercions Georges Elfort, Jean-Louis Smock, Chantal Geniez, Michel Tarcy, Desmo et Wemo Betian, pour leur assistance technique sur le terrain lors des inventaires des arbres, et Pascal Petronelli pour sa contribution aux identifications botaniques. Nous remercions également Olivier Brunaux, Joseph Ateni et Hélène Richard pour leur contribution au recensement des noms vernaculaires utilisés par les forestiers et leurs informateurs en nengee tongo, français, créole et portugais du Brésil. Nous remercions chaleureusement Clarisse Ansoe-Tareau, Marc-Alexandre Tareau et Jammes Panapuy

Wemo Betian, for technical assistance in the field during tree inventories, and Pascal Petronelli for his contribution to botanical identifications. We also thank Olivier Brunaux, Joseph Ateni and Hélène Richard for their contribution to the census of vernacular names used by foresters and tree spotters in Nengee tongo, French, Creole and Brazilian Portuguese languages. Clarisse Ansoe-Tareau, Marc-Alexandre Tareau and Jammes Panapuy are warmly thanked for checking the spelling of, respectively, Nengee tongo, French Guianese Creole, and Teko names. We thank Odile Poncy, Corinne Sarthou and the editor-in-chief Thierry Deroin for their careful reading of the manuscript and helpful suggestions. CAY specimens have been digitized thanks to the “Investissement d’Avenir” program e-RECOLNAT (RECOLNAT-ANR-11-IN-BS-0004). This work was supported by two other “Investissement d’Avenir” grants managed by the *Agence nationale de la Recherche*: Labex DRIIHM/IRDHEI (ANR-11-LABX-0010/11-LABX-0010) and Labex CEBA (ANR-10-LABX-25-01).

Author contributions

JFM and DS initiated the project with M.-F. Prévost†; DS, MFP†, JFM and JE provided inventory data and established the primary list of tree species with input from PG, and from PGD for Rubiaceae; JFM and CAM computed demographic indices; taxonomic and nomenclatural compilations were done by JFM (all families), DF (several families, including Annonaceae, Chrysobalanaceae, Leguminosae-Caesalpinoideae, Lecythidaceae, Meliaceae), PGD (Rubiaceae), with input from JE, and checked by EJL for Myrtaceae; GO initiated the multilingual compilation of vernacular names; PG provided Kali’na, Palikur and Wayápi names, compiled lists of Creole, French and Brazilian names from various sources, and checked names in all languages; MF provided Wayana and Nengee tongo names; GO and DD provided Teko names; JFM wrote the first draft of the text; DF, EJL and PGD corrected the English; all authors commented on and amended the text •

pour la vérification de l’orthographe des noms en nengee tongo, en créole et en teko. Nous remercions Odile Poncy, Corinne Sarthou et le rédacteur-en-chef Thierry Deroin, pour leur lecture attentive du manuscrit et leurs suggestions pertinentes. Les spécimens de CAY ont été numérisés dans le cadre du programme “Investissement d’Avenir” e-RECOLNAT (RECOLNAT-ANR-11-IN-BS-0004) géré par l’Agence nationale de la Recherche. Ce travail a bénéficié de financements provenant de deux autres programmes “Investissement d’Avenir”, eux aussi gérés par l’ANR : le LabEx DRIIHM/IRDHEI (ANR-11-LABX-0010/11-LABX-0010) et le LabEx CEBA (ANR-10-LABX-25-01).

Contribution des auteurs

JFM et DS ont initié le projet avec M.-F. Prévost† ; DS, MFP†, JFM et JE ont fourni les données d’inventaire et établi la liste primaire des espèces d’arbres, avec la contribution de PG, et de PGD pour les Rubiaceae ; JFM et CAM ont calculé les indicateurs démographiques ; les compilations taxonomiques et nomenclaturales ont été réalisées par JFM (toutes les familles), DF (plusieurs familles, dont les Annonaceae, Chrysobalanaceae, Leguminosae-Caesalpinoideae, Lecythidaceae, Meliaceae), PGD (Rubiaceae), avec la contribution de JE, et vérifiées par EJL pour les Myrtaceae ; GO a initié la compilation multilingue des noms vernaculaires ; PG a fourni les noms Kali’na, Palikur et Wayápi, a compilé des listes de noms créoles, français et brésiliens provenant de diverses sources, et a vérifié les noms dans toutes les langues ; MF a fourni les noms Wayana et Nengee tongo ; GO et DD ont fourni les noms Teko ; JFM a écrit la première version du texte ; DF, EJL et PGD ont corrigé l’anglais ; tous les auteurs ont commenté et modifié le texte •

CHECKLIST

Each accepted name is preceded by a unique number in square brackets, which serves as a reference in the indexes of exsiccata and vernacular names that follow the checklist/*Chaque nom accepté est précédé d'un numéro unique entre crochets, qui sert de référence dans les index des exsiccata et des noms vernaculaires qui suivent le catalogue.*

Family ACANTHACEAE Juss.
Genus *Avicennia* L.

[1] *Avicennia germinans* (L.) L.
(Fig. 6A)

Sp. Pl., ed. 3, 2: 891 (Linnaeus 1764). — *Bontia germinans* L., *Syst. Nat.*, ed. 10, 2: 1122 [7 Jun. 1759] (Linnaeus 1759).

Avicennia nitida Jacq., *Enum. Syst. Pl.*: 25 [Aug.-Sep. 1760] (Jacquin 1760). — *Hilairanthus nitidus* (Jacq.) Tiegh., *J. Bot. [Morot]* 12: 358 (Tieghem 1898). — *Avicennia officinalis* L. var. *nitida* (Jacq.) Kuntze, *Revis. Gen. Pl.* 2: 502 [5 Nov. 1891] (Kuntze 1891).

Avicennia tomentosa Jacq., *Enum. Syst. Pl.*: 25 [Aug.-Sep. 1760] (Jacquin 1760). — *Hilairanthus tomentosus* (Jacq.) Tiegh., *J. Bot. [Morot]* 12: 358 (Tieghem 1898).

Avicennia africana P.Beauv., *Fl. Oware* 1: 80 (Palisot 1806).

Avicennia tomentosa var. *campechensis* Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 284 [8 June 1818] (Kunth 1818).

Avicennia tomentosa var. *cumanensis* Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 283 [8 June 1818] (Kunth 1818). — *Avicennia germinans* var. *cumanensis* (Kunth) Moldenke, *Phytologia* 30 (1): 15 [23 Jan. 1975] (Moldenke 1975).

Avicennia tomentosa var. *guayaquilensis* Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 284 [8 June 1818] (Kunth 1818). — *Avicennia germinans* var. *guayaquilensis* (Kunth) Moldenke, *Phytologia* 29 (2): 75 [19 Nov. 1974] (Moldenke 1974).

Avicennia elliptica Thunb., *Pl. Bras.* 3: 37 (Thunberg 1821).

Avicennia floridana Raf., *Atlantic J.*: 148 (Rafinesque 1832).

Avicennia lamarckiana C.Presl, *Abh. Königl. Böhm. Ges. Wiss.*, ser. 5, 3: 529 [Jul.-Dec. 1845] (Presl 1845).

Avicennia meyeri Miq., *Linnaea* 18: 262 ["1844" publ. Feb-(May?) 1845] (Miquel 1845).

Avicennia oblongifolia Nutt. ex Chapm., *Fl. South. U.S.*: 310 [14 Aug. 1860] (Chapman 1860).

Avicennia floridana Gand., *Bull. Soc. Bot. France* 65: 64 (Gandoger 1918), nom. illeg. hom., non Raf. (Rafinesque 1832).

Avicennia nitida var. *trinitensis* Moldenke, *Phytologia* 1 (2): 96 [4 Aug. 1934] (Moldenke 1934).

Avicennia germinans f. *aberrans* Moldenke, *Phytologia* 47 (3): 222 [Dec. 1980] (Moldenke 1980).

Avicennia germinans f. *brasiliensis* Moldenke, *Phytologia* 47 (3): 222 [Dec. 1980] (Moldenke 1980).

Avicennia germinans f. *venezuelensis* Moldenke, *Phytologia* 47 (3): 222 [Dec. 1980] (Moldenke 1980).

NOTES. — A species restricted to mangrove forests. Replacement of *Bontia* L. by *Avicennia* Jacq. already appears in *errata* of Species Plantarum, ed. 2 (*Linnaeus* 1763, 2: 1684, typographical error "990" for 890), but as for genus only. It was fully incorporated in ed. 3 for species (*Linnaeus* 1764: 891) (see Compère 1963).

VERNACULAR NAMES. — Pa: payuy • Ka: apali'i, apaliyu • Nt: paluwa • Cr: pativyé-blanc • Fr: palétuvier blanc • Br: siriúba.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *M.-F. Prévost* 3900, 15 m × 18 cm.

Family ACHARIACEAE Harms
Genus *Carpotroche* Endl.

[2] *Carpotroche crispidentata* Ducke

Arch. Inst. Biol. Veg. 4 (1): 55 [June 1938] (Ducke 1938).

VERNACULAR NAMES. — Pa: wakavu-kamwi, waravru-kamwi • Wn: pëmu epit • Br: fruta-de-cutia.

HERBARIUM DATA (FG). — 80 collections at CAY. Sel. exs.: *D. Sabatier* 965.

SIZE. — Up to 15 cm dbh (Sleumer 1980).

[3] *Carpotroche longifolia* (Poepp.) Benth.
(Fig. 6B)

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 82 [May 1861] (Bentham 1861). — *Mayna longifolia* Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 64 [23-25 Jan. 1845] (Poeppig 1845).

Mayna lingulifolia R.E.Schult., *Caldasia* 3 (15): 439 (Schultes 1945). — *Carpotroche lingulifolia* (R.E.Schult.) Cuatrec., *Trop. Woods* 101: 28 (Cuatrecasas 1955).

Mayna longifolia var. *heliocarpa* R.E.Schult., *Caldasia* 3 (15): 441 (Schultes 1945). — *Carpotroche longifolia* var. *heliocarpa* (R.E.Schult.) Cuatrec., *Trop. Woods* 101: 28 (Cuatrecasas 1955).

Mayna longifolia var. *phasmatocarpa* R.E.Schult., *Bot. Mus. Leafl.* 12 (4): 125 [25 Jan. 1946] (Schultes 1946). — *Carpotroche longifolia* var. *phasmatocarpa* (R.E.Schult.) Cuatrec., *Trop. Woods* 101: 28 (Cuatrecasas 1955).

Mayna pacifica Cuatrec. var. *pusilla* R.E.Schult., *Bot. Mus. Leafl.* 13 (9): 285 [18 Apr. 1949] (Schultes 1949).

Mayna muricida R.E.Schult., *Rhodora* 65: 14 (Schultes 1963).

VERNACULAR NAMES. — Wp: ka'a ki, walapulu lá • Wn: tapalaimë • Br: cacao-branco, fruta-de-cutia, pau-de-cachimbo.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *P. Grenand* 645.

SIZE. — Up to 20 m tall (Sleumer 1980).

[4] *Carpotroche surinamensis* Uittien

Recueil Trav. Bot. Néerl. 22: 368 ["1925" publ. Jan. 1926] (Uitten 1926).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2193*.

SIZE. — Guyana. *M.J. Jansen-Jacobs et al. 230*: 10 cm dbh.

Genus *Lindackeria* C.Presl

[5] *Lindackeria paludosa* (Benth.) Gilg

Nat. Pflanzenfam. [Engler & Prantl], ed. 2, 21: 404 (Gilg 1925). — *Mayna paludosa* Benth., *J. Bot. [Hooker]* 4: 114 (Bentham 1842). — *Carpotroche paludosa* (Benth.) Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 203 [18-20 Sep. 1842] (Walpers 1842).

Mayna laxiflora Benth., *J. Bot. [Hooker]* 4: 114 (Bentham 1842). — *Carpotroche laxiflora* (Benth.) Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 203 [18-20 Sep. 1842] (Walpers 1842). — *Oncoba maynensis* var. *laxiflora* (Benth.) Eichler, *Fl. Bras. [Martius]* 13 (1): 441 [1 Oct. 1871] (Eichler 1871). — *Lindackeria maynensis* var. *laxiflora* (Benth.) Mart. ex Pittier, *Cat. Fl. Venez. [Pittier]* 2: 166 (Pittier 1947).

Lindackeria maynensis Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 63 [23-25 Jan. 1845] (Poeppig 1845). — *Oncoba maynensis* (Poepp.) Eichler, *Fl. Bras. [Martius]* 13 (1): 441 [1 Oct. 1871] (Eichler 1871).

VERNACULAR NAMES. — Pa: iwan-etni, iwen-hetni.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori 2116*.

INVENTORY DATA (FG). — 35 trees in 4 plots; $F_{\max} = 5.1\%$; $dbh_{\text{inv}} = 21.6$ cm.

Family ANACARDIACEAE R.Br.
Genus *Anacardium* L.

[6] *Anacardium amapaense* J.D.Mitch.
(Fig. 6C)

Brittonia 44 (3): 331 (Mitchell 1992), “*amapaense*”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier 3146* (para-, CAY[CAY002075]).

INVENTORY DATA (FG). — 3 trees in 2 plots; $dbh_{\text{inv}} = 52.2$ cm.

[7] *Anacardium giganteum* W.Hancock ex Engl.

Fl. Bras. [Martius] 12 (2): 409 [1 Sep. 1876] (Engler 1876).

VERNACULAR NAMES. — Ka: akayuu • Wp: akayu u • Wn: olojimé • Cr: kajou-gran-bwa • Br: cajuaçu, caju-da-mata, cajueiro-do-mato.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier 5139*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 111.4$ cm.

[8] *Anacardium occidentale* L.

Sp. Pl. 1: 383 [1 May 1753] (Linnaeus 1753). — *Acajuba occidentalis* (L.) Gaertn., *Fruct. Sem. Pl.* 1: 192 [Dec. 1788] (Gaertner 1788).

Cassuvium pomiferum Lam., *Encycl. [J. Lamarck et al.]* 1 (1): 22 [2 Dec. 1783] (Lamarck 1783).

Cassuvium solitarium Stokes, *Bot. Mat. Med.* 2: 474 (Stokes 1812).

Cassuvium reniforme Blanco, *Fl. Filip. [F.M. Blanco]*: 322 (Blanco 1837).

Anacardium occidentale var. *gardneri* Engl., *Monogr. Phan. [A.D.C. & C.D.C.]* 4: 220 [Mar. 1883] (Engler 1883).

Anacardium microcarpum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 202 (Ducke 1922).

NOTE. — Introduced from Brazil, naturalised on the western coast of French Guiana.

VERNACULAR NAMES. — Pa: mihitui • Ka: oloi • Te: akadjun • Wp: akayuu • Wn: olojimé • Nt: busi kasu, busi kasun • Cr: kajou-gran-bwa • Fr: noix de cajou (fruit), pomme cajou (pseudo-fruit), pommier-cajou • Br: caju (fruit), cajueiro.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *A. Le Goff 201*.

SIZE. — Up to 40 cm dbh (Mitchell 1997).

[9] *Anacardium spruceanum* Benth. ex Engl.

Fl. Bras. [Martius] 12 (2): 410 [1 Sep. 1876] (Engler 1876).

VERNACULAR NAMES. — Pa: mihitui-kamwi, mihitui-tivarabuyené • Ka: akayuu • Wp: akayu u • Wn: olojimé • Nt: busi kasu, busi kasun • Cr: kajou-gran-bwa • Fr: cajou sauvage • Br: cajuaçu, caju-da-mata, cajueiro-do-mato.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & D.-Y. Alexandre 889*.

INVENTORY DATA (FG). — 153 trees in 83 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 85$ cm.

Genus *Astronium* Jacq.

[10] *Astronium lecointei* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 202 (Ducke 1922).

NOTE. — According to Mitchell (1995), it is very close to *Astronium obliquum* Griseb., and if synonymized, the latter has priority.

VERNACULAR NAMES. — Te: tininipi • Br: aroeira, muiraquatiara.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1991*.

INVENTORY DATA (FG). — 25 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.7$ cm.

[11] *Astronium ulei* Mattick

Notizbl. Bot. Gart. Berlin-Dahlem 11: 996 [20 Jan. 1934] (Mattick 1934).




Fig. 6. — Acanthaceae: **A**, *Avicennia germinans* (L.) L. (M.-F. Prévost 4058); **B**, *Carpotroche longifolia* (Poepp.) Benth.; **C**, *Anacardium amapaense* J.D.Mitch. (D. Sabatier et al. 6109); **D**, *Thrysodium guianense* Sagot ex Marchand (M.-F. Prévost & D. Sabatier 4801). A, © M.-F. Prévost/IRD; B, © J.-F. Molino/IRD; C, D, © D. Sabatier/IRD.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *M.-F. Prévost 2161*.

INVENTORY DATA (FG). — 81 trees in 30 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 38.4\text{ cm}$.

Genus *Loxopterygium* Hook.f.

[¹²] *Loxopterygium sagotii* Hook.f.

Gen. Pl. [Bentham & Hooker f.] 1 (1): 420 [7 Aug. 1862] (Hooker 1862).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: aneku-kamwi • Ka: kuipali.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P.A. Sagot 973*, 1856 (holo-, K[K000572513, K000572514, K000572515]; iso-, BM[BM000793057], BR[BR0000006952877], F[V0047653F], MPU[MPU021092], U[U0000193]).

INVENTORY DATA (FG). — 1 tree; $dbh = 48.3\text{ cm}$.

Genus *Spondias* L.

[¹³] *Spondias mombin* L.

Sp. Pl. 1: 371 [1 May 1753] (Linnaeus 1753). — *Spondias myrobalanus* L., *Syst. Nat.*, ed. 10, 2: 1036 [7 June 1759] (Linnaeus 1759), nom. illeg. superfl. (based on the same description and reference to “Brown Jam. 229” as *S. lutea* L.). — *Spondias lutea* L., *Sp. Pl.*, ed. 2, 1: 613 [Sep. 1762] (Linnaeus 1762), nom. illeg. superfl. (based on the type of *Spondias mombin* L.). — *Spondias lucida* Salisb., *Prodri. Stirp. Chap. Allerton 172* [Nov.-Dec. 1796] (Salisbury 1796), nom. illeg. superfl. (based on *Spondias myrobalanus* L. and *S. lutea* L.)

Spondias cytherea Tussac, *Fl. Antill. 3*: 95 (Tussac 1825), nom. illeg. hom., non Sonn. (Sonnerat 1782).

Spondias aurantiaca Schumach. & Thonn., *Beskr. Guin. Pl. 225* (Schumacher & Thonning 1827).

Spondias pseudomyrobalanus Tussac, *Fl. Antill. 4*: 97 (Tussac 1827).

Spondias dubia A.Rich., *Fl. Seneg. Tent.* 1 (4): 153 [Sep. 1831] (Richard 1831).

Spondias axillaris Roxb., *Fl. Ind.*, ed. 1832, 2: 453 (Roxburgh 1832).

Spondias oghigee G.Don, *Gen. Hist.* 2: 79 [Oct. 1832] (Don 1832).

Spondias zanzee G.Don, *Gen. Hist.* 2: 79 [Oct. 1832] (Don 1832).

Spondias graveolens Macfad., *Fl. Jamaica [Macfadyen]* 1: 228 (Macfadyen 1837).

Spondias lutea var. *glabra* Engl., *Fl. Bras. [Martius]* 12 (2): 374 [1 Sep. 1876] (Engler 1876).

Spondias lutea var. *maxima* Engl., *Fl. Bras. [Martius]* 12 (2): 374 [1 Sep. 1876] (Engler 1876).

Spondias nigrescens Pittier, *Contr. U.S. Natl. Herb.* 18 (2): 75 [1 Apr. 1914] (Pittier 1914).

Spondias purpurea L. var. *venulosa* Engl., *Fl. Bras. [Martius]* 12 (2): 373 [1 Sep. 1876] (Engler 1876). — *Spondias venulosa* (Engl.) Engl., *Monogr. Phan. [A.D.C. & C.D.C.]* 4: 245 [Mar. 1883] (Engler 1883).

Spondias radlkoferi Donn.Sm., *Bot. Gaz.* 16 (7): 194 [20 July 1891] (Donnell Smith 1891).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Semi-domesticated by pre-Columbian Amerindians (Clement, 1999; Levis *et al.*, 2017).

VERNACULAR NAMES. — Pa: kahambag • Ka: mo:pe, mope • Te: bope • Wp: akaya, mope, tapeliwa • Wn: mope • Nt: mope • Cr: monben • Fr: mombin • Br: cajá, taperebá, taperibá.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *M.-F. Prévost* 3330.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 89$ cm.

Genus *Tapirira* Aubl.

[14] *Tapirira bethanniana* J.D.Mitch.

Mem. New York Bot. Gard. 64: 230 (Mitchell 1990).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: ara.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom* 15253 (holo-, NY[00050969]; iso-, CAY[CAY005527], U[U0000217, U0000218]).

INVENTORY DATA (FG). — 77 trees in 53 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 79$ cm.

[15] *Tapirira guianensis* Aubl.

Hist. Pl. Guiane 1: 470 [Jun.-Dec. 1775] (Aublet 1775). — *Jonquieria paniculata* Willd., *Sp. Pl. ed. 4* 2 (1): 750 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Tapirira guianensis*).

Comocladia tapaculo Kunth, *Nova genera et species plantarum [H.B.K.]* 7: 18 [2 Nov. 1824] (Kunth 1824).

Mauria multiflora Mart. ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 14 (Bentham 1852).

Mauria subbijuga Mart. ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 15 (Bentham 1852).

Odina francoana Netto, *Ann. Sci. Nat., Bot. sér. 5*, 5: 85 (Netto 1866).

Tapirira bijuga Marchand, *Rev. Anacard.*: 162 (Marchand 1869), “*Tapirira*”.

Tapirira myriantha Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5*, 14: 295 (Triana & Planchon 1872).

Tapirira pao-pombo Marchand, *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1873: 58 (Marchand 1873).

Tapirira guianensis var. *cuneata* Engl., *Fl. Bras. [Martius]* 12 (2): 378 [1 Sep. 1876] (Engler 1876).

Tapirira guianensis var. *elliptica* Engl., *Fl. Bras. [Martius]* 12 (2): 378 [1 Sep. 1876] (Engler 1876).

Tapirira pearcei Rusby, *Mem. Torrey Bot. Club* 6 (1): 22 (Rusby 1896).

Tapirira fanshawei Sandwith, *Kew Bull.* 10 (3): 470 [20 Dec. 1955] (Sandwith 1955).

Tapirira guianensis subsp. *subandina* Barfod & Holm-Niels., *Nordic J. Bot.* 6 (4): 425 (Barfod & Holm-Nielsen 1986).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Marchand (1869: 162) ascribed to J.D. Hooker (Hook.f.) the transfer to *Tapirira* (“*Tapirira*”) of “*Mauria (Cyrtocarpa) bijuga* Mart.” Since “*Mauria bijuga*” does not exist, as Marchand (1869: 162) actually referred to *Mauria subbijuga* Martius ex Bentham (1852: 15), and since there is no trace of such a transfer by J.D. Hooker, *Tapirira bijuga* is to be ascribed to Marchand alone (not “Hook.f. ex Marchand”).

VERNACULAR NAMES. — Pa: ara, ara-ahavukunō • Ka: atapili:li, tapilili, watapalili • Te: tatapililit • Wp: tata pilili • Wn: ajawaimé • Nt: agandyaymay • Cr: loussé (fruit), monben-blanc, monben-fou, monben-sovaj • Fr: mombin blanc, mombin fou • Br: cedroí, fruta-de-pombo, pau-pombo, tatapiririca.

HERBARIUM DATA (FG). — 151 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000992002]).

INVENTORY DATA (FG). — 101 trees in 55 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 58.4$ cm.

[16] *Tapirira obtusa* (Benth.) J.D.Mitch.

Novon 3 (1): 66 [spring 1993] (Mitchell 1993). — *Mauria obtusa* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 16 (Bentham 1852).

Tapirira pao-pombo var. *major* Marchand, *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1873: 59 (Marchand 1873). — *Tapirira marchandii* Engl., *Fl. Bras. [Martius]* 12 (2): 379 [1 Sep. 1876] (Engler 1876), nom. nov., based on *Tapirira pao-pombo* var. *major*.

Tapirira peckoltiana Engl., *Fl. Bras. [Martius]* 12 (2): 379 [1 Sep. 1876] (Engler 1876).

VERNACULAR NAMES. — Pa: ara • Ka: atapili:li, tapilili, watapalili • Wp: tata pilili • Nt: baaka agandyaymay • Cr: loussé (fruit), monben-blanc, monben-fou, monben-sovaj • Fr: mombin blanc, mombin fou • Br: cedroí, pau-pombo, tatapiririca.

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *J.-F. Molino 1181*.

INVENTORY DATA (FG). — 71 trees in 45 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 48.7\text{ cm}$.

Genus *Thyrsodium* Salzm. ex Benth.

[17] *Thyrsodium guianense* Sagot ex Marchand (Fig. 6D)

Rev. Anacard.: 160 (Marchand 1869). — *Garuga guianensis* (Sagot ex Marchand) Engl., *Fl. Bras. [Martius]* 12 (2): 288 [1 Sep. 1874] (Engler 1874).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: awao, awau • Wp: anila wisi, pana'i wu • Nt: busi kasu, busi kasun • Cr: karapa-oyak, monben-fou • Br: castanha-de-porco.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *P.A. Sagot 1202*, Oct. 1858 (holo-, P[P02440694]; iso-, B[not seen, photo F neg. 13153], BM[BM000793056[1858]], F[V0049806F, V0049807F], G[G00237128], GH[GH00049263], K[K000572524, K000572525], MPU[MPU020465], P[P02440695], S[S-R-10870], U[U0000222]).

INVENTORY DATA (FG). — 204 trees in 80 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 47.1\text{ cm}$.

[18] *Thyrsodium puberulum* J.D.Mitch. & Daly

Brittonia 45 (2): 122 (Mitchell & Daly 1993).

VERNACULAR NAMES. — Pa: ayao, ayau • Te: tudjulata'i • Wp: taitetu kasi ni, taitetu kási ni • Nt: lebi moni • Cr: karapa-oyak • Br: breu-de-leite.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *J.-J. de Granville B-5486* (holo-, NY[00050976]; iso-, CAY[CAY010453, CAY010454], K[K000035326], NY[00004689]).

INVENTORY DATA (FG). — 200 trees in 86 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 53.5\text{ cm}$.

[19] *Thyrsodium spruceanum* Benth.

Hooker's J. Bot. Kew Gard. Misc. 4: 17 (Bentham 1852). — *Garuga spruceana* (Benth.) Engl., *Fl. Bras. [Martius]* 12 (2): 286 [1 Sep. 1874] (Engler 1874).

Thyrsodium schomburgkianum Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 18 (Bentham 1852). — *Garuga schomburgkiana* (Benth.) Engl., *Fl. Bras. [Martius]* 12 (2): 287 [1 Sep. 1874] (Engler 1874).

Thyrsodium salzmannianum Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 18 (Bentham 1852). — *Garuga schomburgkiana* (Benth.) Engl. var. *salzmanniana* (Benth.) Engl., *Fl. Bras. [Martius]* 12 (2): 287 [1 Sep. 1874] (Engler 1874). — *Thyrsodium schomburgkianum* Benth. var. *salzmannianum* (Benth.) Engl., *Monogr. Phan. [A.DC. & C.DC.]* 4: 297 [Mar. 1883] (Engler 1883).

Garuga gigantea Engl., *Fl. Bras. [Martius]* 12 (2): 288 [1 Sep. 1874] (Engler 1874). — *Thyrsodium giganteum* (Engl.) Engl., *Monogr. Phan. [A.DC. & C.DC.]* 4: 297 [Mar. 1883] (Engler 1883).

Thyrsodium paraense Huber, *Bull. Soc. Bot. Genève*, sér. 2, 6: 183 ["1914" publ. 1915] (Huber 1915).

Thyrsodium dasytrichum Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 210 [27 June 1932] (Sandwith 1932).

NOTE. — One of five species whose relative density in tree communities likely indicative of pre-Columbian archeological sites in French Guiana (Molino et al. 2021).

VERNACULAR NAMES. — Pa: wakavu-kamwi • Wp: tāpē lemiū'i • Cr: karapa-oyak • Br: breu-de-leite.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2436*.

INVENTORY DATA (FG). — 38 trees in 24 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 35.1\text{ cm}$.

Family ANNONACEAE Juss.
Genus *Anaxagorea* A.St.-Hil.

[20] *Anaxagorea acuminata* (Dunal) A.DC.

Mém. Soc. Phys. Genève 5: 211 (Candolle 1832). — *Xylopia acuminata* Dunal, *Monogr. Anonac.*: 122 [Aug.-Nov. 1817] (Dunal 1817). — *Xylopicrum acuminatum* (Dunal) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

Anaxagorea multiflora R.E.Fr., *Acta Horti Berg.* 12 (1): 13 (Fries 1934).

VERNACULAR NAMES. — Ka: apelemu, omose tubulu • Wp: pina'i piye, pina'i sili.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (type G-DC[G00201576]).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 11.5\text{ cm}$.

[21] *Anaxagorea brevipedicellata* Timmerman

Proc. Kon. Ned. Akad. Wetensch. C 87 (3): 299 (Timmerman 1984).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3553* (para-, CAY[CAY010466], P[P01960321]).

SIZE. — Up to 15 m tall (Maas & Westra 1985).

[22] *Anaxagorea dolichocarpa* Sprague & Sandwith

Bull. Misc. Inform. Kew 1930 (10): 475 [4 Dec. 1930] (Sprague & Sandwith 1930).

Anaxagorea megalophylla R.E.Fr., *Acta Horti Berg.* 12 (1): 20 (Fries 1934).

Anaxagorea mutica R.E.Fr., *Acta Horti Berg.* 12 (1): 19 (Fries 1934).

VERNACULAR NAMES. — Pa: manig-kamwi, miret-kamwi • Ka: kuwe enekali • Wp: sa'i melu ka'a • Cr: baton-zin • Br: envira-branca.

HERBARIUM DATA (FG). — 180 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 996*.

INVENTORY DATA (FG). — 43 trees in 27 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 13.4\text{ cm}$.

[23] *Anaxagorea phaeocarpa* Mart.

Fl. Bras. [Martius] 13 (1): 40 [1 Jan. 1841] (Martius 1841).

Anaxagorea costaricensis R.E.Fr., *Acta Horti Berg.* 12 (1): 25 (Fries 1934).

VERNACULAR NAMES. — Pa: kigiksau, kiriksau • Wp: sa'i melu ka'a.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *R.A.A. Oldeman 1897*.

SIZE. — Up to 15 cm dbh (Maas *et al.* 2007).

Genus *Annona* L.

[24] *Annona ambotay* Aubl.

Hist. Pl. Guiane 1: 616 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Ka: ambotay (*fide* Aublet 1775) • Te: dzalatay • Wp: iwi tay, iwi tay sili • Nt: busi atuku • Cr: viratay • Br: araticú, envira-caju, envira-jodó, envira-taia.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN[[LINN-HS 983.5](#)]).

INVENTORY DATA (FG). — 28 trees in 26 plots; $F_{\max} < 1\%$; $dbh_{inv} = 18.9$ cm.

[25] *Annona cuspidata* (Mart.) H.Rainer

Ann. Naturhist. Mus. Wien 108B: 193 (Rainer 2007). — *Rollinia cuspidata* Mart., *Fl. Bras. [Martius]* 13 (1): 20 [1 Jan. 1841] (Martius 1841).

Rollinia cardiantha Diels, *Verb. Bot. Vereins Prov. Brandenburg* 47: 134 [1 Oct. 1905] (Diels 1905).

Rollinia pachyptera Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 176 [20 Dec. 1927] (Diels 1927).

Rollinia sphaerantha R.E.Fr., *Acta Horti Berg.* 12 (1): 184 (Fries 1934).

Rollinia uniflora R.E.Fr., *Acta Horti Berg.* 12 (1): 182 (Fries 1934).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *P. Chareyre 30D*.

SIZE. — Up to 20 cm dbh (Maas *et al.* 1992).

[26] *Annona exsucca* DC.

in Dunal, *Monogr. Anonac.*: 77 [Aug.-Nov. 1817] (Candolle 1817). — *Rollinia exsucca* (DC.) A.DC., *Mém. Soc. Phys. Genève* 5: 199 (Candolle 1832).

Rollinia puberula A.DC., *Mém. Soc. Phys. Genève* 5: 200 (Candolle 1832).

Rollinia multiflora Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 97 [Aug.-Sep. 1842] (Splitgerber 1842).

Rollinia tinifolia Klotzsch ex M.R.Schomb., *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1163 ["1848" publ. 7-10 Mar. 1849] (Schomburgk 1849), *nom. subnud.*

Rollinia glaucescens Miq., *Stirp. Surinam. Select.*: 108 ["1850" publ. Mar. 1851] (Miquel 1851).

Rollinia resinosa Spruce ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 9 (Bentham 1853). — *Rollinia exsucca* subsp. *resinosa* (Spruce ex Benth.) R.E.Fr., *Acta Horti Berg.* 12 (1): 144 (Fries 1934).

Rollinia incurva S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 303 ["1894-96" publ. Dec. 1895] (Moore 1895).

Rollinia brevipes R.E.Fr., *Acta Horti Berg.* 12 (1): 147 (Fries 1934).

Rollinia broadwayi R.E.Fr., *Acta Horti Berg.* 12 (1): 145 (Fries 1934).

Rollinia broadwayi var. *cuneata* R.E.Fr., *Acta Horti Berg.* 12 (1): 145 (Fries 1934).

Rollinia exsucca subsp. *elongata* R.E.Fr., *Acta Horti Berg.* 12 (1): 145 (Fries 1934).

Rollinia gardneri R.E.Fr., *Acta Horti Berg.* 12 (1): 157 (Fries 1934).

Rollinia surinamensis R.E.Fr., *Acta Horti Berg.* 12 (1): 147 (Fries 1934).

Rollinia procera R.E.Fr., *Acta Horti Berg.* 12 (3): 565 (Fries 1939).

NOTE. — Dunal (1817: 77) explicitly ascribed the name and description of *Annona exsucca* to "DC. Syst. univ. ined." See above note for *Annona exsucca*.

VERNACULAR NAMES. — Pa: manig-retni • Ka: kasimylan, tagulewe, takulewe • Wp: iwi lū • Nt: busi atuku • Cr: abriba-sovaj • Fr: corossol sauvage • Br: araticú-do-mato, araticú-macho, biribá-brava.

HERBARIUM DATA (FG). — 130 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (holotype of *Rollinia puberula*: G[G00226077]; iso-, B[B 10 0243129], F[F0047921F], S[S fragm. S-R-6899]).

INVENTORY DATA (FG). — 33 trees in 27 plots; $F_{\max} < 1\%$; $dbh_{inv} = 38$ cm.

[27] *Annona foetida* Mart.

(Fig. 7A)

Fl. Bras. [Martius] 13 (1): 16 [1 Jan. 1841] (Martius 1841).

Annona trunciflora R.E.Fr., *Mem. New York Bot. Gard.* 9 (1): 325 (Fries 1957).

VERNACULAR NAMES. — Pa: manig-ahavukune, pakih-em • Wp: iwi lo, iwi tay • Nt: iliwa • Cr: korosol-sovaj • Br: araticú-do-mato, envira-atá, graviola-da-mata.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *G. Cremers 7287*.

INVENTORY DATA (FG). — 11 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{inv} = 18.1$ cm.

[28] *Annona glabra* L.

Sp. Pl. 1: 537 [1 May 1753] (Linnaeus 1753).

Annona palustris L., *Sp. Pl.*, ed. 2, 1: 757 [Sep. 1762] (Linnaeus 1762). — *Guananabana palustris* (L.) M.Gómez, *Fl. Habanera* 114 (Gómez 1897).




Fig. 7. — Annonaceae: **A**, *Annona foetida* Mart.; **B**, *Duguetia paraensis* R.E.Fr. (D. Sabatier & J.-F. Molino 5304); **C**, *Guatteria wachenheimii* Benoist (D. Sabatier & M.-F. Prévost 3951); **D**, *Fusaea longifolia* (M.-F. Prévost & D. Sabatier 4701). © D. Sabatier/IRD.

Annona laurifolia Dunal, *Monogr. Anonac.* 65 [Aug.-Nov. 1817] (Dunal 1817). Based on “*A. glabra* var. β ” in Lam., *Encycl. [J. Lamarck et al.] 2* (1): 125. 1786 [16 Oct. 1786] (Lamarck 1786).

Annona pisonis A.St.-Hil. & Tul., *Ann. Sci. Nat., Bot. sér. 2, 17*: 131 (Saint-Hilaire & Tulasne 1842), “*Anona Pisonis*”, nom. illeg. hom., non Mart. (Martius 1841) [synonym of *A. montana* Macfad.]

Annona peruviana Humb. & Bonpl. ex Dunal, *Monogr. Anonac.* 67 [Aug.-Nov. 1817] (Dunal 1817). — *Annona uliginosa* Kunth, *Nova genera et species plantarum* [H.B.K.J] 5: 56 [28 May 1821] (Kunth 1821), nom. illeg. superfl. (based on *Annona peruviana*).

NOTE. — The epithet *pisonis* comes from *Piso*, the latinized name of Willem Pies, hence is not to be corrected (Turland et al. 2018: Art. 60.9).

Annona australis A.St.-Hil., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (1): 33 [23 Feb. 1824] (Saint-Hilaire 1824). — *Annona palustris* var. *grandifolia* Mart., *Fl. Bras. [Martius]* 13 (1): 11 [1 Jan. 1841] (Martius 1841).

VERNACULAR NAMES. — Ka: alašityulan • Cr: dji-manmen, tchò-bèf • Br: araticú-da-lagoa.

Asimina arborea Raf., *Autik. Bot.* 77 (Rafinesque 1840), nom. illeg. superfl. (based on *Annona glabra* L.)

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost 1137*.

SIZE. — Up to 15 m tall (Maas et al. 2001).

[29] *Annona hypoglauca* Mart.

Fl. Bras. [Martius] 13 (1): 13 [1 Jan. 1841] (Martius 1841).

Annona tessmannii Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 140 [1 Oct. 1924] (Diels 1924).

VERNACULAR NAMES. — Ka: woto kasimyate • Wp: iwi lū • Wn: iliwaimē • Nt: busi atuku.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2627*.

SIZE. — Up to 10 m tall (Steyermark *et al.* 1995).

[30] *Annona montana* Macfad.

Fl. Jamaica [Macfadyen] 1: 7 (Macfadyen 1837).

Annona muricata Vell., *Fl. Flumin. Icon.* 5: t. 126 ["1827" publ. 29 Oct. 1831] (Vellozo 1831), *nom. illeg. hom., non L.* (Linnaeus 1753).

Annona marcgravii Mart., *Fl. Bras. [Martius]* 13 (1): 5 [1 Jan. 1841] (Martius 1841).

Annona pisonis Mart., *Fl. Bras. [Martius]* 13 (1): 5 [1 Jan. 1841] (Martius 1841).

Annona sphaerocarpa Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 96 [Aug.-Sep. 1842] (Splitgerber 1842).

Annona montana f. *marcgravii* (Mart.) Porto, *Rodriguésia* 2: 106 (Porto 1936).

NOTES. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). See above note for *A. glabra*.

VERNACULAR NAMES. — Pa: manig • Ka: alasigun, alasityu, ulusulu • Cr: korosòl-gran-bwa • Br: araticú-açu, araticú-cagão.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P. Grenand 1594*.

SIZE. — Up to 15 m tall (Maas *et al.* 2001).

[31] *Annona mucosa* Jacq.

Observ. Bot. [Jacquin] 1: 16 (Jacquin 1764). — *Annona mucosa* Aubl., *Hist. Pl. Guiane* 1: 618 [Jun.-Dec. 1775] (Aublet 1775), sphalm. — *Annona reticulata* var. *mucosa* (Jacq.) Willd., *Sp. Pl. ed. 4* 2 (2): 1266 [Dec. 1799] (Willdenow 1799). — *Rollinia mucosa* (Jacq.) Baill., *Adansonia [Baillon]* 8: 268 (Baillon 1868).

Annona obtusiflora Tussac, *Fl. Antill.* 1: 191 (Tussac 1813). — *Annona obtusifolia* DC., *Prod. [A. P. de Candolle]* 1: 84 [mid Jan. 1824] (Candolle 1824), orth. var., based on *Annona obtusiflora* Tussac.

Annona squamosa Vell., *Fl. Flumin.* 239 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), "Anona", *nom. illeg. hom., non L.* (Linnaeus 1763) nec Vell. (Vellozo 1831).

Rollinia biflora G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831). — *Annona biflora* Ruiz & Pav. ex G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831), *nom. nud. pro syn.*

Rollinia micrantha G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831). — *Annona parviflora* Ruiz & Pav. ex G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831), *nom. nud. pro syn.* — *Annona*

parviflora Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 17 (1): 427 (Ruiz & Pavón 1960).

Rollinia pterocarpa G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831). — *Annona pterocarpa* G.Don, *Gen. Hist.* 1: 91 [early Aug. 1831] (Don 1831), *nom. nud. pro syn., orth. var.* — *Annona pteropetala* Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 17 (1): 426 (Ruiz & Pavón 1960). — *Annona pteropetala* Ruiz & Pav. ex R.E.Fr., *Acta Horti Berg.* 12 (1): 187 (Fries 1934), *nom. nud.*

Rollinia orthopetala A.DC., *Mém. Soc. Phys. Genève* 5: 200 (Candolle 1832).

Annona reticulata Sieber ex A.DC., *Mém. Soc. Phys. Genève* 5: 200 (Candolle 1832), *nom. illeg. hom., non L.* (Linnaeus 1763) nec Vell. (Vellozo 1831).

Rollinia sieberi A.DC., *Mém. Soc. Phys. Genève* 5: 200 (Candolle 1832).

Rollinia pulchrinervia A.DC., *Mém. Soc. Phys. Genève* 5: 201 (Candolle 1832).

Annona biflora Sessé & Moc., *Fl. Mexic.*, ed. 2, 133 (Sessé & Mociño 1894).

Rollinia deliciosa Saff., *J. Wash. Acad. Sci.* 6: 375 (Safford 1916).

Rollinia jimenezii Saff., *J. Wash. Acad. Sci.* 6: 378 (Safford 1916).

Rollinia permensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4* (8): 208 [24 Oct. 1929] (Standley 1929).

Rollinia mucosa var. *macropoda* R.E.Fr., *Acta Horti Berg.* 12 (1): 122 (Fries 1934).

Rollinia mucosa subsp. *aequatorialis* R.E.Fr., *Acta Horti Berg.* 12 (1): 123 (Fries 1934).

Rollinia mucosa subsp. *portoricensis* R.E.Fr., *Acta Horti Berg.* 12 (1): 124 (Fries 1934).

Rollinia neglecta R.E.Fr., *Acta Horti Berg.* 12 (1): 124 (Fries 1934). — *Rollinia mucosa* var. *neglecta* (R.E.Fr.) R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13* (2.3): 760 [31 Oct. 1938] (Fries 1938).

Rollinia curvipetala R.E.Fr., *Acta Horti Berg.* 12 (1): 178 (Fries 1934).

Rollinia jimenezii var. *nelsonii* R.E.Fr., *Kungl. Svenska Vetenskapsakad. Handl.*, ser. 3, 24 (10): 18 (Fries 1948).

NOTES. — Don (1831: 91) gave in synonymy under *Rollinia pterocarpa* "Annona pterocarpa Ruiz & Pav. fl. per. 5 t. 483". This is a clear reference to *Annona pteropetala* Ruiz & Pav., despite the miscitation of plate 484, giving 483 instead. Similarly, under *Rollinia micrantha* Don mistakenly cited plate 484 rather than 485 for *A. parviflora* (Don 1831: 91).

VERNACULAR NAMES. — Pa: manig-amutri • Ka: kasimya, sokowe • Wp: iwi li • Wn: iliwa • Cr: abriba, korosòl-sovaj • Br: atá-preta, biriba.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost 3379*, 10 m × 12 cm.

[32] *Annona neoelliptica* H.Rainer & Maas

Ann. Naturhist. Mus. Wien 108B: 199 (Rainer & Maas 2007). — *Rollinia elliptica* R.E.Fr., *Acta Horti Berg.* 12 (1): 160 (Fries 1934).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P.J.M. Maas et al. 8062*.

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 42.5 cm.

[33] *Annona paludosa* Aubl.

Hist. Pl. Guiane 1: 611 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Restricted to coastal marsh forests.

VERNACULAR NAMES. — Pa: kasim • Cr: dji-manmen • Br: araticú-do-brejo.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material MPU? [MPU014808]).

SIZE. — Brazil, Maranhão. *N.A. Rosa 2812* (MO), 12 m × 40 cm.

[34] *Annona prevostiae* H.Rainer

Brittonia 54 (2): 136 (Rainer 2002).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *M.-F. Prévost 3742* (holo-, CAY [CAY018189]; iso-, NY [not seen], WU [WU0026227, not seen]).

INVENTORY DATA (FG). — 16 trees in 12 plots; $F_{\max} < 1\%$; dbh_{inv} = 36.6 cm.

[35] *Annona sericea* Dunal

Monogr. Anonac. 69 [Aug.-Nov. 1817] (Dunal 1817).

Annona jenmanii Saff., *Contr. U.S. Natl. Herb.* 16: 267 (Safford 1913).

Annona trinitensis Saff., *Contr. U.S. Natl. Herb.* 16: 268 (Safford 1913).

VERNACULAR NAMES. — Pa: kasim-kamwi • Ka: ulusulu, yapopale • Wp: iwilū e'e • Br: araticú-do-Pará, atá-brava.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (holo-, G-DC [G00201278]).

SIZE. — Up to 20 m tall (Steyermark et al. 1995).

[36] *Annona symplycarpa* Sandwith

Bull. Misc. Inform. Kew 1930 (10): 477 [4 Dec. 1930] (Sandwith 1930).

Annona tenuipes R.E.Fr., *Ark. Bot. n.s.*, 1 (11): 448 (Fries 1950).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 10907*.

SIZE. — Up to 25 cm dbh (Steyermark et al. 1995).

[37] *Annona tenuiflora* Mart.

Fl. Bras. [Martius] 13 (1): 10 [1 Jan. 1841] (Martius 1841).

Raimondia tenuiflora (Mart.) R.E.Fr., *Acta Horti Berg.* 10 (1): 85 (Fries 1930).

Annona humilis Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 113 (Bentham 1851).

Annona poeppigiana Saff. ex R.E.Fr., *Acta Horti Berg.* 10 (1): 85 (Fries 1930), *nom. nud.*

VERNACULAR NAMES. — Pa: manig-ihipinē, manig-kamwi.

HERBARIUM DATA (FG). — A single collection, *P. Grenand & M.-F. Prévost 2040*.

SIZE. — Ecuador, Chimborazo. *W.H. Camp E-3425* (MO), 10 m tall.

Genus *Bocageopsis* R.E.Fr.

[38] *Bocageopsis multiflora* (Mart.) R.E.Fr.

Acta Horti Berg. 10 (2): 145 (Fries 1931). — *Bocagea multiflora* Mart., *Fl. Bras. [Martius]* 13 (1): 45 [1 Jan. 1841] (Martius 1841).

Guatteria multiflora Poepp. ex Baill., *Hist. Pl. [Baillon]* 1: 216 [Aug.-Dec. 1868] (Baillon 1868), *nom. nud.*

Bocageopsis multiflora var. *angustifolia* R.E.Fr., *Acta Horti Berg.* 10 (2): 146 (Fries 1931).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Br: envira-preta, envira-surucucu-folhamiúda.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2290*.

INVENTORY DATA (FG). — 19 trees in 10 plots; $F_{\max} < 1\%$; dbh_{inv} = 34.7 cm.

Genus *Cardiopetalum* Schltdl.

[39] *Cardiopetalum surinamense* R.E.Fr.

Acta Horti Berg. 10 (2): 195 (Fries 1931). — *Froesiodendron surinamense* (R.E.Fr.) R.E.Fr., *Ark. Bot. n.s.*, 3: 442 (Fries 1956).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier 3528*, 15 m × 18 cm.

Genus *Cremastosperma* R.E.Fr.

[40] *Cremastosperma brevipes* (DC.) R.E.Fr.

Acta Horti Berg. 12 (3): 538 (Fries 1939). — *Guatteria brevipes* DC. in Dunal, *Monogr. Anonac.*: 126 [Aug.-Nov. 1817] (Candolle 1817).

Guatteria poiteaui Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 74 [31 Mar. 1931] (Diels 1931). — *Cremastosperma poiteaui* (Diels) R.E.Fr., *Acta Horti Berg.* 10 (2): 328 (Fries 1931).

NOTES. — Known only from French Guiana. Dunal (1817: 126) explicitly ascribed the name and description of *Guatteria brevipes* to "DC. Syst. univ. ined." See above note for *Annona exsucca*.

VERNACULAR NAMES. — Wp: apelemu'i.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material K[K000485530]); *P.A. Poiteau s.n.* (holotype of *Guatteria poiteauii*: G[G00237256]; iso-, B[B 10 0242372], F[F0047724F], G[G00237255], S[S-R-7016]).

INVENTORY DATA (FG). — 12 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.6$ cm.

Genus *Cymbopetalum* Benth.

[41] *Cymbopetalum brasiliense* (Vell.) Benth. ex Baill.

Hist. Pl. [Baillon] 1: 240 [Aug.-Dec. 1868] (Baillon 1868). — *Uvaria brasiliensis* Vell., *Fl. Flumin.* 238 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829). — *Uva brasiliensis* (Vell.) Kuntze, *Revis. Gen. Pl.* 1: 7 [5 Nov. 1891] (Kuntze 1891).

Unona viridiflora Splitg. ex de Vriese, *Ned. Kruidk. Arch.* 1 (3): 224 (de Vriese 1847). — *Uvaria viridiflora* (Splitg. ex de Vriese) Walp., *Ann. Bot. Syst. [Walpers]* 2 (1): 19 [15-16 Dec. 1851] (Walpers 1851), nom. illeg. hom., non Ruiz & Pav. ex G.Don (1831).

Eschweilera simplex Miers, *Trans. Linn. Soc. London* 30 (2): 264 [14 Nov. 1874] (Miers 1874).

Cymbopetalum odoratissimum Barb.Rodr., *Vellozia*, ed. 2, 1: 1 (Barbosa Rodrigues 1891).

Cymbopetalum brasiliense f. *latifolia* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 357 (Huber 1909), nom. nud.

Trigynaea anastomosans Rusby, *Descr. S. Amer. Pl.*: 19 [20 Dec. 1920] (Rusby 1920).

VERNACULAR NAMES. — Pa: pakih-em-purubumna • Ka: kaikushi anyali • Wp: kwata laway, sa'i melu ka'a wu, yawa popita • Wn: alukaju, kaston • Br: envira-da-mata, pirayaaura-kinha.

HERBARIUM DATA (FG). — 66 collections at CAY. Sel. exs.: *R.A.A. Oldeman 3066*.

SIZE. — Brazil, Pará. *J.J. Strudwick et al. 3121* (MO), 8 m × 10 cm.

Genus *Duguetia* A.St.-Hil.

[42] *Duguetia cadaverica* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 356 (Huber 1909). — *Ganthemnum cadavericum* (Huber) Saff., *Contr. U.S. Natl. Herb.* 18 (1): 67 [17 June 1914] (Safford 1914).

Duguetia adiscandra Jans.-Jac., *Proc. Kon. Ned. Akad. Wetensch. C* 73 (4): 339 (Jansen-Jacobs 1970).

Duguetia friesii Jans.-Jac., *Proc. Kon. Ned. Akad. Wetensch. C* 73 (4): 338 (Jansen-Jacobs 1970).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: kaabasi ana, pepee anga sawtu.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3306*.

INVENTORY DATA (FG). — 1 tree, dbh = 10.1 cm.

[43] *Duguetia calycina* Benoist

Notul. Syst. (Paris) 4: 57 (Benoist 1923).

Annona calycina Sagot ex R.E.Fr., *Acta Horti Berg.* 12 (1): 69 (Fries 1934), nom. nud.

Duguetia calycina subsp. *versteegii* R.E.Fr., *Acta Horti Berg.* 12 (1): 72 (Fries 1934).

Duguetia calycina subsp. *jenmanii* R.E.Fr., *Acta Horti Berg.* 12 (1): 73 (Fries 1934).

Duguetia cuspidata R.E.Fr., *Acta Horti Berg.* 12 (1): 73 (Fries 1934).

Duguetia elegans R.E.Fr., *Acta Horti Berg.* 12 (1): 75 (Fries 1934).

VERNACULAR NAMES. — Pa: korosol-kamwi • Ka: kapasi pomiidiyi, yolokan pomiidiyi • Wp: pila siay, pina'i, pina'i piyū, pina'i tay • Wn: iliwaimé • Nt: gaan pepee anga sawtu, gaan uku tiki, pepee anga sawtu • Cr: baton-zin, manmanyawé-piman • Br: biribarana, envira-caitetu.

HERBARIUM DATA (FG). — 157 collections at CAY. Sel. exs.: *P.A. Sagot 1095*, Dec. 1857 (lecto-, P[P00083598]), designated by Fries [1934: 69]; isolecto-, BR[BR000006952372, BR000006952686], P[P00083611], S[S-R-1796], U[U0000257]).

INVENTORY DATA (FG). — 230 trees in 89 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 19.1$ cm.

[44] *Duguetia eximia* Diels

Notizbl. Bot. Gart. Berlin-Dahlem 11: 79 [31 Mar. 1931] (Diels 1931).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: pakih-em-priye • Wp: mitū luway, pina'i u • Nt: pepee anga sawtu • Cr: manmanyawé-piman.

HERBARIUM DATA (FG). — 70 collections at CAY. Sel. exs.: *G.S. Perrottet s.n.* (holo-, G[G00237263], fragm. B[B 10 1093263]).

SIZE. — Up to 10 cm dbh (Maas *et al.* 2003).

[45] *Duguetia granvilleana* Maas

Bot. Jahrb. Syst. 121 (4): 476 (Maas 1999).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 6330* (holo-, U[U0008263]; iso-, CAY[CAY003561]); *O. Poncy et al. 1748*: 18 m × 18 cm.

[46] *Duguetia paraensis* R.E.Fr.

(Fig. 7B)

Acta Horti Berg. 13 (3): 111 (Fries 1941).

VERNACULAR NAMES. — Wp: apelemu'i, pina'i sī, pina'i sili, pina'i to wi'i, uki'i • Cr: manmanyawé-piman • Br: amejó, meijú.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5304*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.1$ cm.

[47] *Duguetia pycnastera* Sandwith

Bull. Misc. Inform. Kew 1930 (10): 471 [4 Dec. 1930] (Sandwith 1930).

VERNACULAR NAMES. — Ka: tipulu alaidya, yolokan pomiydyi • Nt: gaan kaabasi ana, gaan uku tiki, kaabasi ana, pepee anga sawtu • Br: envira-preta.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2956*.

INVENTORY DATA (FG). — 24 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 13.1$ cm.

[48] *Duguetia riparia* Huber

Bol. Mus. Paraense Hist. Nat. Ethnogr. 3: 416 (Huber 1902).

Duguetia insculpta R.E.Fr., *Acta Horti Berg.* 12 (1): 91 (Fries 1934).

VERNACULAR NAMES. — Pa: pakih-em • Wp: pina'i teáká lupiwa • Wn: oka epu • Cr: baton-zin, manmanyawé • Br: envira-preta, envira-taia.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *P. Grenand 1845*.

SIZE. — Up to 15 cm dbh (Maas et al. 2003).

[49] *Duguetia surinamensis* R.E.Fr.

Acta Horti Berg. 12 (1): 50 (Fries 1934).

Duguetia caudata R.E.Fr., *Acta Horti Berg.* 12 (2): 274 (Fries 1937).

VERNACULAR NAMES. — Wp: inámu pita, mitū luway, pila sila, pina'i sī, pina'i to wu • Nt: pepee anga sawtu • Cr: manmanyawé-piman • Br: atá-brava, biribarana, envira-amargosa.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2560*.

INVENTORY DATA (FG). — 195 trees in 61 plots; $F_{\max} = 4.2\%$; $dbh_{\text{inv}} = 35.8$ cm.

[50] *Duguetia trunciflora* Maas & A.H.Gentry

Feddes Repert. 106 (1995): 341 (Maas & Gentry 1996).

VERNACULAR NAMES. — Nt: gaan kaabasi ana.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & B. Riéra 3113*.

INVENTORY DATA (FG). — 12 trees in 2 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 14$ cm.

[51] *Duguetia yeshidan* Sandwith

Bull. Misc. Inform. Kew 1930 (10): 473 [4 Dec. 1930] (Sandwith 1930).

Duguetia obovata R.E.Fr., *Acta Horti Berg.* 12 (1): 66 (Fries 1934).

Duguetia sandwithii R.E.Fr., *Acta Horti Berg.* 12 (1): 80 (Fries 1934).

VERNACULAR NAMES. — Pa: pakih-em, wiratay • Ka: tamunen isyano manduliyi • Wp: pila siay, pina'i, pina'i sī, pina'i to wi'i, uki'i • Nt: kaabasi ana • Cr: manmanyawé-piman • Br: envira-caitetu.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *D. Sabatier 922*.

INVENTORY DATA (FG). — 1 tree, $dbh = 12.3$ cm.

Genus *Ephedranthus* S.Moore

[52] *Ephedranthus guianensis* R.E.Fr.

Acta Horti Berg. 10 (2): 176 (Fries 1931).

Ephedranthus guianensis var. *oligantha* R.E.Fr., *Acta Horti Berg.* 10 (2): 177 (Fries 1931).

VERNACULAR NAMES. — Wn: iliwaimë.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *R. Benoist 1159* (original material of *E. guianensis* var. *oligantha*: P [P00751066, P00751067]).

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 21.3$ cm.

Genus *Fusaea* (Baill.) Saff.

[53] *Fusaea longifolia* (Aubl.) Saff.
(Fig. 7D)

Contr. U.S. Natl. Herb. 18 (1): 65 [17 June 1914] (Safford 1914). — *Annona longifolia* Aubl., *Hist. Pl. Guiane* 1: 615 [Jun.-Dec. 1775] (Aublet 1775). — *Abermoa longifolia* (Aubl.) Baill., *Hist. Pl. [Baillon]* 1: 205 [Aug.-Dec. 1868] (Baillon 1868). — *Duguetia longifolia* (Aubl.) Baill., *Adansonia [Baillon]* 8: 327 (Baillon 1868).

Uvaria spectabilis DC. in Dunal, *Monogr. Anonac.* 92 [Aug.-Nov. 1817] (Candolle 1817).

Annona rhombipetala Ruiz & Pav. ex G.Don, *Gen. Hist.* 1: 87 [early Aug. 1831] (Don 1831). — *Fusaea rhombipetala* (Ruiz & Pav. ex G.Don) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (7): 172 [24 Oct. 1929] (Macbride 1929).

Fusaea decurrens R.E.Fr., *Acta Horti Berg.* 12 (1): 207 (Fries 1934).

NOTES. — Dunal (1817: 92) explicitly ascribed the name and description of *Uvaria spectabilis* to “DC. Syst. univ. ined.” See above note for *Annona exsucca*.

VERNACULAR NAMES. — Pa: kuukumwi-priye • Wp: yáwí kala, yáwí yiki • Nt: maisyi • Cr: manmanyawé • Br: envira-preta.

HERBARIUM DATA (FG). — 101 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000547151]).

INVENTORY DATA (FG). — 78 trees in 53 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.5$ cm.

Genus *Guatteria* Ruiz & Pav.

[54] *Guatteria anteridifera* Scharf & Maas

Blumea 53 (3): 515 (Scharf & Maas 2008).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kigiksau, kiriksau.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 5047 (holo-, U[U0227761, U0073311]; iso-, CAY[CAY073329, CAY073330]).

INVENTORY DATA (FG). — 30 trees in 8 plots; $F_{\max} = 2.7\%$; dbh_{inv} = 49 cm.

[55] *Guatteria blepharophylla* Mart.

Fl. Bras. [Martius] 13 (1): 38 [1 Jan. 1841] (Martius 1841). — *Guatteriopsis blepharophylla* (Mart.) R.E.Fr., *Acta Horti Berg.* 12 (1): 110 (Fries 1934).

Guatteria oxyacarpa Poepp. ex Mart., *Fl. Bras. [Martius]* 13 (1): 38 [1 Jan. 1841] (Martius 1841), nom. nud. pro syn.

Annona sessiliflora Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 8 (Bentham 1853). — *Guatteria sessiliflora* (Benth.) Saff., *Contr. U.S. Natl. Herb.* 18 (1): 6 [17 June 1914] (Safford 1914). — *Guatteriopsis sessiliflora* (Benth.) R.E.Fr., *Acta Horti Berg.* 12 (1): 109 (Fries 1934).

Guatteria ucayaliana Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 138 (Diels 1924).

Guatteria dielsiana R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (2.3)*: 719 (Fries 1938), nom. illeg. superfl. (based on *Guatteria ucayaliana* Diels [synonym of *G. blepharophylla* Mart.].)

Guatteria brevicuspis R.E.Fr., *Acta Horti Berg.* 12 (3): 491 (Fries 1939).

Guatteria cylindrocarpa R.E.Fr., *Ark. Bot., n.s.*, 3 (18): 601 (Fries 1956).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al.* 25177.

SIZE. — Up to 10 cm dbh (Maas & Westra 2011).

[56] *Guatteria citriodora* Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 104 (Ducke 1930).

Guatteria paraensis R.E.Fr., *Acta Horti Berg.* 12 (3): 464 (Fries 1939).

Guatteria duckeana R.E.Fr. var. *subcordata* R.E.Fr., *Kungl. Svenska Vetenskapsakad. Handl.*, ser. 3, 24 (10): 10 (Fries 1948).

VERNACULAR NAMES. — Br: envira-amarela, laranjinha.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *J.-F. Molino* & *D. Sabatier* 1968.

INVENTORY DATA (FG). — 31 trees in 20 plots; $F_{\max} < 1\%$; dbh_{inv} = 39 cm.

[57] *Guatteria conspicua* R.E.Fr.

Ark. Bot., n.s., 1 (11): 445 (Fries 1950).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier* 4682.

INVENTORY DATA (FG). — 7 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 26.3 cm.

[58] *Guatteria discolor* R.E.Fr.

Acta Horti Berg. 12 (3): 509 (Fries 1939).

VERNACULAR NAMES. — Pa: miret • Wp: matau'i • Br: envira-fofa, envira-preta, envira-rolinha.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M.-F. Prévost* & *D. Sabatier* 2773, dbh 40 cm.

[59] *Guatteria foliosa* Benth.

London J. Bot. 2: 360 (Bentham 1843).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori* & *C. Gracie* 23929.

SIZE. — Up to 28 cm dbh (Maas & Maas-van de Kamer 2002).

[60] *Guatteria guianensis* (Aubl.) R.E.Fr.

Acta Horti Berg. 12 (3): 505 (Fries 1939). — *Aberemoa guianensis* Aubl., *Hist. Pl. Guiane* 1: 610 [Jun.-Dec. 1775] (Aublet 1775).

Guatteria aberemoa Dunal, *Monogr. Anonac.* 126 [Aug.-Nov. 1817] (Dunal 1817).

Guatteria aberemoa var. *microcarpa* DC., *Syst. Nat. [A. P. de Candolle]* 1: 502 ["1818" publ. 1-15 Nov. 1817] (Candolle 1817).

Guatteria multivenia Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 171 (Diels 1927).

Guatteria excellens R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (2.3)*: 721 (Fries 1938).

Guatteria calophylla R.E.Fr., *Acta Horti Berg.* 12 (3): 507 (Fries 1939).

Guatteria robusta R.E.Fr., *Mem. New York Bot. Gard.* 9 (1): 328 (Fries 1957).

NOTE. — Dunal (1817: 126) ascribed the description of *Guatteria aberemoa* to "DC. Syst. univ. ined.", but not the name itself.

VERNACULAR NAMES. — Pa: kuuku-ára-purubumna, kuukumwi-purubumna • Wp: pina'i piyū • Cr: manmanyawé • Br: envira-da-mata.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (lectotype BM [BM000547286] designated by Fries (1939)).

INVENTORY DATA (FG). — 29 trees in 26 plots; $F_{\max} < 1\%$; dbh_{inv} = 15.6 cm.

[61] *Guatteria intermedia* Scharf
(Fig. 8A)

Blumea 51 (3): 545 (Scharf 2006).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: payulilan, wayilu • Wp: iwi.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: R.A.A. Oldeman B-4125 (holo-, CAY[CAY100867, CAY100868]; iso-, P[P00115965]).

INVENTORY DATA (FG). — 18 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{inv} = 19.4$ cm.

[62] *Guatteria leucotricha* Scharf & Maas

Blumea 51 (1): 123 (Scharf & Maas 2006).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: S.A. Mori & B.M. Boom 15360 (holo-, NY[00807020]; iso-, CAY[CAY034217], K[K001096423], P[P00115992], U[U0126456], US[01123113]).

SIZE. — Up to 45 cm dbh (Scharf *et al.* 2006).

[63] *Guatteria liesneri* D.M.Johnson & N.A.Murray

Ann. Missouri Bot. Gard. 77 (3): 598 (Johnson & Murray 1990).

Guatteria anthracina Scharf & Maas, *Blumea* 51 (1): 118 (Scharf & Maas 2006).

VERNACULAR NAMES. — Wn: kaloj ēwa, oka epu • Nt: uku tiki.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: D. Sabatier & M.-F. Prévost 3660.

INVENTORY DATA (FG). — 16 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{inv} = 35.7$ cm.

[64] *Guatteria oblonga* R.E.Fr.

Acta Horti Berg. 12 (3): 443 (Fries 1939).

HERBARIUM DATA (FG). — Known only from the type: E.M. Mélinon s.n. (original material B not seen, P[P00115935, P00115936]).

SIZE. — Not recorded on the type specimen, but like all *Guatteria* spp. in French Guiana, it is probably a tree.

[65] *Guatteria ouregou* (Aubl.) Dunal
(Fig. 8B)

Monogr. Anonac. 126 [Aug.-Nov. 1817] (Dunal 1817). — *Cananga ouregou* Aubl., *Hist. Pl. Guiane* 1: 607 [Jun.-Dec. 1775] (Aublet 1775). — *Uvaria ouregou* (Aubl.) Raeusch., *Nomencl. Bot. [Raeusch.]*, ed. 3: 160 (Raeuschel 1797), “Uregou”. — *Uvaria monosperma* Lam., *Encycl. [J. Lamarck et al.]* 7: 596 [6 July 1806] (Lamarck 1806), nom. illeg. superfl. (based on *Cananga ouregou*).

Guatteria podocarpa DC. in Dunal, *Monogr. Anonac.* 127 [Aug.-Nov. 1817] (Candolle 1817).

Unona crassipetala Dunal, *Monogr. Anonac.* 101 [Aug.-Nov. 1817] (Dunal 1817). — *Guatteria crassipetala* (Dunal) DC. ex Lemée, *Fl. Guyane Franç.* 1: 607 (Lemée 1955). — *Unona pachypetala* Spreng., *Syst. Veg. [Sprengel]* 2: 637 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Unona crassipetala*).

Unona fuscata DC. in Dunal, *Monogr. Anonac.* 102 [Aug.-Nov. 1817] (Candolle 1817).

Guatteria podocarpa var. *oligocarpa* DC., *Syst. Nat. [A. P. de Candolle]* 1: 503 [“1818” publ. 1-15 Nov. 1817] (Candolle 1817).

Guatteria podocarpa var. *polycarpa* DC., *Syst. Nat. [A. P. de Candolle]* 1: 503 [“1818” publ. 1-15 Nov. 1817] (Candolle 1817).

Guatteria ouregou var. *latifolia* Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 11: 138 (Sagot 1881).

NOTES. — Known only from French Guiana. *Sagot s.n.* (P00115911) has been suggested as a possible isotype of *G. podocarpa* var. *oligocarpa* by U. Scharf (annotation dated 2008), but this is unrealistic since this collection dates from 1859. Dunal (1817: 127) explicitly ascribed the name and description of *G. podocarpa* to “DC. Syst. univ. ined.” See above note for *Annona exsucca*.

VERNACULAR NAMES. — Pa: kigiksa, kiriksa • Ka: ouregou (*fide* Aublet 1775) • Wp: iwi.

HERBARIUM DATA (FG). — 81 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (original material of *Unona crassipetala*: G-DC[G00201449]).

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{inv} = 34.4$ cm.

[66] *Guatteria pannosa* Scharf & Maas

Blumea 51 (1): 117 (Scharf & Maas 2006).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: S. Barrier & C. Feuillet 2633 (holo-, CAY[CAY104592]).

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 52.5$ cm.

[67] *Guatteria punctata* (Aubl.) R.A.Howard

J. Arnold Arbor. 64 (2): 260 [Apr. 1983] (Howard 1983). — *Annona punctata* Aubl., *Hist. Pl. Guiane* 1: 614 [Jun.-Dec. 1775] (Aublet 1775).

Guatteria glauca Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 145 [late Dec. 1798] (Ruiz & Pavón 1798).

Annona axilliflora DC., *Prodr. [A. P. de Candolle]* 1: 86 [mid Jan. 1824] (Candolle 1824). — *Guatteria axilliflora* (DC.) R.E.Fr., *Acta Horti Berg.* 12 (3): 427 (Fries 1939).

Guatteria poeppigiana Mart., *Fl. Bras. [Martius]* 13 (1): 36 [1 Jan. 1841] (Martius 1841), “*pöppigiana*”.

Guatteria caniflora Mart., *Fl. Bras. [Martius]* 13 (1): 37 [1 Jan. 1841] (Martius 1841).



FIG. 8. — Annonaceae: **A**, *Guatteria intermedia* Scharf (J.-F. Molino & D. Sabatier 2464); **B**, *Guatteria ouregou* (Aubl.) Dunal (J.-F. Molino & D. Sabatier 2736); **C**, *Trigynaea caudata* (R.E.Fr.) R.E.Fr. (J.-F. Molino & D. Sabatier 2158); **D**, *Unonopsis perrottetii* (A.DC.) R.E.Fr. (J.-F. Molino & D. Sabatier 2713). A-C, © J.-F. Molino/IRD; D, © D. Sabatier/IRD.

Guatteria caniflora var. *angustifolia* Mart., *Fl. Bras. [Martius]* 13 (1): 37 [1 Jan. 1841] (Martius 1841).

Guatteria caniflora var. *latifolia* Mart., *Fl. Bras. [Martius]* 13 (1): 37 [1 Jan. 1841] (Martius 1841).

Annona chrysopetala Steud., *Flora* 26 (45): 754 [17 Dec. 1843] (Steudel 1843). — *Guatteria chrysopetala* (Steud.) Miq., *Linnæa* 22: 466 (Miquel 1849).

Guatteria pteropus Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 8 (Bentham 1853).

Guatteria cargadero Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 17: 34 (Triana & Planchon 1862).

Guatteria platyphylla Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 17: 35 (Triana & Planchon 1862).

- Guatteria sylvicola* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 298 ["1894-96" publ. Dec. 1895] (Moore 1895).
- Guatteria coeloneura* Diels, *Bot. Jahrb. Syst.* 37 (4): 408 (Diels 1906).
- Guatteria pleiocarpa* Diels, *Bot. Jahrb. Syst.* 37 (4): 409 (Diels 1906).
- Guatteria guentheri* Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 169 (Diels 1927).
- Guatteria atra* Sandwith, *Bull. Misc. Inform. Kew* 1930 (10): 468 (Sandwith 1930).
- Guatteria acutissima* R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.3): 712 (Fries 1938).
- Guatteria calliantha* R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.3): 715 (Fries 1938).
- Guatteria juninensis* R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.3): 716 (Fries 1938).
- Guatteria collina* R.E.Fr., *Acta Horti Berg.* 12 (3): 381 (Fries 1939).
- Guatteria buchtienii* R.E.Fr., *Acta Horti Berg.* 12 (3): 388 (Fries 1939).
- Guatteria lasiocalyx* R.E.Fr., *Acta Horti Berg.* 12 (3): 388 (Fries 1939).
- Guatteria rhamnoides* R.E.Fr., *Acta Horti Berg.* 12 (3): 389 (Fries 1939).
- Guatteria pteropus* var. *angustior* R.E.Fr., *Acta Horti Berg.* 12 (3): 420 (Fries 1939).
- Guatteria pteropus* var. *cinerea* R.E.Fr., *Acta Horti Berg.* 12 (3): 420 (Fries 1939).
- Guatteria olivacea* R.E.Fr., *Acta Horti Berg.* 12 (3): 423 (Fries 1939).
- Guatteria obliqua* R.E.Fr., *Acta Horti Berg.* 12 (3): 424 (Fries 1939).
- Guatteria ovalifolia* R.E.Fr., *Acta Horti Berg.* 12 (3): 428 (Fries 1939).
- Guatteria occidentalis* R.E.Fr., *Acta Horti Berg.* 12 (3): 430 (Fries 1939).
- Guatteria chrysopetala* var. *major* R.E.Fr., *Acta Horti Berg.* 12 (3): 435 (Fries 1939).
- Guatteria chrysopetala* var. *tenuipes* R.E.Fr., *Acta Horti Berg.* 12 (3): 435 (Fries 1939).
- Guatteria sagotiana* R.E.Fr., *Acta Horti Berg.* 12 (3): 437 (Fries 1939).
- Guatteria sagotiana* var. *gracilior* R.E.Fr., *Acta Horti Berg.* 12 (3): 437 (Fries 1939).
- Guatteria gracilipes* R.E.Fr., *Acta Horti Berg.* 12 (3): 438 (Fries 1939).
- Guatteria longestipitata* R.E.Fr., *Acta Horti Berg.* 12 (3): 438 (Fries 1939).
- Guatteria parviflora* R.E.Fr., *Acta Horti Berg.* 12 (3): 440 (Fries 1939).
- Guatteria lanceolata* R.E.Fr., *Acta Horti Berg.* 12 (3): 443 (Fries 1939).
- Guatteria elliptica* R.E.Fr., *Acta Horti Berg.* 12 (3): 445 (Fries 1939).
- Guatteria umbonata* R.E.Fr., *Acta Horti Berg.* 12 (3): 447 (Fries 1939).
- Guatteria gamosepala* R.E.Fr., *Acta Horti Berg.* 12 (3): 528 (Fries 1939).
- Guatteria leiocarpa* R.E.Fr., *Acta Horti Berg.* 12 (3): 530 (Fries 1939).
- Guatteria macrocalyx* R.E.Fr., *Acta Horti Berg.* 12 (3): 530 (Fries 1939).

Guatteria calimensis R.E.Fr., *Ark. Bot., n.s.*, 1: 332 (Fries 1950).

Guatteria latipetala R.E.Fr., *Ark. Bot., n.s.*, 3: 602 (Fries 1956).

Guatteria asplundiana R.E.Fr., *Ark. Bot., n.s.*, 4: 24 (Fries 1957).

Guatteria wessels-boerii Jans.-Jac., *Proc. Kon. Ned. Akad. Wetensch. C* 73 (4): 336 (Jansen-Jacobs 1970).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kigiksau, kiriksau • Ka: payulilan, wayilu • Te: wiwit hun • Wp: iwi, iwi lū sī, iwi lū sili, pina'i piyū • Wn: ēwok pokan • Nt: podo tiki • Cr: manmanyawé-yanman • Br: envira-preta, envira-surucucu.

HERBARIUM DATA (FG). — 171 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000547301](#)]).

INVENTORY DATA (FG). — 20 trees in 19 plots; $F_{\max} < 1\%$; dbh_{inv} = 35 cm.

[68] *Guatteria richardii* R.E.Fr.

Acta Horti Berg. 12 (3): 440 (Fries 1939).

Guatteria montis-trinitatis Scharf, *Blumea* 51 (3): 548 (Scharf 2006).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holo-, P[[P00115932](#)]).

INVENTORY DATA (FG). — 1 tree, dbh = 38.4 cm.

[69] *Guatteria schomburgkiana* Mart.

Fl. Bras. [Martius] 13 (1): 38 [1 Jan. 1841] (Martius 1841). — *Cananga schomburgkiana* (Mart.) Baill., *Hist. Pl. [Baillon]* 1: 204 [Aug.-Dec. 1868] (Baillon 1868).

Annona hostmannii Steud., *Flora* 26 (45): 754 [17 Dec. 1843] (Steudel 1843).

Guatteria vestita Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 979 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Guatteria vestita Klotzsch var. *angustifolia* Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 979 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Guatteria vestita var. *latifolia* Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 979 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Guatteria schomburgkiana var. *angustifolia* Klotzsch ex R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 17 (Fries 1900).

Guatteria schomburgkiana var. *latifolia* Klotzsch ex R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 17 (Fries 1900).

Guatteria sessilis R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 17 (Fries 1900).

Guatteria schomburgkiana var. *holosericea* R.E.Fr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.3): 709 (Fries 1938).

Guatteria sandwithii R.E.Fr., *Acta Horti Berg.* 12 (3): 466 (Fries 1939).

Guatteria spruceana R.E.Fr., *Acta Horti Berg.* 12 (3): 469 (Fries 1939).

Guatteria flavovirens R.E.Fr., *Kungl. Svenska Vetenskapsakad. Handl.*, ser. 3, 24 (10): 10 (Fries 1948).

VERNACULAR NAMES. — Ka: payulilan, wayilu • Wp: pina'i sili, pina'i to wi'i.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1703*.

INVENTORY DATA (FG). — 35 trees in 26 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 39.6 \text{ cm}$.

[70] *Guatteria scytophylla* Diels

Verh. Bot. Vereins Prov. Brandenburg 47: 127 [1 Oct. 1905] (Diels 1905).

Guatteria hyposericea Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 76 [31 Mar. 1931] (Diels 1931).

Guatteria insignis R.E.Fr., *Acta Horti Berg.* 12 (3): 449 (Fries 1939).

Guatteria krukoffii R.E.Fr., *Acta Horti Berg.* 12 (3): 442 (Fries 1939).

Guatteria micans R.E.Fr., *Acta Horti Berg.* 12 (3): 451 (Fries 1939).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 5649*, dbh 23.1 cm.

[71] *Guatteria wachenheimii* Benoist
(Fig. 8C)

Bull. Mus. Natl. Hist. Nat. 33: 270 (Benoist 1927).

Guatteria microsperma R.E.Fr., *Brittonia* 8 (4): 236 [Jan. 1957] (Fries 1957).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kigksau, kiriksau, kuukumwi-priye • Br: envira-cheirosa.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *G. Wachenheim 201* (holo-, P[P00115810]; iso-, P[P00115809], S[S-R-2643, S05-6707], US[00810934]).

INVENTORY DATA (FG). — 33 trees in 20 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22.4 \text{ cm}$.

Genus *Oxandra* A.Rich.

[72] *Oxandra asbeckii* (Pulle) R.E.Fr.

Acta Horti Berg. 10 (2): 167 (Fries 1931). — *Bocagea asbeckii* Pulle, *Recueil Trav. Bot. Néerl.* 6: 262 (Pulle 1909).

VERNACULAR NAMES. — Pa: kuukumwi-priye • Ka: kowai epi, pililipyo • Wn: meikolo wewe • Nt: mwanba.

HERBARIUM DATA (FG). — 91 collections at CAY. Sel. exs.: *J.-F. Molino 1197*.

INVENTORY DATA (FG). — 2242 trees in 146 plots; $F_{\max} = 15.5\%$; $\text{dbh}_{\text{inv}} = 59.8 \text{ cm}$.

[73] *Oxandra xylopioides* Diels

Notizbl. Bot. Gart. Berlin-Dahlem 10: 172 [20 Dec. 1927] (Diels 1927).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5321*.

INVENTORY DATA (FG). — 272 trees in 16 plots; $F_{\max} = 9.4\%$; $\text{dbh}_{\text{inv}} = 57.8 \text{ cm}$.

Genus *Pseudoxandra* R.E.Fr.

[74] *Pseudoxandra cuspidata* Maas

Proc. Kon. Ned. Akad. Wetensch. C 89 (3): 267 (Maas 1986).

VERNACULAR NAMES. — Pa: kuukumwi-seine • Te: balupete • Wp: malupete • Wn: milimi • Nt: pendyku • Cr: lanmoussé • Fr: lamoussé • Br: lamuci.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *H.S. Irwin et al. 47819* (holo-, U[U0000376]; iso-, F[V0047890F], K[K000485521], MG[MG025661], MICH[MICH1192941], NY[NY00026238], U[U0000377], US[00810790]).

INVENTORY DATA (FG). — 45 trees in 17 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 22.6 \text{ cm}$.

[75] *Pseudoxandra* sp. A

NOTE. — This species looks like *P. lucida* R.E.Fr.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3161* (CAY[CAY029984]).

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 24.6 \text{ cm}$.

Genus *Tetrameranthus* R.E.Fr.

[76] *Tetrameranthus guianensis* Westra & Maas

PhytoKeys 12: 13 (Westra & Maas 2012).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: a teki uman udu.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & S. Gonzalez 5387* (holo-, CAY[CAY111703, CAY111704]; iso-, CAY[CAY109443]).

INVENTORY DATA (FG). — 8 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.5 \text{ cm}$.

Genus *Trigynaea* Schltdl.

[77] *Trigynaea caudata* (R.E.Fr.) R.E.Fr.
(Fig. 8C)

Ark. Bot. 33A (9): 14 (Fries 1947), in clavi. — *Hornschuchia caudata* R.E.Fr., *Lloydia* 2 (3): 180 (Fries 1939).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 2 collections at CAY: *J.-F. Molino & D. Sabatier 2133*, *J.-F. Molino & D. Sabatier 2158* (both collected from the same tree).

INVENTORY DATA (FG). — 1 tree, dbh = 14 cm.

Genus *Unonopsis* R.E.Fr.

[78] *Unonopsis glaukopetala* R.E.Fr.

Bull. Herb. Boissier, sér. 2, 7: 1004 (Fries 1907).

Unonopsis andersonii Sprague ex Hohenkerk, *J. Board Agric. British Guiana* 11: 99 (Hohenkerk 1919).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kigksau, kirksau • Ka: tapiseipyo • Nt: azo udu.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3308*.

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22$ cm.

[79] *Unonopsis guatterioides* (A.DC.) R.E.Fr.

Acta Horti Berg. 12 (2): 241 (Fries 1937). — *Uvaria guatterioides* A.DC., *Mém. Soc. Phys. Genève* 5: 202 (Candolle 1832).

Annona peduncularis Steud., *Flora* 26 (45): 754 [17 Dec. 1843] (Steudel 1843). — *Guatteria peduncularis* (Steud.) Pulle, *Recueil Trav. Bot. Néerl.* 4: 124 (Pulle 1907).

Trigynaea angustifolia Benth., *J. Proc. Linn. Soc., Bot.* 5: 70 ["1861" publ. Nov. 1860] (Bentham 1860). — *Unonopsis angustifolia* (Benth.) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 26 (Fries 1900).

Trigynaea grandis Benth., *J. Proc. Linn. Soc., Bot.* 5: 70 ["1861" publ. Nov. 1860] (Bentham 1860). — *Unonopsis grandis* (Benth.) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 28 (Fries 1900).

Trigynaea matthewsii Benth., *J. Proc. Linn. Soc., Bot.* 5: 69 ["1861" publ. Nov. 1860] (Bentham 1860). — *Unonopsis matthewsii* (Benth.) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 28 (Fries 1900), "Matthews".

Trigynaea boliviensis Britton, *Bull. Torrey Bot. Club* 16 (1): 14 [12 Jan. 1889] (Britton 1889). — *Unonopsis boliviensis* (Britton) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 28 (Fries 1900).

Trigynaea antillana Rolfe, *Bull. Misc. Inform. Kew* 1893 (81): 235 [Sep. 1893] (Rolfe 1893). — *Unonopsis antillana* (Rolfe) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 28 (Fries 1900).

Unonopsis lindmanii R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 27 (Fries 1900), "Lindmani".

Unonopsis guaraya Herzog, *Repert. Spec. Nov. Regni Veg.* 7: 52 (Herzog 1909).

Unonopsis buchtienii R.E.Fr., *Repert. Spec. Nov. Regni Veg.* 24: 247 (Fries 1928).

Unonopsis gracilis R.E.Fr., *Acta Horti Berg.* 12 (2): 247 (Fries 1937).

Unonopsis guatterioides f. *elongata* R.E.Fr., *Acta Horti Berg.* 12 (2): 242 (Fries 1937).

Unonopsis obovata R.E.Fr., *Acta Horti Berg.* 12 (2): 249 (Fries 1937).

Unonopsis williamsii R.E.Fr., *Acta Horti Berg.* 12 (2): 238 (Fries 1937).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: miret-unikweune • Ka: akale pom'i'dyi, mulewalan • Wp: pila siay, pina'i, pina'i tay • Wn: éwok pokan • Cr: baton-zin, manmanyawé • Br: envira-surucucu.

HERBARIUM DATA (FG). — 77 collections at CAY. Sel. exs.: *M.E. Moricand s.n.* (original material G[G00237443], G[G00237444], S[fragm. S07-17262]).

SIZE. — Up to 35 cm dbh (Maas *et al.* 2007).

[80] *Unonopsis perrottetii* (A.DC.) R.E.Fr.
(Fig. 8D)

Kongl. Svenska Vetensk.-Akad. Handl., n.s., 34 (5): 26 (Fries 1900). — *Annona perrottetii* A.DC., *Mém. Soc. Phys. Genève* 5: 197 (Candolle 1832), "Anona". — *Trigynaea perrottetii* (A.D.C.) Baill., *Adansonia [Baillon]* 8: 179 (Baillon 1868), "Trigynaea Perrottetii".

Trigynaea perrottetii var. *lanceolata* Baill., *Adansonia [Baillon]* 8: 179 (Baillon 1868), "Trigynaea Perrottetii". — *Unonopsis perrottetii* var. *lanceolata* (Baill.) R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 27 (Fries 1900).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kuukumwi-seine • Te: wiwit hun.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *G.S. Perrottet 65* (lecto-, G[G00226157]), designated by Maas *et al.* [2007: 513]; isolecto-, B[B 10 0243101], F[F0077257F], P[P00751169].

INVENTORY DATA (FG). — 52 trees in 26 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 22.9$ cm.

[81] *Unonopsis rufescens* (Baill.) R.E.Fr.

Kongl. Svenska Vetensk.-Akad. Handl., n.s., 34 (5): 27 (Fries 1900). — *Trigynaea rufescens* Baill., *Adansonia [Baillon]* 8: 180 (Baillon 1868), "Trigynaea".

VERNACULAR NAMES. — Pa: kuukumwi-priye • Ka: alaidya, amose-lan • Te: wiwit hun • Wn: kaloj éwa • Nt: azo udu, mamanyawe.

HERBARIUM DATA (FG). — 118 collections at CAY. Sel. exs.: *E.M. Mélinon 19*, 1864 (holo-, P[P00682175]); iso-, P[P00682176], P[P00682177]).

INVENTORY DATA (FG). — 485 trees in 140 plots; $F_{\max} = 2.8\%$; $\text{dbh}_{\text{inv}} = 25.5$ cm.

[82] *Unonopsis stipitata* Diels

Verh. Bot. Vereins Prov. Brandenburg 47: 130 [1 Oct. 1905] (Diels 1905).

Unonopsis oblanceolata R.E.Fr., *Ark. Bot., n.s.*, 3: 604 (Fries 1956).

Unonopsis rigida R.E.Fr., *Mem. New York Bot. Gard.* 9 (1): 329 (Fries 1957).

VERNACULAR NAMES. — Pa: kuukumwi-priye, pakih-em-priye • Wp: mitū luway, pila siay, pila sila, pina'i to wu • Wn: pakila apisan, pakila hapisan • Nt: azo udu, mamanyawe • Cr: fimel-manmanyawé • Br: envira-preta.

HERBARIUM DATA (FG). — 185 collections at CAY. Sel. exs.: *M.-F. Prévost 1110*.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 12.1 cm.

Genus *Xylopia* L.

[83] *Xylopia aromaticata* (Lam.) Mart.

Fl. Bras. [Martius] 13 (1): 43 [1 Jan. 1841] (Martius 1841). — *Uvaria aromatica* Lam., *Encycl. [J. Lamarck et al.]* 1 (2): 596 [1 Aug. 1785] (Lamarck 1785). — *Unona aromatica* (Lam.) Dunal, *Monogr. Anonac.* 112 [Aug.-Nov. 1817] (Dunal 1817). — *Habzelia aromatica* (Lam.) A.DC., *Mém. Soc. Phys. Genève* 5: 208 (Candolle 1832).

Uvaria febrifuga Humb. & Bonpl. ex Dunal, *Syst. Nat. [A. P. de Candolle]* 1: 498 ["1818" publ. 1-15 Nov. 1817] (Dunal 1817), *nom. nud. pro syn.*

Unona lucida DC. in Dunal, *Monogr. Anonac.* 116 [Aug.-Nov. 1817] (Candolle 1817). — *Coclocline lucida* (DC.) A.DC., *Mém. Soc. Phys. Genève* 5: 209 (Candolle 1832). — *Xylopia lucida* (DC.) Baill., *Hist. Pl. [Baillon]* 1: 278 [Aug.-Dec. 1868] (Baillon 1868).

Unona xylopioides Humb. & Bonpl. ex Dunal, *Monogr. Anonac.* 117 [Aug.-Nov. 1817] (Dunal 1817). — *Xylopia xylopioides* (Humb. & Bonpl. ex Dunal) Standl., *J. Wash. Acad. Sci.* 15: 285 (Standley 1925).

Xylopia grandiflora A.St.-Hil., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (1): 40 [23 Feb. 1824] (Saint-Hilaire 1824).

Xylopia longifolia A.DC., *Mém. Soc. Phys. Genève* 5: 210 (Candolle 1832).

Habzelia cubensis DC. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 717 (Steudel 1840), *nom. nud.*

Xylopia cubensis A.Rich., *Hist. Phys. Cuba, Pl. Vasc.*: 36 (Richard 1845).

Xylopia dunaliana Planch. & Linden ex Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 17: 37 (Triana & Planchon 1862), *nom. nud. pro syn.*

Xylociprum aromaticum (Lam.) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

Xylociprum grandiflorum (A.St.-Hil.) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

Xylociprum longifolium (A.DC.) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Dunal (1817: 116) explicitly ascribed the name and description of *Unona lucida* to "DC. Syst. univ. ined." See above note for *Annona exsucra*.

VERNACULAR NAMES. — Ka: amose, omose • Wp: yáwi'i sī • Wn: siwiman • Nt: man pendyeku, man pengeku, pendyeku, pengeku • Br: envira-branca.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *P. Grenand 574*.

SIZE. — Up to 15 m tall (Maas *et al.* 2001).

[84] *Xylopia benthamii* R.E.Fr.

Kongl. Svenska Vetensk.-Akad. Handl., n.s., 34 (5): 35 (Fries 1900), "Benthami".

Xylopia ulei Diels, *Verh. Bot. Vereins Prov. Brandenburg* 47: 133 [1 Oct. 1905] (Diels 1927).

Xylopia benthamii var. *subnuda* R.E.Fr., *Acta Horti Berg.* 10 (2): 333 (Fries 1931), "Benthami".

Xylopia benthamii var. *dolichopetala* R.E.Fr., *Ark. Bot.* 33A(9): 8 (Fries 1947).

VERNACULAR NAMES. — Br: envira-amarela.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P.J.M. Maas et al. 9635*.

SIZE. — Up to 25 cm dbh (Maas *et al.* 2007).

[85] *Xylopia cayennensis* Maas

Bot. Jahrb. Syst. 115 (1): 93 (Maas 1993).

Xylopia nitida var. *longifolia* Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 11: 137 (Sagot 1881). — *Xylopia longifolia* (Sagot) R.E.Fr., *Acta Horti Berg.* 10 (1): 111 (Fries 1930), *nom. illeg. hom., non A.DC.* (Candolle 1832).

VERNACULAR NAMES. — Pa: pukuu • Ka: amose, omose • Wp: yáwi'i, yáwi'i pilá • Wn: pisat • Nt: pendyeku, pengeku • Cr: lanmoussé-nwé • Br: envira-preta, pindaiba-de-folha-pequena.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *P.A. Sagot 940*, 1856 (holo-, P[P00202581]; iso-, B[B 10 0249563, photo F neg. 14256], BM[BM000554085], GOET[GOET000135], MPU[MPU026895], P[P00202582, P00202583, P00202584], U[U0000439, U0000440]).

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; dbh_{inv} = 27.1 cm.

[86] *Xylopia crinita* R.E.Fr.

Ark. Bot., n.s., 1 (11): 447 (Fries 1950).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4359*.

INVENTORY DATA (FG). — 22 trees in 17 plots; $F_{\max} < 1\%$; dbh_{inv} = 18.5 cm.

[87] *Xylopia discreta* (L.f.) Sprague & Hutch.

Bull. Misc. Inform. Kew 1916 (6): 160 [24 July 1916] (Sprague & Hutchinson 1916). — *Unona discreta* L.f., *Suppl. Pl. 270* ["1781" publ. Apr. 1782] (Linnaeus 1782). — *Habzelia discreta* (L.f.) A.DC., *Mém. Soc. Phys. Genève* 5: 208 (Candolle 1832).

Xylopia salicifolia Humb. & Bonpl. ex Dunal, *Monogr. Anonac.* 121 [Aug.-Nov. 1817] (Dunal 1817).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Schnell 12236* (P).

SIZE. — Guyana. *A. Chanderbali 90* (MO), dbh 15 cm.

[88] *Xylopia excellens* R.E.Fr.

Acta Horti Berg. 12 (1): 210 (Fries 1934).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5402*.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24.8$ cm.

[89] *Xylopia frutescens* Aubl.

Hist. Pl. Guiane 1: 602 [Jun.-Dec. 1775] (Aublet 1775).

Xylopia setosa Poir., *Encycl. [J. Lamarck et al.]* 8: 812 (Poiret 1808).

Xylopicrum frutescens (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

Xylopia meridensis Pittier, *Arb. Arbust. Venez.* 6-8: ? [Aug.-Sep. 1927] (Pittier 1927).

VERNACULAR NAMES. — Pa: pukuu • Ka: awiyu • Wp: yáwi'i, yáwi'i sili • Wn: pisat • Nt: pendyeku, pengeku, pikin pendyeku • Cr: lamoussé • Fr: lamoussé • Br: envira-preta, pindaiba-de-folha-pequena.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material P[P00202541]).

INVENTORY DATA (FG). — 25 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30.9$ cm.

[90] *Xylopia nervosa* (R.E.Fr.) Maas

Bot. Jahrb. Syst. 115 (1): 87 (Maas 1993). — *Oxandra nervosa* R.E.Fr., *Acta Horti Berg.* 12 (3): 558 (Fries 1939).

VERNACULAR NAMES. — Wn: milimi.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M.-F. Prévost 1873*.

SIZE. — Peru, Loreto. *J.J. Pipoly et al. 14832* (MO), 15 m.

[91] *Xylopia nitida* Dunal

Monogr. Anonac. 122 [Aug.-Nov. 1817] (Dunal 1817). — *Xylopicrum nitidum* (Dunal) Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891).

Xylopia cinerea Sandwith, *Bull. Misc. Inform. Kew* 1930 (10): 478 [4 Dec. 1930] (Sandwith 1930).

VERNACULAR NAMES. — Pa: pukuu, pukuu-seine, pukuu-seinó • Ka: awiyu • Te: djawi uhu • Wp: yáwi'i si • Wn: kaloj éwa, kalojimé, milimi • Nt: weti pendyeku, weti pengeku • Cr: lanmoussé-blanc • Fr: lamoussé blanc • Br: envira-amarela, envira-branca, envira-vermelha.

HERBARIUM DATA (FG). — 81 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, G-DC[G00201468]; iso-, FI[FI004818, FI004819], P[P00202482, P00202483], R[R000060814, R000060814a]).

INVENTORY DATA (FG). — 269 trees in 121 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 53$ cm.

[92] *Xylopia parviflora* Spruce

J. Proc. Linn. Soc., Bot. 5: 6 ["1861" publ. June 1860] (Spruce 1860). — *Xylopicrum neglectum* Kuntze, *Revis. Gen. Pl.* 1: 8 [5 Nov. 1891] (Kuntze 1891), nom. illeg. superfl. (based on *Xylopia parviflora*). — *Xylopia neglecta* R.E.Fr., *Kongl. Svenska Vetensk.-Akad. Handl.*, n.s., 34 (5): 31 (Fries 1900), nom. illeg. superfl. (based on *Xylopia parviflora*).

NOTE. — *Xylopia neglecta* is based on the illegitimate *Xylopicrum neglectum*, thus the authorship is not "(Kuntze) R.E.Fr."

VERNACULAR NAMES. — Wp: yáwi'i pilá • Br: envira-sarasará, pacovi.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P.J.M. Maas et al. 9351*.

INVENTORY DATA (FG). — 3 trees censused, none in plots; $dbh_{\text{inv}} = 45$ cm.

[93] *Xylopia pulcherrima* Sandwith

Bull. Misc. Inform. Kew 1930 (10): 477 [4 Dec. 1930] (Sandwith 1930).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3937*, dbh 27.5 cm.

INVENTORY DATA (FG). — 18 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24.7$ cm.

[94] *Xylopia surinamensis* R.E.Fr.

Acta Bot. Neerl. 1: 243 (Fries 1952).

VERNACULAR NAMES. — Wp: yáwi'i.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4087*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24.9$ cm.

Family APOCYNACEAE Juss.
Genus *Ambelania* Aubl.

[95] *Ambelania acida* Aubl.

Hist. Pl. Guiane 1: 265 [Jun.-Dec. 1775] (Aublet 1775). — *Willughbeia acida* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 434 [late Sep.-Nov. 1791] (Gmelin 1791), “*Willughbeja*”.

Ambelania sagotii Müll.Arg., *Linnæa* 30: 389 (Müller 1860).

Ambelania tenuiflora Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 16 [30 July 1860] (Müller 1860).

Ambelania tenuiflora var. *tenuiramea* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 17 [30 July 1860] (Müller 1860).

Ambelania cucumerina Miers, *Apocyn. S. Am.*: 13 [May-June 1878] (Miers 1878).

NOTES. — Miers (1878: 13) referred to Spruce’s description (Spruce 1853: 185) of a “forest cucumber” (“Pepino do matto”), but the name *A. cucumerina* is his own creation, not that of Spruce; hence not *A. cucumerina* Spruce ex Miers’.

VERNACULAR NAMES. — Pa: inuva, mamatki-arin • Ka: amapapali, ambalali • Te: dzádupalali • Wp: akusi ákange, akusi walapulu, akusi walapulu kanapua, akusi walapulu poko, akusi walapulu tā'i • Wn: witik • Nt: batibati, mapa, pikin mapa • Cr: papay-bich, tchenbyendan • Fr: ambelanier acide, papaye biche • Br: pau-de-leite, pepino-doce, pepino-do-mato.

HERBARIUM DATA (FG). — 121 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00778013] designated by Lanjouw & Uittien [1940: 148]).

INVENTORY DATA (FG). — 276 trees in 122 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 29.1$ cm.

Genus *Aspidosperma* Mart. & Zucc.

[96] *Aspidosperma album* (Vahl) Benoist ex Pichon

Bull. Mus. Natl. Hist. Nat., sér. 2, 19: 367 (Pichon 1947). — *Macaglia alba* Vahl, *Skr. Naturhist.-Selsk.* 6: 107 (Vahl 1810).

Bignonia latisiliqua Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 632 [2 May 1811] (Poirer 1811). — *Aspidosperma latisiliquum* (Poir.) A.DC., *Prodr. [A. P. de Candolle]* 8: 676 [mid Mar. 1844] (Candolle 1844), “*latisiliqua*”. — *Peltospermum latisiliquum* DC. ex A.DC., *Prodr. [A. P. de Candolle]* 8: 676 [mid Mar. 1844] (Candolle 1844), *nom. nud. pro syn.*

Peltospermum patrisii DC. ex A.DC., *Prodr. [A. P. de Candolle]* 8: 676 [mid Mar. 1844] (Candolle 1844), *nom. nud. pro syn.*

Aspidosperma centrale Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 12: 560 [6 Dec. 1935] (Markgraf 1935).

VERNACULAR NAMES. — Ka: apukuitya, wokili apukuitya • Te: alapalapa • Wp: alala kái • Wn: amakui, lapalapa • Nt: kumanti udu • Cr: bwa-makak, bwa-takari • Fr: flambeau rouge • Br: araracanga, araraúba, piquiá-marfim.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holo-, C[C10005729]; iso-, P[P00645147, P00645148, P00645149]).

INVENTORY DATA (FG). — 48 trees in 36 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 73.1$ cm.

[97] *Aspidosperma carapanaba* Pichon

Bull. Mus. Natl. Hist. Nat., sér. 2, 19: 365 (Pichon 1947).

Geissospermum excelsum Kuhl., *Arch. Inst. Biol. Veg.* 2 (1): 89 [Sep. 1935] (Kuhlmann 1935).

VERNACULAR NAMES. — Nt: baaka pali udu • Fr: bois pagae • Br: carapanáuba-preta.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 1856.

INVENTORY DATA (FG). — 39 trees in 18 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 140$ cm.

[98] *Aspidosperma cruentum* Woodson

Amer. J. Bot. 22 (7): 684 (Woodson 1935), “*cruenta*”.

Aspidosperma sanguineum Bartlett, *Publ. Carnegie Inst. Wash.* 461: 16 [10 July 1935] (Bartlett 1935), *nom. nud.*

Aspidosperma matudae Lundell, *Phytologia* 1 (10): 339 [27 Nov. 1939] (Lundell 1939), “*matudai*”.

Aspidosperma chiapense Matuda, *Madroño* 10 (6): 172 [23 Jan. 1950] (Matuda 1950).

Aspidosperma chiapense f. *tenax* Matuda, *Madroño* 10 (6): 173 [23 Jan. 1950] (Matuda 1950).

NOTE. — Listed as a synonym of *A. desmanthum* Benth. ex Müll.Arg. (1860) in Morales (2009), but reinstated in Morales & Zamora (2017).

VERNACULAR NAMES. — Ka: apukuitya, mantiotio • Wp: alala kái • Wn: lapalapa • Nt: kumanti udu • Cr: bwa-makak, bwa-takari • Br: araracanga, araraúba.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *D. Sabatier* 981.

INVENTORY DATA (FG). — 84 trees in 46 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 56.7$ cm.

[99] *Aspidosperma excelsum* Benth.

J. Bot. [Hooker] 3: 245 (Bentham 1841). — *Macaglia excelsa* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 416 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: isuu-ára-kamwi, isuu-gaáha, pwait • Ka: apukuitya tamunen, tipulu apukuitya • Te: palakú'i • Wn: aipawaj, epalai • Nt: pali udu • Cr: sitronél-blán • Fr: bois pagae • Br: carapanáuba.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *S.A. Mori & T.D. Pennington* 18001.

INVENTORY DATA (FG). — 84 trees in 56 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 83$ cm.

[100] *Aspidosperma helstonei* Donsel.

Acta Bot. Neerl. 21: 253 (Donselaar 1972).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wakaba-kaubimna • Wp: pilake'i • Wn: aipawaj • Cr: bwa-angi.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 947*.

INVENTORY DATA (FG). — 11 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 44.4$ cm.

[101] *Aspidosperma marcgravianum* Woodson

Ann. Missouri Bot. Gard. 38 (2): 170 (Woodson 1951).

NOTE. — Synonym of *A. excelsum* in Morales (2005), but no longer in Morales & Zamora (2017).

VERNACULAR NAMES. — Ka: apukuitya, tamunen apukuitya • Wp: palaku'i, palakuta piyū • Wn: pakolo talalan • Nt: pali udu • Cr: sitronél-blán • Br: araróba, carapanaúba.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *H. Jacquemin 1903*.

INVENTORY DATA (FG). — 120 trees in 56 plots; $F_{\max} = 2.8\%$; $\text{dbh}_{\text{inv}} = 130$ cm.

[102] *Aspidosperma oblongum* A.DC.

Prodr. [A. P. de Candolle] 8: 399 [mid Mar. 1844] (Candolle 1844). — *Macaglia oblonga* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 416 [5 Nov. 1891] (Kuntze 1891).

Aspidosperma kuhlmannii Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 12: 555 [6 Dec. 1935] (Markgraf 1935).

VERNACULAR NAMES. — Pa: isuu-ára-kamwi • Ka: palakusinien, palakusinyan, tipulu apukuitya • Wp: palakuta pilá, palakuta piyū • Nt: pali udu • Cr: sitronél • Br: carapanaúba-branca.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, G-DC[G00143998]; iso-, B[photo F neg. 4424], F[V0092471F], RB[RB00535068]).

INVENTORY DATA (FG). — 30 trees in 15 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 115$ cm.

[103] *Aspidosperma sandwithianum* Markgr.

Notizbl. Bot. Gart. Berlin-Dahlem 12: 561 [6 Dec. 1935] (Markgraf 1935).

VERNACULAR NAMES. — Wp: alala kái • Br: araracanga-branca.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *Service Forestier M-63*.

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 68.4$ cm.

[104] *Aspidosperma schultesii* Woodson

Ann. Missouri Bot. Gard. 38 (2): 168 (Woodson 1951).

Aspidosperma macrophyllum Müll.Arg. subsp. *mori* L.Allorge, *Bull. Soc. Bot. France, Lett. Bot.* 138 (4-5): 274 ["1991" publ. 1992] (Allorge 1992).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *S.A. Mori et al. 19167* (holotype of *Aspidosperma macrophyllum* subsp. *mori*: P[P00077205]; iso-, K[K000587703], MG[MG143431], P[P00639549], U[U0000478], US[00386110], VEN[VEN269912], WAG[WAG0000107]).

INVENTORY DATA (FG). — 26 trees in 11 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 75.8$ cm.

[105] *Aspidosperma* sp. A

NOTE. — Six collections at CAY were identified in 2003 by J.F. Morales as *A. spruceanum* Benth. ex Müll.Arg. However, the long petioles and abaxially glabrous, shiny leaves of the specimens do not agree with the original diagnosis and type of *A. spruceanum*, which has very short petioles and leaves bearing a silvery indument abaxially. None of the collections identified as *A. spruceanum* at CAY, whether by J.F. Morales or anyone else, displays this combination of characters.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3016*.

INVENTORY DATA (FG). — 199 trees in 47 plots; $F_{\max} = 4.1\%$; $\text{dbh}_{\text{inv}} = 66.9$ cm.

Genus *Couma* Aubl.

[106] *Couma guianensis* Aubl.
(Fig. 9A)

Hist. Pl. Guiane 2 (Suppl.): 39 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate.

Cerbera triphylla Rudge, *Pl. Guian. [Rudge]* 1 (4): 31 [Apr.-May 1806] (Rudge 1806).

VERNACULAR NAMES. — Pa: ukum • Ka: akuma, kuma • Te: kumá • Wp: yuwa • Wn: éhepuk • Nt: baaka mapá, miliki udu • Cr: bwa-vach, mapá, pwé • Fr: bois vache, coumier • Br: sorva, sorvão.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000952644]).

INVENTORY DATA (FG). — 177 trees in 90 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 84.7$ cm.

Genus *Geissospermum* Allemão

[107] *Geissospermum argenteum* Woodson
(Fig. 9C)

Lloydia 2 (3): 207 (Woodson 1939).




FIG. 9. — Apocynaceae: **A**, *Couma guianensis* Aubl. (*D. Sabatier & M.-F. Prévost* 3756); **B**, *Himatanthus phagedaenicus* (Mart.) Woodson (*M.-F. Prévost & D. Sabatier* 4743); **C**, *Geissospermum argenteum* Woodson; **D**, *Rauvolfia paraensis* Ducke (*D. Sabatier* 1691). © D. Sabatier/IRD.

VERNACULAR NAMES. — Pa: á-gógo, gógo purubumna, kongo, kongo-ama • Ka: mataki • Wp: pelowi • Wn: wataki • Nt: bita udu, bonga bita • Cr: maya-kongo • Fr: maria congo • Br: acariquara-branca, acarirana, pau-pereira, quinarana.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3064.

INVENTORY DATA (FG). — 259 trees in 27 plots; $F_{\max} = 7.4\%$; $dbh_{\text{inv}} = 102\text{ cm}$.

[108] *Geissospermum laeve* (Vell.) Miers

Apocyn. S. Am.: 84 [May-June 1878] (Miers 1878). — *Tabernaemontana laevis* Vell., *Fl. Flumin.* 105 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Geissospermum vellosii Allemão, *Arch. Med. Bras.* 2 (4): 73 [Dec. 1845] (Allemão 1845).

VERNACULAR NAMES. — Pa: gógo-puvemna • Ka: matakí • Wp: pelowi'u • Wn: wataki • Nt: bita udu, bonga bita • Cr: maya-kongo • Fr: maria congo • Br: acariquara-branca, acarirana, pau-pereira, quinarana.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2680*.

INVENTORY DATA (FG). — 109 trees in 50 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 120 \text{ cm}$.

[109] *Geissospermum sericeum* Miers

Apocyn. S. Am.: 86 [May-June 1878] (Miers 1878).

Geissospermum fuscum Markgr., *Acta Bot. Venez.* 13 (1-4): 353 (Markgraf 1978).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: wipitano ityumban wewe • Wn: wataki • Nt: bita udu • Br: acariquara-branca, quinarana.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P.A. Sagot* 966, Sep. 1856 (holo-, BM[BM000548558]; iso-, B[not seen, photo F neg. 4455], K[K000582549, K000582550], MPU[MPU022918], P[P00646503, P00646504, P00646505], S[S-R-2368]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 29.4 \text{ cm}$.

Genus *Himatanthus* Willd.

[110] *Himatanthus articulatus* (Vahl) Woodson

Ann. Missouri Bot. Gard. 25 (1): 196 [27 Dec. 1937, dated 1938] (Woodson 1937), “*articulata*”. — *Plumeria articulata* Vahl, *Elog. Amer.* 2: 20 (Vahl 1798).

Himatanthus rigidus Willd., *Syst. Veg. [Roemer & Schultes]* 5: 221 [Dec. 1819] (Willdenow 1819).

Plumeria floribunda Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 40 [30 July 1860] (Müller 1860).

Plumeria sucuuba Spruce ex Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 40 [30 July 1860] (Müller 1860). — *Himatanthus sucuuba* (Spruce ex Müll.Arg.) Woodson, *Ann. Missouri Bot. Gard.* 25 (1): 198 [27 Dec. 1937, dated 1938] (Woodson 1937).

Plumeria microcalyx Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 254 [24 Oct. 1929] (Standley 1929).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020, as *H. sucuuba*). The name *H. rigidus* is to be ascribed to Willdenow alone (Turland *et al.* 2018: Art. 46.3, Ex. 15; see also IPNI 2020 and Taylor & Gereau 2019); hence the authorship is neither “Willd. ex Roem. & Schult.” nor “Willd. ex Schult.”

VERNACULAR NAMES. — Pa: uwaitya-avain • Ka: ana'i, ana'u • Wp: melekene, melekene u • Wn: epkui, lapalapa • Nt: geebi udu, gyebi udu • Cr: bwa-chini, bwa-dilèt • Br: molongó, sucuúbarana.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *M.-F. Prévost 1082*.

INVENTORY DATA (FG). — 31 trees in 23 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 60.9 \text{ cm}$.

[111] *Himatanthus bracteatus* (A.DC.) Woodson

Ann. Missouri Bot. Gard. 25 (1): 200 [27 Dec. 1937, dated 1938] (Woodson 1937), “*bracteata*”. — *Plumeria bracteata* A.DC., *Prodri. [A. P. de Candolle]* 8: 394 [mid Mar. 1844] (Candolle 1844).

Plumeria speciosa Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 36 [30 July 1860] (Müller 1860). — *Himatanthus speciosus* (Müll.Arg.) Plumel, *Compt. Rend. Séances Soc. Biogéogr.* 66 (3): 114 (Plumel 1990).

Plumeria ambigua Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 37 [30 July 1860] (Müller 1860).

Plumeria martii Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 37 [30 July 1860] (Müller 1860).

Plumeria floribunda var. *acutifolia* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 41 [30 July 1860] (Müller 1860).

Plumeria floribunda var. *calycina* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 41 [30 July 1860] (Müller 1860).

Plumeria floribunda var. *crassipes* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 40 [30 July 1860] (Müller 1860).

Plumeria lancifolia Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 41 [30 July 1860] (Müller 1860). — *Himatanthus lancifolius* (Müll.Arg.) Woodson, *Ann. Missouri Bot. Gard.* 25 (1): 200 [27 Dec. 1937, dated 1938] (Woodson 1937), “*lancifolia*”.

Plumeria lancifolia var. *major* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 41 [30 July 1860] (Müller 1860).

Plumeria lancifolia var. *microphylla* Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 41 [30 July 1860] (Müller 1860).

NOTE. — All CAY specimens identified so far as *H. bracteatus* (A.DC.) Woodson (1937) belong to *H. phagedaenicus* (Spina 2004).

HERBARIUM DATA (FG). — A single collection, the neotype of *Plumeria speciosa*: L. Allorge & J. Rombold 335 (P[P00646529]): 20 m × 15 cm (Spina *et al.* 2013).

[112] *Himatanthus phagedaenicus* (Mart.) Woodson
(Fig. 9B)

Ann. Missouri Bot. Gard. 25 (1): 199 [27 Dec. 1937, dated 1938] (Woodson 1937), “*phagedaenica*”. — *Plumeria phagedaenica* Mart., *Reise Bras. [Spix & Mart.]* 3: 1128 (Martius 1831).

NOTE. — Here belong all CAY specimens identified so far as *H. bracteatus* (Spina 2004).

VERNACULAR NAMES. — Nt: geebi udu, gyebi udu • Br: sucuúbara.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2882*.

INVENTORY DATA (FG). — 93 trees in 41 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 38 \text{ cm}$.

[113] *Himatanthus tarapotensis* (K.Schum. ex Markgr.)
Plumel

Bradea 5 (Supl.): 50 (Plumel 1991). — *Plumeria tarapotensis* K.Schum. ex Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 339 (Markgraf 1932).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 23792*.

SIZE. — Up to 25 m tall (Plumel 1991).

Genus *Lacmellea* H.Karst.

[114] *Lacmellea aculeata* (Ducke) Monach.

Lloydia 7 (4): 292 [“1944” publ. 25 Jan. 1945] (Monachino 1945). — *Zschokkea aculeata* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 240 (Ducke 1922).

VERNACULAR NAMES. — Pa: ten-washiunó • Ka: akumalan, awaladala, awalatala • Te: tapele dju'a • Wp: tapeleyiwa, tapeleyuwa • Wn: siklet • Nt: kapasi amanda, maka mapa, mapa • Cr: grènn-bich • Br: cumáí, pau-de-chicle, pau-de-colher, sorvinha.

HERBARIUM DATA (FG). — 121 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2377*.

INVENTORY DATA (FG). — 371 trees in 158 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 32.8$ cm.

[115] *Lacmellea guyanensis* (Müll.Arg.) Monach.

Lloydia 7 (4): 295 [“1944” publ. 25 Jan. 1945] (Monachino 1945). — *Zschokkea guyanensis* Müll.Arg., *Linnaea* 30: 391 (Müller 1860).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (original material K[K000587558]); *R.A.A. Oldeman B-652*: 16 m × 24 cm.

Genus *Laxoplermeria* Markgr.

[116] *Laxoplermeria baehniana* Monach.

Phytologia 3 (2): 68 [29 Mar. 1949] (Monachino 1949).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori 15537*.

SIZE. — Brazil, Acre. *B.A. Krukoff 5720* (MO), 35 m tall (110 ft).

Genus *Macoubea* Aubl.

[117] *Macoubea guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 18 [Jun.-Dec. 1775] (Aublet 1775), “Guyannensis” on plate. — *Tabernaemontana aubletii* Pulle, *Recueil Trav. Bot. Néerl.* 9: 157 (Pulle 1912), nom. illeg. superfl. (based on *Macoubea guianensis*).

Tabernaemontana macrophylla Lam., *Tabl. Encycl. 2* [5 (2)]: 299 [31 Oct. 1819] (Lamarck 1819). — *Merizadenia amplifolia* Miers, *Apocyn. S. Am.*: 79 [May-June 1878] (Miers 1878), nom. illeg. superfl. (based on *Tabernaemontana macrophylla*).

Tabernaemontana reticulata A.DC., *Prodr. [A. P. de Candolle]* 8: 366 [mid Mar. 1844] (Candolle 1844). — *Rhigospira reticulata* (A.DC.) Miers, *Apocyn. S. Am.*: 69 [May-June 1878] (Miers

1878). — *Macoubea reticulata* (A.DC.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 178 [10 July 1938] (Markgraf 1938). — *Macoubea guianensis* var. *reticulata* (A.DC.) L.Allorge, *Mém. Mus. Natl. Hist. Nat.*, sér. B, *Bot.* 30: 174 (Allorge 1985).

Tabernaemontana paucifolia Spruce ex Müll.Arg., *Fl. Bras. [Marius]* 6 (1): 87 [30 July 1860] (Müller 1860). — *Rhigospira paucifolia* (Spruce ex Müll.Arg.) Miers, *Apocyn. S. Am.*: 69 [May-June 1878] (Miers 1878). — *Macoubea paucifolia* (Spruce ex Müll.Arg.) Markgr. ex L.O.Williams, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 422 (Williams 1936). — *Macoubea sprucei* (Müll.Arg.) Markgr. var. *paucifolia* (Spruce ex Müll.Arg.) Monach., *Lloydia* 8 (4): 299 [“1945” publ. 20 Dec. 1946] (Monachino 1946).

Rhigospira sinuosa Miers, *Apocyn. S. Am.*: 70 [May-June 1878] (Miers 1878). — *Macoubea sinuosa* (Miers) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 178 [10 July 1938] (Markgraf 1938).

Parahancornia tabernaemontana Woodson, *Bull. Torrey Bot. Club* 75 (5): 556 [11 Oct. 1948] (Woodson 1948).

NOTE. — Semi-domesticated by pre-Columbian Amerindians (Levis et al. 2017).

VERNACULAR NAMES. — Pa: ukum-kamwi • Ka: sokosoko • Wp: ka'i ákáni, ka'i láká ni • Wn: wítik • Nt: mapa, weti mapa • Cr: mapa, pwé-blán • Fr: macoubé (*fide* Aublet 1775) • Br: amapá-doce.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000952638]).

INVENTORY DATA (FG). — 114 trees in 75 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 71.6$ cm.

Genus *Malouetia* A.DC.

[118] *Malouetia duckei* Markgr.

Notizbl. Bot. Gart. Berlin-Dahlem 9: 962 [15 Nov. 1926] (Markgraf 1926).

HERBARIUM DATA (FG). — A single collection, *B. Bordenave & Bétian 7946*.

SIZE. — Up to 38 m tall (Endress 1995).

[119] *Malouetia guianensis* (Aubl.) Miers

Apocyn. S. Am.: 87 [May-June 1878] (Miers 1878). — *Cam eraria guianensis* Aubl., *Hist. Pl. Guiane* 1: 262 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: pahe-avan-priyo.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777974] designated by Lanjouw & Uittien [1940: 149]).

INVENTORY DATA (FG). — 28 trees in 9 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 39.2$ cm.

[120] *Malouetia tamaquarina* (Aubl.) A.DC.

Prodr. [A. P. de Candolle] 8: 378 [mid Mar. 1844] (Candolle 1844). — *Cameraria tamaquarina* Aubl., *Hist. Pl. Guiane* 1: 260 [Jun.-Dec. 1775] (Aublet 1775). — *Cameraria lutea* Lam., *Encycl. [J. Lamarck et al.]* 1 (2): 573 [1 Aug. 1785] (Lamarck 1785), nom. illeg. superfl. (based on *Cameraria tamaquarina*).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventory plots in French Guiana.

VERNACULAR NAMES. — Pa: yu-kig • Ka: yalamilulan • Wp: paluke • Wn: kuje, kulisa, kuše • Cr: bwa-kouyè • Br: molongó-de-colher, tamanqueira-de-leite.

HERBARIUM DATA (FG). — 83 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777973] designated by Lanjouw & Uittien [1940: 149]).

SIZE. — Up to 20 m tall (Endress 1995).

Genus *Parahancornia* Ducke

[121] *Parahancornia fasciculata* (Poir.) Benoist

Arch. Bot. 5 (Mém. 1): 274 [27 Mar. 1933] (Benoist 1933). — *Tabernaemontana fasciculata* Poir., *Encycl. [J. Lamarck et al.]* 7: 531 [6 July 1806] (Poiret 1806). — *Thyrsanthus fasciculatus* (Poir.) Miers, *Apocyn. S. Am.*: 100 [May-June 1878] (Miers 1878). — *Couma fasciculata* (Poir.) Benoist, *Arch. Bot.* 5 (Mém. 1): 253 [27 Mar. 1933] (Benoist 1933). — *Macoubea fasciculata* (Poir.) Lemée, *Fl. Guyane Franç.* 3: 298 (Lemée 1954).

Hancornia amapa Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 443 (Huber 1902). — *Parahancornia amapa* (Huber) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 26, 243 (Ducke 1922), “*amapa*”; “*Parahancornea*” on p. 242.

VERNACULAR NAMES. — Pa: amap, pakih-etni • Ka: amaba, amapa • Wp: amapa • Nt: dokali mapa, lebi mapa • Cr: mapa • Fr: amapa • Br: amapá, amapá-amargoso.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (original material P[P00646707]).

INVENTORY DATA (FG). — 70 trees in 52 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 77.2$ cm.

Genus *Rauvolfia* L.

[122] *Rauvolfia paraensis* Ducke
(Fig. 9D)

Arch. Jard. Bot. Rio de Janeiro 4: 167 (Ducke 1925).

VERNACULAR NAMES. — Wp: wila lo, wila to tawa • Br: gogó-de-guariba.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier 1691*.

INVENTORY DATA (FG). — 11 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 36.4$ cm.

[123] *Rauvolfia pentaphylla* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 244 (Ducke 1922).

Couma pentaphylla Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 7: 124 (Huber 1913), nom. nud.

Rauvolfia duckei Markgr., *Repert. Spec. Nov. Regni Veg.* 20: 121 (Markgraf 1924).

VERNACULAR NAMES. — Br: pau-marfim-falso.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2448*.

INVENTORY DATA (FG). — 1 tree, dbh = 23.2 cm.

Genus *Tabernaemontana* Plum. ex L.

[124] *Tabernaemontana attenuata* (Miers) Urb.

Repert. Spec. Nov. Regni Veg. 13: 471 [1 Apr. 1915] (Urban 1915). — *Bonafousia attenuata* Miers, *Apocyn. S. Am.*: 51 [May-June 1878] (Miers 1878). — *Anartia attenuata* (Miers) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 165 [10 July 1938] (Markgraf 1938).

VERNACULAR NAMES. — Pa: pahe-avan-puvemna • Ka: wayamu sasamili.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *A.J.M. Leeuwenberg 11653* (US).

INVENTORY DATA (FG). — 52 trees in 37 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 22.4$ cm.

[125] *Tabernaemontana flavicans* Willd.

Syst. Veg. [Roemer & Schultes] 4: 797 [Mar.-June 1819] (Willdenow 1819). — *Anartia flavicans* (Willd.) Miers, *Apocyn. S. Am.*: 82 [May-June 1878] (Miers 1878).

Tabernaemontana oblongifolia A.DC., *Prodr. [A. P. de Candolle]* 8: 368 [mid Mar. 1844] (Candolle 1844). — *Bonafousia oblongifolia* (A.DC.) Miers, *Apocyn. S. Am.*: 50 [May-June 1878] (Miers 1878). — *Anartia oblongifolia* (A.DC.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 165 [10 July 1938] (Markgraf 1938).

Tabernaemontana olivacea Müll.Arg., *Fl. Bras. [Marius]* 6 (1): 75 [30 July 1860] (Müller 1860). — *Bonafousia olivacea* (Müll.Arg.) Miers, *Apocyn. S. Am.*: 52 [May-June 1878] (Miers 1878). — *Anartia olivacea* (Müll.Arg.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 165 [10 July 1938] (Markgraf 1938).

Anartia glabrata Miers, *Apocyn. S. Am.*: 81 [May-June 1878] (Miers 1878).

Bonafousia latiflora Miers, *Apocyn. S. Am.*: 50 [May-June 1878] (Miers 1878).

Tabernaemontana disparifolia Miers, *Apocyn. S. Am.*: 63 [May-June 1878] (Miers 1878).

NOTE. — The name *T. flavicans* is to be ascribed to Willdenow alone (Turland *et al.* 2018: Art. 46.3, Ex. 15; see also IPNI 2020 and Taylor & Gereau 2019); hence the authorship is neither “Willd. ex Roem. & Schult.” nor “Willd. ex Schult.”

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *C. Moretti 626*.

SIZE. — Up to 20 cm dbh (Leeuwenberg 1994).

[126] *Tabernaemontana lagenaria* Leeuwenb.

Revis. *Tabernaemontana* 2: 315 (Leeuwenberg 1994).

VERNACULAR NAMES. — Ka: sokosoko • Wp: sokosoko • Br: leiteira, pau-de-colher.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-J. de Granville 7000* (holo-, P[P00093825]; iso-, BR[BR000006956783], CAY[CAY088444, CAY088445, CAY088446, CAY088447], G[G00190729], MG[MG126010], MO[MO-1115689], U[U0005301], US[US00385866]).

SIZE. — Up to 30 m tall (Leeuwenberg 1994).

[127] *Tabernaemontana linkii* A.DC.

Prodr. [A. P. de Candolle] 8: 364 [mid Mar. 1844] (Candolle 1844). — *Peschiera linkii* (A.DC.) Miers, *Apocyn. S. Am.*: 47 [May-June 1878] (Miers 1878).

Tabernaemontana multiflora Link ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 4: 431 [Mar.-June 1819] (Roemer & Schultes 1819), nom. illeg. hom., non Sm. (Smith 1817).

Tabernaemontana muricata Willd., *Syst. Veg. [Roemer & Schultes]* 4: 297 [Mar.-June 1819] (Willdenow 1819).

Tabernaemontana benthamiana Müll.Arg., *Fl. Bras. [Martius]* 6 (1): 80 [30 July 1860] (Müller 1860). — *Peschiera multiflora* Spruce ex Miers, *Apocyn. S. Am.*: 45 [May-June 1878] (Miers 1878), nom. illeg. superfl. (based on *Tabernaemontana benthamiana*). — *Peschiera benthamiana* (Müll.Arg.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 171 [10 July 1938] (Markgraf 1938).

Peschiera ochracea Miers, *Apocyn. S. Am.*: 42 [May-June 1878] (Miers 1878).

Tabernaemontana myriantha Britton ex Rusby, *Descr. S. Amer.: Pl.* 84 [20 Dec. 1920] (Rusby 1920). — *Peschiera myriantha* (Britton ex Rusby) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 171 [10 July 1938] (Markgraf 1938). — *Peschiera benthamiana* var. *myriantha* (Britton ex Rusby) L.Allorge, *Mém. Mus. Natl. Hist. Nat., sér. B, Bot.* 30: 154 (Allorge 1985).

Tabernaemontana stenantha Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 1037 [1 Aug. 1930] (Markgraf 1930). — *Peschiera stenantha* (Markgr.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 171 [10 July 1938] (Markgraf 1938). — *Peschiera benthamiana* var. *stenantha* (Markgr.) L.Allorge, *Mém. Mus. Natl. Hist. Nat., sér. B, Bot.* 30: 154 (Allorge 1985).

NOTE. — The name *T. muricata* is to be ascribed to Willdenow alone (Turland *et al.* 2018: Art. 46.3, Ex. 15; see also IPNI 2020 and Taylor & Gereau 2019); hence the authorship is neither “Willd. ex Roem. & Schult.” nor “Willd. ex Schult.”

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville B-4488*.

SIZE. — Up to 40 cm dbh (Leeuwenberg 1994).

[128] *Tabernaemontana sananho* Ruiz & Pav.

Flora Peruviana 2: 22 (Ruiz & Pavón 1799). — *Merizadenia sananho* (Ruiz & Pav.) Miers, *Apocyn. S. Am.*: 78 [May-June 1878] (Miers 1878). — *Bonafousia sananho* (Ruiz & Pav.) Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 166 [10 July 1938] (Markgraf 1938).

Tabernaemontana poeppigii Müll.Arg., *Linnaea* 30: 405 (Müller 1860). — *Taberna poeppigii* (Müll.Arg.) Miers, *Apocyn. S. Am.*: 63 [May-June 1878] (Miers 1878).

VERNACULAR NAMES. — Wp: a'i kálátá'á • Br: guarecoló.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4424*.

SIZE. — Up to 15 cm dbh (Leeuwenberg 1994).

[129] *Tabernaemontana undulata* Vahl

Eclog. Amer. 2: 20 (Vahl 1798). — *Bonafousia undulata* (Vahl) A.DC., *Prodr. [A. P. de Candolle]* 8: 359 [mid Mar. 1844] (Candolle 1844).

Echites brasiliensis Thunb., *Echitis* 5 (Thunberg 1819).

Tabernaemontana meyeri G.Don, *Gen. Hist.* 4 (1): 89 (Don 1837). — *Anartia meyeri* (G.Don) Miers, *Apocyn. S. Am.*: 80 [May-June 1878] (Miers 1878).

Tabernaemontana perrottetii A.DC., *Prodr. [A. P. de Candolle]* 8: 362 [mid Mar. 1844] (Candolle 1844). — *Bonafousia perrottetii* (A.DC.) Miers, *Apocyn. S. Am.*: 51 [May-June 1878] (Miers 1878).

Peschiera surinamensis Miq., *Linnaea* 18: 742 [“1844” publ. Aug.-Oct. 1845] (Miquel 1845).

Bonafousia obliqua Miers, *Apocyn. S. Am.*: 49 [May-June 1878] (Miers 1878). — *Tabernaemontana obliqua* (Miers) Leeuwenb., *Meded. Landbouwhoogeschool Wageningen* 83 (7): 60 [“1983” publ. 1984] (Leeuwenberg 1984).

Bonafousia undulata var. *ovalifolia* Miers, *Apocyn. S. Am.*: 49 [May-June 1878] (Miers 1878).

Tabernaemontana albescens Rusby, *Descr. S. Amer. Pl.* 83 [20 Dec. 1920] (Rusby 1920). — *Anacampa albescens* (Rusby) Markgr., *Fl. Suriname* 4 (1): 452 (Markgraf 1937).

Stemmadenia nervosa Standl. & L.O.Williams, *Ceiba* 3 (2): 126 [1 Oct. 1952] (Standley & Williams 1952).

VERNACULAR NAMES. — Pa: pahe-avan • Ka: alawata emulutano, pelo emulutano • Wp: sokosoko'i • Wn: pakilemu • Nt: miliki tiki • Cr: bwa-dilèt.

HERBARIUM DATA (FG). — 197 collections at CAY. Sel. exs.: *G.S. Perrottet 275* (holotype of *Tabernaemontana perrottetii*: G-DC[G00143420]; iso-, G[G00170000]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 11.9$ cm.

Family AQUIFOLIACEAE Bercht. & J.Presl
Genus *Ilex* Tourn. ex L.

[130] *Ilex guianensis* (Aubl.) Kuntze

Revis. Gen. Pl. 1: 113 [5 Nov. 1891] (Kuntze 1891). — *Macoucoua guianensis* Aubl., *Hist. Pl. Guiane* 1: 88 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Ilex acuminata* Willd., *Sp. Pl.*, ed. 4 1 (2): 711 [July 1798] (Willdenow 1798), nom. illeg. superfl. (based on *Macoucoua guianensis* Aubl.). — *Ilex macoucoua* Pers., *Syn. Pl. [Persoon]* 1: 152 [1 Apr.-15 June 1805] (Persoon 1805), nom. illeg. superfl. (based on *Macoucoua guianensis*).

Ilex celastroides Klotzsch ex Garcke, *Linnaea* 22: 58 (Garcke 1849).

Ilex occidentalis Macfad., *Fl. Brit. W.I. [Grisebach]* 147 [June 1859] (Macfadyen 1859).

Ilex panamensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 221 [24 Oct. 1929] (Standley 1929).

Ilex gentilei Lundell, *Field & Lab.* 13 (1): 5 (Lundell 1945).

VERNACULAR NAMES. — Ka: macoucou (*fide* Aublet 1775).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777519] designated by Lanjouw & Uittien [1940: 152]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.9$ cm.

[131] *Ilex inundata* Poepp. ex Reissek

Fl. Bras. [Martius] 11 (1): 43 [15 Feb. 1861] (Reissek 1861).

Ilex riparia Reissek, *Fl. Bras. [Martius]* 11 (1): 43 [15 Feb. 1861] (Reissek 1861).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2267.

INVENTORY DATA (FG). — 1 tree, $dbh = 19.3$ cm.

[132] *Ilex jenmanii* Loes.

Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur. 78: 412 (Loesener 1901).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Loubry* 2339.

INVENTORY DATA (FG). — 10 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42.7$ cm.

[133] *Ilex laureola* Triana & Planch.

Ann. Sci. Nat., Bot. sér. 5, 16: 377 (Triana & Planchon 1872), “*Laureola*”.

Ilex laureola var. *genuina* Loes., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 78: 440 (Loesener 1901), nom. inval. (Turland et al. 2018: Art. 24.3).

Ilex laureola var. *neglecta* Loes., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 78: 440 (Loesener 1901).

Ilex macrolaurus Loes., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 78: 438 (Loesener 1901).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & E. Fonty* 5597.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 16.3$ cm.

[134] *Ilex martiniana* D.Don

Descr. Pinus, ed. 2, 2: App. 8* (Don 1828).

NOTES. — Known only from the Guiana Shield. Plate 5 in Don (1828) is wrongly labelled “*Ilex paraguensis*”. *I. paraguensis* is described on p. 5*, t. 4.

HERBARIUM DATA (FG). — No collection at CAY. Sel. exs.: *J. Martin s.n.* (original material not traced); *E.M. Mélinon s.n.* (P[P03275721]).

INVENTORY DATA (FG). — 1 tree, $dbh = 16.8$ cm.

[135] *Ilex* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *P. Acevedo-Rodríguez et al.* 6130.

INVENTORY DATA (FG). — 1 tree, $dbh = 56$ cm.

[136] *Ilex* sp. B

(Fig. 10A)

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino* 1155.

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 51.9$ cm.

Family ARALIACEAE Juss.

Genus *Didymopanax* Decne. & Planch.

[137] *Didymopanax decaphyllus*
(Seem.) Fiaschi & G.M.Plunkett

Brittonia 72 (1): 19 [Publ. online 23 Jan. 2020] (Fiaschi & Plunkett 2020). — *Sciodaphyllum decaphyllum* Seem., *J. Bot. 3:* 266 (Seemann 1865). — *Panax decaphyllus* Sagot ex Seem., *J. Bot. 3:* 266 (Seemann 1865), nom. nud. pro syn. — *Schefflera decapilla* (Seem.) Harms, *Nat. Pflanzenfam. [Engler & Prantl]* 3 (8): 37 (Harms 1894).

Schefflera paraensis Huber ex Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 228 (Ducke 1922).

VERNACULAR NAMES. — Pa: ahamban • Ka: piliwa molototo'u, piliwa mulototou • Wp: molototo • Nt: tobitutu, weti tobitutu • Cr: bwa-lasenjan • Fr: bois de la saint-Jean • Br: morototó-branco.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *P.A. Sagot* 916, May 1856 (lecto-, K[K000588148], designated by Fiaschi & Plunkett [2018: 57]; isolecto-, BM[BM001008663], P[P00445504, P00445505], P02286445, P02286446, P02286447, P02286448], W[W 1889-0089019]).

INVENTORY DATA (FG). — 133 trees in 73 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 73.2$ cm.



FIG. 10. — Aquifoliaceae: **A**, *Ilex* sp. B (D. Sabatier et al. 4715). Araliaceae: **B**, *Oreopanax capitatus* (Jacq.) Decne. & Planch. (D. Sabatier & J.-F. Molino 5723). Arecaceae: **C**, *Astrocarum rodriquesii* Trail (D. Sabatier & M.-F. Prévost 4921); **D**, *Manicaria saccifera* Gaertn. © D. Sabatier/IRD.

[138] *Didymopanax morototoni* (Aubl.) Decne. & Planch.

Rev. Hort. (Paris), sér. 4, 3: 109 [16 Mar. 1854] (Decaisne & Planchon 1854). — *Panax morototoni* Aubl., *Hist. Pl. Guiane* 2: 949 [Jun.-Dec. 1775] (Aublet 1775). — *Panax undulatus* Aubl., *Hist. Pl. Guiane* 4: t. 360 [Jun.-Dec. 1775] (Aublet 1775), “*Undulata*”. — *Oreopanax morototoni* (Aubl.) Pittier, *Expl. Bot. Cuenca Maracaibo* 41 (Pittier 1923). — *Schefflera morototoni* (Aubl.) Maguire, Steyermark & Frodin, *Mem. New York Bot. Gard.* 38: 51 (Maguire et al. 1984).

Panax splendens Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 11 [28 May 1821] (Kunth 1821). — *Didymopanax splendens* (Kunth) Decne. & Planch. ex Seem., *J. Bot.* 6: 132 (Seemann 1868). — *Schefflera splendens* (Kunth) Frodin ex Lindeman, *Fl. Suriname* 3 (1-2, addit. & correct.): 352 (Lindeman 1986).

Didymopanax poeppigii Decne. & Planch., *Rev. Hort. (Paris)*, sér. 4, 3: 109 [16 Mar. 1854] (Decaisne & Planchon 1854). — *Didymopanax morototoni* var. *poeppigii* (Decne. & Planch.) Marchal, *Fl. Bras. [Martius]* 11 (1): 241 [1 Feb. 1878] (Marchal 1878).

Didymopanax morototoni var. *sessiliflorus* Marchal, *Fl. Bras. [Martius]* 11 (1): 241 [1 Feb. 1878] (Marchal 1878). — *Schefflera morototoni* var. *sessiliflorus* (Marchal) Frodin, World Checkl. & Bibliogr. Araliaceae 358 [2003 published Feb. 2004] (Frodin 2004).

NOTE. — Although *Panax “undulata”* is an error on the plate of *P. morototoni* (Aublet 1775, 4: t. 360), *Panax undulatus* is a valid name.

VERNACULAR NAMES. — Pa: ahamban • Ka: molototo'u, mulototou • Te: bolototo uhu • Wp: molototo • Wn: malamala • Nt: lebi tobitutu • Cr: bwa-lasenjan • Fr: bois de la saint-Jean • Br: morototó.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[[BM001008661](#)]), designated by Fiaschi & Plunkett [2018: 74]).

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; dbh_{inv} = 70 cm.

Genus *Oreopanax* Decne. & Planch.

[139] *Oreopanax capitatus* (Jacq.) Decne. & Planch. (Fig. 10B)

Rev. Hort. (Paris), sér. 4, 3: 108 [16 Mar. 1854] (Decaisne & Planchon 1854), “capitatum”. — *Aralia capitata* Jacq., *Enum. Syst. Pl.*: 18 [Aug.-Sep. 1760] (Jacquin 1760). — *Hedera capitata* (Jacq.) Sm., *Icon. Pict. Pl. Rar.* 1: t. 4 (Smith 1790). — *Hedera frondosa* Salisb., *Prodr. Stirp. Chap. Allerton* 144 [Nov.-Dec. 1796] (Salisbury 1796), nom. illeg. superfl. (based on *Aralia capitata*). — *Botryodendrum capitatum* (Jacq.) Endl., *Cat. Horti Vindob.* 2: 177 (Endlicher 1842). — *Sciodaphyllum capitatum* (Jacq.) Griseb., *Fl. Brit. W.I. [Grisebach]* 306 [late 1860] (Grisebach 1860). — *Mesopanax capitatus* (Jacq.) R.Vig., *Ann. Sci. Nat., Bot. sér. 9, 4:* 104 (Viguier 1906).

Oreopanax destructor Seem., *J. Bot.* 7: 351 (Seemann 1869).

Oreopanax liebmannii Marchal, *Bull. Acad. Roy. Sci. Belgique*, sér. 2, 47 (1): 87 (Marchal 1879).

Oreopanax oligocarpus Donn.Sm., *Bot. Gaz.* 16 (7): 194 [20 July 1891] (Donnell Smith 1891), “oligocarpum”.

Aralia ovata Sessé & Moc., *Fl. Mexic.*, ed. 2, 79 (Sessé & Mociño 1894).

Oreopanax meiocephalus Donn.Sm., *Bot. Gaz.* 37 (3): 210 [18 Mar. 1904] (Donnell Smith 1904), “meiocephalum”.

Oreopanax capitatus var. *minor* Steyermark., *Fieldiana, Bot.* 28 (2): 442 (Steyermark 1952).

NOTE. — *Aralia ovata* also appears in the first edition of Flora Mexicana (Sessé & Mociño 1892-1898: 86, as “Ouata”), but edition 2 has priority, probably from p. 49 onward (Stafleu & Cowan 1985: [TL-2-11756](#)).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *O. Poncy 2866*.

SIZE. — Up to 30 m tall (Frodin 1997).

Family ARECACEAE Bercht. & J.Presl Genus *Acrocomia* Mart.

[140] *Acrocomia aculeata* (Jacq.) Lodd. ex Mart.

Hist. Nat. Palm. 3 (8): 286 [19 Sept 1845] (Martius 1845). — *Cocos aculeata* Jacq., *Select. Stirp. Amer. Hist.* 278 [5 Jan. 1763] (Jacquin 1763), “aculeatus”.

Palma spinosa Mill., *Gard. Dict.*, ed. 8, n. 3 [16 Apr. 1768] (Miller 1768), nom. illeg. hom., non Loefl. (Loefling 1758).

Palma mocaia Aubl., *Hist. Pl. Guiane* 2: 976 [Jun.-Dec. 1775] (Aublet 1775).

Bactris globosa Gaertn., *Fruct. Sem. Pl.* 1: 22 [Dec. 1788] (Gaertner 1788). — *Bactris minor* Gaertn., *Fruct. Sem. Pl.* 1: t. 9, f. 1 [Dec. 1788] (Gaertner 1788), nom. illeg. hom., non Jacq. (Jacquin 1780).

Cocos fusiformis Sw., *Fl. Ind. Occid.* 1: 616 [Nov. 1797] (Swartz 1797). — *Acrocomia fusiformis* (Sw.) Sweet, *Hort. Brit. [Sweet]* 432 (Sweet 1826).

Acrocomia sclerocarpa Mart., *Hist. Nat. Palm.* 2 (3): 66 [after 13 Apr. 1824] (Martius 1824), nom. illeg. superfl. (based on *Cocos aculeata*).

Acrocomia sphaerocarpa Mart. ex Desf., *Tabl. École Bot.*, ed. 3 [*Cat. Pl. Horti Paris.*]: 30 (Desfontaines 1829).

Acrocomia guianensis Lodd. ex G.Don, *Hort. Brit. [Loudon]*: 382 (Don 1830), nom. nud.

Acrocomia minor Lodd. ex G.Don, *Hort. Brit. [Loudon]*: 382 (Don 1830).

Acrocomia globosa Lodd. ex Mart., *Hist. Nat. Palm.* 3 (8): 286 [19 Sept 1845] (Martius 1845).

Acrocomia horrida Lodd. ex Mart., *Hist. Nat. Palm.* 3 (8): 286 [19 Sept 1845] (Martius 1845).

Acrocomia mexicana Karw. ex Mart., *Hist. Nat. Palm.* 3 (8): 285 [19 Sept 1845] (Martius 1845).

Acrocomia tenuifrons Lodd. ex Mart., *Hist. Nat. Palm.* 3 (8): 286 [19 Sept 1845] (Martius 1845).

Bactris pavoniana Mart., *Voy. Amér. Mér. 7 (3, Palmiers)*: 70 (Martius 1847).

Acrocomia totai Mart., *Voy. Amér. Mér. 7 (3, Palmiers)*: 78 (Martius 1847).

Acrocomia lasiospatha Mart., *Voy. Amér. Mér. 7 (3, Palmiers)*: 81 (Martius 1847).

Acrocomia cubensis Lodd. ex H.Wendl., *Index Palm.*: 1 (Wendland 1854).

Acrocomia vinifera Oerst., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1858: 47 (Oersted 1859).

Acrocomia antioquiensis Posada-Ar., *Bull. Soc. Bot. France* 25: 184 (Posada-Arango 1878).

Acrocomia zapotecis H.Wendl., *Palmiers [Kerchove]*: 230 (Wendland 1878), nom. nud.

Acrocomia glaucocephylla Drude, *Fl. Bras. [Martius]* 3 (2): 392 [1 Nov. 1881] (Drude 1881).

Acrocomia intumescens Drude, *Fl. Bras. [Martius]* 3 (2): 391 [1 Nov. 1881] (Drude 1881).

Acrocomia sclerocarpa var. *wallaceana* Drude, *Fl. Bras. [Martius]* 3 (2): 391 [1 Nov. 1881] (Drude 1881). — *Acrocomia wallaceana* (Drude) Becc., *Pomona Coll. J. Econ. Bot.* 2: 362 (Beccari 1912), nom. illeg. superfl. (*Cocos aculeata* Jacq. in synonymy).

Acrocomia microcarpa Barb.Rodr., *Vellozia*, ed. 2, 1: 107 (Barbosa Rodrigues 1891).

Acrocomia mokayayba Barb.Rodr., *Pl. Jard. Rio de Janeiro* 5: 11 (Barbosa Rodrigues 1896).

Acrocomia odorata Barb.Rodr., *Palm. Matogross.* 48 (Barbosa Rodrigues 1898).

Acrocomia media O.F.Cook, *Bull. Torrey Bot. Club* 28 (10): 566 (Cook 1901).

Acrocomia erioacantha Barb.Rodr., *Contr. Jard. Bot. Rio de Janeiro* 2: 85 (Barbosa Rodrigues 1902).

Acrocomia ulei Dammer, *Notizbl. Königl. Bot. Gart. Berlin* 6: 266 [30 June 1915] (Dammer 1915).

Acrocomia pilosa Léon, *Mem. Soc. Cub. Hist. Nat. "Felipe Poey"* 14: 52 (Léon 1940).

Acrocomia chunta Covas & Ragonese, *Revista Argent. Agron.* 8: 2 (Covas & Ragonese 1941).

Acrocomia belizensis L.H.Bailey, *Gentes Herb.* 4 (12): 445 [30 Sep. 1941] (Bailey 1941).

Acrocomia hospes L.H.Bailey, *Gentes Herb.* 4 (12): 449 [30 Sep. 1941] (Bailey 1941).

Acrocomia ierensis L.H.Bailey, *Gentes Herb.* 4 (12): 473 [30 Sep. 1941] (Bailey 1941).

Acrocomia karukerana L.H.Bailey, *Gentes Herb.* 4 (12): 466 [30 Sep. 1941] (Bailey 1941).

Acrocomia panamensis L.H.Bailey, *Gentes Herb.* 4 (12): 444 [30 Sep. 1941] (Bailey 1941).

Acrocomia quisqueyana L.H.Bailey, *Gentes Herb.* 4 (12): 471 [30 Sep. 1941] (Bailey 1941).

Acrocomia subinermis Léon ex L.H.Bailey, *Gentes Herb.* 4 (12): 474 [30 Sep. 1941] (Bailey 1941).

Acrocomia antiguana L.H.Bailey, *Gentes Herb.* 8: 142 (Bailey 1949).

Acrocomia christopherensis L.H.Bailey, *Gentes Herb.* 8: 140 (Bailey 1949).

Acrocomia grenadana L.H.Bailey, *Gentes Herb.* 8: 144 (Bailey 1949).

Acrocomia viegasii L.H.Bailey, *Gentes Herb.* 8: 139 (Bailey 1949).

Acrocomia spinosa H.E.Moore, *Gentes Herb.* 9: 238 (Moore 1963).

NOTES. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). In French Guiana, restricted to savannas. Wendland (in Kerchove 1878: 230) mistakenly listed “*Acrocomia Zapotecis* Karw.” as a synonym of *A. mexicana* Karw. ex Mart. Actually, Martius’ sentence: “*A Zapotecis Jaçangha [...] dicitur*” (Martius 1824: 286) only means that “Jacangha” is the vernacular name of *A. mexicana* for the Zapotecs. The attribution of this “binomial” to L. B. de Karwinsky (“*Acrocomia zapotecis* Karw.”) (Wendland in Kerchove 1878: 230) is also a mistake. Karwinsky wrote the name *A. mexicana* on the type specimen that he collected. *Bactris globosa* Gaertn. appears as *B. minor* Gaertn. on plate.

VERNACULAR NAMES. — Ka: mokaya, mukaya • Cr: moukaya • Br: macaúba, mucajá.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville 17664*, dbh 28 cm.

Genus *Astrocaryum* G.Mey.

[141] *Astrocaryum jauari* Mart.

Hist. Nat. Palm. 2 (3): 76 [after 13 Apr. 1824] (Martius 1824).

Astrocaryum guara Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 15 [30 Dec. 1930] (Burret 1930).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Ka: sauwalai • Wn: jawala • Nt: liba awaa • Br: jauari.

HERBARIUM DATA (FG). — No specimen seen. The species is present on the banks of Maroni River, according to Granville & Gayot (2014).

SIZE. — Up to 30 cm dbh (Granville & Gayot 2014).

[142] *Astrocaryum minus* Trail

J. Bot. 15: 78 (Trail 1877).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-J. de Granville 17659*.

SIZE. — Up to 12 cm dbh (Granville & Gayot 2014).

[143] *Astrocaryum murumuru* Mart.

Hist. Nat. Palm. 2 (3): 70 [after 13 Apr. 1824] (Martius 1824).

Astrocaryum yauaperyense Barb.Rodr., *Vellosia* 1: 48 (Barbosa Rodrigues 1888).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana. Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Wp: mulumulu sī • Br: murumuru.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-J. de Granville & F. Kahn 11201*.

SIZE. — Up to 27 cm dbh (Granville & Gayot 2014).

[144] *Astrocaryum rodriquesii* Trail
(Fig. 10C)

J. Bot. 15: 79 (Trail 1877), “Rodriguezii”.

VERNACULAR NAMES. — Pa: amuu • Wp: patali • Wn: widajimë • Br: tucumá-branco.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4921*.

INVENTORY DATA (FG). — 18 trees in 6 plots; $F_{\max} = 1.6\%$; $dbh_{inv} = 20.9$ cm.

[145] *Astrocaryum sciophilum* (Miq.) Pulle

Enum. Vasc. Pl. Surinam 73 (Pulle 1906). — *Bactris sciophila* Miq., *Natuurk. Verh. Wet. Haarlem*, ser. 2, 7: 208 (Miquel 1851).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kwasi-muumuuwa, kwaswe-muumuuwa
• Ka: mulumulu • Te: mubulu • Wp: mulumulu • Wn: mumu,
muumu • Nt: bugu • Cr: mouroumourou • Br: murumuru.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *A.H. Gentry & E.M. Zardini 50232*.

INVENTORY DATA (FG). — 1197 trees in 107 plots; $F_{\max} = 19.8\%$; $dbh_{\text{inv}} = 22.3 \text{ cm}$.

[146] *Astrocaryum vulgare* Mart.

Hist. Nat. Palm. 2 (3): 74 [after 13 Apr. 1824] (Martius 1824).

Astrocaryum awarra de Vriese, *Jaarb. Kon. Ned. Maatsch. Aanm. Tuinb.* 1848: 12 (de Vriese 1848).

Astrocaryum guianense Splitg. ex Mart., *Hist. Nat. Palm.* 3 (10): 323 [Jan.-July 1853] (Martius 1853).

Astrocaryum segregatum Drude, *Fl. Bras. [Martius]* 3 (2): 382 [Nov. 1881] (Drude 1881).

Astrocaryum tucumoides Drude, *Fl. Bras. [Martius]* 3 (2): 381 [Nov. 1881] (Drude 1881).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: wahatwi • Ka: awala, awala'i • Te: dzawala • Wp: awala • Wn: awala • Nt: awaa • Cr: awara, wara • Br: tucumá-do-Pará.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-J. de Granville 8340*.

SIZE. — Up to 20 cm dbh (Granville & Gayot 2014).

Genus *Attalea* Kunth

[147] *Attalea maripa* (Aubl.) Mart.

Voy. Amér. Mér. 7 (3, Palmiers): 123 (Martius 1847). — *Palma maripa* Aubl., *Hist. Pl. Guiane* 2: 974 [Jun.-Dec. 1775] (Aublet 1775). — *Maximiliana maripa* (Aubl.) Drude, *Fl. Bras. [Martius]* 3 (2): 452 [1 Nov. 1881] (Drude 1881). — *Englerophoenix maripa* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 728 [5 Nov. 1891] (Kuntze 1891). — *Ethnora maripa* (Aubl.) O.F.Cook, *J. Wash. Acad. Sci.* 30: 297 (Cook 1940).

Maximiliana regia Mart., *Hist. Nat. Palm.* 2 (4): 132 [Jan.-Mar. 1826] (Martius 1826). — *Englerophoenix regia* (Mart.) Kuntze, *Revis. Gen. Pl.* 2: 728 [5 Nov. 1891] (Kuntze 1891). — *Temene regia* (Mart.) O.F.Cook, *Natl. Hort. Mag.* 18: 276 (Cook 1939). — *Attalea regia* (Mart.) Wess.Boer, *Indig. Palms Surin.*: 150 (Wessels Boer 1965).

Maximiliana elegans H.Karst., *Linnaea* 28: 271 ["1856" publ. Jan. 1857] (Karsten 1857).

Maximiliana martiana H.Karst., *Linnaea* 28: 273 ["1856" publ. Jan. 1857] (Karsten 1857).

Maximiliana caribaea Griseb. & H.Wendl., *Fl. Brit. W.I. [Grisebach]* 522 [Oct. 1864] (Grisebach & Wendland 1864). — *Englerophoenix caribaeum* (Griseb. & H.Wendl.) Kuntze, *Revis. Gen. Pl.* 2: 728 [5 Nov. 1891] (Kuntze 1891).

Scheelea maripa hort. ex H.Wendl., *Palmiers [Kerchove]* 256 (Wendland 1878), *nom. nud.*

Maximiliana tetrasticha Drude, *Fl. Bras. [Martius]* 3 (2): 455 [Nov. 1881] (Drude 1881). — *Scheelea tetrasticha* (Drude) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 667 [10 July 1929] (Burret 1929).

Maximiliana longirostrata Barb.Rodr., *Vellozia*, ed. 2, 1: 112 (Barbosa Rodrigues 1891). — *Englerophoenix longirostrata* (Barb.Rodr.) Barb.Rodr., *Sert. Palm. Brasil.* 1: 77 (Barbosa Rodrigues 1903).

Maximiliana macropetala Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 699 [10 July 1929] (Burret 1929). — *Attalea macropetala* (Burret) Wess.Boer, *Indig. Palms Surin.*: 155 (Wessels Boer 1965).

Maximiliana macrogyne Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 692 [10 July 1929] (Burret 1929).

Maximiliana stenocarpa Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 696 [10 July 1929] (Burret 1929).

Attalea cryptantha Wess.Boer, *Pittieria* 17: 310 (Wessels Boer 1988).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: arawa-kagta, kahikti • Ka: malipa • Te: balipa • Wp: inaya, malipa, malipa tay • Wn: malipa • Nt: maypa • Cr: maripa • Br: anajá, inajá.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: the type from French Guiana (not traced).

INVENTORY DATA (FG). — 49 trees in 30 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 32.8 \text{ cm}$.

Genus *Cocos* L.

[148] *Cocos nucifera* L.

Sp. Pl. 2: 1188 [1 May 1753] (Linnaeus 1753). — *Calappa nucifera* (L.) Kuntze, *Revis. Gen. Pl.* 2: 982 [5 Nov. 1891] (Kuntze 1891).

Palma cocos Mill., *Gard. Dict.*, ed. 8, n. 2 [16 Apr. 1768] (Miller 1768).

Cocos mamillaris Blanco, *Fl. Filip. [FM. Blanco]* 722 (Blanco 1837).

Cocos indica Royle, *Ill. Bot. Himal. Mts.*: 395 (Royle 1840).

Cocos nana Griff., *Not. Pl. Asiat.* 3: 166 (Griffith 1851).

Cocos nucifera var. *synphyllica* Becc., *Agric. Colon.* 10: 586 (Beccari 1916).

Diplothemium henryanum F.Br., *Bull. Bernice P. Bishop Mus.* 84: 128 (Brown 1931).

NOTE. — A species native to the South-West Pacific, introduced and naturalised on all tropical shores.

VERNACULAR NAMES. — Pa: koko • Ka: koko, kokoepu • Te: koko • Wp: koko • Wn: koko • Nt: kokonoto bon, kokonto bon • Fr: cocotier • Br: cóco, coqueiro.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville 7203*, dbh 25-28 cm.

Genus *Euterpe* Mart.[149] *Euterpe oleracea* Mart.*Hist. Nat. Palm. 2* (2): 29 [before 13 Apr. 1824] (Martius 1824).*Euterpe brasiliiana* Oken, *Allg. Naturgesch. 3* (1): 674 (Oken 1841).*Catis martiana* O.F.Cook, *Bull. Torrey Bot. Club 28* (10): 557 (Cook 1901), “*Martiniana*”, nom. illeg. superfl. (based on *Euterpe oleracea*).*Euterpe badiocarpa* Barb.Rodr., *Contr. Jard. Bot. Rio de Janeiro 1*: 12 (Barbosa Rodrigues 1901).*Euterpe beardii* L.H.Bailey, *Gentes Herb. 7*: 426 (Bailey 1947).*Euterpe cuatrecasana* Dugand, *Revista Acad. Colomb. Ci. Exact. 8* (31): 393 [Dec. 1951] (Dugand 1951).NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: was • Ka: wasay, wasei • Te: watsey • Wp: wasay, wasey, wasey iipi manu • Wn: wapu • Nt: pina • Cr: pino, wassay • Br: açaí-do-pará.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 8057*.INVENTORY DATA (FG). — 743 trees in 34 plots; $F_{\max} = 48.7\%$; dbh_{inv} = 20.5 cm.[150] *Euterpe precatoria* Mart.*Voy. Amér. Mér. 7* (3, Palmiers): 10 (Martius 1847).*Euterpe oleracea* Engel, *Linnaea 33*: 671 (Engel 1865), nom. illeg. hom., non Mart. (Martius 1824).*Euterpe stenophylla* Trail ex Thurn, *Timebri 3*: 229 (Thurn 1884).*Euterpe jataguensis* Barb.Rodr., *Contr. Jard. Bot. Rio de Janeiro 1*: 12 (Barbosa Rodrigues 1901).*Euterpe subruminata* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 15*: 3 [15 Feb. 1940] (Burret 1940).*Euterpe petiolata* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 15*: 101 [15 Feb. 1940] (Burret 1940).*Euterpe langloisii* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 13*: 346 [30 Dec. 1936] (Burret 1936).*Euterpe confertiflora* L.H.Bailey, *Gentes Herb. 7*: 427 (Bailey 1947).NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana. Incipiently domesticated by pre-Columbian Amerindians (Levis *et al.* 2017).

VERNACULAR NAMES. — Ka: wapu • Wp: kusili ape, wasey iipipë'i, wasey sì • Wn: wapuimë • Nt: pina • Br: açaí-da-mata, juçara.

HERBARIUM DATA (FG). — No specimen seen, present according to (Granville & Gayot 2014).

SIZE. — Up to 25 cm dbh (Granville & Gayot 2014).

Genus *Manicaria* Gaertn.[151] *Manicaria saccifera* Gaertn.

(Fig. 10D)

Fruct. Sem. Pl. 2: 468 [Sep.-Dec. 1791] (Gaertner 1791). — *Pilophora saccifera* (Gaertn.) H.Wendl., *Palmiers [Kerchove] 253* (Wendland 1878).*Pilophora testicularis* Jacq., *Fragm. Bot. 32* (Jacquin 1801).*Manicaria pluknetii* Griseb. & H.Wendl., *Fl. Brit. W.I. [Grisebach] 518* [Oct. 1864] (Grisebach & Wendland 1864). — *Manicaria saccifera* var. *pluknetii* (Griseb. & H.Wendl.) Drude, *Fl. Bras. [Martius] 3* (2): 520 [1 May 1882] (Drude 1882).*Manicaria saccifera* var. *mediterranea* Trail, *J. Bot. 14*: 332 (Trail 1876).*Manicaria atricha* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 10*: 1013 [1 Aug. 1930] (Burret 1930).

VERNACULAR NAMES. — Pa: tuuvan • Ka: tululi • Te: tululi • Wp: tululi • Cr: tourouri • Fr: toulouri • Br: buçú, ubuçú.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-J. de Granville 10319*.

SIZE. — Up to 30 cm dbh (Granville & Gayot 2014).

Genus *Mauritia* L.f.[152] *Mauritia flexuosa* L.f.*Suppl. Pl.*: 454 [“1781” publ. Apr. 1782] (Linnaeus 1782).*Sagus americana* Poir., *Encycl. [J. Lamarck et al.] 6* (2): 395 [28 Aug. 1805] (Poirier 1805).*Mauritia vinifera* Mart., *Hist. Nat. Palm. 2* (2): 42 [before 13 Apr. 1824] (Martius 1824).*Mauritia sagus* Schult. & Schult.f., *Syst. Veg. [Roemer & Schultes] 7* (2): 1321 [Oct.-Dec. 1830] (Schultes & Schultes 1830), nom. illeg. superfl. (based on *Sagus americana*).*Mauritia setigera* Griseb. & H.Wendl., *Fl. Brit. W.I. [Grisebach] 515* [Oct. 1864] (Grisebach & Wendland 1864).*Saguerus americanus* H.Wendl., *Palmiers [Kerchove] 256* (Wendland 1878), “*americana*”, nom. nud.*Mauritia sphaerocarpa* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 10*: 569 [30 Mar. 1929] (Burret 1929).*Mauritia minor* Burret, *Notizbl. Bot. Gart. Berlin-Dahlem 11*: 1 [30 Dec. 1930] (Burret 1930).*Mauritia flexuosa* var. *venezuelana* Steyermark., *Fieldiana, Bot. 28* (1): 90 (Steyermark 1951).NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana. Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). *Saguerus americanus* H.Wendl. (in Kerchove 1878: 256, as “*Saguerus americana*” Poir., vide *Mauritia flexuosa* L. f.) is obviously a mistranscript of *Sagus americana*. Wendland lists “*Sagus americana* Poir.” as a synonym of *M. flexuosa* (Kerchove 1878: 251). *Sagus americana* Poirier (1805: 395) was based on Aublet’s description of the “*Bache*” (Aublet 1775, 2[Suppl.]: 103), which did not come with a Latin name nor a formal diagnosis.

VERNACULAR NAMES. — Pa: isao, isau • Ka: mulisi, mulishi • Te: bilitsi • Wp: milisi • Wn: koai, kuwai • Nt: moysi • Cr: palmié-bache • Br: buriti.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-J. de Granville 3750*.

SIZE. — Up to 60 cm dbh (Granville & Gayot 2014).

Genus *Oenocarpus* Mart.

[153] *Oenocarpus bacaba* Mart.

Hist. Nat. Palm. 2 (1): 24 [Nov. 1823] (Martius 1823).

Areca bacaba Arruda, *Trav. Brazil [H. Koster]*: 490 (Arruda 1816), *nom. nud.*

Oenocarpus hoppii Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 1041 [20 Jan. 1934] (Burret 1934).

Oenocarpus grandis Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 12: 612 [6 Dec. 1935] (Burret 1935). — *Oenocarpus bacaba* var. *grandis* (Burret) Wess.Boer, *Pittieria* 17: 131 (Wessels Boer 1988).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017), and one of five species whose relative density in tree communities likely indicative of pre-Columbian archeological sites in French Guiana (Molino *et al.* 2021).

VERNACULAR NAMES. — Pa: woki • Ka: kumin, kumu • Te: pindo uhu, wil'a • Wp: a'i pino, pino, pino akusiway, pino e'e, pino tamanuwa • Wn: kumu • Nt: komu • Cr: komou • Br: bacaba.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (original material P[P00725199]).

INVENTORY DATA (FG). — 660 trees in 131 plots; $F_{\max} = 10\%$; $dbh_{inv} = 31$ cm.

[154] *Oenocarpus bataua* Mart. var. *bataua*

Hist. Nat. Palm. 2 (1): 23 [Nov. 1823] (Martius 1823). — *Jessenia bataua* (Mart.) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 300 [30 Mar. 1928] (Burret 1928).

Jessenia polycarpa H.Karst., *Linnaea* 28: 388 (Karsten 1857).

Jessenia repanda Engl., *Linnaea* 33: 691 (Engler 1865).

Oenocarpus seje Cuervo, *Trat. Elem. Bot.*: 458 (Cuervo 1913), *nom. nud.*

Jessenia weberbaueri Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 840 [30 Dec. 1929] (Burret 1929).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: patau • Ka: patawa • Te: pataua • Wp: patawa • Nt: afá komu • Cr: patawa • Br: patauá.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *F. Billet & B. Jadin 1232*.

INVENTORY DATA (FG). — 1140 trees in 147 plots; $F_{\max} = 7.5\%$; $dbh_{inv} = 31.8$ cm.

[155] *Oenocarpus bataua* var. *oligocarpus* (Griseb. & H.Wendl.) A.J.Hend.

Palms Amazon: 120 (Henderson 1995), “*oligocarpa*”. — *Jessenia oligocarpa* Griseb. & H.Wendl., *Fl. Brit. W.I. [Grisebach]* 516 [Oct. 1864] (Grisebach & Wendel 1864). — *Oenocarpus oligocarpa* (Griseb. & H.Wendl.) Wess.Boer, *Fl. Suriname* 5 (1): 58 (Wessels Boer 1965). — *Jessenia bataua* subsp. *oligocarpa* (Griseb. & H.Wendl.) Balick, *Advances Econ. Bot.* 3: 126 (Balick 1986).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: tavu, tawu • Ka: sapululi, tabululi, tapululi • Wp: tapululi • Cr: gro-komou.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville 3411*.

SIZE. — Up to 40 cm dbh (Granville & Gayot 2014).

Genus *Socratea* H.Karst.

[156] *Socratea exorrhiza* (Mart.) H.Wendl.

Bonplandia 8: 103 (Wendland 1860). — *Iriartea exorrhiza* Mart., *Hist. Nat. Palm.* 2 (2): 36 [before 13 Apr. 1824] (Martius 1824), “*exhoriza*”.

Iriartea orbigniana Blume ex Mart., *Hist. Nat. Palm.* 3 (7): 187 [23 Sep. 1838] (Martius 1838). — *Socratea orbigniana* (Blume ex Mart.) H.Karst., *Linnaea* 28: 264 [“1856” publ. Jan. 1857] (Karsten 1857), “*Orbignyana*”. — *Iriartea exorrhiza* var. *orbigniana* (Blume ex Mart.) Drude, *Fl. Bras. [Martius]* 3 (2): 540 [1 May 1882] (Drude 1882).

Socratea elegans H.Karst., *Linnaea* 28: 264 [“1856” publ. Jan. 1857] (Karsten 1857). — *Iriartea exorrhiza* var. *elegans* (H.Karst.) Drude, *Fl. Bras. [Martius]* 3 (2): 539 [1 May 1882] (Drude 1882).

Iriartea durissima Oerst., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1858: 30 (Oersted 1859). — *Socratea durissima* (Oerst.) H.Wendl., *Bonplandia* 8: 103 (Wendland 1860).

Iriartea philonotia Barb.Rodr., *Enum. Palm. Nov.*: 13 (Barbosa Rodrigues 1875). — *Socratea philonotia* (Barb.Rodr.) Hook.f., *Gen. Pl. [Bentham & Hooker f.]* 3 (2): 900 [14 Apr. 1883] (Hooker 1883).

Socratea macrochlamys Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 918 [30 Mar. 1930] (Burret 1930).

Socratea microchlamys Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 3 [30 Dec. 1930] (Burret 1930).

Socratea hoppii Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 232 [10 Nov. 1931] (Burret 1931).

Socratea gracilis Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 1 [15 Feb. 1940] (Burret 1940).

Socratea albolineata Steyermark., *Fieldiana, Bot.* 28 (1): 9 (Steyermark 1951).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: pup • Ka: pasiwi • Te: pats'i • Wp: pasi'i • Wn: pëpë • Nt: paasaa • Cr: awara-monpé • Br: paxiúba.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori 2415*.

INVENTORY DATA (FG). — 81 trees in 39 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 20\text{ cm}$.

Genus *Syagrus* Mart.

[157] *Syagrus inajai* (Spruce) Becc.

Agric. Colon. 10: 467 (Beccari 1916). — *Maximiliana inajai* Spruce, *J. Linn. Soc., Bot.* 11: 163 ["1871" publ. Dec. 1869] (Spruce 1869). — *Cocos inajai* (Spruce) Trail, *J. Bot.* 15: 79 (Trail 1877).

Cocos aequatorialis Barb.Rodr., *Enum. Palm. Nov.*: 38 (Barbosa Rodrigues 1875).

Cocos speciosa Barb.Rodr., *Enum. Palm. Nov.*: 38 (Barbosa Rodrigues 1875). — *Syagrus speciosa* (Barb.Rodr.) Barb.Rodr., *Prot.-App. Enum. Palm. Nov.* 49 (Barbosa Rodrigues 1879). — *Calappa speciosa* (Barb.Rodr.) Kuntze, *Revis. Gen. Pl.* 2: 982 [5 Nov. 1891] (Kuntze 1891).

Cocos chavesiana Barb.Rodr. ex Becc., *Malpighia* 1: 445 (Beccari 1887). — *Syagrus chavesiana* (Barb.Rodr. ex Becc.) Barb.Rodr., *Vellozia* 1: 52 (Barbosa Rodrigues 1888).

VERNACULAR NAMES. — Pa: kurip, kwip (adult form), mahayapna (juvenile form) • Te: dzata'i • Wp: malaliapu, malaliapu lo (juvenile form), yata'i (adult form) • Wn: halikuli • Cr: féy-chasèr (juvenile form), parépou-djab (adult form) • Br: inajai, jata, pupunha-de-porco.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 9913*.

INVENTORY DATA (FG). — 17 trees in 5 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 13\text{ cm}$.

[158] *Syagrus stratincola* Wess.Boer

Fl. Suriname 5 (1): 170 (Wessels Boer 1965).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-J. de Granville 17651*.

SIZE. — Up to 10 cm dbh (Granville & Gayot 2014).

Family BIGNONIACEAE Juss.
Genus *Handroanthus* Mattos

[159] *Handroanthus capitatus* (Bureau & K.Schum.) Mattos

Loefgrenia 50: 4 (Mattos 1970). — *Tecoma capitata* Bureau & K.Schum., *Fl. Bras. [Martius]* 8 (2): 337 [15 Feb. 1897] (Bureau & Schumann 1897). — *Tabebuia capitata* (Bureau & K.Schum.) Sandwith, *Recueil Trav. Bot. Néerl.* 34: 226 (Sandwith 1937).

Tecoma leucoxylon (L.) Mart. ex DC. var. *miquelii* A.DC., *Prod. [A. P. de Candolle]* 9: 219 [1 Jan. 1845] (Candolle 1845).

Tabebuia glomerata Urb., *Repert. Spec. Nov. Regni Veg.* 14: 305 (Urban 1916).

Tabebuia hypolepra Sprague & Sandwith, *Bull. Misc. Inform. Kew* 1932 (1): 25 [23 Feb. 1932] (Sprague & Sandwith 1932).

VERNACULAR NAMES. — Pa: kwik-kamwi • Ka: alawani, alawone • Wp: waiw'i • Cr: débenn-vert, lébenn-vert • Br: pau-d'arco.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5025*.

INVENTORY DATA (FG). — 11 trees in 8 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 93.1\text{ cm}$.

[160] *Handroanthus impetiginosus* (Mart. ex DC.) Mattos

Loefgrenia 50: 2 (Mattos 1970). — *Tecoma impetiginosa* Mart., *Syst. Mat. Med. Veg. Bras.* 54 (Martius 1843), nom. nud. — *Tecoma impetiginosa* Mart. ex DC., *Prod. [A. P. de Candolle]* 9: 218 [1 Jan. 1845] (Candolle 1845). — *Tabebuia impetiginosa* (Mart. ex DC.) Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (5): 176 (Standley 1936).

Tabebuia avellaneda Lorentz ex Griseb., *Symb. Fl. Argent.*: 258 (Grisebach 1879). — *Gelsemium avellaneda* (Lorentz ex Griseb.) Kuntze, *Revis. Gen. Pl.* 3 (3): 245 [28 Sep. 1898] (Kuntze 1898). — *Tecoma avellaneda* (Lorentz ex Griseb.) Speg., *Anal. Soc. Rural Argent. (Cat. Descr. Maderas)*: 379 (Spegazzini 1910). — *Handroanthus avellaneda* (Lorentz ex Griseb.) Mattos, *Loefgrenia* 50: 3 (Mattos 1970).

Tabebuia palmeri Rose, *Contr. U.S. Natl. Herb.* 1 (4): 109 [30 June 1891] (Rose 1891).

Tecoma adenophylla K.Schum., *Fl. Bras. [Martius]* 8 (2): 412 [15 Feb. 1897] (Bureau & Schumann 1897).

Tecoma ipe Mart. ex K.Schum. var. *integrifolia* Hassl., *Revista Inst. Parag.* 3: 166 (Hassler 1901), nom. nud.

Tecoma ipe var. *integra* Sprague, *Bull. Herb. Boissier*, sér. 2, 5: 86 (Sprague 1905). — *Tecoma integra* (Sprague) Chodat, *Bull. Soc. Bot. Genève*, sér. 2, 9: 242 (Chodat 1917), "integrum". — *Tabebuia ipe* (Mart. ex K.Schum.) Standl. var. *integra* (Sprague) Sandwith, *Lloydia* 2 (3): 213 (Sandwith 1939).

Tecoma ipe f. *leucotricha* Hassl., *Repert. Spec. Nov. Regni Veg.* 9: 60 (Hassler 1910).

Tabebuia nicaraguensis S.F.Blake, *Contr. Gray Herb.*, n.s., 52: 95 (Blake 1917).

Tecoma avellaneda var. *alba* Lillo, *Seg. Contr. Conoc. Arbol. Argent.*: 13 (Lillo 1917).

Tabebuia dugandii Standl., *Trop. Woods* 36: 17 (Standley 1933).

Tabebuia schunkevigoi D.R.Simpson, *Fieldiana, Bot.* 36 (1): 1 (Simpson 1972).

VERNACULAR NAMES. — Nt: lebi giin ati • Cr: lébenn-ròz • Fr: ébène rose, ébène soufrée • Br: pau-d'arco-roxo.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *S. Gonzalez 1093*.

INVENTORY DATA (FG). — 12 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 120\text{ cm}$.

[161] *Handroanthus obscurus* (Bureau & K.Schum.) Mattos

Loefgrenia 50: 4 (Mattos 1970). — *Tecoma obscura* Bureau & K.Schum., *Fl. Bras. [Martius]* 8 (2): 343 [15 Feb. 1897] (Bureau &

Schumann 1897). — *Tabebuia obscura* (Bureau & K.Schum.) Sandwith, *Recueil Trav. Bot. Néerl.* 34: 226 (Sandwith 1937).

Tabebuia subtilis var. *schultesiana* Sandwith, *Bot. Mus. Leafl.* 17 (3): 96 [21 Nov. 1955] (Sandwith 1955). — *Tabebuia obscura* var. *schultesiana* (Sandwith) Sandwith, *Mutisia* 25: 16 (Sandwith 1956).

HERBARIUM DATA (FG). — A single collection, *C. Feuillet 1122*.

SIZE. — Venezuela, Amazonas. *G. Davide 27581* (MO), 10 m.

[162] *Handroanthus serratifolius* (Vahl) S.O.Grose

Syst. Bot. 32 (3): 666 [1 July 2007] (Grose 2007). — *Bignonia serratifolia* Vahl, *Eclat. Amer.* 2: 46 (Vahl 1798). — *Tabebuia serratifolia* (Vahl) G.Nicholson, *Ill. Dict. Gard.* 7: 1 (Nicholson 1887).

Bignonia flavescens Vell., *Fl. Flumin.* 252 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829). — *Tecoma flavescens* (Vell.) Mart. ex DC., *Prodr. [A. P. de Candolle]* 9: 216 [1 Jan. 1845] (Candolle 1845). — *Handroanthus flavescens* (Vell.) Mattos, *Loefgrenia* 50: 2 (Mattos 1970).

Bignonia araliacea Cham., *Linnæa* 7: 683 (Chamisso 1832). — *Tecoma araliacea* (Cham.) DC., *Prodr. [A. P. de Candolle]* 9: 221 [1 Jan. 1845] (Candolle 1845). — *Gelseminum araliaceum* (Cham.) Kuntze, *Revis. Gen. Pl.* 3 (2): 245 [28 Sep. 1898] (Kuntze 1898). — *Tabebuia araliacea* (Cham.) Morong & Britton, *Ann. New York Acad. Sci.* 7: 190 (Morong & Britton 1893). — *Handroanthus araliaceus* (Cham.) Mattos, *Loefgrenia* 50: 2 (Mattos 1970).

Tecoma speciosa DC., *Prodr. [A. P. de Candolle]* 9: 218 [1 Jan. 1845] (Candolle 1845). — *Gelseminum speciosum* (DC.) Kuntze, *Revis. Gen. Pl.* 3 (3): 245 [28 Sep. 1898] (Kuntze 1898).

Tecoma conspicua DC., *Prodr. [A. P. de Candolle]* 9: 221 [1 Jan. 1845] (Candolle 1845). — *Bignonia conspicua* Rich. ex DC., *Prodr. [A. P. de Candolle]* 9: 221 [1 Jan. 1845] (Candolle 1845), nom. nud. pro syn.

Tecoma patrisiana DC., *Prodr. [A. P. de Candolle]* 9: 221 [1 Jan. 1845] (Candolle 1845).

Tecoma nigricans Klotzsch ex M.R.Schomb., *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1159 ["1848" publ. 7-10 Mar. 1849] (Schomburgk 1849), nom. nud.

Tecoma atractocarpa Bureau & K.Schum., *Fl. Bras. [Martius]* 8 (2): 326 [15 Feb. 1897] (Bureau & Schumann 1897). — *Handroanthus atractocarpus* (Bureau & K.Schum.) Mattos, *Loefgrenia* 50: 2 (Mattos 1970).

Tabebuia monticola Pittier, *Cat. Fl. Venez. [Pittier]* 2: 409 (Pittier 1947), nom. nud.

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: kwik • Ka: alawani, alawone • Te: tadju • Wp: tayi • Wn: alaone, alawone • Nt: giin ati • Cr: débenn-vert, lébenn • Fr: ébène verte • Br: ipê, pau-d'arco-amarelo.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (original material of *Tecoma patrisiana*: G-DC, *G00133858*).

INVENTORY DATA (FG). — 54 trees in 33 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 110$ cm.

[163] *Handroanthus subtilis*
(Sprague & Sandwith) S.O.Grose
(Fig. 11A)

Syst. Bot. 32 (3): 666 [1 July 2007] (Grose 2007). — *Tabebuia subtilis* Sprague & Sandwith, *Bull. Misc. Inform. Kew* 1932 (1): 23 [23 Feb. 1932] (Sprague & Sandwith 1932).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4826*.

INVENTORY DATA (FG). — 19 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.3$ cm.

Genus *Jacaranda* Juss.

[164] *Jacaranda copaia* (Aubl.) D.Don subsp. *copaia*

Edinburgh Philos. J. 9 (18): 267 [Oct. 1823] (Don 1823). — *Bignonia copaia* Aubl., *Hist. Pl. Guiane* 2: 650 [Jun.-Dec. 1775] (Aublet 1775), "Copaiid". — *Bignonia procera* Willd., *Sp. Pl.*, ed. 4 3 (1): 307 (Willdenow 1800), nom. illeg. superfl. (based on *Bignonia copaia*). — *Jacaranda procera* R.Br., *Bot. Mag.* 49: t. 2327 (Brown 1822), nom. illeg. superfl. (based on *Bignonia copaia*).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: pada • Ka: kopaya, kupaiya, kupaya • Te: mee'i • Wp: mapili poá, pala'i • Wn: kupahja, kupaja • Nt: fayaati, feyati, yasimanbo • Cr: koupaya • Fr: copaia, faux simarouba • Br: caroba, pará-pará.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet 32* (original material P-Ad. [*P00320334*]).

INVENTORY DATA (FG). — 374 trees in 146 plots; $F_{\max} = 2.8\%$; $dbh_{\text{inv}} = 69.4$ cm.

[165] *Jacaranda copaia* subsp. *spectabilis*
(Mart. ex DC.) A.H.Gentry

Rhodora 79: 441 (Gentry 1977). — *Jacaranda spectabilis* Mart. ex DC., *Prodr. [A. P. de Candolle]* 9: 229 [1 Jan. 1845] (Candolle 1845). — *Jacaranda copaia* var. *spectabilis* (Mart. ex DC.) Bureau ex Bureau & K.Schum., *Fl. Bras. [Martius]* 8 (2): 387 [15 Feb. 1897] (Bureau & Schumann 1897).

Jacaranda copaia var. *paraensis* Huber, *Bull. Soc. Bot. Genève*, sé. 2, 6: 202 ["1914" publ. 1915] (Huber 1915). — *Jacaranda paraensis* (Huber) Vattimo, *Rodriguésia* 29 (42): 285 (Vattimo 1977).

Jacaranda superba Pittier, *Bol. Soc. Venez. Ci. Nat.* 6 (41): 19 ["1939" publ. 1940] (Pittier 1940).

Jacaranda amazonensis Vattimo, *Rodriguésia* 29 (44): 231 (Vattimo 1978).

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost 1119*.

SIZE. — Up to 45 cm dbh (Gentry 1992).




FIG. 11. — Bignoniaceae: **A**, *Handroanthus subtilis* (Sprague & Sandwith) S.O. Grose (D. Sabatier & E. Fonty 5588); **B**, *Tabebuia insignis* (Miq.) Sandwith (M.-F. Prévost & D. Sabatier 4818). Bixaceae: **C**, *Bixa* sp. A (D. Sabatier & J.-F. Molino 5609); **D**, *Cochlospermum orinocense* (Kunth) Steud. (D. Sabatier & J.-F. Molino 5070). © D. Sabatier/IRD.

[166] *Jacaranda duckei* Vattimo

Rodriguésia 36 (59): 79 (Vattimo 1984).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *A.H. Gentry & C. Feillet 63218*.

SIZE. — Up to 12 m tall (Gentry 1992).

[167] *Jacaranda obtusifolia* Bonpl.
subsp. *rhombifolia* (G.Mey.) A.H.Gentry

Mem. New York Bot. Gard. 29: 257 (Gentry 1978). — *Jacaranda rhombifolia* G.Mey., *Prim. Fl. Esseq.*: 213 [Nov. 1818] (Meyer 1818). — *Jacaranda filicifolia* D.Don, *Edinburgh Philos. J.* 9 (18): 266 [Oct. 1823] (Don 1823), nom. illeg. superfl. (based on *Jacaranda rhombifolia* G.Mey.).

folia). — *Jacaranda obtusifolia* var. *rhombifolia* (G.Mey.) Sandwith, *Kew Bull.* 8 (4): 458 [“1953” publ. 2 Jan. 1954] (Sandwith 1954).

Bignonia filicifolia A.Anderson, *Trans. Soc. London Encour. Arts* 25: 200 (Anderson 1807), *nom. nud.*

Jacaranda filicifolia var. *puberula* K.Schum., *Fl. Bras. [Martius]* 8 (2): 390 [15 Feb. 1897] (Schumann 1897).

VERNACULAR NAMES. — Ka: arisisi’i, woto kolalu.

HERBARIUM DATA (FG). — A single collection, *M.-F. Prévost 1718*.

SIZE. — Up to 15 m tall (Gentry 1992).

Genus *Tabebuia* Gomes ex DC.

[168] *Tabebuia fluviatilis* (Aubl.) DC.

Prodr. [A. P. de Candolle] 9: 215 [1 Jan. 1845] (Candolle 1845). — *Bignonia fluviatilis* Aubl., *Hist. Pl. Guiane* 2: 655 [Jun.-Dec. 1775] (Aublet 1775). — *Zeyheria fluviatilis* (Aubl.) Miq., *Flora* 25 (2): 431 (Miquel 1842). — *Couralia fluviatilis* (Aubl.) Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 15 [Aug.-Sep. 1842] (Splitgerber 1842). — *Tecoma fluviatilis* (Aubl.) Miq., *Stirp. Surinam. Select.*: 121 [“1850” publ. Mar. 1851] (Miquel 1851). — *Potamoxyton fluviatile* (Aubl.) Pichon, *Bull. Soc. Bot. France* 92 (9): 228 (Pichon 1945).

Bignonia aquatilis E.Mey., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 780 (Meyer 1825). — *Tecoma aquatilis* (E.Mey.) DC., *Prodr. [A. P. de Candolle]* 9: 225 [1 Jan. 1845] (Candolle 1845). — *Tabebuia aquatilis* (E.Mey.) Sprague & Sandwith, *Bull. Misc. Inform. Kew* 1932 (1): 21 [23 Feb. 1932] (Sprague & Sandwith 1932).

Bignonia digitata E.Mey., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 782 (Meyer 1825). — *Zeyheria digitata* (E.Mey.) Miq., *Flora* 25 (2): 431 (Miquel 1842).

Tecoma meyeriana A.DC., *Prodr. [A. P. de Candolle]* 9: 221 [1 Jan. 1845] (Candolle 1845).

VERNACULAR NAMES. — Pa: waravru-kamwi-seine • Ka: pandalan • Cr: bwa-blanché.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.B. Aublet 200* (original material P-Ad., not seen); *R.A.A. Oldeman B-884*: 7 m × 23 cm.

[169] *Tabebuia insignis* (Miq.) Sandwith
(Fig. 11B)

Recueil Trav. Bot. Néerl. 34: 224 (Sandwith 1937). — *Tecoma insignis* Miq., *Stirp. Surinam. Select.*: 122 [“1850” publ. Mar. 1851] (Miquel 1851). — *Gelseminum insigne* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 480 [5 Nov. 1891] (Kuntze 1891).

Bignonia dura Klotzsch ex R.Knuth, *Repert. Spec. Nov. Regni Veg. Beih.* 43: 638 (Knuth 1827), *nom. nud.*

Tecoma dura Bureau & K.Schum., *Nat. Pflanzenfam. [Engler & Prantl]* 4 (3b): 238 (Bureau & Schumann 1894). — *Tabebuia dura* (Bureau & K.Schum.) Sprague & Sandwith, *Bull. Misc. Inform. Kew* 1932 (1): 21 [23 Feb. 1932] (Sprague & Sandwith 1932). — *Handroanthus durus* (Bureau & K.Schum.) Mattos, *Loefgrenia* 50: 4 (Mattos 1970).

Tabebuia roraimae Oliv., *Timebri* 5: 201 (Oliver 1886). — *Tecoma roraimae* (Oliv.) K.Schum., *Nat. Pflanzenfam. [Engler & Prantl]* 4 (3b): 238 (Schumann 1894).

Tecoma leucoxylon (L.) Mart. ex DC. var. *salpingophora* Bureau & K.Schum., *Fl. Bras. [Martius]* 8 (2): 342 [15 Feb. 1897] (Bureau & Schumann 1897).

Tecoma albiflora Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 175 (Ducke 1925).

VERNACULAR NAMES. — Pa: kwik-kamwi, kwik-marikasmatgene • Ka: kupayalan, panda, pokata • Cr: bwa-blanché • Br: capitari, ipê-branco-do-brejo.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4818*.

INVENTORY DATA (FG). — 50 trees in 19 plots; $F_{\max} = 1.7\%$; $dbh_{inv} = 85.9$ cm.

[170] *Tabebuia stenocalyx* Sprague & Stapf

Bull. Misc. Inform. Kew 1910 (6): 196 (Sprague & Stapf 1910).

HERBARIUM DATA (FG). — A single collection, *R.S. Cowan 38837* (MO).

SIZE. — Up to 25 m tall (Gentry 1992).

Family BIXACEAE Kunth
Genus *Bixa* L.

[171] *Bixa arborea* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 6: 87 (Huber 1910).

VERNACULAR NAMES. — Pa: ihap-kamwi • Wp: uluku tawa • Br: urucu-da-mata.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & P. Birnbaum 4451*.

INVENTORY DATA (FG). — 5 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{inv} = 22.9$ cm.

[172] *Bixa* sp. A
(Fig. 11C)

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *B. Dutrèvre 489*, $dbh = 14$ cm.

Genus *Cochlospermum* Kunth

[173] *Cochlospermum orinocense* (Kunth) Steud.
(Fig. 11D)

Nomencl. Bot. [Steudel], ed. 2, 1: 393 (Steudel 1840). — *Bombax orinocense* Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 301 [June 1822] (Kunth 1822). — *Wittelsbachia orinocensis* (Kunth) Mart., *Nova genera et species plantarum [Martius]* 1 (4): 83 [“1824” publ. Jan.-Mar. 1826] (Martius 1826). — *Maximilianea orinocensis* (Kunth) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Cochlospermum parkeri Planch., *London J. Bot.* 6: 310 (Planchon 1847). — *Maximiliane parkeri* (Planch.) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Cochlospermum parvifolium Planch., *London J. Bot.* 6: 311 (Planchon 1847), “parviaefolium”. — *Maximiliane parvifolia* (Planch.) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891), “pawiaefolia”.

Cochlospermum wentii Pulle, *Enum. Vasc. Pl. Surinam:* 310 (Pulle 1906).

Cochlospermum williamsii J.F.Macbr., *Candollea* 5: 388 (Macbride 1934), “*Cochleospermus Williamsii*”.

NOTE. — A species found only on the edges of inselbergs.

VERNACULAR NAMES. — Pa: sirisiri, wakāu-dariu • Wp: māū'i sili • Cr: koton-périch • Br: algodão-do-mato, castanha-de-periquito, periquiteira-da-terra-firme, siri-siri.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5070*, dbh 25 cm.

Family BURSERACEAE Kunth
Genus *Dacryodes* Vahl

[174] *Dacryodes cuspidata* (Cuatrec.) Daly

Brittonia 41 (1): 25 (Daly 1989). — *Trattinnickia cuspidata* Cuatrec., *Webbia* 12 (2): 423 (Cuatrecasas 1957).

VERNACULAR NAMES. — Nt: alauau.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2625*.

INVENTORY DATA (FG). — 106 trees in 43 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 86$ cm.

[175] *Dacryodes decidua* Daly & M.C.Martínez

Brittonia 71 (2): 206 (Daly & Martínez 2019).

HERBARIUM DATA (FG). — 1 collection at CAY. Sel. exs.: *D. Loubry 1224* (holo-, NY[01076225, 01076226]; iso-, CAY[CAY115397, CAY115398]), dbh = 50 cm.

[176] *Dacryodes nitens* Cuatrec.

Trop. Woods 106: 64 (Cuatrecasas 1957).

VERNACULAR NAMES. — Pa: araksim, sedri-seine • Te: waluwā'i • Wp: ayawa sili • Nt: gaan moni • Br: caranha.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *D. Sabatier 1196*.

INVENTORY DATA (FG). — 155 trees in 72 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 49.7$ cm.

[177] *Dacryodes oblongipetala* Daly & M.C.Martínez

Brittonia 71 (2): 213 (Daly & Martínez 2019).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3097*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 21.7$ cm.

[178] *Dacryodes roraimensis* Cuatrec.

Trop. Woods 106: 62 (Cuatrecasas 1957).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori & T.D. Pennington 17983*.

SIZE. — Up to 40 m tall (Daly 1997).

[179] *Dacryodes villosa* Daly & M.C.Martínez

Brittonia 71 (2): 223 (Daly & Martínez 2019).

NOTES. — Known only from French Guiana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3382* (holo-, NY[03231016, 03231017]; iso-, CAY[CAY081853]), dbh = 80 cm.

[180] *Dacryodes* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5555*.

INVENTORY DATA (FG). — 25 trees in 15 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 85$ cm.

Genus *Protium* Burm.f.

[181] *Protium altissimum* (Aubl.) Marchand
(Fig. 12A)

Adansonia [Baillon] 8: 51 (Marchand 1867). — *Icica altissima* Aubl., *Hist. Pl. Guiane* 1: 342 [Jun.-Dec. 1775] (Aublet 1775). — *Amyris altissima* (Aubl.) Willd., *Sp. Pl. ed. 4* 2 (1): 336 [Mar. 1799] (Willdenow 1799). — *Elaphrium altissimum* (Aubl.) Spreng. ex D.Dietr., *Syn. Pl. [D. Dietrich]* 2: 1272 [1-20 Dec. 1840] (Dietrich 1840). — *Bursera altissima* (Aubl.) Baill., *Hist. Pl. [Baillon]* 5: 297 [June 1874] (Baillon 1874). — *Tingulonga altissima* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891). — *Tetragastris altissima* (Aubl.) Swart, *Recueil Trav. Bot. Néerl.* 39: 413 (Swart 1942). — *Protium excelsior* Byng & Christenh., *Global Fl.* 4: 140 [9 Feb. 2018] (Byng & Christenhusz 2018), nom. illeg. superfl. (based on *Icica altissima*).

Tetragastris phanerosepala Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 209 [27 June 1932] (Sandwith 1932).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ayay, ayay-wahuyo • Ka: pilika • Te: dzadza'i • Wp: yaya'i, yaya'i sī • Wn: aimalaje, simajali • Nt: lebi sali • Cr: bwa-yaya, lanson-rouj, lanson-roz, sèd-bagas • Fr: cèdre

bagasse, cèdre blanc (*fide* Aublet 1775) • Br: almesclão, breu-grande, breu-manga.

HERBARIUM DATA (FG). — 218 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[not seen], designated by Swart [1942: 415]); *B.M. Boom & S.A. Mori 2222*.

INVENTORY DATA (FG). — 894 trees in 80 plots; $F_{\max} = 24.1\%$; $\text{dbh}_{\text{inv}} = 105\text{ cm}$.

[182] *Protium altsonii* Sandwith

Bull. Misc. Inform. Kew 1928 (9): 369 [25 Oct. 1928] (Sandwith 1928).

Protium puberulentum Steyermark., *Fieldiana, Bot.* 28 (2): 276 (Steyermark 1952).

Protium paraense Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 11: 4 (Cuatrecasas 1961).

VERNACULAR NAMES. — Pa: marinaiwa-wašiunō • Br: breu-branco.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3983*.

INVENTORY DATA (FG). — 22 trees in 8 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 40.4\text{ cm}$.

[183] *Protium amazonicum* (Cuatrec.) Daly

Fl. Venez. Guayana 3: 713 (Daly 1997). — *Paraprotium amazonicum* Cuatrec., *Bot. Mus. Leafl.* 18 (4): 156 [4 Apr. 1958] (Cuatrecasas 1958).

Protium fimbriatum Swart, *Acta Bot. Neerl.* 15: 48 (Swart 1966).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman 1862*.

SIZE. — Brazil, Amazonas. *C.A. Cid Ferreira 6948* (MO), 18 m × 25 cm.

[184] *Protium apiculatum* Swart

Recueil Trav. Bot. Néerl. 39: 201 (Swart 1942).

Protium firmum Swart, *Acta Bot. Neerl.* 15: 49 (Swart 1966).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), and one of five species whose relative density in tree communities likely indicative of pre-Columbian archeological sites in French Guiana (Molino *et al.* 2021).

VERNACULAR NAMES. — Pa: ahuwahu, auwau, auwau-purubumna, ayao, ayau • Wp: ii, waluwai, waluwai pilá, waluwai sī • Nt: lebi moni, sali • Cr: bwa-lansan • Fr: encens • Br: breu-vermelho.

HERBARIUM DATA (FG). — 143 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4426*.

INVENTORY DATA (FG). — 307 trees in 82 plots; $F_{\max} = 6.4\%$; $\text{dbh}_{\text{inv}} = 54.1\text{ cm}$.

[185] *Protium aracouchini* (Aubl.) Marchand

Adansonia [Baillon] 8: 51 (Marchand 1867). — *Icica aracouchini* Aubl., *Hist. Pl. Guiane* 1: 343 [Jun.-Dec. 1775] (Aublet 1775). — *Bursera aracouchini* (Aubl.) Baill., *Traité Bot. Méd. Phan.* 2: 951 [Mar. 1884] (Baillon 1884), “Aracouchili”. — *Tingulonga aracouchini* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891). — *Amyris heterophylla* Willd., *Sp. Pl.*, ed. 4 2 (1): 335 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Icica aracouchini*). — *Icica heterophylla* DC., *Prodr. [A. P. de Candolle]* 2: 77 [mid Nov. 1825] (Candolle 1825), nom. illeg. superfl. (based on *Icica aracouchini*). — *Elaphrium heterophyllum* Spreng. ex D. Dietr., *Syn. Pl. [D. Dietrich]* 2: 1271 [1-20 Dec. 1840] (Dietrich 1840), nom. illeg. superfl. (based on *Amyris heterophylla*, thus indirectly on *Icica aracouchini*).

Protium divaricatum Engl. var. *intermedium* Swart, *Recueil Trav. Bot. Néerl.* 39: 199 (Swart 1942).

Protium aracouchini var. *angustifolium* Swart, *Acta Bot. Neerl.* 15: 45 (Swart 1966).

VERNACULAR NAMES. — Pa: marinaiwa • Ka: alakuseli, alakušini • Wn: alakuseli.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN[LINN-HS 672.2]).

INVENTORY DATA (FG). — 23 trees in 12 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 42.2\text{ cm}$.

[186] *Protium calendulinum* Daly

Brittonia 59 (1): 7 (Daly 2007).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *R.A.A. Oldeman 1912*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 15\text{ cm}$.

[187] *Protium cuneatum* Swart

Recueil Trav. Bot. Néerl. 39: 191 (Swart 1942).

VERNACULAR NAMES. — Pa: ahuwahu, auwau • Ka: pilika • Wp: waluwai sī • Nt: tongi moni • Cr: bwa-lansan • Fr: encens.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2198*, $\text{dbh} = 48\text{ cm}$.

INVENTORY DATA (FG). — 40 trees in 16 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 45.6\text{ cm}$.

[188] *Protium decandrum* (Aubl.) Marchand

Adansonia [Baillon] 8: 51 (Marchand 1867). — *Icica decandra* Aubl., *Hist. Pl. Guiane* 1: 346 [Jun.-Dec. 1775] (Aublet 1775), “*Pentandra*” on plate. — *Amyris decandra* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (1): 336 [Mar. 1799] (Willdenow 1799). — *Elaphrium decandrum* (Aubl.) Spreng. ex D. Dietr., *Syn. Pl. [D. Dietrich]* 2: 1271 [1-20 Dec. 1840] (Dietrich 1840). — *Bursera decandra* (Aubl.) Baill., *Hist. Pl. [Baillon]* 5: 262, 297 [June 1874] (Baillon 1874).

Icica enneandra Aubl., *Hist. Pl. Guiane* 1: 345 [Jun.-Dec. 1775] (Aublet 1775). — *Amyris enneandra* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (1): 335 [Mar. 1799] (Willdenow 1799). — *Elaphrium*




FIG. 12. — Burseraceae: **A**, *Protium altissimum* (Aubl.) Marchand (D. Sabatier & J.-F. Molino 5203); **B**, *Protium morii* Daly (M.-F. Prévost & D. Sabatier 4952); **C**, *Protium trifoliolatum* Engl. (M.-F. Prévost et al. 4484); **D**, *Protium polybotryum* (Turcz.) Engl. (D. Sabatier & J.-F. Molino 5201). A, B, D, © D. Sabatier/IRD; C, © M.-F. Prévost/IRD.

enneandrum (Aubl.) Spreng. ex D. Dietr., *Syn. Pl. [D. Dietrich] 2:* 1271 [1–20 Dec. 1840] (Dietrich 1840). — *Tingulonga enneandra* (Aubl.) Kuntze, *Revis. Gen. Pl. 1:* 107 [5 Nov. 1891] (Kuntze 1891). — *Protium schomburgkianum* Engl., *Fl. Bras. [Martius] 12 (2):* 276 [1 Sep. 1874] (Engler 1874), *nom. illeg. superfl.* (based on *Icica enneandra*).

Protium orinocense Rusby, *Descr. S. Amer. Pl. 35* [20 Dec. 1920] (Rusby 1920).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ahuwahu, auwau, marinaiwa, sirasira, sirasra • Ka: alakuseli, šipyá, sipyó, šipyó • Nt: moni. • Cr: bwa-lasan • Fr: encens • Br: breu-preto, breu-vermelho.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Icica enneandra*: LINN, [LINN-HS 672.4](#)).

INVENTORY DATA (FG). — 204 trees in 49 plots; $F_{\max} = 2.9\%$; $dbh_{\text{inv}} = 48.2$ cm.

[189] *Protium demerarensis* Swart

Acta Bot. Neerl. 15: 46 (Swart 1966).

VERNACULAR NAMES. — Ka: alakuseli, šipyá, sipyó, šipyó • Nt: moni.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *S.A. Mori et al. 24011*.

INVENTORY DATA (FG). — 76 trees in 15 plots; $F_{\max} = 2.4\%$; $dbh_{\text{inv}} = 52$ cm.

[190] *Protium divaricatum* Engl. subsp. *fumarium* Daly
Brittonia 44 (3): 294 (Daly 1992).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Br: breu-cicantá.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *S.A. Mori & T.D. Pennington* 18034 (holo-, NY[00345683]; iso-, CAY, MG[MG126679], MO[MO2071216], P[P02440861]).

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 28.3$ cm.

[191] *Protium gallicum* Daly

Brittonia 50 (4): 520 (Daly 1998).

VERNACULAR NAMES. — Pa: araksim, marinaiwa, sirasira-seine • Wp: sipi, tulii, walakuseli • Nt: tingi moni • Cr: bwa-lansan-blan • Fr: encens.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *Service Forestier* 7073 (holo-, NY[00370252]; iso-, U[U0008214]).

INVENTORY DATA (FG). — 317 trees in 52 plots; $F_{\max} = 7.5\%$; $dbh_{\text{inv}} = 41.7$ cm.

[192] *Protium giganteum* Engl. var. *giganteum*

Fl. Bras. [Martius] 12 (2): 277 [1 Sep. 1874] (Engler 1874). — *Tingulonga gigantea* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

Icica acuminata Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 136 [24 Sep. 1813] (Poirier 1813), syn. nov. — *Amyris acuminata* DC., *Prodri. [A. P de Candolle]* 2: 78 [mid Nov. 1825] (Candolle 1825), nom. nud. pro. syn.

Protium duckei Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 432 (Huber 1909).

Protium cupreatum Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 11: 1 (Cuatrecasas 1961).

Protium pernervatum Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 11: 5 (Cuatrecasas 1961).

HERBARIUM DATA (FG). — No collection at CAY. Sel. exs.: *Unknown coll. s.n.* ("Herb. Poir.") (original material of *Icica acuminata*: P[P05308177]).

SIZE. — Brazil, Amazonas. *A. Vicentini* 1008 (MO), 20 m × 20 cm.

[193] *Protium giganteum* var. *crassifolium* (Engl.) Daly

Brittonia 44 (3): 296 (Daly 1992). — *Protium crassifolium* Engl., *Fl. Bras. [Martius]* 12 (2): 270 [1 Sep. 1874] (Engler 1874). — *Icica crassifolia* (Engl.) Rich. ex Engl., *Monogr. Phan. [A.DC. & C.DC.]* 4: 81 [Mar. 1883] (Engler 1883).

VERNACULAR NAMES. — Pa: ahuwahu, araksim-priyo, auwau.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *L.C. Richards s.n.* (holo-, B[not seen, photo F neg. 12568]; iso-, P[P02440864]).

INVENTORY DATA (FG). — 65 trees in 39 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 55$ cm.

[194] *Protium goudotianum* (Tul.) Byng & Christenb.

Global Fl. 4: 134 [9 Feb. 2018] (Byng & Christenhusz 2018). — *Icica goudotiana* Tul., *Ann. Sci. Nat., Bot. sér. 3, 7:* 372 (Tulasne 1847). — *Crepidospermum goudotianum* (Tul.) Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 14:* 300 (Triana & Planchon 1872).

Crepidospermum sprucei Hook.f., *Gen. Pl. [Bentham & Hooker f.]* 1 (1): 325 [7 Aug. 1862] (Hooker 1862).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori* 1885.

INVENTORY DATA (FG). — 10 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 35.5$ cm.

[195] *Protium guianense* (Aubl.) Marchand

Adansonia [Baillon] 8: 52 (Marchand 1867). — *Icica guianensis* Aubl., *Hist. Pl. Guiane* 1: 340 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate. — *Icica viridiflora* Lam., *Encycl. [J. Lamarck et al.]* 3 (1): 225 [19 Oct. 1789] (Lamarck 1789), nom. illeg. superfl. (based on *Icica guianensis*). — *Amyris guianensis* (Aubl.) Willd., *Sp. Pl., ed. 4* 2 (1): 335 [Mar. 1799] (Willdenow 1799), nom. illeg. hom., non Aubl. (Aublet 1775: 336). — *Elaphrium guianense* (Aubl.) Spreng. ex D. Dietr., *Syn. Pl. [D. Dietrich]* 2: 1271 [1-20 Dec. 1840] (Dietrich 1840), "gujanense". — *Tingulonga guianensis* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 107 [5 Nov. 1891] (Kuntze 1891).

Icica hostmannii Miq., *Stirp. Surinam. Select.*: 66 ["1850" publ. Mar. 1851] (Miquel 1851). — *Protium hostmannii* (Miq.) Engl., *Fl. Bras. [Martius]* 12 (2): 266 [1 Sep. 1874] (Engler 1874), "hostmanni". — *Tingulonga hostmannii* (Miq.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

Protium heptaphyllum var. *puberulum* Engl., *Fl. Bras. [Martius]* 12 (2): 263 [1 Sep. 1874] (Engler 1874).

VERNACULAR NAMES. — Ka: miyuluwa, šipyo • Wn: alakuhele, pakila siku • Nt: tingi moni • Cr: bwa-lansan • Fr: encens.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN[LINN-HS 672.5]).

INVENTORY DATA (FG). — 106 trees in 40 plots; $F_{\max} = 2.5\%$; $dbh_{\text{inv}} = 33.7$ cm.

[196] *Protium heptaphyllum* (Aubl.) Marchand

Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn 1873: 55 (Marchand 1873). — *Icica heptaphylla* Aubl., *Hist. Pl. Guiane* 1: 337 [Jun.-Dec. 1775] (Aublet 1775). — *Amyris ambrosiaca* L.f., *Suppl. Pl.*: 216 ["1781" publ. Apr. 1782] (Linnaeus 1782), nom. illeg. superfl. (based on *Icica heptaphylla*). — *Elaphrium heptaphyllum* (Aubl.) Spreng. ex D. Dietr., *Syn. Pl. [D. Dietrich]* 2: 1271 [1-20 Dec. 1840] (Dietrich 1840). — *Tingulonga heptaphylla* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 107 [5 Nov. 1891] (Kuntze 1891).

Icica surinamensis Miq., *Stirp. Surinam. Select.*: 65 ["1850" publ. Mar. 1851] (Miquel 1851). — *Protium heptaphyllum* var. *surinamense* (Miq.) Swart, *Recueil Trav. Bot. Néerl.* 39: 298 (Swart 1942).

Protium multiflorum Engl., *Fl. Bras. [Martius]* 12 (2): 273 [1 Sep. 1874] (Engler 1874). — *Tingulonga multiflora* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891). — *Protium heptaphyllum* var. *multiporum* (Engl.) Swart, *Recueil Trav. Bot. Néerl.* 39: 297 (Swart 1942).

Protium angustifolium Swart, *Recueil Trav. Bot. Néerl.* 39: 190 (Swart 1942).

Protium heptaphyllum var. *floribundum* Swart, *Recueil Trav. Bot. Néerl.* 39: 191 (Swart 1942).

Protium heptaphyllum var. *unifoliolatum* Swart, *Recueil Trav. Bot. Néerl.* 39: 191 (Swart 1942).

Protium octandrum Swart, *Recueil Trav. Bot. Néerl.* 39: 198 (Swart 1942).

Protium hostmannii var. *brasiliense* Swart, *Acta Bot. Neerl.* 15: 52 (Swart 1966).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: marinaiwa-kamwi • Ka: alakuseli, alakušini, aliwa'u, šipy, šipyo • Wp: sipi, walakuseli • Wn: aawa, alakuede, alakuheke, alakuhele • Nt: tingi moni • Cr: bwa-lansan-blanc • Fr: encens blanc • Br: almecegueira, breu-branco-verdadeiro.

HERBARIUM DATA (FG). — 74 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN, [LINN-HS 672.3](#)).

INVENTORY DATA (FG). — 15 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 30.9$ cm.

[197] *Protium inodorum* Daly

Brittonia 50 (4): 517 (Daly 1998).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15302* (holo-, NY[[0370253](#), [0370254](#)]; iso-, CAY[[CAY034925](#)], MO[[MO-1110102](#)]).

SIZE. — Up to 20 cm dbh (Daly 1998).

[198] *Protium insigne* (Triana & Planch.) Engl.

Monogr. Phan. [A.DC. & C.DC.] 4: 77 [Mar. 1883] (Engler 1883). — *Icica insignis* Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 14:* 299 (Triana & Planchon 1872). — *Icicopsis insignis* (Triana & Planch.) Engl., *Fl. Bras. [Martius]* 12 (2): 256 [1 Sep. 1874] (Engler 1874). — *Tingulonga insignis* (Triana & Planch.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Br: breu-sucuruba.

HERBARIUM DATA (FG). — 86 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5205*.

INVENTORY DATA (FG). — 95 trees in 42 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 66$ cm.

[199] *Protium melinonii* Engl.

Monogr. Phan. [A.DC. & C.DC.] 4: 68 [Mar. 1883] (Engler 1883), “*Melinonis*”.

NOTE. — The epithet “*melinonii*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland *et al.* 2018: Art. 60.8).

HERBARIUM DATA (FG). — Apparently endemic, presence attested by a single collection, *E.M. Mélinon 250*, 1842 (original material P[[P00781031](#)], L[[L0017689](#)]).

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[200] *Protium morii* Daly (Fig. 12B)

Brittonia 44 (3): 287 (Daly 1992).

VERNACULAR NAMES. — Pa: ahuwahu, auwau • Wp: ayawa sili, nemo áká, sipi sili, waluwai, waluwai sī.

HERBARIUM DATA (FG). — 89 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4952*.

INVENTORY DATA (FG). — 206 trees in 54 plots; $F_{\max} = 5.2\%$; $\text{dbh}_{\text{inv}} = 75$ cm.

[201] *Protium occultum* Daly

Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 7 (2): 253 (Daly 1991).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 22719*.

SIZE. — Up to 26.5 cm dbh (Daly 2002).

[202] *Protium opacum* Swart subsp. *rabelianum* Daly

Brittonia 44 (3): 291 (Daly 1992).

VERNACULAR NAMES. — Pa: ahuwahu, auwau, auwau-seine • Ka: aliwa'u • Te: waluwā'i • Wp: waluwai, waluwai pilā • Nt: lebi moni • Cr: bonm-kochon, bwa-lansan-gran-bwa • Fr: baume cochon • Br: almecegueira, breu-preto.

HERBARIUM DATA (FG). — 131 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2602*.

INVENTORY DATA (FG). — 787 trees in 173 plots; $F_{\max} = 4.4\%$; $\text{dbh}_{\text{inv}} = 70$ cm.

[203] *Protium pallidum* Cuatrec.

Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 11: 8 (Cuatrecasas 1961).

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *S.A. Mori et al. 22810*.

INVENTORY DATA (FG). — 17 trees in 5 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 40$ cm.

[204] *Protium pilosum* (Cuatrec.) Daly

Brittonia 44 (1): 24 (Daly 1989). — *Tetragastris pilosa* Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 11: 8 (Cuatrecasas 1961).

VERNACULAR NAMES. — Wp: waluwai sili • Br: breu-branco.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *A.H. Gentry et al. 63129*.

SIZE. — Up to 12 m tall (Daly 2002).

[205] *Protium plagiocarpium* Benoist

Bull. Soc. Bot. France 66 (8): 359 [“1919” publ. 1920] (Benoist 1920).

VERNACULAR NAMES. — Pa: marinaiwa-puvemna • Wp: sipi lá, sipi laánga, sipi sili, sipi towí'i.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *R. Benoist* 368 (original material P, not seen).

INVENTORY DATA (FG). — 33 trees in 27 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.2$ cm.

[206] *Protium polybotryum* (Turcz.) Engl.
(Fig. 12D)

Fl. Bras. [Martius] 12 (2): 278 [1 Sep. 1874] (Engler 1874). — *Icica polybotrya* Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 31 (1): 473 (Turczaninow 1858). — *Tingulonga polybotrya* (Turcz.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: marinaiwa-purubumna • Ka: pakila šipyoli, tamunen pilika.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 3209.

INVENTORY DATA (FG). — 133 trees in 12 plots; $F_{\max} = 8.1\%$; $dbh_{\text{inv}} = 54.5$ cm.

[207] *Protium rhoifolium* (Benth.) Byng & Christenh.

Global Fl. 4: 134 [9 Feb. 2018] (Byng & Christenhusz 2018). — *Hedwigia rhoifolia* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 17 (Bentham 1852). — *Crepidospermum rhoifolium* (Benth.) Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 14*: 300 (Triana & Planchon 1872). — *Hemicrepidospermum rhoifolium* (Benth.) Swart, *Recueil Trav. Bot. Néerl.* 39: 205 (Swart 1942).

VERNACULAR NAMES. — Pa: ara-kamwi • Ka: ayawalan, pakila šipyoli • Wp: ii, kulitu • Nt: moni.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier* 1001.

INVENTORY DATA (FG). — 36 trees in 24 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 49.5$ cm.

[208] *Protium robustum* (Swart) D.M.Porter

Ann. Missouri Bot. Gard. 56 (3): 476 [1969 publ. 17 July 1970] (Porter 1970). — *Protium neglectum* var. *robustum* Swart, *Recueil Trav. Bot. Néerl.* 39: 204 (Swart 1942).

VERNACULAR NAMES. — Ka: alukasi, pakila šipyoli • Te: waluwa'i • Wp: ayawa ilikili, nemo áká • Cr: bwa-lansan-rouj.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *D. Sabatier* & *J.-F. Molino* 5194.

INVENTORY DATA (FG). — 52 trees in 18 plots; $F_{\max} = 3.7\%$; $dbh_{\text{inv}} = 25.1$ cm.

[209] *Protium sagotianum* Marchand

Adansonia [Baillon] 8: 69 (Marchand 1867).

Bursera caudata Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 614 (Turczaninow 1863). — *Icicopsis caudata* (Turcz.) Engl., *Fl. Bras. [Martius]* 12 (2): 257 [1 Sep. 1874] (Engler 1874). — *Tingulonga caudata* (Turcz.) Kuntze, *Revis. Gen. Pl.* 1: 107 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ahuwahu, auwau • Ka: pakila šipyoli • Te: waluwa'i • Wp: nemo áká • Wn: aawa, awa • Nt: lebi moni, moni • Cr: bwa-lansan, chipa • Fr: encens • Br: breu-branco, breu-inhambu.

HERBARIUM DATA (FG). — 80 collections at CAY. Sel. exs.: *P.A. Sagot* 1191, Aug. 1858 (type P[P00781026]).

INVENTORY DATA (FG). — 251 trees in 71 plots; $F_{\max} = 3.3\%$; $dbh_{\text{inv}} = 67$ cm.

[210] *Protium spruceanum* (Benth.) Engl.

Fl. Bras. [Martius] 12 (2): 276 [1 Sep. 1874] (Engler 1874). — *Icica spruceana* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 16 (Bentham 1852). — *Tingulonga spruceana* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

Protium almecega Marchand, *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1873: 56 (Marchand 1873). — *Tingulonga almecega* (Marchand) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891), “almacega”.

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: sirasira, sirasra • Te: waluwa'i • Wp: sipi • Br: almecegueira-do-brejo.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-F. Molino* & *D. Sabatier* 2284.

INVENTORY DATA (FG). — 51 trees in 9 plots; $F_{\max} = 1.8\%$; $dbh_{\text{inv}} = 40.4$ cm.

[211] *Protium stevensonii* (Standl.) Daly

Brittonia 71 (3): 345 [epubl. 29 May 2019] (Daly 2019). — *Tetragastris stevensonii* Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 216 [24 Oct. 1929] (Standley 1929).

Hedwigia panamensis Engl., *Bot. Jahrb. Syst.* 1: 42 (Engler 1881). — *Tetragastris panamensis* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 107 [5 Nov. 1891] (Kuntze 1891). — *Protium picramnioides* Byng & Christenh., *Global Fl.* 4: 134 [9 Feb. 2018] (Byng & Christenhusz 2018), nom. illeg. superfl. (based on *Tetragastris panamensis*).

Tetragastris panamensis var. *hirtella* Swart, *Recueil Trav. Bot. Néerl.* 39: 207 (Swart 1942).

Tetragastris paraensis Cuatrec., *Bol. Mus. Paraense Emílio Goeldi*, N.S., Bot. 11: 7 (Cuatrecasas 1961).

Tetragastris panamensis var. *grandifolia* Swart, *Acta Bot. Neerl.* 15: 56 (Swart 1966).

VERNACULAR NAMES. — Pa: araksim, auwau-purubumna, kawap-kamwi, sirasira, sirasra • Ka: tamunen pilika • Wp: waluwai, waluwai pilá • Wn: aimalaje, simajali • Nt: gaan busi moni, weti sali • Br: barrotinho, breu-preto.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 4274*.

INVENTORY DATA (FG). — 192 trees in 88 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 80.5\text{ cm}$.

[212] *Protium strumosum* Daly

Brittonia 44 (3): 283 (Daly 1992).

HERBARIUM DATA (FG). — A single collection, *C. Feuillet 2263*.

SIZE. — Up to 15 m tall (Daly 1992).

[213] *Protium subserratum* (Engl.) Engl.

Monogr. Phan. [A.DC. & C.DC.] 4: 89 [Mar. 1883] (Engler 1883). — *Icicopsis subserrata* Engl., *Fl. Bras. [Martius]* 12 (2): 259 [1 Sep. 1874] (Engler 1874). — *Tingulonga subserrata* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: sirasira, sirasra.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2927*.

INVENTORY DATA (FG). — 167 trees in 82 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 50.5\text{ cm}$.

[214] *Protium surinamense* Byng & Christenh.

Global Fl. 4: 134 [9 Feb. 2018] (Byng & Christenhusz 2018). — *Hedwigia hostmannii* Engl., *Monogr. Phan. [A.DC. & C.DC.]* 4: 97 [Mar. 1883] (Engler 1883). — *Tetragastris hostmannii* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 106 [5 Nov. 1891] (Kuntze 1891).

NOTES. — Known only from the Guiana Shield. One of five species whose relative density in tree communities likely indicative of pre-Columbian archeological sites in French Guiana (Molino *et al.* 2021).

VERNACULAR NAMES. — Pa: ahuwahu, auwau • Ka: pilika, wipitano pilika • Wp: yaya'i, yaya'i sī • Cr: bois-yaya, yaya.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1354*.

INVENTORY DATA (FG). — 67 trees in 38 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 59\text{ cm}$.

[215] *Protium tenuifolium* (Engl.) Engl.
(Fig. 13A)

Monogr. Phan. [A.DC. & C.DC.] 4: 76 [Mar. 1883] (Engler 1883). — *Icicopsis tenuifolia* Engl., *Fl. Bras. [Martius]* 12 (2): 255 [1 Sep. 1874] (Engler 1874). — *Tingulonga tenuifolia* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

Protium neglectum Swart, *Recueil Trav. Bot. Néerl.* 39: 203 (Swart 1942).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ahuwahu, auwau, auwau-seinō • Wp: waluwai, waluwai pilá • Wn: awa.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4072*.

INVENTORY DATA (FG). — 195 trees in 95 plots; $F_{\max} = 1.9\%$; $\text{dbh}_{\text{inv}} = 72.5\text{ cm}$.

[216] *Protium trifoliolatum* Engl.
(Fig. 12C)

Fl. Bras. [Martius] 12 (2): 266 [1 Sep. 1874] (Engler 1874). — *Tingulonga trifoliolata* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891). — *Tetragastris trifoliolata* (Engl.) Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S.*, Bot. 11: 9 (Cuatrecasas 1961).

Protium martianum Engl., *Fl. Bras. [Martius]* 12 (2): 266 [1 Sep. 1874] (Engler 1874). — *Tingulonga martiana* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 108 [5 Nov. 1891] (Kuntze 1891).

Protium titubans J.F.Macbr., *Candollea* 5: 379 (Macbride 1934).

Protium pauciflorum Swart, *Recueil Trav. Bot. Néerl.* 39: 196 (Swart 1942).

Protium joannis Cuatrec., *Bol. Mus. Paraense Emílio Goeldi, N.S.*, Bot. 11: 2 (Cuatrecasas 1961).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: marinaiwa • Ka: pakila šipyoli • Te: waluwā'i • Wp: ii, sipi lá, sipi laānga, sipi sili, sipi towi'i • Wn: aimalaje • Nt: sali.

HERBARIUM DATA (FG). — 80 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1836*.

INVENTORY DATA (FG). — 133 trees in 37 plots; $F_{\max} = 6.6\%$; $\text{dbh}_{\text{inv}} = 32.1\text{ cm}$.

Genus *Trattinnickia* Willd.

[217] *Trattinnickia boliviiana* (Swart) Daly

Kew Bull. 54 (1): 130 (Daly 1999). — *Trattinnickia lawrancei* Standl. var. *boliviiana* Swart, *Acta Bot. Neerl.* 1: 249 (Swart 1952).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman 2106*.

SIZE. — Brazil, Acre. *D.C. Daly* 6912 (MO), 25 m × 35 cm.

[218] *Trattinnickia burserifolia* Mart.
(Fig. 13B)

Nova genera et species plantarum [Martius] 3 (2): 93 ["1829" publ. Jan.-Mar (?) 1831] (Martius 1831), "burserifolia".

Trattinnickia guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1188 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Trattinnickia schomburgkii Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1188 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Trattinnickia burserifolia var. *obtusa* Engl., *Fl. Bras. [Martius]* 12 (2): 284 [1 Sep. 1874] (Engler 1874).



FIG. 13. — Burseraceae: **A**, *Protium tenuifolium* (Engl.) Engl. (J.-F. Molino & D. Sabatier 2248); **B**, *Trattinnickia burserifolia* Mart. (M.-F. Prévost 4566). Calophyllaceae: **C**, *Calophyllum brasiliense* Cambess. (D. Sabatier & J.-F. Molino 5002); **D**, *Mahurea palustris* Aubl. (D. Sabatier & M.-F. Prévost 3021). A, © J.-F. Molino/IRD; B, © M.-F. Prévost/IRD; C, D, © D. Sabatier/IRD.

Trattinnickia burserifolia var. *quinquejuga* Engl., *Fl. Bras. [Martius]* 12 (2): 284 [1 Sep. 1874] (Engler 1874).

Trattinnickia subchoripetala Swart, *Recueil Trav. Bot. Néerl.* 39: 210 (Swart 1942).

Trattinnickia multiflora Cuatrec., *Webbia* 12 (2): 424 (Cuatrecasas 1957).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: sipi.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *M.-F. Prévost 4566*.

INVENTORY DATA (FG). — 33 trees in 9 plots; $F_{\max} = 2.2\%$; $dbh_{\text{inv}} = 46.8\text{ cm}$.

[219] *Trattinnickia demerarae* Sandwith

Bull. Misc. Inform. Kew 1931 (4): 185 [18 Apr. 1931] (Sandwith 1931).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: ahuwahu, auwau • Ka: apoto ayawa, ayawa, pakila sipyoli • Wp: ayawa, waluwai, waluwai pilá • Nt: gaan moni • Cr: bonm-kochon, bwa-lansan • Fr: baume cochon.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2278*.

INVENTORY DATA (FG). — 63 trees in 42 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 156.9\text{ cm}$.

[220] *Trattinnickia glaziovii* Swart

Recueil Trav. Bot. Néerl. 39: 208 (Swart 1942), “*Glaziovii*”.

Trattinnickia schwackeana Glaz., *Bull. Soc. Bot. France* 52 (Mém. 3a): 92 (Glaziou 1905), *nom. nud.*

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *B. Riéra 1092*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.4 cm.

[221] *Trattinnickia rhoifolia* Willd.

Sp. Pl., ed. 4 4 (2): 975 [Apr. 1806] (Willdenow 1806). — *Hedwigia rhoifolia* (Willd.) Baill., *Hist. Pl. [Baillon]* 5: 266 [June 1874] (Baillon 1874), *nom. illeg. hom., non* Benth. (Bentham 1852).

Trattinnickia ryanii Didr., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1857: 125 (Didrichsen 1857).

Trattinnickia rhoifolia var. *sprucei* Engl., *Fl. Bras. [Martius]* 12 (2): 283 [1 Sep. 1874] (Engler 1874).

Trattinnickia rhoifolia var. *wilddenowii* Engl., *Fl. Bras. [Martius]* 12 (2): 283 [1 Sep. 1874] (Engler 1874).

VERNACULAR NAMES. — Pa: araksim, ayao, ayau • Ka: aiawa, ayawa, pilika • Wp: ayawa • Wn: awa • Nt: gaan moni • Cr: bwa-lansan, bwa-yawa • Fr: encens • Br: breu-sucuruba.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2193*.

INVENTORY DATA (FG). — 10 trees in 9 plots; F_{max} < 1%; dbh_{inv} = 110 cm.

Family CALOPHYLLACEAE J.Agardh
Genus *Calophyllum* L.

[222] *Calophyllum brasiliense* Cambess.
(Fig. 13C)

Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 1 (8): 320 [28 June 1828] (Cambessèdes 1828).

Calophyllum longifolium Willd., *Mag. Neuesten Entdeck. Gesammten Naturk. Ges. Naturf. Freunde Berlin* 5: 80 (Willdenow 1811). — *Calophyllum brasiliense* subsp. *longifolium* (Willd.) Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 592 [Dec. 1893] (Vesque 1893).

Calophyllum lucidum Benth., *London J. Bot.* 2: 370 (Bentham 1843).

Calophyllum revolutum Rich. ex Engl., *Fl. Bras. [Martius]* 12 (1): 398 [1 Apr. 1888] (Engler 1888), *nom. nud. pro syn.*

Calophyllum brasiliense var. *elongatum* Engl., *Fl. Bras. [Martius]* 12 (1): 398 [1 Apr. 1888] (Engler 1888).

Calophyllum brasiliense subsp. *verum* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 591 [Dec. 1893] (Vesque 1893).

Calophyllum brasiliense var. *genuina* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 591 [Dec. 1893] (Vesque 1893), *nom. inval.* (Turland et al. 2018: Art. 24.3).

Calophyllum brasiliense var. *burchellii* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 592 [Dec. 1893] (Vesque 1893).

Calophyllum brasiliense var. *gardneri* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 592 [Dec. 1893] (Vesque 1893).

Calophyllum brasiliense var. *spruceanum* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 592 [Dec. 1893] (Vesque 1893), “*Spruceana*”.

Calophyllum brasiliense subsp. *mariae* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 593 [Dec. 1893] (Vesque 1893).

Calophyllum brasiliense subsp. *wrightii* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 594 [Dec. 1893] (Vesque 1893).

Calophyllum chiapense Standl., *Contr. U.S. Natl. Herb.* 20 (6): 192 [29 Aug. 1919] (Standley 1919).

Calophyllum rekoi Standl., *Contr. U.S. Natl. Herb.* 20 (6): 192 [29 Aug. 1919] (Standley 1919). — *Calophyllum brasiliense* var. *rekoi* (Standl.) Standl., *Trop. Woods* 30: 7 [1 June 1932] (Standley 1932).

Calophyllum antillanum Britton, *Sci. Surv. Porto Rico & Virgin Islands* 5: 584 (Britton 1924). — *Calophyllum brasiliense* var. *antillanum* (Britton) Standl., *Trop. Woods* 30: 7 [1 June 1932] (Standley 1932). — *Calophyllum jacquinii* Fawc. & Rendle, *Fl. Jamaica [Fawcett & Rendle]* 5: 200 (Fawcett & Rendle 1926), *nom. illeg. superfl.* (based on the same “*C. calaba* Jacq.” as *C. antillanum*).

Calophyllum ellipticum Rusby, *Mem. New York Bot. Gard.* 7: 303 (Rusby 1927).

Calophyllum piaroanum An.Castillo & C.Gil, *Ernstia*, ser. 2, 1 (1): 41 (Castillo & Gil 1991).

NOTES. — Hyperdominant in Amazonia (ter Steege et al. 2020). *C. antillanum* Britton and *C. jacquinii* Fawc. & Rendle were each intended as a new name for “*C. calaba* Jacq.” with reference to Jacquin (1763: 269), but this name does not exist. Jacquin (1763: 269) did not publish a new name, he only referred to *C. calaba* L. (Linnaeus 1753: 514). However, he did so by mistake since Linnaeus’ species is Asiatic, whereas the taxon he was treating is from the Caribbean Region.

VERNACULAR NAMES. — Pa: mpitit-wašiunē, pareine-āra • Ka: kula:la • Wp: yakale'i • Cr: bwa-kayman, manni-rouj • Br: cedro-do-pantano, guanambi, jacareúba.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *P. Grenand 637*.

INVENTORY DATA (FG). — 2 trees in 2 plots; F_{max} < 1%; dbh_{inv} = 57.6 cm.

Genus *Caraipa* Aubl.

[223] *Caraipa ampla* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 218 (Ducke 1922).

Caraipa variabilis Cambess. “var. β” (unnamed), *Mém. Mus. Hist. Nat.* 16: 417 (Cambessèdes 1828), *nom. inval.*

VERNACULAR NAMES. — Pa: wakukwa-tiranó-purubumna • Wp: wila pile lu • Cr: pajilé • Br: tamacoaré, tamaquaré.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (type of *Caraipa variabilis* “var. β”: P[P01901082]).

INVENTORY DATA (FG). — 1 tree, dbh = 28.6 cm.

[224] *Caraipa densifolia* Mart.

Nova genera et species plantarum [Martius] 1 (4): 105 ["1824" publ. Jan.-Mar. 1826] (Martius 1826).

Caraipa variabilis Cambess., *Mém. Mus. Hist. Nat.* 16: 416 (Cambessèdes 1828).

Caraipa fasciculata Cambess., *Mém. Mus. Hist. Nat.* 16: 417 (Cambessèdes 1828).

Caraipa laxiflora Benth., *London J. Bot.* 2: 364 (Bentham 1843).

Caraipa laurifolia Spruce ex Choisy, *Mém. Soc. Phys. Genève* 14 (1): 164 (Choisy 1855).

Caraipa myrciifolia Spruce ex Benth., *J. Proc. Linn. Soc., Bot.* 5: 62 ["1861" publ. Nov. 1860] (Bentham 1860), "myrciaefolia", *nom. nud. pro syn.*

Caraipa fasciculata var. *laxiflora* (Benth.) Wawra, *Fl. Bras. [Martius]* 12 (1): 323 [1 Apr. 1886] (Wawra 1886).

Caraipa excelsa Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 219 (Ducke 1922).

Caraipa melhemiana Paula, *Ci. & Cult.* 20 (2): 133 (Paula 1968).

NOTE. — Silva (1971: 25) provided two references for *C. melhemiana*, the first given above, and a second, *Ci. & Cult.* 21 (4): 170 (Paula 1969). None of them could be verified.

VERNACULAR NAMES. — Ka: sepeipyo, sepupi • Wp: wila pile lu • Nt: auman pisi • Cr: pajilé • Br: tamacoaré, tamaquaré.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *J.-F. Molino* 921.

INVENTORY DATA (FG). — 31 trees in 12 plots; $F_{\max} = 1.2\%$; $dbh_{inv} = 47.9$ cm.

[225] *Caraipa parvifolia* Aubl.

Hist. Pl. Guiane 1: 561 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Fr: manche-hache (*fide* Aublet 1775).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material P[P00106271]); *M.-F. Prévost* 3278, $dbh = 40$ cm.

[226] *Caraipa punctulata* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 216 (Ducke 1922).

Caraipa colombiana Ewan, *Nat. Hist. Misc.* 88: 4 (Ewan 1951).

Caraipa ferruginea Steyermark., *Fieldiana, Bot.* 28 (2): 384 (Steyermark 1952).

VERNACULAR NAMES. — Pa: wakukwa-tiranó-purubumna • Cr: pajilé • Br: tamacoaré, tamaquaré.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.-F. Molino et al.* 2083.

INVENTORY DATA (FG). — 31 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{inv} = 34.5$ cm.

[227] *Caraipa racemosa* Cambess.

Mém. Mus. Hist. Nat. 16: 415 (Cambessèdes 1828).

VERNACULAR NAMES. — Wp: tamukwálē áká, wila pile lu • Cr: pajilé • Br: tamacoaré, tamaquaré.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (holo-, P[P00106270]; iso-, M[M0274726]).

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 39$ cm.

[228] *Caraipa richardiana* Cambess.

Mém. Mus. Hist. Nat. 16: 414 (Cambessèdes 1828).

Caraipa richardiana var. *distorta* Wawra, *Fl. Bras. [Martius]* 12 (1): 318 [1 Apr. 1886] (Wawra 1886).

Caraipa psidiifolia Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 214 (Ducke 1922).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (holo-, P[P00106269]; iso-, P[P01901085, P01901086]).

SIZE. — Up to 15 m tall (Kubitzki 1978).

[229] *Caraipa* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3117.

INVENTORY DATA (FG). — 17 trees in 1 plot; $F_{\max} = 2.6\%$; $dbh_{inv} = 29.3$ cm.

[230] *Caraipa* sp. B

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 4936.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{inv} = 22.3$ cm.

Genus *Mahurea* Aubl.

[231] *Mahurea palustris* Aubl.
(Fig. 13D)

Hist. Pl. Guiane 1: 558 [Jun.-Dec. 1775] (Aublet 1775). — *Bonnetia palustris* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 814 [late Sep.-Nov. 1791] (Gmelin 1791).

Bonnetia meridionalis Sw., *Fl. Ind. Occid.* 2: 967 [late 1800] (Swartz 1800), *nom. illeg. superfl.* (based on *Mahurea palustris*).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: atit-kamwi.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000624839], P-LA[P00287390]).

INVENTORY DATA (FG). — 24 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{inv} = 75.9$ cm.

Family CANELLACEAE Mart.
Genus *Cinnamodendron* Endl.

[232] *Cinnamodendron tenuifolium* Uittien
(Fig. 14A)

Recueil Trav. Bot. Néerl. 22: 367 ["1925" publ. Jan. 1926] (Uittien 1926).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 1055*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.1 cm.

Family CANNABACEAE Martinov
Genus *Trema* Lour.

[233] *Trema integerrima* (Beurl.) Standl.
(Fig. 14B)

Contr. Arnold Arbor. 5: 55 (Standley 1933). — *Sponia integerrima* Beurl., *Kongl. Vetensk.-Akad. Handl.* 40: 144 ["1854" publ. 1856] (Beurling 1856).

Trema laxiflora Lundell, *Wrightia* 2 (2): 50 (Lundell 1960).

NOTE. — *Trema* is neuter, but treated as feminine in accordance with botanical tradition (Turland *et al.* 2018: Art. 62).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 6043*, dbh 65 cm.

[234] *Trema micrantha* (L.) Blume

Mus. Bot. 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856). — *Rhamnus micranthus* L., *Syst. Nat.*, ed. 10, 2: 937 [7 June 1759] (Linnaeus 1759). — *Celtis micrantha* (L.) Sw., *Prodri. [Swartz]* 53 [20 Jun.-29 July 1788] (Swartz 1788), "micranthus". — *Sponia micrantha* (L.) Decne. ex Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 333 [Dec. 1848] (Planchon 1848).

Celtis lima Sw., *Prodri. [Swartz]* 53 [20 Jun.-29 July 1788] (Swartz 1788). — *Sponia lima* (Sw.) Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 498 (Decaisne 1834), nom. inval. (genus name and epithet not associated). — *Trema lima* (Sw.) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Celtis mollis Humb. & Bonpl. ex Willd., *Sp. Pl.*, ed. 4 4 (2): 996 [Apr. 1806] (Willdenow 1806). — *Sponia mollis* (Humb. & Bonpl. ex Willd.) Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 498 (Decaisne 1834), nom. inval. (genus name and epithet not associated). — *Trema mollis* (Humb. & Bonpl. ex Willd.) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Celtis rugosa Willd., *Sp. Pl.*, ed. 4 4 (2): 996 [Apr. 1806] (Willdenow 1806).

Celtis canescens Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 28 [28 Apr. 1817] (Kunth 1817). — *Sponia canescens* (Kunth) Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 498 (Decaisne 1834), nom. inval. (genus name and epithet not associated). — *Celtis albicans* Willd. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 316 (Steudel 1840), nom. illeg. superfl. (based on *Sponia canescens*, thus indi-

rectly on *Celtis canescens*). — *Trema canescens* (Kunth) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Celtis riparia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 28 [28 Apr. 1817] (Kunth 1817). — *Sponia riparia* (Kunth) Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 498 (Decaisne 1834), nom. inval. (genus name and epithet not associated). — *Trema riparia* (Kunth) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Celtis macrophylla Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 30 [28 Apr. 1817] (Kunth 1817). — *Sponia macrophylla* (Kunth) Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 498 (Decaisne 1834), nom. inval. (genus name and epithet not associated). — *Trema macrophylla* (Kunth) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Celtis schiedeana Schlehd., *Linnaea* 7: 140 (Schlechtendal 1832). — *Sponia schiedeana* (Schlehd.) Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 335 [Dec. 1848] (Planchon 1848). — *Trema schiedeana* (Schlehd.) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Sponia peruviana Klotzsch, *Linnaea* 20: 536 [Oct. 1847] (Klotzsch 1847).

Celtis microcarpa Salzm. ex Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 333 [Dec. 1848] (Planchon 1848), nom. nud. pro syn.

Celtis rufescens Banks ex Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 334 [Dec. 1848] (Planchon 1848), nom. nud. pro syn. — *Sponia rufescens* Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 334 [Dec. 1848] (Planchon 1848). — *Trema rufescens* (Planch.) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Sponia chichilea Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 334 [Dec. 1848] (Planchon 1848). — *Celtis chichilea* Ruiz & Pav. ex Planch., *Ann. Sci. Nat., Bot. sér. 3*, 10: 334 [Dec. 1848] (Planchon 1848), nom. nud. pro syn. — *Trema chichilea* (Planch.) Blume, *Mus. Bot.* 2 (1-8): 58 ["1852" publ. Feb. 1856] (Blume 1856).

Sponia crassifolia Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 340 (Liebmann 1851).

Sponia grisea Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 340 (Liebmann 1851).

Trema melinonii Blume, *Mus. Bot.* 2 (1-8): 64 ["1852" publ. Feb. 1856], "Melinona".

Urtica alnifolia Bertero ex Griseb., *Fl. Brit. W.I. [Grisebach]* 709 [Oct. 1864] (Grisebach 1864), nom. nud. pro syn.

Celtis curiandiuba A.I.Gomes ex Planch., *Prodri. [A. P. de Candolle]* 17: 204 [16 Oct. 1873] (Planchon 1873), nom. nud.

Trema floridana Britton, *Fl. S.E. U.S. [Small]* 366, 1329 [22 July 1903] (Britton 1903). — *Trema micrantha* var. *floridana* (Britton) Standl. & Steyermark., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23 (2): 40 [14 Feb. 1944] (Standley & Steyermark 1944).

Trema micrantha var. *obtusata* Urb., *Symb. Antill. [Urban]* 4 (2): 195 [15 Feb. 1905] (Urban 1905).

Trema strigillosa Lundell, *Phytologia* 1 (10): 337 [27 Nov. 1939] (Lundell 1939). — *Trema micrantha* var. *strigillosa* (Lundell) Standl. & Steyermark., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23 (2): 40 [14 Feb. 1944] (Standley & Steyermark 1944).

NOTES. — Incipiently domesticated by pre-Columbian Amerindians (Levis *et al.* 2017). Decaisne (1834: 498) was the first to propose the transfer of *Celtis micrantha* (i.e. *Rhamnus micrantha*) into *Sponia*,



FIG. 14. — Canellaceae: **A**, *Cinnamodendron tenuifolium* Uittien (D. Sabatier 1055). Cannabaceae: **B**, *Trema integerrima* (Beurl.) Standl. (D. Sabatier et al. 6043). Capparaceae: **C**, *Capparis osmanthum* (Diels) Cornejo & Iltis (D. Sabatier 6330); **D**, *Neocalyptrocalyx leprieurii* (Briq.) Iltis (D. Sabatier 3048). © D. Sabatier/IRD.

but he did not associate the genus name with the specific epithet, making the combination invalid.

VERNACULAR NAMES. — Pa: maremi-etni • Ka: konolie, kunuliye • Te: kwanali'i • Wp: kulani'i • Wn: kumiklan • Nt: misobisobi • Cr: bwa-lòm, bwa-ranmié • Br: coatindiva, pau-de-polvora, periquitinha.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5276*.

SIZE. — Up to 20 m tall (Berg 1992).

Family CAPPARACEAE Juss.

NOTES

According to Cardinal-McTeague *et al.* (2016) and Tamboli *et al.* (2018), *Capparis* L. s.s. (*Capparis* sect. *Eucapparis* DC., including type) is an Old World taxon. All New World species (“NW capparoids”) form three or more distinct clades, among which two are clearly monophyletic (*Cynophalla* J.Presl and *Crateva* L.). The phylogenetic status of the remaining clades

is still unclear. If they are merged (the conclusion anticipated by Christenhusz *et al.* 2018), then the correct name would be *Morisonia* L. However, we think this too speculative and retain the genera *Calanthea*, *Capparidastrum*, *Neocalyptrocalyx* and *Presianthus*.

Genus *Calanthea* (DC.) Miers

[235] *Calanthea pulcherrima* (Jacq.) Miers

Proc. Roy. Hort. Soc. London 4: 161 (Miers 1864). — *Capparis pulcherrima* Jacq., *Enum. Syst. Pl.*: 24 [Aug.-Sep. 1760] (Jacquin 1760). — *Linnaeobreynia pulcherrima* (Jacq.) Hutch., *Gen. Fl. Pl.* 2: 310 (Hutchinson 1967), in adnot. — *Morisonia pulcherrima* (Jacq.) Christenh. & Byng, *Global Fl.* 4: 142 [9 Feb. 2018] (Christenhusz & Byng 2018).

Colicodendron pulchellum Seem., *Bot. Voy. Herald [Seemann]* 2: 78 [Aug. 1852] (Seemann 1852), *nom. illeg. superfl.* (based on *Capparis pulcherrima*).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *H. Richard & A. Mambe 622*.

INVENTORY DATA (FG). — 1 tree, dbh = 11.5 cm.

Genus *Capparidastrum* (DC.) Hutch.

[236] *Capparidastrum frondosum* (Jacq.) Cornejo & Iltis

Harvard Pap. Bot. 13 (2): 232 [17 Dec. 2008] (Cornejo & Iltis 2008). — *Capparis frondosa* Jacq., *Enum. Syst. Pl.*: 24 [Aug.-Sep. 1760] (Jacquin 1760). — *Pleuteron frondosa* (Jacq.) Raf., *Sylva Tellur.*: 109 (Rafinesque 1838). — *Uterveria frondosa* (Jacq.) Bertol., *Horti Bononi. Pl. Nov.* 2: 8 (Bertoloni 1839). — *Morisonia frondosa* (Jacq.) Christenh. & Byng, *Global Fl.* 4: 141 [9 Feb. 2018] (Christenhusz & Byng 2018).

Capparis elegans Mart., *Flora* 22 (1, Beibl.): 24 (Martius 1839). — *Capparidastrum elegans* (Mart.) Hutch., *Gen. Fl. Pl.* 2: 310 (Hutchinson 1967).

Capparis stenophylla Standl., *J. Wash. Acad. Sci.* 13: 437 (Standley 1923).

VERNACULAR NAMES. — Ka: abalaba, apalaba, mapalaba • Br: feijão-bravo-preto.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *G. Cremer 8448*.

SIZE. — Belize. *S.W. Brewer 8580* (MO), 10 m × 15 cm.

[237] *Capparidastrum osmanthum* (Diels) Cornejo & Iltis (Fig. 14C)

Harvard Pap. Bot. 13 (2): 234 [17 Dec. 2008] (Cornejo & Iltis 2008). — *Capparis osmantha* Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 332 [31 Mar. 1939] (Diels 1939).

Capparis guaguaensis Steyermark., *Fieldiana, Bot.* 28 (1): 238 (Steyermark 1951).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier et al. 6003*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 36.3 cm.

[238] *Capparidastrum solum* (J.F.Macbr.) Cornejo & Iltis

Harvard Pap. Bot. 11 (1): 17 (Cornejo & Iltis 2006), “*sola*”. — *Capparis sola* J.F.Macbr., *Candollea* 5: 359 (Macbride 1934). — *Morisonia sola* (J.F.Macbr.) Christenh. & Byng, *Global Fl.* 4: 142 [9 Feb. 2018] (Christenhusz & Byng 2018).

Capparis acutifolia J.F.Macbr., *Candollea* 5: 358 (Macbride 1934), *nom. illeg. hom.*, *non* Sweet (1830).

Capparis sola var. *longiracemosa* Dugand, *Caldasia* 7 (32): 110 (Dugand 1955).

VERNACULAR NAMES. — Wp: yawalū lemo.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Grenand 715*.

SIZE. — Ecuador, Napo. *J. Korning & K. Thomsen 47802* (MO), 13.7 m × 12.7 cm.

Genus *Crateva* L.

[239] *Crateva tapia* L.

Sp. Pl. 1: 444 [1 May 1753] (Linnaeus 1753).

Crateva gymandra L., *Sp. Pl.*, ed. 2, 1: 636 [Sep. 1762] (Linnaeus 1762).

Cleome arborea Schrad., *Gött. Gel. Anz.* 2: 707 [5 May 1821] (Schrader 1821).

Crateva acuminata DC., *Prodr. [A. P. de Candolle]* 1: 243 [mid Jan. 1824] (Candolle 1824), “*Crataeva*”.

Crateva radiatiflora DC., *Prodr. [A. P. de Candolle]* 1: 243 [mid Jan. 1824] (Candolle 1824).

Crateva tapiooides DC., *Prodr. [A. P. de Candolle]* 1: 243 [mid Jan. 1824] (Candolle 1824), “*Crataeva*”.

Colicodendron obliquifolium Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 27 (3): 328 (Turczaninow 1854).

Crateva benthamii Eichler, *Fl. Bras. [Martius]* 13 (1): 265 [1 Dec. 1865] (Eichler 1865), “*Crataeva*”.

Crateva benthamii var. *leptopetala* Eichler, *Fl. Bras. [Martius]* 13 (1): 265 [1 Dec. 1865] (Eichler 1865), “*Crataeva*”.

Crateva bahiana Ule, *Bot. Jahrb. Syst.* 42 (2-3): 202 [29 Dec. 1908] (Ule 1908), “*Crataeva*”.

Crateva coriacea Herzog, *Repert. Spec. Nov. Regni Veg.* 7: 52 (Herzog 1909), “*Crataeva*”.

Crateva glauca Lundell, *Bull. Torrey Bot. Club* 69 (5): 389 [May 1942] (Lundell 1942), “*Crataeva*”. — *Crateva tapia* var. *glauca* (Lundell) Standl. & Steyermark., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23 (2): 55 [14 Feb. 1944] (Standley & Steyermark 1944), “*Crataeva*”.

Capparis radiatiflora Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 16: 387 (Ruiz & Pavón 1959).

Capparis ternata Tafalla, *Fl. Huayaquilensis* 1: 161 (Tafalla 1989), *nom. nud.*

NOTE. — *Crateva radiatiflora* DC. was based on *Capparis radiatiflora* Ruiz & Pav., but the latter was unpublished at the time, therefore it cannot serve as a basionym.

VERNACULAR NAMES. — Ka: kuleku, kulelu • Br: catauré, trapiá.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: G.S. Perrottet 201 (type of *Crateva acuminata*: G-DC, [G00207195](#)).

SIZE. — Brazil, Acre. D.C. Daly 10186 (MO), 8 m × 18 cm.

Genus *Cynophalla* J.Presl

[[240](#)] *Cynophalla polyantha* (Triana & Planch.) Cornejo & Iltis

Harvard Pap. Bot. 13 (1): 118 [30 June 2008] (Cornejo & Iltis 2008). — *Capparis polyantha* Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4, 17:* 76 (Triana & Planchon 1862). — *Capparis flexuosa* (L.) L. subsp. *polyantha* (Triana & Planch.) Iltis, *Fl. Vénéz. Guayana* 4: 139 (Iltis 1998). — *Morisonia polyantha* (Triana & Planch.) Christenh. & Byng, *Global Fl.* 4: 142 [9 Feb. 2018] (Christenhusz & Byng 2018).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: [J.-F. Molino & D. Sabatier 3313](#).

INVENTORY DATA (FG). — 1 tree, dbh = 15.2 cm.

Genus *Neocalyptrocalyx* Hutch.

[[241](#)] *Neocalyptrocalyx leprieurii* (Briq.) Iltis
(Fig. 14D)

Harvard Pap. Bot. 13 (1): 110 [30 June 2008] (Iltis 2008). — *Capparis leprieurii* Briq., *Annuaire Conserv. Jard. Bot. Genève* 17: 393 (Briquet 1914).

VERNACULAR NAMES. — Pa: korosol-kamwi.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *F.M.R. Leprieur s.n.* (original material F[[V0053118F](#)], P[[P06795080](#)]).

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14\text{ cm}$.

[[242](#)] *Neocalyptrocalyx maroniensis* (Benoist) Cornejo & Iltis

Harvard Pap. Bot. 13 (1): 112 [30 June 2008] (Cornejo & Iltis 2008). — *Capparis maroniensis* Benoist, *Bull. Mus. Natl. Hist. Nat. 25:* 296 (Benoist 1919).

Capparis surinamensis J.C.Went, *Recueil Trav. Bot. Néerl.* 30: 164 [July 1933] (Went 1933). — *Neocalyptrocalyx surinamensis* (J.C.Went) Hutch., *Gen. Fl. Pl.* 2: 308 (Hutchinson 1967).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: á-wakaha • Ka: ombatapo • Wp: akusi luway.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *R. Benoist 339 bis* (lecto-, P[[P04022914](#)]), designated by Cornejo & Iltis [2008: 112]; isolecto-, P[[P04022915](#), [P04022916](#)].

INVENTORY DATA (FG). — 49 trees in 25 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 55.4\text{ cm}$.

[[243](#)] *Neocalyptrocalyx morii* Cornejo & Iltis

J. Bot. Res. Inst. Texas 2 (2): 807 (Cornejo & Iltis 2008).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *S.A. Mori et al. 24021* (holo-, NY[[01095755](#)] ; iso-, CAY[[CAY074676](#)]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.8\text{ cm}$.

Genus *Preslianthus* Iltis & Cornejo

[[244](#)] *Preslianthus pittieri* (Standl.) Iltis & Cornejo

Harvard Pap. Bot. 16 (1): 70 [June 2011] (Iltis & Cornejo 2011). — *Capparis pittieri* Standl., *J. Wash. Acad. Sci.* 17: 253 (Standley 1927). — *Morisonia pittieri* (Standl.) Christenh. & Byng, *Global Fl.* 4: 142 [9 Feb. 2018] (Christenhusz & Byng 2018).

Capparis schunkei J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (7):* 170 [24 Oct. 1929] (Macbride 1929).

Capparis crotonantha Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8):* 210 [24 Oct. 1929] (Standley 1929).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al. 25729*.

INVENTORY DATA (FG). — 1 tree, dbh = 11.3 cm.

Family CARICACEAE Dumort. Genus *Jacaratia* A.DC.

[[245](#)] *Jacaratia spinosa* (Aubl.) A.DC.
(Fig. 15A)

Prodr. [A. P. de Candolle] 15 (1): 419 [May 1864] (Candolle 1864). — *Carica spinosa* Aubl., *Hist. Pl. Guiane* 2: 908 [Jun.-Dec. 1775] (Aublet 1775). — *Papaya spinosa* (Aubl.) DC., *Encycl. J. Lamarck et al.* 5: 3 [9 Jan. 1804] (Candolle 1804).

Carica dodecaphylla Vell., *Fl. Flumin. Icon.* 10: t. 132 ["1827" publ. 29 Oct. 1831] (Vellozo 1831). — *Jacaratia dodecaphylla* (Vell.) A.DC., *Prodr. [A. P. de Candolle]* 15 (1): 420 [May 1864] (Candolle 1864).

Jacaratia actinophylla Pohl ex Solms, *Fl. Bras. [Martius]* 13 (3): 191 [15 Aug. 1889] (Solms 1889), *nom. nud.*

Jacaratia dodecaphylla f. *longiflora* Hassl., *Bull. Herb. Boissier* 7 (App. 1): 74 (Hassler 1899).

Jacaratia dodecaphylla var. *lucida* Hassl., *Bull. Herb. Boissier*, sér. 2, 2: 745 (Hassler 1902), “Eichleri”.

Jacaratia costaricensis I.M.Johnst., *Contr. Gray Herb.* 70: 79 (Johnston 1924).

VERNACULAR NAMES. — Pa: akabdat, kumak-kamwi • Wp: yalakasi • Nt: tukuman udu • Br: mamão-do-mato, mamão-rana.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM001008452]).

INVENTORY DATA (FG). — 14 trees in 12 plots; $F_{\max} < 1\%$; dbh_{inv} = 67.7 cm.

Family CARYOCARACEAE Voigt
Genus *Caryocar* F.Allam.

[246] *Caryocar glabrum* (Aubl.) Pers.

Syn. Pl. [Persoon] 2 (1): 84 [Nov. 1806] (Persoon 1806). — *Saouari glabra* Aubl., *Hist. Pl. Guiane* 1: 599 [Jun.-Dec. 1775] (Aublet 1775). — *Pekeea ternata* Poir., *Encycl. [J. Lamarck et al.] 5*: 148 [9 Jan. 1804] (Poiret 1804), nom. illeg. superfl. (based on *Saouari glabra*). — *Rhizobolus saouvari* Corrêa, *Ann. Mus. Natl. Hist. Nat.* 8: 394 (Corrêa 1806), nom. illeg. superfl. (based on *Saouari glabra*). — *Rhizobolus souari* Steud., *Nomencl. Bot. [Steudel]* 1: 688 (Steudel 1821), “*Souari*”, nom. illeg. superfl. (based on *Saouari glabra*). — *Rhizobolus glaber* (Aubl.) Corrêa ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 449 (Steudel 1841), nom. illeg. superfl. (based on *Caryocar glabrum*, i.e. on the type of *Saouari* Aubl.).

Caryocar toxiferum Barb.Rodr., *Vellozia* 1: 11 (Barbosa Rodrigues 1888).

Caryocar coccineum Pilg., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 127 [10 July 1927] (Pilger 1927).

Caryocar tessmannii Pilg., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 126 [10 July 1927] (Pilger 1927).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: sawa • Ka: alukumalilan, sawali • Wp: peke'a lá • Wn: inuuman • Nt: agugagi, maka sii • Cr: chawari-granbwa, chawari-rouj • Fr: chawari montagne • Br: piquíarana, tatajuba.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, LINN[LINN-HS 968.3] designated by Prance [Prance & Silva 1973: 40]).

INVENTORY DATA (FG). — 271 trees in 144 plots; $F_{\max} = 1.3\%$; dbh_{inv} = 143 cm.

[247] *Caryocar microcarpum* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 133 (Ducke 1925).

Caryocar glabrum var. *edule* Wittm., *Fl. Bras. [Martius]* 12 (1): 349 [1 Apr. 1886] (Wittmack 1886).

Caryocar glabrum var. *pilosum* Wittm., *Fl. Bras. [Martius]* 12 (1): 349 [1 Apr. 1886] (Wittmack 1886).

Caryocar riparium A.C.Sm., *Lloydia* 2 (3): 195 (Smith 1939).

NOTE. — *Caryocar glabrum* var. *edule* is *pro parte* in synonymy (excluding synonymy with *C. edule* Casar.).

VERNACULAR NAMES. — Pa: kwairu • Ka: alukumali • Te: alakubali • Wp: seweli • Wn: ulimé, wulimé • Nt: kasanyan, maka sii • Cr: chawari-dilo, chawari-rivié • Fr: chawari rivière • Br: piquíarana-do-igapó.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *M.-F. Prévost 1361*, 25 m × 80 cm.

INVENTORY DATA (FG). — 1 tree, dbh = 47.4 cm.

[248] *Caryocar villosum* (Aubl.) Pers.
(Fig. 15B)

Syn. Pl. [Persoon] 2 (1): 84 [Nov. 1806] (Persoon 1806), “*villosa*”. — *Saouari villosa* Aubl., *Hist. Pl. Guiane* 1: 601 [Jun.-Dec. 1775] (Aublet 1775). — *Pekeea villosa* (Aubl.) Poir., *Encycl. [J. Lamarck et al.] 5*: 148 [9 Jan. 1804] (Poiret 1804).

Pekeea butyrosa Aubl., *Hist. Pl. Guiane* 1: 594 [Jun.-Dec. 1775] (Aublet 1775), “*butyrosa*”, *pro parte quoad fructus et flores tantum*. — *Rhizobolus butyrosus* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 840 [late Sep.-Nov. 1791] (Gmelin 1791), *pro parte folia exclusa*. — *Caryocar butyrosum* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (2): 1243 [Dec. 1799] (Willdenow 1799).

Caryocar villosum var. *aesculifolium* Wittm., *Fl. Bras. [Martius]* 12 (1): 354 [1 Apr. 1886] (Wittmack 1886).

Caryocar villosum var. *macrophyllum* Wittm., *Fl. Bras. [Martius]* 12 (1): 354 [1 Apr. 1886] (Wittmack 1886).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: pikia • Ka: pekeea • Te: peke'a • Wp: peke'a • Wn: wake • Nt: agugagi, maka sii • Cr: bwa-dibè, pékéya • Fr: arbre à beurre, péquéia • Br: pequiá, piquíá.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material P-JU, P00671713)).

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 200 cm.

Family CELASTRACEAE R.Br.
Genus *Cheiloclinium* Miers

[249] *Cheiloclinium cognatum* (Miers) A.C.Sm.

Brittonia 3 (3): 529 (Smith 1940). — *Kippistia cognata* Miers, *Trans. Linn. Soc. London* 28 (2): 417 [post 17 May 1872] (Miers 1872). — *Salacia cognata* (Miers) Peyr., *Fl. Bras. [Martius]* 11 (1): 144 [1 Feb. 1878] (Peyritsch 1878).

Salacia cognata var. *egensis* Peyr., *Fl. Bras. [Martius]* 11 (1): 144 [1 Feb. 1878] (Peyritsch 1878), “*Egensis*”. — *Tontelea egensis* Poepp. ex Peyr., *Fl. Bras. [Martius]* 11 (1): 144 [1 Feb. 1878] (Peyritsch 1878), nom. nud. pro syn.

Salacia cognata var. *genuina* Peyr., *Fl. Bras. [Martius]* 11 (1): 144 [1 Feb. 1878] (Peyritsch 1878), nom. inval. (Turland *et al.* 2018: Art. 24.3). — *Cheiloclinium lineolatum* (A.C.Sm.) A.C.Sm., *Brittonia* 3 (3): 533 (Smith 1940).




Fig. 15. — Caricaceae: **A**, *Jacaratia spinosa* (Aubl.) A.DC. (M.-F. Prévost et al. 4478). Caryocaraceae: **B**, *Caryocar villosum* (Aubl.) Pers. (M.-F. Prévost & S. Gonzalez 5329). Celastraceae: **C**, *Maytenus* sp. B (D. Sabatier et al. 4786); **D**, *Zinowiewia aymardii* Steyermark. © D. Sabatier/IRD.

Elaeodendron macrophyllum Rusby, *Descr. S. Amer. Pl.* 24 [20 Dec. 1920] (Rusby 1920).

Salacia sphaerocarpa Rusby, *Descr. S. Amer. Pl.* 52 [20 Dec. 1920] (Rusby 1920).

Salacia lineolata A.C.Sm., *Bull. Torrey Bot. Club* 66 (4): 234 [Apr. 1939] (Smith 1939).

Cheiloclinium neglectum A.C.Sm., *Brittonia* 3 (3): 532 (Smith 1940).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020). Sometimes found as a liana.

VERNACULAR NAMES. — Pa: arak-priyu • Ka: aletipi • Wp: ipo átā, tatu mila • Br: pitombinha, uarutama.

HERBARIUM DATA (FG). — 92 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2152*.

INVENTORY DATA (FG). — 69 trees in 48 plots; $F_{\max} < 1\%$; dbh_{inv} = 23.6 cm.

Genus *Maytenus* Molina

[250] *Maytenus guyanensis* Klotzsch ex Reissek

Fl. Bras. [Martius] 11 (1): 19 [15 Feb. 1861] (Reissek 1861). — *Maytenus guianensis* Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1097 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. subnud.*

Maytenus guyanensis f. *crenulata* Steyermark., *Fieldiana, Bot.* 28 (2): 338 (Steyermark 1952), "guianensis".

NOTE. — Reissek cited "*M. guyanensis* Kl." [instead of "guianensis"], but as *M. guianensis* Klotzsch is not validly published, "guyanensis" should be considered as the original spelling.

VERNACULAR NAMES. — Ka: uwato epityi • Br: chichuasca.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2541*.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 57.6 cm.

[251] *Maytenus* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2850*.

INVENTORY DATA (FG). — 1 tree, dbh = 69.6 cm.

[252] *Maytenus* sp. B
(Fig. 15C)

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 4786*.

INVENTORY DATA (FG). — 11 trees in 8 plots; F_{max} < 1%; dbh_{inv} = 40.3 cm.

Genus *Monteverdia* A.Rich.[253] *Monteverdia floribunda* (Reissek) Biral

Syst. Bot. 42 (4): 689 [epubl. 18 Dec. 2017] (Biral 2017). — *Maytenus floribunda* Reissek, *Fl. Bras. [Martius]* 11 (1): 16 [15 Feb. 1861] (Reissek 1861).

Maytenus cardenasi Rusby, *Mem. New York Bot. Gard.* 7: 290 (Rusby 1927).

Maytenus erythrocarpa Rusby, *Mem. New York Bot. Gard.* 7: 290 (Rusby 1927).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1994*.

INVENTORY DATA (FG). — 14 trees in 12 plots; F_{max} < 1%; dbh_{inv} = 24.5 cm.

[254] *Monteverdia myrsinoides* (Reissek) Biral

Syst. Bot. 42 (4): 689 [epubl. 18 Dec. 2017] (Biral 2017). — *Maytenus myrsinoides* Reissek, *Fl. Bras. [Martius]* 11 (1): 21 [15 Feb. 1861] (Reissek 1861). — *Maytenus reissekii* Urb., *Festschr. Ascherson* 58 (Urban 1904), nom. illeg. superfl. (based on *Maytenus myrsinoides* Reissek, non Urb. [Urban 1904]).

NOTE. — By transferring *Myginda myrsinoides* Kunth [= *Crossopetalum myrsinoides* (Kunth) Kuntze (1891), “myrtinodes”] into *Maytenus* Molina, Urban (1904: 58) created a later homonym of *Maytenus myrsinoides* Reissek. Curiously, he treated the latter as if it was itself the later homonym, and created a new name for the corresponding taxon. As a result, *Maytenus reissekii* is superfluous.

VERNACULAR NAMES. — Ka: palakasana • Wp: kawe lá • Cr: bwabich • Br: apiranga, cafezinho.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 910*.

INVENTORY DATA (FG). — 20 trees in 16 plots; F_{max} < 1%; dbh_{inv} = 68.4 cm.

[255] *Monteverdia oblongata* (Reissek) Biral

Syst. Bot. 42 (4): 689 [epubl. 18 Dec. 2017] (Biral 2017). — *Maytenus oblongata* Reissek, *Fl. Bras. [Martius]* 11 (1): 25 [15 Feb. 1861] (Reissek 1861).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2086*.

INVENTORY DATA (FG). — 170 trees in 77 plots; F_{max} = 1.1%; dbh_{inv} = 86 cm.

Genus *Salacia* L.[256] *Salacia elliptica* (Mart.) G.Don

Gen. Hist. 1: 628 [early Aug. 1831] (Don 1831). — *Anthodon ellipticus* Mart., *Mant. 1 [Schultes]*: 348 (Martius 1822), “*Anthodus*”. — *Tonsella elliptica* (Mart.) Spreng., *Syst. Veg. [Sprengel]* 1: 178 [“1825” publ. late 1824] (Sprengel 1824). — *Raddia elliptica* (Mart.) Miers, *Trans. Linn. Soc. London* 28 (2): 395 [post 17 May 1872] (Miers 1872).

Anthodon glomeratus Mart., *Mant. 1 [Schultes]*: 348 (Martius 1822), “*Anthodus*”. — *Tonsella glomerata* (Mart.) Spreng., *Syst. Veg. [Sprengel]* 1: 177 [“1825” publ. late 1824] (Sprengel 1824). — *Salacia glomerata* (Mart.) G.Don, *Gen. Hist.* 1: 628 [early Aug. 1831] (Don 1831). — *Raddia glomerata* (Mart.) Miers, *Trans. Linn. Soc. London* 28 (2): 397 [post 17 May 1872] (Miers 1872).

Anthodon oblongifolius Mart., *Mant. 1 [Schultes]*: 348 (Martius 1822), “*Anthodus*”. — *Tonsella oblongifolia* (Mart.) Spreng., *Syst. Veg. [Sprengel]* 1: 178 [“1825” publ. late 1824] (Sprengel 1824). — *Salacia oblongifolia* (Mart.) G.Don, *Gen. Hist.* 1: 628 [early Aug. 1831] (Don 1831). — *Raddia oblongifolia* (Mart.) Miers, *Trans. Linn. Soc. London* 28 (2): 393 [post 17 May 1872] (Miers 1872). — *Salacia elliptica* var. *oblongifolia* (Mart.) Peyr., *Fl. Bras. [Martius]* 11 (1): 157 [1 Feb. 1878] (Peyritsch 1878).

Salacia obtusifolia Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 105 [10 Oct. 1829] (Cambessèdes 1829). — *Tontelea obtusifolia* (Cambess.) Endl. ex Walp., *Repert. Bot. Syst. [Walpers]* 1 (3): 400 [6-8 Nov. 1842] (Walpers 1842).

Salacia erythroxyloides Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 106 [10 Oct. 1829] (Cambessèdes 1829). — *Tontelea erythroxyloides* (Cambess.) Endl. ex Walp., *Repert. Bot. Syst. [Walpers]* 1 (3): 400 [6-8 Nov. 1842] (Walpers 1842).

Salacia obtusifolia var. *parviflora* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 106 [10 Oct. 1829] (Cambessèdes 1829).

Tontelea guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 2: 9 [12-15 July 1848] (Klotzsch 1848), nom. nud.

Salacia guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1183 [“1848” publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Raddia fasciculata Miers, *Trans. Linn. Soc. London* 28 (2): 391 [post 17 May 1872] (Miers 1872).

Raddia firmifolia Miers, *Trans. Linn. Soc. London* 28 (2): 391 [post 17 May 1872] (Miers 1872).

Raddia lacunosa Miers, *Trans. Linn. Soc. London* 28 (2): 392 [post 17 May 1872] (Miers 1872). — *Salacia lacunosa* (Miers) Peyr., *Fl. Bras. [Martius]* 11 (1): 158 [1 Feb. 1878] (Peyritsch 1878).

Raddia pachyphylla Miers, *Trans. Linn. Soc. London* 28 (2): 396 [post 17 May 1872] (Miers 1872). — *Salacia pachyphylla* (Miers) Peyr., *Fl. Bras. [Martius]* 11 (1): 158 [1 Feb. 1878] (Peyritsch 1878).

Salacia elliptica var. *apiculata* Peyr., *Fl. Bras. [Martius]* 11 (1): 157 [1 Feb. 1878] (Peyritsch 1878).

Salacia affinis Peyr., *Fl. Bras. [Martius]* 11 (1): 159 [1 Feb. 1878] (Peyritsch 1878).

Salacia amygdalina Peyr., *Fl. Bras. [Martius]* 11 (1): 159 [1 Feb. 1878] (Peyritsch 1878).

Salacia duckei A.C.Sm., *Contr. U.S. Natl. Herb.* 29 (8): 331 [23 Jan. 1950] (Smith 1950).

Salacia induita Rizzini, *Rodriguésia* 28 (41): 171 (Rizzini 1976).

NOTE. — Mostly as a liana, rarely a tree.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2028*.

INVENTORY DATA (FG). — 1 tree, dbh = 17 cm.

[257] *Salacia impressifolia* (Miers) A.C.Sm.

Bull. Torrey Bot. Club 66 (4): 247 [Apr. 1939] (Smith 1939). — *Raddia impressifolia* Miers, *Trans. Linn. Soc. London* 28 (2): 392 [post 17 May 1872] (Miers 1872).

Anthodon grandiflorus Benth., *Hooker's J. Bot. Kew Gard. Misc.* 4: 10 (Bentham 1852). — *Salacia grandiflora* (Benth.) Peyer., *Fl. Bras. [Martius]* 11 (1): 157 [1 Feb. 1878] (Peyritsch 1878), nom. illeg. hom., non Kurz (1872).

Salacia polyanthomaniaca Barb.Rodr., *Vellozia* 1: 16 (Barbosa Rodrigues 1888).

VERNACULAR NAMES. — Wp: kusi'u sē'ē, wali lapi'a • Br: tartaruginha, waimiratipi.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 876*.

SIZE. — Up to 12 m tall (Mennega & Hedin 1999).

[258] *Salacia juruana* Loes.

Verb. Bot. Vereins Prov. Brandenburg 48: 181 ["1906" publ. 8 Mar. 1907] (Loesener 1907).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15534*.

SIZE. — Ecuador, Pastaza. *A. Alvarez 2510* (MO), 12 m × 14.5 cm.

Genus *Zinowiewia* Turcz.

[259] *Zinowiewia aymardii* Steyermark.
(Fig. 15D)

Ann. Missouri Bot. Gard. 75 (3): 1064 [19 Oct. 1988] (Steyermark 1988).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5692*.

INVENTORY DATA (FG). — 4 trees in 2 plots; F_{max} < 1 %; dbh_{inv} = 82 cm.

Family CHRYSOBALANACEAE R.Br.
Genus *Acioa* Aubl.

[260] *Acioa guianensis* Aubl.
(Fig. 16A, B)

Hist. Pl. Guiane 2: 698 [Jun.-Dec. 1775] (Aublet 1775). — *Acia dulcis* Willd., *Sp. Pl.*, ed. 4 3 (1): 717 (Willdenow 1800), nom. illeg. superfl. (based on *Acioa guianensis*). — *Acioa dulcis* Steud., *Nomencl. Bot. [Steudel]* 1: 9 (Steudel 1821), nom. illeg. superfl. (based on *Acioa guianensis*). — *Moquilea aubletiana* Blume, *Mus. Bot. 2 (1-8)*: 92 ["1852" publ. Feb. 1856] (Blume 1856), nom. nov. (based on *Acioa guianensis*), non *Moquilea guianensis* Aubl. — *Licania aubletiana* (Blume) Lemée, *Fl. Guyane Franç.* 2: 23 (Lemée 1952).

VERNACULAR NAMES. — Pa: ašiu-aška • Nt: nengee udu • Cr: achiwa • Br: castanha-de-cotia.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000953533]).

INVENTORY DATA (FG). — 3 trees in 2 plots; F_{max} < 1 %; dbh_{inv} = 84.8 cm.

[261] *Acioa somnolens* Maguire

Brittonia 7: 272 [Dec. 1951] (Maguire 1951).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *E.M. Mélinon 230*, 1842 (holo-, P[P00741119]; iso-, P[P00741120]).

Genus *Chrysobalanus* L.

[262] *Chrysobalanus icaco* L.

Sp. Pl. 1: 513 [1 May 1753] (Linnaeus 1753).

Chrysobalanus purpureus Mill., *Gard. Dict.*, ed. 8, n. 2 [16 Apr. 1768] (Miller 1768).

Chrysobalanus pellocarpus G.Mey., *Prim. Fl. Esseq.* 193 [Nov. 1818] (Meyer 1818). — *Chrysobalanus icaco* var. *pellocarpus* (G.Mey.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 7 [17 Apr. 1867] (Hooker 1867). — *Chrysobalanus icaco* subsp. *pellocarpus* (G.Mey.) E.Murray, *Kalmia* 12: 19 (Murray 1982).

Chrysobalanus ellipticus Sol. ex Sabine, *Trans. Hort. Soc. London* 5: 453 (Sabine 1824). — *Chrysobalanus icaco* var. *ellipticus* (Sol. ex Sabine) Hook.f., *Fl. Bras. [Martius]* 14 (2): 7 [17 Apr. 1867] (Hooker 1867). — *Chrysobalanus icaco* subsp. *ellipticus* (Sol. ex Sabine) Souza, *Ann. Univ. Abidjan*, C. 15: 101 (Souza 1980).

Chrysobalanus luteus Sabine, *Trans. Hort. Soc. London* 5: 453 (Sabine 1824). — *Chrysobalanus icaco* var. *luteus* (Sabine) Souza, *Ann. Univ. Abidjan*, C. 15: 104 (Souza 1980).

Chrysobalanus orbicularis Schumach., *Beskr. Guin. Pl.* 232 (Schumacher 1827). — *Chrysobalanus icaco* subsp. *orbicularis* (Schumach.) Souza, *Ann. Univ. Abidjan*, C. 15: 101 (Souza 1980).

Chrysobalanus guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1024 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Chrysobalanus savannarum Britton, *Bull. Torrey Bot. Club* 48 (12): 331 ["1921" publ. 1922] (Britton 1922).

Chrysobalanus interior Small, *Man. S.E. Fl.*: 645 (Small 1933).

Chrysobalanus icaco var. *genuinus* Stehlé, M. Stehlé & Quentin, *Fl. Guadeloupe [Stehlé]* 2 (3): 48 (Stehlé et al. 1949), *nom. inval.* (Turland et al. 2018: Art. 24.3).

Chrysobalanus icaco f. *albus* G. Klotz, *Wiss. Z. Friedrich-Schiller-Univ. Jena, Math.-Naturwiss. Reihe (Beitr. Phytotax.* 6) 29: 461 (Klotz 1980).

Chrysobalanus icaco var. *macrocarpus* Souza, *Ann. Univ. Abidjan*, C. 15: 104 (Souza 1980), *nom. inval.* (type not properly cited).

Chrysobalanus icaco var. *roseus* Souza, *Ann. Univ. Abidjan*, C. 15: 101 (Souza 1980), *nom. inval.* (no type designated).

NOTES. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017). In French Guiana, restricted to coastal sandridges.

VERNACULAR NAMES. — Pa: warahu • Ka: konoto epi • Cr: prin-lans, prin-zikak • Br: guajuru.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *D. Sabatier* 4896, dbh = 12 cm.

Genus *Couepia* Aubl.

[263] *Couepia bracteosa* Benth.

J. Bot. [Hooker] 2: 215 (Bentham 1840). — *Moquilea bracteosa* (Benth.) Walp., *Repert. Bot. Syst. [Walpers]* 2 (1): 7 [5-9 Apr. 1843] (Walpers 1843).

Couepia bracteosa var. *grandifolia* Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 597 (Benoist 1923).

Couepia bracteosa var. *minor* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 117 (Ducke 1930).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017).

VERNACULAR NAMES. — Ka: kuwepi, kwepi • Br: marirana, pajurá-de-racha, pajurá-verdadeiro.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *E.M. Mélinon* s.n. (original material of *Couepia bracteosa* var. *grandifolia*; P[P04848416, P04848419, P04848420]).

INVENTORY DATA (FG). — 107 trees in 46 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 48.1$ cm.

[264] *Couepia caryophylloides* Benoist

Bull. Mus. Natl. Hist. Nat. 28: 253 (Benoist 1922).

VERNACULAR NAMES. — Pa: bukutru-gateau-duwē • Wp: wila yis̄ • Nt: kwebi kookoo • Cr: bwa-golèt, golèt-nwé, grigri-rouj • Fr: galette noire • Br: pajurá-verdadeiro.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *G. Wachenheim* sér. 3, 0.11 (holo-, P[P00741076]; iso-, P[P00741077, P00741078]).

INVENTORY DATA (FG). — 205 trees in 78 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 131$ cm.

[265] *Couepia excelsa* Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 116 (Ducke 1930).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wn: kula waju.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M. Fleury* 1348.

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 24.2$ cm.

[266] *Couepia exflexa* Fanshawe & Maguire

Bull. Torrey Bot. Club 75 (4): 376 [Jul.-Aug. 1948] (Fanshawe & Maguire 1948).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *S.A. Mori* et al. 23464.

SIZE. — Guyana. *P. Mutchnick* et al. 117 (MO), 15 m.

[267] *Couepia guianensis* Aubl. subsp. *guianensis*

Hist. Pl. Guiane 1: 519 [Jun.-Dec. 1775] (Aublet 1775). — *Acia amara* Willd., *Sp. Pl.*, ed. 4 3 (1): 717 (Willdenow 1800), *nom. illeg. superfl.* (based on *Couepia guianensis*). — *Acioa amara* Steud., *Nomencr. Bot. [Steudel]* 1: 9 (Steudel 1821), *nom. illeg. superfl.* (based on *Couepia guianensis*). — *Moquilea couepia* Steud., *Nomencr. Bot. [Steudel]*, ed. 2, 2: 159 (Steudel 1841), *nom. illeg. superfl.* (based on *Couepia guianensis*).

Couepia leptostachya Benth. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 44 [17 Apr. 1867] (Hooker 1867).

Couepia versicolor Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 596 (Benoist 1923).

Couepia surinamensis Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 390 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: kwep-puvemna • Ka: kuwepi, kwepi • Te: alama'i • Nt: kwebi kookoo • Cr: bwa-golèt, golèt-koumaté • Fr: galette • Br: pajurá-verdadeiro.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (lecto-, BM[BM000065023] designated by Prance [1981: 349]).

INVENTORY DATA (FG). — 207 trees in 97 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 72.1$ cm.

[268] *Couepia guianensis* subsp. *glandulosa* (Miq.) Prance

Brittonia 33 (3): 350 (Prance 1981). — *Couepia glandulosa* Miq., *Stirp. Surinam. Select.*: 28 ["1850" publ. Mar. 1851] (Miquel 1851). — *Moquilea glandulosa* (Miq.) Walp., *Ann. Bot. Syst. [Walpers]* 2 (3): 463 [25-28 Feb. 1852] (Walpers 1852).

Couepia myrtifolia Benth. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 44 [17 Apr. 1867] (Hooker 1867).




FIG. 16. — Chrysobalanaceae: **A, B**, *Acioa guianensis* Aubl.; **C**, *Couepia joaquiniae* Prance (*D. Sabatier & J.-F. Molino* 5341); **D**, *Gaulettia elata* (Ducke) Sothers & Prance (*D. Sabatier & M.-F. Prévost* 4932); **E**, *Hirtella davisii* Sandwith (*D. Sabatier & J.-F. Molino* 5708). **A, B**, © M.-F. Prévost/IRD; **C-E**, © D. Sabatier/IRD.

Couepia thrysiflora Hook.f., *Fl. Bras. [Martius]* 14 (2): 45 [17 Apr. 1867] (Hooker 1867).

VERNACULAR NAMES. — Wp: kumalaki sili, pali'i sili • Cr: bwagolèt, golèt-koumaté • Br: pajurá-verdadeiro.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 2027*.

SIZE. — Brazil, Pará. *G.T. Prance 1701* (MO), 12 m × 15 cm.

[269] *Couepia habrantha* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (3): 249 (Standley 1937).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *S.A. Mori et al. 25554*.

INVENTORY DATA (FG). — 45 trees in 26 plots; $F_{\max} = 2\%$; $dbh_{inv} = 46.2$ cm.

[270] *Couepia joaquiniae* Prance
(Fig. 16C)

Kew Bull. 46 (1): 107 (Prance 1991).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: bukutru-ateu, bukutru-gateau.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5341.

INVENTORY DATA (FG). — 27 trees in 13 plots; $F_{\max} = 2.2\%$; $dbh_{inv} = 41.4$ cm.

[271] *Couepia magnoliifolia* Benth. ex Hook.f.

Fl. Bras. [Martius] 14 (2): 43 [17 Apr. 1867] (Hooker 1867).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Loubry 1698*.

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{inv} = 31.2$ cm.

[272] *Couepia martinii* Prance

Fl. Neotrop. Monogr. 9: 238 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — Apparently endemic, presence attested by a single collection, *J. Martin s.n.* (holo-, NY[00428114];

iso-, BM[BM000602473, BM000602474, BM000602475], F[V0053868F], K[K000220967, K000220968]).

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[273] *Couepia obovata* Ducke

Arch. Inst. Biol. Veg. 2 (1): 35 [Sep. 1935] (Ducke 1935).

VERNACULAR NAMES. — Br: macucu-murici, pajurazinho, uchirana.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3125*.

INVENTORY DATA (FG). — 20 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{inv} = 61.1$ cm.

[274] *Couepia rankiniae* Prance

Kew Bull. 47 (4): 640 (Prance 1992), “rankiniae”, “rankinii” on figure.

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 25174*.

SIZE. — Up to 20 m tall (Prance 1992).

[275] *Couepia* sp. A

NOTE. — This species resembles *C. cognata* (Steud.) Fritsch.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5634*.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 29.8$ cm.

Genus *Exelloidendron* Prance

[276] *Exelloidendron barbatum* (Ducke) Prance

Fl. Neotrop. Monogr. 9: 199 [8 June 1972] (Prance 1972). — *Parinari barbata* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 45 (Ducke 1922), “*Parinarium barbatum*”.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 6121*.

INVENTORY DATA (FG). — 15 trees in 6 plots; $F_{\max} = 1.2\%$; $dbh_{inv} = 26.4$ cm.

Genus *Gaulettia* Sothers & Prance

[277] *Gaulettia canomensis* (Mart.) Sothers & Prance

Phytotaxa 172 (3): 184 [18 June 2014] (Sothers & Prance 2014). — *Hirtella canomensis* (Mart.) Spreng., *Syst. Veg. [Sprengel]* 4 (2): 341 [Jan.-June 1827] (Sprengel 1827). — *Moquilea canomensis* Mart., *Nova genera et species plantarum [Martius]* 2 (2): 80 [Jan.-June 1827] (Martius 1827). — *Couepia canomensis* (Mart.) Benth. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 42 [17 Apr. 1867] (Hooker 1867).

Parinari pilosa Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (3): 259 (Standley 1937).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & M.-F. Prévost 3215*.

INVENTORY DATA (FG). — 1 tree, dbh = 17.8 cm.

[278] *Gaulettia elata* (Ducke) Sothers & Prance
(Fig. 16D)

Phytotaxa 172 (3): 186 [18 June 2014] (Sothers & Prance 2014). — *Couepia elata* Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 35 (Ducke 1935).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & S. Gonzalez 5370*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 70$ cm.

[279] *Gaulettia parillo* (DC.) Sothers & Prance

Phytotaxa 172 (3): 188 [18 June 2014] (Sothers & Prance 2014). — *Couepia parillo* DC., *Prod. [A. P. de Candolle]* 2: 526 [mid Nov. 1825] (Candolle 1825). — *Moquilea parillo* (DC.) Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 159 (Steudel 1841), “*parilla*”.

Couepia pauciflora Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 372 (Huber 1909).

Couepia villosa Fanshawe & Maguire, *Bull. Torrey Bot. Club* 75 (4): 378 [Jul.-Aug. 1948] (Fanshawe & Maguire 1948).

VERNACULAR NAMES. — Pa: á-kiavū, á-kiavunó • Ka: palipyo, polipyo, polipyoli • Wp: pali'i lá • Cr: golèt-ti-fèy.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (“Herb. Ventenat”) (type G[G00177583]).

INVENTORY DATA (FG). — 26 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{inv} = 42$ cm.

Genus *Hirtella* L.

[280] *Hirtella araguariensis* Prance

Fl. Neotrop. Monogr. 9: 278 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *C. Delnatte 1445*.

INVENTORY DATA (FG). — 1 tree, dbh = 17.5 cm.

[281] *Hirtella bicornis* Mart. & Zucc. var. *bicornis*

Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. 1: 377 (Martius & Zuccarini 1832).

Hirtella praealta Sagot, *Ann. Sci. Nat., Bot. sér.* 6, 15: 304 (Sagot 1883).

VERNACULAR NAMES. — Pa: bukutru-gateu-duwē, bukutru-gateu-seinō-puvemna • Ka: kumboeta ši, malimiapo • Te: wila pulua • Wp: tulipalapo, weyu'i • Nt: atila kookoo, kookoo • Cr: golèt-ti-fey • Fr: galette à petites feuilles • Br: ajuru, caraipé, caripé.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P.A. Sagot* 792, Aug. 1855 (holotype of *Hirtella praealta*: P[P00741050]; iso-, B[not seen, photo F neg. 3351], BR[BR000006969158], F[V0044063F], G[G00359638], GOET[GOET000875], K[K000220875, K000220876], L[L.1890947], MPU[MPU024071], NY[00335297, 00428266], P[P00741051, P00741052], S[S-R-8150], U[U0001103, U0001104]).

INVENTORY DATA (FG). — 292 trees in 94 plots; $F_{\max} = 5.1\%$; $\text{dbh}_{\text{inv}} = 48.1\text{ cm}$.

[282] *Hirtella bicornis* var. *pubescens* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 269 (Ducke 1922).

Hirtella caudata Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 389 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

VERNACULAR NAMES. — Wp: tulipalapo, weyu'i • Cr: golèt-ti-fèy • Br: ajuru, caraipé, caripé.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2744.

INVENTORY DATA (FG). — 23 trees in 15 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 36.5\text{ cm}$.

[283] *Hirtella ciliata* Mart. & Zucc.

Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. 1: 378 (Martius & Zuccarini 1832).

Grangeria brasiliensis Hoffmanns. ex Mart. & Zucc., *Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.* 1: 378 (Martius & Zuccarini 1832), nom. nud. pro syn.

Hirtella rubra Benth., *J. Bot. [Hooker]* 2: 217 (Bentham 1840).

Hirtella rotundata Pilg., *Notizbl. Königl. Bot. Gart. Berlin* 6: 140 [20 Mar. 1914] (Pilger 1914).

VERNACULAR NAMES. — Ka: kumboeta ši, kumboeta sipyoti, kupešinilan • Br: ajururana, chorão.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-P. Lescure* 708.

SIZE. — Up to 12 m tall (Prance 1986).

[284] *Hirtella davisi* Sandwith
(Fig. 16E)

Bull. Misc. Inform. Kew 1935 (3): 125 [20 May 1935] (Sandwith 1935).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5708.

INVENTORY DATA (FG). — 34 trees in 15 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 55.4\text{ cm}$.

[285] *Hirtella glandulostipula* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 265 (Ducke 1922).

HERBARIUM DATA (FG). — A single collection, *C. Feuillet et al.* 10083 (MO).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 15.8\text{ cm}$.

[286] *Hirtella glandulosa* Spreng.

Neue Entdeck. Pflanzenk. 1: 303 (Sprengel 1820).

Hirtella damaziana Beauverd, *Bull. Herb. Boissier*, sér. 2, 7: 706 (Beauverd 1907).

Hirtella hookeri Pilg., *Notizbl. Königl. Bot. Gart. Berlin* 6: 139 [20 Mar. 1914] (Pilger 1914).

Hirtella velutina Pilg., *Notizbl. Königl. Bot. Gart. Berlin* 6: 141 [20 Mar. 1914] (Pilger 1914).

Hirtella wachenheimii Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 595 (Benoist 1923).

VERNACULAR NAMES. — Pa: bukutru-gateau-seinō • Wp: iwikwisiñ, mukwisiñ • Nt: kookoo, nengee udu • Cr: golèt-blanc • Fr: galette blanche • Br: vermelhão.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *G. Wachenheim* 372 (holotype of *Hirtella wachenheimii*: P[P00741073]; iso-, K[K000220855], P[P00741074, P00741075]).

INVENTORY DATA (FG). — 117 trees in 73 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 59.7\text{ cm}$.

[287] *Hirtella hispidula* Miq.

Stirp. Surinam. Select.: 27 ["1850" publ. Mar. 1851] (Miquel 1851).

Hirtella egensis Fritsch, *Ann. K. K. Naturhist. Hofmus.* 5: 10 (Fritsch 1890), "Egensis".

Hirtella subsetosa Fanshawe & Maguire, *Bull. Torrey Bot. Club* 75 (4): 381 [Jul.-Aug. 1948] (Fanshawe & Maguire 1948).

VERNACULAR NAMES. — Pa: ahavuiye • Wn: kuluné ki.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *J.-F. Molino et al.* 1661.

INVENTORY DATA (FG). — 37 trees in 17 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 21.2\text{ cm}$.

[288] *Hirtella macrophylla* Benth. ex Hook.f.

Fl. Bras. [Martius] 14 (2): 36 [17 Apr. 1867] (Hooker 1867).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *P. Grenand et al.* 3380.

SIZE. — Brazil, Mato Grosso. *C.A. Cid Ferreira* 5668 (MO), 13 m × 12 cm.

[289] *Hirtella macrosepala* Sandwith

Bull. Misc. Inform. Kew 1939 (10): 549 ["1939" publ. 6 Jan. 1940] (Sandwith 1940).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *Service Forestier 7237*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.3$ cm.

[290] *Hirtella margae* Prance

Proc. Kon. Ned. Akad. Wetensch. C 89 (1): 111 (Prance 1986).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2994*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.1$ cm.

[291] *Hirtella obidensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 267 (Ducke 1922).

Hirtella lanceolata Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 388 (Kleinhoonte 1926).

Hirtella coppenamensis Kleinhoonte, *Recueil Trav. Bot. Néerl.* 30: 180 (Kleinhoonte 1933).

VERNACULAR NAMES. — Ka: kwepilan.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5079*.

INVENTORY DATA (FG). — 10 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22$ cm.

[292] *Hirtella paniculata* Sw.
(Fig. 17A)

Prodr. [Swartz] 51 [20 Jun.-29 July 1788] (Swartz 1788). — *Hirtella hirsuta* Lam., *Tabl. Encycl.* 2[4 (2)]: 114 [6 Nov. 1797] (Lamarck 1797), nom. illeg. superfl. (based on *Hirtella paniculata*). — *Hirtella aggregata* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 53 [24 Sep. 1813] (Poirier 1813), nom. illeg. superfl. (based on *Hirtella paniculata*).

Hirtella indecora Schott ex Spreng., *Syst. Veg. [Sprengel]* 4 (2): 404 [Jan.-June 1827] (Sprengel 1827), nom. nud. pro syn.

Hirtella bracteosa Steud., *Flora* 26 (45): 761 [17 Dec. 1843] (Steu-del 1843).

VERNACULAR NAMES. — Pa: ahavuiye, ahavuiye-seine • Ka: alilime'i, kumboeta ši, kumboeta sipyoti, kupešinilan • Wp: yanu'i pilá, yanu'i yowa.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (original material BM[[BM000617503](#)]); *M.-F. Prévost 3856*, $dbh = 12$ cm.

[293] *Hirtella racemosa* Lam. var. *racemosa*

Encycl. [J. Lamarck et al.] 3 (1): 133 [19 Oct. 1789] (Lamarck 1789).

Tachibota guianensis Aubl., *Hist. Pl. Guiane* 1: 287 [Jun.-Dec. 1775] (Aublet 1775). — *Salmasia guianensis* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 500 [late Sep.-Nov. 1791] (Gmelin 1791), nom. illeg. superfl. (based on *Tachibota guianensis*). — *Salmasia racemosa* Willd., *Sp. Pl.*, ed. 4 1 (2): 1502 [July 1798] (Willdenow 1798), nom. illeg. superfl. (based on *Tachibota guianensis*).

Hirtella nemorosa Hoffmanns. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 274 [Dec. 1819] (Roemer & Schultes 1819).

Hirtella scandens Hoffmanns. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 274 [Dec. 1819] (Roemer & Schultes 1819).

Hirtella violacea Steud., *Flora* 26 (45): 761 [17 Dec. 1843] (Steu-del 1843).

Hirtella racemosa var. *metallica* Cuatrec., *Fieldiana, Bot.* 27 (1): 60 (Cuatrecasas 1950).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

Tachibota guianensis Aublet (1775: 287) predates *H. racemosa* Lamarck (1789: 133), but its epithet is no longer transferable because the name is occupied by *Hirtella guyanensis* (Fritsch) Sandwith (1931) (Sandwith 1963).

VERNACULAR NAMES. — Ka: kumboeta ši, kumboeta sipyoti, kupešinilan, piliwa epulu, piliwa pi • Wp: yanu'i, yanu'i pilá • Wn: kulumnë, kulumnë epí, kulunë epku • Nt: asiantefi • Cr: ti-golèt, ti-golèt-rouj • Br: caripérana-branca.

HERBARIUM DATA (FG). — 74 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lectotype of *Tachibota guianensis* [P-JJR, [P00778468](#)] designated by Lanjouw & Uittien [1940: 159]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 18.1$ cm.

[294] *Hirtella racemosa* var. *hexandra* (Willd.) Prance

Fl. Neotrop. Monogr. 9: 328 [8 June 1972] (Prance 1972). — *Hirtella hexandra* Willd., *Syst. Veg. [Roemer & Schultes]* 5: 274 [Dec. 1819] (Willdenow 1819). — *Hirtella americana* L. var. *hexandra* (Willd.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 33 [17 Apr. 1867] (Hooker 1867).

Hirtella nitida Humb. & Bonpl. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 274 [Dec. 1819] (Roemer & Schultes 1819).

Hirtella acayacensis Sessé & Moc. ex DC., *Prodr. [A. P. de Candolle]* 2: 529 [mid Nov. 1825] (Candolle 1825).

Hirtella oblongifolia DC., *Prodr. [A. P. de Candolle]* 2: 529 [mid Nov. 1825] (Candolle 1825). — *Hirtella americana* var. *oblongifolia* (DC.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 34 [17 Apr. 1867] (Hooker 1867). — *Hirtella racemosa* var. *oblongifolia* (DC.) Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (3): 252 (Standley 1937).

Hirtella rosea Sessé & Moc. ex DC., *Prodr. [A. P. de Candolle]* 2: 529 [mid Nov. 1825] (Candolle 1825), nom. nud. pro syn.

Hirtella coriacea Mart. & Zucc., *Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.* 1: 383 (Martius & Zuccarini 1832).

Hirtella filiformis C.Presl, *Symb. Bot. [C. Presl]* 2: 23 [July 1834] (Presl 1834).

NOTES. — The name *Hirtella hexandra* is to be ascribed to Willdenow alone (Turland *et al.* 2018: Art. 46.3, Ex. 15; see also IPNI

2020 and Taylor & Gereau 2019); hence the authorship is neither “Willd. ex Roem. & Schult.” nor “Willd. ex Schult.”

VERNACULAR NAMES. — Wn: kulumnë • Nt: asiantefi.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M. Fleury 238*.

SIZE. — Brazil, Maranhão. *B.A. Krukoff 11867* (MO), 25 ft. × 4 inch. (7.5 m × 10 cm).

[295] *Hirtella silicea* Griseb.

Fl. Brit. W.I. [Grisebach] 229 [late 1860] (Grisebach 1860).

Hirtella melinonii Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 596 (Benoist 1923).

Hirtella manigera Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 387 [“1925” publ. Jan. 1926] (Kleinhoonte 1926).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kupešinilan, palipali • Wp: yanu'i pilá, yanu'i yowa • Cr: golèt-fronmi.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *E.M. Mélénion 132*, 1842 (holotype of *Hirtella melinonii*: P[P00741046]; iso-, NY[00428251], P[P00741045, P00741047], US[00130896]).

INVENTORY DATA (FG). — 9 trees in 3 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 10.8\text{ cm}$.

[296] *Hirtella suffulta* Prance

Fl. Neotrop. Monogr. 9: 308 [8 June 1972] (Prance 1972).

VERNACULAR NAMES. — Pa: inutauviye • Nt: kookoo.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3296*.

INVENTORY DATA (FG). — 13 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 52.2\text{ cm}$.

[297] *Hirtella tenuifolia* Prance

Fl. Neotrop. Monogr. 9: 321 [8 June 1972] (Prance 1972).

VERNACULAR NAMES. — Pa: ahavuiye • Ka: kupešinilan, kumboeta ší, piliwa epulu • Wn: kulumnë epí, kulunë epku • Nt: santo.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *B. Maguire et al. 47110* (holo-, NY[00468701]; iso-, IAN[IAN108482], US[00130913]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 16.9\text{ cm}$.

[298] *Hirtella triandra* Sw.

Prodr. [Swartz] 51 [20 Jun.-29 July 1788] (Swartz 1788). — *Chrysobalanus triandra* (Sw.) Morales, *Ánales Acad. Ci. Med. Habana*

23: 390 (Morales 1887). — *Zamzela racemosa* Raf., *Sylva Tellur.*: 90 (Rafinesque 1838), nom. illeg. superfl. (based on *Hirtella triandra*).

Hirtella paniculata Lam., *Encycl. [J. Lamarck et al.] 3* (1): 133 [19 Oct. 1789] (Lamarck 1789), nom. illeg. hom., non Sw. (Swartz 1788).

Hirtella cosmibuena Lam., *Tabl. Encycl. 2*[4 (2)]: 114 [6 Nov. 1797] (Lamarck 1797).

Hirtella racemosa Ruiz & Pav., *Flora Peruviana* 3: 5 (Ruiz & Pavón 1802), nom. illeg. hom., non Lam. (Lamarck 1789).

Hirtella peruviana Pers., *Syn. Pl. [Persoon]* 1: 250 [1 Apr.-15 June 1805] (Persoon 1805). — *Sphenista peruviana* (Pers.) Raf., *Sylva Tellur.*: 91 (Rafinesque 1838).

Hirtella castanea DC., *Prodri. [A. P. de Candolle]* 2: 528 [mid Nov. 1825] (Candolle 1825).

Hirtella bracteata Mart. & Zucc., *Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.* 1: 384 (Martius & Zuccarini 1832).

Hirtella jamaicensis Urb., *Symb. Antill. [Urban]* 5 (3): 355 [20 May 1908] (Urban 1908).

Hirtella multiflora Urb., *Symb. Antill. [Urban]* 5 (3): 356 [20 May 1908] (Urban 1908).

NOTES. — *Hirtella peruviana* Pers. is a legitimate and valid replacement name for *H. racemosa* Ruiz & Pav. Although not intended as such (the older *H. racemosa* Lam. is not cited in the protologue), it complies with both requirements of Art. 6.13 (Turland *et al.* 2018): i) the potentially replaced name (*H. racemosa* Ruiz & Pav.) is given as synonym, and ii) the conditions for valid publication (description) are independently met. Being legitimate, it has to be treated as the basionym of *Sphenista peruviana* (Pers.) Raf.

HERBARIUM DATA (FG). — No specimen seen, present according to Prance & Sothers (2003b).

SIZE. — Up to 15 m tall (Prance & Sothers 2003b).

Genus *Hymenopus* (Benth.) Sothers & Prance

[299] *Hymenopus amapaensis* (Prance) Sothers & Prance (Fig. 17B)

Kew Bull. 71 (4)-58: 17 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania amapaensis* Prance, *Fl. Neotrop. Monogr.* 9: 174 [8 June 1972] (Prance 1972).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: inutauviye.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3953*.

INVENTORY DATA (FG). — 19 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 21.9\text{ cm}$.

[300] *Hymenopus caudatus* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 17 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania caudata* Prance, *Fl. Neotrop. Monogr.* 9: 100 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *S.A. Mori et al. 24995*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.4\text{ cm}$.



FIG. 17. — Chrysobalanaceae: **A**, *Hirtella paniculata* Sw. (M.-F. Prévost 3856); **B**, *Hymenopus amapaensis* (Prance) Sothers & Prance (M.-F. Prévost & D. Sabatier 3002); **C**, *Hymenopus latistipulus* (Prance) Sothers & Prance (M.-F. Prévost & D. Sabatier 2992); **D**, *Licania ovalifolia* Kleinhoonte (D. Sabatier & J.-F. Molino 4835). A-C, © M.-F. Prévost/IRD; D, © D. Sabatier/IRD.

[301] *Hymenopus divaricatus* (Benth.) Sothers & Prance

Kew Bull. 71 (4)-58: 17 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania divaricata* Benth., *J. Bot. [Hooker]* 2: 221 (Bentham 1840). — *Licania heteromorpha* Benth. var. *divaricata* (Benth.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 45 (Fritsch 1889).

VERNACULAR NAMES. — Pa: inutauviye • Ka: anaula, ityulutano yapopale.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5629*.

INVENTORY DATA (FG). — 33 trees in 3 plots; $F_{\max} = 3.6\%$; $dbh_{\text{inv}} = 50.5\text{ cm}$.

[302] *Hymenopus glabriflorus* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 18 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania glabriflora* Prance, *Fl. Neotrop. Monogr.* 9: 104 [8 June 1972] (Prance 1972).

VERNACULAR NAMES. — Ka: kumboeta šipyoti.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori* 2344.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30.9$ cm.

[303] *Hymenopus heteromorphus* (Benth.) Sothers & Prance var. *heteromorphus*

Kew Bull. 71 (4)-58: 18 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania heteromorpha* Benth., *J. Bot. [Hooker]* 2: 221 (Bentham 1840).

Licania guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1199 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Licania benthamii Hook.f., *Fl. Bras. [Martius]* 14 (2): 12 [17 Apr. 1867] (Hooker 1867).

Licania biglandulosa Griseb. ex Urb., *Symb. Antill. [Urban]* 5 (3): 354 [20 May 1908] (Urban 1908).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: asiu, ašiu, asiu-seinó, ašiu-seinó, inutauviye • Ka: anaula, kumeti, yapopale • Wp: wila imi'i, wila umi'i • Wn: kulawaja, kulawaju • Nt: boliken kookoo • Cr: golèt-endjen, golèt-rouj • Fr: galette rouge • Br: caripérana, macucu-fofo.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2270.

INVENTORY DATA (FG). — 1001 trees in 136 plots; $F_{\max} = 7.5\%$; $dbh_{\text{inv}} = 53$ cm.

[304] *Hymenopus heteromorphus* var. *glabrus* (Mart. ex Hook.f.) Sothers & Prance

Kew Bull. 71 (4)-58: 18 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania glabra* Mart. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 10 [17 Apr. 1867] (Hooker 1867). — *Licania heteromorpha* var. *glabra* (Mart. ex Hook.f.) Prance, *Fl. Neotrop. Monogr.* 9: 108 [8 June 1972] (Prance 1972).

Licania costata Spruce ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 10 [17 Apr. 1867] (Hooker 1867), *nom. nud. pro syn.*

VERNACULAR NAMES. — Wp: wila imi'i, wila umi'i • Cr: golèt-endjen, golèt-rouj • Fr: galette rouge • Br: caripérana, macucu-fofo.

HERBARIUM DATA (FG). — A single collection, *C. Baraloto* *et al.* 3500.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 37.7$ cm.

[305] *Hymenopus intrapetiolaris* (Spruce ex Hook.f.) Sothers & Prance

Kew Bull. 71 (4)-58: 19 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania intrapetiolaris* Spruce ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 11 [17 Apr. 1867] (Hooker 1867).

Licania heteromorpha var. *grandifolia* Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 513 (Benoist 1919).

Licania intrapetiolaris var. *brevis* J.F.Macbr., *Candollea* 5: 369 (Macbride 1934).

HERBARIUM DATA (FG). — A single collection, *J. Martin s.n.* (original material of *Licania heteromorpha* var. *grandifolia*: B[[BM000602310](#)], K[[K000220657](#)], US[[00130810](#)]).

SIZE. — Venezuela, Bolíva. *B.M. Boom* 10418 (MO), 15 m × 15 cm.

[306] *Hymenopus laevigatus* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 19 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania laevigata* Prance, *Fl. Neotrop. Monogr.* 9S: 41 [8 Mar. 1989] (Prance 1989).

VERNACULAR NAMES. — Pa: inutauviye • Wp: wila imi'i, wila umi'i • Br: macucu-sangue.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4234.

INVENTORY DATA (FG). — 7 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 19.3$ cm.

[307] *Hymenopus latifolius* (Benth. ex Hook.f.) Sothers & Prance

Kew Bull. 71 (4)-58: 19 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania latifolia* Benth. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 11 [17 Apr. 1867] (Hooker 1867).

Licania obovata Benth. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 11 [17 Apr. 1867] (Hooker 1867).

VERNACULAR NAMES. — Pa: inutauviye • Br: macucu-sangue.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier* 1539.

INVENTORY DATA (FG). — 22 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 35.7$ cm.

[308] *Hymenopus latistipulus* (Prance) Sothers & Prance (Fig. 17C)

Kew Bull. 71 (4)-58: 20 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania latistipula* Prance, *Fl. Neotrop. Monogr.* 9: 103 [8 June 1972] (Prance 1972).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: wila imi'i wu, wila umi'i wu.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2417.

INVENTORY DATA (FG). — 50 trees in 354 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.7$ cm.

[309] *Hymenopus macrophyllus* (Benth.) Sothers & Prance

Kew Bull. 71 (4)-58: 20 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania macrophylla* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 240 (Bentham 1850).

VERNACULAR NAMES. — Pa: inura • Ka: alauna, atana • Te: anawa • Wp: ana wila • Wn: kalapaimé, makalawalaimé • Nt: busi peesina • Cr: anawa, grigri-rouj • Br: anauera, anauira, macucu-terra, pintadinha.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *M.-F. Prévost* 1364, dbh 20 cm.

[310] *Hymenopus occultans* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 20 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania occultans* Prance, *Fl. Neotrop. Monogr.* 9S: 42 [8 Mar. 1989] (Prance 1989).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 3857.

INVENTORY DATA (FG). — 31 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 34.1$ cm.

[311] *Hymenopus reticulatus* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 21 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania reticulata* Prance, *Fl. Neotrop. Monogr.* 9: 97 [8 June 1972] (Prance 1972).

VERNACULAR NAMES. — Wp: wila im'i, wila um'i.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P. Grenand* 642; *M.-F. Prévost* & *D. Sabatier* 2789, height 18 m.

Genus *Leptobalanus* (Benth.) Sothers & Prance

[312] *Leptobalanus apetalus* (E.Mey.) Sothers & Prance var. *apetalus*

Kew Bull. 71 (4)-58: 24 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Hirtella apetala* E.Mey., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 803 (Meyer 1825). — *Licania apetala* (E.Mey.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 54 (Fritsch 1889).

Licania pendula Benth., *J. Bot. [Hooker]* 2: 218 (Bentham 1840). — *Moquilea pendula* (Benth.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 22 [17 Apr. 1867] (Hooker 1867). — *Licania apetala* var. *pendula* (Benth.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 55 (Fritsch 1889).

Licania floribunda Benth., *J. Bot. [Hooker]* 2: 219 (Bentham 1840). — *Moquilea floribunda* (Benth.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 21 [17 Apr. 1867] (Hooker 1867).

Moquilea orinocensis Rusby, *Descr. S. Amer. Pl.* 27 [20 Dec. 1920] (Rusby 1920).

Licania dahlgrenii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (3): 255 (Standley 1937).

Licania hylaea Cuatrec., *Brittonia* 8 (3): 198 [July 1956] (Cuatrecasas 1956).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: kuwepi, kwepi • Br: caraipé, caripé.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 2780.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42$ cm.

[313] *Leptobalanus apetalus* var. *apertus* (Benth.) Sothers & Prance

Kew Bull. 71 (4)-58: 24 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania aperta* Benth., *J. Bot. [Hooker]* 2: 218 (Bentham 1840). — *Licania apetala* var. *aperta* (Benth.) Prance, *Fl. Neotrop. Monogr.* 9: 68 [8 June 1972] (Prance 1972).

Licania pubiflora Benth., *J. Bot. [Hooker]* 2: 219 (Bentham 1840).

Licania caracasana Klotzsch ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 25 [17 Apr. 1867] (Hooker 1867), *nom. nud. pro syn.*

Licania affinis Kuntze, *Revis. Gen. Pl.* 1: 217 [5 Nov. 1891] (Kuntze 1891), *nom. illeg. hom., non* Fritsch (1889).

Licania kuntzeana Urb., *Symb. Antill. [Urban]* 5 (3): 353 [20 May 1908] (Urban 1908), in obs. — *Moquilea kuntzeana* (Urb.) R.O. Williams, *Fl. Trinidad & Tobago* 1 (5): 314 [Aug. 1932] (Williams 1932).

VERNACULAR NAMES. — Ka: kuwepi, kwepi • Br: caraipé, caripé.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville* 488.

SIZE. — Up to 40 m tall (Prance 1972).

[314] *Leptobalanus granvillei* (Prance) Sothers & Prance

Kew Bull. 71 (4)-58: 26 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania granvillei* Prance, *Proc. Kon. Ned. Akad. Wetensch. C* 89 (1): 114 (Prance 1986).

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *S.A. Mori* & *B.M. Boom* 14764 (holo-, NY[00345874]; iso-, CAY[CAY010621], P[P00745988]).

INVENTORY DATA (FG). — 132 trees in 48 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 47.7$ cm.

[315] *Leptobalanus latus* (J.F.Macbr.) Sothers & Prance

Kew Bull. 71 (4)-58: 26 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania lata* J.F.Macbr., *Candollea* 5: 369 (Macbride 1934).

VERNACULAR NAMES. — Br: caripé, macucu-bobo.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-1468.

SIZE. — Up to 20 m tall (Prance 1986).

[316] *Leptobalanus longistylus* (Hook.f.) Sothers & Prance

Kew Bull. 71 (4)-58: 26 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Moquilea longistyla* Hook.f., *Fl. Bras. [Martius]* 14 (2):

24 [17 Apr. 1867] (Hooker 1867). — *Licania longistyla* (Hook.f.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 56 (Fritsch 1889).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3243*.

INVENTORY DATA (FG). — 77 trees in 27 plots; $F_{\max} = 2.6\%$; $\text{dbh}_{\text{inv}} = 44.8\text{ cm}$.

[317] *Leptobalanus octandrus*

(Hoffmanns. ex Roem. & Schult.) Sothers & Prance

Kew Bull. 71 (4)-58: 27 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Hirtella octandra* Hoffmanns. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 274 [Dec. 1819] (Roemer & Schultes 1819). — *Licania octandra* (Hoffmanns. ex Roem. & Schult.) Kuntze, *Revis. Gen. Pl.* 1: 217 [5 Nov. 1891] (Kuntze 1891). — *Licania turiuva* Cham. & Schltldl., *Linnaea* 2: 550 (Chamisso & Schlechtendal 1827), “*Turiuva*”, nom. illeg. superfl. (based on *Hirtella octandra*). — *Moquilea turiuva* Hook.f., *Fl. Bras. [Martius]* 14 (2): 25 [17 Apr. 1867] (Hooker 1867), nom. illeg. superfl. (based on *Hirtella octandra*).

Licania bothynophylla Mart., *Flora* 24 (2, Beibl.): 15 (Martius 1841). — *Moquilea bothynophylla* (Mart.) Hook.f., *Fl. Bras. [Martius]* 14 (2): 26 [17 Apr. 1867] (Hooker 1867).

Moquilea utilis Hook.f., *Fl. Bras. [Martius]* 14 (2): 24 [17 Apr. 1867] (Hooker 1867). — *Licania utilis* (Hook.f.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 56 (Fritsch 1889).

Licania sellowiana Klotzsch ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 26 [17 Apr. 1867] (Hooker 1867), nom. nud. pro syn.

Licania hookeri Fritsch var. *obtusa* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 368 (Huber 1909).

Licania takutuensis Standl., *Lloydia* 2 (3): 182 (Standley 1939).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *Moquilea turiuva* Hook.f. is based on the illegitimate *Licania turiuva* Cham. & Schltldl. It is thus treated as a new name, not a new combination.

VERNACULAR NAMES. — Ka: kuwepi, kwepi • Br: caraipé, caripé.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5300*.

INVENTORY DATA (FG). — 27 trees in 9 plots; $F_{\max} = 2.6\%$; $\text{dbh}_{\text{inv}} = 52.4\text{ cm}$.

[318] *Leptobalanus persaudii*

(Fanshawe & Maguire) Sothers & Prance

Kew Bull. 71 (4)-58: 28 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Licania persaudii* Fanshawe & Maguire, *Bull. Torrey Bot. Club* 75 (4): 375 [Jul.-Aug. 1948] (Fanshawe & Maguire 1948).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kuwepi, kwepi.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori et al. 24370*.

SIZE. — Up to 30 m tall (Prance 1986).

[319] *Leptobalanus sprucei* (Hook.f.) Sothers & Prance

Kew Bull. 71 (4)-58: 28 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Moquilea sprucei* Hook.f., *Fl. Bras. [Martius]* 14 (2): 22 [17 Apr. 1867] (Hooker 1867). — *Licania sprucei* (Hook.f.) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 55 (Fritsch 1889).

VERNACULAR NAMES. — Pa: kwep • Nt: kookoo • Br: caraipé, caripé.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3657*.

INVENTORY DATA (FG). — 160 trees in 59 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 39.6\text{ cm}$.

Genus *Licania* Aubl.

[320] *Licania affinis* Fritsch

Ann. K. K. Naturhist. Hofmus. 4: 50 (Fritsch 1889).

Licania schomburgkiana Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1199 [“1848” publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

VERNACULAR NAMES. — Wp: pali’i towu • Cr: bwa-golèt, golèt.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 4834*.

INVENTORY DATA (FG). — 23 trees in 16 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 51.6\text{ cm}$.

[321] *Licania alba* (Bernoulli) Cuatrecasas

Contr. U.S. Natl. Herb. 35 (6): 588 [21 Aug. 1964] (Cuatrecasas 1964). — *Theobroma album* Bernoulli, *Neue Denkschr. Schw. Naturf. Ges.* 24 (3): 14 (Bernoulli 1871), “*alba*”.

Caraipa longifolia Aubl., *Hist. Pl. Guiane* 1: 561 [Jun.-Dec. 1775] (Aublet 1775).

Licania venosa Rusby, *Descr. S. Amer. Pl.* 26 [20 Dec. 1920] (Rusby 1920).

Licania longifolia Benoist, *Bull. Mus. Natl. Hist. Nat.* 28: 252 (Benoist 1923).

NOTES. — *Caraipa longifolia* Aubl. is the earliest name for this taxon, but the epithet is no longer transferable to *Licania* because the name is occupied by *L. longifolia* Benoist. Although also a synonym of *L. alba*, *L. longifolia* is not based on Aublet’s taxon, having a distinct type. Cuatrecasas (1964: 588) erroneously gives “Übersicht Theobroma 14. 1869” as the publication reference for *Theobroma album* Bernoulli; no such name is found therein.

VERNACULAR NAMES. — Pa: bukutru-gateu-seinō, kwep • Ka: kolokolo • Wp: iwa tā’iy, payula • Wn: etuwe, kuepi • Nt: baaka kookoo, kookoo • Cr: golèt-blanc-gran-fey • Fr: galette blanche à grandes feuilles • Br: caraipé, caripé.

HERBARIUM DATA (FG). — 83 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (lectotype of *Licania longifolia* [P[P00745942] designated by Prance & Sothers [2003a: 146]]).

INVENTORY DATA (FG). — 2359 trees in 159 plots; $F_{\max} = 9.2\%$; $\text{dbh}_{\text{inv}} = 96.5\text{ cm}$.

[322] *Licania canescens* Benoist

Bull. Mus. Natl. Hist. Nat. 25: 514 (Benoist 1919).

NOTES. — Benoist (1919: 514) cited two collections in the protologue: “Maroni n° 13 (Mélinon); Maroni n° 43 (Wachenheim)”. Prance (1972: 134) cited for the type “Mélinon sn, French Guiana, Maroni, fl (lectotype, P; isolectotypes, GH, K)”; his lectotype corresponds to three specimens at P ([P00745970](#), [P00745972](#), [P00745973](#)). All three are labelled “*Licania canescens* R. Ben.” apparently in Benoist’s hand, and the same is true of other *Mélinon s.n.* at F, GH, K, NY and US (see below). It cannot be excluded that all these Mélinon specimens are isolectotypes. However, there is only one *Mélinon 13* in P ([P00745971](#)), which is therefore here designated as the lectotype.

VERNACULAR NAMES. — Pa: bukutru-ateu, bukutru-gateu • Te: tsibila lipi • Wp: payula • Nt: kookoo • Cr: golèt-nwé, grigri • Fr: galette noire • Br: caraipé, caripé, macucu-farinha-seca.

HERBARIUM DATA (FG). — 101 collections at CAY. Sel. exs.: *E.M. Mélinon 13*, s.d. (lecto-, P [[P00745971](#)]), here designated; possible isolectotypes: F [[V0053974F](#) {1862}], GH [[GH00046252](#) {s.d.}], K [[K00022089](#) {1863}], NY [[00428358](#) {s.d.}], P [[P00745970](#) {s.d.}], [P00745972](#) {1863}, [P00745973](#) {1864}], US [[00130724](#) {s.d.}].

INVENTORY DATA (FG). — 748 trees in 130 plots; $F_{\max} = 4.9\%$; $dbh_{inv} = 44.7$ cm.

[323] *Licania coriacea* Benth.

J. Bot. [Hooker] 2: 221 (Bentham 1840).

Licania pallidula Standl., *Bull. Torrey Bot. Club* 67 (4): 286 [Apr. 1940] (Standley 1940).

HERBARIUM DATA (FG). — No specimen seen, present according to Prance (1986).

SIZE. — Up to 25 m tall (Prance 1986).

[324] *Licania cyathodes* Benoist

Bull. Mus. Natl. Hist. Nat. 25: 513 (Benoist 1919).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: bukutru-ateu-priye, bukutru-gateu-priye • Ka: palipyo, polipyo, polipyoli • Wn: konoimë, watumo.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *P.A. Sagot s.n.*, Oct. 1856 (lecto-, P [[P00745955](#)]), designated by Prance [1972: 113; and on a label dated January 1966 affixed to the specimen]; isolecto-, P [[P00745956](#), [P00745957](#)]).

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 21$ cm.

[325] *Licania davillifolia* Benoist

Bull. Mus. Natl. Hist. Nat. 25: 513 (Benoist 1919).

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (lecto-, P [[P00745952](#)]) designated by Prance [1972: 133];

isolecto-, BM [[BM000602317](#)], E [[E00326668](#)], F [[V0053988F](#)], K [[K000220715](#)]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 24.6$ cm.

[326] *Licania densiflora* Kleinhoonte

Recueil Trav. Bot. Néerl. 22: 383 [“1925” publ. Jan. 1926] (Kleinhoonte 1926).

Licania kanukuensis Standl., *Lloydia* 2 (3): 182 (Standley 1939).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: bukutru-gateu-duwë • Ka: pesisilan • Nt: baaka kookoo, kookoo • Cr: bwa-golèt, golèt • Fr: galette.

HERBARIUM DATA (FG). — 94 collections at CAY. Sel. exs.: *D. Sabatier 925*.

INVENTORY DATA (FG). — 151 trees in 44 plots; $F_{\max} = 4.6\%$; $dbh_{inv} = 50.9$ cm.

[327] *Licania discolor* Pilg.

Notizbl. Königl. Bot. Gart. Berlin 6: 137 [20 Mar. 1914] (Pilger 1914).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *S.A. Mori et al. 23841*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 42.2$ cm.

[328] *Licania elliptica* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (3): 255 (Standley 1937).

Licania paniculata Fanshawe & Maguire, *Bull. Torrey Bot. Club* 75 (3): 323 [May-June 1948] (Fanshawe & Maguire 1948).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3538*.

SIZE. — Ecuador. *R. Valencia et al.* 68749 (MO), 9 m × 11 cm.

[329] *Licania fanshawei* Prance

Fl. Neotrop. Monogr. 9: 112 [8 June 1972] (Prance 1972).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al. 20808*.

INVENTORY DATA (FG). — 10 trees in 2 plots; $F_{\max} = 1.4\%$; $dbh_{inv} = 30.6$ cm.

[330] *Licania hypoleuca* Benth.

Bot. Voy. Sulphur [Bentham]: 91 [26 Oct. 1844] (Bentham 1844).

Licania microcarpa Hook.f., *Fl. Bras. [Martius]* 14 (2): 15 [17 Apr. 1867] (Hooker 1867).

Licania grisea Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 382 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

Licania parvifolia Pittier, *Bol. Soc. Venez. Ci. Nat.* 4: 351 (Pittier 1938), *nom. illeg. hom.*, *non* Huber (1909).

Licania parviflora var. *conduplicata* Maguire, *Fieldiana, Bot.* 28 (2): 253 (Maguire 1952).

VERNACULAR NAMES. — Pa: bukutru-ateu, bukutru-gateu • Ka: ku-wepilan tipulu • Wp: pali'i sī.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2970*.

INVENTORY DATA (FG). — 24 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 49.3$ cm.

[331] *Licania incana* Aubl.

Hist. Pl. Guiane 1: 119 [Jun.-Dec. 1775] (Aublet 1775). — *Hedycrea incana* (Aubl.) J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1)*: 428 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. superfl.* (genus name superfluous, based on the type of *Licania* Aubl.). — *Chrysobalanus incanus* (Aubl.) M.Gómez, *Fl. Cuba [Gómez]*: 39 (Gómez 1887), *nom. illeg. hom.*, *non* Raf. (Rafinesque 1836).

Licania crassifolia Benth., *J. Bot. [Hooker]* 2: 221 (Bentham 1840). — *Licania leptostachya* var. *crassifolia* (Benth.) Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 512 (Benoist 1919).

VERNACULAR NAMES. — Ka: akuli elepali, caligni (*fide* Aublet 1775), kwepilan • Wp: payula piyū • Cr: kaligni-blan • Br: oiticica.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777908] designated by Lanjouw & Uittien [1940: 152]); *H. van der Werff 23493* (MO), height 15 m.

[332] *Licania irwinii* Prance

Fl. Neotrop. Monogr. 9: 113 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2930*.

INVENTORY DATA (FG). — 190 trees in 20 plots; $F_{\max} = 7.9\%$; $\text{dbh}_{\text{inv}} = 53.2$ cm.

[333] *Licania kunthiana* Hook.f.

Fl. Bras. [Martius] 14 (2): 16 [17 Apr. 1867] (Hooker 1867).

Licania hypargyrea Malme, *Ark. Bot.* 23A(4): 12 (Malme 1930).

Licania parviflora var. *submembranacea* Maguire, *Fieldiana, Bot.* 28 (2): 254 (Maguire 1952).

VERNACULAR NAMES. — Pa: á-kiavunó-tivarabuyenē, bukutru-gateu • Wp: pali'i, pali'i sili, payula sili • Nt: kookoo, weti baka • Cr: golèt-grigri • Br: pajurazinho.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2537*.

INVENTORY DATA (FG). — 62 trees in 16 plots; $F_{\max} = 4.2\%$; $\text{dbh}_{\text{inv}} = 71.3$ cm.

[334] *Licania laxiflora* Fritsch

Ann. K. K. Naturhist. Hofmus. 4: 46 (Fritsch 1889).

Licania macrophylla Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1198 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Licania gracilis Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 382 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

VERNACULAR NAMES. — Pa: bukutru-ateu, bukutru-gateu • Ka: ku-wepilan • Wp: pali'i, pali'i pilá • Nt: kookoo, santi kookoo, weti kookoo • Br: caripérana.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15232*.

INVENTORY DATA (FG). — 295 trees in 86 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 56.8$ cm.

[335] *Licania leptostachya* Benth.

J. Bot. [Hooker] 2: 220 (Bentham 1840).

Licania incana var. *axilliflora* Sagot, *Ann. Sci. Nat., Bot. sér. 6, 15:* 305 (Sagot 1883). — *Licania leptostachya* var. *axilliflora* (Sagot) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 46 (Fritsch 1889). — *Licania axilliflora* (Sagot) Hochr., *Bull. New York Bot. Gard.* 6 (21): 273 [27 Sep. 1910] (Hochreutiner 1910).

VERNACULAR NAMES. — Ka: kwepilan • Te: tsibila lipikit • Wp: pali'i sī, payula sili • Cr: bwa-golèt, golèt • Br: caraipé, caripé.

HERBARIUM DATA (FG). — 54 collections at CAY. Sel. exs.: *M.-F. Prévost 1301*.

INVENTORY DATA (FG). — 57 trees in 14 plots; $F_{\max} = 4.8\%$; $\text{dbh}_{\text{inv}} = 18.7$ cm.

[336] *Licania majuscula* Sagot

Ann. Sci. Nat., Bot. sér. 6, 15: 306 (Sagot 1883).

Licania hostmannii Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 42 (Fritsch 1889), "Hostmanni".

VERNACULAR NAMES. — Ka: wokili kuperéni, yapopale membo • Wp: payula.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (holo-, P[P00746050]; iso-, BM[BM000602320], F[F0054025F, V0054026F], GH[GH00046286], K[K000220723], NY[00428473], P[P00746051], US[00130830]).

INVENTORY DATA (FG). — 82 trees in 36 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 70$ cm.

[337] *Licania maxima* Prance

Fl. Neotrop. Monogr. 9: 165 [8 June 1972] (Prance 1972).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *B. Maguire et al. 47081* (holo-, NY[00428477]; iso-, F[V0054027F], GH[GH00046287], IAN[IAN108457, IAN137569], MG[MG026468], P[P00746118], U[U0001143], US[00130832]).

SIZE. — Up to 70 cm dbh (Prance 1972).

[338] *Licania membranacea* Sagot ex Laness.

Pl. Util. Col. Franç. 130 (Lanessan 1886).

Carapa latifolia Aubl., *Hist. Pl. Guiane* 1: 561 [Jun.-Dec. 1775] (Aublet 1775), syn. nov.

Licania galibica Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 515 (Benoist 1919).

NOTES. — *Carapa latifolia* Aubl. is the earliest name for this taxon, but its epithet can no longer be transferred to *Licania* because *L. latifolia* Benth. ex Hook.f. (1867) occupies the name.

VERNACULAR NAMES. — Pa: bukutru-ateu, bukutru-gateu • Ka: polipyo tagulewe, soloma, wokili kupešini • Nt: baaka kookoo, kookoo • Cr: bwa-golèt, golèt • Fr: galette.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Carapa latifolia*: BM[BM000602313]).

INVENTORY DATA (FG). — 1201 trees in 161 plots; $F_{\max} = 7.1\%$; $dbh_{inv} = 132$ cm.

[339] *Licania micrantha* Miq.

Stirp. Surinam. Select.: 29 ["1850" publ. Mar. 1851] (Miquel 1851).

Licania helvola Spruce ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 18 [17 Apr. 1867] (Hooker 1867), nom. nud. pro syn.

Licania anisophylla Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (3)*: 253 (Standley 1937).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: bukutru-gateu-wahuyo • Ka: soloma, wokili kupešini • Te: tsibit • Wp: pal'i'i • Wn: kalipoimë • Nt: baaka kookoo, kookoo • Cr: golèt-marikaj, golèt-nwé • Fr: galette marécage, galette noire • Br: caripérana, pintadinha.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3379*.

INVENTORY DATA (FG). — 326 trees in 101 plots; $F_{\max} = 1.8\%$; $dbh_{inv} = 104.6$ cm.

[340] *Licania ovalifolia* Kleinhoonte
(Fig. 17D)

Recueil Trav. Bot. Néerl. 30: 180 [July 1933] (Kleinhoonte 1933).

Licania stahelii Kleinhoonte, *Recueil Trav. Bot. Néerl.* 30: 181 [July 1933] (Kleinhoonte 1933).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: bukutru-gateu-duwë • Nt: kookoo, santi kookoo • Cr: golèt-azon, grigri • Fr: galette azon.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier 2347*.

INVENTORY DATA (FG). — 189 trees in 77 plots; $F_{\max} = 1.5\%$; $dbh_{inv} = 74$ cm.

[341] *Licania pallida* Spruce ex Sagot

Ann. Sci. Nat., Bot. sér. 6, 15: 306 (Sagot 1883). — *Licania parviflora* var. *pallida* Hook.f., *Fl. Bras. [Martius]* 14 (2): 18 [17 Apr. 1867] (Hooker 1867). — *Licania pallida* Spruce ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 18 [17 Apr. 1867] (Hooker 1867), *nom. nud. pro syn.*

NOTES. — The authorship of *L. parviflora* var. *pallida* is not "Spruce ex Hook.f."; Hooker (1867: 18) referred to "*Licania pallida* Spruce Mss.", thus Spruce's authorship applies only to the species, not the variety.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M. Hoff 5980*.

SIZE. — Up to 30 m tall (Prance 1998).

[342] *Licania parviflora* Benth.

J. Bot. [Hooker] 2: 221 (Bentham 1840).

Licania parviflora var. *subfalcata* Spruce ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 18 [17 Apr. 1867] (Hooker 1867).

Licania parviflora f. *brevifolia* Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 51 (Fritsch 1889), *nom. illeg.* (includes species type).

Licania parviflora f. *longifolia* Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 51 (Fritsch 1889).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1467*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh = 58.3$ cm.

[343] *Licania parvifructa* Fanshawe & Maguire

Bull. Torrey Bot. Club 75 (4): 374 [Jul.-Aug. 1948] (Fanshawe & Maguire 1948).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *S.A. Mori et al. 26472*.

INVENTORY DATA (FG). — 43 trees in 32 plots; $F_{\max} < 1\%$; $dbh_{inv} = 42.7$ cm.

[344] *Licania pruinosa* Benoist

Bull. Mus. Natl. Hist. Nat. 25: 516 (Benoist 1919).

VERNACULAR NAMES. — Wp: payula.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (type P[P00746083]).

SIZE. — Up to 20 m tall (Prance 1986).

[345] *Licania robusta* Sagot

Ann. Sci. Nat., Bot. sér. 6, 15: 306 (Sagot 1883).

Licania pachystachya Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 384 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

VERNACULAR NAMES. — Ka: ityulanano kupašini, solosolo bakoloidyi.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (holo-, P[P00746074]; iso-, BM[BM000602327], F[V0054063F, V0054064F], NY[00428559], P[P00746075], R[R000063240], US[00130857]).

INVENTORY DATA (FG). — 46 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 52.4$ cm.

[346] *Licania rodriquesii* Prance

Fl. Neotrop. Monogr. 9: 168 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori 2362*.

INVENTORY DATA (FG). — 20 trees in 18 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 83.4$ cm.

[347] *Licania silvae* Prance

Fl. Neotrop. Monogr. 9: 115 [8 June 1972] (Prance 1972).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3850*.

SIZE. — Brazil, Acre. *D.C. Daly et al. 11065* (MO), 30 m × 43 cm.

[348] *Licania* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 5750*, $dbh = 44$ cm.

[349] *Licania* sp. B

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3081*.

INVENTORY DATA (FG). — 8 trees in 1 plot; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 37.1$ cm.

[350] *Licania* sp. C

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *B. Riéra & D. Sabatier 1282*.

INVENTORY DATA (FG). — 7 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54$ cm.

Genus *Moquilea* Aubl.

[351] *Moquilea guianensis* Aubl.

Hist. Pl. Guiane 1: 521 [Jun.-Dec. 1775] (Aublet 1775), "Guyanensis" on plate. — *Licania guianensis* (Aubl.) Griseb., *Abh. Königl. Ges. Wiss. Göttingen* 7: 198 (Grisebach 1857).

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000793077]).

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.1$ cm.

[352] *Moquilea minutiflora* Sagot

Ann. Sci. Nat., Bot. sér. 6, 15: 308 (Sagot 1883). — *Licania minutiflora* (Sagot) Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 54 (Fritsch 1889).

Licania rondonii Pilg., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 540 [15 Aug. 1923] (Pilger 1923).

Moquilea riparia Gleason, *Bull. Torrey Bot. Club* 61 (4): 192 [Apr. 1934] (Gleason 1934). — *Licania riparia* (Gleason) Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (3): 258 (Standley 1937).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (holo-, P[P00746107]; iso-, A[A00046290], BM[BM000602298], F[V0054094F, V0054095F, V0054096F], GH[GH00046291], K[K000220600], NY[00428489], US[00130833]).

INVENTORY DATA (FG). — 73 trees in 14 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 47.1$ cm.

[353] *Moquilea* sp. A

NOTE. — This species seems close to *M. minutiflora* Sagot.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5049*.

INVENTORY DATA (FG). — 8 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 73.2$ cm.

Genus *Parinari* Aubl.

[354] *Parinari campestris* Aubl.

Hist. Pl. Guiane 1: 514 [Jun.-Dec. 1775] (Aublet 1775). — *Petrocarya campestris* (Aubl.) Willd., *Sp. Pl., ed. 4* 2 (1): 287 [Mar. 1799] (Willdenow 1799). — *Ferolia campestris* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 216 [5 Nov. 1891] (Kuntze 1891).

Balantium cordifolium Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 34 [Oct. 1825] (Hamilton 1825).

VERNACULAR NAMES. — Pa: á-kiavū, á-kiavunō, kuuku-araybu-seine • Ka: kubašini, kupašini • Wp: pal'i wate'e • Nt: fungutii kookoo • Cr: golèt-blanc, grigri-blanc • Fr: galette blanche • Br: pajurá, parinari.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000797863]).

INVENTORY DATA (FG). — 152 trees in 50 plots; $F_{\max} = 4.1\%$; $dbh_{\text{inv}} = 87$ cm.

[355] *Parinari excelsa* Sabine

Trans. Hort. Soc. London 5: 451 (Sabine 1824). — *Petrocarya excelsa* (Sabine) Steud., *Nomenc. Bot. [Steudel]*, ed. 2, 2: 309 (Steudel 1841), *nom. illeg. hom.*, *non* Jack (1822). — *Ferolia excelsa* (Sabine) Kuntze, *Revis. Gen. Pl.* 1: 216 [5 Nov. 1891] (Kuntze 1891).

Parinari brachystachya Benth., *J. Bot. [Hooker]* 2: 213 (Bentham 1840), “*Parinarium brachystachyum*”. — *Ferolia amazonica* Kuntze, *Revis. Gen. Pl.* 1: 216 [5 Nov. 1891] (Kuntze 1891), *nom. illeg. superfl.* (based on *Parinari brachystachya*).

Parinari amazonica Mart. ex Hook.f., *Fl. Bras. [Martius]* 14 (2): 52 [17 Apr. 1867] (Hooker 1867), “*Parinarium amazonicum*”, *nom. nud. pro syn.*

Parinari glazioviana Warm., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1874: 72 (Warming 1874), “*Parinarium glaziovianum*”.

Parinari salicifolia Engl., *Pflanzenw. Ost-Afrikas C*: 191 [July 1895] (Engler 1895), “*Parinarium salicifolium*”, *nom. illeg. hom.*, *non* (C.Presl) Miq. (Miquel 1855).

Parinari holstii Engl., *Pflanzenw. Ost-Afrikas C*: 423 [Aug. 1895] (Engler 1895), “*Parinarium Holstii*”. — *Parinari excelsa* subsp. *holstii* (Engl.) R.A.Graham, *Kew Bull.* 12 (2): 229 (Graham 1957).

Parinari elliotii Engl., *Bot. Jahrb. Syst.* 26 (3-4): 377 [31 Jan. 1899] (Engler 1899), “*Parinarium elliotii*”.

Parinari whytei Engl., *Bot. Jahrb. Syst.* 26 (3-4): 378 [31 Jan. 1899] (Engler 1899), “*Parinarium whytei*”.

Parinari verdickii De Wild., *Ann. Mus. Congo*, sér. 4, *Bot.* 1: 182 (De Wildeman 1903), “*Parinarium verdickii*”.

Parinari tenuifolia A.Chev., *Vég. Ut. Afr. Trop. Franç.* 5: 225 (Chevalier 1909), “*Parinarium tenuifolium*”.

Parinari excelsa var. *fulvescens* Engl., *Wiss. Erg. Deut. Zentr.-Afr. Exped.*, *Bot.* 2: 227 (Engler 1911), “*Parinarium excelsum*”.

Parinari mildbraedii Engl., *Wiss. Erg. Deut. Zentr.-Afr. Exped.*, *Bot.* 2: 227 (Engler 1911).

Parinari riparia R.E.Fr., *Repert. Spec. Nov. Regni Veg.* 12: 539 (Fries 1913), “*Parinarium riparium*”.

Parinari holstii var. *longifolia* Engl. ex De Wild., *Bull. Jard. Bot. État Bruxelles* 4: 80 (De Wildeman 1914), “*Parinarium holstii, longifolium*”, *nom. nud.*

Parinari laxiflora Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 44 (Ducke 1922), “*Parinarium laxiflorum*”.

Parinari nalaensis De Wild., *Pl. Bequaert.* 5: 289 (De Wildeman 1931), “*Parinarium nalaense*”.

Parinari laxiflora var. *lata* Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 33 [Sep. 1935] (Ducke 1935), “*Parinarium laxiflorum, latum*”.

Parinari sylvestris M.Kuhlm., *Loefgrenia* 13: 1 (Kuhlmann 1964).

VERNACULAR NAMES. — Ka: wokili besuli.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3562*.

INVENTORY DATA (FG). — 19 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 83$ cm.

[356] *Parinari montana* Aubl.

Hist. Pl. Guiane 1: 514 [Jun.-Dec. 1775] (Aublet 1775). — *Petrocarya montana* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (1): 287 [Mar. 1799] (Willdenow 1799). — *Chrysobalanus montanus* (Aubl.) M.Gómez, *Fl. Cuba [Gómez]*: 39 (Gómez 1887). — *Ferolia montana* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 216 [5 Nov. 1891] (Kuntze 1891).

Parinari pajura Benoist, *Bull. Mus. Natl. Hist. Nat.* 28: 252 (Benoist 1922), “*Parinarium pajura*”.

NOTES. — Emended by Ducke (1935: 180). The leaves on type sheet are *P. rodolphii* Huber (1909), and only the fruit represent *P. montana*. Aublet (1775, 1: 514) cited plates 204 & 205, but only plate 205 is correct, the branches and leaves of plate 204 are *P. rodolphii*.

VERNACULAR NAMES. — Pa: pirivri • Ka: kalapapoli, kubeshini, kupeshini, tapowonulen • Wp: iwa tā'iy • Nt: fungutii kookoo • Cr: grénn-rôch • Fr: graine roche • Br: pajurá-grande, pajurá-pedra.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM, fruit only, not seen); *S.A. Mori et al. 20790*.

INVENTORY DATA (FG). — 63 trees in 38 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 78$ cm.

[357] *Parinari parvifolia* Sandwith

Bull. Misc. Inform. Kew 1931 (7): 374 [21 July 1931] (Sandwith 1931).

VERNACULAR NAMES. — Pa: á-kiavū, á-kiavunō.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier 3005*.

INVENTORY DATA (FG). — 14 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 65$ cm.

[358] *Parinari rodolphii* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 6: 77 (Huber 1910), “*Parinarium rodolphi*”.

Parinari lucidissima Standl., *Lloydia* 2 (3): 183 (Standley 1939), “*Parinarium lucidissimum*”.

VERNACULAR NAMES. — Ka: kalapapoli, tapowonulen • Wp: iwa tā'iy sili, tata pota • Cr: kwépi • Br: farinha-seca.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3035*.

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 100$ cm.

[359] *Parinari* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5728*.

INVENTORY DATA (FG). — 1 tree, $dbh = 34$ cm.

Genus *Parinariopsis* (Huber) Sothers & Prance

[360] *Parinariopsis licaniiiflora* (Sagot) Sothers & Prance

Kew Bull. 71 (4)-58: 11 [epubl. 15 Dec. 2016] (Sothers & Prance 2016). — *Moquilea licaniiiflora* Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 15: 308 (Sagot 1883), “licaniaeflora”. — *Licania bracteosa* Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 54 (Fritsch 1889), nom. illeg. superfl. (based on *Moquilea licaniiiflora*). — *Licania licaniiiflora* (Sagot) S.F.Blake, *Contr. Gray Herb.* 52: 66 (Blake 1917), “licaniaeflora”.

Moquilea parviflora Blume, *Mus. Bot.* 2 (1-8): 92 [“1852” publ. Feb. 1856] (Blume 1856). — *Licania parviflora* (Blume) Lemée, *Fl. Guyane Franç.* 2: 23 (Lemée 1952), nom. illeg. hom., non Benth. (Bentham 1840).

Licania obtusifolia Fritsch, *Ann. K. K. Naturhist. Hofmus.* 4: 53 (Fritsch 1889).

Licania parinariooides Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 368 (Huber 1909).

Licania capinensis Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 71 (Huber 1910).

Licania parinariooides var. *latifolia* Maguire, *Brittonia* 7: 397 [Oct. 1952] (Maguire 1952).

Licania huberiana Maguire, *Brittonia* 7: 398 [Oct. 1952] (Maguire 1952).

Licania obovatifolia Maguire, *Brittonia* 7: 398 [Oct. 1952] (Maguire 1952).

Licania wilson-brownei Maguire, *Brittonia* 7: 399 [Oct. 1952] (Maguire 1952).

NOTE. — Prance (1972: 91) made a first-step lectotypification by selecting among the paratypes *J. Martin s.n.*, as “lecto-, P, isolecto-, K”. There are however three specimens of Martin’s original material at P (P00746011, P00746012, P00746013). It thus appears necessary to select one of them as the lectotype.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J. Martin s.n.* (lectotype of *Moquilea licaniiiflora* Sagot: P[P00746011], here designated; isolecto-, P[P00746012, P00746013], K[K000220643]).

INVENTORY DATA (FG). — 33 trees in 18 plots; $F_{\max} < 1\%$; $dbh_{inv} = 55.4$ cm.

Family CLUSIACEAE Lindl.

Genus *Chrysochlamys* Poepp. & Endl.

[361] *Chrysochlamys membranacea* Planch. & Triana

Ann. Sci. Nat., Bot. sér. 4, 14: 260 (Planchon & Triana 1860). — *Tovomitopsis membranacea* (Planch. & Triana) D’Arcy, *Ann. Missouri Bot. Gard.* 67 (4): 1034 [“1980” publ. 1981] (D’Arcy 1981).

VERNACULAR NAMES. — Wp: amá kiya, pasí’i wapo • Cr: pativyé-gran-bwa.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *M.-F. Prévost 1810*, 10 m tall.

Genus *Clusia* Plum. ex L.

[362] *Clusia cuneata* Benth.

London J. Bot. 2: 368 (Bentham 1843).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1645*.

SIZE. — Guyana. *H.D. Clarke 5485*, 10 m.

[363] *Clusia flavidula* (Benth.) Pipoly

Fl. Venez. Guayana 4: 272 (Pipoly 1998). — *Havetia flavidula* Benth., *London J. Bot.* 2: 369 (Bentham 1843). — *Havetiopsis flavidula* (Benth.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 247 (Planchon & Triana 1860).

Havetiopsis flexilis Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 246 (Planchon & Triana 1860). — *Havetia flexilis* Spruce ex Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 246 (Planchon & Triana 1860), nom. nud. pro syn. — *Havetia flexilis* (Planch. & Triana) Vesque, *Epharmosis* 3: 15, 73 (Vesque 1892).

Clusia gentlei Lundell, *Contr. Univ. Michigan Herb.* 6: 48 (Lundell 1941).

NOTES. — A species restricted to the edges of inselbergs. The name *Havetia flexilis* first appears as a *nomen nudum*, as a synonym of *Havetiopsis flexilis*. Vesque (1892: 73) validly published it with reference to “*H. flexilis* Planch. et Triana” under sect. (or subg.) *Havetiopsis*, and on plate 72 with an explicit citation of *Havetiopsis flexilis*.

VERNACULAR NAMES. — Ka: kusapoi, kuwapo’u, kwapo’u • Wp: pelepele sili • Cr: bwa-rwé, zongnon-danbwá.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *P. Grenand 1222*.

SIZE. — Up to 15 m tall (Pipoly et al. 1998).

[364] *Clusia fockeana* Miq.

Tijdschr. Nat. Geschied. Physiol. 10: 82 (Miquel 1843), “*Fockeana*”. — *Androstylium fockeanum* (Miq.) Miq., *Stirp. Surinam. Select.* 93 [“1850” publ. Mar. 1851] (Miquel 1851).

Clusia stahelii Maguire, *Bull. Torrey Bot. Club* 75 (4): 424 [Jul.-Aug. 1948] (Maguire 1948).

VERNACULAR NAMES. — Ka: kunapolan • Nt: kupa.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *D. Sabatier & E. Fonty 5762*.

INVENTORY DATA (FG). — 300 trees in 6 plots; $F_{\max} = 16.8\%$; $dbh_{inv} = 30.2$ cm.

[365] *Clusia grandiflora* Splitg.

Tijdschr. Nat. Geschied. Physiol. 9: 101 [Aug.-Sep. 1842] (Splitgerber 1842).

Clusia maxima Rich. ex Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 13: 325 (Planchon & Triana 1860), nom. nud. pro syn.

Clusia petiolata Klotzsch ex Engl., *Fl. Bras. [Martius]* 12 (1): 423 [1 Apr. 1888] (Engler 1888), nom. nud. pro syn.

NOTE. — Mostly epiphytic, rarely rooted in the ground.

VERNACULAR NAMES. — Pa: patakwik, zoyō-kamwi • Ka: kusapoi, kuwapo'u, kwapo'u, pelepele • Te: pelepele • Wp: pelepele • Wn: lele anapamigín • Nt: ingi waway • Cr: bwa-rwé, zongnon-danbwa • Br: cebola-grande-da-mata, manga-brava, mata-pau.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (P[[P05071041](#)]).

SIZE. — Guyana. *P.J.M. Maas & L.Y.T. Westra 3875* (MO), 7 m × 30 cm.

[366] *Clusia leprantha* Mart.

Nova genera et species plantarum [Martius] 3 (3): 165 ["1829" publ. Sep. 1832] (Martius 1832).

Arrudea purpurea Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 103 [Aug.-Sep. 1842] (Splitgerber 1842). — *Clusia purpurea* (Splitg.) Engl., *Fl. Bras. [Martius]* 12 (1): 416 [1 Apr. 1888] (Engler 1888).

VERNACULAR NAMES. — Nt: mangu • Br: apuí-grande, cebola-brava.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *Service Forestier 7855*.

SIZE. — Up to 25 m tall (Pipoly *et al.* 1998).

[367] *Clusia melchiorii* Gleason

Bull. Torrey Bot. Club 58 (6): 403 [June 1931] (Gleason 1931).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-J. de Granville B-5401*.

SIZE. — Up to 15 m tall (Pipoly *et al.* 1998).

[368] *Clusia minor* L.

Sp. Pl. 1: 510 [1 May 1753] (Linnaeus 1753).

Clusia venosa Jacq., *Enum. Syst. Pl.*: 34 [Aug.-Sep. 1760] (Jacquin 1760).

Clusia venosa L., *Sp. Pl.*, ed. 2, 2: 1495 [Aug. 1763] (Linnaeus 1763).

Clusia galactodendron Desv., *Ann. Sci. Nat., Bot. sér. 2*, 18: 313 (Desvaux 1842), "Galactodendrum".

Clusia pratensis Seem., *Bot. Voy. Herald [Seemann]* 3: 89 [Nov. 1853] (Seemann 1853).

Clusia couletii Duchass. ex Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 13: 333 (Planchon & Triana 1860), "Couleti", nom. nud. pro syn.

Clusia parviflora Vesque, *Epharmosis* 3: 10 (Vesque 1892).

Clusia cartilaginosa Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 97 [Dec. 1893] (Vesque 1893).

Clusia utilis S.F.Blake, *Contr. U.S. Natl. Herb.* 24 (1): 14 [11 Jan. 1922] (Blake 1922).

NOTE. — Restricted to the edges of inselbergs.

VERNACULAR NAMES. — Wp: pelepele sili.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-J. de Granville 2200*.

SIZE. — Venezuela, Bolívar. *J.A. Steyermark 87845* (L), 10 m.

[369] *Clusia nemorosa* G.Mey.

Prim. Fl. Esseq. 203 [Nov. 1818] (Meyer 1818).

Clusia lhotzkyana Schlehd., *Linnaea* 8: 184 (Schlechtendal 1833).

Clusia mammosa Casar., *Nov. Stirp. Bras.* 7: 60 [Sep ("Jul") 1843] (Casaretto 1843).

Clusia lhotzkyana var. *polygama* Choisy, *Descr. Guttif. Inde* 45 (Choisy 1849). — *Clusia nemorosa* var. *lhotzkyana* (Schlehd.) Engl., *Fl. Bras. [Martius]* 12 (1): 422 [1 Apr. 1888] (Engler 1888).

VERNACULAR NAMES. — Pa: wakukwa-tiranó • Ka: kunapolan.

HERBARIUM DATA (FG). — 76 collections at CAY. Sel. exs.: *M.-F. Prévost 1628*.

SIZE. — Up to 15 m tall (Pipoly *et al.* 1998).

[370] *Clusia octandra* (Poep.) Pipoly

Fl. Venez. Guayana 4: 276 (Pipoly 1998). — *Havetia octandra* Poep., *Nova genera ac species plantarum [Poepig & Endlicher]* 3: 11 [15-21 Aug. 1841] (Poepig 1841). — *Oedematopus octandrus* (Poep.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 250 (Planchon & Triana 1860).

NOTE. — Mostly epiphytic, rarely rooted in the ground.

VERNACULAR NAMES. — Te: tsalikwana'i • Wp: ayá pápē, ayá pápē sili, pelepele sili.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori et al. 23532*.

SIZE. — Colombia, Antioquia. *R. Fonnegra G. & W. Rengifo M. 4834* (MO), 8 m × 10 cm.

[371] *Clusia palmicida* Rich. ex Planch. & Triana

Ann. Sci. Nat., Bot. sér. 4, 13: 326 (Planchon & Triana 1860). — *Clusia palmicida* Rich., *Ann. Mus. Natl. Hist. Nat.* 17: 456 (Richard 1811), nom. inval.

Clusia martinii Sagot ex Engl., *Fl. Bras. [Martius]* 12 (1): 425 [1 Apr. 1888] (Engler 1888), nom. nud. pro syn.

NOTES. — Mostly epiphytic, rarely rooted in the ground. *Clusia palmicida* was not validly published by Richard (1811: 17) because his publication was a treatise on embryo development and his seed diagnosis did not distinguish this species from others. Indeed, the characters described are said to be shared with other species of the genus. His reference to the figure ("Pl. VI, fig. 64, a"), is erroneous, and is in fact, pl. 8, fig. 64, 65. J.J. Pipoly III in 2000 annotated the two sheets of *Richard s.n.* at P ([P01901181](#), [P01901182](#)) as holotype and isotype, but this typification was not published. Also, Planchon & Triana (1860a: 326) under this name cited two collections from French Guiana ("L.-C. Rich. in herb. de Franquev., Martin in herb. Mus. Paris") and one from Guyana ("Schomburgk, n° 739, indétermine"). The selection of a lectotype thus appears necessary.

VERNACULAR NAMES. — Pa: patakwik-kamwi • Ka: kusapoi, kuwapo'u, kwapo'u • Cr: bwa-rwé, zongnon-danbwa • Br: cebola-grande-da-mata, manga-brava, mata-pau.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (lectotype of *Clusia palmicida* Rich. ex Planch. & Triana, P[P01901181, here designated; isolecto-, P01901182]).

SIZE. — Up to 15 m tall (Pipoly *et al.* 1998).

[372] *Clusia panapanari* (Aubl.) Choisy

Prodr. [A. P de Candolle] 1: 559 [mid Jan. 1824] (Choisy 1824), “Pana-panari”. — *Quapoya panapanari* Aubl., *Hist. Pl. Guiane* 2: 900 [Jun.-Dec. 1775] (Aublet 1775), “Pana-panari”.

Clusia colorans Engl., *Fl. Bras. [Martius]* 12 (1): 426 [1 Apr. 1888] (Engler 1888). — *Quapoya colorans* Klotzsch ex Engl., *Fl. Bras. [Martius]* 12 (1): 426 [1 Apr. 1888] (Engler 1888), *nom. nud. pro syn.*

Clusia microphylla Engl., *Fl. Bras. [Martius]* 12 (1): 426 [1 Apr. 1888] (Engler 1888). — *Quapoya microphylla* Klotzsch ex Engl., *Fl. Bras. [Martius]* 12 (1): 426 [1 Apr. 1888] (Engler 1888), *nom. nud. pro syn.*

NOTE. — Mostly epiphytic, rarely rooted in the ground.

VERNACULAR NAMES. — Pa: patakwik • Wp: ayá pápē, ayá pápē sili, pelepele sili • Cr: bwa-rwé, zongnon-danbwa • Br: apuizinha.

HERBARIUM DATA (FG). — 144 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000611559]).

SIZE. — Up to 10 m tall (Pipoly *et al.* 1998).

[373] *Clusia scrobiculata* Benoist

Bull. Mus. Natl. Hist. Nat. 30: 511 (Benoist 1924).

VERNACULAR NAMES. — Pa: patakwik, patakwik-ahinē.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *J.-P. Lescure* 839.

SIZE. — Guyana. *J.J. Pipoly* 10414 (MO), 6 m × 10 cm.

Genus *Garcinia* L.

[374] *Garcinia benthamiana* (Planch. & Triana) Pipoly

Nuevo Cat. Fl. Vasc. Venezuela: 332 [after May 2008] (Pipoly 2008). — *Rheedia benthamiana* Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 320 (Planchon & Triana 1860). — *Rheedia macrophylla* var. *benthamiana* (Planch. & Triana) Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 500 [Dec. 1893] (Vesque 1893).

Rheedia sagotiana Engl., *Fl. Bras. [Martius]* 12 (1): 460 [1 Apr. 1888] (Engler 1888).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wakukwa-tiranó • Ka: nopitya, pakulilan • Te: kulupitá • Wp: kulupitá i, tatu apite • Nt: kukundiefu, meku susu • Br: bacuri-de-espinhos, bacuri-selvagem.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *P.A. Sagot* 1182, 1858 (original material of *Rheedia sagotiana*; BR[BR0000008676238], F[V0054522F], K[K000488572], MPU[MPU014348, MPU014349], S[S11-36125], US[00114327]).

INVENTORY DATA (FG). — 182 trees in 73 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 23.2\text{ cm}$.

[375] *Garcinia brasiliensis* Mart.

Flora 24 (2, Beibl.): 34 (Martius 1841). — *Rheedia brasiliensis* (Mart.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 310 (Planchon & Triana 1860).

Garcinia brasiliensis var. *parvifolia* Mart., *Flora* 24 (2, Beibl.): 34 (Martius 1841).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

HERBARIUM DATA (FG). — No collection at CAY. Sel. exs.: *J.M. Pires* 47466 (L), collected on the French Guianan bank of Oyapock river.

SIZE. — Up to 10 m tall (van den Berg 1979).

[376] *Garcinia macrophylla* Mart.

(Fig. 18A)

Flora 24 (2, Beibl.): 35 (Martius 1841). — *Rheedia macrophylla* (Mart.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 309 (Planchon & Triana 1860).

Rheedia macrantha Standl. & Steyermark, *Publ. Field Mus. Nat. Hist., Bot. Ser. 23 (2)*: 65 [14 Feb. 1944] (Standley & Steyermark 1944).

Garcinia megaphylla Verdc., *Kew Bull.* 31 (2): 262 (Verdcourt 1976), *nom. illeg. superfl.* (based on *Garcinia macrophylla*).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: wakukwa-tiranó • Ka: pakulilan • Te: kulupitá • Wp: kulupitá • Wn: mutu pupot eimë • Nt: kukundiefu, meku susu • Cr: konfitir-makak, koupiton • Br: bacuri-liso, bacuripari.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost* & *D. Sabatier* 2742.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 50\text{ cm}$.

[377] *Garcinia madruno* (Kunth) Hammel

Ann. Missouri Bot. Gard. 76 (3): 928 (Hammel 1989). — *Calophyllum madruno* Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 202 [25 Feb. 1822] (Kunth 1822), “madruño”. — *Rheedia madruno* (Kunth) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 315 (Planchon & Triana 1860). — *Verticillaria madruno* (Kunth) Tul. ex Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 315 (Planchon & Triana 1860), *nom. nud. pro syn.*

Verticillaria acuminata Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 140 [late Dec. 1798] (Ruiz & Pavón 1798). — *Rheedia acuminata* (Ruiz & Pav.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 314 (Planchon & Triana 1860).

Chloromyron verticillatum Pers., *Syn. Pl. [Persoon]* 2 (1): 73 [Nov. 1806] (Persoon 1806), *nom. illeg. superfl.* (based on *Verticillaria acuminata*).

Garcinia floribunda Miq., *Stirp. Surinam. Select.*: 89 [“1850” publ. Mar. 1851] (Miquel 1851). — *Rheedia floribunda* (Miq.) Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14*: 319 (Planchon & Triana

1860). — *Rheedia acuminata* var. *floribunda* (Miq.) Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 510 [Dec. 1893] (Vesque 1893).

Rheedia rostrata Vesque, *Epharmosis* 2: 24 (Vesque 1889).

Rheedia kappleri Eyma, *Polygon. Guttif. Lecythid. Surinam:* 33 (Eyma 1932), nom. illeg. superfl. (based on *Rheedia floribunda*).

VERNACULAR NAMES. — Pa: wakukwa-tiranō • Ka: ololome, olome, tapukēn nōpitya • Wp: kulupitā i, tatu apite • Wn: wiipē • Nt: kukundiefu, meku susu • Br: bacuri-de-espinhos.

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *M.-F. Prévost 2169*.

INVENTORY DATA (FG). — 131 trees in 68 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 32.5$ cm.

Genus *Moronoea* Aubl.

[378] *Moronoea coccinea* Aubl.

Hist. Pl. Guiane 2: 789 [Jun.-Dec. 1775] (Aublet 1775). — *Symponia coccinea* (Aubl.) Oken, *Allg. Naturgesch.* 3 (2): 1431 (Oken 1841). — *Moronoea montana* Schlehd., *Linnaea* 8: 189 (Schlechtendal 1833), nom. illeg. superfl. (based on “*Moronoea coccinea* Aubl. t. 313, n. 1-9”).

Moronoea grandiflora Choisy, *Mém. Soc. Hist. Nat. Paris*, sér. 2, 1: 230 (Choisy 1823), syn. nov. — *Symponia grandiflora* Rich. ex Choisy, *Mém. Soc. Hist. Nat. Paris*, sér. 2, 1: 230 (Choisy 1823), nom. nud. pro syn.

NOTES. — *Moronoea coccinea* Aubl. has often been treated as a synonym of *Symponia globulifera* L.f., because Aublet’s plate 313 (1775) mixes elements of *M. coccinea* (habit and fig. 1-9) and *S. globulifera* (fig. a-j). However, the original specimen at BM (BM000611835) is not a mixture, and the name does not apply to the *S. globulifera* elements on plate 313.

VERNACULAR NAMES. — Pa: ti-wašiunu • Ka: ananiyu, mani • Wp: māni, walatiwā • Wn: mani, manimē • Nt: mataki, mongo mataki • Cr: manni-montangn • Fr: manil montagne • Br: anani-da-terra-firme, bacuri-bravo.

HERBARIUM DATA (FG). — 67 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[BM000611835], here designated).

INVENTORY DATA (FG). — 331 trees in 70 plots; $F_{\max} = 2.9\%$; $dbh_{\text{inv}} = 87.9$ cm.

Genus *Platonia* Mart.

[379] *Platonia insignis* Mart.
(Fig. 18B)

Nova genera et species plantarum [Martius] 3 (3): 169 [“1829” publ. Sep. 1832] (Martius 1832), nom. cons.

Moronoea esculenta Arruda, *Trav. Brazil [H. Koster]* 490 (Arruda 1816), nom. rej. — *Symponia esculenta* Arruda ex Choisy, *Prod. [A. P. de Candolle]* 1: 563 [mid Jan. 1824] (Choisy 1824), nom. nud. pro syn. — *Symponia esculenta* (Arruda) Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 654 (Steudel 1841). — *Platonia esculenta* (Arruda) Oken, *Allg. Naturgesch.* 3 (2): 1431 (Oken 1841). — *Aristoclesia esculenta* (Arruda) Stuntz, *Bull. Bur. Pl. Industr. U.S.D.A.*

248: 58 (Stuntz 1912). — *Platonia esculenta* (Arruda) Rickett & Stafleu, *Taxon* 8: 313 (Rickett & Stafleu 1959), isonym.

NOTES. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). Choisy is the author of Gutiferae in Candolle’s *Prodromus*: his name appears at the end of the treatment (p. 564) indicating his authorship.

VERNACULAR NAMES. — Pa: wakukwa-tiranō-kamwi • Ka: pakuli • Wp: wila nānā kweli • Wn: wiipē • Nt: makasu, paykuli • Cr: parkouri • Fr: parcouri • Br: bacuri.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2575*.

INVENTORY DATA (FG). — 104 trees in 56 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 120$ cm.

Genus *Symponia* L.f.

[380] *Symponia globulifera* L.f.
(Fig. 18D)

Suppl. Pl. 302 [“1781” publ. Apr. 1782] (Linnaeus 1782). — *Moronoea globulifera* (L.f.) Schlehd., *Linnaea* 8: 190 (Schlechtendal 1833).

Aneuriscus aubletii C.Presl, *Symb. Bot. [C. Presl]* 1: 72 [Jan.-Feb. 1832] (Presl 1832).

Aneuriscus exserens C.Presl, *Symb. Bot. [C. Presl]* 1: 72 [Jan.-Feb. 1832] (Presl 1832).

Actinostigma speciosum Welw., *Apont.* 560 (Welwitsch 1859).

Symponia globulifera var. *macoubea* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 229 [Dec. 1893] (Vesque 1893). — *Moronoea macoubea* Miq. ex Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 229 [Dec. 1893] (Vesque 1893), nom. nud. pro syn.

Symponia globulifera var. *gabonensis* Vesque, *Monogr. Phan. [A.DC. & C.DC.]* 8: 231 [Dec. 1893] (Vesque 1893). — *Symponia gabonensis* (Vesque) Pierre, *Bull. Mens. Soc. Linn. Paris* 2 (155): 1228 (Pierre 1896).

Symponia microphylla R.E.Schult., *Bot. Mus. Leafl.* 17 (1): 20 [27 May 1955] (Schultes 1955), nom. illeg. hom., non (Hils. & Bojer ex Cambess.) Benth. & Hook.f. ex Vesque (1893).

Symponia utilissima R.E.Schult., *Bot. Mus. Leafl.* 17 (1): 22 [27 May 1955] (Schultes 1955).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ti-marikasmagene • Ka: ananiyu, mani, wesekapo epityi • Te: baytakini • Wp: wanani, wanāni • Wn: mani, mani epu, maniimē • Nt: kiiki mataki, sabana mataki, wataa mataki • Cr: manni-marikaj • Fr: manil marécage • Br: ananí.

HERBARIUM DATA (FG). — 113 collections at CAY. Sel. exs.: *M.-F. Prévost 4056*.

INVENTORY DATA (FG). — 154 trees in 50 plots; $F_{\max} = 2\%$; $dbh_{\text{inv}} = 120$ cm.




FIG. 18. — Clusiaceae: **A**, *Garcinia macrophylla* Mart. (D. Sabatier & J.-F. Molino 5703); **B**, *Platonia insignis* Mart. (D. Sabatier & M.-F. Prévost 2575); **C**, *Tovomita maxima* Molino & J. Engel; **D**, *Symphonia globulifera* L.f. (D. Sabatier & M.-F. Prévost 4381); **E**, *Tovomita schomburgkii* Planch. & Triana (J.-F. Molino et al. 3372). A-D, © D. Sabatier/IRD; E, © J.-F. Molino/IRD.

[381] *Symphonia* sp. A

VERNACULAR NAMES. — Pa: ti-wašiunu.

HERBARIUM DATA (FG). — 54 collections at CAY. Sel. exs.: [M.-F. Prévost 3527](#).

INVENTORY DATA (FG). — 907 trees in 162 plots; $F_{\max} = 4.2\%$; $dbh_{\text{inv}} = 83\text{ cm}$.

Genus *Tovomita* Aubl.

[382] *Tovomita brevistaminea* Engl.

Fl. Bras. [Martius] 12 (1): 446 [1 Apr. 1888] (Engler 1888).

Tovomita jenmanii Engl., *Fl. Bras. [Martius]* 12 (1): 447 [1 Apr. 1888] (Engler 1888), “jenmani”.

Tovomita triflora Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 436 (Huber 1902).

VERNACULAR NAMES. — Pa: kwatri-wašiune-duwē • Ka: alakapuli paindalyi, paipaiyo wokulu • Wp: pasi'i wapo sili • Cr: pativyé-montang • Br: manguerana, paxiubaraná.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: [M.-F. Prévost 4023](#).

INVENTORY DATA (FG). — 233 trees in 47 plots; $F_{\max} = 5.4\%$; $dbh_{\text{inv}} = 20.7\text{ cm}$.

[383] *Tovomita calodictyos* Sandwith

Bull. Misc. Inform. Kew 1936 (3): 219 [27 June 1936] (Sandwith 1936).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kwatri-washiune • Ka: alakapuli, wateipa upadyali • Cr: pativyé-montangn

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *S.A. Mori & T.D. Pennington 18056*.

SIZE. — Up to 28 m tall (Pipoly & Gustafsson 2002).

[384] *Tovomita caloneura* A.C.Sm.

Lloydia 2 (3): 197 (Smith 1939).

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3990*.

INVENTORY DATA (FG). — 193 trees in 62 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 28.3$ cm.

[385] *Tovomita fanshawei* Maguire

Bull. Torrey Bot. Club 75 (4): 436 [Jul.-Aug. 1948] (Maguire 1948).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2898*.

INVENTORY DATA (FG). — 152 trees in 48 plots; $F_{\max} = 3.2\%$; $dbh_{\text{inv}} = 30.2$ cm.

[386] *Tovomita gazelii* Poncy & Offroy

Adansonia, sér. 3, 28 (1): 114 [4 July 2006] (Poncy & Offroy 2006).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: busi mangu • Cr: pativyé-montangn • Fr: palétuvier montagne.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *O. Poncy et al. 1454* (holo-, P[P00266066, P00266068]; iso-, P[P00266067]).

INVENTORY DATA (FG). — 86 trees in 35 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 50.9$ cm.

[387] *Tovomita grata* Sandwith

Bull. Misc. Inform. Kew 1936 (3): 216 [27 June 1936] (Sandwith 1936).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3988*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.4$ cm.

[388] *Tovomita guianensis* Aubl.

Hist. Pl. Guiane 2: 957 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Marialva guianensis* (Aubl.) Choisy, *Mém. Soc. Hist. Nat. Paris*, sér. 2, 1: 224 (Choisy 1823), “*Tomovita* [sic] *guyanensis*”.

VERNACULAR NAMES. — Wp: pasi’i wapo sili • Cr: pativyé-montangn • Br: manguerana, paxiubarana.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000047393]).

INVENTORY DATA (FG). — 1 tree, dbh = 10.5 cm.

[389] *Tovomita longifolia* (Rich.) Hochr.

Annuaire Conserv. Jard. Bot. Genève 21: 66 (Hochreutiner 1919). — *Clusia longifolia* Rich., *Actes Soc. Hist. Nat. Paris* 1: 113 [Oct. 1792] (Richard 1792). — *Micranthera clusioides* Choisy, *Mém. Soc. Hist. Nat. Paris*, sér. 2, 1: 224 (Choisy 1823), excl. descr. masc. et tab. 11; nom. illeg. superfl. (based on *Clusia longifolia*). — *Tovomita clusiifolia* (Choisy) G.Don, *Gen. Hist.* 1: 617 [early Aug. 1831] (Don 1831), “*clusiaefolia*”, nom. illeg. superfl. (based on *Clusia longifolia*). — *Tovomita richardiana* Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 273 (Planchon & Triana 1860), nom. illeg. superfl. (based on *Clusia longifolia*).

Micranthera clusiifolia Choisy, *Prod. [A. P. de Candolle]* 1: 560 [mid Jan. 1824] (Choisy 1824), “*clusiaefolia*”.

Tovomita choisyana Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 14: 281 (Planchon & Triana 1860), syn. nov.

Tovomita bahiensis Engl., *Fl. Bras. [Martius]* 12 (1): 455 [1 Apr. 1888] (Engler 1888), syn. nov.

Tovomita melinonii Vesque, *Epharmosis* 3: 20 (Vesque 1892), “*melinoni*”, syn. nov.

Tovomita excelsa Andrade-Lima & G.Mariz, *Bull. Torrey Bot. Club* 101 (6): 367 [“1974” publ. 1975] (Andrade-Lima & Mariz 1975), syn. nov.

NOTES. — Hochreutiner’s stated (1919: 66) that “*L’original de Richard se trouve dans l’Herbier Delessert [...]*” (i.e. “Richard’s original material is in Delessert’s herbarium”) is misleading; it is not a Richard collection. Hochreutiner’s precise description and transcription of the labels (Hochreutiner 1919: 66) clearly match *Leblond* 34 (G[G00355740; P[P05061606]). *Micranthera clusiifolia* (“*clusiaefolia*”) is obviously a misprint for *M. clusioides* Choisy.

Following Planchon & Triana (1860b), Marinho *et al.* (2016) distinguished *Tovomita choisyana* from *T. longifolia* (called *T. richardiana* by the former authors); yet their analysis demonstrates that the two taxa do not differ in all the distinguishing characters proposed by Planchon & Triana: number of petals, leaf morphology and inflorescence structure. Marinho *et al.* (2016) placed emphasis on differences in stamen length and shape, and in habitat preference: the stamens and staminodes of *T. longifolia* (according to these authors) were only 3–4 mm long compared to 8–10 mm in *T. choisyana*. Furthermore, they considered that *T. choisyana* was “associated with watercourses”. These differences do not hold: Examination of *Leblond* 34 (P[P05061606, annotated by Marinho as “isotype” of *T. longifolia*) shows that some staminodes are 6 mm, conversely, several stamens of the isotype of *T. choisyana* at G-DC (*Unknown coll. s.n.*, G[G00210645) are less than 7 mm. Further, there is no information on the specimens identified as *T. choisyana* to indicate that they were collected in riparian, flooded, or swampy environments. We agree with Marinho *et al.* (2016) that *T. excelsa* Andrade-Lima & G.Mariz and *T. bahiensis* Engl. are conspecific with *T. choisyana*, and we therefore transfer them into synonymy under *T. longifolia*. The type of *T. melinonii* Vesque (*Mélinon* s.n. [holotype: P[P00093864, isotype: P[P00093865]) also perfectly fits in *T. longifolia*.

VERNACULAR NAMES. — Pa: kwatri-washiune • Ka: alakapuli, alaka-puli paindyali, paipaiyo wokulu, wateipa upadyali • Wp: pasi’i

wapo, pasí'i wapo sili • Wn: itukuyen • Nt: busi mangu, mangu • Cr: pativé-montangn • Fr: palétuvier montagne • Br: manguerana, paxiubarana.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *J.B. Leblond* 34 (lecto-, G[G00355740], designated by Marinho et al. [2016: 771]; isolecto-, MPU[MPU014282], P[P05061606]).

INVENTORY DATA (FG). — 204 trees in 65 plots; $F_{\max} = 4.3\%$; $\text{dbh}_{\text{inv}} = 38.5\text{ cm}$.

[390] *Tovomita mangle* G.Mariz

Bull. Torrey Bot. Club 101 (6): 367 ["1974" publ. 1975] (Mariz 1975).

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.-F. Molino* 1556.

INVENTORY DATA (FG). — 66 trees in 34 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 21.8\text{ cm}$.

[391] *Tovomita maxima* Molino & J.Engel
(Fig. 18C)

Adansonia, sér. 3, 44 (16): 166 (Engel et al. 2022).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier* & *E. Fonty* 5595 (holo-, CAY[CAY112114]).

INVENTORY DATA (FG). — 37 trees in 22 plots; $F_{\max} = 4.1\%$; $\text{dbh}_{\text{inv}} = 43\text{ cm}$.

[392] *Tovomita saulensis* J.Engel & Molino

Adansonia 44, sér. 3, (16): 170 (Engel et al. 2022).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori* et al. 24002 (holo-, CAY[CAY167624]; iso-, P[P04897763], NY[NY04204154], U[U0181523], US[US00872988]).

SIZE. — *L.R. Phillippe* et al. 26984, dbh 12 cm.

[393] *Tovomita schomburgkii* Planch. & Triana
(Fig. 18E)

Ann. Sci. Nat., Bot. sér. 4, 14: 274 (Planchon & Triana 1860).

Tovomita hameliifolia Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 996 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), "hameliaefolia", nom. subnud.

Tovomita schomburgkiana Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 996 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. subnud.

VERNACULAR NAMES. — Ka: wateipa upadyali • Wp: pasí'i wapo, pasí'i wapo sili • Cr: pativé-montangn • Br: mangue-da-mata, manguerana, paxiubarana.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *J.-F. Molino* et al. 3372.

INVENTORY DATA (FG). — 37 trees in 17 plots; $F_{\max} = 2.3\%$; $\text{dbh}_{\text{inv}} = 22\text{ cm}$.

[394] *Tovomita tenuiflora* Benth. ex Planch. & Triana

Ann. Sci. Nat., Bot. sér. 4, 14: 272 (Planchon & Triana 1860).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-F. Molino* & *D. Sabatier* 1948.

INVENTORY DATA (FG). — 32 trees in 21 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 38.8\text{ cm}$.

[395] *Tovomita umbellata* Benth.

London J. Bot. 2: 367 (Bentham 1843).

Tovomita stigmatosa Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14:* 275 (Planchon & Triana 1860).

Tovomita nigrescens Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4, 14:* 276 (Planchon & Triana 1860).

Tovomita cephalostigma Vesque, *Epharmosis* 3: 19 (Vesque 1892).

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *E.M. Mélion* s.n. & 435 (possibly the same gathering, original material of *Tovomita cephalostigma*: P[s.n.]: P00093869, P01901223; n° 435: P01901224, P01901225).

INVENTORY DATA (FG). — 109 trees in 35 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 27.7\text{ cm}$.

[396] *Tovomita weddelliana* Planch. & Triana

Ann. Sci. Nat., Bot. sér. 4, 14: 277 (Planchon & Triana 1860). — *Arawakia weddelliana* (Planch. & Triana) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Clusia oblanceolata Rusby, *Descr. S. Amer. Pl.* 58 [20 Dec. 1920] (Rusby 1920). — *Arawakia oblanceolata* (Rusby) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita longicuneata Engl., *Bot. Jahrb. Syst.* 58 (4, Beibl. 130): 7 [1 July 1923] (Engler 1923). — *Arawakia longicuneata* (Engl.) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita sphenophylla Diels, *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 32 [30 Mar. 1938] (Diels 1938). — *Arawakia sphenophylla* (Diels) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita lingulata Cuatrec., *Anales Inst. Biol. Univ. Nac. México* 20: 99. (Cuatrecasas 1950). — *Arawakia lingulata* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita macrocarpa Cuatrec., *Anales Inst. Biol. Univ. Nac. México* 20: 100. (Cuatrecasas 1950). — *Arawakia macrocarpa* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita rhizophoroides Cuatrec., *Anales Inst. Biol. Univ. Nac. México* 20: 101. (Cuatrecasas 1950). — *Arawakia rhizophoroides* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita lanceolata Cuatrec., Anales Inst. Biol. Univ. Nac. México 20: 102. (Cuatrecasas 1950). — *Arawakia lanceolata* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita glossophylla Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (29): 62 [Nov. 1950] (Cuatrecasas 1950). — *Arawakia glossophylla* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Clusia pithecobia Standl. & L.O.Williams, *Ceiba* 1 (4): 244 (Standley & Williams 1951). — *Tovomita pithecobia* (Standl. & L.O.Williams) Gahagen, *Syst. Bot.* 40 (4): 985 [15 Dec. 2015] (Gahagen 2015). — *Arawakia pithecobia* (Standl. & L.O.Williams) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita angustata Steyermark., *Fieldiana, Bot.* 28 (2): 399 (Steyermark 1952). — *Arawakia angustata* (Steyermark.) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita rileyi Cuatrec., *Brittonia* 14 (1): 52 (Cuatrecasas 1962). — *Arawakia rileyi* (Cuatrec.) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita coriacea Maguire, *Phytologia* 36 (4): 406 [22 Aug. 1977] (Maguire 1977). — *Arawakia coriacea* (Maguire) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita caputmonisia Gahagen, *Syst. Bot.* 40 (4): 975 [15 Dec. 2015] (Gahagen 2015). — *Arawakia caputmonisia* (Gahagen) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita divesora Gahagen, *Syst. Bot.* 40 (4): 975 [15 Dec. 2015] (Gahagen 2015). — *Arawakia divesora* (Gahagen) L.Marinho, *Molec. Phylogen. Evol.* 134: 147 [publ. online Feb. 2019] (Marinho 2019).

Tovomita manchamancha Gahagen, *Syst. Bot.* 40 (4): 982 [15 Dec. 2015] (Gahagen 2015). — *Arawakia manchamancha* (Gahagen) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita panamaea Gahagen, *Syst. Bot.* 40 (4): 983 [15 Dec. 2015] (Gahagen 2015). — *Arawakia panamaea* (Gahagen) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

Tovomita parvifolia Gahagen, *Syst. Bot.* 40 (4): 983 [15 Dec. 2015] (Gahagen 2015). — *Arawakia parvifolia* (Gahagen) L.Marinho, *Molec. Phylogen. Evol.* 134: 148 [publ. online Feb. 2019] (Marinho 2019).

NOTES. — Gahagen, Terbush & Ballard (2016) resurrected several names previously considered synonyms of *T. weddelliana*, and segregated 5 new species from what they called the “*T. weddelliana* species complex”. Their study relied on “quantitative” analyses of morphological characters, based on two to 11 herbarium specimens per species. In our opinion, the number of studied specimens is too low to capture the natural variability of the morphological characters (e.g. twig diameter, pedicel length or leaf blade size). Moreover, the results of their statistical analyses are unconvincing and do not clearly distinguish their 11 so-called species, whose morphological differences could simply reflect the plasticity of a single species found from the Pacific coast to the Tepuis of the Guyana and slopes of the Andes, from 0 to 1300 m altitude. Subsequently, Marinho *et al.* (2019) segregated the “*T. weddelliana* complex” in a new genus, *Arawakia*, on the basis of plastid and nuclear ITS genome analyses of 30 species out of the c. 500 that form the tribe Clusieae. However, their splitting of *Tovomita* into two main groups (one being *Arawakia*) appears much less supported (85-86 bootstrap values and 0.51-

0.66 posterior probabilities) than the monophyly of other genera (99-100 bootstrap/0.85-1 pp). Further, two of the five diagnostic characters that Marinho *et al.* (2019) claim to distinguish *Arawakia* from *Tovomita* do not hold: - white latex, said to be characteristic of *Arawakia* (vs yellow in *Tovomita*), has been observed by the first two authors of the present publication in *T. umbellata* (*J.-F Molino* 731 and *J.-F Molino* 936, CAY), *T. schomburgkii* (*D. Sabatier* & *M.-F. Prévost* 1345, *D. Sabatier* & *M.-F. Prévost* 3354 and *D. Sabatier* & *J.-F Molino* 4679, CAY) and *T. caloneura* (*D. Sabatier* & *M.-F. Prévost* 1363, CAY); - the sepals, said to be always persistent and adpressed on fruit in *Arawakia* (vs not adpressed in *Tovomita*) are actually not adpressed on fruits of the types of *A. lingulata*, *A. macrocarpa* and *A. rileyi*. The remaining characters Marinho *et al.* (2019) present do not warrant the creation of a new genus: two are purely vegetative (leaf shape and colour of the interpetiolar line, black in *Arawakia* vs not black in *Tovomita*), and the third is epicarp colour (red in *Arawakia* vs green to brown in *Tovomita*).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier* 5799.

SIZE. — Up to 30 m tall (Cuello 1998).

[397] *Tovomita* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier* 4872.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 18 cm.

Family COMBRETACEAE R.Br.
Genus *Conocarpus* L.

[398] *Conocarpus erectus* L. (Fig. 19A)

Sp. Pl. 1: 176 [1 May 1753] (Linnaeus 1753), “erecta”. — *Terminalia erecta* (L.) Baill., *Hist. Pl. [Baillon]* 6: 266, 275 [Jan.-May 1876] (Baillon 1876). — *Conocarpus erectus* var. *arboreus* DC., *Prodr. [A. P. de Candolle]* 3: 16 [mid Mar. 1828] (Candolle 1828), “erecta var. *arborea*”, nom. illeg. superfl. (based on the type of *Conocarpus erectus*).

Conocarpus procumbens L., *Sp. Pl.* 1: 177 [1 May 1753] (Linnaeus 1753). — *Conocarpus supinus* Crantz, *Inst. Rei Herb.* 1: 355 (Crantz 1766), nom. illeg. superfl. (based on *Conocarpus procumbens*). — *Conocarpus erectus* var. *procumbens* (L.) DC., *Prodr. [A. P. de Candolle]* 3: 16 [mid Mar. 1828] (Candolle 1828). — *Terminalia erecta* var. *procumbens* (L.) M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.*: 73 (Gómez 1889).

Conocarpus acutifolius Humb. & Bonpl. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 574 [Dec. 1819] (Roemer & Schultes 1819).

Conocarpus pubescens Schumach., *Beskr. Guin. Pl.*: 115 (Schumacher 1827).

NOTE. — A species restricted to mangrove forests.

VERNACULAR NAMES. — Br: mangue-de-botão, mangue-negro.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost* & *C. Proisy* 4550.

SIZE. — Up to 20 m tall (Stace 2010).

Genus *Laguncularia* C.F.Gaertn.

[399] *Laguncularia racemosa* (L.) C.F.Gaertn.

Suppl. Carp.: 209 (Gaertner 1807). — *Conocarpus racemosus* L., *Syst. Nat.*, ed. 10, 2: 930 [7 June 1759] (Linnaeus 1759). — *Horau racemosus* (L.) M.Gómez, *Fl. Cuba [Gómez & Roig]*: 75 (Gómez 1914).

Schousboea commutata Spreng., *Syst. Veg. [Sprengel]* 2: 332 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Conocarpus racemosus*).

Rhizaeris alba Raf., *Sylva Tellur.*: 90 (Rafinesque 1838), nom. illeg. superfl. (based on *Conocarpus racemosus*).

Laguncularia obovata Miq., *Linnaea* 18: 752 ["1844" publ. Aug.-Oct. 1845] (Miquel 1845).

Laguncularia racemosa f. *longifolia* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 8* (2): 125 (Macbride 1930).

NOTES. — A species restricted to mangrove forests. Rafinesque (1838: 90) described *Rhizaeris* as a new genus "very diff. from *Conocarpus*", then "*R. alba* Raf., *Conocarpus racemosa* L. auct. The white Mangrove of Antilles". He apparently thought that *C. racemosus* L. was indeed a *Conocarpus*, that this name had been wrongly used for the white mangrove, and accordingly he proposed a new name. He did not realize that *C. racemosus* is actually the white mangrove and had already been transferred to *Laguncularia* (Gaertner 1807).

VERNACULAR NAMES. — Ka: akila'i, ele'u, elewu • Cr: pativé-gri • Fr: palétuvier gris • Br: falso-mangue, mangue-amarelo, mangue-branco.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *D. Sabatier 2259*, dbh 10 cm.

Genus *Terminalia* L.

[400] *Terminalia amazonia* (J.F.Gmel.) Exell
(Fig. 19B)

Fl. Suriname 3 (1): 173 (Exell 1935). — *Chuncoa amazonia* J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 702 [late Sep.-Nov. 1791] (Gmelin 1791).

Gimbernatia obovata Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 274 [late Dec. 1798] (Ruiz & Pavón 1798). — *Chuncoa obovata* (Ruiz & Pav.) Pers., *Syn. Pl. [Persoon]* 1: 486 [1 Apr.-15 June 1805] (Persoon 1805), "Chunchoa". — *Terminalia obovata* (Ruiz & Pav.) Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 668 (Steudel 1841), nom. illeg. hom., non Cambess. (Cambessèdes 1830). — *Myrobalanus obovata* (Ruiz & Pav.) Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891).

Bucida angustifolia DC., *Prodri. [A. P. de Candolle]* 3: 10 [mid Mar. 1828] (Candolle 1828). — *Bucida buceras* L. var. *angustifolia* (DC.) Eichler, *Fl. Bras. [Martius]* 14 (2): 95 [17 Apr. 1867] (Eichler 1867).

Terminalia odontoptera Van Heurck & Müll.Arg., *Observ. Bot. [Van Heurck]* 2: 217 [10 Sep. 1871] (Van Heurck & Müller 1871).

Terminalia excelsa Liebm. ex Hemsl., *Biol. Cent.-Amer., Bot.* 1 (5): 402 [June 1880] (Hemsley 1880), nom. nud.

Myrobalanus excelsa Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: katauma, katumá • Ka: ityutano kwai, katulimya, kwai • Wp: alala munuwi sili, tatulimá • Nt: anangosi, kwata pepee • Cr: angouchi, bwa-blanché, grènn-oko • Br: ciliarana, mirindiba-branca.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *G.S. Perrottet s.n.* (original material of *Bucida angustifolia*: G[G00236066]).

INVENTORY DATA (FG). — 61 trees in 24 plots; $F_{\max} = 1.9\%$; $dbh_{inv} = 71$ cm.

[401] *Terminalia aubletii* Gere & Boatwr.
(Fig. 19C)

Bot. J. Linn. Soc. 184 (3): 319 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Pamea guianensis* Aubl., *Hist. Pl. Guiane* 2: 946 [Jun.-Dec. 1775] (Aublet 1775). — *Terminalia pamea* DC., *Prodri. [A. P. de Candolle]* 3: 13 [mid Mar. 1828] (Candolle 1828), nom. illeg. superfl. (based on *Pamea guianensis*). — *Myrobalanus pamea* Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891), "Pamaea". — *Buchenavia guianensis* (Aubl.) Alwan & Stace, *Nordic J. Bot.* 5 (5): 447 (Alwan & Stace 1985).

NOTES. — *Myrobalanus pamea* Kuntze is based on the illegitimate *Terminalia pamea* DC., but is legitimate as a replacement name for *Pamea guianensis* Aubl. (which is indirectly cited through *T. pamea*), non *M. guianensis* (Aubl.) Kuntze (based on *Tanibouca guianensis* Aubl., and synonym of *Terminalia dichotoma* [1818: 177]).

VERNACULAR NAMES. — Ka: katulimya • Wp: kwata kaya, tatulimá sili • Nt: anangosi • Cr: zanmann-sovaj • Fr: amandier sauvage • Br: tanimbucá.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Pamea guianensis*: BM[BM000953713]).

INVENTORY DATA (FG). — 17 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{inv} = 101$ cm.

[402] *Terminalia carinata* Sabatier & J.Engel
(Fig. 19D)

Adansonia, sér. 3, 42 (16): 262 [2 Nov. 2020] (Engel & Sabatier 2020).

VERNACULAR NAMES. — Pa: karevru-danó • Ka: kalalawa akunepili • Nt: anangosi, kwata pepe, kwata pepee • Cr: angouchi, bwa-blanché, grènn-oko • Fr: amandier sauvage • Br: mirindiba-branca.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *D. Sabatier et al. 4891* (holo-, CAY[CAY182887]; iso-, K, P[P01155925]).

INVENTORY DATA (FG). — 56 trees in 32 plots; $F_{\max} < 1\%$; $dbh_{inv} = 120$ cm.

[403] *Terminalia catappa* L.

Mant. Pl. 1: 128 (Linnaeus 1767). — *Juglans catappa* (L.) Lour., *Fl. Cochinch.* 2: 573 [Sep. 1790] (Loureiro 1790). — *Myrobalanus catappa* (L.) Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891). — *Buceras catappa* (L.) Hitchc., *Rep. (Annual) Missouri Bot. Gard.* 4: 85 (Hitchcock 1893).

Phytolacca javanica Osbeck, *Dagb. Ostind. Resa*: 276 (Osbeck 1757).

Terminalia moluccana Lam., *Encycl. [J. Lamarck et al.] 1 (2)*: 349 [1 Aug. 1785] (Lamarck 1785).

Terminalia ovatifolia Noronha, *Verh. Batav. Genootsch. Kunst. 5 (4)*: 27 (Noronha 1790).

Badamia commersonii Gaertn., *Fruct. Sem. Pl. 2*: 90 [Sep.-Nov. 1790] (Gaertner 1790), “*Commersonii*”.

Terminalia subcordata Humb. & Bonpl. ex Willd., *Sp. Pl., ed. 4 4 (2)*: 968 [Apr. 1806] (Willdenow 1806). — *Terminalia catappa* var. *subcordata* (Humb. & Bonpl. ex Willd.) DC., *Prodr. [A. P. de Candolle] 3*: 11 [mid Mar. 1828] (Candolle 1828).

Myrobalanus badamia Poir., *Encycl. [J. Lamarck et al.] Suppl. 3*: 707 [3 Sep. 1814] (Poiret 1814), “*Mirobolanus*”. — *Terminalia badamia* (Poir.) DC., *Prodr. [A. P. de Candolle] 3*: 12 [mid Mar. 1828] (Candolle 1828).

Myrobalanus terminalia Poir., *Encycl. [J. Lamarck et al.] Suppl. 3*: 707 [3 Sep. 1814] (Poiret 1814), “*Mirobolanus*”.

Terminalia myrobalana Roth, *Nov. Pl. Sp. 378* [Apr. 1821] (Roth 1821).

Terminalia intermedia Bertero ex Spreng., *Syst. Veg. [Sprengel] 2*: 359 [Jan.-May 1825] (Sprengel 1825).

Terminalia kydiana Roxb. ex Wall., *Numer. List no. 3974* (Wallich 1831).

Terminalia latifolia Blanco, *Fl. Filip. [FM. Blanco] 376* (Blanco 1837).

Terminalia paraensis Mart., *Flora 24* (2, Beibl.): 24 (Martius 1841).

Terminalia mauritiana Blanco, *Fl. Filip. [FM. Blanco], ed. 2*: 264 (Blanco 1845).

Terminalia rubrigemmis Tul., *Ann. Sci. Nat., Bot. sér. 4, 6*: 102 (Tulasne 1856). — *Myrobalanus rubrigemmis* (Tul.) Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891).

Myrobalanus commersonii Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891).

NOTE. — A species native to South-West Asia, introduced and naturalised in most tropical regions.

VERNACULAR NAMES. — Pa: zamād, zaman • Ka: amandra • Nt: amanda, amandaa • Cr: zanmann • Fr: amandier pays • Br: amen-doeira-da-praia.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *S.A. Mori et al. 25515*.

SIZE. — Up to 35 m tall (Stace 2010).

[404] *Terminalia dichotoma* G.Mey.

Prim. Fl. Esseq. 177 [Nov. 1818] (Meyer 1818). — *Terminalia latifolia* Sw. var. *dichotoma* (G.Mey.) DC., *Prodr. [A. P. de Candolle] 3*: 12 [mid Mar. 1828] (Candolle 1828).

Tanibouca guianensis Aubl., *Hist. Pl. Guiane 1*: 448 [Jun.-Dec. 1775] (Aublet 1775). — *Myrobalanus guianensis* (Aubl.) Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891). — *Terminalia tanibouca* Rich., *Actes Soc. Hist. Nat. Paris 1*: 109 [Oct. 1792] (Richard 1792), nom. illeg. superfl. (based on *Tanibouca guianensis*).

Catappa guianensis C.F.Gaertn., *Suppl. Carp. 207* (Gaertner 1807).

NOTE. — According to Stace (2010), *Catappa guianensis* Gaertn. is not based on *Tanibouca guianensis* Aubl.

VERNACULAR NAMES. — Ka: kalalawa akunepili • Wp: alala munuwi • Cr: angouchi, grènn-oko • Br: cinzeiro, tanimbucá.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Tanibouca guianensis*: BM[BM000953723]).

INVENTORY DATA (FG). — 13 trees in 5 plots; $F_{\max} < 1\%$; dbh_{inv} = 59 cm.

[405] *Terminalia grandis* (Ducke) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 320 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia grandis* Ducke, *Arch. Jard. Bot. Rio de Janeiro 4*: 148 (Ducke 1925).

Buchenavia huberi Ducke, *Bol. Técn. Inst. Agron. N. 4*: 24 [31 Mar. 1945] (Ducke 1945).

VERNACULAR NAMES. — Nt: anangosi • Cr: zanmann-sovaj • Fr: amandier sauvage • Br: marindiba, tanimbucá.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4829*.

INVENTORY DATA (FG). — 23 trees in 19 plots; $F_{\max} < 1\%$; dbh_{inv} = 60.5 cm.

[406] *Terminalia guyanensis* Eichler

Fl. Bras. [Martius] 14 (2): 88 [17 Apr. 1867] (Eichler 1867).

Myrobalanus eichleriana Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holo-, B[not seen, photo F neg. N° 14333]; iso-, GH[GH00068628], K[K000640657], P[P01901252, P01901253, P01901254], U[U0001197]).

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 80 cm.

[407] *Terminalia lucida* Hoffmanns. ex Mart. & Zucc.

Flora 7 (1, Beibl.): 130 [25 Apr. 1824] (Martius & Zuccarini 1824). — *Myrobalanus lucida* (Hoffmanns. ex Mart. & Zucc.) Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891).

Terminalia firma Mart. & Zucc., *Flora 7 (1, Beibl.)*: 130 [25 Apr. 1824] (Martius & Zuccarini 1824).

Terminalia eriantha Benth., *Hooker's J. Bot. Kew Gard. Misc. 2*: 240 (Bentham 1850).

Terminalia scutifera Planch. ex M.A.Lawson, *Fl. Trop. Afr. 2*: 417 (Lawson 1871). — *Myrobalanus scutifera* (Planch. ex M.A.Lawson) Kuntze, *Revis. Gen. Pl. 1*: 237 [5 Nov. 1891] (Kuntze 1891).

Terminalia hayesii Pittier, *Contr. U.S. Natl. Herb. 18 (6)*: 239 [22 Sep. 1917] (Pittier 1917).

Terminalia nyssifolia Britton, *Bull. Torrey Bot. Club 48 (12)*: 333 [“1921” publ. 1922] (Britton 1922), “*nyssafolia*”.




FIG. 19. — Combretaceae: **A**, *Conocarpus erectus* L. (M.-F. Prévost & S. Gonzalez 5313); **B**, *Terminalia amazonica* (J.F.Gmel.) Exell; **C**, *Terminalia aubletii* Gere & Boatwr. (D. Sabatier 2283); **D**, *Terminalia carinata* Sabatier & J.Engel (D. Sabatier et al. 4891). © D. Sabatier/IRD.

NOTES. — Martius (1824: 44) cited the collection locality of *T. firma* Mart. & Zucc. as “*Habitat in editis montis Serra de Araracuara ad fluvium Japurá, Provinciae flum. nigri.*” Stace (2010) cited the locality as being situated in Brazil (Amazonas), but in January 1820 Martius collected the original gathering in Araracuara, on the Rio Caquetá, which is in Colombia.

VERNACULAR NAMES. — Wp: alala munuwi • Cr: angouchi, bwablanché, grènn-oko • Br: cinzeiro, tanimbucá.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost 1221; M.-F. Prévost & D. Sabatier 4895*, dbh 12 cm.

[408] *Terminalia macrophylla* (Eichler) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 320 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia macrophylla* Eichler, *Flora* 49 (11): 166 (Eichler 1866). — *Terminalia macrophylla* Spruce ex Eichler, *Fl. Bras. [Martius]* 14 (2): 98 [17 Apr. 1867] (Eichler 1867), *nom. nud. pro syn.*

Buchenavia stellae Cuatrec., *Fieldiana, Bot.* 27 (1): 109 (Cuatrecasas 1950).

VERNACULAR NAMES. — Ka: katulimya • Wp: kwata kaya, tatulimā sili • Wn: alimi huhu, ekesima, ekesimē • Nt: kwata bobi • Cr: tapouliman, zanmann-sovaj • Br: tanimbucá.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5017*, dbh 43 cm.

[409] *Terminalia megalophylla* (Van Heurck & Müll.Arg.) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 320 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia megalophylla* Van Heurck & Müll.Arg., *Observ. Bot. [Van Heurck]* 2: 211 [10 Sep. 1871] (Van Heurck & Müller 1871).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville B-4792*.

SIZE. — Up to 45 m tall (Stace 2009).

[410] *Terminalia nitidissima* Rich.

Actes Soc. Hist. Nat. Paris 1: 109 [Oct. 1792] (Richard 1792). — *Myrobalanus nitidissima* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891). — *Buchenavia nitidissima* (Rich.) Alwan & Stace, *Nordic J. Bot.* 5 (5): 449 (Alwan & Stace 1985).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (original material G[G00177930]).

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; dbh_{inv} = 32.9 cm.

[411] *Terminalia oblonga* (Ruiz & Pav.) Steud.

Nomencl. Bot. [Steudel], ed. 2, 2: 668 (Steudel 1841). — *Gimbernacia oblonga* Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 275 [late Dec. 1798] (Ruiz & Pavón 1798). — *Chuncoa oblonga* (Ruiz & Pav.) Pers., *Syn. Pl. [Persoon]* 1: 486 [1 Apr.-15 June 1805] (Persoon 1805), “*Chunchoa*”. — *Myrobalanus oblonga* (Ruiz & Pav.) Kuntze, *Revis. Gen. Pl.* 1: 237 [5 Nov. 1891] (Kuntze 1891). — *Chuncoa diptera* F.Dietr., *Nachtr. Vollst. Lex. Gärtn.* 2: 244 [May 1816] (Dietrich 1816), nom. illeg. superfl. (based on *Gimbernacia oblonga*).

Terminalia tarapotensis Van Heurck & Müll.Arg., *Observ. Bot. [Van Heurck]* 2: 213 [10 Sep. 1871] (Van Heurck & Müller 1871).

Terminalia chiriquensis Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 238 [22 Sep. 1917] (Pittier 1917).

Terminalia obidensis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 147 (Ducke 1925).

VERNACULAR NAMES. — Wp: alala munuwi • Cr: angouchi, grènnoko • Br: cinzeiro, tanimbucá.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Grand 1303*.

SIZE. — Brazil, Pará. *D.C. Daly 1645* (MO), 20 m × 40 cm.

[412] *Terminalia ochroprumna* (Eichler) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 321 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia ochroprumna* Eichler, *Flora* 49 (11): 165 (Eichler 1866).

Buchenavia discolor Diels, *Verb. Bot. Vereins Prov. Brandenburg* 48: 192 [“1906” publ. 8 Mar. 1907] (Diels 1907).

VERNACULAR NAMES. — Br: tanimbucá-do-igapó, tanimbucá-rana.

HERBARIUM DATA (FG). — A single collection, *P. Béna 1313*.

SIZE. — Up to 30 m tall (Stace 2009).

[413] *Terminalia parvifolia* (Ducke) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 321 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia parvifolia* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 150 (Ducke 1925).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier 2300*.

SIZE. — Brazil, Rondônia. *G.T. Prance et al. 8551* (P), 15 m × 10 cm.

[414] *Terminalia tetraphylla* (Aubl.) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 322 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Cordia tetraphylla* Aubl., *Hist. Pl. Guiane* 1: 224 [Jun.-Dec. 1775] (Aublet 1775). — *Firensia lutea* Raf., *Sylva Tellur.* 40 (Rafinesque 1838), nom. illeg. superfl. (based on *Cordia tetraphylla*). — *Lithocardium tetraphyllum* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Buchenavia tetraphylla* (Aubl.) R.A.Howard, *J. Arnold Arbor.* 64 (2): 266 [Apr. 1983] (Howard 1983). — *Gerascanthus tetraphyllus* (Aubl.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

Bucida capitata Vahl, *Ectog. Amer.* 1: 50 (Vahl 1797). — *Buchenavia capitata* (Vahl) Eichler, *Flora* 49 (11): 165 (Eichler 1866). — *Terminalia capitata* (Vahl) C.Wright, *Anales Acad. Ci. Med. Habana* 5: 410 (Wright 1869).

Hudsonia arborea A.Rob. ex Lunan, *Hort. Jamaic.* 2: 310 (Lunan 1814).

Terminalia obovata Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (17): 241 [“1829” publ. 4 Dec. 1830] (Cambessèdes 1830). — *Terminalia hilariana* Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 668 (Steudel 1841), nom. illeg. superfl. (based on *Terminalia obovata*).

Bucida angustifolia Spruce ex Eichler, *Flora* 49 (11): 165 (Eichler 1866), nom. illeg. hom., non DC. (Candolle 1828).

Pseudolmedia bucidifolia Bello, *Anales Soc. Esp. Hist. Nat.* 12: 109 (Bello 1883), “*bucidaefolia*”.

Buchenavia gracilis Glaz., *Bull. Soc. Bot. France* 54 (Mém. 3c): 203 [“1907” publ. 22 Feb. 1908] (Glaziou 1908), nom. nud.

Buchenavia macabensis Glaz., *Bull. Soc. Bot. France* 54 (Mém. 3c): 203 [“1907” publ. 22 Feb. 1908] (Glaziou 1908), nom. nud.

Buchenavia vaupesana Cuatrec., *Fieldiana, Bot.* 27 (1): 108 (Cuatrecasas 1950).

Buchenavia ptariensis Steyermark., *Fieldiana, Bot.* 28 (2): 423 (Steyermark 1952).

VERNACULAR NAMES. — Ka: katulimya • Wp: kwata kaya, tatulimá sili • Cr: tapouliman, zanmann-sovaj • Br: cuarana, mirindiba.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Loutry 1934*.

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; dbh_{inv} = 75 cm.

[415] *Terminalia viridiflora* (Ducke) Gere & Boatwr.

Bot. J. Linn. Soc. 184 (3): 322 [epubl. 22 June 2017] (Gere & Boatwright 2017). — *Buchenavia viridiflora* Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 63 [Sep. 1935] (Ducke 1935).

Buchenavia amazonia Alwan & Stace, *Ann. Missouri Bot. Gard.* 76 (4): 1127 (Alwan & Stace 1989).

VERNACULAR NAMES. — Pa: wakaba • Wp: tatulimá • Cr: zanmann-dan-bwa.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P. Grand 1776*.

INVENTORY DATA (FG). — 2 trees in 2 plots; dbh_{inv} = 90 cm.

Family CORDIACEAE R.Br. ex Dumort.
Genus *Cordia* L.

[416] *Cordia alliodora* (Ruiz & Pav.) Oken

Allg. Naturgesch. 3 (2): 1098 (Oken 1841). — *Cerdana alliodora* Ruiz & Pav., *Flora Peruviana* 2: 47 (Ruiz & Pavón 1799). — *Cordia alliodora* (Ruiz & Pav.) Cham., *Linnaea* 8 (2): 121 (Chamisso 1833), nom. inval. (genus name and specific epithet not associated [Turland et al. 2018: Art. 35.2]). — *Lithocardium alliodorum* (Ruiz & Pav.) Kuntze, *Revis. Gen. Pl.* 2: 976 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus alliodorus* (Ruiz & Pav.) Kuhlmann & Mattos, *Loefgrenia* 47: 1 (Kuhlmann & Mattos 1970), “*alliodiora*”.

VERNACULAR NAMES. — Nt: wataa kanu • Cr: sèd-sam • Fr: cèdre sam • Br: freijó, freijó-branco, louro-alho.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *S.A. Mori* et al. 24798.

SIZE. — Up to 40 m tall (Gaviria 1997).

[417] *Cordia bicolor* A.DC.

Prod. [A. P. de Candolle] 9: 485 [1 Jan. 1845] (Candolle 1845). — *Lithocardium bicolor* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 976 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus bicolor* (A.DC.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 399 (Borhidi 1988).

Lithocardium lockhartii Kuntze, *Revis. Gen. Pl.* 2: 438 [5 Nov. 1891] (Kuntze 1891). — *Cordia lockhartii* Kuntze, *Revis. Gen. Pl.* 2: 438 [5 Nov. 1891] (Kuntze 1891), “*lockartii*”, nom. nud. pro syn.

Cordia trichostyla Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 252 [22 Sep. 1917] (Pittier 1917).

Cordia carnosa Rusby, *Descr. S. Amer. Pl.* 104 [20 Dec. 1920] (Rusby 1920).

Cordia belizensis Lundell, *Amer. Midl. Naturalist* 29 (2): 488 [5 Apr. 1943] (Lundell 1943).

NOTE. — Although the “Boragineae” of the *Prodromus* were treated by A. P. de Candolle himself, note 1 on p. 466 states that his son Alphonse provided annotations and additions, among them several species whose names and descriptions are explicitly ascribed to “Alph. DC.”

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost* 169.

INVENTORY DATA (FG). — 40 trees in 21 plots; F_{max} < 1 %; dbh_{inv} = 83 cm.

[418] *Cordia dentata* Poir.

Encycl. [J. Lamarck et al.] 7: 48 [6 July 1806] (Poiret 1806). — *Carpiphea dentata* (Poir.) Raf., *Sylva Tellur.*: 39 (Rafinesque 1838).

Cordia calyprata Bertero ex Spreng., *Syst. Veg. [Sprengel]* 1: 649 [“1825” publ. late 1824] (Sprengel 1824). — *Varronia calyprata* (Bertero ex Spreng.) DC., *Prod. [A. P. de Candolle]* 9: 469 [1 Jan. 1845] (Candolle 1845).

Varronia calyprata var. *hartweggii* DC., *Prod. [A. P. de Candolle]* 9: 469 [1 Jan. 1845] (Candolle 1845).

Cordia tenuifolia Bertol., *Misc. Bot. [Bertol.]* 21: 13 (Bertoloni 1860). — *Lithocardium tenuifolium* (Bertol.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia leptopoda K.Krause, *Bot. Jahrb. Syst.* 37 (5): 628 [30 Oct. 1906] (Krause 1906).

Cordia ovata Brandegee, *Univ. Calif. Publ. Bot.* 10: 187 (Brandegee 1922).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost* 4476.

SIZE. — Up to 25 m tall (Nowicke 1969).

[419] *Cordia exaltata* Lam.

Tabl. Encycl. 1[2 (1)]: 422 [30 July 1792] (Lamarck 1792). — *Lithocardium exaltatum* (Lam.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus exaltatus* (Lam.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 399 (Borhidi 1988).

Cordia melanoneura Klotzsch ex M.R.Schomb., *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 960 [“1848” publ. 7-10 Mar. 1849] (Schomburgk 1849), nom. nud.

Cordia scabrida Mart. ex Fresen., *Fl. Bras. [Martius]* 8 (1): 11 [28 Feb. 1857] (Fresenius 1857). — *Lithocardium scabridum* (Mart. ex Fresen.) Kuntze, *Revis. Gen. Pl.* 2: 439 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus scabridus* (Mart. ex Fresen.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

Cordia exaltata var. *melanoneura* I.M.Johnst., *J. Arnold Arbor.* 16 (1): 30 [25 Jan. 1935] (Johnston 1935).

VERNACULAR NAMES. — Wp: kuluwa’i, kuluwa’i sili • Wn: mojoi • Nt: busi kiikii • Cr: bwa-sip • Br: cauarú-caá, chapéu-de-sol.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *L.C. Richards* s.n. (holo-, P[P00633485]; iso-, P[P00633484, P03892075, P03892077]).

INVENTORY DATA (FG). — 4 trees in 2 plots; F_{max} < 1 %; dbh_{inv} = 37 cm.

[420] *Cordia fallax* I.M.Johnst.

J. Arnold Arbor. 16 (1): 10 [25 Jan. 1935] (Johnston 1935).

Cordia guianensis Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 960 [“1848” publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. illeg. hom., non (Desv.) Roem. & Schult. (Roemer & Schultes 1819).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *G. Cremers* 8385.

SIZE. — Up to 35 m tall (Gaviria 1997).

[421] *Cordia fulva* I.M.Johnst.

J. Arnold Arbor. 16 (1): 20 [25 Jan. 1935] (Johnston 1935).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *E.M. Mélinton* 137, 1877 (holo-, GH[[GH00095576](#)]; iso-, GH[[GH00095577](#)]).

INVENTORY DATA (FG). — 1 tree, dbh = 27.7 cm.

[422] *Cordia goeldiana* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 6: 89 (Huber 1910). — *Gerascanthus goeldianus* (Huber) M.Kuhlm. & Mattos, *Loefgrenia* 47: 2 (Kuhlmann & Mattos 1970).

VERNACULAR NAMES. — Wp: ka'i tuli (?) • Br: freijó, frei-jorge.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Larpin* 946.

INVENTORY DATA (FG). — 22 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 85$ cm.

[423] *Cordia laevifrons* I.M.Johnst.
(Fig. 20A)

J. Arnold Arbor. 16 (1): 16 [25 Jan. 1935] (Johnston 1935).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5647.

INVENTORY DATA (FG). — 7 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.1$ cm.

[424] *Cordia lomatoloba* I.M.Johnst.

J. Arnold Arbor. 18 (1): 12 [22 Jan. 1937] (Johnston 1937). — *Gerascanthus lomatolobus* (I.M.Johnst.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 400 (Borhidi 1988).

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4221.

INVENTORY DATA (FG). — 35 trees in 18 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 83$ cm.

[425] *Cordia naidophila* I.M.Johnst.

J. Arnold Arbor. 17 (1): 32 [24 Jan. 1936] (Johnston 1936). — *Gerascanthus naidophilus* (I.M.Johnst.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 400 (Borhidi 1988).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost* 178.

SIZE. — Up to 15 m tall (Gaviria 1997).

[426] *Cordia nervosa* Lam.

Tabl. Encycl. 1[2 (1)]: 422 [30 July 1792] (Lamarck 1792). — *Lithocardium nervosum* (Lam.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia calophylla Vahl, *Eclog. Amer.* 3: 5 (Vahl 1807). — *Gerascanthus calophyllus* (Vahl) Borhidi, *Acta Bot. Hung.* 34 (3-4): 399 (Borhidi

1988). — *Lithocardium calophyllum* (Vahl) Kuntze, *Revis. Gen. Pl.* 2: 976 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (original material P[[P03892659](#)]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 18.8$ cm.

[427] *Cordia nodosa* Lam.

Tabl. Encycl. 1[2 (1)]: 422 [30 July 1792] (Lamarck 1792). — *Cordia collococa* Aubl., *Hist. Pl. Guiane* 1: 219 [Jun.-Dec. 1775] (Aublet 1775), “*collo-cocus*” on plate, *nom. illeg. hom., non Cordia collococa* L. (Linnaeus 1759). — *Cordia hirsuta* Willd., *Sp. Pl.*, ed. 4 1 (2): 1076 [July 1798] (Willdenow 1798), *nom. illeg. superfl.* (based on the illegitimate *Cordia collococa*, for which the replacement name *C. nodosa* already existed). — *Cordia formicarum* Roem. & Schult., *Syst. Veg.* [*Roemer & Schultes*] 4: 800 [Mar.-June 1819] (Roemer & Schultes 1819). — *Cordia nodosa* var. *hispidissima* Fresen., *Fl. Bras.* [*Martius*] 8 (1): 17 [28 Feb. 1857] (Fresenius 1857). — *Cordia nodosa* var. *glabrior* Fresen., *Fl. Bras.* [*Martius*] 8 (1): 16 [28 Feb. 1857] (Fresenius 1857). — *Lithocardium nodosum* (Lam.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Firensia hirsuta Raf., *Sylva Tellur.*: 40 (Rafinesque 1838), *nom. illeg. superfl.* (based on *C. hirsuta* and *Cordia collococa*, for which the replacement name *C. nodosa* already existed).

Cordia hispidissima DC., *Prodr. [A. P. de Candolle]* 9: 475 [1 Jan. 1845] (Candolle 1845). — *Lithocardium hispidissimum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia miranda DC., *Prodr. [A. P. de Candolle]* 9: 475 [1 Jan. 1845] (Candolle 1845). — *Lithocardium mirandum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia nodosa var. *angustifolia* Fresen., *Fl. Bras.* [*Martius*] 8 (1): 17 [28 Feb. 1857] (Fresenius 1857).

Cordia umbrosa Spruce ex Rusby, *Bull. Torrey Bot. Club* 26 (3): 147 [18 Mar. 1899] (Rusby 1899).

Cordia volubilis Pittier, *J. Wash. Acad. Sci.* 19: 184 (Pittier 1929).

NOTE. — Var. *glabrior* is based on *C. formicarum*, and var. *hispidissima* is based on *C. collococa*.

VERNACULAR NAMES. — Pa: kasiu-van, wiwis-kasiuvan • Ka: aluko wonale, awale emulutano, awale mulu, awata epi • Wp: yawatai • Wn: jenékapo, siktukahmit • Nt: wasiwasí uwii, wasiwasí wiwii • Cr: bwa-fronmi, lanmoussé-fronmi • Br: arua-felpudo, graó-de-galo, pau-de-formiga.

HERBARIUM DATA (FG). — 145 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material P-JJR, [P00777834](#)).

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 11.5$ cm.

[428] *Cordia panicularis* Rudge

Pl. Guian. [Rudge] 1 (4): 30 [Apr.-May 1806] (Rudge 1806). — *Lithocardium paniculare* (Rudge) Kuntze, *Revis. Gen. Pl.* 2: 976 [5 Nov. 1891] (Kuntze 1891). — *Gerascanthus panicularis* (Rudge)

Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988), *nom. inval.* (basionym citation incomplete).

VERNACULAR NAMES. — Te: tapi' ilupabi hun.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J. Martin* 151 (holo-, K[K000583350]; iso-, BM[BM000953118], GH[GH00095780], P[P00634041, P00634042]).

INVENTORY DATA (FG). — 13 trees in 6 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 19.4\text{ cm}$.

[429] *Cordia sagotii* I.M.Johnst.
(Fig. 20B)

J. Arnold Arbor. 16 (1): 27 [25 Jan. 1935] (Johnston 1935). — *Gerasanthus sagotii* (I.M.Johnst.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988).

Cordia coriacea Sagot ex Benoist, *Arch. Bot.* 5 (Mém. 1): 257 [27 Mar. 1933] (Benoist 1933), *nom. illeg. hom.*, *non* Killip (1927).

Cordia hirta I.M.Johnst., *J. Arnold Arbor.* 16 (1): 29 [25 Jan. 1935] (Johnston 1935). — *Gerasanthus hirtus* (I.M.Johnst.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988).

NOTE. — A label affixed by J. Gaviria in 1987 on a K specimen, K000583352, wrongly claims that it is the holotype of *Cordia sagotii*. Actually, Johnston (1935: 28) wrote: “Acarouany [...], Dec. 1856, Sagot 447 (TYPE of *C. coriacea* Sagot and *C. Sagotii*, Paris)”. This unambiguously designates as holotype the specimen P00634059. This is furthermore confirmed by two labels in Johnston’s hand on P00634059. All other Sagot material bearing the same number 447, but dated “1855” or “1857” (including K000583352), are probably syntypes.

VERNACULAR NAMES. — Pa: imavui-kamwi, wiwis-ahavukunō, wiwis-priyo, wiwis-purubumna • Ka: alatuluka, kulebogo tikalayi, tagulewe itudyano • Wp: kuluwa'i • Nt: busi kiikii • Cr: bwa-parasol • Fr: arbre parasol, bois parasol.

HERBARIUM DATA (FG). — 59 collections at CAY. Sel. exs.: *P.A. Sagot* 447, Dec. 1856 (holo-, P[P00634059]; iso-, BM[BM000906178], K[K000583351], P[P03860113]).

INVENTORY DATA (FG). — 170 trees in 93 plots; $F_{\max} = 2.6\%$; $\text{dbh}_{\text{inv}} = 53.7\text{ cm}$.

[430] *Cordia sericicalyx* A.DC.

Prodr. [A. P. de Candolle] 9: 485 [1 Jan. 1845] (Candolle 1845). — *Lithocardium sericicalyx* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus sericicalyx* (A.DC.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

Cordia sericicalyx A.DC. var. *latifolia* Miq., *Stirp. Surinam. Select.*: 140 [1850 publ. Mar. 1851] (Miquel 1851).

Cordia opaca Rusby, *Descr. S. Amer. Pl.* 104 [20 Dec. 1920] (Rusby 1920). — *Gerasanthus opacus* (Rusby) Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988).

Bourreria viridis Rusby, *Descr. S. Amer. Pl.* 106 [20 Dec. 1920] (Rusby 1920). — *Gerasanthus viridis* (Rusby) Borhidi, *Acta Bot. Hung.* 34 (3-4): 403 (Borhidi 1988). — *Cordia viridis* (Rusby) I.M.Johnst., *J. Arnold Arbor.* 33 (1): 64 [Jan. 1952] (Johnston 1952).

Cordia ierensis Britton, *Bull. Torrey Bot. Club* 50 (1): 54 [Jan. 1923] (Britton 1923).

Cordia coriacea Killip, *J. Wash. Acad. Sci.* 17: 329 (Killip 1927).

VERNACULAR NAMES. — Wp: kuluwa'i, kuluwa'i sili.

HERBARIUM DATA (FG). — 1 collections at CAY. Sel. exs.: *P. Grenand* 237.

SIZE. — Guyana. *P. Mutchnick* 773, 15 m.

[431] *Cordia sprucei* Mez

Bot. Jahrb. Syst. 12: 549 [23 Dec. 1890] (Mez 1890). — *Lithocardium sprucei* (Mez) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus sprucei* (Mez) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

VERNACULAR NAMES. — Wp: kuluwa'i.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *M.-F. Prévost & C. Feuillet* 3968.

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.1\text{ cm}$.

[432] *Cordia tetrandra* Aubl.
(Fig. 20C)

Hist. Pl. Guiane 1: 222 [Jun.-Dec. 1775] (Aublet 1775). — *Lithocardium tetrandrum* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Borellia aspera* Raf., *Sylva Tellur.*: 41 (Rafinesque 1838), *nom. illeg. superfl.* (based on *Cordia tetrandra*). — *Gerasanthus tetrandrus* (Aubl.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

Cordia cordifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 70 [Late Sep. 1818] (Kunth 1818). — *Lithocardium cordifolium* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 976 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus cordifolius* (Kunth) Borhidi, *Acta Bot. Hung.* 34 (3-4): 399 (Borhidi 1988).

Cordia muneco Kunth, *Nova genera et species plantarum [H.B.K.]* 7: 207 [21 Feb. 1825] (Kunth 1825). — *Lithocardium muneco* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Gerasanthus muneco* (Kunth) Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988).

Cordia umbraculifera DC., *Prod. [A. P. de Candolle]* 9: 484 [1 Jan. 1845] (Candolle 1845).

VERNACULAR NAMES. — Ka: alatuluka • Wp: kuluwa'i u • Br: chapéu-de-sol.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777832] designated by Lanjouw & Uittien [1940: 150]).

SIZE. — Up to 18 m tall (Gaviria 1997).

[433] *Cordia toqueve* Aubl.

Hist. Pl. Guiane 1: 228 [Jun.-Dec. 1775] (Aublet 1775). — *Lithocardium toqueve* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov.




FIG. 20. — Cordiaceae: **A**, *Cordia laevifrons* I.M.Johnst. (*D. Sabatier & J.-F. Molino* 5647); **B**, *Cordia sagotii* I.M.Johnst. (*M.-F. Prévost & D. Sabatier* 4804); **C**, *Cordia tetrandra* Aubl. (*M.-F. Prévost* 1500). Dichapetalaceae: **D**, *Tapura capitulifera* Baill. A, D © D. Sabatier/IRD; B, C, © M.-F. Prévost/IRD.

1891] (Kuntze 1891). — *Gerascanthus toquene* (Aubl.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 402 (Borhidi 1988).

Cordia heterophylla Poir., *Dict. Sci. Nat., ed. 2 [F. Cuvier]*, 10: 409 (Poiret 1818). — *Lithocardium heterophyllum* (Poir.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891). — *Gerascanthus heterophyllus* (Poir.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 400 (Borhidi 1988).

Toquera tomentosa Raf., *Sylva Tellur.*: 40 (Rafinesque 1838).

Cordia hebecarpa DC., *Prodri. [A. P. de Candolle]* 9: 488 [1 Jan. 1845] (Candolle 1845). — *Lithocardium hebecarpum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia pubescens Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 4: 800 [Mar.-June 1819] (Roemer & Schultes 1819). — *Gerascanthus pubescens* (Roem. & Schult.) Borhidi, *Acta Bot. Hung.* 34 (3-4): 401 (Borhidi 1988). — *Lithocardium pubescens* (Roem. & Schult.) Kuntze, *Revis. Gen. Pl.* 2: 977 [5 Nov. 1891] (Kuntze 1891).

Cordia expansa Lingelsh., *Repert. Spec. Nov. Regni Veg.* 7: 244 (Lingelshiem 1909).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777830]) designated by Lanjouw & Uittien [1940: 150]).

INVENTORY DATA (FG). — 1 tree, dbh = 15.5 cm.

Family DICHAPETALACEAE Baill.
Genus *Tapura* Aubl.

[434] *Tapura amazonica* Poepp.

Nova genera ac species plantarum [Poeppig & Endlicher] 3: 41 [8-11 Mar. 1843] (Poeppig 1843).

Tapura ciliata Gardner, *Icon. Pl.* 5 [n.s., 1]: t. 466 [Jan.-Oct. 1842] (Gardner 1842). — *Tapura amazonica* var. *ciliata* (Gardner) Baill., *Fl. Bras. [Martius]* 12 (1): 375 [1 Apr. 1886] (Baillon 1886).

Tapura amazonica var. *cuspidata* Baill., *Fl. Bras. [Martius]* 12 (1): 375 [1 Apr. 1886] (Baillon 1886).

Tapura amazonica var. *dasyphylla* Baill., *Fl. Bras. [Martius]* 12 (1): 375 [1 Apr. 1886] (Baillon 1886).

Tapura amazonica var. *sublanceolata* Baill., *Fl. Bras. [Martius]* 12 (1): 375 [1 Apr. 1886] (Baillon 1886).

VERNACULAR NAMES. — Pa: á-kasiuminio.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *E.M. Mélinon* 170 (P[P04764399]), syntype of *Tapura amazonica* var. *dasyphylla*.

INVENTORY DATA (FG). — 20 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.4$ cm.

[435] *Tapura capitulifera* Baill.
(Fig. 20D)

Adansonias [Baillon] 11: 112 (Baillon 1873). — *Chailletia capitulifera* Spruce ex Baill., *Adansonias [Baillon]* 11: 112 (Baillon 1873), *nom. nud. pro syn.*

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *D. Sabatier* 5792.

INVENTORY DATA (FG). — 455 trees in 76 plots; $F_{\max} = 5.5\%$; $\text{dbh}_{\text{inv}} = 95.5$ cm.

[436] *Tapura guianensis* Aubl.

Hist. Pl. Guiane 1: 126 [Jun.-Dec. 1775] (Aublet 1775). — *Rohria tapura* J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 93 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. superfl.* (based on *Tapura guianensis*). — *Rohria petioliflora* Willd., *Sp. Pl.*, ed. 41 (1): 186 [June 1797] (Willdenow 1797), *nom. illeg. superfl.* (based on *Tapura guianensis*).

Rohria schreberi J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 93 [late Sep.-Nov. 1791] (Gmelin 1791).

Chailletia sessiliflora DC., *Nouv. Bull. Sci. Soc. Philom. Paris* 2 (40): 205 [Jan. 1811] (Candolle 1811).

Tapura cucullata Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 292 (Bentham 1853).

Tapura negrensis Suess., *Repert. Spec. Nov. Regni Veg.* 51: 199 (Suessenguth 1942).

VERNACULAR NAMES. — Ka: wasakau • Wp: awalapuna sili, ka'a sili, paa lo'o, wila towakape si.

HERBARIUM DATA (FG). — 144 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (lecto-, P-JJR[P00777925] designated by Lanjouw & Uittien [1940: 156]).

INVENTORY DATA (FG). — 117 trees in 31 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 22.9$ cm.

[437] *Tapura singularis* Ducke

Trop. Woods 90: 21 (Ducke 1947).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier* 2530.

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 38.2$ cm.

Family DILLENIACEAE Salisb.
Genus *Curatella* Loefl.

[438] *Curatella americana* L.

Syst. Nat., ed. 10, 2: 1079 [7 June 1759] (Linnaeus 1759), “*americ.*”

Curatella cambaiba A.St.-Hil., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (1): 22 [23 Feb. 1824] (Saint-Hilaire 1824), “*Cambaiba*”.

Curatella grisebachiana Eichler, *Fl. Bras. [Martius]* 13 (1): 69 [15 Jan. 1863] (Eichler 1863).

Curatella americana var. *pentagyna* Donn.Sm., *Bot. Gaz.* 46 (2): 109 [22 Aug. 1908] (Donnell Smith 1908).

NOTE. — Restricted to savannas.

VERNACULAR NAMES. — Pa: sawu-anen-kamwi • Ka: kulata • Cr: féy-sab • Br: caimbé, lixeira.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *M.-F. Prévost* 1636.

SIZE. — Up to 16 m tall (Aymard C. 1998).

Family EBENACEAE Gürke
Genus *Diospyros* L.

[439] *Diospyros capreifolia* Mart. ex Hiern

Trans. Cambridge Philos. Soc. 12 (1): 254 (Hiern 1873), “*caprifolia*”.

Maba melinonii Hiern, *Trans. Cambridge Philos. Soc.* 12 (1): 143 (Hiern 1873), “*Mellinoni*”. — *Diospyros melinonii* (Hiern) A.C.Sm., *Bull. Torrey Bot. Club* 60 (6): 390 [1 June 1933] (Smith 1933), “*Melinoni*”.

VERNACULAR NAMES. — Pa: á-seiminio-priyo, á-seiminio-priyu, miret • Nt: bita tiki • Cr: bwa-krayon • Br: altinha.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *E.M. Mélinon* s.n. (original material of *Maba melinonii*: F[V0063306F, V0063307F, V0063308F]).

INVENTORY DATA (FG). — 31 trees in 25 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 28.6$ cm.

[440] *Diospyros carbonaria* Benoist

Arch. Bot. 5 (Mém. 1): 249 [27 Mar. 1933] (Benoist 1933).

Diospyros duckei Sandwith, *Kew Bull.* 4 (4): 488 [“1949” publ. 16 Feb. 1950] (Sandwith 1950).

Diospyros capimnensis Pires & Cavalcante, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 9: 1 (Pires & Cavalcante 1960).

VERNACULAR NAMES. — Pa: miret • Nt: bita tiki • Cr: bwa-charbon, bwa-krayon.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *R. Benoist* 448 (original material P[P00721344, P00721346]).

INVENTORY DATA (FG). — 137 trees in 73 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 35.3$ cm.

[441] *Diospyros cavalcantei* Sothers

Kew Bull. 55 (2): 471 (Sothers 2000).

VERNACULAR NAMES. — Pa: miret • Cr: bwa-krayon.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *M.-F. Prévost* et al. 4671.

INVENTORY DATA (FG). — 13 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 76.1$ cm.

[442] *Diospyros cayennensis* A.DC.

Prodr. [A. P. de Candolle] 8: 224 [mid Mar. 1844] (Candolle 1844).

Diospyros ierensis Britton, *Bull. Torrey Bot. Club* 48 (12): 336 ["1921" publ. 1922] (Britton 1922).

VERNACULAR NAMES. — Ka: palala, simyali epi, talala.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J. Martin s.n.* (lecto-, G-DC[G00142182] designated by Wallnöfer [2001: 888]; isolecto-, P[P00721345, P00721347]).

INVENTORY DATA (FG). — 30 trees in 24 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.7$ cm.

[443] *Diospyros dichroa* Sandwith
(Fig. 21A)

Bull. Misc. Inform. Kew 1931 (10): 482 [31 Dec. 1931] (Sandwith 1931).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: á-kabebetye, miret, pukuu-kamwi • Ka: yukutuna • Nt: bita tiki • Cr: bwa-krayon, bwa-zoranj.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier et al. 4457*.

INVENTORY DATA (FG). — 35 trees in 9 plots; $F_{\max} = 3.5\%$; $dbh_{\text{inv}} = 45$ cm.

[444] *Diospyros guianensis* (Aubl.) Gürke
(Fig. 21B)

Nat. Pflanzenfam. [Engler & Prantl] 4 (1): 163 (Gürke 1891). — *Paralea guianensis* Aubl., *Hist. Pl. Guiane* 1: 576 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Pa: miret • Ka: palala, simyali epi, talala • Wp: miku lapi'a u • Wn: meikolo muhunu, meikolo muhupu, mekolo nohunu • Nt: baaka tiki • Cr: bwa-krayon.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material FI[FI012223]).

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 39.6$ cm.

[445] *Diospyros martinii* Benoist

Arch. Bot. 5 (Mém. 1): 248 [27 Mar. 1933] (Benoist 1933), "Martini".

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5705*.

INVENTORY DATA (FG). — 6 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.5$ cm.

[446] *Diospyros sericea* A.DC.

Prodr. [A. P. de Candolle] 8: 236 [mid Mar. 1844] (Candolle 1844). — *Maba sericea* (A.DC.) Hiern, *Trans. Cambridge Philos. Soc.* 12 (1): 140 (Hiern 1873).

Diospyros vestita Benoist, *Arch. Bot. 5 (Mém. 1)*: 248 [27 Mar. 1933] (Benoist 1933), *in clavi; non* Bakh. [May 1933] (Bakhuizen 1933).

Diospyros praetermissa Sandwith, *Kew Bull. 4 (4)*: 483 ["1949" publ. 16 Feb. 1950] (Sandwith 1950).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *R. Benoist 520* (original material of *Diospyros vestita*: P[P00721401]).

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 51.9$ cm.

[447] *Diospyros tetrandra* Hiern

Trans. Cambridge Philos. Soc. 12 (1): 210 (Hiern 1873), *non* Span. (1841, *nom. nud.*)

VERNACULAR NAMES. — Ka: palala, simyali epi, talala • Wp: miku lapi'a sówí • Nt: baaka tiki.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J. Martin s.n.* (lecto-, K[K000644356] designated by Wallnöfer [2020: 231]).

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23$ cm.

[448] *Diospyros* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier et al. 5803*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 47.7$ cm.

Family ELAEOCARPACEAE Juss.

NOTE

The presence of *Sloanea usurpatrix* Sprague & Riley in French Guiana seems plausible given its known trans-Amazonian distribution. According to Pennington & Wise (2017), it is attested by a single sterile collection (Nouragues Nature Reserve, 40 km SW of Regina, Mori et al. 26678, CAY, K). However, the leaf morphology of this specimen clearly differs from that of *S. usurpatrix*, while fitting perfectly with *S. synandra*.

Genus *Sloanea* L.[449] *Sloanea acutiflora* Uittien
(Fig. 22A)

Recueil Trav. Bot. Néerl. 22: 357 ["1925" publ. Jan. 1926] (Uittien 1926).

NOTES. — Known only from the Guiana Shield. Pennington & Wise (2017: 77) placed *S. acutiflora* in synonymy under *S. latifolia* (Rich.) K.Schum., arguing that they have the same floral morphology, and that there are intermediate states throughout the range of the species (i.e. most of Amazonia) in fruit spine development and leaf shape; the typical form of *S. latifolia* has long petiole and large broad leaves with rounded bases while other “forms”, typified by *S. acutiflora* in the Guiana Shield and *S. inermis* Ducke in Southern and Western Amazonia, have smaller leaves with acute or cuneate base and short petioles. Our observations contradict these statements. In French Guiana, all specimens that correspond to the Pennington and Wise concept of *S. latifolia* can easily be separated into two very distinct groups. In the first group, which includes the original material (see Notes under *S. latifolia*), trees have both buttresses and stilt roots, petioles of very variable length on the same branchlet, the longest 3–15 cm, and large leaves drying yellowish brown, of variable size but frequently 20 cm long, with rounded to obtuse, rarely cuneate base and a densely reticulate venation. Inflorescences are large, well-developed many-flowered corymbs, flower buds are slender, the base gradually attenuated on the pedicel, and covered with a dense yellow tomentum. Fruit is up to 4.5 cm long, brown, smooth, and seeds are surrounded by a white to pink aril (Fig. 22D). Trees of the second group have buttresses but no stilt roots. Petioles are shorter (usually < 1 cm, rarely 1–1.5 cm). Leaves dry to a light green, and are much smaller (rarely up to 15 cm long), with an acute-attenuated base, never round or obtuse, and a much looser reticulum. Inflorescences are short, more or less cymose, bearing 1–3 (–5) flowers. Buds are subglobose to ovoid, the base abruptly reduced at the junction with the pedicel, the whitish, much less dense pilosity making them appear greyish rather than yellow. Fruit is red or pink, less than 2.5 cm long, sometimes covered with sparse, short conic spines, and the aril is orange (Fig. 22A). This last group corresponds with the type of *S. acutiflora*. Pennington & Wise (2017: 77) designated the lectotype of *S. acutiflora* as “[...] BW 6298 (lectotype, U, chosen here; isolectotypes, K, L)”. At U there are three sheets of *C.J. Zaandam-Boschwezen Suriname* 6298; these sheets (barcodes U0001682, U0001683, U0001684) are all kept together, and according to Art. 8.3 of the ICN (Turland *et al.* 2018), they are here treated as a single specimen with multiple preparations. Isolectotypes are at K(not seen), L(L0013589), RB(RB00538334) and US (00098537).

Although the distributions of these two groups overlap throughout much of French Guiana (west, center and south), no intermediate states have yet been found between the character syndromes defining these two groups. It is also interesting to note that, except in the extreme south of the territory (Tumuc-Humac Mountains), the “typical” *S. latifolia* form has never been found within 50 km of the border with Suriname. In view of the above, the reinstatement of *S. acutiflora* appears necessary.

VERNACULAR NAMES. — Pa: waaduk-seine.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *D. Sabatier* 1234.

INVENTORY DATA (FG). — 39 trees in 31 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 65$ cm.

[450] *Sloanea brachytelpala* Ducke

Trop. Woods 76: 22 (Ducke 1943).

Sloanea calva Pal.-Duque & Fern.Alonso, *Revista Acad. Colomb. Ci. Exact.* 29 (111): 180 [June 2005] (Palacios-Duque & Fernández Alonso 2005).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier* & M.-F. Prévost 3568.

INVENTORY DATA (FG). — 16 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54.4$ cm.

[451] *Sloanea brevipes* Benth.

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 68 [May 1861] (Bentham 1861).

VERNACULAR NAMES. — Pa: ihap-kamwi-duwē • Ka: kusewelan • Wn: inuuman • Nt: busi kusuwe • Cr: chateng, roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier* 844.

INVENTORY DATA (FG). — 104 trees in 55 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 80$ cm.

[452] *Sloanea conferta* T.D.Penn.

Opusc. Neotrop. 2: 6 [13 July 2016] (Pennington 2016).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier* 4897 (holo-, K, not seen; iso-, CAY[CAY065610], US[00943331]).

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.1$ cm.

[453] *Sloanea echinocarpa* Uittien

Recueil Trav. Bot. Néerl. 22: 354 [“1925” publ. Jan. 1926] (Uitien 1926).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier* & M.-F. Prévost 2268.

SIZE. — Up to 65 cm dbh (Pennington & Wise 2017).

[454] *Sloanea eichleri* K.Schum.

Fl. Bras. [Martius] 12 (3): 183 [1 Nov. 1886] (Schumann 1886).

Sloanea sinemariensis var. *melinonii* Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 11: 156 (Sagot 1881), “*Melinonis*”.

NOTE. — The epithet “*melinonis*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland *et al.* 2018: Art. 60.8).

VERNACULAR NAMES. — Pa: ihap-kamwi, ihap-kamwi-seine • Wp: uluku panali sili, ulukupalani sili • Cr: roukou-gran-bwa, roukou-sovaj • Br: urucurana.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.-F. Molino* & *D. Sabatier* 2693.

INVENTORY DATA (FG). — 36 trees in 28 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 32$ cm.



FIG. 21.—Ebenaceae: **A**, *Diospyros dichroa* Sandwith (*D.* Sabatier & M.-F. Prévost 5246); **B**, *Diospyros guianensis* (Aubl.) Gürke. Erythroxylaceae: **C**, *Erythroxylum fimbriatum* Poir. (*D.* Sabatier & J.-F. Molino 5229); **D**, *Erythroxylum suberosum* A.St.-Hil. et al. (*D.* Sabatier & J.-F. Molino 5198). © *D. Sabatier/IRD*.

[455] *Sloanea erythrocarpa* T.D.Penn.
(Fig. 22B)

Opusc. Neotrop. 2: 10 [13 July 2016] (Pennington 2016).

NOTE.—Known only from French Guiana.

HERBARIUM DATA (FG).—13 collections at CAY. Sel. exs.: *D. Sabatier 3540*.

INVENTORY DATA (FG).—21 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 45$ cm.

Sloanea longipes Ducke, *Arch. Inst. Biol. Veg.* 2 (2): 166 [Dec. 1935] (Ducke 1935).

Sloanea maroana Steyermark., *Pittieria* 7: 14 (Steyermark 1978).

Sloanea sippapana Steyermark., *Ann. Missouri Bot. Gard.* 75 (4): 1580 [“1988” publ. 8 Mar. 1989] (Steyermark 1989).

HERBARIUM DATA (FG).—No collections at CAY. Sel. exs.: *S.A. Mori 26549* (K, *fide* Pennington & Wise [2017], not seen).

SIZE.—Up to 120 cm dbh (Pennington & Wise 2017).

[456] *Sloanea floribunda* Spruce ex Benth.

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 66 [May 1861] (Bentham 1861).

Sloanea paniculata Spruce ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 66 [May 1861] (Bentham 1861).

[457] *Sloanea garckeana* K.Schum.

Fl. Bras. [Martius] 12 (3): 177 [1 Nov. 1886] (Schumann 1886).

HERBARIUM DATA (FG).—8 collections at CAY. Sel. exs.: *M.-F. Prévost 1893*.

INVENTORY DATA (FG).—15 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 73.5$ cm.

[458] *Sloanea grandiflora* Sm.

Cycl. [Rees] 33: (Sloanea no. 2) [17 May 1816] (Smith 1816).

Sloanea amplifrons I.M.Johnst., *Contr. Gray Herb.*, n.s., 75: 29 (Johnston 1925).

Sloanea linderi I.M.Johnst. ex Kribs, *Trop. Woods* 13: 16 (Kribs 1928), *nom. nud.*

VERNACULAR NAMES. — Pa: ihap-kamwi • Ka: kusewelan, polo ali • Te: tokolo-pipinatsi • Wp: tawisowiso, uluku panali, ulukupalani • Wn: inuuman • Nt: busi kusuwe • Cr: chatengn, roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4341.

INVENTORY DATA (FG). — 28 trees in 16 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 73 \text{ cm}$.

[459] *Sloanea granulosa* Ducke

Bol. Técn. Inst. Agron. N. 19: 13 (Ducke 1950).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4126.

INVENTORY DATA (FG). — 13 trees in 12 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 100 \text{ cm}$.

[460] *Sloanea guianensis* (Aubl.) Benth. subsp. *guianensis*

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 69 [May 1861] (Bentham 1861). — *Ablania guianensis* Aubl., *Hist. Pl. Guiane* 1: 585 [Jun.-Dec. 1775] (Aublet 1775). — *Trichocarpus guianensis* (Aubl.) Raeusch., *Nomencl. Bot. [Raeusch.]*, ed. 3, 158 (Raeuschel 1797), *nom. illeg. superfl.* (genus name superfluous, based on the type of *Ablania* Aubl.). — *Trichocarpus laurifolius* Willd., *Sp. Pl.*, ed. 4 2 (2): 1224 [Dec. 1799] (Willdenow 1799), “*laurifolia*”, *nom. illeg. superfl.* (genus name superfluous, based on the type of *Ablania* Aubl.)

Ablania laurifolia Pers., *Syn. Pl. [Persoon]* 2 (1): 81 [Nov. 1806] (Persoon 1806).

Dasynema pubescens Schott, *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 408 [Jan.-June 1827] (Schott 1827). — *Sloanea pubescens* (Schott) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 329 (Radlkofer 1882), *nom. illeg. hom.*, *non* Benth. (Bentham 1861).

Sloanea alnifolia Mart., *Flora* 20 (2, Beibl.): 94 (Martius 1837). — *Dasynema alnifolium* (Mart.) Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 352 [18-20 Sep. 1842] (Walpers 1842). — *Dasynema cuneifolium* (Mart.) Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 352 [18-20 Sep. 1842] (Walpers 1842).

Sloanea cuneifolia Mart., *Flora* 20 (2, Beibl.): 94 (Martius 1837).

Dasynema pubescens Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 31 (1): 224 (Turczaninow 1858), *nom. illeg. hom.*, *non* Schott (1827) *nec* Poepp. (Poepig 1845).

Sloanea microcarpa Planch. ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 69 [May 1861] (Bentham 1861). — *Sloanea guianensis* var. *microcarpa* (Planch. ex Benth.) K.Schum., *Fl. Bras. [Martius]* 12 (3): 191 [1 Nov. 1886] (Schumann 1886).

Sloanea regelii K.Schum., *Fl. Bras. [Martius]* 12 (3): 189 [1 Nov. 1886] (Schumann 1886).

Sloanea alnifolia var. *lancea* K.Schum., *Fl. Bras. [Martius]* 12 (3): 194 [1 Nov. 1886] (Schumann 1886).

Sloanea alnifolia var. *ovalis* K.Schum., *Fl. Bras. [Martius]* 12 (3): 194 [1 Nov. 1886] (Schumann 1886).

Sloanea ptariana Steyermark., *Fieldiana, Bot.* 28 (2): 360 (Steyermark 1952).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ihap-kamwi • Ka: kusewelan • Wp: uluku panali sili, ulukupalani sili • Wn: inuuman • Nt: busi kusuwe • Cr: chatengn, roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier* 1237.

INVENTORY DATA (FG). — 42 trees in 28 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 60 \text{ cm}$.

[461] *Sloanea guianensis* subsp. *purdiae*
(Griseb. ex R.O.Williams & Cheesman) T.D.Penn.

Opusc. Neotrop. 2: 13 [13 July 2016] (Pennington 2016), “*purdiae*”. — *Sloanea purdiei* Griseb. ex Benth. & Hook.f., *Gen. Pl. [Bentham & Hooker]* 2 (2): 653 [May 1876] (Bentham & Hooker 1876), “*Purdiae*”, in obs., *nom. nud.* — *Sloanea purdiei* Griseb. ex R.O.Williams & Cheesman, *Fl. Trinidad & Tobago* 1 (2): 109 [Nov. 1929] (Williams & Cheesman 1929), “*purdiae*”.

Sloanea breviseta Steyermark., *Fieldiana, Bot.* 28 (2): 357 (Steyermark 1952).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville B-5392*.

SIZE. — Up to 25 m tall (Pennington & Wise 2017).

[462] *Sloanea guianensis* subsp. *stipitata*
(Spruce ex Benth.) T.D.Penn.

Opusc. Neotrop. 2: 6 [13 July 2016] (Pennington 2016). — *Sloanea stipitata* Spruce ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 68 [May 1861] (Bentham 1861).

Sloanea maximowicziana K.Schum., *Fl. Bras. [Martius]* 12 (3): 192 [1 Nov. 1886] (Schumann 1886).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori & Y. Veyret* 8972.

SIZE. — Up to 80 cm dbh (Pennington & Wise 2017).

[463] *Sloanea latifolia* (Rich.) K.Schum.
(Fig. 22D)

Fl. Bras. [Martius] 12 (3): 173 [1 Nov. 1886] (Schumann 1886). — *Blondea latifolia* Rich., *Actes Soc. Hist. Nat. Paris* 1: 110 [Oct. 1792] (Richard 1792).

Sloanea corymbiflora DC., *Prod. [A. P. de Candolle]* 1: 516 [mid Jan. 1824] (Candolle 1824).

Sloanea inermis Ducke, *Arch. Inst. Biol. Veg.* 4 (1): 48 [June 1938] (Ducke 1938).

Sloanea gentryi Pal.-Duque & C.M.Baeza, *Novon* 23 (1): 70 [8 Apr. 2014] (Palacios-Duque & Baeza 2014).

NOTE. — Richard (1792: 110) along with the description of *Blondea latifolia* did not cite any specimen. Pennington (2016: 15) and Pennington and Wise (2017: 76) cited the type of this name as “French Guiana (Cayenne), fl., *LeBlond* 219 (holotype, P; isotype, G). These citations cannot be treated as an inadvertent lectotypification because, being published after 2001, they should have been accompanied by “here designated” or a similar expression. In addition, there are three specimens of *Blondea latifolia* at P that were collected by Leblond, with the locality “*ad amnem Kourou*” (Kourou River), each with a different label, no collector number, and with barcodes P02440486, P02440487, and P02440488. Another specimen of *Blondea latifolia* collected by Leblond, with the collector number 209 (not “219”) is at G (G00104361). Therefore, a lectotype is hereafter designated.

VERNACULAR NAMES. — Pa: ihap-kamwi-purubumna, kunau, sitrukamwi • Ka: tokuhsa • Wp: tapi'ilo wapa'a, wila kalyulu, wila kayulu • Nt: busi kusuwe • Cr: chateng, roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.B. Leblond* 209 (lecto-, P[P02440487], here designated; isolecto-, G[G00104361], P[P02440486, P02440488]).

INVENTORY DATA (FG). — 25 trees in 20 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54$ cm.

[464] *Sloanea morii* T.D.Penn.

Opusc. Neotrop. 2: 17 [13 July 2016] (Pennington 2016).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 2263.

INVENTORY DATA (FG). — 1 tree, $dbh = 11.1$ cm.

[465] *Sloanea nitida* G.Don

Gen. Hist. 1: 555 [early Aug. 1831] (Don 1831).

Sloanea bracteosa Ducke, *Arch. Inst. Biol. Veg.* 2 (2): 168 [Dec. 1935] (Ducke 1935).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, BM[BM000907458, BM000907459]; iso-, K[K000380070]).

SIZE. — Up to 60 cm dbh (Pennington & Wise 2017).

[466] *Sloanea obtusifolia* (Moric.) K.Schum.

Fl. Bras. [Martius] 12 (3): 181 [1 Nov. 1886] (Schumann 1886). — *Adenobasium obtusifolium* Moric., *Pl. Nouv. Amér.*: 83 (Moricand 1840). — *Dasynema obtusifolium* (Moric.) Walp., *Ann. Bot. Syst.*

[Walpers] 1 (1): 110 [6-7 Nov. 1848] (Walpers 1848), “*obtusum*”. — *Sloanea obtusifolia* (Moric.) Planch. ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 63, 68 [May 1861] (Bentham 1861), “*obtusa*”.

Sloanea fernando-costae Hoehne, *Arq. Bot. Estado São Paulo* 1 (4): 93 (Hoehne 1942).

NOTES. — *Dasynema “obtusum”* (Walpers 1848: 110) is treated here as a misspelling of *obtusifolium*, since Walpers unambiguously referenced *Adenobasium obtusifolium* Moric.; therefore it is not an illegitimate homonym of *D. obtusum* Splitg. Similarly, as Bentham (1861: 63) explicitly based his *Sloanea “obtusa”* on *A. obtusifolium*, the epithet is also to be corrected.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost* & *D. Sabatier* 4662.

INVENTORY DATA (FG). — 11 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.2$ cm.

[467] *Sloanea parviflora* Planch. ex Benth.

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 63, 68 [May 1861] (Bentham 1861).

Dasynema obtusum Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 98 [Aug.-Sep. 1842] (Splitgerber 1842). — *Sloanea obtusa* (Splitg.) K.Schum., *Fl. Bras. [Martius]* 12 (3): 181 [1 Nov. 1886] (Schumann 1886), nom. illeg. hom., non Planch. ex Benth. (1861) [synonym of *S. obtusifolia* (Moric.) K.Schum.]

Sloanea parviflora var. *pedicellata* Sagot, *Ann. Sci. Nat., Bot. sér. 6, 11:* 157 (Sagot 1881).

Sloanea kappleriana Pulle, *Enum. Vasc. Pl. Surinam* 279 (Pulle 1906).

Sloanea obidensis Ducke, *Arch. Inst. Biol. Veg.* 2 (2): 160 [Dec. 1935] (Ducke 1935).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 4429.

INVENTORY DATA (FG). — 9 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 26.4$ cm.

[468] *Sloanea pubescens* Benth.

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 69 [May 1861] (Bentham 1861). — *Dasynema pubescens* Poepp., *Nova genera ac species planarum [Poeppig & Endlicher]* 3: 74 [23-25 Jan. 1845] (Poeppig 1845), nom. illeg. hom., non Schott (1827) nec Turcz. (Turczaninow 1858). — *Sloanea egensis* Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 329 (Radlkofer 1882).

Sloanea gracilis Uittien, *Recueil Trav. Bot. Néerl.* 22: 356 [“1925” publ. Jan. 1926] (Uittien 1926).

Sloanea pseudodentata Ducke, *Arch. Inst. Biol. Veg.* 2 (2): 159 [Dec. 1935] (Ducke 1935).

NOTES. — Although based on *Dasynema pubescens* Poepp., *S. pubescens* Benth. is to be treated as a new name because the former is illegitimate.

VERNACULAR NAMES. — Pa: ihap-kamwi-washiune • Wp: uluku panali, ulukupalani • Cr: roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *S. Barrier 5018*.

INVENTORY DATA (FG). — 12 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 40 \text{ cm}$.

[469] *Sloanea rojasiae* Vásquez

Arnaldoa 16 (2): 57 [“2009” publ. Feb. 2010] (Vásquez 2010).

HERBARIUM DATA (FG). — A single collection, *D. Loubry 833*.

SIZE. — Up to 25 cm dbh (Pennington & Wise 2017).

[470] *Sloanea rufa* Planch. ex Benth.

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 68 [May 1861] (Bentham 1861).

Sloanea longicaudata Ducke, *Bol. Técn. Inst. Agron. N.* 19: 14 (Ducke 1950).

Sloanea filiformis D.Samp. & V.C.Souza, *Phytotaxa* 16: 45 [4 Feb. 2011] (Sampaio & Souza 2011).

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, K[K000380085, K000380086, K000380087]).

INVENTORY DATA (FG). — 19 trees in 18 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 34.3 \text{ cm}$.

[471] *Sloanea sinemariensis* Aubl.

Hist. Pl. Guiane 1: 534 [Jun.-Dec. 1775] (Aublet 1775). — *Sloanea aubletii* Sw., *Fl. Ind. Occid.* 2: 940 [late 1800] (Swartz 1800), nom. illeg. superfl. (based on *Sloanea sinemariensis*).

VERNACULAR NAMES. — Pa: ihap-kamwi-priye • Cr: roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4421*.

INVENTORY DATA (FG). — 13 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 63 \text{ cm}$.

[472] *Sloanea synandra* Spruce ex Benth.
(Fig. 22C)

J. Proc. Linn. Soc., Bot. 5 (Suppl. 2): 66 [May 1861] (Bentham 1861).

Sloanea macrantha Ducke, *Arch. Inst. Biol. Veg.* 2 (2): 162 [Dec. 1935] (Ducke 1935).

Sloanea megacarpa Steyermark. & Marc.-Berti, *Bol. Soc. Venez. Ci. Nat.* 26: 467 (Steyermark & Marcano-Berti 1966).

VERNACULAR NAMES. — Wp: wila kalayulu, wilakayulu • Cr: roukou-gran-bwa, roukou-sovaj • Fr: roucou grand bois • Br: urucurana.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4848*.

INVENTORY DATA (FG). — 22 trees in 16 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 100 \text{ cm}$.

[473] *Sloanea trichosticha* R.O.Williams & Sandwith

Fl. Trinidad & Tobago 1 (2): 110 [Nov. 1929] (Williams & Sandwith 1929).

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.-F. Molino & M.-F. Prévost 2531*.

INVENTORY DATA (FG). — 31 trees in 23 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 62.9 \text{ cm}$.

[474] *Sloanea wurdackii* Steyermark.

Ann. Missouri Bot. Gard. 75 (4): 1583 [“1988” publ. 8 Mar. 1989] (Steyermark 1989).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *B. Riéra 1257*.

SIZE. — Up to 45 cm dbh (Pennington & Wise 2017).

[475] *Sloanea* sp. A

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4792*.

INVENTORY DATA (FG). — 17 trees in 15 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 80 \text{ cm}$.

[476] *Sloanea* sp. B

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15498*.

INVENTORY DATA (FG). — 28 trees in 24 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 100 \text{ cm}$.

[477] *Sloanea* sp. C

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 5014*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 50 \text{ cm}$.

[478] *Sloanea* sp. D

NOTE. — This is possibly *S. caribaea* Krug & Urb. ex Duss.

HERBARIUM DATA (FG). — A single collection, *H. Richard & Bagadi 674*, $\text{dbh} = 35 \text{ cm}$.

[479] *Sloanea* sp. E

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 5806*, $\text{dbh} = 35 \text{ cm}$.

[480] *Sloanea* sp. F

NOTES. — According to Pennington & Wise (2017), this fruiting collection could belong to *S. ferruginea* T.D.Penn., a species whose fruit is not known. However, the leaves of this collection are quite different from those of *S. ferruginea*. This specimen does not fit in any other French Guianan species.

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 25669*, 35 m × 75 cm.

[481] *Sloanea* sp. G

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2990*.

INVENTORY DATA (FG). — 51 trees in 36 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 70 \text{ cm}$.

[482] *Sloanea* sp. H

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2947*.

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 70 \text{ cm}$.

Family ERYTHROXYLACEAE Kunth
Genus *Erythroxylum* P.Browne

[483] *Erythroxylum amazonicum* Peyr.

Fl. Bras. [Martius] 12 (1): 167 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum mucronatum var. *majus* Sagot, *Ann. Sci. Nat., Bot. sér. 6, 11*: 179 (Sagot 1881), “*major*”.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P.A. Sagot s.n.*, 1858 (P[P05483039]).

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 15.7 \text{ cm}$.

[484] *Erythroxylum citrifolium* A.St.-Hil.

Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 2 (13): 94 [10 Oct. 1829] (Saint-Hilaire 1829).

Erythroxylum gomphoides Planch. & Linden ex Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4, 18*: 341 (Triana & Planchon 1862).

Erythroxylum micranthum Bong. ex Peyr., *Fl. Bras. [Martius] 12 (1)*: 164 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum paraense Peyr., *Fl. Bras. [Martius] 12 (1)*: 164 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum acutifolium Steud. ex Peyr., *Fl. Bras. [Martius] 12 (1)*: 166 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum citrifolium var. *latifolium* O.E.Schulz, *Pflanzenr. [Engler] IV.134 (Heft 29)*: 37 [13 Aug. 1907] (Schulz 1907).

Erythroxylum citrifolium var. *minus* O.E.Schulz, *Pflanzenr. [Engler] IV.134 (Heft 29)*: 37 [13 Aug. 1907] (Schulz 1907).

Erythroxylum duckei Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2)*: 416 (Huber 1909).

VERNACULAR NAMES. — Pa: agagut, ararut • Wp: pikau ki’iy, pikau lemiū’i, tukā nákū sili • Br: pimenta-de-nambú.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *S.A. Mori et al. 15394*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 13 \text{ cm}$.

[485] *Erythroxylum fimbriatum* Peyr.
(Fig. 21C)

Fl. Bras. [Martius] 12 (1): 162 [1 Dec. 1878] (Peyritsch 1878).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2424*, 12 m × 25 cm.

[486] *Erythroxylum kapplerianum* Peyr.

Fl. Bras. [Martius] 12 (1): 159 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum mapuerae Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2)*: 421 (Huber 1909).

NOTE. — *E. grisebachii* Peyr and *E. coelophlebium* Mart. var. *grisebachii* (Peyr.) O.E.Schulz are possible synonyms of this taxon (Plowman & Hensold 2004).

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-663*.

SIZE. — Up to 10 m tall (Plowman & Berry 1999).

[487] *Erythroxylum leptoneurum* O.E.Schulz

Pflanzenr. [Engler] IV.134 (Heft 29): 101 [13 Aug. 1907] (Schulz 1907).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 6480*, 6 m × 10 cm.

[488] *Erythroxylum ligustrinum* DC.

Prod. [A. P. de Candolle] 1: 574 [mid Jan. 1824] (Candolle 1824).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (holo-, G-DC[G00211017]; iso-, F[V0055692F], G[G00353856], P[P00723895]).

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 20.1 \text{ cm}$.

[489] *Erythroxylum lineolatum* DC.

Prod. [A. P. de Candolle] 1: 575 [mid Jan. 1824] (Candolle 1824).



FIG. 22. — Elaeocarpaceae: **A**, *Sloanea acutiflora* Uittinn. (*D. Sabatier* 1234); **B**, *Sloanea erythrocarpa* T.D.Penn. (*D. Sabatier* 3540); **C**, *Sloanea synandra* Spruce ex Benth. (*D. Sabatier & M.-F. Prévost* 4964); **D**, *Sloanea latifolia* (Rich.) K.Schum. (*D. Sabatier* 2088). © D. Sabatier/IRD.

HERBARIUM DATA (FG). — A single collection, *J.-B. Patris s.n.* (holo-, G-DC[G00211384]; iso-, G[G00353858]; probable iso-, P[P00723900]).

INVENTORY DATA (FG). — 1 tree, dbh = 11.1 cm.

[490] *Erythroxylum macrophyllum* Cav.

Diss. 10: 401 (Cavanilles 1789).

Erythroxylum lucidum Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 179 [25 Feb. 1822] (Kunth 1822).

Erythroxylum floribundum Mart., *Beitr. Erythroxylon:* 118 (Martius 1840).

Erythroxylum laurinum Planch. & Linden ex Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4, 18:* 341 (Triana & Planchon 1862), *nom. illeg. hom., non A.Massal.* (Massalongo & Scarabelli 1859).

Erythroxylum costaricense Donn.Sm., *Bot. Gaz.* 23 (4): 240 [22 Apr. 1897] (Donnell Smith 1897). — *Erythroxylum lucidum* var. *costaricense* (Donn.Sm.) O.E.Schulz, *Pflanzren. [Engler]* IV.134 (Heft 29): 25 [13 Aug. 1907] (Schulz 1907).

Erythroxylum ellipticum Ramírez, *Anales Inst. Méd.-Nac. México* 3: 37 (Ramírez 1897), “*Erythroxylon*”, *nom. illeg. hom., non R.Br. ex Benth. (Bentham 1863) nec Peyr. (Peyritsch 1878)*.

Erythroxylum tabascense Britton, *N. Amer. Fl.* 25 (1): 66 (Britton 1907).

Erythroxylum filipes Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 415 (Huber 1909).

VERNACULAR NAMES. — Pa: yauknabwi-kiabumna • Ka: akale ta-pulala kili • Wp: tuká nákü, tuká nákü sili • Wn: okajpu epeli • Br: pimenta-de-nambú.

HERBARIUM DATA (FG). — 74 collections at CAY. Sel. exs.: *D. Stoupy s.n.* (original material MA[MA475675?]); *M.-F. Prévost & D. Sabatier* 2839, dbh = 10 cm.

[491] *Erythroxylum mucronatum* Benth.

London J. Bot. 2: 372 (Bentham 1843).

Erythroxylum comosum O.E.Schulz, *Pflanzenr. [Engler]* IV.134 (Heft 29): 32 [13 Aug. 1907] (Schulz 1907).

Erythroxylum kirkianum O.E.Schulz, *Repert. Spec. Nov. Regni Veg.* 30: 179 (Schulz 1932).

Erythroxylum venezuelense Steyermark., *Fieldiana, Bot.* 28 (2): 271 (Steyermark 1952).

Erythroxylum albertianum Kuhlmann & W.A.Rodrigues, *Publ. Inst. Nac. Pesq. Amaz., Bot.* 5: 3 (Kuhlmann & Rodrigues 1957).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2230*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15\text{ cm}$.

[492] *Erythroxylum roraimae* Klotzsch ex O.E.Schulz

Pflanzenr. [Engler] IV.134 (Heft 29): 104 [13 Aug. 1907] (Schulz 1907).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman 2238*.

SIZE. — Up to 10 m tall (Plowman & Berry 1999).

[493] *Erythroxylum squamatum* Sw.

Prodri. [Swartz]: 75 [20 Jun.-29 July 1788] (Swartz 1788).

Erythroxylum aristigerum Peyr., *Fl. Bras. [Martius]* 12 (1): 157 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum bahiense Peyr., *Fl. Bras. [Martius]* 12 (1): 160 [1 Dec. 1878] (Peyritsch 1878). — *Erythroxylum aristigerum* var. *bahiense* (Peyr.) O.E.Schulz, *Pflanzenr. [Engler]* IV.134 (Heft 29): 103 [13 Aug. 1907] (Schulz 1907).

Erythroxylum squamatum var. *microcarpum* O.E.Schulz, *Symb. Antill. [Urban]* 5 (2): 192 [20 May 1907] (Schulz 1907).

Erythroxylum trinerve Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 417 (Huber 1909).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P.A. Sagot 1229*, Mar. 1859 (holotype of *Erythroxylum squamatum* var. *microcarpum*; B? [not seen]; iso-, K[K000407437], P[P00723997], P00723998, P00723999], U[U0099038].

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.1\text{ cm}$.

[494] *Erythroxylum suberosum* A.St.-Hil.
(Fig. 21D)

A.Juss. & Cambess., *Pl. Usuel. Bras.* 14: t. 69, fig. A (Saint-Hilaire et al. 1828).

Erythroxylum areolatum Vell., *Fl. Flumin.* 194 ["1825" publ. 7 Sep. 28 Nov. 1829] (Vellozo 1829), "Erythroxylon", nom. illeg. hom., non L. (Linnaeus 1759).

Erythroxylum testaceum Peyr., *Fl. Bras. [Martius]* 12 (1): 170 [1 Dec. 1878] (Peyritsch 1878).

Erythroxylum suberosum f. *brevipetiolatum* O.E.Schulz, *Pflanzenr. [Engler]* IV.134 (Heft 29): 27 [13 Aug. 1907] (Schulz 1907).

VERNACULAR NAMES. — Br: cabelo-de-negro, galinha-choca.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5198*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 65.4\text{ cm}$.

Family EUPHORBIACEAE Juss.

Genus *Adenophaedra* (Müll.Arg.) Müll.Arg.

[495] *Adenophaedra grandifolia* (Klotzsch) Müll.Arg.

Fl. Bras. [Martius] 11 (2): 385 [1 May 1874] (Müller 1874).

Tragia grandifolia Klotzsch, *London J. Bot.* 2: 46 (Klotzsch 1843). — *Bernardia grandifolia* (Klotzsch) Müll.Arg., *Linnaea* 34: 173 (Müller 1865).

Cleidion denticulatum Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 218 [24 Oct. 1929] (Standley 1929). — *Bernardia denticulata* (Standl.) G.L.Webster, *Ann. Missouri Bot. Gard.* 54 (2): 200 [27 Oct. 1967] (Webster 1967).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 10656*.

SIZE. — Guyana. *H.D. Clarke et al. 9046* (MO), 12 m.

Genus *Alchornea* Sw.

[496] *Alchornea discolor* Poepp.

Nova genera ac species plantarum [Poeppig & Endlicher] 3: 19 [15-21 Aug. 1841] (Poeppig 1841).

Alchornea schomburgkii Klotzsch, *London J. Bot.* 2: 46 (Klotzsch 1843).

Alchornea gardneri Müll.Arg., *Flora* 47: 435 (Müller 1864).

Alchornea glaziovii Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.VII (Heft 63): 238 [10 Nov. 1914] (Pax & Hoffmann 1914), in nota.

Alchornea brachygyna Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XIV (Heft 68): 20 [6 June 1919] (Pax & Hoffmann 1919).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: wakavu-kamwi, waravru-kamwi • Wp: pesupi • Wn: kunui, uluimë • Nt: afatu adeeten • Br: amor-seco, sardinheiro, supiārana, tapiá, tartaruginha.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4830*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 52.4\text{ cm}$.

[497] *Alchornea triplinervia* (Spreng.) Müll.Arg.

Prodr. [A. P. de Candolle] 15 (2.2): 909 [late Aug. 1866] (Müller 1866). — *Antidesma triplinervium* Spreng., *Neue Entdeck. Pflanzenk.* 2: 116 (Sprengel 1821). — *Alchornea triplinervia* var. *genuina* Müll. Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866), *nom. inval.* (Turland et al. 2018: Art. 24.3).

Alchornea nemoralis Mart., *Flora 24 (2, Beibl.): 31* (Martius 1841). — *Alchornea triplinervia* var. *nemoralis* (Mart.) Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.VII (Heft 63): 228* [10 Nov. 1914] (Pax & Hoffmann 1914).

Alchornea janeirensis Casar., *Atti Riunione Sci. Ital. 3: 515* [“1841” publ. June 1842] (Casaretto 1842). — *Alchornea nemoralis* var. *janeirensis* (Casar.) Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865). — *Alchornea triplinervia* var. *janeirensis* (Casar.) Müll. Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866).

Alchornea parvifolia Miq., *Linnaea 22: 797* [“1849” publ. May 1850] (Miquel 1850). — *Alchornea triplinervia* var. *parvifolia* (Miq.) Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 910* [late Aug. 1866] (Müller 1866).

Alchornea glandulosa Poepp. var. *parvifolia* Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 330* (Bentham 1854). — *Alchornea triplinervia* var. *laevigata* Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 910* [late Aug. 1866] (Müller 1866), *nom. illeg. superf.* (*A. glandulosa* var. *parvifolia* in synonymy).

Alchornea intermedia Klotzsch ex Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 331* (Bentham 1854), *nom. nud.*

Alchornea parvifolia Klotzsch ex Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 331* (Bentham 1854), *nom. nud.*

Alchornea psilorhachis Klotzsch ex Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 331* (Bentham 1854), *nom. nud.*

Alchornea rotundifolia Moric. ex Baill., *Étude Euphorb.: 447* (Baillon 1858), *nom. inval.* (not accepted by the author).

Alchornea nemoralis var. *lanceolata* Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865). — *Alchornea triplinervia* var. *lanceolata* (Baill.) Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 910* [late Aug. 1866] (Müller 1866).

Alchornea nemoralis var. *parvifolia* Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865), *nom. nud.*

Alchornea nemoralis var. *psilorhachis* Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865), *nom. nud.*

Alchornea nemoralis var. *rotundifolia* Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865), *nom. nud.*

Alchornea triplinervia f. *intermedia* Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866). — *Alchornea nemoralis* var. *intermedia* Baill. ex Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866), *nom. nud. pro syn.*

Alchornea triplinervia f. *psilorhachis* Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866).

Alchornea triplinervia var. *crassifolia* Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 909* [late Aug. 1866] (Müller 1866).

Alchornea triplinervia var. *tomentella* Müll.Arg., *Fl. Bras. [Martius] 11 (2): 380* [1 May 1874] (Müller 1874).

Alchornea triplinervia var. *iricuranoides* Chodat & Hassl., *Bull. Herb. Boissier, sér. 2, 5: 603* (Chodat & Hassler 1905).

Alchornea acroneura Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.VII (Heft 63): 229* [10 Nov. 1914] (Pax & Hoffmann 1914).

Alchornea brevistyla Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.VII (Heft 63): 227* [10 Nov. 1914] (Pax & Hoffmann 1914).

Alchornea obovata Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.VII (Heft 63): 223* [10 Nov. 1914] (Pax & Hoffmann 1914).

Alchornea triplinervia var. *boliviiana* Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.VII (Heft 63): 227* [10 Nov. 1914] (Pax & Hoffmann 1914).

Alchornea triplinervia var. *trinitatis* L.Riley, *Bull. Misc. Inform. Kew 1925 (3): 141* [17 Apr. 1925] (Riley 1925).

VERNACULAR NAMES. — Pa: wakavu-kamwi, waravru-kamwi • Ka: alepa weli, kiyelapolan • Wp: yási leá̄y • Wn: pakutpé • Br: tapiá, tinteiro.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier 2346*.

INVENTORY DATA (FG). — 35 trees in 11 plots; $F_{\max} = 3.4\%$; $\text{dbh}_{\text{inv}} = 66.8$ cm.

Genus *Alchorneopsis* Müll.Arg.

[498] *Alchorneopsis floribunda* (Benth.) Müll.Arg.

Linnaea 34: 156 (Müller 1865). — *Alchornea glandulosa* var. *floribunda* Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 331* (Bentham 1854). — *Alchornea nemoralis* var. *floribunda* (Benth.) Baill., *Adansonia [Baillon] 5: 239* (Baillon 1865).

Alchorneopsis trimera Lanj., *Euphorb. Surinam 23* (Lanjouw 1931).

VERNACULAR NAMES. — Pa: wakavu-kamwi, waravru-kamwi • Ka: atapilipyo, wala ekone • Wp: mulei • Wn: pakutpé • Br: canelaraná.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *M.-F. Prévost 3456*.

INVENTORY DATA (FG). — 24 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 45$ cm.

Genus *Aparisthium* Endl.

[499] *Aparisthium cordatum* (A.Juss.) Baill.

Adansonia [Baillon] 5: 307 (Baillon 1865). — *Conceveibum cordatum* A.Juss., *Euphorb. Gen.: 43* [21 Feb. 1824] (Jussieu 1824), “*Conceveibum*”. — *Alchornea cordata* (A.Juss.) Müll.Arg., *Prodr. [A. P. de Candolle] 15 (2.2): 901* [late Aug. 1866] (Müller 1866), *nom. illeg. hom.*, *non Benth.* (Bentham 1849).

Alchornea macrophylla Mart., *Flora 24 (2, Beibl.): 31* (Martius 1841). — *Aparisthium macrophyllum* (Mart.) Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 333* (Bentham 1854). — *Conceveiba macrophylla* Klotzsch ex Benth., *Hooker's J. Bot. Kew Gard. Misc. 6: 333* (Bentham 1854), *nom. nud. pro syn.*

Alchornea latifolia Klotzsch, *London J. Bot.* 2: 46 (Klotzsch 1843), *nom. illeg. hom.*, non Sw. (Swartz 1788).

Styloceras macrostachyum Poepp. ex Baill., *Monog. Bux.*: 81 (Baillon 1859).

Aparisthium spruceanum Baill., *Adansonia [Baillon]* 5: 307 (Baillon 1865).

Alchornea orinocensis Croizat, *J. Arnold Arbor.* 26 (2): 191 [16 Apr. 1945] (Croizat 1945).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ahakyu-kamwi, ihap-kamwi, waikawavia-kamwi, wakavu-kamwi, waravru-kamwi • Ka: saulanani • Te: a'i mididju • Wp: a'i meyu • Nt: busi kusuwe • Br: mameleiro, morocototó, urucurana.

HERBARIUM DATA (FG). — 70 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (original material P[P05574226]).

INVENTORY DATA (FG). — 58 trees in 22 plots; $F_{\max} = 2.4\%$; $dbh_{\text{inv}} = 17.5$ cm.

Genus *Caryodendron* H.Karst.

[500] *Caryodendron amazonicum* Ducke

Trop. Woods 76: 18 (Ducke 1943).

VERNACULAR NAMES. — Br: castanha-de-porco.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier 4707*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 33.9$ cm.

Genus *Conceveiba* Aubl.

[501] *Conceveiba guianensis* Aubl.

Hist. Pl. Guiane 2: 924 [Jun.-Dec. 1775] (Aublet 1775).

Conceveibum ovatum A.Juss., *Euphorb. Gen.*: 43 [21 Feb. 1824] (Jussieu 1824), “*Conceveibum*”.

Conceveiba trigonocarpa Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 372 [1 May 1874] (Müller 1874).

Conceveiba simulata Steyermark., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (5)*: 416 [27 May 1938] (Steyermark 1938).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ihap-kamwi, wakavu-kamwi, waravru-kamwi • Ka: alemenango wewe, mapiwalan, yawalelan • Te: a'i mididju • Wp: a'i miniyu, aminiyu lá, miniyu lá • Wn: palulumuli • Nt: kusisi, lebi udu • Cr: oubarouna • Br: arara-seringa, arraieira, tapichó.

HERBARIUM DATA (FG). — 142 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000947432]).

INVENTORY DATA (FG). — 575 trees in 139 plots; $F_{\max} = 4.5\%$; $dbh_{\text{inv}} = 50$ cm.

[502] *Conceveiba krukoffii* Steyermark.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (5): 414 [27 May 1938] (Steyermark 1938).

HERBARIUM DATA (FG). — A single collection, *F. Billiet & B. Jadin 1230* (BR) (not seen, *fide* Secco 2004).

SIZE. — Up to 25 cm dbh (Secco 2004).

[503] *Conceveiba martiana* Baill.

(Fig. 23A)

Adansonia [Baillon] 5: 221 (Baillon 1865). — *Conceveibastrum martianum* (Baill.) Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147. VII (Heft 63)*: 217 [10 Nov. 1914] (Pax & Hoffmann 1914). — *Alchornea martiana* (Baill.) Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 375 [1 May 1874] (Müller 1874).

Conceveiba megalophylla Müll.Arg., *Linnaea* 34: 167 (Müller 1865).

VERNACULAR NAMES. — Wp: a'i meyu • Br: arraieira-branca.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *P. Grenand 3502*.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40.7$ cm.

Genus *Croton* L.

[504] *Croton cajucara* Benth.

Hooker's J. Bot. Kew Gard. Misc. 6: 376 (Bentham 1854). — *Oxydectes cajucara* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 611 [5 Nov. 1891] (Kuntze 1891).

Croton seputubensis Hoehne, *Exped. Roosevelt-Rondon. Bot.*: 53 (Hoehne 1914).

Croton motilonorum Croizat, *Mem. Soc. Ci. Nat. La Salle* 10: 126 (Croizat 1950).

VERNACULAR NAMES. — Br: sacaca.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *H. Richard & J. Ateni 432*.

SIZE. — Brazil, Maranhão. *J. Jangoux 1051* (MO), 10 m × 12 cm.

[505] *Croton cuneatus* Mart. ex Klotzsch

London J. Bot. 2: 49 (Klotzsch 1843). — *Macrocroton cuneatus* (Mart. ex Klotzsch) Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1186 [“1848” publ. 7-10 Mar. 1849] (Klotzsch 1849). — *Oxydectes cuneatus* (Mart. ex Klotzsch) Kuntze, *Revis. Gen. Pl.* 2: 611 [5 Nov. 1891] (Kuntze 1891).

Croton surinamensis Müll.Arg., *Linnaea* 34 (1): 82 [pre-Mar. 1865] (Müller 1865). — *Oxydectes surinamensis* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 613 [5 Nov. 1891] (Kuntze 1891).

Croton martii Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 88 [1 Feb. 1873] (Müller 1873). — *Oxydectes martii* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 613 [5 Nov. 1891] (Kuntze 1891).

Croton martii var. *latifolius* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 88 [1 Feb. 1873] (Müller 1873).

Croton martii var. *longifolius* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 88 [1 Feb. 1873] (Müller 1873).

Croton mimeticus S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 453 ["1894-96" publ. Dec. 1895] (Moore 1895).

Croton kaieteuri Jabl., *Mem. New York Bot. Gard.* 12 (3): 155 (Jablonszky 1965).

Croton subcoriaceus Jabl., *Mem. New York Bot. Gard.* 12 (3): 156 (Jablonszky 1965).

Croton monachinoensis Jabl., *Mem. New York Bot. Gard.* 12 (3): 157 (Jablonszky 1965).

NOTES. — *C. surinamensis* Müll.Arg. is based on *C. cuneatus* sensu Miquel (1848: 477); hence it is not a replacement name (see Turland et al. 2018: Art. 41.7, note 3). However, it is validly published since it refers to Miquel's description and types.

VERNACULAR NAMES. — Ka: tasi • Wp: mayawa'i • Wn: ehnajap, enajop, mojoi • Nt: ingi udu • Br: catinga-de-porco.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *M.-F. Prévost* 3847.

INVENTORY DATA (FG). — 1 tree, dbh = 24.8 cm.

[506] *Croton draconoides* Müll.Arg.
(Fig. 23B)

Linnaea 34 (1): 90 [pre-Mar. 1865] (Müller 1865). — *Oxydectes draconoides* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 611 [5 Nov. 1891] (Kuntze 1891), "draconodes".

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost* 185.

INVENTORY DATA (FG). — 3 trees in 1 plot; F_{max} < 1 %; dbh_{inv} = 20 cm.

[507] *Croton icabarui* Jabl.

Mem. New York Bot. Gard. 12 (3): 158 (Jablonszky 1965).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J. Engel et al.* 14, dbh = 10 cm.

[508] *Croton matourensis* Aubl.

Hist. Pl. Guiane 2: 879 [Jun.-Dec. 1775] (Aublet 1775), "Matourense". — *Croton sericeus* Lam., *Encycl. J.J. Lamarck et al. J2* (1): 210 [16 Oct. 1786] (Lamarck 1786), nom. illeg. superfl. (based on *Croton matourensis*). — *Oxydectes matourensis* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 612 [5 Nov. 1891] (Kuntze 1891).

Croton caryophyllus Benth., *Hooker's J. Bot. Kew Gard. Misc.* 6: 374 (Bentham 1854). — *Oxydectes caryophyllus* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 611 [5 Nov. 1891] (Kuntze 1891).

Croton matourensis var. *benthamianus* Müll.Arg., *Linnaea* 34 (1): 95 [pre-Mar. 1865] (Müller 1865).

Croton matourensis var. *poeppigianus* Müll.Arg., *Linnaea* 34 (1): 95 [pre-Mar. 1865] (Müller 1865).

Croton matourensis var. *sericeus* Müll.Arg., *Linnaea* 34 (1): 95 [pre-Mar. 1865] (Müller 1865).

Croton kavanayensis Steyermark., *Fieldiana, Bot.* 28 (2): 313 (Steyermark 1952).

Croton lanjouwianus Jabl., *Mem. New York Bot. Gard.* 12 (3): 158 (Jablonszky 1965), "lanjouwensis".

Croton pakaraimae Jabl., *Mem. New York Bot. Gard.* 12 (3): 159 (Jablonszky 1965).

VERNACULAR NAMES. — Pa: á-seiminio-kamwi, á-seiminio-kamwipriyu, mariuvra-wašiunó • Ka: kunapolan, mulewabo • Wp: mayawa'i • Wn: enajop, mojoi • Nt: ingi udu • Cr: bwa-ranmié • Fr: bois indien • Br: dima, maravuvuiá.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material S, *S-R-1308*).

INVENTORY DATA (FG). — 57 trees in 11 plots; F_{max} = 2.7 %; dbh_{inv} = 38.8 cm.

[509] *Croton palanostigma* Klotzsch

London J. Bot. 2: 48 (Klotzsch 1843). — *Oxydectes palanostigma* (Klotzsch) Kuntze, *Revis. Gen. Pl.* 2: 612 [5 Nov. 1891] (Kuntze 1891).

Croton benthamianus Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 105 [1 May 1874] (Müller 1874).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al.* 24778.

SIZE. — Up to 15 m tall (Gillespie 2002).

[510] *Croton schiedeanus* Schlechtendal

Linnaea 19: 243 ["1847" publ. Feb. 1846] (Schlechtendal 1846). — *Oxydectes schiedeana* (Schlechtendal) Kuntze, *Revis. Gen. Pl.* 2: 613 [5 Nov. 1891] (Kuntze 1891).

Croton perobtusus Lundell, *Phytologia* 1 (12): 405 [Oct. 1940] (Lundell 1940).

VERNACULAR NAMES. — Wp: pesupi.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4423.

INVENTORY DATA (FG). — 58 trees in 9 plots; F_{max} < 1 %; dbh_{inv} = 22.7 cm.

[511] *Croton* sp. A

NOTE. — This entity regroups specimens that are wrongly identified at CAY as *Croton tricolor* Klotzsch ex Baill., a species not present in French Guiana (Ricarda Riina, personal communication to JE, 2019).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-J. de Granville* 13403, dbh = 20 cm.



FIG. 23. — Euphorbiaceae: **A**, *Conceveiba martiana* Baill. (J.-F. Molino & D. Sabatier 2365); **B**, *Croton draconoides* Müll.Arg. (D. Sabatier 6013); **C**, *Glycycdendron amazonicum* Ducke (D. Sabatier & E. Fonty 5760); **D**, *Sapium glandulosum* (L.) Morong (J.-F. Molino & D. Sabatier 2255). A, D, © J.-F. Molino/IRD; B, C, © D. Sabatier/IRD.

Genus *Dodecastigma* Ducke

Dodecastigma mazarunense Croizat, *Bull. Torrey Bot. Club* 75 (4): 404 [Jul.-Aug. 1948] (Croizat 1948).

[512] *Dodecastigma integrifolium* (Lanj.) Lanj. & Sandwith

Kew Bull. 5 (1): 134 (Lanjouw & Sandwith 1950). — *Pausandra integrifolia* Lanj., *Bull. Misc. Inform. Kew* 1932 (4): 183 [13 June 1932] (Lanjouw 1932).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 22811*.

INVENTORY DATA (FG). — 7 trees in 2 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 23.4$ cm.

[513] *Dodecastigma uleanum* (Pax & K.Hoffm.)
G.L.Webster

Contr. Univ. Michigan Herb. 25: 237 [13 Aug. 2007] (Webster 2007). — *Cunuria uleana* Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.XIV* (Heft 68): 51 [6 June 1919] (Pax & Hoffmann 1919). — *Anomalocalyx uleanus* (Pax & K.Hoffm.) Ducke, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 344 [30 Mar. 1932] (Ducke 1932).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman B-3139*.

SIZE. — Up to 25 cm dbh (Secco 2005).

Genus *Glycydendron* Ducke

[514] *Glycydendron amazonicum* Ducke
(Fig. 23C)

Arch. Jard. Bot. Rio de Janeiro 3: 199 (Ducke 1922).

VERNACULAR NAMES. — Pa: paig-seine, pairi-seine • Nt: wandebole • Br: castanha-de-porco, mirindiba-doce, pau-doce.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *D. Sabatier & E. Fonty 5760*.

INVENTORY DATA (FG). — 106 trees in 82 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 92.3$ cm.

Genus *Hevea* Aubl.

[515] *Hevea guianensis* Aubl.

Hist. Pl. Guiane 2: 871 [Jun.-Dec. 1775] (Aublet 1775), “peruviana” on plate. — *Siphonia guianensis* (Aubl.) Juss. ex Baill., *Étude Euphorb.* 326 (Baillon 1858). — *Caoutchoua guianensis* (Aubl.) O.F.Cook, *J. Wash. Acad. Sci.* 31: 58 (Cook 1941).

Hevea peruviana Aubl., *Hist. Pl. Guiane* 4: t. 335 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: akabdat, katatu-avain • Ka: apalaba, mapalaba • Nt: afasi boyti, efea • Cr: évéa • Fr: hévéa • Br: seringueira-itaúba.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000812056]).

INVENTORY DATA (FG). — 214 trees in 70 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 63.9$ cm.

Genus *Hura* L.

[516] *Hura crepitans* L.

Sp. Pl. 2: 1008 [1 May 1753] (Linnaeus 1753).

Hura brasiliensis Willd., *Enum. Pl. [Willdenow]* 2: 997 [June 1809] (Willdenow 1809).

Hura strepens Willd., *Enum. Pl. [Willdenow]* 2: 997 [June 1809] (Willdenow 1809). — *Hura crepitans* var. *strepens* (Willd.) Müll. Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1230 [late Aug. 1866] (Müller 1866).

Hura senegalensis Baill., *Adansonia [Baillon]* 1: 77 (Baillon 1860). — *Hura crepitans* var. *senegalensis* (Baill.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1230 [late Aug. 1866] (Müller 1866).

Hura crepitans f. *oblongifolia* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1230 [late Aug. 1866] (Müller 1866).

Hura crepitans f. *orbicularis* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1229 [late Aug. 1866] (Müller 1866).

Hura crepitans f. *ovata* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1229 [late Aug. 1866] (Müller 1866).

Hura crepitans var. *genuina* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1229 [late Aug. 1866] (Müller 1866), nom. inval. (Turland et al. 2018: Art. 24.3).

Hura crepitans var. *membranacea* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1229 [late Aug. 1866] (Müller 1866).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: tenway • Ka: asiwakala, kumakalan • Wp: wasaku • Wn: mutata • Nt: posenti, postii • Cr: bwa-dyab • Fr: bois diable, pet du diable, sablier • Br: açau.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4857*.

INVENTORY DATA (FG). — 1 tree, dbh = 10.4 cm.

Genus *Mabea* Aubl.

[517] *Mabea piriri* Aubl.

Hist. Pl. Guiane 2: 867 [Jun.-Dec. 1775] (Aublet 1775).

Mabea maynensis Spruce ex Benth., *J. Proc. Linn. Soc., Bot.* 5: 7 [“1861” publ. June 1860] (Bentham 1860).

VERNACULAR NAMES. — Ka: yukuyapoi • Wp: awalapuna • Wn: isoimë watkí • Nt: busi bisangula • Cr: bwa-lélé • Fr: bois à calumet • Br: seringá, taquari.

HERBARIUM DATA (FG). — 122 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000799363]).

INVENTORY DATA (FG). — 143 trees in 43 plots; $F_{\max} = 3.9\%$; $\text{dbh}_{\text{inv}} = 28.3$ cm.

[518] *Mabea salicoides* Esser

Novon 3 (4): 349 [winter 1993] (Esser 1993).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15146*.

SIZE. — Up to 18 cm dbh (Gillespie 2002).

[519] *Mabea speciosa* Müll.Arg.

Fl. Bras. [Martius] 11 (2): 520 [1 May 1874] (Müller 1874).

Mabea caudata Pax & K.Hoffm., *Pflanzenr. [Engler] IV.147.IV* (Heft 52): 282 [13 Feb. 1912] (Pax & Hoffmann 1912).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: awalapuna, awalapuna sili, paa lo'o • Wn: tihstili epit • Cr: bwa-lélé • Br: seringá, taquari.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 4470*.

INVENTORY DATA (FG). — 48 trees in 20 plots; $F_{\max} = 2.2\%$; $\text{dbh}_{\text{inv}} = 29 \text{ cm}$.

[520] *Mabea subsessilis* Pax & K.Hoffm.

Pflanzenr. [Engler] IV.147.IV (Heft 52): 282 [13 Feb. 1912] (Pax & Hoffmann 1912).

Mabea argutissima Croizat, *Bull. Torrey Bot. Club* 67 (4): 288 [Apr. 1940] (Croizat 1940).

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2307*.

INVENTORY DATA (FG). — 14 trees in 4 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 28 \text{ cm}$.

[521] *Mabea taquari* Aubl.

Hist. Pl. Guiane 2: 870 [Jun.-Dec. 1775] (Aublet 1775).

Mabea schomburgkii Benth., *Hooker's J. Bot. Kew Gard. Misc.* 6: 365 (Bentham 1854).

Mabea taquari var. *angustifolia* Müll.Arg., *Prodri. [A. P. de Candolle] 15* (2.2): 1150 [late Aug. 1866] (Müller 1866).

VERNACULAR NAMES. — Ka: akale loway • Wn: isoimë watki.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *J.B. Aublets.s.n.* (original material A[A00055264], BM[BM000799374], F[V0056865F], P[P00716889]; possible original material LINN[LINN-HS 1470.2], MPU[MPU015282]).

SIZE. — Guyana. *H.D. Clarke 3146*, 12 m.

Genus *Maprounea* Aubl.

[522] *Maprounea guianensis* Aubl.

Hist. Pl. Guiane 2: 895 [Jun.-Dec. 1775] (Aublet 1775), “*Mapronnea Guyannensis*” on plate. — *Stillingia guianensis* (Aubl.) Baill., *Adansonia [Baillon]* 5: 332 (Baillon 1865). — *Excoecaria guianensis* (Aubl.) Baill., *Hist. Pl. [Baillon]* 5: 133 [Jan.-Apr. 1874] (Baillon 1874).

Aegopricum betulinum L.f., *Suppl. Pl.*: 413 [“1781” publ. Apr. 1782] (Linnaeus 1782), “*Aegopricon*”.

VERNACULAR NAMES. — Pa: iduras-veiti, iduras-vey, kagegut, karegut • Ka: alisigii, pilapisi • Wp: ka'a ki, kasila, yalakasila, yalakasilo • Nt: ayee wiwii, kusangula • Cr: radjé-chank • Br: vaquinha.

HERBARIUM DATA (FG). — 59 collections at CAY. Sel. exs.: *J.B. Aublets.s.n.* (original material BM[BM000073601, BM000073602], S[S-R-10787]).

INVENTORY DATA (FG). — 14 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 54.1 \text{ cm}$.

Genus *Micrandra* R.Br.

[523] *Micrandra elata* (Didr.) Müll.Arg.

Linnaea 34: 142 (Müller 1865). — *Pogonophyllum elatum* Didr., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1857: 145 (Didrichsen 1857).

Micrandra bracteosa Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 290 [1 Feb. 1873] (Müller 1873).

Micrandra glaziovii Pax, *Pflanzenr. [Engler] IV.147.I* (Heft 42): 20 [3 May 1910] (Pax 1910).

Micrandra brownsbergensis Lanj., *Euphorb. Surinam* 34 (Lanjouw 1931).

Micrandra santanderensis Croizat, *J. Arnold Arbor.* 24 (2): 169 [15 Apr. 1943] (Croizat 1943).

NOTE. — *Micrandra bracteosa* Müll.Arg. is not the basionym of *Cunuria bracteosa* Ducke nor that of *Cunuria spruceana* var. *bracteosa* (Ducke) R.E.Schult. Both names are synonyms of *Micrandra spruceana* (Baill.) R.E.Schult.

VERNACULAR NAMES. — Ka: mule'ilan, tipulu alome.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier et al. 4781*.

INVENTORY DATA (FG). — 151 trees in 13 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 47.1 \text{ cm}$.

[524] *Micrandra rossiana* R.E.Schult.

Bot. Mus. Leafl. 15 (8): 211 [29 May 1952] (Schultes 1952).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier et al. 5802*.

INVENTORY DATA (FG). — 16 trees in 2 plots; $F_{\max} = 1.9\%$; $\text{dbh}_{\text{inv}} = 38.2 \text{ cm}$.

Genus *Pausandra* Radlk.

[525] *Pausandra fordii* Secco

Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 3 (1): 60 (Secco 1987).

VERNACULAR NAMES. — Pa: á-agag.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *R.S. Cowan 38370* (holo-, NY[00272986] ; iso-, NY[00272985],).

INVENTORY DATA (FG). — 256 trees in 31 plots; $F_{\max} = 4.6\%$; $\text{dbh}_{\text{inv}} = 27.5 \text{ cm}$.

[526] *Pausandra martinii* Baill.

Adansonia [Baillon] 11: 92 (Baillon 1873), “*Martini*”.

Pausandra flagellarhachis Lanj., *Euphorb. Surinam* 30 (Lanjouw 1931).

VERNACULAR NAMES. — Pa: Kawap.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, P[P00648720]; iso-, P[P00648721]).

SIZE. — Up to 20 m tall (Gillespie 2002).

[527] *Pausandra trianae* (Müll.Arg.) Baill.

Adansonina [Baillon] 11: 92 (Baillon 1873). — *Pogonophora trianae* Müll.Arg., *Flora* 47: 434 (Müller 1864).

Pausandra quadriglandulosa Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XIV (Heft 68): 43 [6 June 1919] (Pax & Hoffmann 1919).

Pausandra extorris Standl., *Trop. Woods* 17: 24 [Mar. 1929] (Standley 1929).

Pausandra densiflora Lanj., *Recueil Trav. Bot. Néerl.* 33: 766 (Lanjouw 1936).

Pausandra sericea Lanj., *Recueil Trav. Bot. Néerl.* 33: 767 (Lanjouw 1936).

Clavija septentrionalis L.O.Williams, *Fieldiana, Bot.* 32 (12): 205 (Williams 1970).

VERNACULAR NAMES. — Br: orelha-de-burro.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *H. Jacquemin 2435*.

SIZE. — Up to 30 m tall (Webster & Huft 1988).

Genus *Sagotia* Baill.

[528] *Sagotia brachysepala* (Müll.Arg.) Secco

Acta Amazonica Supl. 15 (1-2): 81, 85 (Secco 1985). — *Sagotia racemosa* var. *brachysepala* Müll.Arg., *Flora* 47: 516 (Müller 1864).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost 2719*.

SIZE. — Up to 35 m tall (Secco 1999).

[529] *Sagotia racemosa* Baill.

Adansonina [Baillon] 1: 54 (Baillon 1860).

Sagotia racemosa var. *genuina* Müll.Arg., *Flora* 47: 516 (Müller 1864), nom. inval. (Turland *et al.* 2018: Art. 24.3).

Sagotia racemosa var. *ligularis* Müll.Arg., *Flora* 47: 516 (Müller 1864).

Sagotia racemosa var. *microsepala* Müll.Arg., *Flora* 47: 517 (Müller 1864).

Sagotia tafelbergii Croizat, *Bull. Torrey Bot. Club* 75 (4): 404 [Jul.-Aug. 1948] (Croizat 1948).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: tatu mila • Nt: makwebo.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *J. Martin 27* (original material P[P00648674]).

INVENTORY DATA (FG). — 251 trees in 52 plots; $F_{\max} = 4.2\%$; $dbh_{\text{inv}} = 38.2$ cm.

Genus *Sandwithia* Lanj.

[530] *Sandwithia guyanensis* Lanj.

Bull. Misc. Inform. Kew 1932 (4): 184 [13 June 1932] (Lanjouw 1932).

VERNACULAR NAMES. — Wp: wila tuwi • Br: urucurana-branca.

HERBARIUM DATA (FG). — 59 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3589*.

INVENTORY DATA (FG). — 677 trees in 75 plots; $F_{\max} = 6.2\%$; $dbh_{\text{inv}} = 40.7$ cm.

Genus *Sapium* Jacq.

[531] *Sapium argutum* (Müll.Arg.) Huber

Bull. Herb. Boissier, sér. 2, 6: 439 (Huber 1906). — *Excoecaria arguta* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 614 [1 May 1874] (Müller 1874), “*Excaecaria*”.

Sapium montanum Lanj., *Euphorb. Surinam* 47 (Lanjouw 1931).

VERNACULAR NAMES. — Pa: itey-avain, itey-avan.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *S.A. Mori et al. 25536*.

SIZE. — Up to 10 m tall (Santos & Sales 2009).

[532] *Sapium ciliatum* Hemsl.

Hooker's Icon. Pl. 27 [ser. 4, 7]: t. 2683 [Feb. 1901] (Hemsley 1901).

VERNACULAR NAMES. — Wp: melekene, melekene sili • Br: murupita.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 1964*.

SIZE. — Brazil, Pará. *B.V. Rabelo 3741* (MO), 15 m.

[533] *Sapium glandulosum* (L.) Morong
(Fig. 23D)

Ann. New York Acad. Sci. 7: 227 (Morong 1893). — *Hippomane glandulosa* L., *Sp. Pl.* 2: 1191 [1 May 1753] (Linnaeus 1753). — *Sapium jamaicense* Sw., *Adnot. Bot.*: 62 (Swartz 1829), nom. illeg. superfl. (includes the type of *Hippomane glandulosa*).

Sapium aucuparium Jacq., *Enum. Syst. Pl.*: 21 [Aug.-Sep. 1760] (Jacquin 1760). — *Hippomane aucuparia* (Jacq.) Crantz, *Inst. Rei Herb.* 1: 169 (Crantz 1766). — *Stillingia aucuparia* (Jacq.) Oken, *Allg. Naturgesch.* 3 (3): 1607 (Oken 1841). — *Sapium biglandulosum* var. *aucuparium* (Jacq.) Müll.Arg., *Linnaea* 32: 119 (Müller 1863). — *Excoecaria biglandulosa* var. *aucuparia* (Jacq.) Müll.Arg., *Prodri. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Hippomane biglandulosa L., *Sp. Pl.*, ed. 2, 2: 1431 [Aug. 1763] (Linnaeus 1763), nom. illeg. superfl. (based on *Sapium aucuparium*). — *Sapium hippomane* G.Mey., *Prim. Fl. Esseq.* 275 [Nov. 1818] (Meyer 1818), nom. illeg. superfl. (based on *Hippomane*

biglandulosum). — *Sapium biglandulosum* Müll.Arg., *Linnaea* 32: 116 (Müller 1863). — *Stillingia biglandulosa* Baill., *Adansonia [Baillon]* 5: 320 (Baillon 1865). — *Sapium montevidense* Klotzsch ex Baill., *Adansonia [Baillon]* 5: 320 (Baillon 1865), nom. nud. pro syn. — *Excoecaria biglandulosa* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1204 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium salicifolium Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 65 [28 Apr. 1817] (Kunth 1817). — *Sapium biglandulosum* var. *salicifolium* (Kunth) Müll.Arg., *Linnaea* 32: 119 (Müller 1863). — *Stillingia salicifolia* (Kunth) Klotzsch ex Baill., *Adansonia [Baillon]* 5: 320 (Baillon 1865), nom. illeg. hom., non (Torr.) Raf. (Rafinesque 1832). — *Excoecaria biglandulosa* var. *salicifolia* (Kunth) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1207 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Omphalea glandulata Vell., *Fl. Flumin. Icon.* 10: t. 14 [“1827” publ. 29 Oct. 1831] (Vellozo 1831). — *Sapium glandulatum* (Vell.) Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 229 [13 Feb. 1912] (Pax 1912). — *Excoecaria biglandulosa* var. *glandulata* (Vell.) Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 621 [1 May 1874] (Müller 1874), “*Excaecaria*”.

Sapium prunifolium Klotzsch, *London J. Bot.* 2: 45 (Klotzsch 1843). — *Stillingia prunifolia* (Klotzsch) Baill., *Étude Euphorb.* 513 (Baillon 1858). — *Excoecaria biglandulosa* var. *prunifolia* (Klotzsch) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1205 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium moritzianum Klotzsch, *Bot. Voy. Herald [Seemann]* 3: 100 [Nov. 1853] (Klotzsch 1853). — *Sapium biglandulosum* var. *moritzianum* (Klotzsch) Müll.Arg., *Linnaea* 32: 119 (Müller 1863). — *Excoecaria biglandulosa* var. *moritziana* (Klotzsch) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”, “*Moritzianum*”. — *Sapium aucuparium* subsp. *moritzianum* (Klotzsch) Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 128 [18 June 1918] (Pittier 1918).

Stillingia cremostachya Baill., *Étude Euphorb.* 512 (Baillon 1858). — *Sapium cremostachyum* (Baill.) I.M.Johnst., *Contr. Gray Herb.* 68: 91 (Johnston 1923).

Stillingia hippomane Baill., *Étude Euphorb.* 513 (Baillon 1858).

Sapium aucuparium var. *hippomane* Griseb., *Fl. Brit. W.I. [Grisebach]* 49 [Dec. 1859] (Grisebach 1859).

Sapium aereum Klotzsch ex Müll.Arg., *Linnaea* 32: 119 (Müller 1863). — *Excoecaria aerea* (Klotzsch ex Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1207 [late Aug. 1866] (Müller 1866), “*Excaecaria*”. — *Sapium aureum* H.Buek, *Gen. Sp. Synon. Cand.* 4: 364 [Jul.-Aug. 1874] (Buek 1874), sphalm.

Sapium biglandulosum f. *obovata* Müll.Arg., *Linnaea* 32: 116 (Müller 1863). — *Excoecaria biglandulosa* f. *obovata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1205 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium biglandulosum var. *aubletianum* Müll.Arg., *Linnaea* 32: 116 (Müller 1863). — *Excoecaria biglandulosa* var. *aubletiana* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1205 [late Aug. 1866] (Müller 1866), “*Excaecaria*”, “*Aubletianum*”. — *Sapium aubletianum* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 362 (Huber 1906).

Sapium biglandulosum var. *hamatum* Müll.Arg., *Linnaea* 32: 116 (Müller 1863). — *Excoecaria biglandulosa* var. *hamata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1204 [late Aug. 1866] (Müller 1866). — *Sapium hamatum* (Müll.Arg.) Pax &

K.Hoffm., *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 229 [13 Feb. 1912] (Pax & Hoffmann 1912).

Sapium biglandulosum f. *pavonianum* Müll.Arg., *Linnaea* 32: 116 (Müller 1863), “*Peruvianum*”. — *Excoecaria biglandulosa* var. *pavoniana* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1204 [late Aug. 1866] (Müller 1866), “*Excaecaria*”, “*Pavoniana*”. — *Sapium pavonianum* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 356 (Huber 1906).

Sapium obtusilobum Müll.Arg., *Linnaea* 32: 116 (Müller 1863). — *Excoecaria obtusiloba* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1203 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium biglandulosum var. *meyerianum* Müll.Arg., *Linnaea* 32: 116 (Müller 1863).

Sapium biglandulosum f. *oblongatum* Müll.Arg., *Linnaea* 32: 117 (Müller 1863). — *Excoecaria biglandulosa* f. *oblongata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1205 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium biglandulosum var. *cuneatum* Müll.Arg., *Linnaea* 32: 117 (Müller 1863). — *Excoecaria biglandulosa* var. *cuneata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium biglandulosum var. *klotzschianum* Müll.Arg., *Linnaea* 32: 117 (Müller 1863). — *Excoecaria biglandulosa* var. *klotzschiana* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1205 [late Aug. 1866] (Müller 1866), “*Excaecaria*”, “*Klotzschianum*”. — *Sapium klotzschianum* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 438 (Huber 1906).

Sapium biglandulosum f. *minus* Müll.Arg., *Linnaea* 32: 117 (Müller 1863).

Sapium biglandulosum var. *lanceolatum* Müll.Arg., *Linnaea* 32: 118 (Müller 1863). — *Excoecaria biglandulosa* var. *lanceolata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”. — *Excoecaria marginata* var. *lanceolata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1208 [late Aug. 1866] (Müller 1866), “*Excaecaria*”. — *Sapium lanceolatum* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 441 (Huber 1906).

Sapium biglandulosum var. *serratum* Müll.Arg., *Linnaea* 32: 118 (Müller 1863). — *Sapium serratum* (Müll.Arg.) Klotzsch ex Baill., *Adansonia [Baillon]* 5: 320 (Baillon 1865). — *Excoecaria biglandulosa* var. *serrata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium marginatum Müll.Arg., *Linnaea* 32: 120 (Müller 1863). — *Excoecaria marginata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1208 [late Aug. 1866] (Müller 1866), nom. illeg. hom., non Kunze ex Baill. (Baillon 1858). — *Stillingia marginata* (Müll.Arg.) Baill., *Adansonia [Baillon]* 5: 321 (Baillon 1865).

Sapium marginatum f. *majus* Müll.Arg., *Linnaea* 32: 120 (Müller 1863). — *Excoecaria marginata* f. *major* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1208 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium marginatum var. *spathulatum* Müll.Arg., *Linnaea* 32: 120 (Müller 1863). — *Excoecaria marginata* var. *spathulata* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1208 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.

Sapium marginatum var. *lanceolatum* Müll.Arg., *Linnaea* 32: 120 (Müller 1863).

- Stillingia dracunculoides* Baill., *Adansonia [Baillon]* 5: 321 (Baillon 1865), in obs. — *Excoecaria biglandulosa* var. *dracunculoides* (Baill.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1206 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.
- Sapium suberosum* Müll.Arg., *Linnaea* 34: 217 (Müller 1865). — *Excoecaria suberosa* (Müll.Arg.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1202 [late Aug. 1866] (Müller 1866).
- Excoecaria biglandulosa* var. *hippomane* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 1204 [late Aug. 1866] (Müller 1866), “*Excaecaria*”.
- Excoecaria biglandulosa* var. *clauseniana* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 618 [1 May 1874] (Müller 1874), “*Excaecaria*”, “*Clauseniana*”. — *Sapium clausenianum* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 436 (Huber 1906).
- Excoecaria biglandulosa* var. *intercedens* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 620 [1 May 1874] (Müller 1874). — *Sapium intercedens* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 437 (Huber 1906).
- Excoecaria biglandulosa* var. *grandifolia* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 622 [1 May 1874] (Müller 1874), “*Excaecaria*”.
- Excoecaria biglandulosa* var. *leptadenia* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 620 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium leptadenium* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 436 (Huber 1906).
- Excoecaria biglandulosa* var. *montevidensis* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 621 [1 May 1874] (Müller 1874).
- Excoecaria biglandulosa* var. *petiolaris* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 621 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium aucuparium* var. *petiolare* (Müll.Arg.) Griseb., *Abh. Königl. Ges. Wiss. Göttingen* 24: 62 (Grisebach 1879). — *Sapium petiolare* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 434 (Huber 1906).
- Excoecaria marginata* var. *conjugens* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 617 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium marginatum* var. *conjugens* (Müll.Arg.) Müll.Arg., *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 224 [13 Feb. 1912] (Müller 1912).
- Excoecaria marginata* var. *grandifolia* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 618 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium marginatum* var. *grandifolium* (Müll.Arg.) Müll.Arg., *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 224 [13 Feb. 1912] (Müller 1912).
- Excoecaria marginata* var. *longifolia* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 618 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium marginatum* var. *longifolium* (Müll.Arg.) Müll.Arg., *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 224 [13 Feb. 1912] (Müller 1912).
- Excoecaria occidentalis* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 615 [1 May 1874] (Müller 1874), “*Excaecaria*”. — *Sapium occidentale* (Müll.Arg.) Huber, *Bull. Herb. Boissier*, sér. 2, 6: 441 (Huber 1906).
- Sapium poeppigii* Hemsl., *Hooker's Icon. Pl.* 27 [ser. 4, 7]: t. 2678 [Feb. 1901] (Hemsley 1901).
- Sapium caribaeum* Urb., *Symb. Antill. [Urban]* 3 (2): 308 [15 Aug. 1902] (Urban 1902).
- Sapium taburu* Ule, *Bot. Jahrb. Syst.* 35 (5): 671 [18 Apr. 1905] (Ule 1905).
- Sapium pittieri* Huber, *Bull. Herb. Boissier*, sér. 2, 6: 350 (Huber 1906).
- Sapium bogotense* Huber, *Bull. Herb. Boissier*, sér. 2, 6: 355 (Huber 1906). — *Sapium biglandulosum* var. *bogotense* (Huber) Monach., *Bull. Torrey Bot. Club* 67 (9): 772 [Dec. 1940] (Monachino 1940).
- Sapium hemsleyanum* Huber, *Bull. Herb. Boissier*, sér. 2, 6: 362 (Huber 1906).
- Sapium submarginatum* Huber, *Bull. Herb. Boissier*, sér. 2, 6: 443 (Huber 1906).
- Sapium oligoneurum* K.Schum. & Pittier, *Contr. U.S. Natl. Herb.* 12: 168 (Schumann & Pittier 1908). — *Sapium biglandulosum* var. *oligoneurum* (K.Schum. & Pittier) Monach., *Bull. Torrey Bot. Club* 67 (9): 772 [Dec. 1940] (Monachino 1940).
- Sapium sulciferum* Pittier, *Contr. U.S. Natl. Herb.* 12: 169 (Pittier 1908). — *Sapium biglandulosum* var. *sulciferum* (Pittier) Monach., *Bull. Torrey Bot. Club* 67 (9): 772 [Dec. 1940] (Monachino 1940).
- Sapium fendleri* Hemsl., *Hooker's Icon. Pl.* 29 [ser. 4, 9]: sub t. 2888 [June 1909] (Hemsley 1909).
- Sapium muelleri* Hemsl., *Hooker's Icon. Pl.* 29 [ser. 4, 9]: t. 2884 [June 1909] (Hemsley 1909).
- Hippomane zeocca* L. ex B.D.Jacks., *Index Linn. Herb.*: 86 (Jackson 1912), nom. nud.
- Sapium endlicherianum* Klotzsch ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 228 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium obtusatum* Klotzsch ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 224 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium poblianum* Klotzsch ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 228 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium punctatum* Klotzsch ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 223 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium pycnostachys* K.Schum. ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 208 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium subserratum* Klotzsch ex Pax, *Pflanzenr. [Engler]* IV.147.IV (Heft 52): 226 [13 Feb. 1912] (Pax 1912), nom. nud. pro syn.
- Sapium caudatum* Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 127 [18 June 1918] (Pittier 1918).
- Sapium giganteum* Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 128 [18 June 1918] (Pittier 1918).
- Sapium albomarginatum* Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XVII (Heft 85): 203 [16 May 1924] (Pax & Hoffmann 1924).
- Sapium fragile* Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XVII (Heft 85): 202 [16 May 1924] (Pax & Hoffmann 1924).
- Sapium paranaense* Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XVII (Heft 85): 203 [16 May 1924] (Pax & Hoffmann 1924).
- Sapium guaricense* Pittier, *J. Wash. Acad. Sci.* 19: 355 (Pittier 1929).
- Sapium naiguatense* Pittier, *J. Wash. Acad. Sci.* 19: 355 (Pittier 1929).
- Sapium paucistamineum* Pittier, *J. Wash. Acad. Sci.* 19: 356 (Pittier 1929).
- Sapium integrifolium* Splitg. ex Lanj., *Euphorb. Surinam* 176 (Lanjouw 1931), nom. nud. pro syn.
- Sapium klotzschianum* var. *glaziovii* Lanj., *Euphorb. Surinam* 46 (Lanjouw 1931).
- Stillingia haematantha* Standl., *Ann. Missouri Bot. Gard.* 27: 314 (Standley 1940).

Sapium haematospermum Müll.Arg. var. *saltense* O'Donell & Lourteig, *Lilloa* 8: 586 (O'Donell & Lourteig 1942), “*saltensis*”. — *Sapium saltense* (O'Donell & Lourteig) Jabl., *Phytologia* 14 (7): 451 [15 May 1967] (Jablonszky 1967).

Sapium schippii Croizat, *Amer. Midl. Naturalist* 29 (2): 477 [5 Apr. 1943] (Croizat 1943).

Sapium contortum Croizat, *J. Arnold Arbor.* 26 (2): 193 [16 Apr. 1945] (Croizat 1945).

Sapium nitidum Alain, *Contr. Ocas. Mus. Hist. Nat. Colegio De La Salle* 11: 8 (Alain 1952), *nom. illeg. hom.*, *non* (Monach.) Lundell (1943) [synonym of *S. lateriflorum* Hemsl.]

Sapium moaense Alain, *Revista Soc. Cub. Bot.* 10: 27 (Alain 1953).

Sapium ixiamasense Jabl., *Phytologia* 14 (7): 450 [15 May 1967] (Jablonszky 1967).

Sapium itzanum Lundell, *Wrightia* 5 (4): 77 (Lundell 1975).

Sapium izabalense Lundell, *Wrightia* 5 (9): 346 (Lundell 1977).

Sapium alainianum P.T.Li, *Guibaea* 14 (2): 130 (Li 1994).

NOTES. — *Excoecaria biglandulosa* Müll.Arg., *Sapium biglandulosum* Müll.Arg. and *Stillingia biglandulosa* Baill. are based on the illegitimate *Hippomane biglandulosa* L. *S. biglandulosum* is itself illegitimate, because it includes the type of *H. glandulosa*.

VERNACULAR NAMES. — Pa: itey-avain, itey-avan • Ka: alekosin, kumakalan • Nt: meliki udu • Cr: bwa-dilèt • Br: burra-leiteira, murupita.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2255*.

INVENTORY DATA (FG). — 1 tree, dbh = 28.5 cm.

[534] *Sapium paucinervium* Hemsl.

Hooker's Icon. Pl. 27 [ser. 4, 7]: t. 2648 [“1901” publ. May 1900] (Hemsley 1900).

Sapium microdentatum Lanj., *Euphorb. Surinam* 46 (Lanjouw 1931).

VERNACULAR NAMES. — Br: murupita.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2970*.

INVENTORY DATA (FG). — 21 trees in 12 plots; F_{max} = 1 %; dbh_{inv} = 43 cm.

Family GOUPIACEAE Miers
Genus *Gouphia* Aubl.

[535] *Gouphia glabra* Aubl.
(Fig. 24A)

Hist. Pl. Guiane 1: 296 [Jun.-Dec. 1775] (Aublet 1775). — *Glossopetalum glabrum* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 510 [late Sep.-Nov. 1791] (Gmelin 1791).

Gouphia tomentosa Aubl., *Hist. Pl. Guiane* 1: 296 [Jun.-Dec. 1775] (Aublet 1775). — *Glossopetalum tomentosum* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 510 [late Sep.-Nov. 1791] (Gmelin 1791).

Gouphia paraensis Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 86 (Huber 1910).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: pasis • Ka: kubi'i, kupi'i • Te: patsitsi • Wp: pasisi • Wn: walima • Nt: kopi • Cr: bwa-kaka, goupi • Fr: bois caca, goupi • Br: cupiúba.

HERBARIUM DATA (FG). — 151 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00778564] designated by Lanjouw & Uittien [1940: 151]).

INVENTORY DATA (FG). — 382 trees in 134 plots; F_{max} = 26.2 %; dbh_{inv} = 158 cm.

Family HERNANDIACEAE Blume
Genus *Hernandia* Plum. ex L.

[536] *Hernandia guianensis* Aubl.
(Fig. 24B)

Hist. Pl. Guiane 2: 848 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Pa: maoksi-adudu • Ka: ayowo, tokolo enulu • Wp: wali wowo • Nt: popolika • Cr: bwa-amadou, bwa-bannan • Br: ventosa.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000993866], G[G00355749]).

INVENTORY DATA (FG). — 11 trees censused, 1 in 1 plot; dbh_{inv} = 57 cm.

Family HUMIRIACEAE A.Juss.
Genus *Humiria* J.St.-Hil.

[537] *Humiria balsamifera* Aubl.

Hist. Pl. Guiane 1: 564 [Jun.-Dec. 1775] (Aublet 1775), “*Houmiri*”. — *Myrodendrum balsamiferum* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 817 [late Sep.-Nov. 1791] (Gmelin 1791).

Myrodendrum amplexicaule Willd., *Sp. Pl.*, ed. 4 2 (2): 1171 [Dec. 1799] (Willdenow 1799), *nom. illeg. superfl.* (based on *Humiria balsamifera*). — *Humiria amplexicaulis* Mart. ex Urb., *Fl. Bras. [Marius]* 12 (2): 440 [1 Dec. 1877] (Urban 1877), “*Humirium amplexicaule*”, *nom. nud. pro syn.*

Humiria arenaria Guill. ex Baill., *Adansonia [Baillon]* 1: 209 (Baillon 1860), “*Humirium arenarium*”.

Humiria balsamifera f. *attenuata* Cuatrec., *Contr. U.S. Natl. Herb.* 35 (2): 97 [14 Apr. 1961] (Cuatrecasas 1961).

NOTE. — In order to limit the inflation of binomials, we have chosen not to list here the orthographic variants “*Humirium*” and “*Myrodendron*”, as well as all binomial based on them.

VERNACULAR NAMES. — Pa: psum, umeg • Ka: meli • Wp: weli • Nt: aneisi udu • Cr: bwa-rouj • Fr: houmiri baumier • Br: turamira, umiri.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000795994]).

INVENTORY DATA (FG). — 28 trees in 16 plots; F_{max} < 1 %; dbh_{inv} = 144.2 cm.

Genus *Humiriastrum* (Urb.) Cuatrec.

[538] *Humiriastrum excelsum* (Ducke) Cuatrec.

Contr. U.S. Natl. Herb. 35 (2): 133 [14 Apr. 1961] (Cuatrecasas 1961). — *Sacoglottis excelsa* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 178 (Ducke 1922).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4066.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 63.3$ cm.

[539] *Humiriastrum subcrenatum* (Benth.) Cuatrec.

Contr. U.S. Natl. Herb. 35 (2): 138 [14 Apr. 1961] (Cuatrecasas 1961). — *Humiria subcrenata* Benth., *London J. Bot.* 2: 374 (Bentham 1843), “*Humirium subcrenatum*”. — *Sacoglottis subcrenata* (Benth.) Urb., *Sitzungsber. Ges. Naturf. Freunde Berlin* 5 (Urban 1878).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: ašiu, ašiwa, tumau-atamwa • Nt: lebi kookoo.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J. Martin s.n.* (type K[K000407329]).

INVENTORY DATA (FG). — 112 trees in 53 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 75.8$ cm.

Genus *Sacoglottis* Mart.

[540] *Sacoglottis cydonioides* Cuatrec.
(Fig. 24C)

Contr. U.S. Natl. Herb. 35 (2): 183 [14 Apr. 1961] (Cuatrecasas 1961).

VERNACULAR NAMES. — Pa: ašiu, ašiwa, tumau-atamwa • Ka: pelulu • Wp: asiwa • Cr: bwa-kochon, bwa-maypouri.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *D. Sabatier & J.-L. Smock* 5782.

INVENTORY DATA (FG). — 31 trees in 22 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 73$ cm.

[541] *Sacoglottis guianensis* Benth.

Hooker's J. Bot. Kew Gard. Misc. 5: 104 (Bentham 1853).

Sacoglottis amazonica Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 104 (Bentham 1853), nom. illeg. hom., non Mart. (Martius 1827).

Sacoglottis guianensis f. *dolichocarpa* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 179 (Ducke 1922).

Sacoglottis guianensis var. *maior* Ducke, *Arch. Inst. Biol. Veg.* 4 (1): 27 [June 1938] (Ducke 1938).

Sacoglottis guianensis var. *hispidula* Cuatrec., *Contr. U.S. Natl. Herb.* 35 (2): 180 [14 Apr. 1961] (Cuatrecasas 1961).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: ašiu, ašiwa, tumau-atamwa • Ka: apesiya, yapopali • Nt: bofo udu, bofoo udu • Cr: bwa-kochon, bwa-maypouri • Br: achuá, uchirana.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5202.

INVENTORY DATA (FG). — 182 trees in 72 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 61.3$ cm.

Genus *Schistostemon* (Urb.) Cuatrec.

[542] *Schistostemon dichotomus* (Urb.) Cuatrec.

Contr. U.S. Natl. Herb. 35 (2): 159 [14 Apr. 1961] (Cuatrecasas 1961). — *Sacoglottis dichotoma* Urb., *Fl. Bras. [Martius]* 12 (2): 446 [1 Dec. 1877] (Urban 1877), “*Saccoglottis*”.

NOTE. — The Greek root “*stemon*” is masculine, hence *dichotomus*.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *C. Sastre et al.* 4125.

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[543] *Schistostemon sylvaticus* Sabatier

Proc. Kon. Ned. Akad. Wetensch. C 90 (2): 206 (Sabatier 1987).

NOTES. — Known only from French Guiana. The Greek root “*stemon*” is masculine, hence *sylvaticus*.

VERNACULAR NAMES. — Nt: bofo udu, bofoo udu.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier* 842 (holo-, P[P00077220]; iso-, P[not seen], U[U0008171]).

INVENTORY DATA (FG). — 17 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 56$ cm.

Genus *Vantanea* Aubl.

[544] *Vantanea guianensis* Aubl.

Hist. Pl. Guiane 1: 572 [Jun.-Dec. 1775] (Aublet 1775). — *Lemniscia guianensis* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 817 [late Sep.-Nov. 1791] (Gmelin 1791). — *Lemniscia floribunda* Willd., *Sp. Pl.*, ed. 4 2 (2): 1172 [Dec. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Vantanea guianensis*).

VERNACULAR NAMES. — Pa: ašiu-kamwi • Te: tepotsi átā, wila pikikit • Nt: bofo udu, bofoo udu • Br: uchirana.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000795993], S[S-R-9886]).

INVENTORY DATA (FG). — 34 trees in 17 plots; $F_{\max} = 2.3\%$; $\text{dbh}_{\text{inv}} = 73$ cm.




FIG. 24. — Goupiaceae: **A**, *Gouphia glabra* Aubl. Hernandiaceae: **B**, *Hernandia guianensis* Aubl. (M.-F. Prévost 3500). Humiriaceae: **C**, *Sacoglossis cydonioides* Cuatrec. (D. Sabatier & J.-L. Smock 5782); **D**, *Vantanea maculicarpa* Sabatier & J.Engel (D. Sabatier 5574). © D. Sabatier/IRD.

[545] *Vantanea maculicarpa* Sabatier & J.Engel
(Fig. 24D)

Phytotaxa 338 (1): 130 [09 Feb. 2018] (Sabatier & Engel 2018).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: [D. Sabatier 5574](#) (holo-, CAY[CAY111685]; iso-, CAY[CAY109439], P[P01156374]).

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1 \%$; dbh_{inv} = 66.5 cm.

[546] *Vantanea ovicarpa* Sabatier

Brittonia 54 (4): 233 ["2002" publ. 16 Apr. 2003] (Sabatier 2003).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: [S.A. Mori & N.P. Smith 25047](#) (holo-, NY[00718019, 00718020]; iso-, CAY[CAY015333], P[P00710334], US[00810841]).

SIZE. — Up to 50 cm dbh (Sabatier 2002).

[547] *Vantanea parviflora* Lam.

J. Hist. Nat. 1: 145 (Lamarck 1792).

Vantanea cupularis Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 83 (Huber 1910).

VERNACULAR NAMES. — Pa: ahavuiye-washiune • Nt: adugwe • Br: macucá-murici, quebra-machado, uchirana.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (holo-, P-LA?, not seen; iso-, G[G00368596]).

INVENTORY DATA (FG). — 87 trees in 49 plots; $F_{\max} = 3.1\%$; $dbh_{\text{inv}} = 88\text{ cm}$.

Family HYPERICACEAE Juss.
Genus *Vismia* Vand.

[548] *Vismia cayennensis* (Jacq.) Pers.

Syn. Pl. [Persoon] 2 (1): 86 [Nov. 1806] (Persoon 1806). — *Hypericum cayennense* Jacq., *Enum. Syst. Pl.*: 28 [Aug.-Sep. 1760] (Jacquin 1760). — *Caopia cayennensis* (Jacq.) Kuntze, *Revis. Gen. Pl.* 1: 58 [5 Nov. 1891] (Kuntze 1891).

Hypericum eugeniiifolium Willd. ex Spreng., *Syst. Veg. [Sprengel] 3*: 350 [Jan.-Mar. 1826] (Sprengel 1826), “*eugeniaeifolium*”, nom. nud. pro syn.

Vismia floribunda Sprague, *Trans. & Proc. Bot. Soc. Edinburgh* 22 (4): 428 (Sprague 1905).

VERNACULAR NAMES. — Pa: suwimba, suwimpa • Ka: siwinyani, suwinyani enekan • Wp: suwilani • Wn: osi epít • Nt: lebi pindya udu, pindya udu • Cr: bwa-dat • Br: lacre-branco.

HERBARIUM DATA (FG). — 108 collections at CAY. Sel. exs.: The type from French Guiana (not traced); *W.J. Hahn 3534*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.6\text{ cm}$.

[549] *Vismia gracilis* Hieron.

Bot. Jahrb. Syst. 20 (3, Beibl. 49): 52 [9 Apr. 1895] (Hieronymus 1895). — *Caopia gracilis* Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 52 [9 Apr. 1895] (Hieronymus 1895).

Vismia glaziovii Ruhland, *Bot. Jahrb. Syst.* 30 (Beibl. 67): 27 [12 Mar. 1901] (Ruhland 1901).

Vismia amazonica Ewan, *Contr. U.S. Natl. Herb.* 35 (5): 355 [5 June 1962] (Ewan 1962).

Vismia buchtienii Ewan, *Contr. U.S. Natl. Herb.* 35 (5): 353 [5 June 1962] (Ewan 1962).

NOTE. — *Vismia gracilis* Hieron. and *Caopia gracilis* Hieron. were published simultaneously, but before 1953. Although alternative, both names are valid.

VERNACULAR NAMES. — Pa: suwimba, suwimpa • Wp: suwilani • Cr: bwa-dat • Br: lacre, pau-lacre.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier 2317*, dbh 10 cm.

[550] *Vismia guianensis* (Aubl.) Choisy

Prodr. Monogr. Hypéric.: 34 [9 Mar. 1821] (Choisy 1821), “*guyanensis*”. — *Hypericum guianense* Aubl., *Hist. Pl. Guiane* 2: 784 [Jun.-Dec. 1775] (Aublet 1775), “*Guianense*”, “*Hypericum Guyanense*” on plate. — *Vismia guttifera* Pers. var. *guianensis* (Aubl.) Pers., *Syn. Pl. [Persoon] 2 (1)*: 86 [Nov. 1806] (Persoon 1806). — *Caopia guianensis* (Aubl.) A.Lyons, *Pl. Nam.*, ed. 2: 94 (Lyons 1907), “*guianensis*”.

Hypericum acuminatum Lam., *Encycl. [J. Lamarck et al.] 4 (1)*: 150 [9 Feb. 1797] (Lamarck 1797). — *Vismia acuminata* (Lam.) Pers., *Syn. Pl. [Persoon] 2 (1)*: 86 [Nov. 1806] (Persoon 1806). — *Caopia acuminata* (Lam.) Kuntze, *Revis. Gen. Pl.* 1: 59 [5 Nov. 1891] (Kuntze 1891). — *Vismia guianensis* var. *acuminata* (Lam.) M.E.Berg, *Bol. Mus. Paraense Emílio Goeldi, N.S.*, Bot. 40: 7 (Berg 1971).

Vismia caparosa Kunth, *Nova genera et species plantarum [H.B.K.] 5*: 182 [25 Feb. 1822] (Kunth 1822).

Vismia ferruginea Kunth, *Nova genera et species plantarum [H.B.K.] 5*: 183 [25 Feb. 1822] (Kunth 1822).

VERNACULAR NAMES. — Ka: siwinyani, suwinyani, tamunen suwinyani • Wn: osi epít • Nt: pindya udu, weti pindya udu • Cr: bwa-dat • Fr: bois baptiste, bois dartre, bois de sang (*fide* Aublet 1775) • Br: lacre-vermelho, uchirana.

HERBARIUM DATA (FG). — 73 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000624909]).

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} = 5.8\%$; $dbh_{\text{inv}} = 38.8\text{ cm}$.

[551] *Vismia latifolia* (Aubl.) Choisy
(Fig. 25A)

Prodr. Monogr. Hypéric. 36 [9 Mar. 1821] (Choisy 1821). — *Hypericum latifolium* Aubl., *Hist. Pl. Guiane* 2: 787 [Jun.-Dec. 1775] (Aublet 1775). — *Caopia latifolia* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 58 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Ka: saipyalala • Wp: suwilani pilá • Wn: osi epít • Nt: lebi pindya udu, pindya udu • Cr: bwa-dat • Br: lacre, pau-lacre.

HERBARIUM DATA (FG). — 84 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4822*.

INVENTORY DATA (FG). — 12 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.2\text{ cm}$.

[552] *Vismia macrophylla* Kunth

Nova genera et species plantarum [H.B.K.] 5: 184 [25 Feb. 1822] (Kunth 1822).

Hypericum reticulatum Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 694 [3 Sep. 1814] (Poirier 1814). — *Vismia reticulata* (Poir.) Choisy, *Prodr. Monogr. Hypéric.* 34 [9 Mar. 1821] (Choisy 1821).

Vismia angusta Miq., *Linnaea* 18: 27 [24-26 Oct. 1844] (Miquel 1844).

Vismia macrophylla var. *glabrescens* Hochr., *Annuaire Conserv. Jard. Bot. Genève* 21: 53 (Hochreutiner 1919).

VERNACULAR NAMES. — Pa: suwimba-purubumna, suwimpa-purubumna • Ka: saipyalala • Cr: bwa-dat • Br: lacre, pau-lacre.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4068*, 18 m × 15 cm.

[553] *Vismia ramuliflora* Miq.

Stirp. Surinam. Select.: 88 ["1850" publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Wn: pëmulimë.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2810*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 32.8$ cm.

[554] *Vismia sandwithii* Ewan

Contr. U.S. Natl. Herb. 35 (5): 309 [5 June 1962] (Ewan 1962).

VERNACULAR NAMES. — Pa: suwimba-ahavukune, suwimpa-ahau-wukune • Wp: suwilani, suwilani wu • Wn: osi epít • Br: lacre-vermelho-do-mato.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 890*.

SIZE. — Up to 10 m tall (Pipoly *et al.* 1998).

[555] *Vismia sessilifolia* (Aubl.) Choisy

Prodr. [A. P. de Candolle] 1: 542 [mid Jan. 1824] (Choisy 1824). — *Hypericum sessilifolium* Aubl., *Hist. Pl. Guiane* 2: 787 [Jun.-Dec. 1775] (Aublet 1775). — *Caopia sessilifolia* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 59 [5 Nov. 1891] (Kuntze 1891). — *Vismia cayennensis* var. *sessilifolia* (Aubl.) M.E.Berg, *Acta Amazonica* 4 (3): 16 (Berg 1974).

Hypericum rufescens Lam., *Encycl. [J. Lamarck et al.]* 4 (1): 150 [9 Feb. 1797] (Lamarck 1797). — *Vismia rufescens* (Lam.) Pers., *Syn. Pl. [Persoon]* 2 (1): 86 [Nov. 1806] (Persoon 1806).

VERNACULAR NAMES. — Pa: suwimba-kamwi, suwimpa-kamwi • Ka: talukuwa epi • Wp: kuya lá • Fr: bois baptiste, bois d'artre, bois de sang (*fide* Aublet 1775).

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000624924](#)], G-DC[[G00209682](#)]).

INVENTORY DATA (FG). — 21 trees in 14 plots; $F_{\max} = 3.8\%$; $dbh_{\text{inv}} = 33.7$ cm.

Family IXONANTHACEAE Planch. ex Miq.
Genus *Cyrillopsis* Kuhlm.

[556] *Cyrillopsis paraensis* Kuhlm.
(Fig. 25B)

Arch. Jard. Bot. Rio de Janeiro 4: 357 (Kuhlmann 1925).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.-F. Molino 3429*.

INVENTORY DATA (FG). — 53 trees in 20 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 63.4$ cm.

Family LACISTEMATACEAE Mart.
Genus *Lacistema* Sw.

[557] *Lacistema aggregatum* (P.J.Bergius) Rusby

Bull. New York Bot. Gard. 4 (14): 447 [7 Dec. 1907] (Rusby 1907). — *Piper aggregatum* P.J.Bergius, *Acta Helv. Phys.-Math.* 7: 131 (Bergius 1772). — *Lacistema myricoides* Sw., *Prodr. [Swartz]* 12 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. superfl. (based on *Piper aggregatum*).

Nematospermum laevigatum Rich., *Actes Soc. Hist. Nat. Paris* 1: 105 [Oct. 1792] (Richard 1792).

Synzyganthera purpurea Ruiz & Pav., *Fl. Peruv. Prodr.*: 137 [early Oct. 1794] (Ruiz & Pavón 1794). — *Didymandra purpurea* (Ruiz & Pav.) Willd., *Sp. Pl.*, ed. 4 4 (2): 971 [Apr. 1806] (Willdenow 1806), nom. illeg. superfl. (genus name superfluous, based on the type of *Synzyganthera* Ruiz & Pav.). — *Lacistema purpureum* (Ruiz & Pav.) A.DC., *Prodr. [A. P. de Candolle]* 16 (2.2): 593 [mid July 1868] (Candolle 1868).

Piper fasciculare Rudge, *Pl. Guian. [Rudge]* 1 (1): t. 4 [Apr. 1805] (Rudge 1805).

Lacistema floribundum Miq., *Linnaea* 18: 24 [24-26 Oct. 1844] (Miquel 1844), "floribunda".

Lacistema elongatum Schnizl., *Fl. Bras. [Martius]* 4 (1): 282 [28 Feb. 1857] (Schnizlein 1857).

Lacistema angustum Schnizl., *Fl. Bras. [Martius]* 4 (1): 283 [28 Feb. 1857] (Schnizlein 1857).

Lacistema recurvum Schnizl., *Fl. Bras. [Martius]* 4 (1): 283 [28 Feb. 1857] (Schnizlein 1857).

Lacistema myricoides var. *stipitatum* A.DC., *Prodr. [A. P. de Candolle]* 16 (2.2): 592 [mid July 1868] (Candolle 1868).

Lacistema coriaceum A.DC., *Prodr. [A. P. de Candolle]* 16 (2.2): 593 [mid July 1868] (Candolle 1868).

Lacistema poeppigii A.DC., *Prodr. [A. P. de Candolle]* 16 (2.2): 593 [mid July 1868] (Candolle 1868).

Lacistema boliviianum Gand., *Bull. Soc. Bot. France* 66 (7): 288 ["1919" publ. 1920] (Gandoger 1920).

Lacistema guyanense Gand., *Bull. Soc. Bot. France* 66 (7): 288 ["1919" publ. 1920] (Gandoger 1920).

Lacistema rosidiscum J.F.Macbr., *Candollea* 5: 392 (Macbride 1934).

Lacistema curtum J.F.Macbr., *Candollea* 5: 393 (Macbride 1934).

Lacistema orinocense Baehni, *Candollea* 8: 43 (Baehni 1940).

Lacistema weberbaueri Baehni, *Candollea* 8: 46 (Baehni 1940).

Lacistema aggregatum var. *elongatum* Maguire, *Bull. Torrey Bot. Club* 75 (3): 293 [May-June 1948] (Maguire 1948).

Lacistema occidentale Cuatrec., *Fieldiana, Bot.* 27 (1): 93 (Cuatrecasas 1950).

Lacistema pacificum Cuatrec., *Fieldiana, Bot.* 27 (1): 93 (Cuatrecasas 1950).

NOTE. — The type of *Didymandra* Willd. is that of *Synzyganthera* Ruiz & Pav., thus both *Didymandra* and *D. purpurea* (Ruiz & Pav.) Willd. are superfluous.

VERNACULAR NAMES. — Pa: agagut, ararut • Wp: ināmu sī, ināmusī wila, tatumila sili, yape'a pitā sili.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2419*.

INVENTORY DATA (FG). — 41 trees in 18 plots; $F_{\max} = 1\%$; dbh_{inv} = 24.6 cm.

[558] *Lacistema grandifolium* Schnizl.

Fl. Bras. [Martius] 4 (1): 284 [28 Feb. 1857] (Schnizlein 1857).

VERNACULAR NAMES. — Pa: tahuma • Wp: mulei sili.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (original material G[G00165292, G00202465]).

INVENTORY DATA (FG). — 41 trees in 28 plots; $F_{\max} < 1\%$; dbh_{inv} = 18.5 cm.

[559] *Lacistema polystachyum* Schnizl.
(Fig. 25C)

Fl. Bras. [Martius] 4 (1): 284 [28 Feb. 1857] (Schnizlein 1857).

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2367*.

INVENTORY DATA (FG). — 1 tree, dbh = 11.6 cm.

[560] *Lacistema pubescens* Mart.

Nova genera et species plantarum [Martius] 1 (4): 155 ["1824" publ. Jan.-Mar. 1826] (Martius 1826).

Lacistema ellipticum Schnizl., *Fl. Bras. [Martius]* 4 (1): 285 [28 Feb. 1857] (Schnizlein 1857).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4891*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 15.8 cm.

Family LAMIACEAE Martinov
Genus *Aegiphila* Jacq.

[561] *Aegiphila integrifolia* (Jacq.) B.D.Jacks.

Index Kew. 1 (1): 46 [6 Sep. 1893] (Jackson 1893). — *Callicarpa integrifolia* Jacq., *Enum. Syst. Pl.* 12 [Aug.-Sep. 1760] (Jacquin 1760).

Manabea arborescens Aubl., *Hist. Pl. Guiane* 1: 64 [Jun.-Dec. 1775] (Aublet 1775). — *Aegiphila arborescens* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 259 [late Sep.-Nov. 1791] (Gmelin 1791).

Callicarpa globiflora Ruiz & Pav., *Flora Peruviana* 1: 49 (Ruiz & Pavón 1798).

Callicarpa discolor Willd. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 257 (Steudel 1840), *nom. nud. pro syn.*

Aegiphila arborescens var. *longiflora* Schauer, *Prodr. [A. P. de Candolle]* 11: 649 [25 Nov. 1847] (Schauer 1847).

Aegiphila arborescens var. *breviflora* Schauer, *Prodr. [A. P. de Candolle]* 11: 649 [25 Nov. 1847] (Schauer 1847).

Aegiphila guianensis Moldenke, *Repert. Spec. Nov. Regni Veg.* 33: 125 (Moldenke 1933).

Aegiphila parviflora Moldenke, *Repert. Spec. Nov. Regni Veg.* 33: 135 (Moldenke 1933).

Aegiphila arborea Spruce ex Moldenke, *Phytologia* 1 (5): 206 [1 Mar. 1937] (Moldenke 1937), *nom. nud. pro syn.*

Aegiphila amazonica Moldenke, *Phytologia* 1 (11): 378 [6 Sept 1940] (Moldenke 1940).

Aegiphila cowanii Moldenke, *Phytologia* 27 (1): 63 [Oct. 1973] (Moldenke 1973).

Aegiphila integrifolia var. *lopez-palaci* Moldenke, *Phytologia* 36: 53 [June 1977] (Moldenke 1977).

VERNACULAR NAMES. — Pa: aigvan-kamwi • Cr: bwa-tabak, féy-tabak.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lectotype of *Manabea arborescens* P-JJR[P00777561] designated by Lanjouw & Uittien [1940: 152]).

SIZE. — Venezuela. *L. Marcano-Berti 244-981* (MO), 12 m.

[562] *Aegiphila villosa* (Aubl.) J.F.Gmel.

Syst. Nat., ed. 13[bis], 2 (1): 259 [late Sep.-Nov. 1791] (Gmelin 1791). — *Manabea villosa* Aubl., *Hist. Pl. Guiane* 1: 61 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Pa: aigvan, audika-retni-seine • Wp: a'i makule • Cr: bwa-tabak, féy-tabak • Br: camaá, cambara.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000598566]); *R.A.A. Oldeman B-2398*, 11 m × 25 cm

Genus *Vitex* L.

[563] *Vitex compressa* Turcz.

Bull. Soc. Imp. Naturalistes Moscou 36 (3): 224 (Turczaninow 1863).

Vitex brittoniana Moldenke, *Torreya* 33: 67 (Moldenke 1933).

Vitex compressa f. *angustifolia* Moldenke, *Phytologia* 47 (1): 17 [13 Nov. 1980] (Moldenke 1980).

VERNACULAR NAMES. — Ka: kwa'yan, kwailan • Nt: pakila udu.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M. Fleury 567*.

SIZE. — Up to 30 m tall (Jansen-Jacobs 1988).



FIG. 25. — Hypericaceae: **A**, *Vismia latifolia* (Aubl.) Choisy. Ixonanthaceae: **B**, *Cyrillopsis paraensis* Kuhl. (J.-F. Molino 3429). Lacistemataceae: **C**, *Lacistema polystachyum* Schnizl. (J.-F. Molino & D. Sabatier 2367). Lamiaceae: **D**, *Vitex guianensis* Moldenke (D. Sabatier 2332). A, D, © D. Sabatier/IRD; B, C, © J.-F. Molino/IRD.

[564] *Vitex guianensis* Moldenke
(Fig. 25D)

Phytologia 1 (15): 486 [Jan. 1941] (Moldenke 1941).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: [D. Sabatier 1220](#).

INVENTORY DATA (FG). — 19 trees in 13 plots; $F_{\max} < 1\%$; dbh_{inv} = 86.6 cm.

[565] *Vitex orinocensis* Kunth var. *multiflora* (Miq.) Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (1): 217 (Huber 1908). — *Vitex multiflora* Miq., *Linnaea* 18: 739 ["1844" publ. Aug.-Oct. 1845] (Miquel 1845).

HERBARIUM DATA (FG). — A single collection, [M.-F. Prévost 1460](#).

SIZE. — Up to 20 m tall (Jansen-Jacobs 1988).

[566] *Vitex stabelii* Moldenke

Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2, 30: 310 (Moldenke 1940).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: tayi lá.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: [F. Billet & B. Jadin 1431](#).

SIZE. — Up to 40 m tall (Jansen-Jacobs 1988).

[567] *Vitex triflora* Vahl

Eclog. Amer. 2: 49 (Vahl 1798).

Pyrostoma ternatum G.Mey., *Prim. Fl. Esseq.* 220 [Nov. 1818] (Meyer 1818), "ternata".

Casarettoa diversifolia Walp., *Repert. Bot. Syst. [Walpers]* 4 (1): 92 [25-28 June 1845] (Walpers 1847), nom. illeg. superfl. (based on *Vitex triflora*).

Macrostegia ruiziana Nees, *Prodri. [A. P. de Candolle]* 11: 218 [25 Nov. 1847] (Nees 1847).

Ruellia macrocalyx Ruiz ex Nees, *Prodri. [A. P. de Candolle]* 11: 218 [25 Nov. 1847] (Nees 1847), nom. nud. pro syn.

Vitex triflora var. *angustiloba* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (1): 215 (Huber 1908).

Vitex triflora var. *coriacea* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (1): 215 (Huber 1908).

Vitex triflora var. *floribunda* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (1): 215 (Huber 1908).

Vitex triflora var. *kraatzii* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (1): 216 (Huber 1908).

Vitex triflora var. *tenuifolia* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (1): 214 (Huber 1908).

Vitex triflora var. *quinquefoliolata* Moldenke, *Phytologia* 1 (2): 104 [4 Aug. 1934] (Moldenke 1934). — *Vitex triflora* f. *quinquefoliolata* (Moldenke) Moldenke, *Phytologia* 44 (6): 384 [11 Dec. 1979] (Moldenke 1979).

Vitex triflora var. *hirsuta* Moldenke, *Phytologia* 23 (3): 315 [31 May 1972] (Moldenke 1972).

VERNACULAR NAMES. — Pa: kwairu-kamwi • Wp: tayi lá • Wn: wilimé • Nt: madyo uman wiwii, madyoma uwii • Br: pau-d'arco-rana, tarumá.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (original material B, B-W 11701 -02 0).

INVENTORY DATA (FG). — 30 trees in 28 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 17\text{ cm}$.

Family LAURACEAE Juss.
Genus *Aioea* Aubl.

[568] *Aioea guianensis* Aubl.
(Fig. 26A)

Hist. Pl. Guiane 1: 311 [Jun.-Dec. 1775] (Aublet 1775). — *Laurus hexandra* Willd., *Sp. Pl., ed. 4* 2 (1): 482 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Aioea guianensis*). — *Douglasia laurina* Sm., *Cycl. [Rees]* 12: (Douglasia no. 1) [14 Feb. 1809] (Smith 1809), nom. illeg. superfl. (based on *Aioea guianensis*).

Laurus difformis Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792).

Aioea tenella Nees, *Linnaea* 21: 512 (Nees 1848).

Aioea schomburgkii Meisn., *Prodri. [A. P. de Candolle]* 15 (1): 82 [May 1864] (Meisner 1864).

Aioea rubra A.C.Sm., *Phytologia* 1 (3): 115 [21 Jan. 1935] (Smith 1935).

Aioea demerarensis Kosterm., *Recueil Trav. Bot. Néerl.* 35: 77 (Kostermans 1938).

NOTES. — *L.C. Richards* s.n. (B-W, B-W 07804 -00 0; LE[LE00012694; P[P02004886]) was used by Willdenow (1799: 482) to illustrate his *Laurus hexandra*. It has a label with exactly the same description as in the protologue (“*Laurus hexandra foliis lanceolatis, racemis axillaribus, floribus hexandris. Sp. pl. 2 p. 482*”). Nonetheless, this is a specimen of *Aniba guianensis* Aubl., not *Aioea guianensis* Aubl. (see notes under *Aniba guianensis*).

VERNACULAR NAMES. — Pa: panauna-puvemna.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material P[P00128374, P00662841]).

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 46.6\text{ cm}$.

[569] *Aioea impressa* (Meisn.) Kosterm.

Recueil Trav. Bot. Néerl. 35: 75 (Kostermans 1938). — *Phoebe impressa* Meisn., *Prodri. [A. P. de Candolle]* 15 (1): 33 [May 1864] (Meisner 1864). — *Endlicheria impressa* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 132 (Mez 1889). — *Cinnamomum impressum* (Meisn.) Kosterm., *Reinwardtia* 10 (5): 444 (Kostermans 1988).

Aioea minutiflora Coe-Tex., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 34: 1 (Coe-Teixeira 1970).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* “*Cayenne 31*” (original material G[G00368709], L[L0035523]).

SIZE. — Up to 20 m tall (Renner 1982).

[570] *Aioea longipetiolata* van der Werff

Novon 4 (1): 61 [spring 1994] (van der Werff 1994).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: panauna-kamwi • Wp: yakami ka'a, yakami'i.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *S.A. Mori et al. 20758* (holo-, MO[MO-247078]; iso-, CAY[CAY027837], NY[00022599], P[P00745701], U[U0002590], US[00754392]).

INVENTORY DATA (FG). — 26 trees in 16 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 55.4\text{ cm}$.

[571] *Aioea montana* (Sw.) R.Rohde

Taxon 66 (5): 1102 [epubl. 24 Oct. 2017] (Rohde 2017). — *Laurus montana* Sw., *Prodri. [Swartz]* 65 [20 Jun.-29 July 1788] (Swartz 1788). — *Cinnamomum montanum* (Sw.) J.Presl, *Přir. Rostlin* 2: 36 (Presl 1825). — *Persea montana* (Sw.) Spreng., *Syst. Veg. [Sprengel]* 2: 268 [Jan.-May 1825] (Sprengel 1825). — *Phoebe montana* (Sw.) Griseb., *Pl. Wright. [Grisebach]* 1: 187 [Dec. 1860] (Grisebach 1860).

Laurus triplinervis Ruiz & Pav., *Flora Peruviana* 4: t. 363 [1804-1830] (Ruiz & Pavón 1804). — *Phoebe peruviana* Meisn., *Prodri. [A. P. de Candolle]* 15 (1): 32 [May 1864] (Meisner 1864), nom. illeg. superfl. (based on *Laurus triplinervis*). — *Phoebe triplinervis* (Ruiz & Pav.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 211 (Mez 1889), nom. illeg. hom., non Griseb. (Grisebach 1860) [synonym of *Cinnamomum grisebachii* Lorea-Hern.]. — *Cinnamomum trip-*

linerve (Ruiz & Pav.) Kosterm., *Reinwardtia* 6 (1): 24 (Kostermans 1961), “*triplinervis*”.

Persea cinnamomifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 160 [8 Dec. 1817] (Kunth 1818). — *Phoebe cinnamomifolia* (Kunth) Nees, *Linnaea* 21: 488 (Nees 1848). — *Phoebe granatensis* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 32 [May 1864] (Meisner 1864), *nom. illeg. superfl.* (based on *Persea cinnamomifolia*). — *Cinnamomum cinnamomifolium* (Kunth) Kosterm., *Reinwardtia* 6 (1): 20 (Kostermans 1961).

Phoebe elongata Nees, *Syst. Laur.*: 116 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Laurus elongata* Vahl ex Nees, *Syst. Laur.*: 116 [30 Oct.-5 Nov. 1836] (Nees 1836), *nom. nud. pro syn.* — *Cinnamomum elongatum* (Nees) Kosterm., *Reinwardtia* 6 (1): 21 (Kostermans 1961), *nom. illeg. hom., non* Saporta (1889) *nec* Usnade (1949).

Phoebe maynensis Nees, *Syst. Laur.*: 118 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Phoebe peruviana* var. *glabriflora* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 32 [May 1864] (Meisner 1864), *nom. illeg. superfl.* (based on *Phoebe maynensis*). — *Cinnamomum maynense* (Nees) Kosterm., *Reinwardtia* 6 (1): 22 (Kostermans 1961).

Phoebe cubensis Nees, *Syst. Laur.*: 120 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Phoebe antillana* Meisn. var. *cubensis* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 31 [May 1864] (Meisner 1864). — *Phoebe triplinervis* var. *cubensis* (Nees) C.Wright, *Anales Acad. Ci. Med. Habana* 7: 347 [15 Nov. 1870] (Wright 1870). — *Cinnamomum cubense* (Nees) Kosterm., *Reinwardtia* 6 (1): 21 (Kostermans 1961), “*cubensis*”.

Phoebe valenzuelana A.Rich., *Hist. Fís. Cuba, Bot.* 11: 185 (Richard 1850). — *Persea triplinervis* (Griseb.) M.Gómez var. *valenzuelana* (A.Rich.) M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.*: 25 (Gómez 1889).

Oreodaphne alba A.Rich., *Hist. Fís. Cuba, Bot.* 11: 189 (Richard 1850).

Phoebe mexicana Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 31 [May 1864] (Meisner 1864). — *Persea mexicana* (Meisn.) Hemsl., *Biol. Cent.-Amer. Bot.* 3 (14): 72 [Oct. 1882] (Hemsley 1882). — *Cinnamomum mexicanum* (Meisn.) Kosterm., *Reinwardtia* 6 (1): 22 (Kostermans 1961).

Phoebe poeppigii Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 33 [May 1864] (Meisner 1864).

Phoebe brasiliensis Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 198 (Mez 1889).

Phoebe mexicana var. *bourgeauana* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 214 (Mez 1889). — *Cinnamomum brasiliense* (Mez) Kosterm., *Reinwardtia* 6 (1): 20 (Kostermans 1961), “*brasiliensis*”.

Phoebe heterotepala Mez, *Repert. Spec. Nov. Regni Veg.* 3: 67 (Mez 1906). — *Cinnamomum heterotepalum* (Mez) Kosterm., *Reinwardtia* 6 (1): 21 (Kostermans 1961).

Phoebe paraguariensis Hassl., *Annuaire Conserv. Jard. Bot. Genève* 21: 79 (Hassler 1919). — *Cinnamomum paraguariense* (Hassl.) Kosterm., *Reinwardtia* 6 (1): 22 (Kostermans 1961), “*paraguariensis*”.

Ocotea flavescens Rusby, *Descr. S. Amer. Pl.*: 20 [20 Dec. 1920] (Rusby 1920).

Phoebe pichisensis A.C.Sm., *Bull. Torrey Bot. Club* 58 (2): 103 [Feb. 1931] (Smith 1931). — *Cinnamomum pichisense* (A.C.Sm.) Kosterm., *Reinwardtia* 6 (1): 23 (Kostermans 1961), “*pichisensis*”.

Phoebe johnstonii C.K.Allen, *J. Arnold Arbor.* 26 (4): 433 [15 Oct. 1945] (Allen 1945). — *Cinnamomum johnstonii* (C.K.Allen) Kosterm., *Reinwardtia* 6 (1): 21 (Kostermans 1961).

Cinnamomum chana Vattimo-Gil, *Arch. Jard. Bot. Rio de Janeiro* 17: 223 (Vattimo-Gil 1962).

Cinnamomum australe Vattimo-Gil, *Arch. Jard. Bot. Rio de Janeiro* 17: 224 (Vattimo-Gil 1962).

Cinnamomum xinguense Vattimo-Gil, *Arch. Jard. Bot. Rio de Janeiro* 17: 224 (Vattimo-Gil 1962).

Phoebe filamentosa C.K.Allen, *Mem. New York Bot. Gard.* 15: 69 (Allen 1966). — *Cinnamomum filamentosum* (C.K.Allen) Kosterm., *Reinwardtia* 7 (5): 461 (Kostermans 1969).

Cinnamomum portosecurianum Vattimo-Gil, *Anais 15 Congr. Soc. Bot. Brasil*: 170 (Vattimo-Gil 1967).

Phoebe pickellii Coe-Teix., *Hoehnea* 1: 187 (Coe-Teixeira 1971), “*pickelli*”. — *Cinnamomum pickellii* (Coe-Teix.) Kosterm., *Reinwardtia* 10 (5): 448 (Kostermans 1988).

Phoebe fruticosa Lundell, *Wrightia* 5 (9): 342 (Lundell 1977). — *Cinnamomum fruticosum* (Lundell) Kosterm., *Reinwardtia* 10 (5): 443 (Kostermans 1988).

Cinnamomum peruvianum Kosterm., *Reinwardtia* 10 (5): 448 (Kostermans 1988).

Cinnamomum phoebe Doweld, *Phytotaxa* 243 (2): 191 [epubl. 12 Jan. 2016] (Doweld 2016).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *S.A. Mori et al. 15024*.

INVENTORY DATA (FG). — 1 tree, dbh = 15.4 cm.

[572] *Aiofea opaca* van der Werff

Brittonia 47 (4): 372 (van der Werff 1995).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori et al. 20928* (holo-, MO[MO-247076]; iso-, CAY[CAY027842], NY[00214355], P[P00745702], U[U0002592], US[00478967]).

INVENTORY DATA (FG). — 16 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 37.7$ cm.

Genus *Aniba* Aubl.

[573] *Aniba affinis* (Meisn.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 63 (Mez 1889). — *Aydendron affine* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 92 [May 1864] (Meisner 1864).

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville et al. 10796*.

SIZE. — Up to 15 m tall (Kubitzki 1982).




FIG. 26. — Lauraceae: **A**, *Aioea guianensis* Aubl. (J.-F. Molino et al. 3371); **B**, *Aniba megaphylla* Mez (J.-F. Molino & D. Sabatier 2834); **C**, *Aniba williamsii* O.C.Schmidt (D. Sabatier & E. Fonty 5662); **D-F**, *Nectandra matogrossensis* Coe-Teix. (J.-F. Molino et al. 3406). A, B, D-F, © J.-F. Molino/IRD; C, © D. Sabatier/IRD.

[574] *Aniba citrifolia* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 74 (Mez 1889). — *Aydendron citrifolium* Nees, *Syst. Laur.* 257 [30 Oct.-5 Nov. 1836] (Nees 1836).

Aydendron trinitatis Meisn., *Prod. [A. P. de Candolle]* 15 (1): 91 [May 1864] (Meisner 1864). — *Aniba trinitatis* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 68 (Mez 1889).

Aniba muelleriana Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 72 (Mez 1889), “Mülleriana”.

Cryptocarya pachycarpa Gleason, *Bull. Torrey Bot. Club* 54 (8): 607 [Nov. 1927] (Gleason 1927).

Aniba opaca A.C.Sm., *Bull. Torrey Bot. Club* 61 (4): 191 [Apr. 1934] (Smith 1934).

VERNACULAR NAMES. — Pa: kudagman, kudagman-purubumna • Te: uwahé pitág • Wp: áyú'i tawa, iwa'e • Nt: bamba apisi • Cr: sèd • Fr: cèdre.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: F.M.R. Leprieur 224 (original material of *Aniba muelleriana*: G[G00368670], P[P00745591, P00745592]).

INVENTORY DATA (FG). — 33 trees in 27 plots; $F_{\max} < 1\%$; dbh_{inv} = 33.4 cm.

[575] *Aniba guianensis* Aubl.

Hist. Pl. Guiane 1: 327 [Jun.-Dec. 1775] (Aublet 1775). — *Cedrota guianensis* (Aubl.) Raeusch., *Nomenc. Bot. [Raeusch.] ed. 3:* 111 (Raeuschel 1797). — *Cedrota longifolia* Willd., *Sp. Pl., ed. 4* 2 (1): 338 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Aniba guianensis*).

Aniba salicifolia Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 71 (Mez 1889).

Aniba tessmannii O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 226 [30 Mar. 1928] (Schmidt 1928).

NOTES. — Excluded synonyms: *Aydendron salicifolium* (Sw.) Nees [syn. nov. of *Licaria parvifolia* (Lam.) Kosterm.] and its basionym *Laurus salicifolia* Sw. Kubitzki (1982: 45) placed “*Aydendron salicifolium* Nees” and “*Aniba salicifolia* (Nees) Mez” in synonymy under *Aniba guianensis* Aubl. Actually, although Nees misinterpreted *L. salicifolia* (synonym of *Licaria parvifolia*, a Caribbean taxon), and based his description of *Aydendron salicifolium* on a French Guianan collection of *Aniba guianensis* (L.C. Richard s.n. [B-W, B-W07804-000; LE[LE00012694; P[P02004886], see also notes under *Aioea guianensis*]), he explicitly referred to *L. salicifolia* and to its type. Therefore, he effectively transferred *L. salicifolia* into *Aydendron*. Later, when transferring *Aydendron salicifolium* into

Aniba, Mez (1889: 71) excluded *L. salicifolia* from synonymy. He thereby created a new name for the Guianan species.

VERNACULAR NAMES. — Wp: áyū'i sili, áyū'i tawa • Cr: sèd-kannel.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000617483](#)]).

INVENTORY DATA (FG). — 39 trees in 30 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.8$ cm.

[576] *Aniba hostmanniana* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 67 (Mez 1889). — *Aydendron hostmannianum* Nees, *Linnaea* 21: 499 (Nees 1848).

Aniba gigantifolia O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 225 [30 Mar. 1928] (Schmidt 1928).

Aniba pittieri O.C.Schmidt, *Repert. Spec. Nov. Regni Veg.* 31: 168 (Schmidt 1933).

VERNACULAR NAMES. — Ka: apotono ali, siduwapali, sipilupipyo, waikyala • Wp: kule'i sī, pilapuku'i • Wn: kaikui amoman • Nt: bamba apisi • Cr: sèd-marikaj • Fr: cèdre marécage • Br: louro-capitiu.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *Service Forestier 7816*.

SIZE. — Up to 20 m tall (Kubitzki 1982).

[577] *Aniba jenmanii* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 70 (Mez 1889), “Jenmani”.

Aniba castanea C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 45 (Allen 1964).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier 4370*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.9$ cm.

[578] *Aniba kappleri* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 52 (Mez 1889).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: siduwapali.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3333*.

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 19.4$ cm.

[579] *Aniba megaphylla* Mez
(Fig. 26B)

Jahrb. Königl. Bot. Gart. Berlin 5: 67 (Mez 1889).

Aniba anisosepala Sandwith, *Bull. Misc. Inform. Kew* 1933 (7): 336 [25 Sep. 1933] (Sandwith 1933).

Aniba koumaroucapa Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 24 (Kostermans 1936). — *Laurus koumaroucapa* Rich. ex Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 24 (Kostermans 1936), *nom. nud. pro syn.*

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *F.M.R. Leprieur 225* (holo-, P[[P00745605](#)]; iso-, F[[V0061345F](#)], G[[G00368662](#)], P[[P00745604](#)]).

INVENTORY DATA (FG). — 1 tree, $dbh = 10.5$ cm.

[580] *Aniba panurensis* (Meisn.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 58 (Mez 1889). — *Aydendron panurense* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 89 [May 1864] (Meisner 1864).

Aniba gonggrijpii Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 31 (Kostermans 1936).

Aniba mas Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 31 (Kostermans 1936).

VERNACULAR NAMES. — Ka: sipilulan.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier et al. 4604*.

INVENTORY DATA (FG). — 19 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43$ cm.

[581] *Aniba parviflora* (Meisn.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 56 (Mez 1889). — *Aydendron parviflorum* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 90 [May 1864] (Meisner 1864).

VERNACULAR NAMES. — Pa: sedri-kamwi-seine • Nt: bwadiwosu • Br: louro-puxuri.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3352*.

INVENTORY DATA (FG). — 33 trees in 21 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 50.3$ cm.

[582] *Aniba riparia* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 62 (Mez 1889). — *Aydendron riparium* Nees, *Linnaea* 21: 497 (Nees 1848).

VERNACULAR NAMES. — Ka: siduwapali.

HERBARIUM DATA (FG). — A single collection, *C. Feuillet 10338* (US).

SIZE. — Up to 16 m tall (Kubitzki 1982).

[583] *Aniba rosodora* Ducke

Rev. Bot. Appl. Agric. Colon. 88: 845 [Dec. 1928] (Ducke 1928), “rosaeodora”.

Aniba rosodora var. *amazonica* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 83 (Ducke 1930), “rosaeodora”.

Aniba duckei Kosterm., *Recueil Trav. Bot. Néerl.* 35: 924 (Kostermans 1938).

NOTE. — Epithet corrected according to Art. 60.10 and Rec. 60G.1 (a), of the Shenzhen Code (Turland et al. 2018).

VERNACULAR NAMES. — Pa: sedri-eminyo • Wp: áyū'i tawa, pauteleso • Wn: matolosi, mijetaimē • Nt: bwadiwosu • Cr: bwa-dé-ròz • Fr: bois de rose • Br: pau-rosa.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.F. Prévost 3301*.

INVENTORY DATA (FG). — 15 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 36.8$ cm.

[584] *Aniba taubertiana* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 65 (Mez 1889).

Aniba simulans C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 51 (Allen 1964).

VERNACULAR NAMES. — Ka: tamunen wonu • Wp: áyū'i piye.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *E.M. Mélénan s.n.* (holo-, B[not seen, photo F neg. № 3809]; iso-, BM[BM000617490], CAS[CAS0026940], F[FV0061352F], G[G00368692], GH[GH00056912], K[K000601899, K000601900], NY[00354852, 00354853], P[P00745619, P00745620, P00745621], US[0009455]).

INVENTORY DATA (FG). — 67 trees in 54 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 32.9$ cm.

[585] *Aniba terminalis* Ducke

Rev. Bot. Appl. Agric. Colon. 88: 846 [Dec. 1928] (Ducke 1928).

VERNACULAR NAMES. — Ka: siduwapali.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5099*.

INVENTORY DATA (FG). — 1 tree, $dbh = 12.3$ cm.

[586] *Aniba williamsii* O.C.Schmidt
(Fig. 26C)

Repert. Spec. Nov. Regni Veg. 31: 169 (Schmidt 1933).

Aniba muricana C.K.Allen, *Mem. New York Bot. Gard.* 15: 57 (Allen 1966).

VERNACULAR NAMES. — Wp: kule'i • Br: louro-abacate.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & E. Fonty 5662*.

INVENTORY DATA (FG). — 20 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.1$ cm.

[587] *Aniba* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & E. Fonty 5591*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.3$ cm.

[588] *Aniba* sp. B

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & M.-F. Prévost 1917*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.5$ cm.

[589] *Aniba* sp. C

HERBARIUM DATA (FG). — A single collection, *S.A. Mori & T.D. Pennington 18052*.

SIZE. — Up to 18 m tall (Werff 2002, as *Rhodostemonodaphne* sp. 1).

Genus *Beilschmiedia* Nees

[590] *Beilschmiedia hexanthera* van der Werff

Brittonia 47 (4): 374 (van der Werff 1995).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *H. van der Werff et al. 12951* (holo-, MO[MO-276584]; iso-, B[B100247471], BM[BM000588156, BM000588157], CAY[CAY184925], MO[MO-276583], NY[00214380] , U[U0008155], XAL[XAL0091566]).

INVENTORY DATA (FG). — 1 tree, $dbh = 37.6$ cm.

Genus *Cryptocarya* R.Br.

[591] *Cryptocarya guianensis* Meisn.

Prodr. [A. P. de Candolle] 15 (1): 75 [May 1864] (Meisner 1864).

Cryptocarya maroniensis Benoist, *Bull. Mus. Natl. Hist. Nat.* 30: 510 (Benoist 1924).

Cryptocarya nigropunctata Vattimo-Gil, *Rodriguésia* 25: 222 (Vattimo-Gil 1966).

VERNACULAR NAMES. — Pa: wen-kamwi • Wp: iwa pane • Cr: sèdmarikaj • Br: caa-xiò, caiaxiò.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *M.E. Moricand 113* (lecto-, G-DC[not seen], designated by Moraes [2005: 791]; isolecto-, G[G00418990, G00418997]).

INVENTORY DATA (FG). — 1 tree, $dbh = 22.4$ cm.

Genus *Damburneya* Raf.

[592] *Damburneya purpurea* (Ruiz & Pav.) Trofimov

Taxon 65 (5): 991 [27 Oct. 2016] (Trofimov 2016). — *Laurus purpurea* Ruiz & Pav., *Flora Peruviana* 4: t. 351 [1804-1830] (Ruiz & Pavón 1804). — *Nectandra purpurea* (Ruiz & Pav.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 443 (Mez 1889).

Ocotea latifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 165 [8 Dec. 1817] (Kunth 1818). — *Persea latifolia* (Kunth) Spreng., *Syst. Veg. [Sprengel]* 2: 270 [Jan.-May 1825] (Sprengel 1825). — *Nectandra latifolia* (Kunth) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 454 (Mez 1889).

Nectandra polita Nees & Mart., *Syst. Laur.* 325 [30 Oct.-5 Nov. 1836] (Nees & Martius 1836).

Nectandra polita var. *oerstedii* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 164 [May 1864] (Meisner 1864).

Ocotea flexuosa Rusby, *Descr. S. Amer. Pl.* 21 [20 Dec. 1920] (Rusby 1920).

VERNACULAR NAMES. — Pa: akamna, akamnā, akapna.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 24258*.

SIZE. — Peru, Amazonas. *T.D. Pennington & A. Daza Yomona 17839* (MO), 10 m × 20 cm.

Genus *Endlicheria* Nees

[593] *Endlicheria bracteolata* (Meisn.) C.K.Allen

Mem. New York Bot. Gard. 10 (5): 64 (Allen 1964). — *Goeppertia sericea* (Nees) Nees var. *bracteolata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 174 [May 1864] (Meisner 1864).

VERNACULAR NAMES. — Wp: áyū'i alapitá milá • Br: louro-peludo.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3580*.

SIZE. — Up to 15 m tall (Chanderbali 2004).

[594] *Endlicheria canescens* Chanderb.

Novon 6 (4): 328 [winter 1996] (Chanderbali 1996).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2438*.

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 29.6$ cm.

[595] *Endlicheria chalisea* Chanderb.

Novon 6 (4): 329 [winter 1996] (Chanderbali 1996).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S. Barrier 4026*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 25.9$ cm.

[596] *Endlicheria melinonii* Benoist

Arch. Bot. 5 (Mém. 1): 63 [27 Mar. 1933] (Benoist 1933), “*Melinonii*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: migukat, mihiukat, mirukat.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *E.M. Mélénion s.n.* (holo-, P[P00077245]; iso-, P[P00745669]).

INVENTORY DATA (FG). — 51 trees in 43 plots; $F_{\max} < 1\%$; $dbh_{inv} = 67.8$ cm.

[597] *Endlicheria punctulata* (Mez) C.K.Allen

Mem. New York Bot. Gard. 15: 68 (Allen 1966). — *Ocotea punctulata* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 379 (Mez 1889).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: áyū'i alapitá milá.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *E.M. Mélénion 204*, 1842 (lecto-, P[P00745657], designated by Allen [1966: 68]; isolecto-, P[P00745658]).

INVENTORY DATA (FG). — 10 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 46.7$ cm.

[598] *Endlicheria pyriformis* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 116 (Mez 1889), “*piriformis*”. — *Cryptocarya pyriformis* Nees, *Syst. Laur.*: 220 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Mespilodaphne pyriformis* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 108 [May 1864] (Meisner 1864).

Endlicheria glaberrima Mez, *Bull. Herb. Boissier*, sér. 2, 5: 236 (Mez 1905).

Aniba flexuosa A.C.Sm., *Phytologia* 1 (3): 117 [21 Jan. 1935] (Smith 1935).

VERNACULAR NAMES. — Pa: kudagman • Ka: woko mapili • Wp: amákiya sili, áyū'i • Wn: alukamata • Cr: sèd • Br: louro.

HERBARIUM DATA (FG). — 69 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holo-, B[not seen, photo F neg. N° 3820]; iso-, BM[BM000993875], G-DC[G00201070], LE[LE00012615, LE00012616, LE00012728], NY[00099497], P[P00745654]).

SIZE. — Up to 10 m tall (Chanderbali 2004).

[599] *Endlicheria rubriflora* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 494 (Mez 1889).

Aniba reticulata A.C.Sm., *Bull. Torrey Bot. Club* 58 (2): 99 [Feb. 1931] (Smith 1931).

Endlicheria trianae O.C.Schmidt, *Repert. Spec. Nov. Regni Veg.* 31: 175 (Schmidt 1933).

Endlicheria wurdackiana C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 67 (Allen 1964).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 23320*.

SIZE. — Brazil, Amazonas. *G.T. Prance et al. 14498* (NY), 10 m × 15 cm.

[600] *Endlicheria szyszlowiczii* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 121 (Mez 1889).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman B-2312*, 12 m × 35 cm.

[601] *Endlicheria* sp. A

NOTE. — This is possibly *E. glomerata* Mez.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2803*.

INVENTORY DATA (FG). — 2 trees in 2 plots; dbh_{inv} = 19.5 cm.

Genus *Kubitzkia* van der Werff

[602] *Kubitzkia mezii* (Kosterm.) van der Werff

Taxon 35 (1): 165 [Feb. 1986] (van der Werff 1986). — *Systemonodaphne mezii* Kosterm., *Recueil Trav. Bot. Néerl.* 33: 756 (Kostermans 1936), in obs.

VERNACULAR NAMES. — Wp: kwata pili.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *J. Martin* 45 (holotype not traced; iso-, K[K000601987], P[P00745651, P00745652, P00745653], U[U0002743]).

INVENTORY DATA (FG). — 7 trees in 6 plots; F_{max} < 1 %; dbh_{inv} = 16.6 cm.

Genus *Licaria* Aubl.

[603] *Licaria cannella* (Meisn.) Kosterm.

Recueil Trav. Bot. Néerl. 34: 583 (Kostermans 1937), “*Canella*”. — *Ayndendron cannella* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 90 [May 1864] (Meisner 1864). — *Acrodielidium cannella* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 90 (Mez 1889), “*Canella*”. — *Misanteca cannella* (Meisn.) Lundell, *Wrightia* 4 (3): 99 (Lundell 1969), “*Canella*”.

Aniba megacarpa Hemsl., *Hooker's Icon. Pl.* 28 [ser. 4, 8]: tt. 2751, 2752 [Nov. 1903] (Hemsley 1903).

VERNACULAR NAMES. — Ka: ayiwi, ayu'i, ikanikanali, sitaipyo, wa'elan, wayaka • Wp: áyú'i tawa • Wn: mijetaimë, watau pokan • Nt: kaneli apisi • Cr: bwa-kannel, sèd-jonn, sèd-kannel • Fr: cèdre cannelle • Br: louro-chumbo, louro-pirarucu, louro-preto.

HERBARIUM DATA (FG). — 71 collections at CAY. Sel. exs.: *P.A. Sagot* 1190, Aug. 1858 (holo-, B[not seen, photo F neg. № 3837]; iso-, BM[BM000993917], K[K000602022], P[P00128461, P00128462], NY[00354870]).

INVENTORY DATA (FG). — 197 trees in 96 plots; F_{max} = 3.8 %; dbh_{inv} = 92 cm.

[604] *Licaria chrysophylla* (Meisn.) Kosterm.

Recueil Trav. Bot. Néerl. 34: 601 (Kostermans 1937). — *Acrodielidium chrysophyllum* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 87 [May 1864] (Meisner 1864). — *Misanteca chrysophylla* (Meisn.) Lundell, *Wrightia* 4 (3): 99 (Lundell 1969).

Acrodielidium guianense var. *oppositifolium* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 85 [May 1864] (Meisner 1864).

Acrodielidium rigidum Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 38 (Kostermans 1936). — *Licaria rigida* (Kosterm.) Kosterm., *Recueil Trav. Bot. Néerl.* 34: 584 (Kostermans 1937). —

Misanteca rigida (Kosterm.) Lundell, *Wrightia* 4 (3): 101 (Lundell 1969).

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *P.A. Sagot* 956, 1858 (holo-, B[not seen, photo F neg. № 3838]; iso-, MEL[MEL2386590]; possible iso-, P[P00128467, P00128468]).

INVENTORY DATA (FG). — 58 trees in 33 plots; F_{max} < 1 %; dbh_{inv} = 73 cm.

[605] *Licaria crassifolia* (Poir.) P.L.R.Moraes

Komarovia 6 (1): 55 (Moraes 2008). — *Laurus crassifolia* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 323 [24 Sep. 1813] (Poiret 1813).

Ayndendron cayennense Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 95 [May 1864] (Meisner 1864). — *Acrodielidium cayennense* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 91 (Mez 1889). — *Licaria cayennensis* (Meisn.) Kosterm., *Recueil Trav. Bot. Néerl.* 34: 583 (Kostermans 1937). — *Misanteca cayennensis* (Meisn.) Lundell, *Wrightia* 4 (3): 99 (Lundell 1969).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holo-, P[P00128465]; iso-, P[P00128463, P00128464, P00662838]).

SIZE. — > 10 cm dbh (Cardoso et al. 2017).

[606] *Licaria debilis* (Mez) Kosterm.

Recueil Trav. Bot. Néerl. 34: 596 (Kostermans 1937). — *Acrodielidium debile* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 99 (Mez 1889). — *Misanteca debilis* (Mez) Lundell, *Wrightia* 4 (3): 100 (Lundell 1969).

VERNACULAR NAMES. — Wp: ka'i piye.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-F. Molino* 907.

INVENTORY DATA (FG). — 3 trees in 3 plots; F_{max} < 1 %; dbh_{inv} = 14.3 cm.

[607] *Licaria guianensis* Aubl.

Hist. Pl. Guiane 1: 313 [Jun.-Dec. 1775] (Aublet 1775), “*Guyanensis*” on plate. — *Acrodielidium aubletii* Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 34 (Kostermans 1936).

Acrodielidium appeltii Mez, *Bot. Jahrb. Syst.* 17 (5): 519 [7 Nov. 1893] (Mez 1893). — *Licaria appeltii* (Mez) Kosterm., *Recueil Trav. Bot. Néerl.* 34: 601 (Kostermans 1937). — *Misanteca appeltii* (Mez) Lundell, *Wrightia* 4 (3): 99 (Lundell 1969).

VERNACULAR NAMES. — Br: louro-aritu, louro-cravo.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material P[P00128472, P00128471]).

INVENTORY DATA (FG). — 25 trees in 17 plots; F_{max} < 1 %; dbh_{inv} = 19.4 cm.

[608] *Licaria martiniana* (Mez) Kosterm.

Recueil Trav. Bot. Néerl. 34: 601 (Kostermans 1937). — *Acroclidium martinianum* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 93 (Mez 1889). — *Misanteca martiniana* (Mez) Lundell, *Wrightia* 4 (3): 100 (Lundell 1969).

Licaria foveolata Lemée, *Fl. Guyane Franç.* 1: 648 (Lemée 1955).

VERNACULAR NAMES. — Pa: migukat, miukat, mirukat, sedrikamwi, sedri-priyo • Ka: yokolan pomiidy • Te: uwahe pitág • Nt: geli apisi, mal bwadiwosu.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D. Sabatier 2306*.

INVENTORY DATA (FG). — 100 trees in 53 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 69.8$ cm.

[609] *Licaria pachycarpa* (Meisn.) Kosterm.

Recueil Trav. Bot. Néerl. 34: 604 (Kostermans 1937). — *Ayndendron pachycarpum* Meisn., *Prod. [A. P. de Candolle]* 15 (1): 95 [May 1864] (Meisner 1864). — *Acroclidium pachycarpum* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 91 (Mez 1889). — *Misanteca pachycarpa* (Meisn.) Lundell, *Wrightia* 4 (3): 101 (Lundell 1969).

Acroclidium amarum Mez, *Repert. Spec. Nov. Regni Veg.* 16: 305 [31 Apr. 1920] (Mez 1920). — *Licaria amara* (Mez) Kosterm., *Recueil Trav. Bot. Néerl.* 34: 583 (Kostermans 1937). — *Misanteca amara* (Mez) Lundell, *Wrightia* 4 (3): 99 (Lundell 1969).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3780*.

INVENTORY DATA (FG). — 11 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.6$ cm.

[610] *Licaria polyphylla* (Nees) Kosterm.

Recueil Trav. Bot. Néerl. 34: 584 (Kostermans 1937). — *Nectandra polyphylla* Nees, *Syst. Laur.*: 332 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Misanteca polyphylla* (Nees) Lundell, *Wrightia* 4 (3): 101 (Lundell 1969).

Acroclidium guianense Nees, *Syst. Laur.*: 268 [30 Oct.-5 Nov. 1836] (Nees 1836).

Acroclidium guianense var. *caudatum* Meisn., *Prod. [A. P. de Candolle]* 15 (1): 85 [May 1864] (Meisner 1864). — *Nectandra caudata* Nees & Mart. ex Meisn., *Prod. [A. P. de Candolle]* 15 (1): 85 [May 1864] (Meisner 1864), nom. nud. pro syn.

Acroclidium guianense var. *reticulatum* Meisn., *Prod. [A. P. de Candolle]* 15 (1): 85 [May 1864] (Meisner 1864).

Acroclidium meissneri Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 98 (Mez 1889). — *Licaria meissneri* (Mez) Kosterm., *Recueil Trav. Bot. Néerl.* 34: 597 (Kostermans 1937). — *Misanteca meissneri* (Mez) Lundell, *Wrightia* 4 (3): 100 (Lundell 1969).

NOTE. — *Acroclidium guianense* var. *caudatum* is based on the type of *Acroclidium meissneri*.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1205*, 15 m × 35 cm.

[611] *Licaria rufotomentosa* van der Werff

Novon 13 (3): 342 [autumn 2003] (van der Werff 2003).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *B. Dutrèvre & F. Crozier 7* (holo-, MO[MO-104256, MO-104257]; iso-, CAY[CAY014160, CAY014161]).

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 32.5$ cm.

[612] *Licaria subbulbata* Kosterm.

Reinwardtia 6 (3): 286 (Kostermans 1962). — *Misanteca subbulbata* (Kosterm.) Lundell, *Wrightia* 4 (3): 101 (Lundell 1969).

Licaria wilhelminensis C.K.Allen, *Mem. New York Bot. Gard.* 15: 65 (Allen 1966). — *Misanteca wilhelminensis* (C.K.Allen) Lundell, *Wrightia* 4 (3): 102 (Lundell 1969).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2559*, 15 m × 12 cm.

[613] *Licaria vernicosa* (Mez) Kosterm.

Recueil Trav. Bot. Néerl. 34: 604 (Kostermans 1937). — *Ocotea vernicosa* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 331 (Mez 1889). — *Acroclidium vernicosum* (Mez) Kosterm., *Recueil Trav. Bot. Néerl.* 33: 753 (Kostermans 1936). — *Misanteca vernicosa* (Mez) Lundell, *Wrightia* 4 (3): 102 (Lundell 1969).

Licaria simulans C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 55 (Allen 1964).

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.-F. Molino 1363*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 13.5$ cm.

Genus *Mezilaurus* Kuntze ex Taub.

[614] *Mezilaurus itauba* (Meisn.) Taub. ex Mez

Arbeiten Königl. Bot. Gart. Breslau 1: 112 (Mez 1892), “*Ita-Uba*”. — *Acroclidium itauba* Meisn., *Prod. [A. P. de Candolle]* 15 (1): 86 [May 1864] (Meisner 1864), “*Ita-uba*”. — *Endiandra itauba* (Meisn.) Benth. & Hook.f., *Gen. Pl. [Bentham & Hooker f.]* 3 (1): 154 [7 Feb. 1880] (Bentham & Hooker 1880), “*Ita-uba*”. — *Silvia itauba* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 107 (Mez 1889), “*Ita-Uba*”. — *Mezia itauba* (Meisn.) Kuntze, *Revis. Gen. Pl.* 2: 574 [5 Nov. 1891] (Kuntze 1891), “*Ita-Uba*”.

Acroclidium anacardiooides Spruce ex Meisn., *Prod. [A. P. de Candolle]* 15 (1): 86 [May 1864] (Meisner 1864). — *Misanteca anacardiooides* (Spruce ex Meisn.) Benth., *Hooker's Icon. Pl.* 13 [ser. 3, 3] (2): 47 [June 1878] (Bentham 1878). — *Silvia anacardiooides* (Spruce ex Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 108 (Mez 1889). — *Mezia anacardiooides* (Spruce ex Meisn.) Kuntze, *Revis. Gen. Pl.* 2: 574 [5 Nov. 1891] (Kuntze 1891). — *Mezilaurus anacardiooides* (Spruce ex Meisn.) Taub. ex Mez, *Arbeiten Königl. Bot. Gart. Breslau* 1: 112 (Mez 1892).

Oreodaphne hookeriana Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 131 [May 1864] (Meisner 1864).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). The specific epithet refers to the Tupi-Guarani word *itaúba* meaning “forest on rock”, from *ita* (rock) and *úba* (forest). The hyphen was artificially introduced (“Ita-úba”) by Meissner (in Candolle 1864: 86) and has therefore to be suppressed according to Art. 60.9 of the Shenzhen Code (Turland *et al.* 2018).

VERNACULAR NAMES. — Ka: ikalikanali • Br: itaúba, louro-itaúba.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2716*.

INVENTORY DATA (FG). — 33 trees in 22 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 55.7$ cm.

Genus *Nectandra* Rol. ex Rottb.

[615] *Nectandra amazonum* Nees

Syst. Laur.: 282 [30 Oct.-5 Nov. 1836] (Nees 1836).

Nectandra pallida Nees, *Linnaea* 21: 510 (Nees 1848).

Nectandra urophylla Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 157 [May 1864] (Meisner 1864).

Nectandra ambigua Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 158 [May 1864] (Meisner 1864).

Nectandra bombycinia S. Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 449 [“1894-96” publ. Dec. 1895] (Moore 1895).

VERNACULAR NAMES. — Br: louro-do-igapó.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *O. Poncy et al. 1680*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $dbh_{\text{inv}} = 33.6$ cm.

[616] *Nectandra cissiflora* Nees

Syst. Laur.: 296 [30 Oct.-5 Nov. 1836] (Nees 1836).

Nectandra myriantha Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 163 [May 1864] (Meisner 1864).

Nectandra myriantha var. *attenuata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 163 [May 1864] (Meisner 1864).

Nectandra myriantha var. *glabrata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 163 [May 1864] (Meisner 1864).

Nectandra surinamensis Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 454 (Mez 1889), *nom. illeg. hom.*, non Rottb. ex Bercht. & J. Presl (Berchtold & Presl 1825).

Nectandra kuntzeana Mez in Kuntze, *Revis. Gen. Pl.* 3 (3): 277 [28 Sep. 1898] (Mez 1898).

Nectandra capanahuensis O.C. Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 228 [30 Mar. 1928] (Schmidt 1928).

Nectandra steinbachii O.C. Schmidt, *Repert. Spec. Nov. Regni Veg.* 31: 186 (Schmidt 1933).

Nectandra paulii C.K. Allen, *J. Arnold Arbor.* 26 (4): 400 [15 Oct. 1945] (Allen 1945).

VERNACULAR NAMES. — Br: canela-fedida, louro-babão.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2484*.

INVENTORY DATA (FG). — 9 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 44$ cm.

[617] *Nectandra globosa* (Aubl.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 415 (Mez 1889). — *Laurus globosa* Aubl., *Hist. Pl. Guiane* 1: 364 [Jun.-Dec. 1775] (Aublet 1775). — *Persea globosa* (Aubl.) Spreng., *Syst. Veg. [Sprengel]* 2: 269 [Jan.-May 1825] (Sprengel 1825). — *Ocotea globosa* (Aubl.) Schlechtdl. & Cham., *Linnaea* 6: 366 (Schlechtdal & Chamisso 1831).

Nectandra pisi Miq., *Stirp. Surinam. Select.*: 199 [“1850” publ. Mar. 1851] (Miquel 1851).

Nectandra vaga Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 153 [May 1864] (Meisner 1864).

Nectandra vaga var. *major* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 154 [May 1864] (Meisner 1864).

Nectandra vaga var. *sprucei* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 154 [May 1864] (Meisner 1864).

Nectandra vaga var. *vulgaris* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 154 [May 1864] (Meisner 1864).

Nectandra globosa var. *barbeyana* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 416 (Mez 1889).

VERNACULAR NAMES. — Wp: mulei.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000947248]).

INVENTORY DATA (FG). — 8 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23.4$ cm.

[618] *Nectandra hibua* (Ruiz & Pav.) Rohwer

Fl. Neotrop. Monogr. 60: 196 [13 Aug. 1993] (Rohwer 1993). — *Laurus hibua* Ruiz & Pav., *Flora Peruviana* 4: t. 366 [1804-1830] (Ruiz & Pavón 1804).

Nectandra lucida Nees, *Syst. Laur.*: 334 [30 Oct.-5 Nov. 1836] (Nees 1836), *nom. illeg. hom.*, non Nees (*Linnaea* 8: 47. 1833), *nec* Nees (1836: 334).

Nectandra willdenoviana Nees, *Syst. Laur.*: 290 [30 Oct.-5 Nov. 1836] (Nees 1836), “*Nectandra (Pomatiopsis) willdenoviana*”, *non* Nees (1836: 321).

Nectandra glabrescens Benth., *Bot. Voy. Sulphur [Bentham]*: 161 [8 May 1846] (Bentham 1846).

Nectandra bredemeyeriana Nees, *Linnaea* 21: 505 (Nees 1848).

Nectandra magnoliifolia A.Rich., *Hist. Fís. Cuba, Bot.* 11: 188 (Richard 1850), “*magnoliaefolia*”.

Nectandra leucantha Nees & Mart. var. *attenuata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 151 [May 1864] (Meisner 1864).

Nectandra leucantha var. *guianensis* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 151 [May 1864] (Meisner 1864).

Nectandra leucantha var. *peruviana* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 151 [May 1864] (Meisner 1864).

Nectandra schomburgkii Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 151 [May 1864] (Meisner 1864).

Nectandra grandiflora Nees & Mart. var. *latifolia* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 152 [May 1864] (Meisner 1864), *nom. illeg. hom.*, *non* Nees (1836: 323).

Nectandra antillana Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 153 [May 1864] (Meisner 1864).

Sassafridium macrophyllum Rose, *Contr. U.S. Natl. Herb.* 1 (9): 355 [31 Jan. 1895] (Rose 1895).

Nectandra guanaiensis Rusby, *Bull. New York Bot. Gard.* 6 (22): 508 [30 Nov. 1910] (Rusby 1910).

Nectandra megaphylla Hassl., *Annuaire Conserv. Jard. Bot. Genève* 21: 92 (Hassler 1919).

Nectandra maranonensis O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 229 [30 Mar. 1928] (Schmidt 1928).

Nectandra tessmannii O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 229 [30 Mar. 1928] (Schmidt 1928).

Nectandra albiflora Lundell, *Wrightia* 1 (2): 147 (Lundell 1946).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori & C. Gracie 24212*.

SIZE. — Up to 35 m tall (Rohwer 1993).

[619] *Nectandra matogrossensis* Coe-Teix.
(Fig. 26D-F)

Acta Amazonica 5 (2): 158 (Coe-Teixeira 1975).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3406*.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 14.4$ cm.

[620] *Nectandra pulverulenta* Nees

Syst. Laur.: 283 [30 Oct.-5 Nov. 1836] (Nees 1836).

Nectandra purusensis Coe-Teix., *Acta Amazonica* 5 (2): 158 (Coe-Teixeira 1975).

VERNACULAR NAMES. — Wp: áyū'i sili, áyū'i tawa, áyū'i witowa • Br: louro-abacate.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 1000*.

SIZE. — Up to 30 m tall (Rohwer 1993).

[621] *Nectandra reticulata* (Ruiz & Pav.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 404 (Mez 1889). — *Laurus reticulata* Ruiz & Pav., *Flora Peruviana* 4: t. 348 [1804-1830] (Ruiz & Pavón 1804).

Ocotea mollis Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 164 [8 Dec. 1817] (Kunth 1818). — *Persea mollis* (Kunth) Spreng., *Syst. Veg. [Sprengel]* 2: 270 [Jan.-May 1825] (Sprengel 1825). — *Nectandra mollis* (Kunth) Nees, *Hufeland.* III: 14 (Nees 1833).

Persea incana Schott, *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 405 [Jan.-June 1827] (Schott 1827).

Laurus aestivalis Vell., *Fl. Flumin.*: 164 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), *nom. illeg. hom.*, *non* L. (Linnaeus 1753).

Nectandra villosa Nees & Mart., *Linnaea* 8: 47 (Nees & Martius 1833). — *Nectandra mollis* var. *villosa* (Nees & Mart.) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 148 [May 1864] (Meisner 1864).

Nectandra villosa var. *venosa* Nees, *Syst. Laur.*: 291 [30 Oct.-5 Nov. 1836] (Nees 1836), "Venosa". — *Nectandra mollis* var. *venosa* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 149 [May 1864] (Meisner 1864).

Nectandra discolor (Kunth) Nees var. *subvenosa* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 157 [May 1864] (Meisner 1864).

Nectandra laurel Klotzsch ex Nees var. *triquetra* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 147 [May 1864] (Meisner 1864).

Nectandra mollis var. *attenuata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 148 [May 1864] (Meisner 1864).

Nectandra mollis var. *intermedia* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 149 [May 1864] (Meisner 1864).

Nectandra pittieri Lasser, *Bol. Técn. Minist. Agric.* 3: 13 (Lasser 1942).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3309*.

INVENTORY DATA (FG). — 6 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{inv} = 55.5$ cm.

[622] *Nectandra* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2816*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 28.7$ cm.

Genus *Ocotea* Aubl.

[623] *Ocotea amazonica* (Meisn.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 359 (Mez 1889). — *Oreodaphne amazonica* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 123 [May 1864] (Meisner 1864).

VERNACULAR NAMES. — Nt: fiiman • Br: louro-preto.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3593*.

INVENTORY DATA (FG). — 15 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{inv} = 67$ cm.

[624] *Ocotea argyrophylla* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 193 (Ducke 1925).

VERNACULAR NAMES. — Wp: áyú'i alapítá milá • Wn: taihem • Nt: weti apisi, weti sikin apisi • Cr: sèd-nwé • Fr: cèdre argenté • Br: folha-de-prata.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2147.

INVENTORY DATA (FG). — 93 trees in 43 plots; $F_{\max} = 2.4\%$; $dbh_{\text{inv}} = 64.1\text{ cm}$.

[625] *Ocotea canaliculata* (Rich.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 361 (Mez 1889). — *Laurus canaliculata* Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792).

Oreodaphne cayennensis Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 133 [May 1864] (Meisner 1864), *syn. nov.*

Ocotea scrobiculata Benoist, *Bull. Soc. Bot. France* 75 (5): 979 ["1928 publ. 1929"] (Benoist 1929), *nom. nud.*

VERNACULAR NAMES. — Wp: áyú'i witowa • Br: louro-pimenta.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (holo-, P[P00756974]; P[P00756975], P00756976).

INVENTORY DATA (FG). — 10 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 47.4\text{ cm}$.

[626] *Ocotea ceanothifolia* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 365 (Mez 1889). — *Mespilodaphne ceanothifolia* Nees, *Syst. Laur.*: 236 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Persea ceanothifolia* Mart. ex Nees, *Syst. Laur.*: 236 [30 Oct.-5 Nov. 1836] (Nees 1836), *nom. nud. pro syn.*

VERNACULAR NAMES. — Wp: áyú'i tawa • Nt: likanau apisi.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4200.

INVENTORY DATA (FG). — 25 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 103\text{ cm}$.

[627] *Ocotea cernua* (Nees) Mez

Mitt. Bot. Vereins Kreis Freiburg 47-48: 422 (Mez 1888). — *Oreodaphne cernua* Nees, *Syst. Laur.*: 424 [30 Oct.-5 Nov. 1836] (Nees 1836).

Laurus nitida Ruiz & Pav., *Flora Peruviana* 4: t. 353 [1804-1830] (Ruiz & Pavón 1804). — *Mespilodaphne nitida* (Ruiz & Pav.). Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 106 [May 1864] (Meisner 1864).

Laurus leptobotra Ruiz & Pav., *Flora Peruviana* 4: t. 368 [1804-1830] (Ruiz & Pavón 1804). — *Oreodaphne leptobotra* (Ruiz & Pav.). Nees, *Linnæa* 21: 520 (Nees 1848). — *Ocotea leptobotra* (Ruiz & Pav.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 376 (Mez 1889).

Leptodaphne tenuiflora Nees, *Syst. Laur.*: 359 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Persea tenuiflora* Mart. ex Nees, *Syst. Laur.*: 359 [30 Oct.-5 Nov. 1836] (Nees 1836), *nom. nud. pro syn.*

Leptodaphne subalpina Nees, *Syst. Laur.*: 361 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Oreodaphne subalpina* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 137 [May 1864] (Meisner 1864). — *Ocotea subalpina* (Nees) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 375 (Mez 1889).

Oreodaphne caudata Nees, *Linnæa* 21: 519 (Nees 1848). — *Camphoromoea surinamensis* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 145 [May 1864] (Meisner 1864), *nom. illeg. superfl.* (one of the syntypes [Hostmann 964] is a syntype of *Oreodaphne caudata*). — *Ocotea caudata* (Nees) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 378 (Mez 1889). — *Camphoromoea tenuiflora* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 144 [May 1864] (Meisner 1864).

Oreodaphne marowynensis Miq., *Stirp. Surinam. Select.*: 201 ["1850" publ. Mar. 1851] (Miquel 1851). — *Ocotea marowynensis* (Miq.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 380 (Mez 1889).

Oreodaphne urophylla Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 145 [May 1864] (Meisner 1864). — *Ocotea urophylla* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 379 (Mez 1889).

Ocotea marmellensis Mez var. *acrensis* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.3): 903 [31 Oct. 1938] (Macbride 1938).

NOTES. — Excluded synonym: *Nectandra revoluta* Miq. ex Rohwer (1993: 302), *nom. nud.* Duplicates of *Pl. Surinam*. 629bb distributed under A. Kappler's name (M, MPU, P[P00756833], S, U[U0002941]) bear on the original label "*Nectandra revoluta* Miq." (This is not the case for the duplicates distributed under Hostmann's name). A label by C.K. Allen dated 1970 affixed to the S duplicate wrongly states that this specimen is an isotype of *N. revoluta*, and that this name is a synonym of *Oreodaphne marowynensis* Miquel (1851: 201). With respect to the latter name, Miquel originally wrote (1851: 201): "*Nectandra revoluta* a me perperam dicta" ("which I wrongly called *N. revoluta*"). Therefore, A. Kappler 629bb is not the type of *N. revoluta*, which is a *nomen nudum* and is not a synonym of *O. marowynensis*.

VERNACULAR NAMES. — Ka: wa'elan, wayaka • Te: uwahe hun • Wp: áyú'i tawa, áyú'i witowa.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2719.

INVENTORY DATA (FG). — 22 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23.9\text{ cm}$.

[628] *Ocotea cinerea* van der Werff

Novon 10 (3): 268 [autumn 2000] (van der Werff 2000).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: migukat, miukat, mirukat • Wp: kwata pili • Nt: baaka sikin apisi.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *S.A. Mori et al.* 23898.

INVENTORY DATA (FG). — 159 trees in 58 plots; $F_{\max} = 4.9\%$; $dbh_{\text{inv}} = 98.7\text{ cm}$.

[629] *Ocotea commutata* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 327 (Mez 1889). — *Oreodaphne commutata* Nees, *Syst. Laur.*: 428 [30 Oct.-5 Nov. 1836] (Nees 1836).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Wp: áyú'i.

HERBARIUM DATA (FG). — 11 collections at CAY. *J. Martin s.n.* (holo-, B[B100247389]; iso-, B[B100247390], B100247391], GZU[GZU000254344], K[K000602279, K000602280], P[P00756980, P00756981, P00756982], U[U0002942]).

INVENTORY DATA (FG). — 1 tree, dbh = 14.3 cm.

[630] *Ocotea cujumary* Mart.

Repert. Pharm. [J.A. Buchner] 35: 178 (Martius 1830). — *Alseodaphne cujumary* (Mart.) Kostel., *Allg. Med.-Pharm. Fl.* 2: 484 [Jan.-June 1833] (Kosteletzky 1833). — *Ayndendron cujumary* (Mart.) Nees, *Syst. Laur.*: 247 [30 Oct.-5 Nov. 1836] (Nees 1836).

Oreodaphne macrothrysus Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 115 [May 1864] (Meisner 1864).

Aniba ovalifolia Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 77 (Mez 1889).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5707*.

INVENTORY DATA (FG). — 16 trees in 9 plots; $F_{\max} = 1\%$; $dbh_{inv} = 30.6$ cm.

[631] *Ocotea diffusa* van der Werff

Brittonia 49 (2): 189 (van der Werff 1997).

VERNACULAR NAMES. — Pa: uvayan-akig • Wp: ka'i tulí.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.-J. de Granville B-5195* (holo-, P[P00098348]; iso-, CAY[CAY027845, CAY171665, CAY171666]).

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.5$ cm.

[632] *Ocotea endlicheriopsis* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 300 (Mez 1889). — *Endlicheria endlicheriopsis* (Mez) Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 43 (Kostermans 1936).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* ("605" *fide* Mez 2005: 300), 1862 (original material B[not seen, photo F neg. N° 3651], BM[BM000993946], K[K000602244], P[P00745675, P00048601]).

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[633] *Ocotea fasciculata* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 248 (Mez 1889). — *Oreodaphne fasciculata* Nees, *Linnaea* 21: 521 (Nees 1848). — *Mespilodaphne fasciculata* (Nees) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 108 [May 1864] (Meisner 1864).

Mespilodaphne complicata Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 103 [May 1864] (Meisner 1864). — *Ocotea complicata* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 247 (Mez 1889).

Ocotea nunesii Vattimo-Gil, *Arch. Jard. Bot. Rio de Janeiro* 15: 142 (Vattimo-Gil 1957).

Ocotea duckei Vattimo-Gil, *Rodriguésia* 23 (35): 243 (Vattimo-Gil 1961).

Cinnamomum babiense Vattimo-Gil, *Rodriguésia* 29 (42): 128 (Vattimo-Gil 1977).

Ocotea scrobiculifera Vattimo-Gil, *Rodriguésia* 29 (42): 129 (Vattimo-Gil 1977).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2979*.

INVENTORY DATA (FG). — 27 trees in 5 plots; $F_{\max} = 3.3\%$; $dbh_{inv} = 39.2$ cm.

[634] *Ocotea fendleri* (Meisn.) Rohwer

Mitt. Inst. Allg. Bot. Hamburg 20: 152 (Rohwer 1986). — *Gymnobalanus fendleri* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 142 [May 1864] (Meisner 1864).

Ocotea glandulosa Lasser, *Bol. Técn. Minist. Agric.* 3: 9 (Lasser 1942).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al. 20862*.

INVENTORY DATA (FG). — 1 tree, dbh = 12.6 cm.

[635] *Ocotea floribunda* (Sw.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 325 (Mez 1889). — *Laurus floribunda* Sw., *Prod. [Swartz]* 65 [20 Jun.-29 July 1788] (Swartz 1788). — *Nectandra floribunda* (Sw.) Nees, *Syst. Laur.*: 339 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Strychnodaphne floribunda* (Sw.) Griseb., *Pl. Wright. [Grisebach]* 1: 188 [Dec. 1860] (Grisebach 1860).

Laurus cerifera Vahl, *Skr. Naturhist.-Selsk.* 6: 130 (Vahl 1810).

Laurus divaricata Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 322 [24 Sep. 1813] (Poiret 1813). — *Oreodaphne divaricata* (Poir.) Nees, *Syst. Laur.*: 393 [30 Oct.-5 Nov. 1836] (Nees 1836).

Laurus retroflexa Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 323 [24 Sep. 1813] (Poiret 1813). — *Persea retroflexa* (Poir.) Spreng., *Syst. Veg. [Sprengel]* 2: 269 [Jan.-May 1825] (Sprengel 1825). — *Oreodaphne retroflexa* (Poir.) Nees, *Syst. Laur.*: 432 [30 Oct.-5 Nov. 1836] (Nees 1836).

Oreodaphne willdenoviana Nees, *Syst. Laur.*: 412 [30 Oct.-5 Nov. 1836] (Nees 1836).

Oreodaphne domingensis Nees, *Syst. Laur.*: 439 [30 Oct.-5 Nov. 1836] (Nees 1836).

Ocotea botryophylla Klotzsch & H.Karst. ex Nees, *Linnaea* 21: 524 (Nees 1848).

Ocotea wachenheimii Benoist, *Bull. Mus. Natl. Hist. Nat.* 30: 510 (Benoist 1924).

Ocotea arenaensis R.L.Brooks, *Bull. Misc. Inform. Kew* 1933 (5): 217 [12 July 1933] (Brooks 1933).

VERNACULAR NAMES. — Ka: wa'e, waye • Wp: amu'a iwa u, áyū'i pino, áyū'i sili • Cr: sèd-gri.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5347.

INVENTORY DATA (FG). — 26 trees in 18 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 37.1$ cm.

[636] *Ocotea glomerata* (Nees) Mez
(Fig. 27A, B)

Jahrb. Königl. Bot. Gart. Berlin 5: 294 (Mez 1889). — *Oreodaphne glomerata* Nees, *Linnaea* 21: 515 (Nees 1848).

Oreodaphne moritziana Nees, *Linnaea* 21: 515 (Nees 1848).

VERNACULAR NAMES. — Ka: wa'elan, walili pipyo, wayaka • Te: uwahé • Wp: áyū'i alapitá milá, áyū'i piyú • Nt: baaka apisi, baaka sikin • Cr: sèd-bagas, sèd-gri • Fr: cèdre bagasse, cèdre noir • Br: louro-preto.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2262.

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 52.4$ cm.

[637] *Ocotea guianensis* Aubl.

Hist. Pl. Guiane 2: 781 [Jun.-Dec. 1775] (Aublet 1775), “*Guyanensis*” on plate. — *Laurus ocotea* Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792), nom. illeg. superfl. (based on *Ocotea guianensis*). — *Oreodaphne guianensis* (Aubl.) Nees, *Linnaea* 21: 516 (Nees 1848).

Nectandra surinamensis Rottb. ex Bercht. & J.Presl, *Přir. Rostlin* 2: 60 (Berchtold & Presl 1825), nom. nud. pro syn.

Oreodaphne guianensis var. *argentea* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 112 [May 1864] (Meisner 1864), nom. illeg. superfl. (includes the type of the typical variety).

Oreodaphne guianensis var. *aurea* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 112 [May 1864] (Meisner 1864). — *Ocotea guianensis* var. *aurea* (Meisn.) R.Knuth, *Repert. Spec. Nov. Regni Veg. Beih.* 43: 337 (Knuth 1927).

Ocotea guianensis var. *subsericea* Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 15 (Kostermans 1936).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). One specimen of *O. guianensis* at P[P00756908] is erroneously labelled type: it bears fruits, whereas Aublet's plate and the original BM material have flowers.

VERNACULAR NAMES. — Pa: wen, wen-ethni • Ka: tokowe • Wp: áyū'i sili • Nt: weti apisi, weti sikin apisi • Cr: féy-darjan, sèd-darjan • Fr: bois fribustier, cèdre argenté, cèdre d'argent, feuille d'argent • Br: folha-de-prata, louro-branco, louro-prata, louro-seda.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000947302]).

INVENTORY DATA (FG). — 12 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 64.3$ cm.

[638] *Ocotea leucoxylon* (Sw.) Laness.

Pl. Util. Col. Franç. 158 (Lanessan 1886). — *Laurus leucoxylon* Sw., *Prodr. [Swartz]* 65 [20 Jun.-29 July 1788] (Swartz 1788). — *Persea leucoxylon* (Sw.) Spreng., *Syst. Veg. [Sprengel]* 2: 268 [Jan.-May 1825] (Sprengel 1825). — *Oreodaphne leucoxylon* (Sw.) Nees, *Syst. Laur.*: 413 [30 Oct.-5 Nov. 1836] (Nees 1836).

Laurus parviflora Sw., *Fl. Ind. Occid.* 2: 717 [Jan.-June 1798] (Swartz 1798). — *Persea parviflora* (Sw.) Spreng., *Syst. Veg. [Sprengel]* 2: 270 [Jan.-May 1825] (Sprengel 1825). — *Oreodaphne parviflora* (Sw.) Nees, *Syst. Laur.*: 415 [30 Oct.-5 Nov. 1836] (Nees 1836).

Laurus exaltata Sw. var. *lancifolia* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 319 [24 Sep. 1813] (Poirier 1813).

Nectandra acutangula Miq., *Linnaea* 22: 806 [“1849” publ. May 1850] (Miquel 1850). — *Oreodaphne acutangula* (Miq.) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 121 [May 1864] (Meisner 1864). — *Ocotea acutangula* (Miq.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 330 (Mez 1889).

Oreodaphne lindeniana A.Rich., *Hist. Fís. Cuba, Bot.* 11: 189 (Richard 1850).

Nectandra longifolia A.Rich., *Hist. Fís. Cuba, Bot.* 11: 188 (Richard 1850), nom. illeg. hom., non (Ruiz & Pav.) Nees (1848).

Oreodaphne leucoxylon var. *elongata* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 121 [May 1864] (Meisner 1864). — *Ocotea leucoxylon* var. *elongata* (Meisn.) M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.*: 12 (Gómez 1889).

Ocotea subracemosa Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 360 (Mez 1889).

Ocotea froesii A.C.Sm., *Bull. Torrey Bot. Club* 61 (4): 191 [Apr. 1934] (Smith 1934).

Ocotea subsericea Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 18 (2): 456 [20 Oct. 1937] (Standley 1937).

Ocotea lasseriana C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 90 (Allen 1964).

Ocotea duotincta C.K.Allen, *Mem. New York Bot. Gard.* 12 (3): 119 (Allen 1965).

Ocotea lenticellata Lundell, *Wrightia* 5 (3): 54 (Lundell 1974).

VERNACULAR NAMES. — Wp: amu'a iwa, áyū'i sili.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2767.

INVENTORY DATA (FG). — 7 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31$ cm.

[639] *Ocotea longifolia* Kunth

Nova genera et species plantarum [H.B.K.] 2: 164 [8 Dec. 1817] (Kunth 1818). — *Persea longifolia* (Kunth) Spreng., *Syst. Veg. [Sprengel]*

2: 270 [Jan.-May 1825] (Sprengel 1825). — *Oreodaphne longifolia* (Kunth) Nees, *Syst. Laur.*: 391 [30 Oct.-5 Nov. 1836] (Nees 1836).

Ocotea opifera Mart., *Repert. Pharm. [J.A. Buchner]* 35: 179 (März 1830). — *Oreodaphne opifera* (Mart.) Nees, *Syst. Laur.*: 390 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Mespilodaphne opifera* (Mart.) Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 510 [May 1864] (Meisner 1864).

Oreodaphne grandifolia Nees, *Linnaea* 21: 517 (Nees 1848). — *Ocotea grandifolia* (Nees) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 290 (Mez 1889).

Ocotea rusbyana Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 303 (Mez 1889).

Ocotea trianae Rusby, *Bull. New York Bot. Gard.* 6 (22): 506 [30 Nov. 1910] (Rusby 1910).

Ocotea dielsiana O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 232 [30 Mar. 1928] (Schmidt 1928).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2417.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 15$ cm.

[640] *Ocotea montis-insulae* van der Werff

Blumea 56 (3): 214 (van der Werff 2011).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *G. Cremers & F. Crozier* 15215 (holo-, CAY[CAY101097]; iso-, K[K001057503], MO[MO-359579], NY[01365077], U[U.1359518], US[00618793]), 10 m.

[641] *Ocotea mucronata* (Poir.) Kosterm.

Reinwardtia 5 (4): 395 (Kostermans 1961). — *Laurus mucronata* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 323 [24 Sep. 1813] (Poiret 1813). — *Cryptocarya mucronata* (Poir.) Spreng., *Syst. Veg. [Sprengel]* 2: 271 [Jan.-May 1825] (Sprengel 1825).

HERBARIUM DATA (FG). — Known only from the type: *Unknown coll. s.n.* (holo-, FI[FI005245]; iso-, FI[FI005244]).

SIZE. — Not recorded on the type specimen, but most probably a tree.

[642] *Ocotea nigra* Benoist

Bull. Mus. Natl. Hist. Nat. 30: 511 (Benoist 1924).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: takana • Wp: áyú'i.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *P. Grenand et al.* 3425.

INVENTORY DATA (FG). — 42 trees in 30 plots; $F_{\max} < 1\%$; $dbh_{inv} = 62$ cm.

[643] *Ocotea oblonga* (Meisn.) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 367 (Mez 1889). — *Mespilodaphne oblonga* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 107 [May 1864] (Meisner 1864).

VERNACULAR NAMES. — Pa: akamna-seinó, akamná-seinó, akapna-seinó • Wp: áyú'i, iwa'e • Cr: sèd-blanc, sèd-gri, sèd-kalalou.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *P.A. Sagot* 491, 1857 (holo-, G-DC, not seen; iso-, B[B100247523], BM[BM000993958], BR[BR0000005119950], G[G00369312], K[K000602355], P[P00706273, P00756749, P00756750]).

INVENTORY DATA (FG). — 23 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{inv} = 87$ cm.

[644] *Ocotea pauciflora* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 370 (Mez 1889). — *Oreodaphne pauciflora* Nees, *Syst. Laur.*: 409 [30 Oct.-5 Nov. 1836] (Nees 1836).

Mespilodaphne laxiflora Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 107 [May 1864] (Meisner 1864). — *Ocotea laxiflora* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 371 (Mez 1889).

Oreodaphne diospyrifolia Meisn. var. *incompta* Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 126 [May 1864] (Meisner 1864).

Oreodaphne paraensis Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 126 [May 1864] (Meisner 1864).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.C. Ek* 1835.

SIZE. — Guyana. *B. Hoffman & M.T. Rodrigues* 4705 (MO), 12 m × 10 cm.

[645] *Ocotea percurrens* Vicent.

Novon 10 (3): 281 [autumn 2000] (Vicentini 2000).

VERNACULAR NAMES. — Pa: migukat-kamwi, miukat-kamwi, miukat-kamwi-wahuyo, mirukat-kamwi • Te: uwahe tsing.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5127.

INVENTORY DATA (FG). — 111 trees in 65 plots; $F_{\max} = 1.1\%$; $dbh_{inv} = 83$ cm.

[646] *Ocotea persulcata* C.K.Allen

Mem. New York Bot. Gard. 12 (3): 120 (Allen 1965).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori* 25053.

INVENTORY DATA (FG). — 20 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{inv} = 49$ cm.




FIG. 27. — Lauraceae: **A, B**, *Ocotea glomerata* (Nees) Mez (J.-F. Molino & D. Sabatier 2262); **C, D**, *Ocotea* sp. D (J.-F. Molino & D. Sabatier 2773); **E**, *Rhodostemonodaphne grandis* (Mez) Rohwer; **F**, *Rhodostemonodaphne revolutifolia* Madriñán (D. Sabatier & J.-F. Molino 5078). A-D, © J.-F. Molino/IRD; E, F, © D. Sabatier/IRD.

[647] *Ocotea puberula* (Rich.) Nees

Syst. Laur.: 472 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Laurus puberula* Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792). — *Strychnodaphne puberula* (Rich.) Nees & Mart., *Linnaea* 8: 39 (Nees & Martius 1833).

Persea macropoda Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 160 [8 Dec. 1817] (Kunth 1818). — *Ocotea macropoda* (Kunth) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 348 (Mez 1889). — *Ocotea martiniana* (Nees) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 344 (Mez 1889).

Persea richardiana Cham. & Schlehd., *Linnaea* 6: 366 (Chamisso & Schlechtdal 1831).

Oreodaphne acutifolia Nees var. *latifolia* Nees, *Linnaea* 8: 42 (Nees 1833). — *Strychnodaphne puberula* var. *angustata* Meisn., *Prod. [A. P. de Candolle]* 15(1): 142 [May 1864] (Meisner 1864), nom. illeg. superfl. (based on *Oreodaphne acutifolia* var. *latifolia*).

Oreodaphne martiniana Nees, *Syst. Laur.*: 415 [30 Oct.-5 Nov. 1836] (Nees 1836).

Oreodaphne hostmanniana Miq., *Stirp. Surinam. Select.*: 202 [“1850” publ. Mar. 1851] (Miquel 1851).

Strychnodaphne puberula var. *truncata* Meisn., *Prod. [A. P. de Candolle]* 15 (1): 143 [May 1864] (Meisner 1864). — *Ocotea*

puberula var. *truncata* (Meisn.) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 344 (Mez 1889).

Ocotea pyramidata S.F.Blake ex Brandegee, *Univ. Calif. Publ. Bot.* 7: 326 (Brandegee 1920).

Ocotea ucalayensis O.C.Schmidt, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 234 [30 Mar. 1928] (Schmidt 1928).

Ocotea subglabra Benoist, *Arch. Bot. Bull. Mens.* 3: 10 (Benoist 1929).

Ocotea baturitensis Vattimo-Gil, *Rodriguésia* 23 (35): 246 (Vattimo-Gil 1961).

Ocotea paraensis Coe-Teix., *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 34: 11 (Coe-Teixeira 1970).

Ocotea paranapiacabensis Coe-Teix., *Rodriguésia* 32 (52): 107 (Coe-Teixeira 1980).

VERNACULAR NAMES. — Pa: akamna, akamnā, akapna • Ka: kisipyululan, sipilulan • Wp: amu'a iwa • Cr: séd • Br: canela-pimentá, guaicá.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *J.B. Leblond* 215 (type G[G00020781]).

INVENTORY DATA (FG). — 11 trees in 8 plots; $F_{\max} < 1\%$; dbh_{inv} = 59.4 cm.

[648] *Ocotea scabrella* van der Werff

Novon 10 (3): 287 [autumn 2000] (van der Werff 2000).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *S.A. Mori* 14824.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.1$ cm.

[649] *Ocotea schomburgkiana* (Nees) Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 337 (Mez 1889). — *Oreodaphne schomburgkiana* Nees, *Linnaea* 21: 269 (Nees 1848).

Ocotea castanea C.K.Allen, *Mem. New York Bot. Gard.* 12 (3): 114 (Allen 1965).

NOTES. — Four French Guianan specimens (*D. Sabatier* & *M.-F. Prévost* 2344, 2510, 4432, and 3946) are wrongly identified as *Ocotea schomburgkiana* (Nees) Mez (1889: 337) on the MO duplicates. The first three actually belong to *O. cinerea* van der Werff (van der Werff & Vicentini 2000: 268), the latter to our *O.* sp. B.

VERNACULAR NAMES. — Pa: sedri-priyu • Ka: yapui, yolokan topulukali • Nt: baaka sikan apisi.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *H. van der Werff* 23500 (MO), height 15 m.

[650] *Ocotea splendens* (Meisn.) Baill.

Hist. Pl. [Baillon] 2: 466 (Baillon 1870). — *Oreodaphne splendens* Meisn., *Prodri. [A. P. de Candolle]* 15 (1): 129 [May 1864] (Meisner 1864).

Ocotea citrifolia Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 280 (Mez 1889).

Ocotea globifera Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 328 (Mez 1889).

Ocotea dissimilis C.K.Allen, *Mem. New York Bot. Gard.* 15: 86 (Allen 1966).

VERNACULAR NAMES. — Ka: kulukai, wa'e, waye • Wp: áyú'i piye, áyú'i sili, áyú'i tawa, áyú'i witowa • Cr: mal-bwa-dérose • Br: cedro-pardo.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *P.A. Sagot* 978, Oct. 1856 (holo-, G-DC, not seen; iso-, B[B100185434, photo F neg N° 3719], K[K000602221], P[P00711099, P00711100, P00711101, P00711102]).

INVENTORY DATA (FG). — 50 trees in 21 plots; $F_{\max} = 4.3\%$; $dbh_{\text{inv}} = 52.2$ cm.

[651] *Ocotea subterminalis* van der Werff

Novon 10 (3): 289 [autumn 2000] (van der Werff 2000).

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *D. Sabatier et al.* 5252.

INVENTORY DATA (FG). — 109 trees in 40 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 37.9$ cm.

[652] *Ocotea tomentella* Sandwith

Bull. Misc. Inform. Kew 1935 (3): 130 [20 May 1935] (Sandwith 1935).

VERNACULAR NAMES. — Ka: mila kulan • Te: uwahé tsing • Wp: alamá'i • Wn: toto, wai • Nt: apisi, baaka sikan apisi, geli apisi • Cr: sèd-gro-lapo, sèd-jonn, sèd-nwé • Fr: cèdre jaune • Br: louro.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 4434.

INVENTORY DATA (FG). — 60 trees in 35 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 110$ cm.

[653] *Ocotea* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino* & *D. Sabatier* 2147.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 72$ cm.

[654] *Ocotea* sp. B

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 2248.

INVENTORY DATA (FG). — 73 trees in 29 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 73.2$ cm.

[655] *Ocotea* sp. C

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier* & *J.-F. Molino* 5135.

INVENTORY DATA (FG). — 17 trees in 5 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 79.7$ cm.

[656] *Ocotea* sp. D
(Fig. 27C, D)

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino* & *D. Sabatier* 2773.

INVENTORY DATA (FG). — 1 tree, $dbh = 21$ cm.

[657] *Ocotea* sp. E

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 2495.

INVENTORY DATA (FG). — 27 trees in 18 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 115$ cm.

[658] *Ocotea* sp. F

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier 1512*.

INVENTORY DATA (FG). — 1 tree, dbh = 11 cm.

[659] *Ocotea* sp. G

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 3375*.

INVENTORY DATA (FG). — 1 tree, dbh = 10.2 cm.

[660] *Ocotea* sp. H

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 3397*, 38.2 cm.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.2$ cm.

[661] *Ocotea* sp. I

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2958*.

INVENTORY DATA (FG). — 11 trees in 5 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 29.6$ cm.

[662] *Ocotea* sp. J

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2751*.

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42.6$ cm.

Genus *Persea* Mill.

[663] *Persea nivea* Mez

Jahrb. Königl. Bot. Gart. Berlin 5: 148 (Mez 1889). — *Mutisiopersexa nivea* (Mez) Kosterm., *Rheedia* 3 (2): 135 [31 Dec. 1993] (Kostermans 1993).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: migukat, miukat, mirukat.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4910*, dbh 43.6 cm.

[664] *Persea* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2096*.

INVENTORY DATA (FG). — 1 tree, dbh = 16.8 cm.

Genus *Rhodostemonodaphne* Rohwer & Kubitzki

[665] *Rhodostemonodaphne elephantopus* Madriñán

Brittonia 48 (1): 46 (Madriñán 1996).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *S.A. Mori et al. 20774* (holo-, NY[00076770]; iso-, CAY[CAY027849], GH[GH00061477], MO[MO-287214], P[P00644242], U[U0249275], US[US00902047]).

INVENTORY DATA (FG). — 10 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40.6$ cm.

[666] *Rhodostemonodaphne grandis* (Mez) Rohwer
(Fig. 27E)

Mitt. Inst. Allg. Bot. Hamburg 20: 84 (Rohwer 1986). — *Endlicheria grandis* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 124 (Mez 1889). — *Nectandra grandis* (Mez) Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 17 (Kostermans 1936).

NOTES. — The indications given by Mez (1889: 125) on the (fruiting) material he studied are: “Hab. in Guyana gallica ad fl. Maroni: Mélinon. (V. s. in herb. Berol. et Paris.)” There is only one Mélinon specimen of *R. grandis* at B, with a label in Mez’ hand reading: “*Endlicheria ? grandis* Mez nov. spec., det. Carl Mez”; its original label bears the date 1865 and does not mention the Maroni River, or any other location, within French Guiana. Among the eight Mélinon specimen of *R. grandis* at P (all in fruit), only three bear the same identification label by Mez: P00745439, with an original label reading “Guyane française (Maroni). M. Mélinon 1862”; P00745443, with a label “Herbier de la Guyane française. 1865 – Mélinon”; and P00078423, labelled “Guyane française – M. Mélinon 1863”. The five other specimens (P00745440, P00745441, P00745442, P00745444, P00745445) are all dated 1863, and only one of them (P00745441) has a label mentioning the Maroni. Another Mélinon specimen at NY (00355136) is also dated 1863 and the only geographical indication is “Guyane fse”.

Only the four specimens annotated by Mez can be considered to be indisputably part of the original material. However, they could represent separate collections made in 1862, 1863 and 1864, therefore they must be treated as syntypes.

Madriñán (2004: 85) designated a lectotype as: “French Guiana. Fleuve Maroni [St Laurent?], 1863 (fr), Mélinon 1863 [1865 in B] (lectotype, here designated: B; isolectotypes: NY, P”. This lectotypification is correct in that it unambiguously refers to a single specimen at B, that is part of the original material. However, the date, place and collection number given for this specimen are incorrect, leading to syntypes (at P and NY) being mistaken for isolectotypes. The correct citation for the lectotype is given below.

VERNACULAR NAMES. — Ka: waikyalalan • Wp: áyú'i to u, kwata pili • Wn: toto • Nt: geli apisi • Cr: sèd-jonn • Fr: cèdre jaune.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* 1865 (fr) (lecto-, B[B100247403]; iso-, P[P00745443]).

INVENTORY DATA (FG). — 136 trees in 92 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 58.6$ cm.

[667] *Rhodostemonodaphne kunthiana* (Nees) Rohwer

Mitt. Inst. Allg. Bot. Hamburg 20: 84 (Rohwer 1986). — *Acroclidium kunthianum* Nees, *Syst. Laur.*: 269 [30 Oct.-5 Nov. 1836] (Nees 1836). — *Nectandra kunthiana* (Nees) Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 19 (Kostermans 1936). — *Ocotea kunthiana* (Nees) Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 291 (Mez 1889).

Ayndendron aciphyllum Nees, *Linnaea* 21: 496 (Nees 1848).

Pleurothyrium chrysotyrsus Meisn., *Prodr. [A. P. de Candolle]* 15 (1): 169 [May 1864] (Meisner 1864).

Ocotea cooperi C.K.Allen, *J. Arnold Arbor.* 26 (3): 335 [16 July 1945] (Allen 1945).

Nectandra meyeriana Lasser, *Bol. Soc. Venez. Ci. Nat.* 11: 184 (Lasser 1948).

Pleurothyrium cowanianum C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 121 (Allen 1964), “*cowaniana*”.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holo-, B[B 10 0247364]; iso-, B[B 10 0247365], G[G00368893]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 39.5$ cm.

[668] *Rhodostemonodaphne leptoclada* Madriñán

Brittonia 48 (1): 49 (Madriñán 1996).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: sedri-puvemna-priyo • Wp: áyū’í.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori et al. 14899* (holo-, NY[00076788]; iso-, CAY[CAY027852], U[U0008013], US[00589056]).

INVENTORY DATA (FG). — 1 tree, $dbh = 16.2$ cm.

[669] *Rhodostemonodaphne morii* Madriñán

Brittonia 48 (1): 52 (Madriñán 1996).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: migukat, miukat, mirukat, sedri-puvemna.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *S.A. Mori et al. 20850* (holo-, NY[00076796]; iso-, CAY[CAY027861], MO[MO-287433], P[P00745438], US[00589054]).

INVENTORY DATA (FG). — 28 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{inv} = 51.6$ cm.

[670] *Rhodostemonodaphne revolutifolia* Madriñán
(Fig. 27F)

Brittonia 48 (1): 55 (Madriñán 1996).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier 3500* (holo-, NY[00008290]; iso-, B[B 10 0247362], CAY[CAY027874], K[K000642263], MO[MO-287481], P[P00745437], U[U0003015]).

INVENTORY DATA (FG). — 25 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{inv} = 29.9$ cm.

[671] *Rhodostemonodaphne rufovirgata* Madriñán

Brittonia 48 (1): 58 (Madriñán 1996).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: geli apisi • Cr: sèd-jonn • Fr: cèdre jaune.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori 1811* (holo-, NY[00076794]; iso-, CAY[CAY027866], COL[COL000001404], HBG[HBG508010], K[K000642275], MG[MG152546], MO[MO-287491], P[P00074147], US[00512805], VEN[VEN302994]).

INVENTORY DATA (FG). — 72 trees in 50 plots; $F_{\max} = 1.2\%$; $dbh_{inv} = 45$ cm.

[672] *Rhodostemonodaphne saulensis* Madriñán

Brittonia 48 (1): 60 (Madriñán 1996).

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *S.A. Mori et al. 15014* (holo-, NY[00076795]; iso-, CAY[CAY027877], HBG[HBG508009], K[K000642262], MO[MO-287499], U[U0008017], US[00589055]).

INVENTORY DATA (FG). — 9 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 53$ cm.

Genus *Sextonia* van der Werff

[673] *Sextonia rubra* (Mez) van der Werff

Novon 7 (4): 438 [“1997” publ. 1998] (van der Werff 1998). — *Ocotea rubra* Mez, *Jahrb. Königl. Bot. Gart. Berlin* 5: 258 (Mez 1889). — *Nectandra rubra* (Mez) C.K.Allen, *Mem. New York Bot. Gard.* 10 (5): 120 (Allen 1964).

NOTE. — Several Mélinon specimens of *Sextonia rubra* that are labelled isotypes at P are dated 1862 (P00711138, P00711139), 1864 (P00711130, P00711134, P00711135, P00711136) or 1865 (P00711137), therefore are probably not from the same gathering as the specimen labelled as holotype (P00711131), which is dated 1863. But the fact that one of them (P00711137) bears the same label in Mez’s hand as the “holotype” implies that the original material is composed of at least two distinct gatherings (from 1863 and 1865, respectively), making it necessary to designate a lectotype.

VERNACULAR NAMES. — Pa: á-griyō, griyō • Ka: wana, wonu • Te: atsemá • Wp: asemá • Wn: wana • Nt: wana • Cr: grignon-rouj • Fr: grignon franc • Br: louro-gamela, louro-vermelho.

HERBARIUM DATA (FG). — 108 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.*, 1863 (lecto-, P[P00711131], here designated; isolecto-, B[B100244360, B100244361], P[P00711132, P00711133]).

INVENTORY DATA (FG). — 378 trees in 134 plots; $F_{\max} = 1.5\%$; $dbh_{inv} = 135$ cm.

[674] Lauraceae sp. A

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P. Grenand* et al. 3521.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 17.2\text{ cm}$.

[675] Lauraceae sp. B

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2708*.

INVENTORY DATA (FG). — 13 trees in 6 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 35.7\text{ cm}$.

[676] Lauraceae sp. C

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5727*.

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42.5\text{ cm}$.

[677] Lauraceae sp. D

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2751*.

INVENTORY DATA (FG). — 1 tree, $dbh = 42.5\text{ cm}$.

[678] Lauraceae sp. E

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2869*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 11.6\text{ cm}$.

[679] Lauraceae sp. F

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 3366*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.8\text{ cm}$.

[680] Lauraceae sp. G

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & M.-F. Prévost 1973*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.7\text{ cm}$.

[681] Lauraceae sp. H

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 3423*.

INVENTORY DATA (FG). — 39 trees in 5 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 35\text{ cm}$.

[682] Lauraceae sp. I

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2319*.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24\text{ cm}$.

[683] Lauraceae sp. J

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5072*.

INVENTORY DATA (FG). — 19 trees in 5 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 105\text{ cm}$.

Family LECYTHIDACEAE A.Rich.
Genus *Bertholletia* Bonpl.

[684] *Bertholletia excelsa* Bonpl.

Pl. Aequinoct. [Humboldt & Bonpland] 1 (5): 122 [Apr. 1807] (Bonpland 1807).

Barthollesia excelsa Silva Manso, *Enum. Subst. Braz.*: 45 (Silva Manso 1836), orth. var.

Bertholletia nobilis Miers, *Trans. Linn. Soc. London* 30 (2): 197 (Miers 1874).

NOTE. — Probably not native to French Guiana, where it is mainly found as a cultivated tree. However, three wild individuals collected near pre-Columbian archaeological sites are indicative of naturalization.

VERNACULAR NAMES. — Ka: totoka, tutukaa • Te: titikö uhu • Wp: yá • Wn: tutukë • Nt: ingi noto • Cr: touka • Fr: noix de Pará, noix du Brésil • Br: castanha-do-Pará (fruit), castanheira (tree).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *B. Dutrèze 569*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.3\text{ cm}$.

Genus *Corythophora* R.Knuth

[685] *Corythophora amapaensis* Pires ex S.A.Mori & Prance

Brittonia 33 (3): 365 (Mori & Prance 1981).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Te: watili tsing • Nt: lebi luabi • Br: matamatá-gameleira.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.M. Pires 48579* (holo-, IAN[IAN126599]; iso-, IAN[IAN109580], NY[00000735]).

INVENTORY DATA (FG). — 57 trees in 11 plots; $F_{\max} = 3.7\%$; $dbh_{\text{inv}} = 50.3\text{ cm}$.

[686] *Corythophora labriculata* (Eyma) S.A.Mori & Prance

Brittonia 33 (3): 365 (Mori & Prance 1981). — *Eschweilera labriculata* Eyma, *Polygon. Guttif. Lecythid. Surinam:* 75 (Eyma 1932). — *Chytroma labriculata* (Eyma) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 80 [22 Aug. 1939] (Knuth 1939).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *O. Tostain et al. 1499*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 13.1$ cm.

[687] *Corythophora rimosa* W.A.Rodrigues subsp. *rimosa*

Mem. New York Bot. Gard. 44: 31 (Rodrigues 1987).

VERNACULAR NAMES. — Pa: kwatri-puvemna • Ka: kuwateli, kwatili • Wp: yapukay sili • Nt: lebi luabi • Cr: mao-blan • Br: castanha-jacaré.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3370*.

INVENTORY DATA (FG). — 52 trees in 19 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 85$ cm.

[688] *Corythophora rimosa* subsp. *rubra* S.A.Mori

Mem. New York Bot. Gard. 44: 31 (Mori 1987).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kwatri-puvemna • Ka: kuwateli, kwatili • Cr: mao-blan • Br: castanha-jacaré.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 14746* (holo-, NY[00232379]; iso-, CAY[CAY027879], K[K000600143], M[M0146449], MG[MG136431], P[P01900037], U[U0003039], WIS[v 0255887 WIS]).

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.2$ cm.

Genus *Couratari* Aubl.

[689] *Couratari calycina* Sandwith
(Fig. 28A)

J. Bot. 78: 253 (Sandwith 1940).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: ingi pipa, ingii pipa.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2961*.

INVENTORY DATA (FG). — 86 trees in 23 plots; $F_{\max} = 2.4\%$; $\text{dbh}_{\text{inv}} = 44.2$ cm.

[690] *Couratari gloria* Sandwith

Kew Bull. 10 (3): 472 [20 Dec. 1955] (Sandwith 1955).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wn: momaj • Nt: ingi pipa, ingii pipa • Cr: mao-sigar • Fr: mahot cigare.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *S.A. Mori et al. 15690*.

INVENTORY DATA (FG). — 34 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 79.9$ cm.

[691] *Couratari guianensis* Aubl.

Hist. Pl. Guiane 2: 724 [Jun.-Dec. 1775] (Aublet 1775). — *Lecythis couratari* Spreng., *Syst. Veg. [Sprengel]* 4 (2): 208 [Jan.-June 1827] (Sprengel 1827), “*Lecythin Couratari*”, nom. illeg. superfl. (based on *Couratari guianensis*).

Couratari paraensis Mart. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 510 [1 Feb. 1858] (Berg 1858). — *Cariniana paraensis* (Mart. ex O.Berg) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 37 [22 Aug. 1939] (Knuth 1939).

Couratari panamensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 239 [24 Oct. 1929] (Standley 1929).

Couratari pulchra Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 217 [27 June 1932] (Sandwith 1932).

Couratari bragancae R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 128 [22 Aug. 1939] (Knuth 1939).

VERNACULAR NAMES. — Pa: wakukwa-rimwa • Ka: kwatili, tikalaye ulemali, watala • Te: ponó • Wp: yemi'i • Wn: okalat, ukalat • Nt: ingi pipa, ingii pipa • Cr: mao-sigar • Fr: mahot cigare • Br: tauarí-folha-grande.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM, the fruit in carpological collection only, leaf specimen excluded *BM000953836* [synonym of *Lecythis poiteauii*]); *S.A. Mori et al. 20973*.

INVENTORY DATA (FG). — 99 trees in 68 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 117$ cm.

[692] *Couratari multiflora* (Sm.) Eyma

Polygon. Guttif. Lecythid. Surinam: 60 (Eyma 1932). — *Lecythis multiflora* Sm., *Cycl. [Rees]* 20: (Lecythis no. 8) [19 Mar. 1812] (Smith 1812). — *Allantoma multiflora* (Sm.) Miers, *Trans. Linn. Soc. London* 30 (2): 291 [14 Nov. 1874] (Miers 1874).

Lecythis fagifolia Miq. ex O.Berg, *Linnaea* 27 (4): 451 [“1854” publ. Feb. 1856] (Berg 1856). — *Allantoma fagifolia* (Miq. ex O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 298 [14 Nov. 1874] (Miers 1874). — *Couratari fagifolia* (Miq. ex O.Berg) Eyma, *Polygon. Guttif. Lecythid. Surinam* 62 (Eyma 1932).

Couratari coriacea Mart. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 510 [1 Feb. 1858] (Berg 1858).

Couratari vriesii Miers, *Trans. Linn. Soc. London* 30 (2): 284 [14 Nov. 1874] (Miers 1874).

Allantoma subramosa Miers, *Trans. Linn. Soc. London* 30 (2): 292 [14 Nov. 1874] (Miers 1874), *pro parte folia exclusa*.

VERNACULAR NAMES. — Pa: irimwi • Ka: kisipyulu ulemali, ulemali • Te: tawali • Wp: akiki poa, kwata poa, tawali pilá, tawali sí • Wn: ulemali, waluwalu • Nt: ingi pipa, ingii pipa • Cr: lapo-sigar, mao-sigar • Fr: mahot cigare • Br: tauarí-amarello, tauarí-folha-miuda.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[[BM000953792](#)]).

INVENTORY DATA (FG). — 395 trees in 91 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 83.9$ cm.

[693] *Couratari oblongifolia* Ducke & R.Knuth

Pflanzenr. [Engler] IV.219a (Heft 105): 134 [22 Aug. 1939] (Ducke & Knuth 1939).

VERNACULAR NAMES. — Pa: irimwi, wakukwa-rimwa • Ka: toko • Nt: ingi pipa, ingii pipa • Cr: mao-sigar • Fr: mahot cigare • Br: tauarí.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *D. Sabatier 3573*.

INVENTORY DATA (FG). — 57 trees in 36 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 128.9$ cm.

[694] *Couratari oligantha* A.C.Sm.

Amer. J. Bot. 26 (6): 411 (Smith 1939).

HERBARIUM DATA (FG). — 1 collection at CAY: *D. Loubry 1122*.

SIZE. — Up to 30 m tall (Mori & Prance 1993).

[695] *Couratari stellata* A.C.Sm.
(Fig. 28B)

Amer. J. Bot. 26 (6): 410 (Smith 1939).

Couratari reticulata A.C.Sm., *Amer. J. Bot. 26 (6)*: 410 (Smith 1939).

VERNACULAR NAMES. — Ka: wokili kisipyulu ulemali • Wp: ka'i ka'i nu'a, yemi'i • Wn: okalatimë, tamujom, ulemali • Nt: ingi pipa, ingii pipa • Cr: mao-kaka, mao-nwé • Fr: mahot noir • Br: tauarí.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5340*.

INVENTORY DATA (FG). — 160 trees in 35 plots; $F_{\max} = 3.1\%$; $\text{dbh}_{\text{inv}} = 137$ cm.

Genus *Couroupita* Aubl.

[696] *Couroupita guianensis* Aubl.
(Fig. 28C)

Hist. Pl. Guiane 2: 708 [Jun.-Dec. 1775] (Aublet 1775). — *Lecythis bracteata* Willd., *Sp. Pl., ed. 4 2 (2)*: 1174 [Dec. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Couroupita guianensis*).

Pekea couroupita Juss. ex DC., *Prodri. [A. P de Candolle]* 3: 294 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Couroupita surinamensis Mart. ex O.Berg, *Linnaea* 27 (4): 462 [[“]1854” publ. Feb. 1856] (Berg 1856). — *Couroupita guianensis* var. *surinamensis* (Mart. ex O.Berg) Eyma, *Polygon. Guttif. Lecythid. Surinam*: 65 (Eyma 1932).

Couroupita peruviana O.Berg, *Linnaea* 31: 261 (Berg 1861-1862).

Couroupita antillana Miers, *Trans. Linn. Soc. London* 30 (2): 191 [14 Nov. 1874] (Miers 1874).

Couroupita membranacea Miers, *Trans. Linn. Soc. London* 30 (2): 194 [14 Nov. 1874] (Miers 1874).

Couroupita froesii R.Knuth, *Repert. Spec. Nov. Regni Veg.* 35: 341 (Knuth 1934).

Couroupita saintcroixiana R.Knuth, *Repert. Spec. Nov. Regni Veg.* 35: 341 (Knuth 1934), “*St. Croixiana*”.

Couroupita venezuelensis R.Knuth, *Repert. Spec. Nov. Regni Veg.* 35: 341 (Knuth 1934).

Couroupita acreensis R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 47 [22 Aug. 1939] (Knuth 1939).

Couroupita idolica Dwyer, *Ann. Missouri Bot. Gard.* 52: 358 (Dwyer 1965).

Couroupita pedicellaris Rizzini, *Rodriguésia* 28 (41): 178 (Rizzini 1976).

VERNACULAR NAMES. — Ka: kaupi, kopu • Nt: bakuu kokonoto, kaabasi bon • Cr: boulé-kannon, kouy-serjan • Fr: arbre à boulet de canons • Br: abrico-de-macaco, cuia-de-macaco, macacarecuia.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000953857](#)]); *M.-F. Prévost & D. Sabatier 4720*, $\text{dbh} = 63$ cm.

Genus *Eschweilera* Mart. ex DC.

[697] *Eschweilera alata* A.C.Sm.
(Fig. 29A)

Amer. J. Bot. 26 (6): 407 (Smith 1939).

VERNACULAR NAMES. — Ka: ara-a.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier et al. 4393*.

INVENTORY DATA (FG). — 187 trees in 25 plots; $F_{\max} = 5.8\%$; $\text{dbh}_{\text{inv}} = 58.8$ cm.

[698] *Eschweilera apiculata* (Miers) A.C.Sm.

Phytologia 1 (3): 124 [21 Jan. 1935] (Smith 1935). — *Chytroma apiculata* Miers, *Trans. Linn. Soc. London* 30 (2): 245 [14 Nov. 1874] (Miers 1874).

VERNACULAR NAMES. — Pa: avun-priye, avun-seinó • Wp: ka'i ka'i nu'a sili • Nt: luabi • Br: matamatá-jibóia.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier 1277*.

INVENTORY DATA (FG). — 70 trees in 38 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 76.1$ cm.

[699] *Eschweilera chartaceifolia* S.A.Mori

Fl. Neotrop. Monogr. 21 (2): 228 [10 Apr. 1990] (Mori 1990).

VERNACULAR NAMES. — Pa: avun-seinō, kwatri.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3797*.

INVENTORY DATA (FG). — 134 trees in 55 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 34.6\text{ cm}$.

[700] *Eschweilera collina* Eyma

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 4: 63 (Eyma 1932). — *Chytroma collina* (Eyma) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 78 [22 Aug. 1939] (Knuth 1939).

VERNACULAR NAMES. — Br: ripeiro-branco.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier 3521*.

INVENTORY DATA (FG). — 192 trees in 56 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 45.5\text{ cm}$.

[701] *Eschweilera coriacea* (DC.) Mart. ex S.A.Mori

Fl. Neotrop. Monogr. 21 (2): 203 [10 Apr. 1990] (Mori 1990). — *Lecythis coriacea* DC., *Prodr. [A. P. de Candolle]* 3: 291 [mid Mar. 1828] (Candolle 1828). — *Eschweilera coriacea* (DC.) Mart. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 491 [1 Feb. 1858] (Berg 1858), nom. nud. pro syn. — *Jugastrum coriaceum* (DC.) Miers, *Trans. Linn. Soc. London* 30 (2): 275 [14 Nov. 1874] (Miers 1874).

Eschweilera grandifolia Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 293 [mid Mar. 1828] (Candolle 1828). — *Lecythis grandifolia* (Mart. ex DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 494 [1 Feb. 1858] (Berg 1858). — *Chytroma grandifolia* (Mart. ex DC.) Miers, *Trans. Linn. Soc. London* 30 (2): 237 [14 Nov. 1874] (Miers 1874).

Lecythis odora Poepp. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 492 [1 Feb. 1858] (Berg 1858). — *Eschweilera odora* (Poepp. ex O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 273 [14 Nov. 1874] (Miers 1874).

Lecythis acuminatissima O.Berg, *Fl. Bras. [Martius]* 14 (1): 494 [1 Feb. 1858] (Berg 1858). — *Eschweilera acuminatissima* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 273 [14 Nov. 1874] (Miers 1874).

Chytroma cincturata Miers, *Trans. Linn. Soc. London* 30 (2): 237 [14 Nov. 1874] (Miers 1874).

Eschweilera pallida Miers, *Trans. Linn. Soc. London* 30 (2): 267 [14 Nov. 1874] (Miers 1874).

Eschweilera vageleri R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 96 [22 Aug. 1939] (Knuth 1939).

Eschweilera eymaana R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 100 [22 Aug. 1939] (Knuth 1939).

Eschweilera fractiflexa R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 110 [22 Aug. 1939] (Knuth 1939).

Lecythis retroflexa Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 595 (Benoist 1923). — *Eschweilera retroflexa* (Benoist) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 106 [22 Aug. 1939] (Knuth 1939).

Neohuberia matamata Ledoux, *Lecointea* 1: 3 (Ledoux 1963). — *Eschweilera matamata* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 211 (Huber 1910), nom. nud.

Lecythis peruviana L.O.Williams, *Fieldiana, Bot.* 31 (1-2): 30 [25 Nov. 1964] (Williams 1964).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-priye • Ka: kwatili tabulu, wipitano kuwateli • Te: watili tsing • Wp: walime i • Wn: pukut • Nt: baaka aki • Cr: mao-nwé • Fr: mahot noir • Br: matamatá-preto, matamatá-verdadeira, ripeiro.

HERBARIUM DATA (FG). — 239 collections at CAY. Sel. exs.: *P.A. Sagot 269*, 1854 (lectotype of *Eschweilera pallida*: BM[[BM000795573](#)], designated by Mori *et al.* [1990: 203]; isolecto-, S[[S09-27917](#)]).

INVENTORY DATA (FG). — 1787 trees in 145 plots; $F_{\max} = 9\%$; $\text{dbh}_{\text{inv}} = 88.5\text{ cm}$.

[702] *Eschweilera decolorans* Sandwith
(Fig. 29B)

Bull. Misc. Inform. Kew 1932 (5): 214 [27 June 1932] (Sandwith 1932). — *Chytroma decolorans* (Sandwith) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 79 [22 Aug. 1939] (Knuth 1939).

Eschweilera sandwithiana A.C.Sm. & Beard, *J. Arnold Arbor.* 27 (3): 310 [15 July 1946] (Smith & Beard 1946).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: ulana elepali • Nt: baaka luabi • Br: ripeiro.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1725*.

INVENTORY DATA (FG). — 386 trees in 90 plots; $F_{\max} = 4.7\%$; $\text{dbh}_{\text{inv}} = 69.6\text{ cm}$.

[703] *Eschweilera grandiflora* (Aubl.) Sandwith
(Fig. 28D)

Kew Bull. 10 (3): 472 [20 Dec. 1955] (Sandwith 1955). — *Lecythis grandiflora* Aubl., *Hist. Pl. Guiane* 2: 712 [Jun.-Dec. 1775] (Aublet 1775).

Eschweilera alba R.Knuth, *Repert. Spec. Nov. Regni Veg.* 38: 115 (Knuth 1935).

Eschweilera fracta R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 110 [22 Aug. 1939] (Knuth 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-purubumna • Wp: mani'i • Nt: luabi • Br: castanhara, matamatá-rósea.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[[BM000953843](#)]) designated by Mori *et al.* [1990: 216].

INVENTORY DATA (FG). — 146 trees in 47 plots; $F_{\max} = 2.3\%$; $\text{dbh}_{\text{inv}} = 55.5\text{ cm}$.



FIG. 28. — Lecythidaceae: **A**, *Couratari calycina* Sandwith; **B**, *Couratari stellata* A.C.Sm. (D. Sabatier & J.-F. Molino 5340); **C**, *Couroupita guianensis* Aubl.; **D**, *Eschweilera grandiflora* (Aubl.) Sandwith (M.-F. Prévost & D. Sabatier 4826). © D. Sabatier/IRD.

[704] *Eschweilera laevicarpa* S.A.Mori

Mem. New York Bot. Gard. 44: 32 (Mori 1987).

VERNACULAR NAMES. — Nt: luabi • Br: matamatá-azulada.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3391*.

INVENTORY DATA (FG). — 18 trees in 14 plots; $F_{\max} < 1\%$; dbh_{inv} = 43.3 cm.

[705] *Eschweilera micrantha* (O.Berg) Miers

Trans. Linn. Soc. London 30 (2): 260 [14 Nov. 1874] (Miers 1874). — *Lecythis micrantha* O.Berg, *Linnaea* 27 (4): 454 ["1854" publ. Feb. 1856] (Berg 1856).

Lecythis gracilipes Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20:* 203 (Sagot 1885). — *Eschweilera gracilipes* (Sagot) R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 101 [22 Aug. 1939] (Knuth 1939).

Eschweilera floribunda Eyma, *Polygon. Guttif. Lecythid. Surinam* 74 (Eyma 1932).

Eschweilera polyantha A.C.Sm., *Bull. Torrey Bot. Club* 60 (6): 381 [1 June 1933] (Smith 1933).

NOTES. — Mori et al. (1990: 262) designated "Mélinon 18" as lectotype of *Lecythis gracilipes* Sagot. It is actually an unnumbered collection of Mélinon. "18" is not a collection number, but the beginning of the collecting year (1865), printed in advance on some 18th century labels at P. The last two digits have been added by hand only on some duplicates.

VERNACULAR NAMES. — Pa: avun-priye, avun-priyo • Ka: kuwateli, kwatili • Wp: ayá leme sili • Wn: ponoimé • Cr: mao-nwé • Br: matamatá.

HERBARIUM DATA (FG). — 110 collections at CAY. Sel. exs.: *E.M. Melinon s.n.* (lectotype of *Lecythis gracilipes*, P[P01900083] designated by Mori *et al.* [1990: 262]; isolecto-, F[V0044332F], G[G00369113], K[K000600250], MPU[MPU011047], P[P01900084, P01900085, P01900086, P01900087, P01900088, P01900089, P01900090, P01900091], US[00117432]).

INVENTORY DATA (FG). — 1310 trees in 111 plots; $F_{\max} = 10.4\%$; $\text{dbh}_{\text{inv}} = 117.1$ cm.

[706] *Eschweilera parviflora* (Aubl.) Miers

Trans. Linn. Soc. London 30 (2): 260 [14 Nov. 1874] (Miers 1874). — *Lecythis parviflora* Aubl., *Hist. Pl. Guyane* 2: 717 [Jun.-Dec. 1775] (Aublet 1775).

Eschweilera grata Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 216 [27 June 1932] (Sandwith 1932).

Eschweilera montana A.C.Sm., *Amer. J. Bot.* 26 (6): 407 (Smith 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-seinō • Ka: kwatili, tamunen kuwateli • Wp: ayá leme, ayá leme sili • Nt: baaka aki • Cr: mao-blán, mao-nwé • Fr: mahot noir • Br: matamatá-branco, matamatá-roxa.

HERBARIUM DATA (FG). — 169 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (lectotype of *Lecythis platycarpa*: K[K000600210] designated by Mori *et al.* [1990: 243]).

INVENTORY DATA (FG). — 222 trees in 74 plots; $F_{\max} = 4.5\%$; $\text{dbh}_{\text{inv}} = 28.7$ cm.

[707] *Eschweilera pedicellata* (Rich.) S.A.Mori
(Fig. 29C)

Mem. New York Bot. Gard. 44: 34 (Mori 1987). — *Lecythis pedicellata* Rich., *Actes Soc. Hist. Nat. Paris* 1: 111 [Oct. 1792] (Richard 1792).

Lecythis longipes Poit., *Mém. Mus. Hist. Nat.* 13: 144 (Poiteau 1825). — *Eschweilera longipes* (Poit.) Miers, *Trans. Linn. Soc. London* 30 (2): 253 [14 Nov. 1874] (Miers 1874).

Lecythis platycarpa Poit., *Mém. Mus. Hist. Nat.* 13: 146 (Poiteau 1825), “*platicarpa*”. — *Lecythis longipes* var. *platycarpa* (Poit.) DC., *Prodri. [A. P. de Candolle]* 3: 292 [mid Mar. 1828] (Candolle 1828). — *Eschweilera platycarpa* (Poit.) Miers, *Trans. Linn. Soc. London* 30 (2): 258 [14 Nov. 1874] (Miers 1874). — *Lecythis longipes* f. *platycarpa* (Poit.) Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 20: 205 (Sagot 1885). — *Eschweilera longipes* f. *platycarpa* (Poit.) R.Knuth, *Pflanzren. [Engler]* IV.219a (Heft 105): 101 [22 Aug. 1939] (Knuth 1939).

Lecythis wulsschlaegeliana O.Berg, *Fl. Bras. [Martius]* 14 (1): 493 [1 Feb. 1858] (Berg 1858).

Lecythis idatimonoides O.Berg, *Fl. Bras. [Martius]* 14 (1): 496 [1 Feb. 1858] (Berg 1858). — *Eschweilera idatimonoides* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 252 [14 Nov. 1874] (Miers 1874).

Lecythis pilosa Poepp. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 500 [1 Feb. 1858] (Berg 1858). — *Eschweilera pilosa* (Poepp. ex O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 274 [14 Nov. 1874] (Miers 1874).

Lecythis macrophylla O.Berg, *Fl. Bras. [Martius]* 14 (1): 616 [15 Jan. 1859] (Berg 1859). — *Eschweilera macrophylla* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 263 [14 Nov. 1874] (Miers 1874).

Lecythis basizone O.Berg, *Fl. Bras. [Martius]* 14 (1): 617 [15 Jan. 1859] (Berg 1859).

Lecythis longipes f. *genuina* Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 20: 205 (Sagot 1885), nom. inval. (Turland *et al.* 2018): Art. 24.3. — *Eschweilera longipes* f. *genuina* (Sagot) R.Knuth, *Pflanzren. [Engler]* IV.219a (Heft 105): 104 [22 Aug. 1939] (Knuth 1939), nom. inval. (Turland *et al.* 2018): Art. 24.3.

Chytroma perspicua Miers, *Trans. Linn. Soc. London* 30 (2): 246 [14 Nov. 1874] (Miers 1874).

Eschweilera flaccida Miers, *Trans. Linn. Soc. London* 30 (2): 264 [14 Nov. 1874] (Miers 1874).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-seinō • Ka: kwatili, tamunen kuwateli • Wp: ayá leme, ayá leme sili • Nt: baaka aki • Cr: mao-blán, mao-nwé • Fr: mahot noir • Br: matamatá-branco, matamatá-roxa.

HERBARIUM DATA (FG). — 169 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (lectotype of *Lecythis platycarpa*: K[K000600210] designated by Mori *et al.* [1990: 243]).

INVENTORY DATA (FG). — 222 trees in 74 plots; $F_{\max} = 4.5\%$; $\text{dbh}_{\text{inv}} = 28.7$ cm.

[708] *Eschweilera piresii* S.A.Mori
subsp. *viridipetala* S.A.Mori

Mem. New York Bot. Gard. 64: 229 (Mori 1990).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 20800* (holo-, NY[00000819]; iso-, CAY[CAY027880]).

INVENTORY DATA (FG). — 6 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 17.8$ cm.

[709] *Eschweilera sagotiana* Miers

Trans. Linn. Soc. London 30 (2): 262 [14 Nov. 1874] (Miers 1874).

Lecythis parviflora Sagot ex Miers, *Trans. Linn. Soc. London* 30 (2): 262 [14 Nov. 1874] (Miers 1874), nom. nud. pro syn., nom. illeg. hom., non Aubl. (Aublet 1775).

Lecythis melinonii Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 20: 202 (Sagot 1885), “*Melinonis*”. — *Eschweilera melinonii* (Sagot) R.Knuth, *Pflanzren. [Engler]* IV.219a (Heft 105): 104 [22 Aug. 1939] (Knuth 1939), “*Melinonis*”.

Lecythis jucunda Benoist, *Notul. Syst. (Paris)* 3: 180 [25 Dec. 1915] (Benoist 1915). — *Eschweilera jucunda* (Benoist) R.Knuth, *Pflanzren. [Engler]* IV.219a (Heft 105): 102 [22 Aug. 1939] (Knuth 1939).

Lecythis tapuya Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 594 (Benoist 1923). — *Eschweilera tapuya* (Benoist) R.Knuth, *Pflanzren. [Engler]* IV.219a (Heft 105): 106 [22 Aug. 1939] (Knuth 1939).

Lecythis sagotiana Benoist, *Arch. Bot.* 5 (Mém. 1): 214 [27 Mar. 1933] (Benoist 1933), in clavi, nom. subnud.

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). The epithet “*melinonii*” (in *Lecythis* and *Eschweilera*) is to be automatically corrected to “*melinonii*” (Turland *et al.* 2018: Art. 60.8). *L. sagotiana* Benoist is apparently not based on *E. sagotiana* Miers.



FIG. 29. — Lecythidaceae: **A**, *Eschweilera alata* A.C.Sm. (M.-F. Prévost & D. Sabatier 4615); **B**, *Eschweilera decolorans* Sandwith (M.-F. Prévost 4214); **C**, *Eschweilera pedicellata* (Rich.) S.A.Mori (M.-F. Prévost 4257); **D**, *Lecythis persistens* Sagot subsp. *aurantiaca* S.A.Mori (D. Sabatier et al. 4404). A-C, © M.-F. Prévost/IRD; D, © J.-F. Molino/IRD.

VERNACULAR NAMES. — Pa: avun-duwē, avun-seinō • Ka: kuwateli, ulana elepali • Nt: baaka aki, luabi.

HERBARIUM DATA (FG). — 92 collections at CAY. Sel. exs.: *P.A. Sagot 1104*, 1858 (lecto-, BM[B000953880], designated by Mori et al. [1990: 263], isolecto-, BR[BR0000005175567], F[V0044327F, V0061898F], G[G00369047], GH[GH00075280], K[K000600236, K000600237, K000600238], P[P01900102, P01900103, P01900104, P01900105, P01900106], S[S04-1253], U[U00079776]).

INVENTORY DATA (FG). — 2806 trees in 175 plots; $F_{\max} = 16\%$; $dbh_{inv} = 154$ cm.

^[710] *Eschweilera squamata* S.A.Mori

Mem. New York Bot. Gard. 44: 33 (Mori 1987).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kwatri-washiune-priye • Nt: baaka aki.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15288* (holo-, NY[00000832]; iso-, CAY[CAY027882], K[K000600218], MG[MG126668], MO[MO-313122], NY[00000831], P[P01900115], U[U0003095]).

INVENTORY DATA (FG). — 42 trees in 19 plots; $F_{\max} = 1.4\%$; $dbh_{inv} = 77.3$ cm.

^[711] *Eschweilera subglandulosa* (Steud. ex O.Berg) Miers

Trans. Linn. Soc. London 30 (2): 266 [14 Nov. 1874] (Miers 1874). — *Lecythis subglandulosa* Steud. ex O.Berg, *Linnaea* 27 (4): 459 ["1854" publ. Feb. 1856] (Berg 1856).

Lecythis laevifolia Griseb., *Fl. Brit. W.I. [Grisebach]* 711 [Oct. 1864] (Grisebach 1864), *nom. nud.*

Eschweilera laevifolia Miers, *Trans. Linn. Soc. London* 30 (2): 256 [14 Nov. 1874] (Miers 1874).

VERNACULAR NAMES. — Wp: siliwi, sulu iwi.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier* 996.

SIZE. — Up to 35 m tall (Mori & Prance 1993).

[712] *Eschweilera wachenheimii* (Benoist) Sandwith

Bull. Misc. Inform. Kew 1932 (5): 215 [27 June 1932] (Sandwith 1932). — *Lecythis wachenheimii* Benoist, *Bull. Mus. Natl. Hist. Nat.* 29: 594 (Benoist 1923).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: kuwateli, kватили • Br: matamatá-mirim.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *G. Wachenheim* 359 (lecto-, P[P02284923, P02284924, P02284925], designated by Mori *et al.* [1990: 230]).

INVENTORY DATA (FG). — 92 trees in 34 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 36.9\text{ cm}$.

[713] *Eschweilera* sp. A

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier* 4861.

INVENTORY DATA (FG). — 41 trees in 16 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 31.2\text{ cm}$.

[714] *Eschweilera* sp. B

NOTE. — A species close to *Eschweilera grandiflora* (Aubl.) Sandwith.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier* & J.-F. Molino 4945.

INVENTORY DATA (FG). — 56 trees in 17 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 115\text{ cm}$.

Genus *Gustavia* L.

[715] *Gustavia augusta* L.

Pl. Surin.: 17 [23 June 1775] (Linnaeus 1775). — *Gustavia angusta* J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (2): 1038 ["1791" publ. late Apr.-Oct. 1792] (Gmelin 1792), sphalm. — *Japarandiba augusta* (L.) Kuntze, *Revis. Gen. Pl.* 1: 240 [5 Nov. 1891] (Kuntze 1891).

Pirigara tetrapetala Aubl., *Hist. Pl. Guiane* 1: 487 [Jun.-Dec. 1775] (Aublet 1775). — *Gustavia tetrapetala* (Aubl.) DC., *Prodr. [A. P. de Candolle]* 3: 290 [mid Mar. 1828] (Candolle 1828). — *Grias tetrapetala* (Aubl.) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 37 [Feb. 1892] (Niedenzu 1892).

Gustavia meizocarpa Gaertn., *Fruct. Sem. Pl.* 2: 264 [Apr.-May 1791] (Gaertner 1791).

Gustavia urceolata Poit., *Mém. Mus. Hist. Nat.* 13: 156 (Poiteau 1825).

Gustavia augusta var. *guianensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 469 [1 Feb. 1858] (Berg 1858).

Gustavia augusta var. *brasiliensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 471 [1 Feb. 1858] (Berg 1858).

Gustavia augusta var. *calycaris* O.Berg, *Fl. Bras. [Martius]* 14 (1): 471 [1 Feb. 1858] (Berg 1858). — *Gustavia calycaris* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 185 [14 Nov. 1874] (Miers 1874).

Gustavia augusta var. *verrucosa* Mart. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 471 [1 Feb. 1858] (Berg 1858).

Gustavia insignis Willd. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 471 [1 Feb. 1858] (Berg 1858), *nom. nud. pro syn.*

Gustavia poeppigiana O.Berg var. *rigida* O.Berg, *Fl. Bras. [Martius]* 14 (1): 471 [1 Feb. 1858] (Berg 1858).

Gustavia membrillo Appun, *Bot. Sam. Venez.* 34 (Appun 1858).

Gustavia insignis Linden ex Hook., *Bot. Mag.* 84: t. 5069 (Hooker 1858).

Gustavia brasiliensis E.Morren, *Ann. Hort. Belge Étrangère* 16: 201 (Morren 1866).

Gustavia theophrasta Linden, *Ill. Hort.* 20: 70 (Linden 1873).

Gustavia laciniosa Miers, *Trans. Linn. Soc. London* 30 (2): 178 [14 Nov. 1874] (Miers 1874).

Gustavia antillana Miers, *Trans. Linn. Soc. London* 30 (2): 179 [14 Nov. 1874] (Miers 1874). — *Japarandiba antillana* (Miers) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 37 [Feb. 1892] (Niedenzu 1892).

Gustavia marcgravia Miers, *Trans. Linn. Soc. London* 30 (2): 183 [14 Nov. 1874] (Miers 1874). — *Japarandiba marcgravia* (Miers) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 37 [Feb. 1892] (Niedenzu 1892).

Gustavia mexicana R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 136 [22 Aug. 1939] (Knuth 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wakukwa-adava • Ka: akuli elepali, alepawana, pirigamepé (*fide* Aublet 1775) • Te: tapakadja'i • Wp: a'i walipi • Wn: pisusukimë • Nt: man tapupa • Cr: bwa-pian • Br: castanha-fedorenta, jenipaporana, mucurão.

HERBARIUM DATA (FG). — 103 collections at CAY. Sel. exs.: *S.A. Mori* *et al.* 25667.

INVENTORY DATA (FG). — 81 trees in 29 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 21.3\text{ cm}$.

[716] *Gustavia hexapetala* (Aubl.) Sm.

Cycl. [Rees] 17: (Gustavia no. 2) [8 Mar. 1811] (Smith 1811). — *Pirigara hexapetala* Aubl., *Hist. Pl. Guiane* 1: 490 [Jun.-Dec. 1775] (Aublet 1775). — *Gustavia fastuosa* Willd., *Sp. Pl.*, ed. 4 3 (1): 847 (Willdenow 1800), *nom. illeg. superfl.* (based on *Pirigara hexapetala*). — *Japarandiba fastuosa* Kuntze, *Revis. Gen. Pl.* 1: 240 [5 Nov.

1891] (Kuntze 1891). — *Japarandiba hexapetala* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 240 [5 Nov. 1891] (Kuntze 1891).

Gustavia pterocarpa Poit., *Mém. Mus. Hist. Nat.* 13: 158 (Poiteau 1825). — *Japarandiba pterocarpa* (Poit.) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 37 [Feb. 1892] (Niedenzu 1892).

Gustavia brasiliiana DC., *Prod. [A. P. de Candolle]* 3: 290 [mid Mar. 1828] (Candolle 1828). — *Japarandiba brasiliiana* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 240 [5 Nov. 1891] (Kuntze 1891).

Gustavia brasiliiana var. *minor* DC., *Prod. [A. P. de Candolle]* 3: 290 [mid Mar. 1828] (Candolle 1828).

Gustavia fastuosa var. *angustisepala* O.Berg, *Fl. Bras. [Martius]* 14 (1): 473 [1 Feb. 1858] (Berg 1858).

Gustavia fastuosa var. *latisepala* O.Berg, *Fl. Bras. [Martius]* 14 (1): 473 [1 Feb. 1858] (Berg 1858).

Gustavia microcarpa Pilg., *Verb. Bot. Vereins Prov. Brandenburg* 47: 164 [1 Oct. 1905] (Pilger 1905).

Gustavia eximia Pittier, *Arb. Arbust. Venez.* 1: 9 (Pittier 1921).

Gustavia fustis-mortui Pittier, *Bol. Ci. Técn. Mus. Comercial Venezuela* 1: 64 (Pittier 1925).

Eschweilera carrii Standl., *Trop. Woods* 29: 7 (Standley 1932).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *Japarandiba fastuosa* Kuntze is based on the illegitimate *Gustavia fastuosa* Willd., hence authorship is not "(Willd.) Kuntze".

VERNACULAR NAMES. — Pa: wakukwa-adava-kamwi • Ka: alepawana enekan • Wp: a'i walipi sili • Nt: man tapupa • Cr: mal-bwa-pian • Br: jenipaporana, matamataica.

HERBARIUM DATA (FG). — 103 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000583750]).

INVENTORY DATA (FG). — 1082 trees in 177 plots; $F_{\max} = 6.3\%$; $dbh_{\text{inv}} = 37.6$ cm.

Genus *Lecythis* Loefl.

[717] *Lecythis chartacea* O.Berg

Linnaea 27 (4): 450 ["1854" publ. Feb. 1856] (Berg 1856). — *Chytroma chartacea* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 231 [14 Nov. 1874] (Miers 1874). — *Eschweilera chartacea* (O.Berg) Eyma, *Polygon. Guttif. Lecythid. Surinam* 70 (Eyma 1932).

Lecythis monosperma Mart. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 485 [1 Feb. 1858] (Berg 1858). — *Chytroma monosperma* (Mart. ex O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 239 [14 Nov. 1874] (Miers 1874).

Lecythis spruceana O.Berg, *Fl. Bras. [Martius]* 14 (1): 487 [1 Feb. 1858] (Berg 1858). — *Eschweilera spruceana* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 40 [Feb. 1892] (Niedenzu 1892). — *Chytroma spruceana* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 235 [14 Nov. 1874] (Miers 1874).

Lecythis rorida O.Berg, *Fl. Bras. [Martius]* 14 (1): 488 [1 Feb. 1858] (Berg 1858). — *Chytroma rorida* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 243 [14 Nov. 1874] (Miers 1874). — *Eschweilera rorida* (O.Berg) A.C.Sm., *Phytologia* 1 (3): 124 [21 Jan. 1935] (Smith 1935).

Lecythis marauynensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 489 [1 Feb. 1858] (Berg 1858). — *Chytroma marauynensis* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 245 [14 Nov. 1874] (Miers 1874). — *Eschweilera marauynensis* (O.Berg) Nied., *Enum. Vasc. Pl. Surinam* 332 (Niedenzu 1906).

Lecythis langsdorffii O.Berg, *Fl. Bras. [Martius]* 14 (1): 615 [15 Jan. 1859] (Berg 1859). — *Chytroma langsdorffii* (O.Berg) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 87 [22 Aug. 1939] (Knuth 1939).

Lecythis cupularis Miers, *Trans. Linn. Soc. London* 30 (2): 224 [14 Nov. 1874] (Miers 1874). — *Chytroma cupularis* (Miers) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 85 [22 Aug. 1939] (Knuth 1939).

Lecythis laevicula Miers, *Trans. Linn. Soc. London* 30 (2): 224 [14 Nov. 1874] (Miers 1874). — *Chytroma laevicula* (Miers) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 80 [22 Aug. 1939] (Knuth 1939).

Chytroma cistella Miers, *Trans. Linn. Soc. London* 30 (2): 233 [14 Nov. 1874] (Miers 1874).

Chytroma urceolata Miers, *Trans. Linn. Soc. London* 30 (2): 237 [14 Nov. 1874] (Miers 1874).

Chytroma pilacarpa Miers, *Trans. Linn. Soc. London* 30 (2): 238 [14 Nov. 1874] (Miers 1874).

Eschweilera rigida Miers, *Trans. Linn. Soc. London* 30 (2): 249 [14 Nov. 1874] (Miers 1874). — *Lecythis rigida* (Miers) R.Knuth, *Repert. Spec. Nov. Regni Veg. Beih.* 43: 512 (Knuth 1927).

Cercophora anomala Miers, *Trans. Linn. Soc. London* 30 (2): 301 [14 Nov. 1874] (Miers 1874).

Eschweilera brancoensis R.Knuth, *Repert. Spec. Nov. Regni Veg.* 38: 115 (Knuth 1935).

NOTES. — Mori (Mori *et al.* 1990: 321) listed a name "*Lecythis chartacea* Berg. var. *calyce* Sagot" that has no botanical standing. Indeed, Sagot (1885: 203) wrote "*Lecythis chartacea* Berg., var. *calyce* et *axi racemi griseo-puberulis*; (*L. marauynensis* Berg, tab Flor. Bras.)". Therefore, "*calyce*" is not an epithet, but the first word of a descriptive sentence. Sagot's intention was apparently to make *L. marauynensis* ("marauynensis") Berg a variety of *L. chartacea*, but he did not formally do so.

VERNACULAR NAMES. — Ka: toko • Wp: ka'i ka'i nu'a • Cr: mao-blán, mao-kwátarí, mao-pílon • Br: cachimbo, jaraná-mirim.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *S.A. Mori* *et al.* 15702.

INVENTORY DATA (FG). — 52 trees in 26 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 94$ cm.

[718] *Lecythis confertiflora* (A.C.Sm.) S.A.Mori

Mem. New York Bot. Gard. 44: 31 (Mori 1987). — *Eschweilera confertiflora* A.C.Sm., *Amer. J. Bot.* 26 (6): 409 (Smith 1939).

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *D. Sabatier* 2529.

INVENTORY DATA (FG). — 191 trees in 35 plots; $F_{\max} = 3.8\%$; $dbh_{\text{inv}} = 54.1$ cm.

[719] *Lecythis congestiflora* Benoist

Notul. Syst. (Paris) 3: 177 [25 Dec. 1915] (Benoist 1915). — *Eschweilera congestiflora* (Benoist) Eyma, *Polygon. Guttif. Lecythid. Surinam:* 71 (Eyma 1932). — *Chytroma congestiflora* (Benoist) R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 78 [22 Aug. 1939] (Knuth 1939).

NOTES. — Known only from the Guiana Shield. According to Mori *et al.* (2017), this species is better placed in *Lecythis* than in *Eschweilera*.

VERNACULAR NAMES. — Ka: kuwateli, kwatili • Nt: baaka aki.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *R. Benoist* 578 (lecto-, P[P01900098], designated by Mori *et al.* [1990: 213]; isolecto-, P[P02284914]).

INVENTORY DATA (FG). — 332 trees in 80 plots; $F_{\max} = 4.1\%$; $dbh_{\text{inv}} = 71.3$ cm.

[720] *Lecythis corrugata* Poit.

Mém. Mus. Hist. Nat. 13: 146 (Poiteau 1825). — *Eschweilera corrugata* (Poit.) Miers, *Trans. Linn. Soc. London 30 (2):* 253 [14 Nov. 1874] (Miers 1874). — *Chytroma corrugata* (Poit.) R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 78 [22 Aug. 1939] (Knuth 1939).

Lecythis salebrosa O.Berg, *Fl. Bras. [Martius] 14 (1):* 488 [1 Feb. 1858] (Berg 1858). — *Chytroma salebrosa* (O.Berg) Miers, *Trans. Linn. Soc. London 30 (2):* 240 [14 Nov. 1874] (Miers 1874). — *Eschweilera salebrosa* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl] 3 (7):* 40 [Feb. 1892] (Niedenzu 1892).

Lecythis martinii O.Berg, *Fl. Bras. [Martius] 14 (1):* 616 [15 Jan. 1859] (Berg 1859), “martini”.

Lecythis venusta Miers, *Trans. Linn. Soc. London 30 (2):* 214 [14 Nov. 1874] (Miers 1874).

Lecythis cognata Miers, *Trans. Linn. Soc. London 30 (2):* 215 [14 Nov. 1874] (Miers 1874).

Lecythis rubicunda Miers, *Trans. Linn. Soc. London 30 (2):* 226 [14 Nov. 1874] (Miers 1874).

Chytroma basilaris Miers, *Trans. Linn. Soc. London 30 (2):* 239 [14 Nov. 1874] (Miers 1874).

Chytroma rubriflora Miers, *Trans. Linn. Soc. London 30 (2):* 241 [14 Nov. 1874] (Miers 1874).

Cymbopetalum venosum Gleason, *Bull. Torrey Bot. Club 54 (8):* 607 [Nov. 1927] (Gleason 1927), *pro parte quoad folia tantum*.

Eschweilera jenmanii R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 102 [22 Aug. 1939] (Knuth 1939).

Eschweilera patrisii R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 105 [22 Aug. 1939] (Knuth 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-priye, avun-priyo • Ka: kuwateli, kwatili, kwatili tamunen, tapilen kuwatili • Wp: talaiwi • Wn: pukut • Nt: kwanda • Cr: mao-rouj • Br: ripeiro.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *PA. Poiteau s.n.* (Lectotype K[K000600133] designated by Mori [Mori *et al.* 1990: 277]).

INVENTORY DATA (FG). — 293 trees in 55 plots; $F_{\max} = 5.2\%$; $dbh_{\text{inv}} = 76$ cm.

[721] *Lecythis holcogyne* (Sandwith) S.A.Mori

Brittonia 33 (3): 363 (Mori 1981). — *Eschweilera holcogyne* Sandwith, *Bull. Misc. Inform. Kew 1935 (3):* 126 [20 May 1935] (Sandwith 1935). — *Chytroma holcogyne* (Sandwith) R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 79 [22 Aug. 1939] (Knuth 1939).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: avun-duwē, avun-seine, kwatri-seinō • Nt: luabi.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4177*.

INVENTORY DATA (FG). — 155 trees in 65 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 90$ cm.

[722] *Lecythis idatimon* Aubl.

Hist. Pl. Guiane 2: 721 [Jun.-Dec. 1775] (Aublet 1775). — *Chytroma idatimon* (Aubl.) Miers, *Trans. Linn. Soc. London 30 (2):* 239 [14 Nov. 1874] (Miers 1874). — *Eschweilera idatimon* (Aubl.) Nied., *Nat. Pflanzenfam. [Engler & Prantl] 3 (7):* 40 [Feb. 1892] (Niedenzu 1892).

Lecythis amara Aubl., *Hist. Pl. Guiane 2:* 716 [Jun.-Dec. 1775] (Aublet 1775), *pro parte quoad folia tantum*. — *Chytroma amara* (Aubl.) Miers, *Trans. Linn. Soc. London 30 (2):* 231 [14 Nov. 1874] (Miers 1874). — *Eschweilera amara* (Aubl.) Nied., *Nat. Pflanzenfam. [Engler & Prantl] 3 (7):* 40 [Feb. 1892] (Niedenzu 1892).

Lecythis lutea Aubl., *Hist. Pl. Guiane 2:* 721 [Jun.-Dec. 1775] (Aublet 1775). — *Eschweilera lutea* (Aubl.) Miers, *Trans. Linn. Soc. London 30 (2):* 254 [14 Nov. 1874] (Miers 1874).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avun-duwō • Ka: toko • Te: watili • Wp: siliwi, sulu iwi, talaiwi sī • Wn: pukut • Nt: lebi luabi, luabi • Cr: mao-rouj • Fr: mahot rouge • Br: jatereu, matamatá.

HERBARIUM DATA (FG). — 123 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (Lectotype BM[BM000953839] designated by Mori [1990: 282]; isolecto-, BM[BM000953840], S[S04-1247]).

INVENTORY DATA (FG). — 1900 trees in 122 plots; $F_{\max} = 19.4\%$; $dbh_{\text{inv}} = 120$ cm.

[723] *Lecythis persistens* Sagot subsp. *persistens*

Ann. Sci. Nat., Bot. sér. 6, 20: 201 (Sagot 1885).

Eschweilera ovalis R.Knuth, *Pflanzenr. [Engler] IV.219a* (Heft 105): 104 [22 Aug. 1939] (Knuth 1939).

Lecythis alba Aubl. ex S.A.Mori, *Brittonia 30:* 120 (Mori 1978), *nom. nud.*

VERNACULAR NAMES. — Pa: avun-seine, kwatri-washiune-seinō • Ka: kwatili tabulu • Te: watili tsing • Wp: talaiwi sī • Nt: lebi luabi • Cr: mao-rouj • Fr: mahot rouge • Br: matamatá.

HERBARIUM DATA (FG). — 156 collections at CAY. Sel. exs.: *E.M. Mélinon 59* (lecto-, P[P00723498] designated by Mori [1990: 286]).

INVENTORY DATA (FG). — 4566 trees in 206 plots; $F_{\max} = 14\%$; $dbh_{\text{inv}} = 65.6$ cm.

[724] *Lecythis persistens* subsp. *aurantiaca* S.A.Mori
(Fig. 29D)

Mem. New York Bot. Gard. 44: 32 (Mori 1987).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: avun-duwó • Te: watili tsing • Cr: mao-blan • Br: matamatá.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *S.A. Mori et al. 15075* (holo-, NY[00312044, 00312103]; iso-, CAY[CAY027885], K[K000600123], U[U0003108], WIS[v 0255904 WIS]).

INVENTORY DATA (FG). — 41 trees in 15 plots; $F_{\max} = 2.7\%$; $dbh_{\text{inv}} = 60\text{ cm}$.

[725] *Lecythis pneumatophora* S.A.Mori

Fl. Neotrop. Monogr. 21 (2): 288 [10 Apr. 1990] (Mori 1990).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Nt: sabana luabi • Cr: mao-marikaj • Fr: mahot marécage.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *S.A. Mori & Y. Veyret 8983* (holo-, NY[00000884, 00285206]; iso-, P[P00723487]).

INVENTORY DATA (FG). — 7 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 50\text{ cm}$.

[726] *Lecythis poiteauii* O.Berg

Fl. Bras. [Martius] 14 (1): 615 [15 Jan. 1859] (Berg 1859). — *Jugastrum poiteauii* (O.Berg) Miers, *Trans. Linn. Soc. London* 30 (2): 279 [14 Nov. 1874] (Miers 1874). — *Eschweilera poiteauii* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 40 [Feb. 1892] (Niedenzu 1892).

Lecythis racemiflora Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20:* 200 (Sagot 1885).

Chytroma foetida R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 85 [22 Aug. 1939] (Knuth 1939).

VERNACULAR NAMES. — Pa: kwatri-duwó, kwatri-purubumna, kwatri-wasiune • Te: watili pitág • Wp: maní'i • Nt: luabi, meli • Cr: mao-jonn • Fr: mahot jaune • Br: jaraná-amarela.

HERBARIUM DATA (FG). — 91 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material P[P00723483, P00723484, P00723485, P00723486, P00723488]).

INVENTORY DATA (FG). — 766 trees in 171 plots; $F_{\max} = 5.8\%$; $dbh_{\text{inv}} = 83.1\text{ cm}$.

[727] *Lecythis praecula* (Sandwith)

S.A.Mori ex Molino & Sabatier, comb. nov.

Eschweilera praecula Sandwith, *Bull. Misc. Inform. Kew* 1935 (3): 127 [20 May 1935] (Sandwith 1935).

NOTES. — Known only from the Guiana Shield. Mori & Prance (Mori et al. 1990: 313) hesitantly placed *Eschweilera praecula* in

synonymy under *L. holcogyne*, specifying that it has “flowers and fruits intermediate in size between *L. holcogyne* and *L. chartacea*”. They further hypothesized that *E. praecula* could be a hybrid between these two *Lecythis* species. Since then, Scott Mori has become convinced that it was indeed a distinct species, whether or not it was the result of hybridization, as proven by the determinants “*L. praecula* (Sandwith) Mori” that he affixed to several specimens, notably at CAY. More recently, he has personally and repeatedly reasserted this position to the first two authors of the present work, at the same time explaining his reluctance to publish this new combination because of the uncertainty of the generic limits of *Lecythis*, which has been shown by genetic studies to be polyphyletic (Mori et al. 2017). Even so, these studies have provided evidence that *E. praecula*, as well as *E. congestiflora* and *E. simiorum* are distant from the core *Eschweilera*, and grouped together with several *Lecythis* species (including *L. holcogyne*) in a “Chartacea clade” (Mori et al. 2017). Furthermore, *L. praecula* appeared distinct from *L. holcogyne* (Mori et al. 2017). We are therefore in the presence of a taxon whose valid name is still *E. praecula* although it does not belong to this genus. Even if it is on a provisional basis (the provisional often lasts decades when it comes to botanical nomenclature), it seems to us necessary to remove this taxon from the genus *Eschweilera*, and this can only be done, for the time being, by a transfer into *Lecythis*.

VERNACULAR NAMES. — Pa: kwatri-seinó.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *S.A. Mori et al. 15701*.

INVENTORY DATA (FG). — 32 trees in 25 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 72.3\text{ cm}$.

[728] *Lecythis simiorum* Benoit

Notul. Syst. (Paris) 3: 178 [25 Dec. 1915] (Benoist 1915). — *Eschweilera simiorum* (Benoist) Eyma, *Polygon. Guttif. Lecythid. Surinam*: 81 (Eyma 1932). — *Chytroma simiorum* (Benoist) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 81 [22 Aug. 1939] (Knuth 1939).

Eschweilera simiorum var. *latifolia* Eyma, *Polygon. Guttif. Lecythid. Surinam* 81 (Eyma 1932). — *Chytroma simiorum* var. *latifolia* (Eyma) R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 82 [22 Aug. 1939] (Knuth 1939).

Chytroma crenata R.Knuth, *Pflanzenr. [Engler]* IV.219a (Heft 105): 84 [22 Aug. 1939] (Knuth 1939).

NOTE. — According to Mori et al. (2017), this species is better placed in *Lecythis* than in *Eschweilera*.

VERNACULAR NAMES. — Pa: kwatri-wasiune-seine • Ka: meku kuwaili • Wp: a'i amiya • Wn: pukut.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *R. Benoist 1065* (lecto-, P[P01900116], designated by Mori et al. [1990: 215]; isolecto-, K[K000600162], P[P04561962]).

INVENTORY DATA (FG). — 74 trees in 54 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 37.7\text{ cm}$.

[729] *Lecythis zabucajo* Aubl.

Hist. Pl. Guiane 2: 718 [Jun.-Dec. 1775] (Aublet 1775), *pro parte quoad fructum tantum* (*Hist. Pl. Guiane* 4: pl. 283 in pl. 285 [Jun.-Dec. 1775] (Aublet 1775)).

Lecythis validissima Miers, *Trans. Linn. Soc. London* 30 (2): 205 [14 Nov. 1874] (Miers 1874).

Lecythis crassinoda Miers, *Trans. Linn. Soc. London* 30 (2): 216 [14 Nov. 1874] (Miers 1874), *pro parte quoad fructum tantum*.

Lecythis tumefacta Miers, *Trans. Linn. Soc. London* 30 (2): 216 [14 Nov. 1874] (Miers 1874).

Lecythis lecomtei Pamp., *Agric. Colon.* 10: 633 (Pampanini 1916).

Lecythis davisii Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 213 [27 June 1932] (Sandwith 1932).

Lecythis davisii var. *gracilipes* Eyma, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 4: 77 (Eyma 1932).

Lecythis hians A.C.Sm., *Phytologia* 1 (3): 123 [21 Jan. 1935] (Smith 1935).

VERNACULAR NAMES. — Pa: wakukwa-adava • Ka: kuata tumali, yalakalu tumali epi • Te: titikö • Wp: ka'i nu'a, tukuke, yapukay, ýapukay • Wn: tula, tulaimë • Nt: kwata patu, kwatan patu • Cr: kanari-makak • Fr: canari macaque, marmite de singe (*fide* Aublet 1775) • Br: castanha-sapucaia, marmita-de-macaco, sapucaia.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *S.A. Mori et al. 25034*.

INVENTORY DATA (FG). — 120 trees in 83 plots; $F_{\max} < 1\%$; $dbh_{inv} = 105$ cm.

Family LEGUMINOSAE Juss.
(alternative name: FABACEAE Lindl.)

NOTE. — According to LPWG *et al.* (2017), the genera listed here belong to five of the six subfamilies of Leguminosae:

- subfam. Caesalpinioideae (underlined genera belong to the mimosoid clade): *Abarema*, *Acacia*, *Albizia*, *Anadenanthera*, *Batesia*, *Calliandra*, *Cassia*, *Cedrelinga*, *Chamaecrista*, *Chloroleucon*, *Drimorphandra*, *Dinzia*, *Enterolobium*, *Hydrochorea*, *Inga*, *Leucaena*, *Macrosamanea*, *Parkia*, *Pentaclethra*, *Pseudopiptadenia*, *Recordoxylon*, *Senegalia*, *Senna*, *Stryphnodendron*, *Tachigali*, *Vouacapoua*, *Zygia*.
- subfam. Cercidoideae: *Bauhinia*.
- subfam. Detarioideae: *Coparia*, *Crudia*, *Cynometra*, *Eperua*, .3, *Macrolobium*, *Paloue*, *Peltogyne*.
- subfam. Dialioideae: *Dialium*, *Dicorynia*, *Martiodendron*.
- subfam. Papilioideae: *Alexa*, *Amphidodon*, *Andira*, *Bocoa*, *Can-dolleodendron*, *Clathrotropis*, *Diplostropis*, *Dipteryx*, *Dussia*, *Erythrina*, *Guianodendron*, *Hymenolobium*, *Leptolobium*, *Limadendron*, *Lonchocarpus*, *Monopteryx*, *Ormosia*, *Paramachaerium*, *Platymiscium*, *Pterocarpus*, *Spirotropis*, *Swartzia*, *Taralea*, *Vatairea*, *Vataireopsis*, *Zollernia*.

Genus *Abarema* Pittier

[730] *Abarema barbouriana*
(Standl.) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 70 (Barneby & Grimes 1996). — *Pithecellobium barbourianum* Standl., *Contr. Arnold Arbor.* 5: 74 (Standley 1933), “*Pithecolobium*”.

Albizia dubia Britton & Killip, *Ann. New York Acad. Sci.* 35: 132 [1 Apr. 1936] (Britton & Killip 1936), “*Albizia* (?) *dubia*”.

Pithecellobium fanshawei Sandwith, *Kew Bull.* 3 (2): 314 [20 Nov. 1948] (Sandwith 1948).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *O. Poncy 901*.

INVENTORY DATA (FG). — 27 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{inv} = 24.1$ cm.

[731] *Abarema curvicarpa*
(H.S.Irwin) Barneby & J.W.Grimes

Acta Amazonica 14 (1-2, Suppl.): 95 (“1984” publ. 1986) (Barneby & Grimes 1986). — *Pithecellobium curvicarpum* H.S.Irwin, *Mem. New York Bot. Gard.* 15: 107 (Irwin 1966).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: kabanafoo.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.W. Grimes et al. 3312*.

INVENTORY DATA (FG). — 18 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{inv} = 130$ cm.

[732] *Abarema gallorum* Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 51 (Barneby & Grimes 1996).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-P. Lescure 877* (holo-, P[P00199787]; iso-, CAY[CAY028065, CAY028066], NY[00001428]).

INVENTORY DATA (FG). — 44 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{inv} = 35.1$ cm.

[733] *Abarema jupunba* (Willd.) Britton & Killip
var. *jupunba*

Ann. New York Acad. Sci. 35: 126 [1 Apr. 1936] (Britton & Killip 1936). — *Acacia jupunba* Willd., *Sp. Pl.*, ed. 4 4 (2): 1067 [Apr. 1806] (Willdenow 1806). — *Feuilleea jupunba* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891), “*jupumba*”. — *Pithecellobium jupunba* (Willd.) Urb., *Symb. Antill.* [Urban] 2 (2): 257 [20 Oct. 1900] (Urban 1900), “*Pithecolobium*”. — *Jupunba jupunba* (Willd.) Britton & Rose, *N. Amer. Fl.* 23 (1): 27 [11 Feb. 1928] (Britton & Rose 1928).

Pithecellobium micradenium Benth., *London J. Bot.* 3: 217 (Bentham 1844). — *Feuilleea micradenia* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Pithecellobium brongniartii Duchass. & Walp., *Flora* 36: 232 (Duchassaing & Walpers 1853).

NOTES. — *Feuilleea jupunba* (Willd.) Kuntze is based on “*Acacia jupumba* W.”, but includes *Mimosa trapezifolia* Vahl in synonymy. Barneby & Grimes (1996: 69) cited it twice, as a synonym of var. *jupunba* as “*F. jupumba* [sic] (Willdenow) O. Kuntze”, and as a synonym of var. *trapezifolia* as “*F. jupumba* (Vahl) Kuntze”. The latter does not exist, and should not be cited.

VERNACULAR NAMES. — Pa: yuumwi-washiunu • Ka: kalaip'e'u, kalaipiū, kalape'u • Te: akiki nami • Wp: kalaip'e'i, kwata luway • Wn: kalaujalan • Nt: busi kabana foo, tamalen • Cr: akasya, bougouni-ti'fey • Fr: bois macaque des bancs de sable • Br: angelim-falso, pau-bicho, saboeiro.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2805*.

INVENTORY DATA (FG). — 190 trees in 109 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 101.2$ cm.

[734] *Abarema jupunba* var. *trapezifolia*
(Vahl) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 69 (Barneby & Grimes 1996). — *Mimosa trapezifolia* Vahl, *Eclog. Amer.* 3: 36 (Vahl 1807). — *Inga trapezifolia* (Vahl) DC., *Prodr. [A. P. de Candolle] 2:* 441 [mid Nov. 1825] (Candolle 1825). — *Pithecellobium trapezifolium* (Vahl) Benth., *J. Bot. [Hooker] 2:* 142 (Bentham 1840), “*Pithecolobium*”. — *Abarema trapezifolia* (Vahl) Pittier, *Arb. Arbusc. Venez.* 6-8: ? [Aug.-Sep. 1927] (Pittier 1927). — *Jupunba trapezifolia* (Vahl) Moldenke, *Bull. Torrey Bot. Club* 59 (3): 155 [Mar. 1932] (Moldenke 1932).

Mimosa atakta Steud., *Flora* 26 (45): 758 [17 Dec. 1843] (Steudel 1843).

Pithecellobium benthamianum Miq., *Linnaea* 18: 592 [“1844” publ. prob. Aug. 1845] (Miquel 1845).

VERNACULAR NAMES. — Wp: kalaip'e'i.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *S.A. Mori et al. 23938*.

INVENTORY DATA (FG). — 6 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54$ cm.

[735] *Abarema laeta* (Benth.) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 79 (Barneby & Grimes 1996). — *Pithecellobium laetum* Benth., *London J. Bot.* 3: 203 (Bentham 1844), “*Pithecolobium*”. — *Inga laeta* Poepp., *Nova genera ac species plantarum [Poepig & Endlicher] 3:* 80 [23-25 Jan. 1845] (Poeppig 1845). — *Feuilleea laeta* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891). — *Klugiodendron laetum* (Benth.) Britton & Killip, *Ann. New York Acad. Sci.* 35: 126 [1 Apr. 1936] (Britton & Killip 1936).

Pithecellobium polycarpum Poepp., *Nova genera ac species plantarum [Poepig & Endlicher] 3:* 81 [23-25 Jan. 1845] (Poeppig 1845), “*Pithecolobium*”.

Klugiodendron umbrianum Britton & Killip, *Ann. New York Acad. Sci.* 35: 126 [1 Apr. 1936] (Britton & Killip 1936).

NOTES. — Although *Pithecellobium laetum* Benth. is based on Poeppig's manuscript and on the same type as *Inga laeta* Poepp., the latter was described independently, as a new species (see annotation by Ernst Vitek dated 2007 on the holotype of *I. laeta*, *E.F. Poeppig 2367* [W[W0013462]]). Poeppig is the sole author of the vol. 3 of *Nova Genera ac species plantarum* (Stafleu & Cowan, 1983: *TL-2-8091*).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.W. Grimes & P. Acevedo-Rodríguez 3322*, height 30 m.

[736] *Abarema mataybifolia* (Sandwith) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 78 (Barneby & Grimes 1996). — *Pithecellobium mataybifolium* Sandwith, *Kew Bull.* 3 (2): 313 [20 Nov. 1948] (Sandwith 1948).

VERNACULAR NAMES. — Wp: kalaip'e'i.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier 3536*.

INVENTORY DATA (FG). — 73 trees in 40 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 46.6$ cm.

[737] *Abarema* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4425*.

INVENTORY DATA (FG). — 1 tree, dbh = 11.6 cm.

Genus *Acacia* Mill.

[738] *Acacia mangium* Willd.

Sp. Pl., ed. 44 (2): 1053 [Apr. 1806] (Willdenow 1806). — *Racosperma mangium* (Willd.) Pedley, *Austrobaileya* 2 (4): 352 (Pedley 1987).

NOTE. — A species native to East Malesia (from Maluku to N. Australia), introduced in most tropical regions, now invasive in some northern French Guiana savannas.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *C. Delnatté 1556*.

Genus *Albizia* Durazz.

[739] *Albizia niopoides* (Spruce ex Benth.) Burkart

Legum. Argent., ed. 2: 542 (Burkart 1952). — *Pithecellobium niopoides* Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 591 [10 Apr. 1875] (Bentham 1875), “*Pithecolobium*”. — *Feuilleea niopoides* (Spruce ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Gagnebina richardiana Wall. ex Voigt, *Hort. Suburb. Calcutt.* 257 (Voigt 1845), *nom. nud.*

Pithecellobium caribaeum Urb., *Symb. Antill. [Urban]* 2 (2): 260 [20 Oct. 1900] (Urban 1900), “*Pithecolobium*”. — *Albizia caribaea* (Urb.) Britton & Rose, *N. Amer. Fl.* 23 (1): 44 [11 Feb. 1928] (Britton & Rose 1928), “*Albizzia*”.

Pithecellobium hassleri Chodat, *Bull. Herb. Boissier*, sér. 2, 4: 483 (Chodat 1904), “*Pithecolobium*”. — *Albizia hassleri* (Chodat) Burkart, *Darwiniana* 7 (4): 517 (Burkart 1947), “*Albizzia*”.

Senegalnia liebmannii Britton & Rose, *N. Amer. Fl.* 23 (2): 116 [25 Sep. 1928] (Britton & Rose 1928).

Senegalnia guacamayo Britton & Killip, *Ann. New York Acad. Sci.* 35: 142 [1 Apr. 1936] (Britton & Killip 1936). — *Acacia guacamayo* (Britton & Killip) Standl., *Trop. Woods* 52: 26 (Standley 1937). — *Albizia guacamayo* (Britton & Killip) L.Cárdenas, *Ernstia*, sér. 2, 1 (4): 133 (Cárdenas 1992).

HERBARIUM DATA (FG). — A single collection, *D. Loubry 1579*.

INVENTORY DATA (FG). — 1 tree, dbh = 25.3 cm.

[740] *Albizia pedicellaris* (DC.) L.Rico

Novon 9 (4): 555 [winter 1999] (Rico 1999). — *Inga pedicellaris* DC., *Prodr. [A. P. de Candolle]* 2: 441 [mid Nov. 1825] (Candolle 1825). — *Pithecellobium pedicellare* (DC.) Benth., *London J. Bot.* 3: 219 (Bentham 1844), “*Pithecellobium*”. — *Feuilleea pedicellaris* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891). — *Samanea pedicellaris* (DC.) Killip, *Trop. Woods* 63; 4 (Killip 1940). — *Macrosamanea pedicellaris* (DC.) Kleinhoonte, *Fl. Suriname* 2 (2): 329 (Kleinhoonte 1940). — *Balizia pedicellaris* (DC.) Barneby & J.W.Grimes, *Mem. New York Bot. Gard.* 74 (1): 37 (Barneby & Grimes 1996).

Mimosa terminalis Vell., *Fl. Flumin. Icon.* 11: t. 30 [“1827” publ. 29 Oct. 1831] (Vellozo 1831), *nom. illeg. hom., non Salisb.* (Salisbury 1796). — *Feuilleea terminalis* Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: avukun-kamwi • Ka: alawata muleli, alesikyi'i • Wp: yapu'i, yapu'i pilá • Nt: asao • Cr: bougouni-blanc, bwa-makak, bwa-pagód-ti-féy • Fr: bois pagode • Br: apuxiqui, gambui, jurema-branca, mapuxiqui-vermelho, sobreiro.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *L. Barrabé 232*.

INVENTORY DATA (FG). — 104 trees in 67 plots; F_{max} = 2.3 %; dbh_{inv} = 102.8 cm.

Genus *Alexa* Moq.

[741] *Alexa grandiflora* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 33 (Ducke 1915).

NOTE. — *P. Grenand & M.-F. Prévost 2017*, identified by R. S. Cowan in 1977 at CAY as *A. grandiflora*, actually belongs to *Alexa wachenheimii*.

VERNACULAR NAMES. — Wp: ani, takulawa'i • Cr: sen-martin-blanc • Br: melancieira.

HERBARIUM DATA (FG). — A single collection, *P. Grenand 774*.

SIZE. — Up to 30 m tall (Ducke 1915: 33).

[742] *Alexa wachenheimii* Benoist

Bull. Mus. Natl. Hist. Nat. 27: 115 (Benoist 1921), “*Wachenheimi*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: karavia-kamwi • Ka: kulekulau • Te: tapaka'i • Cr: sen-martin-blanc • Br: melancieira.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *G. Wachenheim 93* (original material P[P01817846, P01817847, P03104012]).

INVENTORY DATA (FG). — 44 trees in 19 plots; F_{max} = 1.2 %; dbh_{inv} = 96 cm.

Genus *Amphiodon* Huber

[743] *Amphiodon effusus* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 399 (Huber 1909). — *Poecilanthe effusa* (Huber) Ducke, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 4 (6): 733 (Ducke 1932).

Poecilanthe ovalifolia Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 398 (Kleinhoonte 1926).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Br: cumarú-de-rato.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier 4815*.

INVENTORY DATA (FG). — 107 trees in 21 plots; F_{max} = 2.5 %; dbh_{inv} = 29.9 cm.

Genus *Anadenanthera* Speg.

[744] *Anadenanthera peregrina* (L.) Speg.

Physis (Buenos Aires) 6: 314 (Spegazzini 1923). — *Mimosa peregrina* L., *Sp. Pl.* 1: 520 [1 May 1753] (Linnaeus 1753). — *Acacia peregrina* (L.) Willd., *Sp. Pl.*, ed. 4 4 (2): 1073 [Apr. 1806] (Willdenow 1806). — *Piptadenia peregrina* (L.) Benth., *J. Bot. [Hooker]* 4: 340 (Bentham 1842). — *Niopa peregrina* (L.) Britton & Rose, *Addisonia* 12: 37 (Britton & Rose 1927).

Inga niopo Humb. & Bonpl. ex Willd., *Sp. Pl.*, ed. 4 4 (2): 1027 [Apr. 1806] (Willdenow 1806). — *Mimosa niopo* (Humb. & Bonpl. ex Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 48 [3 Sep. 1810] (Poirier 1810). — *Acacia niopo* (Humb. & Bonpl. ex Willd.) Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 282 [Apr. 1824] (Kunth 1824). — *Piptadenia niopo* (Humb. & Bonpl. ex Willd.) Spruce, *Notes Bot. Amaz.* 2: 426 (Spruce 1908).

Acacia microphylla Willd., *Sp. Pl.*, ed. 4 4 (2): 1073 [Apr. 1806] (Willdenow 1806).

Mimosa parvifolia Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 74 [3 Sep. 1810] (Poirier 1810), *nom. illeg. hom., non Sw.* (Swartz 1800).

Acacia angustiloba DC., *Prodr. [A. P. de Candolle]* 2: 470 [mid Nov. 1825] (Candolle 1825).

Mimosa acacioides Benth., *J. Bot. [Hooker]* 2: 132 (Bentham 1840), “*M.? acacioides*”.

NOTE. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Br: paricá.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 16138*.

SIZE. — Up to 40 cm dbh (Barneby *et al.* 2011).

Genus *Andira* Lam.

[745] *Andira coriacea* Pulle

Recueil Trav. Bot. Néerl. 6: 267 (Pulle 1909).

Andira wachenheimii Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 296 (Benoist 1919), “Wachenheimi”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wanetun • Ka: akuli kiyeleli, lele elepali • Wn: aliamëlë, pokololomili • Nt: lebi tyabisi • Cr: sen-marten-rouj • Fr: saint-martin rouge • Br: andira-uxi, angelim-coco, morcegueira.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2281*.

INVENTORY DATA (FG). — 167 trees in 68 plots; $F_{\max} = 2.2\%$; $\text{dbh}_{\text{inv}} = 102.5$ cm.

[746] *Andira inermis* (W.Wright) Kunth ex DC.

Prodr. [A. P. de Candolle] 2: 475 [mid Nov. 1825] (Candolle 1825). — *Geoffroea inermis* W.Wright, *London Med. J.* 8 (3): 256 (Wright 1787). — *Geoffroea acutifolia* Stokes, *Bot. Mat. Med.* 4: 46 (Stokes 1812), “*Geoffrea*”, nom. illeg. superfl. (based on *Geoffroea inermis* (as “Sw.”). — *Vouacapoua inermis* (W.Wright) A.Lyons, *Pl. Nam.*: 396 (Lyons 1900). — *Andira jamaicensis* Urb., *Symb. Antill. [Urban]* 4 (2): 298 [15 Feb. 1905] (Urban 1905), nom. illeg. superfl. (based on *Geoffroea inermis* (as “Sw.”)).

Andira excelsa Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 385 [12 July 1824] (Kunth 1824).

Andira riparia Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 386 [12 July 1824] (Kunth 1824).

Pterocarpus sapindoides DC., *Prodr. [A. P. de Candolle]* 2: 419 [mid Nov. 1825] (Candolle 1825). — *Andira sapindoides* (DC.) Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 123 [Mar. 1860] (Bentham 1860). — *Andira inermis* var. *sapindoides* (DC.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 202 [late 1860] (Grisebach 1860). — *Vouacapoua sapindoides* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 212 [5 Nov. 1891] (Kuntze 1891), “*Vuacapua*”, “*sapindoides*”. — *Andira jamaicensis* var. *sapindoides* (DC.) Stehlé, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 18: 116 (Stehlé 1946), “*sapiendioidea*”.

Amerimnon affine Spreng., *Syst. Veg. [Sprengel]* 3: 192 [Jan.-Mar. 1826] (Sprengel 1826), “*Ámerimnum*”.

Glycyrrhiza undulata Ruiz & Pav. ex G.Don, *Gen. Hist.* 2: 227 [Oct. 1832] (Don 1832).

Andira grandiflora Guill. & Perr., *Fl. Seneg. Tent.* 1 (7): 254 [22 Oct. 1832] (Guillemin & Perrottet 1832). — *Andira inermis* subsp. *grandiflora* (Guill. & Perr.) J.B.Gillett ex Polhill, *Kew Bull.* 23 (3): 490 (Polhill 1969).

Andira acuminata Benth., *Commentat. Legum. Gen.*: 45 [June 1837] (Bentham 1837).

Andira inermis var. *riedelii* Benth., *J. Proc. Linn. Soc., Bot. 4 (suppl.)*: 123 [Mar. 1860] (Bentham 1860), “*Riedeli*”.

Lonchocarpus staudtii Harms, *Bot. Jahrb. Syst.* 26 (3-4): 301 [31 Jan. 1899] (Harms 1899).

Andira chiricana Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 234 [22 Sep. 1917] (Pittier 1917).

NOTES. — The “*Geoffroea jamaicensis inermis*” of W. Wright (1777: 512) is an invalid trinomial (Turland et al. 2018: Art. 23.6a) not to be regarded as a species name. Therefore, “*Geoffroea jamaicensis* W.Wright” does not exist, and *Andira jamaicensis* Urb. has no basionym.

VERNACULAR NAMES. — Pa: wanetun • Ka: eleyulu, malipa wewe • Wp: kule’i • Nt: lebi tyabisi.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *M.-F. Prévost 3840*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 49.7$ cm.

[747] *Andira surinamensis* (Bondt) Splitg. ex Pulle

Enum. Vasc. Pl. Surinam 229 (Pulle 1906). — *Geoffroea surinamensis* Bondt, *Cort. Geogr. Surinam.*: 12 (Bondt 1788). — *Geoffroea obtusifolia* Stokes, *Bot. Mat. Med.* 4: 46 (Stokes 1812), “*Geoffrea*”, nom. illeg. superfl. (based on *Geoffroea surinamensis*). — *Andira retusa* var. *surinamensis* (Bondt) DC., *Prodr. [A. P. de Candolle]* 2: 476 [mid Nov. 1825] (Candolle 1825). — *Vouacapoua surinamensis* (Bondt) Kuntze, *Revis. Gen. Pl.* 1: 212 [5 Nov. 1891] (Kuntze 1891), “*Vuacapua*”.

Geoffroea pubescens Rich., *Actes Soc. Hist. Nat. Paris* 1: 111 [Oct. 1792] (Richard 1792), “*Geoffroya*”.

Geoffroea retusa Poir., *Encycl. [J. Lamarck et al.]* 8: 182 [22 Aug. 1808] (Poiret 1808), “*Geoffraea*”. — *Andira retusa* (Poir.) Kunth ex DC., *Prodr. [A. P. de Candolle]* 2: 475 [mid Nov. 1825] (Candolle 1825). — *Vouacapoua retusa* (Poir.) A.Lyons, *Pl. Nam.* 396 (Lyons 1900).

Andira retusa var. *oblonga* Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 121 [Mar. 1860] (Bentham 1860).

Andira oblonga Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 121 [Mar. 1860] (Bentham 1860), nom. nud. pro syn.

Andira surinamensis var. *ovatifoliolata* N.F.Mattos, *Loefgrenia* 58: 3 (Mattos 1973).

VERNACULAR NAMES. — Pa: wanetun • Ka: lele elepali • Wp: anila wisi, pana’i wu • Wn: haka • Nt: lebi tyabisi • Cr: sen-marten-rouj • Fr: saint-martin rouge • Br: andira-uxi, angelim-coco, morcegueira.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2672*.

INVENTORY DATA (FG). — 36 trees in 21 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 62.2$ cm.

Genus *Batesia* Spruce ex Benth.

[748] *Batesia floribunda* Spruce ex Benth.

Trans. Linn. Soc. London 25 (2): 303 [“1866” publ. 30 Nov. 1865] (Bentham 1865).

VERNACULAR NAMES. — Pa: wanaku-duwē, wanaku-umutinō • Br: tento-miúdo, tento-verdadeiro.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier 2308*.

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 89.1 \text{ cm}$.

Genus *Bauhinia* Plum. ex L.

[749] *Bauhinia cinnamomea* DC.

Prod. [A. P. de Candolle] 2: 517 [mid Nov. 1825] (Candolle 1825).

Bauhinia versteegii Pulle, *Enum. Vasc. Pl. Surinam* 213 (Pulle 1906).

VERNACULAR NAMES. — Wp: paná miwi • Wn: omole, wapaimé • Cr: ankono.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, P, [P00798577]; iso-, B[not seen, photo F neg. N° 1572]).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 10.5 \text{ cm}$.

[750] *Bauhinia eilertsii* Pulle

Recueil Trav. Bot. Néerl. 6: 269 (Pulle 1909).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 16705*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 43.3 \text{ cm}$.

Genus *Bocoa* Aubl.

[751] *Bocoa prouacensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 38 [Jun.-Dec. 1775] (Aublet 1775), “*Prouassensis*” on plate. — *Gajanus prouacensis* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891), “*provacensis*”. — *Swartzia prouacensis* (Aubl.) Amshoff, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 52: 40 (Amshoff 1939).

Swartzia minutiflora Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 408 [“1925” publ. Jan. 1926] (Kleinhoonte 1926), “*Swartsia*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: á-danó, á-sivari, wanaku-danó • Ka: aliyaná-i • Te: tatulimá • Wp: tunu'i pilá • Wn: kutali, wapiku • Nt: ayee udu • Cr: boko, bwa-fé • Fr: boco, bois de fer • Br: muira-jibóia.

HERBARIUM DATA (FG). — 83 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000588689]).

INVENTORY DATA (FG). — 1459 trees in 163 plots; $F_{\max} = 9.8\%$; $\text{dbh}_{\text{inv}} = 63.3 \text{ cm}$.

[752] *Bocoa viridiflora* (Ducke) R.S.Cowan

Proc. Biol. Soc. Washington 87: 120 (Cowan 1974). — *Swartzia viridiflora* Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 44 [Sep. 1935] (Ducke 1935).

VERNACULAR NAMES. — Nt: ayee udu • Cr: boko, bwa-fé • Fr: boco, bois de fer • Br: muira-jibóia.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *S.A. Mori et al. 24136*.

INVENTORY DATA (FG). — 111 trees in 17 plots; $F_{\max} = 5.5\%$; $\text{dbh}_{\text{inv}} = 39.4 \text{ cm}$.

Genus *Calliandra* Benth.

[753] *Calliandra hymenaeodes* (Rich.) Benth.
(Fig. 30A)

Trans. Linn. Soc. London 30 (3): 537 (Bentham 1875), “*hymenaeoides*”. — *Mimosa hymenaeodes* Rich., *Syn. Pl. [Persoon]* 2 (1): 262 [Nov. 1806] (Richard 1806). — *Inga hymenaeodes* (Rich.) Desv., *J. Bot. Agric.* 3: 70 (Desvaux 1814).

Calliandra patrisii Sagot, *Ann. Sci. Nat., Bot. sér. 6, 13:* 324 (Sagot 1882).

NOTES. — Known only from the Guiana Shield. It is clear from the protologue of the basionym, *Mimosa hymenaeodes* Rich., that it is authored by L. C. Richard alone: the text reads “23. *Hymenaeodes*, fol. *uniparibus*. Richard.” (Stafleu & Cowan, 1983: TL-2-7732). Richard’s original spelling of the epithet on the holotype label is “*hymaeneoides*”. He naturally corrected the misspelling (“*hymaene-*” for “*hymenae-*”) in the protologue. However, there is no way to know whether the deletion of the “*i*” (“-odes” instead of “-oides”) was intentional or a typographical error. Anyway, both terminations are correct and have the same meaning, so there is no reason to change the original spelling *hymenaeodes*.

VERNACULAR NAMES. — Pa: asiru-wašiunó.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (type P[P00199195]).

SIZE. — French Guiana. *O. Poncy et al. 1430*, 10 cm.

[754] *Calliandra surinamensis* Benth.

London J. Bot. 3: 105 (Bentham 1844).

Inga fasciculata Willd., *Sp. Pl., ed. 4* (2): 1022 [Apr. 1806] (Willdenow 1806). — *Feuilleea fasciculata* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891). — *Anneslia fasciculata* (Willd.) Kleinhoonte, *Fl. Suriname* 2 (2): 322 (Kleinhoonte 1940).

Calliandra tenuiflora Benth., *Trans. Linn. Soc. London* 30 (3): 547 [10 Apr. 1875] (Bentham 1875). — *Feuilleea tenuiflora* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: sakeg-kamwi • Br: balão-chinês.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 1042*.

SIZE. — Up to 12 m tall (Barneby 1998).

[755] *Calliandra trinervia* Benth.

London J. Bot. 3: 94 (Bentham 1844). — *Feuilleea trinervia* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R.A.A. Oldeman 1314*.

SIZE. — Up to 30 m tall (Berry & Barneby 1999).

[756] *Calliandra* sp. A.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 4695*.

INVENTORY DATA (FG). — 23 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.6$ cm.

Genus *Candolleodendron* R.S.Cowan

[757] *Candolleodendron brachystachyum* (DC.) R.S.Cowan

Rhodora 68: 429 [28 Dec. 1966] (Cowan 1966). — *Swartzia brachystachya* DC., *Prod. [A. P. de Candolle]* 2: 423 [mid Nov. 1825] (Candolle 1825). — *Tounatea brachystachya* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 211 [5 Nov. 1891] (Kuntze 1891), “*Tunatea*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: timuvuyen • Wp: tunu'i • Wn: kíwiliúmë • Nt: lebi nenba • Cr: boko-blán.

HERBARIUM DATA (FG). — 79 collections at CAY. Sel. exs.: *H. Richard 70*.

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24$ cm.

Genus *Cassia* L.

[758] *Cassia cowanii* H.S.Irwin & Barneby

Mem. New York Bot. Gard. 35: 24 (Irwin & Barneby 1982).

VERNACULAR NAMES. — Wp: malimali u.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 22034*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 45.6$ cm.

[759] *Cassia fastuosa* Willd. ex Benth.

Fl. Bras. [Martius] 15 (2): 95 [1 Dec. 1870] (Bentham 1870).

VERNACULAR NAMES. — Wp: malimali • Br: baratinha, faveirinha, marimari.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5026*, dbh 56 cm.

[760] *Cassia grandis* L.f.
(Fig. 30B)

Suppl. Pl.: 230 [“1781” publ. Apr. 1782] (Linnaeus 1782). — *Cassia brasiliiana* Lam., *Encycl. [J. Lamarck et al.]* 1 (2): 649 [1 Aug. 1785] (Lamarck 1785), *nom. illeg. superfl.* (based on *Cassia grandis*). — *Cathartocarpus grandis* (L.f.) Pers., *Syn. Pl. [Persoon]* 1: 459 [1 Apr.-15 June 1805] (Persoon 1805). — *Bactyriobium grande* (L.f.) Hornem., *Suppl. Hort. Bot. Hafn.* 1: 135 (Hornemann 1819).

Cassia brasiliensis Buc'hoz, *Hist. Univ. Règne Vég.*, Planches, Cent. 10, Decad. 4: pl. 2 (Buc'hoz 1775).

Cassia mollis Vahl, *Symb. Bot. [Vahl]* 3: 57 (Vahl 1794).

Cathartocarpus brasilianus Jacq., *Fragm. Bot.* 58 (Jacquin 1806).

Bactyriobium molle Schrad., *Gött. Gel. Anz.* 2: 713 [5 May 1821] (Schrader 1821).

Cassia brasiliiana var. *tomentosa* Miq., *Linnaea* 18: 578 [“1844” publ. prob. Aug. 1845] (Miquel 1845).

Cathartocarpus erubescens Ham., *Pharm. J. Trans.* 5: 119 (Hamilton 1845).

Cassia regia Standl., *Contr. U.S. Natl. Herb.* 18 (3): 103 [11 Feb. 1916] (Standley 1916).

Cassia pachycarpa de Wit, *Webbia* 11: 259 (de Wit 1955).

VERNACULAR NAMES. — Br: chuva-de-ouro.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4766*, dbh 40 cm.

[761] *Cassia spruceana* Benth.

Fl. Bras. [Martius] 15 (2): 92 [1 Dec. 1870] (Bentham 1870).

Cassia sagotiana Benth., *Fl. Bras. [Martius]* 15 (2): 93 [1 Dec. 1870] (Bentham 1870).

VERNACULAR NAMES. — Br: fava-marimari.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *P.A. Sagot 802*, 1856 (holotype of *Cassia sagotiana*: K[K000504594]; iso-, F[V0057675F, V0057676F]; possible isotype (no date) G[G00364747]).

INVENTORY DATA (FG). — 7 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 58.6$ cm.

Genus *Cedrelinga* Ducke

[762] *Cedrelinga cateniformis* (Ducke) Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 70 (Ducke 1922), “*catenaeformis*”. — *Piptadenia cateniformis* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 17 (Ducke 1915), “*catenaeformis*”. — *Pithecellobium cateniformis* (Ducke) L.Cárdenas, *Revista Fac. Agron. (Maracay)* 7 (3): 124 (Cárdenas 1974).

VERNACULAR NAMES. — Nt: donsede • Cr: akajou-lan • Br: cedrorana.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier 2316*.

INVENTORY DATA (FG). — 23 trees in 11 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 89.8$ cm.

Genus *Chamaecrista* Moench

[763] *Chamaecrista apoucouita*
(Aubl.) H.S.Irwin & Barneby

Mem. New York Bot. Gard. 35: 642 (Irwin & Barneby 1982). — *Cassia apoucouita* Aubl., *Hist. Pl. Guiane* 1: 379 [Jun.-Dec. 1775] (Aublet 1775). — *Cassia acuminata* Willd., *Sp. Pl.*, ed. 4 2 (1): 517 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Cassia apoucouita*) and nom. illeg. hom., non Moench (1974) [synonym of *Senna marilandica* (L.) Link].

Cassia solimoesensis H.S.Irwin, *Mem. New York Bot. Gard.* 16: 106 (Irwin 1967).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000952149](#)]), LINN[[LINN-HS 736.2](#)]).

INVENTORY DATA (FG). — 75 trees in 20 plots; $F_{\max} = 3.9\%$; $dbh_{\text{inv}} = 55.7$ cm.

Genus *Chloroleucon* (Benth.) Record

[764] *Chloroleucon acacioides*
(Ducke) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 141 (Barneby & Grimes 1996). — *Pithecellobium acacioides* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 69 (Ducke 1922), “*Pithecolobium*”.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F. Prévost 1135*.

INVENTORY DATA (FG). — 1 tree, $dbh = 30$ cm.

Genus *Clathrotropis* (Benth.) Harms

[765] *Clathrotropis brachypetala* (Tul.) Kleinhoonte

Recueil Trav. Bot. Néerl. 22: 398 [“1925” publ. Jan. 1926] (Kleinhooonte 1926). — *Diplotropis brachypetala* Tul., *Arch. Mus. Hist. Nat.* 4: 111 (Tulasne 1844). — *Bowdichia brachypetala* (Tul.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 133 (Ducke 1922).

VERNACULAR NAMES. — Ka: inyekulan.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *Service Forestier 368M* (P), $dbh = 40$ cm.

Genus *Copaifera* L.

[766] *Copaifera guyanensis* Desf.

Mém. Mus. Hist. Nat. 7: 376 (Desfontaines 1821). — *Copaiba guyanensis* (Desf.) Kuntze, *Revis. Gen. Pl.* 1: 172 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: maraura • Ka: apa'uwa • Te: kupawa • Wp: kupaipa • Wn: kupajwa, kupeiwa • Nt: paansu miti • Cr: bwakapayou, koupawa • Fr: copahu • Br: copaíba-branca.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *C. Sastre 4009*.

INVENTORY DATA (FG). — 1 tree, $dbh = 22.8$ cm.

[767] *Copaifera* sp. A

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2124*.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 75.8$ cm.

Genus *Crudia* Schreb.

[768] *Crudia aromatica* (Aubl.) Willd.

Sp. Pl., ed. 4 2 (1): 450 [Mar. 1799] (Willdenow 1799). — *Touchiroa aromatica* Aubl., *Hist. Pl. Guiane* 1: 384 [Jun.-Dec. 1775] (Aublet 1775). — *Apalatoa aromatica* (Aubl.) Lam., *Encycl. J. Lamarck et al.* 4 (2): 559 [1 Nov. 1798] (Lamarck 1798), “*Opalatoa*”. — *Apalatoa aromatica* (Aubl.) Taub., *Bot. Centralbl.* 47: 394 [31 Sept 1891] (Taubert 1891), isonym.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kuyali tapaiko • Wp: kupeiwa • Nt: pikin paanga.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000826529](#)]).

INVENTORY DATA (FG). — 97 trees in 18 plots; $F_{\max} = 2.5\%$; $dbh_{\text{inv}} = 66.3$ cm.

[769] *Crudia bracteata* Benth.

J. Bot. [Hooker] 2: 101 (Bentham 1840), “*Crudyā*”.

VERNACULAR NAMES. — Wp: kupeiwa.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, K[[K000555173](#)]; P[[P03111132](#), [P03111134](#)]).

INVENTORY DATA (FG). — 420 trees in 43 plots; $F_{\max} = 6.2\%$; $dbh_{\text{inv}} = 33.6$ cm.

[770] *Crudia oblonga* Benth.

Bot. Voy. Sulphur [Bentham]: 89 [26 Oct. 1844] (Bentham 1844), “*Crudyā*”. — *Apalatoa oblonga* (Benth.) Taub., *Bot. Centralbl.* 47: 394 [31 Sept 1891] (Taubert 1891). — *Touchiroa oblonga* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 211 [5 Nov. 1891] (Kuntze 1891), “*Touchiroa*”.

Crudia pubescens Spruce ex Benth., *Fl. Bras. [Martius]* 15 (2): 240 [1 Dec. 1870] (Bentham 1870), “*Crudyā*”. — *Apalatoa pubescens* (Spruce ex Benth.) Taub., *Bot. Centralbl.* 47: 394 [31 Sept 1891] (Taubert 1891).

VERNACULAR NAMES. — Pa: kasigu, wap-kamwi, wap-puvemna • Ka: atapalan • Wp: ipewi pilá • Nt: kiabisi • Br: arapari-rana, ipêda-várzea, ipê-rana, rim-de-paca.



FIG. 30. — Leguminosae: **A**, *Calliandra hymenaeodes* (Rich.) Benth. (M.-F. Prévost et al. 4695); **B**, *Cassia grandis* L.f. (M.-F. Prévost & D. Sabatier 4766); **C**, *Dimorphandra ignea* Ducke (D. Sabatier & S. Gonzalez 5891); **D**, *Erythrina poeppigiana* (Walp.) O.F.Cook; **E**, *Inga pilosula* (Rich.) J.F.Macbr. (D. Sabatier & M.-F. Prévost 4868); **F**, *Inga retinocarpa* Poncey (J.-F. Molino 3328). A-E, © D. Sabatier/IRD; F, © J.-F. Molino/IRD.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, K[K000555166, K000555167]; iso-, K[K000555168]).

INVENTORY DATA (FG). — 1 tree, dbh = 22.9 cm.

[771] *Crudia spicata* (Aubl.) Forsyth f.

Bot. Nomencl.: 247 (Forsyth 1794). — *Apalatoa spicata* Aubl., *Hist. Pl. Guiane* 1: 382 [Jun.-Dec. 1775] (Aublet 1775), “*Opalatoa*” on plate.

NOTE. — *Apalatoa spicata* Aubl. is the conserved type of *Crudia* Schreb.

HERBARIUM DATA (FG). — A single collection, *J.B. Aublet s.n.* (original material BM[BM000826536]).

SIZE. — Up to 40 cm dbh (“1,5 pied de diamètre”) (Aublet 1775, 1: 383).

[772] *Crudia tomentosa* (Aubl.) J.F.Macbr.

Contr. Gray Herb. 59: 20 (Macbride 1919). — *Parivoa tomentosa* Aubl., *Hist. Pl. Guiane* 2: 759 [Jun.-Dec. 1775] (Aublet 1775). — *Crudia parivoa* DC., *Prodr. [A. P. de Candolle]* 2: 520 [mid Nov. 1825] (Candolle 1825), “*Crudyā*”, nom. illeg. superfl. (based on *Parivoa tomentosa*). — *Touchiroa parivoa* Rich. ex DC., *Prodr. [A. P. de Candolle]* 2: 520 [mid Nov. 1825] (Candolle 1825), nom. nud. pro syn. — *Apalatoa tomentosa* (Aubl.) Taub., *Bot. Centralbl.* 47: 394 [31 Sept 1891] (Taubert 1891). — *Touchiroa tomentosa* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 211 [5 Nov. 1891] (Kuntze 1891), “*Touchiroa*”.

VERNACULAR NAMES. — Pa: á-seiminio-priyo, á-seiminio-priyu • Nt: pikin paanga • Br: rim-de-paca.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000826533]).

SIZE. — Brazil, Pará. *D.C. Daly 1601* (MO), 30 m.

Genus *Cynometra* L.

[773] *Cynometra baubiniifolia* Benth.

J. Bot. [Hooker] 2: 99 (Bentham 1840), “*bauhiniaefolia*”. — *Cynometra crassifolia* Benth., *J. Bot. [Hooker]* 2: 100 (Bentham 1840).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *C. Sastre 4146*.

SIZE. — Up to 25 m tall (Silva *et al.* 1998).

[774] *Cynometra hostmanniana* Tul.

Arch. Mus. Hist. Nat. 4: 180 (Tulasne 1844).

VERNACULAR NAMES. — Ka: malako • Wp: kumalaki • Nt: mukulakula, mukwalakwala • Cr: bwa-bago • Br: jutaí-rana.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3315*.

INVENTORY DATA (FG). — 1 tree, dbh = 10 cm.

[775] *Cynometra marginata* Benth.

J. Bot. [Hooker] 2: 100 (Bentham 1840).

Cynometra marginata var. *guianensis* Dwyer, *Ann. Missouri Bot. Gard.* 45 (4): 329 [6 Jan. 1959] (Dwyer 1959).

Cynometra marginata var. *laevis* Dwyer, *Ann. Missouri Bot. Gard.* 45 (4): 329 [6 Jan. 1959] (Dwyer 1959).

VERNACULAR NAMES. — Wn: pokololomilii • Nt: man tsaka.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-J. de Granville 4866*.

SIZE. — Up to 21 m tall (Silva *et al.* 1998).

[776] *Cynometra parvifolia* Tul.

Arch. Mus. Hist. Nat. 4: 181 (Tulasne 1844).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *Service Forestier 4436; D. Sabatier 1009*, “Arbre moyen”

Genus *Dialium* L.

[777] *Dialium guianense* (Aubl.) Sandwith

Lloydia 2 (3): 184 (Sandwith 1939). — *Arouna guianensis* Aubl., *Hist. Pl. Guiane* 1: 16 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wap-kamwi-puvemna • Te: takulu wila • Wp: wila takulu • Nt: pikin loka • Cr: ti-koubari • Br: beiju-de côco, cururu, jutaí-mirim, parajuba, pau-ferro, pororoca.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00680458] designated by Lanjouw & Uititten [1940: 148]; isolecto-, BM[BM000952212], P[P00798483]).

INVENTORY DATA (FG). — 56 trees in 38 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 77$ cm.

Genus *Dicorynia* Benth.

[778] *Dicorynia guianensis* Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 28 (Amshoff 1939).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: ájelik • Ka: ki'elei, kiyelewu, tipulu kiyel'e'u • Te: wadi'i • Wp: aisili • Wn: kuwaluk, kwaluk • Nt: anzili, dobono so, singapeetu • Cr: anjélik, lanjélik • Fr: angélique • Br: angélica-do-Pará, tapaiúna.

HERBARIUM DATA (FG). — 147 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2946*.

INVENTORY DATA (FG). — 1520 trees in 211 plots; $F_{\max} = 6.8\%$; $dbh_{\text{inv}} = 115$ cm.

Genus *Dimorphandra* Schott

[779] *Dimorphandra coccinea* Ducke

J. Wash. Acad. Sci. 25: 197 (Ducke 1935).

VERNACULAR NAMES. — Wp: alapokwa'i • Br: arariúba.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *G. Cremer 14582*.

SIZE. — Brazil, Amazonas. *J.W. Grimes 3126* (MO), “Emergent tree”.

[780] *Dimorphandra cuprea* Sprague & Sandwith

Bull. Misc. Inform. Kew 1932 (8): 402 [1 Nov. 1932] (Sprague & Sandwith 1932).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 4836*.

INVENTORY DATA (FG). — 1 tree, dbh = 47.7 cm.

[781] *Dimorphandra ignea* Ducke
(Fig. 30C)

J. Wash. Acad. Sci. 25: 196 (Ducke 1935).

VERNACULAR NAMES. — Nt: ayee weko.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5055*.

INVENTORY DATA (FG). — 4 trees in 1 plot; dbh_{inv} = 43.6 cm.

[782] *Dimorphandra macrostachya* Benth.
subsp. *glabrifolia* (Ducke) M.F.Silva

Fl. Neotrop. Monogr. 44: 106 [2 Oct. 1986] (Silva 1986).

Dimorphandra glabrifolia Ducke, *J. Wash. Acad. Sci.* 25: 196 (Ducke 1935).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 21691*.

SIZE. — Up to 41 m tall (Barneby & Heald 2002).

[783] *Dimorphandra multiflora* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 85 (Ducke 1922).

VERNACULAR NAMES. — Pa: yit-ára • Br: arariúba.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier 1242*.

SIZE. — Brazil, Amazonas. *W.A. Rodrigues s.n.* (MO), 27 m × 50 cm.

[784] *Dimorphandra polyandra* Benoist

Notul. Syst. (Paris) 3: 272 [7 May 1917] (Benoist 1917).

Dimorphandra hohenkerkii Sprague & Sandwith, *Bull. Misc. Inform. Kew* 1932 (8): 403 [1 Nov. 1932] (Sprague & Sandwith 1932).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: alaulama, aweiko • Nt: ayee weko.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *R. Benoist 138* (holo-, P[P03090205, P03090207, P03090208]; iso-, K[K000555304]).

INVENTORY DATA (FG). — 81 trees in 4 plots; F_{max} = 4.3 %; dbh_{inv} = 114.6 cm.

[785] *Dimorphandra pullei* Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 8 (Amshoff 1939).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: alapokwa'i • Nt: makwebo, makweeblo • Br: fava-uim-amarela.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P. Grenand 1116*.

INVENTORY DATA (FG). — 4 trees in 4 plots; F_{max} < 1 %; dbh_{inv} = 45.4 cm.

Genus *Dinizia* Ducke

[786] *Dinizia excelsa* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 76 (Ducke 1922).

VERNACULAR NAMES. — Wp: peilili • Br: angelim, angelim-pedra (verdadeiro), angelim-vermelho.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville et al. 7911*.

SIZE. — Brazil, Amazonas. *S.A. Mori 21327* (MO), 35 m × 80 cm.

Genus *Diplotropis* Benth.

[787] *Diplotropis martiusii* Benth.

Commentat. Legum. Gen. 24 [June 1837] (Bentham 1837), “*Martiusii*”. — *Boudichia martiusii* (Benth.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 131 (Ducke 1922), “*Martiusii*”.

Dibrachion riparium Spruce ex Benth., *Fl. Bras. [Martius]* 15 (1): 321 [15 Jan. 1862] (Bentham 1862), “*Dibrachium*”, nom. nud. pro syn.

VERNACULAR NAMES. — Br: sapupira-da-várzea.

HERBARIUM DATA (FG). — A single collection, *P.A. Sagot 1037* (P[P03101996]).

SIZE. — Up to 25 m tall (Lima & Aymard, 1999).

[788] *Diplotropis purpurea* (Rich.) Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 43 (Amshoff 1939). — *Tachigali purpurea* Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792).

Dibrachion guianense Tul., *Ann. Sci. Nat., Bot. sér. 2, 20*: 139 (Tulasne 1843), “guianensis”. — *Diplotropis guianensis* (Tul.) Benth., *Fl. Bras. [Martius]* 15 (1): 321 [15 Jan. 1862] (Bentham 1862), “guyanensis”. — *Bowdichia guianensis* (Tul.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 32 (Ducke 1915), “guyanensis”.

VERNACULAR NAMES. — Ka: eleyulu, kinoto epi, malipa wewe, woko isyale • Wp: akusi'i, kusi'i • Wn: jawala mili • Nt: baaka tyabisi, tyabisi • Cr: tchò-dérò • Fr: cœur dehors • Br: sapupira, sucupira-açu, sucupira-preta.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *M.-F. Prévost 4065*.

INVENTORY DATA (FG). — 122 trees in 81 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 75$ cm.

Genus *Dipteryx* Schreb.

[789] *Dipteryx odorata* (Aubl.) Forsyth f.

Bot. Nomencl. 391 (Forsyth 1794). — *Coumarouna odorata* Aubl., *Hist. Pl. Guiane* 2: 740 [Jun.-Dec. 1775] (Aublet 1775), “*Odora*” on plate.

Dipteryx tetraphylla Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 125 [Mar. 1860] (Bentham 1860). — *Coumarouna tetraphylla* (Spruce ex Benth.) Taub., *Bot. Centralbl.* 47: 389 [31 Sept 1891] (Taubert 1891), “*Cumaruna*”. — *Coumarouna odorata* var. *tetraphylla* (Spruce ex Benth.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 322 (Ducke 1925).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Levis *et al.* 2017).

VERNACULAR NAMES. — Ka: coumarou (*fide* Aublet 1775), kalapabosi, kalapapo:si • Te: budu'i • Wp: munu'i e'e • Wn: mapiku • Nt: tonka • Cr: gayak • Fr: coumarou odorant, fève de tonka, gaïac de Cayenne • Br: cumarú, cumarú-verdadeiro, muira-pajé.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.B. Aublets.s.n.* (original material BM[BM000931959], P[P02771302]).

INVENTORY DATA (FG). — 33 trees in 31 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 100$ cm.

[790] *Dipteryx punctata* (S.F.Blake) Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 60 (Amshoff 1939). — *Coumarouna punctata* S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (13): 525 [19 July 1924] (Blake 1924).

Coumarouna trifoliolata Ducke, *Notizbl. Bot. Gart. Berlin-Dahlem* 14: 124 [30 Mar. 1938] (Ducke 1938). — *Dipteryx trifoliolata* (Ducke) Ducke, *Trop. Woods* 61: 7 (Ducke 1940).

VERNACULAR NAMES. — Pa: waikwimna • Ka: kalapapo:si, katulimia • Wp: munu'i sī • Wn: mapiku • Nt: tonka • Cr: gayak-blan • Fr: faux gaïac blanc • Br: cumarú-amarela, cumarurana.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3550*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 95.5$ cm.

Genus *Dussia* Krug & Urb. ex Taub.

[791] *Dussia discolor* (Benth.) Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 50 (Amshoff 1939). — *Geoffroea discolor* Benth., *J. Bot. [Hooker]* 2: 69 (Bentham 1840), “*Geoffroya*”.

Vexillifera micranthera Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 140 (Ducke 1922).

Dussia micranthera (Ducke) Harms, *Repert. Spec. Nov. Regni Veg.* 19: 291 (Harms 1924).

Dussia cayennensis Harms, *Repert. Spec. Nov. Regni Veg.* 19: 293 (Harms 1924).

VERNACULAR NAMES. — Wp: inga láli, inga láni • Cr: moutouchi-marikaj.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J. Martin 6* (lecto-, P[P01817936]), designated by Amshoff [1939: 50]; isolecto-, K[K000535276], P[P01817937, P01817938]).

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 90$ cm.

[792] *Dussia tessmannii* Harms

Notizbl. Bot. Gart. Berlin-Dahlem 9: 972 [15 Nov. 1926] (Harms 1926).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4124*.

INVENTORY DATA (FG). — 11 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 150$ cm.

Genus *Enterolobium* Mart.

[793] *Enterolobium oldemanii* Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 249 (Barneby & Grimes 1996).

VERNACULAR NAMES. — Pa: yuumwi-seine • Nt: titim batibataa • Cr: mal-akasya • Fr: acacia mâle.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2242* (holo-, P[P00199424]; iso-, CAY[CAY028067, CAY028068], G[G00365207], K[K000117958], MO[MO-176370], NY[00002032, 00452497], U[U0142265], US[US00432893, 00610742]).

INVENTORY DATA (FG). — 43 trees in 35 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 69.1$ cm.

[794] *Enterolobium schomburgkii* (Benth.) Benth.

Trans. Linn. Soc. London 30 (3): 599 [10 Apr. 1875] (Bentham 1875). — *Pithecellobium schomburgkii* Benth., *London J. Bot.* 3: 219 (Bentham 1844), “*Pithecolobium? Schomburgkii?*”. — *Feuilleea schomburgkii* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Mimosa wilsonii Standl., *Contr. Arnold Arbor.* 5: 73 (Standley 1933).

VERNACULAR NAMES. — Pa: sakeg-kamwi • Ka: apakanilan • Wp: akiki nami • Wn: alawata uhmot, aliwe puman • Nt: bugu batibataa, kasa • Cr: akasya-fran, bwa-lanmori, préfontenn • Fr: acacia franc • Br: fava-de-rosca, orelha-de-macaco, sucupira-amarela, timbaúba.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2157*.

INVENTORY DATA (FG). — 49 trees in 39 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 99.9$ cm.

[795] *Enterolobium* sp. A

NOTE. — This species resembles *E. schomburgkii*, but has distinctive soft fruit and smaller glabrous, non-papillose leaflets.

HERBARIUM DATA (FG). — A single collection, *M.-F. Prévost & D. Sabatier 4976*.

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 91.2$ cm.

Genus *Eperua* Aubl.

[796] *Eperua bijuga* Mart. ex Benth.

Fl. Bras. [Martius] 15 (2): 226 [1 Dec. 1870] (Bentham 1870).

Eperua bijuga f. *typica* Ducke, *Trop. Woods* 62: 26 (Ducke 1940), nom. inval. (Turland et al. 2018): Art. 24.3.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (P[P03465179]).

SIZE. — Up to 12 m tall (Cowan 1975).

[797] *Eperua falcata* Aubl.

Hist. Pl. Guiane 1: 369 [Jun.-Dec. 1775] (Aublet 1775). — *Panzera falcata* (Aubl.) Willd., *Sp. Pl., ed. 4* 2 (1): 540 [Mar. 1799] (Willdenow 1799). — *Dimorpha falcata* (Aubl.) Sm., *Cycl. [Rees]* 11: (Dimorpha no. 3) [3 Dec. 1808] (Smith 1808).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020), although known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wap, wap-duwó • Ka: paliwi, paliwu, wapa, tyoto amote, vouapa-tabaca (*fide* Aublet 1775), watapa • Te: tapaka'i • Wp: tapaka • Wn: wapa • Nt: bii udu, wapa • Cr: pwa-sab, wapa, wapa-gra • Fr: wapa • Br: apa, apazeiro, espadeira, muirapiranga.

HERBARIUM DATA (FG). — 292 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000952284]).

INVENTORY DATA (FG). — 3623 trees in 150 plots; $F_{\max} = 14.7\%$; $\text{dbh}_{\text{inv}} = 101$ cm.

[798] *Eperua grandiflora* (Aubl.) Baill.

Hist. Pl. [Baillon] 2: 110 [Jan.-Feb. 1870] (Baillon 1870). — *Parivoa grandiflora* Aubl., *Hist. Pl. Guiane* 2: 756 [Jun.-Dec. 1775] (Aublet 1775).

Eperua kourouensis Benoist, *Notul. Syst. (Paris)* 3: 273 [7 May 1917] (Benoist 1917).

NOTES. — Known only from the Guiana Shield. The previous transfer by Bentham (1865: 580) is invalid because he only cited *Parivoa grandiflora* Aubl. under *Eperua*, without associating the genus name and the epithet. Although Baillon also cited this species under *Parivoa grandiflora* (Baillon 1870: 111), he clearly made the combination “[*Eperua (Parivoa) grandiflora*]” on figs. 81 & 82. Eventually, Bentham (1870: 226) also published the new combination, but a few months after Baillon.

VERNACULAR NAMES. — Pa: wap-seinô, wap-waikmiunô • Wp: kupeiwa, tapaka • Nt: gaan bii udu, loka bii udu • Cr: wapa-koubari • Fr: wapa courbaril • Br: apa.

HERBARIUM DATA (FG). — 116 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (holo-, BM[BM000952282]; iso-, P-JU[P00835933], P[P00835934]).

INVENTORY DATA (FG). — 1219 trees in 100 plots; $F_{\max} = 5.5\%$; $\text{dbh}_{\text{inv}} = 86.3$ cm.

[799] *Eperua jenmanii* Oliv.

Hooker's Icon. Pl. 20 [ser. 3, 10]: t 1955 [Apr. 1891] (Oliver 1891).

Eperua stipulata Kleinhoonte, *Recueil Trav. Bot. Néerl.* 30: 171 [July 1933] (Kleinhoonte 1933).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *J. Martin s.n.* (BM[BM000065079]).

SIZE. — Up to 70 cm dbh (Cowan 1975).

[800] *Eperua rubiginosa* Miq.

Stirp. Surinam. Select.: 12 [“1850” publ. Mar. 1851] (Miquel 1851).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wap-seinô • Ka: paliwi, paliwu, wapa, watapa • Te: tapaka • Wp: tapaka • Wn: wapa • Nt: tetey bii udu, wataa bii udu • Cr: wapa-larivyé, wapa-rouj • Fr: wapa rivière • Br: apa, apazeiro, espadeira, muirapiranga.

HERBARIUM DATA (FG). — 62 collections at CAY. Sel. exs.: *D. Sabatier 1018*.

INVENTORY DATA (FG). — 390 trees in 31 plots; $F_{\max} = 9.9\%$; $\text{dbh}_{\text{inv}} = 94.2$ cm.

[801] *Eperua schomburgkiana* Benth.

Fl. Bras. [Martius] 15 (2): 226 [1 Dec. 1870] (Bentham 1870).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: tapaka.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *B.M. Boom & S.A. Mori 1805*.

SIZE. — Up to 37 m tall (Barneby & Heald 2002).

Genus *Erythrina* L.

[802] *Erythrina fusca* Lour.

Fl. Cochinch. 2: 427 [Sep. 1790] (Loureiro 1790). — *Corallodendron fuscum* (Lour.) Kuntze, *Revis. Gen. Pl.* 1: 172 [5 Nov. 1891] (Kuntze 1891).

Erythrina glauca Willd., *Neue Schriften Ges. Naturf. Freunde Berlin* 3: 428 (Willdenow 1801). — *Duchassaingia glauca* (Willd.) Walp., *Linnaea* 23: 742 ["1850" publ. Jan. 1851] (Walpers 1851). — *Corallodendron glaucum* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 172 [5 Nov. 1891] (Kuntze 1891).

Erythrina patens Moc. & Sessé ex DC., *Prodri. [A. P. de Candolle]* 2: 414 [mid Nov. 1825] (Candolle 1825). — *Corallodendron patens* (Moc. & Sessé ex DC.) Kuntze, *Revis. Gen. Pl.* 1: 173 [5 Nov. 1891] (Kuntze 1891).

Erythrina ovalifolia Roxb., *Fl. Ind., ed. 1832*, 3: 254 (Roxburgh 1832). — *Duchassaingia ovalifolia* (Roxb.) Walp., *Linnaea* 23: 742 ["1850" publ. Jan. 1851] (Walpers 1851). — *Corallodendron ovalifolia* (Roxb.) Kuntze, *Revis. Gen. Pl.* 1: 173 [5 Nov. 1891] (Kuntze 1891).

Erythrina caffra Blanco, *Fl. Filip. [FM. Blanco]*, ed. 2: 394 (Blanco 1845), nom. illeg. hom., non Thunb. (Thunberg 1800).

Erythrina atrosanguinea Ridl., *J. Straits Branch Roy. Asiat. Soc.* 59: 93 [July 1911] (Ridley 1911).

Erythrina fusca var. *inermis* Pulle, *Nova Guinea* 8 (2): 651 (Pulle 1912).

Erythrina moelebei Vieill. ex Guillaumin & Beauvis., *Ann. Soc. Bot. Lyon* 38: 13 (Guillaumin & Beauvisage 1914), nom. nud.

Erythrina fusca var. *inermis* Rock, *Legum. Pl. Hawaii*: 188 (Rock 1920), nom. illeg. hom., non Pulle (1912).

NOTES. — A species restricted to wet savannas. *Duchassaingia ovalifolia* (Roxb.) Walp. is indirectly based on *Erythrina ovalifolia* Roxb., through citation of Roxburgh's plate in Wight (1840, 1: pl. 247).

VERNACULAR NAMES. — Pa: mitiku • Ka: asiwakan • Wn: matata • Cr: zimortel • Br: açacurana, mulungu.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F. Prévost 1207*, 15 m tall.

[803] *Erythrina poeppigiana* (Walp.) O.F.Cook
(Fig. 30D)

U.S.D.A. Div. Bot. Bull. 25: 57 (Cook 1901). — *Micropteryx poeppigiana* Walp., *Linnaea* 23: 740 ["1850" publ. Jan. 1851] (Walpers 1851). — *Erythrina micropteryx* Poepp. ex Urb., *Symb. Antill. [Urban]* 1 (2): 327 [10 Apr. 1899] (Urban 1899), nom. illeg. superfl. (based on *Micropteryx poeppigiana*).

Erythrina amasia Spruce, *J. Proc. Linn. Soc., Bot.* 3: 202 [Apr. 1859] (Spruce 1859), "Amasia".

Erythrina pisamo Posada-Ar., *Estudios Científicos*: 122 (Posada-Arango 1909), "Písamo".

Erythrina darienensis Standl., *Contr. U.S. Natl. Herb.* 18 (3): 108 [11 Feb. 1916] (Standley 1916).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *O. Tostain 2818*.

SIZE. — Up to 30 m tall (Neill 1999).

Genus *Guianodendron* Sch.Rodr. & A.M.G.Azevedo

[804] *Guianodendron praeclarum*
(Sandwith) Sch.Rodr. & A.M.G.Azevedo

Novon 16 (1): 130 [25 May 2006] (Schütz Rodrigues & Azevedo 2006). — *Sweetia praeclera* Sandwith, *Contr. Gray Herb.* 165: 25 (Sandwith 1947). — *Acosmum praeclarum* (Sandwith) Yakovlev, *Notes Roy. Bot. Gard. Edinburgh* 29 (3): 355 (Yakovlev 1969).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 24980*.

SIZE. — Up to 50 cm dbh (Rodrigues & Tozzi 2006).

Genus *Hydrochorea* Barneby & J.W.Grimes

[805] *Hydrochorea corymbosa*
(Rich.) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 27 (Barneby & Grimes 1996). — *Mimosa corymbosa* Rich., *Actes Soc. Hist. Nat. Paris* 1: 113 [Oct. 1792] (Richard 1792). — *Pithecellobium corymbosum* (Rich.) Benth., *Trans. Linn. Soc. London* 30 (3): 587 [10 Apr. 1875] (Bentham 1875), "Pithecolobium", nom. illeg. hom., non Benth. (Bentham 1844). — *Feuilleea corymbosa* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891). — *Samanea corymbosa* (Rich.) Pittier, *Arb. Arbust. Venez.* 4-5: 55 (Pittier 1925). — *Arthrosamea corymbosa* (Rich.) Kleinhoonte, *Fl. Suriname* 2 (2): 327 (Kleinhoonte 1940). — *Cathormion corybosum* (Rich.) Burkart, *Darwiniana* 13: 446 (Burkart 1964). — *Albizia corymbosa* (Rich.) G.P.Lewis & P.E.Owen, *Legumes Ilha de Maracá*: 40 (Lewis & Owen 1989).

Pithecellobium corymbosum Benth., *London J. Bot.* 3: 221 (Bentham 1844), "Pithecolobium".

Pithecellobium subcorymbosum Hoehne, *Comm. Lin. Telegr., Bot.* 45 (8): 18 (Hoehne 1919), "Pithecolobium".

Pithecellobium corymbosum var. *longipes* Spruce ex Ducke, *Bol. Técn. Inst. Agron. N.* 18: 37 (Ducke 1949), nom. nud.

NOTES. — Bentham (1844: 221) described *Pithecellobium corymbosum* without reference to *Mimosa corymbosa* Rich. and with a different type: the protologue cites a collection of J. Martin, whereas the type of *M. corymbosa* is a Leblond specimen (the species is described in an article dedicated to Leblond's collections). The type of *P. corymbosum* (1844) could not be located. In 1875, Bentham again listed *P. corymbosum*, with reference to his 1844 taxon (Bentham 1875: 587). However, as he placed *M. corymbosa* in synonymy, this time he created a new combination, as well as a later homonym.

VERNACULAR NAMES. — Pa: karevru • Ka: kalaipe'u, kalape'u • Wp: kalaipe'i • Wn: kaju • Nt: kabanafoo • Cr: akasya, tanmaren-granbwa • Br: paricazinho, saboeiro-da-várzea.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* [in Herb. Richard] (type P[P02142909]).

SIZE. — Guyana. *T.W. Henkel et al. 3133* (MO), 12 m.

Genus *Hymenaea* L.

[806] *Hymenaea courbaril* L.

Sp. Pl. 2: 1192 [1 May 1753] (Linnaeus 1753).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: simig, simii • Ka: simili, šimili, simili tabile, simili tamune, simili tubulu • Te: tsimini • Wp: talala, watala, yita'i • Wn: mepu, mépu • Nt: loka • Cr: kaka-chien, koubari • Fr: courbaril • Br: jatauba, jatobá, jutaí, jutaí-açu.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *D. Sabatier 3575*.

INVENTORY DATA (FG). — 22 trees in 17 plots; $F_{\max} < 1\%$; dbh_{inv} = 111.7 cm.

[807] *Hymenaea* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier 5572*.

INVENTORY DATA (FG). — 1 tree, dbh = 30.6 cm.

Genus *Hymenolobium* Benth.

[808] *Hymenolobium excelsum* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 38 (Ducke 1915).

VERNACULAR NAMES. — Wp: pana'i • Nt: geli tyabisi • Cr: sen-marten-gri • Fr: saint-martin gris • Br: angelim-comum, angelim-da-mata.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.-J. de Granville 16824*.

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 158.5 cm.

[809] *Hymenolobium flavum* Kleinhoonte

Recueil Trav. Bot. Néerl. 22: 400 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

VERNACULAR NAMES. — Pa: sakeg • Ka: eleyulu, malipa wewe • Wp: pana'i • Nt: geli tyabisi • Cr: panao, sen-marten-jonn • Fr: angelin, saint-martin jaune • Br: acapurana, angelim, morcegueira, sapupira-da-várzea.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3262*.

INVENTORY DATA (FG). — 32 trees in 27 plots; $F_{\max} < 1\%$; dbh_{inv} = 120 cm.

[810] *Hymenolobium heterocarpum* Ducke

Trop. Woods 47: 6 (Ducke 1936).

VERNACULAR NAMES. — Br: angelim.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2876*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 65 cm.

[811] *Hymenolobium petraeum* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 36 (Ducke 1915).

VERNACULAR NAMES. — Ka: eleyulu • Nt: geli tyabisi • Cr: sem-marten-jonn • Fr: saint-martin jaune • Br: angelim-amarelo, angelim-da-mata, angelim-pedra, murarena.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman 3188; S.A. Mori et al. 23694*, 20 m × 25 cm.

[812] *Hymenolobium pulcherrimum* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 37 (Ducke 1915).

VERNACULAR NAMES. — Wp: wila paye • Nt: geli tyabisi • Br: angelim-pedra, sapupira-amarela.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3007*.

INVENTORY DATA (FG). — 1 tree, dbh = 18.8 cm.

Genus *Inga* Mill.

[813] *Inga acreana* Harms

Notizbl. Königl. Bot. Gart. Berlin 6: 298 [30 June 1915] (Harms 1915).

Inga myriocephala Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 184 [30 Oct. 1916] (Pittier 1916).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier 1052*.

INVENTORY DATA (FG). — 24 trees in 16 plots; $F_{\max} = 1.4\%$; dbh_{inv} = 34.1 cm.

[814] *Inga acrocephala* Steud.

Flora 26 (45): 759 [17 Dec. 1843] (Steudel 1843). — *Feuilleea acrocephala* (Steud.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga brevipedicellata Harms, *Repert. Spec. Nov. Regni Veg.* 19: 62 (Harms 1923).

VERNACULAR NAMES. — Pa: miumiu-akamá-arib, miumiu-akamá-aribiné • Wp: inga pini, inga u po'i • Nt: weko • Cr: pwa-sikré • Br: ingá.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2464*.

INVENTORY DATA (FG). — 86 trees in 44 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 70\text{ cm}$.

[815] *Inga alata* Benoist

Bull. Mus. Natl. Hist. Nat. 27: 198 (Benoist 1921).

VERNACULAR NAMES. — Pa: avukun • Wp: inga masulapa, masulapa • Cr: pwa-sikré • Br: ingá.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *E.M. Mélinon 8a*, 1845 (holo-, P[P01818410]; iso-, P[P01818411, P01818412]).

INVENTORY DATA (FG). — 29 trees in 18 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 73\text{ cm}$.

[816] *Inga alba* (Sw.) Willd.

Sp. Pl., ed. 4 4 (2): 1013 [Apr. 1806] (Willdenow 1806). — *Mimosa alba* Sw., *Prod. [Swartz]* 85 [20 Jun.-29 July 1788] (Swartz 1788). — *Feuilleea alba* (Sw.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga fraxinea Willd., *Sp. Pl.*, ed. 4 4 (2): 1019 [Apr. 1806] (Willdenow 1806). — *Mimosa fraxinea* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 44 [3 Sep. 1810] (Poirer 1810).

Mimosa alba Vahl, *Eclog. Amer.* 3: 31 (Vahl 1807), nom. illeg. hom., non Sw. (Swartz 1788).

Inga thyrsoides Desv., *J. Bot. Agric.* 3: 71 (Desvaux 1814).

Inga aggregata G.Don, *Gen. Hist.* 2: 391 [Oct. 1832] (Don 1832). — *Feuilleea aggregata* (G.Don) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga spruceana Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 239 (Bentham 1850).

Inga parviflora Sagot ex Benth., *Trans. Linn. Soc. London* 30 (3): 602 [10 Apr. 1875] (Bentham 1875), nom. nud. pro syn.

Inga carachensis Pittier, *Arb. Arbust. Venez.* 9-10: 106 [Dec. 1929] (Pittier 1929).

Inga altissima Ducke, *Arch. Inst. Biol. Veg.* 4 (1): 4 [June 1938] (Ducke 1938).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: avukun, avukun-duwō • Ka: abuluguni, apulukuni • Te: pulukuni • Wp: inga sisi, inga u, sisi, sisi si • Wn: apulukun • Nt: kooponyo, lebi weko • Cr: bougouni, bwa-pagód • Fr: bougouni • Br: ingá-branca, ingá-xixica.

HERBARIUM DATA (FG). — 95 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (type BM[BM000541200]).

INVENTORY DATA (FG). — 228 trees in 104 plots; $F_{\max} = 3.6\%$; $dbh_{\text{inv}} = 100\text{ cm}$.

[817] *Inga auristellae* Harms

Notizbl. Königl. Bot. Gart. Berlin 6: 298 [30 June 1915] (Harms 1915).

VERNACULAR NAMES. — Wp: inga sili • Cr: pwa-sikré.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4282*.

INVENTORY DATA (FG). — 11 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 33.1\text{ cm}$.

[818] *Inga bourgonii* (Aubl.) DC.

Prod. [A. P. de Candolle] 2: 434 [mid Nov. 1825] (Candolle 1825). — *Mimosa bourgonii* Aubl., *Hist. Pl. Guiane* 2: 941 [Jun.-Dec. 1775] (Aublet 1775). — *Feuilleea bourgonii* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga assimilis Miq., *Linnaea* 19: 130 ["1847" publ. Feb. 1846] (Miquel 1846).

Inga tapajozensis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 13 (Ducke 1925).

Inga apta J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (3.1): 15 (Macbride 1943).

NOTE. — "Bourgonii" is a vernacular name, and thus should not be changed to "bourgonii".

VERNACULAR NAMES. — Pa: avukun, miumiu-platno-purubumna • Ka: abuluguni, apulukuni, posindyo • Te: pulukuni • Wp: inga sisi, inga ū, inga u sówî, kulikuli inga, sisi • Wn: tuli • Nt: djinaati • Cr: bougouni, bwa-pagód • Br: ingá-xixi, ingá-xixica.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000541054]).

INVENTORY DATA (FG). — 38 trees in 8 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 43\text{ cm}$.

[819] *Inga brachystachys* Ducke

Trop. Woods 90: 12 (Ducke 1947). — *Inga brachystachya* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 54 (Ducke 1922), nom. illeg. hom., non DC. (Candolle 1825) [synonym of *Abarema brachystachya* (DC.) Barneby & J.W. Grimes].

VERNACULAR NAMES. — Wp: inga mulua ya, mulua ya.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.-F. Molino 1107*.

INVENTORY DATA (FG). — 17 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 67.7\text{ cm}$.

[820] *Inga capitata* Desv.

J. Bot. Agric. 3: 71 (Desvaux 1814). — *Feuilleea capitata* (Desv.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Mimosa pacay Aubl., *Hist. Pl. Guiane* 2: 946 [Jun.-Dec. 1775] (Aublet 1775).

Inga albicans Walp., *Linnaea* 14: 298 (Walpers 1840).

Inga calycina Salzm. ex Benth., *Trans. Linn. Soc. London* 30 (3): 611 [10 Apr. 1875] (Bentham 1875), *nom. nud. pro syn.*

Inga peduncularis Mart. ex Benth., *Trans. Linn. Soc. London* 30 (3): 611 [10 Apr. 1875] (Bentham 1875), *nom. nud. pro syn.*

Inga capitata var. *brevicalyx* Benth., *Fl. Bras. [Martius]* 15 (2): 476 [1 July 1876] (Bentham 1876).

Inga capitata var. *tenuior* Benth., *Fl. Bras. [Martius]* 15 (2): 476 [1 July 1876] (Bentham 1876).

Inga falcistipula Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 56 (Ducke 1922).

Inga capuchoi Standl., *Trop. Woods* 33: 12 (Standley 1933).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Aublet (1775, 2: 946) mistakenly identified his *Mimosa pacay* as Frézier's pre-Linnean "*Inga peruviana*" (Frézier 1732: 155, pl. 24), hence the epithet referring to the vernacular name "pacay" recorded by Frézier. According to Frézier's plate, his plant had 4-angular pods, leaves with terete rachis and 4-5 pairs of leaflets. These features are not found in *Inga capitata*, but rather could correspond to *I. nobilis* Willd., which is among the several *Inga* species that are called "pacae" in Peru.

VERNACULAR NAMES. — Pa: miumiu-platno • Ka: abuluguni, apulukuni • Te: inga átā • Wp: inga kala, inga mulua ya, mulua ya • Wn: ilakíptre, tuli • Nt: baaka weko • Cr: pwa-sikré-krapo • Br: ingá-branca, ingá-costella, ingá-de-veado, ingá-facão, ingá-ferro.

HERBARIUM DATA (FG). — 54 collections at CAY. Sel. exs.: *A.N. Desvaux s.n.* (type P[P01818387]).

INVENTORY DATA (FG). — 115 trees in 54 plots; $F_{\max} = 1.9\%$; $dbh_{inv} = 36$ cm.

[821] *Inga cayennensis* Sagot ex Benth.

Trans. Linn. Soc. London 30 (3): 626 [10 Apr. 1875] (Bentham 1875). — *Feuilleea cayennensis* (Sagot ex Benth.) Kuntze, *Revis. Gen. Pl. 1:* 187 [5 Nov. 1891] (Kuntze 1891).

Inga dysantha Benth., *Trans. Linn. Soc. London* 30 (3): 626 [10 Apr. 1875] (Bentham 1875).

Feuilleea sessiliflora Kuntze, *Revis. Gen. Pl. 1:* 186 [5 Nov. 1891] (Kuntze 1891).

Inga cayennensis var. *sessiliflora* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 60 (Ducke 1922), "forma *sessiliflora* Ducke n. var.".

Inga aria J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.1): 16* (Macbride 1943).

NOTES. — *P.A. Sagot 164* includes specimens dated from 1855 and 1857. Pennington (1997: 631) designated as lectotype a specimen at K dated 1855 (K000328430), but cited an isolectotype at NY[barcode: 00002266] which is dated 1857, so is in fact a syntype. Also, two specimens at P labelled syntypes are dated 1855, so are in fact isolectotypes.

VERNACULAR NAMES. — Ka: padyawa, payawa, waiki • Nt: weko • Cr: pwa-sikré • Fr: pois sucré • Br: ingá-cabeludo, ingá-verdadeiro.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *PA. Sagot 164*, Aug. 1855 (lecto-, K[K000328430], designated by Pen-

nington [1997: 631]; isolecto-, B[not seen, photo F neg. N° 1075], P[P01818384, P01818385]).

INVENTORY DATA (FG). — 54 trees in 39 plots; $F_{\max} = 1.6\%$; $dbh_{inv} = 25.5$ cm.

[822] *Inga cordatoalata* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 53 (Ducke 1922).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 23276*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 21.6$ cm.

[823] *Inga cylindrica* (Vell.) Mart.

Flora 20 (2, Beibl.): 114 (Martius 1837). — *Mimosa cylindrica* Vell., *Fl. Flumin. Icon.* 11: t. 9 ["1827" publ. 29 Oct. 1831] (Vellozo 1831). — *Feuilleea cylindrica* (Vell.) Kuntze, *Revis. Gen. Pl. 1:* 187 [5 Nov. 1891] (Kuntze 1891).

Inga polystachya Benth., *London J. Bot.* 4: 587 (Bentham 1845).

Inga tenuifolia Benth., *London J. Bot.* 4: 587 (Bentham 1845). — *Feuilleea tenuifolia* (Benth.) Kuntze, *Revis. Gen. Pl. 1:* 189 [5 Nov. 1891] (Kuntze 1891).

Inga albicoria Poncy, *Bull. Mus. Natl. Hist. Nat., B, Adansonia* 18 (1-2): 67 (Poncy 1996).

VERNACULAR NAMES. — Br: ingá-branca, ingá-feijão.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2143*.

INVENTORY DATA (FG). — 50 trees in 28 plots; $F_{\max} = 1.3\%$; $dbh_{inv} = 58.9$ cm.

[824] *Inga disticha* Benth.

J. Bot. [Hooker] 2: 143 (Bentham 1840). — *Feuilleea disticha* (Benth.) Kuntze, *Revis. Gen. Pl. 1:* 187 [5 Nov. 1891] (Kuntze 1891).

Inga crevauxii Sagot, *Ann. Sci. Nat., Bot. sér. 6, 13:* 331 (Sagot 1882).

VERNACULAR NAMES. — Pa: miumiu-kasuminio • Ka: alawata bul-sulukulu, alawata posulukulu • Te: inga tupa'it • Wp: inga takwā u, inga tupewi • Wn: alawata pupot • Nt: babun weko, djakaasi • Cr: pwa-sikré • Br: ingaranara.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *J.N. Crevaux s.n.* (type of *Inga crevauxii*: P[P01818367, P01818368]).

INVENTORY DATA (FG). — 1 tree, $dbh = 13.2$ cm.

[825] *Inga edulis* Mart.

Flora 20 (2, Beibl.): 113 (Martius 1837). — *Feuilleea edulis* (Mart.) Kuntze, *Revis. Gen. Pl. 1:* 187 [5 Nov. 1891] (Kuntze 1891).

Mimosa yngae Vell., *Fl. Flumin. Icon.* 11: t. 3 ["1827" publ. 29 Oct. 1831] (Vellozo 1831), *nom. illeg. hom., non M. inga* L. (Linnaeus

1753). — *Inga yngaa* (Vell.) J.W.Moore, *Occas. Pap. Bernice Pauahi Bishop Mus.* 10 (19): 6 (Moore 1934).

Inga scabriuscula Benth., *London J. Bot.* 4: 606 (Bentham 1845). — *Feuilleea scabriuscula* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga conferta Benth., *London J. Bot.* 4: 620 (Bentham 1845). — *Feuilleea conferta* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga benthamiana Meisn., *Linnæa* 21: 253 (Meisner 1848).

Inga uncinata Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 630 [10 Apr. 1875] (Bentham 1875), *nom. nud. pro syn.*

Inga scabriuscula var. *villosior* Benth., *Fl. Bras. [Martius]* 15 (2): 497 [1 July 1876] (Bentham 1876).

Inga edulis var. *parviflora* Mart., *Fl. Bras. [Martius]* 15 (2): 498 [1 July 1876] (Martius 1876).

Inga complanata Amshoff, *Natuuruw. Stud. Suriname & Curaçao* 2: 39 (Amshoff 1948).

Inga chorrerana T.S.Elias, *Bot. Mus. Leafl.* 24 (8): 202 [9 Apr. 1976] (Elias 1976); *pro parte quoad specimina in flore.*

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). *Mimosa yngaa* Vell. and *M. inga* L. have been ruled as confusable by the Nomenclature Committee for Vascular Plants (Applequist 2019). The former is therefore to be treated as a later homonym.

VERNACULAR NAMES. — Pa: miiumiu-akamā-arib, miiumiu-akamā-arinē, miiumiu-maoksi-arib (cult. form) • Ka: padyawa, paidyawa, payawa • Te: pailawa • Wp: inga wasa (cult. form), inga yiwa puku, inga yowa puku • Wn: tuli • Nt: tetey weko • Cr: pwa-sikré • Fr: pois sucré • Br: ingá-cipó, rabo-de-mico.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *T.D. Pennington et al. 13867*.

INVENTORY DATA (FG). — 25 trees in 9 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 41.7\text{ cm}$.

[826] *Inga fanchoniana* Poncy

Bull. Mus. Natl. Hist. Nat., B, Adansonia 5 (1): 103 (Poncy 1983).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F. Prévost 478* (holo-, P[P01818350, P01818351, P01818352]; iso-, CAY[CAY028077, CAY028078, CAY028079, CAY028080], MO[MO-1281471], NY[00002291, 00002292], U[U0008512]).

INVENTORY DATA (FG). — 28 trees in 22 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 67.4\text{ cm}$.

[827] *Inga fastuosa* (Jacq.) Willd.

Sp. Pl., ed. 4 4 (2): 1014 [Apr. 1806] (Willdenow 1806). — *Mimosa fastuosa* Jacq., *Fragm. Bot.*: 15 (Jacquin 1800). — *Feuilleea fastuosa* (Jacq.) Kuntze, *Revis. Gen. Pl.* 1: 184, 187 [5 Nov. 1891] (Kuntze 1891).

Inga venosa Griseb. ex Benth., *Trans. Linn. Soc. London* 30 (3): 623 [10 Apr. 1875] (Bentham 1875). — *Feuilleea venosa* (Griseb. ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 184, 189 [5 Nov. 1891] (Kuntze 1891).

Inga longituba Harms, *Repert. Spec. Nov. Regni Veg.* 13: 420 [31 Dec. 1914] (Harms 1914).

Inga guaremalensis Pittier, *Arb. Arbust. Venez.* 1: 5 (Pittier 1921).

VERNACULAR NAMES. — Nt: uman feysi finga.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5722*.

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 54.5\text{ cm}$.

[828] *Inga flagelliformis* (Vell.) Mart.

Flora 20 (2, Beibl.): 112 (Martius 1837). — *Mimosa flagelliformis* Vell., *Fl. Flumin. Icon.* 11: t. 27 ["1827" publ. 29 Oct. 1831] (Vel-lozo 1831). — *Feuilleea flagelliformis* (Vell.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4226*.

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 30.7\text{ cm}$.

[829] *Inga graciliflora* Benth.

London J. Bot. 4: 582 (Bentham 1845). — *Feuilleea graciliflora* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga graciliflora var. *peruviana* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (3.1): 24 (Macbride 1943).

VERNACULAR NAMES. — Ka: salala.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3346*.

INVENTORY DATA (FG). — 18 trees in 17 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 37.2\text{ cm}$.

[830] *Inga gracilifolia* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 52 (Ducke 1922).

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2611*.

INVENTORY DATA (FG). — 48 trees in 33 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 92\text{ cm}$.

[831] *Inga grandiflora* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 59 (Ducke 1922). — *Inga crassiflora* Ducke, *Trop. Woods* 90: 12 (Ducke 1947), *nom. illeg. superfl.* (based on *Inga grandiflora*).

NOTES. — Ducke proposed *Inga crassiflora* as a replacement name for his *I. grandiflora*, because of the existence of *I. grandiflora* Wal-

lich (1831, n. 5285). However, being a *nomen nudum*, *I. grandiflora* Wallich was not validly published, and thus is not to be taken in consideration for priority purposes.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4939*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 18.3$ cm.

[832] *Inga heterophylla* Willd.

Sp. Pl., ed. 4 4 (2): 1020 [Apr. 1806] (Willdenow 1806). — *Feuilleea heterophylla* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891). — *Mimosa parae* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 44 [3 Sep. 1810] (Poiret 1810), “*Para*”, nom. illeg. superfl. (based on *Inga heterophylla*).

Inga umbellata G.Don, *Gen. Hist.* 2: 391 [Oct. 1832] (Don 1832), nom. illeg. hom., non (Vahl) Willd. (Willdenow 1806).

Inga protracta Steud., *Flora* 26 (45): 758 [17 Dec. 1843] (Steudel 1843).

Inga stenocarpa Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 603 [10 Apr. 1875] (Bentham 1875), nom. nud. pro syn.

Inga vouapifolia Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 603 [10 Apr. 1875] (Bentham 1875), “*vouapaefolia*”, nom. nud. pro syn.

Inga mapiriensis Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 174 [30 Oct. 1916] (Pittier 1916).

VERNACULAR NAMES. — Ka: kulisili wokulu, tulili • Wp: inga sili • Cr: pwa-sikré • Br: ingá-ferradura, ingazinho.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *Service Forestier 4005*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.2 cm.

[833] *Inga huberi* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 49 (Ducke 1922).

VERNACULAR NAMES. — Pa: miumiu-avukun-kamwi, miumiu-maoksi-arib-duwē, miumiu-sababuyu.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2718*.

INVENTORY DATA (FG). — 194 trees in 76 plots; $F_{\max} = 2.3\%$; $\text{dbh}_{\text{inv}} = 66.2$ cm.

[834] *Inga ingoides* (Rich.) Willd.

Sp. Pl., ed. 4 4 (2): 1012 [Apr. 1806] (Willdenow 1806). — *Mimosa ingoides* Rich., *Actes Soc. Hist. Nat. Paris* 1: 113 [Oct. 1792] (Richard 1792). — *Feuilleea ingoides* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891), “*ingoides*”.

Inga merianae Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 113 [Aug.-Sep. 1842] (Splitgerber 1842).

Inga laxiflora Benth., *London J. Bot.* 4: 617 (Bentham 1845). — *Feuilleea sesuya* Kuntze, *Revis. Gen. Pl.* 1: 186 [5 Nov. 1891] (Kuntze 1891), nom. illeg. superfl. (based on *Inga laxiflora*).

Inga babiensis Benth., *London J. Bot.* 4: 618 (Bentham 1845). — *Feuilleea babiensis* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891), nom. illeg. hom., non Kuntze [1891: 184, nom. nov. pro *Pithecellobium foliolosum* Benth. (Bentham 1844)].

Inga galibica Duchass. & Walp., *Linnæa* 23: 747 [“1850” publ. Jan. 1851] (Duchassaing & Walpers 1851).

VERNACULAR NAMES. — Ka: padyawa, paidyawa, payawa, waiki (fruit), wala potili • Wn: tulisimé • Nt: weko • Cr: pwa-sikré • Fr: pois sucré.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (type P-JU, not seen); *M.-F. Prévost 3756*, dbh 12 cm.

[835] *Inga jenmanii* Sandwith

Bull. Misc. Inform. Kew 1931 (7): 368 [21 July 1931] (Sandwith 1931), “*Jenmani*”.

Inga sertulifera var. *minor* Benth., *London J. Bot.* 4: 581 (Bentham 1845).

NOTES. — A synonym of *Inga sertulifera* subsp. *leptopus* (Benth.) T.D.Penn. for Pennington (1997: 234), but we agree with Barneby et al. (2011) that it should be considered a distinct taxon, all the more so because, according to the former author, this subspecies is not supposed to be present in French Guiana.

VERNACULAR NAMES. — Pa: miumiu-asiru.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3305*.

INVENTORY DATA (FG). — 106 trees in 62 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 43.8$ cm.

[836] *Inga lateriflora* Miq.

Linnæa 19: 131 [“1847” publ. Feb. 1846] (Miquel 1846). — *Feuilleea lateriflora* (Miq.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga parviflora Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 602 [10 Apr. 1875] (Bentham 1875), nom. nud. pro syn.

Inga lateriflora var. *latior* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 50 (Ducke 1922).

VERNACULAR NAMES. — Ka: tulili.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2399*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 24.4$ cm.

[837] *Inga leiocalycina* Benth.

London J. Bot. 4: 598 (Bentham 1845). — *Feuilleea leiocalycina* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga multiflora Benth., *London J. Bot.* 4: 598 (Bentham 1845).

Inga yunckeri Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 9 (4): 296 (Standley 1940).

VERNACULAR NAMES. — Ka: apipyoloi • Wp: inga sōwī, masulapa pe u, masulapa pilá, paku inga • Cr: pwa-sikré • Br: ingáí.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2818*.

INVENTORY DATA (FG). — 14 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{inv} = 43$ cm.

[838] *Inga lomatophylla* (Benth.) Pittier

Contr. U.S. Natl. Herb. 18 (5): 195 [30 Oct. 1916] (Pittier 1916). — *Inga speciosa* var. *lomatophylla* Benth., *Trans. Linn. Soc. London* 30 (3): 620 [10 Apr. 1875] (Bentham 1875). — *Inga amazonica* var. *lomatophylla* (Benth.) L.Cárdenas, *Ernstia* 51: 1 (Cárdenas 1989).

Feuilleea speciosa Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891). — *Inga speciosa* Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 620 [10 Apr. 1875] (Bentham 1875), nom. illeg. hom., non M.Martens & Galeotti (1843).

Inga amazonica L.Cárdenas, *Ernstia* 51: 1 (Cárdenas 1989).

Inga speciosa var. *membranacea* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 59 (Ducke 1922). — *Inga amazonica* var. *membranacea* (Ducke) L.Cárdenas, *Ernstia* 51: 1 (Cárdenas 1989).

Inga speciosa var. *bracteifera* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 17 (Ducke 1925). — *Inga amazonica* var. *bracteifera* (Ducke) L.Cárdenas, *Ernstia* 51: 1 (Cárdenas 1989). — *Inga lomatophylla* var. *bracteifera* (Ducke) Poncy, *Bull. Mus. Natl. Hist. Nat.*, B., *Adansonia* 13 (3-4): 152 (Poncy 1991).

NOTES. — Contrary to Pennington's claim (Pennington 1997), *Inga amazonica* L.Cárdenas technically is not illegitimate. *Feuilleea speciosa* Kuntze is based on the illegitimate *I. speciosa* Spruce ex Benth.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *M.-F. Prévost 1148*.

INVENTORY DATA (FG). — 30 trees in 25 plots; $F_{\max} < 1\%$; $dbh_{inv} = 52.7$ cm.

[839] *Inga longiflora* Spruce ex Benth.

Trans. Linn. Soc. London 30 (3): 620 [10 Apr. 1875] (Bentham 1875). — *Feuilleea ignota* Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891), nom. illeg. superfl. (based on *Inga longiflora*).

Inga tubiformis Benoist, *Bull. Soc. Bot. France* 66 (8): 390 [“1919” publ. 1920] (Benoist 1920), “tubaeformis”.

VERNACULAR NAMES. — Pa: miumiu-puveyo • Wp: inga takwā u • Cr: pwa-sikré • Br: ingá.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *R. Benoist 1491* (holotype of *Inga tubiformis*: P[P01818292]; iso-, P[P01818293]).

INVENTORY DATA (FG). — 1 tree, $dbh = 13.6$ cm.

[840] *Inga longipedunculata* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 56 (Ducke 1922).

NOTE. — Synonym of *Inga leiocalyxina* Benth. for Pennington (1997), but leaves are distinctive.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2185*.

INVENTORY DATA (FG). — 75 trees in 36 plots; $F_{\max} = 1.7\%$; $dbh_{inv} = 45$ cm.

[841] *Inga loubryana* Poncy

Adansonia, sér. 3, 29 (2): 250 (Poncy 2007).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: miumiu-platno-mna.

HERBARIUM DATA (FG). — 60 collections at CAY. Sel. exs.: *D. Loubry 1135* (holo-, P[P01818279]; iso-, CAY[CAY086128], US[00955612]).

INVENTORY DATA (FG). — 274 trees in 79 plots; $F_{\max} = 2.5\%$; $dbh_{inv} = 75.9$ cm.

[842] *Inga macrophylla* Humb. & Bonpl. ex Willd.

Sp. Pl., ed. 4 4 (2): 1015 [Apr. 1806] (Willdenow 1806). — *Mimosa macrophylla* (Humb. & Bonpl. ex Willd.) Poir., *Encycl. J. Lamarck et al.* Suppl. 1: 42 [3 Sep. 1810] (Poiret 1810). — *Feuilleea macrophylla* (Humb. & Bonpl. ex Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga calocephala Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 78 [23-25 Jan. 1845] (Poeppig 1845).

Inga bracteosa Benth., *London J. Bot.* 4: 609 (Bentham 1845). — *Feuilleea bracteosa* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga brachyptera Benth., *London J. Bot.* 4: 610 (Bentham 1845). — *Feuilleea brachyptera* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Inga quadrangularis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 60 (Ducke 1922).

Inga ouraphylla L.Uribe, *Caldasia* 4 (20): 405 (Uribe 1947).

Inga alatocarpa T.S.Elias, *Phytologia* 14 (4): 206 [6 Feb. 1967] (Elias 1967).

Inga chorrorana T.S.Elias, *Bot. Mus. Leafl.* 24 (8): 202 [9 Apr. 1976] (Elias 1976). pro parte quoad specimina in fructo.

Inga macrophylla var. *stenoptera* Benth., *Fl. Bras. [Martius]* 15 (2): 490 [1 July 1876] (Bentham 1876).

NOTE. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Wp: inga yowa • Cr: pwa-sikré • Br: ingá-peua.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *M.-F. Prévost 761*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 15.1$ cm.

[843] *Inga marginata* Willd.

Sp. Pl., ed. 4 4 (2): 1015 [Apr. 1806] (Willdenow 1806), nom. et typ. cons. — *Feuilleea marginata* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891). — *Inga fagifolia* (L.) Willd. ex Benth. var. *marginata* (Willd.) Hassl., *Repert. Spec. Nov. Regni Veg.* 16: 154 [15 Nov. 1919] (Hassler 1919).

Inga sapida Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 286 [Apr. 1824] (Kunth 1824).

Mimosa semialata Vell., *Fl. Flumin. Icon.* 11: t. 5 ["1827" publ. 29 Oct. 1831] (Vellozo 1831). — *Inga semialata* (Vell.) Mart., *Flora* 20 (2, Beibl.): 111 (Martius 1837).

Inga odorata G.Don, *Gen. Hist.* 2: 388 [Oct. 1832] (Don 1832).

Inga guayaquilensis G.Don, *Gen. Hist.* 2: 391 [Oct. 1832] (Don 1832).

Inga excelsa Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 78 [23-25 Jan. 1845] (Poeppig 1845), nom. illeg. hom., non Kunth (1823) nec Wight ex Wall. (Wallich 1831).

Inga semialata var. *latifolia* Mart. ex Benth., *London J. Bot.* 4: 589 (Bentham 1845).

Inga puberula Benth., *London J. Bot.* 4: 589 (Bentham 1845).

Inga pycnostachya Benth., *London J. Bot.* 4: 589 (Bentham 1845).

Inga leptostachya Benth., *Trans. Linn. Soc. London* 30 (3): 608 [10 Apr. 1875] (Bentham 1875), nom. nud. pro syn.

Inga fagifolia f. *genuina* Hassl., *Repert. Spec. Nov. Regni Veg.* 16: 155 [15 Nov. 1919] (Hassler 1919), nom. inval. (Turland et al. 2018: Art. 24.3).

Inga fagifolia f. *pedicellaris* Hassl., *Repert. Spec. Nov. Regni Veg.* 16: 155 [15 Nov. 1919] (Hassler 1919).

Inga fagifolia var. *intermedia* Hassl., *Repert. Spec. Nov. Regni Veg.* 16: 155 [15 Nov. 1919] (Hassler 1919).

Inga marginata var. *itayensis* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.1)*: 30 (Macbride 1943).

Inga tysonii T.S.Elias, *Ann. Missouri Bot. Gard.* 53 (3): 377 (Elias 1966).

VERNACULAR NAMES. — Wp: inga sisi pay, sisi pay • Br: ingá-ferro.

HERBARIUM DATA (FG). — 59 collections at CAY. Sel. exs.: *D. Sabatier et al.* 4622.

INVENTORY DATA (FG). — 57 trees in 37 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38$ cm.

[844] *Inga melinonii* Sagot

Ann. Sci. Nat., Bot. sér. 6, 13: 335 (Sagot 1882), "Melinonis".

Inga cyclocarpa Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 14 (Ducke 1925).

NOTE. — The epithet "melinonis", which honours the French botanist E. Mélinon, is to be automatically corrected to "melinonii" (Turland et al. 2018: Art. 60.8).

VERNACULAR NAMES. — Pa: miumiu-maoksi-arib, miumiu-maoksi-aribinē • Ka: aliki enakololi • Wp: akiki inga, inga kala • Cr: pwa-sikré-rouj • Br: ingá.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *E.M. Mélinon* 453, 1862 (lecto-, P[P01818266], designated by Poncy [1985: 48]; isolecto-, F[V0093101F], P[P01818267, P01818268, P01818269], PH[00015899], US[00588635]).

INVENTORY DATA (FG). — 45 trees in 29 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 49.2$ cm.

[845] *Inga mitaraka* Poncy

Bull. Mus. Natl. Hist. Nat., B, Adansonia 18 (1-2): 70 (Poncy 1996).

NOTES. — Known only from the Guiana Shield. A synonym of *Inga alba* (Sw.) Willd. for Pennington (1997), but the fruit is distinctive.

VERNACULAR NAMES. — Wp: inga u • Cr: pwa-sikré • Br: ingá.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.-J. de Granville* 1197 (holo-, P[P01818264]; iso-, CAY[CAY028081, CAY028082]).

INVENTORY DATA (FG). — 19 trees in 10 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 66.8$ cm.

[846] *Inga nobilis* Willd.

Enum. Pl. [Willdenow] 2: 1047 [June 1809] (Willdenow 1809). — *Feuilleea nobilis* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga humboldtiana Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 285 [Apr. 1824] (Kunth 1824).

Inga corymbifera Benth., *J. Bot. [Hooker]* 2: 144 (Bentham 1840).

Inga matthewsiana Benth., *London J. Bot.* 4: 594 (Bentham 1845). — *Feuilleea matthewsiana* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga riedeliana Benth., *London J. Bot.* 4: 595 (Bentham 1845).

Inga riedeliana var. *surinamensis* Benth., *London J. Bot.* 4: 595 (Bentham 1845).

Inga corymbifera var. *brasiliensis* Benth., *London J. Bot.* 4: 596 (Bentham 1845).

Inga sericantha Miq., *Linnaea* 19: 132 ["1847" publ. Feb. 1846] (Miquel 1846).

Inga nobilis var. *pavoniana* Benth., *Trans. Linn. Soc. London* 30 (3): 614 [10 Apr. 1875] (Bentham 1875). — *Inga pavoniana* Benth., *London J. Bot.* 4: 595 (Bentham 1845), nom. illeg. hom., non G.Don (1832).

Inga conglomerata Benoit, *Bull. Mus. Natl. Hist. Nat.* 27: 115 (Benoist 1921).

Inga loretana J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.1)*: 29 (Macbride 1943).

Inga jucunda Ducke, *Bol. Técn. Inst. Agron. N.* 2: 3 (Ducke 1944).

NOTE. — *Inga nobilis* var. *pavoniana* Benth. is based on the illegitimate *Inga pavoniana* Benth., which can not be treated a basionym; hence authorship is Benth., not (Benth.) Benth.

VERNACULAR NAMES. — Pa: miumiu-asukwinō, miumiu-maoksi-
arib, miumiu-maoksi-aribinē • Ka: posindyo • Wp: yáwī inga •
Wn: anekélē, mïkotawa, mïkotowa • Nt: bita weko • Cr: pwa-sikré
• Br: ingá-canela.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *G. Wachenheim* 149 (holotype of *Inga conglomera*: P[P01818244]; iso-, P[P01818245, P01818246]).

INVENTORY DATA (FG). — 7 trees in 4 plots; $F_{\max} < 1\%$; dbh_{inv} = 24.2 cm.

[847] *Inga nouragensis* Poncy

Bull. Mus. Natl. Hist. Nat., B, *Adansonia* 18 (1-2): 72 (Poncy 1996).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2569 (holo-, P[P01818235]); iso-, CAY[CAY000778].

INVENTORY DATA (FG). — 50 trees in 26 plots; $F_{\max} = 2\%$; dbh_{inv} = 44.9 cm.

[848] *Inga nubium* Poncy

Bull. Mus. Natl. Hist. Nat., B, *Adansonia* 13 (3-4): 148 (Poncy 1991).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-J. de Granville* 1475 (holo-, P[P01818236]; CAY[CAY000778], US[00478813]).

INVENTORY DATA (FG). — 1 tree, dbh = 12.5 cm.

[849] *Inga paraensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 12 (Ducke 1925).

VERNACULAR NAMES. — Pa: miumiu-asiru.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4361.

INVENTORY DATA (FG). — 142 trees in 53 plots; $F_{\max} = 2.4\%$; dbh_{inv} = 68.9 cm.

[850] *Inga pezizifera* Benth.

London J. Bot. 4: 587 (Bentham 1845). — *Feuilleea pezizifera* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga subsericantha Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 55 (Ducke 1922).

Inga urnifera Kleinhoonte, *Recueil Trav. Bot. Neerl.* 22: 413 ["1925 publ. Jan. 1926"] (Kleinhoonte 1926).

Inga microstachya Britton & Killip, *Ann. New York Acad. Sci.* 35: 115 [1 Apr. 1936] (Britton & Killip 1936).

Inga riopalenquensis A.H.Gentry, *Selbyana* 2 (1): 39 [Aug. 1977] (Gentry 1977).

VERNACULAR NAMES. — Pa: avukun • Ka: anakala, anakolo, maipuli solapali, padyawa, payawa • Te: pulukuni • Wp: inga kala, inga piyū, inga sisi, masulapa pe u, sisi • Wn: apulukun • Nt: koopogno, lebi weko, weko • Cr: bougouni, pwa-sikré-blanc • Fr: pois sucré blanc • Br: ingá-xixica.

HERBARIUM DATA (FG). — 73 collections at CAY. Sel. exs.: *M.-F. Prévost* 1271.

INVENTORY DATA (FG). — 125 trees in 50 plots; $F_{\max} = 1.2\%$; dbh_{inv} = 80 cm.

[851] *Inga pilosula* (Rich.) J.F.Macbr.

(Fig. 30E)

Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.1): 29 (Macbride 1943). — *Mimosa pilosula* Rich., *Actes Soc. Hist. Nat. Paris* 1: 113 [Oct. 1792] (Richard 1792). — *Inga pilosiuscula* Desv., *J. Bot. Agric.* 3: 71 (Desvaux 1814), nom. illeg. superfl. (based on *Inga pilosula*). — *Feuilleea pilosula* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 186 [5 Nov. 1891] (Kuntze 1891).

Inga quassiaefolia Willd., *Sp. Pl.*, ed. 4 4 (2): 1013 [Apr. 1806] (Willdenow 1806), "quassiaefolia". — *Mimosa quassiaefolia* (Willd.) Poir., *Encycl. J. Lamarck et al. Suppl.* 1: 41 [3 Sep. 1810] (Poiret 1810), "quassiaefolia". — *Feuilleea quassiaefolia* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 186 [5 Nov. 1891] (Kuntze 1891), "quassiaefolia".

Inga nitida Willd., *Sp. Pl.*, ed. 4 4 (2): 1013 [Apr. 1806] (Willdenow 1806). — *Mimosa nitida* (Willd.) Poir., *Encycl. J. Lamarck et al. Suppl.* 1: 41 [3 Sep. 1810] (Poiret 1810), nom. illeg. hom., non Vahl (1791).

Mimosa lucida Vahl, *Ectog. Amer.* 3: 31 (Vahl 1807).

Inga setifera DC., *Prodr. [A. P. de Candolle]* 2: 432 [mid Nov. 1825] (Candolle 1825). — *Feuilleea setifera* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 184, 189 [5 Nov. 1891] (Kuntze 1891).

Inga platycarpa Benth., *J. Bot. [Hooker]* 2: 142 (Bentham 1840).

Inga affinis Steud., *Flora* 26 (45): 758 [17 Dec. 1843] (Steudel 1843), nom. illeg. hom., non DC. (Candolle 1825).

Inga macrophylla Hook., *Bot. Mag.* 84: t. 5075 (Hooker 1858), nom. illeg. hom., non Humb. & Bonpl. ex Willd. (Willdenow 1806).

Inga versicolor Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 616 [10 Apr. 1875] (Bentham 1875), nom. nud. pro syn.

Inga sanctae-annae S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 350 ["1894-96" publ. Dec. 1895] (Moore 1895).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.B. Leblond* 144 (holo-, P-JU[not seen]; iso-, G[G00371331]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 38.2 cm.

[852] *Inga poeppigiana* Benth.

London J. Bot. 4: 602 (Bentham 1845). — *Feuilleea poeppigiana* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga ciliata Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 78 [23-25 Jan. 1845] (Poeppig 1845), *nom. illeg. hom.*, *non* C.Presl (1834).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 4465*.

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.2$ cm.

[853] *Inga punctata* Willd.

Sp. Pl., ed. 4 4 (2): 1016 [Apr. 1806] (Willdenow 1806), excl. syn. — *Feuilleea punctata* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Mimosa sericea Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 42 [3 Sep. 1810] (Poiret 1810).

Inga leptoloba Schlehd., *Linnaea* 12: 560 (Schlechtendal 1838). — *Feuilleea leptoloba* (Schlehd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga rufinervis Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 612 [10 Apr. 1875] (Bentham 1875). — *Feuilleea rufinervis* (Spruce ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga strigillosa Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 612 [10 Apr. 1875] (Bentham 1875). — *Feuilleea strigillosa* (Spruce ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga punctata var. *panamensis* Benth., *Trans. Linn. Soc. London* 30 (3): 613 [10 Apr. 1875] (Bentham 1875).

Inga cycladenia Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 184 [30 Oct. 1916] (Pittier 1916).

Inga popayanensis Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 185 [30 Oct. 1916] (Pittier 1916).

Inga ierensis Britton, *Bull. Torrey Bot. Club* 50 (1): 52 [Jan. 1923] (Britton 1923).

Inga punctata subsp. *chagrensis* Pittier, *J. Dept. Agric. Porto Rico* 13: 135 (Pittier 1929).

Inga punctata var. *elongata* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (3.1): 38 (Macbride 1943).

VERNACULAR NAMES. — Wp: paku inga • Cr: pwa-sikré • Br: ingá-canela, ingá-de-leite.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *P. Grenand 1503*.

SIZE. — Up to 60 cm dbh (Pennington 1997).

[854] *Inga retinocarpa* Poncy
(Fig. 30F)

Bull. Mus. Natl. Hist. Nat., B, *Adansonia* 13 (3-4): 147 (Poncy 1991).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-F. Villiers & C. Feuillet 2086* (holo-, P[P01818207]; iso-, CAY[CAY028096], NY[00005376], P[P01818208], U[U0003391]).

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43.4$ cm.

[855] *Inga rhynchocalyx* Sandwith

Kew Bull. 3 (2): 318 [20 Nov. 1948] (Sandwith 1948).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *O. Poncy & J. Munzinger 1637*.

INVENTORY DATA (FG). — 6 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30.6$ cm.

[856] *Inga rubiginosa* (Rich.) DC.

Prodr. [A. P. de Candolle] 2: 434 [mid Nov. 1825] (Candolle 1825). — *Mimosa rubiginosa* Rich., *Actes Soc. Hist. Nat. Paris* 1: 113 [Oct. 1792] (Richard 1792). — *Feuilleea rubiginosa* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga standleyana Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 204 [30 Oct. 1916] (Pittier 1916).

VERNACULAR NAMES. — Pa: miumiu-āhinó • Ka: alawata busulukulu, alawata posu:lukulu, padyawa, paidyawa, payawa, pololu peta • Wp: akiki inga • Cr: pwa-sikré-rouj • Br: ingá-de-pelo.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-723*.

INVENTORY DATA (FG). — 207 trees in 93 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 51$ cm.

[857] *Inga sarmentosa* Glaz. ex Harms

Notizbl. Königl. Bot. Gart. Berlin 6: 303 [30 June 1915] (Harms 1915).

Inga capitata var. *latifolia* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 120 (Ducke 1930).

NOTE. — A synonym of *Inga capitata* Desv. for Pennington (1997), but leaves are distinctive.

VERNACULAR NAMES. — Pa: simartē-puvemna • Wp: inga mani'o • Nt: weko • Cr: pwa-sikré • Br: ingá.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier 2372*.

INVENTORY DATA (FG). — 55 trees in 33 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 59.5$ cm.

[858] *Inga sertulifera* DC.

Prod. [A. P. de Candolle] 2: 436 [mid Nov. 1825] (Candolle 1825).

Mimosa coriacea Pers., *Syn. Pl. [Persoon]* 2 (1): 262 [Nov. 1806] (Persoon 1806). — *Feuilleea coriacea* (Pers.) Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891). — *Inga coriacea* (Pers.) Desv., *J. Bot. Agric.* 3: 71 (Desvaux 1814), “*coriacea*”, nom. illeg. hom., non Humb. & Bonpl. ex Willd. (Willdenow 1806) [synonym of *Calliandra coriacea* (Humb. & Bonpl. ex Willd.) Benth.]

VERNACULAR NAMES. — Pa: miumiu-akamá-arib, miumiu-akamá-ribiné • Ka: waikimili, wayamu tapulu • Te: inga dju • Wp: inga tawa • Wn: kului ahmit • Nt: adai weko, baaka weko • Cr: bakov-kanoté, pwa-sikré-bakòv • Fr: pois sucré • Br: ingá-chata, ingá-de-baixo.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *X. Cuniberti* 22.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38$ cm.

[859] *Inga splendens* Willd.

Sp. Pl., ed. 4 (2): 1017 [Apr. 1806] (Willdenow 1806). — *Mimosa splendens* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 43 [3 Sep. 1810] (Poirier 1810). — *Feuilleea splendens* (Willd.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga floribunda Benth., *J. Bot. [Hooker]* 2: 143 (Bentham 1840).

Inga hostmannii Pittier, *Contr. U.S. Natl. Herb.* 18 (5): 188 [30 Oct. 1916] (Pittier 1916). — *Inga splendens* var. *hostmannii* (Pittier) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 15 (Ducke 1925).

Inga superba Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 57 (Ducke 1922). — *Inga splendens* var. *superba* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 16 (Ducke 1925).

VERNACULAR NAMES. — Pa: miumiu-wašiunu • Ka: anakala, inyai, inyau, lapalapa • Wp: a’ipopita • Wn: anahkale, tokulojem • Nt: kodya weko • Cr: pwa-sikré-koubari • Br: ingá-duro.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3640.

INVENTORY DATA (FG). — 22 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 61.4$ cm.

[860] *Inga stipularis* DC.

Prod. [A. P. de Candolle] 2: 435 [mid Nov. 1825] (Candolle 1825). — *Feuilleea stipularis* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Ka: padyawa, payawa, waiki • Wp: inga sisi pay, inga u sóví, sisi pay • Nt: weko • Cr: pwa-sikré-gran-bwa • Br: ingá.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *J.P.B. von Rohr* (“*von Rohr ex Patris*”) s.n. (original material BM[[BM000541108](#)]).

INVENTORY DATA (FG). — 64 trees in 44 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 27.1$ cm.

[861] *Inga striata* Benth.

London J. Bot. 4: 608 (Bentham 1845).

Mimosa dulcis Vell., *Fl. Flumin. Icon.* 11: t. 4 [“1827” publ. 29 Oct. 1831] (Vellozo 1831), nom. illeg. hom., non Roxb. (Roxburgh 1795). — *Inga dulcis* Mart., *Flora* 20 (2, Beibl.): 113 (Martius 1837), nom. illeg. hom., non (Roxb.) Willd. (Willdenow 1806). — *Inga arrabidae* Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 809 (Steudel 1840), nom. illeg. superfl. (based on *Inga dulcis*).

Inga catharinæ Benth., *London J. Bot.* 4: 605 (Bentham 1845). — *Feuilleea catharinæ* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891).

Inga nuda Salzm. ex Benth., *London J. Bot.* 4: 605 (Bentham 1845). — *Feuilleea nuda* (Salzm. ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga salzmanniana Benth., *London J. Bot.* 4: 608 (Bentham 1845). — *Feuilleea salzmanniana* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga comewynensis Miq., *Stirp. Surinam. Select.*: 1 [“1850” publ. Mar. 1851] (Miquel 1851).

Inga nuda var. *longiflora* Benth., *Trans. Linn. Soc. London* 30 (3): 618 [10 Apr. 1875] (Bentham 1875).

Inga prieurii Sagot, *Ann. Sci. Nat., Bot. sér.* 6, 13: 332 (Sagot 1882), “*Prieurei*”.

Inga perrottetii Sagot, *Ann. Sci. Nat., Bot. sér.* 6, 13: 335 (Sagot 1882).

Inga canaminensis Rusby, *Mem. New York Bot. Gard.* 7: 249 (Rusby 1927).

Inga ellsworthiana L. Uribe, *Caldasia* 4 (20): 407 (Uribe 1947).

NOTES. — *Inga arrabidae* Steud. is not a synonym of *Inga vera* subsp. *affinis* (DC.) T.D.Penn., and not a *nomen nudum*. Steudel (1840, 1: 809) cited in synonymy “*I. dulcis* Mart. (non Willd.)” and “*Mimosa dulcis* Arrab.” (i.e. *Mimosa dulcis* Vell.) *Inga dulcis* is based on the illegitimate *Mimosa dulcis* Vell., thus the authority for this binomial is “Mart.”, not “(Vell.) Mart.”

VERNACULAR NAMES. — Br: ingá-banana, ingá-caixão.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *F.M.R. Leprieur* s.n. (original material of *Inga prieurii*: G[[G00388527](#)]).

INVENTORY DATA (FG). — 14 trees in 5 plots; $dbh_{\text{inv}} = 54.1$ cm.

[862] *Inga suaveolens* Ducke

Bol. Técn. Inst. Agron. N. 2: 5 (Ducke 1944).

VERNACULAR NAMES. — Wp: paku inga • Nt: bita weko.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *S.A. Mori* et al. 23344.

SIZE. — Up to 30 cm dbh (Pennington 1997).

[863] *Inga thibaudiana* DC.

Prodr. [A. P. de Candolle] 2: 434 [mid Nov. 1825] (Candolle 1825). — *Feuilleea thibaudiana* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga gladiata Desv., *Ann. Sci. Nat. (Paris)* 9: 427 (Desvaux 1826).

Inga macradenia Mart. ex Benth., *London J. Bot.* 4: 596 (Bentham 1845), *nom. nud. pro syn.*

Inga tenuiflora Salzm. ex Benth., *London J. Bot.* 4: 596 (Bentham 1845), “*tenuifolia*”.

Inga tenuiflora var. *glabrior* Benth., *London J. Bot.* 4: 596 (Bentham 1845).

Inga recordii Britton & Rose, *Trop. Woods* 7: 5 (Britton & Rose 1926).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: miumiu-akamā-arib, miumiu-akamā-aribinē • Ka: alawata posu:lukulu, padyawa, payawa, sepundi, tamunen waiki, tanimi • Wp: alákwa inga • Wn: panawilili • Nt: kamina weko • Cr: pwa-sikré • Fr: pois sucré • Br: ingá-cipo-preta.

HERBARIUM DATA (FG). — 121 collections at CAY. Sel. exs.: *A.N. Desvaux s.n.* (type of *Inga gladiata*: P[P01818175]).

INVENTORY DATA (FG). — 139 trees in 60 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 45.2$ cm.

[864] *Inga umbellifera* (Vahl) Steud.

Nomenc. Bot. [Steudel] 1: 431 (Steudel 1821). — *Mimosa umbellifera* Vahl, *Eclog. Amer.* 3: 30 (Vahl 1807). — *Feuilleea umbellifera* (Vahl) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga sciadion Steud., *Flora* 26 (45): 758 [17 Dec. 1843] (Steudel 1843). — *Feuilleea sciadion* (Steud.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga myriantha Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 77 [23-25 Jan. 1845] (Poeppig 1845). — *Feuilleea myriantha* (Poepp.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga rutilans Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 602 [10 Apr. 1875] (Bentham 1875), *nom. nud. pro syn.*

Inga gracilipes Standl., *J. Wash. Acad. Sci.* 15: 101 (Standley 1925).

Inga lawranceana Britton & Killip, *Phytologia* 1 (1): 23 [Dec. 1933] (Britton & Killip 1933).

VERNACULAR NAMES. — Pa: miumiu-asiru • Ka: wayamu tapulu • Wp: asila, kululu inga • Cr: pwa-sikré • Br: ingá-de-macaco.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (type C[C10012170]).

INVENTORY DATA (FG). — 41 trees in 35 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 18.2$ cm.

[865] *Inga umbratica* Poepp.

Nova genera ac species plantarum [Poeppig & Endlicher] 3: 77 [23-25 Jan. 1845] (Poeppig 1845). — *Feuilleea umbratica* (Poepp.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

Inga radiata Rusby, *Mem. New York Bot. Gard.* 7: 252 (Rusby 1927).

Inga brevialata Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 119 (Ducke 1930).

Inga suturalis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 119 (Ducke 1930).

NOTE. — Poeppig is the sole author of vol. 3 of *Nova genera ac species plantarum* (Stafleu & Cowan, 1983: TL-2-8091).

HERBARIUM DATA (FG). — No collection at CAY. Sel. exs.: *H.S. Irwin 48608* (NY[02679450]).

SIZE. — Up to 20 cm dbh (Pennington 1997).

[866] *Inga vera* Willd. subsp. *affinis* (DC.) T.D.Penn.

Gen. Inga, *Bot.* 716 (Pennington 1997). — *Inga affinis* DC., *Prodr. [A. P. de Candolle]* 2: 433 [mid Nov. 1825] (Candolle 1825). — *Feuilleea affinis* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891).

Mimosa umbellata Vell., *Fl. Flumin. Icon.* 11: t. 12 [“1827” publ. 29 Oct. 1831] (Vellozo 1831), *nom. illeg. hom., non Vahl* (1791).

Inga uraguensis Hook. & Arn., *Bot. Misc.* 3: 202 (Hooker & Arnott 1833). — *Feuilleea uraguensis* (Hook. & Arn.) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891), “*uruguensis*”.

Inga velloziana Mart., *Flora* 20 (2, Beibl.): 114, 118 (Martius 1837).

Inga acutifolia Benth., *London J. Bot.* 4: 614 (Bentham 1845).

Inga spuria Humb. & Bonpl. ex Willd. var. *sordida* Benth., *London J. Bot.* 4: 616 (Bentham 1845).

Inga meissneriana Miq., *Stirp. Surinam. Select.*: 2 [“1850” publ. Mar. 1851] (Miquel 1851). — *Feuilleea meissneriana* (Miq.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Inga uraguensis f. *tomentosula* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 481 (Chodat & Hassler 1904).

Inga uraguensis var. *parvifolia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 481 (Chodat & Hassler 1904).

Inga arinensis Hoehne, *Comm. Lin. Telegr. Bot.* 45 (8): 17 (Hoehne 1919).

Inga soluta Pittier, *Cat. Fl. Venez. [Pittier]* 1: 353 (Pittier 1945), *nom. nud.*

VERNACULAR NAMES. — Ka: wala potili • Te: inga katapali • Wp: inga katapali, katapali • Br: ingá-banana, ingá-do-brejo.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier 854*.

SIZE. — Up to 40 cm dbh (Pennington 1997).

[867] *Inga virgultosa* (Vahl) Desv.
(Fig. 31A)

Ann. Sci. Nat. (Paris) 9: 426 (Desvaux 1826). — *Mimosa virgultosa* Vahl, *Eclog. Amer.* 3: 32 (Vahl 1807). — *Feuilleea virgultosa* (Vahl) Kuntze, *Revis. Gen. Pl.* 1: 189 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: pukuu • Wp: kululu inga sili • Nt: anpuku weko.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (type C[[C10012171](#)]).

INVENTORY DATA (FG). — 10 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.1$ cm.

[868] *Inga* sp. A

NOTE. — A species that superficially looks like *I. loubryana* Poncy. Dry leaf colour and venation are quite distinctive.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino 1924*.

INVENTORY DATA (FG). — 17 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 86.3$ cm.

[869] *Inga* sp. B

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2673*.

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.7$ cm.

[870] *Inga* sp. C

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2696*.

INVENTORY DATA (FG). — 7 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.5$ cm.

[871] *Inga* sp. D

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 3325*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 13.2$ cm.

Genus *Leptolobium* Vogel

[872] *Leptolobium nitens* Vogel

Linnaea 11: 394 (Vogel 1837). — *Sweetia nitens* (Vogel) Benth., *J. Proc. Linn. Soc., Bot.* 8: 262 [17 Feb. 1865] (Bentham 1865). — *Acosmium nitens* (Vogel) Yakovlev, *Notes Roy. Bot. Gard. Edinburgh* 29 (3): 353 (Yakovlev 1969).

Leptolobium nitidulum Miq., *Stirp. Surinam. Select.*: 18 ["1850" publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Ka: silityo • Br: itaúba-rana.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J. Martin s.n.* (type B[not seen, photo F neg. N° 1875]).

SIZE. — Up to 30 m tall (Rodrigues & Tozzi 2012).

Genus *Leucaena* Benth.

[873] *Leucaena leucocephala* (Lam.) de Wit

Taxon 10: 54 (de Wit 1961). — *Mimosa leucocephala* Lam., *Encycl. [J. Lamarck et al.] 1 (1)*: 12 [2 Dec. 1783] (Lamarck 1783). — *Acacia leucocephala* (Lam.) Link, *Enum. Hort. Berol. Alt.* 2: 444 (Link 1822).

Galega mimosoides Willd., *Sp. Pl.*, ed. 4 3 (2): 1249 [1-10 Nov. 1802] (Willdenow 1802). — *Tephrosia mimosoides* (Willd.) Pers., *Syn. Pl. [Persoon]* 2 (2): 330 [Sep. 1807] (Persoon 1807).

Acacia biceps Willd., *Sp. Pl.*, ed. 4 4 (2): 1075 [Apr. 1806] (Willdenow 1806). — *Mimosa biceps* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 75 [3 Sep. 1810] (Poiret 1810).

Acacia frondosa Willd., *Sp. Pl.*, ed. 4 4 (2): 1076 [Apr. 1806] (Willdenow 1806). — *Mimosa frondosa* (Willd.) Klein ex Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 76 [3 Sep. 1810] (Poiret 1810).

Mimosa glandulosa C.Sm., *Narr. Exped. Zaire*: 249 (Smith 1818), *nom. nud.*

Tephrosia barclayana Sweet, *Hort. Brit. [Sweet]*, ed. 2: 142 (Sweet 1830).

Acacia caringa Buch.-Ham. ex Wall., *Numer. List no. 5239* (Wallich 1831).

Leucaena glauca Benth., *J. Bot. [Hooker]* 4: 416 (Bentham 1842).

NOTE. — A species native to Central America, widely cultivated in tropical and subtropical regions, naturalised in some French Guiana suburban areas.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *C. Feuillet 2961*.

Genus *Limadendron* Meireles & A.M.G.Azevedo

[874] *Limadendron hostmannii*

(Benth.) Meireles & A.M.G.Azevedo
(Fig. 31D)

Pl. Syst. Evol. 301: 705 [epubl. 4 July 2014] (Meireles & Azevedo 2014). — *Cyclolobium hostmannii* Benth., *J. Proc. Linn. Soc., Bot.* 4 (Suppl.): 52 [Mar. 1860] (Bentham 1860), "Hostmanni". — *Poecilanthe hostmannii* (Benth.) Amshoff, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 52: 61 (Amshoff 1939).

VERNACULAR NAMES. — Pa: miumiu-platno-kamwi • Wp: yapuku-liwa u.

HERBARIUM DATA (FG). — 62 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3086*.

INVENTORY DATA (FG). — 72 trees in 18 plots; $F_{\max} = 3.9\%$; $dbh_{\text{inv}} = 20.5$ cm.

Genus *Lonchocarpus* Kunth

[875] *Lonchocarpus hedyosmus* Miq.

Linnaea 18: 564 [“1844” publ. prob. Aug. 1845] (Miquel 1845). — *Derris hedyosma* (Miq.) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.1)*: 262 (Macbride 1943).

Lonchocarpus macrocarpus Benth. var. *sericophyllus* Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 91 [Mar. 1860] (Bentham 1860), “*sericophylla*”.

Lonchocarpus paniculatus Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 161 (Ducke 1922).

VERNACULAR NAMES. — Wp: pana’i, tatu’i u, wila ki • Cr: pana-koko-marikaj.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1559*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; dbh_{inv} = 100 cm.

[876] *Lonchocarpus heptaphyllus* (Poir.) DC.

Prodr. [A. P. de Candolle] 2: 260 [mid Nov. 1825] (Candolle 1825). — *Dalbergia heptaphylla* Poir., *Encycl. [J. Lamarck et al.] Suppl. 2*: 445 [3 July 1812] (Poirier 1812). — *Amerimnon latifolium* Willd., *Sp. Pl. ed. 4 3 (2)*: 909 [1-10 Nov. 1802] (Willdenow 1802), “*Amerimnum*”, nom. illeg. superfl. [*Amerimnon pinnatum* Jacq., synonym of *Platymiscium pinnatum* (Jacq.) Dugand in syn.]

Robinia sepium Sw., *Fl. Ind. Occid. 3*: 1258 [Oct. 1806] (Swartz 1806), nom. illeg. hom., non Jacq. (Jacquin 1760) [synonym of *Gliricidia sepium* (Jacq.) Kunth ex Walp.]

Dalbergia pentaphylla Poir., *Encycl. [J. Lamarck et al.] Suppl. 2*: 445 [3 July 1812] (Poirier 1812).

Lonchocarpus latifolius DC., *Prodr. [A. P. de Candolle] 2*: 260 [mid Nov. 1825] (Candolle 1825).

Lonchocarpus swartzii DC., *Prodr. [A. P. de Candolle] 2*: 261 [mid Nov. 1825] (Candolle 1825).

Cytisus membranaceus Sessé & Moc., *Fl. Mexic.*, ed. 2, 174 (Sessé & Mociño 1894).

Lonchocarpus discolor Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr. 3*: 421 (Huber 1902).

Derris latifolia Ducke, *Bol. Técn. Inst. Agron. N. 18*: 195 (Ducke 1949), nom. illeg. hom., non Prain (1898).

NOTES. — *Dalbergia pentaphylla* Poir. and *D. heptaphylla* Poir., published simultaneously, are the earliest legitimate names for this species, and they have been synonymized by Silva & Tozzi (2012). Therefore, *Lonchocarpus heptaphyllus* (Poir.) DC. is the name to be used for this species. *Amerimnon (“Amerimnum”) latifolium* Willd. is an illegitimate name, because Willdenow (1802: 909) cited *A. pinnatum* (synonym of *Platymiscium pinnatum*) in synonymy. Kunth (Humboldt et al. 1824: 383, adnot. 2) placed *Amerimnon latifolium* in *Lonchocarpus*, but he did not associate the genus name with the epithet, thus the combination is invalid and the correct authority of *L. latifolius* is “DC.” *Robinia nicou* Aubl., synonymised under *L. latifolius* by Tozzi & Silva (2007), is actually a species of *Deguegia* Aubl. (Silva & Tozzi 2012). Even though *Cytisus membranaceus*

Sessé & Moc. already appears in the first edition of the Flora Mexicana (Sessé & Mociño 1892-1898: 190), ed. 2 has priority, probably from p. 49 onward (Stafleu & Cowan 1985: [TL-2-11756](#)).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1441*.

SIZE. — Up to 20 m tall (Amshoff 1976).

Genus *Macrolobium* Schreb.

[877] *Macrolobium acaciifolium* (Benth.) Benth.

Fl. Bras. [Martius] 15 (2): 224 [1 Dec. 1870] (Bentham 1870), “*acaciæfolium*”. — *Outea acaciifolia* Benth., *J. Bot. [Hooker] 2*: 94 (Bentham 1840), “*acaciæfolia*”. — *Vouapa acaciifolia* (Benth.) Baill., *Hist. Pl. [Baillon] 2*: 109 [Jan.-Feb. 1870] (Baillon 1870), “*acaciæfolia*”. — *Vouapa acaciifolia* (Benth.) Kuntze, *Revis. Gen. Pl. 1*: 213 [5 Nov. 1891] (Kuntze 1891), “*Vuapa*”, isonym (orth. variant).

Macrolobium acaciifolium var. *vestitum* Sandwith, *Kew Bull. 3 (2)*: 312 [20 Nov. 1948] (Sandwith 1948).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: iiru • Ka: alapali, alapali, alatapali • Te: kalaipé’i • Wp: alapali • Wn: elekesi • Nt: paata sii, wasipa • Cr: mapa-ri, rapari • Br: arapari, faveira.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2848*.

SIZE. — Up to 100 cm dbh (Cowan & Barneby 1998).

[878] *Macrolobium angustifolium* (Benth.) R.S.Cowan

Mem. New York Bot. Gard. 8 (4): 314 (Cowan 1953). — *Vouapa angustifolia* Benth., *Hooker’s J. Bot. Kew Gard. Misc. 2*: 239 (Bentham 1850).

Vouapa chrysostachya Miq., *Stirp. Surinam. Select.*: 11 [“1850” publ. Mar. 1851] (Miquel 1851). — *Macrolobium chrysostachyum* (Miq.) Benth., *Fl. Bras. [Martius] 15 (2)*: 220 [1 Dec. 1870] (Bentham 1870).

Macrolobium chrysostachyum var. *parviflorum* Benth., *Fl. Bras. [Martius] 15 (2)*: 220 [1 Dec. 1870] (Bentham 1870), “*parviflora*”.

Macrolobium hymenaeifolium Pittier, *Bol. Soc. Venez. Ci. Nat. 7* (49): 141 (Pittier 1941), “*hymeneaefolium*”.

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Ka: alatapa, atapa.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *A. Vailant s.n.* (P[P03464491]).

SIZE. — Up to 100 cm dbh (Cowan & Barneby 1998).

[879] *Macrolobium bifolium* (Aubl.) Pers.

Syn. Pl. [Persoon] 1: 39 [1 Apr.-15 June 1805] (Persoon 1805). — *Vouapa bifolia* Aubl., *Hist. Pl. Guiane* 1: 25 [Jun.-Dec. 1775] (Aublet 1775). — *Macrolobium vuapa* J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1)*: 93 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. superfl.* (based on *Vouapa bifolia* [typ. cons. of *Macrolobium* Schreb. (*nom. cons.*)]. — *Macrolobium hymenaeoides* Willd., *Sp. Pl., ed. 4 1 (1)*: 186 [June 1797] (Willdenow 1797), *nom. illeg. superfl.* (based on *Vouapa bifolia*).

Macrolobium elegans Miq., *Ann. Sci. Nat., Bot. sér. 3, 1*: 40 (Miquel 1844).

Macrolobium stamineum G.Mey., *Prim. Fl. Esseq.* 18 [Nov. 1818] (Meyer 1818). — *Vouapa staminea* (G.Mey.) DC., *Prodri. [A. P. de Candolle]* 2: 511 [mid Nov. 1825] (Candolle 1825).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wap-purubumna • Ka: atapa, vouapa (*fide* Aublet 1775) • Te: takulu wila • Wp: ipewi • Wn: alawata pana, watapania • Nt: watampana • Cr: wapa-dilo, wapa-larivé, wapa-marikaj, wapa-sek • Fr: wapa marécage • Br: fava-de-tambaqui, ingarana-vermelho, ipê-do-igapó.

HERBARIUM DATA (FG). — 91 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777303] designated by Lanjouw & Uittien [1940: 154]; isolecto-, BM[BM000952224]).

INVENTORY DATA (FG). — 100 trees in 34 plots; $F_{\max} = 1.8\%$; $\text{dbh}_{\text{inv}} = 49.4$ cm.

[880] *Macrolobium campestre* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 389 (Huber 1909).

VERNACULAR NAMES. — Nt: watampana.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 4958*.

INVENTORY DATA (FG). — 3 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 44.6$ cm.

[881] *Macrolobium guianense* (Aubl.) Pulle

Enum. Vasc. Pl. Surinam 211 (Pulle 1906), “guyanense”. — *Outea guianensis* Aubl., *Hist. Pl. Guiane* 1: 29 [Jun.-Dec. 1775] (Aublet 1775), “guyanense”. — *Macrolobium utea* J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1)*: 93 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. superfl.* (based on *Outea guianensis*). — *Macrolobium pinnatum* Willd., *Sp. Pl., ed. 4 1 (1)*: 186 [June 1797] (Willdenow 1797), *nom. illeg. superfl.* (based on *Outea guianensis*). — *Outea guianensis* J.St.-Hil., *Expos. Fam. Nat.* 2: 203 (Saint-Hilaire 1805), “*Utea guyannensis*”, isonym (orth. variant). — *Macrolobium utea* Steud., *Nomencl. Bot. [Steudel]* 1: 503 (Steudel 1821), *nom. illeg. superfl.* (based on *Outea guianensis*). — *Vouapa guianensis* (Aubl.) Taub., *Bot. Centralbl.* 47: 394 [31 Sept 1891] (Taubert 1891), “*guyanensis*”. — *Vouapa guianensis* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 213 [5 Nov. 1891] (Kuntze 1891), “*Vuapa*”, isonym (orth. variant).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: watampana.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777303] designated by Lanjouw & Uittien [1940: 154]; isolecto-, BM[BM000952224]).

INVENTORY DATA (FG). — 127 trees in 4 plots; $F_{\max} = 12.9\%$; $\text{dbh}_{\text{inv}} = 70$ cm.

[882] *Macrolobium huberianum* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 26 (Ducke 1915).

VERNACULAR NAMES. — Pa: sakeg-seinô.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier 5571*.

SIZE. — Up to 10 cm dbh (Cowan 1953).

[883] *Macrolobium multijugum* (DC.) Benth.

Fl. Bras. [Martius] 15 (2): 222 [1 Dec. 1870] (Bentham 1870). — *Outea multijuga* DC., *Prodri. [A. P. de Candolle]* 2: 510 [mid Nov. 1825] (Candolle 1825). — *Vouapa multijuga* (DC.) Taub., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (3): 85 (Taubert 1891). — *Vouapa multijuga* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 213 [5 Nov. 1891] (Kuntze 1891), “*Vuapa*”, isonym (orth. variant).

VERNACULAR NAMES. — Pa: iiru-purubumna • Ka: alapali, alatapali.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BR[BR0000005194605, BR0000005194698, BR0000005195022], FI[FI004958], K[K000264998]).

SIZE. — Guyana. *K.M. Redden et al. 3306*, 15 m.

Genus *Macrosamanea* Britton & Rose[884] *Macrosamanea pubiramea* (Steud.) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (1): 196 (Barneby & Grimes 1996). — *Inga pubiramea* Steud., *Flora* 26 (45): 759 [17 Dec. 1843] (Steudel 1843).

Pithecellobium longiflorum Benth., *London J. Bot.* 5: 107 (Bentham 1846), “*Pithecellobium*”. — *Feuilleea longiflora* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891). — *Macrosamanea longiflora* (Benth.) Pittier, *Cat. Fl. Venez. [Pittier]* 1: 358 (Pittier 1945).

Pithecellobium miquelianum Meisn., *Linnaea* 21: 250 (Meisner 1848), “*Pithecellobium*”.

Pithecellobium adiantifolium (Kunth) Benth. var. *multipinnatum* Benth., *Fl. Bras. [Martius]* 15 (2): 445 [1 July 1876] (Bentham 1876), “*Pithecellobium*”.

VERNACULAR NAMES. — Ka: alamilulan.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-J. de Granville 2902*.

SIZE. — Up to 10 m tall (Barneby *et al.* 2011).



FIG. 31. — Leguminosae: **A**, *Inga virgultosa* (Vahl) Desv. (M.-F. Prévost 4435); **B**, *Monoptyrix inpae* W.A.Rodrigues (D. Sabatier & J.-F. Molino 5671); **C**, *Paloue induita* Sandwith (D. Sabatier 5785); **D**, *Limadendron hostmannii* (Benth.) Meireles & A.M.G.Azevedo; **E**, *Parkia gigantocarpa* Ducke (D. Sabatier et al. 6137); **F**, *Platymiscium trinitatis* Benth. (D. Sabatier 3516). A, © M.-F. Prévost/IRD; B-F, © D. Sabatier/IRD.

Genus *Martiodendron* Gleason

[885] *Martiodendron parviflorum* (Amshoff) R.C.Koeppen

Brittonia 14 (2): 202 [15 Apr. 1962] (Koeppen 1962). — *Martiusia parviflora* Amshoff, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 52: 32 (Amshoff 1939).

VERNACULAR NAMES. — Wn: mëpuimë • Nt: mutyengetyenge • Br: muirá-pixuna.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1987*.

INVENTORY DATA (FG). — 17 trees in 12 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 97\text{ cm}$.

Genus *Monopteryx* Spruce ex Benth.

[886] *Monopteryx impae* W.A.Rodrigues
(Fig. 31B)

Acta Amazonica 5 (2): 153 (Rodrigues 1975).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5671.

INVENTORY DATA (FG). — 135 trees in 27 plots; $F_{\max} = 3.5\%$; $\text{dbh}_{\text{inv}} = 170.3$ cm.

Genus *Ormosia* Jacks.

[887] *Ormosia bolivarensis* (Rudd) C.H.Stirt.

Fl. Venez. Guayana 5: 366 (Stirton 1999). — *Ormosia nobilis* var. *bolivarensis* Rudd, *Contr. U.S. Natl. Herb.* 32 (5): 345 [17 Sep. 1965] (Rudd 1965).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2173.

INVENTORY DATA (FG). — 37 trees in 20 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 56.8$ cm.

[888] *Ormosia cinerea* Benoist

Bull. Mus. Natl. Hist. Nat. 26: 86 (Benoist 1920).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *G. Wachenheim* 88 (original material P[P03101329, P03101332]).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 29.9$ cm.

[889] *Ormosia coarctata* Jacks.

Trans. Linn. Soc. London 10 (2): 363 [7 Sept 1811] (Jackson 1811).

Ormosia cuneata Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 64 (Ducke 1925).

VERNACULAR NAMES. — Ka: anakoko, panakoko • Wp: monokoe u, onokoe u, teposi kili • Wn: onohkoi, onohkowe • Nt: agi, neko udu.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M. Fleury* 2166.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.7$ cm.

[890] *Ormosia coccinea* (Aubl.) Jacks.

Trans. Linn. Soc. London 10 (2): 360 [7 Sept 1811] (Jackson 1811). — *Robinia coccinea* Aubl., *Hist. Pl. Guiane* 2: 773 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Pa: wanaku-kamwi • Ka: anakoko, panakoko • Wn: onohkowe • Nt: agi • Cr: panakoko • Fr: panacoco • Br: muirapiranga, tento.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000931940]).

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 100.6$ cm.

[891] *Ormosia costulata* (Miq.) Kleinhoonte

Recueil Trav. Bot. Néerl. 22: 392 ["1925" publ. Jan. 1926] (Kleinhoonte 1926). — *Leptolobium costulatum* Miq., *Stirp. Surinam. Select.*: 17 ["1850" publ. Mar. 1851] (Miquel 1851).

Ormosia trifoliolata Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 398 (Huber 1909). — *Ormosia costulata* var. *trifoliolata* (Huber) Amshoff, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 52: 48 (Amshoff 1939), "trifoliata".

VERNACULAR NAMES. — Ka: anakoko, panakoko.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *B. Dutrèze & F. Crozier* 19.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 21$ cm.

[892] *Ormosia coutinhoi* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 136 (Ducke 1922). — *Macroule coutinhoi* (Ducke) Pierce, *Trop. Woods* 71: 2 (Pierce 1942).

VERNACULAR NAMES. — Pa: kudagman-wašiunó, wadidga-purubumna, wanaku-marikasmagene • Wn: alakapuli, onohkoimë • Nt: neko udu gaan wiwi • Cr: sen-marten-blanc • Fr: saint-martin blanc grandes feuilles • Br: buiuçu.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4385.

INVENTORY DATA (FG). — 111 trees in 46 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 79.6$ cm.

[893] *Ormosia flava* (Ducke) Rudd

Contr. U.S. Natl. Herb. 32 (5): 298 [17 Sep. 1965] (Rudd 1965). — *Clathrotropis flava* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 134 (Ducke 1922). — *Ormosiopsis flava* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 61 (Ducke 1925).

Clathrotropis surinamensis Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 395 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

VERNACULAR NAMES. — Ka: alesiki'i • Wp: mēwā • Br: breucupira, tento-preto.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2403.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 38.2$ cm.

[894] *Ormosia lignivalvis* Rudd

Contr. U.S. Natl. Herb. 32 (5): 331 [17 Sep. 1965] (Rudd 1965).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M.-F. Prévost 2966*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 105.7$ cm.

[895] *Ormosia melanocarpa* Kleinhoonte

Recueil Trav. Bot. Néerl. 22: 391 ["1925" publ. Jan. 1926] (Kleinhoonte 1926).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4137*.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 69.9$ cm.

[896] *Ormosia nobilis* Tul.

Arch. Mus. Hist. Nat. 4: 106 (Tulasne 1844).

VERNACULAR NAMES. — Ka: anakoko, panakoko • Wp: monokoe, onokoe • Nt: lebi agi • Cr: panakoko-blan • Fr: panacoco blanc • Br: jatobaí-do-igapó, tento-grande.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2817*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30.9$ cm.

[897] *Ormosia paraensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 62 (Ducke 1925).

Ormosia crassicarpa Pierce ex Pittier, *Bol. Soc. Venez. Ci. Nat.* 10: 108 (Pittier 1944).

Ormosia heterophylla Pires, *Bol. Técn. Inst. Agron. N.* 38: 24 (Pires 1960).

VERNACULAR NAMES. — Pa: wanaku-priyo • Ka: anakoko, ityulu anakoko, panakoko • Wp: monokoe u, onokoe u, teposi kili • Cr: panakoko • Br: olho-de-cabra, tento.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3967*.

INVENTORY DATA (FG). — 43 trees in 30 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 63$ cm.

[898] *Ormosia stipularis* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 65 (Ducke 1925).

VERNACULAR NAMES. — Ka: ityulanano anakoko • Wp: monokoe u, onokoe u, teposi kili • Wn: wapotoimé • Cr: panakoko • Br: mulungu-brabo, tento.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2194*.

INVENTORY DATA (FG). — 10 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40.1$ cm.

[899] *Ormosia* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier 4853*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 29.1$ cm.

[900] *Ormosia* sp. B

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3420*.

INVENTORY DATA (FG). — 13 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 66.3$ cm.

[901] *Ormosia* sp. C

HERBARIUM DATA (FG). — One collection at CAY. Sel. exs.: *C. Moretti 358; M. Fleury 911* (P): "Grand arbre ripicole".

Genus *Paloue* Aubl.

[902] *Paloue brasiliensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 27 (Ducke 1915), "Paloveda".

VERNACULAR NAMES. — Pa: makit, makitwa • Nt: baaka wapa.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *R.A.A. Oldeman & C. Sastre 163; R.A.A. Oldeman & C. Sastre 77*, 15 m.

[903] *Paloue guianensis* Aubl.

Hist. Pl. Guiane 1: 365 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate.

VERNACULAR NAMES. — Pa: makit, makitwa • Nt: baaka wapa.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *J.B. Aublet s.n. (lecto-, BM[not seen], designated by Redden et al. [2018: 19])*.

INVENTORY DATA (FG). — 3 trees in 1 plot; $dbh_{\text{inv}} = 15.9$ cm.

[904] *Paloue induta* Sandwith
(Fig. 31C)

Bull. Misc. Inform. Kew 1937 (2): 106 (Sandwith 1937).

NOTE. — *Paloue induta* subsp. *induta* is apparently endemic to the Guiana Shield.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier 5785*.

INVENTORY DATA (FG). — 30 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.6$ cm.

[905] *Paloue leiogyne* (Ducke) Redden

Smithsonian Contr. Bot. 109: 23 (Redden 2018). — *Elizabetahia leiogyne* Ducke, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 4 (6): 727 (Ducke 1932).

HERBARIUM DATA (FG). — A single collection, *B.M. Boom & S.A. Mori* 1795.

SIZE. — Up to 30 m tall (Cowan & Berry 1998).

[906] *Paloue paraensis* (Ducke) Redden

Smithsonian Contr. Bot. 109: 26 (Redden 2018). — *Elizabetha paraensis* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 102 (Ducke 1922).

VERNACULAR NAMES. — Wp: amata'i • Cr: bwa-makak • Br: arapari-vermelho.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Grenand* 649.

SIZE. — Up to 50 m tall (Cowan & Berry 1998).

[907] *Paloue princeps* (R.H.Schomb. ex Benth.) Redden

Smithsonian Contr. Bot. 109: 27 (Redden 2018). — *Elizabetha princeps* R.H.Schomb. ex Benth., *J. Bot. [Hooker]* 2: 92 (Bentham 1840).

NOTE. — In their recent revision of *Paloue*, Redden *et al.* (2018) cite the basionym of *P. princeps*, *Elizabetha princeps* with the authorship “M.R.Schomb. ex Benth.”, and give the type “Guyana. 1839 (fl., fr.). R. Schomburgk s.n. (holotype: K!; isotype: F!)”. However, the original material collected in 1839 and reported by Hooker in 1840 was collected by Robert (Hermann) Schomburgk, not his younger brother (Moritz) Richard who first arrived in Guyana in 1841. Further, the specimen mentioned as isotype by Redden *et al.* (2018) (F, V0057797F) is a fragment of a lost B specimen (photo F neg. № 1560). If this specimen was collected by Richard Schomburgk, as the label seems to indicate, then it cannot be part of the original material. Among the three original specimens of *Elizabetha princeps* R.H.Schomb. ex Benth. at K, two are in Herbarium Benthamianum. One of them (K000264886) has a label “Holotype, *Elizabetha princeps* Schomb. ex Benth., det. R.S. Cowan, 10/1975” and is here designated as lectotype. The other two (K000264887, in Herbarium Benthamianum, and K000264888, in Herbarium Hookerianum) are isolectotypes.

VERNACULAR NAMES. — Wp: tapaka'i • Wn: wapaimë • Cr: wapa-mèg.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 4427.

INVENTORY DATA (FG). — 37 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42.1$ cm.

[908] *Paloue riparia* Pulle

Enum. Vasc. Pl. Surinam 212 (Pulle 1906), “*Paloveda*”.

VERNACULAR NAMES. — Te: amata'i.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 5046.

INVENTORY DATA (FG). — 6 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.1$ cm.

Genus *Paramachaerium* Ducke

[909] *Paramachaerium ormosioides* (Ducke) Ducke

Trop. Woods 41: 6 (Ducke 1935). — *Pterocarpus ormosioides* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 154 (Ducke 1922).

VERNACULAR NAMES. — Pa: muhut-ahavukunó • Nt: mongo gwegwe • Cr: moutouchi-montang • Fr: moutouchi montagne.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3564.

INVENTORY DATA (FG). — 25 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 87.1$ cm.

Genus *Parkia* R.Br.

[910] *Parkia decussata* Ducke

Notizbl. Bot. Gart. Berlin-Dahlem 11: 472 [11 July 1932] (Ducke 1932).

VERNACULAR NAMES. — Pa: sakeg-kamwi.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *S.A. Mori et al.* 25564.

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 77$ cm.

[911] *Parkia gigantocarpa* Ducke
(Fig. 31E)

Arch. Jard. Bot. Rio de Janeiro 1 (1): 19 (Ducke 1915).

VERNACULAR NAMES. — Wp: alamulu sili • Br: coré-grande, fava-ataná.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 1671.

INVENTORY DATA (FG). — 4 trees in 4 plots; $dbh_{\text{inv}} = 95$ cm.

[912] *Parkia nitida* Miq.

Stirp. Surinam. Select.: 7 [“1850” publ. Mar. 1851] (Miquel 1851).

Paryphosphaera arborea H.Karst., *Fl. Columb. [H. Karsten]* 2 (1): 7 [15 Dec. 1862] (Karsten 1862). — *Parkia arborea* (H.Karst.) J.F.Macbr., *Contr. Gray Herb.*, n.s., 59: 19 (Macbride 1919).

Parkia oppositifolia Spruce ex Benth., *Trans. Linn. Soc. London* 30 (3): 363 [10 Apr. 1875] (Bentham 1875).

Parkia paryphosphaera Benth., *Trans. Linn. Soc. London* 30 (3): 363 [10 Apr. 1875] (Bentham 1875).

Parkia sylvatica Pulle, *Recueil Trav. Bot. Néerl.* 4: 128 (Pulle 1907), *pro parte quoad flores tantum; leaves belong to Abarema claviflora* (Spruce) Kleinhoonte, *fide* Kleinhoonte (1940).

Parkia ingens Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 34 (Ducke 1925), *pro parte excl. specim. Ducke s.n.* (MG 16644).

Parkia inundabilis Ducke, *Arch. Inst. Biol. Veg.* 4 (1): 7 (Ducke 1938).

Parkia alliodora Ducke, *Bol. Técn. Inst. Agron. N. 2:* 9 (Ducke 1944).

VERNACULAR NAMES. — Pa: karevru, sakeg-kamwi • Ka: ululu'u • Wp: alamulu, alamulu si • Wn: heli apotpé • Nt: weti dondonisinga • Cr: bwa-ara, mal-akasya • Br: arapari-branco, arara-tucupi, caramuru, faveira-benguê, japacanim.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *M.-F. Prévost 3442*.

INVENTORY DATA (FG). — 157 trees in 106 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 148$ cm.

[913] *Parkia pendula* (Willd.) Benth. ex Walp.

Repert. Bot. Syst. [Walpers] 5 (4): 577 [14–15 Apr. 1846] (Walpers 1846). — *Inga pendula* Willd., *Sp. Pl.*, ed. 4 4 (2): 1025 [Apr. 1806] (Willdenow 1806). — *Mimosa pendula* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 47 [3 Sep. 1810] (Poirier 1810).

VERNACULAR NAMES. — Pa: sakeg • Ka: apakanilan, eleyululan, ululu'u • Te: dzelulu • Wp: yolulu, yululu • Wn: julu • Nt: kwata kama • Cr: bwa-ara, mal-bwa-makak • Fr: acacia mâle • Br: arara-tucupi, fava-bolota, faveira-de-chorão, joérana, visgueiro.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3678*.

INVENTORY DATA (FG). — 31 trees in 28 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 120$ cm.

[914] *Parkia reticulata* Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 126 (Ducke 1930).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & E. Fonty 5589*.

INVENTORY DATA (FG). — 10 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 70.9$ cm.

[915] *Parkia ulei* (Harms) Kuhlm. var. *surinamensis* Kleinhoonte

Recueil Trav. Bot. Néerl. 30: 169 [July 1933] (Kleinhoonte 1933).

Parkia microcephala Kleinhoonte, *Recueil Trav. Bot. Néerl.* 22: 411 [“1925” publ. Jan. 1926] (Kleinhoonte 1926).

NOTE. — Since a name has no priority outside its rank, and despite Rec. 24B.2 of the Code (Turland *et al.* 2018), the new name *P. ulei* var. *surinamensis* Kleinhoonte, created at varietal rank for *Parkia microcephala* Kleinhoonte, is legitimate.

VERNACULAR NAMES. — Ka: kuyali tapityi, kuyali topaitye • Wp: akiki nami, alapokwá'i wu • Nt: dondonisinga • Cr: mal-akasya • Br: esponjeira, paricá-de-esponja.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier 865*.

INVENTORY DATA (FG). — 24 trees in 19 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 71.3$ cm.

[916] *Parkia velutina* Benoist

Notul. Syst. (Paris) 3: 271 [7 May 1917] (Benoist 1917).

VERNACULAR NAMES. — Pa: sakeg-kamwi • Wp: alamulu pilá • Nt: baaka dondonisinga • Br: corezeiro, esponjeira.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *R. Benoist 1005* (original material P[P02436134, P02436135, P02436136]).

INVENTORY DATA (FG). — 63 trees in 51 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 91.4$ cm.

[917] *Parkia* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5101*.

INVENTORY DATA (FG). — 1 tree, $dbh = 32.5$ cm.

Genus *Peltogyne* Vogel

[918] *Peltogyne lecointei* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 96 (Ducke 1922), “*LeCointei*”.

VERNACULAR NAMES. — Pa: á-ivuniat • Wp: wila takulu pilá (for *Peltogyne* sp.) • Nt: dasitan, papaati • Cr: bwa-vyolèt • Fr: amarante • Br: coataquiçaua, guarabu, mulateiro, pau-roxo, violeta.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier 2448*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 88.8$ cm.

[919] *Peltogyne paniculata* Benth.

subsp. *pubescens* (Benth.) M.F.Silva

Acta Amazonica 6 (1, Suppl.): 38 (Silva 1976). — *Peltogyne pubescens* Benth., *J. Bot. [Hooker]* 2: 96 (Bentham 1840).

Peltogyne amplissima Pittier ex R.Knuth, *Repert. Spec. Nov. Regni Vég. Beih.* 43: 370 (Knuth 1927), *nom. nud.*

NOTE. — *Peltogyne paniculata* Benth. is hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: malako, šimililan, wewe pipyo • Wp: wila tátá • Wn: pawe, wapaimé • Nt: dasitan, papaati • Cr: bwa-vyolèt • Fr: amarante, bois violet • Br: escorrega-macaco, guarabu, mulateiro, pau-roxo, violeta.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2279*.

INVENTORY DATA (FG). — 28 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 58$ cm.

[920] *Peltogyne venosa* (Vahl) Benth.

Fl. Bras. [Martius] 15 (2): 233 [1 Dec. 1870] (Bentham 1870). — *Hymenaea venosa* Vahl, *Eclog. Amer.* 2: 31 (Vahl 1798).

VERNACULAR NAMES. — Ka: šimililan, wewe pipyo • Wn: pawe, weweimë • Nt: papaati • Cr: bwa-vyolèt • Fr: amarante, bois violet • Br: guarabu, mulateiro, pau-roxo, violeta.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *P.A. Sagot s.n.* (P[P03110846]).

INVENTORY DATA (FG). — 188 trees in 69 plots; $F_{\max} = 3.2\%$; $\text{dbh}_{\text{inv}} = 120 \text{ cm}$.

[921] *Peltogyne* sp. A

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 2447*.

INVENTORY DATA (FG). — 47 trees in 27 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 43 \text{ cm}$.

Genus *Pentaclethra* Benth.[922] *Pentaclethra macroloba* (Willd.) Kuntze

Revis. Gen. Pl. 1: 201 [5 Nov. 1891] (Kuntze 1891). — *Acacia macroloba* Willd., *Sp. Pl., ed. 4 4 (2)*: 1060 (printed 1054 in error) [Apr. 1806] (Willdenow 1806). — *Mimosa macroloba* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 66 [3 Sep. 1810] (Poiret 1810).

Acacia aspidioides G.Mey., *Prim. Fl. Esseq.* 165 [Nov. 1818] (Meyer 1818).

Pentaclethra filamentosa Benth., *J. Bot. [Hooker]* 2: 127 (Bentham 1840).

Pentaclethra brevifila Benth., *J. Bot. [Hooker]* 2: 128 (Bentham 1840).

Cailliea macrostachya Steud., *Flora* 26 (45): 759 [17 Dec. 1843] (Steu-del 1843). — *Dichrostachys macrostachya* (Steud.) Walp., *Repert. Bot. Syst. [Walpers]* 5 (4): 583 [14-15 Apr. 1846] (Walpers 1846).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: yuumwi • Ka: palawaksi • Cr: wapa-charles • Br: pracaxi.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier 901*; *M.-F. Prévost 4059*, 12 m × 13 cm.

Genus *Platymiscium* Vogel[923] *Platymiscium filipes* Benth.

J. Proc. Linn. Soc., Bot. 4 (Suppl.): 83 [Mar. 1860] (Bentham 1860).

HERBARIUM DATA (FG). — A single collection, *J. Martin s.n.* (original material K[K000530367]).

SIZE. — Brazil, Pará. *G.T. Prance 25763* (MO), 10 m × 15 cm.

[924] *Platymiscium pinnatum* (Jacq.) Dugand

Contr. Hist. Nat. Colomb. 1: 11 (Dugand 1938). — *Amerimnon pinnatum* Jacq., *Select. Stirp. Amer. Hist.*: 200 [5 Jan. 1763] (Jacquin 1763). — *Lonchocarpus amerimum DC.*, *Prodr. [A. P. de Candolle]* 2: 261 [mid Nov. 1825] (Candolle 1825), *nom. illeg. superfl.* (based on *Amerimnon pinnatum*).

Platymiscium polystachyum Benth., *Bot. Voy. Herald [Seemann]* 3: 111 [Nov. 1853] (Bentham 1853). — *Platymiscium pinnatum* subsp. *polystachyum* (Benth.) Klitg., *Kew Bull.* 54 (4): 973 ["1999" publ. 3 Feb. 2000] (Klitgaard 2000).

Platymiscium polystachyum var. *fendleri* Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 83 [Mar. 1860] (Bentham 1860).

Platymiscium dubium Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 125 [18 June 1918] (Pittier 1918).

VERNACULAR NAMES. — Ka: kunatepi • Wp: mutusi, mutusi sī • Nt: neko udu.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2950*.

INVENTORY DATA (FG). — 50 trees in 34 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 92 \text{ cm}$.

[925] *Platymiscium trinitatis* Benth.

(Fig. 31F)

J. Proc. Linn. Soc., Bot. 4 (Suppl.): 82 [Mar. 1860] (Bentham 1860).

Platymiscium duckei Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 83 (Huber 1910). — *Platymiscium trinitatis* var. *duckei* (Huber) Klitg., *Kew Bull.* 54 (4): 970 ["1999" publ. 3 Feb. 2000] (Klitgaard 2000).

Platymiscium nigrum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 157 (Ducke 1922). — *Platymiscium duckei* var. *nigrum* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 87 (Ducke 1925). — *Platymiscium trinitatis* var. *nigrum* (Ducke) Klitg., *Kew Bull.* 54 (4): 971 ["1999" publ. 3 Feb. 2000] (Klitgaard 2000).

VERNACULAR NAMES. — Br: guarapiranga, macacaúba.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier 3516*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 46.5 \text{ cm}$.

Genus *Pseudopiptadenia* Rauschert[926] *Pseudopiptadenia psilostachya* (DC.) G.P.Lewis & M.P.Lima

Arch. Jard. Bot. Rio de Janeiro 30: 55 ["1989-90" publ. 1991] (Lewis & Lima 1991), excluding *P. suaveolens* from synonymy. — *Acacia psilostachya* DC., *Prodr. [A. P. de Candolle]* 2: 457 [mid Nov. 1825] (Candolle 1825). — *Piptadenia psilostachya* (DC.) Benth., *J. Bot. [Hooker]* 4: 336 (Bentham 1842). — *Newtonia psilostachya* (DC.) Brenan, *Kew Bull.* 10 (2): 182 [12 Sep. 1955] (Brenan 1955).

Entada urbaeana C.Presl, *Abb. Königl. Böhm. Ges. Wiss.*, ser. 5, 6: 566 (Presl 1851).

VERNACULAR NAMES. — Ka: šimililan, walekone • Te: akiki'i • Wp: wilá lē, wilá lē pilá, wilá lē sī • Nt: alimiaw, pikimisiki • Cr: alimiao • Br: angico, paricá-branco, timborana-foliolo-graúdo.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *J. Martin* 2 (original material F[V0360975F], K[K000504699], US[00000274]).

INVENTORY DATA (FG). — 128 trees in 50 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 121$ cm.

[927] *Pseudopiptadenia suaveolens* (Miq.) J.W.Grimes

Brittonia 45 (1): 27 (Grimes 1993). — *Piptadenia suaveolens* Miq., *Linnaea* 18: 589 ["1844" publ. prob. Aug. 1845] (Miquel 1845). — *Newtonia suaveolens* (Miq.) Brenan, *Kew Bull.* 10 (2): 182 [12 Sep. 1955] (Brenan 1955).

VERNACULAR NAMES. — Pa: mbagwi-puvemna, mbarui-puvemna • Ka: šimililan, walekone • Wp: wilá lē, wilá lē pilá • Wn: alimiap, alimijaop • Nt: alimiaw, pikimisiki • Cr: alimiao • Br: angico, paricá-branco, timborana-foliolo-fina.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2121.

INVENTORY DATA (FG). — 120 trees in 68 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 160$ cm.

Genus *Pterocarpus* Jacq.

[928] *Pterocarpus officinalis* Jacq.

Select. Stirp. Amer. Hist.: 283 [5 Jan. 1763] (Jacquin 1763). — *Lingoum officinale* (Jacq.) Kuntze, *Revis. Gen. Pl.* 1: 193 [5 Nov. 1891] (Kuntze 1891).

Pterocarpus draco L., *Sp. Pl.*, ed. 2, 2: 1662 [Aug. 1763] (Linnaeus 1763). — *Moutouchi draco* (L.) Benth., *Commentat. Legum. Gen. 30* [June 1837] (Bentham 1837), "Moutouchia".

Moutouchi suberosa Aubl., *Hist. Pl. Guiane* 2: 748 [Jun.-Dec. 1775] (Aublet 1775). — *Pterocarpus suberosus* (Aubl.) Pers., *Syn. Pl. [Persoon] 2* (2): 277 [Sep. 1807] (Persoon 1807). — *Pterocarpus moutouchi* Poir., *Tabl. Encycl.* 3: 161 [22 Feb. 1823] (Poirier 1823), "Montouchi", nom. illeg. superfl. (based on *Moutouchi suberosa*).

Pterocarpus hemipterus Gaertn., *Fruct. Sem. Pl.* 2: 351 [Apr.-May 1791] (Gaertner 1791), "hemiptera".

Pterocarpus crispatus Moc. & Sessé ex DC., *Prodri. [A. P. de Candolle]* 2: 418 [mid Nov. 1825] (Candolle 1825). — *Moutouchi crispata* (Moc. & Sessé ex DC.) Benth., *Commentat. Legum. Gen.*: 30 [June 1837] (Bentham 1837), "Moutouchia".

Pterocarpus belizensis Standl., *Trop. Woods* 7: 6 (Standley 1926).

VERNACULAR NAMES. — Pa: muhut • Ka: mutusi, tiyapotano mutusi • Te: bututsi • Wp: mutusi, mutusi e'e, mutusi pilá • Wn: müümili • Nt: gwegwe, sabana gwegwe • Cr: moutouchi-marikaj, moutouchi-rivié, moutouchi-savann • Fr: moutouchi marécage, moutouchi rivière • Br: corticeira, mututy.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *M.-F. Prévost* 752.

INVENTORY DATA (FG). — 120 trees in 18 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 50$ cm.

[929] *Pterocarpus rohrii* Vahl

Symb. Bot. [Vahl] 2: 79 [Jul.-Dec. 1791] (Vahl 1791). — *Amphymenium rohrii* (Vahl) Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 380 [12 July 1824] (Kunth 1824), nom. inval. (genus name and epithet not associated). — *Lingoum rohrii* (Vahl) Kuntze, *Revis. Gen. Pl.* 1: 193 [5 Nov. 1891] (Kuntze 1891).

Pterocarpus apalatoa Rich., *Actes Soc. Hist. Nat. Paris* 1: 111 [Oct. 1792] (Richard 1792).

Amphymenium villosum Mart. ex Benth., *Commentat. Legum. Gen. 31* [June 1837] (Bentham 1837). — *Pterocarpus villosum* (Mart. ex Benth.) Benth., *J. Proc. Linn. Soc., Bot.* 4 (Suppl.): 79 [Mar. 1860] (Bentham 1860). — *Lingoum villosum* (Mart. ex Benth.) Kuntze, *Revis. Gen. Pl.* 1: 193 [5 Nov. 1891] (Kuntze 1891).

Phellocarpus floridus Benth., *Commentat. Legum. Gen. 42* [June 1837] (Bentham 1837). — *Piscidia florida* Mart. ex Benth., *Commentat. Legum. Gen. 42* [June 1837] (Bentham 1837), nom. nud. pro syn.

Phellocarpus laxiflorus Benth., *Commentat. Legum. Gen. 42* [June 1837] (Bentham 1837).

Pterocarpus violaceus Vogel, *Linnaea* 11: 416 (Vogel 1837). — *Lingoum violaceum* (Vogel) Kuntze, *Revis. Gen. Pl.* 1: 193 [5 Nov. 1891] (Kuntze 1891).

Pterocarpus rufescens Benth., *J. Proc. Linn. Soc., Bot.* 4 (Suppl.): 79 [Mar. 1860] (Bentham 1860). — *Lingoum rufescens* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 193 [5 Nov. 1891] (Kuntze 1891).

Pterocarpus violaceus var. *angustifolia* Benth., *J. Proc. Linn. Soc., Bot.* 4 (Suppl.): 79 [Mar. 1860] (Bentham 1860).

Pterocarpus hayesii Hemsl., *Diagn. Pl. Nov. Mexic.* 1: 8 [July 1878] (Hemsley 1878).

Pterocarpus floribundus Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 123 [18 June 1918] (Pittier 1918), nom. illeg. hom., non Wall. (Wall 1831) nec Kuntze (1891).

Pterocarpus steinbachianus Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 1039 [15 Nov. 1926] (Harms 1926).

Pterocarpus rupestris Pittier, *Arb. Arbust. Venez.* 6-8: 97 [Aug.-Sep. 1927] (Pittier 1927).

Pterocarpus reticulatus Standl., *Trop. Woods* 16: 38 (Standley 1928).

Pterocarpus magnicarpus Schery, *Fieldiana, Bot.* 28 (2): 261 (Schery 1952), "magnicarpa".

Pterocarpus rohrii var. *rubiginosus* Schery, *Fieldiana, Bot.* 28 (2): 261 (Schery 1952).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: sakaugi-kamwi, sakauri-kamwi • Ka: mutusilan • Nt: baaka panga, baaka wapa, gwegwe • Cr: moutouchi-ribanné • Fr: moutouchi rubané, moutouchi savane • Br: mututy-da-terra-firme, sangué-de-galo.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost 3667*.

INVENTORY DATA (FG). — 23 trees in 17 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 57.2$ cm.

[930] *Pterocarpus santalinoides* L'Hér. ex DC.

Prod. [A. P. de Candolle] 2: 419 [mid Nov. 1825] (Candolle 1825).

Pterocarpus esculentus Schumach. & Thonn., *Beskr. Guin. Pl. 330* (Schumacher & Thonning 1827). — *Lingoum esculentum* (Schumach. & Thonn.) Kuntze, *Revis. Gen. Pl. 1*: 193 [5 Nov. 1891] (Kuntze 1891).

Pterocarpus michelii Britton, *Ann. New York Acad. Sci. 7*: 86 (Britton 1893).

Pterocarpus amazonicus Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2)*: 402 (Huber 1909).

Pterocarpus grandis R.S.Cowan, *Mem. New York Bot. Gard. 10 (4)*: 86 (Cowan 1961).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: sakaugi, sakauri • Ka: konomelulan • Wp: mutusi, mutusi sī • Wn: mümili • Cr: moutouchi-ribanné • Br: mututy-da-terra-firme, sangue-de-galo.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 2057*.

INVENTORY DATA (FG). — 6 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 50$ cm.

Genus *Recordoxylon* Ducke

[931] *Recordoxylon speciosum* (Benoist) Gazel ex Barneby (Fig. 32A)

Brittonia 45 (3): 235 (Barneby 1993). — *Melanoxyロン speciosum* Benoist, *Bull. Mus. Natl. Hist. Nat. 26*: 87 (Benoist 1920).

Melanoxyロン amazonicum Ducke, *Trop. Woods 31*: 15 (Ducke 1932). — *Recordoxylon amazonicum* (Ducke) Ducke, *Trop. Woods 39*: 17 (Ducke 1934).

VERNACULAR NAMES. — Nt: bunahati tyabisi, buunaati tyabisi • Cr: wakapu-guitin • Fr: wacapou guittin • Br: barúna.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *G. Wachenheim 99* (holo-, P[P01818107]; iso-, P[P01818108, P01818109]).

INVENTORY DATA (FG). — 290 trees in 57 plots; $F_{\max} = 1.9\%$; $\text{dbh}_{\text{inv}} = 71.8$ cm.

Genus *Senegalia* Raf.

[932] *Senegalia polypylla* (DC.) Britton & Rose ex Britton & Killip (Fig. 32B)

Ann. New York Acad. Sci. 35: 142 [1 Apr. 1936] (Britton & Killip 1936). — *Acacia polypylla* DC., *Cat. Pl. Horti Monsp.*: 74 (Candolle

1813). — *Mimosa polypylla* (DC.) Poir., *Encycl. [J. Lamarck et al.] Suppl. 5: 530* [1 Nov. 1817] (Poirier 1817).

Acacia riparia Bertero ex Spreng., *Syst. Veg. [Sprengel] 3*: 142 [Jan.-Mar. 1826] (Sprengel 1826), *nom. illeg. hom.*, *non* Kunth (1824).

Leucaena boliviiana Rusby, *Bull. New York Bot. Gard. 8 (28)*: 91 [23 Nov. 1912] (Rusby 1912).

Senegalia lobana Britton & Killip, *Ann. New York Acad. Sci. 35*: 142 [1 Apr. 1936] (Britton & Killip 1936).

Acacia glomerosa Benth., *London J. Bot. 1*: 521 (Bentham 1842). — *Senegalia glomerosa* (Benth.) Britton & Rose, *N. Amer. Fl. 23 (2)*: 116 [25 Sep. 1928] (Britton & Rose 1928).

Senegalia langlassae Britton & Rose, *N. Amer. Fl. 23 (2)*: 116 [25 Sep. 1928] (Britton & Rose 1928). — *Acacia langlassae* (Britton & Rose) Bullock, *Bull. Misc. Inform. Kew 1939 (1)*: 2 [15 Feb. 1939] (Bullock 1939).

Senegalia tomentella Britton & Killip, *Ann. New York Acad. Sci. 35*: 145 [1 Apr. 1936] (Britton & Killip 1936).

VERNACULAR NAMES. — Br: juqueiri-guaçu, paricá-branco, paricarana-de-espinho.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 6009*.

INVENTORY DATA (FG). — 5 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 58.6$ cm.

Genus *Senna* Mill.

[933] *Senna multijuga* (Rich.) H.S.Irwin & Barneby

Mem. New York Bot. Gard. 35: 492 (Irwin & Barneby 1982). — *Cassia multijuga* Rich., *Actes Soc. Hist. Nat. Paris 1*: 108 [Oct. 1792] (Richard 1792). — *Peiranisia multijuga* (Rich.) Britton & P.Wilson, *Sci. Surv. Porto Rico & Virgin Islands 6*: 350 (Britton & Wilson 1926). — *Chamaesenna multijuga* (Rich.) Pittier, *Third Conf. Interamer. Agric. Caracas: 376* (Pittier 1945).

Cassia calliantha G.Mey., *Prim. Fl. Esseq. 169* [Nov. 1818] (Meyer 1818).

Cassia richardiana Kunth, *Mimos.: 139* (Kunth 1823).

Cassia fulgens Wall. ex Vogel, *Gen. Cass. Syn.: 44* (Vogel 1837).

Cassia ampliflora Steud., *Flora 26 (45)*: 760 [17 Dec. 1843] (Steudel 1843).

Peiranisia aristulata Britton & Killip, *Ann. New York Acad. Sci. 35*: 182 [1 Apr. 1936] (Britton & Killip 1936).

VERNACULAR NAMES. — Pa: datka-arivra, karegut-seine • Ka: malimali, palalapo • Wp: malimali sī • Wn: ididiju, kotkotoloju • Nt: kabanafoo, sabana foo • Br: acácia, amarelinha.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *J.B. Leblond 168* (holo-, P[P00798376]; iso-, G[G00341799], F[V0057614F], P[P00296957, P00798377]).

SIZE. — Up to 25 m tall (Barneby & Heald 2002).

[934] *Senna reticulata* (Willd.) H.S.Irwin & Barneby

Mem. New York Bot. Gard. 35: 458 (Irwin & Barneby 1982). — *Cassia reticulata* Willd., *Enum. Pl. [Willdenow]*: 443 [Apr. 1809] (Willdenow 1809). — *Chamaesenna reticulata* (Willd.) Pittier, *Trab. Mus. Comercial Venezuela* 3: 160 (Pittier 1928).

Cassia strobilacea Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 347 [12 July 1824] (Kunth 1824).

Cassia tarantan Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 348 [12 July 1824] (Kunth 1824).

Cassia dumetorum Bertero ex DC., *Prodr. [A. P. de Candolle]* 2: 499 [mid Nov. 1825] (Candolle 1825).

*Cassia annunciat*a E.H.L.Krause, *Beih. Bot. Centralbl.* 32 (2): 345 (Krause 1914).

VERNACULAR NAMES. — Pa: wahamuvie • Ka: anatapali • Wp: pole • Wn: hawahawaju • Cr: bwa-dat • Br: mata-pasto-grande.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3602*.

SIZE. — Colombia. *A. Prieto 173* (MO), 8 m × 15 cm.

Genus *Spirotropis* Tul.

[935] *Spirotropis longifolia* (DC.) Baill. (Fig. 32C)

Hist. Pl. [Baillon] 2: 364 [Feb.-May 1870] (Baillon 1870). — *Swartzia longifolia* DC., *Prodr. [A. P. de Candolle]* 2: 423 [mid Nov. 1825] (Candolle 1825). — *Spirotropis candolleana* Tul., *Arch. Mus. Hist. Nat.* 4: 113 (Tulasne 1844), nom. illeg. superfl. (based on *Spirotropis longifolia*). — *Tounatea longifolia* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 211 [5 Nov. 1891] (Kuntze 1891), “*Tunatea*”.

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *M.-F. Prévost 1619*.

INVENTORY DATA (FG). — 1686 trees in 7 plots; $F_{\max} = 65\%$; $dbh_{inv} = 65$ cm.

Genus *Stryphnodendron* Mart.

[936] *Stryphnodendron guianense* (Aubl.) Benth. (Fig. 32D)

Trans. Linn. Soc. London 30 (3): 374 [10 Apr. 1875] (Bentham 1875). — *Mimosa guianensis* Aubl., *Hist. Pl. Guiane* 2: 938 [Jun.-Dec. 1775] (Aublet 1775). — *Acacia guianensis* (Aubl.) Willd., *Sp. Pl. ed. 4 4 (2)*: 1061 [Apr. 1806] (Willdenow 1806). — *Folianthera guianensis* (Aubl.) Raf., *Sylva Tellur.*: 120 (Rafinesque 1838). — *Piptadenia guianensis* (Aubl.) Benth., *J. Bot. [Hooker]* 4: 335 (Bentham 1842).

Stryphnodendron purpureum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 16 (Ducke 1915).

VERNACULAR NAMES. — Pa: iduk-veiti, yuumwi • Br: faveira-camuzé.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM001135589]).

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 62.4$ cm.

[937] *Stryphnodendron moricolor* Barneby & J.W.Grimes *Brittonia* 36 (1): 45 (Barneby & Grimes 1984).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Nt: nyon udu.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15236* (holo-, P[P00077203]; iso-, MO[MO-713028], NY[00003370], P[P00710285]).

INVENTORY DATA (FG). — 27 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{inv} = 79.2$ cm.

[938] *Stryphnodendron polystachyum* (Miq.) Kleinhoonte *Recueil Trav. Bot. Néerl.* 22: 416 [“1925” publ. Jan. 1926] (Kleinhoonte 1926). — *Piptadenia polystachya* Miq., *Linnaea* 18: 590 [“1844” publ. prob. Aug. 1845] (Miquel 1845).

VERNACULAR NAMES. — Ka: tipulu sepeipyo • Wp: tatu'i • Nt: nyon udu • Br: taxirana-branca.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3758*.

INVENTORY DATA (FG). — 31 trees in 27 plots; $F_{\max} < 1\%$; $dbh_{inv} = 87.2$ cm.

[939] *Stryphnodendron pulcherrimum* (Willd.) Hochr.

Bull. New York Bot. Gard. 6 (21): 274 [27 Sep. 1910] (Hochreutiner 1910). — *Acacia pulcherrima* Willd., *Sp. Pl. ed. 4 4 (2)*: 1061 [Apr. 1806] (Willdenow 1806). — *Mimosa pulcherrima* (Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 66 [3 Sep. 1810] (Poiret 1810).

Stryphnodendron floribundum Benth., *J. Bot. [Hooker]* 4: 343 (Bentham 1842). — *Stryphnodendron guianense* f. *floribundum* (Benth.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 250 (Ducke 1925).

Stryphnodendron angustum Benth., *Trans. Linn. Soc. London* 30 (3): 375 [10 Apr. 1875] (Bentham 1875).

Stryphnodendron melinonii Sagot, *Ann. Sci. Nat., Bot. sér. 6, 13*: 322 (Sagot 1882), “*Melinonis*”.

Piptadenia cobi Rizzini & A.Mattos, *Anais Acad. Brasil. Ci.* 40: 233 (Rizzini & Mattos 1968).

NOTE. — The epithet “*melinonis*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland et al. 2018: Art. 60.8).

VERNACULAR NAMES. — Ka: apakanilan, eleyululan • Br: timbaúba.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (type of *Stryphnodendron melinonii*: P[P00199447], P[P00199448]).

INVENTORY DATA (FG). — 10 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{inv} = 44.2$ cm.

[940] *Stryphnodendron* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier* 2302, “Grand arbre de canopée”.

Genus *Swartzia* Schreb.

[941] *Swartzia amshoffiana* R.S.Cowan

Fl. Neotrop. Monogr. 1: 159 [10 May 1968] (Cowan 1968).

Swartzia apetala Radde var. *acuminata* Amshoff, *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 52: 34 (Amshoff 1939).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *F. Hallé* 626.

INVENTORY DATA (FG). — 27 trees in 9 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 32.3$ cm.

[942] *Swartzia aptera* DC.

Prodr. [A. P. de Candolle] 2: 423 [mid Nov. 1825] (Candolle 1825). — *Tounatea aptera* (DC.) Taub., *Bot. Centralbl.* 47: 390 [31 Sept 1891] (Taubert 1891), “*Tunatea*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wap-kamwi-priyu • Wp: kumalaki.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[BM000952072], F[V0059859F], FI[FI004911], LE[LE00013722], P[P01818008, P01818009, P01818010], R[R000069375], US[00037067]).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 12.3$ cm.

[943] *Swartzia arborescens* (Aubl.) Pittier

J. Wash. Acad. Sci. 11: 157 (Pittier 1921). — *Possira arborescens* Aubl., *Hist. Pl. Guiane* 2: 934 [Jun.-Dec. 1775] (Aublet 1775). — *Possira triphylla* Sw., *Prodr. [Swartz]*: 82 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. superfl. (based on *Possira arborescens*). — *Swartzia triphylla* Willd., *Sp. Pl. ed. 4 2 (2)*: 1220 [Dec. 1799] (Willdenow 1799), nom. illeg. superfl. (based on both *Possira arborescens* and *P. triphylla*, the latter being itself illegitimate, because also based on *P. arborescens*). — *Tounatea arborescens* (Aubl.) Britton, *Bull. Torrey Bot. Club* 16 (12): 325 [10 Dec. 1889] (Britton 1889).

Rittera dodecandra Vahl, *Symb. Bot. [Vahl]* 2: 60 [Jul.-Dec. 1791] (Vahl 1791). — *Swartzia dodecandra* (Vahl) Willd., *Sp. Pl. ed. 4 2 (2)*: 1220 [Dec. 1799] (Willdenow 1799). — *Possira dodecandra* (Vahl) Poir., *Encycl. [J. Lamarck et al.]* 5: 571 [9 Jan. 1804] (Poirer 1804). — *Tounatea dodecandra* (Vahl) Taub., *Bot. Centralbl.* 47: 391 [31 Sept 1891] (Taubert 1891), “*Tunatea*”.

Swartzia parviflora DC., *Prodr. [A. P. de Candolle]* 2: 423 [mid Nov. 1825] (Candolle 1825).

Rittera triphylla Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 460 (Steudel 1841), nom. nud. pro syn.

Swartzia bifida Steud., *Flora* 26 (45): 757 [17 Dec. 1843] (Steudel 1843).

Swartzia rariflora Hoehne, *Comm. Lin. Telegr. Bot.* 74 (12): 16 [Jul.-Aug. 1922] (Hoehne 1922).

NOTE. — The mention by Steudel (1841: 460) of “*Rittera triphylla* Sw.” was a mistake. He actually referred to *Possira triphylla* Sw.

VERNACULAR NAMES. — Pa: paig-seine, pairi-seine, timuvuyen-kamwi • Ka: apukuitalan, tipulu apukuityalan • Wp: winâme'i, winâme'i piyû • Nt: adugwe, makwelebo • Cr: moutouchi-nwé • Fr: moutouchi noir • Br: jenipapinho, pacapuá.

HERBARIUM DATA (FG). — 70 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000952073]).

INVENTORY DATA (FG). — 82 trees in 66 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 46.4$ cm.

[944] *Swartzia bannia* Sandwith

Contr. Gray Herb. 165: 28 (Sandwith 1947).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: aliyanâ'i, talala, ulana beta.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *B. Torke & S. Gonzalez* 191.

INVENTORY DATA (FG). — 30 trees in 4 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 77.2$ cm.

[945] *Swartzia benthamiana* Miq.

Stirp. Surinam. Select.: 15 [“1850” publ. Mar. 1851] (Miquel 1851). — *Tounatea benthamiana* (Miq.) Taub., *Bot. Centralbl.* 47: 390 [31 Sept 1891] (Taubert 1891), “*Tunatea*”.

VERNACULAR NAMES. — Ka: aliyanâ'i, ulana beta • Wn: alalawaju • Nt: mutusi • Cr: moutouchi-montangn.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *Service Forestier* 7075.

INVENTORY DATA (FG). — 9 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 43$ cm.

[946] *Swartzia canescens* Torke

Bot. J. Linn. Soc. 153 (3): 344 [15 Mar. 2007] (Torke 2007).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: mutusi piyû, winâme'i, winâme'i sî • Nt: adugwe, makwelebo • Cr: moutouchi-nwé • Fr: moutouchi noir.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *R.A.A. Oldeman & C. Sastre* 167 (holo-, CAY[CAY009778]; iso-, A[A00299105], COL[COL00005984], NY[00990840], PI[P01818030, P01818031], U[U0193371], US[00423115], VEN[VEN103660]).

INVENTORY DATA (FG). — 151 trees in 69 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 84.2$ cm.



FIG. 32. — Leguminosae: **A**, *Recordoxylon speciosum* (Benoist) Gazel ex Barneby (*D. Sabatier & C. Geniez* 6283); **B**, *Senegalia polypylla* (DC.) Britton & Killip (*D. Sabatier et al.* 6009); **C**, *Spirotropis longifolia* (DC.) Baill.; **D**, *Stryphnodendron guianense* (Aubl.) Benth.; **E**, *Zygia sabatieri* Barneby & J.W. Grimes (*D. Sabatier & M.-F. Prévost* 4356); **F**, *Tachigali paniculata* Aubl. (*J.-F. Molino & D. Sabatier* 2246). A-D, © D. Sabatier/IRD; E, © M.-F. Prévost/IRD; F, © J.-F. Molino/IRD.

[947] *Swartzia grandifolia* Bong. ex Benth.

J. Bot. [Hooker] 2: 85 (Bentham 1840). — *Tounatea grandifolia* (Bong. ex Benth.) Taub., *Bot. Centralbl.* 47: 391 [31 Sept 1891] (Taubert 1891), “*Tounatea*”.

VERNACULAR NAMES. — Ka: aliyaná'i, ulana beta • Cr: moutouchi-montangn.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D. Sabatier* 839.

INVENTORY DATA (FG). — 71 trees in 54 plots; $F_{\max} < 1\%$; dbh_{inv} = 27.2 cm.

[948] *Swartzia guianensis* (Aubl.) Urb.

Symb. Antill. [Urban] 5 (3): 365 [20 May 1908] (Urban 1908). — *Tounatea guianensis* Aubl., *Hist. Pl. Guiane* 1: 549 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Swartzia alata* Willd., *Sp. Pl., ed. 4* 2 (2): 1220 [Dec. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Tounatea guianensis*).

Swartzia leblondii R.S.Cowan, *Fl. Neotrop. Monogr.* 1: 145 [10 May 1968] (Cowan 1968).

NOTES. — Known only from the Guiana Shield. *Swartzia leblondii* is placed in synonymy here according to B. Torke (pers. comm. 2013).

VERNACULAR NAMES. — Ka: asemunusi, malay elepali • Nt: adugwe, makwelebo • Fr: moutouchi noir.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000952077](#)]).

INVENTORY DATA (FG). — 216 trees in 94 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 26.9 \text{ cm}$.

[949] *Swartzia hostmannii* Benth.

Fl. Bras. [Martius] 15 (2): 18 [1 Dec. 1870] (Bentham 1870), “hostmanni”. — *Tounatea hostmannii* (Benth.) Taub., *Bot. Centralbl.* 47: 391 [31 Sept 1891] (Taubert 1891), “*Tunatea Hostmanni*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: sopu uwii, sopu wiwii.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *C. Sastre & D.A. Bell 8125*.

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[950] *Swartzia longicarpa* Amshoff

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 52: 38 (Amshoff 1939).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville B-4563*.

SIZE. — Up to 19 cm dbh (Cowan 1968).

[951] *Swartzia oblanceolata* Sandwith

Bull. Misc. Inform. Kew 1934 (9): 363 [17 Dec. 1934] (Sandwith 1934).

VERNACULAR NAMES. — Wp: mēwā • Br: pau-de-sangue.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *M.-F. Prévost 3207*.

INVENTORY DATA (FG). — 31 trees in 22 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 60 \text{ cm}$.

[952] *Swartzia panacoco* (Aubl.) R.S.Cowan var. *panacoco*

Fl. Neotrop. Monogr. 1: 38 [10 May 1968] (Cowan 1968). — *Robinia panacoco* Aubl., *Hist. Pl. Guiane* 2: 768 [Jun.-Dec. 1775] (Aublet 1775). — *Robinia tomentosa* Willd., *Sp. Pl.*, ed. 4 3 (2): 1134 [1-10 Nov. 1802] (Willdenow 1802), nom. illeg. superfl. (based on *Robinia panacoco*). — *Swartzia tomentosa* DC., *Prodr. [A. P. de Candolle]* 2: 423 [mid Nov. 1825] (Candolle 1825), nom. illeg. superfl. (based on *Robinia panacoco* and *R. tomentosa*, the latter also illegitimate). — *Tounatea panacoco* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 211 [5 Nov. 1891] (Kuntze 1891), “*Tunatea Panacoco*”.

Swartzia similis Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 297 (Benoist 1919).

VERNACULAR NAMES. — Pa: wanaku-danō, wanaku-divyo • Ka: aliyanā'i, anakoko, panakoko, ulana beta • Te: tatu tunu • Wp: paa letowape • Wn: onohkowe, wapotoimë • Nt: bwasiman weko, kaka a buku, kokobe weko • Cr: panakoko • Fr: féréol, panacoco • Br: coração-de-negro, moçataíba, paracutaca.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[[BM000931982](#), [BM000931983](#)] designated by Cowan [1968: 38]; isolecto-, P[[P02272924](#)]).

INVENTORY DATA (FG). — 175 trees in 109 plots; $F_{\max} = 1.3\%$; $\text{dbh}_{\text{inv}} = 63.7 \text{ cm}$.

[953] *Swartzia panacoco* var. *sagotii* (Sandwith) R.S.Cowan

Fl. Neotrop. Monogr. 1: 34 [10 May 1968] (Cowan 1968). — *Swartzia tomentosa* var. *sagotii* Sandwith, *Bull. Misc. Inform. Kew* 1934 (9): 357 [17 Dec. 1934] (Sandwith 1934), “*Sagoti*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: paa letowape.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D. Sabatier 824*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 24.4 \text{ cm}$.

[954] *Swartzia peruviana* (R.S.Cowan) Torke

Novon 17 (1): 115 [23 Apr. 2007] (Torke 2007). — *Swartzia brachyrachis* Harms var. *peruviana* R.S.Cowan, *Fl. Neotrop. Monogr.* 1: 194 [10 May 1968] (Cowan 1968).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2287*.

INVENTORY DATA (FG). — 13 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 83.7 \text{ cm}$.

[955] *Swartzia polyphylla* DC.

Prodri. [A. P. de Candolle] 2: 424 [mid Nov. 1825] (Candolle 1825). — *Tounatea polyphylla* (DC.) Taub., *Bot. Centralbl.* 47: 392 [31 Sept 1891] (Taubert 1891), “*Tunatea*”.

Swartzia acuminata Willd. ex Vogel, *Linnaea* 11: 173 (Vogel 1837). — *Tounatea acuminata* (Willd. ex Vogel) Taub., *Bot. Centralbl.* 47: 390 [31 Sept 1891] (Taubert 1891), “*Tunatea*”.

Swartzia acuminata var. *platygynne* Benth., *Fl. Bras. [Martius]* 15 (2): 37 [1 Dec. 1870] (Bentham 1870). — *Swartzia platygynne* (Benth.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 127 (Ducke 1922).

Tounatea acuminata var. *puberula* Taub., *Flora* 75 [n.s. 50]: 81 (Taubert 1892). — *Swartzia acuminata* var. *puberula* (Taub.) Glaz., *Bull. Soc. Bot. France* 53 (Mém. 3b): 155 (Glaziou 1906).

Swartzia acuminata var. *tridynamia* Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 2: 506 (Huber 1898).

Swartzia opacifolia J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13* (3.1): 226 (Macbride 1943).

Swartzia urubuensis Ducke, *Bol. Técn. Inst. Agron. N. 2*: 21 (Ducke 1944).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: isuu-á, isuu-ára • Ka: palakusinian, palakusinien, tipulu apukuityalan • Wp: kupemi • Wn: sihkëpani • Nt: bugu bugu, musansi • Cr: bwa-krobo, bwa-pagay • Fr: bois corbeau, bois pagaie • Br: arabá, paracutaca-da-terra-firme, pau-de-remo, pitaica.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (original material not traced); *P.A. Sagot 1199* (K[K000500935, K000500936, K000500937], MPU[MPU013120]).

INVENTORY DATA (FG). — 332 trees in 144 plots; $F_{\max} = 1.8\%$; $\text{dbh}_{\text{inv}} = 172.8\text{ cm}$.

[956] *Swartzia sericea* Vogel

Linnaea 11: 176 (Vogel 1837). — *Tounatea sericea* (Vogel) Kuntze, *Revis. Gen. Pl. 1*: 211 [5 Nov. 1891] (Kuntze 1891), “*Tunatea*”.

Swartzia erythrocarpa Spruce ex Benth., *Fl. Bras. [Martius]* 15 (2): 28 [1 Dec. 1870] (Bentham 1870), *nom. nud. pro syn.*

VERNACULAR NAMES. — Wp: monokoe u, onokoe u, teposi kili.

HERBARIUM DATA (FG). — A single collection, *J. Martin s.n.* “in Herb. Kth.” (original material, not traced).

SIZE. — Up to 15 m tall (Cowan 1968).

Genus *Tachigali* Aubl.

[957] *Tachigali amplifolia* (Ducke) Barneby

Brittonia 48 (2): 182 (Barneby 1996). — *Sclerolobium amplifolium* Ducke, *Arch. Inst. Biol. Veg. 2* (1): 43 [Sep. 1935] (Ducke 1935).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier 1546*.

INVENTORY DATA (FG). — 30 trees in 10 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 80.9\text{ cm}$.

[958] *Tachigali glauca* Tul.

Arch. Mus. Hist. Nat. 4: 165 (Tulasne 1844).

Sclerolobium myrmecophilum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 30 (Ducke 1915). — *Tachigali myrmecophila* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 91 (Ducke 1922), “*Tachigalia*”.

VERNACULAR NAMES. — Wn: jenejapo • Br: taxi-preto.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *F.M.R. Leprieur 336* (holo-, P[P00571997]; iso-, P[P00572002, P00577072, P00577073]); *D. Sabatier & J.-F. Molino 5653*, 40 m × 80 cm.

[959] *Tachigali guianensis* (Benth.) Zarucchi & Herend.

Monogr. Syst. Bot. Missouri Bot. Gard. 45: 1254 [1 Oct. 1993] (Zarucchi & Herendeen 1993). — *Sclerolobium guianense* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 237 (Bentham 1850).

Sclerolobium radlkoferi Rusby, *Mem. Torrey Bot. Club* 6 (1): 26 (Rusby 1896). — *Sclerolobium guianense* var. *radlkoferi* (Rusby) Dwyer, *Lloydia* 20 (2): 98 (Dwyer 1957).

Sclerolobium subbullatum Ducke, *Arch. Inst. Biol. Veg. 2* (1): 42 [Sep. 1935] (Ducke 1935). — *Tachigali subbullata* (Ducke) L.F.Gomes da Silva & H.C.Lima, *Rodriguésia* 58 (2): 400 (Gomes & Lima 2007), “*subbullata*”.

Sclerolobium uleanum Harms, *Verb. Bot. Vereins Prov. Brandenburg* 48: 168 [“1906” publ. 8 Mar. 1907] (Harms 1907). — *Sclerolobium tinctorium* Benth. var. *uleanum* (Harms) Dwyer, *Lloydia* 20 (2): 95 (Dwyer 1957). — *Tachigali uleana* (Harms) Zarucchi & Herend., *Monogr. Syst. Bot. Missouri Bot. Gard.* 45: 1254 [1 Oct. 1993] (Zarucchi & Herendeen 1993).

VERNACULAR NAMES. — Ka: alaulama, alawata muleli, wa:u • Wp: tulili wala, yapakaní'i • Cr: sèd-rémi, tachi.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *Service Forestier 6096*.

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 64\text{ cm}$.

[960] *Tachigali melanocarpa* (Ducke) van der Werff

Ann. Missouri Bot. Gard. 95 (4): 649 [30 Dec. 2008] (van der Werff 2008). — *Sclerolobium melanocarpum* Ducke, *Arch. Inst. Biol. Veg. 2* (1): 43 [Sep. 1935] (Ducke 1935).

VERNACULAR NAMES. — Pa: miumiu-asiru, miumiu-wašiunu • Wp: tulili • Br: taxi-branco-da-terra-firme.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5556*.

INVENTORY DATA (FG). — 50 trees in 35 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 104.5\text{ cm}$.

[961] *Tachigali melinonii* (Harms) Zarucchi & Herend.

Monogr. Syst. Bot. Missouri Bot. Gard. 45: 1254 [1 Oct. 1993] (Zarucchi & Herendeen 1993). — *Sclerolobium melinonii* Harms, *Bot. Jahrb. Syst. 33* (2, Beibl. 72): 24 [13 Mar. 1903] (Harms 1903).

NOTES. — Possible duplicates of the type collection at A and GH are referenced as “*Melinon 18*”. However, “18” is not a collection number, but the beginning of the collecting year, printed in advance on some 18th century labels at P. The last two digits should have been added by hand, but on some sheets they were not. The date 1863 appears on other possible duplicates at P.

VERNACULAR NAMES. — Pa: arey-avain, arey-avayni • Ka: alibagoli, apakanilan, eleylulan, tipulu alaulama • Te: tatsi inán • Wp: tulili wala, yapakaní'i • Nt: dyagidya • Cr: sèd-rémi, tachi • Br: taxi.

HERBARIUM DATA (FG). — 58 collections at CAY. Sel. exs.: *E.M. Mélénin s.n.* (holo-, B[not seen, photo F neg. N° 1805]; iso-, A[A00375130], F[F0057928F], G[G00367876], GH[GH00375131], K[K000264421, K000555829], S[S-R-9137], US[U0002719, 00411736]).

INVENTORY DATA (FG). — 277 trees in 103 plots; $F_{\max} = 9.7\%$; $\text{dbh}_{\text{inv}} = 86.3\text{ cm}$.

[962] *Tachigali micropetala* (Ducke) Zarucchi & Pipoly

Sida 16: 787 (Zarucchi & Pipoly 1995). — *Sclerolobium micropetalum* Ducke, *Bol. Técn. Inst. Agron. N.* 2: 20 (Ducke 1944).

VERNACULAR NAMES. — Ka: tasi epi, topuwunu.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1684*, dbh 30 cm.

[963] *Tachigali paniculata* Aubl.

(Fig. 32F)

Hist. Pl. Guiane 1: 372 [Jun.-Dec. 1775] (Aublet 1775). — *Cubaea paniculata* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 671 [late Sep.-Nov. 1791] (Gmelin 1791).

Tachigali trigona Aubl., *Hist. Pl. Guiane* 1: 374 [Jun.-Dec. 1775] (Aublet 1775). — *Cubaea trigona* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 671 [late Sep.-Nov. 1791] (Gmelin 1791).

Tachigali sericea Tul., *Arch. Mus. Hist. Nat.* 4: 163 (Tulasne 1844).

Tachigali eriocalyx Tul., *Arch. Mus. Hist. Nat.* 4: 164 (Tulasne 1844).

Tachigali angustifolia Miq., *Stirp. Surinam. Select.*: 13 ["1850" publ. Mar. 1851] (Miquel 1851). — *Tachigali paniculata* var. *angustifolia* (Miq.) Dwyer, *Ann. Missouri Bot. Gard.* 41: 239 (Dwyer 1954), "Tachigalia".

Tachigali rusbyi Harms, *Bot. Jahrb. Syst.* 33 (2, Beibl. 72): 20 [13 Mar. 1903] (Harms 1903).

Tachigali grandiflora Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 388 (Huber 1909), "Tachigalia".

Tachigali ulei Harms, *Notizbl. Königl. Bot. Gart. Berlin* 6: 306 [30 June 1915] (Harms 1915).

Tachigali paniculata var. *comosa* Dwyer, *Ann. Missouri Bot. Gard.* 41: 240 (Dwyer 1954).

Tachigali pulchra Dwyer, *Ann. Missouri Bot. Gard.* 41: 249 (Dwyer 1954).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: arey-avain, arey-avayni • Te: tatsi • Wp: tasi, tasi piyū, tasi sī • Wn: alapalī, piya upo • Nt: mene • Cr: bwa-fronmi, tachi, tasi • Fr: bois fourmi, tachi • Br: taxi, taxi-preto.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *J.B. Aublets.n.* (original material BM[BM000885813, G[G00367853]]).

INVENTORY DATA (FG). — 144 trees in 60 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 66.9$ cm.

[964] *Tachigali paraensis* (Huber) Barneby

Brittonia 48: 182 (Barneby 1996). — *Sclerolobium paraense* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 79 (Huber 1910), "paraënsē".

Sclerolobium albiflorum Benoit, *Bull. Soc. Bot. France* 66 (8): 383 ["1919" publ. 1920] (Benoit 1920). — *Tachigali albiflora* (Benoit) Zarucchi & Herend., *Fl. Venez. Guayana* 4: 115 (Zarucchi & Herendeen 1998).

VERNACULAR NAMES. — Pa: arey-avain, arey-avayni, miumiu-asiru • Ka: kumetilan, tamunen alaulama, tulili • Te: talidi, tatsi • Wp:

ka'a wewe • Nt: matawai gedu • Cr: tachi • Fr: bois fourmi, tachi • Br: passuaré, taxi-branco, taxi-da-terra-firme, taxirana.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *R. Benoit 1074* (original material of *Sclerolobium albiflorum*: P[P00578029, P00578030, P00578031]).

INVENTORY DATA (FG). — 139 trees in 68 plots; $F_{\max} = 2.4\%$; $dbh_{\text{inv}} = 105.8$ cm.

[965] *Tachigali richardiana* Tul.

Arch. Mus. Hist. Nat. 4: 166 (Tulasne 1844).

Tassia ovata Rich. ex Tul., *Arch. Mus. Hist. Nat.* 4: 167 (Tulasne 1844), *nom. nud. pro syn.*

Tachigali sulcata Benoit, *Bull. Mus. Natl. Hist. Nat.* 31: 469 (Benoit 1925).

Tachigali bracteolata Dwyer, *Ann. Missouri Bot. Gard.* 41: 233 (Dwyer 1954), "Tachigalia".

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: arey-avain, arey-avayni • Nt: gangi udu • Cr: tachi • Fr: bois fourmi, tachi • Br: taxi-pitomba-preta.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (type P[P00835780]).

INVENTORY DATA (FG). — 152 trees in 56 plots; $F_{\max} = 3.7\%$; $dbh_{\text{inv}} = 101.5$ cm.

[966] *Tachigali* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2842*.

INVENTORY DATA (FG). — 12 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 85$ cm.

Genus *Taralea* Aubl.

[967] *Taralea oppositifolia* Aubl.

Hist. Pl. Guiane 2: 745 [Jun.-Dec. 1775] (Aublet 1775), "Oppositifolia" on plate. — *Dipteryx oppositifolia* (Aubl.) Forsyth f., *Bot. Nomencl.*: 391 (Forsyth 1794). — *Coumarouna oppositifolia* (Aubl.) Taub., *Bot. Centralbl.* 47: 389 [31 Sept 1891] (Taubert 1891), "Cumaruna".

Swartzia coriacea Desv., *Ann. Sci. Nat. (Paris)* 9: 424 (Desvaux 1826). — *Coumarouna coriacea* (Desv.) Taub., *Bot. Centralbl.* 47: 389 [31 Sept 1891] (Taubert 1891).

Dipteryx nudipes Tul., *Arch. Mus. Hist. Nat.* 4: 100 (Tulasne 1844). — *Coumarouna nudipes* (Tul.) Kuntze, *Revis. Gen. Pl.* 1: 177 [5 Nov. 1891] (Kuntze 1891), "Cumaruna". — *Taralea nudipes* (Tul.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 142 (Ducke 1922).

Dipteryx applanata Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 234 (Bentham 1850).

Dipteryx oppositifolia var. *parviflora* Benth., *J. Proc. Linn. Soc., Bot.* 4 (Suppl.): 126 [Mar. 1860] (Bentham 1860). — *Taralea oppositifolia* var. *parviflora* (Benth.) Dugand, *Caldasia* 5 (21): 73 (Dugand 1948).

Dipteryx parviflora Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 126 [Mar. 1860] (Bentham 1860), *nom. nud. pro syn.*

Coumarouna speciosa Ducke, *Arch. Jard. Bot. Rio de Janeiro 3*: 162 (Ducke 1922). — *Dipteryx speciosa* (Ducke) Ducke, *Trop. Woods 61*: 10 (Ducke 1940).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: á-awainó, waikwimna • Te: aí pane • Wp: palaí wátá • Nt: neko udu • Cr: gayak-rivié • Br: acapurana, cumarurana.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000931956](#)], NY[[00033784](#)]); *D. Sabatier & M.-F. Prévost 3394*, dbh 60 cm.

[968] *Taralea* sp. A

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4962*.

INVENTORY DATA (FG). — 155 trees in 33 plots; $F_{\max} = 2.8\%$; $dbh_{inv} = 78.6$ cm.

Genus *Vatairea* Aubl.

[969] *Vatairea erythrocarpa* (Ducke) Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 139, 192 (Ducke 1930). — *Tipuana erythrocarpa* Ducke, *Arch. Jard. Bot. Rio de Janeiro 3*: 152 (Ducke 1922).

VERNACULAR NAMES. — Pa: wanetun • Wp: tayaúi • Nt: nkasa tyabisi • Br: faveira.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3766*.

INVENTORY DATA (FG). — 29 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{inv} = 97.1$ cm.

[970] *Vatairea guianensis* Aubl.

Hist. Pl. Guiane 2: 755 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

Andira amazonum Mart. ex Benth., *Commentat. Legum. Gen. 43 [June 1837]* (Bentham 1837). — *Vouacapoua amazonum* (Mart. ex Benth.) Kuntze, *Revis. Gen. Pl. 1*: 212 [5 Nov. 1891] (Kuntze 1891), “*Vuacapua*”.

Andira bracteosa Benth., *Commentat. Legum. Gen. 43 [June 1837]* (Bentham 1837).

Ormosia pacimonensis Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 4 (Suppl.)*: 119 [Mar. 1860] (Bentham 1860), *nom. nud. pro syn.*

Vatairea surinamensis Kleinhoonte, *Recueil Trav. Bot. Néerl. 22*: 403 [“1925” publ. Jan. 1926] (Kleinhoonte 1926).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wagu, waru • Wn: etekele • Cr: bwadat • Br: faveira-de-bolacha, faveira-de-empingem.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000931917](#)])

INVENTORY DATA (FG). — 1 tree, $dbh = 11.8$ cm.

[971] *Vatairea paraensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 140 (Ducke 1930).

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2763*.

INVENTORY DATA (FG). — 55 trees in 45 plots; $F_{\max} < 1\%$; $dbh_{inv} = 129$ cm.

Genus *Vataireopsis* Ducke

[972] *Vataireopsis surinamensis* H.C.Lima

Rodriguésia 32 (54): 30 (Lima 1980), “*surinamensis*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: wagu-kamwi, waru-kamwi • Ka: eleyulu, malipa wewe • Wp: wila paye • Nt: nkasa, yongo • Cr: dyango, yango • Br: fava-amargosa.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1672*.

INVENTORY DATA (FG). — 38 trees in 31 plots; $F_{\max} < 1\%$; $dbh_{inv} = 115$ cm.

Genus *Vouacapoua* Aubl.

[973] *Vouacapoua americana* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 9 [Jun.-Dec. 1775] (Aublet 1775). — *Andira aubletii* Benth., *Commentat. Legum. Gen.*: 44 [June 1837] (Bentham 1837), *nom. illeg. superfl.* (based on *Vouacapoua americana*). — *Vouacapoua aubletii* A.Lyons, *Pl. Nam.*: 396 (Lyons 1900), *nom. illeg. superfl.* (based on *Vouacapoua americana*).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wakap, wakap-priye, wakap-seine • Ka: wakapu • Te: akapu • Wp: wakapu • Wn: wakap • Nt: bunahati, buunaati • Cr: wakapou • Fr: wacapou • Br: acapú.

HERBARIUM DATA (FG). — 98 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000603704](#)]).

INVENTORY DATA (FG). — 1187 trees in 139 plots; $F_{\max} = 8\%$; $dbh_{inv} = 130$ cm.

Genus *Zollernia* Wied-Neuw. & Nees

[974] *Zollernia surinamensis* Mansano

A.M.G.Azevedo & G.P.Lewis, *Kew Bull. 59 (4)*: 516 [“2004” publ. Mar. 2005] (Mansano *et al.* 2005).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *Service Forestier 7603* (U).

SIZE. — Up to 15 m tall (Mansano *et al.* 2004).

Genus *Zygia* P.Browne

[975] *Zygia cataractae* (Kunth) L.Rico

Kew Bull. 46 (3): 496 [27 Sep. 1991] (Rico 1991). — *Inga cataractae* Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 297 [Apr. 1824] (Kunth 1824). — *Pithecellobium cataractae* (Kunth) Benth., *London J. Bot.* 3: 213 (Bentham 1844), “*Pithecolobium*”. — *Feuilleea cataractae* (Kunth) Kuntze, *Revis. Gen. Pl.* 1: 185 [5 Nov. 1891] (Kuntze 1891).

Inga glomerata DC., *Prod. [A. P. de Candolle]* 2: 438 [mid Nov. 1825] (Candolle 1825). — *Pithecellobium glomeratum* (DC.) Benth., *London J. Bot.* 3: 213 (Bentham 1844), “*Pithecolobium*”. — *Zygia glomerata* (DC.) Pittier, *Arb. Arbust. Venez.* 6-8: 80? [Aug.-Sep. 1927] (Pittier 1927).

Pithecellobium divaricatum Benth., *London J. Bot.* 3: 213 (Bentham 1844), “*Pithecolobium*”. — *Feuilleea divaricata* (Benth.) Kuntze, *Revis. Gen. Pl.* 1: 187 [5 Nov. 1891] (Kuntze 1891). — *Zygia divaricata* (Benth.) Pittier, *Arb. Arbust. Venez.* 6-8: 79 [Aug.-Sep. 1927] (Pittier 1927).

Inga ramiflora G.Don, *Gen. Hist.* 2: 392 [Oct. 1832] (Don 1832). — *Mimosa ramiflora* Ruiz & Pav. ex G.Don, *Gen. Hist.* 2: 392 [Oct. 1832] (Don 1832), *nom. nud. pro syn.*

Inga divaricata Bong. ex Benth., *Fl. Bras. [Martius]* 15 (2): 448 [1 July 1876] (Bentham 1876), *nom. nud. pro syn.*

Calliandra schwackeana Taub., *Flora* 75 [n.s. 50]: 69 (Taubert 1892).

Pithecellobium cauliflorum Mart. f. *niveum* Lindm., *Bih. Kongl. Svenska Vetensk.-Akad. Handl.* 24 (3/7): 56 (Lindman 1898), “*Pithecolobium*”.

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: asiru • Te: pulelu inga • Wp: asila, kulu- lulu inga, kululu inga sili • Wn: kalaujalan • Nt: bwasiman weko, kokobe weko • Cr: bwa-sira • Br: ararandeua.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost 4675*, 8 m × 10 cm.

[976] *Zygia inaequalis* (Humb. & Bonpl. ex Willd.) Pittier

Arb. Arbust. Venez. 6-8: 79 [Aug.-Sep. 1927] (Pittier 1927). — *Inga inaequalis* Humb. & Bonpl. ex Willd., *Sp. Pl.*, ed. 4 4 (2): 1019 [Apr. 1806] (Willdenow 1806). — *Mimosa inaequalis* (Humb. & Bonpl. ex Willd.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 1: 44 [3 Sep. 1810] (Poirier 1810). — *Pithecellobium inaequale* (Humb. & Bonpl. ex Willd.) Benth., *Trans. Linn. Soc. London* 30 (3): 596 [10 Apr. 1875] (Bentham 1875), “*Pithecolobium*”. — *Feuilleea inaequalis* (Humb. & Bonpl. ex Willd.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Pithecellobium bicolor Spruce ex Benth., *Fl. Bras. [Martius]* 15 (2): 451 [1 July 1876] (Bentham 1876), *nom. nud. pro syn.*

Pithecellobium pilosulum Pittier, *Contr. U.S. Natl. Herb.* 20 (12): 466 [9 Jan. 1922] (Pittier 1922), “*Pithecolobium*”.

Pithecellobium longiramosum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 67 (Ducke 1922), “*Pithecolobium*”. — *Zygia longiramosa* (Ducke) L.Rico, *Kew Bull.* 46 (3): 501 [27 Sep. 1991] (Rico 1991).

Pithecellobium foreroi C.Barbosa, *Caldasia* 14 (68-70): 395 (Barbosa 1986). — *Zygia foreroi* (C.Barbosa) L.Rico, *Kew Bull.* 46 (3): 500 [27 Sep. 1991] (Rico 1991).

VERNACULAR NAMES. — Nt: gaan bwasiman weko • Br: ingarana, jarandeuia.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier 1031*.

SIZE. — Up to 20 cm dbh (Barneby & Grimes 1997).

[977] *Zygia latifolia* (L.) Fawc. & Rendle

Fl. Jamaica [Fawcett & Rendle] 4: 150 (Fawcett & Rendle 1920). — *Mimosa latifolia* L., *Syst. Nat.*, ed. 10, 2: 1312 [7 June 1759] (Linnaeus 1759). — *Inga latifolia* (L.) Willd., *Sp. Pl.*, ed. 4 4 (2): 1020 [Apr. 1806] (Willdenow 1806). — *Pithecellobium latifolium* (L.) Benth., *London J. Bot.* 3: 214 (Bentham 1844), “*Pithecolobium*”. — *Calliandra latifolia* (L.) Griseb., *Fl. Brit. WI. [Grisebach]*: 225 [late 1860] (Grisebach 1860). — *Feuilleea latifolia* (L.) Kuntze, *Revis. Gen. Pl.* 1: 188 [5 Nov. 1891] (Kuntze 1891).

Mimosa zygia L., *Fl. Jamaic. [Linnaeus]* 22 [22 Dec. 1759] (Linnaeus 1759).

Zygia arborescens J.St.-Hil., *Expos. Fam. Nat.* 2: 245 (Saint-Hilaire 1805).

Pithecellobium chagrense Pittier, *Contr. U.S. Natl. Herb.* 20 (12): 465 [9 Jan. 1922] (Pittier 1922).

Pithecellobium huberi Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 29 (Ducke 1925), “*Pithecolobium*”. — *Zygia huberi* (Ducke) L.Rico, *Kew Bull.* 46 (3): 501 (Rico 1991).

Pithecellobium latifolium subsp. *tayronense* C.Barbosa, *Caldasia* 14 (68-70): 397 (Barbosa 1986).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana. *Zygia arborescens* J.St.-Hil. was validly published by reference to the (then monotypic) genus *Zygia* P.Browne (Turland *et al.* 2018: Art. 38.12). However, “*Arborescens*” in Browne’s text was descriptive, not epithetical. Therefore, St.-Hilaire is the sole author of the binomial *Z. arborescens* (not “*P.Browne ex J.St.-Hil.*”).

VERNACULAR NAMES. — Pa: asiru • Ka: ayalani • Wp: asila, kulu- lulu inga • Nt: bwasiman weko, kokobe weko • Br: ingá-do-baixo, ingá-dura.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *P. Grenand 1126*.

SIZE. — Up to 30 cm dbh (Barneby & Grimes 1997).

[978] *Zygia racemosa* (Ducke) Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (2): 71 (Barneby & Grimes 1997). — *Pithecellobium racemosum* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: errata following p. 272 (Ducke 1922), “*Pithecellobium*”. — *Abarema racemosa* (Ducke) Kleinhoonte, *Fl. Suriname* 2 (2): 320 (Kleinhoonte 1940). — *Marmoroxylon racemosum* (Ducke) Killip, *Trop. Woods* 63: 3 (Killip 1940).

Pithecellobium racemiflorum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 14 (Ducke 1915), “*Pithecellobium*”, nom. illeg. hom., non Donn.Sm. (Donnell Smith 1913).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kaibune-á, kaibune-ára • Ka: kwata weli • Wn: ekesaimamé • Nt: sineki udu • Cr: bwa-serpan • Fr: bois serpent • Br: angelim-rajado, ingarana-da-terra-firme, urubuzeiro.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2912.

INVENTORY DATA (FG). — 143 trees in 60 plots; $F_{\max} = 3\%$; $\text{dbh}_{\text{inv}} = 63 \text{ cm}$.

[979] *Zygia sabatieri* Barneby & J.W.Grimes
(Fig. 32E)

Mem. New York Bot. Gard. 74 (2): 83 (Barneby & Grimes 1997).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Wp: tayaú'i • Br: faveira.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3896 (holo-, CAY[CAY097639]; iso-, MO[MO-277933], NY[00003389, 00003390], P[P00208187, P02142986], U[U0003453]).

INVENTORY DATA (FG). — 34 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 42.8 \text{ cm}$.

[980] *Zygia tetragona* Barneby & J.W.Grimes

Mem. New York Bot. Gard. 74 (2): 80 (Barneby & Grimes 1997).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: sineki udu.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2561 (holo-, P[P00077200]; iso-, NY[00003392, 00003393], P[P00208190]).

INVENTORY DATA (FG). — 66 trees in 30 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 105 \text{ cm}$.

Family LEPIDOBOTRYACEAE J.Léonard
Genus *Ruptiliocarpon* Hammel & N.Zamora

[981] *Ruptiliocarpon* sp. A
(Fig. 33A)

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2846.

INVENTORY DATA (FG). — 19 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 84 \text{ cm}$.

Family LINACEAE DC. ex Perleb
Genus *Hebepepetalum* Benth.

[982] *Hebepepetalum humiriifolium*
(Planch.) Benth. & Hook.f. ex B.D.Jacks.

Index Kew. 1 (2): 1097 [14 Dec. 1893] (Jackson 1893). — *Roucheeria humiriifolia* Planch., *London J. Bot.* 6: 143 (Planchon 1847).

Roucheria angulata Gleason, *Bull. Torrey Bot. Club* 58 (6): 373 [June 1931] (Gleason 1931).

NOTE. — The combination by Bentham & Hooker f. (*Gen. Pl.* 1: 245. 1862) is invalid, because they did not associate the genus name to the epithet.

VERNACULAR NAMES. — Pa: á-danó • Ka: kunapolan, pakila sipyoli, silimaipo • Wp: pasí'i wapo, pasí'i wapo sili • Cr: bwa-patagay • Br: abacate-rana, itaúba-abacate.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *J. Martin s.n.* (type K[K000407359]).

INVENTORY DATA (FG). — 284 trees in 110 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 56.5 \text{ cm}$.

Genus *Roucheria* Planch.

[983] *Roucheria laxiflora* H.J.P.Winkl.
(Fig. 33B)

Repert. Spec. Nov. Regni Veg. 7: 109 (Winkler 1909).

VERNACULAR NAMES. — Wp: tukánakú sili.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3165.

INVENTORY DATA (FG). — 33 trees in 4 plots; $F_{\max} = 5.7\%$; $\text{dbh}_{\text{inv}} = 23.6 \text{ cm}$.

Family LOGANIACEAE R.Br. ex Mart.
Genus *Antonia* R.Br.

[984] *Antonia ovata* Pohl
(Fig. 33C)

Pl. Bras. Icon. Descr. 2 (1): 14 [1828 publ. Jan. 1829] (Pohl 1829).

Antonia pilosa Hook., *Icon. Pl.* 1: t. 64 [1 Feb. 1837] (Hooker 1837). — *Antonia ovata* var. *pilosa* (Hook.) Progel, *Fl. Bras. [Marius]* 6 (1): 253 [1 Aug. 1868] (Progel 1868).

Antonia pubescens Bong., *Mém. Acad. Imp. Sci. Saint-Pétersbourg*, Sér. 6, *Sci. Math.*, Seconde Pt. *Sci. Nat.* 5 (2, Bot.): 2 [Oct. 1839] (Bongard 1839).

Antonia ovata var. *excelsa* Paula, *Acta Amazonica* 6 (1): 41 (Paula 1976).

VERNACULAR NAMES. — Ka: palulipo, tamanokwale • Wp: kulupí'i, wila pipé mu.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *D. Sabatier* 4863.

INVENTORY DATA (FG). — 61 trees in 36 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 75.6 \text{ cm}$.

Genus *Strychnos* L.

[985] *Strychnos cayennensis* Krukoff & Barneby
(Fig. 33D)

Phytologia 27 (2): 101 [29 Nov. 1973] (Krukoff & Barneby 1973).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.-F. Molino et al. 3329*.

INVENTORY DATA (FG). — 1 tree, dbh = 13 cm.

Family LYTHRACEAE J.St.-Hil.
Genus *Lafoensia* Vand.

[986] *Lafoensia vandelliana* DC. ex Cham. & Schltld.
(Fig. 34A)

Linnaea 2: 346 (Chamisso & Schlechtendal 1827).

Lafoensia replicata Pohl, *Pl. Bras. Icon. Descr.* 2 (4): 144 [Jan.-Sep. 1833] (Pohl 1833), “*Lafoënsia*”. — *Lafoensia vandelliana* subsp. *replicata* (Pohl) Loureig, *Mem. Soc. Ci. Nat. La Salle* 45 (123): 134 (Loureig 1985).

NOTE. — *Lafoensia vandelliana* was first invalidly published by Candolle (1826: 73), because he cited four species in his description of the genus, without distinguishing them (Turland *et al.* 2018: Art. 38.12).

VERNACULAR NAMES. — Br: dedalinho, louro-da-serra, mirindiba-rosa.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 15000*.

INVENTORY DATA (FG). — 5 trees in 1 plot; dbh_{inv} = 18 cm.

Family MALPIGHIAEAE Juss.
Genus *Bunchosia* Rich. ex Kunth.

[987] *Bunchosia argentea* (Jacq.) DC.
(Fig. 34B)

Prodr. [A. P. de Candolle] 1: 582 [mid Jan. 1824] (Candolle 1824). — *Malpighia argentea* Jacq., *Fragm. Bot.* 57 (Jacquin 1806).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5738*.

INVENTORY DATA (FG). — 5 trees in 5 plots; F_{max} < 1%; dbh_{inv} = 29.4 cm.

[988] *Bunchosia decussiflora* W.R.Anderson

Mem. New York Bot. Gard. 32: 279 (Anderson 1981).

VERNACULAR NAMES. — Wp: wila tawa, yanita.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 934*, 12 m × 15 cm.

Genus *Byrsinima* Rich. ex Kunth

[989] *Byrsinima aerugo* Sagot

Ann. Sci. Nat., Bot. sér. 6, 12: 178 (Sagot 1881).

NOTES. — Anderson (2007: 85) made a first-step lectotypification by choosing among the paratypes “*Sagot 102* (lectotype: Pl; isotypes: BM! G! K! MEL! MICH! NY!”. However, there are four collection dates among the eight specimens labelled “*Sagot 102*” at P (1855, 1856, 1857, 1858). It is therefore necessary to designate one of them as lectotype. We selected the specimen *P05529869* dated 1857 on which Sagot affixed a complete handwritten description of his new species.

VERNACULAR NAMES. — Ka: pelulu • Wn: mulei.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *P.A. Sagot 102*, 1857 (lecto-, P[P05529869] here designated; isolecto-, BM[BM000796094], BR[BR0000008497468], GOET[GOET007083, GOET007084], P[P05529871], S[S08-15040]).

INVENTORY DATA (FG). — 14 trees in 12 plots; F_{max} = 2.8 %; dbh_{inv} = 43.4 cm.

[990] *Byrsinima altissima* (Aubl.) DC.

Prodr. [A. P. de Candolle] 1: 579 [mid Jan. 1824] (Candolle 1824). — *Malpighia altissima* Aubl., *Hist. Pl. Guiane* 1: 455 [Jun.-Dec. 1775] (Aublet 1775).

Byrsinima aubletii Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 10 (Kostermans 1936).

Byrsinima discolor Pilg., *Repert. Spec. Nov. Regni Veg.* 42: 179 (Pilger 1937).

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN[LINN-HS 822.4], P-LA[P00287893], US[00036896]).

INVENTORY DATA (FG). — 8 trees in 7 plots; F_{max} < 1%; dbh_{inv} = 88.2 cm.

[991] *Byrsinima christianeae* W.R.Anderson

Mem. New York Bot. Gard. 32: 112 (Anderson 1981).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: mavinbi-kamwi, mavinvi-kamwi.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2664*.

INVENTORY DATA (FG). — 1 tree, dbh = 16.9 cm.

[992] *Byrsinima crassifolia* (L.) Kunth

Nova genera et species plantarum [H.B.K.] 5: 149 [25 Feb. 1822] (Kunth 1822). — *Malpighia crassifolia* L., *Sp. Pl.* 1: 426 [1 May 1753] (Linnaeus 1753).

Malpighia moureila Aubl., *Hist. Pl. Guiane* 1: 459 [Jun.-Dec. 1775] (Aublet 1775).

Malpighia coriacea Sw., *Prodr. [Swartz]* 74 [20 Jun.-29 July 1788] (Swartz 1788). — *Byrsonima coriacea* (Sw.) DC., *Prodr. [A. P. de Candolle]* 1: 580 [mid Jan. 1824] (Candolle 1824).

Byrsonima laurifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 150 [25 Feb. 1822] (Kunth 1822).

Byrsonima ferruginea Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 151 [25 Feb. 1822] (Kunth 1822).

Byrsonima cotinifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 152 [25 Feb. 1822] (Kunth 1822).

Byrsonima lanceolata DC., *Prodr. [A. P. de Candolle]* 1: 580 [mid Jan. 1824] (Candolle 1824).

Byrsonima pulchra DC., *Prodr. [A. P. de Candolle]* 1: 580 [mid Jan. 1824] (Candolle 1824).

Byrsonima rufescens Bertol., *Nuovi Ann. Sci. Nat.* 414 (Bertoloni 1838).

Byrsonima cumingiana A.Juss., *Ann. Sci. Nat., Bot. sér. 2, 13:* 332 [May 1840] (Jussieu 1840), “*Cumingana*”.

Byrsonima karwinskiana A.Juss., *Ann. Sci. Nat., Bot. sér. 2, 13:* 333 [May 1840] (Jussieu 1840).

Byrsonima panamensis Beurl., *Kongl. Vetensk.-Akad. Handl.* 40: 117 [“1854” publ. 1856] (Beurling 1856).

Byrsonima moritziana Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 31 (1): 391 (Turczaninow 1858).

Byrsonima fendleri Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (2): 582 (Turczaninow 1863).

Malpighia pulchra Sessé & Moc., *Fl. Mexic.*, ed. 2, 115 (Sessé & Mociño 1894).

Byrsonima fagifolia Nied., *Byrsonima* 2: 20 (Niedenzu 1901).

Byrsonima laurifolia var. *guatemalensis* Nied., *Pflanzenr. [Engler] IV.141.III (Heft 94):* 724 [27 Nov. 1928] (Niedenzu 1928).

NOTES. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017). In French Guiana, this species is restricted to savannas and other open vegetations.

VERNACULAR NAMES. — Pa: mariuvra, mariuvra-seine • Ka: mile’i, morelia (*fide* Aublet 1775), mule’i • Wp: alalí lá, palálá ka’á • Cr: mourousi, prin-savann • Fr: maurissi • Br: murici, muruci.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *M.-F. Prévost 3845*.

SIZE. — Up to 15 cm dbh (Cuatrecasas & Croat 1980).

[993] *Byrsonima densa* (Poir.) DC.

Prodr. [A. P. de Candolle] 1: 580 [mid Jan. 1824] (Candolle 1824). — *Malpighia densa* Poir., *Encycl. [J. Lamarck et al.] Suppl. 4:* 7 [29 June 1816] (Poiret 1816).

Byrsonima amazonica Griseb., *Fl. Bras. [Martius]* 12 (1): 14 [1 June 1858] (Grisebach 1858).

Byrsonima lucidula Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 423 (Huber 1902). — *Byrsonima amazonica* var. *lucidula* (Huber) Nied., *Pflanzenr. [Engler] IV.141.III (Heft 94):* 745 [27 Nov. 1928] (Niedenzu 1928).

Byrsonima densa var. *emarginata* Kosterm., *Meded. Bot. Mus. Herb. Rijks Univ. Utrecht* 25: 10 (Kostermans 1936).

NOTE. — *F.M.R. Leprieur s.n.*, 1840 (F, V0062737F) is erroneously marked as isotype at F, given it was collected in 1840, long after the publication of *Malpighia densa* Poir. Leprieur was in French Guiana from 1829 to 1870.

VERNACULAR NAMES. — Ka: mile’i, mule’i • Wp: mulei pilá, mulei sówí • Br: muruci-vermelho.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (“Herb. Poir.”) (type P[P02428546]).

INVENTORY DATA (FG). — 18 trees in 14 plots; $F_{\max} = 3.3\%$; $dbh_{\text{inv}} = 49.3$ cm.

[994] *Byrsonima krukoffii* W.R.Anderson
(Fig. 34D)

Contr. Univ. Michigan Herb. 20: 24 (Anderson 1995).

VERNACULAR NAMES. — Pa: tukuyuy-kamwi • Wp: mulei.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 3526*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 18.5$ cm.

[995] *Byrsonima laevigata* (Poir.) DC.
(Fig. 34E)

Prodr. [A. P. de Candolle] 1: 580 [mid Jan. 1824] (Candolle 1824). — *Malpighia laevigata* Poir., *Encycl. [J. Lamarck et al.] 4 (1):* 332 [9 Feb. 1797] (Poiret 1797).

Byrsonima obversa Miq., *Stirp. Surinam. Select.*: 76 [“1850” publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Pa: murusi • Ka: mule’i ityudano.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (holo-, P-LA[P00287881]; iso-, P[P02428604]).

INVENTORY DATA (FG). — 70 trees in 50 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 61.9$ cm.

[996] *Byrsonima sericea* DC.

Prodr. [A. P. de Candolle] 1: 580 [mid Jan. 1824] (Candolle 1824).

Byrsonima eglandulosa A.Juss., *Arch. Mus. Hist. Nat.* 3: 273 (Jussieu 1843). — *Byrsonima sericea* var. *eglandulosa* (A.Juss.) Griseb., *Fl. Bras. [Martius]* 12 (1): 11 [1 June 1858] (Grisebach 1858).

Byrsonima sericea var. *angustifolia* Griseb., *Fl. Bras. [Martius]* 12 (1): 12 [1 June 1858] (Grisebach 1858).

Byrsonima sericea f. *eglandulosa* Nied., *Pflanzenr. [Engler] IV.141.III (Heft 94):* 703 [27 Nov. 1928] (Niedenzu 1928), nom. illeg. hom., non var. *eglandulosa* (A.Juss.) Griseb. (Grisebach 1858).

Byrsonima sericea var. *pubescens* Nied., *Pflanzenr. [Engler] IV.141.III (Heft 94):* 703 [27 Nov. 1928] (Niedenzu 1928).




FIG. 33.— Lepidobotryaceae: **A**, *Ruptiliocarpon* sp. A (D. Sabatier et al. 6045). Linaceae: **B**, *Roucheria laxiflora* H.J.P.Winkl. (D. Sabatier et al. 5877). Loganiaceae: **C**, *Antonia ovata* Pohl (D. Sabatier 4863); **D**, *Strychnos cayennensis* Krukoff & Barneby (J.-F. Molino et al. 3329). © D. Sabatier/IRD.

NOTE. — Niedenzu (1928: 702) listed var. *eglandulosa* but created a distinct, heterotypic f. *eglandulosa* within his var. *pubescens* (Niedenzu 1928: 703).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost 204*.

SIZE. — Up to 20 m tall (Teixeira & Machado 2000).

[997] *Byrsonima spicata* (Cav.) DC.

Prodri. [A. P. de Candolle] 1: 580 [mid Jan. 1824] (Candolle 1824). — *Malpighia spicata* Cav., *Diss.* 8: 409 [4 May 1789] (Cavallilles 1789). — *Byrsonima coriacea* var. *spicata* (Cav.) Nied., *Pflanzenr. [Engler]* IV.141.III (Heft 94): 700 [27 Nov. 1928] (Niedenzu 1928).

Byrsonima chrysophylla Kunth, *Nova genera et species plantarum [H.B.K.J]* 5: 150 [25 Feb. 1822] (Kunth 1822). — *Galphimia chrysophylla* (Kunth) Spreng., *Syst. Veg. [Sprengel]* 2: 385 [Jan.-May 1825] (Sprengel 1825).

Malpighia guadalupensis Spreng., *Syst. Veg. [Sprengel]* 2: 383 [Jan.-May 1825] (Sprengel 1825).

Byrsonima guadalupensis G.Don, *Gen. Hist.* 1: 637 [early Aug. 1831] (Don 1831).

Byrsonima berteroana A.Juss., *Ann. Sci. Nat., Bot. sér. 2, 13*: 333 [May 1840] (Jussieu 1840).

Byrsonima peruviana A.Juss., *Ann. Sci. Nat., Bot. sér. 2, 13*: 333 [May 1840] (Jussieu 1840).

Byrsonima peruviana var. *eglandulosa* A.Juss., *Arch. Mus. Hist. Nat.* 3: 290 (Jussieu 1843).

Byrsonima hostmannii Benth., *London J. Bot.* 7: 120 (Bentham 1848).

Byrsonima propinqua Benth., *London J. Bot.* 7: 120 (Bentham 1848). — *Byrsonima spicata* f. *propinqua* (Benth.) Nied., *Byrsonima* 2: 5 (Niedenzu 1901). — *Byrsonima coriacea* f. *propinqua* (Benth.) Nied., *Pflanzenr. [Engler]* IV.141.III (Heft 94): 701 [27 Nov. 1928] (Niedenzu 1928).

Byrsonima chrysophylla f. *glandulifera* Griseb., *Fl. Bras. [Martius]* 12 (1): 12 [1 June 1858] (Grisebach 1858).

Byrsonima biacuminata Rusby, *Mem. Torrey Bot. Club* 6 (1): 13 (Rusby 1896).

Byrsonima chrysophylla f. *kunthiana* Nied., *Byrsonima* 2: 7 (Niedenzu 1901).

Byrsonima ophiticola Small ex Britton, *Bull. Torrey Bot. Club* 43 (9): 456 (Britton 1916).

Byrsonima horneana Britton & Small, *Sci. Surv. Porto Rico & Virgin Islands* 5: 447 (Britton & Small 1924).

Byrsonima coriacea f. *angustifolia* Nied., *Pflanzenr. [Engler]* IV.141.III (Heft 94): 701 [27 Nov. 1928] (Niedenzu 1928). — *Byrsonima angustifolia* Benth. ex Nied., *Pflanzenr. [Engler]* IV.141.III (Heft 94): 701 [27 Nov. 1928] (Niedenzu 1928), nom. nud. pro syn.




FIG. 34. — Lythraceae: **A**, *Lafoensia vandelliana* DC. ex Cham. & Schlechl. (*D. Sabatier & J.-F. Molino 5104*). Malpighiaceae: **B**, *Bunchosia argentea* (Jacq.) DC. (*D. Sabatier & J.-F. Molino 5738*); **C**, *Byrsonima stipulacea* A.Juss. (*D. Sabatier 4862*); **D**, *Byrsonima krukoffii* W.R.Anderson (*D. Sabatier 3526*); **E**, *Byrsonima laevigata* (Poir.) DC. (*D. Sabatier 5575*). © D. Sabatier/IRD.

Byrsonima coriacea f. *eglandulosa* Nied., *Pflanzenr. [Engler] IV.141. III (Heft 94)*: 701 [27 Nov. 1928] (Niedenzu 1928).

NOTES.—Kunth (Humboldt *et al.* 1822: 147) transferred *Malpighia spicata* Cav. to *Byrsonima* without associating genus name and epithet, making the combination invalid (Turland *et al.* 2018: Art. 35.2). *B. guadalupensis* G.Don is not based on *M. guadalupensis* Spreng.

VERNACULAR NAMES.—Br: calliandra.

HERBARIUM DATA (FG).—20 collections at CAY. Sel. exs.: *M.-F. Prévost 1205*.

SIZE.—Up to 25 m tall (Anderson 1999).

[998] *Byrsonima stipulacea* A.Juss.
(Fig. 34C)

Ann. Sci. Nat., Bot. sér. 2, 13: 332 [May 1840] (Jussieu 1840).—*Alcoceratothrix stipulacea* (A.Juss.) Nied., *Pflanzenr. [Engler] IV.141. III (Heft 94)*: 762 [27 Nov. 1928] (Niedenzu 1928).

Byrsonima longibracteata A.Juss. ex Mart., *Flora 24* (2, Beibl.): 61 (Martius 1841).—*Alcoceratothrix longibracteata* (A.Juss. ex Mart.) Nied., *Byrsonima* 2: 45 (Niedenzu 1901).

Byrsonima rugosa Benth., *London J. Bot. 7*: 118 (Bentham 1848).—*Alcoceratothrix rugosa* (Benth.) Nied., *Byrsonima* 2: 45 (Niedenzu 1901).

Byrsomima coleostachya Griseb., *Linnaea* 22: 3 (Grisebach 1849).

Alcoceratothrix rugosa var. *minor* Nied., *Byrsomima* 2: 45 (Niedenzu 1901).

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier 4862*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 69.5$ cm.

Genus *Spachea* A.Juss.

[999] *Spachea elegans* (G.Mey.) A.Juss.

Icon. Sel. Pl. 3: 19 [“1837” publ. 1838] (Jussieu 1838). — *Malpighia elegans* G.Mey., *Prim. Fl. Esseq.* 178 [Nov. 1818] (Meyer 1818). — *Byrsomima elegans* (G.Mey.) DC., *Prodr. [A. P. de Candolle]* 1: 580 [mid Jan. 1824] (Candolle 1824).

Byrsomima herbert-smithii Rusby, *Descr. S. Amer. Pl.* 37 [20 Dec. 1920] (Rusby 1920). — *Spachea herbert-smithii* (Rusby) Cuatrec., *Webbia* 13 (2): 546 (Cuatrecasas 1958).

VERNACULAR NAMES. — Ka: noya elepali, pakila sipyoli • Wp: yási leáyy.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *Service Forestier M-23*.

SIZE. — Up to 12 cm dbh (Anderson 1981).

Family MALVACEAE Juss.

Genus *Apeiba* Aubl.

[1000] *Apeiba glabra* Aubl.

Hist. Pl. Guiane 1: 541 [Jun.-Dec. 1775] (Aublet 1775). — *Apeiba laevis* Sw., *Prod. [Swartz]*: 83 [20 Jun.-29 July 1788] (Swartz 1788), *nom. illeg. superfl.* (based on *Apeiba glabra*). — *Aubletia laevis* Willd., *Sp. Pl.*, ed. 4 2 (2): 1156 [Dec. 1799] (Willdenow 1799), *nom. illeg. superfl.* (based on *Apeiba laevis* and *A. glabra*).

Apeiba aspera Aubl., *Hist. Pl. Guiane* 1: 545 [Jun.-Dec. 1775] (Aublet 1775).

Apeiba burchellii Sprague, *Bull. Herb. Boissier*, sér. 2, 5: 703 (Sprague 1905).

VERNACULAR NAMES. — Pa: yit-itaibi-priye, yit-itaibi-puvemna, yit-itaibi-seine • Te: petumo ka’apewat • Wp: ape’i, ape’i sī • Nt: kankan udu • Cr: bwa-amadou, bwa-bouchon, bwa-kalou, pengn-makak • Fr: bois de mèche (*fide* Aublet 1775), peigne macaque • Br: pau-de-jangada, pente-de-macaco.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000795157](#)]).

INVENTORY DATA (FG). — 173 trees in 78 plots; $F_{\max} = 2.2\%$; $dbh_{\text{inv}} = 66.8$ cm.

[1001] *Apeiba petoumo* Aubl.

Hist. Pl. Guiane 1: 543 [Jun.-Dec. 1775] (Aublet 1775). — *Apeiba hispida* Gaertn., *Fruct. Sem. Pl.* 2: 188 [Apr.-May 1791] (Gaertner

1791), *nom. illeg. superfl.* (based on *Apeiba petoumo*). — *Aubletia petoumo* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (2): 1156 [Dec. 1799] (Willdenow 1799).

Apeiba echinata Gaertn., *Fruct. Sem. Pl.* 2: 189 [Apr.-May 1791] (Gaertner 1791).

Apeiba hypoleuca Steud., *Flora* 26 (45): 755 [17 Dec. 1843] (Steedel 1843).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), although known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: yit-itaibi-priyo • Ka: me:ku yongay, patumu • Te: petumo tsing • Nt: kankan udu • Cr: bwa-bouchon, bwa-kalou, pengn-makak • Fr: peigne macaque • Br: pente-de-macaco.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000795155](#)]).

INVENTORY DATA (FG). — 67 trees in 39 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 92.1$ cm.

[1002] *Apeiba tibourbou* Aubl.
(Fig. 35A)

Hist. Pl. Guiane 1: 538 [Jun.-Dec. 1775] (Aublet 1775). — *Aubletia tibourbou* (Aubl.) Willd., *Sp. Pl.*, ed. 4 2 (2): 1156 [Dec. 1799] (Willdenow 1799).

Apeiba tibourbou var. *rugosa* Szyszyl., *Pug. Pl. Amer. Centr.* 2 (Szyszylowicz 1894), “Tibourbon”.

Apeiba albiflora Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 209 (Ducke 1922).

Apeiba tibourbou var. *krukoffii* Uittien, *Recueil Trav. Bot. Néerl.* 32: 248 (Uittien 1935).

VERNACULAR NAMES. — Pa: imavui, yit-itaibi • Te: petumo • Wp: ape’i, ape’i sī • Wn: pēinékë emnamotem, petun • Nt: kankan udu, mifungafunga • Cr: bwa-amadou, bwa-bouchon, bwa-kalou, pengn-makak • Fr: peigne macaque • Br: pau-de-jangada, pente-de-macaco.

HERBARIUM DATA (FG). — 62 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000795154](#)], S[[S13-21705](#)]).

INVENTORY DATA (FG). — 10 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42$ cm.

[1003] *Apeiba* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2679*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.7 cm.

Genus *Ayenia* L.

[1004] *Ayenia morii*
(L.C.Barnett & Dorr) Christenh. & Byng

Global Fl. 4: 137 [9 Feb. 2018] (Christenhusz & Byng 2018). — *Byttneria morii* L.C.Barnett & Dorr, *Brittonia* 42 (4): 271 (Barnett & Dorr 1990).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *S.A. Mori et al. 19160* (holo-, NY[00222207]; iso-, CAY[CAY024967], CTES[CTES0001740], F[V0073514F], K[K000381151], MO[MO-309195], P[P01900193], U[U0006820], US[00324040], VEN[VEN262140]).

INVENTORY DATA (FG). — 1 tree, dbh = 13.5 cm.

Genus *Catostemma* Benth.

[1005] *Catostemma commune* Sandwith

Bull. Misc. Inform. Kew 1931 (1): 51 [20 Jan. 1931] (Sandwith 1931).

VERNACULAR NAMES. — Pa: waikwimna.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier et al. 5250*.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 57.5$ cm.

[1006] *Catostemma fragrans* Benth.

London J. Bot. 2: 365 (Bentham 1843).

Guenetia macrosperma Sagot ex Benoist, *Bull. Mus. Natl. Hist. Nat.* 25: 387 (Benoist 1919). — *Catostemma macrospermum* (Sagot ex Benoist) Lemée, *Fl. Guyane Franç.* 3: 631 (Lemée 1954).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), although known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kulebogo • Nt: aganananga.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *S.A. Mori et al. 23417*.

INVENTORY DATA (FG). — 552 trees in 120 plots; $F_{\max} = 9.6\%$; $\text{dbh}_{\text{inv}} = 54.1$ cm.

[1007] *Catostemma* sp. A

NOTE. — This is possibly *C. grazielae* Paula (*Ci. & Cult.* 21: 703 [Paula 1970]).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5530*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 29.7$ cm.

Genus *Ceiba* Mill.

[1008] *Ceiba pentandra* (L.) Gaertn.

Fruct. Sem. Pl. 2: 244 [Apr.-May 1791] (Gaertner 1791). — *Bombax pentandrum* L., *Sp. Pl.* 1: 511 [1 May 1753] (Linnaeus 1753). — *Eriodendron pentandrum* (L.) Kurz, *Jour. As. Soc. Bengal* 43: 113 (Kurz 1874). — *Xylon pentandrum* (L.) Kuntze, *Revis. Gen.*

Pl. 1: 75 [5 Nov. 1891] (Kuntze 1891). — *Eriodendron anfractuosum* DC., *Prodr. [A. P. de Candolle]* 1: 479 [mid Jan. 1824] (Candolle 1824), nom. illeg. superfl. (based on *Bombax pentandrum*). — *Bombax occidentale* Spreng., *Syst. Veg. [Sprengel]* 3: 124 [Jan.-Mar. 1826] (Sprengel 1826), nom. illeg. superfl. (based on *Bombax pentandrum*, as “*B. pentandrum* Jacq.”). — *Ceiba occidentalis* Burkill, *Bull. Misc. Inform. Kew* 1935 (5): 317 [4 Sep. 1935] (Burkill 1935), nom. illeg. superfl. (based on the illegitimate *Bombax occidentale*, thus indirectly on *B. pentandrum*). — *Bombax orientale* Spreng., *Syst. Veg. [Sprengel]* 3: 124 [Jan.-Mar. 1826] (Sprengel 1826). — *Eriodendron orientale* (Spreng.) Kostel., *Allg. Med.-Pharm. Fl.* 5: 1875 [Jan.-Sep. 1836] (Kosteletzky 1836). — *Gossampinus alba* Buch.-Ham., *Trans. Linn. Soc. London* 15 (1): 126 [9 Feb. 1827] (Buchanan-Hamilton 1827), nom. illeg. superfl. (based on *Ceiba pentandra*).

Ceiba casearia Medik., *Malvenfam.*: 16 (Medikus 1787).

Bombax cumanense Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 300 [June 1822] (Kunth 1822).

Bombax mompoxense Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 300 [June 1822] (Kunth 1822).

Eriodendron anfractuosum var. *caribaeum* DC., *Prodr. [A. P. de Candolle]* 1: 479 [mid Jan. 1824] (Candolle 1824). — *Eriodendron caribaeum* (DC.) G.Don, *Hort. Brit. [Loudon]* 292 (Don 1830). — *Ceiba pentandra* var. *caribaea* (DC.) Bakh., *Bull. Jard. Bot. Buitenzorg*, sér. 3, 6: 196 (Bakhuisen 1924). — *Ceiba caribaea* (DC.) A.Chev., *Rev. Bot. Appl. Agric. Trop.* 17: 266 (Chevalier 1937).

Eriodendron anfractuosum var. *indicum* DC., *Prodr. [A. P. de Candolle]* 1: 479 [mid Jan. 1824] (Candolle 1824). — *Ceiba pentandra* var. *indica* (DC.) Bakh., *Bull. Jard. Bot. Buitenzorg*, sér. 3, 6: 196 (Bakhuisen 1924).

Eriodendron anfractuosum var. *africanum* DC., *Prodr. [A. P. de Candolle]* 1: 479 [mid Jan. 1824] (Candolle 1824).

Bombax guineense Thonn., *Beskr. Guin. Pl.* 302 (Thonning 1827). — *Ceiba guineensis* (Thonn.) A.Chev., *Rev. Bot. Appl. Agric. Trop.* 17: 261 (Chevalier 1937).

Eriodendron guineense G.Don, *Hort. Brit. [Loudon]* 292 (Don 1830).

Eriodendron occidentale G.Don, *Gen. Hist.* 1: 513 [early Aug. 1831] (Don 1831).

Gossampinus rumphii Schott & Endl., *Melet. Bot.*: 35 (Schott & Endlicher 1832).

Eriodendron anfractuosum var. *guianense* Sagot, *Ann. Sci. Nat., Bot.* sér. 6, 11: 153 (Sagot 1881).

Ceiba anfractuosa M.Gómez, *Fl. Habanera* 141 (Gómez 1897).

Ceiba pentandra var. *clausa* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 30 [25 Apr. 1913] (Ulbrich 1913). — *Ceiba guineensis* var. *clausa* (Ulbr.) A.Chev., *Rev. Bot. Appl. Agric. Trop.* 17: 264 (Chevalier 1937).

Ceiba pentandra f. *albolana* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 30 [25 Apr. 1913] (Ulbrich 1913).

Ceiba pentandra f. *grisea* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 30 [25 Apr. 1913] (Ulbrich 1913).

Ceiba pentandra f. *albolana* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 31 [25 Apr. 1913] (Ulbrich 1913), nom. illeg. hom., non ibid. p. 30.

Ceiba pentandra f. *grisea* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 31 [25 Apr. 1913] (Ulbrich 1913), nom. illeg. hom., non ibid. p. 30.

Ceiba pentandra var. *dehiscens* Ulbr., *Notizbl. Königl. Bot. Gart. Berlin* 6: 31 [25 Apr. 1913] (Ulbrich 1913).

Eriodendron inerme A.Chev., *Explor. Bot. Afrique Occ. Franç.* 1: 75 (Chevalier 1920), *nom. nud.*

Ceiba guineensis var. *ampla* A.Chev., *Rev. Bot. Appl. Agric. Trop.* 17: 262 (Chevalier 1937).

Ceiba thonningii A.Chev., *Rev. Bot. Appl. Agric. Trop.* 17: 249 (Chevalier 1937).

NOTE. — Ulbrich (1913) created under each of his varieties *clausa* and *dehiscens* two formas to which he gave the same epithets *albolana* and *grisea*. Those under var. *dehiscens* (Ulbrich 1913: 31) are illegitimate homonyms of those under var. *clausa* (Ulbrich 1913: 30).

VERNACULAR NAMES. — Pa: kumak • Ka: kumaka • Te: kubaka • Wp: kumaka • Wn: kumaka • Nt: kankantii • Cr: frommajé • Fr: fromager, kapokier • Br: samaúma, sumaúma.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4580*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 190$ cm.

Genus *Eriotheca* Schott & Endl.

[1009] *Eriotheca crassa* (Uittien) A.Robyns

Bull. Jard. Bot. État Bruxelles 33 (1): 142 (Robyns 1963). — *Bom-
bax crassum* Uittien, *Recueil Trav. Bot. Néerl.* 22: 365 ["1925" publ.
Jan. 1926] (Uittien 1926).

VERNACULAR NAMES. — Pa: katatru-avain, katatu-avain • Ka: ityutano kilikili maululu • Wp: pilisi aminiyu • Nt: katun udu, yankomini • Cr: koton-siam, mao-koton.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Béna
1232*.

[1010] *Eriotheca globosa* (Aubl.) A.Robyns

Bull. Jard. Bot. État Bruxelles 33 (1): 142 (Robyns 1963). — *Bom-
bax globosum* Aubl., *Hist. Pl. Guiane* 2: 701 [Jun.-Dec. 1775]
(Aublet 1775).

Bombax sclerophyllum Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 58 [Sep.
1935] (Ducke 1935). — *Eriotheca macrophylla* (K.Schum.)
A.Robyns subsp. *sclerophylla* (Ducke) A.Robyns, *Bull. Jard. Bot.
État Bruxelles* 33 (2): 154 (Robyns 1963).

VERNACULAR NAMES. — Ka: kilikili maululu, ulu maululu • Wp:
pilisi aminiyu • Wn: hapèlek • Nt: katun udu, yankomini • Cr:
bwa-koton, mao-koton, tcho-bèf • Fr: mahot coton • Br: algodão-
bravo, mamorana-da-terra-firme, mamorana-grande.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D.
Sabatier & M.-F. Prévost 2578*.

INVENTORY DATA (FG). — 138 trees in 75 plots; $F_{\max} = 2.1\%$;
 $dbh_{\text{inv}} = 117$ cm.

[1011] *Eriotheca longitubulosa* A.Robyns

Bull. Jard. Bot. État Bruxelles 33 (2): 169 (Robyns 1963).

VERNACULAR NAMES. — Pa: katatru-avain, katatu-avain • Te: pitsiwit-
mididju • Nt: katun udu, yankomini • Cr: bwa-koton, koton-siam,
mao-koton • Fr: mahot coton • Br: sumaúma-brava.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F.
Prévost & D. Sabatier 2996*.

INVENTORY DATA (FG). — 82 trees in 53 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 76.4$ cm.

[1012] *Eriotheca surinamensis* (Uittien) A.Robyns

Bull. Jard. Bot. État Bruxelles 33 (2): 155 (Robyns 1963). — *Bom-
bax surinamense* Uittien, *Recueil Trav. Bot. Néerl.* 22: 365 ["1925"
publ. Jan. 1926] (Uittien 1926).

VERNACULAR NAMES. — Wn: wewe lémopka.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D.
Sabatier 3547*.

INVENTORY DATA (FG). — 1 tree, $dbh = 25.9$ cm.

[1013] *Eriotheca* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D.
Sabatier 2173*.

INVENTORY DATA (FG). — 6 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25.1$ cm.

Genus *Guazuma* Mill.

[1014] *Guazuma ulmifolia* Lam.

Encycl. [J. Lamarck et al.] 3 (1): 52 [19 Oct. 1789] (Lamarck
1789). — *Bubroma ulmifolia* (Lam.) Oken, *Allg. Naturgesch.* 3 (2):
1204 (Oken 1841). — *Guazuma guazuma* var. *ulmifolia* (Lam.)
Kuntze, *Revis. Gen. Pl.* 3 (3): 24 [28 Sep. 1898] (Kuntze 1898).

Theobroma guazuma L., *Sp. Pl.* 2: 782 [1 May 1753] (Linnaeus
1753). — *Bubroma guazuma* (L.) Willd., *Sp. Pl.*, ed. 43 (2): 1423
[1-10 Nov. 1802] (Willdenow 1802). — *Guazuma guazuma* (L.)
Cockerell, *Bull. Torrey Bot. Club* 19 (3): 95 (Cockerell 1892),
nom. inval. (tautonym).

Guazuma polybotrya Cav., *Icon. [Cavanilles]* 3 (2): 51 (Cavanilles
1796), "polybotra". — *Bubroma polybotryum* (Cav.) Willd., *Enum.
Pl. [Willdenow]* 2: 806 [June 1809] (Willdenow 1809).

Bubroma invira Willd., *Enum. Pl. [Willdenow]* 2: 806 [June 1809]
(Willdenow 1809). — *Guazuma invira* (Willd.) G.Don, *Gen.
Hist.* 1: 523 [early Aug. 1831] (Don 1831).

Guazuma tomentosa Kunth, *Nova genera et species plantarum [H.B.K.]*
5: 320 [24 Mar. 1823] (Kunth 1823). — *Bubroma tomentosum*
(Kunth) Spreng., *Syst. Veg. [Sprengel]* 3: 332 [Jan.-Mar. 1826]
(Sprengel 1826). — *Guazuma ulmifolia* var. *tomentosa* (Kunth)
K.Schum., *Fl. Bras. [Martius]* 12 (3): 81 [1 Mar. 1886] (Schumann
1886). — *Theobroma tomentosum* (Kunth) M.Gómez, *Anales Soc.
Esp. Hist. Nat.* 19 (2): 217 (Gómez 1890), "tomentosa". — *Gu-*




FIG. 35. — Malvaceae: **A**, *Apeiba tibourbou* Aubl.; **B**, *Pachira flaviflora* (Pulle) Fern. Alonso (D. Sabatier & J.-F. Molino 4820); **C**, *Sterculia speciosa* K.Schum. (D. Sabatier & M.-F. Prévost 5421); **D**, *Theobroma guianense* (Aubl.) J.F.Gmel. (D. Sabatier & M.-F. Prévost 2216). A, © J.-F. Molino/IRD; B-D, © D. Sabatier/IRD.

zuma guazuma var. *tomentosa* (Kunth) Kuntze, *Revis. Gen. Pl. 3* (3): 24 [28 Sep. 1898] (Kuntze 1898).

Guazuma bubroma Tussac, *Fl. Antill. 4*: 69 (Tussac 1827).

Guazuma blumei G.Don, *Gen. Hist. 1*: 523 [early Aug. 1831] (Don 1831), “*Blumii*”.

Guazuma tomentosa var. *cumanensis* G.Don, *Gen. Hist. 1*: 523 [early Aug. 1831] (Don 1831).

Guazuma tomentosa var. *monpoxensis* G.Don, *Gen. Hist. 1*: 523 [early Aug. 1831] (Don 1831).

Guazuma utilis Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher] 3*: 72 [23-25 Jan. 1845] (Poeppig 1845).

Guazuma parvifolia A.Rich., *Hist. Phys. Cuba, Pl. Vasc. 190* (Richard 1845).

Diuroglossum rufescens Turcz., *Bull. Soc. Imp. Naturalistes Moscou 25* (2): 156 (Turczaninow 1852).

Guazuma ulmifolia var. *glabra* K.Schum., *Fl. Bras. [Martius] 12* (3): 81 [1 Mar. 1886] (Schumann 1886).

Guazuma ulmifolia var. *tomentella* K.Schum., *Fl. Bras. [Martius] 12* (3): 81 [1 Mar. 1886] (Schumann 1886).

Guazuma ulmifolia var. *trianae* K.Schum., *Fl. Bras. [Martius] 12* (3): 82 [1 Mar. 1886] (Schumann 1886), in obs.

Guazuma ulmifolia var. *velutina* K.Schum., *Fl. Bras. [Martius] 12* (3): 81 [1 Mar. 1886] (Schumann 1886).

Guazuma coriacea Rusby, *Bull. New York Bot. Gard. 4* (14): 332 [7 Dec. 1907] (Rusby 1907).

Guazuma tomentosa var. *parvifolia* Kitan., *Fitologiya 11*: 48 (Kitanov 1979).

VERNACULAR NAMES. — Cr: bwa-lòm • Br: cabeça-de-negro, mutamba.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost 165*.

SIZE. — Up to 60 cm dbh (Robyns & Cuatrecasas 1964).

Genus *Huberodendron* Ducke[1015] *Huberodendron swietenioides* (Gleason) Ducke

Arch. Inst. Biol. Veg. 2 (1): 72 [Sep. 1935] (Ducke 1935). — *Bernoullia swietenioides* Gleason, *Phytologia* 1 (2): 109 [4 Aug. 1934] (Gleason 1934).

VERNACULAR NAMES. — Pa: waravru-kamwi, waravru-wašiunu • Nt: awalaime, gaan dyaba udu • Br: munguba-da-mata.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4926*.

INVENTORY DATA (FG). — 7 trees in 4 plots; dbh_{inv} = 80 cm.

Genus *Luehea* Willd.[1016] *Luehea speciosa* Willd.

Ges. Naturf. Freunde Berlin Neue Schriften 3: 410 (Willdenow 1801), “*Lühea*”.

Cedrus alternifolia Mill., *Gard. Dict.*, ed. 8, n. 3 [16 Apr. 1768] (Miller 1768), “*Alternifolius*”, nom. rej. — *Cedrela alternifolia* (Mill.) Steud., *Nomencl. Bot. [Steudel]* 1: 170 (Steudel 1821), nom. rej. — *Swietenia alternifolia* (Mill.) Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 653 (Steudel 1841), nom. rej. — *Luehea alternifolia* (Mill.) Mabb., *Gard. Bull. Singapore* 54 (2): 254 [1 Jan. 2002] (Mabberley 2002), nom. rej.

Apeiba ulmifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 347 [24 Mar. 1823] (Kunth 1823).

Broteria maritima Vell., *Fl. Flumin.*: 323 [“1825” publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Luehea ferruginea Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 31 (1): 224 (Turczaninow 1858), “*Lühea*”.

Luehea tarapotina J.F.Macbr., *Candollea* 5: 382 (Macbride 1934).

VERNACULAR NAMES. — Wp: etune, pikau’i • Wn: etuwe • Nt: dyaba udu • Br: açoita-cavallo, ivatingi, mutamba-preta, uaçima-do-campo.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 970*.

INVENTORY DATA (FG). — 16 trees in 2 plots; F_{max} = 2.7 %; dbh_{inv} = 11.3 cm.

Genus *Lueheopsis* Burret[1017] *Lueheopsis althaeiflora* (Spruce ex Benth.) Burret

Notizbl. Bot. Gart. Berlin-Dahlem 9: 840 [22 July 1926] (Burret 1926). — *Luehea althaeiflora* Spruce ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 58 [May 1861] (Bentham 1861), “*althaeiflora*”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2001*.

SIZE. — Colombia, Vaupés. *S. Defler 590* (MO), 45 cm.

[1018] *Lueheopsis rosea* (Ducke) Burret

Notizbl. Bot. Gart. Berlin-Dahlem 9: 837 [22 July 1926] (Burret 1926). — *Luehea rosea* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 208 (Ducke 1922).

Luehea flavescentis Uittien, *Recueil Trav. Bot. Néerl.* 22: 363 [“1925” publ. Jan. 1926] (Uittien 1926). — *Lueheopsis flavescens* (Uittien) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 842 [22 July 1926] (Burret 1926).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: etune, pikau’i • Wn: etuwe.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3219*.

INVENTORY DATA (FG). — 2 trees in 2 plots; F_{max} < 1 %; dbh_{inv} = 26.4 cm.

[1019] *Lueheopsis rugosa* (Pulle) Burret

Notizbl. Bot. Gart. Berlin-Dahlem 9: 840 [22 July 1926] (Burret 1926). — *Luehea rugosa* Pulle, *Recueil Trav. Bot. Néerl.* 6: 274 (Pulle 1909), “*Lühea*”.

Luehea maroniensis Benoist, *Bull. Mus. Natl. Hist. Nat.* 27: 114 (Benoist 1921).

VERNACULAR NAMES. — Ka: apopopano epityi, kusewelan • Wp: etune, pikau’i • Nt: dyaba udu.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4170*.

INVENTORY DATA (FG). — 111 trees in 69 plots; F_{max} < 1 %; dbh_{inv} = 76.4 cm.

Genus *Matisia* Bonpl.[1020] *Matisia lasiocalyx* K.Schum.

Fl. Bras. [Martius] 12 (3): 239 [1 Nov. 1886] (Schumann 1886). — *Quararibea lasiocalyx* (K.Schum.) Vischer, *Bull. Soc. Bot. Genève*, sér. 2, 11: 206 (Vischer 1920).

VERNACULAR NAMES. — Wp: yapu tulu’i, yapu tulu’i sī • Br: inajarana-envira, laranjinha.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *P. Grenand 677*.

SIZE. — Ecuador, Pastaza. *E. Gudiño et al. 1628* (MO), 15 m × 10 cm.

[1021] *Matisia ochrocalyx* K.Schum.

Fl. Bras. [Martius] 12 (3): 238 [1 Nov. 1886] (Schumann 1886). — *Quararibea ochrocalyx* (K.Schum.) Vischer, *Bull. Soc. Bot. Genève*, sér. 2, 11: 206 (Vischer 1920).

Quararibea muricata Cuatrec., *Bot. Mus. Leaf.* 15 (1): 53 [25 May 1951] (Cuatrecasas 1951). — *Matisia muricata* (Cuatrec.) Cuatrec., *Phytologia* 4 (8): 479 [Jan. 1954] (Cuatrecasas 1954).

Matisia apaporiensis Cuatrec., *Bot. Mus. Leaf.* 17 (3): 82 [21 Nov. 1955] (Cuatrecasas 1955). — *Quararibea apaporiensis* (Cuatrec.) A.Robyns & S.Nilsson, *Bull. Jard. Bot. Natl. Belg.* 44 (1-2): 87 (Robyns & Nilsson 1974).

VERNACULAR NAMES. — Pa: ikun-kamwi.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 4450*.

INVENTORY DATA (FG). — 7 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22\text{ cm}$.

Genus *Pachira* Aubl.

[1022] *Pachira aquatica* Aubl.

Hist. Pl. Guiane 2: 726 [Jun.-Dec. 1775] (Aublet 1775). — *Bombax aquaticum* (Aubl.) K.Schum., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (6): 62 (Schumann 1890). — *Carolinea princeps* L.f., *Suppl. Pl.*: 314 ["1781" publ. Apr. 1782] (Linnaeus 1782), *nom. illeg. superfl.* (based on *Pachira aquatica*).

Sophia carolina L., *Pl. Surin.*: 11 [23 June 1775] (Linnaeus 1775).

Carolinea macrocarpa Schlehd. & Cham., *Linnaea* 6: 423 (Schlehd. & Chamisso 1831). — *Pachira macrocarpa* (Schlehd. & Cham.) Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 329 [18-20 Sep. 1842] (Walpers 1842). — *Bombax macrocarpum* (Schlehd. & Cham.) K.Schum., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (6): 62 (Schumann 1890).

Pachira aquatica var. *surinamensis* Decne., *Fl. Serres Jard. Eur.* 23: 46 (Decaisne 1880).

Pachira pustulifera Pittier, *Repert. Spec. Nov. Regni Veg.* 13: 315 [5 May 1914] (Pittier 1914).

Pachira villosula Pittier, *Repert. Spec. Nov. Regni Veg.* 13: 316 [5 May 1914] (Pittier 1914).

Bombax rigidifolium Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 127 (Ducke 1925).

Pachira aquatica var. *occidentalis* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 9 (35): 169 [July 1954] (Cuatrecasas 1954).

VERNACULAR NAMES. — Pa: waravru-kamwi • Ka: mamao, maomao, maumau • Te: dzepapa'i • Wp: māū'i • Wn: akuli tumu, palalagwa • Nt: momow • Cr: kakao-larivéy, kakao-sovaj • Fr: cacao rivière • Br: mamão-rana, mamorana, munguba-rana.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000645671](#)]).

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30\text{ cm}$.

[1023] *Pachira dolichocalyx* A.Robyns

Bull. Jard. Bot. Natl. Belg. 58 (3-4): 535 (Robyns 1988).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: momow.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost 840* (holo-, P[P00077210]; iso-, P[P00077211, P05273658], U[U0000784, U0000785]).

INVENTORY DATA (FG). — 132 trees in 28 plots; $F_{\max} = 2\%$; $dbh_{\text{inv}} = 49.3\text{ cm}$.

[1024] *Pachira flaviflora* (Pulle) Fern.Alonso (Fig. 35B)

Anales Jard. Bot. Madrid 56 (2): 310 (Fernández Alonso 1998). — *Bombax flaviflorum* Pulle, *Recueil Trav. Bot. Néerl.* 9: 150 (Pulle 1912). — *Rhodographalopsis flaviflora* (Pulle) A.Robyns, *Bull. Jard. Bot. État Bruxelles* 33 (2): 285 (Robyns 1963). — *Pochota flaviflora* (Pulle) Steyermark & W.D.Stevens, *Ann. Missouri Bot. Gard.* 75 (1): 397 [31 May 1988] (Steyermark & Stevens 1988).

VERNACULAR NAMES. — Ka: kilikili maululu.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 4820*.

INVENTORY DATA (FG). — 168 trees in 19 plots; $F_{\max} = 5.1\%$; $dbh_{\text{inv}} = 32\text{ cm}$.

[1025] *Pachira insignis* (Sw.) Sw. ex Savigny

Encycl. [J. Lamarck et al.] 4 (2): 690 [1 Nov. 1798] (Savigny 1798). — *Carolinea insignis* Sw., *Prodr. [Swartz]* 101 [20 Jun.-29 July 1788] (Swartz 1788).

Pachira spruceana Decne., *Fl. Serres Jard. Eur.* 23: 46 (Decaisne 1880).

Bombax spruceanum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 126 (Ducke 1925).

NOTE. — *Bombax spruceanum* Ducke is not based on *P. spruceana* Decne.

VERNACULAR NAMES. — Te: dzepapa'i • Wn: nukunuku • Nt: momow.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2378*.

INVENTORY DATA (FG). — 7 tree in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 46\text{ cm}$.

[1026] *Pachira macrocalyx* (Ducke) Fern.Alonso

Anales Jard. Bot. Madrid 56 (2): 310 (Fernández Alonso 1998). — *Bombax macrocalyx* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 124 (Ducke 1925). — *Bombacopsis macrocalyx* (Ducke) A.Robyns, *Bull. Jard. Bot. État Bruxelles* 33 (2): 203 (Robyns 1963).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 25303*.

SIZE. — Up to 15 cm dbh (Esteves 2005).

Genus *Pseudobombax* Dugand

[1027] *Pseudobombax munguba* (Mart.) Dugand

Mutisia 9: 4 (Dugand 1952). — *Bombax munguba* Mart., *Nova genera et species plantarum [Martius]* 1 (4): 93 ["1824" publ. Jan.-Mar. 1826] (Martius 1826).

Pseudobombax amapaense A.Robyns, *Mem. New York Bot. Gard.* 17 (1): 195 (Robyns 1967).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: mōngi • Wn: ēhpaimē • Br: munguba.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *P. Grenand 701*.

SIZE. — Brazil, Amazonas. *G.T. Prance 13371* (MO), 10 m × 40 cm.

Genus *Quararibea* Aubl.

[1028] *Quararibea duckei* Huber

Bull. Soc. Bot. Genève, sér. 2, 6: 186 [“1914” publ. 1915] (Huber 1915).

VERNACULAR NAMES. — Wp: yapu tulu’i • Wn: meikolo tamit • Nt: baaka kiikii • Cr: bwa-jelé, bwa-lélé • Br: inajarana.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2610*.

INVENTORY DATA (FG). — 421 trees in 40 plots; $F_{\max} = 16.1\%$; $dbh_{\text{inv}} = 28.6$ cm.

[1029] *Quararibea guianensis* Aubl.

Hist. Pl. Guiane 2: 692 [Jun.-Dec. 1775] (Aublet 1775).

Quararibea machin J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3A.2)*: 618 (Macbride 1956).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kuuku-ariut-kamwi • Ka: apesulu, maipyuli epu • Wp: yapu tulu’i, yapu tulu’i si • Nt: kiikii tiki, weti kiikii • Cr: bwa-jelé, bwa-lélé • Br: envira-sapotinha.

HERBARIUM DATA (FG). — 69 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material G[G00226718]).

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.5$ cm.

[1030] *Quararibea spatulata* Ducke

Trop. Woods 76: 20 (Ducke 1943).

NOTE. — Four specimens identified at CAY as *Q. amazonica* Ulbr. belong here.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *S.A. Mori 22252*.

INVENTORY DATA (FG). — 25 trees in 8 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 36.9$ cm.

Genus *Sterculia* L.

[1031] *Sterculia frondosa* Rich.

Actes Soc. Hist. Nat. Paris 1: 111 [Oct. 1792] (Richard 1792). — *Clompanus frondosa* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 78 [5 Nov. 1891] (Kuntze 1891).

Sterculia glabrifolia Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 1148 [25 Mar. 1927] (Mildbraed 1927).

Sterculia roseiflora Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 58 [Sep. 1935] (Ducke 1935).

VERNACULAR NAMES. — Pa: waravru-ahavukune, yit-itaibi-ahawukune • Wp: iwi si • Cr: bwa-kalou, mao-kochon • Br: axixá.

HERBARIUM DATA (FG). — 30 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (type, not traced); *M.-F. Prévost 2071*.

INVENTORY DATA (FG). — 48 trees in 33 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 61.8$ cm.

[1032] *Sterculia kayae* P.E.Berry

Fl. Venez. Guayana 9: 266 [5 Oct. 2005] (Berry 2005). — *Sterculia pruriens* var. *parviflora* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 130 (Ducke 1925). — *Sterculia parviflora* (Ducke) E.L.Taylor ex Brako & Zarucchi, *Monogr. Syst. Bot. Missouri Bot. Gard.* 45: 1259 [1 Oct. 1993] (Brako & Zarucchi 1993), *nom. illeg. hom.*, non Roxb. ex G.Don (Don 1831).

VERNACULAR NAMES. — Pa: yit-gataibi, yit-itaibi • Wp: iwi si • Cr: bwa-kalou, mao-kochon.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 944*.

INVENTORY DATA (FG). — 86 trees in 41 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 106$ cm.

[1033] *Sterculia multiovula* E.L.Taylor ex Mondragón

Bol. Centro Invest. Biol. Univ. Zulia 40 (2): 169 (Mondragón 2006).

NOTE. — First invalidly published by E.L. Taylor (1989: 110).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori et al. 26565*.

INVENTORY DATA (FG). — 12 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54.7$ cm.

[1034] *Sterculia pruriens* (Aubl.) K.Schum.

Fl. Bras. [Martius] 12 (3): 8 [1 Mar. 1886] (Schumann 1886). — *Ivira pruriens* Aubl., *Hist. Pl. Guiane* 2: 695 [Jun.-Dec. 1775] (Aublet 1775), “*Iuira Pruriens*” on plate. — *Sterculia ivira* Sw., *Prodri. [Swartz]*: 98 [20 Jun.-29 July 1788] (Swartz 1788), *nom. illeg. superfl.* (based on *Ivira pruriens*). — *Clompanus pruriens* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 78 [5 Nov. 1891] (Kuntze 1891).

Sterculia pruriens var. *grandiflora* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 130 (Ducke 1925).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: kilikili maululu, maipyuli omoseli • Te: tapi’it lupabi • Wp: iwi si • Wn: ēwa • Nt: kobe • Cr: bwa-kalou, mao-kochon • Fr: mahot cochon • Br: axixá, castanha-de-piriquito, tacacazeiro.

HERBARIUM DATA (FG). — 67 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000645788](#)]), LINN[[LINN-HS 1498.15](#)]; *D. Sabatier & M.-F. Prévost 3015*.

INVENTORY DATA (FG). — 506 trees in 163 plots; $F_{\max} = 2.6\%$; $\text{dbh}_{\text{inv}} = 160.1\text{ cm}$.

[1035] *Sterculia speciosa* K.Schum.
(Fig. 35C)

Fl. Bras. [Martius] 12 (3): 7 [1 Mar. 1886] (Schumann 1886). — *Clompanus speciosa* (K.Schum.) Kuntze, *Revis. Gen. Pl.* 1: 78 [5 Nov. 1891] (Kuntze 1891).

Sterculia pilosa Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 212 (Ducke 1922). — *Xylosterculia pilosa* (Ducke) Kosterm., *Bot. Tidsskr.* 67 (4): 317 (Kostermans 1973).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017).

VERNACULAR NAMES. — Pa: yit-gataibi, yit-itaibi, yit-itaibi-purubumna • Wp: tapilupami • Nt: kobe • Cr: mao-kochon • Fr: mahot cochon • Br: axixá, capoteiro, tacacazeiro.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5421*.

INVENTORY DATA (FG). — 191 trees in 100 plots; $F_{\max} = 1.3\%$; $\text{dbh}_{\text{inv}} = 74.8\text{ cm}$.

[1036] *Sterculia villifera* Steud.

Flora 26 (45): 755 [17 Dec. 1843] (Steudel 1843).

Sterculia surinamensis R.Br., *Pl. Jav. Rar. [Bennett]* 3: 229 [Nov. 1844] (Brown 1844).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kilikili amoti, maipyuli italali, patulapo • Wp: iwi sī, tapilupami • Cr: bwa-kalou, mao-kochon • Fr: mahot cochon • Br: axixá, tacacazeiro.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2437*.

INVENTORY DATA (FG). — 31 trees in 12 plots; $F_{\max} = 1.3\%$; $\text{dbh}_{\text{inv}} = 78.5\text{ cm}$.

Genus *Talipariti* Fryxell

[1037] *Talipariti tiliaceum* (L.) Fryxell
var. *pernambucense* (Arruda) Fryxell

Contr. Univ. Michigan Herb. 23: 262 (Fryxell 2001). — *Hibiscus pernambucensis* Arruda, *Diss. Pl. Brazil.* 44 (Arruda 1810). — *Paritium pernambucense* (Arruda) G.Don, *Gen. Hist.* 1: 485 [early Aug. 1831] (Don 1831). — *Hibiscus tiliaceus* L. subsp. *pernambucensis* (Arruda) A.Cast., *Sellowia* 19: 50 (Castellanos 1967). — *Hibiscus tiliaceus* var. *pernambucensis* (Arruda) I.M.Johnst., *Sargentia* 8: 196 (Johnston 1949). — *Talipariti pernambucense* (Arruda) Bovini, *Rodriguésia* 61 (Sup.): S20 (Bovini 2010).

Hibiscus bracteosus DC., *Prodr. [A. P. de Candolle]* 1: 455 [mid Jan. 1824] (Candolle 1824).

Hibiscus arboreus Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 49 [Oct. 1825] (Hamilton 1825).

Hibiscus fragrantissimus Sessé & Moc., *Pl. Nov. Hisp.*: 113 (Sessé & Mociño 1889).

Pariti tiliaceum (L.) A.St.-Hil. f. *immaculatum* O.Deg. & Greenwell, *Fl. Hawaii.*: Fam. 221 (Degener & Greenwell 1956). — *Hibiscus tiliaceus* f. *immaculatus* (O.Deg. & Greenwell) H.St.John, *Mem. Pacific Trop. Bot. Gard.* 1: 230 (St.John 1973).

NOTE. — A species restricted to wet savannas.

VERNACULAR NAMES. — Pa: imavui-kamwi • Ka: waiyen, waye • Wp: uluku lá • Cr: bwa-mao • Br: algodão-do-brejo, envira-do-mangue.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *F. Billet 1286*.

SIZE. — Costa Rica. *R. Aguilar 2961* (MO), 7 m × 10 cm.

Genus *Theobroma* L.

[1038] *Theobroma cacao* L.

Sp. Pl. 2: 782 [1 May 1753] (Linnaeus 1753). — *Cacao sativa* Aubl., *Hist. Pl. Guiane* 2: 689 [Jun.-Dec. 1775] (Aublet 1775), *nom. illeg. superfl.* (based on *Theobroma cacao*). — *Cacao sativa* Lam., *Encycl. J. Lamarck et al.* 1 (2): 533 [1 Aug. 1785] (Lamarck 1785), isonym. et *nom. illeg. superfl.* (based on *Theobroma cacao*). — *Cacao theobroma* Tussac, *Fl. Antill.* 1: 101 (Tussac 1808), *nom. illeg. superfl.* (based on *Theobroma cacao*). — *Theobroma sativum* Lign. & Le Bey, *Bull. Soc. Linn. Normandie*, sér. 5, 8: 233 [15 Oct. 1905] (Lignier & Le Bey 1905), “*sativum*”, *nom. inval.* (based on the illegitimate *Cacao sativa* Aubl.). — *Theobroma cacao* subsp. *sativum* Léon, *Cacao Manual [F. Hardy]*: 312 (Léon 1960), *nom. inval.* (based on the illegitimate *Cacao sativa*).

Cacao minus Gaertn., *Fruct. Sem. Pl.* 2: 190 [Apr.-May 1791] (Gaertner 1791). — *Theobroma minus* (Gaertn.) A.Chev., *Rev. Bot. Appl. Agric. Trop.* 26 (285): 283 (Chevalier 1946).

Theobroma integerrimum Stokes, *Bot. Mat. Med.* 4: 83 (Stokes 1812), “*integerrima*”.

Theobroma caribaeum Sweet, *Hort. Brit. [Sweet]*, ed. 2, 67 (Sweet 1830), “*caribaea*”, *nom. nud.*

Theobroma leiocarpum Bernoulli, *Uebersicht Theobroma* 6 (Bernoulli 1869), “*leiocarpa*”. — *Theobroma cacao* var. *leiocarpum* (Bernoulli) Cif., *Mem. Cl. Sci. Fis. Mat. Nat. Reale Accad. Italia* 4: 604 (Ciferri 1933), “*leiocarpa*”. — *Theobroma cacao* f. *leiocarpum* (Bernoulli) Ducke, *Rodriguésia* 4 (13): 274 (Ducke 1940). — *Theobroma cacao* subsp. *leiocarpum* (Bernoulli) Cuatrec., *Publ. Field Mus. Nat. Hist. Bot. Ser.* 13 (3A.2): 654 (Cuatrecasas 1956).

Theobroma pentagonum Bernoulli, *Uebersicht Theobroma* 6 (Bernoulli 1869), “*pentagona*”. — *Theobroma cacao* f. *pentagonum* (Bernoulli) Ducke, *Rodriguésia* 4 (13): 274 (Ducke 1940). — *Theobroma cacao* subsp. *pentagonum* (Bernoulli) Léon, *Cacao Manual [F. Hardy]*: 312 (Léon 1960), “*pentagona*”.

Theobroma saltzmannianum Bernoulli, *Uebersicht Theobroma* 7 (Bernoulli 1869), “*Saltzmanniana*”.

Theobroma kalagua De Wild., *Bull. Herb. Boissier* 7: 957 (De Wildeman 1899).

Theobroma sphaerocarpum A.Chev., *Vég. Ut. Afr. Trop. Franç.* 4: 12 (Chevalier 1908), “*sphaerocarpa*”. — *Theobroma cacao* subsp. *sphaerocarpum* (A.Chev.) Cuatrec., *Contr. U.S. Natl. Herb.* 35 (6): 515 [21 Aug. 1964] (Cuatrecasas 1964).

Theobroma sapidum Pittier, *Bol. Soc. Venez. Ci. Nat.* 1: 183 (Pittier 1932), *nom. nud.*

Theobroma cacao var. *typicum* Cif., *Mem. Cl. Sci. Fis. Mat. Nat. Reale Accad. Italia* 4: 604 (Ciferri 1933), “*typica*”, *nom. inval.* (Turland et al. 2018: Art. 24.3).

Theobroma sativum var. *leucospermum* A.Chev., *Rev. Bot. Appl. Agric. Trop.* 26 (285): 270, 272 (Chevalier 1946), “*leucosperma*”.

Theobroma sativum var. *melanospermum* A.Chev., *Rev. Bot. Appl. Agric. Trop.* 26 (285): 270, 272 (Chevalier 1946), “*melanosperma*”.

NOTES. — Hyperdominant in Amazonia (ter Steege et al. 2020), and semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017). Probably introduced to French Guiana by pre-Columbian Amerindians. It is now naturalised on or near archaeological sites in southern French Guiana.

VERNACULAR NAMES. — Pa: waravru • Ka: kakao • Te: akau • Wp: walapulu • Wn: walapulu • Nt: kakaw • Cr: kakao • Fr: cacaoyer • Br: cacau.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P. Grenand 1044*.

INVENTORY DATA (FG). — 32 trees in 2 plots; $F_{\max} = 3.9\%$; $dbh_{inv} = 23.2$ cm.

[1039] *Theobroma guianense* (Aubl.) J.F.Gmel.
(Fig. 35D)

Syst. Nat., ed. 13[bis], 2 (2): 1151 [“1791” publ. late Apr.-Oct. 1792] (Gmelin 1792), “*guianensis*”. — *Cacao guianensis* Aubl., *Hist. Pl. Guiane* 2: 683 [Jun.-Dec. 1775] (Aublet 1775), “*Guyanensis*” on plate.

Cacao sylvestris Aubl., *Hist. Pl. Guiane* 2: 687 [Jun.-Dec. 1775] (Aublet 1775). — *Theobroma sylvestre* (Aubl.) Mart., *Repert. Pharm. [J.A. Buchner]* 35: 24 (Martius 1830), “*sylvestris*”.

Theobroma subincanum Mart., *Repert. Pharm. [J.A. Buchner]* 35: 23 (Martius 1830).

Theobroma ferrugineum Bernoulli, *Uebersicht Theobroma*: 13 (Bernoulli 1869).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020, as *T. subincanum* Mart.) Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017, as *T. subincanum*). *Cacao guianensis* Aubl. is the oldest name for this taxon, thus *T. guianense* has priority over *T. subincanum*. Contrary to Cuatrecasas’ statement (Cuatrecasas, 1964: 391, 564), Martius was the first to make the new combination *Theobroma sylvestre* (Aubl.) Mart., before D. Don (1831).

VERNACULAR NAMES. — Pa: kupuh-ahavukune, kupuh-kamwi, waravru-ahavukune • Wp: kapia’i • Nt: busi kakaw, supun udu • Cr: koupu-asou-sovaj • Fr: cupuaçu sauvage • Br: busi momow, cupuai, cupuiá.

HERBARIUM DATA (FG). — 72 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000522439]).

INVENTORY DATA (FG). — 457 trees in 130 plots; $F_{\max} = 1.9\%$; $dbh_{inv} = 38.7$ cm.

[1040] *Theobroma velutinum* Benoist

Bull. Mus. Natl. Hist. Nat. 27: 113 (Benoist 1921).

Herrania guianensis Sagot ex K.Schum., *Fl. Bras. [Martius]* 12 (3): 75 [1 Mar. 1886] (Schumann 1886), *nom. nud. pro syn.*

VERNACULAR NAMES. — Te: akau pepen • Wp: aka’i • Wn: akuli tumu • Nt: busi kakaw • Cr: kakao-gran-bwa • Br: cacauí, cacaurana.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *R. Benoist 516* (type, not traced); *D. Sabatier & M.-F. Prévost 4403*.

INVENTORY DATA (FG). — 28 trees in 25 plots; $F_{\max} < 1\%$; $dbh_{inv} = 20.7$ cm.

Family MELASTOMATACEAE Juss.

Genus *Bellucia* Neck. ex Raf.

[1041] *Bellucia arborescens* (Aubl.) Baill.
(Fig. 36A)

Hist. Pl. [Baillon] 7: 34 [Dec. 1877] (Baillon 1877). — *Melastoma arborescens* Aubl., *Hist. Pl. Guiane* 1: 420 [Jun.-Dec. 1775] (Aublet 1775). — *Loreya arborescens* (Aubl.) DC., *Prod. [A. P. de Candolle]* 3: 179 [mid Mar. 1828] (Candolle 1828). — *Oxisma arborescens* (Aubl.) Raf., *Sylva Tellur.*: 94 (Rafinesque 1838).

Loreya acutifolia O.Berg ex Triana, *Trans. Linn. Soc. London* 28 (1): 142 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

Loreya maguirei Wurdack, *Phytologia* 18 (3): 160 [27 Mar. 1969] (Wurdack 1969).

VERNACULAR NAMES. — Pa: kudumig-seine • Nt: lebi musupu • Cr: bwa-mèl.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN, LINN-HS 782.15).

INVENTORY DATA (FG). — 72 trees in 46 plots; $F_{\max} = 6.7\%$; $dbh_{inv} = 73.5$ cm.

[1042] *Bellucia egensis* (DC.)

Penney, Michelang., Judd & Almeda

Syst. Bot. 35 (4): 796 [1 Oct. 2010] (Penney et al. 2010). — *Myriaspora egensis* DC., *Prod. [A. P. de Candolle]* 3: 165 [mid Mar. 1828] (Candolle 1828).

Myriaspora paulensis DC., *Prod. [A. P. de Candolle]* 3: 165 [mid Mar. 1828] (Candolle 1828). — *Melastoma paulense* Schrank ex DC., *Prod. [A. P. de Candolle]* 3: 165 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.* — *Melastoma egense* Mart. ex DC., *Prod. [A. P. de Candolle]* 3: 165 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Myriaspora surinamensis Steud., *Flora* 27: 722 (Steudel 1844).

Myriaspora decipiens Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 101 (Naudin 1851).

Myriaspora integrifolia Steud. ex Triana, *Trans. Linn. Soc. London 28 (1):* 145 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872), *nom. nud. pro syn.*

VERNACULAR NAMES. — Pa: kudumig-kamwi • Wp: imi'i, umi'i • Nt: mabee siii.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *P. Grenand 756*, "Petit arbre 10-15 m".

[1043] *Bellucia grossularioides* (L.) Triana

Trans. Linn. Soc. London 28 (1): 141 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Melastoma grossularioides* L., *Sp. Pl. 1:* 390 [1 May 1753] (Linnaeus 1753).

Blakea quinquenervia Aubl., *Hist. Pl. Guiane 1:* 525 [Jun.-Dec. 1775] (Aublet 1775). — *Blakea triplinervia* L.f., *Suppl. Pl.:* 246 ["1781" publ. Apr. 1782] (Linnaeus 1782), *nom. illeg. superfl.* (based on *Blakea quinquenervia*). — *Webera quinquenervia* (Aubl.) J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1):* 820 [late Sep.-Nov. 1791] (Gmelin 1791). — *Apatitia blakeoides* Desv. ex Ham., *Prodri. Pl. Ind. Occid. [Hamilton]:* 42 [Oct. 1825] (Hamilton 1825), *nom. illeg. superfl.* (based on *Blakea quinquenervia*). — *Bellucia nervosa* Raf., *Sylva Tellur.:* 93 (Rafinesque 1838), *nom. illeg. superfl.* (based on *Blakea triplinervia* and *B. quinquenervia*). — *Bellucia aubletii* Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 102 (Naudin 1851), *nom. illeg. superfl.* (based on *Blakea quinquenervia*). — *Bellucia quinquenervia* (Aubl.) H.Karst., *Linnaea 30:* 159 (Karsten 1859).

Blakea macrophylla D.Don, *Mem. Wern. Nat. Hist. Soc. 4:* 326 [May 1823] (Don 1823). — *Bellucia macrophylla* (D.Don) Triana, *Trans. Linn. Soc. London 28 (1):* 142 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

Bellucia hostmannii Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 103 (Naudin 1851).

Ischyranthera laevigata Steud. ex Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 103 (Naudin 1851), *nom. nud. pro syn.*

Bellucia brasiliensis Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 104 (Naudin 1851).

Bellucia superba Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 104 (Naudin 1851).

Bellucia multiflora H.Karst., *Linnaea 30:* 158 (Karsten 1859).

Bellucia circumscissa Spruce ex Cogn., *Fl. Bras. [Martius] 14 (4):* 515 [15 Aug. 1888] (Cogniaux 1888).

VERNACULAR NAMES. — Pa: asaki • Ka: askali, mainyapo • Te: put-sulu • Wp: pisulu • Wn: puhu • Nt: baaka musupu • Cr: bwa-mèl, grènn-mèl, mésoupou • Fr: graine merle, nélrier • Br: araçá-de-anta, goiaba-de-anta, mandapuça.

HERBARIUM DATA (FG). — 79 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Blakea quinquenervia*: BM[BM001008261], LINN[LINN-HS 853.1]).

INVENTORY DATA (FG). — 13 trees in 10 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 39.2$ cm.

[1044] *Bellucia mespiloides* (Miq.) J.F.Macbr.

Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (4.1): 498 (Macbride 1941). — *Loreya mespiloides* Miq., *Linnaea 18:* 619 ["1844" publ. prob. Aug. 1845] (Miquel 1845).

Henriettea brunnescens Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8):* 247 [24 Oct. 1929] (Standley 1929). — *Loreya brunnescens* (Standl.) Gleason, *Phytologia 3 (7):* 346 [18 Dec. 1950] (Gleason 1950).

VERNACULAR NAMES. — Ka: pakila potay, pakila wesopotali • Wp: imi'i, umi'i • Nt: musupu.

HERBARIUM DATA (FG). — 53 collections at CAY. Sel. exs.: *M.-F. Prévost 4014*.

INVENTORY DATA (FG). — 1 tree, $dbh = 13.1$ cm.

[1045] *Bellucia subrotundifolia* Wurdack

Phytologia 45 (4): 331 [Apr. 1980] (Wurdack 1980). — *Loreya subrotundifolia* (Wurdack) S.S.Renner, *Mem. New York Bot. Gard. 50:* 46 (Renner 1989).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: musupu.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *B.H.J. de Jong 2* (holo-, WAG[WAG0002329]; iso-, US[US0123663]).

SIZE. — Up to 15 m tall (Renner 1989).

Genus *Henriettea* DC.

[1046] *Henriettea duckeana* (Hoehne)
Penneys, Michelang., Judd & Almeda

Syst. Bot. 35 (4): 797 [1 Oct. 2010] (Penneys et al. 2010). — *Henrietella duckeana* Hoehne, *Anexos Mem. Inst. Butantan, Secc. Bot. 1 (5):* 170 (Hoehne 1922).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost 1910*.

SIZE. — Up to 10 m tall (Wurdack et al. 1993).

[1047] *Henriettea flavescens* (Aubl.) Baill.

Hist. Pl. [Baillon] 7: 34 [Dec. 1877] (Baillon 1877). — *Melastoma flavescens* Aubl., *Hist. Pl. Guiane 1:* 423 [Jun.-Dec. 1775] (Aublet 1775). — *Ossaea flavescens* (Aubl.) DC., *Prodri. [A. P. de Candolle] 3:* 169 [mid Mar. 1828] (Candolle 1828). — *Loreya flavescens* (Aubl.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 18:* 110 [Aug. 1852] (Naudin 1852). — *Henrietella flavescens* (Aubl.) Triana, *Trans. Linn. Soc. London 28 (1):* 143 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

HERBARIUM DATA (FG). — 54 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM001008254], LINN[LINN-HS 782.23]).

INVENTORY DATA (FG). — 53 trees in 29 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 19.6$ cm.

[1048] *Henriettea maroniensis* Sagot

Ann. Sci. Nat., Bot. sér. 6, 15: 330 (Sagot 1883).

VERNACULAR NAMES. — Pa: arakeu, arakeu-seine, arateu, arateu-seine • Ka: akuli wesepotapili, kapuwa nana.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *E.M. Mélinton s.n.* (holo-, P[P01818703]; iso-, BR[BR0000005185344], K[K000537032, K000537033], P[P01818704, P01818705]).

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 10.3$ cm.

[1049] *Henriettea multiflora* Naudin

Ann. Sci. Nat., Bot. sér. 3, 18: 105 [Aug. 1852] (Naudin 1852).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kudumig-kamwi • Ka: nana polan, pakila yuyulu.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-P. Lescure 816*.

SIZE. — Up to 16 m tall (Wurdack *et al.* 1993).

[1050] *Henriettea patrisiana* DC.

Prodr. [A. P. de Candolle] 3: 178 [mid Mar. 1828] (Candolle 1828). — *Henriettella patrisiana* (DC.) Naudin, *Ann. Sci. Nat., Bot., sér. 3, 18:* 107 [Aug. 1852] (Naudin 1852).

Henriettea parviflora Griseb., *Cat. Pl. Cub. [Grisebach] 95* [May-Aug. 1866] (Grisebach 1866). — *Henriettella parviflora* (Griseb.) Triana, *Trans. Linn. Soc. London 28 (1):* 143 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

Henriettea williamii Brade, *Publ. Inst. Nac. Pesq. Amaz., Bot. 8:* 17 (Brade 1958).

VERNACULAR NAMES. — Wp: takalawelu lá.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P. Grenand 1510*.

SIZE. — Brasil, Roraima. *J.A. Ratter et al. 5672* (MO), 8 m × 11 cm.

[1051] *Henriettea ramiflora* (Sw.) DC.

Prodr. [A. P. de Candolle] 3: 178 [mid Mar. 1828] (Candolle 1828). — *Melastoma ramiflorum* Sw., *Prodr. [Swartz]:* 69 [20 Jun.-29 July 1788] (Swartz 1788), "ramiflora". — *Henriettella ramiflora* (Sw.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 18:* 106 [Aug. 1852] (Naudin 1852).

Henriettea surinamensis Miq., *Stirp. Surinam. Select.:* 45 ["1850" publ. Mar. 1851] (Miquel 1851).

Henriettea trinervia Naudin, *Ann. Sci. Nat., Bot. sér. 3, 18:* 105 [Aug. 1852] (Naudin 1852).

Henriettea succosa var. *guianensis* Gleason, *Bull. Misc. Inform. Kew 1939 (10):* 551 ["1939" publ. 6 Jan. 1940] (Gleason 1940).

VERNACULAR NAMES. — Pa: arakeu, arakeu-seine, arateu, arateu-seine • Ka: akuli wesepotapili, kapuwa nana.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier 1249*.

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15$ cm.

[1052] *Henriettea stellaris* O.Berg ex Triana

Trans. Linn. Soc. London 28 (1): 145 ["1873" publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

VERNACULAR NAMES. — Wn: sihkëimë • Nt: mabee sii.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *H. de Foresta 490*.

SIZE. — Up to 10 m tall (Wurdack *et al.* 1993).

[1053] *Henriettea succosa* (Aubl.) DC.

Prodr. [A. P. de Candolle] 3: 178 [mid Mar. 1828] (Candolle 1828). — *Melastoma succosum* Aubl., *Hist. Pl. Guiane 1:* 418 [Jun.-Dec. 1775] (Aublet 1775), "succosa".

Henriettea brasiliensis Casar., *Nov. Stirp. Bras. 10:* 85 [Sep. 1845] (Casaretto 1845).

VERNACULAR NAMES. — Pa: kudumig-duwē • Ka: pakila yuyulu • Wp: takalawelu • Nt: mabee sii • Cr: kaka-enrièt • Br: muiva.

HERBARIUM DATA (FG). — 83 collections at CAY. Sel. exs.: *J.B. Aublets.n.* (original material BM[BM001008259, BM001008260]).

INVENTORY DATA (FG). — 13 trees in 2 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 16.1$ cm.

Genus *Miconia* Ruiz & Pav.[1054] *Miconia acuminata* (Steud.) Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 244 (Naudin 1851). — *Conostegia acuminata* Steud., *Flora 27:* 722 (Steudel 1844).

Miconia axilliflora Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 179 (Naudin 1851). — *Acinodendron axilliflorum* (Naudin) Kuntze, *Revis. Gen. Pl. 2:* 949 [5 Nov. 1891] (Kuntze 1891), "Acinodendrum".

VERNACULAR NAMES. — Nt: musupu • Br: buxixu.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2894*.

INVENTORY DATA (FG). — 62 trees in 23 plots; $F_{\max} = 12\%$; $dbh_{\text{inv}} = 25.5$ cm.

[1055] *Miconia affinis* DC.

Prodr. [A. P. de Candolle] 3: 187 [mid Mar. 1828] (Candolle 1828). — *Acinodendron affine* (DC.) Kuntze, *Revis. Gen. Pl. 2:* 950 [5 Nov. 1891] (Kuntze 1891), "Acinodendrum".




FIG. 36. — Melastomataceae: **A**, *Bellucia arborescens* (Aubl.) Baill. (*D. Sabatier 1424*); **B**, *Miconia holosericea* (L.) DC. (*M.-F. Prévost & D. Sabatier 4928*); **C**, *Miconia minutiflora* (Bonpl.) DC. (*D. Sabatier et al. 4807*); **D**, *Miconia mirabilis* (Aubl.) L.O.Williams (*D. Sabatier & M.-F. Prévost 3788*). © D. Sabatier/IRD.

Miconia microcarpa DC., *Prod. [A. P. de Candolle] 3*: 189 [mid Mar. 1828] (Candolle 1828). — *Acinodendron microcarpum* (DC.) Kuntze, *Revis. Gen. Pl. 2*: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia cecidophora Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16*: 166 (Naudin 1851). — *Acinodendron cecidophorum* (Naudin) Kuntze, *Revis. Gen. Pl. 2*: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia planinervia Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16*: 160 (Naudin 1851). — *Acinodendron planinerve* (Naudin) Kuntze, *Revis. Gen. Pl. 2*: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia pusilliflora Beurl., *Kongl. Vetensk. Akad. Handl. 40*: 130 [“1854” publ. 1856] (Beurling 1856), *nom. illeg. hom., non* (DC.) Naudin (1850).

Miconia beurlingii Triana, *Trans. Linn. Soc. London 28 (1)*: 107 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Acinodendron beurlingii* (Triana) Kuntze, *Revis. Gen. Pl. 2*: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia cayumbensis Gleason, *Bull. Torrey Bot. Club 66 (6)*: 416 [June 1939] (Gleason 1939).

VERNACULAR NAMES. — Pa: arakeu, arateu • Wp: siki lá • Br: casquinho.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holo-, G[G00191194]; iso-, B[not seen, photo F neg. N° 17002], G-DC[G00201734], K[K000536012]).

INVENTORY DATA (FG). — 21 trees in 9 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 22\text{ cm}$.

[1056] *Miconia amacurensis* Wurdack

Acta Bot. Venez. 2: 373 (Wurdack 1967).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P.A. Sagot 987*, Nov. 1856 (P[P05206288]).

SIZE. — Up to 16 m tall (Wurdack *et al.* 1993).

[1057] *Miconia ampla* Triana

Trans. Linn. Soc. London 28 (1): 101 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Acinodendron amplum* (Triana) Kuntze, *Revis. Gen. Pl. 2*: 950 [5 Nov. 1891] (Kuntze 1891).

Miconia involucrata Donn.Sm., *Bot. Gaz. 37 (3)*: 209 [18 Mar. 1904] (Donnell Smith 1904).

Miconia megaphylla Gleason, *Bull. Torrey Bot. Club 59 (6)*: 363 [June 1932] (Gleason 1932).

VERNACULAR NAMES. — Wp: taīwī lenipiā, takalawelu sōwī.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-763*.

SIZE. — Venezuela, Bolívar. *R.L. Liesner & B.K. Holst 18851* (MO), 20 m.

[1058] *Miconia argyrophylla* DC.

Prodr. [A. P. de Candolle] 3: 181 [mid Mar. 1828] (Candolle 1828). — *Melastoma argyrophyllum* Schrank & Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 181 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.* — *Acinodendron argyrophyllum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia longistyla Steud., *Flora* 27: 724 (Steudel 1844).

Miconia argyrophylla var. *attenuata* Cogn., *Fl. Bras. [Martius]* 14 (4): 296 [1 Nov. 1887] (Cogniaux 1887).

Miconia argyrophylla subsp. *gracilis* Wurdack, *Mem. New York Bot. Gard.* 10 (4): 37 (Wurdack 1961).

HERBARIUM DATA (FG). — 59 collections at CAY. Sel. exs.: *M.-F. Prévost* 3127.

INVENTORY DATA (FG). — 1 tree, dbh = 16.6 cm.

[1059] *Miconia chrysophylla* (Rich.) Urb.

Symb. Antill. [Urban] 4 (3): 459 [15 May 1910] (Urban 1910). — *Melastoma chrysophyllum* Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792), “*Chrysophylla*”. — *Acinodendron chrysophylla* (Rich.) Kuntze, *Revis. Gen. Pl.* 2: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”. — *Melastoma fulvum* Rich., *Monogr. Melast.* 1 (5): 23 [July 1807] (Richard 1807), “*fulva*”, *nom. illeg. superfl.* (based on *Melastoma chrysophyllum*).

Miconia fulva DC., *Prodr. [A. P. de Candolle]* 3: 180 [mid Mar. 1828] (Candolle 1828).

Miconia fulva var. *tinctoria* DC., *Prodr. [A. P. de Candolle]* 3: 180 [mid Mar. 1828] (Candolle 1828).

Miconia longifolia (Aubl.) DC. var. *aubletiana* Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 157 (Naudin 1851). — *Miconia fulva* var. *aubletiana* (Naudin) Cogn., *Fl. Bras. [Martius]* 14 (4): 389 [1 Nov. 1887] (Cogniaux 1887).

Miconia fulva var. *angustifolia* Cogn., *Fl. Bras. [Martius]* 14 (4): 389 [1 Nov. 1887] (Cogniaux 1887).

Miconia fulva var. *poeppigii* Cogn., *Fl. Bras. [Martius]* 14 (4): 389 [1 Nov. 1887] (Cogniaux 1887).

NOTE. — *Miconia fulva* is based on the illegitimate *Melastoma fulvum*, thus not “(Rich.) DC.”

VERNACULAR NAMES. — Wp: siki • Nt: nyama suwi.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *J.B. Leblond* 100 (type G[G00353804]).

INVENTORY DATA (FG). — 3 trees in 2 plots; dbh_{inv} = 15.8 cm.

[1060] *Miconia dispar* Benth.

Hooker's J. Bot. Kew Gard. Misc. 2: 241 (Bentham 1850). — *Acinodendron dispar* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 2673.

INVENTORY DATA (FG). — 1 tree, dbh = 11.4 cm.

[1061] *Miconia egensis* Cogn.

Fl. Bras. [Martius] 14 (4): 374 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron egense* (Cogn.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia panicularis Gleason, *Bull. Torrey Bot. Club* 52 (7): 384 [June 1925] (Gleason 1925).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 1908.

SIZE. — Up to 10 m tall (Berry *et al.* 1999).

[1062] *Miconia elata* (Sw.) DC.

Prodr. [A. P. de Candolle] 3: 182 [mid Mar. 1828] (Candolle 1828). — *Melastoma elatum* Sw., *Prodr. [Swartz]*: 70 [20 Jun.-29 July 1788] (Swartz 1788), “*elata*”. — *Acinodendron elatum* (Sw.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia punctata var. *latifolia* Cogn., *Monogr. Phan. [A.DC. & C.DC.]* 7: 793 [June 1891] (Cogniaux 1891), *syn. nov.* — *Eurychaenia punctata* var. *latifolia* Griseb., *Pl. Wright. [Grisebach]* 1: 185 [Dec. 1860] (Grisebach 1860), *nom. nud. pro syn.*

Miconia eurychaenoides Griseb., *Pl. Wright. [Grisebach]* 1: 185 [Dec. 1860] (Grisebach 1860).

NOTES. — Although *Miconia punctata* var. *latifolia* Cogn. is based on *Eurychaenia punctata* var. *latifolia* Griseb., the latter cannot be treated as the basionym because it is an illegitimate superfluous name. Goldenberg *et al.* (2013) cited an incorrect reference for it (“*Fl. Brit. W.I. [Grisebach]* 259” (Grisebach 1860); no such name therein), and placed it in synonymy with *M. punctata* (Desr.) D.Don ex DC.

VERNACULAR NAMES. — Wp: takalawelu.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 3060.

INVENTORY DATA (FG). — 18 trees in 8 plots; F_{max} = 1.1 %; dbh_{inv} = 34.7 cm.

[1063] *Miconia eriodonta* DC.

Prodr. [A. P. de Candolle] 3: 185 [mid Mar. 1828] (Candolle 1828). — *Acinodendron eriodontum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia eriodonta var. *oblongifolia* DC., *Prodr. [A. P. de Candolle]* 3: 185 [mid Mar. 1828] (Candolle 1828).

Miconia bracteolaris Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 150 (Naudin 1851).

Miconia bracteolaris var. *cardiophora* Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 151 (Naudin 1851).

VERNACULAR NAMES. — Wp: siki sī.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *P.A. Poiteau* s.n. (holo-, G[G00300234]; iso-, G-DC[G00310952]).

INVENTORY DATA (FG). — 1 tree, dbh = 14.5 cm.

[1064] *Miconia fragilis* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 171 (Naudin 1851).

Acinodendron naudinii Kuntze, *Revis. Gen. Pl. 2:* 949 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: á-seiminio-priyo, á-seiminio-priyu, kamata-kamwi • Ka: manyapo.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, P[P01903903]; iso-, B[not seen, photo F neg. N° 17062], BR[BR0000005207749], F[V0063766F], FI[FI004734], P[P01903904, P01903905], US[00120979]).

INVENTORY DATA (FG). — 21 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 60 \text{ cm}$.

[1065] *Miconia gratissima* Benth. ex Triana

Trans. Linn. Soc. London 28 (1): 101 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Acinodendron gratissimum* (Benth. ex Triana) Kuntze, *Revis. Gen. Pl. 2:* 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville B-4760*.

SIZE. — Up to 10 m tall (Wurdack *et al.* 1993).

[1066] *Miconia holosericea* (L.) DC.
(Fig. 36B)

Prodr. [A. P. de Candolle] 3: 181 [mid Mar. 1828] (Candolle 1828). — *Melastoma holosericum* L., *Sp. Pl. 1:* 390 [1 May 1753] (Linnaeus 1753), “*holosericea*”. — *Pleroma holosericea* (L.) D. Don, *Mem. Wern. Nat. Hist. Soc. 4:* 295 [May 1823] (Don 1823). — *Acinodendron holosericeum* (L.) Kuntze, *Revis. Gen. Pl. 1:* 244 [5 Nov. 1891] (Kuntze 1891).

Melastoma mucronatum Desr., *Encycl. [J. Lamarck et al.] 4 (1):* 46 [9 Feb. 1797] (Desrousseaux 1797), “*mucronata*”. — *Diplochita mucronata* (Desr.) DC., *Prodr. [A. P. de Candolle] 3:* 177 [mid Mar. 1828] (Candolle 1828). — *Miconia mucronata* (Desr.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 120 (Naudin 1851). — *Miconia holosericea* var. *mucronata* (Desr.) Cogn., *Fl. Bras. [Martius] 14 (4):* 237 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron mucronatum* (Desr.) Kuntze, *Revis. Gen. Pl. 1:* 244 [5 Nov. 1891] (Kuntze 1891).

Diplochita bracteata DC., *Prodr. [A. P. de Candolle] 3:* 176 [mid Mar. 1828] (Candolle 1828). — *Melastoma bracteatum* Mart. & Schrank ex DC., *Prodr. [A. P. de Candolle] 3:* 176 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Miconia holosericea* var. *bracteata* (DC.) Cogn., *Fl. Bras. [Martius] 14 (4):* 237 [1 Nov. 1887] (Cogniaux 1887).

Miconia holosericea var. *subquintuplinervia* Cogn., *Fl. Bras. [Martius] 14 (4):* 237 [1 Nov. 1887] (Cogniaux 1887).

VERNACULAR NAMES. — Ka: manyapo weli.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4819*.

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.5 \text{ cm}$.

[1067] *Miconia hypoleuca* (Benth.) Triana

Trans. Linn. Soc. London 28 (1): 119 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Chaenopleura hypoleuca* Benth., *J. Bot. [Hooker] 2:* 315 (Bentham 1840). — *Acinodendron hypoleucum* (Benth.) Kuntze, *Revis. Gen. Pl. 2:* 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Chaenopleura ferruginea Crueg., *Linnaea 20:* 112 [17-18 May 1847] (Crueger 1847), “*Chaenophora*”.

Miconia blanchetiana Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 202 (Naudin 1851).

Miconia cruegeriana Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 245 (Naudin 1851), “*Crugeriana*”.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-P. Lescure 943*.

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23.2 \text{ cm}$.

[1068] *Miconia kappleri* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 162 (Naudin 1851), “*Kapplerii*”. — *Acinodendron kappleri* (Naudin) Kuntze, *Revis. Gen. Pl. 2:* 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

NOTE. — Auguste Kappler is known as a collector of plants from Suriname, not from French Guiana. Yet, as the original label of the type indicates (“*Ad fl. Carouany*”), he collected it near the Acarouany, a small French Guianan river located a few kilometers from the border with Suriname.

VERNACULAR NAMES. — Wp: amutu’i, mutu’i, wamutu’i.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *A. Kappler 1675* (holo-, P[P02442360]; iso-, GOET[GOET008001], P[P02442361, P02442362]).

SIZE. — Up to 10 m tall (Wurdack *et al.* 1993).

[1069] *Miconia lepidota* DC.

Prodr. [A. P. de Candolle] 3: 180 [mid Mar. 1828] (Candolle 1828). — *Melastoma lepidotum* Schrank & Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 180 [mid Mar. 1828] (Candolle 1828), “*lepidota*”, nom. nud. pro syn. — *Acinodendron lepidotum* (DC.) Kuntze, *Revis. Gen. Pl. 2:* 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia schomburgkii Benth., *J. Bot. [Hooker] 2:* 312 (Bentham 1840).

Miconia bifrons Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 178 (Naudin 1851).

Miconia naudiniana Miq., *Stirp. Surinam. Select.:* 50 [“1850” publ. Mar. 1851] (Miquel 1851).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 3574*.

INVENTORY DATA (FG). — 4 trees in 2 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.9 \text{ cm}$.

[1070] *Miconia longifolia* (Aubl.) DC.

Prodr. [A. P. de Candolle] 3: 184 [mid Mar. 1828] (Candolle 1828). — *Melastoma longifolium* Aubl., *Hist. Pl. Guiane* 1: 432 [Jun.-Dec. 1775] (Aublet 1775), “*longifolia*”. — *Chaenopleura longifolia* (Aubl.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 260 [late 1860] (Grisebach 1860). — *Acinodendron longifolium* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia lambertiana DC., *Prodr. [A. P. de Candolle]* 3: 185 [mid Mar. 1828] (Candolle 1828).

VERNACULAR NAMES. — Pa: kamata, kamata-kamwi • Wp: amutu'i, mutu'i, siki sī, takalawelu lá, wamutu'i • Wn: kuni weju, mamhalju • Nt: musupu, weti musupu • Br: jacaterão, tangaraca.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN, **LINN-HS 782.16**).

SIZE. — Up to 10 m tall (Wurdack *et al.* 1993).

[1071] *Miconia longispicata* Triana

Trans. Linn. Soc. London 28 (1): 117 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872). — *Acinodendron longispicatum* (Triana) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia longipedunculata Cogn., *Fl. Bras. [Martius]* 14 (4): 266 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron longipedunculatum* (Cogn.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia longispicata var. *minor* Cogn., *Fl. Bras. [Martius]* 14 (4): 267 [1 Nov. 1887] (Cogniaux 1887).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *J.-J. de Granville B-4482*.

SIZE. — Brazil, Pará. *G. Martinelli 7294* (MO), 17 m.

[1072] *Miconia matthaei* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 176 (Naudin 1851). — *Acinodendron matthaei* (Naudin) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia matthaei var. *undulata* Cogn., *Monogr. Phan. [A.D.C. & C.D.C.J 7]*: 813 [June 1891] (Cogniaux 1891), “*Matthaei*”.

Miconia wilsonii Cogn., *Symb. Antill. [Urban]* 7 (3): 311 [1 Oct. 1912] (Cogniaux 1912).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3579*.

SIZE. — Bolivia, Pando. *C.R. Sperling 6591* (MO), 7 m × 10 cm.

[1073] *Miconia melinonii* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 125 (Naudin 1851), “*Melinonis*”. — *Acinodendron melinonii* (Naudin) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891).

NOTE. — The epithet “*melinonii*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland *et al.* 2018: Art. 60.8).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *E.M. Mélinon 178*, 1842 (holo-, P[P02442442]; iso-, NY[00229251], P[P02442443, P02442444], U[U0004102], US[00121076]).

INVENTORY DATA (FG). — 1 tree, dbh = 24.2 cm.

[1074] *Miconia minutiflora* (Bonpl.) DC.
(Fig. 36C)

Prodr. [A. P. de Candolle] 3: 189 [mid Mar. 1828] (Candolle 1828). — *Melastoma minutiflorum* Bonpl., *Monogr. Melast.* 1 (9): 50 [Aug. 1809] (Bonpland 1809), “*minutiflora*”. — *Acinodendron minutiflorum* (Bonpl.) Kuntze, *Revis. Gen. Pl.* 1: 245 [5 Nov. 1891] (Kuntze 1891).

Cremanium trinitatis Crueg., *Linnaea* 20: 111 [17-18 May 1847] (Crueger 1847). — *Miconia trinitatis* (Crueg.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 246 (Naudin 1851).

Glossocentrum collinum Crueg., *Linnaea* 20: 111 [17-18 May 1847] (Crueger 1847).

Miconia melanodendron Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 162 (Naudin 1851).

Miconia glossocentra Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 243 (Naudin 1851).

Melastoma pendulum Salzm. ex Griseb., *Fl. Brit. W.I. [Grisebach]* 257 [late 1860] (Grisebach 1860), *nom. nud. pro syn.*

Cremanium paniculatum Mart. ex Triana, *Trans. Linn. Soc. London* 28 (1): 118 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872), *nom. nud. pro syn.*

Miconia minutiflora var. *latifolia* Cogn., *Fl. Bras. [Martius]* 14 (4): 385 [1 Nov. 1887] (Cogniaux 1887).

Miconia borealis Gleason, *Bull. Torrey Bot. Club* 55 (2): 118 [Feb. 1928] (Gleason 1928).

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *M.-F. Prévost 1858*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} = 1.8\%$; $dbh_{\text{inv}} = 30.6$ cm.

[1075] *Miconia mirabilis* (Aubl.) L.O.Williams
(Fig. 36D)

Fieldiana, Bot. 29 (10): 574 (Williams 1963). — *Fothergilla mirabilis* Aubl., *Hist. Pl. Guiane* 1: 441 [Jun.-Dec. 1775] (Aublet 1775), “*admirabilis*” in Index, “*Mirabilia*” on plate. — *Lieautautia mirabilis* (Aubl.) Buc'hoz, *Pl. Nouv. Découv.*: 9 (Buc'hoz 1779). — *Melastoma fothergilla* Desr., *Encycl. J. Lamarck et al.* 4 (1): 45 [9 Feb. 1797] (Desrousseaux 1797), *nom. illeg. superfl.* (based on *Fothergilla mirabilis*). — *Chitonia fothergilla* D.Don, *Mem. Wern. Nat. Hist. Soc.* 4: 318 [May 1823] (Don 1823). — *Diplochitona fothergilla* DC., *Prodr. [A. P. de Candolle]* 3: 176 [mid Mar. 1828] (Candolle 1828). — *Abrophaes mirabilis* (Aubl.) Raf., *Sylva Tellur.*: 99 (Rafinesque 1838). — *Miconia fothergilla* Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 119 (Naudin 1851). — *Tamonea fothergilla*

O.F.Cook & G.N.Collins, *Contr. U.S. Natl. Herb.* 8 (2): 249 [27 June 1903] (Cook & Collins 1903).

Tamonea guianensis Aubl., *Hist. Pl. Guiane* 1: 441 [Jun.-Dec. 1775] (Aublet 1775). — *Melastoma tamonea* Sw., *Prod. [Swartz]*: 70 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. superfl. (based on *Tamonea guianensis*). — *Miconia guianensis* (Aubl.) Cogn., *Jahrb. Königl. Bot. Gart. Berlin* 4: 280 (Cogniaux 1886). — *Acinodendron guianense* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 244 [5 Nov. 1891] (Kuntze 1891).

Melastoma compressum Vahl, *Ectog. Amer.* 3: 23 (Vahl 1807), “*compressa*”.

Melastoma fothergilla var. *lanceolata* Bonpl., *Monogr. Melast.* 1: 71 (Bonpland 1809), nom. illeg. superfl. (based on *Fothergilla mirabilis*).

Melastoma fothergilla var. *ovalis* Bonpl., *Monogr. Melast.* 1: 71 (Bonpland 1809), nom. illeg. superfl. (based on *Melastoma dodecandrum* Desr. and *M. compressum*). — *Miconia guianensis* var. *ovalis* Cogn., *Fl. Bras. [Martius]* 14 (4): 246 [1 Nov. 1887] (Cogniaux 1887).

Diplochita florida DC., *Prod. [A. P. de Candolle]* 3: 176 [mid Mar. 1828] (Candolle 1828). — *Miconia florida* (DC.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 244 (Naudin 1851). — *Acinodendron floridum* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891).

Diplochita rosea Macfad., *Fl. Jamaica [Macfadyen]* 2: 74 (Macfadyen 1850).

Miconia guianensis var. *vulgaris* Cogn., *Fl. Bras. [Martius]* 14 (4): 246 [1 Nov. 1887] (Cogniaux 1887).

NOTES. — According to Williams (1963: 574), while his *Histoire des Plantes* was in press, Aublet realized that he had created two *Tamonea*, one in Verbenaceae (Aublet 1775, 2: 659), the other in Melastomataceae (Aublet 1775, 1: 441). His solution was to rename the latter *Fothergilla*, and he managed to have *Tamonea guianensis* corrected to *Fothergilla mirabilis* in most, but not all, copies. Goldenberg et al. (2013: 78) wrongly listed a “*F. guianensis* Aubl.” with the same reference as *F. mirabilis*. The specific epithet of *Melastoma fothergilla* Desr. refers to the genus *Fothergilla*, not to John Fothergill (1712–1780) to whom the genus was dedicated. Therefore it should not be written as “*fothergillii*”.

VERNACULAR NAMES. — Ka: kuyaken elepali • Wp: siki lá.

HERBARIUM DATA (FG). — 82 collections at CAY. Sel. exs.: *J.B. Aublets.n.* (original material of *Fothergilla mirabilis*: P[P02442466]).

INVENTORY DATA (FG). — 40 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{inv} = 27.1$ cm.

[1076] *Miconia myriantha* Benth.

J. Bot. [Hooker] 2: 314 (Bentham 1840).

VERNACULAR NAMES. — Ka: kumete, kumeti.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *G. Cremer & M. Hoff* 11240.

INVENTORY DATA (FG). — 1 tree, $dbh = 11.1$ cm.

[1077] *Miconia phaeophylla* Triana

Trans. Linn. Soc. London 28 (1): 113 [“1873” publ. 8 Dec. 1871–13 Jan. 1872] (Triana 1871–1872). — *Acinodendron phaeophyllum* (Triana) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891).

Miconia fendlariana Cogn., *Monogr. Phan. [A.DC. & C.DC.]* 7: 822 [June 1891] (Cogniaux 1891). — *Acinodendron fendlarianum* (Cogn.) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum Fendleranum*”.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2112*.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 25.8$ cm.

[1078] *Miconia pluknetii* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 116 (Naudin 1851). — *Acinodendron pluknetii* (Naudin) Kuntze, *Revis. Gen. Pl.* 2: 951 [5 Nov. 1891] (Kuntze 1891).

Diplochita sessilifolia Griseb., *Fl. Brit. W.I. [Grisebach]* 252 [late 1860] (Grisebach 1860).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: maipuli kiyelapoli, sipyalala • Wp: takalawelu ki, takalawelu lá.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2815.

INVENTORY DATA (FG). — 8 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 16$ cm.

[1079] *Miconia poeppigii* Triana

Trans. Linn. Soc. London 28 (1): 107 [“1873” publ. 8 Dec. 1871–13 Jan. 1872] (Triana 1871–1872). — *Acinodendron poeppigii* (Triana) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Oxymeris furfuracea Poepp. ex Triana, *Trans. Linn. Soc. London* 28 (1): 107 [“1873” publ. 8 Dec. 1871–13 Jan. 1872] (Triana 1871–1872), nom. nud. pro syn.

Miconia congesta Cogn., *Repert. Spec. Nov. Regni Veg.* 8: 2 (Cogniaux 1910).

Miconia darienensis Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 248 [22 Sep. 1917] (Pittier 1917).

Miconia surinamensis Gleason, *Recueil Trav. Bot. Néerl.* 32: 212 (Gleason 1935).

VERNACULAR NAMES. — Pa: kamata • Ka: kamala kuyulu • Wp: amutu'i, mutu'i, wamutu'i • Nt: man musupu witi • Br: catuaba.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 1005.

INVENTORY DATA (FG). — 9 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 42.2$ cm.

[1080] *Miconia prasina* (Sw.) DC.

Prodr. [A. P. de Candolle] 3: 188 [mid Mar. 1828] (Candolle 1828). — *Melastoma prasinum* Sw., *Prodr. [Swartz]*: 69 [20 Jun.-29 July 1788] (Swartz 1788), “*prasina*”. — *Miconia aubletiana* Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 139 (Steudel 1841), *nom. illeg. superfl.* (based on *Miconia prasina*). — *Acinodendron prasinum* (Sw.) Kuntze, *Revis. Gen. Pl.* 1: 245 [5 Nov. 1891] (Kuntze 1891).

Melastoma parviflorum Aubl., *Hist. Pl. Guiane* 1: 433 [Jun.-Dec. 1775] (Aublet 1775), “*parviflora*”.

Melastoma pendulifolium Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792), “*Pendulifolia*”.

Melastoma montanum Spreng., *Syst. Veg. [Sprengel]* 2: 296 [Jan.-May 1825] (Sprengel 1825), *nom. illeg. hom.*, *non* Sw. (Swartz 1788).

Conostegia parviflora DC., *Prodr. [A. P. de Candolle]* 3: 175 [mid Mar. 1828] (Candolle 1828).

Miconia collina DC., *Prodr. [A. P. de Candolle]* 3: 185 [mid Mar. 1828] (Candolle 1828). — *Miconia prasina* var. *collina* (DC.) Triana, *Trans. Linn. Soc. London* 28 (1): 109 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

Miconia sepiaria DC., *Prodr. [A. P. de Candolle]* 3: 185 [mid Mar. 1828] (Candolle 1828). — *Melastoma sepiarium* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 185 [mid Mar. 1828] (Candolle 1828), “*sepiaria*”, *nom. nud. pro syn.*

Miconia attenuata DC., *Prodr. [A. P. de Candolle]* 3: 186 [mid Mar. 1828] (Candolle 1828). — *Miconia prasina* var. *attenuata* (DC.) Cogn., *Fl. Bras. [Martius]* 14 (4): 317 [1 Nov. 1887] (Cogniaux 1887).

Miconia attenuata var. *subquintuplinervia* DC., *Prodr. [A. P. de Candolle]* 3: 187 [mid Mar. 1828] (Candolle 1828).

Melastoma suaveolens Raddi, *Melast. Bras.*: 25 (Raddi 1828).

Melastoma pulchrum Vell., *Fl. Flumin.* 178 [“1825” publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), “*pulchra*”.

Miconia revoluta Benth., *J. Bot. [Hooker]* 2: 313 (Bentham 1840).

Miconia pteropoda Benth., *J. Bot. [Hooker]* 2: 314 (Bentham 1840). — *Acinodendron pteropodium* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia fleischeriana Steud., *Flora* 27: 723 (Steudel 1844).

Miconia macrophylla Steud., *Flora* 27: 723 (Steudel 1844).

Miconia repandocrenata Steud., *Flora* 27: 724 (Steudel 1844).

Miconia palustris Macfad., *Fl. Jamaica [Macfadyen]* 2: 89 (Macfadyen 1850).

Miconia crispula Spruce ex Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 172 (Naudin 1851), *nom. nud. pro syn.*

Miconia cristulata Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 177 (Naudin 1851).

Miconia nemoralis Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 178 (Naudin 1851).

Miconia prasina var. *denticulata* Griseb., *Bonplandia* 6 (1): 6 (Grisebach 1858).

Melastoma quinquenervium Salzm. ex Griseb., *Fl. Brit. W.I. [Grisebach]* 257 [late 1860] (Grisebach 1860), *nom. illeg. hom.*, *non* Mill. (Miller 1768).

Melastoma trinervium Salzm. ex Griseb., *Fl. Brit. W.I. [Grisebach]* 257 [late 1860] (Grisebach 1860), *nom. illeg. hom.*, *non* Sw. (Swartz 1788).

Miconia trichotoma Bello, *Anales Soc. Esp. Hist. Nat.* 10: 268 (Bello 1883), *nom. illeg. hom.*, *non* (Desr.) DC. (Candolle 1828).

Miconia prasina var. *angustifolia* Cogn., *Fl. Bras. [Martius]* 14 (4): 317 [1 Nov. 1887] (Cogniaux 1887).

Miconia prasina var. *crispula* Cogn., *Fl. Bras. [Martius]* 14 (4): 317 [1 Nov. 1887] (Cogniaux 1887).

Melastoma acuminatum Sessé & Moc., *Fl. Mexic.*, ed. 2, 105 (Sessé & Mociño 1894), “*Acuminata*”, *nom. illeg. hom.*, *non* Desr. (Desrousseaux 1797).

Melastoma leucanthum Sessé & Moc., *Fl. Mexic.*, ed. 2, 105 (Sessé & Mociño 1894), “*Leucantha*”.

Melastoma glabrum Sessé & Moc., *Fl. Mexic.*, ed. 2, 107 (Sessé & Mociño 1894), “*Glabra*”, *nom. illeg. hom.*, *non* G. Forst. (Forster 1786).

Miconia mucronulata Ule, *Notizbl. Königl. Bot. Gart. Berlin* 6: 358 [20 Sep. 1915] (Ule 1915).

NOTE. — Goldenberg *et al.* (2013) listed “*Miconia parviflora* (Aubl.) Cogn., *Monogr. Phan. [A.D.C. & C.D.C.]* 7: 805. 1891”; no such name is found therein. Cogniaux just cited *Melastoma parviflorum* Aubl. in synonymy under *Miconia prasina* (Sw.) DC.

VERNACULAR NAMES. — Pa: arakeu, arakeu-kamwi, arakeu-priye, arateu, arateu-kamwi, arateu-priye • Ka: kinolo epi, yalipi • Wp: amutu'i, mutu'i, takalawelu lá, takalawelu sili, wamutu'i • Nt: nyama suwi • Br: caiuia, fruto-de-jacu.

HERBARIUM DATA (FG). — 116 collections at CAY. Sel. exs.: *M.-F. Prévost 1206*.

INVENTORY DATA (FG). — 23 trees in 4 plots; $F_{\max} = 2.9 \%$; $dbh_{inv} = 21.9$ cm.

[1081] *Miconia pubipetala* Miq.

Stirp. Surinam. Select.: 50 [“1850” publ. Mar. 1851] (Miquel 1851).

Diplochita parviflora Benth., *J. Bot. [Hooker]* 2: 302 (Bentham 1840). — *Miconia parviflora* (Benth.) Cogn., *Fl. Bras. [Martius]* 14 (4): 249 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron parviflorum* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia verticilliflora Steud. ex Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 122 (Naudin 1851).

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *B. Riéra 715*.

SIZE. — Up to 10 m tall (Wurdack *et al.* 1993).

[1082] *Miconia punctata* (Desr.) D.Don ex DC.

Prodr. [A. P. de Candolle] 3: 184 [mid Mar. 1828] (Candolle 1828). — *Melastoma punctatum* Desr., *Encycl. J. Lamarck et al.* 4 (1): 50 [9 Feb. 1797] (Desrousseaux 1797). — *Eurychaenia punctata* (Desr.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 259 [late 1860] (Grisebach 1860). — *Acinodendron punctatum* (Desr.) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia discolor Macfad., *Fl. Jamaica [Macfadyen]* 2: 85 (Macfadyen 1850).

Miconia lepidota var. *grandifolia* Cogn., *Fl. Bras. [Martius]* 14 (4): 293 [1 Nov. 1887] (Cogniaux 1887).

Miconia punctata var. *brevifolia* Cogn., *Monogr. Phan. [A.D.C. & C.D.C.]* 7: 794 [June 1891] (Cogniaux 1891).

Miconia habrolepis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 319 [24 Oct. 1929] (Standley 1929).

NOTE. — Don (1823: 316) did not associate the epithet *punctata* with the genus name *Miconia*, hence did not validly publish the combination (Turland et al. 2018: Art. 35.2).

VERNACULAR NAMES. — Te: mitū pi, takulu welu • Wp: takalawelu • Nt: weti bee.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *M.-F. Prévost 1076*.

INVENTORY DATA (FG). — 20 trees in 7 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 30.9\text{ cm}$.

[1083] *Miconia pyrifolia* Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 164 (Naudin 1851). — *Acinodendron pyrifolium* (Naudin) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Sastre 4516*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 39.7\text{ cm}$.

[1084] *Miconia ruficalyx* Gleason

Brittonia 1: 181 (Gleason 1932).

Copedesma nitens Gleason, *Bull. Torrey Bot. Club* 52 (6): 331 [June 1925] (Gleason 1925).

VERNACULAR NAMES. — Wp: siki lá • Br: maçandarubinha.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1122*.

INVENTORY DATA (FG). — 26 trees in 12 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 33.1\text{ cm}$.

[1085] *Miconia serialis* DC.

Prodr. [A. P. de Candolle] 3: 182 [mid Mar. 1828] (Candolle 1828). — *Acinodendron seriale* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia tomentella Cogn., *Fl. Bras. [Martius]* 14 (4): 284 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron tomentellum* (Cogn.) Kuntze, *Revis. Gen. Pl.* 2: 953 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier 2270*.

SIZE. — Bolivia, Pando. *A. Jardim 1015* (MO), 10 m.

[1086] *Miconia serrulata* (DC.) Naudin

Ann. Sci. Nat., Bot. sér. 3, 16: 118 (Naudin 1851). — *Diplochita serrulata* DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828). — *Melastoma serrulatum* Rich. ex DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.* — *Miconia macrophylla* var. *serrulata* (DC.) Cogn., *Fl. Bras. [Martius]* 14 (4): 241 [1 Nov. 1887] (Cogniaux 1887).

Chitonia macrophylla D.Don, *Mem. Wern. Nat. Hist. Soc.* 4: 319 [May 1823] (Don 1823). — *Melastoma macrophyllum* Pav. ex D.Don, *Mem. Wern. Nat. Hist. Soc.* 4: 319 [May 1823] (Don 1823), *nom. nud. pro syn.* — *Miconia macrophylla* (D.Don) Triana, *Trans. Linn. Soc. London* 28 (1): 103 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872), *nom. illeg. hom., non* Steud. (Steudel 1844). — *Acinodendron macrophyllum* (D.Don) Kuntze, *Revis. Gen. Pl.* 1: 244 [5 Nov. 1891] (Kuntze 1891). — *Diplochita macrophylla* (D.Don) DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828).

Diplochita leucocephala DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828). — *Miconia leucocephala* (DC.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16*: 244 (Naudin 1851). — *Miconia macrophylla* var. *leucocephala* (DC.) Cogn., *Fl. Bras. [Martius]* 14 (4): 241 [1 Nov. 1887] (Cogniaux 1887).

Diplochita serrulata var. *latifolia* DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828). — *Miconia macrophylla* var. *latifolia* (DC.) Cogn., *Fl. Bras. [Martius]* 14 (4): 240 [1 Nov. 1887] (Cogniaux 1887).

Melastoma balbisianum Ser. ex DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Melastoma lasiopetalum Mart. & Schrank ex DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Melastoma patens Spreng. ex DC., *Prodr. [A. P. de Candolle]* 3: 177 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Melastoma strangulatum Raddi, *Melast. Bras.*: 31 (Raddi 1828).

Melastoma macrophyllum Vell., *Fl. Flumin.* 180 [“1825” publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), “macro-phylla”, *nom. illeg. hom., non* Desr. (Desrousseaux 1797).

Decaraphe hostmannii Steud., *Flora* 27: 722 (Steudel 1844), “*Decaraphe Hostmanni*”. — *Miconia macrophylla* var. *hostmannii* (Steud.) Cogn., *Fl. Bras. [Martius]* 14 (4): 241 [1 Nov. 1887] (Cogniaux 1887), “*Hostmanni*”.

Miconia platyhedra Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16*: 244 (Naudin 1851).

Tamonea lasiopetala DC. ex Triana, *Trans. Linn. Soc. London* 28 (1): 103 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872), *nom. nud. pro syn.*

Miconia boissieriana Cogn., *Monogr. Phan. [A.DC. & C.DC.]* 7: 734 [June 1891] (Cogniaux 1891). — *Acinodendron boissieranum* (Cogn.) Kuntze, *Revis. Gen. Pl.* 2: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Tamonea tomentosa Kuntze var. *auriculata* Jenn., *Ann. Carnegie Mus.* 11 (1-2): 208 (Jennings 1917).

VERNACULAR NAMES. — Pa: kudumig • Ka: kulumoto • Wp: takalawelu lá, takalawelu lá sili • Br: buxixú-canela-de-velho.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *M.-F. Prévost 1506*.

SIZE. — Up to 15 m tall (Wurdack *et al.* 1993).

[1087] *Miconia splendens* (Sw.) Griseb.

Fl. Brit. W.I. [Grisebach] 256 [late 1860] (Grisebach 1860). — *Melastoma splendens* Sw., *Prodri. [Swartz]* 70 [20 Jun.-29 July 1788] (Swartz 1788).

Miconia elliptica Macfad., *Fl. Jamaica [Macfadyen]* 2: 88 (Macfadyen 1850).

Miconia obovalis Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 183 (Naudin 1851).

Acinodendron ellipticum (Macfad.) Kuntze, *Revis. Gen. Pl.* 2: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

VERNACULAR NAMES. — Wp: siki lá, taīwī lenipiá lá • Wn: kuni weju.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2335*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.4 cm.

[1088] *Miconia tetraspermoides* Wurdack

Phytologia 18 (3): 155 [27 Mar. 1969] (Wurdack 1969).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5620*.

INVENTORY DATA (FG). — 1 tree, dbh = 15.3 cm.

[1089] *Miconia tomentosa* (Rich.) D.Don ex DC.

Prodri. [A. P. de Candolle] 3: 183 [mid Mar. 1828] (Candolle 1828). — *Melastoma tomentosum* Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792), “*Tomentosa*”. — *Jucunda tomentosa* (Rich.) Benth., *J. Bot. [Hooker]* 2: 302 (Bentham 1840). — *Diplochita tomentosa* (Rich.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 252 [late 1860] (Grisebach 1860). — *Acinodendron tomentosum* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 245 [5 Nov. 1891] (Kuntze 1891).

Melastoma megalophyllum Mart. ex DC., *Prodri. [A. P. de Candolle]* 3: 183 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Pogonorhynchus amplexans Crueg., *Linnaea* 20: 107 [17-18 May 1847] (Crueger 1847). — *Miconia amplexans* (Crueg.) Cogn., *Fl. Bras. [Martius]* 14 (4): 256 [1 Nov. 1887] (Cogniaux 1887). — *Acinodendron amplexans* (Crueg.) Kuntze, *Revis. Gen. Pl.* 2: 950 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Pogonorhynchus sessillifolius Crueg., *Linnaea* 20: 107 [17-18 May 1847] (Crueger 1847).

Miconia umbrifera Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 116 (Naudin 1851).

Miconia tomentosa var. *ovata* Cogn., *Fl. Bras. [Martius]* 14 (4): 255 [1 Nov. 1887] (Cogniaux 1887).

Miconia symplectocaulos Pilg., *Verh. Bot. Vereins Prov. Brandenburg* 47: 175 [1 Oct. 1905] (Pilger 1905).

VERNACULAR NAMES. — Ka: sipyalala • Wp: takalawelu ki.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *J.B. Leblond 64* (original material of *Melastoma tomentosum*: G[G00227574]; *D. Sabatier 3012*, dbh 10 cm.

[1090] *Miconia trinervia* (Sw.) D.Don ex G.Don

Hort. Brit. [Loudon] 174 (Don 1830). — *Melastoma trinervium* Sw., *Prodri. [Swartz]* 69 [20 Jun.-29 July 1788] (Swartz 1788), “*trinervia*”. — *Acinodendron trinervium* (Sw.) Kuntze, *Revis. Gen. Pl.* 2: 953 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Melastoma scorpioides Schlechtld. & Cham., *Linnaea* 5: 564 (Schlechten-dal & Chamisso 1830). — *Miconia scorpioides* (Schlechtld. & Cham.) Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 243 (Naudin 1851). — *Acinodendron scorpioides* (Schlechtld. & Cham.) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Miconia anceps Naudin, *Ann. Sci. Nat., Bot. sér. 3, 16:* 150 (Naudin 1851).

VERNACULAR NAMES. — Wp: takalawelu ka'a lulu, takalawelu lá, takalawelu lá sili, takalawelu sī.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *P. Grenand 738*.

INVENTORY DATA (FG). — 1 tree, dbh = 11.6 cm.

[1091] *Miconia tschudyoides* Cogn.

Fl. Bras. [Martius] 14 (4): 327 [1 Nov. 1887] (Cogniaux 1887).

Tschudyia robusta Sagot, *Ann. Sci. Nat., Bot. sér. 6, 15:* 329 (Sagot 1883). — *Miconia robusta* (Sagot) Cogn., *Fl. Bras. [Martius]* 14 (4): 611 [15 Aug. 1888] (Cogniaux 1888), nom. illeg. hom., non Cogn. (Cogniaux 1887). — *Acinodendron robustum* (Sagot) Kuntze, *Revis. Gen. Pl.* 2: 952 [5 Nov. 1891] (Kuntze 1891), “*Acinodendrum*”.

Clidemia trichodes Sagot ex Cogn., *Fl. Bras. [Martius]* 14 (4): 612 [15 Aug. 1888] (Cogniaux 1888), nom. nud. pro syn.

VERNACULAR NAMES. — Ka: yoloka pomiidy • Nt: musupu.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *P.A. Sagot 220*, 1858 (original material K[K000536081], U[U0004123]).

INVENTORY DATA (FG). — 68 trees in 40 plots; $F_{\max} < 1\%$; dbh_{inv} = 29 cm.

Genus *Mouriri* Aubl.

[1092] *Mouriri acutiflora* Naudin

Ann. Sci. Nat., Bot. sér. 3, 18: 284 [Aug. 1852] (Naudin 1852).

Mouriri acutiflora var. *amblyodon* Naudin, *Ann. Sci. Nat., Bot. sér. 3*, 18: 284 [Aug. 1852] (Naudin 1852).

Mouriri pendulifolia Cogn., *Fl. Bras. [Martius]* 14 (4): 578 [15 Aug. 1888] (Cogniaux 1888).

VERNACULAR NAMES. — Ka: šipa yepo • Cr: bwa-flèch.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *F.M.R. Leprieur, Cat. no. 206* (original material of *Mouriri acutiflora* var. *amblyodon*: BR[BR0000005209880], G-DC[G00328319]).

INVENTORY DATA (FG). — 1 tree, dbh = 10.6 cm.

[1093] *Mouriri angulicosta* Morley

Phytologia 22 (5): 424 [24 Jan. 1972] (Morley 1972).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2233*.

INVENTORY DATA (FG). — 11 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{inv} = 70.3$ cm.

[1094] *Mouriri collocarpa* Ducke

Anais Reunião Sul-Amer. Bot. 3: 69 [“1938” publ. 1940] (Ducke 1940).

Mouriri acutiflora var. *oligantha* Gleason, *Recueil Trav. Bot. Néerl.* 32: 212 (Gleason 1935).

VERNACULAR NAMES. — Wp: wila kitá • Br: mamãozinho.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier 2294*.

INVENTORY DATA (FG). — 15 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{inv} = 46.8$ cm.

[1095] *Mouriri crassifolia* Sagot
(Fig. 37A)

Ann. Sci. Nat., Bot. sér. 6, 15: 336 (Sagot 1883).

Mouriri anomala Pulle, *Ann. Jard. Bot. Buitenzorg, Suppl.* 3: 123 (Pulle 1910).

VERNACULAR NAMES. — Pa: avitkat-seinō, avitkat-wahuye • Ka: amila’u, amula’i, amulau, atakalilan • Wp: wila kitá • Wn: taluwaho • Nt: topi • Cr: bwa-fé, bwa-flèch • Fr: bois flèche, mouriri • Br: camutim, cruili, miraúba, muiraúba, muriri.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (original material BR[BR0000005208524], F[V0093860F], K[K000329677], NY[00229544], P[P05254750]).

INVENTORY DATA (FG). — 328 trees in 104 plots; $F_{\max} = 2.2\%$; $dbh_{inv} = 62.5$ cm.

[1096] *Mouriri duckeana* Morley

Amer. J. Bot. 40: 253 (Morley 1953).

VERNACULAR NAMES. — Br: muiraúba, pitanga-da-mata.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 23427*.

SIZE. — Up to 35 cm dbh (Morley 1976).

[1097] *Mouriri dumetosa* Cogn.

Fl. Bras. [Martius] 14 (4): 584 [15 Aug. 1888] (Cogniaux 1888).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (original material BR[BR0000005221417]).

INVENTORY DATA (FG). — 1 tree, dbh = 18.5 cm.

[1098] *Mouriri francavillana* Cogn.

Fl. Bras. [Martius] 14 (4): 576 [15 Aug. 1888] (Cogniaux 1888).

VERNACULAR NAMES. — Pa: ahayupná, avitkat, avitkat-purubumna • Cr: bwa-flèch • Br: muiraúba, muriri.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (original material BR[BR0000005221189]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.4$ cm.

[1099] *Mouriri grandiflora* DC.

Prodr. [A. P. de Candolle] 3: 8 [mid Mar. 1828] (Candolle 1828).

Mouriri macrophylla Willd. ex Cham. & Schldl., *Linnæa* 10: 218 (Chamisso & Schlechtendal 1835).

Mouriri princeps Naudin, *Ann. Sci. Nat., Bot. sér. 3*, 18: 283 [Aug. 1852] (Naudin 1852).

VERNACULAR NAMES. — Pa: avitkat-purubumna • Ka: amala’i, woko popi • Te: aliwatso tawa • Wp: aliwa so • Wn: akajot, akojot, muku huhu, taluwaho • Nt: lebi moni, sii payopo, topi • Cr: al-waso, grènn-koumarou • Br: camutim, muriri, remelo-de-cachorro, tucunaré-merecá.

HERBARIUM DATA (FG). — 51 collections at CAY. Sel. exs.: *M.-F. Prévost 3329*.

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 22$ cm.

[1100] *Mouriri guianensis* Aubl.
(Fig. 37B)

Hist. Pl. Guiane 1: 452 [Jun.-Dec. 1775] (Aublet 1775).

Petaloma mouriri Sw., *Prod. [Swartz]*: 73 [20 Jun.-29 July 1788] (Swartz 1788).

Myrtus umbellata Ham., *Prod. Pl. Ind. Occid. [Hamilton]*: 45 [Oct. 1825] (Hamilton 1825).

Eugenia brachybotrya DC., *Prod. [A. P. de Candolle]* 3: 277 [mid Mar. 1828] (Candolle 1828).

Mouriri polyantha Miq., *Linnaea* 18: 290 [“1844” publ. Feb (-May?) 1845] (Miquel 1845).

Mouriri weddelli Naudin, *Ann. Sci. Nat., Bot. sér. 3, 18*: 286 [Aug. 1852] (Naudin 1852).

Mouriri ulei Pilg., *Verb. Bot. Vereins Prov. Brandenburg* 47: 184 [1 Oct. 1905] (Pilger 1905).

VERNACULAR NAMES. — Br: criuri, cruiria.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[[BM001008165](#)] designated by Morley [1976: 171]).

SIZE. — Up to 50 cm dbh (Morley 1976).

[1101] *Mouriri huberi* Cogn.

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 255 (Cogniaux 1909).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2101*.

INVENTORY DATA (FG). — 74 trees in 34 plots; $F_{\max} = 1.7\%$; $dbh_{inv} = 51.6$ cm.

[1102] *Mouriri nervosa* Pilg.

Verb. Bot. Vereins Prov. Brandenburg 47: 183 [1 Oct. 1905] (Pilger 1905).

VERNACULAR NAMES. — Pa: ahayumna, ahayupná, timuvukti-duwé • Wp: talakua'i, wila kitá • Wn: kaliaku miñí, taluwaho • Br: miraúba.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.-J. de Granville B-4989*.

INVENTORY DATA (FG). — 21 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{inv} = 40$ cm.

[1103] *Mouriri nigra* (DC.) Morley

Phytologia 22 (5): 428 [24 Jan. 1972] (Morley 1972). — *Eugenia nigra* DC., *Prod. [A. P. de Candolle]* 3: 268 [mid Mar. 1828] (Candolle 1828).

Mouriri plasschaertii Pulle, *Recueil Trav. Bot. Néerl.* 6: 283 (Pulle 1909), “*Mouriria*”.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-F. Molino 938*.

SIZE. — Up to 36 m tall (Morley 1976).

[1104] *Mouriri oligantha* Pilg.

Verb. Bot. Vereins Prov. Brandenburg 47: 184 [1 Oct. 1905] (Pilger 1905).

VERNACULAR NAMES. — Pa: avitkat-kamwi, boko • Wp: akusi luway, wila kitá u • Cr: gouyavié-sovaj.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-390*.

INVENTORY DATA (FG). — 6 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 23.6$ cm.

[1105] *Mouriri sagotiana* Triana

Trans. Linn. Soc. London 28 (1): 155 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

VERNACULAR NAMES. — Pa: timuvukti • Ka: pikyalaiypo, talukuwa epi • Te: talakwa'i • Wp: talakua'i, talakua'i to wu • Cr: bwa-flèch, panapaná • Br: miraúba.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *P.A. Sagot 1203*, May 1858 (original material BR[[BR000005751051](#)], F[[V0077508F](#)], K[[K000329647](#)]).

INVENTORY DATA (FG). — 21 trees in 11 plots; $F_{\max} = 1.5\%$; $dbh_{inv} = 20$ cm.

[1106] *Mouriri sideroxylon* Sagot ex Triana

Trans. Linn. Soc. London 28 (1): 155 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P.A. Sagot 924*, 1856 (original material B[not seen, photo F neg. N° 17289], BM[[BM001008164](#)], BR[[BR000005221578](#)], F[[V0077509F](#)], K[[K000329660](#), [K000329661](#)], NY[[00229564](#)], S[[S05-3861](#)], U[[U0004144](#)], US[[00123933](#)]).

INVENTORY DATA (FG). — 1 tree, dbh = 35.8 cm.

[1107] *Mouriri subumbellata* Triana

Trans. Linn. Soc. London 28 (1): 154 [“1873” publ. 8 Dec. 1871-13 Jan. 1872] (Triana 1871-1872).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *H.S. Irwin et al. 48582* (US).

SIZE. — Up to 40 cm dbh (Morley 1976).

[1108] *Mouriri vernicosa* Naudin

Ann. Sci. Nat., Bot. sér. 3, 18: 285 [Aug. 1852] (Naudin 1852).

VERNACULAR NAMES. — Ka: apalitono, malakana kuwa • Wp: aliwa so átā, aliwa so u.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[[BM001008169](#)], COL[[COL000003404](#)], F[[V0044368F](#), [V0044369F](#)], GH[[GH00073017](#)], K[[K000329644](#)], NY[[00229567](#)], S[[S05-3865](#)], US[[00604497](#)]).

SIZE. — Up to 35 cm dbh (Morley 1976).

Genus *Votomita* Aubl.

[1109] *Votomita guianensis* Aubl.

Hist. Pl. Guiane 1: 91 [Jun.-Dec. 1775] (Aublet 1775).

Mouriri abnormis Naudin, *Ann. Sci. Nat., Bot. sér. 3, 18:* 286 [Aug. 1852] (Naudin 1852). — *Coryphadenia abnormis* (Naudin) Morley, *Amer. J. Bot.* 40: 252 (Morley 1953).

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777669] designated by Lanjouw & Uittien [1940: 160]; isolecto-, BM[BM001008149]).

INVENTORY DATA (FG). — 83 trees in 35 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 46.8\text{ cm}$.

Family MELIACEAE Juss.
Genus *Cabralea* A.Juss.

[1110] *Cabralea canjerana* (Vell.) Mart.

Syst. Mat. Med. Veg. Bras. 38 (Martius 1843), “*Cabralia*”. — *Trichilia canjerana* Vell., *Fl. Flumin.*: 176 [“1825” publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Cabralea glaberrima A.Juss., *Mém. Mus. Hist. Nat.* 19: 270 (Jussieu 1830).

Cabralea cangerana Saldanha, *Ann. Sci. Nat., Bot. sér. 5, 19:* 210 (Saldanha 1874).

Cabralea burchellii C.DC., *Fl. Bras. [Martius]* 11 (1): 171 [1 Feb. 1878] (Candolle 1878).

Cabralea estrellensis C.DC., *Fl. Bras. [Martius]* 11 (1): 171 [1 Feb. 1878] (Candolle 1878).

Cabralea jussiaeana C.DC., *Fl. Bras. [Martius]* 11 (1): 171 [1 Feb. 1878] (Candolle 1878).

Cabralea sulcata C.DC., *Fl. Bras. [Martius]* 11 (1): 171 [1 Feb. 1878] (Candolle 1878).

Cabralea gaudichaudii C.DC., *Fl. Bras. [Martius]* 11 (1): 172 [1 Feb. 1878] (Candolle 1878).

Cabralea glaziovii C.DC., *Fl. Bras. [Martius]* 11 (1): 172 [1 Feb. 1878] (Candolle 1878).

Cabralea pedunculata C.DC., *Fl. Bras. [Martius]* 11 (1): 172 [1 Feb. 1878] (Candolle 1878).

Cabralea silvatica C.DC., *Fl. Bras. [Martius]* 11 (1): 173 [1 Feb. 1878] (Candolle 1878).

Cabralea eichleriana C.DC., *Fl. Bras. [Martius]* 11 (1): 174 [1 Feb. 1878] (Candolle 1878).

Cabralea macrophylla Fenzl ex C.DC., *Fl. Bras. [Martius]* 11 (1): 174 [1 Feb. 1878] (Candolle 1878).

Cabralea macrophylla var. *decomposita* C.DC., *Fl. Bras. [Martius]* 11 (1): 174 [1 Feb. 1878] (Candolle 1878).

Cabralea multijuga C.DC., *Fl. Bras. [Martius]* 11 (1): 175 [1 Feb. 1878] (Candolle 1878).

Cabralea pallescens C.DC., *Fl. Bras. [Martius]* 11 (1): 175 [1 Feb. 1878] (Candolle 1878).

Cabralea laevis C.DC., *Fl. Bras. [Martius]* 11 (1): 176 [1 Feb. 1878] (Candolle 1878).

Cabralea corcovadensis C.DC., *Fl. Bras. [Martius]* 11 (1): 177 [1 Feb. 1878] (Candolle 1878).

Cabralea lagoensis C.DC., *Fl. Bras. [Martius]* 11 (1): 177 [1 Feb. 1878] (Candolle 1878).

Cabralea pilosa C.DC., *Fl. Bras. [Martius]* 11 (1): 177 [1 Feb. 1878] (Candolle 1878).

Cabralea pilosa var. *glabrior* C.DC., *Fl. Bras. [Martius]* 11 (1): 177 [1 Feb. 1878] (Candolle 1878).

Cabralea warmingiana C.DC., *Fl. Bras. [Martius]* 11 (1): 178 [1 Feb. 1878] (Candolle 1878).

Cabralea warmingiana var. *coriacea* C.DC., *Fl. Bras. [Martius]* 11 (1): 178 [1 Feb. 1878] (Candolle 1878).

Cabralea lundii C.DC., *Fl. Bras. [Martius]* 11 (1): 180 [1 Feb. 1878] (Candolle 1878).

Cabralea riedelii C.DC., *Fl. Bras. [Martius]* 11 (1): 180 [1 Feb. 1878] (Candolle 1878).

Cabralea poeppigii C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 471 [June 1878] (Candolle 1878).

Cabralea lagoensis var. *glabra* C.DC., *Bull. Herb. Boissier*, sér. 2, 1: 364 (Candolle 1901).

Cabralea schwackei C.DC., *Bull. Herb. Boissier*, sér. 2, 1: 365 (Candolle 1901).

Cabralea brachystachya C.DC., *Bull. Herb. Boissier*, sér. 2, 3: 412 (Candolle 1903).

Cabralea villosa C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 134 (Candolle 1907).

Cabralea eichleriana var. *macrantha* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 136 (Candolle 1907).

Cabralea rojasii C.DC., *Bull. Soc. Bot. Genève*, sér. 2, 6: 113 [“1914” publ. 1915] (Candolle 1915), “*Rojasii*”.

Cabralea oblongifoliola C.DC., *Repert. Spec. Nov. Regni Veg.* 14: 403 (Candolle 1916).

Cabralea erismatica A.C.Sm., *Bull. Torrey Bot. Club* 60 (5): 360 [May 1933] (Smith 1933).

Cabralea cauliflora Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 13: 501 [30 June 1937] (Harms 1937).

Cabralea macrantha Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 13: 502 [30 June 1937] (Harms 1937).

Cabralea lacaziana Rizzini, *Leandra* 6: 33 (Rizzini 1975).

NOTE. — *J.-J. de Granville 1220* was collected on an inselberg on the Brazil-French Guiana border, a few meters on the Brazilian side. It is thus expected to occur in French Guiana.

VERNACULAR NAMES. — Br: cajarana, canjarana.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville 1220*.

SIZE. — Up to 40 cm dbh (Pennington *et al.* 1981).




FIG. 37. — Melastomataceae: **A**, *Mouriri crassifolia* Sagot (J.-F. Molino et al. 2070); **B**, *Mouriri guianensis* Aubl. (M.-F. Prévost & D. Sabatier 5001). Metteniusaceae: **C**, *Emmotum fagifolium* Desv. ex Ham. Monimiaceae: **D**, *Mollinedia grazielae* Peixoto (D. Sabatier & J.-F. Molino 5755). © D. Sabatier/IRD.

Genus *Carapa* Aubl.

[1111] *Carapa guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 32 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Granatum guianense* (Aubl.) Kuntze, *Revis. Gen. Pl.* 1: 110 [5 Nov. 1891] (Kuntze 1891).

Persoonia guareoides Willd., *Sp. Pl.*, ed. 4 2 (1): 331 [Mar. 1799] (Willdenow 1799).

Amapa guianensis Steud., *Nomencl. Bot. [Steudel]* 1: 35 (Steudel 1821), nom. nud.

Xylocarpus carapa Spreng., *Syst. Veg. [Sprengel]* 2: 213 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Carapa guianensis*).

Carapa latifolia Willd. ex C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 718 [June 1878] (Candolle 1878), nom. nud. pro syn.

Guarea mucronulata C.DC., *Notizbl. Bot. Gart. Berlin-Dahlem* 7: 499 [15 Jan. 1917] (Candolle 1917).

Carapa macrocarpa Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 91 (Ducke 1922), pro parte fruto excluso.

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: tiviru • Ka: kalapa • Te: dzandi • Wp: yani, yani pilá, yani sī • Wn: kalapa • Nt: kaapa • Cr: karapa • Fr: carapa • Br: andiroba.

HERBARIUM DATA (FG). — 53 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000759977]).

INVENTORY DATA (FG). — 170 trees in 42 plots; $F_{\max} = 3\%$; $dbh_{\text{inv}} = 80.2$ cm.

[1112] *Carapa surinamensis* Miq.

Stirp. Surinam. Select.: 75 [“1850” publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Pa: tiviru-seinô • Ka: ityumban kalapa • Te: dzandi • Wp: yani pilá • Wn: kalapa • Nt: kaapa • Cr: karapa • Fr: carapa • Br: andiroba.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *S.A. Mori et al. 26487*.

INVENTORY DATA (FG). — 994 trees in 187 plots; $F_{\max} = 4.4\%$; $dbh_{\text{inv}} = 78.8$ cm.

- Genus *Cedrela* P.Browne
- [1113] *Cedrela fissilis* Vell.
(Fig. 38A)
- Fl. Flumin.*: 75 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829). — *Surenus fissilis* (Vell.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela brasiliensis* A.Juss., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 86 [10 Oct. 1829] (Jussieu 1829).
- Cedrela brasiliensis* var. *australis* A.Juss., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 86 [10 Oct. 1829] (Jussieu 1829).
- Cedrela fissilis* var. *glabrior* C.DC., *Fl. Bras. [Martius]* 11 (1): 224 [1 Feb. 1878] (Candolle 1878).
- Cedrela fissilis* var. *macrocarpa* C.DC., *Bull. Herb. Boissier* 2: 574 (Candolle 1894).
- Cedrela barbata* C.DC., *Bull. Herb. Boissier* 2: 575 (Candolle 1894).
- Cedrela fuscata* Rojas Acosta, *Cat. Hist. Nat. Corrientes*: 154 (Rojas Acosta 1897), nom. nud.
- Cedrela hirsuta* C.DC., *Bull. Herb. Boissier*, sér. 2, 3: 413 (Candolle 1903).
- Cedrela caldasana* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 170 (Candolle 1907).
- Cedrela longiflora* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 173 (Candolle 1907).
- Cedrela pachyrhachis* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 174 (Candolle 1907).
- Cedrela regnelli* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 174 (Candolle 1907).
- Cedrela pilgeri* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 175 (Candolle 1907).
- Cedrela brunelliooides* Rusby, *Bull. New York Bot. Gard.* 8 (28): 99 [23 Nov. 1912] (Rusby 1912), *pro parte folia exclusa*.
- Cedrela balansae* C.DC., *Bull. Soc. Bot. Genève*, sér. 2, 6: 119 ["1914" publ. 1915] (Candolle 1915).
- Cedrela tubiflora* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 135 (Bertoni 1918).
- Cedrela tubiflora* f. *angustifolia* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 137 (Bertoni 1918).
- Cedrela tubiflora* subsp. *bertoniensis* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 137 (Bertoni 1918).
- Cedrela tubiflora* var. *grandifolia* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 137 (Bertoni 1918). — *Cedrela tubiflora* f. *grandifolia* (Bertoni) M.Buchinger & Falcone, *Darwiniana* 10: 464 (Buchinger & Falcone 1953).
- Cedrela tubiflora* var. *intermedia* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 137 (Bertoni 1918).
- Cedrela tubiflora* var. *lagenaria* Bertoni, *Anales Ci. Parag.*, ser. 2, 2: 138 (Bertoni 1918).
- Cedrela alliacea* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 189 (Ducke 1922), nom. nud. pro syn.
- Cedrela huberi* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 189 (Ducke 1922).
- Cedrela macrocarpa* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 189 (Ducke 1922).
- Cedrela tubiflora* f. *parvifoliola* M.Buchinger & Falcone, *Darwiniana* 10: 464 (Buchinger & Falcone 1953).
- Cedrela elliptica* Rizzini & Heringer, *Anais Acad. Brasil. Ci.* 38 (Suppl.): 111 [31 Dec. 1966] (Rizzini & Heringer 1966), "Cedrella".
- VERNACULAR NAMES. — Br: cedro-balata.
- HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al.* 6161.
- INVENTORY DATA (FG). — 1 tree, dbh = 20.5 cm.
- [1114] *Cedrela odorata* L.
- Syst. Nat.*, ed. 10, 2: 940 [7 June 1759] (Linnaeus 1759).
- Cedrela brownii* Loefl., *Iter Hispan.* 183 (Loefling 1758). — *Surenus brownii* (Loefl.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela guianensis* A.Juss., *Mém. Mus. Hist. Nat.* 19: 295 (Jussieu 1830). — *Surenus guianensis* (A.Juss.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela paraguariensis* Mart., *Flora* 20 (2, Beibl.): 93 (Martius 1837).
- Cedrela mexicana* M.Roem., *Fam. Nat. Syn. Monogr.* 1: 137 (Roemer 1846). — *Surenus mexicana* (M.Roem.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela velloziana* M.Roem., *Fam. Nat. Syn. Monogr.* 1: 137 (Roemer 1846). — *Surenus velloziana* (M.Roem.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela glaziovii* C.DC., *Fl. Bras. [Martius]* 11 (1): 224 [1 Feb. 1878] (Candolle 1878). — *Surenus glaziovii* (C.DC.) Kuntze, *Revis. Gen. Pl.* 1: 111 [5 Nov. 1891] (Kuntze 1891).
- Cedrela adenophylla* Mart. ex C.DC., *Fl. Bras. [Martius]* 11 (1): 226 [1 Feb. 1878] (Candolle 1878), nom. nud. pro syn.
- Cedrela paraguariensis* var. *brachystachya* C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 738 [June 1878] (Candolle 1878). — *Cedrela brachystachya* (C.DC.) C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 168 (Candolle 1907).
- Cedrela paraguariensis* var. *multipijuga* C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 738 [June 1878] (Candolle 1878).
- Cedrela dugesii* S.Watson, *Proc. Amer. Acad. Arts* 18: 190 (Watson 1883).
- Cedrela occidentalis* C.DC. & Rose, *Contr. U.S. Natl. Herb.* 5 (4): 190 [31 Oct. 1899] (Candolle & Rose 1899).
- Cedrela paraguariensis* var. *hassleri* C.DC., *Bull. Herb. Boissier*, sér. 2, 3: 413 (Candolle 1903). — *Cedrela hassleri* (C.DC.) C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 172 (Candolle 1907).
- Cedrela mexicana* var. *puberula* C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 427 (Candolle 1905).

Cedrela sintenisii C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 169 (Candolle 1907).

Cedrela mourae C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 171 (Candolle 1907).

Cedrela rotunda S.F.Blake, *Proc. Biol. Soc. Washington* 33: 109 (Blake 1920).

Cedrela whitfordii S.F.Blake, *Proc. Biol. Soc. Washington* 33: 110 (Blake 1920).

Cedrela yucatana S.F.Blake, *Proc. Biol. Soc. Washington* 33: 110 (Blake 1920).

Cedrela ciliolata S.F.Blake, *Proc. Biol. Soc. Washington* 34: 115 (Blake 1921).

Cedrela longipes S.F.Blake, *Contr. U.S. Natl. Herb.* 24 (1): 9 [11 Jan. 1922] (Blake 1922).

Cedrela longipetiolulata Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 179 [20 Dec. 1927] (Harms 1927).

Cedrela palustris Handro, *Arq. Bot. Estado São Paulo* 3: 223 (Handro 1962).

Cedrela odorata var. *xerogelton* Rizzini & Heringer, *Anais Acad. Brasil. Ci.* 38 (Suppl.): 112 [31 Dec. 1966] (Rizzini & Heringer 1966).

Cedrela cubensis Bisbe, *Feddes Repert.* 85 (9-10): 595 (Bisse 1974).

VERNACULAR NAMES. — Pa: awayo, sedri-wahuyo • Ka: samaliyapo, simaliapo • Te: katsu • Wp: kaisu, kaisu pilá, kaisu sī • Wn: simali • Nt: sede, sedee • Cr: akajou-rouj, sèd-akajou, sèd-rouj • Fr: acajou de Guyane, cèdre rouge • Br: cedro-verdadeiro, cedro-vermelho.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J. Martin s.n.* (holotype of *Cedrela guianensis*: P[P01819654]; iso-, BM[BM000992023], F[V0064067F], K[K000700268, K000700269], NY[00053521], P[P01819655, P01819656], US[00108140, 00603677, 00603678, 00603679]).

INVENTORY DATA (FG). — 15 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 100$ cm.

Genus *Guarea* F.Allam.

[1115] *Guarea carinata* Ducke

Trop. Woods 76: 16 (Ducke 1943).

VERNACULAR NAMES. — Wp: payawalu'i sili.

HERBARIUM DATA (FG). — A single collection, *P. Grenand* 1527.

SIZE. — Brazil, Pará. *G.T. Prance* 1929 (MO), 15 m × 25 cm.

[1116] *Guarea cinnamomea* Harms

Notizbl. Bot. Gart. Berlin-Dahlem 13: 504 [30 June 1937] (Harms 1937).

Guarea quadrangularis M.E.Morales, *Revista Acad. Colomb. Ci. Exact.* 22 (84): 336 (Morales 1998).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman* B-1017.

SIZE. — Brazil, Acre. *C.A. Cid Ferreira et al.* 10505 (MO), 20 m × 35 cm.

[1117] *Guarea convergens* T.D.Penn.

Fl. Neotrop. Monogr. 28: 260 [4 Dec. 1981] (Pennington 1981).

HERBARIUM DATA (FG). — A single collection, *D. Loubry* 1774.

SIZE. — Brazil, Amazonas. *T.D. Pennington* 9967 (MO), 25 m × 35 cm.

[1118] *Guarea costata* A.Juss.

Mém. Mus. Hist. Nat. 19: 241 (Jussieu 1830).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: karavia-kamwi • Ka: pešolowa • Wp: yatoa'i sī.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *G.S. Perrottet s.n.* (type P[P02288018, P02288019]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 11.8$ cm.

[1119] *Guarea gomma* Pulle

Recueil Trav. Bot. Néerl. 6: 271 (Pulle 1909).

VERNACULAR NAMES. — Pa: karavia • Ka: siwaluwa, yukutuna • Wp: payawalu'i sili • Cr: bwa-jako • Br: jatuaúba-preta, jitó-da-terra-firme.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-2784.

INVENTORY DATA (FG). — 92 trees in 23 plots; $F_{\max} = 3.5\%$; $dbh_{\text{inv}} = 58.6$ cm.

[1120] *Guarea guidonia* (L.) Sleumer

Taxon 5 (8): 194 [Nov. 1956] (Sleumer 1956). — *Samyda guidonia* L., *Sp. Pl.* 1: 443 [1 May 1753] (Linnaeus 1753).

Melia guara Jacq., *Enum. Syst. Pl.*: 20 [Aug.-Sep. 1760] (Jacquin 1760). — *Trichilia guara* (Jacq.) L., *Sp. Pl.*, ed. 2, 1: 551 [Sep. 1762] (Linnaeus 1762). — *Guarea guara* (Jacq.) P.Wilson, *N. Amer. Fl.* 25 (4): 272 [10 Mar. 1924] (Wilson 1924).

Guarea trichiliooides L., *Mant. Pl.* 2: 228 (Linnaeus 1771), *nom. illeg. superfl.* (based on *Trichilia guara*).

Guarea brachystachya Moc. & Sessé ex DC., *Prodr. [A. P. de Candolle]* 1: 624 [mid Jan. 1824] (Candolle 1824).

Guarea multijuga A.Juss., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 82 [10 Oct. 1829] (Jussieu 1829).

Guarea purgans A.Juss., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (13): 83 [10 Oct. 1829] (Jussieu 1829). — *Guarea tuberculata* Vell. var. *purgans* (A.Juss.) C.DC., *Fl. Bras. [Martius]* 11 (1): 191 [1 Feb. 1878] (Candolle 1878).

Guarea Aubletii A.Juss., *Mém. Mus. Hist. Nat.* 19: 241 (Jussieu 1830).

Guarea multiflora A.Juss., *Mém. Mus. Hist. Nat.* 19: 241 (Jussieu 1830).

Guarea bahiensis Klotzsch, *Linnaea* 14: 300 (Klotzsch 1840).

- Trichilia Aubletii* Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 700 (Steudel 1841), *nom. nud.*
- Guarea trichilioides* var. *brachystachya* C.DC., *Fl. Bras. [Martius]* 11 (1): 184 [1 Feb. 1878] (Candolle 1878).
- Guarea trichilioides* var. *pallida* C.DC., *Fl. Bras. [Martius]* 11 (1): 184 [1 Feb. 1878] (Candolle 1878).
- Guarea alternans* C.DC., *Fl. Bras. [Martius]* 11 (1): 186 [1 Feb. 1878] (Candolle 1878).
- Guarea francavillana* C.DC., *Fl. Bras. [Martius]* 11 (1): 187 [1 Feb. 1878] (Candolle 1878).
- Guarea alba* C.DC., *Fl. Bras. [Martius]* 11 (1): 189 [1 Feb. 1878] (Candolle 1878), *nom. nud.*
- Guarea surinamensis* Miq. ex C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 543 [June 1878] (Candolle 1878), *nom. nud.*
- Sycocarpus rusbyi* Britton, *Bull. Torrey Bot. Club* 14 (7): 143 [July 1887] (Britton 1887). — *Guarea rusbyi* (Britton) Rusby, *Mem. Torrey Bot. Club* 6 (1): 17 (Rusby 1896).
- Guarea bilibil* C.DC., *Bull. Herb. Boissier* 2: 570 (Candolle 1894).
- Guarea sylvestris* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 335 ["1894-96" publ. Dec. 1895] (Moore 1895).
- Guarea rubricalyx* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 336 ["1894-96" publ. Dec. 1895] (Moore 1895).
- Guarea rubescens* C.DC., *Bull. Herb. Boissier*, sér. 2, 3: 407 (Candolle 1903).
- Guarea xiroresana* C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 418 (Candolle 1905).
- Guarea trichilioides* var. *colombiana* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 138 (Candolle 1907).
- Guarea trichilioides* var. *decandra* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 139 (Candolle 1907).
- Guarea cabirme* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 143 (Candolle 1907).
- Guarea andreana* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 144 (Candolle 1907).
- Guarea eggersii* C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 176 (Candolle 1907).
- Guarea campestris* C.DC., *Bull. Soc. Bot. Genève*, sér. 2, 6: 114 ["1914" publ. 1915] (Candolle 1915).
- Guarea parva* C.DC., *Smithsonian Misc. Collect.* 68 (6): 3 [23 July 1917] (Candolle 1917).
- Guarea racemiformis* S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 241 [9 Sep. 1919] (Blake 1919).
- Guarea leticiana* Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 13: 505 [30 June 1937] (Harms 1937).
- Guarea puberula* Pittier, *Bol. Soc. Venez. Ci. Nat.* 4: 357 (Pittier 1938).
- Guarea rubrisepala* Cuatrec., *Fieldiana, Bot.* 27 (1): 73 (Cuatrecasas 1950).
- NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: atiwa'u, atuwai • Wp: yatoa'i si • Wn: wapiju, wapiu • Nt: dyakoomata, dyankoo mata, kalapa-oyak • Cr: bwa-bal, bwa-gro-bèk, bwa-kalimé • Br: jatuaúba-branca, marinheiro.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holotype of *Guarea aubletii*: P[P02288037]; iso-, P[P02288038, P02288039, P02288040, P02288041]).

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 21\text{ cm}$.

[1121] *Guarea kunthiana* A.Juss.

(Fig. 38B)

Mém. Mus. Hist. Nat. 19: 241 (Jussieu 1830).

Guarea densiflora Poepp., *Nova genera ac species plantarum [Poep-pig & Endlicher]* 3: 40 [8-11 Mar. 1843] (Poeppig 1843), *nom. illeg. hom.*, non Spreng. (Sprengel 1827).

Guarea poeppigii Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 15:* 371 (Triana & Planchon 1872).

Trichilia polyantha Poepp. ex Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 15:* 371 (Triana & Planchon 1872), *nom. nud. pro syn.*

Guarea glauca Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 15:* 372 (Triana & Planchon 1872).

Guarea pohlii C.DC., *Fl. Bras. [Martius]* 11 (1): 195 [1 Feb. 1878] (Candolle 1878).

Guarea pohlii var. *glabra* C.DC., *Fl. Bras. [Martius]* 11 (1): 195 [1 Feb. 1878] (Candolle 1878).

Guarea kunthiana var. *densiflora* C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 562 [June 1878] (Candolle 1878).

Guarea kunthiana var. *hahnii* C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 562 [June 1878] (Candolle 1878), "Hahnii".

Guarea kunthiana var. *hahniana* Krug & Urb. ex Duss, *Fl. Phan. Antill. Franç.* 128 ["1896" publ. 1897] (Duss 1897), "Kunthianum, Hahnianum".

Guarea pohlii var. *glabrior* C.DC., *Bull. Herb. Boissier*, sér. 2, 1: 361 (Candolle 1901).

Guarea grandifoliola C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 141 (Candolle 1907).

Guarea oblongiflora C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 141 (Candolle 1907).

Guarea simplicifolia C.DC., *Notizbl. Königl. Bot. Gart. Berlin* 6: 499 [15 Jan. 1917] (Candolle 1917).

Guarea williamsii C.DC., *Smithsonian Misc. Collect.* 68 (6): 4 [23 July 1917] (Candolle 1917).

Guarea membranacea Rusby, *Mem. New York Bot. Gard.* 7: 279 (Rusby 1927).

Guarea depauperata Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 243 [30 Mar. 1928] (Harms 1928).

Guarea steinbachii Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 348 [20 June 1928] (Harms 1928).




FIG. 38.—Malvaceae: **A**, *Cedrela fissilis* (D. Sabatier et al. 6161); **B**, *Guarea kunthiana* A.Juss. (D. Sabatier et al. 6007); **C**, *Guarea pubescens* (Rich.) A.Juss. subsp. *pubescens* (J.-F. Molino & D. Sabatier 2505); **D**, *Trichilia quadrijuga* Kunth (D. Sabatier & J.-F. Molino 5702). A, B, D, © D. Sabatier/IRD; C, © J.-F. Molino/IRD.

Guarea megaphylla Cuatrec., *Fieldiana, Bot.* 27 (1): 73 (Cuatrecasas 1950).

Guarea macrantha Standl. & L.O.Williams, *Ceiba* 1 (4): 240 (Standley & Williams 1951).

NOTE.—Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES.—Pa: á-kamwi • Wp: yatoa'i • Nt: dyankoi mata • Cr: bwa-pistelé • Br: cedrorana, jatuaúba, jitó.

HERBARIUM DATA (FG).—41 collections at CAY. Sel. exs.: P.A. Poiteau s.n. (holo-, P-JU [not seen]; iso-, K[K000380586, K000380587, K000380646]).

INVENTORY DATA (FG).—129 trees in 36 plots; $F_{\max} = 2.8\%$; $dbh_{inv} = 57$ cm.

[1122] *Guarea macrophylla* Vahl
subsp. *pachycarpa* (C.DC.) T.D.Penn.

Fl. Neotrop. Monogr. 28: 289 [4 Dec. 1981] (Pennington 1981).

Guarea trichiliooides var. *pachycarpa* C.DC., *Fl. Bras. [Martius]* 11 (1): 184 [1 Feb. 1878] (Candolle 1878).

Guarea rosea C.DC., *Fl. Bras. [Martius]* 11 (1): 186 [1 Feb. 1878] (Candolle 1878).

Guarea paraensis C.DC., *Fl. Bras. [Martius]* 11 (1): 187 [1 Feb. 1878] (Candolle 1878).

Guarea punctata C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 575 [June 1878] (Candolle 1878).

Guarea costulata C.DC., *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 237 (Candolle 1901).

Guarea subsessiliflora C.DC., *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 238 (Candolle 1901).

Guarea subsessiliflora var. *polyphyllaria* C.DC., *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 238 (Candolle 1901).

Guarea guedesii C.DC., *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 239 (Candolle 1901).

Guarea bangii Rusby, *Bull. Torrey Bot. Club* 49 (9): 262 [Sep. 1922] (Rusby 1922).

NOTE.—*G. macrophylla* is hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES.—Wp: kaliaku la ka'a, yatoa'i sili • Br: jitó-da-terra-firme.

HERBARIUM DATA (FG).—7 collections at CAY. Sel. exs.: *R.A.A. Oldeman 1221*.

INVENTORY DATA (FG).—32 trees in 3 plots; $F_{\max} = 3\%$; $dbh_{inv} = 43.4$ cm.

[1123] *Guarea megantha* A.Juss.

Mém. Mus. Hist. Nat. 19: 241 (Jussieu 1830). — *Guarea megantha* M.Roem., *Fam. Nat. Syn. Monogr.* 1: 120 (Roemer 1846), *nom. illeg. superfl.* (based on *Guarea megantha*).

Guarea grandifolia DC., *Prodri. [A. P. de Candolle]* 1: 624 [mid Jan. 1824] (Candolle 1824), *nom. illeg.* (*Melia guara* Jacq. in synonymy), *syn. nov.* — *Guarea grandiflora* Steud., *Nomencl. Bot. [Steudel]*, ed. 2: 709 (Steudel 1840), sphalm.

Guarea gigantea Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5*, 15: 370 (Triana & Planchon 1872).

Guarea chichon C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 147 (Candolle 1907).

Guarea trompillo C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 147 (Candolle 1907).

Guarea culebrana C.DC., *Smithsonian Misc. Collect.* 68 (6): 5 [23 July 1917] (Candolle 1917).

Guarea longipetiola C.DC., *Smithsonian Misc. Collect.* 68 (6): 5 [23 July 1917] (Candolle 1917).

Guarea pittieri C.DC., *Smithsonian Misc. Collect.* 68 (6): 6 [23 July 1917] (Candolle 1917).

Guarea borisii Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 13: 503 [30 June 1937] (Harms 1937).

Guarea mancharra Cuatrec., *Fieldiana, Bot.* 27 (1): 71 (Cuatrecasas 1950).

NOTES. — A species known only from French Guiana. *G. megantha* A.Juss. was treated by Pennington & Clarkson (2013) as a synonym of *G. grandifolia* DC., but the latter is illegitimate because Candolle (1824: 62) cited *Melia guara* Jacq. in synonymy. *G. megantha* is the earliest legitimate name for this species. *G. grandifolia* is based on two distinct elements: 1) a specimen from French Guiana in G-DC (“*in Guianâ Gallica*”) and 2) fruits of *Melia guara* Jacq. from Cuba [synonym of *Trichilia guara* (Jacq.) L. and of *G. trichilioides* L., *nom. illeg. superfl.*] Candolle (1824) did not see the material of the latter, and copied Jacquin’s description of the fruit of *Melia guara*. Although illegitimate, *G. trichilioides* is the type of the conserved *Guarea* F.Allem. (not “F.Allem. ex L.”, see Wiersema et al. 2018), and a synonym of *G. guidonia* (L.) Sleumer. *Guarea grandiflora* Steud. is actually an erroneous spelling of *G. grandifolia*. The entry in Steudel (1840: 709) reads “*grandiflora* Dec. *trichilioides*”, but under *trichilioides*, it appears as “*G. grandifolia* Dec. 1”. “Dec.” stands for A. P. de Candolle, and *G. grandifolia* is the first entry in Candolle’s treatment of *Guarea*.

VERNACULAR NAMES. — Wp: payawalu’i • Wn: kalapaimë • Br: carrapeta, jatuaúba-preta.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *J. Martin s.n.* (lectotype of *Guarea megantha*, P[P02288028], here designated; isolepto-, P[P02288029, P02288030, P02288031, P02288032]).

INVENTORY DATA (FG). — 78 trees in 33 plots; $F_{\max} = 1.9\%$; $dbh_{inv} = 99$ cm.

[1124] *Guarea pubescens* (Rich.) A.Juss. subsp. *pubescens* (Fig. 38C)

Mém. Mus. Hist. Nat. 19: 241 (Jussieu 1830). — *Trichilia pubescens* Rich., *Acies Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792).

Guarea affinis A.Juss., *Mém. Mus. Hist. Nat.* 19: 241 (Jussieu 1830).

Guarea richardiana A.Juss., *Mém. Mus. Hist. Nat.* 19: 241 (Jussieu 1830).

Guarea spicata C.DC., *Fl. Bras. [Martius]* 11 (1): 194 [1 Feb. 1878] (Candolle 1878).

Guarea huberi C.DC., *Bull. Herb. Boissier, sér. 2, 6:* 984 (Candolle 1906).

Guarea ulei Harms ex C.DC., *Annuaire Conserv. Jard. Bot. Genève* 10: 148 (Candolle 1907).

Ruagea microsepala Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 242 [30 Mar. 1928] (Harms 1928). — *Guarea microsepala* (Harms) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.2):* 767 (Macbride 1949).

Guarea davisii Sandwith, *Bull. Misc. Inform. Kew* 1933 (7): 330 [25 Sep. 1933] (Sandwith 1933).

Guarea rhabdotocarpa Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 13: 506 [30 June 1937] (Harms 1937).

Guarea concinna Sandwith, *Kew Bull.* 3 (2): 309 [20 Nov. 1948] (Sandwith 1948).

Guarea huberi var. *peruviana* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.2):* 764 (Macbride 1949).

VERNACULAR NAMES. — Pa: isauminyo, kawap-kamwi • Ka: yemelu • Wp: kaliaku la ka’á, yatoa’í paye, yatoa’í sī, yatoa’í sili • Wn: wapiu.

HERBARIUM DATA (FG). — 139 collections at CAY. Sel. exs.: *J.B. Leblond* 61 (type G[G00340026]).

INVENTORY DATA (FG). — 242 trees in 64 plots; $F_{\max} = 7.3\%$; $dbh_{inv} = 41.2$ cm.

[1125] *Guarea pubescens* subsp. *pubiflora* (A.Juss.) T.D.Penn.

Fl. Neotrop. Monogr. 28: 298 [4 Dec. 1981] (Pennington 1981).

Guarea pubiflora A.Juss., *Mém. Mus. Hist. Nat.* 19: 241 (Jussieu 1830).

Guarea pubiflora var. *parvifolia* C.DC., *Fl. Bras. [Martius]* 11 (1): 188 [1 Feb. 1878] (Candolle 1878).

Guarea pubiflora var. *angustifoliola* C.DC., *Bull. Herb. Boissier, sér. 2, 6:* 985 (Candolle 1906).

VERNACULAR NAMES. — Ka: yemelu • Wp: kaliaku la ka’á, yatoa’í paye, yatoa’í sili.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 904, 10 m tall.

[1126] *Guarea scabra* A.Juss.

Mém. Mus. Hist. Nat. 19: 241 (Jussieu 1830).

VERNACULAR NAMES. — Pa: á-kamwi • Wp: payawalu’í sili.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (holo-, P-JU[not seen]; iso-, P[P02288143, P00671684]).

INVENTORY DATA (FG). — 19 trees in 5 plots; $F_{\max} = 1.6\%$; $dbh_{inv} = 29.9$ cm.

[1127] *Guarea silvatica* C.DC.

Fl. Bras. [Martius] 11 (1): 195 [1 Feb. 1878] (Candolle 1878).

Guarea pedicellata C.DC., *Fl. Bras. [Martius]* 11 (1): 196 [1 Feb. 1878] (Candolle 1878).

Trichilia longeracemosa Glaz., *Bull. Soc. Bot. France* 52 (Mém. 3a): 95 (Glaizou 1905), nom. nud.

Guarea duckei C.DC., *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 434 (Candolle 1909).

Guarea bilocularis C.DC., *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 435 (Candolle 1909).

Guarea klugii Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 384 [30 Mar. 1932] (Harms 1932).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). One of five species whose relative density in tree communities likely indicative of pre-Columbian archeological sites in French Guiana (Molino *et al.* 2021).

VERNACULAR NAMES. — Pa: simartē, simatē • Wn: kalapi alekane • Nt: a-nyan-mi-kaasi udu • Br: jitó.

HERBARIUM DATA (FG). — 72 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2332.

INVENTORY DATA (FG). — 154 trees in 39 plots; $F_{\max} = 4.4\%$; $dbh_{inv} = 43.4$ cm.

[1128] *Guarea trunciflora* C.DC.

Monogr. Phan. [A.DC. & C.DC.] 1: 571 [June 1878] (Candolle 1878).

Carapa sericea Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 39 [8-11 Mar. 1843] (Poeppig 1843).

NOTE. — Although *D. Sabatier & S. Gonzalez* 5381 and 5399, from French Guiana, were both identified as *G. trunciflora* C.DC. in Pennington & Clarkson (2013), French Guiana was not mentioned in the species distribution.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & S. Gonzalez* 5381.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 40.3$ cm.

Genus *Trichilia* P.Browne[1129] *Trichilia cipo* (A.Juss.) C.DC.

Fl. Bras. [Martius] 11 (1): 214 [1 Feb. 1878] (Candolle 1878). — *Moschoxylum cipo* A.Juss., *Mém. Mus. Hist. Nat.* 19: 239 (Jussieu 1830).

Trichilia sampoiiana Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 387 [30 Mar. 1932] (Harms 1932).

Trichilia froesii A.C.Sm., *Bull. Torrey Bot. Club* 61 (4): 193 [Apr. 1934] (Smith 1934).

NOTE. — Jussieu (1830: 280) cited the material studied as “*Hab. in Guiâna (v.s.s. in herb. Richard qui florentem legerat in ripis fluvi Kourou et e cuius notis mss. quaedam his excerpti; Mus.; Gay.)*”

Therefore he stated that the original material was collected by L.C. Richard near the Kourou River, and is deposited at P. There are four original sheets with this name at P, that with barcode P02274150 has the original label and drawings by L.C. Richard. Pennington *et al.* (1981: 152) overlooked the information on these specimens and stated that their collector is unknown.

VERNACULAR NAMES. — Wp: iawalo sōwī • Br: louro-í.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holo-, P[P02274150]; iso-, P[P02274151, P02274152, P02274153]).

INVENTORY DATA (FG). — 86 trees in 21 plots; $F_{\max} = 1.8\%$; $dbh_{inv} = 32.8$ cm.

[1130] *Trichilia euneura* C.DC.

Monogr. Phan. [A.DC. & C.DC.] 1: 673 [June 1878] (Candolle 1878).

Trichilia stelligera Radlk., *Repert. Spec. Nov. Regni Veg.* 9: 372 (Radlkofe 1911).

VERNACULAR NAMES. — Wp: walili tuli.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *F.M.R. Leprieur s.n.* (holo-, G-DC[not seen]; iso-, G[G00352502], P[P02274170, P02274171, P02274172]).

INVENTORY DATA (FG). — 34 trees in 22 plots; $F_{\max} = 1\%$; $dbh_{inv} = 42.8$ cm.

[1131] *Trichilia lepidota* Mart.

subsp. *leucastera* (Sandwith) T.D.Penn.

Fl. Neotrop. Monogr. 28: 40 [4 Dec. 1981] (Pennington 1981). — *Trichilia leucastera* Sandwith, *Bull. Misc. Inform. Kew* 1933 (7): 328 [25 Sep. 1933] (Sandwith 1933).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2705.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 29.8$ cm.

[1132] *Trichilia martiana* C.DC.

Fl. Bras. [Martius] 11 (1): 205 [1 Feb. 1878] (Candolle 1878).

Trichilia insignis C.DC., *Fl. Bras. [Martius]* 11 (1): 206 [1 Feb. 1878] (Candolle 1878).

Trichilia gaudichaudii C.DC., *Fl. Bras. [Martius]* 11 (1): 209 [1 Feb. 1878] (Candolle 1878).

Trichilia caucana C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 669 [June 1878] (Candolle 1878).

Trichilia cuneata Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 642 (Radlkofe 1879).

Trichilia fuscescens Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 641 (Radlkofe 1879).

Trichilia heydeana C.DC., *Bot. Gaz.* 19 (1): 3 [17 Jan. 1894] (Candolle 1894).

Trichilia acutanthera C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 422 (Candolle 1905).

Trichilia bolleyi C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 423 (Candolle 1905).

Trichilia obtusanthera C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 423 (Candolle 1905).

Trichilia tonduzii C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 423 (Candolle 1905).

Trichilia bolleyi var. *nicoyensis* C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 424 (Candolle 1905).

Trichilia chiriquina C.DC., *Smithsonian Misc. Collect.* 68 (6): 6 [23 July 1917] (Candolle 1917).

Trichilia oaxacana S.F.Blake, *Contr. Gray Herb.* 53: 58 [26 Feb. 1918] (Blake 1918).

Trichilia izabalana S.F.Blake, *Contr. U.S. Natl. Herb.* 24 (1): 10 [11 Jan. 1922] (Blake 1922).

Trichilia colombiana Cuatrec., *Fieldiana, Bot.* 27 (1): 81 (Cuatrecasas 1950).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-734*.

INVENTORY DATA (FG). — 17 trees in 10 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 47.6$ cm.

[1133] *Trichilia micrantha* Benth.

Hooker's J. Bot. Kew Gard. Misc. 3: 369 (Bentham 1851).

Trichilia roraimana C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 670 [June 1878] (Candolle 1878).

Trichilia harmsii Rusby, *Bull. New York Bot. Gard.* 4 (14): 338 [7 Dec. 1907] (Rusby 1907).

Trichilia acariaeantha Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 431 [31 Oct. 1925] (Harms 1925).

Trichilia buchtienii Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 386 [30 Mar. 1932] (Harms 1932).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kawap-kamwi • Nt: samaati • Br: cedro-bravo, jitó-branco.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3333*.

INVENTORY DATA (FG). — 80 trees in 46 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 25.9$ cm.

[1134] *Trichilia micropetala* T.D.Penn.

Fl. Neotrop. Monogr. 28: 172 [4 Dec. 1981] (Pennington 1981).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Moretti 979*.

INVENTORY DATA (FG). — 6 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 11.5$ cm.

[1135] *Trichilia pallida* Sw.

Prodr. [Swartz]: 67 [20 Jun.-29 July 1788] (Swartz 1788).

Portesia ovata Cav., *Diss.* 7: 369 (Cavanilles 1789).

Guarea obtusifolia Lam., *Encycl. [J. Lamarck et al.]* 3 (1): 6 [19 Oct. 1789] (Lamarck 1789).

Hedwigia simplicifolia Spreng., *Neue Entdeck. Pflanzenk.* 3: 24 (Sprengel 1822). — *Trichilia simplicifolia* (Spreng.) Spreng., *Syst. Veg. [Sprengel]* 3: 69 [Jan.-Mar. 1826] (Sprengel 1826). — *Portesia simplicifolia* (Spreng.) M.Roem., *Fam. Nat. Syn. Monogr.* 1: 115 (Roemer 1846).

Trichilia montana Kunth, *Nova genera et species plantarum [H.B.K.]* 7: 226 [25 Apr. 1825] (Kunth 1825). — *Portesia montana* (Kunth) M.Roem., *Fam. Nat. Syn. Monogr.* 1: 115 (Roemer 1846).

Trichilia portoricensis Spreng., *Syst. Veg. [Sprengel]* 3: 68 [Jan.-Mar. 1826] (Sprengel 1826).

Trichilia diversifolia A.Juss., *Mém. Mus. Hist. Nat.* 19: 237 (Jussieu 1830). — *Portesia diversifolia* (A.Juss.) M.Roem., *Fam. Nat. Syn. Monogr.* 1: 116 (Roemer 1846). — *Pholacilia diversifolia* (A.Juss.) Griseb., *Fl. Brit. WI. [Grisebach]*: 130 [June 1859] (Grisebach 1859).

Trichilia trinitensis A.Juss., *Mém. Mus. Hist. Nat.* 19: 237 (Jussieu 1830). — *Portesia trinitensis* (A.Juss.) M.Roem., *Fam. Nat. Syn. Monogr.* 1: 116 (Roemer 1846). — *Pholacilia trinitensis* (A.Juss.) Griseb., *Fl. Brit. WI. [Grisebach]* 130 [June 1859] (Grisebach 1859).

Portesia echinocarpa de Vriese, *Ned. Kruidk. Arch.* 1 (3): 251 (de Vriese 1847). — *Trichilia echinocarpa* (de Vriese) Walp., *Ann. Bot. Syst. [Walpers]* 2 (2): 227 [21-24 Jan. 1852] (Walpers 1852).

Trichilia excelsa Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 368 (Bentham 1851).

Trichilia macrophylla Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 369 (Bentham 1851).

Trichilia goudotiana Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5,* 15: 370 (Triana & Planchon 1872).

Trichilia weddellii C.DC., *Fl. Bras. [Martius]* 11 (1): 201 [1 Feb. 1878] (Candolle 1878).

Trichilia weddellii var. *parvifolia* C.DC., *Fl. Bras. [Martius]* 11 (1): 201 [1 Feb. 1878] (Candolle 1878).

Trichilia weddellii var. *stylosa* C.DC., *Fl. Bras. [Martius]* 11 (1): 201 [1 Feb. 1878] (Candolle 1878).

Trichilia mollis C.DC., *Fl. Bras. [Martius]* 11 (1): 202 [1 Feb. 1878] (Candolle 1878).

Trichilia riedelii C.DC., *Fl. Bras. [Martius]* 11 (1): 202 [1 Feb. 1878] (Candolle 1878).

Trichilia flava C.DC., *Fl. Bras. [Martius]* 11 (1): 203 [1 Feb. 1878] (Candolle 1878).

Trichilia brachystachya Klotsch ex C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 650 [June 1878] (Candolle 1878).

Trichilia montana var. *fendleriana* C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 654 [June 1878] (Candolle 1878).

Trichilia peruviana C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 654 [June 1878] (Candolle 1878).

Trichilia lobulata C.DC., *Bull. Herb. Boissier*, sér. 2, 1: 364 (Candolle 1901).

Trichilia montana var. *acutivalvis* C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 422 (Candolle 1905).

Trichilia laminensis Barb.Rodr., *Contr. Jard. Bot. Rio de Janeiro* 4: 89 (Barbosa Rodrigues 1907).

Trichilia triphylla S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 243 [Sep. 1919] (Blake 1919).

Trichilia pauciflora Rusby, *Mem. New York Bot. Gard.* 7: 280 (Rusby 1927).

Trichilia truncata Leonard, *J. Wash. Acad. Sci.* 17: 68 (Leonard 1927).

Trichilia gigantophylla Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 246 [30 Mar. 1928] (Harms 1928).

Trichilia davisii Sandwith, *Bull. Misc. Inform. Kew* 1933 (7): 329 [25 Sep. 1933] (Sandwith 1933).

Trichilia subsimplex Steyermark., *Fieldiana, Bot.* 28 (2): 279 (Steyermark 1952).

Trichilia skutchii C.V.Morton & P.H.Allen, *Rain Forests Golfo Dulce*: 347 (Morton & Allen 1956).

VERNACULAR NAMES. — Ka: malakatano • Br: baga-de-morcego, catiguá.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2716*.

INVENTORY DATA (FG). — 187 trees in 42 plots; $F_{\max} = 3\%$; $\text{dbh}_{\text{inv}} = 41.4$ cm.

[1136] *Trichilia quadrijuga* Kunth
(Fig. 38D)

Nova genera et species plantarum [H.B.K.] 5: 215 [25 Feb. 1822] (Kunth 1822). — *Odontandra quadrijuga* (Kunth) Triana & Planch., *Ann. Sci. Nat., Bot.* sér. 5, 15: 374 (Triana & Planchon 1872).

Moschoxylum pentandrum Poepp. & Endl., *Nova genera ac species plantarum* [Poeppig & Endlicher] 3: 39 [8-11 Mar. 1843] (Poeppig & Endlicher 1843), "Moschoxylon".

Moschoxylum propinquum Miq., *Stirp. Surinam. Select.*: 74 ["1850" publ. Mar. 1851] (Miquel 1851). — *Trichilia propinqua* (Miq.) C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 693 [June 1878] (Candolle 1878).

Trichilia paraensis C.DC., *Fl. Bras. [Martius]* 11 (1): 213 [1 Feb. 1878] (Candolle 1878).

Trichilia tocacheana C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 701 [June 1878] (Candolle 1878).

Rourea bakeriana Britton, *Bull. Torrey Bot. Club* 16 (7): 192 [6 July 1889] (Britton 1889).

Trichilia paracaimana C.DC., *Notizbl. Königl. Bot. Gart. Berlin* 6: 501 [15 Jan. 1917] (Candolle 1917).

Trichilia alta S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 241 [9 Sep. 1919] (Blake 1919).

Trichilia curranii S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 242 [Sep. 1919] (Blake 1919).

Trichilia compacta A.C.Sm., *Lloydia* 2 (3): 186 (Smith 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: iwawo • Cr: lanson-ti-fey • Br: breu-maxixe, pau-rosa-branca.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 885*.

INVENTORY DATA (FG). — 51 trees in 19 plots; $F_{\max} = 2.4\%$; $\text{dbh}_{\text{inv}} = 60$ cm.

[1137] *Trichilia schomburgkii* C.DC.

Monogr. Phan. [A.DC. & C.DC.] 1: 695 [June 1878] (Candolle 1878).

Trichilia subsessilifolia C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 685 [June 1878] (Candolle 1878).

Trichilia cuneifolia Pulle, *Recueil Trav. Bot. Néerl.* 6: 272 (Pulle 1909).

Trichilia siqueiraei Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 192 (Ducke 1922).

VERNACULAR NAMES. — Pa: ahuwahu-puvemna, auwau-puvemna, kawap • Ka: alawata pana, inyamu ekunali wewe • Wp: yatoa'i sī • Wn: akalali.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3090*.

INVENTORY DATA (FG). — 181 trees in 64 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 57.3$ cm.

[1138] *Trichilia septentrionalis* C.DC.

Fl. Bras. [Martius] 11 (1): 220 [1 Feb. 1878] (Candolle 1878).

Trichilia moritzii C.DC., *Monogr. Phan. [A.DC. & C.DC.]* 1: 707 [June 1878] (Candolle 1878).

Trichilia polynera C.DC., *Bull. Herb. Boissier*, sér. 2, 5: 426 (Candolle 1905).

Trichilia lanceolata Pittier, *Arb. Arbust. Venez.* 1: 8 (Pittier 1921), *nom. illeg. hom., non* C.DC. (Candolle 1878).

Trichilia eurysepala Harms, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 246 [30 Mar. 1928] (Harms 1928).

Trichilia krukovi A.C.Sm., *Bull. Torrey Bot. Club* 60 (5): 360 [May 1933] (Smith 1933).

Trichilia magnifica Baehni & J.F.Macbr., *Candollea* 5: 380 (Baehni & Macbride 1934).

Trichilia grandis Lasser & Maguire, *Bol. Soc. Venez. Ci. Nat.* 15: 101 (Lasser & Maguire 1954).

VERNACULAR NAMES. — Pa: kawap • Ka: papelilan • Wp: yatoa'i sī, yatoa'i u.

HERBARIUM DATA (FG). — 75 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4914*.

INVENTORY DATA (FG). — 142 trees in 23 plots; $F_{\max} = 6.1\%$; $\text{dbh}_{\text{inv}} = 32.5$ cm.

[1139] *Trichilia surinamensis* (Miq.) C.DC.

Monogr. Phan. [A.DC. & C.DC.] 1: 697 [June 1878] (Candolle 1878). — *Moschoxylum surinamense* Miq., *Stirp. Surinam. Select.* 73 ["1850" publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Wp: wila pisi'u, yatoa'i u.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2797*.

INVENTORY DATA (FG). — 82 trees in 29 plots; $F_{\max} = 2\%$; $dbh_{\text{inv}} = 27.4$ cm.

[1140] *Trichilia* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 5648*.

INVENTORY DATA (FG). — 8 trees in 3 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 35$ cm.

Family METTENIUSACEAE H.Karst.ex Schnizl.

Genus *Dendrobangia* Rusby

[1141] *Dendrobangia boliviiana* Rusby

Mem. Torrey Bot. Club 6 (1): 19 (Rusby 1896).

Clavapetalum surinamense Pulle, *Recueil Trav. Bot. Néerl.* 9: 148 (Pulle 1912).

Asterolepidion elatum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 207 (Ducke 1922). — *Clavapetalum elatum* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 116 (Ducke 1925).

Dendrobangia tenuis Ducke, *Bol. Técn. Inst. Agron. N.* 4: 15 [31 Mar. 1945] (Ducke 1945).

VERNACULAR NAMES. — Pa: aituwu • Ka: melimeli • Wp: wila iki'iy • Nt: taapu tiki • Cr: bwa-léchél • Fr: bois échelle • Br: caferana, ivái, pombinho.

HERBARIUM DATA (FG). — 82 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 946*.

INVENTORY DATA (FG). — 371 trees in 102 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 64.3$ cm.

Genus *Emmotum* Desv.

[1142] *Emmotum fagifolium* Desv. ex Ham.
(Fig. 37C)

Prod. Pl. Ind. Occid. [Hamilton]: 29 [Oct. 1825] (Hamilton 1825).

Pogopetalum acutum Benth., *London J. Bot.* 2: 377 (Bentham 1843).

VERNACULAR NAMES. — Nt: tyentyen udu • Cr: bwa-agouti • Br: muiraximbé, pau-de-sobre.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *A.N. Desvaux s.n.* (holo-, P[P00060476]; iso-, P[P00060477]).

INVENTORY DATA (FG). — 24 trees in 19 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 78.9$ cm.

Genus *Poraqueiba* Aubl.

[1143] *Poraqueiba guianensis* Aubl.

Hist. Pl. Guiane 1: 123 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate.

Poraqueiba surinamensis Miers, *Ann. Mag. Nat. Hist.*, ser. 2, 9 (54): 483 [June 1852] (Miers 1852).

VERNACULAR NAMES. — Pa: ahakyu-kamwi, ahikyu-kamwi, kanegma • Ka: aletepe, walulan • Br: umari-bravo.

HERBARIUM DATA (FG). — 64 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777819] designated by Lanjouw & Uittien [1940: 155]; isolecto-, BM[BM000839695]).

INVENTORY DATA (FG). — 705 trees in 130 plots; $F_{\max} = 4.9\%$; $dbh_{\text{inv}} = 43.4$ cm.

Family MONIMIACEAE Juss.

Genus *Mollinedia* Ruiz & Pav.

[1144] *Mollinedia grazielae* Peixoto
(Fig. 37D)

Bradea 4 (41): 332 (Peixoto 1987).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-J. de Granville 7734*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $dbh_{\text{inv}} = 16$ cm.

[1145] *Mollinedia ovata* Ruiz & Pav.

Syst. Veg. Fl. Peruv. Chil. 1: 143 [late Dec. 1798] (Ruiz & Pavón 1798).

Tetratomia latifolia Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher]* 2: 47 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838). — *Mollinedia latifolia* (Poepp. & Endl.) Tul., *Arch. Mus. Hist. Nat.* 8: 402 (Tulasne 1855).

Mollinedia laurina Tul., *Ann. Sci. Nat., Bot. sér. 4, 3*: 43 [May 1855] (Tulasne 1855).

Mollinedia grosseserrata Perkins, *Bot. Jahrb. Syst.* 27 (5): 673 [23 Feb. 1900] (Perkins 1900).

Mollinedia rusbyana Perkins, *Bot. Jahrb. Syst.* 27 (5): 682 [23 Feb. 1900] (Perkins 1900).

Mollinedia grandifolia Perkins, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 158 [20 Dec. 1927] (Perkins 1927).

Mollinedia tessmannii Perkins, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 159 [20 Dec. 1927] (Perkins 1927).

Mollinedia casca J.F.Macbr., *Candollea* 5: 352 (Macbride 1934).

Mollinedia krukovi A.C.Sm., *Phytologia* 1 (3): 114 [21 Jan. 1935] (Smith 1935).

Mollinedia ptariensis Steyermark., *Fieldiana, Bot.* 28 (1): 235 (Steyermark 1951).

Mollinedia glabricaulis Maguire & Steyermark, *Mem. New York Bot. Gard.* 51: 115 (Maguire & Steyermark 1989).

Mollinedia neblinensis Maguire & Steyermark, *Mem. New York Bot. Gard.* 51: 113 (Maguire & Steyermark 1989).

VERNACULAR NAMES. — Pa: peruvia-kamwi • Wp: alasiku sili.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *G. Cremers 6509*.

SIZE. — Up to 15 m tall (Berry & Peixoto 1999).

Family MORACEAE Gaudich.
Genus *Bagassa* Aubl.

[1146] *Bagassa guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 15 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

Piper tiliifolium Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 4 [Oct. 1825] (Hamilton 1825), “*tiliaefolium*”. — *Laurea tiliifolia* (Desv. ex Ham.) Gaudich., *Voy. Uranie, Bot.*: 513 [“1826” publ. 6 Mar. 1830] (Gaudichaud 1830), “*tiliaefolia*”. — *Bagassa tiliifolia* (Desv. ex Ham.) Benoist, *Arch. Bot. 5 (Mém. 1)*: 31 [27 Mar. 1933] (Benoist 1933), “*tiliaefolia*”.

Bagassa sagotiana Bureau ex Benth. & Hook.f., *Gen. Pl. [Bentham & Hooker f.] 3 (1)*: 362 [7 Feb. 1880] (Bentham & Hooker 1880).

NOTE. — Gaudichaud (1830: 513) based his *Laurea tiliifolia* on “*Piper tiliifolium* Desvaux mss. in herb. Desf. Mus. In Guyanâ (Cayenne) Martin Ded.”. This is undoubtedly the type of *Piper tiliifolium* Desv. ex Ham. The latter name is thus the basionym of both *L. tiliifolia* and *B. tiliifolia*.

VERNACULAR NAMES. — Pa: pakad • Ka: pakasa • Te: pakatsa • Wp: pakasa • Nt: kaw udu, odun • Cr: bagas • Fr: bagasse • Br: bagaceira, tatajuba.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000993272]).

INVENTORY DATA (FG). — 26 trees in 21 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 126$ cm.

Genus *Batocarpus* H.Karst.

[1147] *Batocarpus amazonicus* (Ducke) Fosberg

Proc. Biol. Soc. Washington 55: 101 (Fosberg 1942). — *Anonocarpus amazonicus* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 39 (Ducke 1922).

Batocarpus maranhensis Emygdio & Emmerich, *Bol. Mus. Nac. Rio de Janeiro, Bot.* 37: 9 (Emygdio & Emmerich 1968).

VERNACULAR NAMES. — Pa: amap • Br: guariúba-branca, mururé.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier 28*.

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 36.8$ cm.

Genus *Brosimum* Sw.

[1148] *Brosimum acutifolium* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 6: 66 (Huber 1910). — *Brosimopsis acutifolia* (Huber) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 30 (Ducke 1922). — *Piratinera acutifolia* (Huber) Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 100 [18 June 1918] (Pittier 1918).

VERNACULAR NAMES. — Pa: tauni • Ka: takini • Te: takweni • Wp: takweni • Nt: takinan • Cr: bwa-mondan, takini • Fr: bois tarin, takini • Br: mururé.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2585*.

INVENTORY DATA (FG). — 59 trees in 36 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 83$ cm.

[1149] *Brosimum guianense* (Aubl.) Huber
(Fig. 39A)

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 337 (Huber 1909), “*guyanense*”. — *Piratinera guianensis* Aubl., *Hist. Pl. Guiane* 2: 888 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Brosimum aubletii* Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher]* 2: 34 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838), nom. illeg. superfl. (based on *Piratinera guianensis*). — *Aliastrum guianense* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 623 [5 Nov. 1891] (Kuntze 1891).

Brosimum discolor Schott, *Syst. Veg. [Sprengel]* 4 (2): 403 [Jan.-June 1827] (Schott 1827). — *Piratinera discolor* (Schott) Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 100 [18 June 1918] (Pittier 1918).

Piratinera panamensis Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 100 [18 June 1918] (Pittier 1918). — *Brosimum panamense* (Pittier) Standl. & Steyermark, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23 (2): 40 [14 Feb. 1944] (Standley & Steyermark 1944).

Brosimum lecointei Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 28 (Ducke 1922), “*Le Cointei*”.

Piratinera scabridula S.F.Blake, *J. Wash. Acad. Sci.* 12: 397 (Blake 1922).

Piratinera velutina S.F.Blake, *J. Wash. Acad. Sci.* 12: 398 (Blake 1922). — *Brosimum velutinum* (S.F.Blake) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 1 (Ducke 1925).

Piratinera lemeei Benoist, *Bull. Mus. Natl. Hist. Nat.* 31: 468 (Benoist 1925). — *Brosimum lemeei* (Benoist) Lemée, *Fl. Guyane Franç.* 1: 510 (Lemée 1955).

Brosimum tessmannii Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 190 [20 Dec. 1927] (Mildbraed 1927).

Piratinera mollis Killip, *J. Wash. Acad. Sci.* 26: 358 (Killip 1936).

Brosimum palmarum Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 158 (Standley 1937).

Brosimum rotundatum Standl., *Bull. Torrey Bot. Club* 75 (3): 293 [May-June 1948] (Standley 1948).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: paig, paig-seine, pairi, pairi-priye, pairi-seinô • Ka: paidya, paidya tubulu, paila, tokolo abolimeli, tokolo

pailali • Te: paila • Wp: paila, paila wili • Wn: paila • Nt: bofo paaya, bofoo paaya, paaya, pendé paaya • Cr: bwa-lèt, bwa-lèt-moucheté • Fr: amourette, amourette tachetée, bois d'arc, bois de lettre moucheté • Br: muira-pinima, pau-rainha-roxo, pau-tartaruga.

HERBARIUM DATA (FG). — 94 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM, [BM000993312], P[P00089329], P-JU[P00089328]).

INVENTORY DATA (FG). — 401 trees in 142 plots; $F_{\max} = 4.6\%$; $\text{dbh}_{\text{inv}} = 70.6\text{ cm}$.

[1150] *Brosimum lactescens* (S.Moore) C.C.Berg

Acta Bot. Neerl. 19: 326 (Berg 1970). — *Brosimopsis lactescens* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 473 ["1894-96" publ. Dec. 1895] (Moore 1895).

Brosimopsis diandra S.F.Blake, *Proc. Biol. Soc. Washington* 35: 180 (Blake 1922).

Brosimopsis oblongifolia Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 31 (Ducke 1922).

Brosimopsis amplifolia Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 3 (Ducke 1925).

Brosimum ojoche Woodson, *Ann. Missouri Bot. Gard.* 47: 126 (Woodson 1960).

Brosimum belicense Lundell, *Wrightia* 3 (8): 166 (Lundell 1966).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: api, paila lá • Br: leiteira, muiratinga.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *A.H. Gentry et al. 63044*.

INVENTORY DATA (FG). — 31 trees in 12 plots; $F_{\max} = 2.6\%$; $\text{dbh}_{\text{inv}} = 79.1\text{ cm}$.

[1151] *Brosimum parinariooides* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 27 (Ducke 1922).

VERNACULAR NAMES. — Pa: amap • Te: ulukupi • Wp: amapakuwa, amapala, amapalá • Nt: dokali • Cr: mapa • Br: amapá-doce, amapá-rana, mururérana.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *S.A. Mori et al. 25711*.

INVENTORY DATA (FG). — 45 trees in 34 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 101.2\text{ cm}$.

[1152] *Brosimum rubescens* Taub.

Bot. Jahrb. Syst. 12 (Beibl. 27): 4 [24 June 1890] (Taubert 1890). — *Alicastrum rubescens* (Taub.) Taub., *Bot. Jahrb. Syst.* 15 (Beibl. 34): 16 [5 Apr. 1892] (Taubert 1892). — *Piratinera rubescens* (Taub.) Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 100 [18 June 1918] (Pittier 1918).

Ferolia guianensis Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 7 [Jun.-Dec. 1775] (Aublet 1775). — *Parinari guianensis* (Aubl.) Lemée, *Fl.*

Guyane Franç. 2: 25 (Lemée 1952), *nom. illeg. hom., non P. guyanensis* Fritsch (1890).

Brosimum paraense Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 67 (Huber 1910), "paraense". — *Piratinera paraensis* (Huber) Benoit, *Arch. Bot.* 5 (Mém. 1): 45 [27 Mar. 1933] (Benoit 1933).

Brosimum lanciferum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 24 (Ducke 1922). — *Piratinera lancifera* (Ducke) Benoit, *Arch. Bot.* 5 (Mém. 1): 46 [27 Mar. 1933] (Benoit 1933).

Brosimum angustifolium Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 25 (Ducke 1922).

Brosimum caloxylon Standl., *Trop. Woods* 17: 11 [Mar. 1929] (Standley 1929).

Brosimum platyneurum Ducke, *Arq. Serv. Florest.* 1: 4 (Ducke 1939).

Brosimum longistipulatum Ducke, *Trop. Woods* 90: 7 (Ducke 1947).

Brosimum brevipedunculatum Ducke, *Trop. Woods* 90: 8 (Ducke 1947).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: paig-purubumna, paig-puvemna, paig-seine, pairi-purubumna, pairi-puvemna, pairi-seinô, tubá-pairi • Ka: paidya tabile, tukusi pailali • Wp: api, paila lá • Wn: pailaimé • Nt: dokali, paaya • Cr: bwa-lèt-rouj, satiné-ribanné • Fr: bois de lettre rouge, bois satin (*fide* Aublet 1775), férole (*fide* Aublet 1775), satiné rubané • Br: garrote, muira-piranga, pau-rainha, pau-tartaruga.

HERBARIUM DATA (FG). — 67 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15162*.

INVENTORY DATA (FG). — 234 trees in 123 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 113\text{ cm}$.

[1153] *Brosimum utile* (Kunth) Oken
subsp. *ovatifolium* (Ducke) C.C.Berg

Acta Bot. Neerl. 19: 328 (Berg 1970). — *Brosimum ovatifolium* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 25 (Ducke 1922).

Brosimum rigidum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 27 (Ducke 1922).

Brosimum krukovii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 157 (Standley 1937).

Brosimum pallescens Ducke, *Arq. Serv. Florest.* 1: 3 (Ducke 1939).

NOTE. — *Brosimum utile* (Kunth) Oken is hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: ulukupi, ulukupi sili • Nt: dokali • Br: leiteira, turury.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2780*.

INVENTORY DATA (FG). — 62 trees in 50 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 114.5\text{ cm}$.

Genus *Clarisia* Ruiz & Pav.

[1154] *Clarisia ilicifolia* (Spreng.) Lanj. & Rossberg

Recueil Trav. Bot. Néerl. 33: 717 (Lanjouw & Rossberg 1936). — *Excoecaria ilicifolia* Spreng., *Neue Entdeck. Pflanzenk.* 2: 117 (Sprengel 1821). — *Acanthinophyllum ilicifolium* (Spreng.) W.C.Burger, *Ann. Missouri Bot. Gard.* 49: 27 (Burger 1962), “*ilicifolia*”.

Acanthinophyllum strepitans Allemão, Rev. Braz. 1: 369 (Allemão 1857). — *Sahagunia strepitans* (Allemão) Benth. & Hook., *Gen. Pl. [Bentham & Hooker f.]* 3 (1): 377 [7 Feb. 1880] (Bentham & Hooker 1880), in nota. — *Clarisia strepitans* (Allemão) Lanj., *Recueil Trav. Bot. Néerl.* 33: 272 (Lanjouw 1936).

Pseudosorocea poeppigii Baill., *Adansonia [Baillon]* 11: 297 (Baillon 1875).

Sahagunia racemifera Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 334 (Huber 1909).

Aliteria sagotii Benoist, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 1: 163 (Benoist 1929), “*Sagoti*”.

Sorocea stenophylla Standl., *Trop. Woods* 43: 18 (Standley 1935).

Clarisia strepitans var. *cuyunensis* Lanj., *Recueil Trav. Bot. Néerl.* 33: 275 (Lanjouw 1936). — *Clarisia ilicifolia* var. *cuyunensis* (Lanj.) Lanj. & Rossberg, *Recueil Trav. Bot. Néerl.* 33: 718 (Lanjouw & Rossberg 1936).

Clarisia strepitans var. *guianensis* Lanj., *Recueil Trav. Bot. Néerl.* 33: 274 (Lanjouw 1936). — *Clarisia ilicifolia* var. *guianensis* (Lanj.) Lanj. & Rossberg, *Recueil Trav. Bot. Néerl.* 33: 718 (Lanjouw & Rossberg 1936).

Clarisia strepitans var. *micranthera* Lanj., *Recueil Trav. Bot. Néerl.* 33: 275 (Lanjouw 1936). — *Clarisia ilicifolia* var. *micranthera* (Lanj.) Lanj. & Rossberg, *Recueil Trav. Bot. Néerl.* 33: 718 (Lanjouw & Rossberg 1936).

Clarisia strepitans var. *paraensis* Lanj., *Recueil Trav. Bot. Néerl.* 33: 275 (Lanjouw 1936). — *Clarisia ilicifolia* var. *paraensis* (Lanj.) Lanj. & Rossberg, *Recueil Trav. Bot. Néerl.* 33: 718 (Lanjouw & Rossberg 1936).

Clarisia spruceana Lanj., *Recueil Trav. Bot. Néerl.* 33: 272 (Lanjouw 1936). — *Acanthinophyllum spruceanum* (Lanj.) W.C.Burger, *Ann. Missouri Bot. Gard.* 49: 30 (Burger 1962), “*spruceana*”.

VERNACULAR NAMES. — Wp: ka'a polopi, yanita • Br: folha-de-serra, guaruba, janitá.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *M.-F. Prévost 1281*.

SIZE. — Up to 20 m tall (Berg 1992).

[1155] *Clarisia racemosa* Ruiz & Pav.

Syst. Veg. Fl. Peruv. Chil. 1: 255 [late Dec. 1798] (Ruiz & Pavón 1798).

Soaresia nitida Allemão, *Trab. Soc. Vellos.*: 72. (Allemão 1851). — *Sorocea nitida* (Allemão) Warb., *Bull. Soc. Bot. France* 59 (Mém. 3g): 644 [“1912” publ. 1913] (Warburg 1913). — *Clarisia nitida* (Allemão) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (1): 15 (Macbride 1931).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Br: guaríuba-amarela.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5012*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; dbh_{inv} = 25.6 cm.

Genus *Ficus* L.

[1156] *Ficus albert-smithii* Standl.
(Fig. 39B)

Lloydia 2 (3): 174 (Standley 1939).

Ficus frondosa Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 169 (Standley 1937), *nom. illeg. hom.*, *non* S.Moore (1923).

Ficus hypochrysea Dugand, *Caldasia* 7 (33): 228 (Dugand 1955).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier 3571*.

SIZE. — Up to 15 m tall (Berg 1999).

[1157] *Ficus amazonica* (Miq.) Miq.

Ann. Mus. Bot. Lugduno-Batavi 3 (10): 298 (Miquel 1867). — *Urostigma amazonica* Miq., *London J. Bot.* 6: 541 (Miquel 1847).

Urostigma angustifolium Miq., *London J. Bot.* 6: 539 (Miquel 1847). — *Ficus angustifolia* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867), *nom. illeg. hom.*, *non* Blume (1825).

Ficus surinamensis Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 219 (Miquel 1867).

NOTE. — Specimens at CAY are wrongly identified as *F. citrifolia* Mill.

VERNACULAR NAMES. — Wp: kwapo'i sōwī • Wn: halikwana • Nt: ankatu, katu, nkatu.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *M.-F. Prévost 1146*.

SIZE. — Up to 20 m tall (Berg 1992).

[1158] *Ficus americana* Aubl. subsp. *greiffiana* (Dugand)
C.C.Berg

Blumea 52 (2): 300 (Berg 2007). — *Ficus greiffiana* Dugand, *Caldasia* 1 (4): 49 (Dugand 1942).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3288*.

SIZE. — Up to 10 m tall (Berg 1992).

[1159] *Ficus americana* subsp. *guianensis* (Desv. ex Ham.) C.C.Berg

Blumea 52 (2): 300 (Berg 2007). — *Ficus guianensis* Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 62 [Oct. 1825] (Hamilton 1825).

Ficus clusiifolia Schott, *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 409 [Jan.-June 1827] (Schott 1827), “*clusiaeifolia*”. — *Urostigma clusiifolium* (Schott) Miq., *London J. Bot.* 6: 543 (Miquel 1847), “*clusiaeifolium*”.

Ficus splendens Kunth & C.D.Bouché, *Index Seminum (Berlin)*: 14 (Kunth & Bouché 1846-1847). — *Urostigma splendens* (Kunth & C.D.Bouché) Miq., *London J. Bot.* 6: 587 (Miquel 1847).

Ficus anacardiifolia Kunth & C.D.Bouché, *Index Seminum (Berlin)*: 15 (Kunth & Bouché 1846-1847). — *Urostigma anacardiifolium* (Kunth & C.D.Bouché) Miq., *London J. Bot.* 6: 587 (Miquel 1847).

Urostigma matthewsii Miq., *London J. Bot.* 6: 549 (Miquel 1847). — *Ficus matthewsii* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

Pharmacosycea parkeriana Miq., *London J. Bot.* 7: 71 (Miquel 1848). — *Ficus parkeriana* (Miq.) Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 227 [27 June 1932] (Sandwith 1932).

Ficus martinii Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 219 (Miquel 1867), “*martini*”.

Ficus umbonigera Warb., *Symb. Antill. [Urban]* 3 (3): 480 [1 May 1903] (Warburg 1903).

Ficus grenadensis Warb., *Symb. Antill. [Urban]* 3 (3): 481 [1 May 1903] (Warburg 1903).

Ficus ob lanceolata Rusby, *Bull. New York Bot. Gard.* 4 (14): 446 [7 Dec. 1907] (Rusby 1907).

Ficus gleasonii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 170 (Standley 1937).

Ficus sprucei Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 176 (Standley 1937).

Ficus myriasycea Pittier, *Bol. Soc. Venez. Ci. Nat.* 4 (30): 75 (Pittier 1937).

Ficus niceforoi Dugand, *Caldasia* 1 (4): 59 (Dugand 1942).

Ficus vaupesana Dugand, *Caldasia* 1 (4): 72 (Dugand 1942).

Ficus cor pulenta Pittier, *Bol. Soc. Venez. Ci. Nat.* 8: 257 (Pittier 1943).

Ficus metensis Dugand, *Caldasia* 2 (6): 77 (Dugand 1943).

Ficus chiribiquetensis Dugand, *Caldasia* 2 (9): 375 [6 Jan. 1944] (Dugand 1944).

Ficus maroana Pittier, *Bol. Soc. Venez. Ci. Nat.* 9: 120 (Pittier 1944).

Ficus erratica Standl., *Bull. Torrey Bot. Club* 75 (3): 295 [May-June 1948] (Standley 1948).

Ficus mensalis Standl., *Bull. Torrey Bot. Club* 75 (3): 297 [May-June 1948] (Standley 1948).

NOTE. — The original collection by Desvaux of *Ficus guianensis* Desv. ex Ham. was wrongly cited by Hamilton (1825: 62) as collected in Guyana. Augustin Nicaise Desvaux (1784-1856) only collected in French Guiana.

VERNACULAR NAMES. — Pa: muhu, muhu-kamwi • Wp: kwapo'i • Cr: bwa-fidjé • Br: apuí.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *A.N. Desvaux s.n.* (type P[P00756603]).

SIZE. — Up to 35 m tall (Berg 1999).

[1160] *Ficus americana* subsp. *subapiculata* (Miq.) C.C.Berg

Blumea 52 (2): 302 (Berg 2007).

Urostigma subapiculatum Miq., *Fl. Bras. [Martius]* 4 (1): 101 [1 Dec. 1853] (Miquel 1853). — *Ficus subapiculata* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

VERNACULAR NAMES. — Pa: muhu-puvemna.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori et al. 25427*.

SIZE. — Up to 35 m tall (Berg 1992).

[1161] *Ficus caballina* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 15: 81 (Standley 1936).

Ficus ramiflora Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 175 (Standley 1937).

Ficus tamatamae Pittier, *Bol. Soc. Venez. Ci. Nat.* 8: 260 (Pittier 1943).

VERNACULAR NAMES. — Ka: kuasimyan.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2317*.

SIZE. — Up to 25 m tall (Berg 1999).

[1162] *Ficus catappifolia* Kunth & C.D.Bouché

Index Seminum (Berlin) 14 (Kunth & Bouché 1846-1847). — *Urostigma catappifolium* (Kunth & C.D.Bouché) Miq., *London J. Bot.* 6: 533 (Miquel 1847), “*catappaefolium*”.

VERNACULAR NAMES. — Pa: muhu, muhu-kamwi • Wp: asīngau lemi moay • Cr: bwa-fidjé • Br: apuí.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *M.-F. Prévost 791*.

SIZE. — Up to 15 m tall (Berg 1992).

[1163] *Ficus cremersii* C.C.Berg

Acta Amazonica 14 (1-2, Suppl.): 199 [“1984” publ. 1986] (Berg 1986).

VERNACULAR NAMES. — Nt: dyankatu.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *G. Cremers 8233* (holo-, P[P00077201]; iso-, P[P00077202], U[U0004602]).

SIZE. — Brazil, Amazonas. *A.A. Oliveira 283* (MO), 20 m × 62 cm.




FIG. 39. — Moraceae: **A**, *Brosimum guianense* (Aubl.) Huber; **B**, *Ficus albert-smithii* Standl. (*D. Sabatier 5793*); **C**, *Ficus gomelleira* Kunth & C.D.Bouché; **D**, *Ficus trigona* L.f. (*J.-F. Molino & D. Sabatier 2267*). A-C, © *D. Sabatier/IRD*; D, © *J.-F. Molino/IRD*.

[1164] *Ficus donnell-smithii* Standl.

Contr. U.S. Natl. Herb. 20 (1): 21 [31 May 1917] (Standley 1917).

Ficus florenciana Dugand, *Caldasia* 1 (4): 44 (Dugand 1942).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 26542*.

SIZE. — Up to 25 m tall (Berg 1999).

[1165] *Ficus duartei* C.C.Berg & Carauta

Brittonia 54 (4): 240 ["2002" publ. 16 Apr. 2003] (Berg & Carauta 2003).

HERBARIUM DATA (FG). — A single collection, *M. Blanc 128*.

SIZE. — Up to 10 m tall (Berg & Carauta 2003: 241).

[1166] *Ficus duckeana* C.C.Berg & Ribeiro

Brittonia 54 (4): 241 ["2002" publ. 16 Apr. 2003] (Berg & Ribeiro 2003).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3426* (para-, CAY[CAY034924]).

SIZE. — Up to 20 m tall (Berg 1992).

[1167] *Ficus gomelleira* Kunth & C.D.Bouché
(Fig. 39C)

Index Seminum (Berlin): 18 (Kunth & Bouché 1846-1847). — *Urostigma gomelleira* (Kunth & C.D.Bouché) Miq., *London J. Bot.* 6: 531 (Miquel 1847).

Ficus holosericea Schott, *Syst. Vég. [Sprengel]* 4 (2, Cur. Post.): 410 [Jan.-June 1827] (Schott 1827).

Urostigma doliarium Miq., *London J. Bot.* 6: 527 (Miquel 1847). — *Ficus dolaria* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 297 (Miquel 1867). — *Ficus dolaria* Mart. ex Miq., *London J. Bot.* 6: 527 (Miquel 1847), *nom. nud. pro syn.*

Ficus acarouaniensis Benoist, *Arch. Bot. Bull. Mens.* 3: 170 (Benoist 1929).

VERNACULAR NAMES. — Wp: kwapo'i • Nt: ankatu, katu, nkatu • Cr: bwa-fidjé • Fr: figuier • Br: apuí-preto, gameleira.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *M.-F. Prévost 4523*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 210$ cm.

[1168] *Ficus hebetifolia* Dugand

Caldasia 1 (4): 50 (Dugand 1942).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P. Petronelli* 277.

SIZE. — Up to 20 m tall (Berg 1999).

[1169] *Ficus insipida* Willd. subsp. *scabra* C.C.Berg

Acta Amazonica 14 (1-2, Suppl.): 201 [“1984” publ. 1986] (Berg 1986).

VERNACULAR NAMES. — Pa: muhu • Ka: kuwasini, kwasini • Wp: kwasini • Cr: bwa-fidjé • Br: caxinguba, gameira-branca.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-321* (holo-, P[not seen]; iso-, P[P00076241], U[U.1426125]).

INVENTORY DATA (FG). — 8 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 150$ cm.

[1170] *Ficus krukovi* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2): 171 (Standley 1937).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost* 1472.

SIZE. — Up to 20 cm dbh (Standley 1937b).

[1171] *Ficus leiophylla* C.C.Berg

Acta Amazonica 14 (1-2, Suppl.): 205 (Berg 1986).

VERNACULAR NAMES. — Wp: kwapo'i sili • Nt: pikin katu.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *J.-J. de Granville & F. Kahn* 5410 (holo-, P[P00076239]; iso-, CAY[CAY024862, CAY024863], P[P00076240], U[U0004645, U0004646]).

SIZE. — Up to 30 m tall (Berg & Carauta 2003).

[1172] *Ficus maroniensis* Benoist

Bull. Mus. Natl. Hist. Nat., sér. 2, 1: 164 (Benoist 1929).

Ficus lanjouwii DeWolf, *J. Arnold Arbor.* 50 (3): 478 [15 July 1969] (DeWolf 1969).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *R. Benoist* 788 (holo-, P[P00089302]; iso-, MO[MO-204393], P[P00089303, P00089304]).

INVENTORY DATA (FG). — 1 tree, dbh = 16.1 cm.

[1173] *Ficus maxima* Mill.

Gard. Dict., ed. 8, n. 6 [16 Apr. 1768] (Miller 1768).

Ficus citrifolia Lam., *Encycl. [J. Lamarck et al.]* 2 (2): 494 [14 Apr. 1788] (Lamarck 1788), *nom. illeg. hom., non* Mill. (Miller 1768).

Ficus laurifolia Hort. ex Lam., *Encycl. [J. Lamarck et al.]* 2 (2): 495 [14 Apr. 1788] (Lamarck 1788). — *Urostigma laurifolium* (Hort. ex Lam.) Miq., *London J. Bot.* 6: 539 (Miquel 1847).

Ficus radula Humb. & Bonpl. ex Willd., *Sp. Pl.*, ed. 4 4 (2): 1144 [Apr. 1806] (Willdenow 1806). — *Pharmacosycea radula* (Humb. & Bonpl. ex Willd.) Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 330 (Liebmann 1851).

Ficus anthoneminthica DC., *Essai Propri. Méd. Pl.*, ed. 2, 267 (Candolle 1816), *nom. nud.*

Pharmacosycea guyanensis Miq., *London J. Bot.* 7: 67 (Miquel 1848).

Pharmacosycea grandaeva Miq., *London J. Bot.* 7: 70 (Miquel 1848). — *Ficus grandaeva* Mart. ex Miq., *London J. Bot.* 7: 70 (Miquel 1848), *nom. nud. pro syn.*

Pharmacosycea glaucescens Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 332 (Liebmann 1851). — *Ficus glaucescens* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 300 (Miquel 1867).

Pharmacosycea hernandezii Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 332 (Liebmann 1851). — *Ficus hernandezii* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 300 (Miquel 1867).

Pharmacosycea rigida Miq., *Bot. Voy. Herald [Seemann]* 5: 195 [July 1854] (Miquel 1854).

Urostigma protensum Griseb., *Bonplandia* 6 (1): 4 (Grisebach 1858). — *Ficus protensa* (Griseb.) Hemsl., *Biol. Cent.-Amer., Bot.* 3 (15): 147 [Jan. 1883] (Hemsley 1883).

Pharmacosycea pseudoradula Miq., *Verslagen Meded. Afd. Natuurk. Kon. Akad. Wetensch.* 13: 414 (Miquel 1862). — *Ficus pseudoradula* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 299 (Miquel 1867), “pseudo-radula”.

Pharmacosycea mexicana Miq., *Verslagen Meded. Afd. Natuurk. Kon. Akad. Wetensch.* 13: 415 (Miquel 1862). — *Ficus mexicana* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 299 (Miquel 1867).

Ficus coybana Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 300 (Miquel 1867).

Ficus parkeri Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 300 (Miquel 1867).

Ficus guadalajarana S.Watson, *Proc. Amer. Acad. Arts* 26: 151 (Watson 1891).

Ficus populnea Willd. f. *citrifolia* Warb., *Symb. Antill. [Urban]* 3 (3): 479 [1 May 1903] (Warburg 1903).

Ficus picardae Warb., *Symb. Antill. [Urban]* 3 (3): 484 [1 May 1903] (Warburg 1903).

Ficus subscabrida Warb., *Symb. Antill. [Urban]* 3 (3): 485 [1 May 1903] (Warburg 1903).

Ficus rubricosta Warb., *Symb. Antill. [Urban]* 3 (3): 486 [1 May 1903] (Warburg 1903).

Ficus plumieri Urb., *Repert. Spec. Nov. Regini Veg.* 15: 158 [31 May 1918] (Urban 1918), “Plumerii”.

Ficus bopiana Rusby, *Mem. New York Bot. Gard.* 7: 230 (Rusby 1927).

Ficus ulei Rossberg, *Repert. Spec. Nov. Regni Veg.* 42: 60 (Rossberg 1937).

Ficus sodiroi Rossberg, *Repert. Spec. Nov. Regni Veg.* 42: 61 (Rossberg 1937).

Ficus murilloi Dugand, *Caldasia* 1 (4): 57 (Dugand 1942).

Ficus vicencionis Dugand, *Caldasia* 2 (9): 385 [6 Jan. 1944] (Dugand 1944).

Ficus murilloi var. *cajambrensis* Dugand, *Caldasia* 4 (17): 117 (Dugand 1946).

Ficus chaconiana Standl. & L.O.Williams, *Ceiba* 3 (2): 111 [1 Oct. 1952] (Standley & Williams 1952), “*Choconiana*”.

NOTES. — The epithet *chaconiana* honours Jaime Chacón P. The epithet *vicencionis* is not to be corrected, as it results from an intentional latinization of the locality name “(Villa)vicencio” (Turland et al. 2018: Art. 60.9).

VERNACULAR NAMES. — Pa: muhu • Ka: ityulutano kuwasini, kwasini • Wp: kwasini • Nt: dyankatu • Cr: bwa-fidjé • Fr: figuier • Br: caxinguba, gameleira-branca.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *M.-F. Prévost 4194*.

INVENTORY DATA (FG). — 4 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 70\text{ cm}$.

[1174] *Ficus nympaeifolia* Mill.

Gard. Dict., ed. 8, n. 9 [16 Apr. 1768] (Miller 1768), “*Nympaeifolia*”. — *Urostigma nympaeifolium* (Mill.) Miq., *London J. Bot.* 6: 527 (Miquel 1847).

Ficus nympoides Thunb., *Ficus*: 7 [21 Dec. 1786] (Thunberg 1786).

Urostigma cyclophyllum Miq., *Fl. Bras. [Martius]* 4 (1): 91 [1 Dec. 1853] (Miquel 1853). — *Ficus cyclophylla* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 297 (Miquel 1867).

Ficus urbaniana Warb., *Symb. Antill. [Urban]* 3 (3): 459 [1 May 1903] (Warburg 1903). — *Ficus involuta* (Liebm.) Miq. var. *urbaniana* (Warb.) Dugand, *Caldasia* 2 (8): 275 (Dugand 1943).

Ficus ierensis Britton, *Bull. Torrey Bot. Club* 48 (12): 329 [“1921 publ. 1922”] (Britton 1922).

Ficus anguina Benoist, *Bull. Mus. Natl. Hist. Nat.* 30: 104 (Benoist 1924).

Ficus cabusana Standl. & Steyermark, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 22 (4): 226 (Standley & Steyermark 1940).

Ficus duquei Dugand, *Caldasia* 1 (4): 42 (Dugand 1942).

Ficus duquei var. *obtusiloba* Dugand, *Mutisia* 9: 6 (Dugand 1952).

VERNACULAR NAMES. — Pa: muhu-ivine • Ka: kuasini, kwašini • Te: kwatsindi • Wp: asīngau lemi moay • Nt: ankatu, katu, nkatu • Cr: bwa-fidjé • Fr: figuier étrangleur, figuier blanc • Br: apuí, caxinguba.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *D. Sabatier 2246*.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 180\text{ cm}$.

[1175] *Ficus pakkensis* Standl.

Bull. Torrey Bot. Club 75 (3): 297 [May-June 1948] (Standley 1948).

VERNACULAR NAMES. — Ka: kuasini, kwašini • Wp: asīngau lemi moay sili.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *M.-F. Prévost 854*.

SIZE. — Up to 15 m tall (Berg 1999).

[1176] *Ficus paludica* Standl.

Bull. Torrey Bot. Club 75 (3): 298 [May-June 1948] (Standley 1948).

Urostigma leucostictum Miq., *London J. Bot.* 6: 535 (Miquel 1847). — *Ficus leucosticta* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 297 (Miquel 1867), nom. illeg. hom., non Spreng. (Sprengel 1826).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P.A. Sagot s.n.* (P[06753650]).

SIZE. — Up to 20 m tall (Berg 1999).

[1177] *Ficus panurensis* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2): 174 (Standley 1937).

Ficus arctocarpa Standl., *Bull. Torrey Bot. Club* 75 (3): 294 [May-June 1948] (Standley 1948).

Ficus maguirei Standl., *Bull. Torrey Bot. Club* 75 (3): 296 [May-June 1948] (Standley 1948).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville B-3724*.

INVENTORY DATA (FG). — 1 tree, $dbh = 19.9\text{ cm}$.

[1178] *Ficus paraensis* (Miq.) Miq.

Ann. Mus. Bot. Lugduno-Batavi 3 (10): 298 (Miquel 1867). — *Urostigma paraense* Miq., *London J. Bot.* 6: 534 (Miquel 1847).

Ficus myrmecophila Warb., in *Vegetationsbilder [G. Karsten & H. Schenck]*, ser. 3, 1: t. 2 (Warburg 1905).

Urostigma williamsii Rusby, *Bull. New York Bot. Gard.* 6 (22): 499 [30 Nov. 1910] (Rusby 1910). — *Ficus williamsii* (Rusby) Rossberg, *Repert. Spec. Nov. Regni Veg.* 42: 61 (Rossberg 1937), nom. illeg. hom., non C.B.Rob. (Robinson 1908).

Ficus panamensis Standl., *Contr. U.S. Natl. Herb.* 20 (1): 15 [31 May 1917] (Standley 1917).

Ficus thelephora Benoist, *Arch. Bot. Bull. Mens.* 3: 171 (Benoist 1929).

Ficus haughtii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 170 (Standley 1937).

Ficus uberrima Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 177 (Standley 1937).

Ficus putumayonis Dugand, *Caldasia* 1 (4): 62 (Dugand 1942).

- Ficus hydrophila* Pittier, *Bol. Soc. Venez. Ci. Nat.* 8: 259 (Pittier 1943).
Ficus orinocensis Pittier, *Bol. Soc. Venez. Ci. Nat.* 8: 259 (Pittier 1943).
Ficus peroblonga Dugand, *Caldasia* 4 (17): 118 (Dugand 1946).
Ficus arukensis Standl., *Bull. Torrey Bot. Club* 75 (3): 295 [May–June 1948] (Standley 1948).

NOTE. — The epithet *putumayonis*, which results from an intentional latinization of the geographic name “Putumayo”, is not to be corrected (Turland et al. 2018: Art. 60.8).

VERNACULAR NAMES. — Pa: muhu-ivine • Ka: kuwasimyan • Wp: asíngau lemi moay, kwapo'i, kwapo'i sówí • Wn: kana halale, luwi • Cr: bwa-fidjé • Br: apuí.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 976.

SIZE. — Up to 20 m tall (Berg 1999).

[1179] *Ficus pertusa* L.f.

- Suppl. Pl.*: 442 [“1781” publ. Apr. 1782] (Linnaeus 1782). — *Urostigma pertusum* (L.f.) Miq., *London J. Bot.* 6: 548 (Miquel 1847).
Ficus padifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 47 [28 Apr. 1817] (Kunth 1817). — *Urostigma padifolium* (Kunth) Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 324 (Liebmann 1851).
Ficus complicata Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 48 [28 Apr. 1817] (Kunth 1817). — *Urostigma complicatum* (Kunth) Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 325 (Liebmann 1851).
Ficus myrtifolia Link, *Enum. Hort. Berol. Alt.* 2: 450 [Jan.–June 1822] (Link 1822).

Ficus lancifolia Hook. & Arn., *Bot. Beechey Voy.* 310 [“1841” publ. Dec. 1838] (Hooker & Arnott 1838). — *Urostigma lancifolium* (Hook. & Arn.) Miq., *London J. Bot.* 6: 538 (Miquel 1847).

Ficus subtriplinervia Mart., *Flora* 24 (2, Beibl.): 67 (Martius 1841). — *Urostigma subtriplinervium* (Mart.) Miq., *London J. Bot.* 6: 542 (Miquel 1847).

Ficus radicans Casar., *Atti Riunione Sci. Ital.* 3: 515 [“1841” publ. June 1842] (Casaretto 1842), nom. illeg. hom., non Desf. (Desfontaines 1829).

Ficus arpazusa Casar., *Nov. Stirp. Bras.* 1: 15 [Oct. (“May”) 1842] (Casaretto 1842).

Ficus planicostata Kunth & C.D.Bouché, *Index Seminum (Berlin)* 16 (Kunth & Bouché 1846–1847). — *Ficus populnea* f. *planicostata* (Kunth & C.D.Bouché) Warb., *Symb. Antill. [Urban]* 3 (3): 479 [1 May 1903] (Warburg 1903).

Urostigma populneum (Willd.) Miq. f. *mexicanum* Miq., *London J. Bot.* 6: 537 (Miquel 1847), “Mexicana”.

Urostigma schiedeanum Miq., *London J. Bot.* 6: 539 (Miquel 1847).

Urostigma erythrostictum Miq., *London J. Bot.* 6: 540 (Miquel 1847). — *Ficus erythrostictia* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

Urostigma cestrifolium (Schott) Miq. f. *major* Miq., *London J. Bot.* 6: 541 (Miquel 1847).

Urostigma geminum Miq., *London J. Bot.* 6: 547 (Miquel 1847). — *Ficus gemina* Ruiz ex Miq., *London J. Bot.* 6: 547 (Miquel 1847), nom. nud. pro syn. — *Ficus gemina* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

Pharmacosycea laurifolia Miq., *London J. Bot.* 7: 71 (Miquel 1848).

Pharmacosycea peruviana Miq., *London J. Bot.* 7: 72 (Miquel 1848). — *Ficus peruviana* (Miq.) Rossberg, *Repert. Spec. Nov. Regni Veg.* 42: 61 (Rossberg 1937).

Urostigma baccatum Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 327 (Liebmann 1851). — *Ficus baccata* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 299 (Miquel 1867).

Urostigma sapidum Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 327 (Liebmann 1851). — *Ficus sapida* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

Urostigma turbinatum Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 328 (Liebmann 1851). — *Ficus turbinata* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867), nom. illeg. hom., non Willd. (Willdenow 1806).

Urostigma rolanderi Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.*, ser. 5, 2: 329 (Liebmann 1851). — *Ficus rolanderi* (Liebm.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867), “Rolandri”.

Urostigma subtriplinervium f. *major* Miq., *Fl. Bras. [Martius]* 4 (1): 99 [1 Dec. 1853] (Miquel 1853). — *Ficus prinoides* Humb. & Bonpl. ex Willd. var. *subtriplinervia* (Mart.) Kuntze, *Revis. Gen. Pl.* 3 (3): 294 [28 Sep. 1898] (Kuntze 1898), “prinoides”.

Ficus suffocans Griseb., *Fl. Brit. W.I. [Grisebach]* 150 [June 1859] (Grisebach 1859).

Ficus ochroleuca Griseb., *Fl. Brit. W.I. [Grisebach]* 151 [June 1859] (Grisebach 1859).

Urostigma sulcipes Miq., *Verslagen Meded. Afd. Natuurk. Kon. Akad. Wetensch.* 13: 413 (Miquel 1862). — *Ficus sulcipes* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 298 (Miquel 1867).

Ficus daphniphylla Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 300 (Miquel 1867).

Ficus fasciculata S.Watson, *Proc. Amer. Acad. Arts* 24: 78 (Watson 1889), nom. illeg. hom., non Benth. (Bentham 1873) nec King (1888).

Ficus sonorae S.Watson, *Proc. Amer. Acad. Arts* 24: 78 (Watson 1889).

Ficus grabhamii Britton, *Fl. Jamaica [Fawcett & Rendle]* 3: 51 [Dec. 1914] (Britton 1914).

Ficus halliana Britton, *Fl. Jamaica [Fawcett & Rendle]* 3: 51 [Dec. 1914] (Britton 1914).

Ficus morantensis Britton, *Fl. Jamaica [Fawcett & Rendle]* 3: 51 [Dec. 1914] (Britton 1914).

Urostigma subtriplinervium f. *ellipticum* Herzog, *Meded. Rijks-Herb.* 27: 75 (Herzog 1915). — *Ficus elliptica* (Herzog) Herter, *Revista Sudamer. Bot.* 6: 151 (Herter 1940).

Ficus broadwayi Urb., *Repert. Spec. Nov. Regni Veg.* 15: 110 [31 Dec. 1917] (Urban 1917).

Ficus mendelsonii Britton, *Bull. Torrey Bot. Club* 48 (12): 330 [“1921” publ. 1922] (Britton 1922).

Ficus tapajozensis Standl., *Trop. Woods* 33: 11 (Standley 1933).

Ficus palmicida Pittier, *Bol. Soc. Venez. Ci. Nat.* 4 (30): 69 (Pittier 1937).

Ficus kanukuensis Standl., *Lloydia* 2 (3): 174 (Standley 1939).

Ficus garcesii Dugand, *Caldasia* 1 (4): 45 (Dugand 1942).

Ficus trachelosyce Dugand, *Caldasia* 1 (4): 69 (Dugand 1942).

Ficus savannarum Standl., *Bull. Torrey Bot. Club* 75 (3): 298 [May-June 1948] (Standley 1948).

VERNACULAR NAMES. — Br: apuí.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2007*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 35.3$ cm.

[1180] *Ficus piresiana* Vázq. Avila & C.C.Berg

Acta Amazonica 14 (1-2, Suppl.): 207 [“1984” publ. 1986] (Vázquez Avila & Berg 1986).

VERNACULAR NAMES. — Wn: pêtélimé.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost 1291* (para-, CAY[CAY024874]).

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 76.4$ cm.

[1181] *Ficus popenoei* Standl.
subsp. *malacocarpa* (Standl.) C.C.Berg

Blumea 52 (3): 583 (Berg 2007). — *Ficus malacocarpa* Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 172 (Standley 1937).

Ficus scabrida Pittier, *Bol. Soc. Venez. Ci. Nat.* 4 (30): 68 (Pittier 1937).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2687*.

SIZE. — Up to 10 m tall (Berg 1999).

[1182] *Ficus schumacheri* (Liebm.) Griseb.

Fl. Brit. W.I. [Grisebach] 151 [June 1859] (Grisebach 1859). — *Urostigma schumacheri* Liebm., *Kongel. Danske Vidensk. Selsk. Skr. Naturvidensk. Math. Afđ.*, ser. 5, 2: 328 (Liebmann 1851).

VERNACULAR NAMES. — Wp: kwapo'i sili • Nt: bunsumiki katu, pikin katu.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2254*.

INVENTORY DATA (FG). — 22 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.3$ cm.

[1183] *Ficus trigona* L.f.
(Fig. 39D)

Suppl. Pl.: 441 [“1781” publ. Apr. 1782] (Linnaeus 1782). — *Oluntos trigona* (L.f.) Raf., *Sylva Tellur.*: 58 (Rafinesque 1838), “*trigono*”. — *Urostigma trigonum* (L.f.) Miq., *London J. Bot.* 6: 548 (Miquel 1847).

Urostigma fagifolium Miq., *London J. Bot.* 6: 544 (Miquel 1847). — *Ficus fagifolia* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 299 (Miquel 1867).

Urostigma euomphalum Miq., *Fl. Bras. [Martius]* 4 (1): 103 [1 Dec. 1853] (Miquel 1853). — *Ficus euomphala* (Miq.) Miq., *Ann. Mus. Bot. Lugduno-Batavi* 3 (10): 299 (Miquel 1867).

Urostigma costatum Rusby, *Bull. New York Bot. Gard.* 6 (22): 499 [30 Nov. 1910] (Rusby 1910), “*costata*”.

Ficus vulpina Benoist, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 1: 164 (Benoist 1929).

Ficus llewelynii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.2): 304 [15 Mar. 1937] (Standley 1937), “*Llewelyni*”.

Ficus weberbaueri Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.2): 308 [15 Mar. 1937] (Standley 1937).

Ficus regularis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 175 (Standley 1937).

Ficus wuiiana Rossberg, *Repert. Spec. Nov. Regni Veg.* 42: 61 (Rossberg 1937).

Ficus alvareziana Dugand, *Caldasia* 1 (4): 29 (Dugand 1942).

Ficus llanensis Dugand, *Caldasia* 1 (4): 53 (Dugand 1942).

Ficus plicato-ostiolata Pittier, *Bol. Soc. Venez. Ci. Nat.* 8: 258 (Pittier 1943).

Ficus ajahuensis Dugand, *Caldasia* 2 (10): 439 (Dugand 1944).

Ficus fanshawei Standl., *Bull. Torrey Bot. Club* 75 (3): 295 [May-June 1948] (Standley 1948).

Ficus juruensis Warb. ex Dugand, *Caldasia* 7 (33): 231 (Dugand 1955).

NOTE. — “*Ficus juruensis* Warb.” appears as a *nomen nudum* in Ule (1907: 140), and was invalidly published (no latin description) by Standley (1937a: 303).

VERNACULAR NAMES. — Wp: kwapo'i, kwapo'i sili, kwapo'i sówí.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2267*, $dbh = 35$ cm.

Genus *Helicostylis* Trécul

[1184] *Helicostylis pedunculata* Benoist

Bull. Mus. Natl. Hist. Nat. 25: 298 (Benoist 1919).

VERNACULAR NAMES. — Wn: kaikui amoman, tapatangman • Nt: pulo paaya.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *R. Benoit 667* (original material P[P06823043, P00745498, P00745499]).

INVENTORY DATA (FG). — 108 trees in 68 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 53.8$ cm.

[1185] *Helicostylis tomentosa* (Poepp. & Endl.) Rusby

Mem. Torrey Bot. Club 6 (1): 120 (Rusby 1896). — *Olmedia tomentosa* Poepp. & Endl., *Nova genera ac species plantarum [Poep-pig & Endlicher]* 2: 32 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838).

Olmedia poeppigiana Mart., *Flora* 24 (2, Beibl.): 93 (Martius 1841), “*Pöppigiana*”. — *Helicostylis poeppigiana* (Mart.) Trécul, *Ann. Sci. Nat., Bot. sér. 3, 8:* 134 (Trécul 1847). — *Greeneina poeppigiana* (Mart.) Kuntze, *Revis. Gen. Pl.* 2: 628 [5 Nov. 1891] (Kuntze 1891).

Helicostylis affinis Miq., *Fl. Bras. [Martius]* 4 (1): 118 [1 Dec. 1853] (Miquel 1853). — *Greeneina affinis* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 628 [5 Nov. 1891] (Kuntze 1891).

Helicostylis obtusifolia Standl., *Bull. Torrey Bot. Club* 58 (6): 356 [June 1931] (Standley 1931).

Olmedia polyccephala Pittier, *Bol. Soc. Venez. Ci. Nat.* 7 (51): 304 (Pittier 1942).

Helicostylis podogyne Ducke, *Bol. Técn. Inst. Agron. N.* 4: 3 [31 Mar. 1945] (Ducke 1945).

Helicostylis poeppigiana var. *macrophylla* Trécul, *Ann. Sci. Nat., Bot. sér. 3, 8:* 135 (Trécul 1847).

Olmedia asperula Standl., *Bull. Torrey Bot. Club* 75 (3): 299 [May-June 1948] (Standley 1948).

Helicostylis duckei A.D.Hawkes, *Phytologia* 3 (1): 31 [30 Aug. 1948] (Hawkes 1948).

Olmedia affinis Steud. ex Miq., *Fl. Bras. [Martius]* 4 (1): 118 [1 Dec. 1853] (Miquel 1853), *nom. nud. pro syn.*

Trymatococcus guanabarinus Duarte, *Rodriguésia* 23-24 (35-36): 55 (Duarte 1961), “*Trimatococcus*”.

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: paig-seine, pairi-seine, pairi-seinó, tukwanru • Ka: ombatapo, umpatapu • Wp: iwa pe • Nt: pulo paaya, puyo • Cr: boundou • Br: inháre, inháre-paina, mao-de-gato.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3944.

INVENTORY DATA (FG). — 118 trees in 78 plots; $F_{\max} = 2.1\%$; $dbh_{inv} = 56.7$ cm.

[1186] *Helicostylis* sp. A

NOTE. — A species similar to *H. elegans* (J.F.Macbr.) C.C.Berg.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino* 5625.

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 24$ cm.

Genus *Maquira* Aubl.

[1187] *Maquira calophylla* (Poepp. & Endl.) C.C.Berg

Acta Bot. Neerl. 18 (3): 464 (Berg 1969). — *Olmedia calophylla* Poepp. & Endl., *Nova genera ac species plantarum [Poep-pig & Endlicher]* 2: 32 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838). — *Perebea*

calophylla (Poepp. & Endl.) Benth. ex Pittier, *Contr. U.S. Natl. Herb.* 13: 439 (Pittier 1912). — *Olmelioperebea calophylla* (Poepp. & Endl.) Ducke, *Arq. Serv. Florest.* 1: 14 (Ducke 1939).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier et al.* 5161.

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{inv} = 145$ cm.

[1188] *Maquira guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 36 [Jun.-Dec. 1775] (Aublet 1775). — *Olmedia guianensis* (Aubl.) Trécul, *Ann. Sci. Nat., Bot. sér. 3, 8:* 129 (Trécul 1847). — *Olmedia maquira* Steud., *Nomencl. Bot. [Steudel], ed. 2, 2:* 210 (Steudel 1841), *nom. illeg. superfl.* (based on *Maquira guianensis*).

Perebea laurifolia Trécul, *Ann. Sci. Nat., Bot. sér. 3, 8:* 133 (Trécul 1847).

VERNACULAR NAMES. — Pa: mpitit-washiune • Wp: inámu sī sili, yanu'i pilá, yanu'i yowa, yawasi pita • Nt: man puyo • Br: arirana, muiratinga-de-folha-mitúda.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000993316]).

INVENTORY DATA (FG). — 173 trees in 98 plots; $F_{\max} = 1.7\%$; $dbh_{inv} = 57.5$ cm.

[1189] *Maquira sclerophylla* (Ducke) C.C.Berg

Acta Bot. Neerl. 18 (3): 463 (Berg 1969). — *Olmedioperebea sclerophylla* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 34 (Ducke 1922).

Perebea xinguana Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2):* 180 (Standley 1937).

VERNACULAR NAMES. — Br: pau-de-índio, rapé-de-índio.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3414.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 14.6$ cm.

[1190] *Maquira* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier* 2496.

INVENTORY DATA (FG). — 1 tree, $dbh = 47.8$ cm.

Genus *Naucleopsis* Miq.

[1191] *Naucleopsis guianensis* (Mildbr.) C.C.Berg

Acta Bot. Neerl. 18 (3): 465 (Berg 1969). — *Ogcodeia guianensis* Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 422 [30 Mar. 1932] (Mildbraed 1932).

VERNACULAR NAMES. — Pa: tukwanru • Wp: wa'i.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3692*.

INVENTORY DATA (FG). — 84 trees in 53 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 23.1$ cm.

Genus *Perebea* Aubl.

[1192] *Perebea guianensis* Aubl.
(Fig. 40A)

Hist. Pl. Guiane 2: 953 [Jun.-Dec. 1775] (Aublet 1775).

Olmedia grandifolia Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 128 (Trécul 1847).

Castilla australis Hemsl., *Hooker's Icon. Pl.* 27 [ser. 4, 7]: t. 2676 [Feb. 1901] (Hemsley 1901), “*Castilloa*”. — *Perebea australis* (Hemsl.) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (1): 17 (Macbride 1931).

Olmedia habas Pax, *Repert. Spec. Nov. Regni Veg.* 7: 108 (Pax 1909).

VERNACULAR NAMES. — Pa: amap-purubumna, tukwanru-kamwi • Ka: ombatapo, umpatapu • Wp: yiwa piso, yuwa piso • Nt: abeemu, pulo paaya • Cr: abérémou • Br: cauchorana, caxinguba-rana.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material not traced); *C. Sastre 6178*.

INVENTORY DATA (FG). — 48 trees in 19 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 22$ cm.

[1193] *Perebea rubra* (Trécul) C.C.Berg

Fl. Ecuador 60: 87 (Berg 1998). — *Noyera rubra* Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 136 (Trécul 1847). — *Perebea mollis* (Poepp. & Endl.) Huber subsp. *rubra* (Trécul) C.C.Berg, *Acta Bot. Neerl.* 18 (3): 463 (Berg 1969).

VERNACULAR NAMES. — Wp: iwa pe • Wn: kaikui amoman, tapa-tangman • Nt: abeemu, paaya, pulo paaya • Cr: abérémou.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J. Martin s.n.* (n° 114 at G) (holo-, P[P00745490]; iso-, B[B 10 0244527], F[F0064544F], FI[FI011716, FI011717, FI011718], G[G00438407], P[P00745491, P00745492], U[U0004748]).

INVENTORY DATA (FG). — 31 trees in 27 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 56$ cm.

Genus *Poulsenia* Eggers

[1194] *Poulsenia* sp. A

HERBARIUM DATA (FG). — A single collection, *D. Loubry 2097*.

SIZE. — Up to 25 m tall (Berg 1992).

Genus *Pseudolmedia* Trécul

[1195] *Pseudolmedia laevigata* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 131 (Trécul 1847).

Pseudolmedia guaranitica Hassl., *Annuaire Conserv. Jard. Bot. Genève* 21: 121 (Hassler 1919).

Pseudolmedia mildbraedii J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (2): 61 (Macbride 1931).

Pseudolmedia brosimifolia Ducke, *Arg. Serv. Florest.* 1: 8 (Ducke 1939).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Pa: paig-priye, pairi-priye.

HERBARIUM DATA (FG). — A single collection, *S.A. Mori & J.J. Pipoly 15475*.

SIZE. — Up to 35 m tall (Berg 1972).

[1196] *Pseudolmedia laevis* (Ruiz & Pav.) J.F.Macbr.

Publ. Field Mus. Nat. Hist., Bot. Ser. 11 (1): 16 (Macbride 1931). — *Olmedia laevis* Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 258 [late Dec. 1798] (Ruiz & Pavón 1798).

Olmedia ferruginea Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher]* 2: 31 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838). — *Pseudolmedia ferruginea* (Poepp. & Endl.) Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 131 (Trécul 1847).

Olmediopsis lanceolata H.Karst., *Fl. Columb. [H. Karsten]* 2 (1): 17 [15 Dec. 1862] (Karsten 1862).

Olmediopsis obliqua H.Karst., *Fl. Columb. [H. Karsten]* 2 (1): 17 [15 Dec. 1862] (Karsten 1862). — *Pseudolmedia obliqua* (H.Karst.) Renner, *Bot. Jahrb. Syst.* 39 (3-4): 374 [15 Jan. 1907] (Renner 1907).

Pseudolmedia hirsuta Baill., *Adansonia [Baillon]* 11: 295 (Baillon 1875).

Pseudolmedia sagotii Benoist, *Bull. Mus. Natl. Hist. Nat.* 31: 468 (Benoist 1925), “*Sagoti*”.

Pseudolmedia hirtellifolia Rusby, *Mem. New York Bot. Gard.* 7: 228 (Rusby 1927), “*hirtellaefolia*”.

Pseudolmedia alnifolia Rusby, *Mem. New York Bot. Gard.* 7: 229 (Rusby 1927).

Pseudolmedia multinervis Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 189 [20 Dec. 1927] (Mildbraed 1927).

Olmedia caurensis Pittier, *Bol. Soc. Venez. Ci. Nat.* 7 (51): 306 (Pittier 1942).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: paig-seine, pairi-seinō, tukwanru • Ka: tonolo ipyo, tonolo polipyo • Wp: api, paila lá • Nt: puyo • Br: inharé-folha-miúda, pamá.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2800*.

INVENTORY DATA (FG). — 123 trees in 55 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 39.8$ cm.

Genus *Sorocea* A.St.-Hil.

[1197] *Sorocea muriculata* Miq.
subsp. *uaupensis* (Baill.) C.C.Berg

Proc. Kon. Ned. Akad. Wetensch. C 88 (4): 387 (Berg 1985). — *Pseudosorocea uaupensis* Baill., *Adansonia [Baillon]* 11: 297 (Baillon 1875). — *Sorocea uaupensis* (Baill.) J.F.Macbr., *Candollea* 5: 348 (Macbride 1934).

Sorocea guayanensis W.C.Burger, *Acta Bot. Neerl.* 11 (4): 439 (Burger 1962).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *C. Feuillet* 715.

SIZE. — Up to 12 m tall (Berg 1999).

Genus *Trymatococcus* Poepp. & Endl.

[1198] *Trymatococcus amazonicus* Poepp. & Endl.
(Fig. 40B)

Nova genera ac species plantarum [Poeppig & Endlicher] 2: 30 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838).

Lanessania turbinata Spruce ex Baill., *Adansonia [Baillon]* 11: 298 (Baillon 1875). — *Trymatococcus turbinatus* (Spruce ex Baill.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 23 (Ducke 1922).

Trymatococcus paraensis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 22 (Ducke 1922).

VERNACULAR NAMES. — Pa: paig-seine, pairi-seinō • Ka: yuku'ipyo • Wp: api, paila lá, yanu'i u, yanu'i wate'e • Cr: bwa-lèt-blanc.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *D. Sabatier* & J.-F. Molino 5204.

INVENTORY DATA (FG). — 58 trees in 44 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43$ cm.

[1199] *Trymatococcus oligandrus* (Benoist) Lanj.

Recueil Trav. Bot. Néerl. 32: 271 (Lanjouw 1935). — *Lanessania oligandra* Benoist, *Bull. Mus. Natl. Hist. Nat.* 27: 199 (Benoist 1921).

VERNACULAR NAMES. — Pa: paig, paig-seine, pairi, pairi-seinō • Ka: yuku'ipyo • Wp: api, paila lá, yanu'i u, yanu'i wate'e • Nt: yenge pao • Cr: bwa-lèt-blanc.

HERBARIUM DATA (FG). — 62 collections at CAY. Sel. exs.: R. Benoist 1573 (holo-, P[P00710537]; iso-, P[P00710538, P00710539]).

INVENTORY DATA (FG). — 281 trees in 119 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 40.7$ cm.

Family MYRISTICACEAE R.Br.
Genus *Compsonera* (A.DC.) Warb.

[1200] *Compsonera ulei* Warb.
(Fig. 40C)

Verh. Bot. Vereins Prov. Brandenburg 47: 136 [1 Oct. 1905] (Warburg 1905).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier* & P. Birnbaum 4608.

SIZE. — Up to 12 m tall (Mitchell 2002).

Genus *Iryanthera* (A.DC.) Warb.

[1201] *Iryanthera hostmannii* (Benth.) Warb.

Ber. Deutsch. Bot. Ges. 13: (84) ["1895" publ. 18 Feb. 1896] (Warburg 1896). — *Myristica hostmannii* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 7 (Bentham 1853). — *Palala hostmannii* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 567 [5 Nov. 1891] (Kuntze 1891), *nom. inval.* (genus name illegitimate).

VERNACULAR NAMES. — Pa: wahusi-kamwi, wahusi-wašiunō, wahusi-wašiunō-duwē • Ka: inyamu batí, watolo • Wp: kulupiyī, mukulupiyī • Nt: sabana toso paasaa • Cr: mousigo • Fr: moussigot, tosopassa marécage • Br: ucuúbarana.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *M.-F. Prévost* 3775.

INVENTORY DATA (FG). — 685 trees in 88 plots; $F_{\max} = 4.5\%$; $dbh_{\text{inv}} = 46.2$ cm.

[1202] *Iryanthera paraensis* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 358 (Huber 1909).

Iryanthera elongata Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 68 (Huber 1910).

Iryanthera sessilis Markgr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 236 [30 Mar. 1928] (Markgraf 1928).

VERNACULAR NAMES. — Wp: walimei • Wn: mutuju.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost* & P. Grenand 943, 15 m × 65 cm.

[1203] *Iryanthera sagotiana* (Benth.) Warb.

Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur. 68: 158 (Warburg 1897). — *Myristica sagotiana* Benth., *Hooker's Icon. Pl.* 13 [ser. 3, 3](2): 48 [June 1878] (Bentham 1878).

Myristica mouchico Laness. ex H. Stone, *Ann. Mus. Colon. Marseille*, sér. 3, 8 (2): 3 (Stone 1920).

Myristica mouchigo Baill., *Hist. Pl. [Baillon]* 2: 504 (Baillon 1870), *nom. nud.*

VERNACULAR NAMES. — Pa: wahusi-wašiunō • Ka: malakaipyo, watolo • Wp: kulupiyī, mukulupiyī • Wn: mutuju • Nt: mongo toso paasaa, toso paasaa • Cr: mousigo-rouj • Fr: moussigot rouge, tosopassa montagne • Br: ucuúbarana, ucuúba-vermelha.

HERBARIUM DATA (FG). — 167 collections at CAY. Sel. exs.: *P.A. Sagot* 1195, Sep. 1858 (holo-, K[K000575104, K000575105]; iso-, BM[BM000574366], BR[BR0000005782765], G[G00380955], GH[GH00039827], P[P02441770, P02441771, P02441772, P02441773], S[S-R-7029]).

INVENTORY DATA (FG). — 1475 trees in 171 plots; $F_{\max} = 15.4\%$; $dbh_{\text{inv}} = 55.7$ cm.

[1204] *Iryanthera tessmannii* Markgr.

Notizbl. Bot. Gart. Berlin-Dahlem 10: 236 [30 Mar. 1928] (Markgraf 1928).

Iryanthera microcarpa Ducke, *Bol. Técn. Inst. Agron. N.* 4: 8 [31 Mar. 1945] (Ducke 1945).

VERNACULAR NAMES. — Nt: toso paasaa.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori et al. 23340*.

INVENTORY DATA (FG). — 34 trees in 3 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 41.4$ cm.

Genus *Osteophloeum* Warb.

[1205] *Osteophloeum platyspermum* (Spruce ex A.DC.) Warb.

Ber. Deutsch. Bot. Ges. 13: (89) [“1895” publ. 18 Feb. 1896] (Warburg 1896). — *Myristica platysperma* Spruce ex A.DC., *Prodr. [A. P. de Candolle]* 14 (2): 695 [late Nov. 1857] (Candolle 1857). — *Palala platysperma* (Spruce ex A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 567 [5 Nov. 1891] (Kuntze 1891), *nom. inval.* (genus name illegitimate).

Iryanthera krukovi A.C.Sm., *Brittonia* 2 (2): 151 (Smith 1936).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Wp: a'i yū'i, kulupiyī u • Nt: auma pisi • Fr: moussigot blanc • Br: ucuúba-amarela, ucuúba-chico-de-assis.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2159*.

INVENTORY DATA (FG). — 72 trees in 45 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 90$ cm.

Genus *Virola* Aubl.

[1206] *Virola kwatae* Sabatier
(Fig. 40D)

Adansonia, sér. 3, 19 (2): 273 (Sabatier 1997).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: wahusi-washiunó • Te: waletsi pitág • Nt: gaan busi mulunba • Cr: djadjamadou-montangn • Fr: yayamadou montagne.

HERBARIUM DATA (FG). — 32 collections at CAY. Sel. exs.: *D. Sabatier 931* (holo-, P[P00135215]; iso-, CAY[CAY024880, CAY024881], K[K000575173], MPU[MPU024622], NY[00346038], P[P00135216, P00135217, P00135218], US[00623540]).

INVENTORY DATA (FG). — 153 trees in 66 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 127$ cm.

[1207] *Virola michelii* Heckel

Ann. Inst. Colon. Marseille 5 (2): 118 (Heckel 1898).

Myristica melinonii Benoist, *Bull. Mus. Natl. Hist. Nat.* 30: 104 [Benoist 1924]. — *Virola melinonii* (Benoist) A.C.Sm., *Brittonia* 2 (5): 502 [“1937” publ. 2 May 1938] (Smith 1938).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wahu, wahu-washiunó • Ka: walolo, walusi ityudano, walusilan tamunen • Te: waletsi tsing • Wp: wololo, wololo'i • Wn: mutuju, walolo • Nt: busi mulunba, lunba, mongo babun udu • Cr: djadjamadou-gran-bwa, djadjamadou-montangn • Fr: yayamadou montagne • Br: ucuúba-da-terra-firme, ucuúba-preta.

HERBARIUM DATA (FG). — 154 collections at CAY. Sel. exs.: *Hayes & Michel s.n.* (original material MARS, not seen); *E.M. Mélénin s.n.* (original material of *Myristica melinonii*: P[P00624725]).

INVENTORY DATA (FG). — 606 trees in 174 plots; $F_{\max} = 6.2\%$; $\text{dbh}_{\text{inv}} = 84.8$ cm.

[1208] *Virola multicostata* Ducke

J. Wash. Acad. Sci. 26: 261 (Ducke 1936).

VERNACULAR NAMES. — Pa: wahu, wahu-washiunó • Wp: wololo'u • Br: ucuúba-da-terra-firme.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier 960*.

INVENTORY DATA (FG). — 25 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 31.8$ cm.

[1209] *Virola sebifera* Aubl.

Hist. Pl. Guiane 2: 904 [Jun.-Dec. 1775] (Aublet 1775). — *Myristica sebifera* (Aubl.) Sw., *Prodr. [Swartz]*: 96 [20 Jun.-29 July 1788] (Swartz 1788). — *Palala sebifera* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 567 [5 Nov. 1891] (Kuntze 1891), *nom. inval.* (genus name illegitimate).

Myristica virola Raeusch., *Nomencl. Bot. [Raeusch.]*, ed. 3, 292 (Raeuschel 1797), *nom. nud.*

Myristica mocoia A.DC., *Prodr. [A. P. de Candolle]* 14 (1): 195 [mid Oct. 1856] (Candolle 1856). — *Palala mocoia* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 567 [5 Nov. 1891] (Kuntze 1891), *nom. inval.* (genus name illegitimate). — *Virola mocoia* (A.DC.) Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 183 (Warburg 1897).

Myristica sebifera var. *cordifolia* A.DC., *Prodr. [A. P. de Candolle]* 14 (1): 195 [mid Oct. 1856] (Candolle 1856). — *Myristica cordifolia* Mart. ex A.DC., *Prodr. [A. P. de Candolle]* 14 (1): 195 [mid Oct. 1856] (Candolle 1856), *nom. nud. pro syn.*

Myristica sebifera var. *curvinervia* A.DC., *Prodr. [A. P. de Candolle]* 14 (1): 195 [mid Oct. 1856] (Candolle 1856). — *Virola sebifera* var. *curvinervia* (A.DC.) Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 174 (Warburg 1897).

Myristica panamensis Hemsl., *Biol. Cent.-Amer., Bot.* 3 (14): 67 [Oct. 1882] (Hemsley 1882). — *Palala panamensis* (Hemsl.) Kuntze, *Revis. Gen. Pl.* 2: 567 [5 Nov. 1891] (Kuntze 1891), *nom. inval.* (genus name illegitimate). — *Virola panamensis* (Hemsl.) Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 185 (Warburg 1897).




FIG. 40. — Moraceae: **A**, *Perebea guianensis* Aubl.; **B**, *Trymatococcus amazonicus* Poepp. & Endl. (D. Sabatier & J.-F. Molino 5204). Myristicaceae: **C**, *Compsonaea ulei* Warb. (D. Sabatier & P. Birnbaum 4608); **D**, *Virola kwatae* Sabatier. © D. Sabatier/IRD.

Myristica fulva Rich. ex Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 169 (Warburg 1897), nom. nud. pro syn.

Virola venezuelensis Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 182 (Warburg 1897).

Virola boliviensis Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 184 (Warburg 1897).

Virola peruviana (A.DC.) Warb. var. *tomentosa* Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 189 (Warburg 1897).

Virola mycetis Pulle, *Recueil Trav. Bot. Néerl.* 4: 125 (Pulle 1907), *pro parte inflorescentia exclusa*.

Virola warburgii Pittier, *Contr. U.S. Natl. Herb.* 18 (4): 143 [3 Mar. 1916] (Pittier 1937).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wahusi-purubumna • Ka: virola (*fide* Aublet 1775), walolo, waluši ityudano, walušilan • Te: waletsi lawöt • Wp: suwilani wu, walime • Nt: kawee babun udu, kawee mulunba • Cr: bwa-chandèl, djadjamadou • Fr: bois chandelle • Br: ucuúba-vermelha.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM [BM000993836, BM000993837]).

INVENTORY DATA (FG). — 17 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40.2$ cm.

[1210] *Virola surinamensis* (Rol. ex Rottb.) Warb.

Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur. 68: 208 (Warburg 1897). — *Myristica surinamensis* Rol. ex Rottb., *Descr. Rar. Pl. Surin. [Rottbøll]*: 13 (Rottbøll 1776). — *Palala surinamensis* (Rol. ex Rottb.) Kuntze, *Revis. Gen. Pl.* 2: 568 [5 Nov. 1891] (Kuntze 1891), nom. inval. (genus name illegitimate).

Myristica americana Rottb., *Descr. Rar. Pl. Surin. [Rottbøll]*: 33 (Rottbøll 1776), nom. nud.

Myristica fatua Sw., *Prodr. [Swartz]* 96 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. hom., non Houtt. (Houttuyn 1774), nec Blume (1835).

Myristica sebifera var. *longifolia* Poir. ex Lam., *Encycl. [J. Lamarck et al.]* 4 (1): 390 [9 Feb. 1797] (Lamarck 1797).

Virola glaziovii Warb., *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 68: 219 (Warburg 1897).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: wahusi • Ka: wališi, waluši • Te: waletsi • Wp: walusi • Wn: aluti, waluti • Nt: lunba, mulunba, wata babun udu • Cr: djadjamadou-marikaj • Fr: yayamadou marécage • Br: ucuúba, ucuúba-branca, ucuúba-do-igapó.

HERBARIUM DATA (FG). — 80 collections at CAY. Sel. exs.: *Unknown coll. s.n.* (holotype of *Myristica sebifera* var. *longifolia*: P[P00631979]).

INVENTORY DATA (FG). — 39 trees in 22 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 90$ cm.

Family MYRTACEAE Juss.
Genus *Calycolpus* O.Berg

[1211] *Calycolpus goetheanus* (Mart. ex DC.) O.Berg
(Fig. 41A)

Linnaea 27 (2-3): 381 ["1854" publ. Jan. 1856] (Berg 1856). —
Myrtus goetheana Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 240
[mid Mar. 1828] (Candolle 1828).

Campomanesia glabra Benth., *J. Bot. [Hooker]* 2: 319 (Bentham
1840). — *Calycolpus glaber* (Benth.) O.Berg, *Linnaea* 27 (2-3):
379 ["1854" publ. Jan. 1856] (Berg 1856).

Calycolpus ovalifolius O.Berg, *Linnaea* 27 (2-3): 379 ["1854" publ.
Jan. 1856] (Berg 1856).

Calycolpus schomburgkianus O.Berg, *Linnaea* 27 (2-3): 380 ["1854"
publ. Jan. 1856] (Berg 1856).

Calycolpus schomburgkianus var. *recurvatus* O.Berg, *Linnaea* 27 (2-3):
380 ["1854" publ. Jan. 1856] (Berg 1856).

Calycolpus schomburgkianus var. *speciosus* O.Berg, *Linnaea* 27 (2-3):
380 ["1854" publ. Jan. 1856] (Berg 1856).

Calycolpus angustifolius L.Riley, *Bull. Misc. Inform. Kew* 1926 (4):
151 [20 Apr. 1926] (Riley 1926).

Calycolpus cordatus L.Riley, *Bull. Misc. Inform. Kew* 1926 (4): 152
[20 Apr. 1926] (Riley 1926).

Calycolpus glaber var. *angustilobus* L.Riley, *Bull. Misc. Inform. Kew*
1926 (4): 152 [20 Apr. 1926] (Riley 1926).

VERNACULAR NAMES. — Ka: awale tamipipyo, otonto epityi • Br: soim.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 3986.

INVENTORY DATA (FG). — 20 trees in 11 plots; $F_{\max} = 1.1\%$;
 $dbh_{inv} = 50$ cm.

[1212] *Calycolpus revolutus* (Schauer) O.Berg

Linnaea 27 (2-3): 383 ["1854" publ. Jan. 1856] (Berg 1856). —
Myrtus revoluta Schauer, *Linnaea* 21: 272 (Schauer 1848).

Calycolpus kegelianus O.Berg, *Linnaea* 27 (2-3): 381 ["1854" publ.
Jan. 1856] (Berg 1856).

Calycolpus kegelianus var. *robustus* O.Berg, *Linnaea* 27 (2-3): 381
["1854" publ. Jan. 1856] (Berg 1856).

Calycolpus kegelianus var. *gracilis* O.Berg, *Linnaea* 27 (2-3): 382
["1854" publ. Jan. 1856] (Berg 1856). — *Calycolpus gracilis*
(O.Berg) L.Riley, *Bull. Misc. Inform. Kew* 1926 (4): 150 [20 Apr.
1926] (Riley 1926).

Calycolpus chnoiophyllus L.Riley, *Bull. Misc. Inform. Kew* 1926 (4):
148 [20 Apr. 1926] (Riley 1926).

Calycolpus megalodon L.Riley, *Bull. Misc. Inform. Kew* 1926 (4): 149
[20 Apr. 1926] (Riley 1926).

Calycolpus pyrifera L.Riley, *Bull. Misc. Inform. Kew* 1926 (4): 149
[20 Apr. 1926] (Riley 1926).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: asaikiabe, sadyabe, seityape.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *P.A. Sagot*
215, 1857 (holotype of *Calycolpus megalodon*: K[K000277220];
iso-, W[W18890092620]).

INVENTORY DATA (FG). — 22 trees in 3 plots; $F_{\max} = 1.9\%$;
 $dbh_{inv} = 19.3$ cm.

Genus *Campomanesia* Ruiz & Pav.

[1213] *Campomanesia aromatica* (Aubl.) Griseb.

Fl. Brit. W.I. [Grisebach] 242 [late 1860] (Grisebach 1860). —
Psidium aromaticum Aubl., *Hist. Pl. Guiane* 1: 485 [Jun.-Dec. 1775]
(Aublet 1775). — *Burchardia aromatica* (Aubl.) Raf., *Sylva Tellur.*:
106 (Rafinesque 1838).

Myrtus psidioides Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 44 [Oct.
1825] (Hamilton 1825).

Psidium tenuifolium DC., *Prodr. [A. P. de Candolle]* 3: 236 [mid
Mar. 1828] (Candolle 1828). — *Campomanesia tenuifolia* (DC.)
O.Berg, *Fl. Bras. [Martius]* 14 (1): 452 [15 May 1857] (Berg 1857).

Myrtus fascicularis DC., *Prodr. [A. P. de Candolle]* 3: 240 [mid Mar.
1828] (Candolle 1828).

Eugenia sparsiflora DC., *Prodr. [A. P. de Candolle]* 3: 263 [mid Mar.
1828] (Candolle 1828). — *Myrtus sparsiflora* Mart. ex DC., *Prodr.*
[A. P. de Candolle] 3: 263 [mid Mar. 1828] (Candolle 1828), nom.
nud. pro syn. — *Abbevillea martiana* O.Berg, *Fl. Bras. [Martius]* 14
(1): 435 [15 May 1857] (Berg 1857), nom. illeg. superfl. (based on
Eugenia sparsiflora). — *Campomanesia sparsiflora* (DC.) J.F.Macbr.,
Candollea 5: 394 (Macbride 1934).

Campomanesia coactanea O.Berg, *Fl. Bras. [Martius]* 14 (1): 444
[15 May 1857] (Berg 1857).

Campomanesia synchrona O.Berg, *Fl. Bras. [Martius]* 14 (1): 444
[15 May 1857] (Berg 1857).

Campomanesia ciliata O.Berg, *Fl. Bras. [Martius]* 14 (1): 453 [15
May 1857] (Berg 1857).

Eugenia desvauxiana O.Berg, *Linnaea* 27 (2-3): 198 ["1854" publ.
Jan. 1856] (Berg 1856).

Campomanesia beaurepaireiana Kiaersk., *Enum. Myrt. Bras.*: 15 [Oct.-
Nov. 1893] (Kiaerskov 1893).

Campomanesia glazioviana Kiaersk., *Enum. Myrt. Bras.*: 16 [Oct.-
Nov. 1893] (Kiaerskov 1893).

VERNACULAR NAMES. — Pa: pirimavan • Ka: kalawilulan • Nt: andoya,
madyo uman • Cr: bwa-sitronèl • Fr: citronelle (*fide* Aublet
1775) • Br: aracá-lima.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000953694], LINN[LINN
HS 881.3]).

SIZE. — Up to 20 m tall (Landrum 1986).

[1214] *Campomanesia grandiflora* (Aubl.) Sagot
(Fig. 41B)

Ann. Sci. Nat., Bot. sér. 6, 20: 182 (Sagot 1885). — *Psidium*
grandiflorum Aubl., *Hist. Pl. Guiane* 1: 483 [Jun.-Dec. 1775]

(Aublet 1775). — *Psidium aromaticum* var. *grandiflorum* (Aubl.) Pers., *Syn. Pl. [Persoon] 2 (1)*: 27 [Nov. 1806] (Persoon 1806). — *Burchardia grandiflora* (Aubl.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838), “*Burcardia*”.

Campomanesia poiteauii O.Berg, *Linnaea* 27 (4): 432 [“1854” publ. Feb. 1856] (Berg 1856).

VERNACULAR NAMES. — Pa: pirimavan • Cr: bwa-sitronèl • Br: aracá-lima.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000953691], LINN[LINN HS 881.4]).

INVENTORY DATA (FG). — 12 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27$ cm.

Genus *Eugenia* L.

[1215] *Eugenia albicans* (O.Berg) Urb.

Bot. Jahrb. Syst. 19 (5): 617 [26 Feb. 1895] (Urban 1895). — *Stenocalyx albicans* O.Berg, *Linnaea* 30: 698 [Mar. 1861] (Berg 1861).

VERNACULAR NAMES. — Pa: inam-etri-seinô, inamkat-seinô • Wp: yawayi lû • Cr: bwa-sann.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *B. Dutrèvre 388*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 16.7$ cm.

[1216] *Eugenia anastomosans* DC.

Prodr. [A. P. de Candolle] 3: 269 [mid Mar. 1828] (Candolle 1828).

Eugenia ptariensis Steyermark., *Fieldiana, Bot.* 28 (4): 1013 (Steyermark 1957).

VERNACULAR NAMES. — Ka: kumeti • Wp: kumati, kumati u.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material F[V0065081F], L[L0009403], P[P00784923, P00784924, P05156416]).

INVENTORY DATA (FG). — 29 trees in 24 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 25$ cm.

[1217] *Eugenia armeniaca* Sagot

Ann. Sci. Nat., Bot. sér. 6, 20: 190 (Sagot 1885).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: moyu’i.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *P.A. Sagot 259* (syntypes BR[BR0000005228652 {1857}], F[V0065085F {1857}], K[K000276620 & K000276621 {1857}], P[P01902320 {1855}], P01902321 {Nov. 1856}, P01902322 {1856}], S[S05-2789 {1857}]).

INVENTORY DATA (FG). — 5 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 35.9$ cm.

[1218] *Eugenia aubletiana* Mattos

Loefgrenia 120: 9 [Mar. 2005] (Mattos 2005), excluding synonyms *E. latifolia* Aubl. and *Catinga moschata* Aubl. [synonym of *E. moschata* (Aubl.) Nied. ex T.Durand & B.D.Jacks.]. — *Calycorectes latifolius* O.Berg, *Linnaea* 30: 701 [Mar. 1861] (Berg 1861), excluding synonym *E. latifolia* Aubl. — *Calycorectes bergii* Sandwith, *Bull. Misc. Inform. Kew* 1932 (5): 212 [27 June 1932] (Sandwith 1932).

Catinga oblongifolia Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20*: 197 (Sagot 1885). — *Eugenia oblongifolia* (Sagot) Nied. ex T.Durand & B.D.Jacks., *Index Kew. Suppl.* 1: 164 (Durand & Jackson 1902), *nom. illeg. hom., non* Duthie (1878).

NOTES. — Berg (1861: 701) misinterpreted *E. latifolia* Aubl. and used it as the basionym of *Calycorectes latifolius* O.Berg, while the description and type corresponded to an entirely distinct species. Sandwith (1932: 212) pointed out this mistake and created the new name *C. bergii* Sandwith for the taxon described by Berg. Since molecular data showed that *Calycorectes* is polyphyletic within *Eugenia* (Proença et al. 2020), *C. bergii* has been transferred to *Eugenia*.

VERNACULAR NAMES. — Wp: iwá ū, wila ū.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *P.A. Sagot 272*, Jan. 1857 (lecto-, P[P04722646], designated by Proença et al. [2020: 50]; isolecto-, BR[BR0000005304073], K[K000565050], P[P05229281, P05229285], S[S05-2964]); *B. Bordenave 887*, height 12 m.

[1219] *Eugenia biflora* (L.) DC.

Prodr. [A. P. de Candolle] 3: 276 [mid Mar. 1828] (Candolle 1828). — *Myrtus biflora* L., *Syst. Nat.*, ed. 10, 2: 1056 [7 June 1759] (Linnaeus 1759).

Caryophyllus fruticosus Mill., *Gard. Dict.*, ed. 8, n. 3 [16 Apr. 1768] (Miller 1768).

Eugenia mini Aubl., *Hist. Pl. Guiane* 1: 498 [Jun.-Dec. 1775] (Aublet 1775). — *Eugenia microcarpos* var. *mini* (Aubl.) Lam., *Encycl. [J. Lamarck et al.] 3 (1)*: 201 [19 Oct. 1789] (Lamarck 1789). — *Eugenia mini* var. *microcarpos* (Lam.) Pers., *Syn. Pl. [Persoon] 2 (1)*: 28 [Nov. 1806] (Persoon 1806). — *Myrtus mini* (Aubl.) Spreng., *Syst. Veg. [Sprengel] 2*: 486 [Jan.-May 1825] (Sprengel 1825). — *Myrcia splendens* var. *mini* (Aubl.) DC., *Prodr. [A. P. de Candolle] 3*: 244 [mid Mar. 1828] (Candolle 1828). — *Myrcia mini* (Aubl.) Sweet, *Hort. Brit. [Sweet]*, ed. 2: 211 (Sweet 1830). — *Cumetea mini* (Aubl.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838). — *Myrcia schomburgkiana* O.Berg, *Linnaea* 27 (1): 110 [“1854” publ. Nov. 1855] (Berg 1855), *nom. illeg. superfl.* (based on *Eugenia mini*). — *Eugenia biflora* var. *mini* (Aubl.) Amshoff, *Recueil Trav. Bot. Néerl.* 42: 12 (Amshoff 1950).

Myrtus virgultosa Sw., *Prodr. [Swartz] 70* [20 Jun.-29 July 1788] (Swartz 1788). — *Eugenia virgultosa* (Sw.) DC., *Prodr. [A. P. de Candolle] 3*: 280 [mid Mar. 1828] (Candolle 1828). — *Eugenia biflora* var. *virgultosa* (Sw.) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 628 [26 Feb. 1895] (Krug & Urban 1895).

Eugenia microcarpos Lam., *Encycl. [J. Lamarck et al.] 3 (1)*: 201 [19 Oct. 1789] (Lamarck 1789).

Myrtus pallens Vahl, *Symb. Bot. [Vahl] 2*: 57 [Jul.-Dec. 1791] (Vahl 1791). — *Eugenia pallens* (Vahl) DC., *Prodr. [A. P. de Candolle] 3*: 284 [mid Mar. 1828] (Candolle 1828), *nom. illeg. hom., non* Poir. (Poirier 1813). — *Eugenia biflora* var. *pallens* (Vahl) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 629 [26 Feb. 1895] (Krug & Urban 1895).

- Eugenia lancea* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 123 [24 Sep. 1813] (Poirret 1813). — *Myrtus lancea* (Poir.) Spreng., *Syst. Veg. [Sprengel]* 2: 482 [Jan.-May 1825] (Sprengel 1825). — *Eugenia biflora* var. *lancea* (Poir.) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 631 [26 Feb. 1895] (Krug & Urban 1895).
- Eugenia australis* Colla, *Hortus Ripulensis* 1: 54 [Jun.-July 1824] (Colla 1824), in nota, *nom. illeg. hom.*, *non* Wendl. ex Link (1822).
- Myrtus berteroana* Spreng., *Syst. Veg. [Sprengel]* 2: 482 [Jan.-May 1825] (Sprengel 1825), “*Berteriana*”. — *Eugenia berteroana* (Spreng.) DC., *Prod. [A. P. de Candolle]* 3: 285 [mid Mar. 1828] (Candolle 1828), “*Berteriana*”.
- Eugenia xylopifolia* DC., *Prod. [A. P. de Candolle]* 3: 279 [mid Mar. 1828] (Candolle 1828).
- Eugenia ludibunda* Bertero ex DC., *Prod. [A. P. de Candolle]* 3: 280 [mid Mar. 1828] (Candolle 1828). — *Eugenia biflora* var. *ludibunda* (Bertero ex DC.) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 630 [26 Feb. 1895] (Krug & Urban 1895).
- Eugenia acutiloba* DC., *Prod. [A. P. de Candolle]* 3: 281 [mid Mar. 1828] (Candolle 1828).
- Eugenia racemosa* DC., *Prod. [A. P. de Candolle]* 3: 281 [mid Mar. 1828] (Candolle 1828), *nom. illeg. hom.*, *non* L. (Linnaeus 1753).
- Eugenia sericiflora* Benth., *Bot. Voy. Sulphur [Bentham]*: 98 [14 Apr. 1845] (Bentham 1845).
- Eugenia wallenii* Macfad., *Fl. Jamaica [Macfadyen]* 2: 118 (Macfadyen 1850). — *Eugenia biflora* var. *wallenii* (Macfad.) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 629 [26 Feb. 1895] (Krug & Urban 1895).
- Eugenia virgata* Macfad., *Fl. Jamaica [Macfadyen]* 2: 121 (Macfadyen 1850), *nom. illeg. hom.*, *non* Gardner (1845).
- Eugenia jamaicensis* O.Berg, *Linnaea* 27 (2-3): 237 [“1854” publ. Jan. 1856] (Berg 1856). — *Eugenia virgultosa* var. *jamaicensis* (O.Berg) Proctor, *J. Arnold Arbor.* 63 (3): 276 [Apr. 1982] (Proctor 1982).
- Eugenia macarensis* O.Berg, *Linnaea* 27 (2-3): 283 [“1854” publ. Jan. 1856] (Berg 1856).
- Myrcia erythroxylon* var. *virescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 173 [15 May 1857] (Berg 1857).
- Eugenia freireana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 305 [15 May 1857] (Berg 1857).
- Eugenia freireana* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 305 [15 May 1857] (Berg 1857).
- Eugenia freireana* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 305 [15 May 1857] (Berg 1857).
- Eugenia salicifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 312 [15 May 1857] (Berg 1857), *nom. illeg. hom.*, *non* (Kunth) DC. (Candolle 1828).
- Eugenia modesta* DC. var. *jamaicensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 314 [15 May 1857] (Berg 1857).
- Eugenia acuminatissima* O.Berg, *Fl. Bras. [Martius]* 14 (1): 315 [15 May 1857] (Berg 1857), *nom. illeg. hom.*, *non* Miq. (Miquel 1846).
- Eugenia hoffmannseggii* O.Berg, *Fl. Bras. [Martius]* 14 (1): 315 [15 May 1857] (Berg 1857). — *Eugenia biflora* var. *hoffmannseggii* (O.Berg) Amshoff, *Fl. Surinam* 3 (2): 123 (Amshoff 1951).
- Eugenia hoffmannseggii* var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 315 [15 May 1857] (Berg 1857).
- Eugenia hoffmannseggii* var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 316 [15 May 1857] (Berg 1857).
- Eugenia inundata* DC. var. *acutifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 520 [1 Feb. 1858] (Berg 1858).
- Eugenia fieldingii* O.Berg, *Linnaea* 29 (2): 242 [June 1858] (Berg 1858).
- Eugenia meyeriana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 588 [15 Jan. 1859] (Berg 1859).
- Eugenia meyeriana* var. *depauperata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 588 [15 Jan. 1859] (Berg 1859).
- Eugenia meyeriana* var. *dives* O.Berg, *Fl. Bras. [Martius]* 14 (1): 588 [15 Jan. 1859] (Berg 1859).
- Eugenia richardiana* O.Berg, *Linnaea* 30: 694 [Mar. 1861] (Berg 1861).
- Eugenia xylopifolia* var. *brevipes* O.Berg, *Linnaea* 30: 694 [Mar. 1861] (Berg 1861).
- Eugenia myriostigma* Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20:* 194 (Sagot 1885). — *Eugenia biflora* var. *myriostigma* (Sagot) Amshoff, *Fl. Surinam* 3 (2): 124 (Amshoff 1951).
- Eugenia hartii* Kiaersk., *Bot. Tidsskr.* 17: 271 (Kiaerskov 1890).
- Eugenia alexandri* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 626 [26 Feb. 1895] (Krug & Urban 1895).
- Myrtus biflora* f. *subsericea* Kuntze, *Revis. Gen. Pl.* 3 (3): 89 [28 Sep. 1898] (Kuntze 1898).
- Myrtus biflora* var. *salicifolia* Kuntze, *Revis. Gen. Pl.* 3 (3): 89 [28 Sep. 1898] (Kuntze 1898).
- Myrtus biflora* var. *yapacani* Kuntze, *Revis. Gen. Pl.* 3 (3): 89 [28 Sep. 1898] (Kuntze 1898).
- Eugenia loretensis* Diels, *Bot. Jahrb. Syst.* 37 (5): 597 [30 Oct. 1906] (Diels 1906).
- Eugenia leptophlebia* Diels, *Verh. Bot. Vereins Prov. Brandenburg* 48: 192 [“1906” publ. 8 Mar. 1907] (Diels 1907).
- Eugenia brachythrix* Urb., *Symb. Antill. [Urban]* 6 (1): 23 [15 July 1909] (Urban 1909).
- Eugenia alfaroana* Standl., *J. Wash. Acad. Sci.* 14: 240 (Standley 1924).
- Eugenia nicholsii* Fawc. & Rendle, *J. Bot.* 64: 14 (Fawcett & Rendle 1926).
- Eugenia amanuensis* Steyermark., *Fieldiana, Bot.* 28 (4): 1010 (Steyermark 1957).
- Eugenia caurensis* Steyermark., *Fieldiana, Bot.* 28 (4): 1010 (Steyermark 1957).
- Eugenia jamaicana* Mattos, *Loefgrenia* 123: 2 [Aug. 2006] (Mattos 2006).
- VERNACULAR NAMES. — Pa: kagegut-puvemna, karegut-puvemna
• Br: vassourinha.
- HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 1064*.
- SIZE. — Brazil, Amapá. *B.V. Rabelo et al. 2278* (MO), 10 m × 15 cm.



FIG. 41.—Myrtaceae: **A**, *Calycolpus goetheanus* (Mart. ex DC.) O.Berg (M.-F. Prévost & D. Sabatier 3986); **B**, *Campomanesia grandiflora* (Aubl.) Sagot (J.-F. Molino & D. Sabatier 2249); **C**, *Eugenia mimus* McVaugh (D. Sabatier et al. 5780); **D**, **E**, *Eugenia* sp. O (J.-F. Molino 3430). A, C, © D. Sabatier/IRD; B, D, E, © J.-F. Molino/IRD.

[1220] *Eugenia brownsbergii* Amshoff

Recueil Trav. Bot. Néerl. 42: 14 (Amshoff 1950). — *Eugenia brownsbergii* var. *glaucia* Amshoff, *Recueil Trav. Bot. Néerl.* 42: 14 (Amshoff 1950).

VERNACULAR NAMES. — Pa: arak-kamwi, pakusin-amana • Ka: kuwapityano.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville B-4549*.

SIZE. — Brazil, Amazonas. *W.W. Thomas et al. 5247* (MO), 15 m.

[1221] *Eugenia chrysophyllum* Poir.

Encycl. [J. Lamarck et al.] Suppl. 3: 129 [24 Sep. 1813] (Poiret 1813). — *Myrtus cayennensis* Spreng., *Syst. Veg. [Sprengel]* 2: 481 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Eugenia citrifolia*).

Eugenia chrysophylloides DC., *Prodri. [A. P. de Candolle]* 3: 276 [mid Mar. 1828] (Candolle 1828).

Eugenia llewelynii Steyermark., *Fieldiana, Bot.* 28 (4): 1011 (Steyermark 1957).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J. Martin s.n.* (type FI[FI11503]).

SIZE. — Colombia, Amazonas. *L.E. Urrego et al. 325* (MO), 15 m × 10 cm.

[1222] *Eugenia citrifolia* Poir.

Encycl. [J. Lamarck et al.] Suppl. 3: 129 [24 Sep. 1813] (Poiret 1813). — *Myrtus cayennensis* Spreng., *Syst. Veg. [Sprengel]* 2: 481 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Eugenia citrifolia*).

Eugenia adenocalyx DC., *Prod. [A. P. de Candolle] 3:* 271 [mid Mar. 1828] (Candolle 1828).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J. Martin s.n.* [“*bords de la rivière d’Oyac, Caïenne*”, “*b. Poiret*”] (original material FI[FI011504], P[P05237715]).

SIZE. — Brazil, Amazonas. *M. Pacheco 96* (MO), 12 m × 18.6 cm.

[1223] *Eugenia cowanii* McVaugh

Mem. New York Bot. Gard. 18 (2): 173 (McVaugh 1969).

Eugenia cryptadena var. *gracilis* Amshoff, *Recueil Trav. Bot. Néerl.* 42: 18 (Amshoff 1950).

VERNACULAR NAMES. — Ka: alamilu.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 9455* (CAY[CAY095706], US[00659642]).

SIZE. — Up to 12 m tall (McVaugh 1969).

[1224] *Eugenia cucullata* Amshoff

Recueil Trav. Bot. Néerl. 42: 17 (Amshoff 1950).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *C. Moretti 571*.

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 23.4$ cm.

[1225] *Eugenia cupulata* Amshoff

Recueil Trav. Bot. Néerl. 39: 160 (Amshoff 1942).

VERNACULAR NAMES. — Pa: arak-priyu • Ka: pomiyilan • Wp: kalai ka'a • Nt: busi banda.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 930*.

INVENTORY DATA (FG). — 25 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 21.6$ cm.

[1226] *Eugenia denigrata* McVaugh

Mem. New York Bot. Gard. 18 (2): 176 (McVaugh 1969).

VERNACULAR NAMES. — Pa: paku-amana.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P. Grenand 2119*.

SIZE. — > 10 cm dbh (Cardoso *et al.* 2017).

[1227] *Eugenia densiracemosa* Mazine & Faria

Phytotaxa 151 (1): 53 [6 Dec. 2013] (Mazine & Faria 2013).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-F. Molino 2235* (para-, CAY[CAY073409]).

INVENTORY DATA (FG). — 37 trees in 5 plots; $F_{\max} = 5.2\%$; $\text{dbh}_{\text{inv}} = 21.3$ cm.

[1228] *Eugenia egensis* DC.

Prod. [A. P. de Candolle] 3: 281 [mid Mar. 1828] (Candolle 1828). — *Myrtus egensis* Mart. ex DC., *Prod. [A. P. de Candolle] 3:* 281 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Eugenia sphaerosperma DC., *Prod. [A. P. de Candolle] 3:* 278 [mid Mar. 1828] (Candolle 1828).

Eugenia tenuiramis Miq., *Stirp. Surinam. Select.*: 39 [“1850” publ. Mar. 1851] (Miquel 1851). — *Eugenia egensis* var. *tenuiramis* (Miq.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia perforata O.Berg, *Linnaea* 27 (2-3): 236 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia egensis var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia egensis var. *bimarginata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia egensis var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia egensis var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia egensis var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 296 [15 May 1857] (Berg 1857).

Eugenia maculata O.Berg, *Fl. Bras. [Martius]* 14 (1): 297 [15 May 1857] (Berg 1857).

Eugenia parodiana Morong, *Ann. New York Acad. Sci.* 7: 107 (Morong 1893).

VERNACULAR NAMES. — Br: azeitona-brava.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3228*.

SIZE. — Up to 10 m tall (McVaugh 1958).

[1229] *Eugenia exaltata* Rich. ex O.Berg

Linnaea 30: 687 [Mar. 1861] (Berg 1861).

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *L.C. Richard 25* (original material F[V0065165F], P[P01902431, P01902432]).

INVENTORY DATA (FG). — 102 trees in 42 plots; $F_{\max} = 3.9\%$; $\text{dbh}_{\text{inv}} = 26.9$ cm.

[1230] *Eugenia excelsa* O.Berg

Fl. Bras. [Martius] 14 (1): 277 [15 May 1857] (Berg 1857).

Eugenia cuspidata O.Berg, *Fl. Bras. [Martius]* 14 (1): 286 [15 May 1857] (Berg 1857).

Eugenia obovata O.Berg, *Fl. Bras. [Martius]* 14 (1): 289 [15 May 1857] (Berg 1857), nom. illeg. hom., non Poir. (Poirier 1813).

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Larpin* 702.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{inv} = 15.3$ cm.

[1231] *Eugenia ferreiraeana* O.Berg

Fl. Bras. [Martius] 14 (1): 285 [15 May 1857] (Berg 1857).

Eugenia nemoralis DC., *Prodr. [A. P. de Candolle]* 3: 267 [mid Mar. 1828] (Candolle 1828).

Eugenia ramiflora Desv. ex Ham. var. *montana* Amshoff, *Bull. Torrey Bot. Club* 75 (5): 535 [11 Oct. 1948] (Amshoff 1948).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-2026.

SIZE. — Brazil, Amazonas. *N.T. Silva & U. Brazão* 60788 (MO), 10 m × 15 cm.

[1232] *Eugenia flavescens* DC.

Prodr. [A. P. de Candolle] 3: 272 [mid Mar. 1828] (Candolle 1828). — *Myrtus flavescens* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 272 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Eugenia flavescens var. *guyanensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 272 [15 May 1857] (Berg 1857).

Eugenia flavescens var. *longifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 272 [15 May 1857] (Berg 1857).

Eugenia flavescens var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 272 [15 May 1857] (Berg 1857).

VERNACULAR NAMES. — Pa: arak-priyu.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *Service Forestier* 5118.

SIZE. — Up to 10 cm dbh (Amshoff 1951).

[1233] *Eugenia florida* DC.

Prodr. [A. P. de Candolle] 3: 283 [mid Mar. 1828] (Candolle 1828). — *Myrtus florida* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 283 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Eugenia patula DC., *Prodr. [A. P. de Candolle]* 3: 284 [mid Mar. 1828] (Candolle 1828).

Myrcia patula Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 284 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Eugenia sylvatica Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 337 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832).

Eugenia atropunctata Steud., *Flora* 26 (45): 762 [17 Dec. 1843] (Steudel 1843).

Eugenia atropunctata var. *robusta* O.Berg, *Linnaea* 27 (2-3): 293 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia atropunctata var. *gracilis* O.Berg, *Linnaea* 27 (2-3): 294 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia gardneriana O.Berg, *Fl. Bras. [Martius]* 14 (1): 316 [15 May 1857] (Berg 1857). — *Eugenia moraviana* O.Berg var. *gardneriana* (O.Berg) Mattos, *Loefgrenia* 85: 2 (Mattos 1984).

Eugenia gardneriana var. *dives* O.Berg, *Fl. Bras. [Martius]* 14 (1): 316 [15 May 1857] (Berg 1857).

Eugenia gardneriana var. *depauperata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 317 [15 May 1857] (Berg 1857).

Eugenia oligoneura O.Berg, *Fl. Bras. [Martius]* 14 (1): 321 [15 May 1857] (Berg 1857).

Eugenia membranacea O.Berg, *Fl. Bras. [Martius]* 14 (1): 389 [15 Jan. 1859] (Berg 1859).

Eugenia membranacea O.Berg, *Fl. Bras. [Martius]* 14 (1): 389 [15 Jan. 1859] (Berg 1859).

Eugenia gardneriana var. *rigida* O.Berg, *Linnaea* 30: 694 [Mar. 1861] (Berg 1861).

Eugenia racemifera Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 20: 195 (Sagot 1885), nom. illeg. hom., non O.Berg (1857).

Eugenia seriatoracemosa Kiaersk., *Enum. Myrt. Bras.* 151 [Oct.-Nov. 1893] (Kiaerskov 1893), "seriato-racemosa".

Myrcia doloresensis Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 63 [9 Apr. 1895] (Hieronymus 1895).

Eugenia tinge-lingua S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 357 ["1894-96" publ. Dec. 1895] (Moore 1895).

Eugenia perorebi D.Parodi ex Speg. & Girola, *Anal. Soc. Rural Argent. (Cat. Descr. Maderas)* 361 (Spegazzini & Girola 1910).

Eugenia melanosticta Standl., *J. Arnold Arbor.* 11 (2): 126 [29 Apr. 1930] (Standley 1930), nom. illeg. hom., non (Miq.) Koord. & Valeton (Koorders & Valeton 1900).

Eugenia coloradoensis Standl., *Trop. Woods* 52: 27 (Standley 1937).

VERNACULAR NAMES. — Pa: arak-kamwi, inam-etni-seine, pakusin-amana • Ka: yalami • Te: pakulea • Wp: iwá ū, wila ū • Wn: pailaimë, tēpepulu • Br: guamirim, pitanga.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 2059.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{inv} = 16.7$ cm.

[1234] *Eugenia galbaensis* Mattos

Loefgrenia 120: 4 [Mar. 2005] (Mattos 2005).

Calycorectes guyanensis Mattos, *Loefgrenia* 108: 1 (Mattos 1996).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-J. de Granville* et al. 9032 (holo-, US[00479191]; iso-, CAY[CAY075775, CAY075776, CAY075777, CAY075778], P[P05239223]).

INVENTORY DATA (FG). — 1 tree, dbh = 28.5 cm.

[1235] *Eugenia gerdae* Mazine

Neodiversity 4: 2 [14 Dec. 2009] (Mazine 2009).

Eugenia monticola (Sw.) DC. var. *racemosa* Amshoff, *Fl. Surinamæ* 3 (2): 121 (Amshoff 1951).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 15181*.

INVENTORY DATA (FG). — 3 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 10.2$ cm.

[1236] *Eugenia gomesiana* O.Berg

Fl. Bras. [Martius] 14 (1): 254 [15 May 1857] (Berg 1857).

Eugenia prosoneura O.Berg, *Linnaea* 31: 255 (Berg 1861-1862).

VERNACULAR NAMES. — Wn: olowaimë.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-J. de Granville 1869*, 10-15m tall.

[1237] *Eugenia gongylocarpa* M.L.Kawas. & B.Holst

Brittonia 46 (2): 138 (Kawasaki & Holst 1994).

VERNACULAR NAMES. — Wn: tēpepuimë.

HERBARIUM DATA (FG). — 53 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15554* (original material).

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 13.7$ cm.

[1238] *Eugenia griseiflora* McVaugh

Mem. New York Bot. Gard. 18 (2): 187 (McVaugh 1969).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al. 25434*.

SIZE. — Up to 10 cm dbh (McVaugh 1969).

[1239] *Eugenia lambertiana* DC.

Prod. [A. P. de Candolle] 3: 270 [mid Mar. 1828] (Candolle 1828).

Eugenia schomburgkii Benth., *J. Bot. [Hooker]* 2: 321 (Bentham 1840).

Eugenia smaragdina O.Berg, *Linnaea* 27 (2-3): 218 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia smaragdina var. *angustifolia* O.Berg, *Linnaea* 27 (2-3): 218 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia smaragdina var. *brevipes* O.Berg, *Linnaea* 27 (2-3): 219 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia smaragdina var. *rigida* O.Berg, *Linnaea* 27 (2-3): 219 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia correae O.Berg, *Fl. Bras. [Martius]* 14 (1): 277 [15 May 1857] (Berg 1857).

Eugenia oligophylla Rich. ex O.Berg, *Linnaea* 30: 685 [Mar. 1861] (Berg 1861).

Eugenia flavonigra Rich. ex O.Berg, *Linnaea* 30: 691 [Mar. 1861] (Berg 1861), "flavo-nigra".

Eugenia flavonigra var. *guadalupensis* O.Berg, *Linnaea* 30: 691 [Mar. 1861] (Berg 1861).

Eugenia flavonigra var. *martinicensis* O.Berg, *Linnaea* 30: 692 [Mar. 1861] (Berg 1861).

Eugenia lambertiana var. *hispidula* McVaugh, *Mem. New York Bot. Gard.* 18 (2): 190 (McVaugh 1969).

VERNACULAR NAMES. — Wp: kumati sili • Wn: tēpepu.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-688*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 16.2$ cm.

[1240] *Eugenia latifolia* Aubl.

Hist. Pl. Guiane 1: 502 [Jun.-Dec. 1775] (Aublet 1775). — *Myrtus latifolia* (Aubl.) Spreng., *Syst. Veg. [Sprengel]* 2: 482 [Jan.-May 1825] (Sprengel 1825), nom. illeg. hom., non B.Heyne ex Roth (1821). — *Calycorectes latifolius* (Aubl.) O.Berg, *Linnaea* 30: 701 [Mar. 1861] (Berg 1861), excl. descr. et specim.

VERNACULAR NAMES. — Wp: kumati, kumati u, pakea'i sī • Wn: kalapime, tēpepu, tēpepuimë • Nt: kiinsikin banda.

HERBARIUM DATA (FG). — 65 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000953757]).

INVENTORY DATA (FG). — 12 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.3$ cm.

[1241] *Eugenia luciae* Amshoff

Recueil Trav. Bot. Néerl. 42: 14 (Amshoff 1950).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *O. Poncy et al. 1830*.

SIZE. — Suriname. *R.J. Evans et al. 2513* (MO), 12 m.

[1242] *Eugenia marowynensis* Miq.

Stirp. Surinam. Select.: 39 ["1850" publ. Mar. 1851] (Miquel 1851). — *Myrciaria marowynensis* (Miq.) O.Berg, *Linnaea* 27 (2-3): 335 ["1854" publ. Jan. 1856] (Berg 1856).

VERNACULAR NAMES. — Pa: avitkat-kamwi • Ka: pasami • Wp: kulupitá átā • Wn: waiko epit.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *C. Moretti 1035*.

INVENTORY DATA (FG). — 1 tree, dbh = 17.2 cm.

[1243] *Eugenia minus* McVaugh
(Fig. 41C)

Mem. New York Bot. Gard. 18 (2): 193 (McVaugh 1969).

VERNACULAR NAMES. — Pa: arak-kamwi, pakusin-amana • Wp: akusi yeti, miku ka'a, tamukwálē áká • Br: araçá.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *P. Grenand 3131*.

INVENTORY DATA (FG). — 1 tree, dbh = 12.1 cm.

[1244] *Eugenia morii* B.Holst & M.L.Kawas.

Brittonia 52 (1): 21 (Holst & Kawasaki 2000).

HERBARIUM DATA (FG). — Known only from the type: *S.A. Mori et al. 14900* (holo-, CAY[CAY024885, CAY024886]; iso-, MO, not seen).

INVENTORY DATA (FG). — 6 trees in 3 plots; F_{max} < 1 %; dbh_{inv} = 33.1 cm.

[1245] *Eugenia moschata* (Aubl.) Nied. ex T.Durand & B.D.Jacks.

Index Kew. Suppl. 1: 164. (Durand & Jackson 1902). — *Catinga moschata* Aubl., *Hist. Pl. Guiane* 1: 511 [Jun.-Dec. 1775] (Aublet 1775).

Catinga aromatica Aubl., *Hist. Pl. Guiane* 1: 511 [Jun.-Dec. 1775] (Aublet 1775). — *Eugenia catinga* Baill., *Hist. Pl. [Baillon]* 6: 344 [June 1876] (Baillon 1876), nom. illeg. superfl. (based on *Catinga moschata*).

Eugenia fasciculiflora O.Berg, *Linnaea* 27 (2-3): 233 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia pisonis O.Berg, *Fl. Bras. [Martius]* 14 (1): 228 [15 May 1857] (Berg 1857).

Eugenia leucantha O.Berg, *Fl. Bras. [Martius]* 14 (1): 277 [15 May 1857] (Berg 1857).

Eugenia feijoi O.Berg, *Fl. Bras. [Martius]* 14 (1): 283 [15 May 1857] (Berg 1857).

Eugenia paraensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 301 [15 May 1857] (Berg 1857).

Eugenia costata O.Berg, *Fl. Bras. [Martius]* 14 (1): 577 [15 Jan. 1859] (Berg 1859), nom. illeg. hom., non Cambess. (Cambessèdes 1833).

Acca glazioviana Kjaersk., *Enum. Myrt. Bras.* 115 [Oct.-Nov. 1893] (Kjaerskov 1893). — *Acrandra glazioviana* (Kjaersk.) Burret, *Repert. Spec. Nov. Regni Veg.* 50: 59 (Burret 1941).

Eugenia pleurosiphonea Diels, *Verh. Bot. Vereins Prov. Brandenburg* 48: 191 ["1906" publ. 8 Mar. 1907] (Diels 1907).

Luma bergii Herter, *Revista Sudamer. Bot.* 7: 219 (Herter 1943).

Eugenia kiaerskoviana Mattos & D.Legrand, *Loefgrenia* 67: 30 (Mattos & Legrand 1975), "kiaerskovana".

NOTE. — The epithet *pisonis*, from "Piso", the latinized name of Willem Pies, is not to be corrected (Turland et al. 2018: Art. 60.9).

VERNACULAR NAMES. — Pa: arak-kamwi, pakusin-amana • Wp: akusi yeti, miku ka'a, tamukwálē áká • Br: araçá.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 995*.

INVENTORY DATA (FG). — 3 trees in 3 plots; F_{max} < 1 %; dbh_{inv} = 26.5 cm.

[1246] *Eugenia muricata* DC.

Prodr. [A. P. de Candolle] 3: 283 [mid Mar. 1828] (Candolle 1828). — *Myrtus muricata* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 283 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Myrtus riparia* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 283 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Eugenia rugosa Ruiz & Pav. ex DC., *Prodr. [A. P. de Candolle]* 3: 280 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Eugenia riparia DC., *Prodr. [A. P. de Candolle]* 3: 283 [mid Mar. 1828] (Candolle 1828).

Eugenia rutidocarpa Ruiz & Pav. ex G.Don, *Gen. Hist.* 2: 865 [Oct. 1832] (Don 1832). — *Eugenia ripidocarpa* Ruiz & Pav., *Anal. Inst. Bot. Cavanilles* 15: 219 (Ruiz & Pavón 1958), "rypidocarpa" on plate, nom. illeg. superfl. (based on the type of *E. rutidocarpa*).

Eugenia lugens O.Berg, *Linnaea* 27 (2-3): 299 ["1854" publ. Jan. 1856] (Berg 1856). — *Myrtus lugens* Poepp. ex O.Berg, *Linnaea* 27 (2-3): 300 ["1854" publ. Jan. 1856] (Berg 1856), nom. nud. pro syn.

Eugenia rugosa Ruiz ex O.Berg, *Linnaea* 27 (2-3): 300 ["1854" publ. Jan. 1856] (Berg 1856).

Eugenia amazonica O.Berg, *Fl. Bras. [Martius]* 14 (1): 322 [15 May 1857] (Berg 1857).

Eugenia racemifera O.Berg, *Fl. Bras. [Martius]* 14 (1): 322 [15 May 1857] (Berg 1857).

Eugenia fenziana O.Berg, *Fl. Bras. [Martius]* 14 (1): 323 [15 May 1857] (Berg 1857).

Eugenia casarettiana O.Berg, *Fl. Bras. [Martius]* 14 (1): 520 [1 Feb. 1858] (Berg 1858).

Eugenia muricata var. *guyanensis* O.Berg, *Linnaea* 30: 695 [Mar. 1861] (Berg 1861). — *Eugenia spicata* Rich. ex O.Berg, *Linnaea* 30: 695 [Mar. 1861] (Berg 1861), nom. nud. pro syn.

Eugenia calothysra Diels, *Verh. Bot. Vereins Prov. Brandenburg* 48: 189 ["1906" publ. 8 Mar. 1907] (Diels 1907).

NOTE. — Although based on the unpublished plate of Ruiz & Pavón, *E. rutidocarpa* is validly published, with a description.

VERNACULAR NAMES. — Wp: pila lea.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *L.C. Richard 76* (original material of *E. muricata* var. *guyanensis*: P[P01902488]).

SIZE. — Peru, Loreto. *J.J. Pipoly et al. 13026* (MO), 18 m × 15 cm.

[1247] *Eugenia neograndifolia* Mattos

Loefgrenia 120: 10 [Mar. 2005] (Mattos 2005). — *Calycorectes grandifolius* O.Berg, *Linnaea* 27 (2-3): 317 ["1854" publ. Jan. 1856] (Berg 1856).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: avitkat-purubumna.

HERBARIUM DATA (FG). — 56 collections at CAY. Sel. exs.: *B. Dutrèvre* 399 (CAY); *H.S. Irwin et al.* 47561 (NY), 7 m × 10 cm.

[1248] *Eugenia omissa* McVaugh

Mem. New York Bot. Gard. 18 (2): 197 (McVaugh 1969).

VERNACULAR NAMES. — Pa: aku-aiwut • Wp: kumati, kumati u, yawa poi li sī.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *R.A.A. Oldeman & C. Sastre* 156.

SIZE. — Venezuela, Amazonas. *L. Delgado* 840 (MO), 10 m.

[1249] *Eugenia patrisii* Vahl

Eclog. Amer. 2: 35 (Vahl 1798). — *Myrtus patrisii* (Vahl) Spreng., *Syst. Veg. [Sprengel]* 2: 480 [Jan.-May 1825] (Sprengel 1825). — *Stenocalyx patrisii* (Vahl) O.Berg, *Linnaea* 29 (2): 247 [June 1858] (Berg 1858).

Eugenia inocarpa DC., *Prodr. [A. P. de Candolle]* 3: 264 [mid Mar. 1828] (Candolle 1828). — *Myrtus inocarpa* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 264 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Eugenia berlynensis O.Berg, *Linnaea* 27 (4): 468 ["1854" publ. Feb. 1856] (Berg 1856).

Eugenia teffensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 255 [15 May 1857] (Berg 1857).

Eugenia vellozoi O.Berg, *Fl. Bras. [Martius]* 14 (1): 255 [15 May 1857] (Berg 1857), "vellozii".

Eugenia teffensis var. *subcordata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 256 [15 May 1857] (Berg 1857).

Eugenia teffensis var. *truncata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 256 [15 May 1857] (Berg 1857).

Stenocalyx patrisii var. *parvifolius* O.Berg, *Linnaea* 30: 699 [Mar. 1861] (Berg 1861).

Stenocalyx patrisii var. *grandifolius* O.Berg, *Linnaea* 30: 700 [Mar. 1861] (Berg 1861).

VERNACULAR NAMES. — Pa: inam-etni • Ka: kuapitano, kuwapi-tyano • Te: uwa pitāg ndi • Wp: iwa pitā, yāwī ki'iy • Nt: busi siriz, paawisi besi • Cr: siriz-gran-bwa, siriz-ronde • Fr: cerise de Cayenne • Br: fruta-de-jaboti, pitanga, ubaia.

HERBARIUM DATA (FG). — 74 collections at CAY. Sel. exs.: *J.P.B. von Rohr s.n.* (type C[C10015707]).

INVENTORY DATA (FG). — 115 trees in 68 plots; $F_{\max} = 2.9\%$; $\text{dbh}_{\text{inv}} = 26.8$ cm.

[1250] *Eugenia polystachya* Rich.

Actes Soc. Hist. Nat. Paris 1: 110 [Oct. 1792] (Richard 1792).

Eugenia patens Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 124 [24 Sep. 1813] (Poiret 1813).

Eugenia verruculosa DC., *Prodr. [A. P. de Candolle]* 3: 284 [mid Mar. 1828] (Candolle 1828). — *Myrtus verruculosa* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 285 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Eugenia schlechtendaliana O.Berg, *Fl. Bras. [Martius]* 14 (1): 321 [15 May 1857] (Berg 1857).

Eugenia forsteri O.Berg, *Linnaea* 29 (2): 241 [June 1858] (Berg 1858).

VERNACULAR NAMES. — Pa: kuyau-kamwi • Wp: miku ka'a, moyu'i, tamukwälé áká, wila tipita toto, yawayi lü.

HERBARIUM DATA (FG). — 57 collections at CAY. Sel. exs.: *L.C. Richard* 77 (original material P[P01902610, P01902611]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $\text{dbh}_{\text{inv}} = 26$ cm.

[1251] *Eugenia pseudopsidium* Jacq.

Enum. Syst. Pl.: 23 [Aug.-Sep. 1760] (Jacquin 1760). — *Myrtus pseudopsidium* (Jacq.) Spreng., *Syst. Veg. [Sprengel]* 2: 481 [Jan.-May 1825] (Sprengel 1825).

Eugenia portoricensis DC., *Prodr. [A. P. de Candolle]* 3: 266 [mid Mar. 1828] (Candolle 1828). — *Myrtus willdenowii* Spreng. var. *portoricensis* Spreng. ex DC., *Prodr. [A. P. de Candolle]* 3: 266 [mid Mar. 1828] (Candolle 1828), "Portoricensis", *nom. nud. pro syn.* — *Stenocalyx portoricensis* (DC.) O.Berg, *Linnaea* 29 (2): 246 [June 1858] (Berg 1858). — *Eugenia pseudopsidium* var. *portoricensis* (DC.) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 647 [26 Feb. 1895] (Krug & Urban 1895).

Eugenia portoricensis var. *brevipes* DC., *Prodr. [A. P. de Candolle]* 3: 266 [mid Mar. 1828] (Candolle 1828).

Eugenia psidiooides DC., *Prodr. [A. P. de Candolle]* 3: 268 [mid Mar. 1828] (Candolle 1828).

Stenocalyx pseudopsidium O.Berg, *Linnaea* 27 (2-3): 314 ["1854" publ. Jan. 1856] (Berg 1856), "Pseudo-Psidium". — *Eugenia pseudopsidium* Willd. ex O.Berg, *Linnaea* 27 (2-3): 314 ["1854" publ. Jan. 1856] (Berg 1856), *nom. nud. pro syn., non Jacq.*

Eugenia compta Rich. ex O.Berg, *Linnaea* 30: 677 [Mar. 1861] (Berg 1861).

Eugenia prieurii O.Berg, *Linnaea* 30: 681 [Mar. 1861] (Berg 1861). — *Eugenia prieurii* Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20:* 188 (Sagot 1885), "Prieurei", isonym.

Eugenia prieurii var. *robusta* O.Berg, *Linnaea* 30: 681 [Mar. 1861] (Berg 1861), *nom. inval.*

Eugenia prieurii var. *tenuiramis* O.Berg, *Linnaea* 30: 682 [Mar. 1861] (Berg 1861).

Eugenia willdenowii Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 81 [Jan. 1893] (Niedenzu 1893).

Eugenia pseudopsidium var. *genuina* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 647 [26 Feb. 1895] (Krug & Urban 1895), *nom. inval.* (Turland et al. 2018: Art. 24.3).

Eugenia megalocarpa Urb., *Symb. Antill. [Urban]* 5 (3): 444 [20 May 1908] (*Urban* 1908).

Eugenia cryptadena Amshoff, *Recueil Trav. Bot. Néerl.* 42: 17 (Amshoff 1950).

NOTES. — Since the type of *Eugenia prieurii* var. *robusta* O.Berg is that of *E. prieurii* O.Berg, the former is invalid (it should be var. *prieurii*). *Eugenia prieurii* Sagot is based on the very same type, and is thus an isonym. *Stenocalyx pseudopsidium* O.Berg is not based on *E. pseudopsidium* Jacq., because both taxa are treated as distinct in the same publication (Berg 1856).

VERNACULAR NAMES. — Pa: kagegut-puvemna, kagegut-puveyo, karegut-puvemna • Wp: piki li • Br: araçázhino, murta.

HERBARIUM DATA (FG). — 124 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 956.

INVENTORY DATA (FG). — 56 trees in 46 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.2$ cm.

[1252] *Eugenia sinemariensis* Aubl.

Hist. Pl. Guiane 1: 501 [Jun.-Dec. 1775] (*Aublet* 1775). — *Myrtus sinemariensis* (Aubl.) Spreng., *Syst. Veg. [Sprengel]* 2: 483 [Jan.-May 1825] (*Sprengel* 1825). — *Epleienda sinemariensis* (Aubl.) Raf., *Sylva Tellur.*: 107 (*Rafinesque* 1838).

Eugenia coffeeifolia DC., *Prod. [A. P. de Candolle]* 3: 272 [mid Mar. 1828] (*Candolle* 1828), “*coffeeifolia*”, syn. nov.

Myrciaria ramiflora O.Berg, *Linnaea* 27 (2-3): 334 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia coffeeifolia var. *grandifolia* O.Berg, *Linnaea* 29 (2): 231 [June 1858] (Berg 1858).

Eugenia coffeeifolia var. *parvifolia* O.Berg, *Linnaea* 29 (2): 231 [June 1858] (Berg 1858).

Eugenia melinonii Sagot, *Ann. Sci. Nat., Bot. sér. 6, 20:* 194 (*Sagot* 1885), “*Melinonis*”.

NOTES. — Govaerts *et al.* (2008) provisionally placed *E. sinemariensis* in synonymy under *E. coffeeifolia*. The species known as *E. coffeeifolia* is a very common understorey tree in the forests of Northern French Guiana, and its red, edible and fleshy cherry-sized berries are frequently seen. It would therefore have been surprising if Aublet had never encountered it. Leaves from the type of *E. sinemariensis* at LINN (LINN-HS 883.12) match perfectly with *E. coffeeifolia*, and its fruit was compared by Aublet to the azerola (*Malpighia emarginata* Sessé & Moc. ex DC.), i.e. a red, cherry-sized fleshy berry. Aublet’s specimen filed as *E. arivoa* at BM (BM000953762) and labelled by Amshoff “Type of *Eug. sinemariensis* Aubl. ???”, is actually *Mouriri sagotiana* Triana, probably from the same collection as the lower part of the supposed original material of *Eugenia arivoa* Aubl. (BM000953761). The epithet “*melinonis*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland *et al.* 2018: Art. 60.8).

VERNACULAR NAMES. — Pa: inam-etni-duwō, inam-etni-kamwi, inamkat-duwē, inamkat-wahuyo • Ka: alaidya, pasyalawa • Wp: mila yowa, pila lea, wila yowa • Nt: baaka bee sii, gibomasa • Br: murta.

HERBARIUM DATA (FG). — 175 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN, LINN-HS 883.12).

INVENTORY DATA (FG). — 209 trees in 75 plots; $F_{\max} = 3.8\%$; $dbh_{\text{inv}} = 36.7$ cm.

[1253] *Eugenia spruceana* O.Berg

Fl. Bras. [Martius] 14 (1): 257 [15 May 1857] (Berg 1857).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al.* 8765.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.3$ cm.

[1254] *Eugenia stictopetala* DC.

Prod. [A. P. de Candolle] 3: 270 [mid Mar. 1828] (*Candolle* 1828).

Eugenia martiusiana DC., *Prod. [A. P. de Candolle]* 3: 269 [mid Mar. 1828] (*Candolle* 1828).

Eugenia ochra O.Berg, *Linnaea* 27 (2-3): 216 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia roraimana O.Berg, *Linnaea* 27 (2-3): 219 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia tapacumensis O.Berg, *Linnaea* 27 (2-3): 222 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia tapacumensis var. *angustifolia* O.Berg, *Linnaea* 27 (2-3): 222 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia tapacumensis var. *latifolia* O.Berg, *Linnaea* 27 (2-3): 222 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia piauhiensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 285 [15 May 1857] (Berg 1857).

Eugenia eschholtziana O.Berg, *Fl. Bras. [Martius]* 14 (1): 582 [15 Jan. 1859] (Berg 1859).

Eugenia eschholtziana var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 582 [15 Jan. 1859] (Berg 1859).

Eugenia eschholtziana var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 582 [15 Jan. 1859] (Berg 1859).

NOTES. — Govaerts *et al.* (2008) placed *E. doniana* O.Berg in synonymy under *E. stictopetala* DC. In fact, *E. doniana* is an error in Berg’s index of names (1859: 643-644). Under *Eugenia* is written “*Doniana* Bg, 518”, that is on p. 518, where is found *Myrcia doniana* O.Berg, a species accepted by Govaerts *et al.* (2008) and unrelated to *E. stictopetala*. We therefore exclude *E. doniana* from synonymy.

VERNACULAR NAMES. — Wp: kumati, kuya lá.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 992.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.8$ cm.

[1255] *Eugenia tenuiflora* Mazine

Kew Bull. 64 (1): 149 [May 2009] (Mazine 2009).

HERBARIUM DATA (FG). — A single collection, *S.A. Mori et al.* 24757.

SIZE. — Brazil, Amazonas. *G.T. Prance et al.* 3222 (holo-, MG, not seen; iso-, F[V0109955F], K[K000913195], US[01898006]), 12 m × 10 cm.

[1256] *Eugenia tetramera*
 (McVaugh) M.L.Kawas. & B.Holst
 (Fig. 42A)

Brittonia 46 (2): 142 (Kawasaki & Holst 1994). — *Campomanesia tetramera* McVaugh, *Mem. New York Bot. Gard.* 18 (2): 239 (McVaugh 1969).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: inam-etni-kamwi • Nt: busi banda.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2461*.

INVENTORY DATA (FG). — 62 trees in 36 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 28.6$ cm.

[1257] *Eugenia wentii* Amshoff
 (Fig. 42B)

Recueil Trav. Bot. Néerl. 39: 158 (Amshoff 1942). — *Phyllocalyx wentii* Amshoff, *Recueil Trav. Bot. Néerl.* 39: 158 (Amshoff 1942).

Calycorectes macrocalyx Rusby, *Mem. New York Bot. Gard.* 7: 313 (Rusby 1927). — *Eugenia macrocalyx* (Rusby) McVaugh, *Fiel-diana, Bot.* 29 (3): 212 [30 Nov. 1956] (McVaugh 1956), nom. illeg. hom., non Mart. ex B.D.Jacks. (Jackson 1893).

NOTE. — *Eugenia wentii* and *Phyllocalyx wentii* are alternative names, but as they were published before January 1, 1953, they are both valid (Turland *et al.* 2018: Art. 36.3).

VERNACULAR NAMES. — Pa: kagegut, karegut • Wp: yawayi lū.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5794*.

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 27.5$ cm.

[1258] *Eugenia wullschaegeliana* Amshoff

Natuurw. Stud. Suriname & Curacao 2: 20. (Amshoff 1948).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *Y. Caraglio 508*.

SIZE. — Up to 18 m tall (McVaugh 1969).

[1259] *Eugenia* sp. A

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-10”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4660*.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 27.1$ cm.

[1260] *Eugenia* sp. B

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-12”.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2714*.

INVENTORY DATA (FG). — 6 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 26.4$ cm.

[1261] *Eugenia* sp. C

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-13”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2657*.

INVENTORY DATA (FG). — 6 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 36.3$ cm.

[1262] *Eugenia* sp. D

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-16”.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2309*.

INVENTORY DATA (FG). — 15 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.1$ cm.

[1263] *Eugenia* sp. E

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-19”.

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 2116*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 10.1$ cm.

[1264] *Eugenia* sp. F

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-20”.

HERBARIUM DATA (FG). — 2 collections at CAY: *D. Paget 201; D. Loubry 2392*.

INVENTORY DATA (FG). — 42 trees in 10 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 50$ cm.

[1265] *Eugenia* sp. G

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-21”.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3670*.

INVENTORY DATA (FG). — 39 trees in 34 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 37.7$ cm.

[1266] *Eugenia* sp. H

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-22”.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Paget 204*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 31\text{ cm}$.

[1267] *Eugenia* sp. I

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-27”.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier 858*.

INVENTORY DATA (FG). — 3 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 19.6\text{ cm}$.

[1268] *Eugenia* sp. J

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-29”.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Larpin 687*.

INVENTORY DATA (FG). — 36 trees in 9 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 37.6\text{ cm}$.

[1269] *Eugenia* sp. K

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-31”.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *J.-F. Viliers 1873*, height 15 m.

[1270] *Eugenia* sp. L

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-6”.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 23307*.

INVENTORY DATA (FG). — 12 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.8\text{ cm}$.

[1271] *Eugenia* sp. M

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-7”.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 4780*, $\text{dbh} = 14\text{ cm}$.

[1272] *Eugenia* sp. N

NOTE. — This is the “*Eugenia* sp. B” in Holst & Kawasaki (2002); collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-8”.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15192*, 10 m × 13 cm.

[1273] *Eugenia* sp. O
(Fig. 41D, E)

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Eugenia* FG-9”.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2075*.

INVENTORY DATA (FG). — 14 trees in 14 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 27.7\text{ cm}$.

[1274] *Eugenia* sp. P

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier 828*.

INVENTORY DATA (FG). — 16 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 34.4\text{ cm}$.

[1275] *Eugenia* sp. Q

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & M.-F. Prévost 3102*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 18.3\text{ cm}$.

[1276] *Eugenia* sp. R

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2360*.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 17.2\text{ cm}$.

Genus *Melaleuca* L.

[1277] *Melaleuca leucadendra* (L.) L.

Mant. Pl. 1: 105 (Linnaeus 1767). — *Myrtus leucadendra* L., *Herb. Amb.*: 9 (Linnaeus 1754). — *Leptospermum leucadendrum* (L.) J.R.Forst. & G.Forst., *Char. Gen. Pl.*: 72 [“1776” publ. late Nov.–early Dec. 1775] (Forster & Forster 1775), “*Leucadendron*”. — *Meladendron leucocladum* St.-Lag., *Ann. Soc. Bot. Lyon* 7: 64 (Saint-Lager 1880), *nom. illeg. superfl.* [based on *Melaleuca leucadendra* (L.) L.]. — *Kajuputi leucadendra* (L.) Farw., *Druggists' Circ.* 61: 175 (Farwell 1917), “*Leucadendron*”.

Myrtus saligna Burm.f., *Fl. Indica* [N.L. Burman]: 116 (Burman 1768).

Myrtus alba Noronha, *Verb. Batav. Genootsch. Kunst.* 5 (4): 20 (Noronha 1790).

Metrosideros coriacea K.D.Koenig & Sims, *Prod. Stirp. Chap. Allerton* 352 [Nov.-Dec. 1796] (Koenig & Sims 1796).

Melaleuca rigida Roxb., *Fl. Ind.*, ed. 1832, 3: 399 (Roxburgh 1832).

Melaleuca leucadendra var. *angusta* C.Rivièrè, *Bull. Soc. Natl. Acclim. France*, sér. 3, 9: 537 (Rivièrè 1882).

Melaleuca leucadendra var. *cunninghamii* F.M.Bailey, Syn. Queensl. Fl. 171 (Bailey 1883).

Melaleuca leucadendra var. *lanceifolia* F.M.Bailey, Syn. Queensl. Fl. 170 (Bailey 1883).

Melaleuca mimosoides A.Cunn. ex Schauer, *Repert. Bot. Syst. [Walpers]* 2 (5): 927 [28-30 Dec. 1843] (Schauer 1843). — *Melaleuca leucadendra* var. *mimosoides* (A.Cunn. ex Schauer) Cheel, *Fl. N. Territory*: 295 (Cheel 1917).

Melaleuca amboinensis Gand., *Bull. Soc. Bot. France* 65: 26, (Gandoger 1918), “*abominensis*”.

NOTE. — A species native to East Indonesia and North Australia, introduced in French Guiana and now invasive in a few coastal savannas.

VERNACULAR NAMES. — Fr: niaouli.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *C. Delnatté* 1549.

[1278] *Melaleuca quinquenervia* (Cav.) S.T.Blake

Proc. Roy. Soc. Queensland 69: 76 (Blake 1958). — *Metrosideros quinquenervia* Cav., *Icon. [Cavanilles]* 4: 19 (Cavanilles 1797).

Melaleuca leucadendra var. *angustifolia* L.f., *Suppl. Pl.*: 342 [“1781” publ. Apr. 1782] (Linnaeus 1782), “Leucadendron”. — *Kajuputia leucadendra* var. *angustifolia* (L.f.) Farw., *Druggists' Circ.* 61: 175 (Farwell 1917), “Leucadendron, Angustifolia”. — *Melaleuca viridiflora* Sol. ex Gaertn. var. *angustifolia* (L.f.) Byrnes, *Austrobaileya* 2: 74 (Byrnes 1984).

Metrosideros coriacea Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 685 [3 Sep. 1814] (Poirier 1814). — *Melaleuca leucadendra* var. *coriacea* (Poir.) Cheel, *Fl. N. Territory* 297 (Cheel 1917), “Leucadendron”.

Metrosideros albida Sieber ex DC., *Prod. [A. P. de Candolle]* 3: 212 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Melaleuca viridiflora Sol. ex Gaertn. var. *rubriflora* Brongn. & Gris, *Bull. Soc. Bot. France* 11: 183 (Brongniart & Gris 1864). — *Melaleuca rubriflora* Vieill. ex Brongn. & Gris, *Bull. Soc. Bot. France* 11: 183 (Brongniart & Gris 1864), nom. nud. pro syn.

Melaleuca maidenii R.T.Baker, *Proc. Linn. Soc. New South Wales* 38 (4): 598 [“1913” publ. 23 Mar. 1914] (Baker 1914), “Maideni”.

Melaleuca smithii R.T.Baker, *Proc. Linn. Soc. New South Wales* 38 (4): 599 [“1913” publ. 23 Mar. 1914] (Baker 1914).

Melaleuca leucadendra var. *albida* Cheel, *Fl. N. Territory* 301 (Cheel 1917), “Leucadendron”.

NOTE. — A species native to New Guinea, eastern Australia and New Caledonia, introduced in French Guiana and now invasive in a few coastal savannas.

VERNACULAR NAMES. — Fr: niaouli.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Loubry* 1626.

Genus *Myrcia* DC. ex Guill.

[1279] *Myrcia amazonica* DC.

Prod. [A. P. de Candolle] 3: 250 [mid Mar. 1828] (Candolle 1828). — *Myrtus amazonica* Mart. ex DC., *Prod. [A. P. de Candolle]* 3: 250 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Aulomyrcia amazonica* (DC.) O.Berg, *Linnaea* 27 (1): 41 [“1854” publ. Nov. 1855] (Berg 1855).

Myrcia leptoclada DC., *Prod. [A. P. de Candolle]* 3: 244 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia leptoclada* (DC.) O.Berg, *Linnaea* 27 (1): 40 [“1854” publ. Nov. 1855] (Berg 1855).

Myrcia corymbosa DC., *Prod. [A. P. de Candolle]* 3: 252 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia corymbosa* (DC.) O.Berg, *Linnaea* 27 (1): 41 [“1854” publ. Nov. 1855] (Berg 1855).

Myrcia detergens Miq., *Linnaea* 22: 795 [“1849” publ. May 1850] (Miquel 1850). — *Aulomyrcia detergens* (Miq.) O.Berg, *Linnaea* 27 (1): 46 [“1854” publ. Nov. 1855] (Berg 1855).

Aulomyrcia hostmanniana O.Berg, *Linnaea* 27 (1): 42 [“1854” publ. Nov. 1855] (Berg 1855). — *Myrcia hostmanniana* (O.Berg) Kjaersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kjaerskov 1893).

Aulomyrcia hostmanniana var. *gracilior* O.Berg, *Linnaea* 27 (1): 43 [“1854” publ. Nov. 1855] (Berg 1855).

Aulomyrcia hostmanniana var. *robustior* O.Berg, *Linnaea* 27 (1): 43 [“1854” publ. Nov. 1855] (Berg 1855). — *Myrcia hostmanniana* var. *robustior* (O.Berg) Kjaersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kjaerskov 1893).

Aulomyrcia paraensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 76 [15 May 1857] (Berg 1857). — *Myrcia paraensis* (O.Berg) Kjaersk., *Enum. Myrt. Bras.* 78 [Oct.-Nov. 1893] (Kjaerskov 1893), “Paraënsis”.

Aulomyrcia spruceana O.Berg, *Fl. Bras. [Martius]* 14 (1): 76 [15 May 1857] (Berg 1857). — *Myrcia sprucei* Mattos, *Loefgrenia* 126: 3 [Sep. 2008] (Mattos 2008), nom. illeg. superfl. (based on *Aulomyrcia spruceana*).

Aulomyrcia detergens var. *depauperata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 77 [15 May 1857] (Berg 1857).

Aulomyrcia detergens var. *dives* O.Berg, *Fl. Bras. [Martius]* 14 (1): 77 [15 May 1857] (Berg 1857).

Myrcia oitchi O.Berg, *Fl. Bras. [Martius]* 14 (1): 563 [15 Jan. 1859] (Berg 1859).

Aulomyrcia vauthieriana O.Berg, *Linnaea* 30: 655 [Mar. 1861] (Berg 1861), “Vauthiereana”.

Myrcia leptoclada var. *glazioviana* Kjaersk., *Enum. Myrt. Bras.* 78 [Oct.-Nov. 1893] (Kjaerskov 1893).

Myrcia lundiana Kjaersk., *Enum. Myrt. Bras.* 78 [Oct.-Nov. 1893] (Kjaerskov 1893).

Myrcia gentlei Lundell, *Wrightia* 2 (4): 214 (Lundell 1961). — *Myrcia minoriflora* Mattos, *Loefgrenia* 129: 2 (Mattos 2009), nom. illeg. superfl. (based on *Gomidesia minutiflora*).

Myrcia sagotii Mattos, *Loefgrenia* 125: 4 [July 2008] (Mattos 2008).




FIG. 42. — Myrtaceae: **A**, *Eugenia tetramera* (McVaugh) M.L.Kawas. & B.Holst (J.-F. Molino & D. Sabatier 2461); **B**, *Eugenia wentii* Amshoff (D. Sabatier & J.-F. Molino 5794); **C**, *Myrcia bracteata* (Rich.) DC. (J.-F. Molino & D. Sabatier 2442); **D**, *Myrcia platyclada* DC. (M.-F. Prévost & S. Gonzalez 5314). A, C, © J.-F. Molino/IRD; B, D, © D. Sabatier/IRD.

Gomidesia minutiflora Mattos & D.Legrand, *Loefgrenia* 67: 15 (Mattos & Legrand 1975).

VERNACULAR NAMES. — Pa: arak-kwikwiyo • Ka: asaikiabe, seityape, tapilen seityape • Wp: piki li sili • Br: araçá.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: [D. Sabatier & M.-F. Prévost 2249](#).

INVENTORY DATA (FG). — 18 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{inv} = 41.1$ cm.

[1280] *Myrcia amshoffae* (McVaugh) A.R.Lourenço & E.Lucas

Phytotaxa 373 (1): 73 [23 Oct. 2018] (Lourenço & Lucas 2018). — *Calypranthes amshoffae* McVaugh, *Mem. New York Bot. Gard.* 10 (1): 70 (McVaugh 1958).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: pila lea.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: [D. Larpin 846](#).

INVENTORY DATA (FG). — 6 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.5$ cm.

[1281] *Myrcia attenuata* M.F.Santos

Phytotaxa 234 (2): 163 (Santos 2015).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al.* 6503 (holo-, B, not seen; iso-, BR[BR0000013176433], CAY[CAY097245], P[P05233192]); *R.A.A. Oldeman T-650* [paratype], 25 m × 20 cm.

[1282] *Myrcia bracteata* (Rich.) DC.
(Fig. 42C)

Prodr. [A. P. de Candolle] 3: 245 [mid Mar. 1828] (Candolle 1828). — *Eugenia bracteata* Rich., *Actes Soc. Hist. Nat. Paris* 1: 110 [Oct. 1792] (Richard 1792). — *Myrtus bracteata* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 245 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*

Myrcia lanceolata Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 329 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832).

Myrcia lanceolata var. *angustifolia* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 329 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832).

Myrcia lanceolata var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 155 [15 May 1857] (Berg 1857).

Myrcia lanceolata var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 155 [15 May 1857] (Berg 1857).

Myrcia lanceolata var. *avenia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 156 [15 May 1857] (Berg 1857).

Myrcia lanceolata var. *racemosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 156 [15 May 1857] (Berg 1857).

Myrcia hirtellifolia Gleason, *Bull. Torrey Bot. Club* 58 (7): 411 [Oct. 1931] (Gleason 1931), “*hirtellaefolia*”.

Eugenia hirsuta Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 15: 220 (Ruiz & Pavón 1958).

VERNACULAR NAMES. — Pa: arak-priyu, kagegut, kagegut-ahavukune, kagegut-puvemna, karegut • Wp: kumesi sili, piki li sili, pila lea • Cr: kouamaté • Br: cumaté, murtinha.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J.B. Leblond* 110 (original material BR[BR000005238668], G[G00223176]).

SIZE. — Ecuador. *C. Kajekai* 356 (MO), 10 m.

[1283] *Myrcia coumete* (Aubl.) DC.

Prodr. [A. P. de Candolle] 3: 245 [mid Mar. 1828] (Candolle 1828), “*Coumeta*”. — *Eugenia coumete* Aubl., *Hist. Pl. Guiane* 1: 497 [Jun.-Dec. 1775] (Aublet 1775), “*Coumète*”. — *Myrtus coumete* (Aubl.) Spreng., *Syst. Veg. [Sprengel]* 2: 488 [Jan.-May 1825] (Sprengel 1825), “*Comète*”. — *Cumetea alba* Raf., *Sylva Tellur.*: 106 (Rafinesque 1838), nom. illeg. superfl. (based on *Eugenia coumete*). — *Aulomyrcia coumete* (Aubl.) O.Berg, *Linnaea* 27 (1): 60 [“1854” publ. Nov. 1855] (Berg 1855), “*Coumeta*”.

Myrtus carnea G.Mey., *Prim. Fl. Esseq.* 191 [Nov. 1818] (Meyer 1818), “*Myrthus*”. — *Myrcia carnea* (G.Mey.) DC., *Prodr. [A. P. de Candolle]* 3: 245 [mid Mar. 1828] (Candolle 1828).

VERNACULAR NAMES. — Ka: tapuliyuipo • Wp: kumesi sili • Wn: pasiwt • Nt: dondowku, lebi tongo • Br: cumaté-rana, pitanga-miuda.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *F. Halle* 629.

SIZE. — Costa Rica. *L. Acosta & L.A. González* 14 (MO), 16 m × 11 cm.

[1284] *Myrcia cuprea* (O.Berg) Kiaersk.

Enum. Myrt. Bras. 95 [Oct.-Nov. 1893] (Kiaerskov 1893). — *Aulomyrcia cuprea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 77 [15 May 1857] (Berg 1857).

Aulomyrcia chrysophylla O.Berg, *Fl. Bras. [Martius]* 14 (1): 125 [15 May 1857] (Berg 1857). — *Myrcia chrysophylla* (O.Berg) Mattos, *Loefgrenia* 126: 4 [Sep. 2008] (Mattos 2008).

VERNACULAR NAMES. — Wp: wai piyū sili, wai sili.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-849A, 15 m × 20 cm.

[1285] *Myrcia decorticans* DC.

Prodr. [A. P. de Candolle] 3: 252 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia polymorpha* O.Berg, *Linnaea* 27 (1): 46 [“1854” publ. Nov. 1855] (Berg 1855), nom. illeg. superfl. (based on *Myrcia decorticans*). — *Aulomyrcia polymorpha* var. *decorticans* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 78 [15 May 1857] (Berg 1857).

Myrcia leucophlaea DC., *Prodr. [A. P. de Candolle]* 3: 247 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia polymorpha* var. *leucophloea* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 78 [15 May 1857] (Berg 1857).

Myrcia duriuscula Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia polymorpha* var. *duriuscula* (Mart. ex DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 78 [15 May 1857] (Berg 1857).

Myrcia lasiopus DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828). — *Eugenia lasiopus* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Aulomyrcia polymorpha* var. *lasiopus* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 78 [15 May 1857] (Berg 1857).

Aulomyrcia salzmannii O.Berg, *Fl. Bras. [Martius]* 14 (1): 116 [15 May 1857] (Berg 1857).

Calyptanthes tobagensis Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 593 [26 Feb. 1895] (Krug & Urban 1895). — *Myrcia tobagensis* (Krug & Urb.) Urb., *Repert. Spec. Nov. Regni Veg.* 14: 336 (Urban 1916). — *Aulomyrcia tobagensis* (Krug & Urb.) Amshoff, *Recueil Trav. Bot. Néerl.* 39: 155 (Amshoff 1942).

Aulomyrcia tetramera Amshoff, *Recueil Trav. Bot. Néerl.* 42: 9 (Amshoff 1950). — *Myrcia tetramera* (Amshoff) Lemée, *Fl. Guyane Franç.* 3: 150 (Lemée 1954).

Marlierea gleasonii McVaugh, *Mem. New York Bot. Gard.* 10 (1): 83 (McVaugh 1958).

VERNACULAR NAMES. — Pa: aku-aiwut, inam-etni-duwó • Wp: pakea'i • Nt: lebi tongo • Br: cumaté-rana.

HERBARIUM DATA (FG). — 71 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3762.

INVENTORY DATA (FG). — 93 trees in 49 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 26.7\text{ cm}$.

[1286] *Myrcia deflexa* (Poir.) DC.

Prodr. [A. P. de Candolle] 3: 244 [mid Mar. 1828] (Candolle 1828). — *Eugenia deflexa* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 124 [24 Sep. 1813] (Poirier 1813). — *Myrtus deflexa* (Poir.) Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 142 [6 Aug. 1823] (Kunth 1823). — *Myrcia humboldtiiana* DC., *Prodr. [A. P. de Candolle]* 3: 256 [mid Mar. 1828] (Candolle 1828), nom. illeg. superfl. (based on *Myrtus deflexa*).

Myrcia crassinervia DC., *Prodr. [A. P. de Candolle]* 3: 245 [mid Mar. 1828] (Candolle 1828).

Eugenia paniculata Sieber ex C.Presl, *Isis [Oken]* 21: 274 (Presl 1828).

Myrcia duchassaingiana O.Berg, *Linnaea* 27 (1): 88 [“1854” publ. Nov. 1855] (Berg 1855).

Myrcia ferruginea var. *domingensis* O.Berg, *Linnaea* 27 (1): 90 [“1854” publ. Nov. 1855] (Berg 1855).

Myrcia humboldtiana var. *orinocensis* O.Berg, *Linnaea* 27 (1): 120 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia humboldtiana var. *caribaea* O.Berg, *Linnaea* 27 (1): 121 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia sulcata O.Berg, *Linnaea* 30: 667 [Mar. 1861] (Berg 1861). — *Eugenia sulcata* Rich. ex O.Berg, *Linnaea* 30: 667 [Mar. 1861] (Berg 1861), *nom. nud. pro syn.*

Myrcia deflexa var. *dussii* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 588 [26 Feb. 1895] (Krug & Urban 1895). — *Myrcia dussii* (Krug & Urb.) T.Durand & B.D.Jacks., *Index Kew. Suppl.* 1: 285 [20 Nov. 1903] (Durand & Jackson 1903).

NOTE. — Govaerts et al. (2008) wrongly place *Myrcia humboldtiana* DC. in synonymy under *M. splendens* (Sw.) DC. *M. humboldtiana* is undoubtedly based on *Myrtus deflexa* (Poir.) DC., as both the name and type of the latter are cited by Candolle (1828: 256).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *Service Forestier 5147*.

SIZE. — Up to 30 cm dbh (Amshoff 1948).

[1287] *Myrcia fasciculata* (O.Berg) K.Campbell & K.Samra

Phytotaxa 406 (3): 148 (Campbell & Samra 2019). — *Calyptranthes fasciculata* O.Berg, *Linnaea* 27 (1): 31 ["1854" publ. Nov. 1855] (Berg 1855). — *Chytraculia fasciculata* (O.Berg) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891). — *Calyptranthes fasciculata* var. *genuina* Stehlé & Quentin, *Fl. Guadeloupe [Stehlé]* 2 (3): 62 (Stehlé & Quentin 1949), *nom. inval.* (Turland et al. 2018: Art. 24.3).

Calyptranthes sericea Griseb., *Fl. Brit. W.I. [Grisebach]* 233 [late 1860] (Grisebach 1860).

Calyptranthes sericea var. *hahni* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 602 [26 Feb. 1895] (Krug & Urban 1895). — *Calyptranthes fasciculata* var. *hahni* (Krug & Urb.) Stehlé & Quentin, *Fl. Guadeloupe [Stehlé]* 2 (3): 62 (Stehlé & Quentin 1949).

VERNACULAR NAMES. — Ka: talipi • Wp: yawa poi li.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *P. Grenand 1013*, height 10 m.

[1288] *Myrcia ferruginea* (Poir.) DC.

Prodr. [A. P. de Candolle] 3: 245 [mid Mar. 1828] (Candolle 1828). — *Eugenia ferruginea* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 124 [24 Sep. 1813] (Poiret 1813). — *Myrtus ferruginea* (Poir.) Spreng., *Syst. Veg. [Sprengel]* 2: 487 [Jan.-May 1825] (Sprengel 1825). — *Krugia ferruginea* (Poir.) Urb., *Bot. Jahrb. Syst.* 19 (5): 604 [26 Feb. 1895] (Urban 1895). — *Marlierea ferruginea* (Poir.) McVaugh, *Mem. New York Bot. Gard.* 10 (1): 83 (McVaugh 1958).

Marlierea acuminata O.Berg, *Linnaea* 27 (1): 13 ["1854" publ. Nov. 1855] (Berg 1855).

Marlierea elliptica Griseb., *Fl. Brit. W.I. [Grisebach]* 233 [late 1860] (Grisebach 1860). — *Krugia elliptica* (Griseb.) Urb., *Ber. Deutsch. Bot. Ges.* 11: 376 (Urban 1893).

Eugeniopsis richardiana O.Berg, *Linnaea* 30: 665 [Mar. 1861] (Berg 1861). — *Marlierea richardiana* (O.Berg) Nied., *Nat. Pflanzenfam.*

[*Engler & Prantl*] 3 (7): 76 [Jan. 1893] (Niedenzu 1893), *nom. illeg. hom.*, non O.Berg (1861).

Marlierea guyanensis D.Legrand, *Comun. Bot. Mus. Hist. Nat. Montevideo* 3 (40): 36 (Legrand 1962).

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *Unknown coll. s.n.* ("Herb. Poir.") (original material FI[FI011506], P[P00547993, P01817729]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 13.9$ cm.

[1289] *Myrcia forsteri* (O.Berg) Molino & G.P.Burton

Ann. Missouri Bot. Gard. 107: 108 (Burton et al. 2022). — *Calyptranthes forsteri* O.Berg, *Linnaea* 27 (1): 23 ["1854" publ. Nov. 1855] (Berg 1855). — *Chytraculia forsteri* (O.Berg) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891). — *Myrcia neoforsteri* A.R.Loureiro & E.Lucas, *Phytotaxa* 373 (1): 75 [23 Oct. 2018] (Loureiro & Lucas 2018), *nom. illeg. superfl.* (based on *Calyptranthes forsteri*).

NOTES. — Lourenço et al. (2018) transferred the majority of continental American species of *Calyptranthes* to *Myrcia* to reflect the paraphyly of the former genus and modern infra-generic arrangement of the latter (Lucas et al. 2018). Among these new names, some turned out to be illegitimate as homonyms of older names in *Myrcia* (e.g. Lima et al. 2020) or, as here, because the original preference for a name significantly different from existing *Myrcia fosteri* Croat (Croat 1974) would not be supported by the approach to homonyms of the International Code for Botanical Nomenclature (Turland et al. 2018). The appropriate combination is therefore made here.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2825*, 10 m × 40 cm.

[1290] *Myrcia gigas* McVaugh

Mem. New York Bot. Gard. 18 (2): 88 (McVaugh 1969).

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *S.A. Mori et al. 20960*.

INVENTORY DATA (FG). — 1 tree, dbh = 10.5 cm.

[1291] *Myrcia graciliflora* Sagot

Ann. Sci. Nat., Bot. sér. 6, 20: 185 (Sagot 1885).

Eugenia schaueriana Miq., *Linnaea* 22: 174 (Miquel 1849). — *Aulomyrcia schaueriana* (Miq.) Amshoff, *Recueil Trav. Bot. Néerl.* 39: 155 (Amshoff 1942). — *Myrciaria schaueriana* (Miq.) O.Berg, *Linnaea* 27 (2-3): 323 ["1854" publ. Jan. 1856] (Berg 1856).

VERNACULAR NAMES. — Ka: akami enulu, tutu • Cr: koumaté.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-1825*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.3 cm.

- [1292] *Myrcia guianensis* (Aubl.) DC.
- Prodr. [A. P. de Candolle] 3:* 245 [mid Mar. 1828] (Candolle 1828). — *Eugenia guianensis* Aubl., *Hist. Pl. Guiane* 1: 506 [Jun.-Dec. 1775] (Aublet 1775). — *Myrtus guianensis* (Aubl.) Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 45 [Oct. 1825] (Hamilton 1825). — *Aguava guianensis* (Aubl.) Raf., *Sylva Tellur.* 107 (Rafinesque 1838).
- Myrtus pyrifolia* J.St.-Hil., *Traité Arbr. Arbust.*, ed. 2, 1: 208 (Saint-Hilaire 1803).
- Myrcia daphnoides* DC., *Prodr. [A. P. de Candolle] 3:* 246 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia daphnoides* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 132 [15 May 1857] (Berg 1857).
- Myrcia myoporina* DC., *Prodr. [A. P. de Candolle] 3:* 246 [mid Mar. 1828] (Candolle 1828).
- Myrcia exsucca* DC., *Prodr. [A. P. de Candolle] 3:* 247 [mid Mar. 1828] (Candolle 1828). — *Myrtus exsucca* Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 247 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Aulomyrcia exsucca* (DC.) O.Berg, *Linnaea* 27 (1): 79 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia queimadensis* Mattos, *Loefgrenia* 122: 4 [June 2006] (Mattos 2006), nom. illeg. superfl. (based on *Myrcia exsucca*).
- Myrtus duartii* Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 247 [mid Mar. 1828] (Candolle 1828), "Duarti", nom. nud. pro syn.
- Myrcia schrankiana* DC., *Prodr. [A. P. de Candolle] 3:* 247 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia schrankiana* (DC.) O.Berg, *Linnaea* 27 (1): 79 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia cassinioides* DC., *Prodr. [A. P. de Candolle] 3:* 249 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia cassinioides* (DC.) O.Berg, *Linnaea* 27 (1): 74 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia elaeodendra* DC., *Prodr. [A. P. de Candolle] 3:* 250 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia elaeodendra* (DC.) O.Berg, *Linnaea* 27 (1): 75 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia torta* DC., *Prodr. [A. P. de Candolle] 3:* 250 [mid Mar. 1828] (Candolle 1828). — *Myrtus torta* Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 250 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.
- Myrcia elegans* DC., *Prodr. [A. P. de Candolle] 3:* 251 [mid Mar. 1828] (Candolle 1828). — *Myrtus elegans* Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 251 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Calyptromyrcia elegans* (DC.) O.Berg, *Linnaea* 27 (1): 34 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcianthes elegans* (DC.) Mattos, *Loefgrenia* 128: 2 [Apr. 2009] (Mattos 2009).
- Myrcia leucadendron* DC., *Prodr. [A. P. de Candolle] 3:* 251 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia leucadendron* (DC.) O.Berg, *Linnaea* 27 (1): 64 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia spixiana* DC., *Prodr. [A. P. de Candolle] 3:* 251 [mid Mar. 1828] (Candolle 1828). — *Calyptromyrcia spixiana* (DC.) O.Berg, *Linnaea* 27 (1): 35 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcianthes spixiana* (DC.) Mattos, *Loefgrenia* 128: 2 [Apr. 2009] (Mattos 2009).
- Myrcia lauriflora* DC., *Prodr. [A. P. de Candolle] 3:* 252 [mid Mar. 1828] (Candolle 1828). — *Myrtus lauriflora* Mart. ex DC., *Prodr. [A. P. de Candolle] 3:* 252 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn. — *Aulomyrcia lauriflora* (DC.) O.Berg, *Linnaea* 27 (1): 64 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia alternifolia* Miq., *Linnaea* 22: 534 (Miquel 1849). — *Aulomyrcia alternifolia* (Miq.) O.Berg, *Linnaea* 27 (1): 72 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia obtecta* var. *alternifolia* (Miq.) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 283 (Legrand 1969).
- Myrcia pallens* DC., *Prodr. [A. P. de Candolle] 3:* 252 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia pallens* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 122 [15 May 1857] (Berg 1857).
- Myrcia crassicaulis* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (18): 311 ["1829" publ. 25 Feb. 1832] (Cambessèdes 1832). — *Aulomyrcia crassicaulis* (Cambess.) O.Berg, *Linnaea* 27 (1): 65 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia suaveolens* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (18): 315 ["1829" publ. 25 Feb. 1832] (Cambessèdes 1832). — *Aulomyrcia suaveolens* (Cambess.) O.Berg, *Linnaea* 27 (1): 78 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia microcarpa* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 324 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832). — *Aulomyrcia microcarpa* (Cambess.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 81 [15 May 1857] (Berg 1857).
- Myrcia hiemalis* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 332 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832).
- Myrcia obtusa* Schauer, *Linnaea* 21: 273 (Schauer 1848). — *Aulomyrcia obtusa* (Schauer) O.Berg, *Linnaea* 27 (1): 66 ["1854" publ. Nov. 1855] (Berg 1855).
- Myrcia surinamensis* Miq., *Linnaea* 22: 170 (Miquel 1848). — *Aulomyrcia surinamensis* (Miq.) O.Berg, *Linnaea* 27 (1): 64 ["1854" publ. Nov. 1855] (Berg 1855). — *Aulomyrcia obtusa* var. *surinamensis* (Miq.) Amshoff, *Recueil Trav. Bot. Néerl.* 39: 154 (Amshoff 1942).
- Aulomyrcia dichroma* O.Berg, *Linnaea* 27 (1): 65 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia obtusa* var. *grandifolia* O.Berg, *Linnaea* 27 (1): 66 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia obtusa* var. *longipes* O.Berg, *Linnaea* 27 (1): 67 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia obtusa* var. *panicularis* O.Berg, *Linnaea* 27 (1): 67 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia obtusa* var. *pauciflora* O.Berg, *Linnaea* 27 (1): 67 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia obtusa* var. *tenuifolia* O.Berg, *Linnaea* 27 (1): 67 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia roraimensis* O.Berg, *Linnaea* 27 (1): 68 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia roraimensis* (O.Berg) D.Legrand, *Atas Simp. Biota Amazônica* 4: 151 (Legrand 1967).
- Aulomyrcia cuneata* O.Berg, *Linnaea* 27 (1): 72 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia cuneata* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 76 [Jan. 1893] (Niedenzu 1893). — *Myrcia guianensis* var. *cuneata* (O.Berg) McVaugh, *Mem. New York Bot. Gard.* 18 (2): 93 (McVaugh 1969).
- Aulomyrcia schomburgkiana* O.Berg, *Linnaea* 27 (1): 75 ["1854" publ. Nov. 1855] (Berg 1855). — *Aulomyrcia obtusa* var. *schomburgkiana* (O.Berg) Amshoff, *Fl. Suriname* 3 (2): 79 (Amshoff 1951).
- Aulomyrcia conduplicata* O.Berg, *Linnaea* 27 (1): 76 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia pruinosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 114 [15 May 1857] (Berg 1857).

- Aulomyrcia bimarginata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 115 [15 May 1857] (Berg 1857).
- Aulomyrcia botrys* O.Berg, *Fl. Bras. [Martius]* 14 (1): 116 [15 May 1857] (Berg 1857). — *Myrcia botrys* (O.Berg) N.Silveira, *Loefgrenia* 91: 1 (Silveira 1987).
- Aulomyrcia nigropunctata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 116 [15 May 1857] (Berg 1857), "nigro-punctata". — *Myrcia nigropunctata* (O.Berg) N.Silveira, *Loefgrenia* 88: 1 (Silveira 1985).
- Aulomyrcia fragilis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 117 [15 May 1857] (Berg 1857).
- Aulomyrcia obtecta* O.Berg, *Fl. Bras. [Martius]* 14 (1): 117 [15 May 1857] (Berg 1857). — *Myrcia obtecta* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 89 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia clauseniana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 118 [15 May 1857] (Berg 1857).
- Aulomyrcia mansoni* O.Berg, *Fl. Bras. [Martius]* 14 (1): 121 [15 May 1857] (Berg 1857). — *Myrcia mansoni* (O.Berg) N.Silveira, *Loefgrenia* 88: 1 (Silveira 1985).
- Aulomyrcia pallens* var. *ovalis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 122 [15 May 1857] (Berg 1857). — *Myrcia pallens* var. *ovalis* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia pallens* var. *ovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 123 [15 May 1857] (Berg 1857).
- Aulomyrcia poeppigiana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 123 [15 May 1857] (Berg 1857). — *Myrcia poeppigiana* (O.Berg) Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 65 [9 Apr. 1895] (Hieronymus 1895), nom. illeg. hom., non O.Berg (1857). — *Myrcia poeppigiana* (O.Berg) Mattos, *Loefgrenia* 126: 2 [Sep. 2008] (Mattos 2008), isynom. et nom. illeg. hom., non O.Berg (1857).
- Aulomyrcia rotundifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 123 [15 May 1857] (Berg 1857). — *Myrcia rotundifolia* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 89 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia cassinioides* var. *glabrata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 129 [15 May 1857] (Berg 1857).
- Aulomyrcia cassinioides* var. *velutina* O.Berg, *Fl. Bras. [Martius]* 14 (1): 129 [15 May 1857] (Berg 1857).
- Aulomyrcia gardneriana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 129 [15 May 1857] (Berg 1857).
- Aulomyrcia gardneriana* var. *caeruleascens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 130 [15 May 1857] (Berg 1857).
- Aulomyrcia gardneriana* var. *virescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 130 [15 May 1857] (Berg 1857).
- Aulomyrcia lingua* O.Berg, *Fl. Bras. [Martius]* 14 (1): 130 [15 May 1857] (Berg 1857). — *Myrcia lingua* (O.Berg) Mattos, *Dusenia* 8: 161 (Mattos 1968).
- Aulomyrcia lingua* var. *glabrata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 130 [15 May 1857] (Berg 1857).
- Aulomyrcia lingua* var. *rufa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 130 [15 May 1857] (Berg 1857). — *Myrcia lingua* var. *rufa* (O.Berg) Mattos, *Loefgrenia* 70: 5 (Mattos 1976).
- Aulomyrcia hepatica* O.Berg, *Fl. Bras. [Martius]* 14 (1): 132 [15 May 1857] (Berg 1857). — *Myrcia hepatica* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 86 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia obscura* O.Berg, *Fl. Bras. [Martius]* 14 (1): 132 [15 May 1857] (Berg 1857). — *Myrcia obscura* (O.Berg) N.Silveira, *Roessleria* 7 (1): 66 (Silveira 1985).
- Aulomyrcia emarginata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 134 [15 May 1857] (Berg 1857). — *Myrcia emarginata* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 76 [Jan. 1893] (Niedenzu 1893).
- Aulomyrcia angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 135 [15 May 1857] (Berg 1857). — *Myrcia angustifolia* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 76 [Jan. 1893] (Niedenzu 1893).
- Aulomyrcia decrescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 135 [15 May 1857] (Berg 1857). — *Myrcia decrescens* (O.Berg) Mattos, *Loefgrenia* 126: 4 [Sep. 2008] (Mattos 2008).
- Aulomyrcia suffruticosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 136 [15 May 1857] (Berg 1857).
- Aulomyrcia glandulosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 139 [15 May 1857] (Berg 1857). — *Myrcia glandulosa* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia glandulosa* var. *elliptica* O.Berg, *Fl. Bras. [Martius]* 14 (1): 139 [15 May 1857] (Berg 1857).
- Aulomyrcia glandulosa* var. *longifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 139 [15 May 1857] (Berg 1857).
- Aulomyrcia glandulosa* var. *obovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 139 [15 May 1857] (Berg 1857). — *Myrcia glandulosa* var. *obovata* (O.Berg) N.Silveira, *Roessleria* 7 (1): 67 (Silveira 1985).
- Aulomyrcia pusilla* O.Berg, *Fl. Bras. [Martius]* 14 (1): 140 [15 May 1857] (Berg 1857). — *Myrcia pusilla* (O.Berg) Mattos, *Loefgrenia* 126: 2 [Sep. 2008] (Mattos 2008).
- Aulomyrcia vacciniifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 140 [15 May 1857] (Berg 1857). — *Myrcia vacciniifolia* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 76 [Jan. 1893] (Niedenzu 1893).
- Aulomyrcia bifornis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 141 [15 May 1857] (Berg 1857).
- Aulomyrcia plumbea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 142 [15 May 1857] (Berg 1857). — *Myrcia plumbea* (O.Berg) Mattos, *Loefgrenia* 126: 4 [Sep. 2008] (Mattos 2008).
- Myrcia terebinthacea* Poepp. ex O.Berg, *Fl. Bras. [Martius]* 14 (1): 155 [15 May 1857] (Berg 1857), nom. nud. pro syn.
- Aulomyrcia uaupensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 518 [1 Feb. 1858] (Berg 1858).
- Aulomyrcia maritima* O.Berg, *Fl. Bras. [Martius]* 14 (1): 553 [15 Jan. 1859] (Berg 1859).
- Aulomyrcia pallens* var. *subcordata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 553 [15 Jan. 1859] (Berg 1859). — *Myrcia pallens* var. *subcordata* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Aulomyrcia bracteata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 554 [15 Jan. 1859] (Berg 1859).
- Aulomyrcia desertorum* O.Berg, *Fl. Bras. [Martius]* 14 (1): 556 [15 Jan. 1859] (Berg 1859). — *Myrcia desertorum* (O.Berg) N.Silveira, *Loefgrenia* 88: 1 (Silveira 1985).

- Aulomyrcia obscura* var. *longipes* O.Berg, *Fl. Bras. [Martius]* 14 (1): 556 [15 Jan. 1859] (Berg 1859). — *Myrcia obscura* var. *longipes* (O.Berg) N.Silveira, *Roessleria* 7 (1): 66 (Silveira 1985).
- Aulomyrcia regeliana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 557 [15 Jan. 1859] (Berg 1859). — *Aulomyrcia regeliana* var. *oppositifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 557 [15 Jan. 1859] (Berg 1859), *nom. illeg.* (includes species type).
- Aulomyrcia regeliana* var. *sparsifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 557 [15 Jan. 1859] (Berg 1859).
- Aulomyrcia jequitinhonensis* O.Berg var. *glaucoides* O.Berg, *Fl. Bras. [Martius]* 14 (1): 560 [15 Jan. 1859] (Berg 1859). — *Myrcia torta* f. *glaucoides* (O.Berg) Kieresk., *Enum. Myrt. Bras.* 79 [Oct.-Nov. 1893] (Kiereskov 1893).
- Aulomyrcia velhensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 560 [15 Jan. 1859] (Berg 1859). — *Myrcia velhensis* (O.Berg) N.Silveira, *Loefgrenia* 91: 1 (Silveira 1987).
- Aulomyrcia androsaemoides* O.Berg, *Linnaea* 30: 661 [Mar. 1861] (Berg 1861). — *Myrcia androsaemoides* (O.Berg) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 579 [26 Feb. 1895] (Krug & Urban 1895).
- Aulomyrcia buxizans* O.Berg, *Linnaea* 30: 664 [Mar. 1861] (Berg 1861). — *Eugenia buxizans* Rich. ex O.Berg, *Linnaea* 30: 664 [Mar. 1861] (Berg 1861), *nom. nud. pro syn.*
- Myrcia scrobiculata* O.Berg, *Linnaea* 30: 668 [Mar. 1861] (Berg 1861).
- Myrcia roraimae* Oliv., *Timehri* 5: 192 (Oliver 1886). — *Aulomyrcia roraimae* (Oliv.) Steyermark, *Fieldiana, Bot.* 28 (4): 1007 (Steyermark 1957).
- Myrcia adpresepilosa* Kieresk., *Enum. Myrt. Bras.* 75 [Oct.-Nov. 1893] (Kiereskov 1893), “*adpresso-pilosa*”.
- Myrcia didrichseniana* Kieresk., *Enum. Myrt. Bras.* 82 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia cymosopaniculata* Kieresk., *Enum. Myrt. Bras.* 90 [Oct.-Nov. 1893] (Kiereskov 1893), “*cymoso-paniculata*”.
- Myrcia diaphanosticta* Kieresk., *Enum. Myrt. Bras.* 91 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia fastigiata* Kieresk., *Enum. Myrt. Bras.* 92 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia fastigiata* var. *coriacea* Kieresk., *Enum. Myrt. Bras.* 93 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia paracatuensis* Kieresk., *Enum. Myrt. Bras.* 99 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia rhabdoides* Kieresk., *Enum. Myrt. Bras.* 99 [Oct.-Nov. 1893] (Kiereskov 1893).
- Myrcia yungasensis* Rusby, *Mem. Torrey Bot. Club* 3: 27 (Rusby 1893).
- Myrcia androsaemoides* var. *parvifolia* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 579 [26 Feb. 1895] (Krug & Urban 1895).
- Myrcia lehmannii* Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 65 [9 Apr. 1895] (Hieronymus 1895).
- Myrcia collina* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 356 [“1894-96” publ. Dec. 1895] (Moore 1895).
- Myrcia daphnoides* var. *nervosa* Kieresk. ex Glaz., *Bull. Soc. Bot. France* 54 (Mém. 3c): 218 [“1907” publ. 22 Feb. 1908] (Glaziou 1908).
- Myrcia arimensis* Britton, *Bull. Torrey Bot. Club* 48 (12): 334 [“1921” publ. 1922] (Britton 1922).
- Myrcia incisa* D.Legrand, *Sellowia* 13: 290 (Legrand 1961).
- Myrcia renatoana* Mattos, *Arq. Bot. Estado São Paulo* 4 (2): 62 (Mattos 1966).
- Myrcianthes terminalis* Mattos & D.Legrand, *Loefgrenia* 67: 12 (Mattos & Legrand 1975).
- Myrcia stemmeriana* D.Legrand, *Fl. Ilustr. Catar.* 1 (Mirt., Supl. 1): 4 (Legrand 1977).
- Myrcia camapuana* Mattos, *Loefgrenia* 123: 2 [Aug. 2006] (Mattos 2006).
- NOTE. — Govaerts et al. (2008) erroneously stated that *Aulomyrcia scrobiculata* O.Berg is the basionym of *Myrcia scrobiculata* O.Berg. Actually, there is no reference to the former in the protologue of the latter; they have distinct types which are clearly not conspecific.
- VERNACULAR NAMES. — Ka: palaka pomiidy • Wp: pila lea • Nt: lebi tongo • Cr: kouamaté • Fr: kouamaté • Br: cambuí, guamirim-branco.
- HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material LINN, **LINN-HS 883.22**).
- INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 14.6 cm.
- [1293] *Myrcia inaequiloba* (DC.) Lemée
- Fl. Guyane Franç.* 3: 150 (Lemée 1954). — *Eugenia inaequiloba* DC., *Prod. [A. P. de Candolle]* 3: 282 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia inaequiloba* (DC.) Amshoff, *Recueil Trav. Bot. Néerl.* 42: 7 (Amshoff 1950). — *Myrcia inaequiloba* (DC.) D.Legrand, *Atlas Simp. Biota Amazônica* 4: 149 (Legrand 1967), isonym. — *Myrcia inaequiloba* (DC.) McVaugh, *Taxon* 17: 378 (McVaugh 1968), isonym.
- Eugenia nitida* Benth., *J. Bot. [Hooker]* 2: 322 (Bentham 1840), *nom. illeg. hom.*, *non* Vell. (Vellozo 1829) *nec* Cambess. (Cambessèdes 1832).
- Eugenia polyantha* Miq., *Linnaea* 18: 741 [“1844” publ. Aug.-Oct. 1845] (Miquel 1845). — *Myrciaria polyantha* (Miq.) O.Berg, *Linnaea* 27 (2-3): 322 [“1854” publ. Jan. 1856] (Berg 1856).
- Aulomyrcia pirarensis* O.Berg, *Linnaea* 27 (1): 41 [“1854” publ. Nov. 1855] (Berg 1855).
- Aulomyrcia paniculata* O.Berg, *Linnaea* 27 (1): 49 [“1854” publ. Nov. 1855] (Berg 1855). — *Aulomyrcia inaequiloba* var. *paniculata* (O.Berg) Amshoff, *Recueil Trav. Bot. Néerl.* 42: 7 (Amshoff 1950).
- Myrciaria nitida* O.Berg, *Linnaea* 27 (2-3): 324 [“1854” publ. Jan. 1856] (Berg 1856).
- Myrciaria nitida* var. *chartacea* O.Berg, *Linnaea* 27 (2-3): 324 [“1854” publ. Jan. 1856] (Berg 1856).
- Myrciaria nitida* var. *coriacea* O.Berg, *Linnaea* 27 (2-3): 325 [“1854” publ. Jan. 1856] (Berg 1856).
- Myrciaria nitida* var. *dives* O.Berg, *Linnaea* 27 (2-3): 325 [“1854” publ. Jan. 1856] (Berg 1856).
- Aulomyrcia lancifolia* O.Berg, *Linnaea* 30: 658 [Mar. 1861] (Berg 1861).

Myrcia laevis O.Berg, *Linnaea* 31: 252, (Berg 1861-1862), *nom. illeg. hom., non* G.Don (1832).

Aulomyrcia inaequiloba var. *nitida* Amshoff, *Recueil Trav. Bot. Néerl.* 42: 8 (Amshoff 1950).

Myrcia lucida McVaugh, *Mem. New York Bot. Gard.* 18 (2): 100 (McVaugh 1969).

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (original material G[G00222421]).

SIZE. — Guyana. *J.J. Pipoly et al.* 9345 (MO), 4 m × 20 cm.

INVENTORY DATA (FG). — 4 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12$ cm.

[1294] *Myrcia minutiflora* Sagot

Ann. Sci. Nat., Bot. sér. 6, 20: 185 (Sagot 1885). — *Aulomyrcia minutiflora* (Sagot) Amshoff, *Bull. Torrey Bot. Club* 75 (5): 532 [11 Oct. 1948] (Amshoff 1948).

VERNACULAR NAMES. — Wp: wila wasey.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *E.M. Mélinon* 442 (original material P[P00163092]).

SIZE. — Up to 10 m tall (Holst 2003).

[1295] *Myrcia multiflora* (Lam.) DC.

Prod. [A. P. de Candolle] 3: 244 [mid Mar. 1828] (Candolle 1828). — *Eugenia multiflora* Lam., *Encycl. J. Lamarck et al.* 3 (1): 202 [19 Oct. 1789] (Lamarck 1789). — *Myrtus multiflora* (Lam.) Spreng., *Syst. Veg. [Sprengel]* 2: 486 [Jan.-May 1825] (Sprengel 1825). — *Cumetea multiflora* (Lam.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838). — *Aulomyrcia multiflora* (Lam.) O.Berg, *Linnaea* 27 (1): 47 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia camaraeana DC., *Prod. [A. P. de Candolle]* 3: 251 [mid Mar. 1828] (Candolle 1828). — *Myrtus camaraeana* Mart. ex DC., *Prod. [A. P. de Candolle]* 3: 251 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.* — *Aulomyrcia camaraeana* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 83 [15 May 1857] (Berg 1857).

Myrcia sphaerocarpa DC., *Prod. [A. P. de Candolle]* 3: 251 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia sphaerocarpa* (DC.) O.Berg, *Linnaea* 27 (1): 51 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia ellipticifolia Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (18): 312 ["1829" publ. 25 Feb. 1832] (Cambessèdes 1832).

Aulomyrcia buxifolia O.Berg, *Fl. Bras. [Martius]* 14 (1): 80 [15 May 1857] (Berg 1857).

Aulomyrcia buxifolia var. *elliptica* O.Berg, *Fl. Bras. [Martius]* 14 (1): 80 [15 May 1857] (Berg 1857), *nom. illeg.* (includes species type).

Aulomyrcia buxifolia var. *ovalis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 80 [15 May 1857] (Berg 1857).

Aulomyrcia caeruleascens O.Berg, *Fl. Bras. [Martius]* 14 (1): 80 [15 May 1857] (Berg 1857). — *Myrcia caeruleascens* (O.Berg) Kiersk., *Enum. Myrt. Bras.* 79 [Oct.-Nov. 1893] (Kiersk 1893).

Aulomyrcia caesia O.Berg, *Fl. Bras. [Martius]* 14 (1): 83 [15 May 1857] (Berg 1857).

Aulomyrcia glaucescens O.Berg, *Fl. Bras. [Martius]* 14 (1): 81 [15 May 1857] (Berg 1857). — *Myrcia glaucescens* (O.Berg) Kiersk., *Enum. Myrt. Bras.* 83 [Oct.-Nov. 1893] (Kiersk 1893). — *Myrcia multiflora* var. *glaucescens* (O.Berg) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 305 (Legrand 1969).

Aulomyrcia glaucescens var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 81 [15 May 1857] (Berg 1857).

Aulomyrcia ovalifolia O.Berg, *Fl. Bras. [Martius]* 14 (1): 81 [15 May 1857] (Berg 1857). — *Myrcia ovalifolia* (O.Berg) Kiersk., *Enum. Myrt. Bras.*: 84 [Oct.-Nov. 1893] (Kiersk 1893). — *Myrcia multiflora* f. *ovalifolia* (O.Berg) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 305 (Legrand 1969).

Aulomyrcia glaucescens var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 82 [15 May 1857] (Berg 1857).

Aulomyrcia perforata O.Berg, *Fl. Bras. [Martius]* 14 (1): 83 [15 May 1857] (Berg 1857).

Aulomyrcia goyazensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 85 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *arborescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 85 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *gracilis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 85 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *intermedia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 85 [15 May 1857] (Berg 1857).

Aulomyrcia pallida O.Berg, *Fl. Bras. [Martius]* 14 (1): 86 [15 May 1857] (Berg 1857). — *Myrcia pallida* (O.Berg) N.Silveira, *Loefgrenia* 88: 2 (Silveira 1985).

Aulomyrcia sphaerocarpa var. *complicata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 86 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *obtusata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 86 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *ovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 86 [15 May 1857] (Berg 1857).

Aulomyrcia sphaerocarpa var. *pauciflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 86 [15 May 1857] (Berg 1857).

Aulomyrcia laruotteana (Cambess.) O.Berg var. *peruviana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 91 [15 May 1857] (Berg 1857).

Aulomyrcia multiflora var. *grandifolia* O.Berg, *Linnaea* 30: 660 [Mar. 1861] (Berg 1861).

Myrcia taubatensis Kiersk., *Enum. Myrt. Bras.* 79 [Oct.-Nov. 1893] (Kiersk 1893). — *Myrcia taubatensis* var. *ovalis* (O.Berg) Kiersk., *Enum. Myrt. Bras.* 79 [Oct.-Nov. 1893] (Kiersk 1893).

Myrcia glaberrima Barb.Rodr. ex Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 7: 803 (Chodat & Hassler 1907), *nom. nud.*

Aulomyrcia vinacea Steyermark., *Fieldiana, Bot.* 28 (4): 1008 (Steyermark 1957).

Myrcia multiflora var. *ramulosa* D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 309 (Legrand 1969).

Myrcia heringeriana Mattos, *Loefgrenia* 126: 2 [Sep. 2008] (Mattos 2008).

VERNACULAR NAMES. — Pa: awao, awau • Ka: asaikiabe, sadyabe, seityape • Br: cambúi.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *M.-F. Prévost* 709.

INVENTORY DATA (FG). — 13 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20\text{ cm}$.

[1296] *Myrcia multiglomerata*
(Amshoff) E.Lucas & C.E.Wilson

Ann. Missouri Bot. Gard. 101 (4): 676 (Lucas & Wilson 2016). — *Marlierea multiglomerata* Amshoff, *Recueil Trav. Bot. Néerl.* 42: 3 (Amshoff 1950).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: piliwa epulu, piliwa pi • Wp: kumati sili, pila lea • Wn: wahawe • Br: pitangueira-de-cachorro.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *P. Grenand* 356, height 20 m.

[1297] *Myrcia neolucida* A.R.Lourenço & E.Lucas

Phytotaxa 373 (1): 79 [23 Oct. 2018] (Lourenço & Lucas 2018). — *Calyptranthes lucida* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 258 [mid Mar. 1828] (Candolle 1828). — *Chytraculia lucida* (Mart. ex DC.) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891).

Calyptranthes pobliana O.Berg, *Fl. Bras. [Martius]* 14 (1): 42 [15 May 1857] (Berg 1857). — *Chytraculia pobliana* (O.Berg) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Wp: atimi letimá'i, pila lea sili.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-1913.

SIZE. — Up to 30 cm dbh (Amshoff 1951).

[1298] *Myrcia neomontana* E.Lucas & C.E.Wilson

Ann. Missouri Bot. Gard. 101 (4): 679 (Lucas & Wilson 2016). — *Eugenia montana* Aubl., *Hist. Pl. Guiane* 1: 495 [Jun.-Dec. 1775] (Aublet 1775). — *Cumetea montana* (Aubl.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838). — *Marlierea montana* (Aubl.) Amshoff, *Recueil Trav. Bot. Néerl.* 39: 147 (Amshoff 1942). — *Myrcianthes montana* (Aubl.) C.Nelson, *Anales Jard. Bot. Madrid* 57 (2): 406 ["1999" publ. Jan. 2000] (Nelson 2000).

Marlierea suffruticosa O.Berg, *Linnaea* 27 (1): 16 ["1854" publ. Nov. 1855] (Berg 1855).

Marlierea richardiana O.Berg, *Linnaea* 30: 650 [Mar. 1861] (Berg 1861).

NOTE. — The name "*Calyptranthes obtusa* Miq." listed in Govaerts *et al.* (2008) as an illegitimate synonym of *Myrcia montana*, with the reference "*Linnaea* 22: 172 (1849)", has actually no standing: Miquel explicitly refers to *C. obtusa* Benth., and cites *Hostmann* 1176 as occurring in Surinam, not as the type of a new species.

VERNACULAR NAMES. — Pa: arak, arak-kamwi, pakusin-amana • Ka: alamilu.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[[BM000754877](#)]).

INVENTORY DATA (FG). — 35 trees in 11 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 36.9\text{ cm}$.

[1299] *Myrcia neospeciosa* A.R.Lourenço & E.Lucas

Phytotaxa 373 (1): 79 [23 Oct. 2018] (Lourenço & Lucas 2018). — *Calyptranthes speciosa* Sagot, *Ann. Sci. Nat., Bot. sér.* 6, 20: 187 (Sagot 1885).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *E.M. Mélinon* 488, 1862 [erroneously filed at F as "E.M. Mélinon 18"; "18" actually represents the first two digits of the year of collection, often pre-printed on P labels in the 19th century] (holo-, P[[P00723171](#)]; iso-, BM[[BM001011735](#)], F[[V0092885F](#)], K[[K000331318](#)], P[[P00723172](#)]).

SIZE. — Colombia, Caquetá. *S. Bergeron & S. Roman* 494-556 (MO), 12 m × 13.05 cm.

INVENTORY DATA (FG). — 1 tree, $dbh = 11.9\text{ cm}$.

[1300] *Myrcia paivae* O.Berg

Fl. Bras. [Martius] 14 (1): 179 [15 May 1857] (Berg 1857).

Myrcia frontinensis Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 63 [9 Apr. 1895] (Hieronymus 1895).

Myrcia gamaeana Glaz., *Bull. Soc. Bot. France* 54 (Mém. 3c): 210 ["1907" publ. 22 Feb. 1908] (Glaziou 1908), *nom. inval.* (opus utique oppr.).

Myrcia paivae var. *gracilis* Lingelsh., *Repert. Spec. Nov. Regni Veg.* 7: 243 (Lingelsheim 1909).

NOTE. — *M. gamaeana* erroneously appears as "*Myrcia frontinensis* Hieron. var. *gamaeana* Glaz." in Govaerts *et al.* (2008).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *G. Cremers & P. Petronelli* 11776.

SIZE. — Bolivia, La Paz. *A.F. Fuentes et al.* 6109 (MO), 10 m × 14.6 cm.

[1301] *Myrcia platyclada* DC.
(Fig. 42D)

Prodr. [A. P. de Candolle] 3: 244 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia platyclada* (DC.) Amshoff, *Bull. Torrey Bot. Club* 75 (5): 531 [11 Oct. 1948] (Amshoff 1948), *nom. illeg. hom.*, non O.Berg (1849).

Aulomyrcia dumosa O.Berg, *Linnaea* 30: 656 [Mar. 1861] (Berg 1861). — *Myrcia dumosa* (O.Berg) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 580 [26 Feb. 1895] (Krug & Urban 1895).

Aulomyrcia edulis O.Berg, *Linnaea* 30: 657 [Mar. 1861] (Berg 1861). — *Myrcia edulis* (O.Berg) Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 581 [26 Feb. 1895] (Krug & Urban 1895).

Aulomyrcia platyclada var. *kaieteurensis* Amshoff, *Bull. Torrey Bot. Club* 75 (5): 532 [11 Oct. 1948] (Amshoff 1948).

VERNACULAR NAMES. — Pa: arak-seinō, inam-etni-priyo • Wn: katanom.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: L.C. Richard 63 (holotype of *Aulomyrcia edulis*: P[P00725809]; iso-, P[P00725811]).

INVENTORY DATA (FG). — 15 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23.1$ cm.

[1302] *Myrcia pulchella* (DC.) A.R.Lourenço & E.Lucas

Phytotaxa 373 (1): 80 [23 Oct. 2018] (Lourenço & Lucas 2018). — *Calyptanthes pulchella* DC., *Prodr. [A. P. de Candolle]* 3: 257 [mid Mar. 1828] (Candolle 1828). — *Chytraculia pulchella* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891).

Calyptanthes pulchella var. *cuneata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 43 [15 May 1857] (Berg 1857).

Calyptanthes pulchella var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 43 [15 May 1857] (Berg 1857).

Calyptanthes pulchella var. *grandiflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 44 [15 May 1857] (Berg 1857).

Calyptanthes pulchella var. *parviflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 516 [1 Feb. 1858] (Berg 1858).

VERNACULAR NAMES. — Wp: atimi letimā'i • Br: guaiabinha.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: P. Grenand 2868.

SIZE. — Colombia, Caquetá. M. Sánchez S. et al. 1653 (MO), 12 m × 15 cm.

[1303] *Myrcia pullei* (Burret ex Amshoff)
A.R.Lourenço & E.Lucas

Phytotaxa 373 (1): 81 [23 Oct. 2018] (Lourenço & Lucas 2018). — *Calyptanthes pullei* Burret ex Amshoff, *Recueil Trav. Bot. Néerl.* 42: 4 (Amshoff 1950).

Calyptanthes pullei var. *immaculata* McVaugh, *Mem. New York Bot. Gard.* 10 (1): 78 (McVaugh 1958).

HERBARIUM DATA (FG). — A single collection, P. Grenand 1525.

SIZE. — Venezuela, Bolívar. E. Sanoja 2708 (US), 10 m.

[1304] *Myrcia pyrifolia* (Desv. ex Ham.) Nied.

Nat. Pflanzenfam. [Engler & Prantl] 3 (7): 76 [Jan. 1893] (Niedenzu 1893). — *Eugenia pyrifolia* Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 44 [Oct. 1825] (Hamilton 1825). — *Aulomyrcia pyrifolia* (Desv. ex Ham.) O.Berg, *Linnaea* 27 (1): 44 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia divergens DC., *Prodr. [A. P. de Candolle]* 3: 245 [mid Mar. 1828] (Candolle 1828).

Aulomyrcia pyrifolia var. *gracilis* O.Berg, *Linnaea* 27 (1): 44 ["1854" publ. Nov. 1855] (Berg 1855).

Aulomyrcia pyrifolia var. *robusta* O.Berg, *Linnaea* 27 (1): 45 ["1854" publ. Nov. 1855] (Berg 1855).

Aulomyrcia ovata O.Berg, *Linnaea* 27 (4): 467 ["1854" publ. Feb. 1856] (Berg 1856).

VERNACULAR NAMES. — Wn: pakumila.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: C. Sastre 6387.

SIZE. — Brazil, Pará. G. Martinelli et al. 12268 (MO), 12 m.

[1305] *Myrcia saxatilis* (Amshoff) McVaugh

Mem. New York Bot. Gard. 18 (2): 105 (McVaugh 1969). — *Aulomyrcia saxatilis* Amshoff, *Recueil Trav. Bot. Néerl.* 39: 154 (Amshoff 1942).

VERNACULAR NAMES. — Pa: arak-puvemna • Wp: kumesi sili.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: J.-F. Molino et al. 1981, 15 m × 10 cm.

[1306] *Myrcia servata* McVaugh

Mem. New York Bot. Gard. 18 (2): 139 (McVaugh 1969).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: R. Benoist 540 (P), 15 m tall.

[1307] *Myrcia splendens* (Sw.) DC.

Prodr. [A. P. de Candolle] 3: 244 [mid Mar. 1828] (Candolle 1828). — *Myrtus splendens* Sw., *Prodr. [Swartz]* 79 [20 Jun.-29 July 1788] (Swartz 1788).

Eugenia periplocifolia Jacq., *Collectanea* 2: 108 [Apr. 1789] (Jacquin 1789), "periplocaefolia".

Eugenia divaricata Lam., *Encycl. [J. Lamarck et al.]* 3 (1): 202 [19 Oct. 1789] (Lamarck 1789). — *Myrcia divaricata* (Lam.) DC., *Prodr. [A. P. de Candolle]* 3: 243 [mid Mar. 1828] (Candolle 1828). — *Cumetea divaricata* (Lam.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838).

Eugenia fallax Rich., *Actes Soc. Hist. Nat. Paris* 1: 110 [Oct. 1792] (Richard 1792). — *Myrcia fallax* (Rich.) DC., *Prodr. [A. P. de Candolle]* 3: 244 [mid Mar. 1828] (Candolle 1828).

Myrtus bracteolaris Poir., *Encycl. [J. Lamarck et al.]* 4 (2): 411 [1 Nov. 1798] (Poiret 1798). — *Myrtus stouyi* Spreng., *Syst. Vég. [Sprengel]* 2: 484 [Jan.-May 1825] (Sprengel 1825), "Stoupii", nom. illeg. superfl. (based on *Myrtus bracteolaris*). — *Myrcia bracteolaris* (Poir.) DC., *Prodr. [A. P. de Candolle]* 3: 245 [mid Mar. 1828] (Candolle 1828).

Eugenia laxiflora Poir., *Encycl. [J. Lamarck et al.] Suppl.* 3: 123 [24 Sep. 1813] (Poiret 1813).

Myrtus complicata Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 140 [6 Aug. 1823] (Kunth 1823). — *Myrcia complicata* (Kunth) DC., *Prodr. [A. P. de Candolle]* 3: 255 [mid Mar. 1828] (Candolle 1828).

Myrtus polyantha Kunth, *Nova genera et species plantarum [H.B.K.]* 6: 140 [6 Aug. 1823] (Kunth 1823). — *Myrcia polyantha* (Kunth)

- DC., *Prodr. [A. P. de Candolle] 3*: 255 [mid Mar. 1828] (Candolle 1828).
- Myrtus acuminata* Kunth, *Nova genera et species plantarum [H.B.K.] 6*: 141 [6 Aug. 1823] (Kunth 1823). — *Myrcia acuminata* (Kunth) DC., *Prodr. [A. P. de Candolle] 3*: 256 [mid Mar. 1828] (Candolle 1828).
- Myrcia sororia* DC., *Prodr. [A. P. de Candolle] 3*: 243 [mid Mar. 1828] (Candolle 1828).
- Myrcia hayneana* DC., *Prodr. [A. P. de Candolle] 3*: 246 [mid Mar. 1828] (Candolle 1828). — *Myrtus hayneana* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 246 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*
- Myrcia macrophylla* DC., *Prodr. [A. P. de Candolle] 3*: 248 [mid Mar. 1828] (Candolle 1828).
- Myrcia magnoliifolia* DC., *Prodr. [A. P. de Candolle] 3*: 248 [mid Mar. 1828] (Candolle 1828), “*magnoliaefolia*”.
- Myrcia sepiaria* DC., *Prodr. [A. P. de Candolle] 3*: 249 [mid Mar. 1828] (Candolle 1828). — *Myrtus sepiaria* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 249 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*
- Myrcia costata* DC., *Prodr. [A. P. de Candolle] 3*: 252 [mid Mar. 1828] (Candolle 1828). — *Myrtus costata* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 252 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*
- Myrcia pseudomini* DC., *Prodr. [A. P. de Candolle] 3*: 252 [mid Mar. 1828] (Candolle 1828), “*pseudo-Mini*”. — *Myrcia rostrata f. pseudomini* (DC.) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 243 (Legrand 1969).
- Myrcia berberis* DC., *Prodr. [A. P. de Candolle] 3*: 254 [mid Mar. 1828] (Candolle 1828). — *Myrtus berberis* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 254 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*
- Myrcia formosiana* DC., *Prodr. [A. P. de Candolle] 3*: 255 [mid Mar. 1828] (Candolle 1828).
- Myrcia rostrata* DC., *Prodr. [A. P. de Candolle] 3*: 255 [mid Mar. 1828] (Candolle 1828). — *Myrtus rostrata* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 255 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.*
- Eugenia mikaniiana* DC., *Prodr. [A. P. de Candolle] 3*: 283 [mid Mar. 1828] (Candolle 1828). — *Myrtus mikaniiana* Mart. ex DC., *Prodr. [A. P. de Candolle] 3*: 283 [mid Mar. 1828] (Candolle 1828), *nom. nud. pro syn.* — *Myrcia mikaniiana* (DC.) O.Berg, *Fl. Bras. [Martius] 14 (1)*: 182 [15 May 1857] (Berg 1857).
- Myrcia oocarpa* Cambess., *Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 2 (18)*: 298 [“1829” publ. 25 Feb. 1832] (Cambessèdes 1832).
- Myrcia rostrata* var. *brunea* Cambess., *Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 2 (19)*: 321 [“1829” publ. 27 Oct. 1832] (Cambessèdes 1832).
- Myrcia rufidula* Schltld., *Linnaea 13*: 416 [“1839” publ. Mar.-June 1840] (Schlechtendal 1840).
- Myrcia kunthiana* Steud., *Nomencl. Bot. [Steudel], ed. 2, 2*: 173 (Steudel 1841), *nom. illeg. superfl.* (based on *Myrtus polyantha*).
- Eugenia paniculiflora* Steud., *Flora 26 (45)*: 762 [17 Dec. 1843] (Steudel 1843), “*paniculaeflora*”.
- Myrcia rufula* Miq., *Linnaea 19*: 440 (Miquel 1846).
- Myrcia lindeniana* O.Berg, *Linnaea 27 (1)*: 86 [“1854” publ. Nov. 1855] (Berg 1855).
- Aulomyrcia wulschlaegeliana* O.Berg, *Linnaea 27 (1)*: 90 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia saxicola* O.Berg, *Linnaea 27 (1)*: 92 [“1854” publ. Nov. 1855] (Berg 1855). — *Myrtus saxicola* Poepp. ex O.Berg, *Linnaea 27 (1)*: 92 [“1854” publ. Nov. 1855] (Berg 1855), *nom. nud. pro syn.*
- Myrcia acuminata* var. *bullata* O.Berg, *Linnaea 27 (1)*: 94 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia acuminata* var. *genuina* O.Berg, *Linnaea 27 (1)*: 94 [“1854” publ. Nov. 1855] (Berg 1855), *nom. inval.* (Turland *et al.* 2018: Art. 24.3).
- Myrcia acuminata* var. *meridensis* O.Berg, *Linnaea 27 (1)*: 94 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia acuminata* var. *peruviana* O.Berg, *Linnaea 27 (1)*: 94 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia acuminata* var. *tovarensis* O.Berg, *Linnaea 27 (1)*: 95 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia venezuelensis* O.Berg, *Linnaea 27 (1)*: 96 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia cucullata* O.Berg, *Linnaea 27 (1)*: 97 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia chilensis* O.Berg, *Linnaea 27 (1)*: 99 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrtus ruizii* Poepp. ex O.Berg, *Linnaea 27 (1)*: 99 [“1854” publ. Nov. 1855] (Berg 1855), *nom. nud. pro syn.*
- Myrcia ayresiana* O.Berg, *Linnaea 27 (1)*: 100 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia reticulata* O.Berg, *Linnaea 27 (1)*: 101 [“1854” publ. Nov. 1855] (Berg 1855), *nom. illeg. hom., non Cambess. (Cambessèdes 1832)*.
- Myrcia coumetoides* O.Berg, *Linnaea 27 (1)*: 102 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia costaricensis* O.Berg, *Linnaea 27 (1)*: 104 [“1854” publ. Nov. 1855] (Berg 1855), “*Costa-Ricensis*”.
- Myrcia splendens* var. *genuina* O.Berg, *Linnaea 27 (1)*: 105 [“1854” publ. Nov. 1855] (Berg 1855), *nom. inval.* (Turland *et al.* 2018: Art. 24.3).
- Myrcia splendens* var. *micropora* O.Berg, *Linnaea 27 (1)*: 105 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia splendens* var. *obscura* O.Berg, *Linnaea 27 (1)*: 105 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia discolor* O.Berg, *Linnaea 27 (1)*: 111 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia overstediana* O.Berg, *Linnaea 27 (1)*: 112 [“1854” publ. Nov. 1855] (Berg 1855).
- Myrcia melanoclada* O.Berg, *Linnaea 27 (1)*: 113 [“1854” publ. Nov. 1855] (Berg 1855).

- Myrcia plicatocostata* O.Berg, *Linnaea* 27 (1): 114 ["1854" publ. Nov. 1855] (Berg 1855), "plicato-costata".
- Myrcia sericea* O.Berg, *Linnaea* 27 (1): 114 ["1854" publ. Nov. 1855] (Berg 1855), *nom. illeg. hom., non* G.Don (1832).
- Eugenia mollis* Willd. ex O.Berg, *Linnaea* 27 (1): 121 ["1854" publ. Nov. 1855] (Berg 1855), *nom. nud. pro syn.*
- Aulomyrcia costata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 79 [15 May 1857] (Berg 1857).
- Myrcia costata* var. *bahiensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 157 [15 May 1857] (Berg 1857).
- Myrcia costata* var. *minensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 157 [15 May 1857] (Berg 1857).
- Myrcia elongata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 159 [15 May 1857] (Berg 1857).
- Myrcia martiana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 159 [15 May 1857] (Berg 1857). — *Myrcia rufula* var. *martiana* (O.Berg) Kjaersk., *Enum. Myrt. Bras.* 55 [Oct.-Nov. 1893] (Kjaerskov 1893).
- Myrcia elongata* var. *brunnea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857).
- Myrcia elongata* var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857).
- Myrcia elongata* var. *ochracea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857).
- Myrcia guajavifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857), "guajavaefolia".
- Myrcia guajavifolia* var. *impunctata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857).
- Myrcia guajavifolia* var. *perforata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 160 [15 May 1857] (Berg 1857).
- Myrcia friburgensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 161 [15 May 1857] (Berg 1857).
- Myrcia guajavifolia* var. *bullata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 161 [15 May 1857] (Berg 1857).
- Myrcia magnoliifolia* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 162 [15 May 1857] (Berg 1857).
- Myrcia magnoliifolia* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 162 [15 May 1857] (Berg 1857).
- Myrcia magnoliifolia* var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 162 [15 May 1857] (Berg 1857).
- Myrcia brandamii* O.Berg, *Fl. Bras. [Martius]* 14 (1): 164 [15 May 1857] (Berg 1857), "Brandami".
- Myrcia alagoensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 165 [15 May 1857] (Berg 1857).
- Myrcia alagoensis* var. *intermedia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 165 [15 May 1857] (Berg 1857).
- Myrcia alagoensis* var. *oblongata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 165 [15 May 1857] (Berg 1857).
- Myrcia alagoensis* var. *ovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 165 [15 May 1857] (Berg 1857).
- Myrcia spruceana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 165 [15 May 1857] (Berg 1857).
- Myrcia phaeoclada* O.Berg, *Fl. Bras. [Martius]* 14 (1): 167 [15 May 1857] (Berg 1857).
- Myrcia phaeoclada* var. *alagoensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 167 [15 May 1857] (Berg 1857).
- Myrcia phaeoclada* var. *guyanensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 167 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* var. *longifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* var. *pendula* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia kegeliana* var. *vulgaris* O.Berg, *Fl. Bras. [Martius]* 14 (1): 168 [15 May 1857] (Berg 1857).
- Myrcia klotzschiana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 169 [15 May 1857] (Berg 1857).
- Myrcia micrantha* O.Berg, *Fl. Bras. [Martius]* 14 (1): 169 [15 May 1857] (Berg 1857).
- Myrcia berberis* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 170 [15 May 1857] (Berg 1857).
- Myrcia berberis* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 170 [15 May 1857] (Berg 1857).
- Myrcia latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 170 [15 May 1857] (Berg 1857).
- Myrcia erythroxylon* O.Berg, *Fl. Bras. [Martius]* 14 (1): 173 [15 May 1857] (Berg 1857).
- Myrcia erythroxylon* var. *caeruleascens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 173 [15 May 1857] (Berg 1857).
- Myrcia pellucida* O.Berg, *Fl. Bras. [Martius]* 14 (1): 173 [15 May 1857] (Berg 1857).
- Myrcia gracilis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 174 [15 May 1857] (Berg 1857). — *Myrcia rostrata* f. *gracilis* (O.Berg) D.Legrand, *Fl. Ilustr. Catar. I (Mirt.)*: 240 (Legrand 1969).
- Myrcia gracilis* var. *prasina* O.Berg, *Fl. Bras. [Martius]* 14 (1): 174 [15 May 1857] (Berg 1857).
- Myrcia acutata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 175 [15 May 1857] (Berg 1857).
- Myrcia gracilis* var. *opaca* O.Berg, *Fl. Bras. [Martius]* 14 (1): 175 [15 May 1857] (Berg 1857).
- Myrcia laevigata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 175 [15 May 1857] (Berg 1857).
- Myrcia laevigata* var. *brunnea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 175 [15 May 1857] (Berg 1857).

- Myrcia laevigata* var. *canescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 175 [15 May 1857] (Berg 1857).
- Myrcia catharinae* O.Berg, *Fl. Bras. [Martius]* 14 (1): 176 [15 May 1857] (Berg 1857).
- Myrcia corcovadensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 177 [15 May 1857] (Berg 1857).
- Myrcia opaca* O.Berg, *Fl. Bras. [Martius]* 14 (1): 177 [15 May 1857] (Berg 1857).
- Myrcia opaca* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 177 [15 May 1857] (Berg 1857).
- Myrcia opaca* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 177 [15 May 1857] (Berg 1857).
- Myrcia ciarensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 178 [15 May 1857] (Berg 1857).
- Myrcia nitens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 178 [15 May 1857] (Berg 1857).
- Myrcia sericiflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 178 [15 May 1857] (Berg 1857). — *Myrcia rostrata* f. *sericiflora* (O.Berg) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 243 (Legrand 1969).
- Myrcia riparia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 179 [15 May 1857] (Berg 1857).
- Myrcia schuechiana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 181 [15 May 1857], “Schüchiana”.
- Myrcia mikaniiana* var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 182 [15 May 1857] (Berg 1857).
- Myrcia mikaniiana* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 182 [15 May 1857] (Berg 1857).
- Myrcia velutina* O.Berg, *Fl. Bras. [Martius]* 14 (1): 182 [15 May 1857] (Berg 1857).
- Myrcia communis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 183 [15 May 1857] (Berg 1857). — *Myrcia rostrata* f. *communis* (O.Berg) D.Legrand, *Fl. Ilustr. Catar. 1 (Mirt.)*: 243 (Legrand 1969).
- Myrcia communis* var. *glabrata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 183 [15 May 1857] (Berg 1857).
- Myrcia velutina* var. *canescens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 183 [15 May 1857] (Berg 1857).
- Myrcia velutina* var. *ochracea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 183 [15 May 1857] (Berg 1857).
- Myrcia communis* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 183 [15 May 1857] (Berg 1857).
- Myrcia gardneriana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 184 [15 May 1857] (Berg 1857).
- Myrcia impressa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 184 [15 May 1857] (Berg 1857).
- Myrcia ypanemensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 186 [15 May 1857] (Berg 1857).
- Myrcia barrensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 187 [15 May 1857] (Berg 1857).
- Myrcia hayneana* var. *paraensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 187 [15 May 1857] (Berg 1857).
- Myrcia negrensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 187 [15 May 1857] (Berg 1857).
- Myrcia acutiloba* O.Berg, *Fl. Bras. [Martius]* 14 (1): 189 [15 May 1857] (Berg 1857).
- Myrcia sellowiana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 197 [15 May 1857] (Berg 1857).
- Myrcia sellowiana* var. *bullata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 197 [15 May 1857] (Berg 1857).
- Myrcia sellowiana* var. *costata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 197 [15 May 1857] (Berg 1857).
- Myrcia superba* O.Berg, *Fl. Bras. [Martius]* 14 (1): 198 [15 May 1857] (Berg 1857).
- Myrcia saxicola* var. *grandifolia* O.Berg, *Linnæa* 29 (2): 219 [June 1858] (Berg 1858).
- Myrcia sartoriana* O.Berg, *Linnæa* 29 (2): 220 [June 1858] (Berg 1858).
- Myrcia langsdorffii* O.Berg, *Fl. Bras. [Martius]* 14 (1): 562 [15 Jan. 1859] (Berg 1859).
- Myrcia klotzschiana* var. *impellucida* O.Berg, *Fl. Bras. [Martius]* 14 (1): 563 [15 Jan. 1859] (Berg 1859).
- Myrcia gracilis* var. *sessiliflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 564 [15 Jan. 1859] (Berg 1859).
- Myrcia tingens* O.Berg, *Fl. Bras. [Martius]* 14 (1): 564 [15 Jan. 1859] (Berg 1859).
- Myrcia riedeliana* O.Berg, *Fl. Bras. [Martius]* 14 (1): 565 [15 Jan. 1859] (Berg 1859).
- Myrcia splendens* var. *robustior* Kuntze, *Revis. Gen. Pl.* 1: 241 [5 Nov. 1891] (Kuntze 1891).
- Myrcia augustana* Kiaersk., *Enum. Myrt. Bras.* 52 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Myrcia guajavifolia* f. *grandifolia* Kiaersk., *Enum. Myrt. Bras.* 54 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Myrcia melanosticta* Kiaersk., *Enum. Myrt. Bras.* 54 [Oct.-Nov. 1893] (Kiaerskov 1893).
- Myrcia oxyoentophylla* Kiaersk., *Enum. Myrt. Bras.* 57 [Oct.-Nov. 1893] (Kiaerskov 1893), “oxyoëntophylla”.
- Myrcia martinicensis* Krug & Urb., *Bot. Jahrb. Syst.* 19 (5): 586 [26 Feb. 1895] (Krug & Urban 1895).
- Myrcia dictyoneura* Diels, *Bot. Jahrb. Syst.* 37 (5): 594 [30 Oct. 1906] (Diels 1906).
- Myrcia brachylopadia* Diels, *Bot. Jahrb. Syst.* 37 (5): 595 [30 Oct. 1906] (Diels 1906).
- Myrcia lamprosericea* Diels, *Bot. Jahrb. Syst.* 37 (5): 596 [30 Oct. 1906] (Diels 1906).
- Myrcia coroicensis* Rusby, *Bull. New York Bot. Gard.* 4 (14): 354 [7 Dec. 1907] (Rusby 1907).
- Myrcia luetzelburgii* Burret ex Luetzelb., Estud. Bot. Nordeste Braz. 3: 201 (Luetzelburg 1923), *nom. nud.*

Myrcia aguitensis Gleason, *Bull. Torrey Bot. Club* 58 (7): 409 [Oct. 1931] (Gleason 1931).

Myrcia compressa Gleason, *Bull. Torrey Bot. Club* 58 (7): 410 [Oct. 1931] (Gleason 1931).

Myrcia longicaudata Lundell, *Amer. Midl. Naturalist* 29 (2): 481 [5 Apr. 1943] (Lundell 1943).

Myrcia schippii Lundell, *Amer. Midl. Naturalist* 29 (2): 482 [5 Apr. 1943] (Lundell 1943).

Myrcia splendens var. *chrysocoma* McVaugh, *Fieldiana, Bot.* 29 (3): 193 [30 Nov. 1956] (McVaugh 1956).

Myrcia belizensis Lundell, *Wrightia* 2 (4): 213 (Lundell 1961).

Myrcia splendens var. *guantanamana* Borhidi & O.Muñiz, *Bot. Közlem.* 64 (1): 20 (Borhidi & Muñiz 1977).

Myrcia rostrata f. *flexuosa* Soares-Silva, *Bradea* 8 (48): 323 [22 Oct. 2002] (Soares-Silva 2002).

NOTES. — *Aulomyrcia costata* O.Berg is not based on *Myrcia costata* DC., which is treated separately by Berg in the same work (1857: 157). The type of *Myrtus stoupyi* Spreng. (i.e. that of *Myrtus bracteolaris* Poir.) was given to "citoyen" Lamarck by "citoyen" Stoupy (Poirier 1798: 411), hence the automatic correction of "stoupii".

VERNACULAR NAMES. — Pa: kagegut, karegut • Ka: kumete, kumeti, meli lan • Wp: inámu pita, pakeá'i, pila lea, pila lea u, pila lea wate'e, wila imí'i wu, wila umí'i wu • Nt: baaka bee sii, bee sii • Br: frutin-heira, guamirim, murta-grande, murta-parida.

HERBARIUM DATA (FG). — 94 collections at CAY. Sel. exs.: *J.B. Leblond* 114 (original material of *Eugenia fallax*: G).

INVENTORY DATA (FG). — 25 trees in 20 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 24.7$ cm.

[1308] *Myrcia subobliqua* (Benth.) Nied.

Nat. Pflanzenfam. [Engler & Prantl] 3 (7): 76 [Jan. 1893] (Niedenzu 1893). — *Eugenia subobliqua* Benth., *J. Bot. [Hooker]* 2: 322 (Bentham 1840). — *Aulomyrcia subobliqua* (Benth.) O.Berg, *Linnaea* 27 (1): 57 ["1854" publ. Nov. 1855] (Berg 1855).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: pakeá'i sī, pila lea.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P. Grenand* 1328.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.9$ cm.

[1309] *Myrcia subsessilis* O.Berg

Linnaea 31: 251 (Berg 1861-1862).

Myrcia subsessilis var. *subcordata* O.Berg, *Linnaea* 31: 251 (Berg 1861-1862).

Myrcia subsessilis var. *ovalis* O.Berg, *Linnaea* 31: 252 (Berg 1861-1862).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville* 3327.

SIZE. — Brazil, Amapá. *S.A. Mori* 17488 (MO), 20 m × 30 cm.

[1310] *Myrcia sylvatica* (G.Mey.) DC.

Prodr. [A. P. de Candolle] 3: 244 [mid Mar. 1828] (Candolle 1828). — *Myrtus sylvatica* G.Mey., *Prim. Fl. Esseq.* 191 [Nov. 1818] (Meyer 1818), "Myrthus". — *Myrcia ambigua* var. *sylvatica* (G.Mey.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 180 [15 May 1857] (Berg 1857).

Myrtus lucida L., *Syst. Nat.*, ed. 10, 2: 1056 [7 June 1759] (Linnaeus 1759).

Myrcia ambigua DC., *Prodr. [A. P. de Candolle]* 3: 252 [mid Mar. 1828] (Candolle 1828).

Myrcia ambigua var. *pauciflora* DC., *Prodr. [A. P. de Candolle]* 3: 252 [mid Mar. 1828] (Candolle 1828).

Myrcia ambigua var. *dives* O.Berg, *Fl. Bras. [Martius]* 14 (1): 180 [15 May 1857] (Berg 1857).

Myrcia ambigua var. *multiflora* O.Berg, *Fl. Bras. [Martius]* 14 (1): 180 [15 May 1857] (Berg 1857).

Myrcia ambigua var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 181 [15 May 1857] (Berg 1857).

Myrcia ambigua var. *rostrata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 181 [15 May 1857] (Berg 1857).

VERNACULAR NAMES. — Ka: pikililan • Wp: kasila, pila lea, pila lea sili.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 3064.

SIZE. — Brazil, Amazonas. *M. Pacheco et al.* 43 (MO), 14 m × 15.4 cm.

[1311] *Myrcia tomentosa* (Aubl.) DC.

Prodr. [A. P. de Candolle] 3: 245 [mid Mar. 1828] (Candolle 1828). — *Eugenia tomentosa* Aubl., *Hist. Pl. Guiane* 1: 504 [Jun.-Dec. 1775] (Aublet 1775). — *Myrtus aubletii* Spreng., *Syst. Veg. [Sprengel]* 2: 486 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Eugenia tomentosa*). — *Aguava tomentosa* (Aubl.) Raf., *Sylva Tellur.*: 107 (Rafinesque 1838). — *Cumetea tomentosa* (Aubl.) Raf., *Sylva Tellur.*: 106 (Rafinesque 1838). — *Aulomyrcia tomentosa* (Aubl.) Amshoff, *Recueil Trav. Bot. Néerl.* 39: 153 (Amshoff 1942).

Myrcia hirtiflora DC., *Prodr. [A. P. de Candolle]* 3: 249 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia hirtiflora* (DC.) O.Berg, *Linnaea* 27 (1): 40 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia curatellifolia DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828), "curatellaefolia". — *Myrtus curatellifolia* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828), "curatellaefolia", nom. nud. pro syn. — *Aulomyrcia curatellifolia* (DC.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 95 [15 May 1857] (Berg 1857), "curatellaefolia".

Myrcia prunifolia DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828). — *Aulomyrcia prunifolia* (DC.) O.Berg, *Linnaea* 27 (1): 55 ["1854" publ. Nov. 1855] (Berg 1855).

Myrcia pilosa DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828). — *Myrtus pilosa* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Myrcia prunifolia var. *angustior* DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828).

- Myrcia prunifolia* var. *obovata* DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828).
- Myrcia prunifolia* var. *ovata* DC., *Prodr. [A. P. de Candolle]* 3: 253 [mid Mar. 1828] (Candolle 1828).
- Myrcia puberula* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (18): 316 ["1829" publ. 25 Feb. 1832] (Cambessèdes 1832). — *Aulomyrcia puberula* (Cambess.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 94 [15 May 1857] (Berg 1857). — *Myrcianthes puberula* (Cambess.) Mattos, *Loefgrenia* 128: 1 [Apr. 2009] (Mattos 2009).
- Myrcia capivarhyensis* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19): 328 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832). — *Aulomyrcia capivarhyensis* (Cambess.) O.Berg, *Fl. Bras. [Martius]* 14 (1): 63 [15 May 1857] (Berg 1857).
- Myrcia floribunda* Miq., *Linnaea* 22: 534 (Miquel 1849).
- Aulomyrcia aureolanata* O.Berg, *Linnaea* 27 (1): 52 ["1854" publ. Nov. 1855] (Berg 1855), "aureo-lanata".
- Aulomyrcia alloiota* O.Berg, *Linnaea* 27 (1): 54 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia alloiota* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia rosulans* O.Berg, *Linnaea* 27 (1): 54 ["1854" publ. Nov. 1855] (Berg 1855). — *Myrcia rosulans* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 72 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia ottonis* O.Berg, *Linnaea* 27 (1): 55 ["1854" publ. Nov. 1855] (Berg 1855).
- Aulomyrcia confusa* O.Berg, *Linnaea* 27 (1): 56 ["1854" publ. Nov. 1855] (Berg 1855).
- Calyptromyrcia venosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 57 [15 May 1857] (Berg 1857). — *Myrcia venosa* (O.Berg) Mattos, *Loefgrenia* 128: 3 [Apr. 2009] (Mattos 2009).
- Aulomyrcia lancea* O.Berg, *Fl. Bras. [Martius]* 14 (1): 88 [15 May 1857] (Berg 1857). — *Myrcia lancea* (O.Berg) Mattos, *Arq. Bot. Estado São Paulo* 4 (2): 62 (Mattos 1966).
- Aulomyrcia spathulata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 91 [15 May 1857] (Berg 1857). — *Myrcia spathulata* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 72 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia alloiota* var. *cuneata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 92 [15 May 1857] (Berg 1857).
- Aulomyrcia alloiota* var. *obovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 92 [15 May 1857] (Berg 1857). — *Myrcia alloiota* var. *obovata* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia alloiota* var. *ovalis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 92 [15 May 1857] (Berg 1857).
- Aulomyrcia alloiota* var. *pyramidalis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 92 [15 May 1857] (Berg 1857).
- Aulomyrcia alloiota* var. *subcordata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 92 [15 May 1857] (Berg 1857). — *Myrcia alloiota* var. *subcordata* (O.Berg) N.Silveira, *Loefgrenia* 86: 1 (Silveira 1985).
- Aulomyrcia leucantha* O.Berg, *Fl. Bras. [Martius]* 14 (1): 93 [15 May 1857] (Berg 1857). — *Myrcia leucantha* (O.Berg) N.Silveira, *Roessleria* 7 (1): 66 (Silveira 1985).
- Aulomyrcia longipes* O.Berg, *Fl. Bras. [Martius]* 14 (1): 94 [15 May 1857] (Berg 1857). — *Myrcia longipes* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia longipes* var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 94 [15 May 1857] (Berg 1857).
- Aulomyrcia longipes* var. *obovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 94 [15 May 1857] (Berg 1857). — *Myrcia longipes* f. *obovata* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia longipes* var. *spathulata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 94 [15 May 1857] (Berg 1857).
- Aulomyrcia curatellifolia* var. *grandifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 95 [15 May 1857] (Berg 1857), "curatellaefolia". — *Myrcia curatellifolia* var. *grandifolia* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia curatellifolia* var. *parvifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 95 [15 May 1857] (Berg 1857), "curatellaefolia". — *Myrcia curatellifolia* var. *parvifolia* (O.Berg) Kaaersk., *Enum. Myrt. Bras.* 70 [Oct.-Nov. 1893] (Kaaersk 1893).
- Aulomyrcia prunifolia* var. *brevipes* O.Berg, *Fl. Bras. [Martius]* 14 (1): 95 [15 May 1857] (Berg 1857).
- Aulomyrcia prunifolia* var. *longipes* O.Berg, *Fl. Bras. [Martius]* 14 (1): 96 [15 May 1857] (Berg 1857).
- Aulomyrcia lanuginosa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 102 [15 May 1857] (Berg 1857). — *Myrcia lanuginosa* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 76 [Jan. 1893] (Niedenzu 1893), nom. illeg. hom., non O.Berg (1857).
- Myrcia piauhiensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 196 [15 May 1857] (Berg 1857).
- Aulomyrcia micrantha* O.Berg, *Fl. Bras. [Martius]* 14 (1): 517 [1 Feb. 1858] (Berg 1858).
- Aulomyrcia curatellifolia* var. *australis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 547 [15 Jan. 1859] (Berg 1859), "curatellaefolia".
- Aulomyrcia lanuginosa* var. *pyramidata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 548 [15 Jan. 1859] (Berg 1859).
- Myrcia marahanensis* Kaaersk., *Enum. Myrt. Bras.* 71 [Oct.-Nov. 1893] (Kaaersk 1893).
- Myrcia membranacea* Kaaersk., *Enum. Myrt. Bras.* 71 [Oct.-Nov. 1893] (Kaaersk 1893).
- Myrcia rhodeosepala* Kaaersk., *Enum. Myrt. Bras.* 75 [Oct.-Nov. 1893] (Kaaersk 1893).
- Myrcia verruculata* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 355 ["1894-96" publ. Dec. 1895] (Moore 1895).
- Eugenia vallis* Standl., *Ann. Missouri Bot. Gard.* 27: 323 (Standley 1940).
- HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000953642]).
- SIZE. — Up to 10 m tall (Holst 2003).

[1312] *Myrcia* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-J. de Granville 14611*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 11.6 \text{ cm}$.

[1313] *Myrcia* sp. B

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as “*Marlierea* FG-2”

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2329*, $\text{dbh} = 16.9 \text{ cm}$.

Genus *Myrcianthes* O.Berg

[1314] *Myrcianthes prodigiosa* McVaugh

Fieldiana, Bot. 29 (8): 492 [31 May 1963] (McVaugh 1963).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Barthélémy 3687*.

INVENTORY DATA (FG). — 14 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 38.5 \text{ cm}$.

Genus *Myrciaria* O.Berg

[1315] *Myrciaria floribunda* (H.West ex Willd.) O.Berg

Linnaea 27 (2-3): 330 [“1854” publ. Jan. 1856] (Berg 1856). — *Eugenia floribunda* H.West ex Willd., *Sp. Pl.*, ed. 4 2 (2): 960 [Dec. 1799] (Willdenow 1799). — *Myrtus floribunda* (H.West ex Willd.) Spreng., *Syst. Veg. [Sprengel] 2*: 484 [Jan.-May 1825] (Sprengel 1825).

Myrtus micrantha Nees & Mart., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 51 (Nees & Martius 1824), nom. illeg. hom., non Kunth (1823) [synonym of *Eugenia monticola* (Sw.) DC.]

Eugenia maximiliana DC., *Prodr. [A. P. de Candolle] 3*: 270 [mid Mar. 1828] (Candolle 1828). — *Myrciaria maximiliana* (DC.) O.Berg, *Linnaea* 27 (2-3): 329 [“1854” publ. Jan. 1856] (Berg 1856).

Eugenia ciliolata Cambess., *Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 2 (19)*: 344 [“1829” publ. 27 Oct. 1832] (Cambessèdes 1832). — *Myrciaria ciliolata* (Cambess.) O.Berg, *Linnaea* 27 (2-3): 327 [“1854” publ. Jan. 1856] (Berg 1856). — *Paramyrciaria ciliolata* (Cambess.) Rotman, *Bol. Soc. Argent. Bot.* 24 (3-4): 418 (Rotman 1986).

Eugenia salzmannii Benth., *J. Bot. [Hooker] 2*: 319 (Bentham 1840), “*Salzmanni*”. — *Myrciaria salzmannii* (Benth.) O.Berg, *Linnaea* 27 (2-3): 331 [“1854” publ. Jan. 1856] (Berg 1856), “*Salzmanni*”.

Myrtus verticillata Salzm. ex Benth., *J. Bot. [Hooker] 2*: 320 (Bentham 1840), nom. illeg. hom., non Vell. (Vellozo 1829), nom. nud.

Eugenia protracta Steud., *Flora 26 (45)*: 762 [17 Dec. 1843] (Steudel 1843). — *Myrciaria protracta* (Steud.) O.Berg, *Linnaea* 27 (2-3): 330 [“1854” publ. Jan. 1856] (Berg 1856).

Myrciaria uliginosa O.Berg, *Linnaea* 27 (2-3): 329 [“1854” publ. Jan. 1856] (Berg 1856).

Myrciaria verticillata O.Berg, *Linnaea* 27 (2-3): 332 [“1854” publ. Jan. 1856] (Berg 1856).

Myrciaria leucadendron O.Berg, *Fl. Bras. [Martius]* 14 (1): 364 [15 May 1857] (Berg 1857).

Myrciaria leucophloea O.Berg, *Fl. Bras. [Martius]* 14 (1): 370 [15 May 1857] (Berg 1857). — *Eugenia leucophloea* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 183 [Oct.-Nov. 1893] (Kiaerskov 1893).

Myrciaria leucophloea var. *conferta* O.Berg, *Fl. Bras. [Martius]* 14 (1): 370 [15 May 1857] (Berg 1857).

Myrciaria leucophloea var. *laxa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 370 [15 May 1857] (Berg 1857).

Myrciaria splendens O.Berg, *Fl. Bras. [Martius]* 14 (1): 371 [15 May 1857] (Berg 1857).

Myrciaria tenuiramis O.Berg, *Fl. Bras. [Martius]* 14 (1): 371 [15 May 1857] (Berg 1857).

Myrciaria maragnanensis O.Berg, *Fl. Bras. [Martius]* 14 (1): 372 [15 May 1857] (Berg 1857). — *Myrciaria maranhensis* O.Berg, *Fl. Bras. [Martius]* 14 (1): 652 [15 Jan. 1859] (Berg 1859), sphalm. — *Eugenia maranhensis* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 183 [Oct.-Nov. 1893] (Kiaerskov 1893).

Myrciaria tolypantha O.Berg, *Fl. Bras. [Martius]* 14 (1): 372 [15 May 1857] (Berg 1857).

Myrciaria tolypantha var. *angustifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 372 [15 May 1857] (Berg 1857).

Myrciaria tolypantha var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 372 [15 May 1857] (Berg 1857).

Myrciaria axillaris O.Berg, *Fl. Bras. [Martius]* 14 (1): 373 [15 May 1857] (Berg 1857).

Myrciaria prasina O.Berg, *Fl. Bras. [Martius]* 14 (1): 373 [15 May 1857] (Berg 1857).

Myrciaria schuechiana O.Berg, *Fl. Bras. [Martius]* 14 (1): 373 [15 May 1857] (Berg 1857), “*schüchiana*”.

Myrciaria amazonica O.Berg, *Fl. Bras. [Martius]* 14 (1): 374 [15 May 1857] (Berg 1857).

Myrciaria sellowiana O.Berg, *Fl. Bras. [Martius]* 14 (1): 374 [15 May 1857] (Berg 1857).

Myrciaria ferruginea O.Berg, *Fl. Bras. [Martius]* 14 (1): 597 [15 Jan. 1859] (Berg 1859).

Myrciaria longipes O.Berg, *Fl. Bras. [Martius]* 14 (1): 599 [15 Jan. 1859] (Berg 1859).

Myrciaria schuechiana var. *deflexa* O.Berg, *Fl. Bras. [Martius]* 14 (1): 599 [15 Jan. 1859] (Berg 1859), “*schüchiana*”.

Myrciaria schuechiana var. *latifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 599 [15 Jan. 1859] (Berg 1859), “*schüchiana*”.

Myrciaria longipes var. *opaca* O.Berg, *Fl. Bras. [Martius]* 14 (1): 600 [15 Jan. 1859] (Berg 1859).

Myrciaria longipes var. *pellucida* O.Berg, *Fl. Bras. [Martius]* 14 (1): 600 [15 Jan. 1859] (Berg 1859).

Marlierea brachymischa Kiaersk., *Enum. Myrt. Bras.* 47 [Oct.-Nov. 1893] (Kiaerskov 1893).

Eugenia leucophloea var. *warmingiana* Kiaersk., *Enum. Myrt. Bras.* 183 [Oct.-Nov. 1893] (Kiaerskov 1893). — *Myrciaria ciliolata* var. *warmingiana* (Kiaersk.) Mattos, *Loefgrenia* 78: 3 (Mattos 1983). — *Myrciaria leucophloea* var. *warmingiana* (Kiaersk.) Mattos, *Loefgrenia* 64: 3 (Mattos 1975).

Eugenia polyneura Urb., *Symb. Antill. [Urban]* 5 (3): 446 [20 May 1908] (Urban 1908), nom. illeg. hom., non Koord. & Valeton (Koorders & Valeton 1900).

Eugenia pycnoneura Urb., *Symb. Antill. [Urban]* 6 (1): 25 [15 July 1909] (Urban 1909).

Plinia acutissima Urb., *Ark. Bot.* 22A(10): 25 (Urban 1929).

Eugenia oneillii Lundell, *Bull. Torrey Bot. Club* 64 (8): 555 [Nov. 1937] (Lundell 1937), “O’Neillii”. — *Myrciaria oneillii* (Lundell) I.M.Johnst., *Sargentia* 8: 228 (Johnston 1949), “O’Neillii”.

Myrciaria arborea D.Legrand, *Sellowia* 13: 328 (Legrand 1961).

Siphoneugena cantareirae Mattos, *Ci. & Cult.* 19 (2): 332 (Mattos 1967).

Myrciaria longicaudata Lundell, *Wrightia* 4 (4): 144 (Lundell 1970).

Siphoneugena micrantha Kausel, *Lilloa* 36 (6): 111 (Kausel 1972).

Myrciaria mexicana Lundell, *Wrightia* 5 (2): 44 (Lundell 1974).

Myrciaria arborea var. *rostrata* Mattos, *Loefgrenia* 66: 3 (Mattos 1975).

NOTE. — Excluded synonyms: *Calyptanthes floribunda* Blume and *Caryophyllus floribundus* (Blume) Blume (from Java).

VERNACULAR NAMES. — Pa: awao, awau, awau-seinō • Ka: asaikiabe, seityape, tamunen seityape • Wp: kumesi • Cr: gouyavié-montangn • Br: goiaba-do-mato, goiabinha.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2218*.

INVENTORY DATA (FG). — 39 trees in 32 plots; $F_{\max} < 1\%$; dbh_{inv} = 33.3 cm.

[1316] *Myrciaria vismiifolia* (Benth.) O.Berg

Linnaea 27 (2-3): 336 [“1854” publ. Jan. 1856] (Berg 1856), “*vismeafolia*”. — *Eugenia vismiifolia* Benth., *J. Bot. [Hooker]* 2: 320 (Bentham 1840), “*vismeafolia*”.

NOTE. — The specific epithet refers to the genus *Vismia* Vand., hence it is corrected to *vismiifolia*.

VERNACULAR NAMES. — Wn: hapi, pēneju, takamalakma.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *Service Forestier 7911*, dbh = 20 cm.

Genus *Plinia* L.

[1317] *Plinia rivularis* (Cambess.) Rotman

Bol. Soc. Argent. Bot. 24 (1-2): 195 (Rotman 1985). — *Eugenia rivularis* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (19):

337 [“1829” publ. 27 Oct. 1832] (Cambessèdes 1832). — *Myrciaria rivularis* (Cambess.) O.Berg, *Linnaea* 27 (2-3): 320 [“1854” publ. Jan. 1856] (Berg 1856).

Myrciaria hagendorffii O.Berg, *Fl. Bras. [Martius]* 14 (1): 360 [15 May 1857] (Berg 1857). — *Eugenia hagendorffii* (O.Berg) Kiaersk., *Enum. Myrt. Bras.* 179 [Oct.-Nov. 1893] (Kiaerskov 1893).

Myrcia silvatica Barb.Rodr. ex Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 7: 803 (Chodat & Hassler 1907), nom. nud.

Eugenia variifolia Barb.Rodr. ex Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 7: 806 (Chodat & Hassler 1907), nom. nud.

Myrcia granulata R.O.Williams, *Fl. Trinidad & Tobago* 1 (6): 340 [Oct. 1934] (Williams 1934).

Myrciaria baporeti D.Legrand, *Anales Mus. Hist. Nat. Montevideo* 4 (11): 63 (Legrand 1936). — *Myrciariopsis baporeti* (D.Legrand) Kausel, *Ark. Bot.* n.s., 3: 509 (Kausel 1956). — *Siphoneugena baporeti* (D.Legrand) Kausel, *Lilloa* 32: 345 (Kausel 1966). — *Myrciaria rivularis* var. *baporeti* (D.Legrand) D.Legrand, *Fl. Ilustr. Catar. (Mirt.)*: 768 (Legrand 1978). — *Plinia baporeti* (D.Legrand) Rotman, *Darwiniana* 24 (1-4): 169 (Rotman 1982).

Siphoneugena legrandii Mattos & N.Silveira, *Loefgrenia* 87: 2 (Mattos & Silveira 1985).

VERNACULAR NAMES. — Pa: mpitit-kamwi (for *Plinia* sp.) • Br: quamirim, jaboticaba-rana.

HERBARIUM DATA (FG). — A single collection, *C. Baraloto et al. 8056*.

SIZE. — Up to 18 m tall (Holst 2003).

Genus *Psidium* L.

[1318] *Psidium acutangulum* DC.

Prodr. [A. P. de Candolle] 3: 233 [mid Mar. 1828] (Candolle 1828). — *Guajava acutangula* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 239 [5 Nov. 1891] (Kuntze 1891).

Psidium acutangulum var. *acidum* DC., *Prodr. [A. P. de Candolle]* 3: 233 [mid Mar. 1828] (Candolle 1828). — *Britoa acida* (DC.) O.Berg, *Linnaea* 27 (4): 436 [“1854” publ. Feb. 1856] (Berg 1856). — *Psidium acidum* Mart. ex O.Berg, *Linnaea* 27 (4): 436 [“1854” publ. Feb. 1856] (Berg 1856), nom. nud. pro syn. — *Psidium acidum* (DC.) Landrum, *Brittonia* 68 (4): 411 [epubl. 28 Apr. 2016] (Landrum 2016).

Psidium acutangulum var. *crassirame* O.Berg, *Fl. Bras. [Martius]* 14 (1): 409 [15 May 1857] (Berg 1857).

Psidium acutangulum var. *tenuirame* O.Berg, *Fl. Bras. [Martius]* 14 (1): 409 [15 May 1857] (Berg 1857).

Psidium grandiflorum Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 15: 228 (Ruiz & Pavón 1958), nom. illeg. hom., non Aubl. (Aublet 1775) [synonym of *Campomanesia grandiflora* (Aubl.) Sagot].

Psidium persoonii McVaugh, *Mem. New York Bot. Gard.* 18 (2): 255 (McVaugh 1969).

Psidium acutangulum var. *oblongatum* Mattos, *Loefgrenia* 94: 12 (Mattos 1989), “*oblongata*”.

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017).

VERNACULAR NAMES. — Te: ala'a • Wp: alali • Wn: takamalaimë • Nt: liba gobaya • Cr: gouyave-so • Br: araçá-pera.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *M. Fleury 2176*.

SIZE. — Up to 10 m tall (Holst 2003).

[1319] *Psidium oligospermum* Mart. ex DC.

Prodr. [A. P. de Candolle] 3: 236 [mid Mar. 1828] (Candolle 1828). — *Guajava oligosperma* (Mart. ex DC.) Kuntze, *Revis. Gen. Pl.* 1: 239 [5 Nov. 1891] (Kuntze 1891). — *Mitropsidium oligospermum* (Mart. ex DC.) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 486 [30 Mar. 1941] (Burret 1941).

Calyptranthes eugenoides Cambess., *Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.)* 2 (19): 370 ["1829" publ. 27 Oct. 1832] (Cambessèdes 1832), "Eugenoides". — *Mitrantha eugenoides* (Cambess.) O.Berg, *Linnaea* 27 (2-3): 317 ["1854" publ. Jan. 1856] (Berg 1856). — *Chytraculia eugenoides* (Cambess.) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891). — *Psidium eugenoides* (Cambess.) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 69 [Jan. 1893] (Niedenzu 1893). — *Mitropsidium eugenoides* (Cambess.) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 489 [30 Mar. 1941] (Burret 1941).

Psidium galapageium Hook.f., *Trans. Linn. Soc. London* 20 (2): 224 (Hooker 1847).

Psidium ciliatum O.Berg, *Linnaea* 27 (2-3): 353 ["1854" publ. Jan. 1856] (Berg 1856), nom. illeg. hom., non Benth. (Bentham 1840) [synonym of *P. salutare* (Kunth) O.Berg].

Mitrantha gardneriana O.Berg, *Fl. Bras. [Martius]* 14 (1): 354 [15 May 1857] (Berg 1857). — *Chytraculia gardneriana* (O.Berg) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891). — *Mitropsidium gardnerianum* (O.Berg) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 487 [30 Mar. 1941] (Burret 1941).

Mitrantha eugenoides var. *oblongifolia* O.Berg, *Fl. Bras. [Martius]* 14 (1): 355 [15 May 1857] (Berg 1857), nom. illeg. (species basionym in synonymy).

Mitrantha eugenoides var. *ovata* O.Berg, *Fl. Bras. [Martius]* 14 (1): 355 [15 May 1857] (Berg 1857).

Mitrantha sartoriana O.Berg, *Linnaea* 29 (2): 248 [June 1858] (Berg 1858). — *Chytraculia sartoriana* (O.Berg) Kuntze, *Revis. Gen. Pl.* 1: 238 [5 Nov. 1891] (Kuntze 1891). — *Psidium sartorianum* (O.Berg) Nied., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (7): 69 [Jan. 1893] (Niedenzu 1893). — *Calyptropsisidium sartorianum* (O.Berg) Krug & Urb., *Bot. Jahrb. Syst.* 19 (4): 571 [28 Dec. 1894] (Krug & Urban 1894). — *Mitropsidium sartorianum* (O.Berg) Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 487 [30 Mar. 1941] (Burret 1941).

Calycorectes protractus Griseb., *Cat. Pl. Cub. [Grisebach]* 284 [May-Aug. 1866] (Grisebach 1866). — *Psidium protractum* (Griseb.) Lundell, *Wrightia* 5 (3): 70 (Lundell 1974).

Calyptranthes tonduzii Donn.Sm., *Bot. Gaz.* 23 (4): 245 [22 Apr. 1897] (Donnell Smith 1897).

Psidium paucinerve Urb., *Symb. Antill. [Urban]* 9 (2): 82 [1 Jan. 1923] (Urban 1923). — *Myrtus paucinervia* (Urb.) Landrum, *Canotia* 13: 49 (Landrum 2017), "paucinerve", nom. inval. pro syn.

Psidium claraense Urb., *Symb. Antill. [Urban]* 9 (4): 466 [15 Mar. 1928] (Urban 1928). — *Myrtus claraensis* (Urb.) Bisce, *Ciencias (Havana)*, ser. 10, 12: 10 ["1976" publ. 1977] (Bisce 1977).

Psidium microphyllum Britton, *Sci. Surv. Porto Rico & Virgin Islands* 6: 555 (Britton 1930).

Psidium soccorense I.M.Johnst., *Proc. Calif. Acad. Sci.*, ser. 4, 20: 81 (Johnston 1931).

Mitropsidium ob lanceolatum Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 487 [30 Mar. 1941] (Burret 1941).

Mitropsidium pittieri Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 488 [30 Mar. 1941] (Burret 1941).

Psidium yucatanense Lundell, *Contr. Univ. Michigan Herb.* 7: 35 (Lundell 1942). — *Psidium sartorianum* var. *yucatanense* (Lundell) McVaugh, *Fieldiana, Bot.* 29 (8): 527 [31 May 1963] (McVaugh 1963).

Psidium solisii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23 (3): 133 [14 Feb. 1944] (Standley 1944).

Psidium minutiflorum Amshoff, *Recueil Trav. Bot. Néerl.* 42: 19 (Amshoff 1950).

Psidium molinae Amshoff, *Acta Bot. Neerl.* 5 (3): 277 [Nov. 1956] (Amshoff 1956), "Molinae".

Psidium galapageium var. *howellii* D.M.Porter, *Ann. Missouri Bot. Gard.* 55 (3): 370 (Porter 1969).

NOTE. — McVaugh based *Psidium sartorianum* var. *yucatanense* on *P. yucatanense* Lundell, but argued that the former name should not be treated as a new combination, on the mistaken belief that *P. yucatanense* was a taxonomic synonym of *Calycorectes protractus* Griseb. As *P. yucatanense* was validly published, all requirements were fulfilled for a new combination.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5029*.

SIZE. — Brazil, Pará. *T.C. Plowman et al. 8730* (MO), 15 m × 30 cm.

[1320] *Myrtaceae* sp. A

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as "Myrtaceae FG-2".

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier et al. 5812*.

INVENTORY DATA (FG). — 11 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.5$ cm.

[1321] *Myrtaceae* sp. B

NOTE. — Collections at CAY are provisionally classified by B. K. Holst as "Myrtaceae FG-3".

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5411*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 10.8$ cm.

[1322] *Myrtaceae* sp. C

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & M.-F. Prévost 2956*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.3$ cm.

Family NYCTAGINACEAE Juss.
Genus *Guapira* Aubl.

[1323] *Guapira eggersiana* (Heimerl) Lundell

Wrightia 4 (2): 80 [31 Dec. 1968] (Lundell 1968). — *Pisonia eggersiana* Heimerl, *Bot. Jahrb. Syst.* 21 (5): 627 [12 May 1896] (Heimerl 1896). — *Torrubia eggersiana* (Heimerl) Standl., *Contr. U.S. Natl. Herb.* 18 (3): 100 [11 Feb. 1916] (Standley 1916).

Pisonia broadwayana Heimerl, *Repert. Spec. Nov. Regni Veg.* 17: 1 (Heimerl 1921). — *Torrubia broadwayana* (Heimerl) Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 8 (5): 308 (Standley 1931). — *Guapira broadwayana* (Heimerl) Lundell, *Wrightia* 4 (2): 80 [31 Dec. 1968] (Lundell 1968).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: aaku-gawut • Ka: wayamu sasamili • Wp: inimopo'i piyunga, pikau ki'iy, pikau meyu • Cr: grènn-malenbé • Br: carvoeiro.

HERBARIUM DATA (FG). — 55 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3833.

INVENTORY DATA (FG). — 55 trees in 22 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 51.4$ cm.

[1324] *Guapira guianensis* Aubl.

Hist. Pl. Guiane 1: 308 [Jun.-Dec. 1775] (Aublet 1775), “*Quapira Guyannensis*” on plate. — *Pisonia guianensis* (Aubl.) R.A.Howard, *J. Arnold Arbor.* 64 (2): 270 [Apr. 1983] (Howard 1983).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 1 collection from French Guiana: *J.B. Aublet s.n.* (original material P-JU, P00675567).

SIZE. — Venezuela, Aragua. *O. Huber* 333 (MO), 7 m × 11 cm.

[1325] *Guapira salicifolia* (Heimerl) Lundell

Wrightia 4 (2): 83 [31 Dec. 1968] (Lundell 1968). — *Pisonia salicifolia* Heimerl, *Symb. Antill. [Urban]* 7 (2): 216 [15 June 1912] (Heimerl 1912). — *Torrubia salicifolia* (Heimerl) Standl., *Contr. U.S. Natl. Herb.* 18 (3): 101 [11 Feb. 1916] (Standley 1916).

Pisonia albiflora Heimerl, *Bull. Misc. Inform. Kew* 1932 (5): 219 [27 June 1932] (Heimerl 1932).

Torrubia heimeriana Standl., *Lloydia* 2 (3): 177 (Standley 1939). — *Guapira heimeriana* (Standl.) Lundell, *Wrightia* 4 (2): 81 [31 Dec. 1968] (Lundell 1968).

VERNACULAR NAMES. — Wp: inimopo'i piyunga • Cr: grènn-malenbé • Br: carvoeiro.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P. Grenand* 2107.

INVENTORY DATA (FG). — 1 tree, dbh = 14.6 cm.

[1326] *Guapira* sp. A

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2071*.

INVENTORY DATA (FG). — 19 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.6$ cm.

Genus *Neea* Ruiz & Pav.

[1327] *Neea constricta* Spruce ex J.A.Schmidt

Fl. Bras. [Martius] 14 (2): 368 [1 May 1872] (Schmidt 1872).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *S.A. Mori & C. Gracie* 24205.

SIZE. — Up to 15 cm dbh (DeFilipps 2002).

[1328] *Neea floribunda* Poepp. & Endl. (Fig. 43A)

Nova genera ac species plantarum [Poeppig & Endlicher] 2: 46 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838), “*Neaea*”.

Neea cauliflora Heimerl, *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 16c: 129 (Heimerl 1934).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: miukat-kamwi, mirukat-kamwi • Ka: sakusaku • Wp: inimopo'i piyunga.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom* 15333.

INVENTORY DATA (FG). — 90 trees in 35 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 37.2$ cm.

[1329] *Neea mollis* Spruce ex J.A.Schmidt

Fl. Bras. [Martius] 14 (2): 367 [1 May 1872] (Schmidt 1872).

VERNACULAR NAMES. — Pa: kwepan-wašiunu • Ka: sakusaku • Wp: alasiku, inimopo'i piyunga, ka'a sala • Br: joão-mole.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *D. Sabatier et al. 4623*, dbh = 14.6 cm.

[1330] *Neea ovalifolia* Spruce ex J.A.Schmidt

Fl. Bras. [Martius] 14 (2): 368 [1 May 1872] (Schmidt 1872).

VERNACULAR NAMES. — Wp: kwapo'i sili.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2249*.

SIZE. — Up to 35 cm dbh (DeFilips 2002).

[1331] *Neea spruceana* Heimerl

Notizbl. Königl. Bot. Gart. Berlin 6: 131 [20 Mar. 1914] (Heimerl 1914).

Neea weberbaueri Heimerl, *Bot. Jahrb. Syst.* 54 (1, Beibl. 117): 38 [26 Apr. 1916] (Heimerl 1916).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville 652*.

SIZE. — Up to 20 cm dbh (DeFilips & Maina 2003).

[1332] *Neea* sp. A
(Fig. 43B)

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3385*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 26.9$ cm.

[1333] *Neea* sp. B

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2760*.

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} = 2.8\%$; $\text{dbh}_{\text{inv}} = 22.4$ cm.

[1334] *Neea* sp. C

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5082*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 26.3$ cm.

[1335] Nyctaginaceae sp. A

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5334*.

INVENTORY DATA (FG). — 74 trees in 29 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 57.3$ cm.

[1336] Nyctaginaceae sp. B

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1877*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 60$ cm.

[1337] Nyctaginaceae sp. C

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3169*.

INVENTORY DATA (FG). — 15 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 34.3$ cm.

[1338] Nyctaginaceae sp. D

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 4880*.

INVENTORY DATA (FG). — 21 trees in 2 plots; $F_{\max} = 3\%$; $\text{dbh}_{\text{inv}} = 23.6$ cm.

Family OCHNACEAE DC.
Genus *Elvasia* DC.

[1339] *Elvasia elvasioides* (Planch.) Gilg

Nat. Pflanzenfam. [Engler & Prantl] 3 (6): 145 (Gilg 1893). — *Hostmannia elvasioides* Planch., *Icon. Pl.* 8 [n.s., 4]: t. 709 [May 1845] (Planchon 1845). — *Elvasia hostmannia* Planch., *London J. Bot.* 5: 648 (Planchon 1846), nom. illeg. superfl. (based on *Hostmannia elvasioides*).

Hostmannia sagotii Tiegh., *Ann. Sci. Nat., Bot. sér. 8, 16:* 414 (Tieghem 1902), “*Sagoti*”. — *Elvasia sagotii* (Tiegh.) Lemée, *Fl. Guyane Franç.* 3: 8 (Lemée 1954), “*Sagoti*”.

Elvasia caurensis Pittier, *Bol. Soc. Venez. Ci. Nat.* 6 (41): 17 [“1939” publ. 1940] (Pittier 1940).

VERNACULAR NAMES. — Pa: yauknabui • Ka: pakila eu.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3996*, dbh = 30.3 cm.

INVENTORY DATA (FG). — 12 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 27.9$ cm.

[1340] *Elvasia macrostipularis* Sastre & Lescure
(Fig. 44A)

Caldasia 12 (57): 134 (Sastre & Lescure 1978).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4664*.

INVENTORY DATA (FG). — 292 trees in 31 plots; $F_{\max} = 6.5\%$; $\text{dbh}_{\text{inv}} = 26.6$ cm.

Genus *Lacunaria* Ducke

[1341] *Lacunaria crenata* (Tul.) A.C.Sm.
(Fig. 44B)

Trop. Woods 58: 31 (Smith 1939). — *Quiina crenata* Tul., *Ann. Sci. Nat., Bot. sér. 3, 11:* 163 (Tulasne 1849).

Quiina panamensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 236 [24 Oct. 1929] (Standley 1929). — *Lacunaria panamensis* (Standl.) Standl., *Ann. Missouri Bot. Gard.* 29 (4): 358 (Standley 1942).

Lacunaria pauciflora Ducke, *Arch. Jard. Bot. Rio de Janeiro* 5: 169 (Ducke 1930).

VERNACULAR NAMES. — Ka: woko molokotoli • Wp: tulumele • Wn: kulataju • Br: moella-de-mutum.

HERBARIUM DATA (FG). — 67 collections at CAY. Sel. exs.: G.S. Perrottet s.n. (type P[P05466483]).

INVENTORY DATA (FG). — 67 trees in 49 plots; $F_{\max} < 1\%$; $dbh_{inv} = 21.8$ cm.

[1342] *Lacunaria jenmanii* (Oliv.) Ducke subsp. *jenmanii*

Arch. Jard. Bot. Rio de Janeiro 5: 171 (Ducke 1930). — *Touroulia jenmanii* Oliv., *Hooker's Icon. Pl.* 20 [ser. 3, 10]: t. 1998 [Aug. 1891] (Oliver 1891), “Jenmani”.

Quiina silvatica Pulle, *Recueil Trav. Bot. Néerl.* 6: 277 (Pulle 1909). — *Lacunaria silvatica* (Pulle) A.C.Sm., *Trop. Woods* 58: 31 (Smith 1939).

Lacunaria grandiflora Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 140 (Ducke 1925).

VERNACULAR NAMES. — Ka: woko molokotoli • Wp: mitū'ay, tulumele u • Br: moella-de-mutum.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: D. Sabatier 1423.

INVENTORY DATA (FG). — 52 trees in 40 plots; $F_{\max} < 1\%$; $dbh_{inv} = 31.8$ cm.

[1343] *Lacunaria jenmanii* subsp. *subsessilis*
J.V.Schneid. & Zizka

Syst. Bot. 37 (1): 179 [1 Jan. 2012] (Schneider & Zizka 2012).

VERNACULAR NAMES. — Pa: avakni-awak puverna, avakni-awak-seine.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: J.-F. Villiers & C. Feuillet 1974.

SIZE. — Up to 25 m tall (Schneider & Zizka 2012).

[1344] *Lacunaria umbonata* Pires

Bol. Técn. Inst. Agron. N. 28: 46 [1954] (Pires 1953).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, R.L. Fróes 26671 (type IAN[IAN059825]).

INVENTORY DATA (FG). — 1 tree, $dbh = 20.7$ cm.

Genus *Ouratea* Aubl.

[1345] *Ouratea candollei* (Planch.) Tiegh.

J. Bot. [Morot] 16: 186 [June 1902] (Tieghem 1902). — *Gomphia candollei* Planch., *London J. Bot.* 6: 4 (Planchon 1847).

NOTE. — The binomial *Ouratea candollei* was first invalidly published by Engler (1876: 314) in synonymy under *O. guianensis* Aubl.

VERNACULAR NAMES. — Pa: psuk-awak • Cr: malmani • Br: batiputá.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: J. Martin 148 (holo-, P[P00542206]; iso-, BR[BR0000005281756], K[K000382126, K000382127, K000382128], MO[MO-279755], P[P00542207]).

INVENTORY DATA (FG). — 15 trees in 3 plots; $F_{\max} = 1.8\%$; $dbh_{inv} = 22.6$ cm.

[1346] *Ouratea cerebroidea* Sastre

Bull. Mus. Natl. Hist. Nat., B, Adansonia 8 (4): 413 [“1986” publ. 1987] (Sastre 1987).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: yauknabui.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: C. Sastre 6405 (holo-, P[P00542192]; iso-, P[P00542193], U[U0005264]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 18.9$ cm.

[1347] *Ouratea decagyna* Maguire

Bull. Torrey Bot. Club 75 (6): 644 [Dec. 1948] (Maguire 1948).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: D. Sabatier & M.-F. Prévost 2944.

INVENTORY DATA (FG). — 31 trees in 25 plots; $F_{\max} < 1\%$; $dbh_{inv} = 28$ cm.

[1348] *Ouratea franciniae* Sastre

Bull. Mus. Natl. Hist. Nat., B, Adansonia 8 (4): 417 [“1986” publ. 1987] (Sastre 1987).

VERNACULAR NAMES. — Wp: wai tawa.

HERBARIUM DATA (FG). — Known only from the type: P Grenand 510.

SIZE. — Up to 10 m tall (Sastre 1986).

[1349] *Ouratea guianensis* Aubl.

Hist. Pl. Guiane 1: 397 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: yauknabui • Ka: akale tapulala kili • Wp: tuká nákū • Wn: potipotpiu • Nt: weti tiki • Cr: malmani • Br: batiputá.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000797998](#)]).

INVENTORY DATA (FG). — 17 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 33.7$ cm.

[1350] *Ouratea leblondii* (Tiegh.) Lemée

Fl. Guyane Franç. 3: 8 (Lemée 1954), “Leblondi”. — *Camptouratea leblondii* Tiegh., *Ann. Sci. Nat., Bot. sér. 8, 16:* 205 (Tieghem 1902), “Leblondi”.

Camptouratea sagotii Tiegh., *Ann. Sci. Nat., Bot. sér. 8, 16:* 211 (Tieghem 1902), “Sagoti”. — *Ouratea sagotii* (Tiegh.) Lemée, *Fl. Guyane Franç.* 3: 9 (Lemée 1954).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: ii.

HERBARIUM DATA (FG). — 94 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (lecto-, P[[P00542279](#)], designated by Sastre & Offroy [2016: 60]; isolecto-, K[not seen], P[[P00582165](#)]); *C. Sastre & C. Moretti 4020*, 15 m.

[1351] *Ouratea macrocarpa* Sastre

Bull. Mus. Natl. Hist. Nat., B, Adansonia 8 (4): 414 [“1986” publ. 1987] (Sastre 1987).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-J. de Granville 44* (holo-, P[[P00542348](#)]; iso-, CAY[[CAY024895](#), [CAY024896](#)], COL[[COL000002626](#)], K[[K000382115](#)], NY[[00428959](#)], P[[P00542349](#)], RB[[RB00542384](#)], SI[[SI003068](#), [SI003069](#)], U[[U0008103](#)], US[[00289224](#)]).

SIZE. — Up to 17 m tall (Sastre 1986).

[1352] *Ouratea melinonii* (Tiegh.) Lemée

Fl. Guyane Franç. 3: 12 (Lemée 1954). — *Cercouratea melinonii* Tiegh., *Ann. Sci. Nat., Bot. sér. 8, 16:* 278 (Tieghem 1902), “Melinoni”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: ii.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *E.M. Mélénion s.n.* (holo-, P[[P00542463](#)]; iso-, P[[P00542464](#)]).

INVENTORY DATA (FG). — 40 trees in 21 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 18.8$ cm.

[1353] *Ouratea retrorsa* Sastre

Adansonia, sér. 3, 29 (1): 89 (Sastre 2007).

HERBARIUM DATA (FG). — Known only from the type: *D. Larpin 1021* (holo-, P[[P00542460](#)]; iso-, CAY[[CAY079504](#)]).

SIZE. — Up to 12 m tall (Sastre 2007).

[1354] *Ouratea scottii* Sastre
(Fig. 44C)

Brittonia 46 (4): 309 (Sastre 1994).

VERNACULAR NAMES. — Wp: tuká nili • Cr: malmani • Br: batiputá.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *S.A. Mori et al. 14927* (holo-, NY[[00345875](#)] ; iso-, CAY[[CAY002334](#)], P[[P00542208](#)]).

INVENTORY DATA (FG). — 19 trees in 6 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 20.7$ cm.

Genus *Quiina* Aubl.

[1355] *Quiina berryi* J.V.Schneid. & Zizka

Candollea 58 (2): 464 (Schneider & Zizka 2003).

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville 7755*.

SIZE. — Up to 20 m tall (Schneider & Zizka 2016).

[1356] *Quiina cruegeriana* Griseb.

Fl. Brit. W.I. [Grisebach] 106 [June 1859] (Grisebach 1859).

Quiina guianensis Crueg, ex Griseb., *Fl. Brit. W.I. [Grisebach] 106* [June 1859] (Grisebach 1859), *nom. nud. pro syn.*

Quiina peruviana Engl., *Fl. Bras. [Martius] 12 (1):* 481 [1 Apr. 1888] (Engler 1888).

Quiina tessmannii Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 298 [10 Nov. 1931] (Mildbraed 1931).

Quiina cajambrensis Cuatrec., *Trop. Woods* 96: 41 (Cuatrecasas 1950).

Quiina congesta R.S.Cowan, *Brittonia* 8 (4): 239 [Jan. 1957] (Cowan 1957).

Quiina guaporensis Pires, *Bol. Técn. Inst. Agron. N.* 38: 31 (Pires 1960).

Lacunaria colonensis D'Arcy, *Ann. Missouri Bot. Gard.* 67 (4): 967 [“1980” publ. 1981] (D'Arcy 1981). — *Quiina colonensis* (D'Arcy) D'Arcy, *Monogr. Syst. Bot. Missouri Bot. Gard.* 18: 534 (D'Arcy 1987).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Grand 1902*.

SIZE. — Up to 15 m tall (Schneider & Zizka 2012).

[1357] *Quiina guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 19 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

VERNACULAR NAMES. — Ka: wokunse • Wp: iwaka'i sili, waka'i sili • Nt: dyabatee, pikin busi suku • Br: quinarana.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000073181](#)]).

INVENTORY DATA (FG). — 18 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.2$ cm.




FIG. 43. — Nyctaginaceae: **A**, *Neea floribunda* Poepp. & Endl. (*D. Sabatier & J.-F. Molino* 5616); **B**, *Neea* sp. A (*M.-F. Prévost & D. Sabatier* 3385). Olacaceae: **C**, *Dulacia guianensis* (Engl.) Kuntze; **D**, *Heisteria ovata* Benth. (*D. Sabatier & J.-F. Molino* 5086). A, C, D, © D. Sabatier/IRD; B, © M.-F. Prévost/IRD.

[1358] *Quiina integrifolia* Pulle

Recueil Trav. Bot. Néerl. 9: 153 (Pulle 1912).

Quiina rigidifolia Pires, *Bol. Técn. Inst. Agron. N.* 20: 45 (Pires 1950).

VERNACULAR NAMES. — Pa: yauknabui-puvemna • Br: quinarana.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 4977.

INVENTORY DATA (FG). — 58 trees in 35 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 26.3$ cm.

[1359] *Quiina leptoclada* Tul.

Ann. Sci. Nat., Bot. sér. 3, 11: 159 (Tulasne 1849).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *E.M. Mélinon* 147, 1845 (holo-, P[P02441362]; iso-, K[K000328614], P[P00071950, P02441363]).

INVENTORY DATA (FG). — 5 trees in 4 plots; $\text{dbh}_{\text{inv}} = 16.5$ cm.

[1360] *Quiina obovata* Tul.

Ann. Sci. Nat., Bot. sér. 3, 11: 157 (Tulasne 1849).

Quiina decaisneana Planch. & Triana, *Ann. Sci. Nat., Bot. sér. 4*, 15: 315 (Planchon & Triana 1861).

Quiina albiflora A.C.Sm., *Trop. Woods* 58: 30 (Smith 1939).

Quiina oblanceolata Sandwith, *Bull. Misc. Inform. Kew* 1939 (10): 546 [“1939” publ. 6 Jan. 1940] (Sandwith 1940).

VERNACULAR NAMES. — Pa: kwatri-wašiunô-purubumna • Ka: wok-unse • Wp: iwaka'i, waka'i u • Nt: dyabatee • Br: quinarana.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, P[P00071947]; iso-, B[B 10 0294392], K[K000328613], P[P02441367, P02441368], SI[SI004194], U[U0282664]).

INVENTORY DATA (FG). — 61 trees in 49 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.9$ cm.

[1361] *Quiina oiapocensis* Pires

Bol. Técn. Inst. Agron. N. 38: 32 (Pires 1960).

VERNACULAR NAMES. — Pa: kwatri-duwë • Br: quinarana.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier* 1541.

INVENTORY DATA (FG). — 40 trees in 22 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22.3$ cm.



FIG. 44. — Ochnaceae: **A**, *Elvasia macrostipularis* Sastre & Lescure (J.-F. Molino & D. Sabatier 2099); **B**, *Lacunaria crenata* (Tul.) A.C.Sm. (D. Sabatier et al. 4805); **C**, *Ouratea scottii* Sastre (D. Sabatier & J.-F. Molino 5071); **D**, *Quiina* sp. A (D. Sabatier & J.-F. Molino 5197). © D. Sabatier/IRD.

[1362] *Quiina pteridophylla* (Radlk.) Pires

Bol. Técn. Inst. Agron. N. 20: 48 (Pires 1950). — *Touroulia pteridophylla* Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 19: 218 (Radlkofer 1889).

Quiina acutangula Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 143 (Ducke 1925).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori et al. 23207*.

SIZE. — Up to 12 m tall (Schneider & Zizka 2004).

[1363] *Quiina sessilis* Choisy ex Planch. & Triana

Ann. Sci. Nat., Bot. sér. 4, 15: 312 (Planchon & Triana 1861).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: kwatri-wašiunō, kwatri-wašiunō-duwē.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (holo-, G[G00389893]; iso-, G[G00389894]; possible iso-, MPU[MPU014417]).

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; dbh_{inv} = 12.7 cm.

[1364] *Quiina yatuensis* J.V.Schneid. & Zizka

Novon 7 (4): 406 [“1997” publ. 1998] (Schneider & Zizka 1998).

HERBARIUM DATA (FG). — A single collection, *M.-F. Prévost & D. Sabatier 4623*, dbh 12.9 cm.

[1365] *Quiina* sp. A
(Fig. 44D)

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2661*.

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; dbh_{inv} = 28.3 cm.

Genus *Touroulia* Aubl.[1366] *Touroulia guianensis* Aubl.

Hist. Pl. Guiane 1: 492 [Jun.-Dec. 1775] (Aublet 1775), “*Guyanensis*” on plate. — *Robinsonia guianensis* J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 796 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. superfl.* (based on *Touroulia guianensis*). — *Robinsonia melanthifolia* Willd., *Sp. Pl.*, ed. 4 2 (2): 999 [Dec. 1799] (Willdenow 1799), *nom. illeg. superfl.* (based on *Touroulia guianensis*). — *Touroulia solitaria* Stokes, *Bot. Mat. Med.* 3: 105 (Stokes 1812), “*Touroulia*”, *nom. illeg. superfl.* (based on *Robinsonia melanthifolia*).

VERNACULAR NAMES. — Pa: yit-ára • Wp: wila taiwa • Nt: yooka wiwii • Cr: bwa-flanbo.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P[P00114408] designated by Zizka & Schneider [1999: 232]).

INVENTORY DATA (FG). — 44 trees in 31 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 26.9$ cm.

Family OLACACEAE R.Br.
Genus *Cathedra* Miers

[1367] *Cathedra acuminata* (Benth.) Miers

Ann. Mag. Nat. Hist., ser. 3, 4: 361 (Miers 1859). — *Diplocrater acuminatus* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 367 (Bentham 1851).

Cathedra crassifolia Benth. ex Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 4: 361 (Miers 1859).

Cathedra aestuaria Sleumer, *Repert. Spec. Nov. Regni Veg.* 39: 279 (Sleumer 1936).

Cathedra inaequilatera Sleumer, *Repert. Spec. Nov. Regni Veg.* 39: 280 (Sleumer 1936).

Cathedra oblonga Sleumer, *Repert. Spec. Nov. Regni Veg.* 39: 280 (Sleumer 1936).

Cathedra caurensis Pittier, *Bol. Soc. Venez. Ci. Nat.* 6 (41): 5 [“1939” publ. 1940] (Pittier 1940).

Cathedra guianensis Sandwith, *Kew Bull.* 10 (3): 469 [20 Dec. 1955] (Sandwith 1955).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier 5778*.

INVENTORY DATA (FG). — 14 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 28.8$ cm.

Genus *Chaunochiton* Benth.[1368] *Chaunochiton kappleri* (Sagot ex Engl.) Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 41 (Ducke 1922). — *Heisteria kappleri* Sagot ex Engl., *Fl. Bras. [Martius]* 12 (2): 14 [1 Dec. 1872] (Engler 1872). — *Sagotanthus kappleri* (Sagot ex Engl.) Dyer, *Index Kew. Suppl.* 2: 162 (Dyer 1904).

Chaunochiton breviflorum Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 41 (Ducke 1922).

Chaunochiton purpurascens Rizzini, *Dusenia* 1 (5): 289 (Rizzini 1950).

VERNACULAR NAMES. — Ka: aluku elepali, apesiya • Wp: wila pile lu • Nt: auman pisi, gawenti, tu pikin weko • Br: pau-vermelho.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *P.A. Sagot 1198*, 1858 (holo-, P[P02441958]; iso-, B[B 10 0248312], BM[BM000839877], BR[BR0000005280810], GH[GH00035955], GOET[GOET008319], K[K000580931], L[L0038980], P[P02441959], P02441960, P02441961, P02441962, P02441963], S[S-R-2779], U[U0005296]).

INVENTORY DATA (FG). — 127 trees in 71 plots; $F_{\max} = 1.4\%$; $dbh_{\text{inv}} = 72.6$ cm.

Genus *Dulacia* Vell.[1369] *Dulacia guianensis* (Engl.) Kuntze
(Fig. 43C)

Revis. Gen. Pl. 1: 111 [5 Nov. 1891] (Kuntze 1891). — *Liriosma guianensis* Engl., *Fl. Bras. [Martius]* 12 (2): 25 [1 Dec. 1872] (Engler 1872). — *Olax guianensis* (Engl.) Christenh. & Byng, *Global Fl.* 4: 143 [9 Feb. 2018] (Christenhusz & Byng 2018).

Olax schomburgkii Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1177 [“1848” publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Liriosma cerifera A.C.Sm., *Brittonia* 2 (2): 146 (Smith 1936).

NOTE. — According to Malécot & Nickrent (2008), *Dulacia* Vell. is possibly embedded in *Olax* L.

VERNACULAR NAMES. — Pa: balata-kamwi.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2108*.

INVENTORY DATA (FG). — 18 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38$ cm.

Genus *Heisteria* Jacq.[1370] *Heisteria barbata* Cuatrec.

Trop. Woods 101: 26 (Cuatrecasas 1955).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 5403*.

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.2$ cm.

[1371] *Heisteria densifrons* Engl.

Fl. Bras. [Martius] 12 (2): 17 [1 Dec. 1872] (Engler 1872).

Heisteria microcarpa Spruce ex Engl., *Fl. Bras. [Martius]* 12 (2): 21 [1 Dec. 1872] (Engler 1872).

Heisteria microcalyx Sagot, *Ann. Sci. Nat., Bot. sér. 6, 11:* 159 (Sagot 1881).

Heisteria sessilis Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 8 (Ducke 1925).

Heisteria parvicalyx A.C.Sm., *Brittonia* 2 (2): 147 (Smith 1936).

VERNACULAR NAMES. — Pa: pun-kamwi, sitru-kamwi • Wp: wila munuwi • Wn: mekuhuluimë, mulekju • Cr: bwa-pedri • Br: cafezinho.

HERBARIUM DATA (FG). — 137 collections at CAY. Sel. exs.: *F.M.R. Leprieur s.n.* (holotype of *Heisteria microcalyx*: P[P02441925]; iso-, L[L0038998], P[P02441926]).

INVENTORY DATA (FG). — 71 trees in 51 plots; $F_{\max} = 1.1\%$; $dbh_{\text{inv}} = 22.3$ cm.

[1372] *Heisteria ovata* Benth.
(Fig. 43D)

Hooker's J. Bot. Kew Gard. Misc. 3: 366 (Bentham 1851).

Heisteria subsessilis Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 367 (Bentham 1851).

Heisteria nitida Engl., *Fl. Bras. [Martius]* 12 (2): 18 [1 Dec. 1872] (Engler 1872).

Heisteria flexuosa Engl., *Fl. Bras. [Martius]* 12 (2): 21 [1 Dec. 1872] (Engler 1872).

Heisteria rubricalyx S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 337 ["1894-96" publ. Dec. 1895] (Moore 1895).

Heisteria micrantha Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 340 (Huber 1909).

Heisteria vageleri Burret, *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 50 [1 June 1924] (Burret 1924).

Heisteria krukovi A.C.Sm., *Brittonia* 2 (2): 147 (Smith 1936).

Heisteria surinamensis Amshoff, *Recueil Trav. Bot. Néerl.* 34: 497 (Amshoff 1937).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5086*.

INVENTORY DATA (FG). — 12 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.8$ cm.

Genus *Minquartia* Aubl.

[1373] *Minquartia guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 4 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate.

Secretaria loranthacea Müll.Arg., *Prodri. [A. P. de Candolle]* 15 (2.2): 228 [late Aug. 1866] (Müller 1866).

Endusa punctata Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 16: 313 (Radlkofer 1886). — *Minquartia punctata* (Radlk.) Sleumer, *Repert. Spec. Nov. Regni Veg.* 39: 282 (Sleumer 1936).

Eganthus poeppigii Tiegh., *J. Bot. [Morot]* 13: 77 (Tieghem 1899).

Minquartia macrophylla Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 33 [Sep. 1935] (Ducke 1935).

Minquartia parvifolia A.C.Sm., *Brittonia* 2 (2): 149 (Smith 1936).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: mekwa, yawu • Ka: alataweli, iwiyyu, wiyyu • Te: iwidju • Wp: wakal'i • Wn: kunawa • Nt: alata udu • Cr: mékwa • Fr: mincourt • Br: acariquara, acariúba.

HERBARIUM DATA (FG). — 71 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000028175]).

INVENTORY DATA (FG). — 373 trees in 143 plots; $F_{\max} = 4.4\%$; $dbh_{\text{inv}} = 112.4$ cm.

Genus *Ptychopetalum* Benth.

[1374] *Ptychopetalum olacoides* Benth.

London J. Bot. 2: 377 (Bentham 1843).

VERNACULAR NAMES. — Pa: aneku, aneku-kamwi • Te: wilapilátá • Wp: wila pilatá, wila tai • Cr: bwa-bandé • Fr: bois bandé • Br: marapuama, muira-puama.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *J. Martin s.n.* (type K[K000581314]).

INVENTORY DATA (FG). — 138 trees in 46 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 33.7$ cm.

Genus *Ximenia* Plum. ex L.

[1375] *Ximenia americana* L.

Sp. Pl. 2: 1193 [1 May 1753] (Linnaeus 1753). — *Ximenia spinosa* Salisb., *Prodr. Stirp. Chap. Allerton*: 276 [Nov.-Dec. 1796] (Salisbury 1796), nom. illeg. superfl. (based on *Ximenia americana*). — *Ximenia montana* Macfad., *Fl. Jamaica [Macfadyen]* 1: 121 (Macfadyen 1837), nom. illeg. superfl. (based on *Ximenia americana* and *X. multiflora*).

Amyris arborescens P.Browne, *Civ. Nat. Hist. Jamaica*: 209 (Browne 1756). — *Ximenia inermis* L., *Sp. Pl.*, ed. 2, 1: 497 [Sep. 1762] (Linnaeus 1762), nom. illeg. superfl. (based on *Amyris arborescens*).

Heymassoli inermis Aubl., *Hist. Pl. Guiane* 1: 325 [Jun.-Dec. 1775] (Aublet 1775).

Heymassoli spinosa Aubl., *Hist. Pl. Guiane* 1: 324 [Jun.-Dec. 1775] (Aublet 1775). — *Ximenia americana* var. *oblonga* DC., *Prodri. [A. P. de Candolle]* 1: 533 [mid Jan. 1824] (Candolle 1824), nom. illeg. superfl. (based on *Heymassoli spinosa*).

Pimecaria odorata Raf., *Alsogr. Amer.*: 65 (Rafinesque 1838).

Ximenia aculeata Crantz, *Inst. Rei Herb.* 2: 381 (Crantz 1766).

Ximenia americana f. *inermis* (Aubl.) Engl., *Fl. Bras. [Martius]* 12 (2): 9 [1 Dec. 1872] (Engler 1872).

Ximenia americana f. *inermis* S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 337 ["1894-96" publ. Dec. 1895] (Moore 1895), nom. illeg. hom., non (Aubl.) Engl. (Engler 1872).

Ximenia arborescens Tussac ex Walp., *Repert. Bot. Syst. [Walpers]* 1 (2): 377 [18-20 Sep. 1842] (Walpers 1842).

Ximenia elliptica G.Forst., *Fl. Ins. Austr.* 27 [Oct.-Nov. 1786] (Forster 1786). — *Ximenia verrucosa* M.Roem., *Fam. Nat. Syn.*

Monogr. 1 (Hesperides): 23 (Roemer 1846), *nom. illeg. superf.* (based on *Ximenia elliptica*).

Ximenia fluminensis M.Roem., *Fam. Nat. Syn. Monogr. 1 (Hesperides)*: 22 (Roemer 1846).

Ximenia laurina Delile, *Ann. Sci. Nat., Bot. sér. 2, 20*: 89 (Delile 1843).

Ximenia loranthifolia Span., *Linnaea 15*: 177 (Spanoghe 1841), *nom. nud. pro syn.*

Ximenia multiflora Jacq., *Enum. Syst. Pl.: 19* [Aug.-Sep. 1760] (Jacquin 1760). — *Ximenia americana* var. *ovata* DC., *Prodr. [A. P. de Candolle] 1*: 533 [mid Jan. 1824] (Candolle 1824), *nom. illeg. superf.* (based on *Ximenia multiflora*).

Ximenia oblonga Lam. ex Hemsl., *Biol. Cent.-Amer., Bot. 1 (3)*: 185 [Feb. 1879] (Hemsley 1879).

NOTES. — Restricted to coastal sandridges. *Ximenia americana* f. *inermis* S.Moore is not based on *Heymassoli inermis* Aubl.

VERNACULAR NAMES. — Ka: heymassoli (*fide* Aublet 1775) • Fr: citron de mer.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Heymassoli spinosa*: P-JU, P00671912]).

SIZE. — Up to 12 m tall (Hiepko 1993).

Family OLEACEAE Hoffmanns. & Link
Genus *Chionanthus* D.Royen

[1376] *Chionanthus guianensis* (Aubl.) Pers.

Syn. Pl. [Persoon] 1: 9 [1 Apr.-15 June 1805] (Persoon 1805). — *Mayepea guianensis* Aubl., *Hist. Pl. Guiane* 1: 81 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Chionanthus mayepea* Vahl, *Symb. Bot. [Vahl] 2*: 1 [Jul.-Dec. 1791] (Vahl 1791), *nom. illeg. superf.* (based on *Mayepea guianensis*). — *Chionanthus tetrandrus* Vahl, *Enum. Pl. [Vahl] 1*: 45 [Jul.-Dec. 1804] (Vahl 1804), “*tetrandra*”, *nom. illeg. superf.* (based on *Mayepea guianensis*).

Linociera tetrandra R.Br., *Prodr. Fl. Nov. Holland. 523* [27 Mar. 1810] (Brown 1810), *nom. inval.* (genus name and epithet not associated).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777522] designated by Lanjouw & Uittien [1940: 153]).

SIZE. — Aublet (1775): “Son tronc a [...] environ cinq pouces de diamètre”, i.e. c. 12.5 cm diameter.

Family OPILIACEAE Valeton
Genus *Agonandra* Miers ex Benth.

[1377] *Agonandra silvatica* Ducke
(Fig. 45A)

Arch. Jard. Bot. Rio de Janeiro 3: 41 (Ducke 1922).

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2341*.

INVENTORY DATA (FG). — 33 trees in 24 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43.9$ cm.

Family PENTAPHYLACACEAE Engl.
Genus *Ternstroemia* Mutis ex L.f.

[1378] *Ternstroemia delicatula* Choisy

Mém. Soc. Phys. Genève 14 (1): 106 (Choisy 1855).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material P[P00780885, P00780886, P00780887]).

INVENTORY DATA (FG). — 38 trees in 4 plots; $F_{\max} = 2.8\%$; $dbh_{\text{inv}} = 55.1$ cm.

[1379] *Ternstroemia dentata* (Aubl.) Sw.
(Fig. 45B)

Prodr. [Swartz] 81 [20 Jun.-29 July 1788] (Swartz 1788). — *Taonabo dentata* Aubl., *Hist. Pl. Guiane* 1: 569 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Ka: palipyali, pipi • Wp: walatiwā sili • Wn: maipuli apepiton.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000041973]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.3$ cm.

[1380] *Ternstroemia punctata* (Aubl.) Sw.

Prodr. [Swartz] 81 [20 Jun.-29 July 1788] (Swartz 1788). — *Taonabo punctata* Aubl., *Hist. Pl. Guiane* 1: 571 [Jun.-Dec. 1775] (Aublet 1775).

Ternstroemia revoluta Splitg., *Tijdschr. Nat. Geschied. Physiol. 9*: 99 [Aug.-Sep. 1842] (Splitgerber 1842).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000041974]).

SIZE. — Aublet (1775, 1: 571) wrote: “Cette espèce de palétuvier qui s’élève aussi haut que le précédent, et qui en a le port [...], i.e. similar in height and habit to the preceding. The preceding is *Taonabo dentata* Aubl., which is said to have a trunk “deux pieds de diamètre”, i.e. c. 60 cm diameter.

Family PERACEAE Klotzsch
Genus *Chaetocarpus* Thwaites

[1381] *Chaetocarpus schomburgkianus*
(Kuntze) Pax & K.Hoffm.

Pflanzenr. [Engler] IV.147.IV (Heft 52): 10 [13 Feb. 1912] (Pax & Hoffmann 1912). — *Gaedawakka schomburgkiana* Kuntze, *Revis. Gen. Pl. 2*: 606 [5 Nov. 1891] (Kuntze 1891).

Drypetes spruceana Müll.Arg., *Prod. [A. P. de Candolle]* 15 (2.2): 454 [late Aug. 1866] (Müller 1866).

Discocarpus mazarunensis Croizat, *Bull. Torrey Bot. Club* 75 (4): 400 [Jul.-Aug. 1948] (Croizat 1948).

Drypetes maguireana Monach., *Phytologia* 3 (1): 33 [30 Aug. 1948] (Monachino 1948).

Chaetocarpus williamsii Steyermark., *Fieldiana, Bot.* 28 (2): 306 (Steyermark 1952).

VERNACULAR NAMES. — Ka: kusali yepo, wiye kane • Nt: sipiki udu.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *D. Sabatier 2329*.

INVENTORY DATA (FG). — 386 trees in 123 plots; $F_{\max} = 1.9\%$; $\text{dbh}_{\text{inv}} = 76\text{ cm}$.

[1382] *Chaetocarpus* sp. A

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *R. Benoist 539* (P[P05479470]).

INVENTORY DATA (FG). — 80 trees in 29 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 82\text{ cm}$.

Genus *Pera* Mutis

[1383] *Pera bicolor* (Klotzsch) Müll.Arg.

Prod. [A. P. de Candolle] 15 (2.2): 1028 [late Aug. 1866] (Müller 1866). — *Peridium bicolor* Klotzsch, *London J. Bot.* 2: 44 (Klotzsch 1843).

Peridium schomburgkii Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1089 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), nom. nud.

Peridium schomburgkianum Klotzsch ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 6: 323 (Bentham 1854), in obs. — *Pera schomburgkiana* (Klotzsch ex Benth.) Müll.Arg., *Prod. [A. P. de Candolle]* 15 (2.2): 1027 [late Aug. 1866] (Müller 1866).

VERNACULAR NAMES. — Ka: boko.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Loubry 55*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 46.5\text{ cm}$.

[1384] *Pera glabrata* (Schott) Poepp. ex Baill.

Étude Euphorb.: 434 (Baillon 1858). — *Peridium glabratum* Schott, *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 410 [Jan.-June 1827] (Schott 1827).

Peridium ferrugineum Schott, *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 410 [Jan.-June 1827] (Schott 1827). — *Pera ferruginea* (Schott) Müll.Arg., *Prod. [A. P. de Candolle]* 15 (2.2): 1031 [late Aug. 1866] (Müller 1866).

Peridium oblongifolium Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 243 (Bentham 1850).

NOTE. — "Pera arborea Baill.", listed in IPNI and Tropicos, does not exist. Baillon (*Adansonia [Baillon]* 5: 224 [Baillon 1865]) misinterpreted *P. arborea* Mutis, to which he explicitly referred.

VERNACULAR NAMES. — Ka: pilityalapo, woko popi.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *M.-F. Prévost 649*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 38\text{ cm}$.

Genus *Pogonophora* Miers ex Benth.

[1385] *Pogonophora schomburgkiana* Miers ex Benth. (Fig. 45C)

Hooker's J. Bot. Kew Gard. Misc. 6: 373 (Bentham 1854).

Pogonophora schomburgkiana var. *longifolia* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 6: 373 (Bentham 1854).

Pogonophora schomburgkiana f. *elliptica* Pax, *Pflanzenr. [Engler]* IV.147.III (Heft 47): 108 [14 Feb. 1911] (Pax 1911).

Poraresia anomala Gleason, *Bull. Torrey Bot. Club* 58 (6): 385 [June 1931] (Gleason 1931).

VERNACULAR NAMES. — Ka: tapilen pipyo, tikilin wewe • Nt: geli udu • Br: amarelinho, miatoá.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *D. Sabatier 3556*.

INVENTORY DATA (FG). — 1054 trees in 100 plots; $F_{\max} = 8.8\%$; $\text{dbh}_{\text{inv}} = 38.2\text{ cm}$.

Family PHYLLANTHACEAE Martinov
Genus *Amanoa* Aubl.

[1386] *Amanoa congesta* W.J.Hayden

Brittonia 42 (4): 261 (Hayden 1990).

VERNACULAR NAMES. — Wp: wila ū.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *M.-F. Prévost 3768*.

INVENTORY DATA (FG). — 16 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 51.7\text{ cm}$.

[1387] *Amanoa grandiflora* (Müll.Arg.) Müll.Arg. (Fig. 46A)

Flora 55: 2 (Müller 1872). — *Amanoa guianensis* var. *grandiflora* Müll.Arg., *Prod. [A. P. de Candolle]* 15 (2.2): 910 [late Aug. 1866] (Müller 1866).

Amanoa potamophila Croizat, *Amer. Midl. Naturalist* 29 (2): 475 [5 Apr. 1943] (Croizat 1943).

Amanoa macrocarpa Cuatrec., *Brittonia* 11: 164 (Cuatrecasas 1959).

NOTES.— Hayden (1990) showed that the Aublet collections, which could have served as the type of *Amanoa guianensis* Aubl., actually belong to two distinct species. One is known only from a few specimens from French Guiana, Amapá and Suriname; the other is widespread from Central America to northern South America (Hayden 1990). He therefore selected among these specimens a lectotype (W[[W-1889-0012576](#)]; isolectotypes G[not seen], F[[V0055866F](#)], A[not seen], MO[not seen]) which corresponds to the current usage of Aublet's name, that is to the widespread species, and described a new species, *Amanoa neglecta* W.J.Hayden, based on the second entity found in Aublet collections (holo-, P-JJR[[P00777979](#)], iso-, BM[[BM000645699](#)]) (Hayden 1990). This was a laudable choice to preserve the current usage of the name *A. guianensis*, but unfortunately Hayden (1990) overlooked the previous lectotypification of *A. guianensis* Aubl. by Lanjouw & Uittien (1940: 148), who designated the P-JJR specimen. Given the anteriority of this lectotypification (Delprete 2015), there is no choice but to change the name of the most widespread taxon, and to adopt the oldest available name, *Amanoa grandiflora* (Müll.Arg.) Müll.Arg.

VERNACULAR NAMES.— Ka: kuata mopeli • Wp: wila tī.

HERBARIUM DATA (FG).— 35 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (W[[W-1889-0012576](#)], F[[V0055866F](#)]).

INVENTORY DATA (FG).— 31 trees in 20 plots; $F_{\max} < 1\%$; $dbh_{inv} = 66.8$ cm.

[1388] *Amanoa guianensis* Aubl.

Hist. Pl. Guiane 1: 256 [Jun.-Dec. 1775] (Aublet 1775).— *Amanoa neglecta* W.J.Hayden, *Brittonia* 42 (4): 267 (Hayden 1990), nom. illeg. superfl. (based on the lectotype of *A. guianensis*).

NOTES.— Known only from the Guiana Shield. As stated above [see notes under *Amanoa grandiflora* (Müll.Arg.) Müll.Arg.], the lectotypification of *A. guianensis* Aubl. by Lanjouw & Uittien (1940: 148) predates that by Hayden (1990). As a consequence, *A. neglecta* W.J.Hayden, which is based on the lectotype of *A. guianensis*, is an illegitimate superfluous name.

HERBARIUM DATA (FG).— 2 collections known from FG, none at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[[P00777979](#)] designated by Lanjouw & Uittien [1940: 148]).

SIZE.— Brazil, Amapá. *N.A. Rosa & M.R. dos Santos 1834* (MO), 20 m × 48 cm (“1,50 m de circunferencia”).

Genus *Hieronyma* Allemão

[1389] *Hieronyma alchorneoides* Allemão

Hieronima alchorneoides [shortly after 30 Apr. 1848] (Allemão 1848), “*Hyeronima*”.

Stilaginella amazonica Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 241 (Tulasne 1851).

Stilaginella laxiflora Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 244 (Tulasne 1851).— *Hieronyma laxiflora* (Tul.) Müll.Arg., *Linnaea* 34 (1): 67 [pre-Mar. 1865] (Müller 1865).

Stilaginella ferruginea Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 250 (Tulasne 1851).— *Hieronyma ferruginea* (Tul.) Tul., *Fl. Bras. [Martius]* 4 (1): 334 [15 Feb. 1861] (Tulasne 1861).

Hieronyma mollis Müll.Arg., *Prod. [A. P. de Candolle]* 15 (2.2): 269 [late Aug. 1866] (Müller 1866).

Hieronyma caribaea Urb., *Repert. Spec. Nov. Regni Veg.* 16: 139 [15 Nov. 1919] (Urban 1919).

Hieronyma heterotricha Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XV (Heft 81): 39 [19 Sep. 1922] (Pax & Hoffmann 1922).

Hieronyma mattogrossensis Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XV (Heft 81): 39 [19 Sep. 1922] (Pax & Hoffmann 1922).

Hieronyma choocoensis Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 7: 52 [Dec. 1946] (Cuatrecasas 1946).

Hieronyma tectissima Standl. & L.O.Williams, *Rain Forests Golfo Dulce*: 222 (Standley & Williams 1956), nom. nud.

Hieronyma ovatifolia Lundell, *Wrightia* 4 (4): 134 (Lundell 1970).

Hieronyma alchorneoides var. *stipulosa* P.Franco, *Bot. Jahrb. Syst.* 111 (3): 321 [27 Apr. 1990] (Franco 1990).

VERNACULAR NAMES.— Pa: waikwavia • Ka: ako, kasili wewe, tapilen katulimya, tokolo enulu • Wp: minua'i • Wn: okomë öki, palulumili • Nt: win udu • Cr: bwa-diven • Br: aricurqua, muiragonçalo, urucurana.

HERBARIUM DATA (FG).— 42 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2703*.

INVENTORY DATA (FG).— 21 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{inv} = 63.8$ cm.

[1390] *Hieronyma oblonga* (Tul.) Müll.Arg.

Linnaea 34 (1): 66 [pre-Mar. 1865] (Müller 1865).— *Stilaginella oblonga* Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 248 (Tulasne 1851).

Stilaginella benthamii Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 247 (Tulasne 1851).— *Hieronyma oblonga* var. *benthamii* (Tul.) Müll.Arg., *Linnaea* 34 (1): 66 [pre-Mar. 1865] (Müller 1865).

Stilaginella blanchetiana Tul., *Ann. Sci. Nat., Bot. sér. 3, 15:* 249 (Tulasne 1851).— *Hieronyma blanchetiana* (Tul.) Tul., *Fl. Bras. [Martius]* 4 (1): 333 [15 Feb. 1861] (Tulasne 1861).— *Hieronyma oblonga* var. *blanchetiana* (Tul.) Müll.Arg., *Linnaea* 34 (1): 66 [pre-Mar. 1865] (Müller 1865).

Hieronyma oblonga var. *obtusa* Müll.Arg., *Linnaea* 34 (1): 66 [pre-Mar. 1865] (Müller 1865).

Hieronyma guatemalensis Donn.Sm., *Bot. Gaz.* 54 (3): 241 [21 Sep. 1912] (Donnell Smith 1912).

Hieronyma andina Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XV (Heft 81): 37 [19 Sep. 1922] (Pax & Hoffmann 1922).

Hieronyma poasana Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 18 (2):* 611 [20 Oct. 1937] (Standley 1937).

Hieronyma oblonga var. *crassifolia* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 300 [Dec. 1951] (Cuatrecasas 1951).

Hieronyma oblonga var. *nervata* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 301 [Dec. 1951] (Cuatrecasas 1951).

Hieronyma oblonga f. *glabra* Steyermark., *Fieldiana, Bot.* 28 (4): 951 (Steyermark 1957).

Hieronyma gentlei Lundell, *Wrightia* 5 (7): 248 (Lundell 1976).

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *B. Riéra & D. Sabatier 1925*.

INVENTORY DATA (FG). — 32 trees in 22 plots; $F_{\max} < 1\%$; $dbh_{inv} = 169.7$ cm.

Genus *Margaritaria* L.f.

[1391] *Margaritaria nobilis* L.f.
(Fig. 46B)

Suppl. Pl.: 428 ["1781" publ. Apr. 1782] (Linnaeus 1782). — *Phyllanthus nobilis* (L.f.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 414 [late Aug. 1866] (Müller 1866).

Cicca antillana A.Juss., *Euphorb. Gen.* 108 [21 Feb. 1824] (Jussieu 1824). — *Phyllanthus antillanus* (A.Juss.) Müll.Arg., *Linnaea* 32: 51 (Müller 1863). — *Phyllanthus nobilis* var. *antillanus* (A.Juss.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 415 [late Aug. 1866] (Müller 1866). — *Diasperus antillanus* (A.Juss.) Kuntze, *Revis. Gen. Pl.* 2: 597 [5 Nov. 1891] (Kuntze 1891). — *Margaritaria nobilis* var. *antillana* (A.Juss.) Stehlé & Quentin, *Fl. Guadeloupe* [Stehlé] 2 (1): 47 [May 1938] (Stehlé & Quentin 1938).

Cicca surinamensis Miq., *Linnaea* 21: 479 (Miquel 1848).

Cicca pavoniana Baill., *Étude Euphorb.* 618 (Baillon 1858). — *Phyllanthus nobilis* var. *pavonianus* (Baill.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 415 [late Aug. 1866] (Müller 1866).

Margaritaria adelioides Rich. ex Baill., *Étude Euphorb.* 618 (Baillon 1858), nom. nud.

Cicca antillana var. *pedicularis* Griseb., *Pl. Wright. [Grisebach]* 1: 158 [Dec. 1860] (Grisebach 1860).

Phyllanthus antillanus var. *pedicularis* Müll.Arg., *Linnaea* 32: 51 (Müller 1863).

Phyllanthus nobilis var. *guyanensis* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 414 [late Aug. 1866] (Müller 1866).

Phyllanthus nobilis var. *peruvianus* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 414 [late Aug. 1866] (Müller 1866).

Phyllanthus nobilis var. *brasiliensis* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 415 [late Aug. 1866] (Müller 1866).

Phyllanthus nobilis var. *riedelianus* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 415 [late Aug. 1866] (Müller 1866).

Phyllanthus nobilis var. *martii* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 70 [1 Feb. 1873] (Müller 1873).

Phyllanthus nobilis var. *panamensis* Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 70 [1 Feb. 1873] (Müller 1873).

Phyllanthus nobilis var. *hypomalacus* Standl., *Publ. Carnegie Inst. Wash.* 46: 68 (Standley 1935). — *Phyllanthus antillanus* var. *hypomalacus* (Standl.) Lundell, *Phytologia* 1 (10): 337 [27 Nov. 1939] (Lundell 1939). — *Margaritaria nobilis* var. *hypomalaca* (Standl.) Dugand, *Phytologia* 13 (6): 387 [19 Sep. 1966] (Dugand 1966).

VERNACULAR NAMES. — Br: cabelo-de-cotia, fruto-de-jacamin.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 4951*.

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{inv} = 37.5$ cm.

Genus *Phyllanthus* L.

[1392] *Phyllanthus attenuatus* Miq.

Linnaea 21: 479 (Miquel 1848).

Meborea guianensis Aubl., *Hist. Pl. Guiane* 2: 825 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate. — *Phyllanthus guianensis* (Aubl.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 376 [late Aug. 1866] (Müller 1866), nom. illeg. hom., non Klotzsch (1843). — *Diasperus guianensis* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 599 [5 Nov. 1891] (Kuntze 1891), "guyanensis".

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Larpin 670*.

SIZE. — Up to 12 m tall (Webster 2004).

[1393] *Phyllanthus juglandifolius* Willd.

Enum. Pl. Suppl. [Willdenow]: 64 [Jul.-Dec. 1814] (Willdenow 1814).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman B-3261*.

SIZE. — Brazil, Amazonas. *G.T. Prance 10376* (MO), 15 m × 20 cm.

Genus *Richeria* Vahl

[1394] *Richeria grandis* Vahl

Eclog. Amer. 1: 29 (Vahl 1797).

Amanoa divaricata Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 22 [15-21 Aug. 1841] (Poeppig 1841). — *Richeria grandis* var. *divaricata* (Poepp.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 467 [late Aug. 1866] (Müller 1866).

Amanoa racemosa Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 23 [15-21 Aug. 1841] (Poeppig 1841). — *Richeria grandis* var. *racemosa* (Poepp.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 467 [late Aug. 1866] (Müller 1866). — *Richeria racemosa* (Poepp.) Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XV (Heft 81): 28 [19 Sep. 1922] (Pax & Hoffmann 1922).

Guarania ramiflora Wedd. ex Baill., *Étude Euphorb.*: 598 (Baillon 1858), nom. nud.

Guarania laurifolia Baill., *Adansonia [Baillon]* 5: 348 (Baillon 1865). — *Richeria laurifolia* (Baill.) Baill., *Adansonia [Baillon]* 6: 16 (Baillon 1865). — *Richeria grandis* var. *laurifolia* (Baill.) Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 468 [late Aug. 1866] (Müller 1866).

Richeria grandis var. *genuina* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 468 [late Aug. 1866] (Müller 1866), nom. inval. (Turland et al. 2018: Art. 24.3).

Richeria grandis var. *obovata* Müll.Arg., *Prodr. [A. P. de Candolle]* 15 (2.2): 468 [late Aug. 1866] (Müller 1866). — *Richeria obovata* (Müll.Arg.) Pax & K.Hoffm., *Pflanzenr. [Engler]* IV.147.XV (Heft 81): 29 [19 Sep. 1922] (Pax & Hoffmann 1922).

Richeria australis Müll.Arg., *Fl. Bras. [Martius]* 11 (2): 17 [1 Feb. 1873] (Müller 1873).




FIG. 45. — Opiliaceae: **A**, *Agonandra silvatica* Ducke (D. Sabatier & M.-F. Prévost 3065). Pentaphylacaceae: **B**, *Ternstroemia dentata* (Aubl.) Sw. Peraceae: **C**, *Polygonophora schomburgkiana* Miers ex Benth. (D. Sabatier & J.-F. Molino 4841). Piperaceae: **D**, *Piper aduncum* L. (M.-F. Prévost 4081). A-C, © D. Sabatier/IRD; D, © M.-F. Prévost/IRD.

Richeria submembranacea Steyermark., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (5)*: 419 [27 May 1938] (Steyermark 1938).

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost 4262*.

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; dbh_{inv} = 63 cm.

Family PICRAMNIACEAE Fernando & Quinn
Genus *Picramnia* Sw.

[1395] *Picramnia guianensis* (Aubl.) Jans.-Jac.

Fl. Suriname 5 (1): 329 (Jansen-Jacobs 1979). — *Tariri guianensis* Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 37 [Jun.-Dec. 1775] (Aublet 1775), “Guyannensis” on plate. — *Picramnia tariri* DC., *Prodr. [A. P. de Candolle]* 2: 66 [mid Nov. 1825] (Candolle 1825), nom. illeg. superfl. (based on *Tariri guianensis*).

VERNACULAR NAMES. — Pa: tarii, tarit • Ka: tariri (*fide* Aublet 1775) • Wp: paipayo namipay • Wn: talili • Br: caferana, tariri.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JU, **P00678733**, here designated; isolectotypes *fide* Howard [1983: 287]: BM, P[not seen]).

SIZE. — Brazil, Pará. *N.T. Silva* 2389 (MO), 10 cm.

[1396] *Picramnia latifolia* Tul.

Ann. Sci. Nat., Bot. sér. 3, 7: 258 (Tulasne 1847).

Picramnia longissima Tul., *Ann. Sci. Nat., Bot. sér. 3, 7*: 257 (Tulasne 1847).

Picramnia umbrosa Seem., *Bot. Voy. Herald [Seemann]* 3: 97 [Nov. 1853] (Seemann 1853).

Picramnia macrostachys Klotzsch ex Engl., *Fl. Bras. [Martius]* 12 (2): 238 [1 Sep. 1874] (Engler 1874).

Picramnia eosina J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.2)*: 697 (Macbride 1949).

Picramnia cooperi D.M.Porter, *J. Arnold Arbor.* 54 (2): 317 [26 Apr. 1973] (Porter 1973).

VERNACULAR NAMES. — Wp: paipayo namipay • Br: caferana, tariri.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-4007*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 10 \text{ cm}$.

[1397] *Picramnia spruceana* Engl.

Fl. Bras. [Martius] 12 (2): 238 [1 Sep. 1874] (Engler 1874). — *Picramnia sellowii* Planch. subsp. *spruceana* (Engl.) Pirani, *Bol. Bot. Univ. São Paulo* 12: 132 ["1990" publ. 1991] (Pirani 1991).

Picramnia tenuis J.F.Macbr., *Candollea* 5: 375 (Macbride 1934).

Picramnia krukovi A.C.Sm., *Brittonia* 2 (2): 153 (Smith 1936).

VERNACULAR NAMES. — Wp: paipayo namipay • Br: caferana, tariri.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *P. Grenand 543*.

SIZE. — Up to 20 m tall (Pirani 2005).

Family PIPERACEAE Giseke
Genus *Piper* L.

[1398] *Piper aduncum* L.
(Fig. 45D)

Sp. Pl. 1: 29 [1 May 1753] (Linnaeus 1753). — *Artanthe adunca* (L.) Miq., *Comm. Phytogr.* 14 (2): 49 [16-21 Mar. 1840] (Miquel 1840). — *Steffensia adunca* (L.) Kunth, *Linnaea* 13: 633 ["1839" publ. Mar.-June 1840] (Kunth 1840).

Piper angustifolium Ruiz & Pav., *Flora Peruviana* 1: 38 (Ruiz & Pavón 1798), nom. illeg. hom., non Lam. (Lamarck 1791) [synonym of *P. consanguineum* (Kunth) Steud.]

Piper elongatum Vahl, *Enum. Pl. [Vahl]* 1: 312 [Jul.-Dec. 1804] (Vahl 1804). — *Steffensia elongata* (Vahl) Kunth, *Linnaea* 13: 637 ["1839" publ. Mar.-June 1840] (Kunth 1840). — *Artanthe elongata* (Vahl) Miq., *Syst. Piperac. [F.A.W. Miquel]*: 434 ["1843" publ. 4 Apr. 1844] (Miquel 1844).

Piper celtidifolium Kunth, *Nova genera et species plantarum [H.B.K.] 1:* 50 [29 Jan. 1816] (Kunth 1816). — *Steffensia celtidifolia* (Kunth) Kunth, *Linnaea* 13: 635 ["1839" publ. Mar.-June 1840] (Kunth 1840).

Piper purpurascens D.Dietr., *Syn. Pl. [D. Dietrich]* 1: 119 [July 1839] (Dietrich 1839), nom. illeg. hom., non Desf. (Desfontaines 1829).

Piper multinervium M.Martens & Galeotti, *Bull. Acad. Roy. Sci. Bruxelles* 10: 130 (Martens & Galeotti 1843).

Artanthe galleottii Miq., *Syst. Piperac. [F.A.W. Miquel]*: 451 ["1843" publ. 4 Apr. 1844] (Miquel 1844).

Artanthe adunca f. *angustifolia* Miq., *London J. Bot.* 4: 456 (Miquel 1845). — *Artanthe celtidifolia* (Kunth) Miq., *Syst. Piperac. [F.A.W. Miquel]*: 452 ["1843" publ. 4 Apr. 1844] (Miquel 1844).

Piper angustifolium var. *cordulatum* C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 286 [mid Nov. 1869] (Candolle 1869). — *Piper elongatum* var. *cordulatum* (C.DC.) C.DC., *Repert. Spec. Nov. Regni Veg.* 9: 230 (Candolle 1911). — *Piper aduncum* var. *cordulatum* (C.DC.) Yunck., *Lilloa* 27: 129 (Yuncker 1955).

Piper pseudovelutinum C.DC. var. *flavescens* C.DC., *Bull. Soc. Roy. Bot. Belgique* 30 (1): 203 ["1891" publ. 25 May 1892] (Candolle 1892), "pseudo-velutinum". — *Piper flavescens* (C.DC.) Trel., *Contr. U.S. Natl. Herb.* 26 (4): 184 [31 Dec. 1929] (Trelease 1929).

Piper kuntzei C.DC. in Kuntze, *Revis. Gen. Pl.* 3 (3): 274 [28 Sep. 1898] (Candolle 1898).

Piper guanaianum C.DC., *Repert. Spec. Nov. Regni Veg.* 9: 232 (Candolle 1911).

Piper herzogii C.DC., *Meded. Rijks-Herb.* 27: 7 (Candolle 1915).

Piper fatoanum C.DC., *Smithsonian Misc. Collect.* 71 (6): 7 [12 Feb. 1920] (Candolle 1920).

Piper aduncum var. *laevisolum* C.DC., *Smithsonian Misc. Collect.* 71 (6): 8 [12 Feb. 1920] (Candolle 1920). — *Piper elongatum* var. *laevisolum* (C.DC.) Trel., *Contr. U.S. Natl. Herb.* 26 (2): 37 [21 May 1927] (Trelese 1927).

Piper cardenasii Trel., *Mem. New York Bot. Gard.* 7: 223 (Trelese 1927).

Piper elongatum var. *brachyarthrum* Trel., *Contr. U.S. Natl. Herb.* 26 (2): 37 [21 May 1927] (Trelese 1927). — *Piper aduncum* var. *brachyarthrum* (Trel.) Yunck., *Ann. Missouri Bot. Gard.* 37: 32 [31 Mar. 1950] (Yuncker 1950).

Piper nonconformans Trel., *J. Wash. Acad. Sci.* 19: 334 (Trelese 1929).

Piper disparispicum Trel., *Contr. U.S. Natl. Herb.* 26 (4): 170 [31 Dec. 1929] (Trelese 1929).

Piper aduncifolium Trel., *Contr. U.S. Natl. Herb.* 26 (4): 171 [31 Dec. 1929] (Trelese 1929).

Piper anguillispicum Trel., *Contr. U.S. Natl. Herb.* 26 (4): 175 [31 Dec. 1929] (Trelese 1929), "anguillaespicum".

Piper ob lanceolatum Trel. var. *fragilicaule* Trel., *Contr. U.S. Natl. Herb.* 26 (4): 175 [31 Dec. 1929] (Trelese 1929).

Piper submolle Trel., *Contr. U.S. Natl. Herb.* 26 (4): 178 [31 Dec. 1929] (Trelese 1929).

Piper multinervium Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 10: 160 (Trelese 1931), nom. illeg. hom., non M.Martens & Galeotti (1843).

Piper stevensonii Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 12: 104 (Trelese 1936), nom. inval. (anglice).

Piper elongatifolium Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.1): 161 [10 June 1936] (Trelese 1936).

Piper elongatum var. *pampayacusum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.1): 162 [10 June 1936] (Trelese 1936).

Piper lineatum Ruiz & Pav. var. *hirtipetiolatum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.1): 184 [10 June 1936] (Trelese 1936).




FIG. 46. — Phyllanthaceae: **A**, *Amanoa grandiflora* (Müll.Arg.) Müll.Arg. (*D. Sabatier & M.-F. Prévost 3007*); **B**, *Margaritaria nobilis* L.f. (*D. Sabatier & J.-F. Molino 4951*). Polygonaceae: **C**, *Coccoloba mollis* Casar. (*M.-F. Prévost & D. Sabatier 3010*). Primulaceae: **D**, *Cybianthus potiae* (Mez) G.Agostini (*D. Sabatier & J.-F. Molino 5545*). © D. Sabatier/IRD.

Piper cuatrecasasii Trel., *Trab. Mus. Nac. Ci. Nat., Ser. Bot.* 33: 48 [30 July 1936] (Trelease 1936), “Cuatrecasasi”, nom. inval. (anglice).

Piper cumbricola Trel., *Trab. Mus. Nac. Ci. Nat., Ser. Bot.* 33: 48 [30 July 1936] (Trelease 1936), nom. inval. (anglice).

Piper illudens Trel., *Trab. Mus. Nac. Ci. Nat., Ser. Bot.* 33: 50 [30 July 1936] (Trelease 1936), nom. inval. (anglice).

Piper multinervium var. *amplum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 347 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *cayoense* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 347 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *flavicans* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 348 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *hirsuticaule* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 348 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *kantelolense* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 348 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *paralense* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 348 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *peracutum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 349 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *productipes* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 349 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *protractifolium* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 349 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *puberulipedunculum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 349 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *pubescenticaule* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 350 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *skutchii* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 350 [31 Jan. 1938] (Trelease 1938), “Skutchii”.

Piper multinervium var. *tamashense* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 350 [31 Jan. 1938] (Trelease 1938).

Piper multinervium var. *telanum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (4): 350 [31 Jan. 1938] (Trelease 1938).

Piper intersitum Trel., *Caldasia* 1 (1): 86 (Trelease 1940).

Piper intersitum f. *porcecitense* Trel., *Caldasia* 1 (1): 87 (Trelease 1940).

Piper reciprocum Trel., Ciencia (Mexico) 2 (5): 207 [25 May 1941] (Trelease 1941).

VERNACULAR NAMES. — Wn: pëtpëjot • Nt: gaanman udu ana • Br: pimenta-longa.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *M.-F. Prévost 3567*.

SIZE. — Up to 10 m tall (Steyermark & Callejas-Posada 2003).

[1399] *Piper cernuum* Vell.

Fl. Flumin. 25 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829). — *Artanthe cernua* (Vell.) C.Presl, *Abh. Königl. Böhm. Ges. Wiss.*, ser. 5, 6: 585 (Presl 1851).

Steffensia eximia Kunth, *Linnaea* 13: 665 ["1839" publ. Mar.-June 1840] (Kunth 1840). — *Piper eximium* Kunth, *Linnaea* 13: 665 ["1839" publ. Mar.-June 1840] (Kunth 1840), nom. nud. pro syn. — *Artanthe eximia* (Kunth) Miq., *Syst. Piperac.* [F.A.W. Miquel]: 393 ["1843" publ. 4 Apr. 1844] (Miquel 1844). — *Piper obliquum* Ruiz & Pav. var. *eximium* (Kunth) C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 307 [mid Nov. 1869] (Candolle 1869).

Steffensia cocoloboides Kunth, *Linnaea* 13: 666 ["1839" publ. Mar.-June 1840] (Kunth 1840). — *Piper cocoloboides* Kunth, *Linnaea* 13: 666 ["1839" publ. Mar.-June 1840] (Kunth 1840), nom. nud. pro syn.

Steffensia richardiifolia Kunth, *Linnaea* 13: 668 ["1839" publ. Mar.-June 1840] (Kunth 1840), "richardiaeifolia". — *Piper richardiifolium* Kunth, *Linnaea* 13: 668 ["1839" publ. Mar.-June 1840] (Kunth 1840), "richardiaeifolium", nom. nud. pro syn. — *Artanthe richardiifolia* (Kunth) Miq., *Syst. Piperac.* [F.A.W. Miquel] 395 ["1843" publ. 4 Apr. 1844] (Miquel 1844), "richardiaeifolia".

Artanthe spectabilis Miq., *Linnaea* 20: 138 [17-18 May 1847] (Miquel 1847).

Artanthe spectabilis var. *glabrior* Miq., *Linnaea* 20: 139 [17-18 May 1847] (Miquel 1847).

Piper boliviannum C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 280 [mid Nov. 1869] (Candolle 1869).

Piper ovalifolium C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 305 [mid Nov. 1869] (Candolle 1869).

Piper cabellense C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 306 [mid Nov. 1869] (Candolle 1869).

Piper gigantifolium C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 306 [mid Nov. 1869] (Candolle 1869).

Piper araguense Trel. & Yunck., *Piperac. N. South Amer.* 1: 138 (Trelease & Yuncker 1950). — *Piper cernuum* var. *araguense* (Trel. & Yunck.) Steyermark., *Fl. Venez.* 2 (2): 372 (Steyermark 1984).

Piper arbelaezii Trel. & Yunck. var. *subglabrirameum* Yunck., *Fidiana, Bot.* 28 (1): 204 (Yuncker 1951).

Piper perlongispicum Yunck., *Bol. Soc. Venez. Ci. Nat.* 23 (101): 89 (Yuncker 1962). — *Piper cernuum* var. *perlongispicum* (Yunck.) Steyermark., *Fl. Venez.* 2 (2): 372 (Steyermark 1984).

Piper cernuum var. *glabricaule* Yunck., *Bol. Inst. Bot. (São Paulo)* 3: 20 [23 Dec. 1966] (Yuncker 1966).

Piper cernuum var. *biformipilum* Yunck., *Bol. Inst. Bot. (São Paulo)* 3: 21 [23 Dec. 1966] (Yuncker 1966).

NOTES. — Kunth published simultaneously *Piper cocoloboides* and *Steffensia cocoloboides* (Kunth 1840: 666) as well as *P. eximium* and *S. eximia* (Kunth 1840: 665), and *P. richardiifolium* and *S. richardiifolia* (Kunth 1840: 668). As these names were published before 1953, all of them are valid; hence both *Artanthe eximia* (Kunth) Miq. and *A. richardiifolia* (Kunth) Miq. have alternative basionyms, respectively, *P. eximium* Kunth and *S. eximia* Kunth, and *P. richardiifolium* Kunth and *S. richardiifolia* Kunth.

VERNACULAR NAMES. — Wp: taīwī lenipi'ā, yali taku'ā sili.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville B-5436*.

SIZE. — Up to 12 m tall (Steyermark & Callejas-Posada 2003).

[1400] *Piper reticulatum* L.

Sp. Pl. 1: 29 [1 May 1753] (Linnaeus 1753). — *Enckea reticulata* (L.) Miq., *Syst. Piperac.* [F.A.W. Miquel]: 365 ["1843" publ. 4 Apr. 1844] (Miquel 1844). — *Discipiper reticulatum* (L.) Trel. & Stehlé, *Candollea* 10: 283 (Trelease & Stehlé 1946).

Pipersmilacifolium Kunth, *Nova genera et species plantarum [H.B.K.]* 1: 56 [29 Jan. 1816] (Kunth 1816). — *Enckea smilacifolia* (Kunth) Kunth, *Linnaea* 13: 605 ["1839" publ. Mar.-June 1840] (Kunth 1840), "Enckia".

Piper latum Kunth, *Nova genera et species plantarum [H.B.K.]* 1: 57 [29 Jan. 1816] (Kunth 1816). — *Enckea lata* (Kunth) Kunth, *Linnaea* 13: 606 ["1839" publ. Mar.-June 1840] (Kunth 1840), "Enckia". — *Macropiper latum* (Kunth) C.Presl, *Abh. Königl. Böhm. Ges. Wiss.*, ser. 5, 6: 583 (Presl 1851).

Artanthe ruiziana Miq., *London J. Bot.* 4: 444 (Miquel 1845).

Piper tarapotianum C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 246 [mid Nov. 1869] (Candolle 1869).

Piper duchassaingii C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 251 [mid Nov. 1869] (Candolle 1869).

Piper pangoense C.DC., *Prodr. [A. P. de Candolle]* 16 (1): 368 [mid Nov. 1869] (Candolle 1869).

Piper discophorum C.DC., *Bull. Soc. Roy. Bot. Belgique* 30 (1): 201 ["1891" publ. 25 May 1892] (Candolle 1892).

Piper reticulatum var. *santarosanum* Trel., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.1): 223 [10 June 1936] (Trelease 1936), "santa-rosanum".

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *C. Feuillet 616*.

SIZE. — Up to 15 m tall (Görts van Rijn 2007).

Family POLYGONACEAE Juss.
Genus *Coccoloba* P.Browne

[1401] *Coccoloba latifolia* Poir.

Encycl. [J. Lamarck et al.] 6 (1): 61 [2 Oct. 1804] (Poiret 1804). — *Uvifera latifolia* (Poir.) Kuntze, *Revis. Gen. Pl.* 2: 561 [5 Nov. 1891] (Kuntze 1891).

Coccoloba rheifolia Desf., *Tabl. École Bot.*, ed. 2, 46 (Desfontaines 1815).

Coccoloba grandis Benth., *London J. Bot.* 4: 624 (Bentham 1845).

VERNACULAR NAMES. — Ka: matula, patula • Br: cabeça-de-macaco.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost 1579*.

SIZE. — Up to 15 m tall (Aymard C. & Howard 2004).

[1402] *Coccoloba mollis* Casar.
(Fig. 46C)

Nov. Stirp. Bras. 8: 72 [June 1844] (Casaretto 1844). — *Coccoloba polystachya* var. *mollis* (Casar.) Meisn., *Prodr. [A. P. de Candolle]* 14 (1): 151 [mid Oct. 1856] (Meisner 1856).

Coccoloba polystachya Wedd., *Ann. Sci. Nat., Bot. sér. 3, 13:* 261 (Weddell 1849). — *Uvifera polystachya* (Wedd.) Kuntze, *Revis. Gen. Pl.* 2: 561 [5 Nov. 1891] (Kuntze 1891).

Coccoloba paniculata Meisn., *Fl. Bras. [Martius]* 5 (1): 43 [1 Jan. 1855] (Meisner 1855).

Coccoloba polystachya var. *glabra* Lindau, *Bot. Jahrb. Syst.* 13 (2): 133 [28 Oct. 1890] (Lindau 1890).

Coccoloba polystachya var. *pubescens* Lindau, *Bot. Jahrb. Syst.* 13 (2): 133 [28 Oct. 1890] (Lindau 1890).

Coccoloba polystachya var. *jamaicensis* Fawc. & Rendle, *J. Bot.* 51: 125 (Fawcett & Rendle 1913).

Coccoloba dugandiana A.Fernández, *Mutisia* 5: 1 (Fernández 1952).

Coccoloba standleyana P.H.Allen, *Rain Forests Golfo Dulce* 409 (Allen 1956).

VERNACULAR NAMES. — Pa: tivu-arigni • Ka: matula, patula • Nt: doon udu, doonudu.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3010*.

INVENTORY DATA (FG). — 53 trees in 30 plots; $F_{\max} < 1\%$; dbh_{inv} = 38.4 cm.

[1403] *Coccoloba uvifera* (L.) L.

Syst. Nat., ed. 10, 2: 1007 [7 June 1759] (Linnaeus 1759). — *Polygonum uvifera* L., *Sp. Pl.* 1: 365 [1 May 1753] (Linnaeus 1753). — *Guaiabara uvifera* (L.) House, *Amer. Midl. Naturalist* 8: 64 (House 1922).

NOTE. — Restricted to coastal sandy beaches.

VERNACULAR NAMES. — Cr: rézen-bòdlanmé.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *H. de Foresta 760*.

SIZE. — Panama, Colón, 17 May 1974. *M.H. Nee 11694* (MO), 7 m × 25 cm.

Genus *Ruprechtia* C.A.Mey.

[1404] *Ruprechtia brachysepala* Meisn.

Fl. Bras. [Martius] 5 (1): 57 [1 Jan. 1855] (Meisner 1855). — *Magonia brachysepala* (Meisn.) Kuntze, *Revis. Gen. Pl.* 2: 553 [5 Nov. 1891] (Kuntze 1891).

Ruprechtia amentacea Meisn., *Fl. Bras. [Martius]* 5 (1): 56 [1 Jan. 1855] (Meisner 1855). — *Magonia amentacea* (Meisn.) Kuntze, *Revis. Gen. Pl.* 2: 553 [5 Nov. 1891] (Kuntze 1891).

Ruprechtia latifolia Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 345 (Huber 1909).

Ruprechtia marowynensis Eyma, *Recueil Trav. Bot. Néerl.* 32: 224 (Eyma 1935).

Ruprechtia nitida Brandbyge, *Nordic J. Bot.* 10 (2): 158 (Brandbyge 1990).

VERNACULAR NAMES. — Nt: pindya.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3358*.

SIZE. — Up to 12 m tall (Pendry 2004).

Genus *Triplaris* Loefl.

[1405] *Triplaris americana* L.

Syst. Nat., ed. 10, 2: 881 [7 June 1759] (Linnaeus 1759), “*americ.*”.

Triplaris pyramidalis Jacq., *Select. Stirp. Amer. Hist.* 13 [5 Jan. 1763] (Jacquin 1763).

Triplaris brasiliiana Cham., *Linnaea* 8: 139 (Chamisso 1833).

Triplaris schomburgkiana Benth., *London J. Bot.* 4: 628 (Bentham 1845).

Triplaris felipensis Wedd., *Ann. Sci. Nat., Bot. sér. 3, 13:* 263 (Weddell 1849).

Triplaris noli-tangere Wedd., *Ann. Sci. Nat., Bot. sér. 3, 13:* 264 (Weddell 1849).

Ruprechtia martii Meisn., *Fl. Bras. [Martius]* 5 (1): 58 [1 Jan. 1855] (Meisner 1855).

Triplaris pavonii Meisn., *Prodr. [A. P. de Candolle]* 14 (1): 172 [mid Oct. 1856] (Meisner 1856).

Triplaris formicosa S.Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 444 [“1894-96” publ. Dec. 1895] (Moore 1895).

Triplaris guanaiensis Rusby, *Mem. Torrey Bot. Club* 6 (1): 111 (Rusby 1896).

Triplaris estriata Kuntze, *Revis. Gen. Pl.* 3 (3): 271 [28 Sep. 1898] (Kuntze 1898).

Triplaris boliviiana Britton, *Bull. Torrey Bot. Club* 27 (3): 130 (Britton 1900).

Triplaris euryphylla S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 239 [9 Sep. 1919] (Blake 1919).

Triplaris laxa S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (7): 240 [9 Sep. 1919] (Blake 1919).

Triplaris williamsii Rusby, *Mem. New York Bot. Gard.* 7: 235 (Rusby 1927).

HERBARIUM DATA (FG). — Not yet collected, but the presence of this very widespread species is expected.

SIZE. — Up to 27 m tall (Aymard C. & Howard 2004).

[1406] *Triplaris weigeltiana* (Rchb.) Kuntze

Revis. Gen. Pl. 3 (3): 270 [28 Sep. 1898] (Kuntze 1898). — *Blochmannia weigeltiana* Rchb., *Surinam Exsic. [Weigelt]* s.n. (Reichenbach 1828).

Triplaris surinamensis Cham., *Linnaea* 8: 138 (Chamisso 1833).

Triplaris vahliana Fisch. & C.A.Mey. ex C.A.Mey., *Mém. Acad. Imp. Sci. Saint-Pétersbourg*, Sér. 6, *Sci. Math., Seconde Pt. Sci. Nat.* 6 (2, Bot.): 148 [Nov. 1840] (Meyer 1840).

Triplaris surinamensis var. *crassifolia* Benth., *London J. Bot.* 4: 628 (Bentham 1845).

Triplaris martiana Fisch. & C.A.Mey. ex C.A.Mey. var. *oblongifolia* Meisn., *Fl. Bras. [Martius]* 5 (1): 50 [1 Jan. 1855] (Meisner 1855).

Triplaris surinamensis var. *benthamiana* Meisn., *Fl. Bras. [Martius]* 5 (1): 50 [1 Jan. 1855] (Meisner 1855), “*Benthamiana*”.

Triplaris surinamensis var. *chamissoana* Meisn., *Fl. Bras. [Martius]* 5 (1): 50 [1 Jan. 1855] (Meisner 1855).

Triplaris siphonopetala H.Gross, *Bot. Jahrb. Syst.* 49 (2): 347 [14 Jan. 1913] (Gross 1913).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Weigelt's *Surinam exciccatae* were sent to several herbaria in 1828, together with printed labels bearing full descriptions (Stafleu & Cowan 1988). Therefore, all requirement are met for the effective publication of *Blochmannia weigeltiana* Rchb. (see Turland *et al.* 2018: Art. 30.8, Ex. 13).

VERNACULAR NAMES. — Pa: á-wareuni • Ka: tasi • Wp: moyu tasi • Wn: mekulak, mukulap • Nt: doon udu, doonudu • Cr: bwa-fronmi • Br: taxi-preto-da-várzea, taxiseiro.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *M.-F. Prévost* 750.

INVENTORY DATA (FG). — 1 tree, dbh = 48 cm.

Family PRIMULACEAE Batsch ex Borkh.
Genus *Cybianthus* Mart.

[1407] *Cybianthus amplus* (Mez) G.Agostini

Acta Biol. Venez. 10 (2): 152 (Agostini 1980). — *Conomorpha ampla* Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 257 [6 May 1902] (Mez 1902).

Conomorpha macrophylla Mart., *Flora* 24 (2, Beibl.): 20 [28 Aug. 1841] (Martius 1841).

Conomorpha utiarityi Hoehne, *Comm. Lin. Telegr., Bot.* 6: 64 [Sep. 1915] (Hoehne 1915).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman* T-909.

SIZE. — Brazil, Amazonas. *C.A. Cid Ferreira* 7600 (MO), 15 m × 25 cm.

[1408] *Cybianthus guyanensis* (A.DC.) Miq. subsp. *multipunctatus* (A.DC.) Pipoly

Ann. Missouri Bot. Gard. 79 (4): 944 (Pipoly 1992). — *Cybianthus multipunctatus* A.DC., *Ann. Sci. Nat., Bot. sér. 2,* 16: 94 (Candolle 1841). — *Conomorpha multipunctata* (A.DC.) Miq., *Stirp. Surinam. Select.*: 111 [“1850” publ. Mar. 1851] (Miquel 1851).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *J. Martin* s.n. (original material G-DC, G00139364).

INVENTORY DATA (FG). — 1 tree, dbh = 19.4 cm.

[1409] *Cybianthus leprieurii* G.Agostini

Acta Biol. Venez. 10 (2): 159 (Agostini 1980). — *Weigelia parviflora* Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 288 [6 May 1902] (Mez 1902).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *F.M.R. Leprieur* s.n. (original material P[P00715205]); *R.A.A. Oldeman* B-521, height 11 m.

[1410] *Cybianthus microbotrys* A.DC.

Ann. Sci. Nat., Bot. sér. 2, 16: 94 (Candolle 1841). — *Peckia microbotrys* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 402 [5 Nov. 1891] (Kuntze 1891). — *Weigelia microbotrys* (A.D.C.) Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 285 [6 May 1902] (Mez 1902).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: pune-etni-kamwi • Wp: ka'a pelele • Br: capororoquinha.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *S.A. Mori* & *J.J. Pipoly* 15528.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{inv} = 14.5$ cm.

[1411] *Cybianthus potiae* (Mez) G.Agostini
(Fig. 46D)

Acta Biol. Venez. 10 (2): 161 (Agostini 1980). — *Weigelia potiae* Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 285 [6 May 1902] (Mez 1902).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *P.A. Poiteau* s.n. (original material P[P00715218]).

INVENTORY DATA (FG). — 1 tree, dbh = 11.8 cm.

[1412] *Cybianthus prieurii* A.DC.

Ann. Sci. Nat., Bot. sér. 2, 16: 93 (Candolle 1841), “*Prieurei*”. — *Peckia prieurii* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 402 [5 Nov. 1891] (Kuntze 1891).

Cybianthus nitidus Miq., *Fl. Bras. [Martius]* 10: 295 [15 Mar. 1856] (Miquel 1856). — *Peckia nitida* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 402 [5 Nov. 1891] (Kuntze 1891).

Cybianthus subspicatus Benth. ex Miq., *Fl. Bras. [Martius]* 10: 296 [15 Mar. 1856] (Miquel 1856). — *Peckia subspicata* (Benth. ex Miq.) Kuntze, *Revis. Gen. Pl.* 2: 402 [5 Nov. 1891] (Kuntze 1891).

Cybianthus comatus Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 219 [6 May 1902] (Mez 1902).

Cybianthus viridiflorus A.C.Sm., *Lloydia* 2 (3): 204 (Smith 1939).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *N. Marshall & J. Rombold 146*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; dbh_{inv} = 12.7 cm.

[1413] *Cybianthus resinosus* Mez

Pflanzenr. [Engler] IV.236 (Heft 9): 219 [6 May 1902] (Mez 1902).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 8915*.

SIZE. — Up to 15 m tall (Pipoly & Gustafsson 2002).

[1414] *Cybianthus surinamensis* (A.Spreng.) G.Agostini

Acta Biol. Venez. 10 (2): 161 (Agostini 1980). — *Salvadora surinamensis* A.Spreng., *Tent. Suppl.*: 7 [Sep. 1828] (Sprengel 1828). — *Peckia surinamensis* (A.Spreng.) Kuntze, *Revis. Gen. Pl.* 2: 402 [5 Nov. 1891] (Kuntze 1891). — *Weigeltia surinamensis* (A.Spreng.) Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 289 [6 May 1902] (Mez 1902).

Weigeltia myrianthos A.DC., *Trans. Linn. Soc. London* 17 (2): 102 [“1837” publ. 26 Apr.-8 May 1834] (Candolle 1834). — *Wallenia myrianthos* Rchb. ex A.DC., *Trans. Linn. Soc. London* 17 (2): 102 [“1837” publ. 26 Apr.-8 May 1834] (Candolle 1834), nom. nud. pro syn. — *Cybianthus myrianthos* (A.DC.) Miq., *Fl. Bras. [Martius]* 10: 300 [15 Mar. 1856] (Miquel 1856).

Weigeltia capitellata Miq., *Fl. Bras. [Martius]* 10: 300 [15 Mar. 1856] (Miquel 1856), nom. nud. pro syn.

Weigeltia schomburgkiana Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 285 [6 May 1902] (Mez 1902). — *Cybianthus schomburgkianus* (Mez) G.Agostini, *Acta Biol. Venez.* 10 (2): 165 (Agostini 1980).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-J. de Granville B-4467*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 13.8 cm.

[1415] *Cybianthus venezuelanus* Mez

Pflanzenr. [Engler] IV.236 (Heft 9): 221 [6 May 1902] (Mez 1902).

Cybianthus egensis Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 222 [6 May 1902] (Mez 1902).

Peckia purpurea Rusby, *Bull. New York Bot. Gard.* 4 (14): 405 [7 Dec. 1907] (Rusby 1907).

Cybianthus brownii Gleason, *Bull. Torrey Bot. Club* 53 (5): 293 [May 1926] (Gleason 1926).

VERNACULAR NAMES. — Wp: wila ka’alulu wipē’i.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-3045*.

SIZE. — Brazil, Acre. *C.A. Cid Ferreira 10202* (MO), 10 m × 25 cm.

[1416] *Cybianthus* sp. A

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino et al. 2130*.

INVENTORY DATA (FG). — 1 tree, dbh = 13.1 cm.

Genus *Myrsine* L.[1417] *Myrsine guianensis* (Aubl.) Kuntze

Revis. Gen. Pl. 2: 402 [5 Nov. 1891] (Kuntze 1891). — *Rapanea guianensis* Aubl., *Hist. Pl. Guiane* 1: 121 [Jun.-Dec. 1775] (Aublet 1775). — *Myrsine rapanea* Roem. & Schult., *Syst. Veg. [Roemer & Schultes/4:* 509 [Mar.-June 1819] (Roemer & Schultes 1819), nom. illeg. superfl. (based on *Rapanea guianensis*). — *Myrsine rapanea* f. *communis* Miq., *Fl. Bras. [Martius]* 10: 308 [15 Mar. 1856] (Miquel 1856), nom. illeg. superfl. (based on *Rapanea guianensis*).

Myrsine rapanea var. *major* Hook. & Arn., *J. Bot. [Hooker]* 1: 283 (Hooker & Arnott 1834).

Myrsine rapanea var. *minor* Hook. & Arn., *J. Bot. [Hooker]* 1: 283 (Hooker & Arnott 1834).

Myrsine rapanea var. *subacuta* A.DC., *Prodr. [A. P. de Candolle]* 8: 98 [mid Mar. 1844] (Candolle 1844).

Myrsine rapanea f. *angustata* Miq., *Fl. Bras. [Martius]* 10: 308 [15 Mar. 1856] (Miquel 1856).

Myrsine rapanea f. *paviflora* Miq., *Fl. Bras. [Martius]* 10: 308 [15 Mar. 1856] (Miquel 1856).

Myrsine rapanea f. *robusta* Miq., *Fl. Bras. [Martius]* 10: 308 [15 Mar. 1856] (Miquel 1856). — *Rapanea oblonga* Pohl ex Miq., *Fl. Bras. [Martius]* 10: 308 [15 Mar. 1856] (Miquel 1856), nom. nud. pro syn.

Myrsine ovalifolia Miq., *Fl. Bras. [Martius]* 10: 313 [15 Mar. 1856] (Miquel 1856), nom. illeg. hom., non (A.DC.) D.Dietr. (Dietrich 1839).

Rapanea guianensis var. *andicola* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 325 [Dec. 1951] (Cuatrecasas 1951), “*guyanensis*”.

Rapanea guianensis var. *bogotensis* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 325 [Dec. 1951] (Cuatrecasas 1951), “*guyanensis*”.

Rapanea guianensis var. *calensis* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 325 [Dec. 1951] (Cuatrecasas 1951), “*guyanensis*”.

NOTE. — Roemer and Schultes mistakenly ascribed the binomial *Myrsine rapanea* to R.Br., referencing Brown (1810: 533). There Brown simply (and incorrectly) synonymised *Rapanea* Aubl. (i.e. *R. guianensis*) under *Samara floribunda* Willd.

VERNACULAR NAMES. — Ka: kunapolan, kusapoi.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777907] designated by Lanjouw & Uittien [1940: 156]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.4$ cm.

Genus *Stylogyne* A.DC.

[1418] *Stylogyne orinocensis* (Kunth) Mez

Pflanzenr. [Engler] IV.236 (Heft 9): 270 [6 May 1902] (Mez 1902). — *Ardisia orinocensis* Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 244 [9 July 1819] (Kunth 1819). — *Tinus orinocensis* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 974 [5 Nov. 1891] (Kuntze 1891).

Badula schomburgkiana A.DC., *Ann. Sci. Nat., Bot. sér. 2*, 16: 90 (Candolle 1841). — *Ardisia schomburgkiana* (A.DC.) Miq., *Fl. Bras. [Martius]* 10: 289 [15 Mar. 1856] (Miquel 1856). — *Tinus schomburgkiana* (A.DC.) Kuntze, *Revis. Gen. Pl.* 2: 975 [5 Nov. 1891] (Kuntze 1891). — *Stylogyne schomburgkiana* (A.DC.) Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 270 [6 May 1902] (Mez 1902).

Badula schomburgkiana var. *brasiliensis* A.DC., *Prodr. [A. P. de Candolle]* 8: 111 [mid Mar. 1844] (Candolle 1844). — *Stylogyne brasiliensis* (A.DC.) Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 269 [6 May 1902] (Mez 1902).

Ardisia hostmannii Miq., *Fl. Bras. [Martius]* 10: 288 [15 Mar. 1856] (Miquel 1856), “hostmannii”. — *Tinus hostmannii* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 974 [5 Nov. 1891] (Kuntze 1891).

Ardisia surinamensis Miq., *Fl. Bras. [Martius]* 10: 288 [15 Mar. 1856] (Miquel 1856). — *Tinus surinamensis* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 975 [5 Nov. 1891] (Kuntze 1891). — *Stylogyne surinamensis* (Miq.) Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 271 [6 May 1902] (Mez 1902).

Stylogyne micans Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 272 [6 May 1902] (Mez 1902).

Stylogyne poeppigii Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 274 [6 May 1902] (Mez 1902).

Stylogyne amazonica Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 275 [6 May 1902] (Mez 1902).

Stylogyne kappleri Mez, *Pflanzenr. [Engler]* IV.236 (Heft 9): 275 [6 May 1902] (Mez 1902).

VERNACULAR NAMES. — Pa: pune-etni-priye (for *Stylogyne* sp.) • Ka: payawalu.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Moretti* 606.

SIZE. — Up to 14 m tall (Ricketson & Pipoly 2010).

Family PROTEACEAE Juss.
Genus *Euplassa* Salisb. ex Knight

[1419] *Euplassa pinnata* (Lam.) I.M.Johnst.

Contr. Gray Herb. 73: 42 (Johnston 1924). — *Roupala pinnata* Lam., *Tabl. Encycl.* 1[1 (2)]: 243 [13 Feb. 1792] (Lamarck 1792). — *Adenostephanus guyanensis* Meissn., *Fl. Bras. [Martius]* 5 (1): 95 [1 Jan. 1855] (Meissner 1855), *nom. illeg. superfl.* (based on *Roupala pinnata*).

Euplassa meridionalis Salisb. ex Knight, *Cult. Prot.* 101 [Dec. 1809] (Knight 1809).

VERNACULAR NAMES. — Wp: tatu kāsī • Nt: liba apisi • Cr: bwa-graj • Fr: bois grage blanc • Br: facira, louro-faia.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J. Martins* s.n. (original material BM[BM000080820, BM000080821, BM000080822]).

INVENTORY DATA (FG). — 17 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 42$ cm.

Genus *Panopsis* Salisb. ex Knight

[1420] *Panopsis rubescens* (Pohl) Rusby

Mem. Torrey Bot. Club 6 (1): 116 (Rusby 1896). — *Andriapetalum rubescens* Pohl, *Pl. Bras. Icon. Descr.* 1 (4): 114 [May-Dec. 1828] (Pohl 1828).

Andriapetalum rubescens var. *acuminatum* Meissn., *Prodr. [A. P. de Candolle]* 14 (1): 346 [mid Oct. 1856] (Meissner 1856), “*Andriapetalum*”. — *Panopsis acuminata* (Meissn.) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (2): 66 (Macbride 1931).

Andriapetalum sprucei Meissn., *Prodr. [A. P. de Candolle]* 14 (1): 347 [mid Oct. 1856] (Meissner 1856). — *Panopsis rubescens* var. *sprucei* (Meissn.) Sleumer, *Bot. Jahrb. Syst.* 76 (2): 180 [Jan. 1954] (Sleumer 1954).

Roupala yauaperyensis Barb.Rodr., *Vellozia*, ed. 2, 1: 66 (Barbosa Rodrigues 1891).

Panopsis sprucei Rusby, *Mem. Torrey Bot. Club* 6 (1): 116 (Rusby 1896), *nom. illeg. superfl.*

Panopsis rubescens var. *simulans* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (2): 67 (Macbride 1931).

Panopsis cuaensis Steyermark, *Ann. Missouri Bot. Gard.* 74 (3): 612 (Steyermark 1987).

NOTES. — Rusby (1896: 116) did not cite *Andriapetalum sprucei* Meissn. However, he ascribed the binomial “*Panopsis sprucei*” to “Meissn. Ms. in Herb. Kew”, while stating that the original material (*Bang* 1686 [NY, 01364973]) is “the same as Spruce’s no. 1817”. Spruce 1817 is the type of *Andriapetalum sprucei*, and the K specimens (K000634341, K000634342) bear the handwritten annotation (not in Meissner’s hand) “*Andriapetalum Sprucei* Meissn.”, which is presumably what Rusby was referring to. As Meissner did not use *Panopsis sprucei*, the authorship of this name is “Rusby”, not “Meissn. ex Rusby”. Furthermore, *P. sprucei* is an illegitimate superfluous name because although independent from *A. sprucei*, the type of the latter is cited in the protologue.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori et al. 20968*.

INVENTORY DATA (FG). — 8 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54.1$ cm.

[1421] *Panopsis sessilifolia* (Rich.) Sandwith
(Fig. 47A)

Bull. Misc. Inform. Kew 1932 (5): 226 [27 June 1932] (Sandwith 1932). — *Roupala sessilifolia* Rich., *Actes Soc. Hist. Nat. Paris* 1: 106 [Oct. 1792] (Richard 1792). — *Andriapetalum sessilifolium* (Rich.) Klotzsch, *Linnaea* 15: 53 (Klotzsch 1841), “*Andripetalum sessilifolia*”.

Roupala hameliiifolia Rudge, *Pl. Guian. [Rudge]* 1 (3): 22 [June 1805] (Rudge 1805), “*hameliaefolia*”. — *Panopsis hameliiifolia* (Rudge) Knight, *Cult. Prot.*: 104 [Dec. 1809] (Knight 1809).

Andriapetalum cayennense Klotzsch ex Meisn., *Fl. Bras. [Martius]* 5 (1): 78 [1 Jan. 1855] (Meisner 1855), “*Andripetalum*”.

VERNACULAR NAMES. — Wp: kuya'i lá • Wn: meliju • Cr: bwa-grajroz • Br: faeira, louro-faia.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.B. Leblond 224* (original material P[P00750531]).

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 34.1$ cm.

Genus *Roupala* Aubl.

[1422] *Roupala montana* Aubl.
(Fig. 47B)

Hist. Pl. Guiane 1: 83 [Jun.-Dec. 1775] (Aublet 1775).

Embothrium pinnatum Ruiz & Pav., *Flora Peruviana* 1: 63 (Ruiz & Pavón 1798). — *Roupala diversifolia* R.Br., *Trans. Linn. Soc. London* 10 (1): 193 [“1811” publ. 1810] (Brown 1810), “*Rhopala*”, nom. illeg. superfl. (based on *Embothrium pinnatum*). — *Roupala pinnata* (Ruiz & Pav.) Diels ex J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (2.2)*: 374 [15 Mar. 1937] (Macbride 1937), nom. illeg. hom., non Lam. (Lamarck 1791) [synonym of *Euplassa pinnata*].

Roupala pyrifolia Knight, *Cult. Prot.*: 102 [Dec. 1809] (Knight 1809).

Roupala media R.Br., *Trans. Linn. Soc. London* 10 (1): 191 [“1811” publ. 1810] (Brown 1810), “*Rhopala*”.

Roupala dentata R.Br., *Trans. Linn. Soc. London* 10 (1): 192 [“1811” publ. 1810] (Brown 1810).

Roupala complicata Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 153 [8 Dec. 1817] (Kunth 1818), “*Rhopala*”.

Roupala ovalis Pohl, *Pl. Bras. Icon. Descr.* 1 (4): 107 [May-Dec. 1828] (Pohl 1828), “*Rhopala*”.

Roupala affinis Pohl, *Pl. Bras. Icon. Descr.* 1 (4): 112 [May-Dec. 1828] (Pohl 1828), “*Rhopala*”.

Roupala macropoda Klotzsch & H.Karst., *Linnaea* 20: 473 [Oct. 1847] (Klotzsch & Karsten 1847), “*Rhopala*”.

Roupala tomentosa Pohl var. *sellowii* Meisn., *Fl. Bras. [Martius]* 5 (1): 82 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”. — *Roupala ovalis*

Klotzsch ex Meisn., *Fl. Bras. [Martius]* 5 (1): 82 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”, nom. nud. pro syn.

Roupala gardneri Meisn., *Fl. Bras. [Martius]* 5 (1): 83 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala gardneri var. *angustata* Meisn., *Fl. Bras. [Martius]* 5 (1): 83 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala gardneri var. *dentata* Meisn., *Fl. Bras. [Martius]* 5 (1): 83 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala gardneri var. *integrifolia* Meisn., *Fl. Bras. [Martius]* 5 (1): 83 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala glabrata Klotzsch ex Meisn., *Fl. Bras. [Martius]* 5 (1): 84 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala brasiliensis Klotzsch var. *macropoda* Meisn., *Fl. Bras. [Martius]* 5 (1): 86 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala martii Meisn., *Fl. Bras. [Martius]* 5 (1): 87 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala martii var. *pinnata* Meisn., *Fl. Bras. [Martius]* 5 (1): 87 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”.

Roupala martii var. *simplicifolia* Meisn., *Fl. Bras. [Martius]* 5 (1): 87 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”. — *Roupala veraguensis* Klotzsch ex Meisn., *Fl. Bras. [Martius]* 5 (1): 87 [1 Jan. 1855] (Meisner 1855), “*Rhopala*”, nom. nud. pro syn.

Roupala boissieriana Meisn., *Prodr. [A. P. de Candolle]* 14 (1): 430 [mid Oct. 1856] (Meisner 1856), “*Rhopala*”.

Roupala montana var. *complicata* (Kunth) Griseb., *Fl. Brit. W.I. [Grisebach]*: 277 [late 1860] (Grisebach 1860).

Roupala montana var. *heterophylla* Griseb., *Fl. Brit. W.I. [Grisebach]*: 278 [late 1860] (Grisebach 1860).

Roupala borealis Hemsl., *Biol. Cent.-Amer. Bot.* 3 (14): 78 [Oct. 1882] (Hemsley 1882).

Roupala acuminata Glaz., *Bull. Soc. Bot. France* 59 (Mém. 3g): 602 [“1912” publ. 1913] (Glaziou 1913), nom. nud.

Roupala darienensis Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 228 [22 Sep. 1917] (Pittier 1917).

Roupala panamensis Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 229 [22 Sep. 1917] (Pittier 1917).

Roupala discolor Rusby, *Descr. S. Amer. Pl.* 12 [20 Dec. 1920] (Rusby 1920).

Roupala dissimilis Pittier, *Bol. Soc. Venez. Ci. Nat.* 5: 303 (Pittier 1939).

Roupala repanda Lundell, *Amer. Midl. Naturalist* 29 (2): 472 [5 Apr. 1943] (Lundell 1943).

Roupala sphenophylla Diels ex Sleumer, *Bot. Jahrb. Syst.* 76 (2): 166 [Jan. 1954] (Sleumer 1954).

Roupala montana var. *dentata* (R.Br.) Sleumer, *Bot. Jahrb. Syst.* 76 (2): 173 [Jan. 1954] (Sleumer 1954).

Roupala mayana Lundell, *Wrightia* 5 (4): 73 (Lundell 1975).

NOTES. — *Roupala brasiliensis* var. *macropoda* Meisn. is not based on *R. macropoda* Klotzsch & H.Karst. Prance (2009: 134) incorrectly cited the protologue of *R. tomentosa* var. *sellowii* Meisn.: “*Prod. [A. P. de Candolle]* 14 (1): 28. 1856”. It is obvious that the “4” was

omitted, as on p. 428 of Candolle's *Prodromus* this variety is cited in synonymy under *R. tomentosa* var. *dentata* Meisn.

VERNACULAR NAMES. — Wp: tatu kásī • Nt: liba apisi • Cr: bwa-grajrouj • Fr: bois grage rouge • Br: carne-de-vaca, carvalho-vermelha, facira, louro-faia.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777521] designated by Lanjouw & Uittien [1940: 156]).

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 23.2 \text{ cm}$.

[1423] *Roupala nitida* Rudge

Pl. Guian. [Rudge] 1 (3): 26 [June 1805] (Rudge 1805).

Roupala laurifolia Knight, *Cult. Prot.* 102 [Dec. 1809] (Knight 1809), nom. illeg. superfl. (based on *Roupala nitida*).

VERNACULAR NAMES. — Wp: tatu kásī • Wn: awawaju.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J. Martin 52* (original material BM[BM000839785]); *P. Grenand 626*, height 15 m.

Family PUTRANJIVACEAE Endl.
Genus *Drypetes* Vahl

[1424] *Drypetes fanshawei* Sandwith
(Fig. 47C)

Kew Bull. 7 (2): 258 [25 July 1952] (Sandwith 1952).

VERNACULAR NAMES. — Pa: wakukwa-ára • Nt: weti kookoo.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2831*.

INVENTORY DATA (FG). — 295 trees in 89 plots; $F_{\max} = 5\%$; $\text{dbh}_{\text{inv}} = 41.7 \text{ cm}$.

[1425] *Drypetes variabilis* Uittien
(Fig. 47D)

Recueil Trav. Bot. Néerl. 22: 348 ["1925" publ. Jan. 1926] (Uittien 1926).

VERNACULAR NAMES. — Ka: pana akala, sekeleu • Nt: weti kookoo.

HERBARIUM DATA (FG). — 128 collections at CAY. Sel. exs.: *J.-F. Molino 1769*.

INVENTORY DATA (FG). — 369 trees in 126 plots; $F_{\max} = 5.4\%$; $\text{dbh}_{\text{inv}} = 73 \text{ cm}$.

Family RHABDODENDRACEAE Prance
Genus *Rhabdodendron* Gilg & Pilg.

[1426] *Rhabdodendron amazonicum*
(Spruce ex Benth.) Huber
(Fig. 48A)

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 5 (2): 427 (Huber 1909). — *Lecostemon amazonicum* Spruce ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 295 (Bentham 1853).

Lecostemon crassipes Spruce ex Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 295 (Bentham 1853). — *Rhabdodendron crassipes* (Spruce ex Benth.) Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 428 (Huber 1909).

Lecostemon crassipes var. *cayennense* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 5: 296 (Bentham 1853).

Rhabdodendron duckei Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 428 (Huber 1909).

Rhabdodendron longifolium Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 429 (Huber 1909).

Rhabdodendron paniculatum Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 429 (Huber 1909).

Rhabdodendron arirambae Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 5 (2): 430 (Huber 1909).

Lecostemon sylvestre Gleason, *Bull. Torrey Bot. Club* 54 (2): 68 [Feb. 1927] (Gleason 1927). — *Rhabdodendron sylvestre* (Gleason) Maguire, *Bull. Torrey Bot. Club* 75 (4): 397 [Jul.-Aug. 1948] (Maguire 1948).

VERNACULAR NAMES. — Pa: batakivie • Wp: alala kā'i sili, wila to pokó • Br: batiputá.

HERBARIUM DATA (FG). — 54 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3022*.

INVENTORY DATA (FG). — 93 trees in 30 plots; $F_{\max} = 4\%$; $\text{dbh}_{\text{inv}} = 25.5 \text{ cm}$.

Family RHAMNACEAE Juss.
Genus *Sarcomphalus* P.Browne

[1427] *Sarcomphalus cinnamomum*
(Triana & Planch.) Hauenschild
(Fig. 48B)

Taxon 65 (1): 55 (Hauenschild 2016). — *Ziziphus cinnamomum* Triana & Planch., *Ann. Sci. Nat., Bot. sér. 5, 16:* 380 (Triana & Planchon 1872).

Ziziphus itacaiunensis Fróes, *Bol. Técn. Inst. Agron. N.* 36: 151 (Fróes 1959).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2538*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 70 \text{ cm}$.

Family RHIZOPHORACEAE Pers.
Genus *Cassipourea* Aubl.

[1428] *Cassipourea elliptica* (Sw.) Poir.

Encycl. JJ. Lamarck et al.] Suppl. 2: 131 [23 Oct. 1811] (Poirer 1811). — *Legnotis elliptica* Sw., *Prodr. [Swartz]*: 84 [20 Jun.-29 July 1788] (Swartz 1788). — *Cassipourea guianensis* var. *elliptica* (Sw.) M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.*: 84 (Gómez 1889). — *Cassipourea elliptica* var. *typica* Stehlé, *Bull. Soc. Bot. France* 92 (9): 265 (Stehlé 1945), nom. inval. (Turland et al. 2018: Art. 24.3).

Cassipourea alba Griseb., *Abh. Königl. Ges. Wiss. Göttingen* 7: 223 (Grisebach 1857). — *Cassipourea elliptica* var. *alba* (Griseb.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 274 [late 1860] (Grisebach 1860).

Cassipourea elliptica var. *pauciserrata* Griseb., *Fl. Brit. W.I. [Grisebach]*: 274 [late 1860] (Grisebach 1860).

Endosteira oppositifolia Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 576 (Turczaninow 1863).

Cassipourea elliptica var. *ovata* Griseb., *Cat. Pl. Cub. [Grisebach]*: 108 [May-Aug. 1866] (Grisebach 1866).

Cassipourea cubensis Urb., *Symb. Antill. [Urban]* 7 (2): 293 [15 June 1912] (Urban 1912).

Cassipourea brittoniana Fawc. & Rendle, *J. Bot.* 64: 14 (Fawcett & Rendle 1926).

Cassipourea podantha Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 241 [24 Oct. 1929] (Standley 1929).

Cassipourea macrodonta Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 242 [24 Oct. 1929] (Standley 1929).

Cassipourea guildingii Briq., *Candollea* 4: 343 (Briquet 1931).

Cassipourea belizensis Lundell, *Bull. Torrey Bot. Club* 66 (9): 598 [Dec. 1939] (Lundell 1939).

Cassipourea elliptica var. *parvifolia* Stehlé, *Bull. Soc. Bot. France* 92 (9): 265 (Stehlé 1945).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly* 15425.

SIZE. — Venezuela, Táchira, 7 Jul. 1983. *H. van der Werff* 5021 (MO), 15 m.

[1429] *Cassipourea guianensis* Aubl.
(Fig. 48C)

Hist. Pl. Guiane 1: 529 [Jun.-Dec. 1775] (Aublet 1775). — *Legnotis cassipourea* Sw., *Prodr. [Swartz]* 84 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. superfl. (based on *Cassipourea guianensis*).

Cassipourea macrophylla DC., *Prodr. [A. P. de Candolle]* 3: 34 [mid Mar. 1828] (Candolle 1828). — *Legnotis macrophylla* Mart. ex DC., *Prodr. [A. P. de Candolle]* 3: 34 [mid Mar. 1828] (Candolle 1828), nom. nud. pro syn.

Cassipourea serrata Benth., *J. Bot. [Hooker]* 2: 223 (Bentham 1840). — *Cassipourea guianensis* var. *serrata* (Benth.) Engl., *Fl. Bras. [Martius]* 12 (2): 430 [1 Sep. 1876] (Engler 1876).

Cassipourea quadrilocularis Spruce ex Benth., *Fl. Bras. [Martius]* 12 (2): 429 [1 Sep. 1876] (Bentham 1876), nom. nud. pro syn.

Cassipourea latifolia Alston, *Bull. Misc. Inform. Kew* 1925 (6): 266 [16 June 1925] (Alston 1925). — *Cassipourea elliptica* var. *latifolia* (Alston) Stehlé, *Bull. Soc. Bot. France* 92 (9): 265 (Stehlé 1945).

Cassipourea broadwayi Briq., *Candollea* 4: 342 (Briquet 1931).

Cassipourea guianensis var. *trichopoda* Briq., *Candollea* 4: 346 (Briquet 1931).

VERNACULAR NAMES. — Pa: wakáu • Ka: apali'ilan, apaliyulan • Wp: yawa poi li • Cr: pativyé-gri • Br: laranja-do-mato, mangue-d'agua-doce.

HERBARIUM DATA (FG). — 88 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000016972](#)]).

INVENTORY DATA (FG). — 423 trees in 93 plots; $F_{\max} = 4.5\%$; $dbh_{inv} = 23.1$ cm.

Genus *Rhizophora* L.

[1430] *Rhizophora mangle* L.

Sp. Pl. 1: 443 [1 May 1753] (Linnaeus 1753). — *Rhizophora americana* Nutt., *N. Amer. Sylva* 1 (2): 112 [Jul.-Dec. 1842] (Nuttall 1842), nom. illeg. superfl. (based on *Rhizophora mangle*).

Rhizophora mangle var. *samoensis* Hochr., *Candollea* 2: 447 (Hochreutiner 1925).

Rhizophora samoensis (Hochr.) Salvoza, *Nat. Appl. Sci. Bull. Univ. Philipp.* 5: 220 (Salvoza 1936).

NOTE. — A species restricted to mangrove forests.

VERNACULAR NAMES. — Pa: kwatri • Ka: konopo, kunapo • Cr: pativyé-janm, pativyé-rouj • Fr: palétuvier rouge • Br: mangue-vermelho.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Loubry* 2000.

SIZE. — Up to 100 cm dbh (Prance 2018).

[1431] *Rhizophora racemosa* G.Mey.

Prim. Fl. Esseq. 185 [Nov. 1818] (Meyer 1818). — *Rhizophora mangle* var. *racemosa* (G.Mey.) Engl., *Fl. Bras. [Martius]* 12 (2): 427 [1 Sep. 1876] (Engler 1876).

NOTE. — A species restricted to mangrove forests.

VERNACULAR NAMES. — Pa: kwatri • Ka: konopo, kunapo • Cr: pativyé-janm, pativyé-rouj • Fr: palétuvier rouge • Br: mangue-vermelho.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *M.-F. Prévost* 4684.

SIZE. — Up to 15 m tall (Steyermark 2004).

Family ROSACEAE Juss.

Genus *Prunus* L.

[1432] *Prunus accumulans* (Koehne) C.L.Li & Aymard

BioLlania, ed. espec. 6: 449 (Li & Aymard 1997). — *Prunus myrtifolia* var. *accumulans* Koehne, *Bot. Jahrb. Syst.* 52 (4-5): 320 [3 Aug. 1915] (Koehne 1915).

HERBARIUM DATA (FG). — A single collection, *G. Bourdy* 2941.

INVENTORY DATA (FG). — 13 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{inv} = 53$ cm.

[1433] *Prunus myrtifolia* (L.) Urb.
(Fig. 48D)

Bot. Jahrb. Syst. 52 (4-5): 320 [3 Aug. 1915] (Urban 1915). — *Celastrus myrtifolius* L., *Sp. Pl.* 1: 196 [1 May 1753] (Linnaeus 1753). — *Lauro-cerasus myrtifolia* (L.) Britton, *N. Amer. Trees* 510 [9 May 1908] (Britton 1908), “*Laurocerasus*”.

Prunus sphaerocarpa Sw., *Prodri. [Swartz]* 80 [20 Jun.-29 July 1788] (Swartz 1788).

Prunus tikalana Lundell, *Wrightia* 4 (2): 88 [31 Dec. 1968] (Lundell 1968).

VERNACULAR NAMES. — Pa: kunau • Ka: okoipyo • Wp: kunawalu'i • Cr: bwa-noyo • Br: marmelo-bravo, virarú.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 10765*.

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 48.1$ cm.

Family RUBIACEAE Juss.
Genus *Alibertia* A.Rich.

[1434] *Alibertia edulis* (Rich.) A.Rich.

Prodri. [A. P. de Candolle] 4: 443 [late Sep. 1830] (Richard 1830). — *Genipa edulis* Rich., *Actes Soc. Hist. Nat. Paris* 1: 107 [Oct. 1792] (Richard 1792). — *Gardenia edulis* (Rich.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 2: 708 [3 July 1812] (Poiret 1812). — *Alibertia utilis* A.Rich., *Mém. Rubiac.*, t. 21 [Dec. 1830] (Richard 1830), orthographical variant. — *Amaioua utilis* Baill., *Bull. Mens. Soc. Linn. Paris* 1 (28): 220 (Baillon 1879), *nom. illeg. superfl.* (based on *Alibertia edulis*). — *Amaioua edulis* (Rich.) Baill., *Hist. Pl. [Baillon]* 7: 387 [Feb. 1880] (Baillon 1880). — *Cordiera edulis* (Rich.) Kuntze, *Revis. Gen. Pl.* 1: 279 [5 Nov. 1891] (Kuntze 1891). — *Sabicea edulis* (Rich.) Seem., *Hooker's J. Bot. Kew Gard. Misc.* 3: 266 (Seemann 1851).

NOTES. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017). Persson & Delprete (2017) recognized 4 varieties in *Alibertia edulis* (Rich.) A.Rich. Only var. *edulis* is present in French Guiana.

VERNACULAR NAMES. — Pa: tukuyuy • Wp: yanipa lali • Cr: gouyav-kaka-poule, gouyav-nwé • Br: apuruizinho, marmelada, puruhý, purui-pequeno.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *J.B. Leblond s.n.* (lecto-, P[P00870033], designated by Delprete & Persson in Taylor et al. [2011: 136]).

SIZE. — Up to 10 m tall (Taylor et al. 2004).

[1435] *Alibertia latifolia* (Benth.) K.Schum.

Fl. Bras. [Martius] 6 (6): 386 [15 June 1889] (Schumann 1889). — *Cordiera latifolia* Benth., *J. Bot. [Hooker]* 3: 221 (Bentham 1841).

Alibertia oligantha K.Schum., *Fl. Bras. [Martius]* 6 (6): 385 [15 June 1889] (Schumann 1889). — *Cordiera oligantha* (K.Schum.) Kuntze, *Revis. Gen. Pl.* 1: 279 [5 Nov. 1891] (Kuntze 1891).

Alibertia latifolia var. *parvifolia* K.Schum., *Fl. Bras. [Martius]* 6 (6): 386 [15 June 1889] (Schumann 1889).

Alibertia granulosa Rusby, *Descr. S. Amer. Pl.* 133 [20 Dec. 1920] (Rusby 1920).

Alibertia latifolia var. *pargueniana* Steyermark, *Mem. New York Bot. Gard.* 12 (3): 225 (Steyermark 1965).

VERNACULAR NAMES. — Pa: tukuyuy.

HERBARIUM DATA (FG). — A single collection, *P. Grenand* 1808, 12 m.

Genus *Alseis* Schott

[1436] *Alseis longifolia* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 256 (Ducke 1922).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville B-4741*.

SIZE. — Up to 45 cm dbh (Boom & Delprete 2002).

Genus *Amaioua* Aubl.

[1437] *Amaioua glomerulata* (Lam. ex Poir.)
Delprete & C.H.Perss.

Adansonia, sér. 3, 34 (2): 356 [Dec. 2012] (Delprete & Persson 2012). — *Mussaenda glomerulata* Lam. ex Poir., *Encycl. [J. Lamarck et al.]* 4 (1): 393 [9 Feb. 1797] (Poiret 1797). — *Rondeletia glomerulata* (Lam. ex Poir.) Spreng., *Syst. Veg. [Sprengel]* 1: 707 [“1825” publ. late 1824] (Sprengel 1824).

Amaioua fagifolia Desf., *Mém. Mus. Hist. Nat.* 6: 14 (Desfontaines 1820).

Amaioua corymbosa Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 419 [13 Mar. 1820] (Kunth 1820). — *Hexactina corymbosa* (Kunth) Willd. ex Schult.f., *Syst. Veg. [Roemer & Schultes]* 7 (1): 91 (Schultes 1829), *nom. nud. pro syn.*

VERNACULAR NAMES. — Pa: wakavu-kamwi • Ka: wayu, wayun • Wp: kuluwi, puluwi • Cr: bwa-négrès, bwa-pièr • Br: goiabeira-preta, purui-grande.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *H. Puig 12055*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.6$ cm.

[1438] *Amaioua guianensis* Aubl.
(Fig. 49A)

Hist. Pl. Guiane 2 (Suppl.): 13 [Jun.-Dec. 1775] (Aublet 1775). — *Hamelia glabra* Lam., *Encycl. [J. Lamarck et al.]* 3 (1): 65 [19 Oct. 1789] (Lamarck 1789), *nom. illeg. superfl.* (based on *Amaioua guianensis*). — *Hamelia sessiliflora* Willd., *Sp. Pl. ed. 4* 1 (2): 981 [July 1798] (Willdenow 1798), *nom. illeg. superfl.* (based on *Amaioua guianensis*). — *Duhamelia glabra* Pers., *Syn. Pl. [Persoon]* 1: 203 [1 Apr.-15 June 1805] (Persoon 1805), *nom. illeg. superfl.* (based on *Amaioua guianensis*).




FIG. 47. — Proteaceae: **A**, *Panopsis sessilifolia* (Rich.) Sandwith (M.-F. Prévost & D. Sabatier 4000); **B**, *Roupala montana* Aubl. (D. Sabatier & M.-F. Prévost 4925). Putranjivaceae: **C**, *Drypetes fanshawei* Sandwith (D. Sabatier & J.-F. Molino 5001); **D**, *Drypetes variabilis* Uittien. © D. Sabatier/IRD.

Amaiaoua guianensis var. *macrantha* Steyermark., *Mem. New York Bot. Gard.* 12 (3): 216 (Steyermark 1965).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: maras-gamana • Ka: kapasi tuno • Wp: kuluwi, puluwi • Wn: akuli epa • Cr: bwa-négrès, bwa-pièr • Br: goiabeira-preta, purui-grande.

HERBARIUM DATA (FG). — 113 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, BM[BM001008914] designated by Steyermark [1965: 216]).

INVENTORY DATA (FG). — 149 trees in 80 plots; $F_{\max} = 1.6\%$; dbh_{inv} = 26.8 cm.

Genus *Botryarrhena* Ducke

[1439] *Botryarrhena pendula* Ducke

Notizbl. Bot. Gart. Berlin-Dahlem 11: 477 [11 July 1932] (Ducke 1932).

VERNACULAR NAMES. — Pa: iwan-etni, tukuyuy-kamwi.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *C. Feuillet et al. 10067*.

SIZE. — Tree to 20 m tall, dbh to 22.5(-45) cm.

Genus *Capirona* Spruce

[1440] *Capirona macrophylla* (Poepp.) Delporte (Fig. 49B)

Phytotaxa 443 (1): 102 [18 May 2020] (Delporte 2020). — *Condaminea macrophylla* Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 30 [15-21 Aug. 1841] (Poeppig 1841).

Capirona decorticans Spruce, *J. Proc. Linn. Soc., Bot.* 3: 200 [Apr. 1859] (Spruce 1859).

Monadelphanthus floridus H. Karst., *Fl. Columb. [H. Karsten]* 1 (2): 67 [13 Feb. 1860] (Karsten 1860).

Capirona duckei Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 7: 185 (Huber 1913), *nom. nud.*

Capirona leiophloea Benoist, *Bull. Mus. Natl. Hist. Nat.* 27: 367 (Benoist 1921).

Capirona huberiana Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 257 (Ducke 1922).

Capirona surinamensis Bremek., *Recueil Trav. Bot. Néerl.* 31: 261 (Bremekamp 1934).

Loretoa peruviana Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (5): 222 (Standley 1936).

Capirona wurdackii Steyermark., *Mem. New York Bot. Gard.* 10 (5): 190 (Steyermark 1964).

VERNACULAR NAMES. — Pa: wadidga, wadidga-priye, wadidga-wahuyo • Ka: akepiimyo, talipi, tapili • Wp: ayá pili • Wn: ékepüimë • Nt: lisa pao, mutende • Cr: bwa-palika • Br: escorrega-macaco, mulatorana, pau-mulato, perna-de-moça.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *O. Poncy 1048* (P[P00208816]).

INVENTORY DATA (FG). — 31 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 35.5$ cm.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2116*.

INVENTORY DATA (FG). — 207 trees in 93 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 114.6$ cm.

Genus *Cordiera* A.Rich.

[1443] *Cordiera myrciifolia* (K.Schum.) C.H.Perss. & Delprete

Fl. Venez. Guayana 8: 559 (Persson & Delprete 2004). — *Alibertia myrciifolia* K.Schum., *Fl. Bras. [Martius]* 6 (6): 393 [15 June 1889] (Schumann 1889). — *Cordiera myrciifolia* Spruce ex K.Schum., *Fl. Bras. [Martius]* 6 (6): 393 [15 June 1889] (Schumann 1889), *nom. nud. pro syn.*

Alibertia uniflora Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (5): 179 (Standley 1936).

Alibertia myrciifolia var. *tepuiensis* Steyermark., *Mem. New York Bot. Gard.* 12 (3): 226 (Steyermark 1965).

Alibertia triloba Steyermark., *Mem. New York Bot. Gard.* 12 (3): 227 (Steyermark 1965).

VERNACULAR NAMES. — Wn: malipaimë.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1993*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.8$ cm.

[1444] *Cordiera triflora* A.Rich.

Prod. [A. P. de Candolle] 4: 445 [late Sep. 1830] (Richard 1830). — *Alibertia triflora* (A.Rich.) K.Schum., *Fl. Bras. [Martius]* 6 (6): 392 [15 June 1889] (Schumann 1889).

Octavia sessiliflora DC., *Prod. [A. P. de Candolle]* 4: 464 [late Sep. 1830] (Candolle 1830). — *Nonatelia sessiliflora* (DC.) Kuntze, *Revis. Gen. Pl.* 1: 290 [5 Nov. 1891] (Kuntze 1891).

Cordiera trifolia Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 419 (Steudel 1840), orth. var.

Alibertia tenuifolia K.Krause, *Verh. Bot. Vereins Prov. Brandenburg* 50: 106 [30 Sep. 1908] (Krause 1908).

Alibertia steinbachii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 287 [24 Oct. 1929] (Standley 1929).

Alibertia benensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 7 (3): 291 (Standley 1931).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (lecto-, P[P03821428] selected by Persson & Delprete [2010: 263]).

SIZE. — Up to 20 cm dbh (Taylor et al. 2004).

Genus *Chimarrhis* Jacq.

[1441] *Chimarrhis microcarpa* Standl.

Bull. Torrey Bot. Club 53 (7): 471 [Oct. 1926] (Standley 1926).

Chimarrhis longistipulata Bremek., *Recueil Trav. Bot. Néerl.* 31: 260 (Bremekamp 1934).

VERNACULAR NAMES. — Wp: walalu'i • Cr: bwa-chapèl • Fr: bois cathédrale.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15562*.

INVENTORY DATA (FG). — 84 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 80$ cm.

[1442] *Chimarrhis turbinata* DC.

Prod. [A. P. de Candolle] 4: 404 [late Sep. 1830] (Candolle 1830). — *Pseudochimarrhis turbinata* (DC.) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 255 (Ducke 1922). — *Chimarrhis duckei* Rizzini, *Revista Brasil. Biol.* 7: 277 (Rizzini 1947), *nom. illeg. superfl.* (based on *Chimarrhis turbinata*).

Bathysa difformis Benoist, *Bull. Mus. Natl. Hist. Nat.* 26: 185 (Benoist 1920). — *Pseudochimarrhis difformis* (Benoist) Benoist, *Arch. Bot. 5 (Mém. 1)*: 265 [27 Mar. 1933] (Benoist 1933).

Elaeagia brasiliensis Standl., *Bull. Torrey Bot. Club* 60 (6): 395 [1 June 1933] (Standley 1933).

VERNACULAR NAMES. — Pa: kinuwup, warerey • Ka: abalaba, atakali, mapalaba, me:ku ubupo • Te: pukutsa'i • Wp: walalu'i • Wn: epalai, épalaï • Nt: pali udu • Cr: bwa-chapèl, bwa-pagay, sitronèl-rouj • Fr: bois cathédrale • Br: carapanaúba-cinzero, pau-de-remo.

Genus *Cosmibuena* Ruiz & Pav.

[1445] *Cosmibuena grandiflora* (Ruiz & Pav.) Rusby

Bull. New York Bot. Gard. 4 (14): 368 [7 Dec. 1907] (Rusby 1907). — *Cinchona grandiflora* Ruiz & Pav., *Fl. Peruv. [Ruiz & Pavón]* 2: 54 (Ruiz & Pavón 1799). — *Cinchona obtusifolia* D.Dietr., *Syn. Pl. [D. Dietrich]* 1: 721 [July 1839] (Dietrich 1839), nom. illeg. superfl. (based on *Cinchona grandiflora*).

Cosmibuena obtusifolia Ruiz & Pav., *Flora Peruviana* 3: 3 (Ruiz & Pavón 1802). — *Buena obtusifolia* (Ruiz & Pav.) DC., *Prod. [A. P de Candolle]* 4: 356 [late Sep. 1830] (Candolle 1830).

Cinchona quina Ruiz & Pav., *Flora Peruviana* 3: 4 (Ruiz & Pavón 1802).

Cinchona longiflora Mutis ex Steud., *Nomencl. Bot. [Steudel]* 1: 196 (Steudel 1821), nom. nud. pro syn.

Cinchona china Ruiz ex DC., *Prod. [A. P de Candolle]* 4: 356 [late Sep. 1830] (Candolle 1830).

Buena triflora Benth., *J. Bot. [Hooker]* 3: 216 (Bentham 1841). — *Cosmibuena triflora* (Benth.) Klotzsch, *Getreue Darstell. Gew.* 14 (2): t. 15 (Klotzsch 1846).

Buena latifolia Benth., *Pl. Hartw. [Bentham]*: 191 [? 16 Sep. 1845] (Bentham 1845). — *Cosmibuena latifolia* (Benth.) Klotzsch ex Walp., *Repert. Bot. Syst. [Walpers]* 6 (1): 69 [3-5 Sep. 1846] (Walpers 1846). — *Cosmibuena obtusifolia* var. *latifolia* (Benth.) Hook.f., *Bot. Mag.* 102: t. 6239 (Hooker 1876). — *Cosmibuena grandiflora* var. *latifolia* (Benth.) Steyermark., *Mem. New York Bot. Gard.* 23: 296 (Steyermark 1972).

Cosmibuena quinqueflora Klotzsch, *Getreue Darstell. Gew.* 14 (2): t. 15 (Klotzsch 1846).

Buena skinneri Oerst., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1852: 48 (Oersted 1852). — *Cosmibuena skinneri* (Oerst.) Hemsl., *Biol. Cent.-Amer., Bot.* 2 (7): 12 [Apr. 1881] (Hemsl. 1881).

Cosmibuena gardenioides Wernham, *J. Bot.* 51: 321 (Wernham 1913).

Cosmibuena arborea Standl., *Contr. U.S. Natl. Herb.* 17: 447 (Standley 1914).

Cosmibuena ovalis Standl., *Contr. U.S. Natl. Herb.* 18 (3): 137 [11 Feb. 1916] (Standley 1916).

NOTES. — Epiphytic or rarely terrestrial shrub or tree (rarely hemi-epiphytic strangling tree). *Cinchona china* is a misspell of *C. quina*.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *G. Cremers* & *J.-J. de Granville* 13176.

SIZE. — Up to 14 m tall (Taylor *et al.* 2004).

Genus *Coussarea* Aubl.

[1446] *Coussarea albescens* (DC.) Müll.Arg.

Fl. Bras. [Martius] 6 (5): 87 [1 July 1881] (Müller 1881), in obs. — *Faramea albescens* DC., *Prod. [A. P. de Candolle]* 4: 498 [late Sep. 1830] (Candolle 1830).

Coussarea racemosa A.Rich., *Prod. [A. P. de Candolle]* 4: 494 [late Sep. 1830] (Richard 1830).

NOTES. — The description and name of *C. racemosa* came from Richard (Dec. 1830: 177), the text of which Candolle had seen a preprint version. Thereafter Candolle (Sept. 1830: 498) cited Richard as the author of this name, so that in this case, the authority is "A.Rich.", not "A.Rich. ex DC." The names *F. albescens* and *C. racemosa* were published in the same volume and have thus equal priority. When transferring *F. albescens* to *Coussarea* Aubl., Müller (1881: 87) stated that it was synonymous with *C. racemosa*, and he chose *C. albescens* over *C. racemosa*, as indicated in the index of vol. 6 (5) of *Flora Brasilensis* (Müller 1888) where the former appears in normal type and the latter in italics (i.e. as a synonym). Under Art. 11.5 of the ICN (Turland *et al.* 2018), we thus have to use *C. albescens*, relegating *C. racemosa* as its junior synonym.

VERNACULAR NAMES. — Pa: mavinbi, mavinbi-kamwi, mavinvi, mavinvi-kamwi.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (original material of *Coussarea racemosa*: P[P03914186]).

INVENTORY DATA (FG). — 75 trees in 24 plots; $F_{\max} = 2.2\%$; $dbh_{inv} = 18.5$ cm.

[1447] *Coussarea grandis* Müll.Arg.

Fl. Bras. [Martius] 6 (5): 100 [1 July 1881] (Müller 1881).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *G. Cremers* 8397.

INVENTORY DATA (FG). — 11 trees in 2 plots; $F_{\max} = 1.5\%$; $dbh_{inv} = 16.1$ cm.

[1448] *Coussarea granvillei* Delprete & B.M.Boom

Brittonia 51 (4): 403 (Delprete & Boom 1999).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *J.-J. de Granville* B-4744 (holo-, CAY[CAY007384]); *J.-J. de Granville* et al. 8601, 10 m × 10 cm.

[1449] *Coussarea hallei* Steyermark.

Mem. New York Bot. Gard. 23: 387 (Steyermark 1972).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Nt: ayee wato.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *F. Hallei* 1018 (holo-, P[P00836890]; iso-, CAY[CAY028108, CAY028109], P[P00836891]); *P. Petronelli* 300, dbh 10 cm.

[1450] *Coussarea lutea* (Aubl.) Delprete

Taxon 64: 609 (Delprete 2015). — *Nonatelia lutea* Aubl., *Hist. Pl. Guiane* 1: 190, t. 74 [only inflorescence and flowers] [Jun.-Dec. 1775] (Aublet 1775). — *Psychotria lutea* (Aubl.) Willd., *Sp. Pl. ed. 4* 1 (2): 971 [July 1798] (Willdenow 1798).

Coussarea mapouriooides Bremek., *Recueil Trav. Bot. Néerl.* 31: 282 (Bremekamp 1934).



FIG. 48. — Rhabdodendraceae: **A**, *Rhabdodendron amazonicum* (Spruce ex Benth.) Huber (D. Sabatier & M.-F. Prévost 3022). Rhamnaceae: **B**, *Sarcomphalus cinnamomum* (Triana & Planch.) Hauenschild (D. Sabatier & M.-F. Prévost 2538). Rhizophoraceae **C**, *Cassipourea guianensis* Aubl. (J.-F. Molino & D. Sabatier 2844). Rosaceae: **D**, *Prunus myrtifolia* (L.) Urb. (D. Sabatier & J.-F. Molino 5527). A, B, D, © D. Sabatier/IRD; C, © J.-F. Molino/IRD.

NOTE.— Several specimens of *C. lutea* collected in the Guianas have been identified as *Coussarea hirticalyx* Standl. (1930: 175), which is the species most similar to *C. lutea*. According to Taylor (1999), *C. hirticalyx* occurs from Amazonian Colombia to Peru. Taylor & Steyermark (2004: 564) cited a specimen from the Venezuelan Guayana (*A. Fernández* 7544) having a corolla tube up to 20 mm long; they expressed doubts regarding the delimitation of the two species. However, pending a more detailed comparison between the two taxa, *C. lutea* can be distinguished from *C. hirticalyx* by its persistent stipules (vs caducous in *C. hirticalyx*), corolla tube (8.5-)10.3-13(-20) mm long (vs 33-34 mm long), and fruit 9-10 × 8-9 mm (vs c. 18 × 14 mm).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (Lectotype BM[[BM000058142](#)] designated by Delprêtre [2015: 609]).

SIZE. — Up to 10 m tall.

^[1451] *Coussarea machadoana* Standl.
(Fig. 49C)

Bull. Torrey Bot. Club 60 (6): 395 [1 June 1933] (Standley 1933).

VERNACULAR NAMES. — Pa: wakáu, uwakaa.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly* 15459.

INVENTORY DATA (FG). — 40 trees in 26 plots; $F_{\max} < 1\%$; dbh_{inv} = 20.8 cm.

^[1452] *Coussarea paniculata* (Willd.) Standl.
(Fig. 49D)

J. Wash. Acad. Sci. 18: 282 (Standley 1928). — *Froelichia paniculata* Willd., *Sp. Pl.*, ed. 4 1 (2): 607 [July 1798] (Willdenow 1798). — *Billardiera paniculata* Vahl, *Eclog. Amer.* 1: 13 (Vahl 1797), “*Billardiera*”, nom. inval. (combination not definitely accepted by the author). — *Froelichia paniculata* Vahl, *Eclog. Amer.* 3: 56 (Vahl 1807), isonym. — *Coussarea froelichia* A.Rich., *Prodr. [A. P. de Candolle]* 4: 494 [late Sep. 1830] (Richard 1830), nom. illeg. superfl. (based on *Billardiera paniculata*).

Coussarea macrocarpa A.Rich., *Prodr. [A. P. de Candolle]* 4: 494 [late Sep. 1830] (Richard 1830).

Faramea martinii DC., *Prodr. [A. P. de Candolle]* 4: 498 [late Sep. 1830] (Candolle 1830), “*Martini*”. — *Coussarea martinii* (DC.) Benth. & Hook.f. ex B.D.Jacks., *Index Kew.* 1 (1): 631 [6 Sep. 1893] (Jackson 1893), “*Martini*”.

Coussarea darienensis Steyermark., *Ceiba* 3 (1): 20 (Steyermark 1952).

Coussarea morii Dwyer, *Ann. Missouri Bot. Gard.* 67 (1): 130 (Dwyer 1980).

NOTES. — Vahl realised after the printing, but before publication of the first volume of his book, that the name *Billardiera* was already in use for another genus (*Billardiera* Sm. 1793). In his foreword, he therefore proposed the new name *Froelichia* Vahl, wherewith he invalidated all the species' names described under *Billardiera*. Furthermore, none of the new combinations in *Froelichia* were validly published in 1797, because no association was made between genus name and specific epithets. In fact, validation of Vahl's binomials occurred when the index in the last volume was published (1807), but this was after Willdenow's publication of *F. paniculata*. *Coussarea paniculata* is based on the invalid *F. paniculata* Vahl, but this is an indirect reference to the valid *F. paniculata* Willd. The descriptions and names of *C. froelichia* and *C. macrocarpa* in Candolle (Sept. 1830: 494) came from Richard (Dec. 1830: 177), the text of which Candolle had seen a preprint version. Both names are thus to be ascribed to Richard alone (not "A.Rich. ex DC.").

VERNACULAR NAMES. — Pa: mavinbi, mavinvi • Ka: kofilan, maipuli elepali • Wp: tapi'i ka'alulu.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *R.C. Ek 1745*.

INVENTORY DATA (FG). — 15 trees in 6 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 20.9 \text{ cm}$.

Genus *Coutarea* Aubl.

[1453] *Coutarea hexandra* (Jacq.) K.Schum.
(Fig. 49E)

Fl. Bras. [Martius] 6 (6): 185 [15 June 1889] (Schumann 1889). — *Portlandia hexandra* Jacq., *Enum. Syst. Pl.*: 16 [Aug.-Sep. 1760] (Jacquin 1760). — *Coutarea portlandia* Dum.Cours., *Bot. Cult. 2*: 561 [1-4 July 1802] (Dumont 1802), nom. illeg. superfl. (based on *Portlandia hexandra*).

Coutarea speciosa Aubl., *Hist. Pl. Guiane 1*: 314 [Jun.-Dec. 1775] (Aublet 1775). — *Portlandia speciosa* (Aubl.) Baill., *Hist. Pl. [Baillon] 7*: 381 [Feb. 1880] (Baillon 1880). — *Coutarea hexandra* var. *speciosa* (Aubl.) K.Schum., *Fl. Bras. [Martius] 6 (6)*: 197 [15 June 1889] (Schumann 1889).

Portlandia acuminata Willd., *Syst. Veg. [Roemer & Schultes] 5*: 23 [Dec. 1819] (Willdenow 1819).

Exostema souzanum Mart., *Reise Bras. [Spix & Mart.] 2*: 789 (Martius 1828). — *Cinchona souzana* (Mart.) Brign., *Mem. Mat. Fis. Soc. Ital. Sci. Modena, Pt. Mem. Fis. 2* (1): 63 (Brignoli 1862).

Coutarea campanilla DC., *Prodr. [A. P. de Candolle] 4*: 350 [late Sep. 1830] (Candolle 1830). — *Coutarea hexandra* var. *campanilla* (DC.) Steyermark., *Mem. New York Bot. Gard. 23*: 298 (Steyermark 1972).

Coutarea flavescens Sessé & Moc. ex DC., *Prodr. [A. P. de Candolle] 4*: 350 [late Sep. 1830] (Candolle 1830).

Coutarea pubescens Pohl, *Pl. Bras. Icon. Descr. 2 (4)*: 148 [Jan.-Sep. 1833] (Pohl 1833). — *Coutarea hexandra* var. *pubescens* (Pohl) K.Schum., *Fl. Bras. [Martius] 6 (6)*: 198 [15 June 1889] (Schumann 1889). — *Coutarea hexandra* f. *pubescens* (Pohl) Steyermark., *Mem. New York Bot. Gard. 23*: 298 (Steyermark 1972).

Bignonia triflora Pav. ex DC., *Prodr. [A. P. de Candolle] 9*: 148 [Jan. 1845] (Candolle 1845), nom. nud.

Gardenia hexagona Lem., *Ill. Hort. 16*: 92 (Lemaire 1869).

Coutarea scherffiana André, *Ill. Hort. 25*: 120 (André 1878).

Coutarea lindeniana Baill., *Adansonia [Baillon] 12*: 300 (Baillon 1879).

Coutarea hexandra f. *tarapotensis* K.Schum., *Fl. Bras. [Martius] 6 (6)*: 198 [15 June 1889] (Schumann 1889).

Coutarea hexandra var. *amazonica* K.Schum., *Fl. Bras. [Martius] 6 (6)*: 197 [15 June 1889] (Schumann 1889).

Coutarea hexandra var. *fluminensis* K.Schum., *Fl. Bras. [Martius] 6 (6)*: 197 [15 June 1889] (Schumann 1889).

Coutarea hexandra f. *albiflora* Chodat & Hassl., *Bull. Herb. Boissier, sér. 2, 4*: 92 (Chodat & Hassler 1904).

Coutarea hexandra f. *grandiflora* Chodat & Hassl., *Bull. Herb. Boissier, sér. 2, 4*: 92 (Chodat & Hassler 1904).

Coutarea hexandra f. *roseiflora* Chodat & Hassl., *Bull. Herb. Boissier, sér. 2, 4*: 92 (Chodat & Hassler 1904).

Coutarea hexandra var. *calycina* Chodat & Hassl., *Bull. Herb. Boissier, sér. 2, 4*: 92 (Chodat & Hassler 1904).

NOTE. — The name *Portlandia acuminata* Willd. is to be ascribed to Willdenow alone (Turland et al. 2018: Art. 46.3, Ex. 15).

VERNACULAR NAMES. — Pa: kwik-duwē • Ka: pakeli • Cr: quinaquina • Br: capança-vermelha, falsa-quina, quinarana.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (Neotype of *Coutarea speciosa*, P-JU No. 9927a [[P00680269](#)], designated by Delprete [2015: 603], isoneotype P-JU No. 9927b [[P00680270](#)]).

INVENTORY DATA (FG). — 1 tree, dbh = 13 cm.

Genus *Duroia* L.f.

[1454] *Duroia aquatica* (Aubl.) Bremek.

Recueil Trav. Bot. Néerl. 31: 270 (Bremekamp 1934). — *Coupouia aquatica* Aubl., *Hist. Pl. Guiane 2* (Suppl.): 16 [Jun.-Dec. 1775] (Aublet 1775). — *Cupirana aubletiana* Miers, *Apocyn. S. Am.*: 16 [May-June 1878] (Miers 1878), nom. illeg. superfl. (based on *Coupouia aquatica*).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kinuwup • Ka: atagali, atakali, yalakalu pupo • Wp: yanipa u • Wn: mutupupot • Nt: gaan maamadosu, maamadosu • Cr: konfitir-makak, mari-pwèl • Br: purui-grande.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000793078](#)]).

INVENTORY DATA (FG). — 95 trees in 58 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 19.9 \text{ cm}$.

[1455] *Duroia eriopila* L.f.

Suppl. Pl.: 209 ["1781" publ. Apr. 1782] (Linnaeus 1782). — *Genipa eriopila* (L.f.) Oken, *Allg. Naturgesch. 3 (2)*: 886 (Oken 1841). — *Amaioua eriopila* (L.f.) Baill., *Hist. Pl. [Baillon] 7*: 387 [Feb. 1880] (Baillon 1880).

Genipa merianae Rich., *Actes Soc. Hist. Nat. Paris 1*: 107 [Oct. 1792] (Richard 1792). — *Gardenia merianae* (Rich.) Poir., *Encycl. J.*

Lamarek et al.] Suppl. 2: 708 [3 July 1812] (Poiret 1812), *nom. illeg.* (*Duroia eriopila* in synonymy).

Amaioua surinamensis Steud., *Flora* 26 (45): 763 [17 Dec. 1843] (Steudel 1843), “*Amajoua*”. — *Duroia surinamensis* (Steud.) Benth. & Hook.f. ex B.D.Jacks., *Index Kew.* 1 (2): 804 [14 Dec. 1893] (Jackson 1893).

Amaioua grandifolia Miq., *Linnaea* 18: 291 [“1844” publ. Feb. (-May?) 1845] (Miquel 1845).

Amaioua affinis Miq., *Linnaea* 18: 613 [“1844” publ. prob. Aug. 1845] (Miquel 1845).

Amaioua ursina Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 8 (3): 168 (Standley 1930).

Duroia eriopila f. *glabra* Steyermark., *Mem. New York Bot. Gard.* 12 (3): 203 (Steyermark 1965).

VERNACULAR NAMES. — Pa: kinuwup • Ka: atagali, atakali, me:ku ubupo, wayamu wolekotopo • Wp: yanipa u • Cr: konfitir-makak, mari-pwél • Br: purui-grande.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *P. Grenand et al. 3442*.

INVENTORY DATA (FG). — 36 trees in 28 plots; $F_{\max} < 1\%$; $dbh_{inv} = 20$ cm.

[1456] *Duroia longiflora* Ducke

Arch. Jard. Bot. Rio de Janeiro 4: 181 (Ducke 1925).

VERNACULAR NAMES. — Pa: inam-etni-seine, kuyau-kamwi • Nt: maamadosu.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *S.A. Mori et al. 21520*.

INVENTORY DATA (FG). — 305 trees in 96 plots; $F_{\max} = 4.2\%$; $dbh_{inv} = 52.5$ cm.

[1457] *Duroia martiniana* (Miers) Bremek.

Recueil Trav. Bot. Néerl. 31: 270 (Bremekamp 1934). — *Cupirana martiniana* Miers, *Apocyn. S. Am.*: 17 [May-June 1878] (Miers 1878). — *Coupoua martiniana* (Miers) Kuntze, *Revis. Gen. Pl.* 2: 414 [5 Nov. 1891] (Kuntze 1891), “*Cupuia*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: kinuwup • Ka: atakali • Wp: yanipa u • Wn: mutupupot • Nt: maamadosu, pikin maamadosu.

HERBARIUM DATA (FG). — 49 collections at CAY. Sel. exs.: *C. Persson et al. 2069*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 11.6$ cm.

[1458] *Duroia micrantha* (Ladbr.) Zarucchi & J.H.Kirkbr.

Ann. Missouri Bot. Gard. 77 (4): 851 (Zarucchi & Kirkbride 1990). — *Coupoua micrantha* Ladbr., *J. Bot.* 58: 176 (Ladbrook 1920).

Duroia sprucei Rusby, *Descr. S. Amer. Pl.* 133 [20 Dec. 1920] (Rusby 1920).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *O. Poncy et al. 1746*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 29$ cm.

Genus *Eumachia* DC.

[1459] *Eumachia boliviiana* (Standl.) Delporte & J.H.Kirkbr.

J. Bot. Res. Inst. Texas 9: 76 (Delporte & Kirkbride 2015). — *Psychotria boliviiana* Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 7 (3): 302 (Standley 1931). — *Margaritopsis boliviiana* (Standl.) C.M.Taylor, *Syst. Geogr. Pl.* 75: 170 (Taylor 2005).

Psychotria lawrancei Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (3): 281 (Standley 1937).

Psychotria foetidiflora Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 22 (3): 202 (Standley 1940).

Psychotria kukenanensis Steyermark., *Fieldiana, Bot.* 28 (3): 601 (Steyermark 1953).

Psychotria turboensis Standl. ex Steyermark., *Acta Bot. Venez.* 4: 101. (Steyermark 1964)

Psychotria puberulenta Steyermark., *Mem. New York Bot. Gard.* 23: 488 (Steyermark 1972).

Psychotria plowmanii Steyermark., *Ann. Missouri Bot. Gard.* 71 (4): 1178 (Steyermark 1984).

Psychotria ronaldii Steyermark., *Ann. Missouri Bot. Gard.* 74 (2): 111 (Steyermark 1987).

VERNACULAR NAMES. — Pa: mavinvi-kamwi • Wp: tapi’i wapa’a.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Feillet et al. 10354*.

SIZE. — Up to 12 m tall.

Genus *Faramea* Aubl.

[1460] *Faramea corymbosa* Aubl.

Hist. Pl. Guiane 1: 102 [Jun.-Dec. 1775] (Aublet 1775).

Faramea pendula Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 29 [15-21 Aug. 1841] (Poeppig 1841).

Faramea truncata DC., *Prodr. [A. P. de Candolle]* 4: 496 [late Sep. 1830] (Candolle 1830).

VERNACULAR NAMES. — Wp: ka’apelele.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777662] designated by Lanjouw & Uittien [1940: 151]).

INVENTORY DATA (FG). — 1 tree, $dbh = 10.2$ cm.

[1461] *Faramea multiflora* A.Rich.

Prod. [A. P. de Candolle] 4: 497 [late Sep. 1830] (Richard 1830). — *Coussarea multiflora* (A.Rich.) Lemée, *Fl. Guyane Franç.* 3: 542 (Lemée 1954).

Coffea umbellata Vell., *Fl. Flumin.* 63 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), nom. illeg. hom., non Ruiz & Pav. (Ruiz & Pavón 1799).

Psychotria cyanea Vell., *Fl. Flumin.* 68 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Faramea salicifolia C.Presl, *Symb. Bot. [C. Presl]* 2: 24 [July 1834] (Presl 1834). — *Faramea multiflora* var. *salicifolia* (C.Presl) Steyermark., *Mem. New York Bot. Gard.* 17 (1): 394 (Steyermark 1967).

Faramea amazonica Müll.Arg., *Flora* 58: 470 (Müller 1875). — *Faramea multiflora* var. *amazonica* (Müll.Arg.) Steyermark., *Mem. New York Bot. Gard.* 17 (1): 392 (Steyermark 1967).

Faramea egensis Müll.Arg., *Flora* 58: 470 (Müller 1875).

Faramea parviflora Müll.Arg., *Flora* 58: 470 (Müller 1875).

Faramea salicifolia f. *paniculata* Kuntze, *Revis. Gen. Pl.* 1: 282 [5 Nov. 1891] (Kuntze 1891).

Faramea salicifolia f. *subumbellata* Kuntze, *Revis. Gen. Pl.* 1: 282 [5 Nov. 1891] (Kuntze 1891).

Faramea maynensis Spruce ex Rusby, *Bull. New York Bot. Gard.* 4 (13): 300 [25 June 1906] (Rusby 1906). — *Faramea multiflora* var. *maynensis* (Spruce ex Rusby) Steyermark., *Mem. New York Bot. Gard.* 17 (1): 393 (Steyermark 1967).

Rudgea scandens K.Krause, *Bot. Jahrb. Syst.* 40 (3): 342 [24 Jan. 1908] (Krause 1908).

Faramea benensis Rusby, *Mem. New York Bot. Gard.* 7: 379 (Rusby 1927). — *Faramea multiflora* var. *benensis* (Rusby) Steyermark., *Mem. New York Bot. Gard.* 17 (1): 394 (Steyermark 1967).

Faramea hondureae Standl., *J. Wash. Acad. Sci.* 18: 169 (Standley 1928).

Faramea stenura Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 331 [24 Oct. 1929] (Standley 1929).

Faramea talamancae Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 332 [24 Oct. 1929] (Standley 1929).

Faramea cuencana Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 7 (2): 219 (Standley 1931).

Faramea laxula K.Krause, *Notizbl. Bot. Gart. Berlin-Dahlem* 15: 392 [30 Mar. 1941] (Krause 1941).

Faramea cuatrecasasii Standl. ex Steyermark., *Acta Biol. Venez.* 4 (1): 42 (Steyermark 1964).

Faramea multiflora var. *epedunculata* Steyermark., *Mem. New York Bot. Gard.* 17 (1): 392 (Steyermark 1967).

Faramea talamancae f. *acutifolia* Dwyer, *Ann. Missouri Bot. Gard.* 54 (2): 141 [27 Oct. 1967] (Dwyer 1967).

NOTE. — The description and name of *F. multiflora* came from Richard (published in December 1830: 176), the text of which Candolle had seen a preprint version. The name is thus to be ascribed to Richard alone (not "A.Rich. ex DC.").

VERNACULAR NAMES. — Wp: kusili sī ka'a, tapi'i wapa'a sili, uwa kaya • Wn: iliwaimë • Br: chacrona.

HERBARIUM DATA (FG). — 69 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (lecto-, P[P00836846] designated by Jardim [Taylor & Jardim 2020: 121]).

SIZE. — Up to 10 m tall (Taylor *et al.* 2004).

[1462] *Faramea occidentalis* (L.) A.Rich.

Mém. Rubiac. 96 [Dec. 1830] (Richard 1830). — *Ixora occidentalis* L., *Syst. Nat.*, ed. 10, 2: 893 [7 June 1759] (Linnaeus 1759). — *Coffea occidentalis* (L.) Jacq., *Enum. Syst. Pl.*: 16 [Aug.-Sep. 1760] (Jacquin 1760). — *Tetramerium odoratissimum* C.F.Gaertn., *Suppl. Carp.*: 90 [May 1806] (Gaertner 1806), nom. illeg. superfl. (based on several previously published plates and citing several published names in synonymy, notably *Coffea occidentalis*). — *Antoniana laurifolia* Tussac, *Fl. Antill.* 2: 62 (Tussac 1818), nom. illeg. superfl. (refers to the plates of Plumier and Brown on which is based *Ixora occidentalis*). — *Tetramerium occidentale* (L.) Nees & Mart., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 13 (Nees & Martius 1824). — *Faramea odoratissima* DC., *Prod. [A. P. de Candolle]* 4: 496 [late Sep. 1830] (Candolle 1830), nom. illeg. superfl. (based on *Ixora occidentalis*). — *Coussarea odoratissima* M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.*: 28 (Gómez 1889), nom. illeg. superfl. (based on *Tetramerium odoratissimum*, i.e. on *Coffea occidentalis*).

Faramea sertulifera DC., *Prod. [A. P. de Candolle]* 4: 496 [late Sep. 1830] (Candolle 1830). — *Coussarea sertulifera* (DC.) M.Gómez, *Anales Soc. Esp. Hist. Nat.* 23: 292 (Gómez 1894).

Darluca prostrata Raf., *New Fl. [Rafinesque]* 4: 56 (Rafinesque 1838), nom. inval. (simultaneously published with *Neleixa prostrata* Raf.)

Neleixa prostrata Raf., *New Fl. [Rafinesque]* 4: 56 (Rafinesque 1838), nom. inval. (simultaneously published with *Darluca prostrata* Raf.)

Faramea tenuifolia Rusby, *Mem. New York Bot. Gard.* 7: 380 (Rusby 1927).

Faramea zetekii Standl., *Contr. Arnold Arbor.* 5: 146 (Standley 1933).

Faramea belizensis Standl., *Publ. Carnegie Inst. Wash.* 461: 90 [26 Nov. 1935] (Standley 1935).

Faramea occidentalis subsp. *lonchocalyx* Steyermark., *Mem. New York Bot. Gard.* 17 (1): 385 (Steyermark 1967).

Faramea occidentalis var. *brachycalyx* Steyermark., *Mem. New York Bot. Gard.* 17 (1): 385 (Steyermark 1967).

Faramea occidentalis var. *meridionalis* Steyermark., *Mem. New York Bot. Gard.* 17 (1): 384 (Steyermark 1967).

Faramea standleyana L.O.Williams, *Phytologia* 26 (6): 490 [11 Oct. 1973] (Williams 1973).

NOTE. — Jacquin did not explicitly cite *Ixora occidentalis* L. as the basionym of his *Coffea occidentalis*, but the use of the same epithet and iconotype ("Plum. ic. 156, f. 2") is an indirect reference to it.

VERNACULAR NAMES. — Pa: waaduk-purubumna.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand* 3778, dbh 10 cm.




FIG. 49. — Rubiaceae: **A**, *Amaioua guianensis* Aubl. (*D. Sabatier et al. 4792*); **B**, *Capirona macrophylla* (Poepp.) Delprete (*D. Sabatier 3554*); **C**, *Coussarea machadoana* Standl. (*D. Sabatier & J.-F. Molino 4839*); **D**, *Coussarea paniculata* (Willd.) Standl. (*D. Sabatier & J.-F. Molino 5307*); **E**, *Coutarea hexandra* (Jacq.) K.Schum. (*J.-F. Molino et al. 3382*). A, © M.-F. Prévost/IRD; B-D, © D. Sabatier/IRD; E, © J.-F. Molino/IRD.

[1463] *Faramea pedunculata* (Bremek.) Delprete

Blumea 51 (2): 361 (Delprete 2006). — *Faramea sessiliflora* Aubl. var. *pedunculata* Bremek., *Brittonia* 8 (4): 241 [Jan. 1957] (Bremekamp 1957).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: yanipa lali sili.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *R.S. Cowan 38808* (original material US [00129992]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 11.5$ cm.

[1464] *Faramea polytriadophora* Bremek.

Brittonia 8 (4): 241 [Jan. 1957] (Bremekamp 1957).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & E. Fonty 5590*.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 30.7$ cm.

[1465] *Faramea sessilifolia* (Kunth) DC.

Prodr. [A. P. de Candolle] 4: 497 [late Sep. 1830] (Candolle 1830). — *Tetramerium sessilifolium* Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 374 [21 Nov. 1819] (Kunth 1819). — *Ixora sessilifolia* (Kunth) Spreng., *Syst. Veg. [Sprengel]* 1: 409 ["1825" publ. late 1824] (Sprengel 1824).

Faramea longifolia Benth., *J. Bot. [Hooker]* 3: 234 (Bentham 1841).

Faramea coarinensis Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 133 [1 July 1881] (Müller 1881).

Faramea longifolia var. *petiolaris* Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 133 [1 July 1881] (Müller 1881).

Faramea planitiarum Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 22 (3)*: 187 (Standley 1940).

Faramea costata Steyermark., *Mem. New York Bot. Gard.* 17 (1): 389 (Steyermark 1967).

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville 638*.

SIZE. — Suriname. *S. Koemar & K. McDonnell 46* (MO), 15 m.

[1466] *Faramea* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino 2144*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 10.2 \text{ cm}$.

Genus *Ferdinandusa* Pohl

[1467] *Ferdinandusa goudotiana* K.Schum.

Fl. Bras. [Martius] 6 (6): 211 [15 June 1889] (Schumann 1889).

Ferdinandusa goudotiana var. *eciliata* Steyermark, *Mem. New York Bot. Gard.* 23: 281 (Steyermark 1972).

Ferdinandusa goudotiana var. *psilocarpa* Steyermark, *Mem. New York Bot. Gard.* 23: 281 (Steyermark 1972).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Loubry 1892*.

SIZE. — Up to 18 m tall (Taylor *et al.* 2007).

[1468] *Ferdinandusa paraensis* Ducke

Arch. Jard. Bot. Rio de Janeiro 3: 259 (Ducke 1922).

Ferdinandusa paraensis var. *palustris* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 259 (Ducke 1922).

VERNACULAR NAMES. — Pa: á-wakaha • Wp: wila to tawa • Br: acauá, bacabinha-quina.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *S.A. Mori et al. 21018*.

INVENTORY DATA (FG). — 75 trees in 37 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 40.9 \text{ cm}$.

[1469] *Ferdinandusa* sp. A
(Fig. 50A)

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4928*.

INVENTORY DATA (FG). — 12 trees in 3 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 16.2 \text{ cm}$.

Genus *Genipa* L.

[1470] *Genipa americana* L. var. *americana*

Syst. Nat., ed. 10, 2: 931 [7 June 1759] (Linnaeus 1759), "americ.". — *Gardenia genipa* Sw., *Prod. [Swartz]*: 51 [20 Jun.-29 July 1788] (Swartz 1788), nom. illeg. superfl. (based on *Genipa americana*). — *Genipa grandifolia* Pers., *Syn. Pl. [Persoon]* 1: 198 [1 Apr.-15 June 1805] (Persoon 1805), nom. illeg. superfl. (based on *Genipa americana*).

Genipa oblongifolia Ruiz & Pav., *Flora Peruviana* 2: 67 (Ruiz & Pavón 1799). — *Gardenia oblongifolia* (Ruiz & Pav.) Poir., *Encycl. [J. Lamarck et al.] Suppl.* 2: 708 [3 July 1812] (Poirier 1812).

Genipa humilis Vell., *Fl. Flumin.* 94 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Genipa barbata C.Presl, *Symb. Bot. [C. Presl]* 2: 13 [July 1834] (Presl 1834).

Genipa excelsa K.Krause, *Bot. Jahrb. Syst.* 40 (3): 327 [24 Jan. 1908] (Krause 1908).

Genipa codonocalyx Standl., *Contr. U.S. Natl. Herb.* 17: 446 (Standley 1914).

NOTE. — Domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: arasgu • Ka: tabulupo, tapulupo • Te: dzádupa uhu • Wp: yanipa, yanipa e'e • Wn: kupé • Nt: uman tapupa • Cr: jénipa, jinipa • Fr: génipa • Br: genipapo, jenipapo.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4201*.

SIZE. — Up to 60 cm dbh (Taylor *et al.* 2004).

[1471] *Genipa americana* var. *caruto* (Kunth) K.Schum.

Fl. Bras. [Martius] 6 (6): 352 [15 June 1889] (Schumann 1889). — *Genipa caruto* Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 407 [13 Mar. 1820] (Kunth 1820).

Genipa pubescens DC., *Prod. [A. P. de Candolle]* 4: 378 [late Sep. 1830] (Candolle 1830).

Genipa americana f. *grandifolia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 171 (Chodat & Hassler 1904).

Genipa americana f. *parvifolia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 171 (Chodat & Hassler 1904).

Genipa americana f. *jorgensenii* Steyermark, *Mem. New York Bot. Gard.* 23: 353 (Steyermark 1972).

VERNACULAR NAMES. — Br: genipapo.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *V. Hiquet 159*.

SIZE. — Guyana. *M.J. Jansen-Jacobs et al. 78* (MO), 8 m × 30 cm.

[1472] *Genipa spruceana* Steyermark.

Mem. New York Bot. Gard. 23: 353 (Steyermark 1972).

VERNACULAR NAMES. — Pa: amuwan, arasgu • Te: dzádupa'i • Wp: yanipa i • Wn: pisusuk • Nt: uman tapupa • Cr: jinipa-dilo, jinipa-rivié, ti-jinipa • Fr: petit génipa.

HERBARIUM DATA (FG). — 60 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 2032*.

SIZE. — Brazil, Pará. *C.A. Cid Ferreira 8010* (MO), 10 m × 18 cm.

Genus *Guettarda* L.

[1473] *Guettarda argentea* Lam.

Encycl. [J. Lamarck et al.] 3 (1): 54 [19 Oct. 1789] (Lamarck 1789). — *Matthiola argentea* (Lam.) Kuntze, *Revis. Gen. Pl.* 1: 288 [5 Nov. 1891] (Kuntze 1891).

Guettarda argentea var. *glabrata* Urb., *Symb. Antill. [Urban]* 6 (1): 47 [15 July 1909] (Urban 1909).

VERNACULAR NAMES. — Pa: seivan.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5027*, dbh = 46.5 cm.

Genus *Isertia* Schreb.

[1474] *Isertia coccinea* (Aubl.) J.F.Gmel.

Syst. Nat., ed. 13[bis], 2 (1): 567 [late Sep.-Nov. 1791] (Gmelin 1791). — *Guettarda coccinea* Aubl., *Hist. Pl. Guiane* 1: 317 [Jun.-Dec. 1775] (Aublet 1775). — *Duhamelia coccinea* (Aubl.) Dum. Cours., *Bot. Cult.*, ed. 2, 4: 328 (Dumont 1811). — *Phosanthus coccineus* (Aubl.) Raf., *Ann. Gen. Sci. Phys.* 6: 82 (Rafinesque 1820), “*coccinea*”.

Isertia flava Miq., *Linnaea* 18: 613 [“1844” publ. prob. Aug. 1845] (Miquel 1845).

Isertia coccinea var. *pentamera* Bremek., *Recueil Trav. Bot. Néerl.* 31: 263 (Bremekamp 1934).

VERNACULAR NAMES. — Pa: audik-amana, audika-retni, tahuma • Wp: wainimi susu • Wn: palalipanuimë • Nt: kotika uman gonbe • Cr: lanmou-rouj • Br: coralleira.

HERBARIUM DATA (FG). — 96 collections at CAY. Sel. exs.: *J.B. Aublets.s.n.* (lecto-, P[P02273439] designated by Boom [1984: 439]).

INVENTORY DATA (FG). — 14 trees in 12 plots; $F_{\max} < 1\%$; dbh_{inv} = 27.6 cm.

[1475] *Isertia hypoleuca* Benth.

J. Bot. [Hooker] 3: 220 (Bentham 1841). — *Isertia coccinea* var. *hypoleuca* (Benth.) K.Schum., *Fl. Bras. [Martius]* 6 (6): 286 [15 June 1889] (Schumann 1889).

Isertia breviflora Mart. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 825 (Steudel 1840), *nom. nud.*

Cassupa scarlatina K.Schum. & K.Krause, *Verb. Bot. Vereins Prov. Brandenburg* 50: 98 [30 Sep. 1908] (Schumann & Krause 1908). — *Isertia scarlatina* (K.Schum. & K.Krause) Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 8 (5): 346 (Standley 1931).

Isertia hoehnei K.Krause, *Arch. Bot. São Paulo* 1: 115 (Krause 1925).

VERNACULAR NAMES. — Ka: melamelulan • Wp: wainimi susu.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *L.E. Skog & C. Feillet 7440*.

SIZE. — Up to 15 m tall, to 35 cm dbh.

[1476] *Isertia parviflora* Vahl

Eclog. Amer. 2: 28 (Vahl 1798).

Brignolia acuminata DC., *Prodri. [A. P. de Candolle]* 4: 444 [late Sep. 1830] (Candolle 1830).

Brignolia pubigera Benth., *J. Bot. [Hooker]* 3: 219 (Bentham 1841).

Bruinsmania isertioides Miq., *Linnaea* 17: 73 (Miquel 1843).

Isertia glabra Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 179 (Ducke 1925).

Isertia parviflora var. *hirta* Steyermark., *Mem. New York Bot. Gard.* 17 (1): 295 (Steyermark 1967).

Yutajea liesneri Steyermark., *Ann. Missouri Bot. Gard.* 74 (3): 676 (Steyermark 1987).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *L. Andersson 1993*.

SIZE. — Up to 12 m tall, to 15 cm dbh.

[1477] *Isertia spiciformis* DC.

Prodri. [A. P. de Candolle] 4: 437 [late Sep. 1830] (Candolle 1830), “*spicaeformis*”.

Isertia commutata Miq., *Stirp. Surinam. Select.*: 172 [“1850” publ. Mar. 1851] (Miquel 1851).

Isertia bullata K.Schum., *Fl. Bras. [Martius]* 6 (6): 286 [15 June 1889] (Schumann 1889).

Isertia viscosa Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 260 (Ducke 1922).

Isertia pterantha Bremek., *Brittonia* 8 (4): 242 [Jan. 1957] (Bremekamp 1957).

VERNACULAR NAMES. — Ka: melamelulan • Wp: wainimi susu.

HERBARIUM DATA (FG). — 93 collections at CAY. Sel. exs.: *S.A. Mori et al. 26555*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 11.1 cm.

Genus *Ixora* L.

[1478] *Ixora ferrea* (Jacq.) Benth.

Linnaea 23: 447 (Bentham 1850). — *Sideroxyloides ferreum* Jacq., *Select. Stirp. Amer. Hist.*: 19 [5 Jan. 1763] (Jacquin 1763). — *Siderodendrum ferreum* (Jacq.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 242 [late Sep.-Nov. 1791] (Gmelin 1791). — *Siderodendrum ferreum* (Jacq.) Lam., *Tabl. Encycl.* 1[1 (2)]: 282 [13 Feb. 1792] (Lamarck 1792), isonym. — *Siderodendrum triflorum* Vahl, *Eclog. Amer.* 1: 10 (Vahl 1797), *nom. illeg. superfl.* (based on *Sideroxyloides ferreum*).

VERNACULAR NAMES. — Wp: walaku ki’iy.

HERBARIUM DATA (FG). — 48 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 6268*.

SIZE. — Up to 12 m tall, to 15 cm dbh.

[1479] *Ixora panurensis* Müll.Arg.

Flora 58: 454 (Müller 1875).

Ixora ulei K.Krause, *Notizbl. Königl. Bot. Gart. Berlin* 6: 205 [15 May 1914] (Krause 1914).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Larpin* 882.

INVENTORY DATA (FG). — 5 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 17.7$ cm.

[1480] *Ixora schomburgkiana* Benth.

Linnaea 23: 448 (Bentham 1850).

Siderodendrum macrophyllum Benth., *J. Bot. [Hooker]* 3: 220 (Bentham 1841), “*Siderodendron*”.

Ixora gleasonii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 344 [24 Oct. 1929] (Standley 1929).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier* 1433.

SIZE. — Up to 10 m tall, to 15 cm dbh.

Genus *Kutchubaea* Fisch. ex DC.

[1481] *Kutchubaea insignis* Fisch. ex DC.
(Fig. 50D)

Prod. [A. P. de Candolle] 4: 373 [late Sep. 1830] (Candolle 1830).

Gardenia integra A.Rich., *Mém. Rubiac.* 241 [Dec. 1830] (Richard 1830), nom. illeg. superfl. (includes original material of *Kutchubaea insignis*).

Kutchubaea laxiflora Benth. & Hook.f., *Gen. Pl. [Bentham & Hooker f.]* 2 (1): 98 [7-9 Apr. 1873] (Bentham & Hooker 1873), “*Kotchubaea*”, pro parte quoad specim. in flore, nom. inval. (genus name and epithet not associated).

NOTES. — The original description of *Gardenia integra* (Richard 1830: 241) is followed by: “*Crescit in Guianâ* (Richard). *Kutchubaea insignis* Fischer ex De Cand. annot. in herb. Mus. Paris.” The first sentence refers to *L.C. Richard s.n.* ([P00836504](#)), on whose left label Achille Richard wrote “*Gardenia integra* nob.” The second sentence is an indirect reference to another specimen at *P. J. Martin s.n.* ([P00836503](#)), which bears a label in Candolle’s hand that reads “*Kutchubaea insignis* Fisch.” The original material of *G. integra* thus comprises two collections, making lectotypification of this name necessary. *L.C. Richard s.n.*, barcode [P00836504](#), is here designated as lectotype. Also, the publication of *K. insignis* Fisch. ex DC. predates that of *G. integra*; hence the latter name is superfluous and illegitimate.

VERNACULAR NAMES. — Pa: tukuyuy-kamwi.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material P[\[P00836503\]](#)).

INVENTORY DATA (FG). — 10 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 44.2$ cm.

Genus *Ladenbergia* Klotzsch

[1482] *Ladenbergia lambertiana*

(A.Braun ex Mart.) Klotzsch

Getreue Darstell. Gew. 14 (2): t. 15 (Klotzsch 1846). — *Cinchona lambertiana* A.Braun ex Mart., *Reise Bras. [Spix & Mart.]* 3: 1286 (Martius 1831). — *Remijia lambertiana* (A.Braun ex Mart.) Wedd., *Ann. Sci. Nat., Bot. sér. 3, 10:* 13 [Dec. 1848] (Weddell 1848). — *Cascarilla lambertiana* (A.Braun ex Mart.) Wedd., *Hist. Nat. Quinquinas:* 85 (Weddell 1849). — *Buena lambertiana* (A.Braun ex Mart.) Wedd., *J. Linn. Soc., Bot.* 11: 187 [“1871” publ. Dec. 1869] (Weddell 1869).

Ladenbergia schomburgkii Klotzsch, *Getreue Darstell. Gew.* 14 (2): t. 15 (Klotzsch 1846). — *Cascarilla schomburgkii* (Klotzsch) Triana, *Revista Acad. Colomb. Ci. Exact.* 2: 410 (Triana 1938).

Ladenbergia lucens Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 7 (4): 362 (Standley 1931).

Ladenbergia puberula Steyermark., *Mem. New York Bot. Gard.* 23: 270 (Steyermark 1972).

Ladenbergia venamoensis Steyermark., *Mem. New York Bot. Gard.* 23: 271 (Steyermark 1972).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville & F. Crozier* 17010.

SIZE. — Up to 20 m tall (Steyermark 1974).

Genus *Morinda* L.

[1483] *Morinda citrifolia* L.

Sp. Pl. 1: 176 [1 May 1753] (Linnaeus 1753), nom. cons. — *Samama citrifolia* (L.) Kuntze, *Revis. Gen. Pl.* 1: 297 [5 Nov. 1891] (Kuntze 1891).

Platanocephalus orientalis Crantz, *Inst. Rei Herb.* 1: 349 [Jan.-July 1766] (Crantz 1766).

Morinda tinctoria Noronha, *Verh. Batav. Genootsch. Kunst.* 5 (Art. 4): 20 (Noronha 1790), nom. nud.

Morinda tomentosa B.Heyne ex Roth, *Syst. Veg. [Roemer & Schultes]* 5: 216 [Dec. 1819] (Roth 1819).

Morinda angustifolia Roth, *Nov. Pl. Sp.* 147 [Apr. 1821] (Roth 1821).

Morinda chacha Buch.-Ham., *Trans. Linn. Soc. London* 13 (2): 536 (Buchanan-Hamilton 1822).

Morinda nodosa Buch.-Ham., *Trans. Linn. Soc. London* 13 (2): 537 (Buchanan-Hamilton 1822).

Morinda mudia Buch.-Ham., *Trans. Linn. Soc. London* 13 (2): 536 (Buchanan-Hamilton 1822).

Morinda multiflora Roxb., *Fl. Ind.* 2: 200 (Roxburgh 1824).

Morinda stenophylla Spreng., *Syst. Veg. [Sprengel]* 1: 749 [1825 publ. late 1824] (Sprengel 1824). — *Morinda coreia* Buch.-Ham. var. *stenophylla* (Spreng.) Chandrab., *Fl. Coimbatore [M. Chandrabose & N.C. Nair]*: xxviii, 142 (Chandrabose 1988).

Morinda macrophylla Desf., *Tabl. École Bot.*, ed. 3 [*Cat. Pl. Horti Paris.*], 404 (Desfontaines 1829).

Morinda aspera Wight & Arn., *Prod. Fl. Ind. Orient.* 1: 420 [10 Oct. 1834] (Wight & Arnott 1834). — *Morinda pubescens* Sm. var. *aspera* (Wight & Arn.) M.Gangop., *Nelumbo* 59 (1): 24 (Gangopadhyay 2017).

Morinda quadrangularis G.Don, *Gen. Hist.* 3: 545 [8-15 Nov. 1834] (Don 1834).

Morinda ligulata Blanco, *Fl. Filip. [F.M. Blanco]*, ed. 2: 105 (Blanco 1845).

Morinda littoralis Blanco, *Fl. Filip. [F.M. Blanco]*, ed. 2: 109 (Blanco 1845), “*litoralis*”.

Morinda teysmanniana Miq., *Fl. Ned. Ind.* 2: 244 (Miquel 1857).

Morinda zollingeriana Miq., *Fl. Ned. Ind.* 2: 243 (Miquel 1857).

Sarcocephalus leichhardtii F.Muell., *Essay Pl. Burdekin*: 12 (Mueller 1860).

Morinda citrifolia L. f. *potteri* H.St.John, *Bull. Torrey Bot. Club* 111 (4): 482 (St.John 1984).

NOTE. — A species native to India, West Indies and Australia, cultivated and naturalised throughout the tropics. Cultivated and naturalised in Central America, the Greater Antilles, and in the Guianas; several naturalised populations have recently been observed in French Guiana (Delprete, pers. obs.).

VERNACULAR NAMES. — Fr: noni.

HERBARIUM DATA (FG). — 5 collections. Sel. exs.: *M.-F. Prévost* 4686.

SIZE. — Up to 12 m tall, to 15 cm dbh.

Genus *Pagamea* Aubl.

[1484] *Pagamea guianensis* Aubl.

Hist. Pl. Guiane 1: 113 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

Pagamea guianensis var. *angustifolia* Progel, *Fl. Bras. [Martius]* 6 (1): 285 [1 Aug. 1868] (Progel 1868).

Pagamea guianensis var. *parviflora* Spruce ex Progel, *Fl. Bras. [Martius]* 6 (1): 285 [1 Aug. 1868] (Progel 1868).

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777682] designated by Lanjouw & Uittien [1940: 152]); *P.G. Delprete & F. Crozier* 7150.

INVENTORY DATA (FG). — 11 trees in 3 plots; $F_{\max} = 1.4\%$; $dbh_{inv} = 13.8$ cm.

[1485] *Pagamea thrysiflora* Spruce ex Benth.

J. Proc. Linn. Soc., Bot. 1: 110 [“1857” publ. 1856] (Bentham 1856).

HERBARIUM DATA (FG). — A single collection at CAY, *J.-J. de Granville* 13833.

SIZE. — Up to 10 m tall.

Genus *Palicourea* Aubl.

[1486] *Palicourea brachyloba* (Müll.Arg.) B.M.Boom

Brittonia 37 (2): 209 (Boom 1985). — *Psychotria brachyloba* Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 243 [1 July 1881] (Müller 1881). — *Uragoga brachyloba* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 959 [5 Nov. 1891] (Kuntze 1891).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville* B-5356.

SIZE. — Up to 12 m tall (Boom & Delprete 2002).

[1487] *Palicourea guianensis* Aubl.

Hist. Pl. Guiane 1: 173 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Stephanium guianense* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 368 [late Sep.-Nov. 1791] (Gmelin 1791). — *Psychotria palicourea* Sw., *Fl. Ind. Occid.* 1: 433 [Nov. 1797] (Swartz 1797), nom. illeg. superfl. (based on *Palicourea guianensis*). — *Simira palicourea* Poir., *Encycl. [J. Lamarck et al.]* 7: 196 [6 July 1806] (Poirier 1806), nom. illeg. (based on the illegitimate *Psychotria palicourea*). — *Uragoga palicourea* Kuntze, *Revis. Gen. Pl.* 1: 300 [5 Nov. 1891] (Kuntze 1891), nom. illeg. superfl. (based on *Palicourea guianensis*).

Palicourea barbinervia DC., *Prod. [A. P. de Candolle]* 4: 530 [late Sep. 1830] (Candolle 1830). — *Uragoga barbinervia* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 959 [5 Nov. 1891] (Kuntze 1891). — *Palicourea guianensis* subsp. *barbinervia* (DC.) Steyermark., *Mem. New York Bot. Gard.* 23: 730 (Steyermark 1972).

Psychotria lutea Spreng. ex DC., *Prod. [A. P. de Candolle]* 4: 530 [late Sep. 1830] (Candolle 1830), nom. nud. pro syn.

Psychotria cataractarum Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 226 [1 July 1881] (Müller 1881). — *Nonatelia cataractarum* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 1: 291 [5 Nov. 1891] (Kuntze 1891).

Palicourea guianensis var. *tetramera* Bremek., *Recueil Trav. Bot. Néerl.* 31: 283 (Bremekamp 1934).

Palicourea guianensis var. *trimera* Bremek., *Recueil Trav. Bot. Néerl.* 31: 283 (Bremekamp 1934).

Palicourea guianensis f. *glabra* Steyermark., *Mem. New York Bot. Gard.* 23: 730 (Steyermark 1972).

Palicourea guianensis subsp. *occidentalis* Steyermark., *Mem. New York Bot. Gard.* 23: 729 (Steyermark 1972).

Palicourea guianensis var. *glabrescens* Steyermark., *Mem. New York Bot. Gard.* 23: 730 (Steyermark 1972).

VERNACULAR NAMES. — Pa: audik-amana, audika-retni • Ka: maipuli elepali, maipyuli kiyelapole, matula, padula, patula • Wp: tewitulū u, tewitulū wate'e, tewitutū wate'e • Nt: ayee wato, ayuwato • Cr: bwa-fizi • Br: erva-de-rato.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00778113] designated by Lanjouw & Uittien [1940: 155]).

INVENTORY DATA (FG). — 29 trees in 17 plots; $F_{\max} = 2.7\%$; $dbh_{inv} = 31.5$ cm.

Genus *Posoqueria* Aubl.

[1488] *Posoqueria gracilis* (Rudge) Roem. & Schult.

Syst. Veg. [Roemer & Schultes] 5: 227 [Dec. 1819] (Roemer & Schultes 1819). — *Solena gracilis* Rudge, *Pl. Guian. [Rudge] 1 (3):* 27 [June 1805] (Rudge 1805). — *Tocoyena gracilis* (Rudge) A.Rich., *Mém. Rubiac.*: 167 [Dec. 1830] (Richard 1830). — *Posoqueria latifolia* subsp. *gracilis* (Rudge) Steyermark, *Mem. New York Bot. Gard.* 17 (1): 327 (Steyermark 1967).

NOTE. — Even though volume 5 of the *Systema vegetabilium* was published shortly after Roemer's death, Schultes' foreword clearly confirms Roemer's prominent role. Therefore, contrary to Stafleu & Cowan's recommendation (1983: TL-2-9408), most names should be ascribed to Roemer and Schultes, not Schultes alone.

VERNACULAR NAMES. — Wp: aliwa.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *E. Rudge s.n.* (original material BM[[BM001008906](#)]).

SIZE. — Guyana. *P. Mutchnick* 836 (MO), 20 m.

[1489] *Posoqueria latifolia* (Rudge) Roem. & Schult.

Syst. Veg. [Roemer & Schultes] 5: 227 [Dec. 1819] (Roemer & Schultes 1819). — *Solena latifolia* Rudge, *Pl. Guian. [Rudge] 1 (3):* 26 [June 1805] (Rudge 1805). — *Ucriana insignis* Willd. ex Spreng., *Syst. Veg. [Sprengel] 1:* 761 ["1825" publ. late 1824] (Sprengel 1824), nom. illeg. superfl. (based on *Tocoyena latifolia*).

Tocoyena speciosa Rich., *Actes Soc. Hist. Nat. Paris* 1: 107 [Oct. 1792] (Richard 1792).

Tocoyena latifolia Lam., *Tabl. Encycl. 2[5 (2):* 259 [31 Oct. 1819] (Lamarck 1819).

Tocoyena insignis Roem. & Schult., *Syst. Veg. [Roemer & Schultes] 5:* 229 [Dec. 1819] (Roemer & Schultes 1819). — *Posoqueria insignis* (Roem. & Schult.) Nees, *Flora 4 (1):* 301 [21 May 1821] (Nees 1821).

Tocoyena longifolia Kunth, *Nova genera et species plantarum [H.B.K.] 3:* 411 [13 Mar. 1820] (Kunth 1820). — *Ucriana humboldtii* Spreng., *Syst. Veg. [Sprengel] 1:* 761 ["1825" publ. late 1824] (Sprengel 1824), nom. illeg. superfl. (based on *Tocoyena longifolia*).

Tocoyena mutisii Kunth, *Nova genera et species plantarum [H.B.K.] 3:* 411 [13 Mar. 1820] (Kunth 1820). — *Ucriana mutisii* (Kunth) Spreng., *Syst. Veg. [Sprengel] 1:* 761 ["1825" publ. late 1824] (Sprengel 1824).

Tocoyena macrophylla Kunth, *Nova genera et species plantarum [H.B.K.] 3:* 412 [13 Mar. 1820] (Kunth 1820). — *Ucriana macrophylla* (Kunth) D.Dietr., *Syn. Pl. [D. Dietrich] 1:* 798 [July 1839] (Dietrich 1839). — *Posoqueria macrophylla* (Kunth) Hemsl., *Biol. Cent.-Amer. Bot. 2 (7):* 39 [Apr. 1881] (Hemsley 1881).

Posoqueria revoluta Schrad., *Gött. Gel. Anz. 2:* 714 [5 May 1821] (Schradner 1821). — *Solena revoluta* (Schrad.) D.Dietr., *Syn. Pl. [D. Dietrich] 1:* 799 [July 1839] (Dietrich 1839).

Gardenia suaveolens Vell., *Fl. Flumin.:* 102 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829), "suave-olens".

Posoqueria decora DC., *Prodr. [A. P. de Candolle] 4:* 375 [late Sep. 1830] (Candolle 1830). — *Solena decora* (DC.) D.Dietr., *Syn. Pl. [D. Dietrich] 1:* 799 [July 1839] (Dietrich 1839).

Posoqueria lucida Mart., *Flora 24 (2, Beibl.):* 77 (Martius 1841).

Posoqueria macropus Mart., *Flora 24 (2, Beibl.):* 79 (Martius 1841).

Stannia panamensis Walp. & Duchass., *Linnaea 23:* 755 ["1850" publ. Jan. 1851] (Walpers & Duchassaing 1851). — *Posoqueria panamensis* (Walp. & Duchass.) Walp., *Ann. Bot. Syst. [Walpers] 2 (5):* 797 [21-24 Apr. 1852] (Walpers 1852).

Stannia metensis H.Karst., *Linnaea 28:* 441 (Karsten 1857). — *Posoqueria metensis* (H.Karst.) R.Knuth, *Repert. Spec. Nov. Regni Veg. Beih. 43:* 667 (Knuth 1928).

Stannia grandiflora H.Karst., *Fl. Columb. [H. Karsten] 1 (1):* 31 [6 Apr. 1859] (Karsten 1859). — *Posoqueria panamensis* subsp. *grandiflora* (H.Karst.) Steyermark, *Mem. New York Bot. Gard.* 17 (1): 325 (Steyermark 1967).

Posoqueria multiflora Lem., *Ill. Hort. 16:* t. 597 (Lemaire 1869), nom. illeg. hom., non (Willd.) Blume (1826).

Posoqueria spraguei Wernham, *Bull. Misc. Inform. Kew 1914 (2):* 66 [30 Mar. 1914] (Wernham 1914).

Oxyanthus isthmia W.Mill. & L.H.Bailey, *Stand. Cycl. Hortic. 4:* 2419 [23 Feb. 1916] (Miller & Bailey 1916).

Posoqueria mutisii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 11 (5):* 232 (Standley 1936).

NOTE. — See note under *Posoqueria gracilis* as to Roemer and Schultes authority.

VERNACULAR NAMES. — Pa: mavinvi-kamwi, mbibi-washiune, pune-etni-kamwi • Ka: kuyaken elepali, pipya tamili • Wp: aliwa so áta, aliwa so u, aliwa u • Wn: meku susu • Br: açucena-do-mato, lirio-do-mato.

HERBARIUM DATA (FG). — 96 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[[BM001008911](#)]).

INVENTORY DATA (FG). — 329 trees in 102 plots; $F_{\max} = 2.8\%$; $dbh_{\text{inv}} = 31.8$ cm.

[1490] *Posoqueria longiflora* Aubl.
(Fig. 50B)

Hist. Pl. Guiane 1: 134 [Jun.-Dec. 1775] (Aublet 1775). — *Kyrtanthus longiflorus* (Aubl.) J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1):* 362 [late Sep.-Nov. 1791] (Gmelin 1791). — *Solena longiflora* (Aubl.) Willd., *Sp. Pl., ed. 4 1 (2):* 961 [July 1798] (Willdenow 1798).

Willdenovia schreberi J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1):* 362 [late Sep.-Nov. 1791] (Gmelin 1791).

Tocoyena undulatifolia A.Rich., *Mém. Rubiac.* 167 [Dec. 1830] (Richard 1830).

Posoqueria acuminata Mart., *Flora 24 (2, Beibl.):* 79 (Martius 1841).

VERNACULAR NAMES. — Pa: inuva-kamwi • Ka: aimyala posowepo, kapaya wati • Te: aliwa • Wp: aliwa • Wn: asisi, meku susu, munpatutpë • Nt: agu siton • Cr: grènn-koumarou • Br: açucena-do-mato, papa-terra.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[[P00777902](#)] designated by Lanjouw & Uittien [1940: 156]).

SIZE. — Up to 10 m tall (Taylor *et al.* 2004).

Genus *Psychotria* L.

[1491] *Psychotria anceps* Kunth

Nova genera et species plantarum [H.B.K.] 3: 360 [21 Nov. 1819] (Kunth 1819). — *Uragoga anceps* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 959 [5 Nov. 1891] (Kuntze 1891).

Psychotria salicifolia Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 356 [21 Nov. 1819] (Kunth 1819). — *Uragoga salicifolia* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 962 [5 Nov. 1891] (Kuntze 1891).

Psychotria lucida Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 361 [21 Nov. 1819] (Kunth 1819).

Psychotria viburnoides Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 361 [21 Nov. 1819] (Kunth 1819). — *Uragoga viburnoides* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 963 [5 Nov. 1891] (Kuntze 1891), “viburnodes”.

Psychotria patrisii DC., *Prodr. [A. P. de Candolle]* 4: 510 [late Sep. 1830] (Candolle 1830). — *Uragoga patrisii* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 962 [5 Nov. 1891] (Kuntze 1891). — *Mapouria patrisii* (DC.) Lemée, *Fl. Guyane Franç.* 3: 549 (Lemée 1954).

Psychotria chlorantha Benth., *J. Bot. [Hooker]* 3: 225 (Bentham 1841). — *Uragoga chlorantha* (Benth.) Kuntze, *Revis. Gen. Pl.* 2: 960 [5 Nov. 1891] (Kuntze 1891).

Mapouria corymbifera Müll.Arg., *Flora* 59: 464 (Müller 1876). — *Uragoga corymbifera* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 960 [5 Nov. 1891] (Kuntze 1891).

Mapouria corumbensis S. Moore, *Trans. Linn. Soc. London, Bot.* 4 (3): 380 [“1894–96” publ. Dec. 1895] (Moore 1895). — *Psychotria corumbensis* (S. Moore) Hoehne, *Ind. Bibl. Pl. Col. Com. Rondon*: 387 (Hoehne 1951).

Psychotria familiarifolia Wernham, *J. Bot.* 55: 253 (Wernham 1917).

Mapouria biacuminata Rusby, *Descr. S. Amer. Pl.* 136 [20 Dec. 1920] (Rusby 1920).

Mapouria ornithophila Bremek., *Recueil Trav. Bot. Néerl.* 31: 288 (Bremekamp 1934).

Psychotria paracatuensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (5): 245 (Standley 1936).

Psychotria anceps var. *robustior* Steyermark., *Mem. New York Bot. Gard.* 23: 466 (Steyermark 1972).

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Loubry* 2413.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.4$ cm.

[1492] *Psychotria carthagenaensis* Jacq.

Enum. Syst. Pl.: 16 [Aug.–Sep. 1760] (Jacquin 1760). — *Uragoga carthagenaensis* (Jacq.) Kuntze, *Revis. Gen. Pl.* 2: 959 [5 Nov. 1891] (Kuntze 1891).

Psychotria ardisiifolia Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 359 [21 Nov. 1819] (Kunth 1819), “ardisiifolia”.

Psychotria floribunda Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 359 [21 Nov. 1819] (Kunth 1819). — *Uragoga floribunda* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 960 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: pune-etni-seine • Ka: akami ekunali • Wp: tāngala yiki, tāngala yiki sili, tapi'i ka'alulu sili, uwa ki'iy • Br: rainha.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *M.-F. Prévost* 1106.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 14.4$ cm.

[1493] *Psychotria ficigemma* DC.

Prodr. [A. P. de Candolle] 4: 510 [late Sep. 1830] (Candolle 1830). — *Uragoga ficigemma* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 960 [5 Nov. 1891] (Kuntze 1891). — *Mapouria ficigemma* (DC.) Lemée, *Fl. Guyane Franç.* 3: 549 (Lemée 1954).

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (original material G-DC, not seen); *P.G. Delprete & F. Crozier* 7127.

INVENTORY DATA (FG). — 44 trees in 22 plots; $F_{\max} = 2.1\%$; $dbh_{inv} = 49.6$ cm.

[1494] *Psychotria guianensis* (Aubl.) Clos

Mém. Acad. Sci. Toulouse, sér. 7, 10: 217 (Clos 1878). — *Mapouria guianensis* Aubl., *Hist. Pl. Guiane* 1: 175 [Jun.–Dec. 1775] (Aublet 1775), “Guyannensis” on plate. — *Psychotria nitida* Willd., *Sp. Pl.*, ed. 4 1 (2): 963 [July 1798] (Willdenow 1798), nom. illeg. superfl. (based on *Mapouria guianensis*). — *Simira nitida* Poir., *Encycl. J. Lamarck et al.* 7: 196 [6 July 1806] (Poiret 1806), nom. illeg. (based on the illegitimate *Psychotria nitida*). — *Psychotria mapouria* Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 187 [Dec. 1819] (Roemer & Schultes 1819), “*Mapuria*”, nom. illeg. (based on the illegitimate *Psychotria nitida*). — *Uragoga mapouria* Kuntze, *Revis. Gen. Pl.* 1: 300 [5 Nov. 1891] (Kuntze 1891), nom. illeg. superfl. (based on *Mapouria guianensis*).

Psychotria magnoliifolia Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 360 [21 Nov. 1819] (Kunth 1819), “*magnoliaefolia*”. — *Uragoga magnoliifolia* (Kunth) Kuntze, *Revis. Gen. Pl.* 2: 961 [5 Nov. 1891] (Kuntze 1891), “*magnoliaefolia*”.

Psychotria magnoliifolia Willd. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 190 [Dec. 1819] (Roemer & Schultes 1819), “*magnoliaefolia*”. — *Psychotria willdenowii* DC., *Prodr. [A. P. de Candolle]* 4: 505 [late Sep. 1830] (Candolle 1830).

Palicourea elliptica Kunth, *Nova genera et species plantarum* [H.B.K.] 3: 369 [21 Nov. 1819] (Kunth 1819).

Psychotria sambucina Link ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 188 [Dec. 1819] (Roemer & Schultes 1819). — *Mapouria sambucina* (Link ex Roem. & Schult.) Müll.Arg., *Flora* 59: 458 (Müller 1876). — *Uragoga sambucina* (Link ex Roem. & Schult.) Kuntze, *Revis. Gen. Pl.* 2: 962 [5 Nov. 1891] (Kuntze 1891).

Psychotria elliptica Humb. & Bonpl. ex Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 189 [Dec. 1819] (Roemer & Schultes 1819). — *Uragoga elliptica* (Humb. & Bonpl. ex Roem. & Schult.) Kuntze, *Revis. Gen. Pl.* 1: 300 [5 Nov. 1891] (Kuntze 1891).

Mapouria myriantha Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 400 [1 July 1881] (Müller 1881). — *Uragoga myriantha* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 961 [5 Nov. 1891] (Kuntze 1891).

Psychotria ernestii K.Krause, *Verh. Bot. Vereins Prov. Brandenburg* 50: 109 [30 Sep. 1908] (Krause 1908), "Ernestii".

Psychotria mathewsi Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 342 [24 Oct. 1929] (Standley 1929).

VERNACULAR NAMES. — Pa: pune-etni-seine • Ka: kuyaken elepali • Te: tapi'i ka'a lili • Wp: tapi'i ka'alulu • Wn: talepëpë • Nt: man agu siton • Br: apurú.

HERBARIUM DATA (FG). — 154 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00778114] designated by Lanjouw & Uittien [1940: 153]).

SIZE. — Up to 25 m tall (Boom & Delprete 2002).

[1495] *Psychotria pedunculosa* Rich.

Actes Soc. Hist. Nat. Paris 1: 107 [Oct. 1792] (Richard 1792). — *Uragoga pedunculosa* (Rich.) Kuntze, *Revis. Gen. Pl.* 2: 962 [5 Nov. 1891] (Kuntze 1891).

Psychotria mapouriooides DC., *Prodr. [A. P. de Candolle]* 4: 509 [late Sep. 1830] (Candolle 1830).

Palicourea pedunculosa DC., *Prodr. [A. P. de Candolle]* 4: 526 [late Sep. 1830] (Candolle 1830).

Palicourea chionantha DC., *Prodr. [A. P. de Candolle]* 4: 526 [late Sep. 1830] (Candolle 1830). — *Mapouria chionantha* (DC.) Müll.Arg., *Fl. Bras. [Martius]* 6 (5): 387 [1 July 1881] (Müller 1881). — *Psychotria chionantha* (DC.) Britton, *Bull. Torrey Bot. Club* 18 (4): 109 (Britton 1891). — *Uragoga chionantha* (DC.) Kuntze, *Revis. Gen. Pl.* 2: 959 [5 Nov. 1891] (Kuntze 1891). — *Psychotria mapouriooides* var. *chionantha* (DC.) Steyermark., *Mem. New York Bot. Gard.* 23: 462 (Steyermark 1972).

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *J.B. Leblond* 319 (original material G[G00341845]).

SIZE. — Brazil. *S.A. Mori & T.S. dos Santos* 10131 (MO), 16 m × 20 cm.

Genus *Randia* L.

[1496] *Randia armata* (Sw.) DC.

Prodr. [A. P. de Candolle] 4: 387 [late Sep. 1830] (Candolle 1830). — *Gardenia armata* Sw., *Prodr. [Swartz]*: 51 [20 Jun.-29 July 1788] (Swartz 1788). — *Solena armata* (Sw.) D.Dietr., *Syn. Pl. [D. Dietrich]* 1: 800 [July 1839] (Dietrich 1839).

Mussaenda spinosa Jacq., *Enum. Syst. Pl.*: 16 [Aug.-Sep. 1760] (Jacquin 1760). — *Gardenia tetricantha* Lam., *Encycl. J.J. Lamarck et al.* 2 (2): 609 [14 Apr. 1788] (Lamarck 1788), nom. illeg. superfl. (based on *Mussaenda spinosa*). — *Randia spinosa* (Jacq.) H.Karst., *Fl. Columb. [H. Karsten]* 2 (4): 128 [28 Nov. 1866] (Karsten 1866), nom. illeg. hom., non (Thunb.) Poir. (Poirier 1812). — *Basanacantha spinosa* (Jacq.) K.Schum., *Fl. Bras. [Martius]* 6 (6): 376 [15 June 1889] (Schumann 1889). — *Basanacantha spinosa* var. *typica* K.Schum., *Fl. Bras. [Martius]* 6 (6): 377 [15 June 1889] (Schumann 1889), nom. inval. (Turland et al. 2018: Art. 24.3).

Randia ovata Duchass. ex Griseb., *Bonplandia* 6 (1): 8 (Grisebach 1858).

Randia guianensis Sagot ex K.Schum., *Fl. Bras. [Martius]* 6 (6): 377 [15 June 1889] (Schumann 1889), nom. nud. pro syn.

Basanacantha spinosa var. *polyantha* K.Schum., *Fl. Bras. [Martius]* 6 (6): 377 [15 June 1889] (Schumann 1889).

Basanacantha spinosa f. *pubiloba* K.Schum., *Fl. Bras. [Martius]* 6 (6): 377 [15 June 1889] (Schumann 1889).

Basanacantha spinosa var. *longipedunculata* Kuntze, *Revis. Gen. Pl.* 3 (3): 118 [28 Sep. 1898] (Kuntze 1898).

Basanacantha spinosa f. *grandiflora* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 173 (Chodat & Hassler 1904).

Basanacantha spinosa f. *puberula* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 173 (Chodat & Hassler 1904).

Basanacantha spinosa var. *macrocalyx* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 173 (Chodat & Hassler 1904).

Basanacantha phyllosepala Sprague & R.O.Williams ex R.O.Williams & Cheesman, *Fl. Trinidad & Tobago* 2 (1): 22 [Aug. 1928] (Williams & Cheesman 1928).

VERNACULAR NAMES. — Ka: alemikilan • Wp: alamakulu alami • Br: espinho-de-judeu, esporão-de-galo.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2250.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 11.5 cm.

Genus *Rudgea* Salisb.

[1497] *Rudgea crassiloba* (Benth.) B.L.Rob.

Proc. Amer. Acad. Arts 45: 408 (Robinson 1910). — *Coffea crassiloba* Benth., *J. Bot. [Hooker]* 3: 233 (Bentham 1841).

Rudgea schomburgkiana Benth., *Linnaea* 23: 459 (Bentham 1850), nom. illeg. superfl. (based on *Coffea crassiloba* Benth.)

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-J. de Granville* 5909.

SIZE. — Brazil, Pará. *T.C. Plowman* 9069 (MO), 7 m × 10 cm.

[1498] *Rudgea graciliflora* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 11 (5): 262 (Standley 1936).

Rudgea duckei Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 11 (5): 261 (Standley 1936).

Rudgea standleyana Steyermark., *Mem. New York Bot. Gard.* 17 (1): 409 (Steyermark 1967).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S. Gonzalez* 2334.

SIZE. — Brazil, Amazonas. *S.A. Mori* 20612 (MO), 20 m × 10 cm.

[1499] *Rudgea lanceifolia* Salisb.
(Fig. 50C)

Trans. Linn. Soc. London 8: 327 (Salisbury 1807), “*lanceaefolia*”. — *Uragoga lanceifolia* (Salisb.) Kuntze, *Revis. Gen. Pl.* 2: 961 [5 Nov. 1891] (Kuntze 1891), “*lanceaefolia*”.

Rudgea ovalifolia Salisb., *Trans. Linn. Soc. London* 8: 328 (Salisbury 1807). — *Uragoga ovalifolia* (Salisb.) Kuntze, *Revis. Gen. Pl.* 2: 961 [5 Nov. 1891] (Kuntze 1891).

Rudgea fissistipula Müll.Arg., *Flora* 59: 449 (Müller 1876). — *Uragoga fissistipula* (Müll.Arg.) Kuntze, *Revis. Gen. Pl.* 2: 960 [5 Nov. 1891] (Kuntze 1891).

Rudgea carolina Standl. & Steyerl., *Fieldiana, Bot.* 28 (3): 614 (Standley & Steyermark 1953).

Rudgea dasyantha Bremek., *Brittonia* 8 (4): 244 [Jan. 1957] (Bremekamp 1957).

Rudgea sipapoensis Steyerl., *Mem. New York Bot. Gard.* 17 (1): 411 (Steyermark 1967).

Rudgea oldemanii Steyerl., *Brittonia* 33 (3): 397 (Steyermark 1981).

Rudgea prancei Steyerl., *Brittonia* 33 (3): 398 (Steyermark 1981).

VERNACULAR NAMES. — Pa: pune-etni-kamwi.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[BM000582180, BM000582181]).

INVENTORY DATA (FG). — 9 trees in 3 plots; $F_{\max} < 1\%$; dbh_{inv} = 13.8 cm.

Genus *Simira* Aubl.

[1500] *Simira tinctoria* Aubl.

Hist. Pl. Guiane 1: 170 [Jun.-Dec. 1775] (Aublet 1775). — *Psychotria tinctoria* (Aubl.) Raeusch., *Nomencl. Bot. [Raeusch.]*, ed. 3: 55 (Raeuschel 1797). — *Psychotria parviflora* Willd., *Sp. Pl.*, ed. 4 1 (2): 962 [July 1798] (Willdenow 1798), nom. illeg. superfl. (based on *Simira tinctoria*). — *Psychotria simira* Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 5: 187 [Dec. 1819] (Roemer & Schultes 1819), nom. illeg. superfl. (based on *Simira tinctoria*). — *Mapouria simira* A.Rich., *Mém. Rubiac.*: 94 [Dec. 1830] (Richard 1830), nom. illeg. superfl. (based on *Simira tinctoria*). — *Uragoga simira* Baill., *Hist. Pl. [Baillon]* 7: 374 [Feb. 1880] (Baillon 1880), nom. illeg. superfl. (based on *Simira tinctoria*). — *Uragoga tinctoria* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 958 [5 Nov. 1891] (Kuntze 1891). — *Sickingia tinctoria* (Aubl.) Lemée, *Fl. Guyane Franç.* 3: 505 (Lemée 1954), nom. illeg. hom., non (Willd. ex Schult.) K.Schum. (Schumann 1888).

Chimarrhis clausicorollata J.H.Kirkbr., *BioLlania, ed. espec.* 6: 394 (Kirkbride 1997).

VERNACULAR NAMES. — Pa: adamna • Ka: simira (*fide* Aublet 1775) • Wn: kujeimë, kusi wewe • Nt: kenkii udu • Br: arariba.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00778112] designated by Lanjouw & Uitien [1940: 156]); *D. Sabatier & M.-F. Prévost 3400*, dbh = 35 cm.

Genus *Stenostomum* C.F.Gaertn.

[1501] *Stenostomum acreanum*

(K.Krause) Achille & Delprete

Blumea 55 (2): 167 [17 Aug. 2010] (Achille & Delprete 2010). — *Stenostomum acreanum* (K.Krause) C.M.Taylor, *Novon* 20 (3): 360 [13 Sep. 2010] (Taylor 2010), comb. superfl. — *Guettarda acreana* K.Krause, *Notizbl. Königl. Bot. Gart. Berlin* 6: 204 [15 May 1914] (Krause 1914).

Antirhea panamensis Standl., *N. Amer. Fl.* 32 (4): 264 [21 Dec. 1934] (Standley 1934). — *Chomelia panamensis* (Standl.) Dwyer, *Ann. Missouri Bot. Gard.* 67 (1): 100 (Dwyer 1980).

Antirhea surinamensis Bremek., *Acta Bot. Neerl.* 8: 479 (Bremekamp 1959).

Chomelia chambersii Dwyer & M.V.Hayden, *Ann. Missouri Bot. Gard.* 54 (2): 138 [27 Oct. 1967] (Dwyer & Hayden 1967). — *Chione chambersii* Dwyer, *Ann. Missouri Bot. Gard.* 67 (1): 100 (Dwyer 1980).

Guettarda leiantha Steyerl., *Ann. Missouri Bot. Gard.* 71 (4): 1175 (Steyermark 1984).

NOTE. — *Chione chambersii* is a mispell of *Chomelia chambersii* in the synonymy of *Chomelia panamensis*.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 14691*.

INVENTORY DATA (FG). — 39 trees in 12 plots; $F_{\max} = 1.6\%$; dbh_{inv} = 50 cm.

[1502] *Stenostomum guianensis*
(Bremek.) Delprete & Achille

Blumea 55 (2): 169 [17 Aug. 2010] (Delprete & Achille 2010). — *Antirhea guianensis* Bremek., *Kew Bull.* 7 (2): 260 [25 July 1952] (Bremekamp 1952).

Guettarda fanshawei Steyerl., *Mem. New York Bot. Gard.* 23: 359 (Steyermark 1972).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman B-1936*.

SIZE. — Up to 50 cm dbh (Delprete et al. 2010).

Genus *Tocoyena* Aubl.

[1503] *Tocoyena guianensis* K.Schum.

Fl. Bras. [Martius] 6 (6): 346 [15 June 1889] (Schumann 1889).

Tocoyena guianensis var. *communis* Steyerl., *Mem. New York Bot. Gard.* 12 (3): 194 (Steyermark 1965).

Tocoyena guianensis var. *glabriuscula* Steyerl., *Mem. New York Bot. Gard.* 12 (3): 194 (Steyermark 1965).

VERNACULAR NAMES. — Pa: amuwan, tukuyuy-kamwi • Wp: yanipa lali u • Wn: kuwajimë, pilimokju • Br: jenipapo-do-campo.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *R.A.A. Oldeman 2756*.

INVENTORY DATA (FG). — 8 trees in 1 plot; Fmax< 1 %; dbh_{inv} = 33 cm.

Family RUTACEAE Juss.
Genus *Erythrociton* Nees & Mart.

[1504] *Erythrociton brasiliensis* Nees & Mart.

Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur. 11: 166 (Nees & Martius 1823).

Pentamorpha graveolens Scheidw., *Bull. Acad. Roy. Sci. Bruxelles* 9 (1): 21 (Scheidweiler 1842).

VERNACULAR NAMES. — Wn: kiapoku, pakutpë, pijsa piloman.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M. Fleury 2053*.

SIZE. — Up to 12 m tall (Kallunki 1992).

Genus *Esenbeckia* Kunth

[1505] *Esenbeckia cowanii* Kaastra

Acta Bot. Neerl. 26: 481 (Kaastra 1977).

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *R.S. Cowan 38757* (holo-, US[00101521]; iso-, F[V0071245F], NY[00051857]).

INVENTORY DATA (FG). — 34 trees in 8 plots; F_{max} < 1 %; dbh_{inv} = 35 cm.

[1506] *Esenbeckia grandiflora* Mart.

Nova genera et species plantarum [Martius] 3 (2): 85 ["1829" publ. Jan.-Mar (?) 1831] (Martius 1831). — *Colythrum grandiflorum* (Mart.) Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 399 (Steudel 1840).

Polembryum castanocarpum A.Juss., *Mém. Mus. Hist. Nat.* 12: 541 (Jussieu 1825).

Polembryum jussieui Schott, *Rutac.* t. 6 (Schott 1834).

Esenbeckia attenuata Griseb., *Fl. Brit. W.I. [Grisebach]*: 135 [June 1859] (Grisebach 1859).

Esenbeckia fasciculata Barb.Rodr., *Rev. Engenh.* 5: 1 (Barbosa Rodrigues 1883).

Esenbeckia grandiflora var. *macrophylla* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1285 (Chodat & Hassler 1904).

Esenbeckia obovalifolia Pittier, *Bol. Soc. Venez. Ci. Nat.* 9: 122 (Pittier 1944).

Esenbeckia grandiflora var. *peruviana* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (3.2): 673 (Macbride 1949).

Esenbeckia rigida R.S.Cowan, *Bol. Mus. Nac. Rio de Janeiro, Bot.* 27: 1 (Cowan 1961).

VERNACULAR NAMES. — Br: chupa-ferro, guaxupita, pitaguará-amarelo.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost 3423*.

SIZE. — Up to 20 cm dbh (Kallunki 2005).

Genus *Galipea* Aubl.

[1507] *Galipea davisii* Sandwith

Bull. Misc. Inform. Kew 1935 (3): 118 [20 May 1935] (Sandwith 1935).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Sastre & C. Moretti 3852* (P).

SIZE. — Up to 15 m tall (Kallunki 2005).

[1508] *Galipea trifoliata* Aubl.
(Fig. 51A)

Hist. Pl. Guiane 2: 662 [Jun.-Dec. 1775] (Aublet 1775).

Galipea fissa Miq., *Stirp. Surinam. Select.*: 64 ["1850" publ. Mar. 1851] (Miquel 1851).

VERNACULAR NAMES. — Ka: waikyara • Br: amarelinho, canela-de-velho, folha-de-couro.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000630633]).

SIZE. — Up to 15 m tall (Pirani & Kallunki 2007).

Genus *Hortia* Vand.

[1509] *Hortia excelsa* Ducke
(Fig. 51B)

Arch. Jard. Bot. Rio de Janeiro 3: 182 (Ducke 1922).

VERNACULAR NAMES. — Nt: gaan busi tobitutu • Br: cachaceiro.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 3753*.

INVENTORY DATA (FG). — 1 tree, dbh = 10.8 cm.

[1510] *Hortia superba* Ducke

Arch. Inst. Biol. Veg. 1 (3): 207 [Aug. 1935] (Ducke 1935).

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville 17272*.

SIZE. — Up to 30 cm dbh (Groppo & Pirani 2012).

Genus *Neoraputia* Emmerich ex Kallunki

[1511] *Neoraputia paraensis* (Ducke) Emmerich ex Kallunki

Brittonia 61 (1): 29 [2 Mar. 2009] (Kallunki 2009). — *Raputia paraensis* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 184 (Ducke

1922). — *Neoraputia paraensis* (Ducke) Emmerich, *Rodriguésia* 30 (45): 243 (Emmerich 1978), *nom. inval.* (genus not validly published).

Neoraputia cowanii Emmerich, *Rodriguésia* 30 (45): 259 (Emmerich 1978), *nom. inval.* (genus not validly published).

VERNACULAR NAMES. — Br: capança, caporé.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Moretti* 354.

SIZE. — Up to 10 m tall (Kallunki 2005).

Genus *Pilocarpus* Vahl

[1512] *Pilocarpus racemosus* Vahl

Ectog. Amer. 1: 29 (Vahl 1797).

Pilocarpus latifolius A.St.-Hil. ex Tul., *Ann. Sci. Nat., Bot. sér. 3*, 7: 285 (Tulasne 1847), “*latifolia*”.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *M.-F. Prévost et al.* 4604.

INVENTORY DATA (FG). — 25 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.2$ cm.

Genus *Raputia* Aubl.

[1513] *Raputia* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2174.

INVENTORY DATA (FG). — 19 trees in 6 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 50$ cm.

Genus *Ticorea* Aubl.

[1514] *Ticorea foetida* Aubl.
(Fig. 51C)

Hist. Pl. Guiane 2: 689 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000570872]).

SIZE. — Up to 14 m tall (Kallunki 1998).

[1515] *Ticorea longiflora* DC.

Mém. Mus. Hist. Nat. 9: 146 (Candolle 1822).

VERNACULAR NAMES. — Pa: kwik-washiune, waravru-kamwi • Wp: yapukuli sili • Wn: malihaliimë.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material G-DC, G00310774).

SIZE. — Up to 33 cm dbh (Kallunki 1998).

Genus *Zanthoxylum* L.

[1516] *Zanthoxylum acuminatum* (Sw.) Sw.
subsp. *juniperinum* (Poep.) Reynel

Phytoneuron 2015-22: 6 (Reynel 2015).

Zanthoxylum juniperinum Poep., *Nova genera ac species plantarum [Poepig & Endlicher]* 3: 77 [23-25 Jan. 1845] (Poepig 1845). — *Fagara juniperina* (Poep.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896).

Zanthoxylum minutiflorum Tul., *Ann. Sci. Nat., Bot. sér. 3*, 7: 278 (Tulasne 1847). — *Fagara minutiflora* (Tul.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 118 (Engler 1896).

Zanthoxylum pringlei S.Watson, *Proc. Amer. Acad. Arts* 26: 134 (Watson 1891), “*Xanthoxylum*”. — *Fagara pringlei* (S.Watson) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896).

Fagara warmingiana Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896).

Zanthoxylum procerum Donn.Sm., *Bot. Gaz.* 23 (1): 4 [20 Jan. 1897] (Donnell Smith 1897). — *Fagara procera* (Donn.Sm.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 19a: 223 (Engler 1931).

Fagara acreana K.Krause, *Notizbl. Königl. Bot. Gart. Berlin* 6: 143 [4 Apr. 1914] (Krause 1914). — *Zanthoxylum acreanum* (K.Krause) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.2)*: 664 (Macbride 1949).

Fagara ocmarensis Pittier, *Arb. Arbust. Venez.* 9-10: 120 [Dec. 1929] (Pittier 1929). — *Zanthoxylum ocmarense* (Pittier) Steyermark., *Fieldiana, Bot.* 28 (2): 273 (Steyermark 1952).

VERNACULAR NAMES. — Ka: awaladala, awalatala.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *S.A. Mori et al.* 21661.

INVENTORY DATA (FG). — 7 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 45.2$ cm.

[1517] *Zanthoxylum amapaense* (Albuq.) P.G.Waterman

Taxon 24 (2/3): 363 [6 June 1975] (Waterman 1975). — *Fagara amapaense* Albuq., *Publ. Inst. Nac. Pesq. Amaz., Bot.* 28: 11 (Albuquerque 1968).

Fagara idae Albuq., *Publ. Inst. Nac. Pesq. Amaz., Bot.* 28: 8 (Albuquerque 1968). — *Zanthoxylum idae* (Albuq.) P.G.Waterman, *Taxon* 24 (2/3): 364 [6 June 1975] (Waterman 1975).

Fagara machadoi Albuq., *Acta Amazonica* 1 (1): 24 (Albuquerque 1971). — *Zanthoxylum machadoi* (Albuq.) Albuq., *Acta Amazonica* 6 (3, Suppl.): 55 [Sep. 1976] (Albuquerque 1976).

HERBARIUM DATA (FG). — A single collection, *J.-F. Molino & D. Sabatier* 2349.

INVENTORY DATA (FG). — 1 tree, dbh = 56.7 cm.

[1518] *Zanthoxylum apiculatum* (Sandwith) P.G.Waterman

Taxon 24 (2/3): 363 [6 June 1975] (Waterman 1975). — *Fagara apiculata* Sandwith, *Bull. Misc. Inform. Kew* 1935 (3): 120 [20 May 1935] (Sandwith 1935).

Fagara dellomei Albuq., *Publ. Inst. Nac. Pesq. Amaz.*, Bot. 28: 3 (Albuquerque 1968). — *Zanthoxylum dellomei* (Albuq.) P.G.Waterman, *Taxon* 24 (2/3): 363 [6 June 1975] (Waterman 1975).

Zanthoxylum achidek Sanoja, *Acta Bot. Venez.* 32 (2): 304 (Sanoja 2009).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2901*.

INVENTORY DATA (FG). — 14 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 33.3$ cm.

[1519] *Zanthoxylum caribaeum* Lam.

Encycl. [J. Lamarck et al.] 2 (1): 39 [16 Oct. 1786] (Lamarck 1786). — *Fagara caribaea* (Lam.) Krug & Urb., *Bot. Jahrb. Syst.* 21 (5): 562 [12 May 1896] (Krug & Urban 1896).

Zanthoxylum elephantiasis Macfad., *Fl. Jamaica [Macfadyen]* 1: 193 (Macfadyen 1837). — *Fagara elephantiasis* (Macfad.) Krug & Urb., *Bot. Jahrb. Syst.* 21 (5): 564 [12 May 1896] (Krug & Urban 1896).

Zanthoxylum horridum Sessé & Moc., *Fl. Mexic.*, ed. 2, 231 (Sessé & Mociño 1894), “*Horrida*”.

Zanthoxylum occidentale Rose, *Contr. U.S. Natl. Herb.* 5 (4): 164 [31 Oct. 1899] (Rose 1899). — *Fagara occidentalis* (Rose) Engl., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 19a: 219 (Engler 1931).

Zanthoxylum gentlei Lundell, *Lloydia* 2 (2): 92 (Lundell 1939).

HERBARIUM DATA (FG). — A single collection, *H. de Foresta 125ter*.

SIZE. — Brazil, Pará. *G.T. Prance 1822* (MO), 10 m × 12 cm.

[1520] *Zanthoxylum ekmanii* (Urb.) Alain
(Fig. 51D)

Contr. Ocas. Mus. Hist. Nat. Colegio De La Salle 9: 24 (Alain 1950). — *Fagara ekmanii* Urb., *Repert. Spec. Nov. Regni Veg.* 20: 302 (Urban 1924).

Zanthoxylum belicense Lundell, *Contr. Univ. Michigan Herb.* 6: 35 (Lundell 1941).

Zanthoxylum sobrevielae D.R.Simpson, *Phytologia* 51 (5): 316 [July 1982] (Simpson 1982).

VERNACULAR NAMES. — Wp: sala'i, sala'i sili • Cr: bwa-pidjan, zépini-gran-fey • Br: limãozinho, maminha-de-porco, tamanqueira.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15603*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 27.2$ cm.

[1521] *Zanthoxylum paulae* (Albuq.) P.G.Waterman

Taxon 24 (2/3): 364 [6 June 1975] (Waterman 1975). — *Fagara paulae* Albuq., *Publ. Inst. Nac. Pesq. Amaz.*, Bot. 27: 4 (Albuquerque 1968).

Fagara luizii Albuq., *Publ. Inst. Nac. Pesq. Amaz.*, Bot. 27: 5 (Albuquerque 1968). — *Zanthoxylum luizii* (Albuq.) P.G.Waterman, *Taxon* 24 (2/3): 364 [6 June 1975] (Waterman 1975).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *V. Hequet 3018bis*.

INVENTORY DATA (FG). — 1 tree, $dbh = 66.9$ cm.

[1522] *Zanthoxylum pentandrum* (Aubl.) R.A.Howard

J. Arnold Arbor. 64 (2): 269 [Apr. 1983] (Howard 1983). — *Fagara pentandra* Aubl., *Hist. Pl. Guiane* 1: 78 [Jun.-Dec. 1775] (Aublet 1775). — *Fagara guianensis* Lam., *Encycl. [J. Lamarck et al.]* 2 (2): 446 [14 Apr. 1788] (Lamarck 1788), nom. illeg. superfl. (based on *Fagara pentandra*). — *Zanthoxylum hermaproditum* Willd., *Sp. Pl.*, ed. 4 4 (2): 756 [Apr. 1806] (Willdenow 1806), nom. illeg. superfl. (based on *Fagara pentandra*).

Zanthoxylum trinitense R.O.Williams, *Fl. Trinidad & Tobago* 1 (2): 148 [Nov. 1929] (Williams 1929). — *Fagara trinitensis* (R.O.Williams) Beard, *Empire For. J.* 21: 14 (Beard 1942).

VERNACULAR NAMES. — Pa: puduku-wakáu-purubumna, wahinhu • Ka: alemikilan, awaladala, awalatala • Wp: sala'i u • Nt: kee udu, kee maka udu • Cr: bwa-pidjan, zépini-gran-fey • Fr: bois piquant • Br: limãozinho, maminha-de-porco, tamanqueira.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777615] designated by Lanjouw & Uittien [1940: 151]).

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 31.7$ cm.

[1523] *Zanthoxylum rhoifolium* Lam.

Encycl. [J. Lamarck et al.] 2 (1): 39 [16 Oct. 1786] (Lamarck 1786). — *Fagara rhoifolia* (Lam.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 118 (Engler 1896).

Langsdorfia instrumentaria Leandro, *Denkschr. Königl. Akad. Wiss. München* 7: 242 (Leandro 1819). — *Pohlana langsdorffii* Nees & Mart., *Nova Acta Phys.-Med. Acad. Caes. Leop.-Carol. Nat. Cur.* 12: 17 (Nees & Martius 1824), nom. illeg. superfl. (based on *Langsdorfia instrumentaria*). — *Zanthoxylum langsdorffii* A.St.-Hil., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (2): 76 [26 Mar. 1825] (Saint-Hilaire 1825), nom. illeg. superfl. (based on *Pohlana langsdorffii*, thus indirectly on *Langsdorfia instrumentaria*).

Zanthoxylum perrottetii DC., *Prodri. [A. P. de Candolle]* 1: 726 [mid Jan. 1824] (Candolle 1824), “perrottetii”. — *Fagara perrottetii* (DC.) Lemée, *Fl. Guyane Franç.* 2: 181 (Lemée 1952).

Zanthoxylum sorbifolium A.St.-Hil., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (2): 75 [26 Mar. 1825] (Saint-Hilaire 1825).

Pohlana instrumentaria Mart. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 366, 796, (Steudel 1841) nom. nud.

Zanthoxylum pubescens A.St.-Hil. & Tul., *Ann. Sci. Nat., Bot. sér. 2*, 17: 141 (Saint-Hilaire & Tulasne 1842). — *Schinus pubescens*



FIG. 50. — Rubiaceae: **A**, *Ferdinandusa* sp. A (D. Sabatier & M.-F. Prévost 4928); **B**, *Posoqueria longiflora* Aubl.; **C**, *Rudgea lanceifolia* Salisb.; **D**, *Kutchubaea insignis* Fisch. ex DC. (M.-F. Prévost et al. 4734). A, © M.-F. Prévost/IRD; B, © P. G. Delprêtre/IRD; C, D, © D. Sabatier/IRD.

(A.St.-Hil. & Tul.) Spreng. ex Mart., *Fl. Bras. [Martius]* 12 (2): 176 [1 Sep. 1874] (Martius 1874), *nom. nud. pro syn.* — *Zanthoxylum rhoifolium* var. *pubescens* (A.St.-Hil. & Tul.) Engl., *Fl. Bras. [Martius]* 12 (2): 176 [1 Sep. 1874] (Engler 1874). — *Fagara rhoifolia* subsp. *pubescens* (A.St.-Hil. & Tul.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 118 (Engler 1896). — *Fagara pubescens* (A.St.-Hil. & Tul.) Herzog, *Meded. Rijks-Herb.* 27: 80 (Herzog 1915), *nom. illeg. hom.*, *non* A.Chev. (Chevalier 1912).

Zanthoxylum peltophorum Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 31 (1): 442 (Turczaninow 1858). — *Zanthoxylum rhoifolium* var. *peltophorum* (Turcz.) Engl., *Fl. Bras. [Martius]* 12 (2): 175 [1 Sep. 1874] (Engler 1874). — *Fagara rhoifolia* var. *peltophora* (Turcz.) Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1283 (Chodat & Hassler 1904), “*peltophorum*”.

Zanthoxylum microcarpum Griseb., *Fl. Brit. W.I. [Grisebach]*: 138 [June 1859] (Grisebach 1859). — *Fagara microcarpa* (Griseb.) Krug & Urb., *Bot. Jahrb. Syst.* 21 (5): 570 [12 May 1896] (Krug & Urban 1896).

Zanthoxylum obscurum Engl., *Fl. Bras. [Martius]* 12 (2): 169 [1 Sep. 1874] (Engler 1874). — *Fagara obscura* (Engl.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896).

Zanthoxylum acutifolium Engl., *Fl. Bras. [Martius]* 12 (2): 170 [1 Sep. 1874] (Engler 1874). — *Fagara acutifolia* (Engl.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896).

Zanthoxylum acutifolium var. *petiolatum* Engl., *Fl. Bras. [Martius]* 12 (2): 170 [1 Sep. 1874] (Engler 1874). — *Fagara rhoifolia* var. *petiolata* (Engl.) Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1283 (Chodat & Hassler 1904), “*petiolatum*”.

Zanthoxylum obscurum var. *ruizianum* Klotzsch ex Engl., *Fl. Bras. [Martius]* 12 (2): 170 [1 Sep. 1874] (Engler 1874). — *Fagara ruiziana* (Klotzsch ex Engl.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 117 (Engler 1896). — *Zanthoxylum ruizianum* (Klotzsch ex Engl.) J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser. 13 (3.2)*: 667 (Macbride 1949).

Zanthoxylum regnellianum Engl., *Fl. Bras. [Martius] 12 (2)*: 174 [1 Sep. 1874] (Engler 1874). — *Fagara regnelliana* (Engl.) Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1283 (Chodat & Hassler 1904).

Zanthoxylum rhoifolium var. *petiolatum* Engl., *Fl. Bras. [Martius] 12 (2)*: 175 [1 Sep. 1874] (Engler 1874).

Zanthoxylum rhoifolium var. *sessilifolium* Engl., *Fl. Bras. [Martius] 12 (2)*: 176 [1 Sep. 1874] (Engler 1874).

Fagara regnelliana var. *calvata* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1283 (Chodat & Hassler 1904).

Fagara rhoifolia f. *paucijuga* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1283 (Chodat & Hassler 1904).

Fagara rhoifolia f. *angustifolia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1284vt (Chodat & Hassler 1904).

Fagara rhoifolia f. *intermedia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1284 (Chodat & Hassler 1904).

Fagara rhoifolia f. *latifolia* Chodat & Hassl., *Bull. Herb. Boissier*, sér. 2, 4: 1284 (Chodat & Hassler 1904).

Fagara coco (Gillies ex Hook.f. & Arn.) Engl. var. *formosana* Lillo, *Contr. Conoc. Arb. Argent.*: 89 (Lillo 1910). — *Zanthoxylum rhoifolium* var. *formosanum* (Lillo) P.G.Waterman, *Taxon 25 (5/6)*: 594 [26 Nov. 1976] (Waterman 1976).

Fagara rothschuhii Loes., *Bot. Jahrb. Syst. 60 (4)*: 366 [1 Aug. 1926] (Loesener 1926).

Fagara astrigera R.S.Cowan, *Sellowia 12*: 81 (Cowan 1960). — *Zanthoxylum astrigerum* (R.S.Cowan) P.G.Waterman, *Taxon 24 (2/3)*: 363 [6 June 1975] (Waterman 1975).

Fagara rhoifolia var. *surparanaensis* Najera, Galdeano & Escal., *Bol. Soc. Argent. Bot.* 14 (3): 244 (Najera et al. 1972). — *Zanthoxylum rhoifolium* var. *surparanaense* (Najera, Galdeano & Escal.) P.G.Waterman, *Taxon 25 (5/6)*: 594 [26 Nov. 1976] (Waterman 1976).

Fagara rhoifolia var. *intermedia* R.S.Cowan & L.B.Sm., *Fl. Ilustr. Catar. (Ruta.)*: 13 (Cowan & Smith 1973).

VERNACULAR NAMES. — Pa: puduku-wakáu • Ka: awaladala, awalatala • Te: wila atsi • Wn: kanike • Nt: kee udu, kee maka udu • Cr: bwa-pidjan, zépini-ti'fey • Fr: bois piquant • Br: mamiqueira, tamanqueira, tembetaru.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *M.-F. Prévost 3781*, dbh 10 cm.

Family SABIACEAE Blume
Genus *Meliosma* Blume

[1524] *Meliosma* sp. A
(Fig. 51E)

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5672*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1 \%$; $dbh_{inv} = 31.3$ cm.

Genus *Ophiocaryon* Endl.

[1525] *Ophiocaryon chironectes* Barneby

Mem. New York Bot. Gard. 23: 117 (Barneby 1972).

VERNACULAR NAME. — Wp: tápē lemiū'i.

HERBARIUM DATA (FG). — A single collection, *P. Grenand 1543*. The type is from Clevelandia, Amapá, on the Brazilian border of the Oyapock River: *L.Y.T. Westra 47298* (NY[00387482]).

INVENTORY DATA (FG). — 1 tree, dbh = 10.8 cm.

[1526] *Ophiocaryon paradoxum* R.H.Schomb.

London J. Bot. 4: 377 (Schomburgk 1845).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — A single collection, *C. Sastre 4704*.

SIZE. — Guyana. *B. Maguire 22937* (MO), dbh 30 cm.

Family SALICACEAE Mirb.
Genus *Banara* Aubl.

[1527] *Banara guianensis* Aubl.

Hist. Pl. Guiane 1: 548 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate.

Banara fagifolia Vahl, *Symb. Bot. [Vahl] 3*: 65 (Vahl 1794), nom. illeg. superfl. (based on *Banara guianensis*).

Xyladenius glandulosus Desv. ex Ham., *Prodr. Pl. Ind. Occid. [Hamilton]*: 41 [Oct. 1825] (Hamilton 1825). — *Banara glandulosa* (Desv. ex Ham.) Speg., *Revista Argent. Bot.* 1: 210 (Spegazzini 1926).

Kuhlia mollis Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher] 3*: 74 [23-25 Jan. 1845] (Poeppig 1845). — *Banara mollis* (Poepp.) Tul., *Ann. Sci. Nat., Bot. sér. 3, 7*: 288 (Tulasne 1847). — *Banara guianensis* var. *mollis* (Poepp.) Eichler, *Fl. Bras. [Martius] 13 (1)*: 501 [1 Oct. 1871] (Eichler 1871).

Laetia glandulosa Poepp. ex Tul., *Ann. Sci. Nat., Bot. sér. 3, 7*: 288 (Tulasne 1847), nom. nud. pro syn.

Trilia glandulosa Dombey ex Tul., *Ann. Sci. Nat., Bot. sér. 3, 7*: 288 (Tulasne 1847), “*Trilin*”, nom. nud. pro syn.

Banara pubescens Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 5 (Suppl. 2)*: 92 [May 1861] (Bentham 1861).

Banara pyramidata Rusby, *Mem. Torrey Bot. Club 3*: 33 (Rusby 1893).

Banara guianensis var. *spruceana* Briq., *Annuaire Conserv. Jard. Bot. Genève 2*: 49 (Briquet 1898).

Banara tulasnei J.F.Macbr., *Candollea 5*: 389 (Macbride 1934).

Trilia macrobotrys Ruiz & Pav., *Anales Inst. Bot. Cavanilles 16*: 374 (Ruiz & Pavón 1959).

VERNACULAR NAMES. — Pa: kuuku-ariut-priye, tahuma • Wp: yape'a pitá sili • Wn: kumidan • Nt: bita tiki, boni udu • Cr: mavévé-sikriyé • Br: cavaqueiro-do-baixo, lacre-branco.

HERBARIUM DATA (FG). — 109 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material P-JU, P00672058)).

INVENTORY DATA (FG). — 6 trees in 3 plots; $dbh_{inv} = 16.4$ cm.

Genus *Casearia* Jacq.

[1528] *Casearia arborea* (Rich.) Urb.

Symb. Antill. [Urban] 4 (3): 421 [15 May 1910] (Urban 1910). — *Samyda arborea* Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792). — *Casearia stipularis* Vent., *Choix Pl.* 1: 46 [Apr. 1804] (Ventenat 1804), nom. illeg. superfl. (based on *Samyda arborea*). — *Samyda stipularis* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 5: 32 [1 Nov. 1817] (Poiret 1817), nom. illeg. superfl. (based on *Samyda arborea*). — *Casearia stipularis* Poir. ex Spreng., *Syst. Veg. [Sprengel]* 2: 355 [Jan.-May 1825] (Sprengel 1825), nom. illeg. superfl. (based on *Samyda stipularis*, thus on *Samyda arborea*). — *Guidonia stipularis* M. Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.* 54 (Gómez 1889), nom. illeg. superfl. (based on *Casearia stipularis* Vent., thus on *Samyda arborea*).

Samyda viridiflora Aubl., *Hist. Pl. Guiane* 1: 402 [Jun.-Dec. 1775] (Aublet 1775).

Chaetocrater capitatum Ruiz & Pav., *Syst. Veg. Fl. Peruv. Chil.* 1: 108 [late Dec. 1798] (Ruiz & Pavón 1798). — *Casearia capitata* (Ruiz & Pav.) Pers., *Syn. Pl. [Persoon]* 1: 485 [1 Apr.-15 June 1805] (Persoon 1805).

Samyda niviana Poir., *Encycl. [J. Lamarck et al.]* 6 (2): 493 [28 Aug. 1805] (Poiret 1805).

Casearia incana Bertero ex Spreng., *Syst. Veg. [Sprengel]* 2: 355 [Jan.-May 1825] (Sprengel 1825).

Casearia serrata Macfad., *Fl. Jamaica [Macfadyen]* 1: 216 (Macfadyen 1837).

Casearia lanceolata Miq., *Linnaea* 18: 753 ["1844" publ. Aug.-Oct. 1845] (Miquel 1845).

Casearia hostmanniana Steud. ex Griseb., *Fl. Brit. W.I. [Grisebach]* 23 [Dec. 1859] (Grisebach 1859), nom. nud. pro syn.

Casearia poeppigii Eichler, *Fl. Bras. [Martius]* 13 (1): 475 [1 Oct. 1871] (Eichler 1871).

Casearia cambessedesii var. *angustifolia* Eichler, *Fl. Bras. [Martius]* 13 (1): 476 [1 Oct. 1871] (Eichler 1871).

Casearia cambessedesii var. *parvifolia* Eichler, *Fl. Bras. [Martius]* 13 (1): 476 [1 Oct. 1871] (Eichler 1871).

Casearia oligantha Eichler, *Fl. Bras. [Martius]* 13 (1): 476 [1 Oct. 1871] (Eichler 1871).

Casearia brasiliensis Eichler, *Fl. Bras. [Martius]* 13 (1): 477 [1 Oct. 1871] (Eichler 1871).

Casearia bangii Rusby, *Mem. Torrey Bot. Club* 3: 34 (Rusby 1893).

Samyda virgata Sessé & Moc., *Fl. Mexic.*, ed. 2, 113 (Sessé & Mociño 1894).

Casearia glaziovii Briq., *Annuaire Conserv. Jard. Bot. Genève* 2: 72 (Briquet 1898).

Casearia umbellifera Benoist, *Bull. Mus. Natl. Hist. Nat.* 26: 353 (Benoist 1920).

Casearia glaberrima Uittien, *Recueil Trav. Bot. Néerl.* 22: 371 ["1925" publ. Jan. 1926] (Uittien 1926).

NOTE. — *Samyda virgata* Sessé & Moc. appears also in the first edition of *Flora mexicana* (p. 123, as "Uirgata"), but edition 2 has

priority, probably from p. 49 onward (Stafleu & Cowan, 1985: TL-2-11756]).

VERNACULAR NAMES. — Ka: akukuwa elepali, pilitalaipyo • Wp: yape'a pirá sili • Wn: luwelue • Br: capança-braba, carniceiro, erva-de-lagarto.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *J.B. Leblond* s.n. (in herb. L.C. Richard) (type P[P00371634]).

INVENTORY DATA (FG). — 1 tree, dbh = 14 cm.

[1529] *Casearia bicolor* Urb.

Symb. Antill. [Urban] 1 (2): 372 [10 Apr. 1899] (Urban 1899).

Laetia procera (Poepp.) Eichler, *Fl. Bras. [Martius]* 13 (1): 453 [1 Oct. 1871] (Eichler 1871). — *Samyda procera* Poepp., *Nova genera ac species plantarum [Poeppig & Endlicher]* 3: 67 [23-25 Jan. 1845] (Poeppig 1845). — *Casinga procera* (Poepp.) Griseb., *Fl. Brit. W.I. [Grisebach]* 710 [Oct. 1864] (Grisebach 1864). — *Guidonia procera* (Poepp.) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Laetia caseariooides Sagot ex Benth., *J. Proc. Linn. Soc., Bot.* 5 (Suppl. 2): 85 [May 1861] (Bentham 1861).

Samyda obtusifolia Rich. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 454 [1 Oct. 1871] (Eichler 1871), in obs.

Laetia obtusifolia Eichler ex Benoist, *Bull. Mus. Natl. Hist. Nat.* 26: 351 (Benoist 1920), nom. nud.

Casearia belizensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 12: 412 (Standley 1936).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). The genus *Laetia* Loefl. ex L. was included in *Casearia* Jacq. by Samarakoon & Alford (2019).

VERNACULAR NAMES. — Pa: ahakyu, ahikyu • Ka: akale wewe, alokoyulu, auluko yulu, kayman udu, mainyapo • Te: tale'i li • Wp: miyú'i pepo, muyú'i pepo, tale'i li • Wn: mijelemlíi • Nt: kayman udu • Cr: bwa-jako, bwa-zanmand • Fr: bois caïman • Br: apijó, casinga-cheirosa, muira-pucu, pau-jacaré, tuchauá.

HERBARIUM DATA (FG). — 79 collections at CAY. Sel. exs.: *P.A. Sagot* 1137, 1857 (original material of *Laetia caseariooides*: K[K000471402, K000471403], P[P02442044, P02442045, P02442046, P02442047]).

INVENTORY DATA (FG). — 153 trees in 88 plots; F_{max} = 1.2 %; dbh_{inv} = 94.2 cm.

[1530] *Casearia decandra* Jacq.

Enum. Syst. Pl. 21 [Aug.-Sep. 1760] (Jacquin 1760). — *Casearia parviflora* Jacq., *Select. Stirp. Amer. Hist.*, ed. 1780-1781, 66 (Jacquin 1780), nom. illeg. superfl. (based on the illustration of *Casearia decandra*). — *Casearia parvifolia* Willd., *Sp. Pl.*, ed. 4 2 (1): 627 [Mar. 1799] (Willdenow 1799), nom. illeg. superfl. (based on *Casearia decandra*). — *Samyda parvifolia* Poir., *Encycl. [J. Lamarck et al.]* 6 (2): 492 [28 Aug. 1805] (Poiret 1805), nom. illeg. superfl. (based on *Casearia decandra*). — *Anavinga parvifolia* (Poir.) Lam., *Tabl. Encycl.* 2[5] (2): 484 [31 Oct. 1819] (Lamarck 1819), nom. illeg. superfl. (based on *Samyda parvifolia*, thus on *Casearia decandra*). — *Guidonia parvifolia* M. Gómez, *Dicc. Bot. Nombres Vulg.*

Cub. Puerto-Riq.: 93 (Gómez 1889), nom. illeg. superfl. (based on *Casearia parvifolia*, thus on *Casearia decandra*).

Casearia adamantium Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (16): 230 ["1829" publ. Oct.-Nov. 1830] (Cambessèdes 1830).

Casearia adstringens Mart., *Syst. Mat. Med. Veg. Bras.* 51 (Martius 1843), nom. nud.

Casearia nitida Sieber ex Griseb., *Fl. Brit. W.I. [Grisebach]* 23 [Dec. 1859] (Grisebach 1859), nom. nud. pro syn.

Casearia serrulata Sieber ex Griseb., *Fl. Brit. W.I. [Grisebach]* 23 [Dec. 1859] (Grisebach 1859), nom. nud. pro syn.

Casearia parvifolia var. *microcarpa* Eggers, *Bull. U.S. Natl. Mus.* 13: 26 (Eggers 1879).

Guidonia adstringens Baill., *Traité Bot. Méd. Phan.* 2: 827 [Mar. 1884] (Baillon 1884).

Samyda lancifolia Sessé & Moc., *Fl. Mexic.*, ed. 2, 113 (Sessé & Mociño 1894).

Casearia floribunda Briq., *Bull. Herb. Boissier*, sér. 2, 7: 672 (Briquet 1907).

Casearia parvifolia var. *paraguariensis* Briq., *Bull. Herb. Boissier*, sér. 2, 7: 672 (Briquet 1907).

Casearia albicalvis Rusby, *Mem. New York Bot. Gard.* 7: 307 (Rusby 1927).

Casearia pavoniana Sleumer, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 958 [20 Jan. 1934] (Sleumer 1934).

Casearia reflexa Sleumer, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 959 [20 Jan. 1934] (Sleumer 1934).

Chaetocrater reflexus Ruiz & Pav., *Anales Inst. Bot. Cavanilles* 15: 130 (Ruiz & Pavón 1958), "reflexum".

NOTES.—Jacquin (1780: 66) cited *Samyda parviflora* L. "Syst. 340", in synonymy under *Casearia parviflora* Jacq., but his plate 127 is the same as plate 85 in the first edition of *Selectarum stirpium Americanarum historia* (Jacquin 1763), which illustrates *C. decandra* Jacq. (p. 133). Although *Samyda lancifolia* Sessé & Moc. already appeared in the first edition of *Flora mexicana* (Sessé & Mociño 1892-1898: 123), edition 2 has priority, probably from p. 49 onward (Stafleu & Cowan, 1985: [TL-2-11756](#)). *Guidonia adstringens* Baill. is based on the invalid *Casearia adstringens* Mart.

VERNACULAR NAMES.—Br: caferana, café-do-mato.

HERBARIUM DATA (FG).—17 collections at CAY. Sel. exs.: [M.-F. Prévost 3064](#).

INVENTORY DATA (FG).—6 trees in 6 plots; $F_{\max} = 1.3\%$; dbh_{inv} = 16.9 cm.

[1531] *Casearia grandiflora* Cambess.
(Fig. 52A)

Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.) 2 (16): 232 ["1829" publ. Oct.-Nov. 1830] (Cambessèdes 1830).

Casearia hypoleuca Mart., *Flora* 20 (2, Beibl.): 128 (Martius 1837).—*Casearia grandiflora* var. *hypoleuca* (Mart.) Eichler, *Fl. Bras. [Martius]* 13 (1): 479 [1 Oct. 1871] (Eichler 1871).

Casearia grandiflora var. *obtusifolia* Eichler, *Fl. Bras. [Martius]* 13 (1): 479 [1 Oct. 1871] (Eichler 1871).

Casearia grandiflora var. *pauciflora* Eichler, *Fl. Bras. [Martius]* 13 (1): 479 [1 Oct. 1871] (Eichler 1871).

Casearia anavinga A.St.-Hil. ex Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 13: 285 (Sagot 1882).

HERBARIUM DATA (FG).—25 collections at CAY. Sel. exs.: [D. Sabatier & J.-F. Molino 5187](#).

INVENTORY DATA (FG).—5 trees in 4 plots; $F_{\max} < 1\%$; dbh_{inv} = 21.7 cm.

[1532] *Casearia guianensis* (Aubl.) Urb.

Symb. Antill. [Urban] 3 (2): 322 [15 Aug. 1902] (Urban 1902).—*Iroucana guianensis* Aubl., *Hist. Pl. Guiane* 1: 329 [Jun.-Dec. 1775] (Aublet 1775), "Guyannensis" on plate. — *Athenaea guianensis* (Aubl.) J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 629 [late Sep.-Nov. 1791] (Gmelin 1791).—*Casearia ramiflora* Vahl, *Symb. Bot. [Vahl]* 2: 50 [Jul.-Dec. 1791] (Vahl 1791), nom. illeg. superfl. (based on *Iroucana guianensis*).—*Samyda iroucana* Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792), nom. illeg. superfl. (based on *Iroucana guianensis*).—*Guidonia ramiflora* M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.* 64 (Gómez 1889), nom. illeg. superfl. (based on *Casearia ramiflora*, thus indirectly on *Iroucana guianensis*).

Casearia fallax Miq., *Linnaea* 19: 128 ["1847" publ. Feb. 1846] (Miquel 1846).

Samyda octandra Sessé & Moc., *Fl. Mexic.*, ed. 2, 113 (Sessé & Mociño 1894).

NOTE.—*Samyda octandra* Sessé & Moc. appears also in the first edition of *Flora mexicana* (Sessé & Mociño 1892-1898: 123), but edition 2 has priority, probably from p. 49 onward (Stafleu & Cowan, 1985: [TL-2-11756](#)).

VERNACULAR NAMES.—Ka: alekikolan.

HERBARIUM DATA (FG).—30 collections at CAY. Sel. exs.: [J.B. Aublet s.n.](#) (original material P[P00689737]).

INVENTORY DATA (FG).—1 tree, dbh = 16.2 cm.

[1533] *Casearia mariquitensis* Kunth

Nova genera et species plantarum [H.B.K.] 5: 363 [24 Mar. 1823] (Kunth 1823).

Casearia fockeana Miq., *Ann. Sci. Nat., Bot. sér. 3*, 1: 39 (Miquel 1844).

Casearia pubiflora Benth., *Bot. Voy. Sulphur [Bentham]*: 66 [16 Aug. 1844] (Bentham 1844).

Casearia platyphylla Briq., *Annuaire Conserv. Jard. Bot. Genève* 2: 68 (Briquet 1898), pro parte quoad floram tantum (Sleumer 1980).

Casearia camporum Sprague, *Trans. & Proc. Bot. Soc. Edinburgh* 22 (4): 427 (Sprague 1905).

Casearia tarapotina Pilg., *Verh. Bot. Vereins Prov. Brandenburg* 47: 161 [1 Oct. 1905] (Pilger 1905).



FIG. 51. — Rutaceae: **A**, *Galipea trifoliata* Aubl. (J.-F. Molino & D. Sabatier 2164); **B**, *Hortia excelsa* Ducke (D. Sabatier & M.-F. Prévost 4935); **C**, *Ticorea foetida* Aubl. (J.-F. Molino et al. 2041); **D**, *Zanthoxylum ekmanii* (Urb.) Alain (D. Sabatier & J.-F. Molino 5747). Sabiaceae: **E**, *Meliosma* sp. A (D. Sabatier & J.-F. Molino 5672). A, © J.-F. Molino/IRD; B-E, © D. Sabatier/IRD.

HERBARIUM DATA (FG). — A single collection *fide* Sleumer (1980): *L.C. Richard s.n.* (P). This collection could not be located with certainty, but could correspond to P04821776 or P04821777.

SIZE. — Up to 60 cm dbh (Sleumer 1980).

[1534] *Casearia negrensis* Eichler

Fl. Bras. [Martius] 13 (1): 466 [1 Oct. 1871] (Eichler 1871).

VERNACULAR NAMES. — Pa: waaduk.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 3191.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} 14.3 cm.

[1535] *Casearia pitumba* Sleumer

Blumea 24: 118 (Sleumer 1978). — *Pitumba guianensis* Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 29 [Jun.-Dec. 1775] (Aublet 1775). — *Samyda pitumba* Poir., *Encycl. [J. Lamarck et al.]* 6 (2): 492 [28 Aug. 1805] (Poiret 1805), nom. illeg. superfl. (based on *Pitumba guianensis*). — *Casearia macrophylla* Vahl, *Eclog. Amer.* 2: 32 (Vahl 1798), nom. illeg. superfl. (based on *Pitumba guianensis*). — *Casearia microphylla* Dennst., *Nomencl. Bot. [Dennstedt]* 1: 94 (Dennstedt 1810), nom. illeg. superfl. (based on *Pitumba guianensis*).

Pitumba edulis A.Rich. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 470 [1 Oct. 1871] (Eichler 1871), nom. nud. pro syn.

Casearia timbuchi J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 358 (Macbride 1936), nom. inval. (anglice).

Casearia macrophylla var. *barbatula* J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (4.1): 45 (Macbride 1941).

NOTE. — *Casearia microphylla* is an orthographic variant that possibly originated from a typographical error (*macrophylla* intended).

VERNACULAR NAMES. — Pa: agagut-aška, ararut-aška, tahuma-pu-rubumna, tahuma-seine • Ka: pelepele apotokon • Wp: yape'a pitá, yape'a pitá u • Wn: luweluve • Nt: kwasikwasi tiki • Cr: grénn-oko, grénn-toti • Br : cabelo-de-cotia.

HERBARIUM DATA (FG). — 156 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000611049]).

INVENTORY DATA (FG). — 12 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22.3 \text{ cm}$.

[1536] *Casearia prunifolia* Kunth

Nova genera et species plantarum [H.B.K.] 5: 364 [24 Mar. 1823] (Kunth 1823).

Casearia matheusii Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 608 (Turczaninow 1863).

HERBARIUM DATA (FG). — A single collection, *C. Feuillet et al. 10319*.

SIZE. — Up to 20 m tall (Olson *et al.* 1999).

[1537] *Casearia rusbyana* Briq.

Annuaire Conserv. Jard. Bot. Genève 2: 73 (Briquet 1898).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: ahakyu, ahikyu.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier 2321*.

INVENTORY DATA (FG). — 6 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 29.4 \text{ cm}$.

[1538] *Casearia singularis* Eichler

Fl. Bras. [Martius] 13 (1): 473 [1 Oct. 1871] (Eichler 1871).

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5642*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 11.5 \text{ cm}$.

[1539] *Casearia sylvestris* Sw.

Fl. Ind. Occid. 2: 752 [Jan.-June 1798] (Swartz 1798). — *Samyda sylvestris* (Sw.) Poir., *Encycl. [J. Lamarck et al.] 6 (2)*: 492 [28 Aug. 1805] (Poirier 1805), “*silvestris*”. — *Guidonia sylvestris* (Sw.) M.Gómez, *Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq.* 95 (Gómez 1889).

Samyda parviflora L., *Syst. Nat.*, ed. 10, 2: 1025 [7 June 1759] (Linnaeus 1759), *nom. illeg. hom.*, *non Loefl.* (Loefling 1758, *nom. rej.*) *nec Sessé & Moc.* (Sessé & Mociño 1894).

Casearia parviflora J.F.Gmel., *Syst. Nat.*, ed. 13[bis], 2 (1): 700 [late Sep.-Nov. 1791] (Gmelin 1791), *nom. illeg. hom.*, *non Jacq.* (Jacquin 1780).

Casearia serrulata Sw., *Fl. Ind. Occid.* 2: 754 [Jan.-June 1798] (Swartz 1798).

Casearia parviflora Willd., *Sp. Pl.*, ed. 4 2 (1): 627 [Mar. 1799] (Willdenow 1799), *nom. illeg. hom.*, *non Jacq.* (Jacquin 1780).

Anavinga samyda C.F.Gaertn., *Suppl. Carp.* 241 (Gaertner 1807). — *Casearia samyda* (C.F.Gaertn.) DC., *Prodr. [A. P. de Candolle]* 1: 51 [mid Jan. 1824] (Candolle 1824).

Casearia punctata Spreng., *Neue Entdeck. Pflanzenk.* 2: 154 (Sprengel 1821).

Casearia sylvestris var. *platyphylla* DC., *Prodr. [A. P. de Candolle]* 2: 49 [mid Nov. 1825] (Candolle 1825).

Casearia parviflora var. *microphylla* Schlechtld., *Linnaea* 4: 90 (Schlechtdal 1829).

Casearia affinis Gardner, *London J. Bot.* 1: 529 (Gardner 1842).

Casearia benthamiana Miq., *Ann. Sci. Nat., Bot. sér. 3*, 1: 38 (Miquel 1844). — *Casearia sylvestris* var. *benthamiana* (Miq.) Uittien, *Recueil Trav. Bot. Néerl.* 22: 373 [“1925” publ. Jan. 1926] (Uittien 1926).

Casearia sylvestris var. *myricoides* Griseb., *Cat. Pl. Cub. [Grisebach]* 9 [May-Aug. 1866] (Grisebach 1866).

Samyda parviflora Sessé & Moc., *Fl. Mexic.*, ed. 2, 112 (Sessé & Mociño 1894), *nom. illeg. hom.*, *non Loefl.* (Loefling 1758, *nom. rej.*) *nec L.* (Linnaeus 1759).

Casearia attenuata Rusby, *Mem. Torrey Bot. Club* 6 (1): 41 (Rusby 1896).

Casearia sylvestris var. *tomentella* Rusby, *Mem. Torrey Bot. Club* 6 (1): 41 (Rusby 1896).

Casearia sylvestris var. *wydleri* Briq., *Annuaire Conserv. Jard. Bot. Genève* 2: 74 (Briquet 1898).

Casearia sylvestris var. *eichleri* Briq., *Bull. Herb. Boissier*, sér. 2, 7: 672 (Briquet 1907).

Casearia onacaensis Rusby, *Descr. S. Amer. Pl.* 62 [20 Dec. 1920] (Rusby 1920).

Casearia chlorophoroidea Rusby, *Descr. S. Amer. Pl.* 63 [20 Dec. 1920] (Rusby 1920). — *Casearia sylvestris* var. *chlorophoroidea* (Rusby) Sleumer, *Fl. Suriname* 3 (1): 298 (Sleumer 1935).

Casearia herbert-smithii Rusby, *Descr. S. Amer. Pl.* 63 [20 Dec. 1920] (Rusby 1920).

Casearia formosa Urb., *Repert. Spec. Nov. Regni Veg.* 22: 91 (Urban 1925).

Casearia lindeniana Urb., *Repert. Spec. Nov. Regni Veg.* 22: 92 (Urban 1925).

Casearia caudata Uittien, *Recueil Trav. Bot. Néerl.* 22: 373 [“1925” publ. Jan. 1926] (Uittien 1926).

Casearia sylvestris var. *angustifolia* Uittien, *Recueil Trav. Bot. Néerl.* 22: 373 [“1925” publ. Jan. 1926] (Uittien 1926).

Casearia sylvestris var. *paraensis* Uittien, *Recueil Trav. Bot. Néerl.* 22: 373 [“1925” publ. Jan. 1926] (Uittien 1926).

Casearia schulziana O.C.Schmidt, *Repert. Spec. Nov. Regni Veg.* 32: 87 (Schmidt 1933).

Casearia ekmanii Sleumer, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 954 [20 Jan. 1934] (Sleumer 1934).

Casearia ovoidea Sleumer, *Notizbl. Bot. Gart. Berlin-Dahlem* 11: 958 [20 Jan. 1934] (Sleumer 1934).

Casearia sylvestris var. *martinicensis* J.F.Macbr. ex Ll.Williams, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 357 (Williams 1936), *nom. inval.* (*anglice*).

Casearia subsessiliflora Lundell, *Contr. Univ. Michigan Herb.* 6: 50 (Lundell 1941).

NOTE. — *Samyda parviflora* Sessé & Moc. also appears in the first edition of *Flora mexicana* (Sessé & Mociño 1892–1898: 123, as “*Paruiflora*”), but edition 2 has priority, probably from p. 49 onward (Stafleu & Cowan, 1985: [TL-2-11756](#)].

VERNACULAR NAMES. — Wp: yape'a pitā, yape'a pitā sili • Nt: kwasikwasi tiki • Br: cafézeiro-do-mato, erva-de-bugre, erva-de-lagarto, guaçatunga.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Sabatier 2100*.

INVENTORY DATA (FG). — 35 trees in 22 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22$ cm.

[1540] *Casearia ulmifolia* Vahl ex Vent.

(Fig. 52C)

Choix Pl. 1: 46, verso [Apr. 1804] (Ventenat 1804). — *Guidonia ulmifolia* (Vahl ex Vent.) Baill., *Traité Bot. Méd. Phan.* 2: 827 [Mar. 1884] (Baillon 1884).

Casearia celtidifolia Kunth, *Nova genera et species plantarum [H.B.K.] 5:* 363 [24 Mar. 1823] (Kunth 1823).

Casearia petraea Benth., *J. Bot. [Hooker]* 4: 111 (Bentham 1842).

Casearia blanchetiana Miq., *Linnaea* 22: 801 [“1849” publ. May 1850] (Miquel 1850).

Casearia cambessedesii Eichler, *Fl. Bras. [Martius]* 13 (1): 475 [1 Oct. 1871] (Eichler 1871).

Casearia celtidifolia Poepp. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 477 [1 Oct. 1871] (Eichler 1871), *nom. illeg. hom.*, *non* Kunth (1823).

Casearia cotticensis Uittien, *Recueil Trav. Bot. Néerl.* 22: 369 [“1925” publ. Jan. 1926] (Uittien 1926).

Casearia reginae J.F.Macbr. ex Ll.Williams, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 356 (Williams 1936), *nom. inval.* (*anglice*).

Casearia tremifolia J.F.Macbr., *Candollea* 8: 23 (Macbride 1940).

VERNACULAR NAMES. — Wp: yape'a pitā sili • Br: caferana.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5006bis*.

INVENTORY DATA (FG). — 16 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 23.3$ cm.

[1541] *Casearia zizyphoides* Kunth

Nova genera et species plantarum [H.B.K.] 5: 362 [24 Mar. 1823] (Kunth 1823).

Casearia celastroides Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1167 [“1848” publ. 7–10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Casearia laevigata Willd. ex Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1167 [“1848” publ. 7–10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

HERBARIUM DATA (FG). — A single collection *fide* Sleumer (1980): *J. Martin s.n.* (K, not seen).

SIZE. — Up to 10 m tall (Sleumer 1980).

[1542] *Casearia* sp. A

NOTE. — This species is also the “*Casearia* sp. A” in Mori & Fischer (2002).

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 14721*.

INVENTORY DATA (FG). — 13 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 16.4$ cm.

[1543] *Casearia* sp. B

(Fig. 52B)

NOTE. — This species is also the “*Casearia* sp. B” in Mori & Fischer (2002).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *S.A. Mori & T.D. Pennington 18124*.

INVENTORY DATA (FG). — 25 trees in 20 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 28.2$ cm.

[1544] *Casearia* sp. C

NOTES. — This species, represented at CAY by specimens formerly identified as *Casearia acuminata* DC., differs from the latter in: 1) habit (trees vs treelets/shrubs of 5 cm dbh maximum in *C. acuminata*), 2) leaf pilosity (hairs on midvein on abaxial side vs glabrous), 3) leaf colour and size (brownish when dried and relatively large vs yellowish and smaller), 4) fruit pilosity (very short white hairs on the lower half of fruit vs glabrous except for a few long hairs at apex), 5) fruit colour (dull black vs shiny reddish).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4064*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.3$ cm.

[1545] *Casearia* sp. D

HERBARIUM DATA (FG). — A single collection, *M.-F. Prévost & D. Sabatier 2280*.

INVENTORY DATA (FG). — 1 tree, $dbh = 10$ cm.

Genus *Hasseltia* Kunth

[1546] *Hasseltia floribunda* Kunth

Nova genera et species plantarum [H.B.K.] 7: 232 [25 Apr. 1825] (Kunth 1825).

Hasseltia pubescens Benth., *Pl. Hartw. [Bentham]*: 164 [before mid-Nov. 1845] (Bentham 1845).

Banara laxiflora Benth., *J. Proc. Linn. Soc., Bot. 5 (Suppl. 2)*: 91 [May 1861] (Bentham 1861). — *Hasseltia laxiflora* (Benth.) Eichler, *Fl. Bras. [Martius] 13 (1)*: 502 [1 Oct. 1871] (Eichler 1871).

Hasseltia peruviana Pilg., *Verb. Bot. Vereins Prov. Brandenburg 47*: 160 [1 Oct. 1905] (Pilger 1905).

Hasseltia tomentulosa Cuatrec., *Fieldiana, Bot. 27 (1)*: 98 (Cuatrecasas 1950).

Hasseltia monagensis Steyermark., *Fieldiana, Bot. 28 (2)*: 407 (Steyermark 1952).

Hasseltia panamensis Standl. ex Dayton, *Phytologia 4 (4)*: 253 [June 1953] (Dayton 1953), *nom. nud.*

Hasseltia micrantha L.O.Williams, *Fieldiana, Bot. 29 (6)*: 361 (Williams 1961).

Hasseltia psittacarum L.O.Williams, *Fieldiana, Bot. 31 (1-2)*: 28 [25 Nov. 1964] (Williams 1964).

Hasseltia rigida Woodson ex A.Robyns, *Ann. Missouri Bot. Gard. 55 (1)*: 77 (Robyns 1968).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5683*.

INVENTORY DATA (FG). — 28 trees in 9 plots; $F_{\max} = 3\%$; $dbh_{\text{inv}} = 47.7$ cm.

Genus *Homalium* Jacq.

[1547] *Homalium guianense* (Aubl.) Oken

Allg. Naturgesch. 3 (2): 810 (Oken 1841). — *Racoubea guianensis* Aubl., *Hist. Pl. Guiane 1*: 590 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Homalium spicatum* Lam., *Encycl. [J. Lamarck et al.] 1 (1)*: 32 [2 Dec. 1783] (Lamarck 1783), *nom. illeg. superfl.* (based on *Racoubea guianensis*). — *Homalium racoubea* Sw., *Prod. [Swartz] 86 [20 Jun.-29 July 1788]* (Swartz 1788), *nom. illeg. superfl.* (based on *Racoubea guianensis*).

Napimoga guianensis Aubl., *Hist. Pl. Guiane 1*: 592 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Homalium napimoga* Spreng., *Syst. Veg. [Sprengel] 4 (2, Cur. Post.)*: 210 [Jan.-June 1827] (Sprengel 1827), *nom. illeg. superfl.* (based on *Napimoga guianensis*).

Homalium surinamense Steud., *Flora 26 (45)*: 756 [17 Dec. 1843] (Steudel 1843).

Homalium densiflorum Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 4*: 36 [“1860” publ. July 1859] (Bentham 1859).

Homalium puberulum Klotzsch ex Eichler, *Fl. Bras. [Martius] 13 (1)*: 507 [1 Oct. 1871] (Eichler 1871).

Homalium mattogrossense Malme, *Ark. Bot. 23A(4)*: 20 (Malme 1930).

Homalium choocoense Cuatrec., *Fieldiana, Bot. 27 (1)*: 99 (Cuatrecasas 1950).

VERNACULAR NAMES. — Pa: tahuma • Ka: alekikolan, masalaipyo • Wp: yape'a pitā sili, yawa luway • Cr: bwa-anmē.

HERBARIUM DATA (FG). — 46 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Racoubea guianensis*: BM[[BM000624532](#)]).

SIZE. — Up to 12 cm dbh (Sleumer 1980).

[1548] *Homalium racemosum* Jacq.

Enum. Syst. Pl. 24 [Aug.-Sep. 1760] (Jacquin 1760).

Homalium senarium Sessé & Moc. ex DC., *Prodr. [A. P. de Candolle] 2*: 54 [mid Nov. 1825] (Candolle 1825).

Homalium obtusatum Turcz., *Bull. Soc. Imp. Naturalistes Moscou 31 (1)*: 465 (Turczaninow 1858).

Homalium pedicellatum Spruce ex Benth., *J. Proc. Linn. Soc., Bot. 4*: 36 [“1860” publ. July 1859] (Bentham 1859).

Homalium hondurensse Donn.Sm., *Bot. Gaz. 20 (1)*: 4 [18 Jan. 1895] (Donnell Smith 1895).

Homalium integrifolium Britton, *Bull. Torrey Bot. Club 37 (7)*: 354 (Britton 1910), *nom. illeg. hom., non* (Lam.) Baill. (Baillon 1886).

Homalium trichostemon S.F.Blake, *Contr. Gray Herb. 53*: 60 [26 Feb. 1918] (Blake 1918).

Homalium nicaraguense S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 225 [9 Sep. 1919] (Blake 1919).

Homalium mollicellum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 226 [9 Sep. 1919] (Blake 1919).

Homalium pleiandrum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 227 [9 Sep. 1919] (Blake 1919).

Homalium hemistylum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 228 [9 Sep. 1919] (Blake 1919).

Homalium leiogynum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 228 [9 Sep. 1919] (Blake 1919).

Homalium racemosum subsp. *barbellatum* S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 229 [9 Sep. 1919] (Blake 1919).

Homalium pittieri S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 230 [9 Sep. 1919] (Blake 1919).

Homalium trichocladum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 231 [9 Sep. 1919] (Blake 1919).

Homalium eleutherostylum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 232 [9 Sep. 1919] (Blake 1919).

Homalium columbianum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 233 [9 Sep. 1919] (Blake 1919).

Homalium eurypetalum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 234 [9 Sep. 1919] (Blake 1919).

Homalium stenosepalum S.F.Blake, *Contr. U.S. Natl. Herb. 20 (7)*: 234 [9 Sep. 1919] (Blake 1919).

Homalium riparium Standl., *Publ. Carnegie Inst. Wash. 461*: 74 [26 Nov. 1935] (Standley 1935), *nom. illeg. hom., non* Gilg (1908).

Homalium schippii Standl., *Trop. Woods 52*: 27 (Standley 1937).

Homalium anzoateguiense Steyermark., *Fieldiana, Bot. 28 (2)*: 410 (Steyermark 1952).

Homalium mituense Cuatrec., *Trop. Woods 101*: 19 (Cuatrecasas 1955).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J. Martin s.n.* ([P\[P04822073\]](#)).

SIZE. — Brazil, Maranhão. *D.C. Daly D649* (MO), 15 m × 25 cm.

Genus *Piparea* Aubl.

[1549] *Piparea dentata* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 31 [Jun.-Dec. 1775] (Aublet 1775). — *Casearia dentata* (Aubl.) Eichler, *Fl. Bras. [Martius]* 13 (1): 484 [1 Oct. 1871] (Eichler 1871), nom. illeg. hom., non Sessé & Moc. ex DC. (Candolle 1825). — *Alsodeia piparea* Spreng., *Syst. Veg. [Sprengel]* 1: 807 [1825 publ. late 1824] (Sprengel 1824), nom. illeg. superfl. (based on *Piparea dentata*). — *Conohoria aubletii* D.Dietr., *Syn. Pl. [D. Dietrich]* 1: 831 [July 1839] (Dietrich 1839), nom. illeg. superfl. (based on *Piparea dentata*).

Casearia commersoniana Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 2 (16): 235 ["1829" publ. Oct.-Nov. 1830] (Cambessèdes 1830).

Casearia densiflora Benth., *J. Bot. [Hooker]* 4: 113 (Bentham 1842).

Casearia laurifolia Benth., *J. Bot. [Hooker]* 4: 113 (Bentham 1842).

Casearia lasiosperma Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 17: 113 (Triana & Planchon 1862).

Casearia congestiflora Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 608 (Turczaninow 1863).

Casearia myriantha Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 609 (Turczaninow 1863). — *Casearia javitensis* var. *myriantha* (Turcz.) L.O.Williams, *Fieldiana, Bot.* 29 (6): 359 (Williams 1961).

Casearia densiflora var. *parviflora* Eichler, *Fl. Bras. [Martius]* 13 (1): 487 [1 Oct. 1871] (Eichler 1871).

Casearia maximilianii Eichler, *Fl. Bras. [Martius]* 13 (1): 487 [1 Oct. 1871] (Eichler 1871), "maximiliani".

Casearia miradorensis Eichler, *Fl. Bras. [Martius]* 13 (1): 487 [1 Oct. 1871] (Eichler 1871).

Casearia brighamii Watson, *Proc. Amer. Acad. Arts* 21: 459 (Watson 1886), "Brighami".

Casearia piparea Sleumer, *Lilloa* 23: 249 (Sleumer 1950).

NOTE. — The genus *Piparea* Aubl., once sunked into *Casearia* Jacq., has been resurrected by Samarakoon & Alford (Samarakoon & Alford 2019).

VERNACULAR NAMES. — Pa: tahuma • Wp: kasilikita'i sili, kasilikita'i sili, kwasilikita'i sili • Br: cafezinho-do-mato, canela-de-veado.

HERBARIUM DATA (FG). — 87 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material of *Piparea dentata*: BM[[BM000624376](#)]).

SIZE. — Up to 20 cm dbh (Sleumer 1980).

[1550] *Piparea multiflora* C.F.Gaertn.

Suppl. Carp. 231 (Gaertner 1807). — *Casearia gaertneri* Donn. Sm., *Enum. Pl. Guatem.*: 25 (Donnell Smith 1891), nom. illeg. superfl. (based on *Piparea multiflora*, excl. specim. cit. Donn. Sm., which is *Casearia commersoniana* Cambess.).

Casearia javitensis Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 366 [24 Mar. 1823] (Kunth 1823), in nota, under *Lindleya glabra*. — *Chaetocrater javitensis* (Kunth) Raf., *Sylva Tellur.* 149 (Rafinesque 1838).

Casearia iquitosensis J.F.Macbr., *Candollea* 8: 22 (Macbride 1940).

NOTE. — The genus *Piparea* Aubl., once sunked into *Casearia* Jacq., has been resurrected by Samarakoon & Alford (Samarakoon & Alford 2019).

VERNACULAR NAMES. — Pa: kanegma, yawu-seine • Ka: alawata muleli • Wp: kasilikita'i, kasilikita'i, kwasilikita'i • Wn: sieuju • Nt: kwasikwasi tiki • Cr: kafé-sovaj • Br: capança.

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *J.-B. Patris* s.n. (original material of *Piparea multiflora*: G-DC, [G00208434](#)).

INVENTORY DATA (FG). — 162 trees in 100 plots; $F_{\max} = 1.7\%$; $dbh_{inv} = 29.6$ cm.

Genus *Ryania* Vahl

[1551] *Ryania pyrifera* (Rich.) Uittien & Sleumer

Fl. Suriname 3 (1): 286 (Uittien & Sleumer 1935). — *Patrisia pyrifera* Rich., *Actes Soc. Hist. Nat. Paris* 1: 110 [Oct. 1792] (Richard 1792).

Ryania sagotiana Eichler, *Fl. Bras. [Martius]* 13 (1): 491 [1 Oct. 1871] (Eichler 1871). — *Patrisia sagotiana* (Eichler) Kuntze, *Revis. Gen. Pl.* 1: 45 [5 Nov. 1891] (Kuntze 1891).

NOTE. — Excluded synonym: *Ryania speciosa* Vahl.

VERNACULAR NAMES. — Pa: tahuma seine.

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *J.B. Leblond* s.n. (original material P[[P02442081](#)]).

SIZE. — Up to 10 cm dbh (Sleumer 1980).

[1552] *Ryania speciosa* Vahl var. *bicolor* (DC.) Monach. (Fig. 52D)

Lloydia 12: 19 (Monachino 1949). — *Patrisia bicolor* DC., *Prodri. [A. P. de Candolle]* 1: 256 [mid Jan. 1824] (Candolle 1824). — *Ryania candollei* Miq., *Ann. Mag. Nat. Hist.* 11: 16 (Miquel 1843), nom. illeg. superfl. (based on *Patrisia bicolor*). — *Ryania bicolor* (DC.) Eichler, *Fl. Bras. [Martius]* 13 (1): 491 [1 Oct. 1871] (Eichler 1871).

VERNACULAR NAMES. — Pa: kuuku-ariut-seine • Ka: tomoipo • Wp: iwkwiši, mukwiši.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-B. Patris* s.n. (original material G-DC, [G00207421](#)).

INVENTORY DATA (FG). — 1 tree, dbh = 10.2 cm.

[1553] *Ryania speciosa* var. *subuliflora* (Sandwith) Monach.

Lloydia 12: 14 (Monachino 1949). — *Ryania pyrifera* var. *subuliflora* Sandwith, *J. Arnold Arbor.* 24 (2): 219 [15 Apr. 1943] (Sandwith 1943).

VERNACULAR NAMES. — Pa: kuuku-ariut • Wp: kasilikita'i, kasilikita'i, kwasilikita'i • Br: capança, mata-calado.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *R.A.A. Oldeman* 2386.

SIZE. — Brazil, Amazonas. *L.O. Adão Teixeira* 943 (MO), 8 m × 10 cm.

[1554] *Rymania speciosa* var. *tomentosa* (Miq.) Monach.

Lloydia 12: 16 (Monachino 1949). — *Rymania tomentosa* Miq., *Ann. Mag. Nat. Hist.* 11: 15 (Miquel 1843). — *Patrisia tomentosa* (Miq.) M. Roem., *Fam. Nat. Syn. Monogr.* 2: 136 (Roemer 1846). — *Rymania pyrifera* var. *tomentosa* (Miq.) Sleumer ex Sleumer & Uittien, *Fl. Suriname* 3 (1): 286 (Sleumer & Uittien 1935).

Patrisia parviflora DC., *Prod. [A. P. de Candolle]* 1: 256 [mid Jan. 1824] (Candolle 1824). — *Rymania patrisii* Miq., *Ann. Mag. Nat. Hist.* 11: 16 (Miquel 1843), nom. illeg. superfl. (based on *Patrisia parviflora*). — *Rymania parviflora* (DC.) Griseb., *Fl. Brit. W.I. [Grisebach]*: 296 [late 1860] (Grisebach 1860).

Rymania casiquiarensis Steyermark., *Fieldiana, Bot.* 28 (2): 411 (Steyermark 1952).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J.-B. Patris s.n.* (original material of *Patrisia parviflora*: P[P02442059]).

SIZE. — Colombia, Amazonas. *A. Rudas L. 3391* (MO), 10 m × 15.9 cm.

Genus *Xylosma* G. Forst.

[1555] *Xylosma benthamii* (Tul.) Triana & Planch.

Ann. Sci. Nat., Bot. sér. 4, 17: 99 (Triana & Planchon 1862). — *Flacourtia benthamii* Tul., *Ann. Sci. Nat., Bot. sér. 3*, 7: 291 (Tulasne 1847), “*benthamii*”. — *Hisingera benthamii* (Tul.) Clos, *Ann. Sci. Nat., Bot. sér. 4*, 8: 225 (Clos 1857). — *Myroxylon benthamii* (Tul.) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Flacourtia nitida Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 119 (Bentham 1851). — *Hisingera lucens* Clos, *Ann. Sci. Nat., Bot. sér. 4*, 8: 225 (Clos 1857), nom. illeg. superfl. (based on *Flacourtia nitida*). — *Myroxylon lucens* Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Xylosma digyna Eichler, *Fl. Bras. [Martius]* 13 (1): 447 [1 Oct. 1871] (Eichler 1871), “*digynum*”. — *Flacourtia digyna* Benth. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 447 [1 Oct. 1871] (Eichler 1871), nom. nud. pro syn.

Myroxylon digynum Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Xylosma armata J.F. Macbr., *Candollea* 5: 391 (Macbride 1934).

Xylosma pilosa J.F. Macbr., *Candollea* 5: 391 (Macbride 1934).

Xylosma pallidifolia Steyermark & O. Huber, *Fl. Avila*: 425 (Steyermark & Huber 1978), “*pallidifolium*”, nom. nud. pro syn.

NOTES. — Being based on *Flacourtia nitida*, *Hisingera lucens* is superfluous and illegitimate. Therefore it cannot serve as the basionym of *Myroxylon lucens*.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (P[P04734493]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.5$ cm.

[1556] *Xylosma ciliatifolia* (Clos) Eichler

Fl. Bras. [Martius] 13 (1): 449 [1 Oct. 1871] (Eichler 1871), “*ciliatifolium*”. — *Hisingera ciliatifolia* Clos, *Ann. Sci. Nat., Bot. sér. 4*, 8: 223 (Clos 1857). — *Myroxylon ciliatifolium* (Clos) Kuntze, *Revis. Gen. Pl.* 1: 44 [5 Nov. 1891] (Kuntze 1891).

Xylosma pou-orfilae Herter, *Revista Sudamer. Bot.* 7: 214 (Herter 1943), nom. nud.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P. Grenand & M.-F. Prévost 1959*.

INVENTORY DATA (FG). — 1 tree, $dbh = 10.2$ cm.

[1557] *Xylosma tessmannii* Sleumer

Notizbl. Bot. Gart. Berlin-Dahlem 12: 477 [30 June 1935] (Sleumer 1935).

NOTE. — According to Sleumer (1980), this species is close to *X. benthamii*, perhaps only a luxuriant form of it.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost 3302*.

SIZE. — Up to 40 cm dbh (Sleumer 1980).

Family SAPINDACEAE Juss.

Genus *Allophylus* L.

[1558] *Allophylus edulis* (A.St.-Hil., A.Juss. & Cambess.)

Hieron. ex Niederl.

(Fig. 53A)

Bol. Mens. Mus. Prod. Argent. 3 (29): 180 [Oct. 1890] (Niederlein 1890). — *Schmidelia edulis* A.St.-Hil., A.Juss. & Cambess., *Pl. Usuel. Bras.* 14: t. 67 [4 Aug. 1828] (Saint-Hilaire et al. 1828).

Allophylus edulis var. *gracilis* Radlk., *Fl. Bras. [Martius]* 13 (3): 485 [1 Apr. 1900] (Radlkofe 1900).

Allophylus edulis var. *subsessilis* Huber, *Bull. Soc. Bot. Genève*, sér. 2, 6: 185 [“1914” publ. 1915] (Huber 1915).

NOTES. — Acevedo-Rodríguez stated (pers. comm. 2015) that he mistakenly treated this taxon as *A. angustatus* (Triana & Planch.) Radlk. in (Acevedo-Rodríguez 2012). A. Jussieu and Cambessèdes are co-authors of parts 9–14 (pl. 41–70) of *Plantes usuelles des Brésiliens* (Saint-Hilaire et al. 1828).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2356*.

INVENTORY DATA (FG). — 19 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{inv} = 22.3$ cm.

[1559] *Allophylus latifolius* Huber

Bull. Soc. Bot. Genève, sér. 2, 6: 186 [“1914” publ. 1915] (Huber 1915).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 14752*.

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{inv} = 13.5$ cm.



FIG. 52. — Salicaceae: **A**, *Casearia grandiflora* Cambess. (*D. Sabatier & J.-F. Molino* 5187); **B**, *Casearia* sp. B (*J.-F. Molino & D. Sabatier* 2469); **C**, *Casearia ulmifolia* Vahl ex Vent. (*D. Sabatier & J.-F. Molino* 5006bis); **D**, *Ryania speciosa* Vahl var. *bicolor* (DC.) Monach. (*J.-F. Molino et al.* 1852). A, D, © D. Sabatier/IRD; B, C, © J.-F. Molino/IRD.

[1560] *Allophylus leucoclados* Radlk.

Fl. Bras. [Martius] 13 (3): 473 [1 Apr. 1900] (Radlkofe 1900).

VERNACULAR NAMES. — Wp: tapi'i lo wapa'a.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *C. Sastre* 1621, height 15 m.

[1561] *Allophylus robustus* Radlk.

Monogr. Paullinia: 185 (Radlkofe 1895).

HERBARIUM DATA (FG). — 27 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (original material G-DC, G00008481).

INVENTORY DATA (FG). — 4 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 17$ cm.

Genus *Cupania* L.

[1562] *Cupania diphylla* Vahl

Eclog. Amer. 3: 9 (Vahl 1807).

Cupania geminata Poir., *Encycl. [J. Lamarck et al.] Suppl.* 2: 419 [3 July 1812] (Poiret 1812).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *Unknown coll. s.n.* ("Herb. Poir.") (probable original material of *Cupania geminata*: P[P06671081]).

INVENTORY DATA (FG). — 10 trees in 9 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.6$ cm.

[1563] *Cupania hirsuta* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 565 (Radlkofe 1879).

Cupania velutina Klotzsch, *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1180 ["1848" publ. 7-10 Mar. 1849] (Klotzsch 1849), *nom. nud.*

Cupania lanuginosa Sagot ex Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 566 (Radlkofe 1879).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *P.A. Sagot* 1214, Jan. 1859 (holotype of *Cupania lanuginosa*: P[P06671233]; iso-, P[P06671232, P06671234, P06671235]; possible iso-, F[V0071577F, V0071578F], NY[00337808]).

INVENTORY DATA (FG). — 27 trees in 18 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 19\text{ cm}$.

[1564] *Cupania macrostylis* (Radlk.) Acev.-Rodr.

Fl. Guianas, ser. A, 29: 25 (Acevedo-Rodríguez 2012). — *Matayba macrostylis* Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 625 (Radlkofe 1879).

Matayba macrolepis Radlk., *Fl. Bras. [Martius]* 13 (3): 605 [1 Apr. 1900] (Radlkofe 1900).

Trichilia caudata Killip & Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 5: 36 [Jan.-June 1942] (Killip & Cuatrecasas 1942).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier* 2401.

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 15.5\text{ cm}$.

[1565] *Cupania rubiginosa* (Poir.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 566 (Radlkofe 1879). — *Robinia rubiginosa* Poir., *Encycl. J. Lamarck et al.* 6 (1): 227 [2 Oct. 1804] (Poiret 1804).

Cupania subrepanda Mart., *Flora* 21 (2, Beibl.): 70 (Martius 1838).

VERNACULAR NAMES. — Pa: mbagwi, mbagwi-seine, mbarui, mbarui-seine • Ka: tohmopala, tonolo ipyo, tonolo polipyro • Te: tononipö • Wp: kalima'i, kalima'i sili, kalima'i yowa • Nt: singabusu • Cr: gangouti-blanc • Br: pau-de-arapuce.

HERBARIUM DATA (FG). — 70 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2792.

INVENTORY DATA (FG). — 24 trees in 19 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 19.9\text{ cm}$.

[1566] *Cupania scrobiculata* Rich.
(Fig. 53B)

Actes Soc. Hist. Nat. Paris 1: 109 [Oct. 1792] (Richard 1792).

Cupania reticulata Cambess., *Mém. Mus. Hist. Nat.* 18: 41 (Cambessèdes 1829). — *Cupania scrobiculata* f. *reticulata* (Cambess.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 565 (Radlkofe 1879). — *Cupania scrobiculata* var. *reticulata* (Miq.) Uittien, *Fl. Suriname* 2 (1): 389 (Uittien 1937).

Cupania frondosa Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 199 (Bentham 1851). — *Cupania scrobiculata* f. *frondosa* (Benth.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.*

München 9: 565 (Radlkofe 1879). — *Cupania scrobiculata* var. *frondosa* (Benth.) Uittien, *Fl. Suriname* 2 (1): 390 (Uittien 1937).

Cupania porosa Miq., *Stirp. Surinam. Select.*: 70 ["1850" publ. Mar. 1851] (Miquel 1851).

Cupania guianensis Miq., *Stirp. Surinam. Select.*: 71 ["1850" publ. Mar. 1851] (Miquel 1851). — *Cupania scrobiculata* f. *guianensis* (Miq.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 565 (Radlkofe 1879).

Cupania obovata Gleason, *Bull. Torrey Bot. Club* 54 (8): 616 [Nov. 1927] (Gleason 1927).

Cupania auriculata Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 8* (1): 20 (Standley 1930).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: mbagwi, mbagwi-seine, mbarui, mbarui-seine • Ka: tohmopala, tonolo ipyo, tonolo polipyro • Te: tononipö • Wp: kalima'i, kalima'i sili, kalima'i yowa • Nt: singabusu • Cr: gangouti-blanc • Br: pau-de-arapuce.

HERBARIUM DATA (FG). — 113 collections at CAY. Sel. exs.: *L.C. Richard* s.n. (holo-, P[P00117101; iso-, P00117102, P00117103, P00117104, P00117105, P00117106]).

INVENTORY DATA (FG). — 331 trees in 120 plots; $F_{\max} = 9.7\%$; $\text{dbh}_{\text{inv}} = 38\text{ cm}$.

Genus *Matayba* Aubl.

[1567] *Matayba arborescens* (Aubl.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 353 (Radlkofe 1879). — *Sapindus arborescens* Aubl., *Hist. Pl. Guiane* 1: 357 [Jun.-Dec. 1775] (Aublet 1775).

Cupania laevigata Rich., *Actes Soc. Hist. Nat. Paris* 1: 109 [Oct. 1792] (Richard 1792), "Laevigata".

Cupania aubletii Miq., *Stirp. Surinam. Select.*: 67 ["1850" publ. Mar. 1851] (Miquel 1851).

Matayba reducta Steyermark., *Fieldiana, Bot.* 28 (2): 351 (Steyermark 1952).

NOTE. — The original material of *Sapindus arborescens* at BM (BM000884118) is labelled "TYPE SPECIMEN *Sapindus frutescens* Aubl. Pl. Gui. 1: 355. T 138", while that of *S. frutescens* Aubl. [synonym of *Pseudima frutescens* (Aubl.) Radlk.] (BM000884091) is labelled "TYPE SPECIMEN *Sapindus arborescens* Aubl. Pl. Gui. 1: 357. T 137" (Acevedo-Rodríguez 2012: 38).

VERNACULAR NAMES. — Ka: tonolo ipyo, tonolo polipyro • Te: tononipö • Wp: kalima'i, kalima'i sili, yatoa'i sili • Nt: singabusu • Cr: sanvonié • Br: breu-de-tucano.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (original material BM[BM000884118]).

INVENTORY DATA (FG). — 10 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 27.1\text{ cm}$.

[1568] *Matayba guianensis* Aubl.

Hist. Pl. Guiane 1: 331 [Jun.-Dec. 1775] (Aublet 1775). — *Matayba guianensis* f. *genuina* Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 631 (Radlkofer 1879), nom. inval. (Turland et al. 2018: Art. 24.3).

VERNACULAR NAMES. — Wn: asisimë, sihkëimëimë.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000884102]).

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 30.1 \text{ cm}$.

[1569] *Matayba inelegans* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 535 (Radlkofer 1879). — *Cupania inelegans* Spruce ex Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 517 (Radlkofer 1879), nom. nud. pro syn.

Matayba sororopaniana Steyerl., *Fieldiana, Bot.* 28 (2): 348 (Steyermark 1952).

NOTE. — Although Spruce's label on type reads “*Cupania inelegans* Spruce”, Spruce never made the combination *Matayba inelegans*. Authorship is thus Radlk., not “Spruce ex Radlk.”

VERNACULAR NAMES. — Pa: kwepan-wašiunu • Ka: tonolo ipyo, tonolo polipyo • Nt: geli udu.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 3348.

INVENTORY DATA (FG). — 126 trees in 27 plots; $F_{\max} = 10.9\%$; $\text{dbh}_{\text{inv}} = 33.9 \text{ cm}$.

[1570] *Matayba laevigata* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 628 (Radlkofer 1879). — *Cupania laevigata* Miq., *Stirp. Surinam. Select.*: 72 [“1850” publ. Mar. 1851] (Miquel 1851), nom. illeg. hom., non Rich. (Richard 1792) [synonym of *Matayba guianensis*].

Matayba guianensis subf. *subovalis* Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 9: 632 (Radlkofer 1879).

Matayba miquelii Uittien, *Fl. Suriname* 2 (1): 384 (Uittien 1937).

NOTES. — *Cupania laevigata* Miq. is illegitimate, thus cannot be the basionym of *Matayba laevigata* Radlk. Although published as a subforma of the invalid forma *genuina*, *M. guianensis* subf. *subovalis* Radlk. is legitimate.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *F.M.R. Leprieur* 333 (original material of *Matayba guianensis* subf. *subovalis*; L[L0014432, L0014433]).

INVENTORY DATA (FG). — 9 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 32.7 \text{ cm}$.

[1571] *Matayba opaca* Radlk.
(Fig. 53C)

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 628 (Radlkofer 1879).

Matayba fallax Radlk., *Fl. Bras. [Martius]* 13 (3): 612 [1 Apr. 1900] (Radlkofer 1900). — *Matayba opaca* var. *fallax* (Radlk.) Uittien, *Fl. Suriname* 2 (1): 384 (Uittien 1937).

Matayba affinis Steyerl., *Ann. Missouri Bot. Gard.* 75 (3): 1067 [19 Oct. 1988] (Steyermark 1988).

VERNACULAR NAMES. — Ka: tikalayen tonoloipyo • Wp: wila pisi'u • Wn: jala.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier* 2102.

INVENTORY DATA (FG). — 50 trees in 8 plots; $F_{\max} = 3.4\%$; $\text{dbh}_{\text{inv}} = 28.3 \text{ cm}$.

[1572] *Matayba peruviana* Radlk. subsp. *oligandra*
(Sandwith) T.D.Penn. ex Acev.-Rodr.

Fl. Guianas, ser. A, 29: 46 (Acevedo-Rodríguez 2012). — *Matayba oligandra* Sandwith, *Bull. Misc. Inform. Kew* 1935 (3): 123 [20 May 1935] (Sandwith 1935).

Trichilia ptarihana Steyerl., *Fieldiana, Bot.* 28 (2): 278 (Steyermark 1952).

Matayba oligandra var. *ptarihana* (Steyerl.) Steyerl., *Bol. Soc. Venez. Ci. Nat.* 26: 426 (Steyermark 1966).

Matayba chimantensis Steyerl. & Maguire, *Mem. New York Bot. Gard.* 17 (1): 447 (Steyermark & Maguire 1967).

Matayba jauaensis Steyerl., *Bol. Soc. Venez. Ci. Nat.* 33: 347 (Steyermark 1976).

Matayba oligandra var. *occidentalis* Steyerl., *Ann. Missouri Bot. Gard.* 75 (3): 1069 [19 Oct. 1988] (Steyermark 1988).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori* & *T.D. Pennington* 17937.

SIZE. — Up to 20 m tall (Acevedo-Rodríguez 2012).

[1573] *Matayba stenodictya* Radlk.

Fl. Bras. [Martius] 13 (3): 629 [1 Apr. 1900] (Radlkofer 1900).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier* et al. 4712.

INVENTORY DATA (FG). — 12 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 44.6 \text{ cm}$.

Genus *Melicoccus* P.Browne

[1574] *Melicoccus pedicellaris* (Sagot ex Radlk.) Acev.-Rodr.

Fl. Neotrop. Monogr. 87: 30 [22 May 2003] (Acevedo-Rodríguez 2003). — *Talisia pedicellaris* Sagot ex Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 8: 342 (Radlkofer 1878).

Talisia pulverulenta Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 8: 342 (Radlkofer 1878).

NOTE. — Radlkofe (1878) did ascribe the binomial *Talisia pedicellaris* to Sagot, by stating: “Der speciesbeiname ist aus dem von Sagot der Pflanze beigelegten Namen [...]”.

VERNACULAR NAMES. — Wp: tuli átā u.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *P.A. Sagot 1188*, Aug. 1858 (lecto-, P, designated by Acevedo-Rodríguez [2003: 31], not seen, fragm. M[M-0224883]; isolecto-, BM[BM000884044], BR[BR000005417018], F[V0071843F, V0071844F, V0071845F], GOET[GOET010791], K[K000035428, K000035429], MPU[MPU010928], P[P00117093, P00117094, P00117095, P00117096, P00117097, P00117098], U[U0006556]; possible isolecto-, P[P02428274, P02428275]).

INVENTORY DATA (FG). — 80 trees in 52 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 27.7\text{ cm}$.

Genus *Pentascyphus* Radlk.

[1575] *Pentascyphus thyrsiflorus* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 539 (Radlkofe 1879).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *F.M.R. Leprieur 335* (holo-, P[P00214699]; iso-, L[L0014517]).

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 24.8\text{ cm}$.

Genus *Pseudima* Radlk.

[1576] *Pseudima frutescens* (Aubl.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 300 (Radlkofe 1878). — *Sapindus frutescens* Aubl., *Hist. Pl. Guiane* 1: 355 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — The original material of *Sapindus frutescens* at BM (BM000884091) is labelled “TYPE SPECIMEN Sapindus arborescens Aubl. Pl. Gui. 1: 357. T 137”, while that of *S. arborescens* Aubl. [synonym of *Matayba arborescens* (Aubl.) Radlk.] (BM000884118) is labelled “TYPE SPECIMEN Sapindus frutescens Aubl. Pl. Gui. 1: 355. T 138” (Acevedo-Rodríguez 2012: 38).

VERNACULAR NAMES. — Pa: tuu-ariut, tuu-kamwi-seine • Ka: tiyawasisyen • Wp: yikisiñi, yikisiñi lo • Cr: sanvonié • Br: camaá, fruta-de-anel, pau-de-gafanhoto, pitomba-de-macaco.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000884091]).

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 44.1\text{ cm}$.

Genus *Scyphonymchium* Radlk.

[1577] *Scyphonymchium multiflorum* (Mart.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 9: 519 (Radlkofe 1879). — *Cupania multiflora* Mart., *Flora* 21 (2, Beibl.): 75 (Martius 1838).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 3479*.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 19.3\text{ cm}$.

Genus *Talisia* Aubl.

[1578] *Talisia clathrata* Radlk. subsp. *canescens* Acev.-Rodr.

Fl. Neotrop. Monogr. 87: 56 [22 May 2003] (Acevedo-Rodríguez 2003).

VERNACULAR NAMES. — Nt: baala wiwii.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *P. Acevedo-Rodríguez et al. 5012* (holo-, US[00614239]); iso-, A, CAY[CAY001980, CAY001981], CTES, F, HUA, K, NY, P[P04857357], L[exU]).

INVENTORY DATA (FG). — 16 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 25.6\text{ cm}$.

[1579] *Talisia furfuracea* Sandwith

Bull. Misc. Inform. Kew 1935 (3): 121 [20 May 1935] (Sandwith 1935).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3663*.

INVENTORY DATA (FG). — 46 trees in 39 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 47.1\text{ cm}$.

[1580] *Talisia guianensis* Aubl.

Hist. Pl. Guiane 1: 349 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Talisia rosea* Vahl, *Ectog. Amer.* 2: 30 (Vahl 1798), nom. illeg. superfl. (based on *Talisia guianensis*).

Talisia glabra DC., *Prodr. [A. P. de Candolle]* 1: 609 [mid Jan. 1824] (Candolle 1824).

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000838014]); *P. Acevedo-Rodríguez et al. 4871* (MO), 10 m tall.

[1581] *Talisia hemidasya* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 349 (Radlkofe 1878).

Talisia glandulifera Steyermark., *Ann. Missouri Bot. Gard.* 75 (3): 1072 [19 Oct. 1988] (Steyermark 1988).

VERNACULAR NAMES. — Ka: sepupi, supupi, tonolo ipyo • Wp: tuli átā • Cr: bwa-flanbo, payakousa • Br: pitomba-do-norte, pitombeira.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *S.A. Mori et al. 15027* (holotype of *Talisia glandulifera*: MO[MO-256425]); iso-, NY[00039423]).

INVENTORY DATA (FG). — 9 trees in 4 plots; $F_{\max} = 1\%$; $\text{dbh}_{\text{inv}} = 26.5\text{ cm}$.

[1582] *Talisia hexaphylla* Vahl

Ectog. Amer. 2: 29 (Vahl 1798).

Talisia panamensis Pittier, *Contr. U.S. Natl. Herb.* 20 (3): 129 [18 June 1918] (Pittier 1918).

Talisia cararensis Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (31): 306 [Dec. 1951] (Cuatrecasas 1951).

VERNACULAR NAMES. — Pa: miumiu-asiru-washiunu, tuu-ariut • Ka: wa:u • Te: tuli • Wn: kälépukaju • Nt: gaan tatu tiki, tatu, tatu udu • Cr: bwa-flanbo, payakousa • Br: pitomba-do-norte, pitombeira.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4631*.

INVENTORY DATA (FG). — 198 trees in 107 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 46.5$ cm.

[1583] *Talisia longifolia* (Benth.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 348 (Radlkofer 1878). — *Cupania longifolia* Benth., *Hooker's J. Bot. Kew Gard. Misc.* 2: 211 (Bentham 1850).

VERNACULAR NAMES. — Wp: tuli áta • Wn: täpu • Nt: tatu • Cr: bwa-flanbo, payakousa • Br: pitomba-do-norte, pitombeira.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2872*.

INVENTORY DATA (FG). — 7 trees in 6 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.5$ cm.

[1584] *Talisia macrophylla* (Mart.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 347 (Radlkofer 1878). — *Cupania macrophylla* Mart., *Flora* 21 (2, Beibl.): 74 (Martius 1838).

Talisia elephantipes Sandwith, *J. Bot.* 78: 256 (Sandwith 1940).

Talisia allenii Croat, *Ann. Missouri Bot. Gard.* 63: 527 (Croat 1976).

Talisia pentantha Steyermark., *Ann. Missouri Bot. Gard.* 75 (3): 1075 [19 Oct. 1988] (Steyermark 1988).

Talisia sancarlosiana Steyermark., *Ann. Missouri Bot. Gard.* 75 (3): 1075 [19 Oct. 1988] (Steyermark 1988).

VERNACULAR NAMES. — Te: tuli • Wp: tuli áta u • Cr: bwa-flanbo, payakousa.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3253*.

INVENTORY DATA (FG). — 8 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15$ cm.

[1585] *Talisia megaphylla* Sagot ex Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 350 (Radlkofer 1878).

VERNACULAR NAMES. — Pa: tuu-aška, tuu-kamwi • Ka: malay wokili, tulisi, tulusili.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *P.A. Sagot 1194*, 1857 (lecto-, P[P06694733], designated by Acevedo-Rodríguez [2003: 136]; isolecto-, BR[BR000005417100], K[K000035531, K000035532]).

INVENTORY DATA (FG). — 8 trees in 8 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.5$ cm.

[1586] *Talisia microphylla* Uittien

Recueil Trav. Bot. Néerl. 34: 483 (Uittien 1937).

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3855*.

INVENTORY DATA (FG). — 46 trees in 34 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 41.1$ cm.

[1587] *Talisia mollis* Kunth ex Cambess.

Mém. Mus. Hist. Nat. 18: 48 (Cambessèdes 1829).

Tapirocarpus talisia Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 13: 292 (Sagot 1882), *pro parte vegetativa*.

VERNACULAR NAMES. — Wp: tuli áta, tuli áta u, tuli áta yowa • Nt: tatu • Cr: bwa-flanbo, payakousa • Br: pitomba-brava.

HERBARIUM DATA (FG). — 52 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-4183*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 12.4$ cm.

[1588] *Talisia pachycarpa* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 350 (Radlkofer 1878).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holo-, G[G00008241, photo F neg. 23635]; iso-, F[F0071842F], possible iso-, K[K000542729]).

SIZE. — Up to 15 cm dbh (Acevedo-Rodríguez 2003).

[1589] *Talisia praealta* Sagot ex Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 345 (Radlkofer 1878). — *Cupania praealta* (Sagot ex Radlk.) Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 12: 199 (Sagot 1881).

VERNACULAR NAMES. — Pa: bukutru-ateu • Ka: tulisi, tulusili • Te: tuli • Wn: jala, täpuimë • Cr: bwa-flanbo, bwa-golët, payakousa • Br: pitomba-do-norte, pitombeira.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *P.A. Sagot 1047*, 1857 (holo-, B[not seen, photo F neg 5679]; lecto-, P[P04857336], designated by Acevedo-Rodríguez [2003: 76]; isolecto-, BM[BM001042336], BR[BR000005416639], F[F0071846F, V0071847F], GOET[GOET010870], K[K000035414, K000035415], M[M0224877], MPU[MPU010929], NY[00658344], P[P04857336], U[U0006555]).

INVENTORY DATA (FG). — 114 trees in 74 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 79.5$ cm.

[1590] *Talisia simaboides* K.U.Kramer

Acta Bot. Neerl. 21: 676 (Kramer 1972).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Nt: gaan tatu tiki, tatu udu • Cr: bwa-flanbo, payakousa.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3658*.

INVENTORY DATA (FG). — 132 trees in 93 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 70 \text{ cm}$.

[1591] *Talisia squarrosa* Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 346 (Radlkofe 1878).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *M.-F. Prévost 2170*.

INVENTORY DATA (FG). — 36 trees in 4 plots; $F_{\max} = 2.4\%$; $\text{dbh}_{\text{inv}} = 54.7 \text{ cm}$.

[1592] *Talisia sylvatica* (Aubl.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 341 (Radlkofe 1878). — *Racaria sylvatica* Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 24 [Jun.-Dec. 1775] (Aublet 1775).

Talisia carinata Radlk. f. *acutisepala* Radlk., *Fl. Bras. [Martius]* 13 (3): 549 [1 Apr. 1900] (Radlkofe 1900).

Talisia micrantha Radlk., *Repert. Spec. Nov. Regni Veg.* 9: 375 (Radlkofe 1911).

Talisia reticulata Radlk., *Repert. Spec. Nov. Regni Veg.* 9: 376 (Radlkofe 1911).

VERNACULAR NAMES. — Pa: tuu-aška • Te: tuli • Wp: tuli áta • Wn: tépu.

HERBARIUM DATA (FG). — 64 collections at CAY. Sel. exs.: *P. Acevedo-Rodríguez et al. 4911*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 25.6 \text{ cm}$.

Genus *Toulacia* Aubl.

[1593] *Toulacia elliptica* Radlk.

(Fig. 53D)

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 8: 371 (Radlkofe 1878).

VERNACULAR NAMES. — Pa: tuu-kamwi.

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 5076*.

INVENTORY DATA (FG). — 41 trees in 7 plots; $F_{\max} = 2.5\%$; $\text{dbh}_{\text{inv}} = 47.4 \text{ cm}$.

[1594] *Toulacia guianensis* Aubl.

Hist. Pl. Guiane 1: 359 [Jun.-Dec. 1775] (Aublet 1775).

Toulacia pulvinata Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 8: 371 (Radlkofe 1878).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana.

VERNACULAR NAMES. — Ka: tulisi, tulusili • Wp: tuli sī.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000629523](#)]).

SIZE. — Up to 30 m tall (Acevedo-Rodríguez 2012).

Genus *Vouarana* Aubl.

[1595] *Vouarana guianensis* Aubl.

Hist. Pl. Guiane 2 (Suppl.): 12 [Jun.-Dec. 1775] (Aublet 1775).

VERNACULAR NAMES. — Wp: tāpē lemiū'i • Br: maraximbé.

HERBARIUM DATA (FG). — 28 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000645706](#)]); D. Sabatier 1538 (epitype US[[00725189](#)] ; isoepitypes CAY[CAY015974], P[[P06671280](#)]).

INVENTORY DATA (FG). — 47 trees in 30 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 30.9 \text{ cm}$.

Family SAPOTACEAE Juss.

Genus *Chromolucuma* Ducke

[1596] *Chromolucuma congestifolia* (Pilz) Alves-Araújo & M.Alves

Brittonia 64 (1): 28 (Alves-Araújo & Alves 2012). — *Pouteria congestifolia* Pilz, *Ann. Missouri Bot. Gard.* 68 (1): 191 [17 Nov. 1981] (Pilz 1981).

HERBARIUM DATA (FG). — A single collection, *T.D. Pennington et al. 13840* (K).

SIZE. — Costa Rica. *T.D. Pennington 11539* (MO), 12 m × 30 cm.

Genus *Chrysophyllum* L.

[1597] *Chrysophyllum argenteum* Jacq.
subsp. *auratum* (Miq.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 545 [26 Apr. 1990] (Pennington 1990). — *Chrysophyllum auratum* Miq., *Fl. Bras. [Martius]* 7: 97 [15 Jan. 1863] (Miquel 1863). — *Cynodendron auratum* (Miq.) Baehni, *Boissiera* 11: 143 (Baehni 1965).

Chrysophyllum sericeum A.DC., *Prodri. [A. P. de Candolle]* 8: 158 [mid Mar. 1844] (Candolle 1844), *non* Salisb. (1796, nom. nud.)

Chrysophyllum richardii Klotsch ex Miq., *Fl. Bras. [Martius]* 7: 97 [15 Jan. 1863] (Miquel 1863), *nom. nud. pro syn.*

Chrysophyllum auratum var. *majus* Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863).

Chrysophyllum guyanense Klotzsch ex Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863), *nom. nud. pro syn.*

Chrysophyllum hostmannianum Klotzsch ex Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863), *nom. nud. pro syn.*

Chrysophyllum nitidissimum Klotzsch ex Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863), *nom. nud. pro syn.*

Chrysophyllum sericeum var. *acutifolium* Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863).

Chrysophyllum sericeum var. *obtusifolium* Miq., *Fl. Bras. [Martius]* 7: 98 [15 Jan. 1863] (Miquel 1863).

VERNACULAR NAMES. — Ka: kalu melei, kameli • Wp: kalamūlī sili, wai piyū, wai piyū sili, wai to wilā • Cr: zoliv • Br: ajará, guajará, uajara-branco.

HERBARIUM DATA (FG). — 60 collections at CAY. Sel. exs.: *A. Gabriel* s.n. (holotype of *Chrysophyllum sericeum*: G-DC[G00139532]; iso-, F[V0071948F], G[G00434796, G00434797]).

INVENTORY DATA (FG). — 28 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 61.4$ cm.

[1598] *Chrysophyllum argenteum*
subsp. *nitidum* (G.Mey.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 548 [26 Apr. 1990] (Pennington 1990). — *Chrysophyllum nitidum* G.Mey., *Prim. Fl. Esseq.*: 116 [Nov. 1818] (Meyer 1818). — *Cynodendron nitidum* (G.Mey.) Aubrév., *Adansonia*, n.s., 7 (4): 466 [Dec. 1967] (Aubréville 1967).

Chrysophyllum psilophyllum A.DC., *Prodr. [A. P. de Candolle]* 8: 160 [mid Mar. 1844] (Candolle 1844).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: kalu melei, kameli • Wp: wai • Wn: wapolimē • Nt: kwata bobi, nyanboka • Cr: zoliv • Br: ajará, guajará.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J. Martin* s.n. (holotype of *Chrysophyllum psilophyllum*: P[P00649300]; iso-, F[V0071943F], P[P00649301, P00649302, P00649303]; possible iso-, US[00323625]).

INVENTORY DATA (FG). — 250 trees in 108 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 43.9$ cm.

[1599] *Chrysophyllum cuneifolium* (Rudge) A.DC.

Prodr. [A. P. de Candolle] 8: 160 [mid Mar. 1844] (Candolle 1844). — *Bumelia cuneifolia* Rudge, *Pl. Guian. [Rudge]* 1 (4): 30 [Apr.-May 1806] (Rudge 1806). — *Prieurella cuneifolia* (Rudge) Pierre ex Aubrév., *Adansonia*, n.s., 1 (1): 19 [Nov. 1961] (Aubréville 1961), *nom. inval.* — *Ecclinusa cuneifolia* (Rudge) Aubrév., *Adansonia*, n.s., 1 (1): 20 [Nov. 1961] (Aubréville 1961), *nom. inval.*

NOTES. — The new combinations *Prieurella cuneifolia* and *Ecclinusa cuneifolia* were not validly published because the basionym was not explicitly cited (Turland et al. 2018: Art. 41.3). *Prieurella cuneifolia* (Rudge) Pierre ex Aubrév. appeared on p. 69 of Pierre's *Notes Botaniques: Sapotaceae*, but pp. 69-83 of that work were never published (they exist only as proofs at P).

VERNACULAR NAMES. — Pa: uu-kamwi-wahuyo, wakukwa-adudu • Ka: atakamala, kusili palatali • Te: wila pulu'a • Wp: wila pele • Nt: kwata bobi • Cr: tété-kwata • Br: abiurana, maçarandúbarana.

HERBARIUM DATA (FG). — 47 collections at CAY. Sel. exs.: *J. Martin* s.n. (original material BM[BM000952593]).

INVENTORY DATA (FG). — 51 trees in 36 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 22.1$ cm.

[1600] *Chrysophyllum durifructum* (W.A.Rodrigues)
T.D.Penn.

Fl. Neotrop. Monogr. 52: 604 [26 Apr. 1990] (Pennington 1990). — *Achrouteria durifructa* W.A.Rodrigues, *Acta Amazonica* 4 (3): 15 (Rodrigues 1974).

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori* et al. 24828.

INVENTORY DATA (FG). — 16 trees in 4 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 84$ cm.

[1601] *Chrysophyllum eximium* Ducke

Bull. Mus. Natl. Hist. Nat., sér. 2, 4 (6): 744 (Ducke 1932). — *Ecclinusa eximia* (Ducke) Cronquist, *Bull. Torrey Bot. Club* 73 (3): 310 [June 1946] (Cronquist 1946).

Chrysophyllum rufocupreum Ducke, *Arch. Inst. Biol. Veg.* 2 (1): 67 [Sep. 1935] (Ducke 1935). — *Ecclinusa rufocuprea* (Ducke) Cronquist, *Bull. Torrey Bot. Club* 73 (3): 311 [June 1946] (Cronquist 1946).

VERNACULAR NAMES. — Nt: busi apaa.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier* & M.-F. Prévost 2568.

INVENTORY DATA (FG). — 107 trees in 30 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 66.3$ cm.

[1602] *Chrysophyllum lucentifolium* Cronquist
subsp. *pachycarpum* Pires & T.D.Penn.

Fl. Neotrop. Monogr. 52: 606 [26 Apr. 1990] (Pires & Pennington 1990).

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *D. Sabatier* & M.-F. Prévost 2696.

INVENTORY DATA (FG). — 109 trees in 46 plots; $F_{\max} = 2.4\%$; $dbh_{\text{inv}} = 125$ cm.

[1603] *Chrysophyllum manaosense* (Aubrév.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 598 [26 Apr. 1990] (Pennington 1990). — *Prieurella manaosensis* Aubrév., *Adansonia*, n.s., 4 (3): 370 [Dec. 1964] (Aubréville 1964).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier* & J.-F. Molino 5134.

INVENTORY DATA (FG). — 6 trees in 6 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 34$ cm.

[1604] *Chrysophyllum pomiferum* (Eyma) T.D.Penn.

Fl. Neotrop. Monogr. 52: 602 [26 Apr. 1990] (Pennington 1990). — *Achrouteria pomifera* Eyma, *Recueil Trav. Bot. Néerl.* 33: 193 (Eyma 1936). — *Pouteria pomifera* (Eyma) Baehni, *Candollea* 9: 353 (Baehni 1942).

Planchonella guianensis P.Royen, *Blumea* 8: 395 (Royen 1957).

VERNACULAR NAMES. — Pa: aku-aiwut • Wn: tēpepuluimë • Nt: mongii soke • Cr: balata-ponm • Fr: jaune d'œuf à grandes feuilles • Br: abiurana.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Sabatier* 984.

INVENTORY DATA (FG). — 81 trees in 58 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 79.1$ cm.

[1605] *Chrysophyllum prieurii* A.DC.
(Fig. 54A)

Prodr. [A. P. de Candolle] 8: 161 [mid Mar. 1844] (Candolle 1844), “*Prieurei*”. — *Ecclinusa prieurii* (A.DC.) Aubrév., *Adansonia*, n.s., 1 (1): 20 [Nov. 1961] (Aubréville 1961). — *Prieurella prieurii* (A.DC.) Aubrév., *Adansonia*, n.s., 4 (3): 370 [Dec. 1964] (Aubréville 1964).

Chrysophyllum cyanoogenum Ducke, *Trop. Woods* 71: 18 (Ducke 1942). — *Ecclinusa cyanogena* (Ducke) Aubrév., *Adansonia*, n.s., 1 (1): 20 [Nov. 1961] (Aubréville 1961).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: panauna-priye, uu-kamwi • Wp: takama • Nt: pepee boyti • Cr: apisi-jonn, balata-jonndef • Fr: balata indien, balata jaune d'œuf • Br: abiurana-mocambo, abiurana-vermelha, maçarandúbarana.

HERBARIUM DATA (FG). — 44 collections at CAY. Sel. exs.: *E.M.R. Leprieur s.n.* (holo-, G-DC[G00139685]; iso-, GH[GH00075582], P[P00649386, P00649387]).

INVENTORY DATA (FG). — 350 trees in 138 plots; $F_{\max} = 1.9\%$; $dbh_{\text{inv}} = 71.9$ cm.

[1606] *Chrysophyllum sanguinolentum* (Pierre) Baehni

Boissiera 11: 74 (Baehni 1965). — *Ragala sanguinolenta* Pierre, *Notes Bot. Sapot.* 2: 60 (Pierre 1891). — *Ecclinusa sanguinolenta* Pierre, *Notes Bot. Sapot.* 2: 58 (Pierre 1891), nom. nud. pro syn. — *Ecclinusa sanguinolenta* (Pierre) Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachtr.* 1: 278 (Engler 1897).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Nt: switi amini • Cr: balata-senj-rouj • Fr: balata pommier • Br: balata-brava, ucuquirana.

HERBARIUM DATA (FG). — 42 collections at CAY. Sel. exs.: *E.M. Mélinon* 42, 1862 (holo-, P[P00649392]; iso-, BM[BM000952597], F[F0072216F, V0072217F], US[00067565, 00067566, 00067567]).

INVENTORY DATA (FG). — 399 trees in 123 plots; $F_{\max} = 5.3\%$; $dbh_{\text{inv}} = 76.9$ cm.

[1607] *Chrysophyllum sparsiflorum* Klotzsch ex Miq.

Fl. Bras. [Martius] 7: 90 [15 Jan. 1863] (Miquel 1863).

Chrysophyllum sparsiflorum var. *fagifolium* Miq., *Fl. Bras. [Martius]* 7: 90 [15 Jan. 1863] (Miquel 1863).

Chrysophyllum steyermarkii Monach., *Fieldiana, Bot.* 28 (3): 480 (Monachino 1953).

VERNACULAR NAMES. — Br: mangabarana.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *C. Feuillet* 3936.

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.4$ cm.

[1608] *Chrysophyllum venezuelanense* (Pierre) T.D.Penn.

Fl. Neotrop. Monogr. 52: 607 [26 Apr. 1990] (Pennington 1990). — *Cornuella venezuelanensis* Pierre, *Notes Bot. Sapot.* 2: 68 (Pierre 1891).

Chrysophyllum excelsum Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr.* 3: 55 (Huber 1900). — *Gambeya excelsa* (Huber) Aubrév., *Notul. Syst. (Paris)* 16: 248 (Aubréville 1961).

Lucuma lucentifolia Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 251 [24 Oct. 1929] (Standley 1929). — *Pouteria lucentifolia* (Standl.) Baehni, *Candollea* 9: 424 (Baehni 1942).

Lucuma pentasperma Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 251 [24 Oct. 1929] (Standley 1929). — *Pouteria pentasperma* (Standl.) Baehni, *Candollea* 9: 353 (Baehni 1942).

Pouteria petenensis Lundell, *Wrightia* 5 (1): 6 (Lundell 1972).

Pouteria dibrachiata Lundell, *Wrightia* 5 (4): 95 (Lundell 1975).

Pouteria mayana Lundell, *Wrightia* 5 (4): 100 (Lundell 1975). — *Achnas mayana* (Lundell) Lundell, *Wrightia* 5 (9): 350 (Lundell 1977).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Levis *et al.* 2017).

VERNACULAR NAMES. — Wp: wila kea.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Prevost* 4166.

INVENTORY DATA (FG). — 1 tree, $dbh = 12.6$ cm.

[1609] *Chrysophyllum* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier* & *J.-F. Molino* 5332.

INVENTORY DATA (FG). — 15 trees in 13 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 20.7$ cm.

[1610] *Chrysophyllum* sp. B

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier* 1464.

INVENTORY DATA (FG). — 24 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 92$ cm.



Fig. 53. — Sapindaceae: **A**, *Allophylus edulis* (A.St.-Hil., A.Juss. & Cambess.) Hieron. ex Niederl. (*D. Sabatier & M.-F. Prévost 1648*); **B**, *Cupania scrobiculata* Rich. (*J.-F. Molino & D. Sabatier 2484*); **C**, *Matayba opaca* Radlk. (*D. Sabatier 2102*); **D**, *Toulicia elliptica* Radlk. (*D. Sabatier & J.-F. Molino 5076*). A, C, D, © D. Sabatier/IRD; B, © J.-F. Molino/IRD.

[1611] *Chrysophyllum* sp. C

HERBARIUM DATA (FG). — A single collection, *D. Sabatier 3535*, dbh = 35 cm.

[1612] *Chrysophyllum* sp. D

HERBARIUM DATA (FG). — A single collection, *D. Sabatier et al. 5262*.

INVENTORY DATA (FG). — 24 trees in 5 plots; $F_{\max} = 1.9 \%$; $dbh_{\text{inv}} = 38.5 \text{ cm}$.

[1613] *Chrysophyllum* sp. E

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5651*.

INVENTORY DATA (FG). — 5 trees in 5 plots; $F_{\max} < 1 \%$; $dbh_{\text{inv}} = 79.3 \text{ cm}$.

[1614] *Chrysophyllum* sp. F

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2815*.

INVENTORY DATA (FG). — 1 tree, dbh = 11 cm.

Genus *Diploön* Cronquist

[1615] *Diploön cuspidatum* (Hoehne) Cronquist

Bull. Torrey Bot. Club 73 (5): 466 [Oct. 1946] (Cronquist 1946). — *Chrysophyllum cuspidatum* Hoehne, *Ostenia*: 302 (Hoehne 1933).

Diploön venezuelanum Aubrév., *Mem. New York Bot. Gard.* 23: 225 (Aubréville 1972), “venezuelana”.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *J.-F. Molino et al. 2076*.

INVENTORY DATA (FG). — 68 trees in 30 plots; $F_{\max} = 1.4 \%$; $dbh_{\text{inv}} = 57.2 \text{ cm}$.

Genus *Ecclinusa* Mart.

[1616] *Ecclinusa guianensis* Eyma

Recueil Trav. Bot. Néerl. 33: 203 (Eyma 1936). — *Chrysophyllum guianense* (Eyma) Baehni, *Boissiera* 11: 74 (Baehni 1965), “guyanense”.

Ecclinusa bacuri Aubrév. & Pellegr., *Adansonnia*, n.s., 1 (1): 21 [Nov. 1961] (Aubréville & Pellegrin 1961).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: balatabali, kulumoto, matamata wewe, palatabali • Wp: kalamüli, kalamüli sili, tamanuwa • Nt: busi apaa, malobi • Cr: balata-blanc, bataballi • Fr: balata bali • Br: abiurana-abiu, abiurana-bacuri, abiurana-caju, caramuri.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-762*.

INVENTORY DATA (FG). — 329 trees in 87 plots; $F_{\max} = 24.1\%$; $\text{dbh}_{\text{inv}} = 71\text{ cm}$.

[1617] *Ecclinusa lanceolata* (Mart. & Eichler) Pierre

Notes Bot. Sapot. 2: 57 (Pierre 1891). — *Passaveria lanceolata* Mart. & Eichler, *Fl. Bras. [Martius]* 7: 86 [15 Jan. 1863] (Martius & Eichler 1863).

VERNACULAR NAMES. — Nt: malobi.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier 3519*.

INVENTORY DATA (FG). — 8 trees in 5 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22\text{ cm}$.

[1618] *Ecclinusa ramiflora* Mart.

Flora 22 (1, Beibl.): 2 (Martius 1839). — *Chrysophyllum rami-florum* (Mart.) A.DC., *Prodr. [A. P. de Candolle]* 8: 158 [mid Mar. 1844] (Candolle 1844). — *Passaveria obovata* Mart. & Eichler, *Fl. Bras. [Martius]* 7: 87 [15 Jan. 1863] (Martius & Eichler 1863), nom. illeg. superfl. (based on *Ecclinusa ramiflora*).

Passaveria obovata var. *tomentosa* Miq., *Fl. Bras. [Martius]* 7: 87 [15 Jan. 1863] (Miquel 1863).

Lucuma tomentosa Poepp. ex Miq., *Fl. Bras. [Martius]* 7: 87 [15 Jan. 1863] (Miquel 1863), nom. nud. pro syn.

Ecclinusa costata Pierre, *Notes Bot. Sapot.* 2: 56 (Pierre 1891).

Ecclinusa abbreviata Ducke, *Bull. Mus. Natl. Hist. Nat.*, sér. 2, 4 (6): 743 (Ducke 1932). — *Chrysophyllum abbreviatum* (Ducke) Baehni, *Boissiera* 11: 75 (Baehni 1965).

VERNACULAR NAMES. — Wp: miku lapi'a u • Nt: malobi • Br: abi-fofo, coquirana-brava.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier 1027*.

INVENTORY DATA (FG). — 74 trees in 54 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 31.8\text{ cm}$.

Genus *Elaeolumma* Baill.

[1619] *Elaeolumma nuda* (Baehni) Aubrév.

Mem. New York Bot. Gard. 23: 224 (Aubréville 1972). — *Pouteria nuda* Baehni, *Candollea* 14: 72 (Baehni 1952).

Pouteria auyantepuiensis Steyermark., *Acta Bot. Venez.* 2 (5-8): 278 (Steyermark 1967).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2889*.

INVENTORY DATA (FG). — 34 trees in 23 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 53.5\text{ cm}$.

[1620] *Elaeolumma* sp. A

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 4647*.

INVENTORY DATA (FG). — 9 trees in 6 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 29.6\text{ cm}$.

Genus *Manilkara* Adans.

[1621] *Manilkara bidentata* (A.DC.) A.Chev.
subsp. *bidentata*
(Fig. 54B)

Rev. Bot. Appl. Agric. Trop. 12: 270 (Chevalier 1932). — *Mimusops bidentata* A.DC., *Prodr. [A. P. de Candolle]* 8: 204 [mid Mar. 1844] (Candolle 1844). — *Manilkara balata* Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 19 (Dubard 1915), nom. illeg. superfl. (*Mimusops bidentata* in synonymy).

Mimusops sieberi A.DC., *Prodr. [A. P. de Candolle]* 8: 204 [mid Mar. 1844] (Candolle 1844). — *Mimusops balata* (Aubl.) C.F.Gaertn. var. *sieberi* (A.DC.) Pierre, *Bull. Mens. Soc. Linn. Paris* 1 (64): 508 (Pierre 1885). — *Manilkara balata* var. *sieberi* (A.DC.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 20 (Dubard 1915).

Sapota mulleri Blume, *Ann. Sci. Nat., Bot.* sér. 4, 7: 225 (Blume 1857).

Mimusops balata var. *gutta* Pierre, *Bull. Mens. Soc. Linn. Paris* 1 (64): 507 (Pierre 1885). — *Manilkara balata* var. *gutta* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 21 (Dubard 1915).

Mimusops balata var. *melinonii* Pierre, *Bull. Mens. Soc. Linn. Paris* 1 (64): 507 (Pierre 1885), “*Melinonis*”. — *Manilkara balata* var. *melinonii* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 21 (Dubard 1915), “*Melinonis*”.

Mimusops balata var. *schomburgkii* Pierre, *Bull. Mens. Soc. Linn. Paris* 1 (64): 507 (Pierre 1885). — *Manilkara balata* var. *schomburgkii* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 20 (Dubard 1915).

Mimusops balata var. *cruegeri* Pierre, *Symb. Antill. [Urban]* 5 (1): 165 [20 May 1904] (Pierre 1904). — *Manilkara balata* var. *cruegeri* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 20 (Dubard 1915). — *Manilkara bidentata* var. *cruegeri* (Pierre) A.Chev., *Rev. Bot. Appl. Agric. Trop.* 12: 275 (Chevalier 1932).

Mimusops balata var. *hartii* Pierre, *Symb. Antill. [Urban]* 5 (1): 166 [20 May 1904] (Pierre 1904). — *Manilkara balata* var. *hartii* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 21 (Dubard 1915).

Mimusops darienensis Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 249 [22 Sep. 1917] (Pittier 1917). — *Manilkara darienensis* (Pittier) Standl., *Trop. Woods* 31: 45 (Standley 1932).

Manilkara williamsii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 22 (3): 165 (Standley 1940).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *Mimusops balata* sensu Pierre, non (Aubl.) C.F.Gaertn., is a misinterpretation of *Achras balata* Aubl. The latter was described from a tree imported from the Mauritius Island (“Île de France”) and cultivated at Cayenne (Aublet 1775). The type of *A. balata* is in P-JU, and was obtained from a plant growing in the Mascarene Islands (Chevalier 1932, Pennington 1990). Although Aublet in his *Histoire des plantes* did not mention the Guianan “Balata”, he

collected specimens of this species, which are in P and P-JU (Chevalier 1932). Pierre (1885) correctly stated that these 2 specimens were conspecific with the type of *Mimusops bidentata* A.DC., but wrongly concluded that they were type material of *A. balata*. Although aware of the previous transfer of *A. balata* to *Mimusops* L. by Gaertner (1807: 133), Pierre repropose the same combination, as “*Mimusops balata* non Gaertner”. Notwithstanding, all varieties created by Pierre (1885) under *Mimusops balata* are legitimate under *Mimusops balata* (Aubl.) C.F.Gaertn. *Manilkara balata* is similarly confused: it is based on “*Mimusops balata* Pierre” and *Mimusops bidentata* is a synonym, which accords with the Guianan plant, even though *A. balata* was also cited in synonymy. When the latter is excluded from synonymy, then *Manilkara balata* (Aubl.) Dubard is superfluous (because *Mimusops bidentata* is in synonymy) and is therefore illegitimate. Bleekrode (1857) ascribed both the name and description of *Sapota mulleri* to Blume, hence authorship is “Blume”, not “Blume ex Bleekrode”; by the way, although the author of the “*Notice sur la gutta-percha de Surinam*”, as it appears on the first page, is “M. le Professeur Bleekrod de l’Académie de Delft”, he is actually Salomon Abraham Bleekrode (not “Bleekrod”, see https://nl.wikipedia.org/wiki/Salomon_Abraham_Bleekrode). The epithet “*melinonis*”, which honours the French botanist E. Mélinon, is to be automatically corrected to “*melinonii*” (Turland et al. 2018: Art. 60.8); hence *Manilkara balata* var. *melinonii*. and *Mimusops balata* var. *melinonii*.

VERNACULAR NAMES. — Pa: balata • Ka: palata, polata • Te: balata'i • Wp: iwií, iwilá, masaláni • Wn: palakta, palaktaimé • Nt: boyti • Cr: balata-fran, balata-gonm • Fr: balata franc • Br: balateira, maçarandúba, maparajúba.

HERBARIUM DATA (FG). — 53 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (holo-, P[P00645496]; iso-, P[P00645494, P00645495]).

INVENTORY DATA (FG). — 649 trees in 139 plots; $F_{\max} = 7.4\%$; $dbh_{inv} = 111.4$ cm.

[1622] *Manilkara bidentata*
subsp. *surinamensis* (Miq.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 61 [26 Apr. 1990] (Pennington 1990). — *Mimusops surinamensis* Miq., *Fl. Bras. [Martius]* 7: 43 [15 Jan. 1863] (Miquel 1863). — *Kaukenia surinamensis* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 406 [5 Nov. 1891] (Kuntze 1891). — *Manilkara surinamensis* (Miq.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 22 (Dubard 1915).

Mimusops balata var. *domingensis* Pierre, *Bull. Mens. Soc. Linn. Paris* 1 (64): 508 (Pierre 1885). — *Manilkara balata* var. *domingensis* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 21 (Dubard 1915). — *Mimusops domingensis* (Pierre) Moscoso, Cat. Fl. Domingensis 1: 484 (Moscoso 1943).

Mimusops riedeliana Pierre ex Baill., *Bull. Mens. Soc. Linn. Paris* 2 (116): 922 (Baillon 1891).

Acras nitida Sessé & Moc., *Fl. Mexic.*, ed. 2: 85 (Sessé & Mociño 1894). — *Mimusops nitida* (Sessé & Moc.) Urb., *Symb. Antill. [Urban]* 5 (1): 167 [20 May 1904] (Urban 1904). — *Manilkara nitida* (Sessé & Moc.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 18 (Dubard 1915).

Mimusops riedleiana Pierre ex Duss, *Ann. Inst. Colon. Marseille* 3: 387 [“1896” publ. 1897] (Duss 1897). — *Manilkara riedleiana* (Pierre ex Duss) Dubard, *Ann. Mus. Colon. Marseille*, sér. 3, 3: 17 (Dubard 1915).

Mimusops amazonica Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 4: 433 (Huber 1906). — *Manilkara amazonica* (Huber) Standl., *Trop. Woods* 34: 41 (Standley 1933).

Mimusops maparajuba Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 4: 434 (Huber 1906).

Manilkara longiciliata Ducke, *Trop. Woods* 71: 22 (Ducke 1942).

Mimusops longiciliata Ducke, *Trop. Woods* 71: 22 (Ducke 1942).

Manilkara siqueiraei Ducke, *Trop. Woods* 71: 24 (Ducke 1942).

Mimusops siqueiraei Ducke, *Trop. Woods* 71: 24 (Ducke 1942).

NOTES. — Ducke (1942: 22) simultaneously published *Manilkara longiciliata* and *Mimusops longiciliata* as alternative names, but both are nevertheless valid, because they were published before 1953 (Turland et al. 2018: Art. 36.3). The same applies to *Manilkara siqueiraei* and *Mimusops siqueiraei* (Ducke 1942: 24).

VERNACULAR NAMES. — Wp: iwilá, iwilá pilá.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *P. Grenand* 1488.

SIZE. — Up to 10 cm dbh (Pennington 1990).

[1623] *Manilkara huberi* (Ducke) A.Chev.

Rev. Bot. Appl. Agric. Trop. 12: 276 (Chevalier 1932). — *Mimusops huberi* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 2: 14 (Ducke 1917).

NOTE. — Incipiently domesticated by pre-Columbian Amerindians (Clement 1999; Levis et al. 2017).

VERNACULAR NAMES. — Pa: balata-duwé • Nt: boyti, kayman buba boyti • Cr: balata • Fr: balata franc • Br: maçarandúba-da-terra-firme, maparajúba.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *B. Riéra* & *D. Sabatier* 1904.

INVENTORY DATA (FG). — 76 trees in 34 plots; $F_{\max} = 1.9\%$; $dbh_{inv} = 123.3$ cm.

[1624] *Manilkara paraensis* (Huber) Standl.

Trop. Woods 34: 41 (Standley 1933). — *Mimusops paraensis* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 4: 435 (Huber 1906).

Mimusops paraensis var. *densiflora* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 4: 435 (Huber 1906).

Mimusops paraensis var. *discolor* Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 4: 435 (Huber 1906).

VERNACULAR NAMES. — Pa: balata-puvemna • Wp: masaláni • Br: maçarandúba-de-folha miuda, maçarandúba-jacaré.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 3422.

INVENTORY DATA (FG). — 1 tree, dbh = 48.7 cm.

Genus *Micropholis* (Griseb.) Pierre

[1625] *Micropholis acutangula* (Ducke) Eyma
(Fig. 54C)

Recueil Trav. Bot. Néerl. 33: 198 (Eyma 1936). — *Sideroxylon acutangulum* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 159 (Ducke 1925). — *Pouteria acutangula* (Ducke) Baehni, *Candollea* 9: 246 (Baehni 1942). — *Paramicropholis acutangula* (Ducke) Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 171 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962). — *Paralabatia acutangula* (Ducke) Baehni, *Boissiera* 11: 141 (Baehni 1965).

Micropholis paraensis Eyma, *Recueil Trav. Bot. Néerl.* 33: 198 (Eyma 1936), nom. nud. — *Sideroxylon paraense* Huber ex Eyma, *Recueil Trav. Bot. Néerl.* 33: 198 (Eyma 1936), nom. nud. pro syn.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & S. Gonzalez* 5371.

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 46.7$ cm.

[1626] *Micropholis cayennensis* T.D.Penn.

Fl. Neotrop. Monogr. 52: 220 [26 Apr. 1990] (Pennington 1990).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: uu-kamwi, uu-kamwi-purubumna, yauknabwi-wahuye • Nt: bakuman, nyanboka • Cr: zoliv.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *B. Maguire et al.* 47074 (holo-, NY[01199690]; iso-, G[G00439289], GH[GH00589970], K[K000641500], R[R000112723], S[S05-10441], SP[SP001602], US[US00323728]).

INVENTORY DATA (FG). — 100 trees in 36 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 89.1$ cm.

[1627] *Micropholis egensis* (A.DC.) Pierre

Symb. Antill. [Urban] 5 (1): 127 [20 May 1904] (Pierre 1904). — *Sideroxylon egense* A.DC., *Prodr. [A. P. de Candolle]* 8: 182 [mid Mar. 1844] (Candolle 1844). — *Bumelia egensis* Poepp. ex A.DC., *Prodr. [A. P. de Candolle]* 8: 182 [mid Mar. 1844] (Candolle 1844), nom. nud. pro syn. — *Sideroxylon rugosum* (Sw.) Roem. & Schult. var. *egense* (A.DC.) Miq., *Fl. Bras. [Martius]* 7: 51 [15 Jan. 1863] (Miquel 1863), "Sideroxylum". — *Pouteria egensis* (A.DC.) Baehni, *Candollea* 9: 207 (Baehni 1942).

Micropholis martiana Pierre, *Symb. Antill. [Urban]* 5 (1): 126 [20 May 1904] (Pierre 1904).

Sideroxylon ulei K.Krause, *Verh. Bot. Vereins Prov. Brandenburg* 50: 95 [10 June 1908] (Krause 1908). — *Micropholis ulei* (K.Krause) Eyma, *Recueil Trav. Bot. Néerl.* 33: 198 (Eyma 1936). — *Pouteria ulei* (K.Krause) Baehni, *Candollea* 9: 217 (Baehni 1942).

Sideroxylon quinilla Standl. ex Ll.Williams, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 414 (Williams 1936), nom. inval. (anglise).

Pouteria setosa Baehni, *Candollea* 18: 168 (Baehni 1962).

Micropholis wurdackii Aubrév., *Mem. New York Bot. Gard.* 23: 212 (Aubréville 1972).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: paivan • Wp: waa • Wn: tamaniwa • Nt: mamanten • Cr: zoliv • Br: mangabarana.

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *D. Sabatier* 1267.

INVENTORY DATA (FG). — 162 trees in 84 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 85$ cm.

[1628] *Micropholis guyanensis* (A.DC.) Pierre
subsp. *guyanensis*

Notes Bot. Sapot. 2: 40 (Pierre 1891), "Guyanensis".

Sideroxylon guyanense A.DC., *Prodr. [A. P. de Candolle]* 8: 182 [mid Mar. 1844] (Candolle 1844), "Guyanense".

Sideroxylon rufum Mart. & Eichler, *Fl. Bras. [Martius]* 7: 52 [15 Jan. 1863] (Martius & Eichler 1863). — *Micropholis rufa* (Mart. & Eichler) Pierre, *Symb. Antill. [Urban]* 5 (1): 130 [20 May 1904] (Pierre 1904). — *Pouteria rufa* (Mart. & Eichler) Baehni, *Candollea* 9: 199 (Baehni 1942).

Sideroxylon cyrtobotryum Mart. ex Miq., *Fl. Bras. [Martius]* 7: 57 [15 Jan. 1863] (Miquel 1863). — *Sprucella cyrtobotrya* (Mart. ex Miq.) Pierre, *Notes Bot. Sapot.* 1: 27 (Pierre 1890). — *Micropholis cyrtobotrya* (Mart. ex Miq.) Baill., *Hist. Pl. [Baillon]* 11: 282 [Sep.-Oct. 1891] (Baillon 1891), nom. inval. (genus name and epithet not associated). — *Pouteria cyrtobotrya* (Mart. ex Miq.) Baehni, *Candollea* 9: 199 (Baehni 1942).

Chrysophyllum melinonii Engl., *Bot. Jahrb. Syst.* 12: 521 [23 Dec. 1890] (Engler 1890), "Melinoni". — *Pouteria melinonii* (Engl.) Baehni, *Candollea* 9: 200 (Baehni 1942). — *Planchonella melinonii* (Engl.) Baehni, *Boissiera* 11: 67 (Baehni 1965).

Micropholis chrysophylloides Pierre, *Notes Bot. Sapot.* 2: 38 (Pierre 1891). — *Pouteria chrysophylloides* (Pierre) Stehlé, *Caribbean Forest.* 4 (3): 118 (Stehlé 1943).

Micropholis cruegeriana Pierre, *Notes Bot. Sapot.* 2: 38 (Pierre 1891). — *Pouteria cruegeriana* (Pierre) Baehni, *Candollea* 9: 206 (Baehni 1942).

Micropholis portoricensis Pierre, *Notes Bot. Sapot.* 2: 38 (Pierre 1891).

Micropholis imrayana Pierre, *Notes Bot. Sapot.* 2: 39 (Pierre 1891).

Micropholis portoricensis var. *curvata* Pierre, *Notes Bot. Sapot.* 2: 39 (Pierre 1891). — *Micropholis curvata* (Pierre) Urb., *Symb. Antill. [Urban]* 5 (1): 124 [20 May 1904] (Urban 1904), in obs.

Micropholis portoricensis var. *mesuifolia* Pierre, *Notes Bot. Sapot.* 2: 39 (Pierre 1891).

Micropholis achradiformis Pierre, *Symb. Antill. [Urban]* 5 (1): 118 [20 May 1904] (Pierre 1904). — *Pouteria achradiformis* (Pierre) Baehni, *Candollea* 9: 207 (Baehni 1942), "achradiformis".

Micropholis eggarsiana Pierre, *Symb. Antill. [Urban]* 5 (1): 118 [20 May 1904] (Pierre 1904).

Micropholis truncata Pierre, *Symb. Antill. [Urban]* 5 (1): 119 [20 May 1904] (Pierre 1904). — *Pouteria truncata* (Pierre) Baehni, *Candollea* 9: 216 (Baehni 1942).

Micropholis chrysophylloides var. *truncata* Pierre, *Symb. Antill. [Urban]* 5 (1): 119 [20 May 1904] (Pierre 1904), nom. nud. pro syn.

Micropholis balata Pierre, *Symb. Antill. [Urban]* 5 (1): 120 [20 May 1904] (Pierre 1904). — *Pouteria balata* (Pierre) Baehni, *Candollea* 9: 351 (Baehni 1942).

Micropholis discolor Pierre, *Symb. Antill. [Urban]* 5 (1): 121 [20 May 1904] (Pierre 1904). — *Chrysophyllum discolor* Walp. & Duchass. ex Pierre, *Symb. Antill. [Urban]* 5 (1): 121 [20 May 1904] (Pierre 1904), nom. nud. pro syn. — *Pouteria discolor* (Pierre) Stehlé, *Caribbean Forest*, 4 (3): 118 (Stehlé 1943), nom. illeg. hom., non (Baill.) Baehni (1942).

Micropholis dominicensis Pierre, *Symb. Antill. [Urban]* 5 (1): 122 [20 May 1904] (Pierre 1904). — *Pouteria dominicensis* (Pierre) Stehlé, *Fl. Guadeloupe [Stehlé]* 4: 121 (Stehlé 1943).

Micropholis grandifolia Pierre ex Glaz., *Bull. Soc. Bot. France* 57 (Mém. 3e): 441 (Glaziou 1910), nom. nud.

Sideroxylon williamsii Baehni, *Candollea* 7: 135 (Baehni 1936). — *Mastichodendron williamsii* (Baehni) Baehni ex Bernardi, *Candollea* 22 (2): 231 (Bernardi 1967).

Pouteria aeranea Baehni, *Candollea* 9: 205 (Baehni 1942).

Pouteria silvicola Baehni, *Candollea* 18: 161 (Baehni 1962).

Pouteria bopiensis Baehni, *Candollea* 18: 164 (Baehni 1962).

Pouteria piresii Baehni, *Candollea* 18: 168 (Baehni 1962).

Micropholis cowanii Aubrév., *Mem. New York Bot. Gard.* 23: 211 (Aubréville 1972).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *Micropholis curvata* is a valid recombination, by Urban alone (not “Pierre & Urban” as in Pennington, 1990), of *M. portoricensis* var. *curvata* Pierre.

VERNACULAR NAMES. — Pa: aku-aiwut, uu-kamwi, waaduk-seine, yauknabui • Ka: asepoko, asepuku, kusili palatali, wesepuku, woko wokulu • Wp: wamisalu, watuliya • Wn: kanawa popalin, palakta jalan • Nt: bakuuman, gaan wiwii bakuuman, tapooka mapa • Cr: balata-blan, balata-endjen • Fr: balata blanc à grandes feuilles • Br: balata-rosada, chicle-bravo, maparajúba.

HERBARIUM DATA (FG). — 69 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, P[P00649234]).

INVENTORY DATA (FG). — 382 trees in 129 plots; $F_{\max} = 2.3\%$; $dbh_{\text{inv}} = 96$ cm.

[1629] *Micropholis guyanensis* subsp. *duckeana* (Baehni)
T.D.Penn.

Fl. Neotrop. Monogr. 52: 180 [26 Apr. 1990] (Pennington 1990). — *Pouteria duckeana* Baehni, *Candollea* 18: 161 (Baehni 1962).

Micropholis rosadinho-brava Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 178 [“Dec. 1961” publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

VERNACULAR NAMES. — Pa: yauknabui • Br: abiu-balatarana, balata-rosadinho.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2973*.

INVENTORY DATA (FG). — 73 trees in 29 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 53$ cm.

[1630] *Micropholis longipedicellata* Aubrév.

Mem. New York Bot. Gard. 23: 209 (Aubréville 1972).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: impitit-ahavukune, mpitit-ahavukune.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *D. Sabatier 2241*.

INVENTORY DATA (FG). — 55 trees in 26 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 37.6$ cm.

[1631] *Micropholis melinoniana* Pierre

Notes Bot. Sapot. 2: 40 (Pierre 1891). — *Pouteria melinoniana* (Pierre) Baehni, *Candollea* 9: 212 (Baehni 1942).

Sideroxylon rugosum var. *parviflorum* Miq., *Fl. Bras. [Martius]* 7: 51 [15 Jan. 1863] (Miquel 1863).

Stephanoluma rugosa Baill., *Hist. Pl. [Baillon]* 11: 283 [Sep.-Oct. 1891] (Baillon 1891).

Sideroxylon calophylloides Lundell, *Contr. Univ. Michigan Herb.* 6: 56 (Lundell 1941), nom. illeg. hom., non (Pierre) Engl. (Engler 1897).

Micropholis mexicana Gilly ex Cronquist, *Lloydia* 9 (4): 257 [“1945” publ. 20 Dec. 1946] (Cronquist 1946).

Micropholis guatemalensis Lundell, *Wrightia* 5 (4): 92 (Lundell 1975).

NOTE. — Baillon based his *Stephanoluma rugosa* on “*Sideroxylon rugosum* A.DC. [...] (non Roem. et Sch.)”. Actually, “*Sideroxylon rugosum* A.DC.” is not a name; it is a misinterpretation of *Sideroxylon rugosum* (Sw.) Roem. & Schult. by Candolle. Therefore, *Stephanoluma rugosa* is a new name, and its authorship is Baill., not “(A.DC.) Baill.”

VERNACULAR NAMES. — Pa: muhut-wašiunó • Wp: waa • Wn: kanawa popalin, wopoli • Nt: mamanten • Cr: balata-blan, zoliv-maypouri • Br: bacuri-de-anta, balata-rosada, caramuri-da-várzea, corrupixá.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *E.M. Mélion s.n.* (holo-, P[P00649246]).

INVENTORY DATA (FG). — 142 trees in 59 plots; $F_{\max} = 2.9\%$; $dbh_{\text{inv}} = 155$ cm.

[1632] *Micropholis mensalis* (Baehni) Aubrév.

Adansonia, n.s., 3 (1): 21 [Jan.-Mar. 1963] (Aubréville 1963). — *Pouteria mensalis* Baehni, *Candollea* 14: 64 (Baehni 1952).

VERNACULAR NAMES. — Br: abiurana-goiabinha, abiurana-roxa.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier et al. 6122*.

INVENTORY DATA (FG). — 24 trees in 11 plots; $F_{\max} = 1.3\%$; $dbh_{\text{inv}} = 16.9$ cm.

[1633] *Micropholis obscura* T.D.Penn.
(Fig. 54D)

Fl. Neotrop. Monogr. 52: 227 [26 Apr. 1990] (Pennington 1990).

VERNACULAR NAMES. — Nt: akwansiba • Cr: balata-blan, balata-endjen, zoliv • Fr: balata blanc • Br: balata-rosada.

HERBARIUM DATA (FG). — 37 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom* 14784 (holo-, K, not seen; iso-, MA[MA400083, MA400083-2], NY[00901963], U[U0006611]).

INVENTORY DATA (FG). — 152 trees in 62 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 95\text{ cm}$.

[1634] *Micropholis porphyrocarpa* (Baehni) Monach.

Brittonia 7: 409 [Oct. 1952] (Monachino 1952). — *Pouteria porphyrocarpa* Baehni, *Candollea* 9: 214 (Baehni 1942).

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 1853.

INVENTORY DATA (FG). — 24 trees in 13 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 63.8\text{ cm}$.

[1635] *Micropholis sanctae-rosae* (Baehni) T.D.Penn.

Fl. Neotrop. Monogr. 52: 215 [26 Apr. 1990] (Pennington 1990). — *Pouteria sanctae-rosae* Baehni, *Candollea* 9: 211 (Baehni 1942).

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier et al.* 4794.

INVENTORY DATA (FG). — 30 trees in 12 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 21.6\text{ cm}$.

[1636] *Micropholis trunciflora* Ducke

Bol. Técn. Inst. Agron. N. 19: 19 (Ducke 1950).

Pouteria klugii Baehni, *Candollea* 14: 76 (Baehni 1952).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3298.

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 44.2\text{ cm}$.

[1637] *Micropholis venulosa* (Mart. & Eichler) Pierre

Notes Bot. Sapot. 2: 40 (Pierre 1891). — *Sideroxylon venulosum* Mart. & Eichler, *Fl. Bras. [Martius]* 7: 52 [15 Jan. 1863] (Martius & Eichler 1863). — *Lucuma venulosa* Spruce ex Mart. & Eichler, *Fl. Bras. [Martius]* 7: 52 [15 Jan. 1863] (Martius & Eichler 1863), nom. nud. pro syn. — *Pouteria venulosa* (Mart. & Eichler) Baehni, *Candollea* 9: 195 (Baehni 1942). — *Xantolis venulosa* (Mart. & Eichler) Baehni, *Boissiera* 11: 24 (Baehni 1965).

Micropholis calophylloides Pierre, *Notes Bot. Sapot.* 2: 40 (Pierre 1891). — *Platyluma calophylloides* (Pierre) Baill., *Hist. Pl. [Baillon]* 11: 282 [Sep.-Oct. 1891] (Baillon 1891). — *Sideroxylon calophylloides* (Pierre) Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachr.* 1: 276 (Engler 1897).

Meioloma guianensis Baill., *Hist. Pl. [Baillon]* 11: 282 [Sep.-Oct. 1891] (Baillon 1891). — *Sideroxylon guianense* (Baill.) Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachr.* 1: 276 (Engler 1897), nom. illeg. hom., non A.DC. (Candolle 1844).

Micropholis mucronata Pierre, *Symb. Antill. [Urban]* 5 (1): 112 [20 May 1904] (Pierre 1904).

Micropholis rigida Pierre, *Symb. Antill. [Urban]* 5 (1): 125 [20 May 1904] (Pierre 1904).

Pouteria polyneura Baehni, *Candollea* 7: 133 (Baehni 1936).

Pouteria rivularis Baehni, *Candollea* 9: 208 (Baehni 1942).

Pouteria flava Baehni, *Candollea* 18: 164 (Baehni 1962).

Pouteria cataractae Baehni, *Candollea* 18: 168 (Baehni 1962).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: inam-etni-seinō, muhut-wašiunō • Ka: tabuku, woko popi • Te: kutsili palatahan • Wp: wai tawa • Nt: bakuuman • Cr: zoliv • Br: abiurana-branca, mulungu-rosa-verde.

HERBARIUM DATA (FG). — 63 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (original material of *Meioloma guianensis*: P[P00649260]).

INVENTORY DATA (FG). — 127 trees in 74 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 61.6\text{ cm}$.

[1638] *Micropholis* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier* 2339.

INVENTORY DATA (FG). — 7 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 26.8\text{ cm}$.

Genus *Pouteria* Aubl.

[1639] *Pouteria ambelaniifolia* (Sandwith) T.D.Penn.

Fl. Neotrop. Monogr. 52: 362 [26 Apr. 1990] (Pennington 1990). — *Chrysophyllum ambelaniifolium* Sandwith, *Bull. Misc. Inform. Kew* 1931 (10): 476 [31 Dec. 1931] (Sandwith 1931). — *Oxythecia ambelaniifolia* (Sandwith) Cronquist, *Bull. Torrey Bot. Club* 73 (3): 310 [June 1946] (Cronquist 1946). — *Pseudoxylethea ambelaniifolia* (Sandwith) Aubrév., *Mem. New York Bot. Gard.* 23: 219 (Aubréville 1972).

VERNACULAR NAMES. — Pa: waaduk-seine.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2973.

INVENTORY DATA (FG). — 138 trees in 73 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 57.9\text{ cm}$.

[1640] *Pouteria anomala* (Pires) T.D.Penn.

Fl. Neotrop. Monogr. 52: 344 [26 Apr. 1990] (Pennington 1990). — *Chrysophyllum anomalum* Pires, *Bol. Técn. Inst. Agron. N.* 38: 34 (Pires 1960).

VERNACULAR NAMES. — Br: abiurana-roxa.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Loubry 621*.

SIZE. — Up to 70 cm dbh (Pennington 1990).

[1641] *Pouteria aubrevillei* Bernardi

Candollea 22 (2): 231 (Bernardi 1967).

Eremoluma krukoffii Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 169 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962), "Krukoffii".

Eremoluma wurdackii Aubrév., *Adansonia*, n.s., 5 (2): 197 [July 1965] (Aubréville 1965).

VERNACULAR NAMES. — Pa: uu-kamwi • Wp: ei li • Cr: bwa-kay • Br: abiurana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5130*.

INVENTORY DATA (FG). — 30 trees in 17 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 36.6$ cm.

[1642] *Pouteria bangii* (Rusby) T.D.Penn.

Fl. Neotrop. Monogr. 52: 288 [26 Apr. 1990] (Pennington 1990). — *Sideroxylon bangii* Rusby, *Bull. New York Bot. Gard.* 4 (14): 407 [7 Dec. 1907] (Rusby 1907).

Pseudocladia melinonii Baill., *Hist. Pl. [Baillon]* 11: 291 [Sep.-Oct. 1891] (Baillon 1891), "Melinoni", nom. nud. — *Lucuma melinonii* Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachtr.* 1: 275 (Engler 1897), "Melinoni", nom. illeg., based on the illegitimate *Pseudocladia melinonii*.

Podoluma glaziovii Baill. ex Glaz., *Bull. Soc. Bot. France* 57 (Mém. 3e): 442 (Glaziou 1910).

Pouteria scytalophora Eyma, *Recueil Trav. Bot. Néerl.* 33: 181 (Eyma 1936). — *Pseudocladia scytalophora* (Eyma) Aubrév., *Adansonia*, n.s., 1 (2): 165 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

Pouteria micans Baehni, *Candollea* 9: 236 (Baehni 1942).

Pseudocladia colombiana Aubrév., *Adansonia*, n.s., 7 (2): 148 [Jul.-Sep. 1967] (Aubréville 1967).

VERNACULAR NAMES. — Ka: lemu epi • Wp: wila tawa • Cr: balata-pwé • Br: abiu-rosadinha, maparajúba-branca.

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier 4462*.

INVENTORY DATA (FG). — 86 trees in 49 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 54.3$ cm.

[1643] *Pouteria benai* (Aubrév. & Pellegr.) T.D.Penn.
(Fig. 55A)

Fl. Neotrop. Monogr. 52: 274 [26 Apr. 1990] (Pennington 1990). — *Podoluma benai* Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 182 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962), "Benai".

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *Service Forestier M-207* (holo-, P[P00640533]; iso-, P[P00640534], U[U0006626]).

INVENTORY DATA (FG). — 60 trees in 30 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 49.7$ cm.

[1644] *Pouteria bilocularis* (H.J.P.Winkl.) Baehni

Candollea 9: 229 (Baehni 1942). — *Labatia bilocularis* H.J.P.Winkl., *Repert. Spec. Nov. Regni Veg.* 7: 112 (Winkler 1909).

Pseudocladia neblinaensis Aubrév., *Mem. New York Bot. Gard.* 23: 204 (Aubréville 1972).

VERNACULAR NAMES. — Pa: impitit-wašiunō, mpitit-wašiunō • Br: abiurana.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *D. Sabatier 3524*.

INVENTORY DATA (FG). — 91 trees in 61 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43.6$ cm.

[1645] *Pouteria brachyandra* (Aubrév. & Pellegr.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 404 [26 Apr. 1990] (Pennington 1990). — *Radlkofarella brachyandra* Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 186 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

VERNACULAR NAMES. — Wp: pakuku, yá laāngá.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5005*.

INVENTORY DATA (FG). — 16 trees in 10 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 35.3$ cm.

[1646] *Pouteria caimito* (Ruiz & Pav.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 12: 312 (Radlkofe 1882). — *Achras caimito* Ruiz & Pav., *Flora Peruviana* 3: 18 (Ruiz & Pavón 1802). — *Lucuma caimito* (Ruiz & Pav.) Roem. & Schult., *Syst. Veg. [Roemer & Schultes]* 4: 701 [Mar.-June 1819] (Roemer & Schultes 1819). — *Caleatia caimito* (Ruiz & Pav.) Mart. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 2: 76. (Steudel 1841) nom. nud. pro syn. — *Labatia caimito* (Ruiz & Pav.) Mart., *Flora* 21 (2, Beibl.): 90 (Martius 1838). — *Guapeba caimito* (Ruiz & Pav.) Pierre, *Notes Bot. Sapot.* 2: 42 (Pierre 1891).

Guapeba laurifolia Gomes, *Mem. Math. Phis. Acad. Real Sci. Lisboa* 3 (Mem. Corresp. 1): 21 (Gomes 1812). — *Lucuma laurifolia* (Gomes) A.DC., *Prodr. [A. P. de Candolle]* 8: 166 [mid Mar. 1844] (Candolle 1844). — *Pouteria laurifolia* (Gomes) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 333 (Radlkofe 1882). — *Pouteria caimito* var. *laurifolia* (Gomes) Baehni, *Candollea* 9: 260 (Baehni 1942).

Lucuma temare Kunth, *Nova genera et species plantarum [H.B.K.]* 3: 241 [9 July 1819] (Kunth 1819). — *Richardella temare* (Kunth) Pierre, *Notes Bot. Sapot.* 1: 20 (Pierre 1890). — *Pouteria temare* (Kunth) Aubrév., *Adansonia*, n.s., 1 (2): 157 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

Labatia reticulata Mart., *Flora* 21 (2, Beibl.): 90 (Martius 1838). — *Lucuma laurifolia* var. *reticulata* (Mart.) A.DC., *Prodr. [A. P. de*

*Candolle] 8: 166 [mid Mar. 1844] (Candolle 1844). — *Guapeba brasiliensis* Steud., *Nomenc. Bot. [Steudel]*, ed. 2, 1: 708 (Steudel 1840), nom. illeg. superfl. (based on *Labatia reticulata*).*

Labatia lasiocarpa Mart., *Flora* 21 (2, Beibl.): 91 (Martius 1838). — *Lucuma lasiocarpa* (Mart.) A.DC., *Prod. [A. P. de Candolle]* 8: 166 [mid Mar. 1844] (Candolle 1844). — *Pouteria lasiocarpa* (Mart.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 333 (Radlkofe 1882). — *Guapeba lasiocarpa* (Mart.) Pierre, *Notes Bot. Sapot.* 2: 42 (Pierre 1891).

Acras guapeba Casar., *Nov. Stirp. Bras.* 7: 61 [Sep ("Jul") 1843] (Casaretto 1843).

Pouteria leucophaea Baehni, *Candollea* 18: 176 (Baehni 1962).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Ka: atakamala, kusili palatali • Wp: wila kea • Wn: alimihmo • Cr: zoliv • Br: abiurana-acariquara.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier* 2564.

INVENTORY DATA (FG). — 19 trees in 7 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 32.8$ cm.

[1647] *Pouteria cayennensis* (A.DC.) Eyma

Recueil Trav. Bot. Néerl. 33: 174 (Eyma 1936). — *Chrysophyllum cayennense* A.DC., *Prod. [A. P. de Candolle]* 8: 160 [mid Mar. 1844] (Candolle 1844), "Cayennense". — *Richardella cayennensis* (A.DC.) Aubrév., *Adansonia*, sér. 2, 11 (2): 300 [July 1971] (Aubréville 1971).

Chrysophyllum sessiliflorum Poir., *Encycl. [J. Lamarck et al.] Suppl.* 2: 16 [23 Oct. 1811] (Poiret 1811).

Lucuma pulverulenta Mart. & Eichler, *Fl. Bras. [Martius]* 7: 70 [15 Jan. 1863] (Martius & Eichler 1863). — *Vitellaria pulverulenta* (Mart. & Eichler) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 326 (Radlkofe 1882).

NOTE. — Pennington (1990: 366) states that the holotype is at P, and that there is an isotype at G-DC. However, there is at G-DC a specimen (G00139668) bearing a label in the hand of Alphonse de Candolle, which reads "Chrysophyllum Cayennense A. DC.", as well as another which reads "Sapotaceae. Cayenne ou Guyane françoise. Museum de Paris 1821". There is no reason not to admit that it is the very specimen to which Candolle refers in the protologue of *Chrysophyllum cayennense*, in the form "v. s. comm. a Mus. par." (i.e., "an exsiccatum that I saw, given by the Paris Museum"). In other words, the specimen at G-DC (G00139668) must be the holotype.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, G-DC[G00139668]; iso-, P[P00640548, P00640549, P00640550], US[00037030]).

INVENTORY DATA (FG). — 85 trees in 26 plots; $F_{\max} = 3.8\%$; $dbh_{\text{inv}} = 87$ cm.

[1648] *Pouteria cladantha* Sandwith

Bull. Misc. Inform. Kew 1931 (10): 480 [31 Dec. 1931] (Sandwith 1931). — *Neoxythecia cladantha* (Sandwith) Aubrév., *Adansonia*, n.s., 1 (2): 183 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville

1962). — *Richardella cladantha* (Sandwith) Baehni, *Boissiera* 11: 97 (Baehni 1965).

Pouteria loretensis Baehni, *Candollea* 9: 235 (Baehni 1942).

VERNACULAR NAMES. — Ka: kusali yepo • Wp: wai piyū • Cr: zoliv • Br: abiurana-seca, caramuri-preto.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *J.-J. de Granville et al.* 8099.

INVENTORY DATA (FG). — 36 trees in 11 plots; $F_{\max} = 2.2\%$; $dbh_{\text{inv}} = 125$ cm.

[1649] *Pouteria coriacea* (Pierre) Pierre

Symb. Antill. [Urban] 5 (1): 109 [20 May 1904] (Pierre 1904). — *Guapeba coriacea* Pierre, *Notes Bot. Sapot.* 2: 42 (Pierre 1891).

Chrysophyllum minutiflorum Britton, *Bull. Torrey Bot. Club* 48 (12): 337 ["1921" publ. 1922] (Britton 1922). — *Pouteria minutiflora* (Britton) Sandwith, *Bull. Misc. Inform. Kew* 1931 (10): 478 [31 Dec. 1931] (Sandwith 1931). — *Pseudocladiella minutiflora* (Britton) Aubrév., *Adansonia*, n.s., 1 (2): 165 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

VERNACULAR NAMES. — Pa: uu-kamwi-duwē • Wn: tépuklem.

HERBARIUM DATA (FG). — 61 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3886.

INVENTORY DATA (FG). — 70 trees in 28 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 55.3$ cm.

[1650] *Pouteria cuspidata* (A.DC.) Baehni
subsp. *dura* (Eyma) T.D.Penn.

Fl. Neotrop. Monogr. 52: 352 [26 Apr. 1990] (Pennington 1990). — *Pouteria dura* Eyma, *Recueil Trav. Bot. Néerl.* 33: 187 (Eyma 1936). — *Neoxythecia dura* (Eyma) Aubrév. & Pellegr., *Adansonia*, n.s., 1 (1): 17 [Nov. 1961] (Aubréville & Pellegrin 1961).

Sideroxylon cuspidatum A.DC. var. *crassifolium* Miq., *Fl. Bras. [Martius]* 7: 56 [15 Jan. 1863] (Miquel 1863).

Sideroxylon durum Klotzsch ex Miq., *Fl. Bras. [Martius]* 7: 56 [15 Jan. 1863] (Miquel 1863), nom. nud. pro syn.

Neoxythecia schulzii Aubrév., *Adansonia*, n.s., 4 (2): 231 [July 1964] (Aubréville 1964).

Neoxythecia wurdackii Aubrév., *Adansonia*, n.s., 5 (2): 201 [July 1965] (Aubréville 1965). — *Chrysophyllum wurdackii* (Aubrév.) Bernardi, *Candollea* 22 (2): 231 (Bernardi 1967).

NOTE. — *P. cuspidata* is hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Nt: nyanboka • Br: abiurana-arana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *D. Sabatier* 1430.

INVENTORY DATA (FG). — 22 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43$ cm.



FIG. 54. — Sapotaceae: **A**, *Chrysophyllum prieurii* A.DC. (D. Sabatier et al. 4620); **B**, *Manilkara bidentata* (A.DC.) A.Chev. subsp. *bidentata* (D. Sabatier 3553); **C**, *Micropholis acutangula* (D. Sabatier & S. Gonzalez 5371); **D**, *Micropholis obscura* T.D.Penn. (D. Sabatier 1221). A-C, © D. Sabatier/IRD; D, © M.-F. Prévost/IRD.

[1651] *Pouteria cuspidata* subsp. *robusta*
(Mart. & Eichler) T.D.Penn.

Fl. Neotrop. Monogr. 52: 351 [26 Apr. 1990] (Pennington 1990). — *Sideroxylon robustum* Mart. & Eichler, *Fl. Bras. [Martius]* 7: 56 [15 Jan. 1863] (Martius & Eichler 1863). — *Oxythecia robusta* (Mart. & Eichler) Pierre, *Symb. Antill. [Urban]* 5 (1): 161 [20 May 1904] (Pierre 1904), “*robustum*”. — *Pouteria robusta* (Mart. & Eichler) Eyma, *Recueil Trav. Bot. Néerl.* 33: 188 (Eyma 1936). — *Neoxythecia robusta* (Mart. & Eichler) Aubrév. & Pellegr., *Adansonia*, sér. 2, 1 (1): 17 [Nov. 1961] (Aubréville & Pellegrin 1961).

Pouteria robusta var. *longifolia* Eyma, *Recueil Trav. Bot. Néerl.* 33: 188 (Eyma 1936).

Pouteria robusta var. *typica* Baehni, *Candollea* 9: 213 (Baehni 1942), nom. inval. (Turland et al. 2018: Art. 24.3).

Oxythecia robusta var. *brevipetiolata* Monach., *Fieldiana, Bot.* 28 (3): 486 (Monachino 1953).

NOTES. — *P. cuspidata* is hyperdominant in Amazonia (ter Steege et al. 2020). Although Miquel provided the treatment of Sapotaceae for the *Flora Brasiliensis* (Miquel 1863), Martius clearly claims, in the footnote at the beginning of the treatment of this family (Miquel 1863: 37), his coauthorship with Eichler of some new species, among which is *Sideroxylon robustum* Mart. & Eichler.

VERNACULAR NAMES. — Ka: atakamala, kusili palatali.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & S. Gonzalez 5385*.

INVENTORY DATA (FG). — 4 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 29 cm.

[1652] *Pouteria decorticans* T.D.Penn.

Fl. Neotrop. Monogr. 52: 489 [26 Apr. 1990] (Pennington 1990).

VERNACULAR NAMES. — Pa: kuyau-kamwi-duwē, kuyau-kamwi-seinō, uu-kamwi-wahuyo • Wp: yawayi lū • Nt: nyanboka • Cr: zoliv • Br: abiu-giboa.

HERBARIUM DATA (FG). — 33 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2617*.

INVENTORY DATA (FG). — 147 trees in 38 plots; $F_{\max} = 3.3\%$; dbh_{inv} = 41.1 cm.

[1653] *Pouteria deliciosa* T.D.Penn.

Fl. Neotrop. Monogr. 52: 446 [26 Apr. 1990] (Pennington 1990).

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier 1213*.

INVENTORY DATA (FG). — 24 trees in 15 plots; $F_{\max} < 1\%$; dbh_{inv} = 75 cm.

[1654] *Pouteria durlandii* (Standl.) Baehni

Candollea 9: 422 (Baehni 1942). — *Lucuma durlandii* Standl., *Trop. Woods* 4: 5 (Standley 1925). — *Paralabatia durlandii* (Standl.) Aubrév., *Adansonia*, n.s., 3 (1): 21 [Jan.-Mar. 1963] (Aubréville 1963). — *Peteniodendron durlandii* (Standl.) Lundell, *Wrightia* 5 (7): 255 (Lundell 1976).

Pouteria suffulta Baehni, *Candollea* 14: 70 (Baehni 1952).

Pouteria potosina Lundell, *Wrightia* 5 (4): 101 (Lundell 1975). — *Peteniodendron potosinum* (Lundell) Lundell, *Wrightia* 5 (7): 255 (Lundell 1976). — *Paralabatia potosina* (Lundell) Aubrév., *Wrightia* 5 (9): 350 (Aubréville 1977).

Peteniodendron belizense Lundell, *Wrightia* 5 (7): 254 (Lundell 1976). — *Paralabatia belizensis* (Lundell) Lundell, *Wrightia* 5 (9): 350 (Lundell 1977).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *T.D. Pennington & S.A. Mori 12177*.

INVENTORY DATA (FG). — 14 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 19.4$ cm.

[1655] *Pouteria egregia* Sandwith

Bull. Misc. Inform. Kew 1931 (10): 479 [31 Dec. 1931] (Sandwith 1931). — *Sandwithiodoxa egregia* (Sandwith) Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 163 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

VERNACULAR NAMES. — Ka: olomelan • Wp: wila náná, wila náná sili • Br: abiu-pitomba, abiurana, cururu.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier 1219*.

INVENTORY DATA (FG). — 74 trees in 38 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 70$ cm.

[1656] *Pouteria engleri* Eyma

Recueil Trav. Bot. Néerl. 33: 178 (Eyma 1936). — *Nemaluma engleri* (Eyma) Aubrév. & Pellegr., *Adansonia*, n.s., 1 (1): 31 [Nov. 1961] (Aubréville & Pellegrin 1961).

Chrysophyllum alnifolium Engl., *Bot. Jahrb. Syst.* 12: 522 [23 Dec. 1890] (Engler 1890), nom. illeg. hom., non Baker (1877).

VERNACULAR NAMES. — Ka: atakamala, kusili palatali • Nt: nyanboka.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (original material P[P00640595, P00640596, P00640597]).

INVENTORY DATA (FG). — 71 trees in 44 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 72.9$ cm.

[1657] *Pouteria ephedrantha* (A.C.Sm.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 508 [26 Apr. 1990] (Pennington 1990). — *Lucuma ephedrantha* A.C.Sm., *Brittonia* 2 (2): 158 (Smith 1936).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier & J.-F. Molino 5717*.

INVENTORY DATA (FG). — 46 trees in 11 plots; $\text{dbh}_{\text{inv}} = 115$ cm.

[1658] *Pouteria eugenifolia* (Pierre) Baehni
(Fig. 55B)

Candollea 9: 218 (Baehni 1942). — *Micropholis eugenifolia* Pierre, *Notes Bot. Sapot.* 2: 40 (Pierre 1891), "eugeniaeifolia". — *Sideroxylon eugenifolium* (Pierre) Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachr.* 1: 276 (Engler 1897). — *Myrtiluma eugenifolia* (Pierre) Aubrév., *Adansonia*, n.s., 1 (2): 180 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962), "eugeniaeifolia".

Micropholis steyermarkii Monach., *Fieldiana, Bot.* 28 (3): 481 (Monachino 1953), nom. inval. (simultaneously published with *Sideroxylon steyermarkii*).

Pouteria steyermarkii Monach., *Fieldiana, Bot.* 28 (3): 481 (Monachino 1953), nom. inval. (simultaneously published with *Micropholis steyermarkii*).

VERNACULAR NAMES. — Ka: mandasabay • Nt: nyanboka • Br: batanga.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (holo, P[P00647901]; iso-, G[G00439472], P[P00640598, P00640599, P00647902]).

INVENTORY DATA (FG). — 177 trees in 78 plots; $F_{\max} = 1.2\%$; $\text{dbh}_{\text{inv}} = 98$ cm.

[1659] *Pouteria filipes* Eyma

Recueil Trav. Bot. Néerl. 33: 180 (Eyma 1936). — *Pseudolabatia filipes* (Eyma) Aubrév., *Adansonia*, n.s., 3 (1): 21 [Jan.-Mar. 1963] (Aubréville 1963).

VERNACULAR NAMES. — Pa: kuyau-kamwi-puvemna.

HERBARIUM DATA (FG). — 23 collections at CAY. Sel. exs.: *D. Sabatier 1516*.

INVENTORY DATA (FG). — 89 trees in 31 plots; $F_{\max} = 3.5\%$; $\text{dbh}_{\text{inv}} = 87$ cm.

[1660] *Pouteria fimbriata* Baehni

Candollea 14: 68 (Baehni 1952).

Pseudolabatia raoulantonis Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 168 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & B. Riéra 1741*.

INVENTORY DATA (FG). — 123 trees in 81 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 63$ cm.

[1661] *Pouteria flavilatex* T.D.Penn.

Rodriguésia 57 (2): 308 (Pennington 2006).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5128*.

INVENTORY DATA (FG). — 41 trees in 33 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 58.1$ cm.

[1662] *Pouteria franciscana* Baehni

Candollea 9: 262 (Baehni 1942).

VERNACULAR NAMES. — Wp: waa pitā, wapitā • Wn: palakwa • Nt: nyanboka • Cr: zoliv • Br: abiurana-branca.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1637*.

INVENTORY DATA (FG). — 24 trees in 7 plots; $F_{\max} = 1.9\%$; $\text{dbh}_{\text{inv}} = 104\text{ cm}$.

[1663] *Pouteria glomerata* (Miq.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 12: 333 (Radlkofer 1882). — *Lucuma glomerata* Miq., *Fl. Bras. [Martius]* 7: 81 [15 Jan. 1863] (Miquel 1863). — *Labatia glomerata* (Miq.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 14: 451 (Radlkofer 1884). — *Guapeba glomerata* (Miq.) Pierre, *Notes Bot. Sapot.* 2: 43 (Pierre 1891). — *Richardella glomerata* (Miq.) Baehni, *Boissiera* 11: 97 (Baehni 1965). — *Neolabatia glomerata* (Miq.) Aubrév., *Mem. New York Bot. Gard.* 23: 203 (Aubréville 1972).

Labatia parinariooides Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 14: 451 (Radlkofer 1884). — *Guapeba parinariooides* (Radlk.) Pierre, *Notes Bot. Sapot.* 2: 43 (Pierre 1891).

Pouteria weddelliana Pierre, *Notes Bot. Sapot.* 2: 45 (Pierre 1891).

Pouteria glomerata var. *glabrescens* Huber, *Bull. Soc. Bot. Genève*, sér. 2, 6: 197 ["1914" publ. 1915] (Huber 1915), "Ponteria".

Labatia parviflora Pittier, *Arb. Arbust. Venez.* 2-3: 41 (Pittier 1923). — *Neolabatia parviflora* (Pittier) Aubrév., *Mem. New York Bot. Gard.* 23: 203 (Aubréville 1972).

Lucuma hypoglauca Standl., *Lista Pl. Salvador [Standley & S. Calderon]*: 170 (Standley 1925). — *Pouteria hypoglauca* (Standl.) Baehni, *Candollea* 9: 250 (Baehni 1942). — *Richardella hypoglauca* (Standl.) Baehni, *Boissiera* 11: 97 (Baehni 1965). — *Neolabatia hypoglauca* (Standl.) Aubrév., *Mem. New York Bot. Gard.* 23: 203 (Aubréville 1972).

Pouteria glomerata var. *typica* Baehni, *Candollea* 9: 243 (Baehni 1942), nom. inval. (Turland et al. 2018: Art. 24.3).

Pouteria pittieri Baehni, *Candollea* 9: 254 (Baehni 1942).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *D. Sabatier 1488*.

INVENTORY DATA (FG). — 7 trees in 3 plots; $F_{\max} = 1.1\%$; $\text{dbh}_{\text{inv}} = 43\text{ cm}$.

[1664] *Pouteria gonggrijpii* Eyma

Recueil Trav. Bot. Néerl. 33: 185 (Eyma 1936), "Gonggrijpii". — *Franchetella gonggrijpii* (Eyma) Aubrév., *Adansonia*, n.s., 1 (2): 183 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962), "Gonggrijpii".

NOTE. — This species is dedicated to the Dutch collector Justus Willem Gonggrijp. Eyma himself wrote "Pouteria gonggrijpii" on the type. The spelling "gonggrijpii" should therefore be taken as an

unintentional typographical error, and corrected (Turland et al. 2018: Art. 60.1).

VERNACULAR NAMES. — Pa: kuyau-kamwi, tukuyuy-kamwi, uu-kamwi, uu-kamwi-duwē • Ka: ololome, olome • Te: wila pitāg tsing • Nt: nyanboka • Cr: balata-pwé • Fr: balata poirier • Br: abiurana-vermelha.

HERBARIUM DATA (FG). — 100 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 4304*.

INVENTORY DATA (FG). — 611 trees in 126 plots; $F_{\max} = 8.6\%$; $\text{dbh}_{\text{inv}} = 51\text{ cm}$.

[1665] *Pouteria grandis* Eyma

Recueil Trav. Bot. Néerl. 33: 170 (Eyma 1936). — *Radlkoferella grandis* (Eyma) Aubrév., *Adansonia*, n.s., 1 (2): 183 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: uu-aška.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *D. Sabatier 2450*.

INVENTORY DATA (FG). — 101 trees in 41 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 70.7\text{ cm}$.

[1666] *Pouteria guianensis* Aubl.

Hist. Pl. Guiane 1: 85 [Jun.-Dec. 1775] (Aublet 1775). — *Labatia pedunculata* Willd., *Sp. Pl.*, ed. 4 1 (2): 624 [July 1798] (Willdenow 1798), nom. illeg. superfl. (based on *Pouteria guianensis*). — *Pouteria pedunculata* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 4: 546 [29 June 1816] (Poirer 1816).

Lucuma psammophila (Mart.) A.DC. var. *macrophylla* Raunk., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1889: 6 (Raunkiaer 1889).

Labatia tovarensis Engl., *Bot. Jahrb. Syst.* 12: 515 [23 Dec. 1890] (Engler 1890). — *Pouteria tovarensis* (Engl.) Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 4 (1): 142 (Engler 1891).

Guapeba glazioviana Pierre, *Notes Bot. Sapot.* 2: 43 (Pierre 1891), "Glazioveana", nom. nud. — *Pouteria glazioviana* Dubard, *Notul. Syst. (Paris)* 1: 380 (Dubard 1911), "Glazioveana", nom. nov., based on *Lucuma psammophila* var. *macrophylla*.

Krugella hartii Pierre, *Notes Bot. Sapot.* 2: 52 (Pierre 1891). — *Pouteria hartii* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, 10: 34 (Dubard 1912).

Lucuma hartii Hemsl., *Hooker's Icon. Pl.* 26 [ser. 4, 6]: t. 2565 ["1899" publ. Apr. 1898] (Hemslay 1898).

Lucuma glazioviana Pierre ex Glaz., *Bull. Soc. Bot. France* 57 (Mem. 3e): 438 (Glaziou 1910), nom. nud.

Pouteria obidensis Huber, *Bull. Soc. Bot. Genève*, sér. 2, 6: 197 ["1914" publ. 1915] (Huber 1915).

Pouteria demerarae Sandwith, *Bull. Misc. Inform. Kew* 1931 (10): 477 [31 Dec. 1931] (Sandwith 1931).

Lucuma huallagae Standl. ex Ll.Williams, *Publ. Field Mus. Nat. Hist., Bot. Ser.* 15: 411 (Williams 1936).

Pouteria caimito var. *strigosa* Monach., *Fieldiana, Bot.* 28 (3): 487 (Monachino 1953).

Pouteria cuatrecasasii Aubrév., *Adansonia, n.s.*, 7 (2): 141 [Jul.-Sep. 1967] (Aubréville 1967).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *Pouteria hartii* (Pierre) Dubard is based on both *Krugella hartii* Pierre and *Lucuma hartii* Hemsl. These names have distinct types, which are both cited by Dubard (1912: 35).

VERNACULAR NAMES. — Pa: balata-kamwi, uu-ahavukune, uu-kamwi-seine • Ka: asepoko, asepuku, kulumoto, wasepuku • Te: wila pitág piñág • Wp: wau, wawiyu • Wn: alimihmo • Nt: nyanboka • Cr: grènn-kwata, jonndèf • Fr: jaune d'œuf • Br: abiurana-abiu, abiurana-branca, abiurana-casca-fina.

HERBARIUM DATA (FG). — 79 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777520] designated by Lanjouw & Uittien [1940: 156]).

INVENTORY DATA (FG). — 411 trees in 147 plots; $F_{\max} = 2.6\%$; $dbh_{\text{inv}} = 70$ cm.

[1667] *Pouteria hispida* Eyma

Recueil Trav. Bot. Néerl. 33: 177 (Eyma 1936).

Pouteria trichopoda Baehni, *Candollea* 9: 245 (Baehni 1942).

Pouteria solimoensis Aubrév. & Pellegr., *Adansonia, n.s.*, 1 (2): 157 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

Pouteria benoistii Aubrév., *Adansonia, n.s.*, 4 (2): 228 [July 1964] (Aubréville 1964).

VERNACULAR NAMES. — Pa: uu-kamwi • Ka: asepuku, tuwonule, wasepuku • Wp: wawiyu • Nt: nyanboka • Br: abiurana-braba, abiurana-jarani.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *E.M. Mélinon 108*, 1863 (holotype of *Pouteria benoistii*: P[P00647949]; iso-, P[P00647950, P00647951]).

INVENTORY DATA (FG). — 100 trees in 62 plots; $F_{\max} = 1\%$; $dbh_{\text{inv}} = 110$ cm.

[1668] *Pouteria jariensis* Pires & T.D.Penn.
(Fig. 55D)

Fl. Neotrop. Monogr. 52: 331 [26 Apr. 1990] (Pires & Pennington 1990).

VERNACULAR NAMES. — Pa: tukuyuy-kamwi • Cr: zoliv • Br: abiurana.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *J.-F. Molino 1198*.

INVENTORY DATA (FG). — 95 trees in 64 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 67$ cm.

[1669] *Pouteria laevigata* (Mart.) Radlk.

Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München 14: 457 (Radlkofler 1884). — *Labatia laevigata* Mart., *Flora* 21 (2, Beibl.): 92 (Martius 1838). — *Lucuma laevigata* (Mart.) A.DC., *Prod. [A. P. de Candolle]* 8: 167 [mid Mar. 1844] (Candolle 1844). — *Gua-peba laevigata* (Mart.) Pierre, *Notes Bot. Sapot.* 2: 42 (Pierre 1891).

Pouteria bouffardiana Bernardi, *Candollea* 22 (2): 228 (Bernardi 1967).

VERNACULAR NAMES. — Wp: wau átā • Br: abiurana-de-casca-grossa, abiurana-sapota.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *D. Sabatier 1098*.

INVENTORY DATA (FG). — 18 trees in 15 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 85.1$ cm.

[1670] *Pouteria macrocarpa* (Mart.) D.Dietr.

Syn. Pl. [D. Dietrich] 1: 431 [July 1839] (Dietrich 1839). — *Labatia macrocarpa* Mart., *Nova genera et species plantarum [Martius]* 2 (2): 71 [Jan.-June 1827] (Martius 1827). — *Richardella macrocarpa* (Mart.) Baehni, *Boissiera* 11: 97 (Baehni 1965). — *Neolabatia macrocarpa* (Mart.) Aubrév., *Mem. New York Bot. Gard.* 23: 203 (Aubréville 1972).

Lucuma paraensis Standl., *Trop. Woods* 33: 13 (Standley 1933). — *Pouteria paraensis* (Standl.) Baehni, *Candollea* 9: 424 (Baehni 1942).

NOTE. — Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: uu-kamwi-duwē, uu-purubumna • Wp: pakuku, yá laánga • Br: abiu-grande, cutite-grande.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *P. Grenand 699*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 30.2$ cm.

[1671] *Pouteria macrophylla* (Lam.) Eyma

Recueil Trav. Bot. Néerl. 33: 164 (Eyma 1936). — *Chrysophyllum macrophyllum* Lam., *Tabl. Encycl. 2[5 (2)]*: 44 [1 Sep. 1794] (Lamarck 1794). — *Richardella macrophylla* (Lam.) Aubrév., *Adansonia, n.s.*, 1 (2): 175 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

Bumelia nervosa Vahl, *Eclog. Amer.* 1: 28 (Vahl 1797).

Lucuma rivicoa C.F.Gaertn., *Suppl. Carp.* 130 (Gaertner 1807). — *Vitellaria rivicoa* (C.F.Gaertn.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 326 (Radlkofler 1882). — *Richardella rivicoa* (C.F.Gaertn.) Pierre, *Notes Bot. Sapot.* 1: 19 (Pierre 1890).

Achras glaucescens Willd. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 15 (Steudel 1840), *nom. nud.*

Vitellaria glaucophylla Engl., *Bot. Jahrb. Syst.* 12: 513 [23 Dec. 1890] (Engler 1890). — *Lucuma rivicoa* var. *glaucophylla* (Engl.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, 10: 15 (Dubard 1912).

Lucuma acreana K.Krause, *Notizbl. Königl. Bot. Gart. Berlin* 6: 169 [4 Apr. 1914] (Krause 1914).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020), but not yet found in inventories of trees above 10 cm dbh in French Guiana. Semi-domesticated by pre-Columbian Amerindians (Clement 1999; Levis *et al.* 2017).

VERNACULAR NAMES. — Pa: uu • Wp: akusi yeti • Nt: sooke • Cr: jonndèf • Br: abiurana-cutiti, acará-uba, cutite, cutitiribá.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *J.P.B. von Rohr* 125 (original material of *Bumelia nervosa*: C[[C10018776](#), [C10018777](#)]).

SIZE. — Brazil, Amazonas. *G.T. Prance et al.* [22998](#) (MO), 15 m × 15 cm.

[1672] *Pouteria maxima* T.D.Penn.

Rodriguésia [57](#) (2): 319 (Pennington 2006).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *T.D. Pennington & S.A. Mori* [12137](#).

INVENTORY DATA (FG). — 5 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 110$ cm.

[1673] *Pouteria melanopoda* Eyma

Recueil Trav. Bot. Néerl. [33](#): 174 (Eyma 1936).

VERNACULAR NAMES. — Ka: asepokolan • Br: abiurana-goiabinha.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* [1843](#).

INVENTORY DATA (FG). — 61 trees in 49 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 61.3$ cm.

[1674] *Pouteria oblanceolata* Pires

Bol. Técn. Inst. Agron. N. [38](#): 38 (Pires 1960).

VERNACULAR NAMES. — Nt: nyanboka • Br: abiurana.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier* [1106](#).

INVENTORY DATA (FG). — 4 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 47$ cm.

[1675] *Pouteria platyphylla* (A.C.Sm.) Baehni

Candollea [9](#): 274 (Baehni 1942). — *Lucuma platyphylla* A.C.Sm., *Bull. Torrey Bot. Club* [60](#) (6): 388 [1 June 1933] (Smith 1933). — *Franchetella platyphylla* (A.C.Sm.) Aubrév., *Adansonia*, n.s., [1](#) (2): 183 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

VERNACULAR NAMES. — Pa: uu-kamwi • Br: abiurana-vermelha.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier* [5843](#).

INVENTORY DATA (FG). — 11 trees in 11 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40$ cm.

[1676] *Pouteria procera* (Mart.) K.Hammer

Verz. Landwirtsch. Gartn. Kulturpfl., ed. 2, 2: 1046 (Hammer 1986). — *Lucuma procera* Mart., *Flora* [22](#) (1, Beibl.): 57 (Martius 1839). — *Vitellaria procera* (Mart.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* [12](#): 325 (Radlkofer 1882). — *Urbanella procera* (Mart.) Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890). — *Calocarpum procerum* (Mart.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, [10](#): 8 (Dubard 1912).

Lucuma procera var. *cuspidata* Miq., *Fl. Bras. [Martius]* [7](#): 72 [15 Jan. 1863] (Miquel 1863). — *Urbanella cuspidata* (Miq.) Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890). — *Calocarpum procerum* var. *cuspidatum* (Miq.) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, [10](#): 8 (Dubard 1912).

Urbanella buchananiiifolia Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890). — *Lucuma buchananiiifolia* Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890), "buchananiaeifolia", nom. nud. pro syn. — *Calocarpum buchananiiifolium* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, [10](#): 9 (Dubard 1912), "buchananiaeifolium". — *Achras buchananiiifolia* (Pierre) Bernardi, *Candollea* [22](#) (2): 230 (Bernardi 1967).

Urbanella oblonga Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890). — *Lucuma oblonga* Pierre, *Notes Bot. Sapot.* [1](#): 25 (Pierre 1890), nom. nud. pro syn. — *Calocarpum procerum* var. *oblongum* (Pierre) Dubard, *Ann. Mus. Colon. Marseille*, sér. 2, [10](#): 9 (Dubard 1912), "oblongifolium".

Lucuma excelsa A.C.Sm., *Brittonia* [2](#) (2): 159 (Smith 1936). — *Pouteria excelsa* (A.C.Sm.) Baehni, *Candollea* [9](#): 350 (Baehni 1942). — *Urbanelia excelsa* (A.C.Sm.) Aubrév., *Adansonia*, n.s., [3](#) (1): 21 [Jan.-Mar. 1963] (Aubréville 1963).

VERNACULAR NAMES. — Br: maparajúba-vermelha.

HERBARIUM DATA (FG). — A single collection, *J.-J. de Granville et al.* [7632](#) (P).

SIZE. — Up to 100 cm dbh (Pennington 1990).

[1677] *Pouteria putamen-ovi* T.D.Penn.

Fl. Neotrop. Monogr. [52](#): 270 [26 Apr. 1990] (Pennington 1990).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* [3407](#).

INVENTORY DATA (FG). — 1 tree, dbh = 44.4 cm.

[1678] *Pouteria reticulata* (Engl.) Eyma

Recueil Trav. Bot. Néerl. [33](#): 183 (Eyma 1936). — *Chrysophyllum reticulatum* Engl., *Bot. Jahrb. Syst.* [12](#): 522 [23 Dec. 1890] (Engler 1890). — *Franchetella reticulata* (Engl.) Aubrév., *Adansonia*, n.s., [1](#) (2): 183 ["Dec. 1961" publ. Jan.-Mar. 1962] (Aubréville 1962).

Lucuma minutiflora Allemão, Trab. Comm. Sci. Expl., Bot., Rio de Janeiro [1](#): 57 (Allemão 1866).

Sideroxylon uniloculare Donn.Sm., *Bot. Gaz.* [35](#) (1): 5 [20 Jan. 1903] (Donnell Smith 1903). — *Pouteria unilocularis* (Donn.Sm.) Baehni, *Candollea* [9](#): 273 (Baehni 1942). — *Franchetella unilocularis* (Donn.Sm.) Aubrév., *Adansonia*, n.s., [3](#) (1): 21 [Jan.-Mar. 1963] (Aubréville 1963).

Sideroxylon hondurensis Pittier, *Contr. U.S. Natl. Herb.* 20 (12): 483 [9 Jan. 1922] (Pittier 1922). — *Pouteria hondurensis* (Pittier) Baehni, *Candollea* 9: 228 (Baehni 1942).

Sideroxylon meyeri Standl., *Trop. Woods* 31: 45 (Standley 1932). — *Pouteria meyeri* (Standl.) Baehni, *Candollea* 9: 272 (Baehni 1942).

Lucuma anibifolia A.C.Sm., *Bull. Torrey Bot. Club* 60 (6): 389 [June 1933] (Smith 1933), “*anibifolia*”. — *Pouteria anibifolia* (A.C.Sm.) Baehni, *Candollea* 9: 269 (Baehni 1942). — *Franchetella anibifolia* (A.C.Sm.) Aubrév., *Adansonia*, n.s., 1 (2): 183 [“Dec. 1961” publ. Jan.-Mar. 1962] (Aubréville 1962).

Pouteria cearensis Baehni, *Candollea* 9: 279 (Baehni 1942).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kuyau-kamwi-duwē • Ka: olomelan • Br: abiurana-cascuda.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5322bis.

INVENTORY DATA (FG). — 86 trees in 27 plots; $F_{\max} = 3.7\%$; $\text{dbh}_{\text{inv}} = 53.8$ cm.

[1679] *Pouteria retinervis* T.D.Penn.

Fl. Neotrop. Monogr. 52: 308 [26 Apr. 1990] (Pennington 1990).

VERNACULAR NAMES. — Br: abiurana-grande.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom* 15239 (holo-, NY[00335312]; iso-, CAY[CAY044794], K, NO[NO0109877]).

INVENTORY DATA (FG). — 9 trees in 7 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 51.9$ cm.

[1680] *Pouteria rodriquesiana* Pires & T.D.Penn.

Fl. Neotrop. Monogr. 52: 379 [26 Apr. 1990] (Pires & Pennington 1990).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: uu-kamwi, uu-kamwi-seine • Te: wila pitág hun • Nt: nyanboka • Br: abiu-maparajúba.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *B. Maguire et al.* 47097 (holo-, K[K000994418, K000994419]; iso-, NY[01212902], US[00516534]).

INVENTORY DATA (FG). — 70 trees in 29 plots; $F_{\max} = 2.4\%$; $\text{dbh}_{\text{inv}} = 80$ cm.

[1681] *Pouteria sagotiana* (Baill.) Eyma
(Fig. 55C)

Recueil Trav. Bot. Néerl. 33: 170 (Eyma 1936). — *Eremoluma sagotiana* Baill., *Hist. Pl. [Baillon]* 11: 292 [Sep.-Oct. 1891] (Baillon 1891). — *Lucuma sagotiana* (Baill.) Engl., *Nat. Pflanzenfam. [Engler & Prantl] Nachtr.* 1: 274 (Engler 1897).

VERNACULAR NAMES. — Pa: inam-etni-seinō, waaduk • Ka: asinadan.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *P.A. Sagot* 1189, 1858 (holo-, P[P00648047]).

INVENTORY DATA (FG). — 26 trees in 16 plots; $F_{\max} = 2\%$; $\text{dbh}_{\text{inv}} = 30.9$ cm.

[1682] *Pouteria singularis* T.D.Penn.

Fl. Neotrop. Monogr. 52: 459 [26 Apr. 1990] (Pennington 1990).

VERNACULAR NAMES. — Pa: kuyau-kamwi-duwō, uu-kamwi • Nt: nyanboka • Cr: jonndèf, zoliv • Fr: jaune d’œuf.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 2645.

INVENTORY DATA (FG). — 149 trees in 62 plots; $F_{\max} = 2.8\%$; $\text{dbh}_{\text{inv}} = 60$ cm.

[1683] *Pouteria speciosa* (Ducke) Baehni

Candollea 9: 398 (Baehni 1942). — *Lucuma speciosa* Ducke, *Arch. Mus. Nac. Rio de Janeiro* 22: 67 (Ducke 1919). — *Englerella speciosa* (Ducke) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 230 (Ducke 1922), nom. inval. (new combination not formally accepted by the author). — *Richardella speciosa* (Ducke) Aubrév., *Adansonia*, n.s., 1 (2): 175 [“Dec. 1961” publ. Jan.-Mar. 1962] (Aubréville 1962).

Englerella macrocarpa Pierre, *Notes Bot. Sapot.* 2: 47 (Pierre 1891).

VERNACULAR NAMES. — Wn: kanapupali • Nt: taka, tyaka • Br: pajurá, pajurá-de-obidos.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino* 5117.

INVENTORY DATA (FG). — 30 trees in 9 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 88.4$ cm.

[1684] *Pouteria stipulifera* T.D.Penn.

Rodriguésia 57 (2): 283 (Pennington 2006).

HERBARIUM DATA (FG). — A single collection, *J. Engel & P. Petronelli* 19 (CAY).

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 29.3$ cm.

[1685] *Pouteria tenuisepala* Pires & T.D.Penn.

Fl. Neotrop. Monogr. 52: 435 [26 Apr. 1990] (Pires & Pennington 1990).

VERNACULAR NAMES. — Wp: ei li.

HERBARIUM DATA (FG). — 15 collections at CAY. Sel. exs.: *W.A. Egler* 47673 (holo-, MG, not seen; iso-, K[K000641152]).

INVENTORY DATA (FG). — 24 trees in 7 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 38$ cm.

[1686] *Pouteria torta* (Mart.) Radlk.
subsp. *glabra* T.D.Penn.

Fl. Neotrop. Monogr. 52: 484 [26 Apr. 1990] (Pennington 1990).

Lucuma jenmanii Pittier, *Contr. U.S. Natl. Herb.* 13: 458 (Pittier 1912). — *Pouteria jenmanii* (Pittier) Sandwith, *Bull. Misc. Inform. Kew* 1931 (10): 478 [31 Dec. 1931] (Sandwith 1931).

Lucuma gutta Ducke, *Ann. Acad. Brasil Sci.* 6: 208 (Ducke 1934). — *Pouteria gutta* Ducke, *Ann. Acad. Brasil Sci.* 6: 208; (Ducke 1934) in nota.

Pouteria gutta Baehni, *Candollea* 9: 261 (Baehni 1942), isonym.

Lucuma dolichophylla Standl. ex Ll.Williams, *Bot. Mus. Leafl.* 13 (9): 289 [18 Apr. 1949] (Williams 1949).

NOTES. — Schultes (1949: 289) stated that he only translated into latin Ll. Williams' English description of *Lucuma dolichophylla*, and he unequivocally ascribed authorship to Ll. Williams. *Lucuma gutta* Ducke and *Pouteria gutta* Ducke are alternative names (published simultaneously for the same taxon), but both are nevertheless valid, because they were published before 1953 (Turland *et al.* 2018: Art. 36.3). Therefore, the transfer of *Lucuma gutta* to *Pouteria* by Baehni is superfluous.

VERNACULAR NAMES. — Pa: uu-kamwi, uu-kamwi-wahuyo • Wp: wamisalu, wau • Nt: nyanboka • Cr: zoliv • Br: abiurana-camazal, curiola, grão-de-galo, guapeva.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1351*.

INVENTORY DATA (FG). — 203 trees in 92 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 55.4$ cm.

[1687] *Pouteria torta* subsp. *tuberculata*
(Sleumer) T.D.Penn.

Fl. Neotrop. Monogr. 52: 483 [26 Apr. 1990] (Pennington 1990). — *Lucuma tuberculata* Sleumer, *Repert. Spec. Nov. Regni Veg.* 45: 18 (Sleumer 1938). — *Xantolis tuberculata* (Sleumer) Baehni, *Boissiera* 11: 22 (Baehni 1965).

Pouteria neglecta Cronquist, *Lloydia* 9 (4): 286 [“1945” publ. 20 Dec. 1946] (Cronquist 1946).

Pouteria cooperi Cronquist, *Lloydia* 9 (4): 291 [“1945” publ. 20 Dec. 1946] (Cronquist 1946). — *Calocarpum cooperi* (Cronquist) Lundell, *Wrightia* 5 (7): 252 (Lundell 1976).

Pouteria wurdackii Aubrév., *Adansonia*, n.s., 5 (2): 201 [July 1965] (Aubréville 1965).

Pouteria echinocarpa W.A.Rodrigues, *Acta Amazonica* 4 (3): 9 (Rodrigues 1974).

VERNACULAR NAMES. — Nt: nyanboka.

HERBARIUM DATA (FG). — 22 collections at CAY. Sel. exs.: *D. Larpin 885*.

INVENTORY DATA (FG). — 78 trees in 44 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 51.3$ cm.

[1688] *Pouteria venosa* (Mart.) Baehni subsp. *amazonica*
T.D.Penn.

Fl. Neotrop. Monogr. 52: 399 [26 Apr. 1990] (Pennington 1990).

Lucuma littoralis Mart., *Fl. Bras. [Martius]* 7: 65 [15 Jan. 1863] (Martius 1863). — *Vitellaria littoralis* (Mart.) Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 326 (Radlkofe 1882). — *Radlkofeella littoralis* (Mart.) Pierre, *Notes Bot. Sapot.* 1: 21 (Pierre 1890).

Vitellaria dissepala K.Krause, *Notizbl. Königl. Bot. Gart. Berlin* 6: 171 [4 Apr. 1914] (Krause 1914). — *Lucuma dissepala* (K.Krause) Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 233 (Ducke 1922). — *Pouteria dissepala* (K.Krause) Rizzini, *Revista Brasil. Biol.* 33 (4): 599 (Rizzini 1973), nom. inval. (no reference to the basionym).

Lucuma duckei Huber, *Bull. Soc. Bot. Genève*, sér. 2, 6: 194 [“1914” publ. 1915] (Huber 1915).

Pouteria marginata (Mart. & Eichler) Rizzini var. *impressa* Rizzini, *Revista Brasil. Biol.* 33 (4): 599 (Rizzini 1973).

Pouteria dasystyla Rizzini, *Revista Brasil. Biol.* 33 (4): 599 (Rizzini 1973).

Pouteria dasystyla var. *abaetensis* Rizzini, *Revista Brasil. Biol.* 33 (4): 600 (Rizzini 1973).

VERNACULAR NAMES. — Ka: wasepuku • Wp: pakuku, wamisalu, yá laángá • Nt: nyanboka • Br: abiurana, cutitiriba-rana.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 1995*.

INVENTORY DATA (FG). — 55 trees in 30 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 44.4$ cm.

[1689] *Pouteria virescens* Baehni

Candollea 14: 66 (Baehni 1952).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier 1528*.

INVENTORY DATA (FG). — 7 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 107$ cm.

[1690] *Pouteria williamii* (Aubrév. & Pellegr.) T.D.Penn.

Fl. Neotrop. Monogr. 52: 338 [26 Apr. 1990] (Pennington 1990). — *Eremoloma williamii* Aubrév. & Pellegr., *Adansonia*, n.s., 1 (2): 169 [“Dec. 1961” publ. Jan.-Mar. 1962] (Aubréville & Pellegrin 1962).

VERNACULAR NAMES. — Pa: á-maruibika, balata-kamwi • Wp: kalamúli • Br: abiurana-balatatinha.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2787*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $dbh_{\text{inv}} = 26.2$ cm.

[1691] *Pouteria* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori & J.J. Pipoly 15515*.

INVENTORY DATA (FG). — 1 tree, $dbh = 17$ cm.

[1692] *Pouteria* sp. B

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3651*.

INVENTORY DATA (FG). — 14 trees in 14 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 54.6 \text{ cm}$.

[1693] *Pouteria* sp. C

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2343*.

INVENTORY DATA (FG). — 15 trees in 13 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 33.5 \text{ cm}$.

[1694] *Pouteria* sp. D

HERBARIUM DATA (FG). — 11 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 3648*.

INVENTORY DATA (FG). — 51 trees in 35 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 53.3 \text{ cm}$.

[1695] *Pouteria* sp. E

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 1266*.

INVENTORY DATA (FG). — 131 trees in 29 plots; $F_{\max} = 6\%$; $\text{dbh}_{\text{inv}} = 52.2 \text{ cm}$.

[1696] *Pouteria* sp. F

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-F. Molino 1617*.

INVENTORY DATA (FG). — 12 trees in 10 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 57.9 \text{ cm}$.

[1697] *Pouteria* sp. G

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5554*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 86 \text{ cm}$.

[1698] *Pouteria* sp. H

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5550*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 33.4 \text{ cm}$.

[1699] *Pouteria* sp. I

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5628*.

INVENTORY DATA (FG). — 14 trees in 3 plots; $F_{\max} = 2.1\%$; $\text{dbh}_{\text{inv}} = 34.9 \text{ cm}$.

Genus *Pradosia* Liais[1700] *Pradosia cochlearia* (Lecomte) T.D.Penn.

Fl. Neotrop. Monogr. 52: 656 [26 Apr. 1990] (Pennington 1990). — *Chrysophyllum cochlearium* Lecomte, *Notul. Syst. (Paris)* 4: 63 (Lecomte 1923). — *Ecclinusa cochlearia* (Lecomte) Aubrév., *Adansonia*, n.s., 1 (1): 20 [Nov. 1961] (Aubréville 1961).

VERNACULAR NAMES. — Pa: uu-kamwi-puvemna • Nt: gaan kinboto • Br: casca-doce, miraceem.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *R. Benoist 224* (holo-, P[P00649438]; iso-, P[P00649439]).

INVENTORY DATA (FG). — 449 trees in 78 plots; $F_{\max} = 4.1\%$; $\text{dbh}_{\text{inv}} = 129.9 \text{ cm}$.

[1701] *Pradosia huberi* (Ducke) Ducke

Trop. Woods 71: 16 (Ducke 1942). — *Glycoxyylon huberi* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 3: 235 (Ducke 1922).

VERNACULAR NAMES. — Pa: á-maruibika • Br: casca-doce, pau-doce, pracuuba-de-leite.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2895*.

INVENTORY DATA (FG). — 7 trees in 4 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 50 \text{ cm}$.

[1702] *Pradosia ptychandra* (Eyma) T.D.Penn.

(Fig. 55E)

Fl. Neotrop. Monogr. 52: 648 [26 Apr. 1990] (Pennington 1990). — *Pouteria ptychandra* Eyma, *Recueil Trav. Bot. Néerl.* 33: 189 (Eyma 1936). — *Neopometia ptychandra* (Eyma) Aubrév., *Adansonia*, n.s., 1 (1): 26 [Nov. 1961] (Aubréville 1961).

Voyara montana Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 26 [Jun.-Dec. 1775] (Aublet 1775), pro parte fruto excluso. — *Capparis montana* (Aubl.) Lemée, *Fl. Guyane Franç.* 1: 666 (Lemée 1955).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: inam-etni-seinô • Wp: tasiasiy, wila ē̄s • Wn: tamaniwa • Nt: busi amanda, busi amandaa, kinboto • Cr: balata-ponm, grènn-kwata • Fr: balata pommier • Br: casca-doce.

HERBARIUM DATA (FG). — 78 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material of *Voyara montana*: BM[BM000952572]).

INVENTORY DATA (FG). — 229 trees in 97 plots; $F_{\max} = 2.2\%$; $\text{dbh}_{\text{inv}} = 58.6 \text{ cm}$.




FIG. 55. — Sapotaceae: **A**, *Pouteria benai* (Aubrév. & Pellegr.) T.D.Penn. (*D. Sabatier & M.-F. Prévost* 3533); **B**, *Pouteria eugeniifolia* (Pierre) Baehni (*D. Sabatier* 5045); **C**, *Pouteria sagotiana* (Baill.) Eyma; *Pradosia ptychandra* (Eyma) T.D.Penn. (*D. Sabatier* 823); **D**, *Pouteria jariensis* Pires & T.D.Penn. (*D. Sabatier & J.-F. Molino* 5541); **E**, *Pradosia ptychandra* (Eyma) T.D.Penn. (*D. Sabatier* 823). A, B, D, © D. Sabatier/IRD; C, © J.-F. Molino/IRD; E, © M.-F. Prévost/IRD.

[1703] *Pradosia surinamensis* (Eyma) T.D.Penn.

Fl. Neotrop. Monogr. 52: 652 [26 Apr. 1990] (Pennington 1990). — *Pouteria surinamensis* Eyma, *Recueil Trav. Bot. Néerl.* 33: 189 (Eyma 1936).

VERNACULAR NAMES. — Ka: ololome, olome • Wn: tamaniwa • Nt: kinboto • Br: abiurana-branca.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost* 3319.

INVENTORY DATA (FG). — 3 trees in 1 plots; $F_{\max} < 1\%$; dbh_{inv} = 35 cm.

[1704] *Pradosia verticillata* Ducke

Trop. Woods 71: 12 (Ducke 1942).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *T.D. Pennington & S.A. Mori* 12134.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; dbh_{inv} = 14.2 cm.

[1705] *Pradosia* sp. A

NOTE. — A species resembling *P. subverticillata* Ducke.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *D. Sabatier* 1772, dbh = 80 cm.

Genus *Sarcaulus* Radlk.

[1706] *Sarcaulus brasiliensis* (A.DC.) Eyma

Recueil Trav. Bot. Néerl. 33: 192 (Eyma 1936). — *Chrysophyllum brasiliense* A.DC., *Prodr. [A. P. de Candolle]* 8: 156 [mid Mar. 1844] (Candolle 1844).

Chrysophyllum macrophyllum Mart., *Flora* 21 (2, Beibl.): 95 (Martius 1838), nom. illeg. hom., non Lam. (Lamarck 1794) nec C.F.Gaertn. (Gartner 1806) nec Sabine (1824), nec Desf. (Desfontaines 1829). — *Acras brasiliensis* Willd. ex Steud., *Nomencl. Bot. [Steudel]*, ed. 2, 1: 15 (Steudel 1840), nom. illeg. superfl. (based on *Chrysophyllum macrophyllum*).

Chrysophyllum brasiliense var. *minus* Miq., *Fl. Bras. [Martius]* 7: 103 [15 Jan. 1863] (Miquel 1863).

Sarcaulus macrophyllus Radlk., *Sitzungsber. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. München* 12: 311 (Radlkofer 1882).

VERNACULAR NAMES. — Pa: ukum-kamwi, uu-kamwi-wahyō • Wp: wai • Br: ajará, guajara-do-igapó, jaraí.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier 2338*.

INVENTORY DATA (FG). — 8 trees in 7 plots; $F_{\max} < 1\%$; dbh_{inv} = 31.5 cm.

Family SIMAROUBACEAE DC.
Genus *Homalolepis* Turcz.

[1707] *Homalolepis cedron* (Planch.) Devecchi & Pirani

Phytotaxa 366 (1): 67 [epubl. 28 Aug. 2018] (Devecchi & Pirani 2018). — *Simaba cedron* Planch., *London J. Bot.* 5: 566 (Planchon 1846). — *Quassia cedron* (Planch.) Baill., *Hist. Pl. [Baillon]* 4: 406 [Jul.-Sep. 1873] (Baillon 1873). — *Aruba cedron* (Planch.) Kuntze, *Revis. Gen. Pl.* 1: 103 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: timaa-avain-purubumna • Nt: wan ede • Cr: wan-édé • Br: manga-de-anta, pau-para-tudo.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2255*.

INVENTORY DATA (FG). — 485 trees in 77 plots; $F_{\max} = 6.9\%$; dbh_{inv} = 29.8 cm.

[1708] *Homalolepis morettii* (Feuillet) Devecchi & Pirani
(Fig. 56A)

Phytotaxa 366 (1): 43 [epubl. 28 Aug. 2018] (Devecchi & Pirani 2018). — *Simaba morettii* Feuillet, *Candollea* 38 (2): 746 (Feuillet 1983).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Nt: busi kananbuli • Cr: koubari-savann, mal-simarouba.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *C. Moretti 1027* (holo-, P, not seen; iso-, CAY[CAY024960, CAY024961, CAY024962], P[P01817225]).

INVENTORY DATA (FG). — 77 trees in 50 plots; $F_{\max} = 1\%$; dbh_{inv} = 86.3 cm.

Genus *Picrolemma* Hook.f.

[1709] *Picrolemma sprucei* Hook.f.

Gen. Pl. [Bentham & Hooker f.] 1 (1): 312 [7 Aug. 1862] (Hooker 1862).

Picrolemma pseudocoffea Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 196 (Ducke 1925).

VERNACULAR NAMES. — Pa: tuu-ariut, tuu-kamwi • Cr: kafé-lane • Br: caferana.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M.-F. Prévost et al. 4440*.

SIZE. — Up to 10 m tall (Thomas & Franceschinelli 2005).

Genus *Quassia* L.

[1710] *Quassia amara* L.

Sp. Pl., ed. 2, 1: 553 [Sep. 1762] (Linnaeus 1762).

Quassia amara var. *paniculata* Engl., *Fl. Bras. [Martius]* 12 (2): 207 [1 Sep. 1874] (Engler 1874). — *Quassia amara* f. *paniculata* (Engl.) Cronquist, *Brittonia* 5: 146 (Cronquist 1944).

NOTE. — A species native to Central America and the Caraibes, probably introduced to French Guiana during the 18th century by French Jesuit priests (Odonne et al. 2020).

VERNACULAR NAMES. — Pa: kwašvan • Ka: ki'elepun, kuasi bita, kuwasi, pekui • Nt: bita kwasi, kwasi • Cr: bwa-kwachi, kwachi • Fr: couachi • Br: quina, quina-de-Caiena.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *P. Grenand 2118*.

SIZE. — Costa Rica. *K. Martínez Sequeira 280* (MO), 9 m × 10 cm.

Genus *Simaba* Aubl.

[1711] *Simaba guianensis* Aubl.

Hist. Pl. Guiane 1: 400 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Zwingera amara* Willd., *Sp. Pl.*, ed. 4 2 (1): 569 [Mar. 1799] (Willdenow 1799), *nom. illeg. superfl.* (based on *Simaba guianensis*). — *Quassia crocea* Vahl, *Eclog. Amer.* 3: 12 (Vahl 1807), *nom. illeg. superfl.* (based on *Simaba guianensis*). — *Quassia guianensis* (Aubl.) D.Dietr., *Syn. Pl. [D. Dietrich]* 2: 1416 [1-20 Dec. 1840] (Dietrich 1840).

Aruba guianensis Aubl., *Hist. Pl. Guiane* 1: 294 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Simaba guianensis* (Aubl.) Engl., *Fl. Bras. [Martius]* 12 (2): 211 [1 Sep. 1874] (Engler 1874), *nom. illeg. hom.*, non Aubl. (Aublet 1775).

Zwingera aruba Spreng., *Syst. Veg. [Sprengel]* 2: 319 [Jan.-May 1825] (Sprengel 1825). — *Quassia aruba* (Spreng.) D.Dietr., *Syn. Pl. [D. Dietrich]* 2: 1416 [1-20 Dec. 1840] (Dietrich 1840).

Simaba cuspidata Spruce ex Engl., *Fl. Bras. [Martius]* 12 (2): 212 [1 Sep. 1874] (Engler 1874). — *Quassia cuspidata* (Spruce ex Engl.) Noot., *Blumea* 11: 522 [15 Dec. 1962] (Nooteboom 1962).

NOTE. — *Zwingera amara* Willd. is based on *Simaba guianensis* Aubl., not on *Simarouba amara* Aubl.

VERNACULAR NAMES. — Wp: wasaku sili • Br: cajuirana, maruparana.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lectotype of *Aruba guianensis* Aubl. [P-JJR, P00778459] designated by Lanjouw & Uittien [1940: 148]).

INVENTORY DATA (FG). — 17 trees in 12 plots; $F_{\max} < 1\%$; dbh_{inv} = 36.1 cm.

[1712] *Simaba orinocensis* Kunth

Nova genera et species plantarum [H.B.K.] 6: 18 [14 Apr. 1823] (Kunth 1823). — *Zwingera orinocensis* (Kunth) Spreng., *Syst. Veg. [Sprengel]* 2: 319 [Jan.-May 1825] (Sprengel 1825). — *Quassia orinocensis* (Kunth) D.Dietr., *Syn. Pl. [D. Dietrich]* 2: 1416 [1-20 Dec. 1840] (Dietrich 1840). — *Aruba orinocensis* (Kunth) Kuntze, *Revis. Gen. Pl.* 1: 103 [5 Nov. 1891] (Kuntze 1891).

Simaba multiflora A.Juss., *Mém. Mus. Hist. Nat.* 12: t. 27, f. 45a (Jussieu 1825). — *Quassia multiflora* (A.Juss.) Noot., *Blumea* 11: 522 [15 Dec. 1962] (Nooteboom 1962).

Simaba foetida Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 370 (Bentham 1851).

Simaba angustifolia Benth., *Hooker's J. Bot. Kew Gard. Misc.* 3: 371 (Bentham 1851).

Simaba crustacea Engl., *Fl. Bras. [Martius]* 12 (2): 211 [1 Sep. 1874] (Engler 1874). — *Aruba crustacea* (Engl.) Kuntze, *Revis. Gen. Pl.* 1: 103 [5 Nov. 1891] (Kuntze 1891). — *Quassia crustacea* (Engl.) Noot., *Blumea* 11: 522 [15 Dec. 1962] (Nooteboom 1962).

Simaba alata Maguire, *Bull. Torrey Bot. Club* 75 (4): 397 [Jul.-Aug. 1948] (Maguire 1948).

VERNACULAR NAMES. — Pa: kwepan, kwepan-wašiunu • Ka: kuyali, pakoweni • Wp: paku akayu, sokolo • Wn: kanapali • Nt: kananbuli • Cr: kananbouli • Br: cajurana, papacuaió.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *P. Grenand 1291*.

SIZE. — Up to 20 cm dbh (Cavalcante 1983).

[1713] *Simaba polyphylla* (Cavalcante) W.W.Thomas

Acta Amazonica 15 (1-2, Suppl.): 76 ["1985" publ. 1987] (Thomas 1987). — *Simaba guianensis* subsp. *polyphylla* Cavalcante, *Publ. Avulsas Mus. Paraense Emílio Goeldi* 37: 40 (Cavalcante 1983).

VERNACULAR NAMES. — Pa: timaa-avain • Br: marupazinho.

HERBARIUM DATA (FG). — 19 collections at CAY. Sel. exs.: *J.-F. Molino & D. Sabatier 2788*.

INVENTORY DATA (FG). — 57 trees in 40 plots; $F_{\max} < 1\%$; $dbh_{inv} = 41.4$ cm.

Genus *Simarouba* Aubl.

[1714] *Simarouba amara* Aubl.

Hist. Pl. Guiane 2: 860 [Jun.-Dec. 1775] (Aublet 1775).

Quassia dioica P.J.Bergius, *Mater. Med.*: 355 (Bergius 1778).

Quassia simaruba L.f., *Suppl. Pl.*: 234 ["1781" publ. Apr. 1782] (Linnaeus 1782).

Quassia officinalis Rich., *Actes Soc. Hist. Nat. Paris* 1: 108 [Oct. 1792] (Richard 1792).

Simarouba glauca DC., *Ann. Mus. Natl. Hist. Nat.* 17: 424 (Candolle 1811), "Simaruba". — *Quassia glauca* (DC.) Spreng., *Syst. Veg. [Sprengel]* 2: 319 [Jan.-May 1825] (Sprengel 1825).

Quassia alatifolia Stokes, *Bot. Mat. Med.* 2: 491 (Stokes 1812).

Simarouba amara var. *opaca* Engl., *Nat. Pflanzenfam. [Engler & Prantl]* 3 (4): 213 (Engler 1896). — *Simarouba opaca* (Engl.) Radlk. ex F.Boas, *Beih. Bot. Centralbl.* 29 (1): 332 (Boas 1913).

Simarouba amara var. *typica* Cronquist, *Bull. Torrey Bot. Club* 71 (3): 229 [May 1944] (Cronquist 1944), nom. inval. (Turland et al. 2018: Art. 24.3).

Simarouba glauca var. *latifolia* Cronquist, *Bull. Torrey Bot. Club* 71 (3): 231 [May 1944] (Cronquist 1944).

Simarouba glauca var. *typica* Cronquist, *Bull. Torrey Bot. Club* 71 (3): 231 [May 1944] (Cronquist 1944), nom. inval. (Turland et al. 2018): Art. 24.3).

Simarouba amara var. *puberula* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 8 (32): 466 [June 1952] (Cuatrecasas 1952).

NOTE. — Hyperdominant in Amazonia (ter Steege et al. 2020).

VERNACULAR NAMES. — Pa: timaa-avain • Ka: šimaluba, simalupa, sumaliba • Wp: iwé'i • Wn: pëünpë • Nt: sumaypa • Cr: akajou-blán, simarouba • Fr: simarouba • Br: marupa, paraiba.

HERBARIUM DATA (FG). — 61 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material not traced); *R.A.A. Oldeman B-1925* (P).

INVENTORY DATA (FG). — 77 trees in 62 plots; $F_{\max} < 1\%$; $dbh_{inv} = 90$ cm.

Family SIPARUNACEAE Schodde
Genus *Siparuna* Aubl.

[1715] *Siparuna cristata* (Poepp. & Endl.) A.DC.

Prodr. [A. P. de Candolle] 16 (2): 655 [mid July 1868] (Candolle 1868). — *Citrosma cristata* Poepp. & Endl., *Nova genera ac species plantarum [Poepig & Endlicher]* 2: 47 [Jan.-Sep. 1838] (Poepig & Endlicher 1838), "cristatum".

Siparuna cristata var. *macrophylla* A.DC., *Prodr. [A. P. de Candolle]* 16 (2): 655 [mid July 1868] (Candolle 1868).

Siparuna cristata var. *petiolaris* A.DC., *Prodr. [A. P. de Candolle]* 16 (2): 655 [mid July 1868] (Candolle 1868).

Siparuna williamsii J.F.Macbr., *Candollea* 5: 353 (Macbride 1934).

Siparuna monogyna Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 7 (1): 133 (Jangoux 1991).

Siparuna obconica Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 7 (1): 135 (Jangoux 1991).

VERNACULAR NAMES. — Pa: yahiwemna-kamwi, yariwapna-kamwi • Wp: mila suku, wila suku.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *J. Martin s.n.* (holotype of *Siparuna cristata* var. *petiolaris*: P[P016912]).

INVENTORY DATA (FG). — 31 trees in 13 plots; $F_{\max} = 1.3\%$; $dbh_{inv} = 19.4$ cm.

[1716] *Siparuna cuspidata* (Tul.) A.DC.

Prodr. [A. P. de Candolle] 16 (2.2): 655 [mid July 1868] (Candolle 1868). — *Citrosma cuspidata* Tul., *Ann. Sci. Nat., Bot. sér. 4, 3*: 40 [May 1855] (Tulasne 1855), “*Citriosma*”.

Citrosma myristicoidea Spruce, *J. Proc. Linn. Soc., Bot. 5*: 5 [“1861” publ. June 1860] (Spruce 1860). — *Siparuna crassiflora* Perkins, *Bot. Jahrb. Syst. 28 (5)*: 702 [11 Jan. 1901] (Perkins 1901), *nom. illeg. superfl.* (based on the type of *Citrosma myristicoidea*).

Siparuna argyrochrysea Perkins, *Bot. Jahrb. Syst. 28 (5)*: 702 [11 Jan. 1901] (Perkins 1901).

Siparuna lepidiflora Perkins, *Notizbl. Bot. Gart. Berlin-Dahlem 10*: 166 [20 Dec. 1927] (Perkins 1927).

Siparuna ucayaliensis Perkins, *Notizbl. Bot. Gart. Berlin-Dahlem 10*: 167 [20 Dec. 1927] (Perkins 1927).

HERBARIUM DATA (FG). — 80 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (original material of *Siparuna argyrochrysea*: holo-, B (not seen, photo F neg 13478, fragm. GH[GH00046549]); possible iso-, P[P00053272, P00053273, P00053274]).

INVENTORY DATA (FG). — 36 trees in 19 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 18.8$ cm.

[1717] *Siparuna decipiens* (Tul.) A.DC.

Prodr. [A. P. de Candolle] 16 (2): 643 [mid July 1868] (Candolle 1868). — *Citrosma decipiens* Tul., *Arch. Mus. Hist. Nat. 8*: 368 (Tulasne 1855), “*Citriosma*”.

Conuleum guyannense Rich. ex A.Rich., *Mém. Soc. Hist. Nat. Paris, sér. 2, 1*: 391 (Richard 1823).

Siparuna lepidantha Perkins, *Bot. Jahrb. Syst. 28 (5)*: 701 [11 Jan. 1901] (Perkins 1901).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: á-kamwi, kwatri-purubumna • Ka: piyawan wewe • Wp: mila suku, wila suku • Wn: awalepokan • Nt: faya pao • Br: amarelinho, quariquara-branca.

HERBARIUM DATA (FG). — 122 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material K[K000587966], MEL[MEL2384184], P[P016899, P00053052, P00053053], S[S-R-7098]).

INVENTORY DATA (FG). — 329 trees in 125 plots; $F_{\max} = 3.3\%$; $dbh_{\text{inv}} = 73.5$ cm.

[1718] *Siparuna guianensis* Aubl.
(Fig. 56B)

Hist. Pl. Guiane 2: 865 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Citrosma guianensis* (Aubl.) Tul., *Arch. Mus. Hist. Nat. 8*: 361 (Tulasne 1855), “*Citriosma*”.

Citrosma discolor Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher] 2*: 47 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838). — *Siparuna discolor* (Poepp. & Endl.) A.DC., *Prodr. [A. P. de Candolle] 16 (2.2)*: 656 [mid July 1868] (Candolle 1868).

Citrosma camporum Tul., *Ann. Sci. Nat., Bot. sér. 4, 3*: 39 [May 1855] (Tulasne 1855), “*Citriosma*”. — *Siparuna camporum* (Tul.)

A.DC., *Prodr. [A. P. de Candolle] 16 (2.2)*: 654 [mid July 1868] (Candolle 1868).

Citrosma guianensis var. *divergentifolia* Tul., *Arch. Mus. Hist. Nat. 8*: 362 (Tulasne 1855), “*Citriosma*”. — *Siparuna guianensis* var. *divergentifolia* (Tul.) A.DC., *Prodr. [A. P. de Candolle] 16 (2.2)*: 654 [mid July 1868] (Candolle 1868).

Citrosma guianensis var. *nuda* Tul., *Arch. Mus. Hist. Nat. 8*: 362 (Tulasne 1855), “*Citriosma*”.

Siparuna panamensis A.DC., *J. Bot. 3*: 219 (Candolle 1865).

Siparuna guianensis var. *glabrescens* A.DC., *Prodr. [A. P. de Candolle] 16 (2.2)*: 654 [mid July 1868] (Candolle 1868), *nom. illeg. superfl.* (one of the syntypes is also a syntype of *S. guianensis* var. *nuda*).

Siparuna guianensis var. *longifolia* A.DC., *Prodr. [A. P. de Candolle] 16 (2.2)*: 654 [mid July 1868] (Candolle 1868).

Siparuna foetida Barb.Rodr., *Vellozia*, ed. 2, 1: 68 (Barbosa Rodrigues 1891).

Siparuna guianensis var. *nitens* Kuntze, *Revis. Gen. Pl. 3 (3)*: 276 [28 Sep. 1898] (Kuntze 1898).

Siparuna archeri A.C.Sm., *Bull. Torrey Bot. Club 59 (9)*: 517 [Dec. 1932] (Smith 1932).

Siparuna arianeae V.Pereira, *Bradea 4* (36): 291 (Pereira 1986).

Siparuna cavalcantei Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 7 (1)*: 120 (Jangoux 1991).

Siparuna duckeana Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 7 (1)*: 125 (Jangoux 1991).

Siparuna itacaiunensis Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 7 (1)*: 129 (Jangoux 1991).

Siparuna savanicola Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot. 7 (1)*: 144 (Jangoux 1991).

VERNACULAR NAMES. — Pa: yahiwemna, yariwapna • Ka: idyakopi, ilakopi, yalakopi • Te: dzawat móböhá • Wp: enēmi'o, nēmi'o, wainimi'o • Wn: jaléké, jawiléké • Nt: kapasi wiwii • Cr: vénéré, viniré • Br: caá-pitiú.

HERBARIUM DATA (FG). — 140 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000611207]).

INVENTORY DATA (FG). — 1 tree, $dbh = 25$ cm.

[1719] *Siparuna pachyantha* A.C.Sm.

J. Arnold Arbor. 20 (3): 293 [19 July 1939] (Smith 1939).

Siparuna emarginata R.S.Cowan, *Brittonia 8 (4)*: 237 [Jan. 1957] (Cowan 1957).

VERNACULAR NAMES. — Pa: avakni-avak, avatni-awak • Wp: a'i makule • Cr: féy-tabak • Br: caá-pitiú-de-folha-grande.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5552*.

INVENTORY DATA (FG). — 24 trees in 16 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 70$ cm.

[1720] *Siparuna poeppigii* (Tul.) A.DC.

Prodr. [A. P. de Candolle] 16 (2): 653 [mid July 1868] (Candolle 1868). — *Citrosma poeppigii* Tul., *Ann. Sci. Nat., Bot. sér. 4*, 3: 39 [May 1855] (Tulasne 1855), “*Citriosma*”.

Tetratome lepidota Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher]* 2: 47 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838).

Siparuna sprucei A.DC., *J. Bot.* 3: 219 (Candolle 1865).

Siparuna amazonica Mart. ex A.DC., *Prodr. [A. P. de Candolle]* 16 (2.2): 653 [mid July 1868] (Candolle 1868).

Siparuna espinhacensis Jangoux, *Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.* 7 (1): 127 (Jangoux 1991).

VERNACULAR NAMES. — Pa: yahiemna, yariwpana • Wp: enēmi'o, nēmi'o, wainimi'o • Cr: vénéré, viniré • Br: caá-pitiú.

HERBARIUM DATA (FG). — 71 collections at CAY. Sel. exs.: *R.A.A. Oldeman 1916*.

SIZE. — Up to 20 cm dbh (Renner & Hausner 2005).

Family SOLANACEAE Juss.
Genus *Solanum* L.

[1721] *Solanum crinitum* Lam.

Tabl. Encycl. 2[4 (1)]: 20 [21 Nov. 1796] (Lamarck 1796).

Solanum formosum Humb. & Bonpl. ex Dunal, *Solan. Syn.*: 35 [Oct. 1816] (Dunal 1816).

Solanum jubatum Willd., *Syst. Veg. [Roemer & Schultes]* 4: 669 [Mar.-June 1819] (Willdenow 1819). — *Solanum cyananthum* var. *jubatum* (Willd.) Dunal, *Prodr. [A. P. de Candolle]* 13 (1): 316 [10 May 1852] (Dunal 1852).

Solanum cyananthum Dunal, *Prodr. [A. P. de Candolle]* 13 (1): 316 [10 May 1852] (Dunal 1852).

NOTE. — *Solanum jubatum* Willd. is to be ascribed to Willdenow alone (Turland et al. 2018: Art. 46.3, Ex. 15; see also IPNI, 2020 and Taylor & Gereau, 2019).

VERNACULAR NAMES. — Pa: puduku • Ka: kwasisi, kwašisi, pal-emulu, poto:no palemulu • Wp: yū ásiši • Nt: mantuga maka • Cr: ponm-sousouri, tonmat-sousouri • Br: jurubeba.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: *P. Grenand 15*.

SIZE. — Colombia, Nariño. *M.I. Barreto 141* (MO), 14 m × 30 cm.

[1722] *Solanum endopogon* (Bitter) Bohs
subsp. *guianense* (Bohs) Bohs

J. Bot. Res. Inst. Texas 3 (2): 509 [25 Nov. 2009] (Bohs 2009). — *Cyphomandra endopogon* Bitter subsp. *guianensis* Bohs, *Fl. Neotrop. Monogr.* 63: 90 [5 July 1994] (Bohs 1994).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: awala kāsī, uluwu kāsī • Cr: mavévé-chien • Br: pau-mirim.

HERBARIUM DATA (FG). — 25 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 14709* (holo-, F[V0072836F]; iso-, CAY[CAY070876], NY[00721521]); *S.A. Mori et al. 15028*, 15 m × 20 cm.

[1723] *Solanum splendens* (Dunal) Bohs
(Fig. 56C)

Monogr. Syst. Bot. Missouri Bot. Gard. 131: 320 (Bohs 2015). — *Cyphomandra splendens* Dunal, *Prodr. [A. P. de Candolle]* 13 (1): 395 (Dunal 1852).

Pionandra hartwegii Miers, *London J. Bot.* 4: 363 (Miers 1845). — *Cyphomandra hartwegii* (Miers) Sendtn. ex Walp., *Repert. Bot. Syst. [Walpers]* 6 (4): 579 [19-22 May 1847] (Walpers 1847).

Cyphomandra artocarpophylloides H.J.P.Winkl., *Repert. Spec. Nov. Regni Veg.* 7: 246 (Winkler 1909).

Cyphomandra mollicella Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 322 [24 Oct. 1929] (Standley 1929).

Solanum circinatum Bohs, *Taxon* 44 (4): 585 [Nov. 1995] (Bohs 1995).

VERNACULAR NAMES. — Pa: isuu-aveya, isuu-vey • Wp: awala kāsī, uluwu kāsī • Cr: mavévé-chien.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *M.-F. Prévost & C. Feuillet 3964*.

SIZE. — Up to 12 m tall (Bohs 1994).

Family STEMONURACEAE Kårehed
Genus *Discophora* Miers

[1724] *Discophora guianensis* Miers
(Fig. 56D)

Ann. Mag. Nat. Hist., ser. 2, 10 (56): 119 [Aug. 1852] (Miers 1852). — *Kümmereria brasiliensis* Mart. ex Engl., *Fl. Bras. [Martius]* 12 (2): 52 [1 Dec. 1872] (Engler 1872), nom. illeg. superfl. (based on *Discophora guianensis*).

Lasianthera amazonica Barb.Rodr., *Vellozia*, ed. 2, 1: 12 (Barbosa Rodrigues 1891).

Discophora panamensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 4 (8)*: 222 [24 Oct. 1929] (Standley 1929).

Discophora froesii Pires, *Bol. Técn. Inst. Agron. N.* 38: 28 (Pires 1960).

VERNACULAR NAMES. — Pa: kanegma • Wp: inámu sī, inámušī wila, mulei sī, yawasi pita • Nt: boni udu • Cr: bwa-léchél • Br: caferana, ivaí, pombinho.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *B. Riéra & D. Sabatier 1808*.

INVENTORY DATA (FG). — 37 trees in 25 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 15.5$ cm.

Family STYRACACEAE DC. & Spreng.
Genus *Styrax* L.

[1725] *Styrax glabratus* Schott

Syst. Veg. [Sprengel] 4 (2): 406 [Jan.-June 1827] (Schott 1827). — *Strigilia glabrata* (Schott) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 283 [Apr. 1859] (Miers 1859).

Epigenia integerrima Vell., *Fl. Flumin.*: 183 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).

Styrax erymophyllus Pohl, *Pl. Bras. Icon. Descr.* 2 (2): 57 [Sep. 1830] (Pohl 1830).

Styrax psilophyllus A.DC., *Prodri. [A. P. de Candolle]* 8: 266 [mid Mar. 1844] (Candolle 1844), "psilophyllum". — *Strigilia psilophylla* (A.DC.) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 283 [Apr. 1859] (Miers 1859).

Styrax leiophyllus Miers, *Veg. Kingd.*, ed. 3: 593a (Miers 1853), "leiophyllum". — *Strigilia leiophylla* (Miers) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 283 [Apr. 1859] (Miers 1859).

Styrax lauraceus Perkins, *Bot. Jahrb. Syst.* 31 (3): 478 [10 Dec. 1901] (Perkins 1901).

Styrax squamulosus M.F.Silva, *Acta Amazonica* 1 (1): 23 (Silva 1971), "squamulosa".

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *Service Forestier 7445*.

INVENTORY DATA (FG). — 1 tree, dbh = 15 cm.

[1726] *Styrax guyanensis* A.DC.

Prodri. [A. P. de Candolle] 8: 261 [mid Mar. 1844] (Candolle 1844), "Guyanense". — *Strigilia guyanensis* (A.DC.) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 282 [Apr. 1859] (Miers 1859), "Guayanensis".

Styrax guyanensis var. *japurensis* Seub., *Fl. Bras. [Martius]* 7: 188 [1 Aug. 1868] (Seubert 1868), "guianense", "japurense".

Styrax tessmannii Perkins, *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 459 [1 Dec. 1928] (Perkins 1928).

Styrax longifolius Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 22 (3): 166 (Standley 1940).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 962*.

INVENTORY DATA (FG). — 3 trees in 3 plots; F_{max} < 1%; dbh_{inv} = 17.8 cm.

[1727] *Styrax macrophyllus* Schott ex Pohl

Pl. Bras. Icon. Descr. 2 (2): 61 [Sep. 1830] (Pohl 1830). — *Strigilia macrophylla* (Schott ex Pohl) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 283 [Apr. 1859] (Miers 1859).

Styrax heteroclitus J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (5.1): 230 (Macbride 1959).

VERNACULAR NAMES. — Pa: suwimba-ahavukune.

HERBARIUM DATA (FG). — a single specimen, *P. Grenand 1933*.

SIZE. — Suriname. *R.S. Cowan & T.R. Soderstrom 1973* (MO), dbh 10 cm.

[1728] *Styrax pallidus* A.DC.
(Fig. 57A)

Prodri. [A. P. de Candolle] 8: 261 [mid Mar. 1844] (Candolle 1844), "pallidum". — *Strigilia pallida* (A.DC.) Miers, *Ann. Mag. Nat. Hist.*, ser. 3, 3 (16): 282 [Apr. 1859] (Miers 1859).

VERNACULAR NAMES. — Wp: suluku'a.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *F.M.R. Leprieur 272* (holo-, G-DC[G00142381]; iso-, B[not seen, photo neg. F[F0BN00431], US[00112566]]; possible iso-, F[V0076078F, V0076079F]).

INVENTORY DATA (FG). — 10 trees in 9 plots; F_{max} < 1%; dbh_{inv} = 35 cm.

[1729] *Styrax sieberi* Perkins

Pflanzenr. [Engler] IV.241 (Heft 30): 44 [3 Sep. 1907] (Perkins 1907).

Styrax burchellii Perkins, *Pflanzenr. [Engler]* IV.241 (Heft 30): 46 [3 Sep. 1907] (Perkins 1907).

Styrax pearcei Perkins, *Pflanzenr. [Engler]* IV.241 (Heft 30): 46 [3 Sep. 1907] (Perkins 1907).

Styrax pearcei var. *bolivianus* Perkins, *Pflanzenr. [Engler]* IV.241 (Heft 30): 46 [3 Sep. 1907] (Perkins 1907).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *S.A. Mori et al. 22781*.

INVENTORY DATA (FG). — 2 trees in 2 plots; F_{max} < 1%; dbh_{inv} = 22 cm.

Family SYMPLOCACEAE Desf.
Genus *Symplocos* Jacq.

[1730] *Symplocos guianensis* (Aubl.) Gürke

Nat. Pflanzenfam. [Engler & Prantl] 4 (1): 172 (Gürke 1891), "guyanensis". — *Ciponima guianensis* Aubl., *Hist. Pl. Guiane* 1: 566 [Jun.-Dec. 1775] (Aublet 1775). — *Symplocos ciponima* L'Hér., *Trans. Linn. Soc. London* 1: 175 [13 Aug. 1791] (L'Héritier 1791), nom. illeg. superfl. (based on *Ciponima guianensis*). — *Ciponima scabridula* Miers, *J. Linn. Soc., Bot.* 17: 288 ["1880" publ. May 1879] (Miers 1879). — *Eugeniodes guianense* (Aubl.) Kuntze, *Revis. Gen. Pl.* 2: 975 [5 Nov. 1891] (Kuntze 1891).

Symplocos paraensis Endl. ex Miq., *Fl. Bras. [Martius]* 7: 24 [15 Mar. 1856] (Miquel 1856), "paraënsis", nom. nud. pro syn. — *Symplocos guianensis* var. *paraensis* Brand, *Pflanzenr. [Engler]* IV.242 (Heft 6): 84 [3 Sep. 1907] (Brand 1901), "paraënsis".

Eugeniodes paraense Kuntze, *Revis. Gen. Pl.* 2: 975 [5 Nov. 1891] (Kuntze 1891).

VERNACULAR NAMES. — Pa: waravru-puvemna • Ka: alilime'i.

HERBARIUM DATA (FG). — 40 collections at CAY. Sel. exs.: *Y. Caraglio 802*.

SIZE. — Brazil, Pará. *T.C. Plowman et al. 8668* (MO), 8 m × 10 cm.

[1731] *Symplocos martinicensis* Jacq.
(Fig. 57B)

Enum. Syst. Pl. 5 [Aug.-Sep. 1760] (Jacquin 1760). — *Eugeniodes martinicense* (Jacq.) Kuntze, *Revis. Gen. Pl.* 2: 975 [5 Nov. 1891] (Kuntze 1891).

Eugeniodes martinicense var. *paraguayense* Kuntze, *Revis. Gen. Pl.* 3 (2): 196 [28 Sep. 1898] (Kuntze 1898).

VERNACULAR NAMES. — Wp: mulei, mulei pilá, mulei sówí • Br: pau-ferro.

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier 1051*.

INVENTORY DATA (FG). — 17 trees in 12 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 29.9$ cm.

Family THEACEAE Mirb.
Genus *Gordonia* J.Ellis

[1732] *Gordonia fruticosa* (Schrad.) H.Keng
(Fig. 57C)

Gard. Bull. Singapore 33 (2): 310 [1 Jan. 1980] (Keng 1980). — *Wikstroemia fruticosa* Schrad., *Gött. Gel. Anz.* 2: 711 [5 May 1821] (Schrader 1821), “*fruticosa*”. — *Lindleya semiserrata* Nees, *Flora* 4 (1): 328 [21 May 1821] (Nees 1821), *nom. illeg. superfl.* (based on *Wikstroemia fruticosa*). — *Haemocharis semiserrata* Mart., *Nova genera et species plantarum [Martius]* 1 (4): 107 [“1824” publ. Jan.-Mar. 1826] (Martius 1826), *nom. illeg. superfl.* (based on *Wikstroemia fruticosa*). — *Gordonia semiserrata* Spreng., *Syst. Veg. [Sprengel]* 4 (2, Cur. Post.): 260, 408 [Jan.-June 1827] (Sprengel 1827), *nom. illeg. superfl.* (based on *Wikstroemia fruticosa* and *Haemocharis semiserrata*). — *Laplacea semiserrata* Cambess., *Fl. Bras. Merid. [A. St.-Hil.]* (quarto ed.) 1 (8): 300 [28 June 1828] (Cambessèdes 1828), *nom. illeg. superfl.* (based on *Haemocharis semiserrata*, thus indirectly on *Wikstroemia fruticosa*). — *Laplacea fruticosa* (Schrad.) Kobuski, *J. Arnold Arbor.* 28 (4): 437 [15 Oct. 1947] (Kobuski 1947).

Laplacea speciosa Kunth, *Nova genera et species plantarum [H.B.K.]* 5: 209 [25 Feb. 1822] (Kunth 1822).

Laplacea parviflora Spix & Mart., *Reise Bras. [Spix & Mart.]* 1: 207 [Nov. 1823] (Spix & Martius 1823), *nom. nud.*

Laplacea praemorsa Splitg., *Tijdschr. Nat. Geschied. Physiol.* 9: 100 [Aug.-Sep. 1842] (Splitgerber 1842). — *Haemocharis praemorsa* (Splitg.) Kuntze, *Revis. Gen. Pl.* 1: 62 [5 Nov. 1891] (Kuntze 1891).

Laplacea intermedia Benth., *Pl. Hartw. [Bentham]* 126 [late Dec. 1843-early Jan. 1844] (Bentham 1843). — *Haemocharis intermedia* (Benth.) Choisy, *Mém. Soc. Phys. Genève* 14 (1): 144 (Choisy 1855). — *Wikstroemia intermedia* (Benth.) S.F.Blake, *Contr. Gray Herb.* 53: 40 [26 Feb. 1918] (Blake 1918). — *Laplacea speciosa* var. *intermedia* (Benth.) Kobuski, *J. Arnold Arbor.* 31 (4): 421 [15 Oct. 1950] (Kobuski 1950).

Laplacea quinoderma Wedd., *Hist. Nat. Quinquinas* 33 (Weddell 1849).

Laplacea camellioides Sond., *Linnaea* 22: 549 (Sonder 1849). — *Haemocharis camellioides* (Sond.) Kuntze, *Revis. Gen. Pl.* 1: 62 [5 Nov. 1891] (Kuntze 1891), “*camellioides*”.

Haemocharis parviflora Choisy, *Mém. Soc. Phys. Genève* 14 (1): 144 (Choisy 1855). — *Wikstroemia parviflora* (Choisy) S.F.Blake, *Contr. Gray Herb.* 53: 40 [26 Feb. 1918] (Blake 1918).

Laplacea pubescens Planch. & Linden ex Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 18: 269 (Triana & Planchon 1862). — *Haemocharis pubescens* (Planch. & Linden ex Triana & Planch.) Hieron., *Bot. Jahrb. Syst.* 20 (3, Beibl. 49): 48 [9 Apr. 1895] (Hieronymus 1895).

Laplacea symplocoidea Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 18: 269 (Triana & Planchon 1862). — *Haemocharis symplocoidea* (Triana & Planch.) Kuntze, *Revis. Gen. Pl.* 1: 62 [5 Nov. 1891] (Kuntze 1891), “*symplocoidea*”. — *Wikstroemia symplocoidea* (Triana & Planch.) S.F.Blake, *Contr. Gray Herb.* 53: 41 [26 Feb. 1918] (Blake 1918). — *Laplacea fruticosa* var. *symplocoidea* (Triana & Planch.) Kobuski, *J. Arnold Arbor.* 31 (4): 414 [15 Oct. 1950] (Kobuski 1950).

Laplacea camelliifolia Triana & Planch., *Ann. Sci. Nat., Bot. sér. 4*, 18: 270 (Triana & Planchon 1862), “*camelliifolia*”. — *Wikstroemia camelliifolia* (Triana & Planch.) S.F.Blake, *Contr. Gray Herb.* 53: 39 [26 Feb. 1918] (Blake 1918), “*camelliifolia*”. — *Laplacea pubescens* var. *camelliifolia* (Triana & Planch.) Kobuski, *J. Arnold Arbor.* 31 (4): 427 [15 Oct. 1950] (Kobuski 1950), “*camelliifolia*”.

Haemocharis caracasana Linden & Planch., *Pl. Columb. [Linden]* 1: 59 [“1863” publ. 1874-1875] (Linden & Planchon 1874-1875).

Laplacea caracasana Klotzsch & H.Karst. ex Wawra, *Fl. Bras. [Martius]* 12 (1): 289 [1 Apr. 1886] (Wawra 1886), *nom. nud. pro syn.*

Laplacea pulcherrima Melch., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 21: 136 (Melchior 1925). — *Laplacea fruticosa* var. *pulcherrima* (Melch.) Kobuski, *J. Arnold Arbor.* 31 (4): 414 [15 Oct. 1950] (Kobuski 1950).

Laplacea raimondiana Melch., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 21: 136 (Melchior 1925).

Laplacea pubescens var. *subcaudata* Kobuski, *J. Arnold Arbor.* 31 (4): 427 [15 Oct. 1950] (Kobuski 1950).

Gordonia humboldtii H.Keng, *Gard. Bull. Singapore* 33 (2): 310 [1 Jan. 1980] (Keng 1980).

Gordonia planchonii H.Keng, *Gard. Bull. Singapore* 33 (2): 310 [1 Jan. 1980] (Keng 1980).

Laplacea fruticosa var. *chimantae* Steyermark., *Ann. Missouri Bot. Gard.* 74 (3): 650 (Steyermark 1987).

Laplacea pubescens var. *minor* Steyermark., *Ann. Missouri Bot. Gard.* 74 (3): 652 (Steyermark 1987).

VERNACULAR NAMES. — Wp: pasí’i wapo sili.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *V. Hequet 3003*.

SIZE. — Brazil, Pará. *A.S. Tavares 289* (MO), 30 m.

Family ULMACEAE Mirb.
Genus *Ampelocera* Klotzsch

[1733] *Ampelocera edentula* Kuhlm.
(Fig. 57D)

Anais Reunião Sul-Amer. Bot. 3: 75 ["1938" publ. 1940] (Kuhmann 1940).

Ampelocera latifolia Ducke, *Trop. Woods* 76: 15 (Ducke 1943).

VERNACULAR NAMES. — Ka: watoipyo, watoyati • Wp: inámu sī, inámusī wila • Br: cafezinho, paracanaúba.

HERBARIUM DATA (FG). — 18 collections at CAY. Sel. exs.: *S.A. Mori et al. 15115*.

INVENTORY DATA (FG). — 41 trees in 27 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 47.1$ cm.

Family URTICACEAE Juss.
Genus *Cecropia* Loefl.

[1734] *Cecropia distachya* Huber

Bol. Mus. Goeldi Hist. Nat. Ethnogr. 6: 65 (Huber 1910).

Cecropia riparia Warb. ex Snethl., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 363 [30 June 1915] (Snethlage 1923).

Cecropia richardii Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 9 (36/37): 336 [May 1956] (Cuatrecasas 1956).

HERBARIUM DATA (FG). — 14 collections at CAY. Sel. exs.: *D. Sabatier et al. 5805*.

INVENTORY DATA (FG). — 1 tree, $dbh = 23.6$ cm.

[1735] *Cecropia granvilleana* C.C.Berg
(Fig. 58A)

Bull. Mus. Natl. Hist. Nat., B, Adansonia 7 (3): 255 ["1985" publ. 1986] (Berg 1986).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 6000* (holo-, U[U0004467]; iso-, BR[BR0000005296460], CAY[CAY010617], G[G00438028], NY[00284199], P[P00077213]); *D. Sabatier & J.-F. Molino 5016*, $dbh = 27$ cm.

[1736] *Cecropia latiloba* Miq.

Fl. Bras. [Martius] 4 (1): 147 [1 Dec. 1853] (Miquel 1853). — *Ambaiba latiloba* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia paraensis Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 64 (Huber 1910).

Cecropia orinocensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 22 (1): 14 (Standley 1940).

VERNACULAR NAMES. — Te: kulekule ilebe, kulekule ilupiwat • Br: imbaúba-branca, imbaúba-da-várzea.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *M.-F. Prévost 1693*.

INVENTORY DATA (FG). — 7 trees in 1 plot; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 36$ cm.

[1737] *Cecropia obtusa* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 79 (Trécul 1847). — *Ambaiba obtusa* (Trécul) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891). — *Coilotapalus obtusa* (Trécul) M.Gómez, *Fl. Cuba [Gómez & Roig]* 12 (Gómez 1914).

Cecropia lisboana Snethl., *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 171 [30 Dec. 1924] (Snethlage 1924).

VERNACULAR NAMES. — Pa: tukuwi • Ka: tapilen salasala, tuleke, yalayala • Te: kule kule • Wp: ama'i • Wn: kulegli, kulekle • Nt: uman papay • Cr: bwa-kannon • Fr: bois canon • Br: embaúba, imbaúba.

HERBARIUM DATA (FG). — 82 collections at CAY. Sel. exs.: *F.M.R. Leprieur 195* (lecto-, P[P00089163], designated by Berg [Berg & Dewolf 1975: 288]; isolecto-, F[V0053465F], P[P00089164]).

INVENTORY DATA (FG). — 187 trees in 107 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 48.5$ cm.

[1738] *Cecropia palmata* Willd.

Spp. Pl., ed. 4 4 (2): 652 [Apr. 1806] (Willdenow 1806). — *Ambaiba palmata* (Willd.) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia bureauiiana V.A.Richt., *Biblioth. Bot.* 43: 19 (Richter 1897).

VERNACULAR NAMES. — Ka: yalayala • Br: imbaúba-vermelha, torém.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *E.M. Mélinon s.n.* (holotype of *Cecropia bureauiiana*: P[P00089172]; iso-, P[P00089173], P[P00089174], P[P00089175]).

SIZE. — Up to 20 m tall (Berg *et al.* 2005).

[1739] *Cecropia peltata* L.

Syst. Nat., ed. 10, 2: 1286 [7 June 1759] (Linnaeus 1759). — *Ambaiba peltata* (L.) Kuntze, *Revis. Gen. Pl.* 2: 623 [5 Nov. 1891] (Kuntze 1891).

Cecropia humboldtiana Klotzsch, *Linnaea* 20: 530 [Oct. 1847] (Klotzsch 1847). — *Ambaiba humboldtiana* (Klotzsch) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia schiedeana Klotzsch, *Linnaea* 20: 531 [Oct. 1847] (Klotzsch 1847). — *Ambaiba schiedeana* (Klotzsch) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia surinamensis Miq., *Fl. Bras. [Martius]* 4 (1): 143 [1 Dec. 1853] (Miquel 1853). — *Ambaiba surinamensis* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia digitata Ten. ex Miq. var. *grisea* Miq., *Fl. Bras. [Martius]* 4 (1): 149 [1 Dec. 1853] (Miquel 1853).



FIG. 56. — Simaroubaceae: **A**, *Homalolepis moretti* (Feuillet) Devecchi & Pirani (M.-F. Prévost & D. Sabatier 2987). Siparunaceae: **B**, *Siparuna guianensis* Aubl. (M.-F. Prévost 4517). Solanaceae: **C**, *Solanum splendens* (Dunal) Bohs (D. Sabatier 5852). Stemonuraceae: **D**, *Discophora guianensis* Miers (B. Ríera & D. Sabatier 1808). © D. Sabatier/IRD.

Cecropia propinqua Miq., *Fl. Bras. [Martius]* 4 (1): 149 [1 Dec. 1853] (Miquel 1853).

Cecropia scabrifolia V.A.Richt., *Biblioth. Bot.* 43: 15 (Richter 1897).

Cecropia dielsiana Snethl., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 362 [30 June 1915] (Snethlage 1923), “Dielsana”.

Cecropia arachnoidea Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 226 [22 Sep. 1917] (Pittier 1917).

Cecropia asperrima Pittier, *Contr. U.S. Natl. Herb.* 18 (6): 227 [22 Sep. 1917] (Pittier 1917).

Cecropia hondurensis Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 4 (8): 302 [24 Oct. 1929] (Standley 1929).

Cecropia goodspeedii Cuatrec., *Notas Fl. Colombia* 6: 41 [30 Mar. 1944] (Cuatrecasas 1944).

VERNACULAR NAMES. — Ka: salasala, tapilen salasala, tuleke • Cr: bwa-kannon • Br: embaúba, imbaúba.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *R.A.A. Oldeman T-737*.

SIZE. — Guyana. *M.J. Jansen-Jacobs et al. 2658* (P), 10 m.

[1740] *Cecropia sciadophylla* Mart.

Flora 24 (2, Beibl.): 93 (Martius 1841). — *Ambaiba sciadophylla* (Mart.) Kuntze, *Revis. Gen. Pl.* 2: 624 [5 Nov. 1891] (Kuntze 1891).

Cecropia juranyiana V.A.Richt., *Biblioth. Bot.* 43: 13 (Richter 1897). — *Cecropia sciadophylla* var. *juranyiana* (V.A.Richt.) Snethl., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 358 [15 Apr. 1923] (Snethlage 1923).

Cecropia sciadophylla var. *decurrans* Snethl., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 358 [15 Apr. 1923] (Snethlage 1923).

Cecropia sciadophylla var. *guamuesensis* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 6 (22/23): 299 [Jan.-Aug. 1945] (Cuatrecasas 1945).

Cecropia sciadophylla var. *pedroa* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 6 (22/23): 299 [Jan.-Aug. 1945] (Cuatrecasas 1945).

Cecropia sciadophylla var. *subsessilis* Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 6 (22/23): 299 [Jan.-Aug. 1945] (Cuatrecasas 1945).

Cecropia inchuensis Cuatrec., *Revista Acad. Colomb. Ci. Exact.* 9 (36/37): 329 [May 1956] (Cuatrecasas 1956).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: tukuwi-awaig • Ka: tamunen salasala, tuleke, yalayala tupulu • Te: kule kule tapelelapör • Wp: ama'i áta

• Wn: makamakalu • Nt: diya papay, man papay • Cr: mal-bwa-kannon • Fr: bois canon • Br: imbaúba-da-mata, imbaúba-vermelha.

HERBARIUM DATA (FG). — 43 collections at CAY. Sel. exs.: PA. Sagot 861, 1855 (holotype of *Cecropia juranyiana*: P[P00089178]; isotype, P[P00089179]).

INVENTORY DATA (FG). — 44 trees in 27 plots; $F_{\max} = 3.6\%$; $\text{dbh}_{\text{inv}} = 50.9\text{ cm}$.

[1741] *Cecropia silvae* C.C.Berg
(Fig. 58B)

Acta Bot. Neerl. 21: 695 (Berg 1972).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: D. Sabatier & J.-F. Molino 5137, dbh 30.6 cm.

Genus *Coussapoa* Aubl.

[1742] *Coussapoa angustifolia* Aubl.

Hist. Pl. Guiane 2: 956 [Jun.-Dec. 1775] (Aublet 1775).

NOTE. — Known only from the Guiana Shield. The original material at BM is a leaf base on the same sheet as the original material of *Coussapoa latifolia* Aubl. and a barcode is attributed to each sample. *C. angustifolia* (BM000993415) is erroneously referenced in JSTOR's Global Plants under the *C. latifolia* (BM000993414) barcode.

VERNACULAR NAMES. — Wp: kwapo'i.

HERBARIUM DATA (FG). — 41 collections at CAY. Sel. exs.: J.B. Aublet s.n. (original material BM[BM000993415]).

INVENTORY DATA (FG). — 18 trees in 4 plots; $F_{\max} = 3.6\%$; $\text{dbh}_{\text{inv}} = 30.6\text{ cm}$.

[1743] *Coussapoa asperifolia* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 96 (Trécul 1847).

Coussapoa ficina Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2):* 161 (Standley 1937).

Coussapoa cayennensis A.D.Hawkes, *Phytologia* 3 (1): 30 [30 Aug. 1948] (Hawkes 1948).

VERNACULAR NAMES. — Pa: sawu-anen, wiwis-kamwi • Ka: puluma • Wp: kalate to táláwáwá, kwapo'i kalate • Wn: okopiaphawin • Br: caimbérana.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: WE. Broadway 880 (holotype of *Coussapoa cayennensis*: NY[00284221]; iso-, GH[GH00034414], US[00090305]).

INVENTORY DATA (FG). — 1 tree, dbh = 17.2 cm.

[1744] *Coussapoa ferruginea* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 93 (Trécul 1847).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: F.M.R. Leprieur s.n. (holo-, P[P00756698]; iso-, F[V0053601F], G[G00438122], GH[GH00034421], L[L0039918], P[P00756699], P00756700], R[R000056202], U[U0004507]).

SIZE. — Ecuador, Zamora-Chinchipe. H. van der Werff et al. 19273 (MO), 25 m.

[1745] *Coussapoa latifolia* Aubl.

Hist. Pl. Guiane 2: 955 [Jun.-Dec. 1775] (Aublet 1775). — *Euosma latifolia* (Aubl.) Sterler, *Hort. Nympheb.*: 46 (Sterler 1821), “*Euosma*”.

Coussapoa obovata Miq., *Het Instituut* [2] 1842: 200 (Miquel 1843). — *Coussapoa latifolia* var. *obovata* (Miq.) Miq., *Fl. Bras. [Martius]* 4 (1): 135 [1 Dec. 1853] (Miquel 1853).

Coussapoa froesii Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2):* 162 (Standley 1937).

VERNACULAR NAMES. — Pa: kawukwine-awak • Wp: kwapo'i átā • Br: apuí-grande, muirapenima.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: J.B. Aublet s.n. (original material BM[BM000993414], see note under *C. angustifolia*); R.A.A. Oldeman B-1338, 9.5 m × 23 cm.

[1746] *Coussapoa leprieurii* Benoist

Bull. Mus. Natl. Hist. Nat. 30: 103 (Benoist 1924).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: F.M.R. Leprieur s.n. (holo-, P[P00757098]; iso-, P[P00757099]); R.A.A. Oldeman B-1946, dbh 25 cm.

[1747] *Coussapoa microcephala* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 96 (Trécul 1847).

Coussapoa fagifolia Klotzsch, *Linnaea* 20: 528 [Oct. 1847] (Klotzsch 1847).

Coussapoa cuneata Miq., *Fl. Bras. [Martius]* 4 (1): 138 [1 Dec. 1853] (Miquel 1853).

HERBARIUM DATA (FG). — A single collection, S.A. Mori et al. 8864.

SIZE. — Suriname. B. Maguire 24588 (MO), 15 m × 20 cm.

[1748] *Coussapoa parvifolia* Standl.

Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2): 165 (Standley 1937).

Coussapoa cornifolia Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2):* 160 (Standley 1937). — *Coussapoa microcephala* subsp. *cornifolia* (Standl.) Akkermans & C.C.Berg, *Proc. Kon. Ned. Akad. Wetensch. C* 85 (4): 457 (Akkermans & Berg 1982).

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: S.A. Mori et al. 23682.

SIZE. — Up to 20 m tall (Berg et al. 1990).




FIG. 57. — Styracaceae: **A**, *Styrax pallidus* A.DC. Symplocaceae: **B**, *Symplocos martinicensis* Jacq. Theaceae: **C**, *Gordonia fruticosa* (Schrad.) H.Keng. Ulmaceae: **D**, *Ampelocera edentula* Kuhlm. (*D. Sabatier & J.-F. Molino 5641*). A, B, D, © D. Sabatier/IRD; C, © C. Girod.

Genus *Pourouma* Aubl.

[1749] *Pourouma bicolor* Mart. subsp. *bicolor*

Syst. Mat. Med. Veg. Bras. 34 (Martius 1843).

Pourouma aspera Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 102 (Trécul 1847).

Pourouma crassivenosa Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 419 [1 Dec. 1928] (Mildbraed 1928).

Pourouma lawrancei Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser. 17 (2)*: 183 (Standley 1937).

Pourouma schultesii Cuatrec., *Caldasia* 7 (34): 303 (Cuatrecasas 1956).

Pourouma camaratana Cuatrec., *Acta Bot. Venez.* 2 (5-8): 202 (Cuatrecasas 1967).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Pa: kwem-purubumna • Wp: kulumá tololo • Wn: ulumapi • Nt: busi papay • Cr: mal-bwa-kannon • Fr: pourouma • Br: imbaúbarana.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (holotype of *Pourouma aspera*: P[P00757092]; iso-, K[K000512459], LE[LE00011416, LE00011417], P[P00757091]).

INVENTORY DATA (FG). — 63 trees in 41 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38$ cm.

[1750] *Pourouma bicolor* subsp. *digitata* (Trécul) C.C.Berg & Heusden

Proc. Kon. Ned. Akad. Wetensch. C 91 (2): 106 (Berg & Heusden 1988). — *Pourouma digitata* Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 106 (Trécul 1847).

VERNACULAR NAMES. — Pa: kwem, kwem-purubumna • Wp: a'i lea, kulumá tololo • Cr: mal-bwa-kannon • Br: imbaúbarana, mapatirana.

HERBARIUM DATA (FG). — 31 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (lecto-, P[P00753657], designated by Berg & van Heusden [Berg *et al.* 1990: 137]; isolecto-, B[B100248462]).

INVENTORY DATA (FG). — 39 trees in 22 plots; $F_{\max} = 1.2\%$; $dbh_{\text{inv}} = 46.9$ cm.

[1751] *Pourouma guianensis* Aubl.

Hist. Pl. Guiane 2: 892 [Jun.-Dec. 1775] (Aublet 1775).

Pourouma palmata Poepp. & Endl., *Nova genera ac species plantarum [Poeppig & Endlicher]* 2: 29 [Jan.-Sep. 1838] (Poeppig & Endlicher 1838).

Pourouma acutiflora Trécul, *Ann. Sci. Nat., Bot. sér. 3*, 8: 105 (Trécul 1847).

Pourouma cinerascens Mart. ex Miq., *Fl. Bras. [Martius]* 4 (1): 125 [1 Dec. 1853] (Miquel 1853).

Pourouma heterophylla Mart. ex Miq., *Fl. Bras. [Martius]* 4 (1): 125 [1 Dec. 1853] (Miquel 1853).

Pourouma fuliginea Miq., *Fl. Bras. [Martius]* 4 (1): 129 [1 Dec. 1853] (Miquel 1853).

Pourouma scabra Rusby, *Bull. New York Bot. Gard.* 6 (22): 498 [30 Nov. 1910] (Rusby 1910).

Pourouma radula Benoist, *Bull. Mus. Natl. Hist. Nat.* 28: 320 (Benoist 1922).

Pourouma subtriloba Rusby, *Mem. New York Bot. Gard.* 7: 232 (Rusby 1927).

Pourouma substrigosa Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 192 [20 Dec. 1927] (Mildbraed 1927).

Pourouma mildbraediana Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 183 (Standley 1937).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: puluma, yalayala • Wp: kulumā kalate, kulumā sī • Cr: mal-bwa-kannon • Br: embaúba-bengué.

HERBARIUM DATA (FG). — 21 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000993430](#)]).

INVENTORY DATA (FG). — 28 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 38.5$ cm.

[1752] *Pourouma melinonii* Benoist

Bull. Mus. Natl. Hist. Nat. 28: 318 (Benoist 1922).

Pourouma apaporiensis Cuatrec., *Caldasia* 7 (34): 297 (Cuatrecasas 1956).

Pourouma apaporiensis var. *macrophylla* Cuatrec., *Caldasia* 7 (34): 298 (Cuatrecasas 1956), “fma. *macrophylla* Cuatr, var. nov.”.

VERNACULAR NAMES. — Pa: kwem purubumna • Ka: puluma, yalayala • Te: pulumá • Wp: kule pi, kulumá tololo • Wn: ulumapi • Nt: busi papay • Cr: mal-bwa-kannon • Fr: pourouma • Br: imbaúba-da-mata, imbaúbarana, tararanga.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *P.A. Sagot 990*, Nov. 1856 (lecto-, P[[P06855173](#)], designated by Berg [Berg & Dewolf 1975: 274]; isolecto-, B[[B100248451](#)], K[[K000512469](#)]).

INVENTORY DATA (FG). — 147 trees in 63 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 50.3$ cm.

[1753] *Pourouma minor* Benoist

Bull. Mus. Natl. Hist. Nat. 30: 103 (Benoist 1924).

Pourouma aurea Mildbr., *Notizbl. Bot. Gart. Berlin-Dahlem* 10: 418 [1 Dec. 1928] (Mildbraed 1928).

Pourouma folleata J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 8 (2): 114 (Macbride 1930).

Pourouma isophlebia Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 182 (Standley 1937).

Pourouma subplicata Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 184 (Standley 1937).

Pourouma umbellata Standl., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 17 (2): 185 (Standley 1937).

Coussapoa emarginata Killip ex J.F.Macbr., *Publ. Field Mus. Nat. Hist., Bot. Ser.* 13 (2.2): 296 [15 Mar. 1937] (Macbride 1937).

Pourouma cuatrecasasii Standl., *Revista Acad. Colomb. Ci. Exact.* 9 (36/37): 339 [May 1956] (Standley 1956).

Pourouma umbellifera W.C.Burger, *Phytologia* 26 (6): 430 [11 Oct. 1973] (Burger 1973).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). Benoist (1924: 104) cited three collections in the protologue of *P. minor*: *R. Benoist 960*, *R. Benoist 978*, and *Wachenheim, 3e série, 18. Berg & van Heusden (1990: 186)* cited for the type “*Benoist 960* (lectotype, P, chosen here)”; their lectotype corresponds to two specimens at P ([P00756799](#) and [P00756800](#)). Both are labelled “*Pourouma minor* R. Ben.” by Benoist, but only one of them ([P00756800](#)) is labelled “Revised for *Flora Neotropicica*. *Pourouma minor* Benoist. Lectotype” by Berg and van Heusden. It is therefore hereafter chosen as the lectotype.

VERNACULAR NAMES. — Pa: kwem-seinō, kwem-tino • Wp: kulumā sī • Cr: mal-bwa-kannon • Br: mapatirana, purumaí, tourém.

HERBARIUM DATA (FG). — 64 collections at CAY. Sel. exs.: *R. Benoist 960* (lecto-, P[[P00756800](#), here designated; isolecto-, P[[P00756799](#)]).

INVENTORY DATA (FG). — 250 trees in 51 plots; $F_{\max} = 4.9\%$; $dbh_{\text{inv}} = 45$ cm.

[1754] *Pourouma mollis* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 102 (Trécul 1847).

VERNACULAR NAMES. — Pa: kwem-duwō, kwem-puvemna • Wp: wilau piyu'ā • Cr: mal-bwa-kannon • Br: tararanga-vermelha, uva-de-macaco.

HERBARIUM DATA (FG). — 29 collections at CAY. Sel. exs.: *F.M.R. Leprieur 141* (lecto-, P[[P00757085](#)], designated by Berg [Berg & Dewolf 1975: 272]; isolecto-, G[[G00438436](#)]).

INVENTORY DATA (FG). — 115 trees in 49 plots; $F_{\max} = 1.5\%$; $dbh_{\text{inv}} = 45.5$ cm.

[1755] *Pourouma saulensis* C.C.Berg & Kooy (Fig. 58C)

Brittonia 34 (1): 36 (Berg & Kooy 1982).

NOTE. — Known only from the Guiana Shield.

HERBARIUM DATA (FG). — 10 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-4132* (holo-, U[[U0004771](#)]; iso-, CAY[[CAY010613](#)], P[[P00756778](#)], VEN[[VEN179488](#)]).

INVENTORY DATA (FG). — 7 trees in 4 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 40.1$ cm.

[1756] *Pourouma tomentosa* Mart. ex Miq.
subsp. *maroniensis* (Benoist) C.C.Berg & Heusden
(Fig. 58D)

Proc. Kon. Ned. Akad. Wetensch. C 91 (2): 109 (Berg & Heusden 1988). — *Pourouma maroniensis* Benoist, *Bull. Mus. Natl. Hist. Nat.* 28: 318 (Benoist 1922).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: ayiwilan, yalayala • Wp: a'i lea, kulumá pilá • Cr: mal-bwa-kannon • Br: imbaúba-do-vinho.

HERBARIUM DATA (FG). — 24 collections at CAY. Sel. exs.: G. Wachenheim sér. 2, 392 (holo-, P[P06855698]; iso-, K[K000512468], P[P06855696, P06855697]).

INVENTORY DATA (FG). — 28 trees in 18 plots; $F_{\max} = 1.6\%$; $\text{dbh}_{\text{inv}} = 59.2\text{ cm}$.

[1757] *Pourouma velutina* Mart. ex Miq.

Fl. Bras. [Martius] 4 (1): 130 [1 Dec. 1853] (Miquel 1853).

Pourouma steyermarkii Standl. & Cuatrec., *Fieldiana, Bot.* 28 (1): 210 (Standley & Cuatrecasas 1951).

VERNACULAR NAMES. — Pa: kwem, kwem-duwó • Wp: wilau piyúá • Cr: mal-bwa-kannon • Br: mapatirana.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: D. Sabatier & M.-F. Prévost 2189.

INVENTORY DATA (FG). — 18 trees in 11 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 22\text{ cm}$.

[1758] *Pourouma villosa* Trécul

Ann. Sci. Nat., Bot. sér. 3, 8: 103 (Trécul 1847).

Pourouma laevis Benoist, *Bull. Mus. Natl. Hist. Nat.* 28: 319 (Benoist 1922).

VERNACULAR NAMES. — Pa: kwem • Wp: kulumá sī • Cr: mal-bwa-kannon • Br: purumá, tararanga.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: F.M.R. Leprieur s.n. (holo-, P, not seen; iso-, G[G00438449], P[P00756786, P00756787]).

INVENTORY DATA (FG). — 87 trees in 50 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 66.8\text{ cm}$.

[1759] *Pourouma* sp. A

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: J.-F. Molino & D. Sabatier 2337.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 17.6\text{ cm}$.

[1760] *Pourouma* sp. B

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: D. Sabatier 4873.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 48.7\text{ cm}$.

Genus *Urera* Gaudich.

[1761] *Urera caracasana* (Jacq.) Gaudich. ex Griseb.

Fl. Brit. W.I. [Grisebach] 154 [June 1859] (Grisebach 1859). — *Urtica caracasana* Jacq., *Pl. Rar. Hort. Schoenbr.* 3: 71 (Jacquin 1798). — *Urera jacquinii* Wedd., *Ann. Sci. Nat., Bot. sér. 3, 18:* 200 [Aug. 1852] (Weddell 1852), *nom. illeg. superfl.* (based on *Urtica caracasana*).

Urtica acuminata Poir., *Encycl. [J. Lamarck et al.] Suppl.* 4: 226 [29 June 1816] (Poirer 1816). — *Urera acuminata* (Poir.) Gaudich. ex Decne., *Nouv. Ann. Mus. Hist. Nat.* 3: 490 (Decaisne 1834). — *Urera jacquinii* var. *miquelii* Wedd., *Arch. Mus. Hist. Nat.* 9: 145 (Weddell 1856), *nom. illeg. superfl.* (based on *Urtica acuminata*). — *Urera caracasana* var. *miquelii* Wedd., *Prod. [A. P. de Candolle]* 16 (1): 90 [mid Nov. 1869] (Weddell 1869), *nom. illeg. superfl.* (based on *Urtica acuminata*).

Urtica alceifolia Poir., *Encycl. [J. Lamarck et al.] Suppl.* 4: 227 [29 June 1816] (Poirer 1816), “*alceaefolia*”. — *Urera alceifolia* (Poir.) Gaudich., *Voy. Uranie, Bot.* 497 [“1826” publ. 6 Mar. 1830] (Gaudichaud 1830), “*alceaefolia*”, *nom. inval.* (genus name and epithet not associated).

Urtica tiliifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 42 [28 Apr. 1817] (Kunth 1817), “*tiliaefolia*”.

Urtica ulmifolia Kunth, *Nova genera et species plantarum [H.B.K.]* 2: 42 [28 Apr. 1817] (Kunth 1817). — *Urera jacquinii* var. *ulmifolia* (Kunth) Wedd., *Arch. Mus. Hist. Nat.* 9: 145 (Weddell 1856).

Urtica mitis Vell., *Fl. Flumin. Icon.* 10: t. 19 [“1827” publ. 29 Oct. 1831] (Vellozo 1831). — *Urera mitis* (Vell.) Miq., *Fl. Bras. [Martius]* 4 (1): 191 [1 Dec. 1853] (Miquel 1853). — *Urera caracasana* var. *mitis* (Vell.) Wedd., *Prod. [A. P. de Candolle]* 16 (1): 90 [mid Nov. 1869] (Weddell 1869).

Urtica corallina Liebm., *Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afsl.*, ser. 5, 2: 295 (Liebmann 1851). — *Urera jacquinii* var. *corallina* (Liebm.) Wedd., *Arch. Mus. Hist. Nat.* 9: 145 (Weddell 1856). — *Urera corallina* (Liebm.) Wedd., *Prod. [A. P. de Candolle]* 16 (1): 90 [mid Nov. 1869] (Weddell 1869).

Urera subpeltata Miq., *Fl. Bras. [Martius]* 4 (1): 189 [1 Dec. 1853] (Miquel 1853). — *Urera jacquinii* var. *subpeltata* (Miq.) Wedd., *Arch. Mus. Hist. Nat.* 9: 145 (Weddell 1856). — *Urera caracasana* var. *subpeltata* (Miq.) Wedd., *Prod. [A. P. de Candolle]* 16 (1): 90 [mid Nov. 1869] (Weddell 1869).

Urera acuminata Miq., *Fl. Bras. [Martius]* 4 (1): 190 [1 Dec. 1853] (Miquel 1853), *nom. illeg. hom., non* (Poir.) Gaudich. ex Decne. (Decaisne 1834).

Urera subpeltata var. *morifolia* Miq., *Fl. Bras. [Martius]* 4 (1): 190 [1 Dec. 1853] (Miquel 1853).

Urera caracasana var. *tomentosa* Wedd., *Prod. [A. P. de Candolle]* 16 (1): 90 [mid Nov. 1869] (Weddell 1869).




FIG. 58. — Urticaceae: **A**, *Cecropia granvilleana* C.C.Berg (D. Sabatier & J.-F. Molino 5016); **B**, *Cecropia silvae* C.C.Berg (D. Sabatier & J.-F. Molino 5137); **C**, *Pourouma saulensis* C.C.Berg & Kooy (D. Sabatier & M.-F. Prévost 4846); **D**, *Pourouma tomentosa* subsp. *maroniensis* (Benoist) C.C.Berg & Heusden. © D. Sabatier/IRD.

VERNACULAR NAMES. — Wp: kelekele, pinô • Cr: zouti-montangn • Br: cansançao, urtiga-brava.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *M.-F. Prévost 3655*.

SIZE. — Up to 10 m tall (Berg 1992).

Family VERBENACEAE J.St.-Hil.
Genus *Citharexylum* B.Juss.

[1762] *Citharexylum macrophyllum* Poir.
(Fig. 59A)

Encycl. [J. Lamarck et al.] Suppl. 2: 367 [23 Oct. 1811] (Poiret 1811).

Citharexylum amazonicum Moldenke, *Repert. Spec. Nov. Regni Veg.* 37: 216 (Moldenke 1934).

VERNACULAR NAMES. — Pa: ivery • Ka: uluyatu • Wp: ka'i pimâ • Cr: bwa-kotlêt • Br: calango-cego.

HERBARIUM DATA (FG). — 16 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material FI[FI011051], P[P00689427]).

SIZE. — Up to 20 m tall (Jansen-Jacobs 1988).

[1763] *Citharexylum spinosum* L.

Sp. Pl. 2: 625 [1 May 1753] (Linnaeus 1753).

Citharexylum fruticosum L., *Syst. Nat.*, ed. 10, 2: 1115 [7 June 1759] (Linnaeus 1759), “fruticos.”. — *Citharexylum cinereum* L., *Sp. Pl.*, ed. 2, 2: 872 [Aug. 1763] (Linnaeus 1763), nom. illeg. superfl.

- (based on the same references as *Citharexylum fruticosum* ["Pluk. alm. 108. t. 162. f. 1" & "Plum. ic. 157. f. 1"]).
- Citharexylum quadrangulare* Jacq., *Enum. Syst. Pl.*: 26 [Aug.-Sep. 1760] (Jacquin 1760).
- Citharexylum teres* Jacq., *Enum. Syst. Pl.*: 26 [Aug.-Sep. 1760] (Jacquin 1760).
- Citharexylum villosum* Jacq., *Icon. Pl. Rar. [Jacquin]* 1: t. 118 (Jacquin 1784). — *Citharexylum fruticosum* var. *villosum* (Jacq.) O.E.Schulz, *Symb. Antill. [Urban]* 6 (1): 63 [15 July 1909] (Schulz 1909). — *Citharexylum spinosum* f. *villosum* (Jacq.) I.E.Méndez, *Willdenowia* 31 (2): 422 [17 Dec. 2001] (Méndez 2001).
- Citharexylum subserratum* Sw., *Prod. [Swartz]* 91 [20 Jun.-29 July 1788] (Swartz 1788). — *Citharexylum fruticosum* var. *subserratum* (Sw.) Moldenke, *Phytologia* 1 (1): 17 [Dec. 1933] (Moldenke 1933). — *Citharexylum fruticosum* f. *subserratum* (Sw.) Moldenke, *Phytologia* 36 (3): 164 [22 July 1977] (Moldenke 1977). — *Citharexylum spinosum* f. *subserratum* (Sw.) I.E.Méndez, *Willdenowia* 31 (2): 421 [17 Dec. 2001] (Méndez 2001).
- Citharexylum caudatum* Sw., *Observ. Bot. [Swartz]*: 234 [May-July 1791] (Swartz 1791), nom. illeg. hom., non L. (Linnaeus 1759).
- Citharexylum molle* Salisb., *Prodr. Stirp. Chap. Allerton* 108 [Nov.-Dec. 1796] (Salisbury 1796), nom. illeg. superf. (based on *Citharexylum subserratum* and *C. villosum*).
- Citharexylum pentandrum* Vent., *Descr. Pl. Nouv.*: pl. 47 (Ventenat 1800).
- Citharexylum pulverulentum* Pers., *Syn. Pl. [Persoon]* 2 (1): 142 [Nov. 1806] (Persoon 1806).
- Citharexylum tomentosum* Poir., *Encycl. [J. Lamarck et al.] Suppl.* 2: 368 [23 Oct. 1811] (Poirier 1811).
- Citharexylum molle* Jacq. ex Spreng., *Syst. Veg. [Sprengel]* 2: 764 [Jan.-May 1825] (Sprengel 1825), nom. illeg. hom., non Salisb. (Salisbury 1796).
- Citharexylum coriaceum* Desf., *Tabl. École Bot.*, ed. 3 [*Cat. Pl. Hort. Paris.*], 392 (Desfontaines 1829).
- Citharexylum cinereum* Moc. & Sessé ex D.Don, *Edinburgh New Philos. J.* 10: 238 [Oct. 1830-Mar. 1831] (Don 1831), nom. nud. pro syn.
- Citharexylum laevigatum* Hostmann ex Griseb., *Fl. Brit. W.I. [Grisebach]* 497 [prob. May 1862] (Grisebach 1862), nom. nud. pro syn.
- Citharexylum surrectum* Griseb., *Fl. Brit. W.I. [Grisebach]* 497 [prob. May 1862] (Grisebach 1862).
- Citharexylum polystachyum* Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (3): 209 (Turczaninow 1863).
- Citharexylum bahamense* Millsp., *Bull. New York Bot. Gard.* 3 (11): 450 [14 Apr. 1905] (Millspaugh 1905). — *Citharexylum fruticosum* f. *bahamense* (Millsp.) Moldenke, *Phytologia* 13 (4): 242 [16 July 1966] (Moldenke 1966).
- Citharexylum broadwayi* O.E.Schulz, *Symb. Antill. [Urban]* 7 (3): 354 [1 Oct. 1912] (Schulz 1912).
- Citharexylum fruticosum* var. *smallii* Moldenke, *Repert. Spec. Nov. Regni Veg.* 37: 223 (Moldenke 1934). — *Citharexylum spinosum* f. *smallii* (Moldenke) I.E.Méndez, *Willdenowia* 31 (2): 421 [17 Dec. 2001] (Méndez 2001).
- Citharexylum fruticosum* var. *subvillosum* Moldenke, *Repert. Spec. Nov. Regni Veg.* 37: 223 (Moldenke 1934). — *Citharexylum fruticosum* f. *subvillosum* (Moldenke) Moldenke, *Phytologia* 36 (3): 164 [22 July 1977] (Moldenke 1977). — *Citharexylum spinosum* f. *subvillosum* (Moldenke) I.E.Méndez, *Willdenowia* 31 (2): 422 [17 Dec. 2001] (Méndez 2001).
- Citharexylum fruticosum* var. *brittonii* Moldenke, *Lilloa* 4 (2): 311 (Moldenke 1939). — *Citharexylum spinosum* f. *brittonii* (Moldenke) I.E.Méndez, *Willdenowia* 31 (2): 422 [17 Dec. 2001] (Méndez 2001).
- Citharexylum hybridum* Moldenke, *Lilloa* 4 (2): 313 (Moldenke 1939).
- HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *C. Sastre* 265.
- SIZE. — Up to 16 m tall (Jansen-Jacobs 1988).
- Family VIOLACEAE Batsch
Genus *Amphirrhox* Spreng.
- [1764] *Amphirrhox longifolia* (A.St.-Hil.) Spreng.
Syst. Veg. [Sprengel] 4 (2, Cur. Post.): 99 [Jan.-June 1827] (Sprengel 1827). — *Spathularia longifolia* A.St.-Hil., *Mém. Mus. Hist. Nat.* 11: 492 (Saint-Hilaire 1824).
- Braddleya legalis* Vell., *Fl. Flumin.* 94 ["1825" publ. 7 Sep.-28 Nov. 1829] (Vellozo 1829).
- Amphirrhox latifolia* Mart. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 376 [1 Oct. 1871] (Eichler 1871).
- Amphirrhox surinamensis* Eichler, *Fl. Bras. [Martius]* 13 (1): 377 [1 Oct. 1871] (Eichler 1871). — *Hybanthus surinamensis* Miq. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 377 [1 Oct. 1871] (Eichler 1871), nom. nud. pro syn. — *Leonia surinamensis* (Eichler) Byng & Christenh., *Global Fl.* 4: 128 [9 Feb. 2018] (Byng & Christenhusz 2018).
- Amphirrhox juruana* Ule, *Verb. Bot. Vereins Prov. Brandenburg* 47: 156 [1 Oct. 1905] (Ule 1905).
- NOTE. — As the relationships between *Amphirrhox* Spreng. and *Leonia* Ruiz & Pav. are still unclear (Wahlert et al. 2014), the transfer of *A. surinamensis* Eichler to *Leonia* (Christenhusz et al. 2018): 128 seems premature.
- VERNACULAR NAMES. — Pa: mavinbi-kamwi, mavinvi-kamwi • Ka: kulali wepo.
- HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *J.-F. Molino* 1766.
- INVENTORY DATA (FG). — 139 trees in 24 plots; $F_{\max} = 3.1\%$; $dbh_{inv} = 26.7$ cm.
- Genus *Gloeospermum* Triana & Planch.
- [1765] *Gloeospermum sphaerocarpum* Triana & Planch.
Ann. Sci. Nat., Bot. sér. 4, 17: 129 (Triana & Planchon 1862). — *Leonia sphaerocarpum* (Triana & Planch.) Byng & Christenhusz., *Global Fl.* 4: 128 [9 Feb. 2018] (Byng & Christenhusz 2018).

Gloeospermum sphaerocarpum var. *latifolium* Melch., *Notizbl. Bot. Gart. Berlin-Dahlem* 8: 622 [15 Nov. 1923] (Melchior 1923).

NOTE. — According to Wahlert *et al.* (2014), there is a close relationship between *Gloeospermum* Triana & Planch. and *Leonia* Ruiz & Pav., but further studies are needed.

HERBARIUM DATA (FG). — 26 collections at CAY. Sel. exs.: *J.-J. de Granville B-3833*.

SIZE. — Colombia, Antioquia. *D. Cárdenas L. 2850* (MO), 8 m × 32.4 cm.

Genus *Leonia* Ruiz & Pav.

[1766] *Leonia glycycarpa* Ruiz & Pav.
(Fig. 59B, C)

Flora Peruviana 2: 69 (Ruiz & Pavón 1799). — *Theophrasta glycycarpa* (Ruiz & Pav.) Spreng., *Syst. Veg. [Sprengel]* 1: 671 [“1825” publ. late 1824] (Sprengel 1824). — *Leonia racemosa* Mart., *Nova genera et species plantarum [Martius]* 2 (2): 86 [Jan.–June 1827] (Martius 1827), nom. illeg. superfl. (based on *Leonia glycycarpa*).

Steudelia racemosa Mart., *Nova genera et species plantarum [Martius]* 2 (2): t. 168 [Jan.–June 1827] (Martius 1827), nom. nud.

Clavija sparsifolia Miq., *Fl. Bras. [Martius]* 10: 278 [15 Mar. 1856] (Miquel 1856). — *Theophrasta sparsifolia* (Miq.) Kuntze, *Revis. Gen. Pl.* 2: 404 [5 Nov. 1891] (Kuntze 1891).

Leonia melinoniana Baill. ex Sagot, *Ann. Sci. Nat., Bot. sér. 6*, 11: 146 (Sagot 1881).

Leonia glycycarpa var. *racemosa* L.B.Sm. & A.Fernández, *Caldasia* 6 (28): 169 (Smith & Fernández 1954).

NOTES. — Hyperdominant in Amazonia (ter Steege *et al.* 2020). *L. glycycarpa* var. *racemosa* L.B.Sm. is based on the illegitimate *L. racemosa* Mart., thus authorship is not “(Mart.) L.B.Sm. & A.Fernández”, but otherwise it complies with the requirements for a new name (it was published before 1958, therefore designation of a type is not mandatory).

VERNACULAR NAMES. — Pa: inam-etni-ayeweyo • Wp: eilowakulu, inámu sī, inámu sī wila • Br: gogó-de-guariba.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2753*.

INVENTORY DATA (FG). — 145 trees in 78 plots; $F_{\max} = 1.7\%$; $dbh_{\text{inv}} = 37.5$ cm.

Genus *Paypayrola* Aubl.

[1767] *Paypayrola bordenavei* (H.E.Ballard & Munzinger)
Byng & Christenb.

Global Fl. 4: 128 [9 Feb. 2018] (Byng & Christenhusz 2018). — *Hekkingia bordenavei* H.E.Ballard & Munzinger, *Syst. Bot.* 28 (2): 345 [15 May 2003] (Ballard & Munzinger 2003).

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *B. Bordenave 5040* (holo-, P[P00217081]; iso-, K[K001070514, K001070515], U[U0098667, U0098668], US[00902493]).

SIZE. — Up to 15 cm dbh (Munzinger & Ballard 2003).

[1768] *Paypayrola grandiflora* Tul.

Ann. Sci. Nat., Bot. sér. 3, 7: 371 (Tulasne 1847).

Paypayrola ventricosa Tul., *Ann. Sci. Nat., Bot. sér. 3, 11:* 153 (Tulasne 1849).

HERBARIUM DATA (FG). — A single collection, *R.A.A. Oldeman B-3110*.

SIZE. — Brazil, Amazonas. *B.A. Krukoff 8023* (MO), 50 ft × 5 inch. (15 m × 12.6 cm).

[1769] *Paypayrola guianensis* Aubl.

Hist. Pl. Guiane 1: 249 [Jun.–Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Lignonia monodynema* J.F.Gmel., *Syst. Nat., ed. 13[bis], 2 (1):* 373 [late Sep.–Nov. 1791] (Gmelin 1791), nom. illeg. superfl. (based on *Lignonia* Scop., thus indirectly on *Paypayrola guianensis*). — *Payrola guianensis* (Aubl.) Lam., *Tabl. Encycl. 2[4 (1):* 72 [21 Nov. 1796] (Lamarck 1796), nom. illeg. superfl. (genus name superfluous, based on the type of *Paypayrola*). — *Wibelia guianensis* (Aubl.) Pers., *Syn. Pl. [Persoon]* 1: 210 [1 Apr.–15 June 1805] (Persoon 1805), nom. illeg. superfl. (genus name superfluous, based on the type of *Paypayrola*).

Periclistia latifolia Benth., *J. Bot. [Hooker]* 4: 109 (Bentham 1842).

VERNACULAR NAMES. — Pa: yauknabui • Ka: kakaulan, kupali elepalí • Wp: wila poloelaa • Cr: bwa-poul • Br: manaca-rana.

HERBARIUM DATA (FG). — 132 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-PJR[P00777978] designated by Lanjouw & Uittien [1940: 155]).

INVENTORY DATA (FG). — 60 trees in 14 plots; $F_{\max} = 3.3\%$; $dbh_{\text{inv}} = 21.3$ cm.

[1770] *Paypayrola bulkiana* Pulle
(Fig. 59D)

Recueil Trav. Bot. Néerl. 9: 155 (Pulle 1912).

VERNACULAR NAMES. — Pa: mavinbi-priyu, mavinvi-priyu • Wp: tatu kāsī • Cr: bwa-poul • Br: manaca-rana.

HERBARIUM DATA (FG). — 60 collections at CAY. Sel. exs.: *J.-P. Lescure 723*.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 13.8$ cm.

[1771] *Paypayrola longifolia* Tul.

Ann. Sci. Nat., Bot. sér. 3, 7: 372 (Tulasne 1847).

VERNACULAR NAMES. — Ka: palipyo, polipyo, polipyoli • Wp: kelekele sili • Cr: bwa-poul • Br: manaca-rana.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *H. Jacquin* 1763.

SIZE. — Suriname. *J.C. Lindeman et al.* 555 (MO), 10 m × 11 cm.

Genus *Rinorea* Aubl.

[1772] *Rinorea amapensis* Hekking

Phytologia 43 (5): 476 [24 Sep. 1979] (Hekking 1979).

VERNACULAR NAMES. — Pa: inam-etni, kuuku-ariut-kamwi • Ka: pel-pele apotokon • Wp: iáyü • Cr: ti-bwa-lélé • Br: canela-de-jacamim.

HERBARIUM DATA (FG). — 96 collections at CAY. Sel. exs.: *M.-F. Prévost* 3075.

INVENTORY DATA (FG). — 6 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 14.8$ cm.

[1773] *Rinorea babiensis* (Moric.) Kuntze

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Alsodeia babiensis* Moric., *Pl. Nouv. Amér.* 68 (Moricand 1839).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *R.A.A. Oldeman* B-813.

INVENTORY DATA (FG). — 1 tree, $dbh = 11.1$ cm.

[1774] *Rinorea brevipes* (Benth.) S.F.Blake

Contr. U.S. Natl. Herb. 20 (13): 512 [19 July 1924] (Blake 1924). — *Alsodeia brevipes* Benth., *J. Bot. [Hooker]* 4: 106 (Bentham 1842). — *Conohoria brevipes* (Benth.) Miq., *Linnæa* 22: 556 (Miquel 1849). — *Alsodeia guianensis* (Aubl.) Eichler var. *brevipes* (Benth.) Eichler, *Fl. Bras. [Martius]* 13 (1): 387 [1 Oct. 1871] (Eichler 1871).

Alsodeia laxiflora Benth., *J. Bot. [Hooker]* 4: 107 (Bentham 1842). — *Conohoria laxiflora* (Benth.) Miq. *Linnæa* 22: 556 (Miquel 1849). — *Alsodeia guianensis* var. *laxiflora* (Benth.) Eichler, *Fl. Bras. [Martius]* 13 (1): 387 [1 Oct. 1871] (Eichler 1871). — *Rinorea laxiflora* (Benth.) Melch., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 21: 352 (Melchior 1925).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *C. Feillet et al.* 10115.

INVENTORY DATA (FG). — 28 trees in 4 plots; $dbh_{inv} = 19.7$ cm.

[1775] *Rinorea falcata* (Mart. ex Eichler) Kuntze
(Fig. 59E)

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Alsodeia falcata* Mart. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 386 [1 Oct. 1871] (Eichler 1871).

Alsodeia guianensis var. *parviflora* Eichler, *Fl. Bras. [Martius]* 13 (1): 387 [1 Oct. 1871] (Eichler 1871).

Rinorea surinamensis Melch., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 21: 352 (Melchior 1925), nom. nud.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier* 5573.

SIZE. — Up to 10 m tall (Hekking 1988).

[1776] *Rinorea flavescentis* (Aubl.) Kuntze

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Conohoria flavescentis* Aubl., *Hist. Pl. Guiane* 1: 239 [Jun.-Dec. 1775] (Aublet 1775). — *Alsodeia flavescentis* (Aubl.) Spreng., *Syst. Veg. [Sprengel]* 1: 806 ["1825" publ. late 1824] (Sprengel 1824), "Alsodeia".

VERNACULAR NAMES. — Pa: aaku-priyo, pahe-avan-kamwi • Ka: palipyo, polipyo, polipyoli • Wp: iáyü • Nt: bofo tiki, bofoo tiki • Cr: ti-bwa-lélé • Br: canela-de-jacamim.

HERBARIUM DATA (FG). — 50 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (lecto-, P-JJR[P00777786] designated by Lanjouw & Uittien [1940: 150]).

INVENTORY DATA (FG). — 24 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{inv} = 19.1$ cm.

[1777] *Rinorea guianensis* Aubl.

Hist. Pl. Guiane 1: 235 [Jun.-Dec. 1775] (Aublet 1775). — *Conohoria rinorea* A.St.-Hil., *Mém. Mus. Hist. Nat.* 11: 492 (Saint-Hilaire 1824), nom. illeg. superfl. (based on *Rinorea guianensis*). — *Alsodeia rinorea* Spreng., *Syst. Veg. [Sprengel]* 1: 807 ["1825" publ. late 1824] (Sprengel 1824), nom. illeg. superfl. (based on *Rinorea guianensis*).

Alsodeia floribunda Moric., *Pl. Nouv. Amér.* 70 (Moricand 1839).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.B. Aublet* s.n. (lecto-, P-JJR[P00777787] designated by Lanjouw & Uittien [1940: 156]).

INVENTORY DATA (FG). — 98 trees in 19 plots; $F_{\max} = 4.1\%$; $dbh_{inv} = 35$ cm.

[1778] *Rinorea macrocarpa* (Mart. ex Eichler) Kuntze

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Alsodeia macrocarpa* Mart. ex Eichler, *Fl. Bras. [Martius]* 13 (1): 385 [1 Oct. 1871] (Eichler 1871).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *J.-J. de Granville* 1169.

SIZE. — Up to 10 m tall (Hekking 1988).

[1779] *Rinorea neglecta* Sandwith

Kew Bull. 10 (3): 371 [20 Dec. 1955] (Sandwith 1955).

HERBARIUM DATA (FG). — 34 collections at CAY. Sel. exs.: *D. Sabatier* & *M.-F. Prévost* 2203.

INVENTORY DATA (FG). — 2 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 10.5$ cm.

[1780] *Rinorea paniculata* (Mart.) Kuntze

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Alsodeia paniculata* Mart., *Nova genera et species plantarum [Martius]* 1 (2): 30 [late 1823 or Jan.-Feb. 1824] (Martius 1823-1824), “*Alsodeia*”.

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *R.A.A. Oldeman B-2084*.

SIZE. — Up to 25 m tall (Hekking 1988).

[1781] *Rinorea pectinosquamata* Hekking

Phytologia 53 (4): 252 [2 June 1983] (Hekking 1983), “*pectino-squamata*”.

NOTE. — Known only from French Guiana.

HERBARIUM DATA (FG). — 45 collections at CAY. Sel. exs.: *B. Riéra* 470 (holo-, U[U0007081]; iso-, CAY[CAY024975, CAY024976, CAY024977, CAY024978, CAY024979]).

INVENTORY DATA (FG). — 211 trees in 48 plots; $F_{\max} = 3.4\%$; $dbh_{\text{inv}} = 21\text{ cm}$.

[1782] *Rinorea pubiflora* (Benth.) Sprague & Sandwith var. *pubiflora*

Bull. Misc. Inform. Kew 1931 (4): 171 [18 Apr. 1931] (Sprague & Sandwith 1931). — *Passoura guianensis* Aubl., *Hist. Pl. Guiane* 2 (Suppl.): 21 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Conohoria passoura* DC. ex Ging., *Prodr. [A. P. de Candolle]* 1: 312 [mid Jan. 1824] (Gingins 1824), *nom. illeg. superfl.* (based on *Passoura guianensis*). — *Alsodeia pubiflora* Benth., *J. Bot. [Hooker]* 4: 106 (Bentham 1842). — *Alsodeia guianensis* (Aubl.) Eichler, *Fl. Bras. [Martius]* 13 (1): 387 [1 Oct. 1871] (Eichler 1871). — *Rinorea passoura* Kuntze, *Revis. Gen. Pl.* 1: 42 [5 Nov. 1891] (Kuntze 1891), “*Passura*”, *nom. illeg. superfl.* (based on *Passoura guianensis*).

Alsodeia martinii Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 36 (1): 557 (Turczaninow 1863), “*Alsodeja, Martinii*”. — *Rinorea martinii* (Turcz.) S.F.Blake, *Contr. U.S. Natl. Herb.* 20 (13): 506 [19 July 1924] (Blake 1924), “*martini*”.

VERNACULAR NAMES. — Wp: iāyū • Nt: bonbekiidya • Cr: ti-bwa-lélé • Br: canela-de-jacamim.

HERBARIUM DATA (FG). — 86 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 893*.

INVENTORY DATA (FG). — 2 trees in 1 plot; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 14.7\text{ cm}$.

[1783] *Rinorea pubiflora* var. *grandifolia* (Eichler) Hekking

Fl. Neotrop. Monogr. 46: 108 [19 Jan. 1988] (Hekking 1988). — *Alsodeia falcata* var. *grandifolia* Eichler, *Fl. Bras. [Martius]* 13 (1): 386 [1 Oct. 1871] (Eichler 1871). — *Rinorea passoura* var. *grandifolia* (Eichler) Hekking, *Phytologia* 53 (4): 257 [2 June 1983] (Hekking 1983). — *Rinorea passoura* f. *grandifolia* (Eichler) Hekking, *Phytologia* 53 (4): 257 [2 June 1983] (Hekking 1983). — *Rinorea pubiflora* f. *grandifolia* (Eichler) Hekking, *Fl. Neotrop. Monogr.* 46: 109 [19 Jan. 1988] (Hekking 1988).

Rinorea passoura var. *andersonii* Sandwith ex Hekking, *Phytologia* 43 (5): 479 [24 Sep. 1979] (Hekking 1979).

Rinorea passoura f. *andersonii* Sandwith ex Hekking, *Phytologia* 43 (5): 479 [24 Sep. 1979] (Hekking 1979).

Rinorea pubiflora f. *andersonii* (Sandwith ex Hekking) Hekking, *Fl. Neotrop. Monogr.* 46: 110 [19 Jan. 1988] (Hekking 1979).

Rinorea passoura f. *leiosperma* Hekking, *Phytologia* 43 (5): 480 [24 Sep. 1979] (Hekking 1979).

Rinorea pulleana Melch., *Nat. Pflanzenfam. [Engler & Prantl]*, ed. 2, 21: 352 (Melchior 1925), *nom. nud.*

Rinorea scandens Ule, *Verb. Bot. Vereins Prov. Brandenburg* 47: 157 [1 Oct. 1905] (Ule 1905).

VERNACULAR NAMES. — Ka: tona, tonawewe • Cr: ti-bwa-lélé.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *J.-J. de Granville et al. 92294*.

SIZE. — Up to 20 m tall (Hekking 1988).

[1784] *Rinorea riana* Kuntze

Revis. Gen. Pl. 1: 42 [5 Nov. 1891] (Kuntze 1891). — *Riana guianensis* Aubl., *Hist. Pl. Guiane* 1: 237 [Jun.-Dec. 1775] (Aublet 1775), “*Guyannensis*” on plate. — *Conohoria riana* DC. ex Ging., *Prodr. [A. P. de Candolle]* 1: 312 [mid Jan. 1824] (Gingins 1824), *nom. illeg. superfl.* (based on *Riana guianensis*). — *Alsodeia prunifolia* Spreng., *Syst. Veg. [Sprengel]* 1: 807 [“1825” publ. late 1824] (Sprengel 1824), *nom. illeg. superfl.* (based on *Riana guianensis*).

NOTES. — The name “*Alsodeia riana* Turcz.”, sometimes given in synonymy, does not exist. Turczaninow (1863): 557 did not associate the genus name with the specific epithet. The text reads: “*Alsodeja* (*Conohoria Riana* Aubl. *nec* Steud. in Hostm. et Kappl. pl. Surin. No 1126) [...].” This statement does not refer to *Conohoria riana* DC. ex Ging., but rather to a manuscript annotation by Steudel on specimen *Hostmann and Kappler's 1126*.

VERNACULAR NAMES. — Wp: iāyū, yu'i li • Wn: pakumleimë • Cr: ti-bwa-lélé.

HERBARIUM DATA (FG). — 131 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00777787]), designated by Lanjouw & Uittien [1940: 156].

SIZE. — Suriname. *S.A. Mori & A. Bolten 8398* (P), 6 m × 10 cm.

Genus *Rinoreocarpus* Ducke[1785] *Rinoreocarpus ulei* (Melch.) Ducke

Arch. Jard. Bot. Rio de Janeiro 5: 173 (Ducke 1930). — *Gloeospermum ulei* Melch., *Notizbl. Bot. Gart. Berlin-Dahlem* 9: 164 [30 Dec. 1924] (Melchior 1924). — *Rinoreocarpus salmoneus* Ducke, *Arch. Jard. Bot. Rio de Janeiro* 4: 144 (Ducke 1925), *nom. illeg. superfl.* (includes the type of *Gloeospermum ulei*). — *Leonia ulei* (Melch.) Byng & Christenh., *Global Fl.* 4: 128 [9 Feb. 2018] (Byng & Christenhusz 2018).

NOTES. — Hyperdominant in Amazonia (ter Stege *et al.* 2020). The position of *Rinoreocarpus* Ducke is still unresolved (Wahlert *et al.* 2014). It is therefore premature to place *R. ulei* in *Leonia*.

VERNACULAR NAMES. — Pa: agagut-aška, ararut-aška • Br: pau-estalador.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *M.-F. Prévost & P. Grenand 4324*.

INVENTORY DATA (FG). — 18 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{inv} = 19.6$ cm.

Family VOCHysiaceae A.St.-Hil.
Genus *Erisma* Rudge

[1786] *Erisma floribundum* Rudge

Pl. Guian. [Rudge] 1 (1): 7 [Apr. 1805] (Rudge 1805). — *Debraea floribunda* (Rudge) Roem. & Schult., *Syst. Veg. [Roemer & Schultes] 1*: 34 [Jan.-June 1817] (Roemer & Schultes 1817). — *Ditmaria floribunda* (Rudge) Spreng., *Syst. Veg. [Sprengel] 1*: 16 ["1825" publ. late 1824] (Sprengel 1824).

Erisma parvifolium Gleason, *Bull. Torrey Bot. Club* 60 (5): 362 [May 1933] (Gleason 1933).

Erisma pallidiflorum Ducke, *Arch. Inst. Biol. Veg. 2 (1)*: 54 [Sep. 1935] (Ducke 1935). — *Erisma parvifolium* var. *pallidiflorum* (Ducke) Ducke, *Arch. Inst. Biol. Veg. 4 (1)*: 43 [June 1938] (Ducke 1938).

VERNACULAR NAMES. — Ka: kuwalilan • Wp: kwali sili • Nt: feli kwali, jaboti, manhunti kwali • Cr: grignon-fou • Br: jaboti-da-terra-firme.

HERBARIUM DATA (FG). — 12 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material BM[[BM000566722](#), [BM000938998](#)], BR, [[BR0000005529711](#)], LINN[LINN-HS 17.1], M[[M0239626](#)], MO[[MO-251856](#)]).

INVENTORY DATA (FG). — 8 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{inv} = 90$ cm.

[1787] *Erisma nitidum* DC.

Prod. [A. P. de Candolle] 3: 30 [mid Mar. 1828] (Candolle 1828).

Qualea lutea Desf. ex Link, *Jahrb. Gewächsk. 1 (3)*: 24 (Link 1820), *nom. nud.*

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: makuk • Nt: manhunti kwali.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *J. Martin s.n.* (holo-, FI[[FI005833](#)]; iso-, FI[[FI005832](#)], P[[P00733810](#), [P00733811](#), [P00733812](#)]; probable iso-, B[B -W 00040 -01 0]).

INVENTORY DATA (FG). — 8 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{inv} = 27.8$ cm.

[1788] *Erisma uncinatum* Warm.

Fl. Bras. [Martius] 13 (2): 110 [1 Mar. 1875] (Warming 1875).

Erisma pulverulentum Poepp. ex Warm., *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1889: 28 (Warming 1889).

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Ka: kusili yepo, palulu ipyo, walapa kuwalini • Wp: kwali sili • Wn: kalapi alikan • Nt: feli kwali, manhunti

kwali • Cr: grignon-fou, kwali • Fr: grignon fou • Br: jaboti-da-terra-firme, quarubarana.

HERBARIUM DATA (FG). — 39 collections at CAY. Sel. exs.: *D. Sabatier 2312*.

INVENTORY DATA (FG). — 69 trees in 30 plots; $F_{\max} < 1\%$; $dbh_{inv} = 116.5$ cm.

Genus *Qualea* Aubl.

[1789] *Qualea acuminata* Spruce ex Warm.

Fl. Bras. [Martius] 13 (2): 40 [1 Mar. 1875] (Warming 1875).

Qualea speciosa Huber, *Bol. Mus. Paraense Hist. Nat. Ethnogr. 3*: 425 (Huber 1902).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *F.M.R. Leprieur 284* (P[[P04660120](#)]).

SIZE. — Up to 20 m tall (Marcano-Berti 1998).

[1790] *Qualea amapaensis* Balslev & S.A.Mori
(Fig. 60A)

Brittonia 33 (1): 5 (Balslev & Mori 1981), "amapaënsis".

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Wp: kulawa'i, kwali sili, pilikolo'i.

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *M.-F. Prévost & D. Sabatier 2755*.

INVENTORY DATA (FG). — 1 tree, $dbh = 20.4$ cm.

[1791] *Qualea caerulea* Aubl.

Hist. Pl. Guiane 1: 7 [Jun.-Dec. 1775] (Aublet 1775), "coerulea".

NOTES. — Maas & Maas-van de Kamer (1993: 30) concluded that the specific epithet "coerulea" should be corrected to "caerulea". They stated that for epithets printed in italics in Aublet (1775), "one can see no difference between 'æ' and 'œ'". However, these signs are clearly distinct. The latter is used in the heading of the description of all species with epithets "foetida" (*Agave foetida* L., *Passiflora foetida* L. and *Ticorea foetida* Aubl.), "foetidum" (*Eryngium foetidum* L.), "foetidissima" (*Tournefortia foetidissima* L.) or "stoechadifolia" (*Verbena stoechadifolia* L.), as well as in the index for the same species, except for *Tournefortia "stæcadifolia"* (obviously a double typographical error). The sign for "æ" is used both in the heading and in the index for *aestuans* (*Urtica aestuans* L.), *chamaecrista* (*Cassia chamaecrista* L.), *hypogaea* (*Arachis hypogaea* L.), *caribaicum* (*Pancratium caribaeum* L.), *sagittafolium* (*Arum sagittafolium* L.), *pes-caprae* (*Convolvulus pes-caprae* L.), *laevis* (*Manabea laevis* Aubl. and *Trigonia laevis* Aubl.), *laevigata* (*Melastoma laevigata* L.), as well as for 4 out of 6 epithets based on the "caerul" root: *caerulea* (*Besleria caerulea* Aubl.), *caerulescens* (*Lisanthius caerulescens* Aubl. and *Spermacoce caerulescens* Aubl.) and *caerulea* (*Voyria caerulea* Aubl.) For the remaining two (*Qualea caerulea* Aubl. and *Xiphidium caeruleum* Aubl.), the "æ" sign is also used in the index, although "œ" is used for the heading in the text. It thus seems clear that the latter two should be treated as typographical errors. Meanwhile, the "æ" sign never appears on plates (only nine of the above species are illustrated). However, the




Fig. 59. — Verbenaceae: **A**, *Citharexylum macrophyllum* Poir. (M.-F. Prévost 1404). Violaceae: **B**, **C**, *Leonia glycycarpa* Ruiz & Pav. (D. Sabatier 3502); **D**, *Paypayrola hulkiana* Pulle (M.-F. Prévost et al. 4587); **E**, *Rinorea falcata* (Mart. ex Eichler) Kuntze (D. Sabatier 5573). A, C, © M.-F. Prévost/IRD; B, D, E, © D. Sabatier/IRD.

fact that “*guianensis*” is almost always misspelled “*Guyannensis*” on plates suggests that the figure legends were not checked by Aublet.

VERNACULAR NAMES. — Pa: á-seiminio-puvemna • Ka: kumetilan, kuwali, kwali, tamunen ilakopi • Te: kwadi • Wp: pilima'i • Nt: sabana kwali • Cr: grignon-fou, grignon-sentmarie • Fr: grignon fou • Br: mandioqueira-azul, quaruba-azul.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[[BM000566635](#)]).

INVENTORY DATA (FG). — 56 trees in 8 plots; $F_{\max} = 4.1\%$; $\text{dbh}_{\text{inv}} = 92.6\text{ cm}$.

[1792] *Qualea dinizii* Ducke

Arch. Jard. Bot. Rio de Janeiro 1 (1): 49 (Ducke 1915).

VERNACULAR NAMES. — Wp: pilima'i piyū, pilima'i wu.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *G. Angel 29*.

INVENTORY DATA (FG). — 3 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 50.6\text{ cm}$.

[1793] *Qualea moriboomiorum* Marc.-Berti
(Fig. 60C)

Pittieria 18: 5 (Marcano-Berti 1989), “*mori-boomii*”.

NOTE. — Known only from French Guiana. Automatic correction of “*mori-boomii*”, according to Art. 60.8 of the ICN (Turland *et al.* 2018).

VERNACULAR NAMES. — Fr: gonfolo bleu.

HERBARIUM DATA (FG). — 6 collections at CAY. Sel. exs.: *S.A. Mori & B.M. Boom 15225* (holo-, NY[[00001003](#)] ; iso-, CAY[[CAY024985](#)], P[[P00162580](#)]).

INVENTORY DATA (FG). — 44 trees in 18 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 86.1\text{ cm}$.

[1794] *Qualea polychroma* Stafleu

Acta Bot. Neerl. 2 (2): 182 (Stafleu 1953).

HERBARIUM DATA (FG). — 3 collections at CAY. Sel. exs.: *D. Sabatier 1462*.

INVENTORY DATA (FG). — 6 trees in 3 plots; $F_{\max} < 1\%$; $\text{dbh}_{\text{inv}} = 50.4\text{ cm}$.

[1795] *Qualea rosea* Aubl.

Hist. Pl. Guiane 1: 5 [Jun.-Dec. 1775] (Aublet 1775).

Qualea violacea Mart. & Zucc. ex Schult., *Mant.* 3 [Schultes] Add. 2: 99 (Schultes 1822), *nom. nud. pro syn.*

Qualea melinonii P.Beckm., *Bot. Jahrb. Syst.* 40 (3): 280 [24 Jan. 1908] (Beckmann 1908), “*Mélinonii*”.

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Pa: á-seiminio-seine • Ka: ilakopi • Wp: pilima'i piyū, pilima'i wu • Nt: gonfolo kwali • Cr: grignon-fou • Fr: gonfolo rose • Br: mandioqueira-rosa.

HERBARIUM DATA (FG). — 53 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (original material BM[BM000566650]).

INVENTORY DATA (FG). — 955 trees in 104 plots; $F_{\max} = 11.8\%$; $\text{dbh}_{\text{inv}} = 125\text{ cm}$.

[1796] *Qualea tricolor* Benoist
(Fig. 60B)

Notul. Syst. (Paris) 3: 176 [25 Dec. 1915] (Benoist 1915).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: á-seiminio-seine • Nt: gonfolo kwali • Br: mandioqueira.

HERBARIUM DATA (FG). — 7 collections at CAY. Sel. exs.: *R. Benoist 1564* (holo-, P[P00733870]; iso-, P[P00733871]).

INVENTORY DATA (FG). — 37 trees in 14 plots; $F_{\max} = 1.7\%$; $\text{dbh}_{\text{inv}} = 98.4\text{ cm}$.

Genus *Ruizterania* Marc.-Berti

[1797] *Ruizterania albiflora* (Warm.) Marc.-Berti

Pittieria 2: 9 (Marcano-Berti 1969). — *Qualea albiflora* Warm., *Fl. Bras. [Martius]* 13 (2): 36 [1 Mar. 1875] (Warming 1875).

Qualea glaberrima Ducke, *Arch. Jard. Bot. Rio de Janeiro* 1 (1): 46 (Ducke 1915).

VERNACULAR NAMES. — Pa: á-seiminio-puvemna • Ka: tiyapotano ilakopi • Nt: gonfolo kwali • Cr: grignon-boni, grignon-fou • Fr: gonfolo gris • Br: mandioqueira-lisa, mandioqueira-preta.

HERBARIUM DATA (FG). — 35 collections at CAY. Sel. exs.: *M.-F. Prévost 3198*.

INVENTORY DATA (FG). — 174 trees in 94 plots; $F_{\max} = 1.5\%$; $\text{dbh}_{\text{inv}} = 143.2\text{ cm}$.

[1798] *Ruizterania ferruginea* (Steyermark.) Marc.-Berti

Pittieria 2: 11 (Marcano-Berti 1969). — *Qualea ferruginea* Steyermark., *Fieldiana, Bot.* 28 (2): 295 (Steyermark 1952).

Qualea rubiginosa Stafleu, *Acta Bot. Neerl.* 2 (2): 154 (Stafleu 1953). — *Ruizterania rubiginosa* (Stafleu) Marc.-Berti, *Pittieria* 2: 14 (Marcano-Berti 1969).

Qualea apodocarpa Steyermark., *Bol. Soc. Venez. Ci. Nat.* 26: 473 (Steyermark 1966). — *Ruizterania apodocarpa* (Steyermark.) Marc.-Berti, *Pittieria* 2: 10 (Marcano-Berti 1969).

Qualea rubiginosa var. *angustior* Steyermark., *Acta Bot. Venez.* 2 (5-8): 239 (Steyermark 1967). — *Ruizterania rubiginosa* var. *angustior* (Steyermark.) Marc.-Berti, *Pittieria* 2: 15 (Marcano-Berti 1969).

HERBARIUM DATA (FG). — A single collection, *D. Sabatier 4874*, $\text{dbh} = 90\text{ cm}$.

Genus *Vochysia* Aubl.

[1799] *Vochysia cayennensis* Warm.

Fl. Bras. [Martius] 13 (2): 80 [1 Mar. 1875] (Warming 1875).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Wp: kwali sī, malakapuli'i • Nt: wana kwali • Fr: grignon sainte marie.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *J. Martin s.n.* (original material B[B -W 00038 -01 0], BM[BM000566734], E[E00285635], F[V0074840F, V0074841F], K[K000566091], NY[00001041], P[P00733830, P00733831, P00733832], US[00108791]).

SIZE. — Up to 80 cm dbh (Marcano-Berti 1998).

[1800] *Vochysia densiflora* Spruce ex Warm.

(Fig. 60D)

Fl. Bras. [Martius] 13 (2): 101 [1 Mar. 1875] (Warming 1875).

VERNACULAR NAMES. — Ka: tapilen kuwali • Wp: kwali pilá, kwali pitá • Wn: kanawaimë • Nt: wana kwali • Cr: kwali.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier et al. 6272*.

INVENTORY DATA (FG). — 20 trees in 7 plots; $F_{\max} = 1.4\%$; $\text{dbh}_{\text{inv}} = 112.7\text{ cm}$.

[1801] *Vochysia glaberrima* Warm.

Fl. Bras. [Martius] 13 (2): 78 [1 Mar. 1875] (Warming 1875).

Vochysia lucida Klotsch ex M.R.Schomb., *Reis. Br.-Guiana [Ri. Schomburgk]* 3: 1099 [“1848” publ. 7-10 Mar. 1849] (Schomburgk 1849), *nom. illeg. hom., non* C.Presl (1834).

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *D. Sabatier & J.-F. Molino 5136*.

INVENTORY DATA (FG). — 1 tree, $\text{dbh} = 111.4\text{ cm}$.

[1802] *Vochysia guianensis* Aubl.

Hist. Pl. Guiane 1: 18 [Jun.-Dec. 1775] (Aublet 1775) “*Vochy*”, “*Guyannensis*” on plate. — *Cucullaria excelsa* Willd., *Sp. Pl.*, ed. 4 1 (1): 17 [June 1797] (Willdenow 1797), *nom. illeg. superfl.* (based on *Vochysia guianensis*).

Vochysia excelsa A.Dietr., *Sp. Pl.*, ed. 6, 1: 103 (Dietrich 1831).

Vochysia melinonii P.Beckm., *Bot. Jahrb. Syst.* 40 (3): 280 [24 Jan. 1908] (Beckmann 1908), “*Mélinonii*”.

Vochysia paraensis Huber, *Bol. Mus. Goeldi Hist. Nat. Ethnogr.* 6: 214 (Huber 1910), “paraënsis”, nom. nud.

VERNACULAR NAMES. — Pa: á-seiminio • Ka: kulalu, kumetilan, kuwali, kwali, wosiwosi • Wp: malakapuli'i • Nt: mutende kwali • Cr: bwa-kruzo, kwali • Fr: bois cruzeau • Br: quaruba-branca, quaruba-rosa, quaruba-vermelha, rabo-de-tucano.

HERBARIUM DATA (FG). — 38 collections at CAY. Sel. exs.: *J.B. Aublet s.n.* (lecto-, P-JJR[P00680428] designated by Lanjouw & Uittien [1940: 160]).

INVENTORY DATA (FG). — 45 trees in 24 plots; $F_{\max} = 1.6\%$; $dbh_{\text{inv}} = 107.3$ cm.

[1803] *Vochysia neyratii* Normand

Adansonia, sér. 2, 17 (1): 11 (Normand 1977).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: á-seiminio • Wp: kwali pilá, kwali pitá, kwali sili • Nt: asiwa kwali • Cr: achiwa-kwali • Fr: couali albert.

HERBARIUM DATA (FG). — 20 collections at CAY. Sel. exs.: *I. Petrov 185* (holo-, P[P00733889]; iso-, CAY[CAY024986, CAY024987], P[P00733887, P00733888]).

INVENTORY DATA (FG). — 1 tree, $dbh = 43.6$ cm.

[1804] *Vochysia rufescens* W.A.Rodrigues

Acta Amazonica 1 (2): 33 (Rodrigues 1971).

HERBARIUM DATA (FG). — 9 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 4419*.

INVENTORY DATA (FG). — 5 trees in 3 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 57.9$ cm.

[1805] *Vochysia sabatieri* Marc.-Berti
(Fig. 60E)

Pittieria 18: 9 (Marcano-Berti 1989).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Nt: kopi kwali.

HERBARIUM DATA (FG). — 13 collections at CAY. Sel. exs.: *D. Sabatier 1170* (original material CAY[CAY024988, CAY024989], NY[00927585]).

INVENTORY DATA (FG). — 3 trees in 2 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 87$ cm.

[1806] *Vochysia sofiae* Marc.-Berti & Poncy

Pittieria 40: 176 (Marcano-Berti & Poncy 2016).

NOTE. — Known only from French Guiana.

VERNACULAR NAMES. — Pa: á-seiminio-purubumna • Cr: kwali • Br: quaruba.

HERBARIUM DATA (FG). — 4 collections at CAY. Sel. exs.: *O. Tostain et al. 2725* (holo-, P[P04776153]; iso-, CAY[CAY101513, CAY101514, CAY101515], P[P04776154]), 35 m.

[1807] *Vochysia speciosa* Warm.

Fl. Bras. [Martius] 13 (2): 79 [1 Mar. 1875] (Warming 1875).

HERBARIUM DATA (FG). — 5 collections at CAY. Sel. exs.: *P.A. Poiteau s.n.* (original material G[G00343253, G00343255], K[K000565944], M[M0239555], P[P00733917, P00733918]).

SIZE. — Up to 45 m tall (Marcano-Berti 1998).

[1808] *Vochysia surinamensis* Stafleu

Meded. Bot. Mus. Herb. Rijks Univ. Utrecht 95: 439 (Stafleu 1948).

VERNACULAR NAMES. — Ka: wosiwosi • Wp: malakapuli'i • Nt: mutende kwali • Cr: kwali • Fr: couali.

HERBARIUM DATA (FG). — 17 collections at CAY. Sel. exs.: *D. Sabatier & M.-F. Prévost 2942*.

INVENTORY DATA (FG). — 29 trees in 14 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 96.8$ cm.

[1809] *Vochysia tetraphylla* (G.Mey.) DC.

Prodr. [A. P. de Candolle] 3: 27 [mid Mar. 1828] (Candolle 1828). — *Cucullaria tetraphylla* G.Mey., *Prim. Fl. Esseq.* 12 [Nov. 1818] (Meyer 1818).

Vochysia arcuata Garcke, *Linnaea* 22: 58 (Garcke 1849).

NOTE. — Known only from the Guiana Shield.

VERNACULAR NAMES. — Ka: woto kuwalili.

HERBARIUM DATA (FG). — 8 collections at CAY. Sel. exs.: *D. Sabatier 990*.

SIZE. — Up to 100 cm dbh (Marcano-Berti 1998).

[1810] *Vochysia tomentosa* (G.Mey.) DC.

Prodr. [A. P. de Candolle] 3: 26 [mid Mar. 1828] (Candolle 1828). — *Cucullaria tomentosa* G.Mey., *Prim. Fl. Esseq.* 13 [Nov. 1818] (Meyer 1818).

VERNACULAR NAMES. — Ka: kuwali, kwali, walapa kuwalili, wosiwosi tamunen • Wp: kwali, kwali e'e, kwali pilá, kwali pitá • Nt: wana kwali • Cr: kwali • Fr: couali • Br: quaruba.

HERBARIUM DATA (FG). — 36 collections at CAY. Sel. exs.: *L.C. Richard s.n.* (original material P[P00733963, P00733965]).

INVENTORY DATA (FG). — 74 trees in 28 plots; $F_{\max} = 2.1\%$; $dbh_{\text{inv}} = 133$ cm.

[1811] *Vochysia vismiifolia* Spruce ex Warm.

Fl. Bras. [Martius] 13 (2): 99 [1 Mar. 1875] (Warming 1875), “*vismiaefolia*”.

NOTE. — Hyperdominant in Amazonia (ter Steege *et al.* 2020).

VERNACULAR NAMES. — Br: quaruba-vermelha.

HERBARIUM DATA (FG). — 2 collections at CAY. Sel. exs.: *J.-J. de Granville 5550*.

INVENTORY DATA (FG). — 10 trees in 5 plots; $F_{\max} < 1\%$; $dbh_{\text{inv}} = 43.6$ cm.




FIG. 60. — Vochysiaceae: **A**, *Qualea amapaensis* Balslev & S.A.Mori (M.-F. Prévost & D. Sabatier 2755); **B**, *Qualea tricolor* Benoist (D. Sabatier 6341); **C**, *Qualea moriboomiorum* Marc.-Berti; **D**, *Vochysia densiflora* Spruce ex Warm.; **E**, *Vochysia sabatieri* Marc.-Berti (D. Sabatier & M.-F. Prévost 4850). © D. Sabatier/IRD.

REFERENCES

- ACEVEDO-RODRÍGUEZ P. 2003. — *Melicocceae (Sapindaceae)*: Melicoccus and Talisia. The New York Botanical Garden Press, Bronx, 179 p. (Flora Neotropica Monograph; 87).
- ACEVEDO-RODRÍGUEZ P. 2012. — *Sapindaceae*. Royal Botanic Gardens, Kew, 196 p. (Flora of the Guianas. Series A, Phanerogams; Fascicle 29).
- AHLBRINCK W. G. 1931 [transl. van Herwijnen D. 1956]. — *L'Encyclopédie des Caraïbes*. Institut géographique national, Paris, 544 p.
- ALLEN C. K. 1966. — Contributions to the botany of Guyana II. Lauraceae. *Memoirs of the New York Botanical Garden* 15: 53-93.
- AMSHOFF G. J. H. 1939. — On South American Papilionaceae. *Medeelingen van het Botanisch Museum en Herbarium van de Rijks Universiteit te Utrecht* 52: 1-78.
- AMSHOFF G. J. H. 1948. — Myrtaceae. *Bulletin of the Torrey Botanical Club* 75 (5): 528-538. <https://doi.org/10.2307/2481787>
- AMSHOFF G. J. H. 1951. — Myrtaceae, in LANJOUW J. & STOFFERS A. L. (eds), *Flora of Suriname*. Vol. 3 (2). Brill, Leiden, the Netherlands: 56-158.
- AMSHOFF G. J. H. 1976. — Papilionaceae, in LANJOUW J. & STOFFERS A. L. (eds.), *Flora of Suriname*. Vol. 2 (2). Brill, Leiden, the Netherlands: 1-257.
- ANDERSON W. R. 1981. — *Malpighiaceae*, in MAGUIRE B. (ed.), *The Botany of the Guyana Highland*. Part XI. The New York Botanical Garden Press, New York: 21-305 (Memoirs of the New York Botanical Garden; 32).
- ANDERSON W. R. 1999. — *Malpighiaceae*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 6. Missouri Botanical Garden Press, Saint Louis: 82-185.
- ANDERSON W. R. 2007. — Lectotypification of names of *Malpighiaceae* – I. *Contributions of the University of Michigan Herbarium* 25: 83-93. <https://www.biodiversitylibrary.org/page/12697665>
- ANGIOSPERM PHYLOGENY GROUP 2016. — An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181 (1): 1-20. <https://doi.org/10.1111/bj.12385>
- AUBLET J.-B. C. F. 1775. — *Histoire des plantes de la Guyane françoise*. Didot, Paris, 4 vol. <https://doi.org/10.5962/bhl.title.674>
- AYMARD C. G. A. 1998. — Dilleniaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 609-610.
- AYMARD C. G. A. & HOWARD R. A. 2004. — Polygonaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 8. Missouri Botanical Garden Press, Saint Louis: 347-370.
- BAILLON H. E. 1870. — *Histoire des plantes*. Vol. 2. Hachette, Paris, 512 p. <https://doi.org/10.5962/bhl.title.40796>
- BARNEBY R. C. 1998. — *Silk Tree, Guanacaste, Monkey's Earring*. Part III. Calliandra. New York Botanical Garden Press, Bronx, 223 p. (Memoirs of the New York Botanical Garden; 77).

- BARNEBY R. C. & GRIMES J. W. 1996. — *Silk Tree, Guanacaste, Monkey's Earring: a Generic System of the Synandrous Mimosaceae of the Americas*. Part I. Abarema, Albizia, and allies. New York Botanical Garden Press, Bronx, 292 p. (Memoirs of the New York Botanical Garden; 74 [1]).
- BARNEBY R. C. & GRIMES J. W. 1997. — *Silk Tree, Guanacaste, Monkey's Earring*. Part II. Pithecellobium, Cojoba, and Zygia. New York Botanical Garden Press, Bronx, 161 p. (Memoirs of the New York Botanical Garden; 74 [2]).
- BARNEBY R. C. & HEALD S. V. 2002. — Caesalpiniaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx: 167-183. (Memoirs of the New York Botanical Garden; 76 [2]).
- BARNEBY R. C., GRIMES J. W. & PONCY O. 2011. — *Leguminosae subfamily Mimosoideae*. Royal Botanic Gardens, Kew, 381 p. (Flora of the Guianas. Series A, Phanerogams; Fascicle 28).
- BARRÈRE P. 1741. — *Essai sur l'histoire naturelle de la France Equinoxiale*. Piget, Paris, 215 p. <https://doi.org/10.5962/bhl.title.131289>
- BARTHÉLEMI G. 2007. — *Dictionnaire créole guyanais-français*. Ibis Rouge Editions, Cayenne, 461 p.
- BENOIST R. 1919. — Les Licania (Chrysobalanacées) de la Guyane française. *Bulletin du Muséum national d'Histoire naturelle* 25 (1): 512-516. <https://www.biodiversitylibrary.org/page/5027185>
- BENOIST R. 1933. — *Les bois de la Guyane Française*. Éditions des Archives de Botanique, Caen, 292 p., 58 pl. h.t. (Archives de Botanique tome V, Mémoire n°1).
- BENTHAM G. 1844. — Notes on Mimoseae, with a synopsis of species. Tribe III. Acacieae (continued). *The London Journal of Botany* 3: 195-226.
- BENTHAM G. 1852. — Second Report on Mr. Spruce's Collections of Dried Plants from North Brazil. *Hooker's Journal of Botany and Kew Garden Miscellany* 4: 8-18.
- BENTHAM G. 1861. — Notes on Tiliaceae. *Journal of the Proceedings of the Linnean Society. Botany* 5(Suppl. 2): 52-74.
- BENTHAM G. 1865. — Ordo LVII. Leguminosae, in BENTHAM G. & HOOKER J. D. (eds), *Genera Plantarum*. Vol. 1 (2). Lovell Reeve, London: 434-600.
- BENTHAM G. 1870. — Leguminosae II. et III, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 15 (2). F. Fleischer, Leipzig: 1-528.
- BENTHAM G. 1875. — Revision of the suborder Mimoseae. *Transactions of the Linnean Society of London* 30 (3): 335-664.
- BERG C. C. 1972. — *Olmedieae & Brosimeae (Moraceae)*. The New York Botanical Garden Press, Bronx, 228 p. (Flora Neotropica Monograph; 7).
- BERG C. C. 1992. — *Ulmaceae, Moraceae, Cecropiaceae, Urticaceae, Casuarinaceae*. Koeltz Scientific Books, Koenigstein, Germany, 224 p. (Flora of the Guianas. Series A, Phanerogams; Fascicle 11).
- BERG C. C. 1999. — Moraceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 6. Missouri Botanical Garden Press, Saint Louis: 693-729.
- BERG C. C. & CARAUTA J. P. P. 2003. — New species of *Ficus* (Moraceae) from Brazil. *Brittonia* 54 (4): 236-250. <https://doi.org/cmjhck>
- BERG C. C. & DEWOLF G. P. 1975. — Moraceae, in LANJOUW J. & STOFFERS A. L. (eds), *Flora of Suriname*. Vol. 5. Brill, Leiden, the Netherlands: 173-319.
- BERG C. C., AKKERMANS R. W. A. P. & HEUSDEN E. C. H. VAN 1990. — *Cecropiaceae: Coussapoa and Pourouma, with an Introduction to the Family*. The New York Botanical Garden Press, Bronx, 208 p. (Flora Neotropica Monograph; 51).
- BERG C. C., FRANCO ROSELLI P. & DAVIDSON D. W. 2005. — *Cecropia*. The New York Botanical Garden Press, Bronx, 230 p. (Flora Neotropica Monograph; 94).
- BERG O. 1859. — Myrtaceae. III, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 14(1). F. Fleischer, Leipzig: 529-656.
- BERG O. 1861. — Mantissa II ad revisionem Myrtacearum Americae. *Linnæa* 30: 647-713.
- BERRY P. E. & BARNEBY R. C. 1999. — Calliandra, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 6. Missouri Botanical Garden Press, Saint Louis: 600-607.
- BERRY P. E. & PEIXOTO A. L. 1999. — Monimiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 6. Missouri Botanical Garden Press, Saint Louis: 689-691.
- BERRY P. E., GRÖGER A., HOLST B. K., MORLEY T., MICHELANGELI F. A., LUCKANA N. G., ALMEDA F., RENNER S. S., FREIRE-FIERRO A., ROBINSON O. R. & YATSKIEVYCH K. 1999. — Melastomataceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 6. Missouri Botanical Garden Press, Saint Louis: 263-528.
- BHL 2022. — Biodiversity Heritage Library. Available from <https://www.biodiversitylibrary.org/> [accessed 7 March 2022].
- BLEEKRODE S. A. 1857. — Notice sur la Gutta-Percha de Surinam. *Annales des Sciences naturelles, Botanique*, sér. 4, 7: 220-228.
- BNF 2022. — Gallica. Available from <https://gallica.bnf.fr/> [accessed 7 March 2022].
- BOHS L. 1994. — Cyphomandra (Solanaceae). The New York Botanical Garden Press, Bronx, 175 p. (Flora Neotropica Monograph; 63).
- BOOM B. M. 1984. — A Revision of *Isertia* (Isertiae: Rubiaceae). *Brittonia* 36 (4): 425. <https://doi.org/10.2307/2806603>
- BOOM B. M. & DELPRETE P. G. 2002. — Rubiaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx: 606-649 (Memoirs of the New York Botanical Garden; 76 [2]).
- BRUMMITT R. K. & POWELL C. E. 1992. — *Authors of Plant Names*. Royal Botanic Gardens, Kew, 732 p.
- CAMARGO E. & TAPINKILI 2009. — *Dictionnaire bilingue wayana-français*. CELIA, DRAC-Guyane, TEKUREMAI, Cayenne, 25 p.
- CANDOLLE A. P. DE 1824. — *Prodromus Systematis Naturalis Regni Vegetabilis*. Vol. 1. Treuttel & Würtz, Paris, 747 p. <https://www.biodiversitylibrary.org/page/153953>
- CANDOLLE A. P. DE 1826. — Revue de la famille des Lythraires. *Mémoires de la Société de Physique et d'Histoire naturelle de Gênevèse* 3 (2): 65-96.
- CANDOLLE A. P. DE 1830. — *Prodromus Systematis Naturalis Regni Vegetabilis*. Vol. 4. Treuttel & Würtz, Paris, 683 p. <https://www.biodiversitylibrary.org/page/152139>
- CANDOLLE A. P. DE 1844. — *Prodromus Systematis Naturalis Regni Vegetabilis*. Vol. 8. Fortin, Masson & ass., Paris, 684 p. <https://www.biodiversitylibrary.org/page/162945>
- CANDOLLE A. P. DE 1864. — *Prodromus Systematis Naturalis Regni Vegetabilis*. Vol. 15 (1). Masson, Paris, 522 p. <https://www.biodiversitylibrary.org/page/157758>
- CARDINAL-MCTEAGUE W. M., SYTSMA K. J. & HALL J. C. 2016. — Biogeography and diversification of Brassicales: A 103million year tale. *Molecular Phylogenetics and Evolution* 99: 204-224. <https://doi.org/10.1016/j.ympev.2016.02.021>
- CARDOSO D., SÄRKINEN T., ALEXANDER S., AMORIM A. M., BITTRICH V., CELIS M., DALY D. C., FIASCHI P., FUNK V. A., GIACOMIN L. L., GOLDENBERG R., HEIDEN G., IGANCI J., KELLOFF C. L., KNAPP S., CAVALCANTE DE LIMA H., MACHADO A. F. P., DOS SANTOS R. M., MELLO-SILVA R., MICHELANGELI F. A., MITCHELL J., MOONLIGHT P., DE MORAES P. L. R., MORI S. A., NUNES T. S., PENNINGTON T. D., PIRANI J. R., PRANCE G. T., DE QUEIROZ L. P., RAPINI A., RIINA R., RINCON C. A. V., ROQUE N., SHIMIZU G., SOBRAL M., STEHMANN J. R., STEVENS W. D., TAYLOR C. M., TROVÓ M., VAN DEN BERG C., VAN DER WERFF H., VIANA P. L., ZARTMAN C. E. & FORZZA R. C. 2017. — Amazon plant diversity revealed by a taxonomically verified species list. *Proceedings of the National Academy of Sciences* 114 (40): 10695-10700. <https://doi.org/10.1073/pnas.1706756114>

- CAVALCANTE P. B. 1983. — *Revisão taxonômica do gênero Simaba Aubl. (Simaroubaceae) na América do Sul*. Museu Goeldi, Manaus, 85 p. (Publicações Avulsas do Museu Goeldi; 37).
- CHANDERBALI A. S. 2004. — *Endlicheria (Lauraceae)*. The New York Botanical Garden Press, Bronx, 141 p. (Flora Neotropica Monograph; 91).
- CHEVALIER A. 1932. — Les vrais et les faux Balatas. *Journal d'agriculture traditionnelle et de botanique appliquée* 12 (128): 261-282. <https://doi.org/10.3406/jatba.1932.5122>
- CHRISTENHUSZ M. J. M., FAY M. F. & BYNG J. W. (eds) 2018. — *Plant Gateway's the Global Flora: A Practical Flora to Vascular Plant Species of the World*. Vol. 4. GLOVAP Nomenclature Part 1. Plant Gateway Ltd., Bradford, United Kingdom, 155 p.
- CLEMENT C. R. 1999. — 1492 and the loss of Amazonian crop genetic resources. I. The relation between domestication and human population decline. *Economic Botany* 53: 188-202.
- COMPÈRE P. 1963. — The correct name of the Afro-American black mangrove. *Taxon* 12 (4): 150-152. <https://doi.org/10.2307/1216183>
- CORNEJO X. & ILTIS H. H. 2008. — Two new genera of Capparaceae: *Sarcotoxicum* and *Mesocapparis* stat. nov., and the reinstatement of *Neocalyptrocalyx*. *Harvard Papers in Botany* 13 (1): 103-116. <https://doi.org/c7mxhq>
- COURTZ H. 2008. — *A Carib Grammar and Dictionary*. Magonia books, Toronto, 501 p.
- COWAN R. S. 1953. — A taxonomic revision of the genus *Macrolobium* (Leguminosae-Caesalpinoideae). *Memoirs of the New York Botanical Garden* 8 (4): 257-407.
- COWAN R. S. 1968. — *Swartzia* (Leguminosae, Caesalpinoideae Swartzieae). The New York Botanical Garden Press, Bronx, 228 p. (Flora Neotropica Monograph; 1).
- COWAN R. S. 1975. — A monograph of the genus *Eperua* (Leguminosae: Caesalpinoideae). *Smithsonian Contributions to Botany* 28: 1-45.
- COWAN R. S. & BARNEY R. C. 1998. — *Macrolobium*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 71-82.
- COWAN R. S. & BERRY P. E. 1998. — *Elizabetha*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 61-63.
- CROAT T. B. 1974. — A new species of *Myrcia* (Myrtaceae) for Panama. *Annals of the Missouri Botanical Garden* 61 (3): 886-888. <https://doi.org/10.2307/2395035>
- CUATRECASAS J. 1964. — Cacao and its allies: a taxonomic revision of the genus *Theobroma*. *Contributions of the U.S. National Herbarium* 35: 379-614.
- CUATRECASAS J. & CROAT T. B. 1980. — Flora of Panama, Part VI. Family 93. *Malpighiaceae*. *Annals of the Missouri Botanical Garden* 67 (4): 851-945.
- CUELLO N. L. 1998. — *Tovomita*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 290-298.
- DALY D. C. 1992. — New taxa and combinations in *Protium* Burm. f. Studies in neotropical Burseraceae VI. *Brittonia* 44 (3): 280-299. <https://doi.org/10.2307/2806927>
- DALY D. C. 1997. — Burseraceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 3. Missouri Botanical Garden Press, Saint Louis: 688-728.
- DALY D. C. 1998. — Two new species of *Protium* from French Guiana. Studies in Neotropical Burseraceae VIII. *Brittonia* 50 (4): 517-523. <https://doi.org/10.2307/2807763>
- DALY D. C. 2002. — Burseraceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx: 151-165. (Memoirs of the New York Botanical Garden; 76 [2]).
- DECAISNE J. 1834. — Description d'un herbier de l'île de Timor, faisant partie des collections botaniques du Muséum d'histoire naturelle (Herbarii Timorensis descriptio). *Nouvelles Annales du Muséum d'Histoire naturelle* 3: 333-501. <https://www.biodiversitylibrary.org/page/10294502>
- DEFILIPPS R. A. 2002. — *Nyctaginaceae*, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx: 551-553. (Memoirs of the New York Botanical Garden; 76 [2]).
- DEFILIPPS R. A. & MAINA S. L. 2003. — *Nyctaginaceae*. Royal Botanic Garden, Kew: 23-55. (Flora of the Guianas. Series A, Phanerogams; Fascicle 22).
- DELPRETE P. G. 2015. — Typification and etymology of Aublet's Rubiaceae names. *Taxon* 64 (3): 595-624. <https://doi.org/10.12705/643.13>
- DELPRETE P. G. & PERSSON C. 2012. — *Octavia sessiliflora* DC. and *Mussaenda glomerulata* Lam. ex Poir., two obscure taxa from French Guiana synonymous with members of the *Alibertia* group (Rubiaceae, Gardeniae). *Adansonia* 34 (2): 353-363. <https://doi.org/10.5252/a2012n2a9>
- DELPRETE P. G., ACHILLE F. & MOULY A. 2010. — Four new combinations in Chomelia and Stenostomum (Rubiaceae, Guettardeae) from Central America, the Guianas and the Amazon Basin. *Blumea – Biodiversity, Evolution and Biogeography of Plants* 55 (2): 164-170. <https://doi.org/10.3767/000651910X526906>
- DON D. 1823. — An Illustration of the Natural Family of Plants called Melastomaceae. *Memoirs of the Wernerian Natural History Society* 4 (2): 276-329.
- DON D. 1828. — *Description of the genus Pinus*. 2nd ed., Vol. 1. Weddell et al., London, 42 p.
- DON G. 1831. — *A General History of the Dichlamydeous Plants*. Vol. 1. Rivington et al., London, 818 p.
- DORR L. J. & NICOLSON D. H. 2008-2009. — *Taxonomic Literature. A Selective Guide to Botanical Publications and Collections with Dates, Commentaries and Types*. Suppl. 7-8. Koelz Scientific Books, Königstein, 2 vols.
- DUBARD M. 1912. — Les Sapotacées du groupe des Sidéroxylinées. *Annales du Musée colonial de Marseille* 10: 1-90.
- DUCKE A. 1915. — Plantes nouvelles ou peu connues de la région amazonienne. *Archivos do Jardim Botânico do Rio de Janeiro* 1 (1): 7-57.
- DUCKE A. 1935. — Le « Pajurá » et le « Parinary » d'Amazonie. *Revue de Botanique appliquée et d'Agriculture coloniale* 15: 179-182.
- DUCKE A. 1942. — New and noteworthy Sapotaceae of Brazilian Amazonia. *Tropical Woods* 71: 7-25.
- DUNAL M.-F. 1817. — *Monographie de la famille des Anonacées*. Treuttel & Würtz, Paris, 145 p.
- ENDRESS M. E. 1995. — *Malouetia* (Apocynaceae), in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 2. Missouri Botanical Garden Press, Saint Louis: 369-372.
- ENGEL J. 2015. — *Plot Networks & Teams*. Available from <https://atdnmorphospecies.myspecies.info/node/781> [accessed 14 December 2020].
- ENGEL J. & SABATIER D. 2020. — *Terminalia carinata* Sabatier & J. Engel, sp. nov. (Combretaceae), a new large tree species from the Guiana shield revealed by re-examination of material previously identified as *T. guyanensis* Eichler. *Adansonia*, sér. 3, 42 (16): 261-271. <https://doi.org/10.5252/adansonia2020v42a16>. <http://adansonia.com/42/16>
- ENGEL J., MOLINO J.-F. & MARINHO L. 2022. — The maximum and the minimum: two new species of *Tovomita* Aubl. (Clusiaceae) from the Guiana Shield with an unusual number of stamens. *Adansonia* 44 (16): 165-174. <https://doi.org/10.5252/adansonia2022v44a16>. <http://adansonia.com/44/16>

- ENGLER H. G. A. 1876. — Ochnaceae, Anacardiaceae, Sabiaceae, Rhizophoraceae, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 12(2). F. Fleischer, Leipzig: 297-432.
- E-ReColNat 2022. — Réseau des Collections naturalistes. Available from <https://www.recolnat.org/> [accessed 11 April 2022].
- ESTEVES G. L. 2005. — Flora da Reserva Ducke, Amazonas, Brasil. Part I. Bombacaceae. *Rodriguésia* 56 (86): 115-124. <https://doi.org/10.1590/2175-78602005568624>
- Europeana 2021. — Available from <https://www.europeana.eu/> [accessed 2 December 2021].
- FEUILLET C. 2009. — Checklist of the Plants of the Guiana Shield. 1. An update to the angiosperms. *Journal of the Botanical Research Institute of Texas* 3 (2): 799-814. <https://archive.org/details/bistor-157820>
- FIASCHI P. & PLUNKETT G. M. 2018. — Revision of the Didymopanax Group of Neotropical Schefflera (Araliaceae) 1. *Annals of the Missouri Botanical Garden* 103 (1): 24-105. <https://doi.org/10.3417/2017031>
- FLEURY M. 1991. — 'Busi nenge', les Hommes-Forêt. *Essai d'ethnobotanique chez les Aluku (Boni) en Guyane française*, Thèse de Doctorat, Univ. Paris 6, Paris, 357 p.
- FORTE J. 1996. — *The Fanshaw/Boyan Glossary of Arawak Names in Natural History*. Amerindian Research Unit, University of Guyana, Georgetown, Guyana, 68 p.
- FRÉZIER A.-F. 1732. — *Relation du voyage de la Mer du Sud aux côtes du Chily et du Pérou, fait pendant les années 1712, 1713 & 1714*. Nyon, Didot, Quillau, Paris, 298 p.
- FRIES R. E. 1934. — Revision der Arten einiger Annonaceen-Gattungen III. *Acta Horti Bergiani* 12 (1): 28-105.
- FRIES R. E. 1939. — Revision der Arten einiger Annonaceen-Gattungen V. *Acta Horti Bergiani* 12 (3): 289-577.
- FRODIN D. G. 1997. — Araliaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 3. Missouri Botanical Garden Press, Saint Louis: 1-18.
- FUNK V., HOLLOWELL T., BERRY P. E., KELLOFF C. L. & ALEXANDER S. N. 2007. — Checklist of the plants of the Guiana Shield (Venezuela; Amazonas, Bolívar, Delta Amacuro; Guyana, Surinam, French Guiana). *Contributions from the United States National Herbarium* 55: 1-584. <https://repository.si.edu/handle/10088/34962>
- GAERTNER C. F. VON 1807. — *Supplementum carpologiae*. C. F. Richter, Leipzig, 256 p.
- GAHAGEN B., TERBUSH M. & BALLARD H. 2015. — A taxonomic synthesis of the *Tovomita weddelliana* (Clusiaceae) species complex. *Systematic Botany* 40 (4): 968-988. <https://doi.org/10.1600/036364415X690012>
- GAUDICHAUD C. 1830. — *Voyage autour du monde, entrepris par ordre du roi. Exécuté sur les corvettes de S.M. l'Uranie et la Physicienne, pendant les années 1817, 1818, 1819 et 1820*. Vol. 4. Botanique. Pillet Aîné, Paris, 522 p.
- GAVIRIA J. 1997. — *Cordia*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 3. Missouri Botanical Garden Press, Saint Louis: 297-304.
- GENTRY A. H. 1992. — *Bignoniaceae*. Part II. *Tribe Tecomeae*. The New York Botanical Garden Press, Bronx, 335 p. (Flora Neotropica Monograph; 25 [2]).
- GILLESPIE L. J. 2002. — Euphorbiaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx: 206-298. (Memoirs of the New York Botanical Garden; 76 [2]).
- GOLDENBERG R., ALMEDA F., CADDAA M. K., MARTINS A. B., MEIRELLES J., MICHELANGELI F. A. & WEISS M. 2013. — Nomenclator botanicus for the neotropical genus *Miconia* (Melastomataceae: Miconieae). *Phytotaxa* 106 (1): 1-171. <https://doi.org/10.11646/phytotaxa.106.1.1>
- GOND V., FREYCON V., MOLINO J.-F., BRUNAUX O., INGRASSIA F., JOUBERT P., PEKEL J.-F., PRÉVOST M.-F., THIERRON V., TROMBE P.-J. & SABATIER D. 2011. — Broad-scale spatial pattern of forest landscape types in the Guiana Shield. *International Journal of Applied Earth Observation and Geoinformation* 13 (3): 357-367. <https://doi.org/10.1016/j.jag.2011.01.004>
- GONZALEZ S., BILOT-GUÉRIN V., DELPRETE P. G., GENIEZ C., MOLINO J.-F., SMOCK J.-L. & THEVENY F. 2021. — *L'herbier IRD de Guyane*. <https://doi.org/10.23708/herbier-guyane-ird>
- GÖRTS VAN RIJN A. R. A. 2007. — *Piperaceae*. Royal Botanic Garden, Kew: 15-215. (Flora of the Guianas. Series A, Phanerogams; Fascicle 24).
- GOVAERTS R., SOBRAL N., ASHTON P., BARRIE F., HOLST B. K., LANDRUM L. L., MATSUMOTO K., MAZINE F. F., NIC LUGHADHA E. M., PROENÇA C. E. B. et al. 2008. — *World Checklist of Myrtaceae*. Kew Publishing, Kew, United Kingdom, 455 p.
- GRANVILLE J.-J. DE & GAYOT M. 2014. — *Guide des palmiers de Guyane*. ONF, Cayenne, 272 p.
- GREEN D. & GREEN H. 1998. — *Yuwit kawihka: Dicionário experimental Palikur Português*. Sociedad international de lingüística, Belem, Pará, 361 p.
- GRENNAND F. 1989. — *Dictionnaire Wayápi Français; Lexique Français Wayápi (Guyane Française)*. Peeters/Selaf, Paris, 538 p. (Langues et Sociétés d'Amérique traditionnelle).
- GRENNAND P., MORETTI C., JACQUEMIN H. & PRÉVOST M.-F. 2004. — *Pharmacopées traditionnelles en Guyane : Créoles, Wayápi, Palikur*. 2^{me} éd. IRD Éditions, Paris, Nogent-sur-Marne, 816 p.
- GROPPONI M. & PIRANI J. R. 2012. — A revision of *Hertia* (Rutaceae). *Systematic Botany* 37 (1): 197-212. <https://doi.org/10.1600/036364412X616765>
- GUITET S., SABATIER D., BRUNAUX O., HÉRAULT B., AUBRY-KIENTZ M., MOLINO J.-F. & BARALOTO C. 2014. — Estimating tropical tree diversity indices from forestry surveys: A method to integrate taxonomic uncertainty. *Forest Ecology and Management* 328 (0): 270-281. <https://doi.org/10.1016/j.foreco.2014.05.045>
- GUITET S., SABATIER D., BRUNAUX O., COUTERON P., DENIS T., FREYCON V., GONZALEZ S., HÉRAULT B., JAOUEN G., MOLINO J.-F., PÉLISSIER R., RICHARD-HANSEN C. & VINCENT G. 2018. — Disturbance regimes drive the diversity of regional floristic pools across Guianan rainforest landscapes. *Scientific Reports* 8 (1): 3872. <https://doi.org/10.1038/s41598-018-22209-9>
- HAMILTON W. 1825. — *Prodromus plantarum indiae occidentalis*. Treuttel & Würtz, London, 67 p. <https://doi.org/10.5962/bhl.title.426>
- HARIPERSAUD P., TER STEEGE H., GRANVILLE J.-J. DE, CHEVILLOTTE H. & HOFF M. 2010. — Species abundance, distribution and diversity in time and space after centuries of botanical collecting in the Guianas. *Taxon* 59 (2): 592-597. <https://doi.org/10.1002/tax.592024>
- HAYDEN W. J. 1990. — Notes on neotropical *Amanoa* (Euphorbiaceae). *Brittonia* 42 (4): 260. <https://doi.org/10.2307/2806814>
- HEKKING W. H. A. 1988. — *Violaceae*. Part I. *Rinorea* and *Rinoreocarpus*. The New York Botanical Garden Press, Bronx, 207 p. (Flora Neotropica Monograph; 46).
- HÉRITIER P. 2011. — *Le climat guyanais: petit atlas climatique de la Guyane française*. Matoury, 57 p.
- HIEPKO P. 1993. — *Olaraceae*. Koeltz Scientific Books, Koenigstein, Germany (Flora of the Guianas. Series A, Phanerogams; Fascicle 14): 3-35.
- HOCHREUTINER B. P. G. 1919. — *Guttiferae novae vel minus cognitae*. *Annuaire du Conservatoire et du Jardin botaniques de Genève* (21): 49-68.
- HOLST B. K. 2003. — Myrtaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 7. Missouri Botanical Garden Press, Saint Louis: 54-73.
- HOOKER J. D. 1867. — Rosaceae, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 14(2). F. Fleischer, Leipzig: 1-76.
- HOWARD R. A. 1983. — The plates of Aublet's *Histoire des plantes de la Guyane françoise*. *Journal of the Arnold Arboretum* 64 (2): 255-292. <https://www.jstor.org/stable/43782107>

- HUBER J. 1909. — Novitates Floraes Amazonicae. *Boletim do Museu Goeldi de Historia Natural e Ethnographia* 6: 60-90.
- HUMBOLDT F. W. H. A. VON, BONPLAND A. J. A. & KUNTH K. S. 1822. — *Nova genera et species plantarum. Quarto ed. Vol. 5* (21-22). N. Maze, Paris, 176 p.
- HUMBOLDT F. W. H. A. VON, BONPLAND A. J. A. & KUNTH K. S. 1824. — *Nova genera et species plantarum. Quarto ed. Vol. 6* (26-29). Gide fils, Paris, 374 p.
- HURault J. 1965. — *La vie matérielle des Noirs Réfugiés Boni et des Indiens Wayana du Haut-Maroni*. ORSTOM, Paris, 142 p. (Mémoires ORSTOM; 3).
- IDIGBIO 2022. — *Integrated Digitized Biocollections*. Available from <https://www.idigbio.org/> [accessed 21 February 2022].
- ILDIS 2018. — *International Legume Database & Information Service. Version 10.01*. Available from <https://ildis.org/LegumeWeb/> [accessed 11 December 2020].
- INDEX HERBARIORUM 2020. — Available from <http://sweetgum.nybg.org/science/ih/> [accessed 1 October 2020].
- IPNI 2022. — *International Plant Names Index*. Available from <http://www.ipni.org> [accessed 16 February 2022].
- IUCN 2021. — *The IUCN Red List of Threatened Species. Version 2021-2*. Available from <https://www.iucnredlist.org/> [accessed 7 June 2021].
- JACQUIN N. J. 1763. — *Nicolai Josephi Jacquin Selectarum stirpium Americanarum historia*. Kraus, Vienna, 379 p.
- JACQUIN N. J. 1780. — *Nicolai Josephi Jacquin Selectarum stirpium Americanarum historia, ed. 1780-1781*. Kraus, Vienna, 379 p.
- JANSEN-JACOBS M. J. 1988. — *Verbenaceae*. Koeltz Scientific Books, Koenigstein, Germany, 116 p. (Flora of the Guianas. Series A, Phanerogams; Fascicle 4).
- JSTOR 2022. — Global Plants. Available from <https://plants.jstor.org/> [accessed 27 April 2022].
- JISSIEU A. DE 1830. — Mémoire sur le groupe des Méliacées. *Mémoires du Muséum d'histoire naturelle* 19: 153-304. <https://www.biodiversitylibrary.org/page/26229796>
- KALLUNKI J. A. 1992. — A revision of Erythrociton sensu lato (Cuspariinae, Rutaceae). *Brittonia* 44 (2): 107-139. <https://doi.org/10.2307/2806828>
- KALLUNKI J. A. 1998. — Revision of Ticorea Aubl. (Rutaceae, Galipeinae). *Brittonia* 50 (4): 500-513. <https://doi.org/10.2307/2807761>
- KALLUNKI J. A. 2005. — Rutaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 9. Missouri Botanical Garden Press, Saint Louis: 1-39.
- KEARNS D. M. 1998. — Crudia, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 48-49.
- KERCHOVE O. DE D. 1878. — *Les palmiers; histoire iconographique, géographie – paléontologie – botanique – description – culture – emploi, etc., avec index général des noms et synonymes des espèces connues [...] Vol. 1*. J. Rothschild, Paris, 443 p.
- KLOOS P. 1971. — *The Maroni River Caribs of Surinam*. Royal van Gorcum, Assen, The Nederlands, 303 p.
- KUBITZKI K. 1978. — *Caraipa* and *Mahurea* (Bonnetiaceae), in MAGUIRE B. (ed.), *The Botany of the Guyana Highland*. Part X: 82-138 (Memoirs of the New York Botanical Garden; 29).
- KUBITZKI K. 1982. — Aniba, in KUBITZKI K. & RENNER S. (eds), *Lauraceae I* (Aniba and Aiouea). The New York Botanical Garden Press, Bronx (Flora Neotropica Monograph; 31): 1-84.
- LAMARCK J.-B. P. A. 1789. — *Encyclopédie méthodique. Botanique*. Vol. 3. Panckoucke, Liège, 759 p.
- LANDRUM L. R. 1986. — Campomanesia, Pimenta, Blepharocalyx, Legrandia, Acca, Myrrhinium, and Luma (Myrtaceae). The New York Botanical Garden Press, Bronx, 179 p. (Flora Neotropica Monograph; 45).
- LANJOUW J. & UTTIEN H. 1940. — Un nouvel herbier de Fusée Aublet découvert en France. *Recueil des Travaux botaniques néerlandais* 37: 133-170.
- LEEUWENBERG A. J. M. 1994. — *A Revision of Tabernaemontana 2: the New World Species and Stemmadenia*. Royal Botanic Gardens, Kew, 450 p. (Series of Revisions of Apocynaceae; 36).
- LEMÉE A. 1953. — *Flore de la Guyane française. Tome III. Dilléniacées à Composées*. Paul Lechevallier, Paris, 655 p.
- LEVIS C., COSTA F. R. C., BONGERS F., PEÑA-CLAROS M., CLEMENT C. R., JUNQUEIRA A. B., NEVES E. G., TAMANHA E. K., FIGUEIREDO F. O. G., SALOMÃO R. P., CASTILHO C. V., MAGNUSSON W. E., PHILLIPS O. L., GUEVARA J. E., SABATIER D., MOLINO J.-F., LÓPEZ D. C., MENDOZA A. M., PITMAN N. C. A., DUQUE A., VARGAS P. N., ZARTMAN C. E., VASQUEZ R., ANDRADE A., CAMARGO J. L., FELDPAUSCH T. R., LAURANCE S. G. W., LAURANCE W. F., KILLEEN T. J., NASCIMENTO H. E. M., MONTERO J. C., MOSTACEDO B., AMARAL I. L., GUIMARÃES VIEIRA I. C., BRIENEN R., CASTELLANOS H., TERBORGH J., CARIM M. DE J. V., GUIMARÃES J. R. DA S., COELHO L. DE S., MATOS F. D. DE A., WITTMANN F., MOGOLLÓN H. F., DAMASCO G., DÁVILA N., GARCÍA-VILLACORTA R., CORONADO E. N. H., EMILIO T., FILHO D. DE A. L., SCHIETTI J., SOUZA P., TARGHETTA N., COMISKEY J. A., MARIMON B. S., MARIMON B.-H., NEILL D., ALONSO A., ARROYO L., CARVALHO F. A., DE SOUZA F. C., DALLMEIER F., PANSONATO M. P., DUVENVOORDEN J. F., FINE P. V. A., STEVENSON P. R., ARAUJO-MURAKAMI A., AYMARD C. G. A., BARALOTO C., DO AMARAL D. D., ENGEL J., HENKEL T. W., MAAS P., PETRONELLI P., REVILLA J. D. C., STROPP J., DALY D., GRIBEL R., PAREDES M. R., SILVEIRA M., THOMAS-CAESAR R., BAKER T. R., DA SILVA N. F., FERREIRA L. V., PERES C. A., SILMAN M. R., CERÓN C., VALVERDE F. C., DI FIORE A., JIMENEZ E. M., MORA M. C. P., TOLEDO M., BARBOSA E. M., BONATES L. C. DE M., ARBOLEDA N. C., FARIA E. DE S., FUENTES A., GUILLAUMET J.-L., JØRGENSEN P. M., MALHI Y., DE ANDRADE MIRANDA I. P., PHILLIPS J. F., PRIETO A., RUDAS A., RUSCHEL A. R., SILVA N., VON HILDEBRAND P., VOS V. A., ZENT E. L., ZENT S., CINTRA B. B. L., NASCIMENTO M. T., OLIVEIRA A. A., RAMIREZ-ANGULO H., RAMOS J. F., RIVAS G., SCHÖNGART J., SIERRA R., TIRADO M., VAN DER HEIJDEN G., TORRE E. V., WANG O., YOUNG K. R., BAIDER C., CANO A., FARFAN-RIOS W., FERREIRA C., HOFFMAN B., MENDOZA C., MESONES I., TORRES-LEZAMA A., MEDINA M. N. U., VAN ANDEL T. R., VILLAROEL D., ZAGT R., ALEXIADES M. N., BALSLEV H., GARCIA-CABRERA K., GONZALES T., HERNANDEZ L., HUAMANTUPA-CHUQUIMACO I., MANZATTO A. G., MILLIKEN W., CUENCA W. P., PANSINI S., PAULETTO D., AREVALO F. R., REIS N. F. C., SAMPAIO A. F., GIRALDO L. E. U., SANDOVAL E. H. V., GAMARRA L. V., VELA C. I. A. & TER STEEGE H. 2017. — Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science* 355 (6328): 925-931. <https://doi.org/10.1126/science.aal0157>
- LIMA A. C. & AYMARD C. G. A. 1999. — *Diplotropis*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 315-318.
- LIMA D. F., LUCAS E. J., LOURENÇO A. R. L. & MAZINE F. F. 2020. — New names in *Myrcia* sect. *Calyptranthes* (Myrtaceae). *Phytotaxa* 433 (3): 251-252. <https://doi.org/10.11646/phytotaxa.433.3.7>
- LINNAEUS C. 1753. — *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*. Vol. 1. Lars Salvius, Stockholm (Holmia), 560 p. <https://doi.org/10.5962/bhl.title.37656>
- LINNAEUS C. VON 1763. — *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*, ed. 2. Vol. 2. Lars Salvius, Stockholm (Holmia), 1684 p. + index.
- LINNAEUS C. VON 1764. — *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*, ed. 3. Vol. 2. J. T. Trattner, Vienna, 1682 p. + index.

- LORENZI H. 2002. — *Arvores Brasileiras: Manual de Identificação e Cultivo de plantas arbóreas nativas do Brasil*. Instituto Plantarum de Estudos da Flora Ltda, Nova Odessa, SP, 2 vols.
- LOURENÇO A. R. L., PARRA-O C., SÁNCHEZ-CHÁVEZ E. & LUCAS E. 2018. — New combinations and names for continental American *Calyptranthes* (Myrtaceae: Myrcia s.l.). *Phytotaxa* 373 (1): 71-85. <https://doi.org/10.11646/phytotaxa.373.1.3>
- LPWG, AZANI N., BABINEAU M., BAILEY C. D., BANKS H., BARBOSA A. R., PINTO R. B., BOATWRIGHT J. S., BORGES L. M., BROWN G. K., BRUNEAU A., CANDIDO E., CARDOSO D., CHUNG K.-F., CLARK R. P., CONCEIÇÃO A. DE S., CRISP M., CUBAS P., DELGADO-SALINAS A., DEXTER K. G., DOYLE J. J., DUMINIL J., EGAN A. N., ESTRELLA M. DE LA, FALCÃO M. J., FILATOV D. A., FORTUNA-PEREZ A. P., FORTUNATO R. H., GAGNON E., GASSON P., RANDO J. G., TOZZI A. M. G. DE A., GUNN B., HARRIS D., HASTON E., HAWKINS J. A., HERENDEEN P. S., HUGHES C. E., IGANCI J. R. V., JAVADI F., KANU S. A., KAZEMPOUR-OSALOO S., KITE G. C., KLITGAARD B. B., KOCHANOVSKI F. J., KOENEN E. J. M., KOVAR L., LAVIN M., LE ROUX M. LE, LEWIS G. P., LIMA H. C. DE, LÓPEZ-ROBERTS M. C., MACKINDER B., MAIA V. H., MALÉCOT V., MANSANO V. F., MARAZZI B., MATTAPHA S., MILLER J. T., MILLER C., MOURA T., MURPHY D. J., NAGESWARA-RAO M., NEVADO B., NEVES D., OJEDA D. I., PENNINGTON R. T., PRADO D. E., PRENNER G., QUEIROZ L. P. DE, RAMOS G., RANZATO FILARDI F. L., RIBEIRO P. G., RICO-ARCE M. DE L., SANDERSON M. J., SANTOS-SILVA J., SÃO-MATEUS W. M. B., SILVA M. J. S., SIMON M. F., SINOU C., SNAK C., SOUZA É. R. DE, SPRENT J., STEELE K. P., STEIER J. E., STEEVES R., STIRTON C. H., TAGANE S., TORKE B. M., TOYAMA H., CRUZ D. T. DA, VATANPARAST M., WIERINGA J. J., WINK M., WOJCIECHOWSKI M. F., YAHARA T., YI T. & ZIMMERMAN E. 2017. — A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny. *Taxon* 66 (1): 44-77. <https://doi.org/10.12705/661.3>
- LUCAS E. J., AMORIM B. S., LIMA D. F., LIMA-LOURENÇO A. R., NIC LUGHADHA E. M., PROENÇA C. E. B., ROSA P. O., ROSÁRIO A. S., SANTOS L. L., SANTOS M. F., SOUZA M. C., STAGGEMEIER V. G., VASCONCELOS T. N. C. & SOBRAL M. 2018. — A new infra-generic classification of the species-rich Neotropical genus *Myrcia* s.l. *Kew Bulletin* 73 (1): 9. <https://doi.org/10.1007/s12225-017-9730-5>
- MAAS P. J. M. & MAAS-VAN DE KAMER H. 1993. — *Haemodoraceae*. The New York Botanical Garden Press, Bronx, 44 p. (Flora Neotropica Monograph; 61).
- MAAS P. J. M. & MAAS-VAN DE KAMER H. 2002. — Annonaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx (Memoirs of the New York Botanical Garden; 76 [2]): 53-67.
- MAAS P. J. M. & WESTRA L. Y. T. 1985. — Studies in Annonaceae. II. A monograph of the genus *Anaxagorea* A. St. Hil. Part 1. *Botanische Jahrbücher Systematik* 105 (1): 73-134.
- MAAS P. J. M. & WESTRA L. Y. T. 2011. — A taxonomic survey of Guatteria section Mecocarpus including the genera Guatteriopsis and Guatteriella p.p. (Annonaceae). *Blumea* 56 (2): 113-145. <https://doi.org/10.3767/000651911X588844>
- MAAS P. J. M., WESTRA L. Y. T., BROWN K. S., TER WELLE B. J. H., WEBBER A. C., LE THOMAS A., WAHA M., HEIJDEN E. VAN DER, BOUMAN F., CAVÉ A., LEBOEUF M., LAPRÉVOTE O., KOEK-NOORMAN J., MORAWETZ W. & HEMMER W. 1992. — *Rollinia* (Annonaceae). The New York Botanical Garden Press, Bronx, 188 p. (Flora Neotropica Monograph; 57).
- MAAS P. J. M., MAAS-VAN DE KAMER H., JUNIKKA L., MELLO-SILVA R. DE & RAINER H. 2001. — Annonaceae from Central-eastern Brazil. *Rodriguésia* 52 (80): 65-98. <http://dx.doi.org/10.1590/2175-78602001528005>
- MAAS P. J. M., WESTRA L. Y. T., CHATROU L. W., BAKKER M. A., KOEK-NOORMAN J., TER WELLE B. J. H., BENKO-ISEPPON A. M., GOTTSBERGER G., JUNIKKA L. M. K., LE THOMAS A., MORAWETZ W., ROBLOT F., HOCQUEMILLER R., LEBOEUF M., CAVÉ A., ROOSMALEN M. G. M. VAN, SVOMA E. & WEBBER A. C. 2003. — *Duguetia* (Annonaceae). The New York Botanical Garden Press, Bronx, 274 p. (Flora Neotropica Monograph; 88).
- MAAS P. J. M., MAAS-VAN DE KAMER H., MIRALHA J. M. S. & JUNIKKA L. 2007a. — Flora da Reserva Ducke, Amazonas, Brasil: Annonaceae. *Rodriguésia* 58 (3): 617-662. <https://doi.org/10.1590/2175-7860200758307>
- MAAS P. J. M., WESTRA L. Y. T. & VERMEER M. 2007b. — Revision of the Neotropical genera *Bocageopsis*, *Onychopetalum*, and *Unonopsis* (Annonaceae). *Blumea* 52 (3): 413-554. <https://doi.org/10.3767/000651907X608909>
- MADRINAN S. 2004. — *Rhodostemonodaphne* (Lauraceae). The New York Botanical Garden Press, Bronx, 102 p. (Flora Neotropica Monograph; 92).
- MALÉCOT V. & NICKRENT D. L. 2008. — Molecular phylogenetic relationships of Olacaceae and related santalales. *Systematic Botany* 33 (1): 97-106. <https://doi.org/10.1600/036364408783887384>
- MANSANO V. DE F., TOZZI A. M. G. DE A. & LEWIS G. P. 2004. — A revision of the South American genus *Zollernia* Wied-Neuw. & Nees (Leguminosae, Papilionoideae, Swartzieae). *Kew Bulletin* 59 (4): 497-520. <https://doi.org/10.2307/4110905>
- MARCANO-BERTI L. 1998. — *Vochysiaceae*. Royal Botanic Garden, Kew (Flora of the Guianas. Series A, Phanerogams; Fascicle 21): 1-44.
- MARCHAND L. 1869. — *Révision du groupe des anacardiacées*. Bailière & Fils, Paris, 191 p.
- MARINHO L. C., FIASCHI P., GAHAGEN B., SANTOS F. DE A. R. DOS & AMORIM A. M. 2016. — *Tovomita* (Clusiaceae) from the Brazilian Atlantic forest: Taxonomy and utility of leaf venation characters at the species level. *Systematic Botany* 41 (3): 758-774. <https://doi.org/10.1600/036364416x692406>
- MARINHO L. C., CAI L., DUAN X., RUHFEL B. R., FIASCHI P., AMORIM A. M., VAN DEN BERG C. & DAVIS C. C. 2019. — Plastomes resolve generic limits within tribe Clusiae (Clusiaceae) and reveal the new genus Arawakia. *Molecular Phylogenetics and Evolution* 134: 142-151. <https://doi.org/10.1016/j.ympev.2019.02.005>
- MARTIUS C. F. P. VON 1824a. — *Historia naturalis palmarum*. Vol. 2. Weigel, Leipzig, 152 p. <https://www.biodiversitylibrary.org/page/281694>
- MARTIUS C. F. P. VON 1824b. — *Nova genera et species plantarum*. Vol. 1. Lindauer, München, 158 p. <https://www.biodiversitylibrary.org/page/281352>
- MAUREL D., CACHINE J.-M., COUCHILI L., MONERVILLE E. & PANAPUY J. 2020. — *Kaleta Teko'awu – Panaits'i awu ipope maë*. *Dictionnaire Teko-Français*. Available from <https://corporan.huma-num.fr/Lexiques/dicoTeko.html> [accessed 15 July 2021].
- MCVAUGH R. 1958. — Myrtaceae. Flora of Peru. *Publications of the Field Museum of Natural History, Botanical Series* 13 (4, 2): 569-818.
- MCVAUGH R. 1969. — The Botany of the Guyana Highland, Part VIII. Myrtaceae. *Memoirs of the New York Botanical Garden* 18 (2): 55-286.
- MENNEGA A. M. W. & HEDIN J. P. 1999. — Hippocrateaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 594-617.
- MEYER G. F. W. 1818. — *Primitiae florae Essequeboensis adjectis descriptionibus centum circiter stirpium novarum, observationibusque criticis*. H. Dieterich, Göttingen, 316 p.
- MEZ C. 1889. — Lauraceae Americanae monographice descriptae. *Jahrbuch des Königlichen Botanischen Gartens und des Botanischen Museums zu Berlin* 5: 1-556.
- MIERS J. 1878. — *On the Apocynaceae of South America*. Williams & Norgate, London, 291 p. <https://doi.org/10.5962/bhl.title.54802>

- MIQUEL F. A. W. 1848. — *Symbolae ad floram Surinamensem. Linnaea* (21): 473-479.
- MIQUEL F. A. W. 1851. — *Stirpes Surinamenses selectae*. Arnz. & Soc., Leiden, 234 p.
- MIQUEL F. A. W. 1863. — Sapoteae, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis*. Vol. 7. F. Fleischer, Leipzig: 37-118.
- MITCHELL J. D. 1995. — Anacardiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 2. Missouri Botanical Garden Press, Saint Louis: 291-300.
- MITCHELL J. D. 2002. — Burseraceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx (Memoirs of the New York Botanical Garden; 76[2]): 605-606.
- MOLINO J.-F., SABATIER D., PRÉVOST M.-F., FRAME D., GONZALEZ S. & BILOT-GUÉRIN V. 2009. — *Établissement d'une liste des espèces d'arbres de la Guyane Française. Rapport final de la Convention E 24/08 entre le Ministère de l'Agriculture et de la Pêche et l'IRD, dans le cadre du Programme 149 Forêt BOP 14901C, sous-action 15*. IRD, Cayenne, 59 p.
- MOLINO J.-F., BALEÉ W., ENGEL J., MARTIN C. A. & SABATIER D. 2021. — Forest tree inventories, in ODONNE G. & MOLINO J.-F. (eds), *Methods in Historical Ecology – Insights from Amazonia*. Routledge, London: 94-103.
- MORAES P. L. R. DE 2005. — Lectotypification of names of Brazilian species of *Cryptocarya* (Lauraceae). *Taxon* 54 (3): 789-795. <https://doi.org/10.2307/25065435>
- MORALES J. F. 2005. — Estudios en las Apocynaceae Neotropicales XIX: la familia Apocynaceae s. str. (Apocynoideae, Rauvolfioideae) de Costa Rica. *Darwiniana* 43 (1-4): 90-191. <https://doi.org/10.14522/darwiniana.2014.431-4.139>
- MORALES J. F. 2009. — La familia Apocynaceae (Apocynoideae, Rauvolfioideae) en Guatemala. *Darwiniana* 47 (1): 140-184. <https://doi.org/10.14522/darwiniana.2014.471.50>
- MORALES J. F. & ZAMORA N. A. 2017. — A synopsis of Aspidosperma (Apocynaceae) in Mexico and Central America with a taxonomic clarification of Aspidosperma cruentum and a new cryptic species. *Phytoneuron* (68): 1-13.
- MORI S. A. & FISCHER B. 2002. — Flacourtiaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, 319-328 (Memoirs of the New York Botanical Garden; 76 [2]): 206-298.
- MORI S. A. & PRANCE G. T. 1993. — *Lecythidaceae*. Royal Botanic Garden, Kew (Flora of the Guianas. Series A, Phanerogams; Fascicle 12): 1-141.
- MORI S. A., PRANCE G. T. & ZEEUW C. H. DE 1990. — *Lecythidaceae*. Part II. *The Zygomorphic-Flowered New World Genera (Couroupita, Corythophora, Bertholletia, Couratari, Eschweilera, & Lecythis), with a Study of Secondary Xylem of Neotropical Lecythidaceae*. The New York Botanical Garden Press, Bronx. 373 p. (Flora Neotropica Monograph; 21 [2]).
- MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. 2002. — *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx. 944 p. (Memoirs of the New York Botanical Garden; 76 [2]).
- MORI S. A., KIERNAN E. A., SMITH N. P., KELLY L. M., HUANG Y.-Y., PRANCE G. T. & THIERS B. 2017. — Observations on the phytogeography of the Lecythidaceae clade (Brazil nut family). *Phytoneuron* 2017-30: 1-86.
- MORLEY T. 1976. — *Memecyleae (Melastomataceae)*. The New York Botanical Garden Press, Bronx. 295 p. (Flora Neotropica Monograph; 15).
- MÜLLER J. 1881. — Rubiaceae, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 6 (5). F. Fleischer, Leipzig: 1-470.
- MÜLLER J. 1888. — Rubiaceae - Index, in MARTIUS C. F. P. VON (ed.), *Flora Brasiliensis* 6 (5). F. Fleischer, Leipzig: 471-486.
- MUNZINGER J. K. & BALLARD H. E. 2003. — *Hekkingia* (Violaceae), a new arborescent violet genus from French Guiana, with a key to genera in the family. *Systematic Botany* 28 (2): 345-351. <https://doi.org/10.1043/0363-6445-28.2.345>
- NEILL D. A. 1999. — *Erythrina*, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 322-324.
- NIEDENZU F. 1928. — Fam. IV.141. *Malpighiaceae*. Pars III, in ENGLER H. G. A. (ed.), *Das Pflanzenreich. Regni vegetabilis conspectus*. Vol. 94. W. Engelmann, Leipzig: 573-870.
- NOWICKE J. W. 1969. — Boraginaceae, in WOODSON R. W., SCHERY R. W. et al. (eds), *Flora of Panama*. Part IX. *Annals of the Missouri Botanical Garden* 56 (1): 33-69.
- ODONNE G., TAREAU M.-A. & VAN ANDEL T. 2020. — Geopolitics of bitterness: Deciphering the history and cultural biogeography of *Quassia amara* L. *Journal of Ethnopharmacology*: 113546. <https://doi.org/10.1016/j.jep.2020.113546>
- ODERON C., ODONNE G., CRISTINOI A., ENGEL J., GRENDAND P., BEAUCHÈNE J., CLAIR B. & DAVY D. 2018. — Palikur traditional roundwood construction in eastern French Guiana: ethnobotanical and cultural perspectives. *Journal of Ethnobiology and Ethnomedicine* 14 (1): 28. <https://doi.org/10.1186/s13002-018-0226-7>
- OLSON M., BERRY P. E. & AYMAR C. G. A. 1999. — Flacourtiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 434-474.
- ONF 2004. — *Guide de Reconnaissance des Arbres de Guyane*. 2nd ed. Office national des forêts, Cayenne, 374 p.
- PENDRY C. A. 2004. — Monograph of Ruprechtia (Polygonaceae). *Systematic Botany Monographs* 67: 1-113. <https://doi.org/10.2307/25027911>
- PENNINGTON T. D. 1990. — *Sapotaceae*. The New York Botanical Garden Press, Bronx, 770 p. (Flora Neotropica Monograph; 52).
- PENNINGTON T. D. 1997. — *The genus Inga – Botany*. Royal Botanic Gardens, Kew, Kew, UK, 844 p.
- PENNINGTON T. D. 2016. — *Prodromus Sloanearium Americanarum*. David Hunt, Sherborne, UK, 24 p. (Opuscula Neotropica; 2/2016).
- PENNINGTON T. D. & CLARKSON J. J. 2013. — A revision of Guarea (Meliaceae). *Edinburgh Journal of Botany* 70 (2): 179-362. <https://doi.org/10.1017/S0960428613000036>
- PENNINGTON T. D. & WISE R. 2017. — *The genus Sloanea (Elaeocarpaceae) in America*. DH Books, Milborne Port, United Kingdom, 428 p.
- PENNINGTON T. D., STYLES B. T. & TAYLOR D. A. H. 1981. — *Meliaceae, with Accounts of Swietenioideae and Chemotaxonomy*. The New York Botanical Garden Press, Bronx, 470 p. (Flora Neotropica Monograph; 28).
- PERSSON C. & DELPRETE P. G. 2010. — Cordiera. Rubiaceae – Parte 1: Introdução, Gêneros A-H, in RIZZO J. A. (ed.), *Flora dos Estados de Goiás e Tocantins*. Vol. 40. IRD/Universidade Federal de Goiás, Goiânia: 212-265.
- PERSSON C. & DELPRETE P. G. 2017. — *The Alibertia Group (Gardenieae-Rubiaceae)*. Part I (Agouticarpa, Alibertia, Cordiera, Melanopsisidium, and Stenosepala). The New York Botanical Garden Press, Bronx, 241 p. (Flora Neotropica Monographs; 119).
- PIERRE J. B. L. 1885. — Plantes à Gutta-percha. *Bulletin mensuel de la Société linnaéenne de Paris* 1 (63): 497-499.
- PIO CORRÊA M. 1926-1975. — *Dicionário das plantas úteis do Brasil e das exóticas cultivadas*. Imprensa Nacional, Rio de Janeiro, 6 vol.
- PIPOLY J. J. I. & GUSTAFSSON M. H. G. 2002. — Apocynaceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx (Memoirs of the New York Botanical Garden; 76 [2]): 212-224..

- PIPOLY J. J. I., KEARNS D. M. & BERRY P. E. 1998. — Clusiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 243-277.
- PIRANI J. R. 2005. — Picramniaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 9. Missouri Botanical Garden Press, Saint Louis: 177-179.
- PIRANI J. R. & KALLUNKI J. A. 2007. — Two new species of *Galipea* (Rutaceae, Galipeae) from Bolivia, Ecuador, and Peru. *Brittonia* 59 (4): 343-349.
- PLANCHON J. É. & TRIANA J. J. 1860a. — Mémoire sur la famille des Guttifères. 1. *Annales des Sciences naturelles, Botanique*, sér. 4, 13: 306-376.
- PLANCHON J. É. & TRIANA J. J. 1860b. — Mémoire sur la famille des Guttifères. 2. *Annales des Sciences Naturelles, Botanique* sér. 4 14: 226-367.
- POWO 2022. — *Plants of the World Online*. Published on the Internet. Available from <https://powo.science.kew.org/> [accessed 15 April 2022].
- PLOWMAN T. & BERRY P. E. 1999. — Erythroxylaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 59-71.
- PLOWMAN T. & HENSOLD N. 2004. — Names, types, and distribution of neotropical species of *Erythroxylum* (Erythroxylaceae). *Brittonia* 56 (1): 1-53.
- PLUMEL M. M. 1991. — Le genre *Himatanthus* (Apocynaceae): révision taxonomique. *Bradea* 5(Suppl.): 1-118.
- POIRET J. L. M. 1798. — *Encyclopédie méthodique. Botanique, par le citoyen Lamarck*. Vol. 4. Agasse, Paris, 764 p. <https://doi.org/10.5962/bhl.title.824>
- PONCY O. 1985. — Le genre *Inga* (Légumineuses, Mimosoideae) en Guyane française. *Mémoires du Muséum national d'Histoire naturelle*, série B, *Botanique* 31: 1-124. <https://www.biodiversitylibrary.org/page/58138295>
- PONCY O. 2007. — A new species of *Inga* Mill. (Fabaceae, Mimosoideae) from the Guianas. *Adansonia*, sér. 3, 29 (2): 249-254.
- PONCY O. & OFFROY B. 2006. — Une nouvelle espèce de *Tovomita* Aubl. (Clusiaceae) de Guyane française. *Adansonia*, sér. 3, 28 (1): 113-117.
- PRANCE G. T. 1972. — *Chrysobalanaceae*. The New York Botanical Garden Press, Bronx, 409 p. (Flora Neotropica Monograph; 9).
- PRANCE G. T. 1981. — Notes on *Couepia* and *Hirtella* (Chrysobalanaceae). *Brittonia* 33 (3): 347-356. <https://doi.org/10.2307/2806425>
- PRANCE G. T. 1986. — *Chrysobalanaceae*. Royal Botanic Garden, Kew (Flora of the Guianas. Series A, Phanerogams; Fascicle 19): 1-146.
- PRANCE G. T. 1992. — New Species and New Records of Neotropical Chrysobalanaceae. *Kew Bulletin* 47 (4): 633-646. <https://doi.org/10.2307/4110701>
- PRANCE G. T. 1998. — Chrysobalanaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 187-232.
- PRANCE G. T. 2009. — Proteaceae. Royal Botanic Garden, Kew (Flora of the Guianas. Series A, Phanerogams; Fascicle 27): 18-40.
- PRANCE G. T. 2018. — Rhizophoraceae. The New York Botanical Garden Press, Bronx, 61 p. (Flora Neotropica Monograph; 120).
- PRANCE G. T. & SILVA M. F. DA 1973. — Caryocaraceae. The New York Botanical Garden Press, Bronx, 75 p. (Flora Neotropica Monograph; 12).
- PRANCE G. T. & SOTHERS C. A. 2003a. — *Chrysobalanaceae I: Chrysobalanus to Parinari*. Australian Biological Resources Study, Canberra, 319 p. (Species Plantarum: Flora of the World; 9).
- PRANCE G. T. & SOTHERS C. A. 2003b. — *Chrysobalanaceae II: Acioa to Magnistipula*. Australian Biological Resources Study, Canberra, 268 p. (Species Plantarum: Flora of the World; 10).
- PROENÇA C. E. B., FARIA J. E. Q., GIARETTA A., LUCAS E. J., STAGGEMEIER V. G., TULER A. C. & VASCONCELOS T. N. C. 2020. — Nomenclatural and taxonomic changes in tribe Myrtae (Myrtaceae) spurred by molecular phylogenies. *Herlingeriana* 14 (1): 49-61.
- RADLKOFER L. A. T. 1878. — Ueber *Sapindus* und damit in Zusammenhang stehende Pflanzen. *Sitzungsberichte der Mathematisch-Physikalischen Classe der Königlich Bayerische Akademie der Wissenschaften zu München* 8: 221-408.
- REDDEN K. M., HERENDEEN P. S. & LEWIS G. P. 2018. — *Understanding Paloue (Leguminosae: Detarioideae)*. Revision of a predominantly Guiana Shield endemic. Smithsonian Institution Press, Washington, D.C., 44 p. (Smithsonian Contributions to Botany; 109).
- RENAULT-LESCURE O. 2009. — La langue kali'na, in RENAULT-LESCURE O. & GOURY L. (eds), *Langues de Guyane, cultures en Guyane*. Vents d'ailleurs/IRD, Marseille: 66-77.
- RENAULT-LESCURE O. & GOURY L. (eds) 2009. — *Langues de Guyane, cultures en Guyane*. Vents d'ailleurs/IRD, Marseille, 188 p.
- RENNER S. S. 1982. — Aiouea, in KUBITZKI K. & RENNER S. (eds), *Lauraceae I (Aniba and Aiouea)*. The New York Botanical Garden Press, Bronx (Flora Neotropica Monograph; 31): 85-116.
- RENNER S. S. 1989. — Systematic studies in the Melastomataceae: *Bellucia*, *Loreya* and *Macairea*. *Memoirs of the New York Botanical Garden* 50: 1-112.
- RENNER S. S. & HAUSNER G. 2005. — *Siparunaceae*. The New York Botanical Garden Press, Bronx, 247 p. (Flora Neotropica Monograph; 95).
- RIBEIRO J. E. L. D. S., HOPKINS M. J. G., VICENTINI A., SOTHERS C. A., COSTA M. A. DA S., BRITO J. M. DE, SOUZA M. A. D. DE, MARTINS L. H. P., LOHMANN L. G., ASSUNÇÃO P. A. C. L., PEREIRA E. DA C., SILVA C. F. DA, MESQUITA M. R. & PROCOPPIO L. C. 1999. — *Flora da Reserva Ducke: Guia de identificação das plantas vasculares de uma floresta de terra-firme na Amazônia Central*. INPA/DFID, Manaus, 800 p.
- RIBEIRO V. M. L., LIMA COSTA E. DE & LIMA BARROSO M. A. 1979. — Catalógo de nomes científicos e vulgares de plantas de porte arbóreo ocorrentes no Brasil. *Rodriguésia* 31 (49): 155-233.
- RICHARD L. C. 1830. — *Mémoire sur la famille des Rubiaceae*. J. Tastu, Paris, 224 p.
- RICKETSON J. M. & PIPOLY J. J. I. 2010. — A Synopsis of Neotropical *Stylogyne* (Myrsinaceae). *Novon* 20 (4): 437-447. <https://doi.org/10.3417/2010027>
- RJB 2022. — *Biblioteca Digital del Real Jardín Botánico*. Available from <https://bibdigital.rjb.csic.es> [accessed 10 January 2022].
- ROBYNS A. & CUATRECASAS J. 1964. — *Flora of Panamá. Part VI. Family 117, Sterculiaceae*. *Annals of the Missouri Botanical Garden* 51 (1/4): 69-107.
- RODRIGUES R. S. & TOZZI A. M. G. DE A. 2006. — *Guianodendron*, a new genus of Leguminosae (Papilionoideae) from South America. *Novon* 16 (1): 129-132.
- RODRIGUES R. S. & TOZZI A. M. G. DE A. 2012. — Revisão taxonómica de *Leptolobium* (Papilionoideae, Leguminosae). *Acta Botanica Brasilica* 26 (1): 146-164.
- ROHWER J. G. 1993. — *Lauraceae: Nectandra*. The New York Botanical Garden Press, Bronx, 332 p. (Flora Neotropica Monograph; 60).
- ROOSMALEN M. G. M. VAN 1985. — *Fruits of the Guianan Flora*. Institute of Systematic Botany, Utrecht University, Utrecht, The Netherlands, 481 p.
- SABATIER D. 1993. — Diversité des arbres et du peuplement forestier guyanais, in *Forêt Guyanaise. Gestion de l'écosystème forestier guyanais et aménagement de l'espace régional (Actes du 2e Congrès Régional de l'Environnement, Cayenne 1990)*. SEPANGUY-CCEE, Cayenne (Guyane Française): 41-47.
- SABATIER D. 1997. — Description et biologie d'une nouvelle espèce de *Virola* (Myristicaceae) de Guyane. *Adansonia*, sér. 3, 19 (2): 273-278.

- SABATIER D. 2002. — *Vantanea ovicarpa* (Humiriaceae), a new species from French Guiana. *Brittonia* 54 (4): 233-235. <https://doi.org/bv8chx>
- SAGOT P. A. 1885. — Catalogue des plantes phanérogames et cryptogames vasculaires de la Guyane française. *Annales des Sciences naturelles, Botanique*, sér. 6, 20: 181-216.
- SAINT-HILAIRE A. DE, JUSSIEU A. DE & CAMBESSEDES J. 1828. — *Plantes usuelles des Brasiliens. 14^e livraison*. Grimbert, Paris, pls 66-70.
- SAMARAKOON T. & ALFORD M. H. 2019. — New Names and Combinations in Neotropical Samydaceae. *Novon* 27 (1): 65-71. <https://doi.org/10.3417/2018307>
- SANDWITH N. Y. 1932. — Contributions to the flora of Tropical America: XII. New and noteworthy species from British Guiana, mainly collected by the Oxford University Expedition, 1929 (continued). *Bulletin of Miscellaneous Information, Royal Botanic Gardens, Kew* 5: 209-229.
- SANDWITH N. Y. 1963. — Contributions to the flora of tropical America: LXX: Notes on some Aublet types in the Paris Herbarium. *Kew Bulletin* 17 (2): 257-262. <https://doi.org/10.2307/4118953>
- SANTOS V. DE J. & SALES M. F. DE 2009. — A tribo Hippomaneae A. Juss. ex Spach. (Euphorbiaceae Juss.) no estado de Pernambuco, Brasil. *Acta Botanica Brasilica* 23 (4): 976-990.
- ASTRE C. 1986. — Deux espèces nouvelles d’Ochnaceae du Venezuela. *Bulletin du Muséum national d’Histoire naturelle, section B, Adansonia, Botanique, Phytochimie* 8 (1): 13-16. <https://www.biodiversitylibrary.org/page/13739412>
- ASTRE C. 2007. — Six nouvelles espèces d’*Ouratea* (Ochnaceae) des Guyanes. *Adansonia*, sér. 3, 29 (1): 77-91.
- ASTRE C. & OFFROY B. 2016. — Révision nomenclaturale des binômes du genre néotropical *Ouratea* Aublet (Ochnaceae) décrits par Van Tieghem. *Adansonia* 38 (1): 55-98. <https://doi.org/10.5252/a2016n1a5>
- SCHARF U., MAAS P. J. M. & MORAWETZ W. 2006. — Five new species of *Guatteria* (Annonaceae) from French Guiana, Guyana and Surinam. *Blumea* 51 (1): 117-130. <https://doi.org/10.3767/000651906X622373>
- SCHNEIDER J. V. & ZIZKA G. 2004. — Quiinaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 8. Missouri Botanical Garden Press, Saint Louis: 393-407.
- SCHNEIDER J. V. & ZIZKA G. 2012. — Taxonomic Revision of the Neotropical Genus *Lacunaria* (Quiinaceae/Ochnaceae s.l.). *Systematic Botany* 37 (1): 165-188. <https://doi.org/10.1600/036364412X616747>
- SCHNEIDER J. V. & ZIZKA G. 2016. — *Quiinaceae*. The New York Botanical Garden Press, Bronx, 162 p. (Flora Neotropica Monograph; 115).
- SCHULTES R. E. 1949. — Plantae Austro-Americanae V. De plantis principaliter Colombiae observationes. *Botanical Museum leaflets* 13 (9): 261-292.
- SECCO R. DE S. 1999. — Sagotia, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 5. Missouri Botanical Garden Press, Saint Louis: 215-217.
- SECCO R. DE S. 2004. — *Alchorneae* (Euphorbiaceae) (Alchornea, Aparisthium e Conceveiba). The New York Botanical Garden Press, Bronx, 194 p. (Flora Neotropica Monograph; 93).
- SECCO R. DE S. 2005. — Flora da Reserva Ducke, Amazonas, Brasil – Part I: Euphorbiaceae. *Rodriguésia* 56 (86): 143-168. <https://doi.org/10.1590/2175-78602005568624>
- SESSÉ M. & MOCIÓN J. M. 1892. — *Flora mexicana*. Vol. 1. I. Escalante, Mexico, 263 p.
- SIL (SUMMER INSTITUTE OF LINGUISTICS) 2003. — *Languages of Suriname*. Available from <http://www.suriname-languages.sil.org/Aukan/English/AukanEngDictIndex.html> [accessed 21 April 2020].
- SILVA M. F. DA, LISBOA P. L. B. & LISBOA R. C. L. 1977. — *Nomes vulgares de plantas amazonicas*. INPA, Manaus, 222 p.
- SILVA M. F. DA 1971. — Os Tipos do Herbário do Instituto Nacional de Pesquisas da Amazônia – I. *Acta Amazonica* 1 (2): 19-32.
- SILVA M. F. DA, TAVARES A. S. & BERRY P. E. 1998. — Cynometra, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 4. Missouri Botanical Garden Press, Saint Louis: 49-51.
- SILVA M. J. DA & TOZZI A. M. G. DE A. 2012. — Revisão taxonômica de *Lonchocarpus* s. str. (Leguminosae, Papilionoideae) do Brasil. *Acta Botanica Brasilica* 26 (2): 357-377. <https://doi.org/10.1590/S0102-33062012000200012>
- SLEUMER H. O. 1980. — *Flacourtiaceae*. The New York Botanical Garden Press, Bronx, 499 p. (Flora Neotropica Monograph; 22).
- SPINA A. P. 2004. — *Estudos taxonomico, micro-morfológico e filogenético do genero Himatanthus Willd. ex Schult. (Apocynaceae: Rauvolfioideae – Plumerieae)*, Tese de Doutorado, Universidade Estadual de Campinas, Instituto de Biologia, Campinas, SP, 191 p.
- SPINA A. P., BITTRICH V. & KINOSHITA L. S. 2013. — Typifications, new synonyms and a new combination in *Himatanthus* (Apocynaceae). *Taxon* 62 (6): 1304-1307. <https://doi.org/10.12705/626.16>
- SPRUCE R. 1853. — Edible fruits of the Rio Negro, South America. *Hooker’s Journal of Botany and Kew Garden Miscellany* 5: 183-187.
- STACE C. A. 2009. — *Combretaceae*. Royal Botanic Garden, Kew (Flora of the Guianas. Series A, Phanerogams; Fascicle 27): 41-98.
- STACE C. A. 2010. — *Combretaceae*. The New York Botanical Garden Press, Bronx, 370 p. (Flora Neotropica Monograph; 107).
- STAFLEU F. A. & COWAN R. S. 1976-1988. — *Taxonomic Literature. A Selective Guide to Botanical Publications and Collections with Dates, Commentaries and Types*. Bohn, Scheltema and Holkema, Utrecht, 7 vols.
- STAFLEU F. A. & MENNEGA E. A. 1992-2000. — *Taxonomic Literature. A Selective Guide to Botanical Publications and Collections with Dates, Commentaries and Types*. Suppl. 1-6. Koeltz Scientific Books, Königstein, 5 vols.
- STANLEY P. C. 1930. — Studies of American plants IV. *Publications of the Field Museum of Natural History, Botanical Series* 8 (3): 133-236.
- STANLEY P. C. 1937a. — Flora of Peru. *Publications of the Field Museum of Natural History, Botanical Series* 13 (2): 257-661.
- STANLEY P. C. 1937b. — Studies of American plants VII. *Publications of the Field Museum of Natural History, Botanical Series* 17 (2): 155-224.
- STEUDEL E. G. VON 1840. — *Nomenclator botanicus*, ed. 2. G. Cotta, Stuttgart, 810 p. <https://doi.org/10.5962/bhl.title.655>
- STEVENS P. F. 2019. — APweb. *The Angiosperm Phylogeny website*. Available from www.mobot.org/MOBOT/research/APweb [accessed 11 December 2020].
- STEYERMARK J. A. 1965. — Rubiaceae, in MAGUIRE B. & WURDACK J. J. (eds), *Memoirs of the New York Botanical Garden* 12 (3): 178-285.
- STEYERMARK J. A. 1974. — Rubiaceae, in LASSER T. & STEYERMARK J. A. (eds), *Flora de Venezuela*. Vol. 9. Ediciones Fundación Educación Ambiental, Caracas: 1-2070.
- STEYERMARK J. A. 2004. — Rhizophoraceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 8. Missouri Botanical Garden Press, Saint Louis: 484-490.
- STEYERMARK J. A. & CALLEJAS-POSADA R. 2003. — Piperaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 7. Missouri Botanical Garden Press, Saint Louis: 681-738.
- STEYERMARK J. A., MAAS P. J. M., BERRY P. E., JOHNSON D. M., MURRAY N. A. & RAINER H. 1995. — Annonaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 2. Missouri Botanical Garden Press, Saint Louis: 300-339.
- SWART J. J. 1942. — A monograph of the genus *Protium* and some allied genera (Burseraceae). *Recueil des Travaux notaniques néerlandais* 39: 212-446.

- TAMBOLI A. S., YADAV P. B., GOTHE A. A., YADAV S. R. & GOVINDWAR S. P. 2018. — Molecular phylogeny and genetic diversity of genus *Capparis* (Capparaceae) based on plastid DNA sequences and ISSR markers. *Plant Systematics and Evolution* 304 (2): 205-217. <https://doi.org/10.1007/s00606-017-1466-z>
- TAYLOR C. M. 1999. — 162(20). Rubiaceae-Coussareae, *Flora of Ecuador* 62. Council for Nordic Publications in Botany, Copenhagen: 245-320.
- TAYLOR C. M. & GEREAU R. E. 2019. — *Nomenclature Notes on Rubiaceae*. Available from http://www.tropicos.org/projectweb-portal.aspx?pagename=NomenclatureNotesonRubiaceae&proj_eid=34 [accessed 11 May 2020].
- TAYLOR C. M. & JARDIM J. G. 2020. — Rubiacearum Americanarum magna hama pars XLVI: New species and taxonomic changes in *Faramea* of Central and South America (Rubiaceae, Coussareae). *Novon* 28 (2): 108-142. <https://doi.org/10.3417/2019552>
- TAYLOR C. M. & STEYERMARK J. A. 2004. — Rubiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 8. Missouri Botanical Garden Press, Saint Louis: 579-587.
- TAYLOR C. M., CAMPOS M. T. V. A. & ZAPPI D. 2007. — Flora da Reserva Ducke, Amazonas, Brasil: Rubiaceae. *Rodriguésia* 58 (3): 549-616. <https://doi.org/10.1590/2175-7860200758306>
- TAYLOR C. M., SÁNCHEZ-GONZÁLEZ J., HAMMEL B., LORENCE D. H., PERSSON C., DELPRETE P. G. & GEREAU R. E. 2011. — Rubiacearum Americanarum magna hama pars XXVIII: New taxa, new combinations, new names, and lectotypification for several species found in Mexico and Central America. *Novon* 21 (1): 133-148. <https://doi.org/10.3417/2009129>
- TAYLOR C. M., STEYERMARK J. A., DELPRETE P. G., VINCENTINI A., CORTÉS R., ZAPPI D., PERSSON C., COSTA C. B. & ANUNCIAÇÃO E. 2004. — Rubiaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 8. Missouri Botanical Garden Press, Saint Louis: 497-847.
- TAYLOR E. L. 1989. — *Systematic Studies in the Tribe Sterculiae: a Taxonomic Revision of the Neotropical Species of Sterculia L. (Sterculiaceae)*, Ph.D. Thesis, Harvard University, Cambridge, MA, 508 p.
- TEIXEIRA L. A. G. & MACHADO I. C. 2000. — Pollination and reproductive system of *Byrsonima sericea* DC (Malpighiaceae). *Acta Botanica Brasiliensis* 14 (3): 347-357. <https://doi.org/10.1590/S0102-33062000000300011>
- TER STEEGE H., PRADO P. I., LIMA R. A. F. DE, POS E., DE SOUZA COELHO L., DE ANDRADE LIMA FILHO D., SALOMÃO R. P., AMARAL I. L., DE ALMEIDA MATOS F. D., CASTILHO C. V., PHILLIPS O. L., GUEVARA J. E., DE JESUS VEIGA CARIM M., CÁRDENAS LÓPEZ D., MAGNUSSON W. E., WITTMANN F., MARTINS M. P., SABATIER D., IRUME M. V., DA SILVA GUIMARÃES J. R., MOLINO J.-F., BÁNKI O. S., PIEDADE M. T. F., PITMAN N. C. A., RAMOS J. F., MONTEAGUDO MENDOZA A., VENTICINQUE E. M., LUIZE B. G., NÚÑEZ VARGAS P., SILVA T. S. F., DE LEÃO NOVO E. M. M., REIS N. F. C., TERBORGH J., MANZATTO A. G., CASULA K. R., HONORIO CORONADO E. N., MONTERO J. C., DUQUE A., COSTA F. R. C., CASTAÑO ARBOLEDA N., SCHÖNGART J., ZARTMAN C. E., KILLEEN T. J., MARIMON B. S., MARIMON-JUNIOR B. H., VASQUEZ R., MOSTACEDO B., DEMARCHI L. O., FELDPAUSCH T. R., ENGEL J., PETRONELLI P., BARALOTO C., ASSIS R. L., CASTELLANOS H., SIMON M. F., DE MEDEIROS M. B., QUARESMA A., LAURANCE S. G. W., RINCÓN L. M., ANDRADE A., SOUSA T. R., CAMARGO J. L., SCHIETTI J., LAURANCE W. F., DE QUEIROZ H. L., NASCIMENTO H. E. M., LOPES M. A., DE SOUSA FARIAS E., MAGALHÃES J. L. L., BRIENEN R., AYMARD C. G. A., REVILLA J. D. C., VIEIRA I. C. G., CINTRA B. B. L., STEVENSON P. R., FEITOSA Y. O., DUVENVOORDEN J. F., MOGOLLÓN H. F., ARAUJO-MURAKAMI A., FERREIRA L. V., LOZADA J. R., COMISKEY J. A., DE TOLEDO J. J., DAMASCO G., DÁVILA N., LOPES A., GARCÍA-VILLACORTA R., DRAPER F., VICENTINI A., CORNEJO VALVERDE F., LLOYD J., GOMES V. H. F., NEILL D., ALONSO A., DALLMEIER F., DE SOUZA F. C., GRIBEL R., ARROYO L., CARVALHO F. A., DE AGUIAR D. P. P., DO AMARAL D. D., PANSONATO M. P., FEELEY K. J., BERENGUER E., FINE P. V. A., GUEDES M. C., BARLOW J., FERREIRA J., VILLA B., PENUELA MORA M. C., JIMENEZ E. M., LICONA J. C., CERÓN C., THOMAS R., MAAS P., SILVEIRA M., HENKEL T. W., STROPP J., PAREDES M. R., DEXTER K. G., DALY D., BAKER T. R., HUAMANTUPA-CHUQUIMACO I., MILLIKEN W., PENNINGTON T., TELLO J. S., PENA J. L. M., PERES C. A., KLITGAARD B., FUENTES A., SILMAN M. R., DI FIORE A., VON HILDEBRAND P., CHAVE J., VAN ANDEL T. R., HILÁRIO R. R., PHILLIPS J. F., RIVAS-TORRES G., NORONHA J. C., PRIETO A., GONZALES T., DE SÁ CARPANEDO R., GONZALES G. P. G., GÓMEZ R. Z., DE JESUS RODRIGUES D., ZENT E. L., RUSCHEL A. R., VOS V. A., FONTY É., JUNQUEIRA A. B., DOZA H. P. D., HOFFMAN B., ZENT S., BARBOSA E. M., MALHI Y., DE MATOS BONATES L. C., DE ANDRADE MIRANDA I. P., SILVA N., BARBOSA F. R., VELA C. I. A., PINTO L. F. M., RUDAS A., ALBUQUERQUE B. W., UMAÑA M. N., CARRERO MÁRQUEZ Y. A., VAN DER HEIJDEN G., YOUNG K. R., TIRADO M., CORREA D. F., SIERRA R., COSTA J. B. P., ROCHA M., VILANOVA TORRE E., WANG O., OLIVEIRA A. A., KALAMANDEEN M., VRIESENDORP C., RAMIREZ-ANGULO H., HOLMGREN M., NASCIMENTO M. T., GALBRAITH D., FLORES B. M., SCUDELLER V. V., CANO A., AHUITE REATEGUI M. A., MESONES I., BAIDER C., MENDOZA C., ZAGT R., URREGO GIRALDO L. E., FERREIRA C., VILLAROEL D., LINARES-PALOMINO R., FARFAN-RIOS W., FARFAN-RIOS W., CASAS L. F., CÁRDENAS S., BALSLEV H., TORRES-LEZAMA A., ALEXIADES M. N., GARCIA-CABRERA K., VALENZUELA GAMARRA L., VALDERRAMA SANDOVAL E. H., RAMIREZ AREVALO F., HERNANDEZ L., SAMPAIO A. F., PANSINI S., PALACIOS CUENCA W., DE OLIVEIRA E. A., PAULETTO D., LEVESLEY A., MELGAÇO K. & PICKAVANCE G. 2020. — Biased-corrected richness estimates for the Amazonian tree flora. *Scientific Reports* 10 (1): 10130. <https://doi.org/10.1038/s41598-020-66686-3>
- TER STEEGE H., SABATIER D., CASTELLANOS H., VAN ANDEL T., DUVENVOORDEN J. F., OLIVEIRA A. A. DE, EK R. C., LILWAH R., MAAS P. J. M. & MORI S. A. 2000. — An analysis of the floristic composition and diversity of Amazonian forests including those of the Guiana shield. *Journal of Tropical Ecology* 16: 801-828.
- THE PLANT LIST 2013. — *The Plant List. Version 1.1*. Published on the Internet. Available from <http://www.theplantlist.org> [accessed 11 December 2020].
- THOMAS W. W. & FRANCESCHINELLI E. V. 2005. — Simaroubaceae, in BERRY P. E., YATSKIEVYCH K. & HOLST B. K. (eds), *Flora of the Venezuelan Guayana*. Vol. 9. Missouri Botanical Garden Press, Saint Louis: 168-176.
- TOZZI A. M. G. DE A. & SILVA M. J. 2007. — Sinonimizações em *Lonchocarpus* Kunth (Leguminosae-Papilionoideae-Millettieae). *Rodriguésia* 58 (2): 275-282.
- Tropicos 2022. — Tropicos.org. Missouri Botanical Garden. Available from <http://www.tropicos.org> [accessed 25 April 2022].
- TURCZANINOW N. S. 1863. — *Animadversiones ad catalogum primum et secundum herbarii Universitatis Charkoviensis. Bulletin de la Société impériale des Naturalistes de Moscou* 36 (1): 545-615.
- TURLAND N. J., WIERSEMA J. H., BARRIE F. R., GREUTER W., HAWKSWORTH D. L., HERENDEEN P. S., KNAPP S., KUSBER W.-H., LI D.-Z., MARHOLD K., MAY T. W., MCNEILL J., MONRO A. M., PRADO J., PRICE M. J. & SMITH G. F. 2018. — *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. Koeltz Botanical Books, Glashütten. (Regnum Vegetabile; 159). <https://doi.org/10.12705/Code.2018>
- ULBRICH O. E. 1913. — Die Kapok liefernden Baumwollbäume der deutschen Kolonien im tropischen Afrika. *Notizblatt des Königlichen botanischen Gartens und Museums zu Dahlem bei Steglitz (Berlin)* 51 (6): 1-34.

- ULE E. 1907. — Die Pflanzenformationen des Amazonas-Gebietes. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 40 (2): 114-172.
- URBAN I. 1904. — Ueber einige Celastraceen-Gattungen, in URBAN I. & GRAEBNER P. (eds), *Festschrift zur Feier des siebzigsten Geburtstages des Herrn Professor Dr. Paul Ascherson*. Gebrüder Borntraeger, Leipzig: 48-58.
- VAN ANDEL T. R. 2000. — *Non-Timber Forest Products of the North-West District of Guyana. Part I*. Tropenbos, Georgetown, Guyana, 320 p. (Tropenbos Guyana Series; 8A).
- VAN DEN BERG M. E. 1979. — Revisão das espécies brasileiras do gênero *Rheedia* L. (Guttiferae). *Acta Amazonica* 9 (1): 43-74. <https://doi.org/10.1590/1809-43921979091043>
- VAN DER WERFF H. 2002. — Lauraceae, in MORI S. A., CREMERS G., GRACIE C. A., GRANVILLE J.-J. DE, HEALD S. V., HOFF M. & MITCHELL J. D. (eds), *Guide to the Vascular Plants of Central French Guiana*. Part 2. *Dicotyledons*. New York Botanical Garden Press, Bronx (Memoirs of the New York Botanical Garden; 76 [2]): 370-384.
- VAN DER WERFF H. & VICENTINI A. 2000. — New species of Lauraceae from central Amazonia, Brazil. *Novon* 10: 264-297. <https://doi.org/10.2307/3393111>
- VAN'T KLOOSTER C. I. E. A., LINDEMAN J. C. & JANSEN-JACOBS M. J. 2003. — Index of vernacular plant names of Suriname. *Blumea Suppl.* 15: 1-322.
- VESQUE J. 1892. — *Epharmosis, sive, Materiae ad instruendam anatomiam systematis naturalis*. Delapierre, Vincennes, 24 p. + illustr.
- WAHLERT G. A., MARCUSSEN T., DE PAULA-SOUZA J., FENG M. & BALLARD H. E. 2014. — A phylogeny of the Violaceae (Malpighiales) inferred from plastid DNA sequences: implications for generic diversity and intrafamilial classification. *Systematic Botany* 39 (1): 239-252. <https://doi.org/10.1600/036364414X678008>
- WALLICH N. 1828. — *A Numerical List of Dried Specimens in the Museum of the Honl. East India Company*. Manuscript, London, 306 p.
- WALLNÖFER B. 2001. — Lectotypification of *Diospyros cayennensis* A. DC. (Ebenaceae). *Taxon* 50 (3): 887-889. <https://doi.org/10.2307/1223718>
- WALLNÖFER B. 2020. — A revision of neotropical *Diospyros* (Ebenaceae): part 13. *Annalen des Naturhistorischen Museums in Wien* 122B: 205-243.
- WALPERS W. G. 1848. — *Annales Botanices Systematicae*. Vol. 1. F. Hofmeister, Leipzig, 1127 p.
- WCSP 2022. — *World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew*. Published on the Internet. Available from <http://wcsp.science.kew.org/> [accessed 11 February 2022].
- WEBSTER G. L. 2004. — A revision of *Phyllanthus* section *Hylaeanthus* (Euphorbiaceae). *Lundellia* 2004 (7): 11-27. <https://doi.org/10.25224/1097-993X-7.1.11>
- WEBSTER G. L. & HUFT M. J. 1988. — Revised Synopsis of Panamanian Euphorbiaceae. *Annals of the Missouri Botanical Garden* 75 (3): 1087-1144. <https://doi.org/10.2307/2399381>
- WIERSEMA J. H., TURLAND N. J., BARRIE F. R., GREUTER W., HAWKSWORTH D. L., HERENDEEN P. S., KNAPP S., KUSBER W.-H., LI D.-Z., MARHOLD K., MAY T. W., MCNEILL J., MONRO A. M., PRADO J., PRICE M. J. & SMITH G. F. 2018. — *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017: Appendices I-VII*. Available from <https://naturalhistory2.si.edu/botany/codes-proposals/> [accessed 2 November 2020].
- WIGHT R. 1840. — *Icones plantarum Indiae Orientalis, or figures of Indian plants*. Vol. 1. J.B. Pharoah, Madras, 696 p.
- WILLDENOW C. L. 1799. — *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*, ed. 4. Vol. 2(1). G.C. Nauk, Berlin, 823 p.
- WILLDENOW C. L. 1802. — *Species plantarum, exhibentes plantas rite cognitas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas*, ed. 4. Vol. 3 (2). G.C. Nauk, Berlin, 625 p. (849-1474).
- WILLIAMS L. O. 1963. — Tropical American Plants, V. *Fieldiana, Botany* 29 (10): 545-597.
- WRIGHT W. 1777. — Description and use of the cabbage-bark tree of Jamaica. *Philosophical transactions of the Royal Society of London* 67: 507-513.
- WURDACK J. J., MORLEY T. & RENNER S. S. 1993. — *Melastomataceae*. Royal Botanic Garden, Kew, 425 p. (Flora of the Guianas. Series A, Phanerogams; Fascicle 13).
- ZIZKA G. & SCHNEIDER J. V. 1999. — The genus *Touroulia* Aubl. (Quiinaceae). *Willdenowia* 29 (1/2): 227-234.

Submitted on 1 September 2021;
accepted on 9 May 2022;
published on 13 December 2022.

APPENDIX 1. — List of Exsiccatae. The numbers between square brackets (**in green**) refer to accepted taxa, as numbered in the checklist/Liste des Exsiccatae. Les numéros entre crochets (**en vert**) font référence aux taxons acceptés, tels que numérotés dans le catalogue.

- Acevedo-Rodríguez P. et al.* 4871 [1580], 4911 [1592], 5012 [1578], 6130 [135]
Acosta L. & González L.A. 14 [1283]
Adão Teixeira L.O. 943 [1553]
Aguilar R. 2961 [1037]
Allorge L. & Rombold J. 335 [111]
Alvarez A. 2510 [258]
Andersson L. 1993 [1476]
Angel G. 29 [1792]
Aublet J.B. s.n. (BM000016972) [1429], s.n. (BM000028175) [1373], s.n. (BM000041973) [1379], s.n. (BM000041974) [1380], s.n. (BM000047393) [388], s.n. (BM000058142) [1450], s.n. (BM000065023) [267], s.n. (BM000073181) [1357], s.n. (BM000073601) [522], s.n. (BM000522439) [1039], s.n. (BM000541054) [818], s.n. (BM000547151) [53], s.n. (BM000547286) [60], s.n. (BM000547301) [67], s.n. (BM000566635) [1791], s.n. (BM000566650) [1795], s.n. (BM000570872) [1514], s.n. (BM000583750) [716], s.n. (BM000588689) [751], s.n. (BM000598566) [562], s.n. (BM000602313) [338], s.n. (BM000603704) [973], s.n. (BM000611049) [1535], s.n. (BM000611207) [1718], s.n. (BM000611559) [372], s.n. (BM000611835) [378], s.n. (BM000617483) [575], s.n. (BM000624376), s.n. (BM000624532) [1547], s.n. (BM000624909) [550], s.n. (BM000624924) [555], s.n. (BM000629523) [1594], s.n. (BM000630633) [1508], s.n. (BM000645671) [1022], s.n. (BM000645699) [1388], s.n. (BM000645706) [1595], s.n. (BM000645788) [1034], s.n. (BM000754877) [1298], s.n. (BM000759977) [1111], s.n. (BM000793077) [351], s.n. (BM000793078) [1454], s.n. (BM000795154) [1002], s.n. (BM000795155) [1001], s.n. (BM000795157) [1000], s.n. (BM000795575) [706], s.n. (BM000795993) [544], s.n. (BM000795994) [537], s.n. (BM000797863) [354], s.n. (BM000797998) [1349], s.n. (BM000799363) [517], s.n. (BM000812056) [515], s.n. (BM000826529) [768], s.n. (BM000826533) [772], s.n. (BM000826536) [771], s.n. (BM000838014) [1580], s.n. (BM000884091) [1576] (see [1567]), s.n. (BM000884102) [1568], s.n. (BM000884118) [1567] (see [1576]), s.n. (BM000885813) [963], s.n. (BM000931917) [970], s.n. (BM000931940) [890], s.n. (BM000931956) [967], s.n. (BM000931959) [789], s.n. (BM000931983) [952], s.n. (BM000947248) [617], s.n. (BM000947302) [637], s.n. (BM000947432) [501], s.n. (BM000952073) [943], s.n. (BM000952077) [948], s.n. (BM000952149) [763], s.n. (BM000952284) [797], s.n. (BM000952572) [1702], s.n. (BM000952638) [117], s.n. (BM000952644) [106], s.n. (BM000953533) [260], s.n. (BM000953642) [1311], s.n. (BM000953691) [1214], s.n. (BM000953694) [1213], s.n. (BM000953713) [401], s.n. (BM000953723) [404], s.n. (BM000953757) [1240], s.n. (BM000953761) (see [1252]), s.n. (BM000953762) [1105] (see [1252]), s.n. (BM000953836) [726] (see [726]), s.n. (BM000953839) [722], s.n. (BM000953843) [703], s.n. (BM000953857) [696], s.n. (BM000992002) [15], s.n. (BM000993272) [1146], s.n. (BM000993312) [1149], s.n. (BM000993316) [1188], s.n. (BM000993414) [1745], s.n. (BM000993415) [1742], s.n. (BM000993430) [1751], s.n. (BM000993837) [1209], s.n. (BM000993866) [536], s.n. (BM001008165) [1100], s.n. (BM001008254) [1047], s.n. (BM001008259) [1053], s.n. (BM001008452) [245], s.n. (BM001008661) [138], s.n. (BM001008914) [1438], s.n. (BM001135589) [936], s.n. (FI012223) [444], s.n. (G00226718) [1029], s.n. (LINN-HS 672.2) [185], s.n. (LINN-HS 672.3) [196], s.n. (LINN-HS 672.4) [188], s.n. (LINN-HS 672.5) [195], s.n. (LINN-HS 782.15) [1041], s.n. (LINN-HS 782.16) [1070], s.n. (LINN-HS 853.1) [1043], s.n. (LINN-HS 883.12) [1252], s.n. (LINN-HS 883.22) [1292], s.n. (LINN-HS 968.1) [248], s.n. (LINN-HS 968.3) [246], s.n. (LINN-HS 983.5) [24], s.n. (MPU014808) [33], s.n. (P00106271) [225], s.n. (P00114408) [1366], s.n. (P00128471) [607], s.n. (P00202541) [89], s.n. (P00287390) [231], s.n. (P00287893) [990], s.n. (P00662841) [568], s.n. (P00671713) [248], s.n. (P00671912) [1375], s.n. (P00672058) [1527], s.n. (P00675567) [1324], s.n. (P00678733) [1395], s.n. (P00680269) [1453], s.n. (P00680428) [1802], s.n. (P00689737) [1532], s.n. (P00716889) [521], s.n. (P00777268) [777], s.n. (P00777303) [881], s.n. (P00777305) [879], s.n. (P00777519) [130], s.n. (P00777520) [1666], s.n. (P00777521) [1422], s.n. (P00777522) [1376], s.n. (P00777561) [561], s.n. (P00777615) [1522], s.n. (P00777662) [1460], s.n. (P00777669) [1109], s.n. (P00777682) [1484], s.n. (P00777786) [1776], s.n. (P00777787) [1784], s.n. (P00777787) [1777], s.n. (P00777819) [1143], s.n. (P00777830) [433], s.n. (P00777832) [432], s.n. (P00777834) [427], s.n. (P00777902) [1490], s.n. (P00777907) [1417], s.n. (P00777908) [331], s.n. (P00777925) [436], s.n. (P00777973) [120], s.n. (P00777974) [119], s.n. (P00777978) [1769], s.n. (P00778013) [95], s.n. (P00778112) [1500], s.n. (P00778113) [1487], s.n. (P00778114) [1494], s.n. (P00778459) [1711], s.n. (P00778468) [293], s.n. (P00778564) [535], s.n. (P00835933) [798], s.n. (P02273439) [1474], s.n. (P02442466) [1075], s.n. (S-R-1308) [508], s.n. (W-1889-0012576) [1387], 32 [164], 200 [168]
Bang M. 1686 [1420]
Baraloto C. et al. 3500 [304], 8056 [1317]
Barrabé L. 232 [740]
Barreto M.I. 141 [1721]
Barrier S. 2633 [66], 4026 [595], 5018 [468]
Béna P. 1232 [1009], 1313 [412]
Benoist R. 138 [784], 224 [1700], 339 [242], 368 [205], 448 [440], 516 [1040], 520 [446], 539 [1382], 540 [1306], 578 [719], 667 [1184], 788 [1172], 960 [1753], 1005 [916], 1065 [728], 1074 [964], 1159 [52], 1491 [839], 1564 [1796], 1573 [1199]
Bergeron S. & Roman S. 494-556 [1299]
Billiet F. (et Jadin B.) 1230 [502], 1232 [154], 1286 [1037], 1431 [566]
Blanc M. 128 [1165]

Appendix 1. — Continuation.

- Boom B.M. (et Mori S.A.)* 1795 [905], 1805 [801], 1811 [671], 1885 [194], 2116 [5], 2222 [181], 2344 [302], 2362 [346], 2415 [156], 10418 [305]
- Bordenave B. (et Bétian)* 887 [1218], 5040 [1767], 7946 [118]
- Bourdy G.* 2941 [1432]
- Brewer S.W.* 8580 [236]
- Broadway W.E.* 880 [1743]
- Camp W.H.* E-3425 [37]
- Caraglio Y.* 508 [1258], 802 [1730]
- Cárdenas L.D.* 2850 [1765]
- Chanderbali A.* 90 [87]
- Chareyre P.* 30D [25]
- Cid Ferreira C.A. (et al.)* 5668 [288], 6948 [183], 7600 [1407], 8010 [1472], 10202 [1415], 10505 [1116]
- Clarke H.D. (et al.)* 3146 [521], 5485 [362], 9046 [495]
- Cowan R.S. (et al.)* 1973 [1727], 38370 [525], 38757 [1505], 38808 [1463], 38837 [170]
- Cremers G. (et al.)* 6509 [1145], 7287 [27], 8233 [1163], 8385 [420], 8397 [1447], 8448 [236], 11240 [1076], 11776 [1300], 13176 [1445], 14582 [779], 15215 [640]
- Crevaux J.N. s.n. (P01818367)* [824]
- Cuniberri X.* 22 [858]
- Daly D.C.* D649 [1548], 1601 [772]
- Davidse G.* 27581 [161]
- de Jong B.H.J.* 2 [1045]
- Defler S.* 590 [1017]
- Delgado L.* 840 [1248]
- Delnatte C.* 1445 [280], 1549 [1277], 1556 [738]
- Delprete P.G. & Crozier F.* 7127 [1493], 7150 [1484]
- Desvaux A.N. s.n. (P00060476, P00060477)* [1142], s.n.
(P00756603) [1159], s.n. (P01818175) [863], s.n. (P01818387) [820]
- Dutrèze B. (et al.)* 7 [611], 19 [891], 388 [1215], 399 [1247], 489 [172], 569 [684]
- Egler W.A.* 47673 [1685]
- Ek R.C.* 1745 [1452], 1835 [644]
- Engel J. (et al.)* 14 [507], 19 [1684]
- Evans R.J. et al.* 2513 [1241]
- Fernández A.* 7544 (see [1450])
- Feuillet C. (et al.)* 616 [1400], 715 [1197], 1122 [161], 2263 [212], 2961 [873], 3936 [1607], 10067 [1439], 10083 [285], 10115 [1774], 10319 [1536], 10338 [582], 10354 [1459]
- Fleury M.* 238 [294], 567 [563], 911 [901], 1348 [265], 2053 [1504], 2166 [889], 2176 [1318]
- Fonnegra G.R. & Rengifo M.W.* 4834 [370]
- Forestal H. de 125ter* [1519], 490 [1052], 760 [1403]
- Fróes R.L.* 26671 [1344]
- Fuentes A.F. et al.* 6109 [1300]
- Gabriel A.s.n. (G00139532)* [1597]
- Gentry A.H. (et al.)* 50232 [145], 63044 [1150], 63129 [204], 63218 [166]
- Gonzalez S.* 1093 [160], 2334 [1498]
- Granville J.-J. de (et al.)* 44 [1351], 488 [313], 638 [1465], 652 [1331], 1169 [1778], 1197 [845], 1475 [848], 1869 [1236], 2200 [368], 2902 [884], 3327 [1309], 3411 [155], B-3724 [1177], 3750 [152], B-3833 [1765], B-4467 [1414], B-4482 [1071], B-4488 [127], B-4549 [1220], B-4563 [950], B-4741 [1436], B-4744 [1448], B-4760 [1065], B-4792 [409], 4866 [775], B-4989 [1102], B-5195 [631], B-5356 [1486], B-5392 [461], B-5401 [367], 5410 [1171], B-5436 [1399], B-5486 [18], 5550 [1811], 5909 [1497], 6000 [1735], 6121 [276], 6268 [1478], 6330 [45], 6480 [487], 6503 [1281], 7000 [126], 7203 [148], 7632 [1676], 7734 [1144], 7755 [1355], 7911 [786], 8057 [149], 8099 [1648], 8340 [146], 8601 [1448], 8765 [1253], 8915 [1413], 9032 [1234], 9229A [1783], 9455 [1223], 9913 [157], 10319 [151], 10656 [495], 10765 [1433], 10796 [573], 10907 [36], 11201 [143], 13403 [511], 13833 [1485], 14611 [1312], 15000 [986], 15181 [1235], 16138 [744], 16705 [750], 16824 [808], 17010 [1482], 17272 [1510], 17651 [158], 17659 [142], 17664 [140]
- Grenand P. (et al.)* 15 [1721], 237 [430], 356 [1296], 510 [1348], 543 [1397], 574 [83], 626 [1423], 637 [222], 642 [311], 645 [3], 649 [906], 677 [1020], 699 [1670], 701 [1027], 715 [238], 738 [1090], 756 [1042], 774 [741], 1013 [1287], 1044 [1038], 1116 [785], 1126 [977], 1222 [363], 1291 [1712], 1303 [411], 1328 [1308], 1488 [1622], 1503 [853], 1510 [1050], 1525 [1303], 1527 [1115], 1543 [1525], 1594 [30], 1776 [415], 1808 [1435], 1845 [48], 1902 [1356], 1933 [1727], 1959 [1556], 2017 [742] (see [741]), 2040 [37], 2107 [1325], 2118 [1710], 2119 [1226], 2868 [1302], 3131 [1243], 3380 [288], 3425 [642], 3442 [1455], 3502 [503], 3521 [674]
- Grimes (et al.)* 3126 [779], 3312 [731], 3322 [735]
- Gudiño E. et al.* 1628 [1020]
- Hahn W.J.* 3534 [548]
- Hallé F.* 626 [941], 629 [1283], 1018 [1449]
- Henkel T.W. et al.* 3133 [805]
- Hequet V.* 159 [1471], 3003 [1732], 3018bis [1521]
- Hoff M.* 5980 [341]
- Hoffman B. & Rodrigues M.T.* 4705 [644]
- Hostmann F.W.R.* 1176 [1298]
- Huber O.* 333 [1324]
- Irwin H.S. (et al.)* 47561 [1247], 47819 [74], 48582 [1107], 48608 [865]
- Jacquemin H.* 1763 [1771], 1903 [101], 2435 [527]
- Jangoux J.* 1051 [504]
- Jansen-Jacobs M.J. et al.* 78 [1471], 230 [4], 2658 [1739]
- Jardim A.* 1015 [1085]
- Kajekai C.* 356 [1282]
- Kappler A.* 629bb (see [627]), 1675 [1068]
- Koemar S. & McDonnell K.* 46 [1465]
- Korning J. & Thomsen K.* 47802 [238]
- Krukoff B.A.* 5720 [116], 8023 [1768], 11867 [294]
- Larpin D.* 670 [1392], 687 [1268], 702 [1230], 846 [1280], 882 [1479], 885 [1687], 946 [422], 1021 [1353]
- Le Goff A.* 201 [8]
- Leblond J.B. s.n. (1434), s.n. (834), s.n. (G00177930) [410], s.n. (G00368596) [547], s.n. (P00371634) [1528], s.n. (P00542279)*

Appendix 1. — Continuation.

- [1350], s.n. (P00756974) [625], s.n. (P02428604) [995], s.n. (P02442081) [1551], 34 [389], 61 [1124], 64 [1089], 100 [1059], 110 [1282], 114 [1307], 144 [851], 168 [933], 209 [463], 215 [647], 224 [1421], 319 [1495]
- Leeuwenberg A.J.M.* 11653 [124]
- Leprieur F.M.R.* s.n. (F0053118F) [241], s.n. (F0062737F) [993], s.n. (G00352502) [1130], s.n. (G00388527) [861], s.n. (P00649386) [1605], s.n. (P00715205) [1409], s.n. (P00756698) [1744], s.n. (P00756786) [1758], s.n. (P00757098) [1746], s.n. (P02441925) [1371], s.n. (US00112566) [1728], 141 [1754], 195 [1737], 206 [1092], 224 [574], 225 [579], 284 [1789], 333 [1570], 335 [1575], 336 [958]
- Lescure J.-P.* 708 [283], 723 [1770], 816 [1049], 839 [373], 877 [732], 943 [1067]
- Liesner R.L. & Holst B.K.* 18851 [1057]
- Lindeman J.C. et al.* 555 [1771]
- Loubry D.* 55 [1383], 621 [1640], 833 [469], 1122 [694], 1135 [841], 1579 [739], 1626 [1278], 1698 [271], 1774 [1117], 1892 [1467], 1934 [414], 2000 [1430], 2097 [1194], 2339 [132], 2392 [1264], 2413 [1491]
- Maas P.J.M. (et al.)* 3875 [365], 8062 [32], 9351 [92], 9635 [84]
- Maguire B. (et al.)* 22937 [1526], 24588 [1747], 47074 [1626], 47081 [337], 47097 [1680], 47110 [297]
- Marcano-Berti L.* 244-981 [561]
- Marshall N. & Rombold J.* 146 [1412]
- Martin J.* 6 [791], s.n. ("in Herb. Kth.") [956], s.n. (BM000065079) [799], s.n. (BM000080822) [1419], s.n. (BM000582180) [1499], s.n. (BM000952593) [1599], s.n. (BM000953792) [692], s.n. (BM001008911) [1489], s.n. (BR0000005195022) [883], s.n. (BR0000005529711) [1786], s.n. (F0BN001875) [872], s.n. (F0BN017062) [1064], s.n. (FI005833) [1787], s.n. (FI011051) [1762], s.n. (FI011503) [1221], s.n. (G00139364) [1408], s.n. (G00143998) [102], s.n. (G00201468) [91], s.n. (G00310774) [1515], s.n. (K) [1541], s.n. (K000220643, P00746011, P00746012, P00746013) [360], s.n. (K000220657) [305], s.n. (K000329644) [1108], s.n. (K000380070) [465], s.n. (K000380085) [470], s.n. (K000407329) [539], s.n. (K000407359) [982], s.n. (K000530367) [923], s.n. (K000555167) [770], s.n. (K000555173) [769], s.n. (K000581314) [1374], s.n. (L0009403) [1216], s.n. (NY00428114) [272], s.n. (P00053053) [1717], s.n. (P00071947) [1360], s.n. (P00542206) [1345], s.n. (P00640548) [1647], s.n. (P00648720) [526], s.n. (P00649234) [1628], s.n. (P00649300) [1598], s.n. (P00721345, P00721347) [442], s.n. (P00721391) [447], s.n. (P00723485) [726], s.n. (P00733832) [1799], s.n. (P00745490) [1193], s.n. (P00780886) [1378], s.n. (P00798577) [749], s.n. (P00836503) [1481], s.n. (P016912) [1715], s.n. (P01818008) [942], s.n. (P01819656) [1114], s.n. (P02288028, P02288029, P02288030, P02288031, P02288032) [1123], s.n. (P04822073) [1548], s.n. (P05237715) [1222], 2 [926], 27 [529], 45 [602], 52 [1423], 151 [428]
- Martinelli G. (et al.)* 7294 [1071], 12268 [1304]
- Martínez Sequeira K.* 280 [1710]
- Mélinon E.M.* s.n. (F0063306F) [439], s.n. (G00367876) [961], s.n. (K000329677) [1095], s.n. (P00053272) [1716], s.n. (P00077245) [596], s.n. (P00089172) [1738], s.n. (P00093864, P00093865) [389], s.n. (P00093869) [395], s.n. (P00115935) [64], s.n. (P00163092) [1294], s.n. (P00199448) [939], s.n. (P00542463) [1352], s.n. (P00624725) [1207], s.n. (P00640597) [1656], s.n. (P00647901) [1658], s.n. (P00649246) [1631], s.n. (P00649260) [1637], s.n. (P00711133) [673], s.n. (P00745439) [666], s.n. (P00745621) [584], s.n. (P00745675) [632], s.n. (P00745952) [325], s.n. (P00746050) [336], s.n. (P00746074) [345], s.n. (P00746107) [352], s.n. (P01818703) [1048], s.n. (P01900083) [705], s.n. (P03275721) [134], s.n. (P04848420) [263], 8a (1845) [815], 13 [322], 19 (1864) [81], 42 (1862) [1606], 59 [723], 108 (1863) [1667], 132 (1842) [295], 137 (1877) [421], 147 (1845) [1359], 170 [434], 178 (1842) [1073], 204 (1842) [597], 230 (1842) [261], 250 (1842) [199], 453 (1862) [844], 488 (1862) [1299]
- Molino J.-F. (et al.)* 731 [395] (see [396]), 907 [606], 921 [224], 936 [395] (see [396]), 938 [1103], 1107 [819], 1155 [136], 1181 [16], 1197 [72], 1198 [1668], 1363 [613], 1556 [390], 1617 [1696], 1661 [287], 1703 [69], 1725 [702], 1766 [1764], 1769 [1425], 1852 [1552] (Fig. 52D), 1924 [868], 1948 [394], 1968 [56], 1981 [1305], 1987 [885], 1991 [10], 1993 [1443], 1994 [253], 1995 [1688], 2007 [1179], 2041 [1514] (Fig. 51C), 2070 [1095] (Fig. 37A), 2071 [1326], 2075 [1273], 2076 [1615], 2083 [226], 2086 [255], 2096 [664], 2099 [1340] (Fig. 44A), 2108 [1369], 2112 [1077], 2116 [1263], 2124 [767], 2130 [1416], 2133 [77], 2143 [823], 2144 [1466], 2147 [653], 2158 [77] (Fig. 8C), 2164 [1508] (Fig. 51A), 2173 [1013], 2174 [1513], 2218 [1315], 2230 [491], 2235 [1227], 2246 [963] (Fig. 32F), 2248 [215] (Fig. 13A), 2249 [1214] (Fig. 41B), 2250 [1496], 2254 [1182], 2255 [533] (Fig. 23D), 2262 [636] (Fig. 27A, B), 2267 [1183] (Fig. 39D), 2270 [1085], 2284 [210], 2287 [954], 2290 [38], 2307 [520], 2309 [1262], 2319 [682], 2329 [1313], 2332 [1127], 2335 [1087], 2337 [1759], 2349 [1517], 2365 [503] (Fig. 23A), 2367 [559] (Fig. 25C), 2399 [836], 2401 [1564], 2403 [893], 2417 [639], 2419 [557], 2436 [19], 2437 [1036], 2438 [594], 2442 [1282] (Fig. 42C), 2448 [123], 2461 [1256] (Fig. 42A), 2464 [61] (Fig. 8A), 2469 [1543] (Fig. 52B), 2484 [616] (Fig. 53B), 2496 [1190], 2505 [1124] (Fig. 38C), 2531 [473], 2657 [1261], 2661 [1365], 2664 [991], 2673 [869], 2679 [1003], 2693 [454], 2696 [870], 2713 [80] (Fig. 8D), 2708 [675], 2714 [1260], 2716 [614], 2719 [627], 2736 [65] (Fig. 8B), 2751 [677], 2773 [656] (Fig. 27C, D), 2788 [1713], 2797 [1139], 2803 [601], 2810 [553], 2815 [1078], 2816 [622], 2834 [579] (Fig. 26B), 2844 [1428] (Fig. 48C), 2846 [981], 2850 [251], 2869 [678], 3305 [835], 3309 [621], 3313 [240], 3315 [774], 3325 [871], 3328 [854] (Fig. 30F), 3329 [985] (Fig. 33D), 3333 [578], 3352 [581], 3366 [679], 3371 [568] (Fig. 26A), 3372 [393] (Fig. 18E) (see [396]), 3375 [659], 3382 [1453] (Fig. 49E), 3397 [660], 3406 [619] (Fig. 26D-F), 3423 [681], 3429 [556] (Fig. 25B), 3430 [1273] (Fig. 41D, E).

Appendix 1. — Continuation.

- Moretti C.* 354 [1511], 358 [901], 571 [1224], 606 [1418], 626 [125], 979 [1134], 1027 [1708], 1035 [1242]
- Mori S.A. (et al.)* 8398 [1784], 8864 [1747], 8972 [462], 8983 [725], 10131 [1495], 14691 [1501], 14709 [1722], 14721 [1542], 14746 [688], 14752 [1559], 14764 [314], 14784 [1633], 14824 [648], 14899 [668], 14900 [1244], 14927 [1354], 15014 [672], 15024 [571], 15027 [1581], 15028 [1722], 15075 [724], 15115 [1733], 15146 [518], 15162 [1152], 15192 [1272], 15225 [1793], 15232 [334], 15236 [937], 15239 [1679], 15253 [14], 15288 [710], 15302 [197], 15333 [1328], 15360 [62], 15394 [484], 15425 [1428], 15459 [1451], 15475 [1195], 15498 [476], 15515 [1691], 15528 [1410], 15534 [258], 15537 [116], 15554 [1237], 15562 [1441], 15603 [1520], 15690 [690], 15701 [727], 15702 [717], 17488 [1309], 17937 [1572], 17983 [178], 18001 [99], 18034 [190], 18052 [589], 18056 [383], 18124 [1543], 19160 [1004], 19167 [104], 20612 [1498], 20758 [570], 20774 [665], 20790 [356], 20800 [708], 20808 [329], 20850 [669], 20862 [634], 20928 [572], 20960 [1290], 20968 [1420], 20973 [691], 21018 [1468], 21327 [786], 21520 [1456], 21661 [1516], 21691 [782], 22034 [758], 22252 [1030], 22719 [201], 22781 [1729], 22810 [203], 22811 [512], 23207 [1362], 23276 [822], 23307 [1270], 23320 [599], 23340 [1204], 23344 [862], 23417 [1006], 23427 [1096], 23464 [266], 23532 [370], 23682 [1748], 23694 [811], 23792 [113], 23841 [327], 23898 [628] (see [649]), 23929 [59], 23938 [734], 24002 [392], 24011 [189], 24021 [243], 24136 [752], 24205 [1327], 24212 [618], 24258 [592], 24370 [318], 24757 [1255], 24778 [509], 24798 [416], 24828 [1600], 24980 [804], 24995 [300], 25034 [729], 25047 [546], 25053 [646], 25174 [274], 25177 [55], 25303 [1026], 25427 [1160], 25434 [1238], 25515 [403], 25536 [531], 25554 [269], 25564 [910], 25667 [715], 25669 [480], 25711 [1151], 25729 [244], 26472 [343], 26487 [1112], 26542 [1164], 26549 [456], 26555 [1477], 26565 [1033]
- Moricand M.E. s.n. (G00237443)* [79], 113 [591]
- Mutchnick P. et al.* 117 [266], 773 [430]
- Nee M.H.* 11694 [1403]
- Oldeman R.A.A. (et al.)* 77 [902], 156 [1248], 163 [902], 167 [946], T-321 [1169], T-390 [1104], B-521 [1409], T-650 [1281], B-652 [115], T-663 [486], B-688 [1239], B-723 [856], T-734 [1132], T-737 [1739], T-762 [1616], T-763 [1057], B-813 [1773], B-849A [1284], B-884 [168], T-909 [1407], B-1017 [1116], B-1122 [1084], B-1205 [610], 1221 [1122], 1314 [755], B-1338 [1745], B-1441 [876], B-1467 [342], B-1468 [315], B-1645 [362], B-1825 [1291], 1862 [183], 1897 [23], 1908 [1061], 1912 [186], B-1913 [1297], 1916 [1720], B-1925 [1714], B-1936 [1502], B-1946 [1746], 2001 [1017], B-2026 [1231], B-2028 [256], B-2084 [1780], B-2101 [1101], 2106 [217], 2193 [4], 2238 [492], 2242 [793], B-2312 [600], B-2317 [1161], 2386 [1553], B-2398 [562], B-2424 [485], 2627 [29], 2672 [747], 2673 [1060], B-2687 [1181], 2756 [1503], B-2784 [1119], B-2825 [1289], B-3045 [1415], 3060 [1062], 3064 [1310], 3066 [41], B-3097 [177], B-3110 [1768], B-3139 [513], 3188 [811], 3191 [1534], B-3228 [1228], B-3261 [1393], B-3288 [1158], B-3358 [1404], B-3538 [328], B-3553 [21], B-3579 [1072], B-3580 [593], B-3602 [934], B-4007 [1396], B-4125 [61], B-4132 [1755], B-4183 [1587]
- Oliveira A.A.* 283 [1163]
- Pacheco M. (et al.)* 43 [1310], 96 [1222]
- Paget D.* 201 [1264], 204 [1266]
- Patris J.-B. s.n. (G00133858)* [162], s.n. (G00201576) [20], s.n. (G00207421) [1552], s.n. (G00208434), s.n. (G00211017) [488], s.n. (G00211384) [489], s.n. (G00222421) [1293], s.n. (G00368709) [569], s.n. (G00389893) [1363], s.n. (G-DC) [1493], s.n. (P02442059) [1554]
- Pennington T.D. (et al.)* 9967 [1117], 11539 [1596], 12134 [1704], 12137 [1672], 12177 [1654], 13840 [1596], 13867 [825], 17839 [592]
- Perrottet G.S. s.n. (G00236066)* [400], s.n. (G00237263) [44], s.n. (P02288018) [1118], s.n. (P05466483) [1341], 65 [80], 201 [239], 275 [129]
- Persson C. et al.* 2069 [1457]
- Petronelli P.* 277 [1168], 300 [1449]
- Petrov I.* 185 [1803]
- Phillippe L.R. et al.* 26984 [392]
- Pipoly J.J. (et al.)* 10414 [373], 13026 [1246], 14832 [90]
- Pires J.M.* 47466 [375], 48579 [685]
- Plouman T.C. (et al.)* 8668 [1730], 8730 [1319], 9069 [1497]
- Poeppig E.F.* 2367 [735]
- Poiteau P.A. s.n. (G00008241)* [1588], s.n. (G00008481) [1561], s.n. (G00191194) [1055], s.n. (G00202465) [558], s.n. (G00237256) [40], s.n. (G00300234) [1063], s.n. (G00368893) [667], s.n. (K000380586) [1121], s.n. (K000587558) [115], s.n. (K000600133) [720], s.n. (K000600210) [707], s.n. (K00640657) [406], s.n. (P00715218) [1411], s.n. (P00725199) [153], s.n. (P00733918) [1807], s.n. (P00745654) [598], s.n. (P00753657) [1750], s.n. (P00757092) [1749]
- Poncy O. (et al.)* 901 [730], 1048 [1440], 1430 [753], 1454 [386], 1637 [855], 1680 [615], 1746 [1458], 1748 [45], 1830 [1241], 2866 [139]
- Prance G.T. (et al.)* 1701 [268], 1822 [1519], 1929 [1115], 3222 [1255], 8551 [413], 10376 [1393], 13371 [1027], 14498 [599], 22998 [1671], 25763 [923]
- Prévost M.-F. (et al.)* 165 [1014], 169 [417], 178 [425], 185 [506], 204 [996], 478 [826], 649 [1384], 709 [1295], 750 [1406], 752 [928], 761 [842], 791 [1162], 840 [1023], 854 [1175], 885 [1136], 890 [554], 893 [1782], 904 [1125], 910 [254], 930 [1225], 934 [988], 943 [1204], 944 [1032], 946 [1141], 947 [100], 956 [1251], 962 [1726], 970 [1016], 976 [1178], 992 [1254], 995 [1245], 996 [22], 1000 [620], 1005 [1079], 1042 [754], 1064 [1219], 1076 [1082], 1082 [110], 1106 [1492], 1110 [82], 1119 [165], 1135 [764], 1137 [28], 1146 [1157], 1148 [838], 1205 [997], 1206 [1080], 1207 [802], 1221 [407], 1271 [850], 1281 [1154], 1291 [1180], 1301 [335], 1361 [247], 1364 [309], 1404 [1762] (Fig. 59A), 1460 [565], 1472 [1170], 1500 [432]

Appendix 1. — Continuation.

- (Fig. 20C), 1506 [1086], 1579 [1401], 1619 [935], 1628 [369], 1636 [438], 1693 [1736], 1718 [167], 1810 [361], 1858 [1074], 1873 [90], 1893 [457], 1910 [1046], 1964 [532], 2027 [268], 2032 [1472], 2057 [930], 2059 [1233], 2071 [1031], 2161 [11], 2169 [377], 2170 [1591], 2193 [221], 2249 [1330], 2280 [1545], 2356 [1558], 2360 [1276], 2377 [114], 2378 [1025], 2464 [814], 2564 [1646], 2610 [1028], 2625 [174], 2680 [108], 2705 [1131], 2719 [528], 2742 [376], 2744 [282], 2755 [1790] (Fig. 60A), 2763 [971], 2767 [638], 2773 [58], 2780 [1153], 2787 [1690], 2789 [311], 2792 [1565], 2805 [733], 2815 [1614], 2817 [896], 2818 [837], 2839 [490], 2848 [877], 2912 [978], 2946 [778], 2950 [924], 2956 [47], 2961 [689], 2966 [894], 2970 [534], 2973 [1629], 2987 [1708] (Fig. 56A), 2990 [481], 2992 [308] (Fig. 17C), 2994 [290], 2996 [1011], 3002 [299] (Fig. 17B), 3007 [812], 3010 [1402] (Fig. 46C), 3064 [1530], 3075 [1772], 3127 [1058], 3198 [1797], 3207 [951], 3278 [225], 3301 [583], 3302 [1557], 3329 [1099], 3330 [13], 3379 [31], 3385 [1332] (Fig. 43B), 3423 [1506], 3442 [912], 3456 [498], 3479 [1577], 3500 [536] (Fig. 24B), 3527 [381], 3567 [1398], 3655 [1761], 3667 [929], 3687 [1314], 3742 [34], 3753 [1509], 3756 [834], 3768 [1386], 3775 [1201], 3778 [1462], 3781 [1523], 3840 [746], 3845 [992], 3847 [505], 3856 [292] (Fig. 17A), 3900 [1], 3964 [1723], 3968 [431], 3986 [1211] (Fig. 41A), 3988 [387], 3990 [384], 3996 [1339], 4000 [1421] (Fig. 47A), 4014 [1044], 4023 [382], 4056 [380], 4058 [1] (Fig. 6A), 4059 [922], 4065 [788], 4081 [1398] (Fig. 45D), 4166 [1608], 4194 [1173], 4214 [702] (Fig. 29B), 4257 [707] (Fig. 29C), 4262 [1394], 4274 [211], 4304 [1664], 4324 [1785], 4435 [867] (Fig. 31A), 4341 [458], 4440 [1709], 4450 [1021], 4465 [852], 4470 [519], 4476 [418], 4478 [245] (Fig. 15A), 4484 [216] (Fig. 12C), 4517 [1718] (Fig. 56B), 4523 [1167], 4550 [398], 4566 [218] (Fig. 13B), 4580 [1008], 4587 [1770] (Fig. 59D), 4604 [1512], 4615 [697] (Fig. 29A), 4623 [1364], 4647 [1620], 4660 [1259], 4662 [466], 4664 [1340], 4671 [441], 4684 [1431], 4686 [1483], 4695 [756] (Fig. 30A), 4701 [53] (Fig. 7D), 4720 [696], 4734 [1499] (Fig. 50D), 4743 [112] (Fig. 9B), 4766 [760] (Fig. 30B), 4792 [475], 4801 [17] (Fig. 6D), 4804 [429] (Fig. 20B), 4818 [169] (Fig. 11B), 4819 [1066], 4822 [551], 4826 [703] (Fig. 28D), 4891 [560], 4895 [407], 4928 [1066] (Fig. 36B), 4936 [230], 4939 [831], 4952 [200] (Fig. 12B), 4976 [795], 5001 [1100] (Fig. 37B), 5014 [477], 5313 [398] (Fig. 19A), 5314 [1301] (Fig. 42D), 5329 [248] (Fig. 15B)
- Prieto A. 173 [934]
- Puig H. 12055 [1437]
- Rabelo B.V. (et al.) 2278 [1219], 3741 [532]
- Ratter J.A. et al. 5672 [1050]
- Redden K.M. et al. 3306 [883]
- Richard H. (et al.) 70 [757], 432 [504], 622 [235], 674 [478]
- Richard L.C. s.n. (B-W07804-000) [575] (see [568]), s.n. (BR0000005221189) [1098], s.n. (BR0000005221417) [1097], s.n. (P00106269) [228], s.n. (P00106270) [227], s.n. (P00115932) [68], s.n. (P00117101) [1566], s.n. (P00199195) [753], s.n. (P00645147) [96], s.n. (P00645496) [1621], s.n. (P00662838) [605], s.n. (P00733963) [1810], s.n. (P00745942) [321], s.n. (P00746083) [344], s.n. (P00835780) [965], s.n. (P00836504) [1481], s.n. (P00836846) [1461], s.n. (P01901082) [223], s.n. (P01901181, P01901182) [371], s.n. (P02142909) [805], s.n. (P02274150) [1129], s.n. (P02288037) [1120], s.n. (P02440864) [193], s.n. (P03465179) [796], s.n. (P03821428) [1444], s.n. (P03892077) [419], s.n. (P03892659) [426], s.n. (P03914186) [1446], s.n. (P04734493) [1555], s.n. (P04821776 ?) [1533], s.n. (P05071041) [365], s.n. (P05574226) [499], 25 [1229], 63 [1301], 76 [1246], 77 [1250]
- Riéra B. (et al.) 470 [1781], 715 [1081], 1092 [220], 1257 [474], 1282 [350], 1808 [1724] (Fig. 56D), 1904 [1623], 1925 [1390]
- Rodrigues W.A. s.n. [783]
- Rohr J.P.B. vons.n. (B-W11701-020) [567], s.n. (BM000541108) [860], s.n. (BM000541200) [816], s.n. (BM000617503) [292], s.n. (C10012170) [864], s.n. (C10012171) [867], s.n. (C10015707) [1249], 125 [1671]
- Rosa N.A. (et al.) 1834 [1388], 2812 [33]
- Rudas L. A. 3391 [1554]
- Rudge E. s.n. (BM001008906) [1488]
- Sabatier D. (et al.) 28 [1147], 823 [1702] (Fig. 55E), 824 [953], 828 [1274], 839 [947], 842 [543], 844 [451], 854 [866], 858 [1267], 865 [915], 876 [257], 889 [9], 922 [51], 925 [326], 931 [1206], 960 [1208], 965 [2], 981 [98], 984 [1604], 990 [1809], 996 [711], 1001 [207], 1009 [776], 1018 [800], 1027 [1618], 1031 [976], 1051 [1731], 1052 [813], 1055 [232] (Fig. 14A), 1098 [1669], 1106 [1674], 1170 [1805], 1196 [176], 1213 [1653], 1219 [1655], 1220 [564], 1221 [1633] (Fig. 54D), 1234 [449] (Fig. 22A), 1237 [460], 1242 [783], 1249 [1051], 1266 [1695], 1267 [1627], 1277 [698], 1345 [393] (see [396]), 1351 [1686], 1354 [214], 1363 [384] (see [396]), 1423 [1342], 1424 [1041] (Fig. 36A), 1430 [1650], 1433 [1480], 1462 [1794], 1464 [1610], 1488 [1663], 1512 [658], 1516 [1659], 1528 [1689], 1539 [307], 1541 [1361], 1546 [957], 1559 [875], 1637 [1662], 1648 [1558] (Fig. 53A), 1671 [911], 1672 [972], 1684 [962], 1691 [122] (Fig. 9D), 1741 [1660], 1772 [1705], 1836 [216], 1843 [1673], 1853 [1634], 1856 [97], 1877 [1336], 1917 [588], 1973 [680], 2088 [463] (Fig. 22D), 2100 [1539], 2102 [1571] (Fig. 53C), 2116 [1442], 2121 [927], 2147 [624], 2152 [249], 2157 [794], 2159 [1205], 2173 [887], 2185 [840], 2189 [1757], 2194 [898], 2198 [187], 2203 [1779], 2216 [1039] (Fig. 35D), 2233 [1093], 2241 [1630], 2246 [1174], 2248 [654], 2249 [1279], 2255 [1707], 2259 [399], 2263 [464], 2267 [131], 2268 [453], 2270 [303], 2278 [219], 2279 [919], 2281 [745], 2283 [401] (Fig. 19C), 2294 [1094], 2300 [413], 2302 [940], 2306 [608], 2308 [748], 2312 [1788], 2316 [762], 2317 [549], 2321 [1537], 2329 [1381], 2332 [564] (Fig. 25D), 2338 [1706], 2339 [1638], 2341 [1377], 2343 [1693], 2344 [628], 2346 [497], 2347 [340], 2372 [857],

Appendix 1. — Continuation.

2417 [308], 2447 [921], 2448 [918], 2450 [1665], 2495 [657], 2510, 2529 [718], 2530 [437], 2537 [333], 2538 [1427] (Fig. 48B), 2541 [250], 2559 [612], 2560 [49], 2561 [980], 2568 [1601], 2569 [847], 2575 [379] (Fig. 18B), 2578 [1010], 2585 [1148], 2602 [202], 2611 [830], 2617 [1652], 2645 [1682], 2696 [1602], 2703 [1389], 2716 [1135], 2718 [833], 2751 [662], 2753 [1766], 2760 [1333], 2780 [312], 2800 [1196], 2831 [1424], 2842 [966], 2872 [1583], 2876 [810], 2882 [112], 2889 [1619], 2894 [1054], 2895 [1701], 2898 [385], 2901 [1518], 2927 [213], 2930 [332], 2942 [1808], 2944 [1347], 2947 [482], 2956 [1322], 2958 [661], 2970 [330], 2973 [1639], 2979 [633], 3005 [357], 3007 [1387] (Fig. 46A), 3012 [1089], 3016 [105], 3021 [231] (Fig. 13D), 3022 [1426] (Fig. 48A), 3035 [358], 3048 [241] (Fig. 14D), 3064 [107], 3065 [1377] (Fig. 45A), 3081 [349], 3086 [874], 3090 [1137], 3102 [1275], 3113 [50], 3117 [229], 3125 [273], 3146 [6], 3161 [75], 3165 [983], 3169 [1337], 3209 [206], 3215 [277], 3219 [1018], 3243 [316], 3253 [1584], 3262 [809], 3296 [296], 3298 [1636], 3306 [42], 3308 [78], 3319 [1703], 3333 [1133], 3346 [829], 3348 [1569], 3354 [393] (see [396]), 3370 [687], 3379 [339], 3382 [179], 3391 [704], 3394 [967], 3400 [1500], 3407 [1677], 3414 [1189], 3420 [900], 3422 [1624], 3426 [1166], 3500 [670], 3502 [1766] (Fig. 59B, C), 3516 [925] (Fig. 31F), 3519 [1617], 3521 [700], 3524 [1644], 3526 [994] (Fig. 34D), 3528 [39], 3533 [1643] (Fig. 55A), 3535 [1611], 3536 [736], 3540 [455] (Fig. 22B), 3547 [1012], 3550 [790], 3553 [1621] (Fig. 54B), 3554 [1440] (Fig. 49B), 3556 [1385], 3562 [355], 3564 [909], 3568 [450], 3571 [1156], 3573 [693], 3574 [1069], 3575 [806], 3756 [106] (Fig. 9A), 3589 [530], 3593 [623], 3640 [859], 3648 [1694], 3651 [1692], 3657 [319], 3658 [1590], 3660 [63], 3663 [1579], 3670 [1265], 3678 [913], 3692 [1191], 3758 [938], 3762 [1285], 3766 [969], 3780 [609], 3788 [1075] (Fig. 36D), 3797 [699], 3833 [1323], 3850 [347], 3855 [1586], 3857 [310], 3886 [1649], 3896 [979], 3937 [93], 3944 [1185], 3946 [654] (see [649]), 3951 [71] (Fig. 7C), 3953 [299], 3967 [897], 3983 [182], 4064 [1544], 4066 [538], 4068 [552], 4072 [215], 4087 [94], 4124 [792], 4126 [459], 4137 [895], 4170 [1019], 4177 [721], 4200 [626], 4201 [1470], 4221 [424], 4226 [828], 4234 [306], 4282 [817], 4356 [979] (Fig. 32E), 4359 [86], 4361 [849], 4370 [577], 4381 [380] (Fig. 18D), 4385 [892], 4393 [697], 4403 [1040], 4404 [724] (Fig. 29D), 4419 [1804], 4421 [471], 4423 [510], 4424 [128], 4425 [737], 4426 [184], 4427 [907], 4429 [467], 4432 [628] (see [649]), 4432, 4434 [652], 4451 [171], 4457 [443], 4462 [1642], 4604 [580], 4608 [1200] (Fig. 40C), 4620 [1605] (Fig. 54A), 4622 [843], 4623 [1329], 4631 [1582], 4675 [975], 4682 [57], 4707 [500], 4712 [1573], 4715 [136] (Fig. 10A), 4780 [1271], 4781 [523], 4786 [252] (Fig. 15C), 4792 [1438] (Fig. 49A), 4794 [1635], 4805 [1341] (Fig. 44B), 4807 [1074] (Fig. 36C), 4815 [743], 4820 [1024] (Fig. 35B), 4826 [163], 4829 [405], 4830 [496], 4834 [320], 4835 [340] (Fig. 17D), 4836 [780], 4839 [1451] (Fig. 49C), 4841 [1385] (Fig. 45C), 4846 [1755] (Fig. 58C), 4848 [472], 4850 [1805] (Fig. 60E), 4853 [899], 4857 [516], 4861 [713], 4862 [998] (Fig. 34C), 4863 [984] (Fig. 33C), 4868 [851] (Fig. 30E), 4872 [397], 4873 [1760], 4874 [1798], 4880 [1338], 4891 [402] (Fig. 19D), 4896 [262], 4897 [452], 4910 [663], 4914 [1138], 4921 [144] (Fig. 10C), 4925 [1422] (Fig. 47B), 4926 [1015], 4928 [1469] (Fig. 50A), 4932 [278] (Fig. 16D), 4935 [1509] (Fig. 51B), 4945 [714], 4951 [1391] (Fig. 46B), 4958 [880], 4962 [968], 4964 [472] (Fig. 22C), 4977 [1358], 5001 [1424] (Fig. 47C), 5002 [222] (Fig. 13C), 5005 [1645], 5006bis [1540] (Fig. 52C), 5012 [1155], 5016 [1735] (Fig. 58A), 5017 [408], 5025 [159], 5026 [759], 5027 [1473], 5029 [1319], 5045 [1658] (Fig. 55B), 5046 [908], 5047 [54], 5049 [353], 5055 [781], 5070 [173] (Fig. 11D), 5071 [1354] (Fig. 44C), 5072 [683], 5076 [1593] (Fig. 53D), 5078 [670] (Fig. 27F), 5079 [291], 5082 [1334], 5086 [1372] (Fig. 43D), 5099 [585], 5101 [917], 5104 [986] (Fig. 34A), 5117 [1683], 5127 [645], 5128 [1661], 5130 [1641], 5134 [1603], 5135 [655], 5136 [1801], 5137 [1741] (Fig. 58B), 5139 [7], 5161 [1187], 5187 [1531] (Fig. 52A), 5194 [208], 5197 [1365] (Fig. 44D), 5198 [494] (Fig. 21D), 5201 [206] (Fig. 12D), 5202 [541], 5203 [181] (Fig. 12A), 5204 [1198] (Fig. 40B), 5205 [198], 5229 [485] (Fig. 21C), 5246 [443] (Fig. 21A), 5250 [1005], 5252 [651], 5262 [1612], 5276 [234], 5300 [317], 5304 [46] (Fig. 7B), 5307 [1452] (Fig. 49D), 5321 [73], 5322bis [1678], 5332 [1609], 5334 [1335], 5340 [695] (Fig. 28B), 5341 [270] (Fig. 16C), 5347 [635], 5370 [278], 5371 [1625] (Fig. 54C), 5381 [1128], 5385 [1651], 5387 [76], 5399 [1128], 5402 [88], 5403 [1370], 5411 [1321], 5421 [1035] (Fig. 35C), 5527 [1433] (Fig. 48D), 5530 [1007], 5541 [1668] (Fig. 55D), 5545 [1411] (Fig. 46D), 5550 [1698], 5552 [1719], 5554 [1697], 5555 [180], 5556 [960], 5571 [882], 5572 [807], 5573 [1775] (Fig. 59E), 5574 [545] (Fig. 24D), 5575 [995] (Fig. 34E), 5588 [163] (Fig. 11A), 5589 [914], 5590 [1464], 5591 [587], 5595 [391], 5597 [133], 5609 [172] (Fig. 11C), 5616 [1328] (Fig. 43A), 5620 [1088], 5625 [1186], 5628 [1699], 5629 [301], 5634 [275], 5641 [1733] (Fig. 57D), 5642 [1538], 5647 [423] (Fig. 20A), 5648 [1140], 5649 [70], 5651 [1613], 5653 [958], 5662 [586] (Fig. 26C), 5671 [886] (Fig. 31B), 5672 [1524] (Fig. 51E), 5683 [1546], 5692 [259], 5702 [1136] (Fig. 38D), 5703 [376] (Fig. 18A), 5705 [445], 5707 [630], 5708 [284] (Fig. 16E), 5717 [1657], 5722 [827], 5723 [139] (Fig. 10B), 5727 [676], 5728 [359], 5738 [987] (Fig. 34B), 5747 [1520] (Fig. 51D), 5750 [348], 5755 [1144] (Fig. 37D), 5760 [514] (Fig. 23C), 5762 [364], 5778 [1367], 5780 [1243] (Fig. 41C), 5782 [540] (Fig. 24C), 5785 [904] (Fig. 31C), 5792 [435], 5793 [1156] (Fig. 39B), 5794 [1257] (Fig. 42B), 5799 [396], 5802 [524], 5803 [448], 5805 [1734], 5806 [479], 5812 [1320], 5843 [1675], 5852 [1723] (Fig. 56C), 5877 [983] (Fig. 33B), 5891 [781] (Fig. 30C), 6003 [237], 6007 [1121] (Fig. 38B), 6009 [932] (Fig. 32B), 6013 [506] (Fig. 23B), 6043 [233] (Fig. 14B), 6045 [981] (Fig. 33A), 6109 [6] (Fig. 6C), 6122 [1632], 6137 [911] (Fig. 31E), 6161 [1113] (Fig. 38A), 6272 [1800], 6283 [931] (Fig. 32A), 6330 [237] (Fig. 14C), 6341 [1796] (Fig. 60B).

Appendix 1. — Continuation.

- Sagot P.A. s.n. (P00115911) [65], s.n. (P00745955) [324], s.n. (P03110846) [920], s.n. (P05483039) [483], s.n. (P06753650) [1176], 102 (1857) [989], 164 (Aug. 1855) [821], 215 (1857) [1212], 220 (1858) [1091], 259 [1217], 269 (1854) [701], 272 (Jan. 1857) [1218], 447 (Dec. 1856) [429], 491 (1857) [643], 792 (Aug. 1855) [281], 802 (1856) [761], 861 (1855) [1740], 916 (May 1856) [137], 924 (1856) [1106], 940 (1856) [85], 956 (1858) [604], 966 (Sep. 1856) [109], 973 (1956) [12], 978 (Oct. 1856) [650], 987 (P05206288) [1056], 990 (Nov. 1856) [1752], 1037 [787], 1047 (1857) [1589], 1095 (Oct. 1857) [43], 1104 (1858) [709], 1137 (1857), 1182 (1858) [374], 1188 (Aug. 1858) [1574], 1189 (1858) [1681], 1190 (Aug. 1858) [603], 1191 (Aug. 1858) [209], 1194 (1857) [1585], 1195 (Sep. 1858) [1203], 1198 (1858) [1368], 1199 [955], 1202 (Oct. 1858) [17], 1203 (May 1858) [1105], 1214 (Jan. 1859) [1563], 1229 (Mar. 1859) [493]*
- Sánchez S. M. et al. 1653 [1302]*
- Sanoja E. 2708 [1303]*
- Sastre C. (et al.) 265 [1763], 1621 [1560], 3852 [1507], 4009 [766], 4020 [1350], 4125 [542], 4146 [773], 4516 [1083], 4704 [1526], 6178 [1192], 6387 [1304], 6405 [1346], 8125 [949]*
- Schnell R.A.A. 12236 [87]*
- Schomburgk M.R.(?) s.n. (V0057797F) [907]*
- Schomburgk R.H. s.n. (K000264886, K000264887, K000264888) [907]*
- Service Forestier M-23 [999], M-63 [103], M-207 [1643], 368M [765], 4005 [832], 4436 [776], 5118 [1232], 5147 [1286], 6096 [959], 7073 [191], 7075 [945], 7237 [289], 7445 [1725], 7603 [974], 7816 [576], 7855 [366], 7911 [1316]*
- Silva N.T. & Brazão U. 60788 [1231]*
- Skog L.E. & Feuillet C. 7440 [1475]*
- Sperling C.R. 6591 [1072]*
- Spruce R. 1817 [1420]*
- Steyermark J.A. 87845 [368]*
- Stoupy D. s.n. (MA475675) [490]*
- Strudwick J.J. et al. 3121 [41]*
- Tavares A.S. 289 [1732]*
- Thomas W.W. et al. 5247 [1220]*
- Torke B. & Gonzalez S. 191 [944]*
- Tostain O. (et al.) 1499 [686], 2725 [1806], 2818 [803]*
- Unknown coll. s.n. (FI005245) [641], s.n. (G00177583) [279], s.n. (G00201278) [35], s.n. (G00201449) [65], s.n. (G00210645) [389], s.n. (G00226077) [26], s.n. (P00631979) [1210], s.n. (P00646707) [121], s.n. (P01817729) [1288], s.n. (P02288143) [1126], s.n. (P02428546) [993], s.n. (P05308177) [192], s.n. (P06671081) [1562]*
- Urrego L.E. et al. 325 [1221]*
- Vaillant A. s.n. (P03464491) [878]*
- van der Werff H. (et al.) 5021 [1428], 12951 [590], 19273 [1744], 23493 [331], 23500 [649]*
- Vicentini A. 1008 [192]*
- Villiers J.-F. (et al.) 1873 [1269], 1974 [1343], 2086 [854]*
- Wachenheim G., sér. 2, 392 [1756], sér. 3, 11 [264], 88 [888], 93 [742], 99 [931], 149 [846], 201 [71], 359 [712], 372 [286]*
- Westra L.Y.T. 47298 [1525]*
- Zaandam C.J. 6298 [449]*

APPENDIX 2. — Index of vernacular names. The numbers between brackets refer to accepted taxa, as numbered in the checklist/Index des noms vernaculaires. Les chiffres entre parenthèses renvoient aux taxons acceptés, tels que numérotés dans le catalogue.

A

- | | | | |
|--------------------------------|---|-------------------------|----------------------------|
| a teki uman udu | [76] | abiurana-vermelha | [1605], [1664], [1675] |
| a'i amiya | [728] | abiu-rosadinha | [1642] |
| a'i kálátá'a | [128] | abriba | [31] |
| a'i lea | [1750], [1756] | abriba-sovaj | [26] |
| a'i makule | [562], [1719] | abrico-de-macaco | [696] |
| a'i meyu | [499], [503] | abuluguni | [816], [818], [820] |
| a'i mididju | [499], [501] | acácia | [933] |
| a'i miniyu | [501] | acacia franc | [794] |
| a'i pane | [967] | acacia mâle | [793], [913] |
| a'i pino | [153] | açacu | [516] |
| a'i walipi | [715] | açacurana | [802] |
| a'i walipi sili | [716] | açaí-da-mata | [150] |
| a'i yú'i | [1205] | açaí-do-pará | [149] |
| a'ipopita | [859] | acajou de Guyane | [1114] |
| á-agag | [525] | acapú | [973] |
| aaku-gawut | [1323] | acapurana | [809], [967] |
| aaku-priyo | [1776] | acará-uba | [1671] |
| aawa | [196], [209] | acariquara | [1373] |
| á-awainó | [967] | acariquara-branca | [107], [108], [109] |
| abacate-rana | [982] | acarirana | [107], [108] |
| abalaba | [236], [1442] | acariúba | [1373] |
| abeemu | [1192], [1193] | acauá | [1468] |
| abérémou | [1192], [1193] | achiwa | [260] |
| abiu | [1646] | achiwa-kwali | [1803] |
| abiu-balatarana | [1629] | achuá | [541] |
| abiu-fofo | [1618] | açoita-cavalo | [1016] |
| abiu-giboa | [1652] | açucena-do-mato | [1489], [1490] |
| abiu-grande | [1670] | adai weko | [858] |
| abiu-maparajúba | [1680] | adamna | [1500] |
| abiu-pitomba | [1655] | á-danó | [751], [982] |
| abiurana | [1599], [1604], [1641], [1644], [1655], [1668],
..... [1674], [1688] | adugwe | [547], [943], [946], [948] |
| abiurana-abiu | [1616], [1666] | afa komu | [154] |
| abiurana-acariquara | [1646] | afasi boyti | [515] |
| abiurana-arana | [1650] | afatu adeeten | [496] |
| abiurana-bacuri | [1616] | agagut | [484], [557] |
| abiurana-balatatinha | [1690] | agagut-aška | [1535], [1785] |
| abiurana-braba | [1667] | aganananga | [1006] |
| abiurana-branca | [1637], [1662], [1666], [1703] | agandyaymay | [15] |
| abiurana-caju | [1616] | agi | [889], [890] |
| abiurana-camazal | [1686] | á-gógo | [107] |
| abiurana-casca-fina | [1666] | á-griyó | [673] |
| abiurana-cascuda | [1678] | agu siton | [1490] |
| abiurana-cutiti | [1671] | agugagi | [246], [248] |
| abiurana-de-casca-grossa | [1669] | ahakyu | [1529], [1537] |
| abiurana-goiabinha | [1632], [1673] | ahakyu-kamwi | [499], [1143] |
| abiurana-grande | [1679] | ahamban | [137], [138] |
| abiurana-jarani | [1667] | ahavuiye | [287], [292], [297] |
| abiurana-mocambo | [1605] | ahavuiye-seine | [292] |
| abiurana-roxa | [1632], [1640] | ahavuiye-wašiune | [547] |
| abiurana-sapota | [1669] | ahayumna | [1102] |
| abiurana-seca | [1648] | ahayupná | [1098], [1102] |
| | | ahikyu | [1529], [1537] |

Appendix 2. — Continuation.

ahikyu-kamwi	[1143]	akiki inga.....	[844], [856]
ahuwahu	[184], [187], [188], [193], [200], [202], [209], [214], [215], [219]	akiki nami	[733], [794], [915]
ahuwahu-puvemna.....	[1137]	akiki poa.....	[692]
aigvan.....	[562]	akiki'í	[926]
aigvan-kamwi	[561]	akila'í.....	[399]
aimalaje.....	[181], [211], [216]	ako	[1389]
aimyala posowepo	[1490]	akojo.....	[1099]
aipawaj.....	[99], [100]	aku-aiwut	[1248], [1285], [1604], [1628]
aisili.....	[778]	akukuwa elepali	[1528]
aituwu.....	[1141]	akuli elepali	[331], [715]
á-ivuniat.....	[918]	akuli epa.....	[1438]
aiyawa	[221]	akuli kiyeleli	[745]
ajará	[1597], [1598], [1706]	akuli tumu	[1022], [1040]
ajawaimé	[15]	akuli weseapotapili	[1051]
ájelik	[778]	akuma	[106]
ajuru	[281], [282]	akumalan	[114]
ajururana.....	[283]	akusi ákánge	[95]
aka'í.....	[1040]	akusi luway.....	[242], [1104]
akabdat	[245], [515]	akusi walapulu.....	[95]
á-kabebetye	[443]	akusi walapulu kanapua.....	[95]
akadjun	[8]	akusi walapulu pokó	[95]
akajot	[1099]	akusi walapulu tâ'í	[95]
akajou-blan	[1714]	akusi yeti	[1245], [1671]
akajou-lan	[762]	akusi'í	[788]
akajou-rouj.....	[1114]	akwansiba.....	[1633]
akalali.....	[1137]	ala'a	[1318]
akale loway	[521]	alaídya	[81], [1252]
akale pomí'idyi	[79]	alakapuli	[383], [389], [892]
akale tapulala kili	[490], [1349]	alakapuli paindyali	[382], [389]
akale wewe	[1529]	alakubali	[247]
akami ekunali	[1492]	alakuede	[196]
akami enulu	[1291]	alakuheke	[196]
akamna.....	[592], [647]	alakuhele	[195], [196]
akamná.....	[592], [647]	alakuseli	[185], [188], [189], [196]
akamna-seinó	[643]	alakušini	[185], [196]
akamná-seinó	[643]	alákwá inga.....	[863]
á-kamwi	[1121], [1126], [1717]	alala ká'í	[96], [98], [103]
akapna.....	[592], [647]	alala ká'í sili	[1426]
akapna-seinó	[643]	alala munuwi	[404], [407], [411]
akapu	[973]	alala munuwi sili	[400]
á-kasiuminio	[434]	alalawaju	[945]
akasya.....	[733], [805]	alali	[1318]
akasya-fran	[794]	alali lá	[992]
akau	[1038]	alamá'í	[267]
akau pepen	[1040]	alamá'í	[652]
akaya	[13]	alamakulu alami	[1496]
akayu.....	[8]	alamilu	[1223], [1298]
akayu u.....	[7], [9]	alamilulan	[884]
akayuu.....	[7], [9]	alamulu	[912]
akepiimyo.....	[1440]	alamulu pilá	[916]
á-kiavú.....	[279], [354], [357]	alamulu sī	[912]
á-kiavunó	[279], [354], [357]	alamulu sili	[911]
á-kiavunó-tivarabuyenë	[333]	alaone	[162]
		alapalapa	[96]

Appendix 2. — Continuation.

alapali.....	[877]	aliwa so u	[1108], [1489]
alapalí.....	[963]	aliwa u	[1489]
alapali.....	[877], [883]	aliwa'ú	[188], [196], [202]
alapokwa'i.....	[779], [785]	aliwatso tawa	[1099]
alapokwa'i wu.....	[915]	aliwe puman.....	[794]
alasigun.....	[30]	aliyana'i	[751], [944], [945], [947], [952]
alasiku.....	[1329]	almecegueira.....	[196], [202]
alasiku sili.....	[1145]	almecegueira-do-brejo	[210]
alašityu.....	[30]	almesclão	[181]
alašityulan.....	[28]	alokoyulu	[1529]
alata udu.....	[1373]	altinha	[439]
alatapa.....	[878]	aluau	[174]
alatapali.....	[877], [883]	alukaju	[41]
alataweli.....	[1373]	alukasi	[208]
alatuluka.....	[429], [432]	aluko wonale	[427]
alaulama.....	[784], [959]	aluku elepali	[1368]
alauna.....	[309]	alukumali	[247]
alawani.....	[159], [162]	alukumalilan	[246]
alawata busulukulu.....	[824], [856]	alupkamata.....	[598]
alawata emulutano.....	[129]	aluti.....	[1210]
alawata muleli.....	[740], [959], [1550]	alwaso	[1099]
alawata pana.....	[879], [1137]	amā kiya	[361]
alawata posu:lukulu.....	[824], [856], [863]	ama'i.....	[1737]
alawata pupot.....	[824]	ama'i átā	[1740]
alawata uhmot.....	[794]	amaba.....	[121]
alawone.....	[159], [162]	amākiya sili.....	[598]
alekikolan.....	[1532], [1547]	amakui	[96]
alekosin.....	[533]	amala'i	[1099]
alemenango wewe.....	[501]	amanda	[403]
alemikilan.....	[1496], [1522]	amandaa.....	[403]
alepa weli.....	[497]	amandier pays	[403]
alepawana.....	[715]	amandier sauvage	[401], [402], [405]
alepawana enekan.....	[716]	amandra	[403]
alesiki'i.....	[893]	amap	[121], [1147], [1151]
alesikyi'i.....	[740]	amapa.....	[121]
aletepe.....	[1143]	amapá.....	[121]
aletipi.....	[249]	amapá-amargoso.....	[121]
algodão-bravo.....	[1010]	amapá-doce	[117], [1151]
algodão-do-brejo	[1037]	amapakuwa	[1151]
algodão-do-mato	[173]	amapala	[1151]
aliamélë.....	[745]	amapalá	[1151]
alibagoli.....	[961]	amapapali	[95]
aliki enakololi.....	[844]	amapá-rana	[1151]
alilime'i.....	[292], [1730]	amap-purubumna	[1192]
alimi huhu.....	[408]	amarante	[918], [919], [920]
alimiao.....	[926], [927]	amarelinha	[933]
alimiap.....	[927]	amarelinho	[1385], [1508], [1717]
alimiaw.....	[926], [927]	á-maruibika	[1690], [1701]
alimihmo.....	[1646], [1666]	amata'i	[906], [908]
alimijaop.....	[927]	ambalali	[95]
alisigii.....	[522]	ambelanier acide	[95]
aliwa.....	[1488], [1490]	ambotay	[24]
aliwa so	[1099]	amejó	[46]
aliwa so átā	[1108], [1489]	amendoeira-da-praia.....	[403]

Appendix 2. — Continuation.

amila'u.....	[1095]	anjélik	[778]
aminiyu lá	[501]	ankatu.....	[1157], [1167], [1174]
amor-seco	[496]	ankono.....	[749]
amose	[83], [85]	anpuku weko.....	[867]
amoselan	[81]	a-nyan-mi-kaasi udu.....	[1127]
amourette	[1149]	anzili	[778]
amourette tachetée	[1149]	apa	[797], [798], [800]
amu'a iwa	[638], [647]	apa'uwa	[766]
amu'a iwa u	[635]	apakanilan	[794], [913], [939], [961]
amula'i.....	[1095]	apalaba	[236], [515]
amulau	[1095]	apali'i	[1]
amutu'i.....	[1068], [1070], [1079], [1080]	apali'ilan.....	[1429]
amuu.....	[144]	apalitono	[1108]
amuwan	[1472], [1503]	apaliyu	[1]
ana wila	[309]	apaliyulan.....	[1429]
ana'i.....	[110]	apazeiro	[797], [800]
ana'u	[110]	ape'i.....	[1000], [1002]
anahkale	[859]	ape'i sī	[1000], [1002]
anajá.....	[147]	apelemu.....	[20]
anakala	[850], [859]	apelemu'i	[40], [46]
anakoko	[889], [890], [891], [896], [897], [952]	apesiya	[541], [1368]
anakolo	[850]	apesulu	[1029]
anangosi	[400], [401], [402], [405]	api.....	[1150], [1152], [1196], [1198], [1199]
ananí	[380]	apijó.....	[1529]
anani-da-terra-firme	[378]	apipyoloi	[837]
ananiyu	[378], [380]	apiranga	[254]
anatapali	[934]	apisi.....	[652]
anauera	[309]	apisi-jonn	[1605]
anauira	[309]	apopopano eptyi	[1019]
anaula.....	[301], [303]	apoto ayawa	[219]
anawa	[309]	apotono ali	[576]
andira-uxi.....	[745], [747]	apuí.....	[1159], [1162], [1174], [1178], [1179]
andiroba	[1111], [1112]	apuí-grande	[366], [1745]
andoya	[1213]	apuí-preto	[1167]
aneisi udu	[537]	apuizinha	[372]
anekélë	[846]	apukuitalan	[943]
aneku	[1374]	apukuitya	[96], [98], [101]
aneku-kamwi	[12], [1374]	apukuitya tamunen	[99]
angélica-do-Pará	[778]	apulukun	[816], [850]
angelim	[786], [809], [810]	apulukuni	[816], [818], [820]
angelim-amarelo	[811]	apuruí	[1494]
angelim-coco	[745], [747]	apuruizinho	[1434]
angelim-comum	[808]	apuxiqui	[740]
angelim-da-mata	[808], [811]	ara	[14], [15], [16]
angelim-falso	[733]	ara-a	[697]
angelim-pedra	[786], [811], [812]	ara-ahavukunō	[15]
angelim-rajado	[978]	arabá	[955]
angelim-vermelho	[786]	araçá	[1245], [1279]
angelin	[809]	araçá-de-anta	[1043]
angélique	[778]	araçá-lima	[1213], [1214]
angico	[926], [927]	araçá-pera	[1318]
angouchi	[400], [402], [404], [407], [411]	araçázinho	[1251]
ani	[741]	arak	[1243], [1298]
anila wisi	[17], [747]	arak-ahavukune	[1243]

Appendix 2. — Continuation.

ara-kamwi	[207]	asakali	[1043]
arakeu	[1051], [1055], [1080]	asaki	[1043]
arakeu-kamwi.....	[1080]	asao	[740]
arakeu-priye	[1080]	á-seiminio	[1802], [1803]
arakeu-seine.....	[1048], [1051]	á-seiminio-kamwi.....	[508]
arak-kamwi	[1220], [1233], [1245], [1298]	á-seiminio-kamwi-priyu	[508]
arak-kwikwiyo	[1279]	á-seiminio-priyo	[439], [772], [1064]
arak-priyu.....	[249], [1225], [1232], [1282]	á-seiminio-priyu	[439], [772], [1064]
arak-puvemna.....	[1305]	á-seiminio-purubumna.....	[1806]
arak-seinô.....	[1301]	á-seiminio-puvemna	[1791], [1797]
araksim.....	[176], [191], [211], [221]	á-seiminio-seine.....	[1795], [1796]
araksim-priyo	[193]	asemâ	[673]
arapari	[877]	asemunusi	[948]
arapari-branco	[912]	asepoko	[1628], [1666]
arapari-rana	[770]	asepokolan	[1673]
arapari-vermelho	[906]	asepuku	[1628], [1666], [1667]
araracanga	[96], [98]	asiantefi	[293], [294]
araracanga-branca.....	[103]	asila	[864], [975], [977]
ararandeu.....	[975]	asinadan	[1681]
arara-seringa	[501]	asîngau lemi moay	[1162], [1174], [1178]
arara-tucupi.....	[912], [913]	asîngau lemi moay sili.....	[1175]
araraúba	[96], [98]	asiru	[975], [977]
arariba	[1500]	asiru-washiunô	[753]
arariúba	[779], [783]	asisi	[1490]
araróba	[101]	asisimê	[1568]
ararut	[484], [557]	asiu.....	[303]
ararut-aška.....	[1535], [1785]	ashi	[303], [539], [540], [541]
arasgu	[1470], [1472]	ashi-aška	[260]
arateu	[1051], [1055], [1080]	ashi-kamwi	[544]
arateu-kamwi	[1080]	asiu-seinô	[303]
arateu-priye	[1080]	ashi-seinô	[303]
arateu-seine	[1048], [1051]	á-sivari	[751]
araticú	[24]	asiwa	[540]
araticú-açu	[30]	ashiwa	[539], [540], [541]
araticú-cagão	[30]	asiwa kwali	[1803]
araticú-da-lagoa.....	[28]	asiwakalan	[516], [802]
araticú-do-brejo.....	[33]	atá-brava.....	[35], [49]
araticú-do-mato.....	[26], [27]	atagalí	[1454], [1455]
araticú-do-Pará.....	[35]	atakali	[1442], [1454], [1455], [1457]
araticú-macho	[26]	atakalilan	[1095]
arawa-kagta	[147]	atakamala	[1599], [1646], [1651], [1656]
arbre à beurre	[248]	atana	[309]
arbre à boulet de canons	[696]	atapa	[878], [879]
arbre parasol.....	[429]	atapalan	[770]
arey-avain	[961], [963], [964], [965]	atapili:li	[15], [16]
arey-avayni	[961], [963], [964], [965]	atapilipyo	[498]
aricurqua	[1389]	atá-preta	[31]
arirana	[1188]	atila kookoo	[281]
arisis'i	[167]	atimi letimá'i.....	[1297], [1302]
aroeira	[10]	atit-kamwi	[231]
arraieira	[501]	atiwa'u	[1120]
arraieira-branca.....	[503]	atsemâ	[673]
arua-felpudo.....	[427]	atuwai	[1120]
asaikiabe	[1212], [1279], [1295], [1315]	audik-amana.....	[1474], [1487]

Appendix 2. — Continuation.

- audika-retni..... [1474], [1487]
 audika-retni-seine..... [562]
 auluko yulu [1529]
 auma pisi..... [1205]
 auman pisi..... [224], [1368]
 auwau..... [184], [187], [188], [193], [200],
 [202], [209], [214], [215], [219]
 auwau-purubumna..... [184], [211]
 auwau-puvemna [1137]
 auwau-seine..... [202]
 auwau-seinó [215]
 avakni-avak [1719]
 avakni-awak puvemna [1343]
 avakni-awak-seine..... [1343]
 avatni-awak [1719]
 avitkat [1098]
 avitkat-kamwi..... [1104], [1242]
 avitkat-purubumna [1098], [1099], [1247]
 avitkat-seinó [1095]
 avitkat-wahuye [1095]
 avukun [815], [816], [818], [850]
 avukun-duwó [816]
 avukun-kamwi [740]
 avun-duwé [709], [721]
 avun-duwó [722], [724]
 avun-priye [698], [701], [705], [720]
 avun-priyo [705], [720]
 avun-purubumna [703]
 avun-seine [721], [723]
 avun-seinó [698], [699], [706], [707], [709]
 awa..... [209], [215], [221]
 awaa [146]
 á-wakaha [242], [1468]
 awala [146]
 awala kásí [1722], [1723]
 awala'i [146]
 awaladala [114], [1516], [1522], [1523]
 awalaime [1015]
 awalapuna [517], [519]
 awalapuna sili [436], [519]
 awalatala [114], [1516], [1522], [1523]
 awale emulutano [427]
 awale mulu [427]
 awale tamipipyo [1211]
 awalepokan [1717]
 awao [17], [1295], [1315]
 awara [146]
 awara-monpé [156]
 á-wareuni [1406]
 awata epi [427]
 awau [17], [1295], [1315]
 awau-seinó [1315]
 awawaju [1423]
 awayo [1114]
- aweiko [784]
 awiyu [89], [91]
 axixá [1031], [1034], [1035], [1036]
 ayá leme [707]
 ayá leme sili [705], [707]
 ayá pápē [370], [372]
 ayá pápē sili [370], [372]
 ayá pili [1440]
 ayalani [977]
 ayao [18], [184], [221]
 ayau [18], [184], [221]
 ayawa [219], [221]
 ayawa ilikili [208]
 ayawa sili [176], [200]
 ayawalan [207]
 ayay [181]
 ayay-wahuyo [181]
 ayee udu [751], [752]
 ayee wato [1449], [1487]
 ayee weko [781], [784]
 ayee wiwii [522]
 ayiwi [603]
 ayiwilan [1756]
 ayowo [536]
 ayu'i [603]
 áyú'i [598], [629], [642], [643], [668]
 áyú'i alapítá milá [593], [597], [624], [636]
 áyú'i pino [635]
 áyú'i piye [584], [650]
 áyú'i piyú [636]
 áyú'i sili [575], [620], [635], [637], [638], [650]
 áyú'i tawa [574], [575], [583], [603], [620],
 [626], [627], [650]
 áyú'i to u [666]
 áyú'i witowa [620], [625], [627], [650]
 ayuwato [1487]
 azeitona-brava [1228]
 azo udu [78], [81], [82]

B

- baaka agandyaymay [16]
 baaka aki [701], [707], [709], [710], [719]
 baaka apisi [636]
 baaka bee sii [1252], [1307]
 baaka dondomisinga [916]
 baaka kiikii [1028]
 baaka kookoo [321], [326], [338], [339]
 baaka luabi [702]
 baaka mapa [106]
 baaka musupu [1043]
 baaka pali udu [97]
 baaka panga [929]

Appendix 2. — Continuation.

baaka sikin	[636]	baratinha	[759]
baaka sikin apisi	[628], [649], [652]	barrotinho	[211]
baaka tiki.....	[444], [447]	barúna	[931]
baaka tyabisi.....	[788]	bataballi	[1616]
baaka wapa.....	[902], [903], [929]	batakivie.....	[1426]
baaka weko.....	[820], [858]	batibati.....	[95]
baala wiwii	[1578]	batinga	[1658]
babun weko.....	[824]	batiputá.....	[1345], [1349], [1354], [1426]
bacaba	[153]	baton-zin	[22], [43], [48], [79]
bacabinha-quina.....	[1468]	baume cochon	[202], [219]
bacuri.....	[379]	baytakini	[380]
bacuri-bravo	[378]	bee sii.....	[1307]
bacuri-de-anta	[1631]	beiju-de côco	[777]
bacuri-de-espinhos	[374], [377]	yii udu	[797]
bacuri-liso	[376]	bilitsi	[152]
bacuripari	[376]	biriba.....	[31]
bacuri-selvagem	[374]	biribá-brava	[26]
bagaceira	[1146]	biribarana	[43], [49]
baga-de-morcego	[1135]	bita kwasi	[1710]
bagas	[1146]	bita tiki	[439], [440], [443], [1527]
bagasse	[1146]	bita udu	[107], [108], [109]
bakov-kanotié.....	[858]	bita weko	[846], [862]
bakuu kokonoto	[696]	boco	[751], [752]
bakuuman	[1626], [1628], [1637]	bofo paaya	[1149]
balão-chinês	[754]	bofo tiki	[1776]
balata	[1621], [1623]	bofo udu	[541], [543], [544]
balata bali.....	[1616]	bofoo paaya	[1149]
balata blanc	[1633]	bofoo tiki	[1776]
balata blanc à grandes feuilles	[1628]	bofoo udu	[541], [543], [544]
balata franc.....	[1621], [1623]	bois à calumet	[517]
balata indien.....	[1605]	bois bandé	[1374]
balata jaune d'œuf	[1605]	bois baptiste	[550], [555]
balata poirier	[1664]	bois caca	[535]
balata pommier	[1606], [1702]	bois caïman	[1529]
balata'i.....	[1621]	bois canon	[1737], [1740]
balatabali	[1616]	bois cathédrale	[1441], [1442]
balata-blan	[1616], [1628], [1631], [1633]	bois chandelle	[1209]
balata-brava	[1606]	bois corbeau	[955]
balata-duwē	[1623]	bois cruzeau	[1802]
balata-endjen	[1628], [1633]	bois d'arc	[1149]
balata-fran	[1621]	bois dartre	[550], [555]
balata-gonm	[1621]	bois de fer	[751], [752]
balata-jonndèf	[1605]	bois de la saint-Jean	[137], [138]
balata-kamwi	[1369], [1666], [1690]	bois de lettre moucheté	[1149]
balata-ponm	[1604], [1702]	bois de lettre rouge	[1152]
balata-puvemna	[1624]	bois de mèche	[1000]
balata-pwé	[1642], [1664]	bois de rose	[583]
balata-rosada	[1628], [1631], [1633]	bois de sang	[550], [555]
balata-rosadinha	[1629]	bois diable	[516]
balata-senj-rouj.....	[1606]	bois échelle	[1141]
balateira	[1621]	bois flèche	[1095]
balipa	[147]	bois flibustier	[637]
balupete	[74]	bois fourmi	[963], [964], [965]
bamba apisi	[574], [576]	bois grage blanc	[1419]

Appendix 2. — Continuation.

bois grage rouge.....	[1422]	bukutru-gateu-wahuyo	[339]
bois indien	[508]	bunahati.....	[973]
bois macaque des bancs de sable	[733]	bunahati tyabisi.....	[931]
bois pagae.....	[97], [99], [955]	bunsumiki katu	[1182]
bois pagode	[740]	buriti.....	[152]
bois parasol	[429]	burra-leiteira.....	[533]
bois piquant	[1522], [1523]	busi amanda	[1702]
bois satin	[1152]	busi amandaa	[1702]
bois serpent	[978]	busi apaa	[1601], [1616]
bois tarin	[1148]	busi atuku	[24], [26], [29]
bois vache.....	[106]	busi banda.....	[1225], [1256]
bois violet.....	[919], [920]	busi bisangula.....	[517]
bois-yaya	[214]	busi kabana foo	[733]
boko.....	[751], [752], [1104], [1383]	busi kakaw	[1039], [1040]
boko-blan.....	[757]	busi kananbuli.....	[1708]
boliken kookoo	[303]	busi kasu	[9], [17]
bolototo uhu	[138]	busi kasun	[9], [17]
bonbekiyya.....	[1782]	busi kiikii	[419], [429]
bonga bita	[107], [108]	busi kusuwe.....	[451], [458], [460], [463], [499]
boni udu	[1527], [1724]	busi mangu	[386], [389]
bonm-kochon	[202], [219]	busi momow	[1039]
bope	[13]	busi mulunba	[1207]
bougouni.....	[816], [818], [850]	busi papay	[1749], [1752]
bougouni-blan.....	[740]	busi peesina	[309]
bougouni-tí'fey.....	[733]	busi siriz	[1249]
boulé-kannon.....	[696]	bututsi	[928]
boundou	[1185]	buunaati	[973]
boyti.....	[1621], [1623]	buunaati tyabisi	[931]
breu-branco.....	[182], [204], [209]	buxixu	[1054]
breu-branco-verdadeiro	[196]	buxixú-canela-de-velho.....	[1086]
breu-cicantá.....	[190]	bwa-agouti	[1142]
breu-de-leite	[18], [19]	bwa-amadou.....	[536], [1000], [1002]
breu-de-tucano	[1567]	bwa-angi	[100]
breu-grande	[181]	bwa-anmé	[1547]
breu-inhambu	[209]	bwa-ara	[912], [913]
breu-manga	[181]	bwa-bago	[774]
breu-maxixe.....	[1136]	bwa-bal	[1120]
breu-preto	[188], [202], [211]	bwa-bandé	[1374]
breu-sucupira	[893]	bwa-bannan	[536]
breu-sucuruba	[198], [221]	bwa-bich	[254]
breu-vermelho	[184], [188]	bwa-blanché	[168], [169], [400], [402], [407]
buçú.....	[151]	bwa-bouchon	[1000], [1001], [1002]
budu'i.....	[789]	bwa-chandèl	[1209]
bugu.....	[145]	bwa-chapél	[1441], [1442]
bugu batibataa.....	[794]	bwa-charbon	[440]
bugu bugu.....	[955]	bwa-chini	[110]
buiuçu.....	[892]	bwa-dat	[548], [549], [550], [551], [552], [934], [970]
bukutru-ateu	[270], [322], [330], [334], [338], [1589]	bwa-dé-ròz	[583]
bukutru-ateu-priye	[324]	bwa-dibè	[248]
bukutru-gateu	[270], [322], [330], [333], [334], [338]	bwa-dilèt	[110], [129], [533]
bukutru-gateu-duwē.....	[264], [281], [326], [340]	bwa-diven	[1389]
bukutru-gateu-priye	[324]	bwadiwosu	[581], [583]
bukutru-gateu-seinó	[286], [321]	bwa-dyab	[516]
bukutru-gateu-seinó-puvemna.....	[281]	bwa-fé	[751], [752], [1095]

Appendix 2. — Continuation.

bwa-fidjé	[1159], [1162], [1167], [1169],	bwa-pagód-ti-féy	[740]
.....	[1173], [1174], [1178]	bwa-palika	[1440]
bwa-fizi	[1487]	bwa-parasòl	[429]
bwa-flanbo	[1366], [1581], [1582], [1583], [1584],	bwa-patagay	[982]
.....	[1587], [1589], [1590]	bwa-pedri	[1371]
bwa-flèch	[1092], [1095], [1098], [1105]	bwa-pian	[715]
bwa-fronmi	[427], [963], [1406]	bwa-pidjan	[1520], [1522], [1523]
bwa-golèt ..	[264], [267], [268], [320], [326], [335], [338],	bwa-pièr	[1437], [1438]
.....	[1589]	bwa-pistelé	[1121]
bwa-graj	[1419]	bwa-poul	[1769], [1770], [1771]
bwa-graj-rouj	[1422]	bwa-ranmié	[234], [508]
bwa-graj-roz	[1421]	bwa-rouj	[537]
bwa-gro-bèk	[1120]	bwa-rwé	[363], [365], [371], [372]
bwa-jako	[1119], [1529]	bwa-sann	[1215]
bwa-jelé	[1028], [1029]	bwa-serpan	[978]
bwa-kaka	[535]	bwasiman weko	[952], [975], [977]
bwa-kalimé	[1120]	bwa-sip	[419]
bwa-kalou	[1000], [1001], [1002], [1031],	bwa-sira	[975]
.....	[1032], [1034], [1036]	bwa-sitronél	[1213], [1214]
bwa-kannel	[603]	bwa-tabak	[561], [562]
bwa-kannon	[1737], [1739]	bwa-takari	[96], [98]
bwa-kapayou	[766]	bwa-vach	[106]
bwa-kay	[1641]	bwa-vyolèt	[918], [919], [920]
bwa-kayman	[222]	bwa-yawa	[221]
bwa-kochon	[540], [541]	bwa-yaya	[181]
bwa-kotlèt	[1762]	bwa-zanmand	[1529]
bwa-koton	[1010], [1011]	bwa-zoranj	[443]
bwa-kouyè	[120]		
bwa-krayon	[439], [440], [441], [443], [444]		
bwa-krobo	[955]		
bwa-kruzo	[1802]		
bwa-kwachi	[1710]		
bwa-lamori	[794]		
bwa-lansan	[184], [187], [188], [195], [209], [219], [221]		
bwa-lansan-blan	[191], [196]		
bwa-lansan-gran-bwa	[202]		
bwa-lansan-rouj	[208]		
bwa-lasenjan	[137], [138]		
bwa-léchél	[1141], [1724]		
bwa-lélé	[517], [519], [1028], [1029]		
bwa-lèt	[1149]		
bwa-lèt-blan	[1198], [1199]		
bwa-lèt-moucheté	[1149]		
bwa-lèt-rouj	[1152]		
bwa-lòm	[234], [1014]		
bwa-makak	[96], [98], [740], [906]		
bwa-mao	[1037]		
bwa-maypouri	[540], [541]		
bwa-mèl	[1041], [1043]		
bwa-mondan	[1148]		
bwa-négrès	[1437], [1438]		
bwa-noyo	[1433]		
bwa-pagay	[955], [1442]		
bwa-pagód	[816], [818]		

C

caá-pitiú	[1718], [1720]
caá-pitiú-de-folha-grande	[1719]
caa-xió	[591]
cabeça-de-macaco	[1401]
cabeça-de-negro	[1014]
cabelo-de-cotia	[1391], [1535]
cabelo-de-negro	[494]
cacao rivière	[1022]
cacao-branco	[3]
cacaoyer	[1038]
cacau	[1038]
cacauí	[1040]
cacaurana	[1040]
cachaceiro	[1509]
cachimbo	[717]
café-do-mato	[1530]
caferana	[1141], [1395], [1396], [1397], [1530], [1540],
.....	[1709], [1724]
cafézeiro-do-mato	[1539]
cafezinho	[254], [1371], [1733]
cafezinho-do-mato	[1549]
caiaxió	[591]
caimbé	[438]

Appendix 2. — Continuation.

caimbérana.....	[1743]	caroba	[164]
caiuia.....	[1080]	carrapeta.....	[1123]
cajá.....	[13]	carvalho-vermelha	[1422]
cajarana.....	[1110]	carvoeiro	[1323], [1325]
cajou sauvage.....	[9]	casca-doce	[1700], [1701], [1702]
caju	[8]	casinga-cheirosa.....	[1529]
cajuaçu.....	[7], [9]	casquinho.....	[1055]
caju-da-mata	[7], [9]	castanha-de-cotia.....	[260]
cajueiro	[8]	castanha-de-periquito	[173]
cajueiro-do-mato.....	[7], [9]	castanha-de-piriquito.....	[1034]
cajurana.....	[1711]	castanha-de-porco	[17], [500], [514]
cajurana.....	[1712]	castanha-do-Pará	[684]
calango-cego.....	[1762]	castanha-fedorenta.....	[715]
caligni	[331]	castanha-jacaré	[687], [688]
calliandra.....	[997]	castanharaña.....	[703]
camaá.....	[562], [1576]	castanha-sapucaia	[729]
cambara.....	[562]	castanheira	[684]
cambuí.....	[1292], [1295]	catauré.....	[239]
camutim.....	[1095], [1099]	catiguá.....	[1135]
canari macaque.....	[729]	catinga-de-porco.....	[505]
canela-de-jacamim.....	[1772], [1776], [1782]	catuaba.....	[1079]
canela-de-veado	[1549]	cauarú-caá	[419]
canela-de-velho.....	[1508]	cauchorana	[1192]
canela-fedida	[616]	cavaqueiro-do-baixo	[1527]
canela-pimentá	[647]	caxinguba	[1169], [1173], [1174]
canelarana	[498]	caxinguba-rana	[1192]
canjarana.....	[1110]	cebola-brava	[366]
cansanção	[1761]	cebola-grande-da-mata	[365], [371]
capança	[1511], [1550], [1553]	cèdre	[574]
capança-braba.....	[1528]	cèdre argenté	[624], [637]
capança-vermelha	[1453]	cèdre bagasse	[181], [636]
capitari.....	[169]	cèdre blanc	[181]
caporé	[1511]	cèdre cannelle	[603]
capororoquinha	[1410]	cèdre d'argent	[637]
capoteiro	[1035]	cèdre jaune	[652], [666], [671]
caraipé.....	[281], [282], [312], [313], [317],[319], [321], [322], [335]	cèdre marécage	[576]
caramuri	[1616]	cèdre noir	[636]
caramuri-da-várzea	[1631]	cèdre rouge	[1114]
caramuri-preto	[1648]	cèdre sam	[416]
caramuru.....	[912]	cedro-balata	[1113]
caranha.....	[176]	cedro-bravo	[1133]
carapa.....	[1111], [1112]	cedro-do-pantano	[222]
carapanauá	[99], [101]	cedroí	[15], [16]
carapanauá-branca	[102]	cedro-pardo	[650]
carapanauá-cinzero	[1442]	cedrorana	[762], [1121]
carapanaúba-preta	[97]	cedro-verdadeiro	[1114]
caripé	[281], [282], [312], [313], [315],[317], [319], [321], [322], [335]	cedro-vermelho	[1114]
caripérana.....	[303], [304], [334], [339]	cerise de Cayenne	[1249]
caripérana-branca	[293]	chacrona	[1461]
carne-de-vaca.....	[1422]	chapéu-de-sol	[419], [432]
carniceiro	[1528]	chatengn	[451], [458], [460], [463]
		chawari montagne	[246]
		chawari rivière	[247]

Appendix 2. — Continuation.

chawari-dilo	[247]	curiola	[1686]
chawari-gran-bwa	[246]	cururu	[777], [1655]
chawari-rivié	[247]	cutite	[1671]
chawari-rouj	[246]	cutite-grande	[1670]
chichuasca	[250]	cutitiribá	[1671]
chicle-bravo	[1628]	cutitiriba-rana	[1688]
chipa	[209]		
chorão	[283]		
chupa-ferro	[1506]	D	
chuva-de-ouro	[760]	dasitan	[918], [919]
cinzeiro	[404], [407], [411]	datka-arivra	[933]
citron de mer	[1375]	débenn-vert	[159], [162]
citronelle	[1213]	dedalinho	[986]
coataquiçaua	[918]	dima	[508]
coatindiva	[234]	diya papay	[1740]
cóco	[148]	djadjamadou	[1209]
cocotier	[148]	djadjamadou-gran-bwa	[1207]
cœur dehors	[788]	djadjamadou-marikaj	[1210]
copahu	[766]	djadjamadou-montangn	[1206], [1207]
cpaia	[164]	djawi uhu	[91]
cpaíba-branca	[766]	dji-manmen	[28], [33]
coqueiro	[148]	djinaati	[818]
coquirana-brava	[1618]	dobono so	[778]
coração-de-negro	[952]	dokali	[1151], [1152], [1153]
coralleira	[1474]	dokali mapa	[121]
coré-grande	[911]	dondomisinga	[915]
corezeiro	[916]	dondowku	[1283]
corosol sauvage	[26]	donsede	[762]
corrupixá	[1631]	doon udu	[1402], [1406]
corticeira	[928]	doonudu	[1402], [1406]
couachi	[1710]	dyaba udu	[1016], [1019]
couali	[1808], [1810]	dyabatee	[1357], [1360]
couali albert	[1803]	dyagidya	[961]
coumarou	[789]	dyakasi weko	[824]
coumarou odorant	[789]	dyakoomata	[1120]
courmier	[106]	dyango	[972]
courbaril	[806]	dyankatu	[1163], [1173]
criuri	[1100]	dyankoo mata	[1120], [1121]
cruili	[1095]	dzádupa uhu	[1470]
cruiri	[1100]	dzádupa'i	[1472]
cuia-de-macaco	[696]	dzádupalali	[95]
cuiarana	[400], [414]	dzadza'i	[181]
cumaí	[114]	dzalatay	[24]
cumarú	[789]	dzandi	[1111], [1112]
cumarú-amarela	[790]	dzata'i	[157]
cumarú-de-rato	[743]	dzawala	[146]
cumarurana	[790], [967]	dzawat móböha	[1718]
cumarú-verdadeiro	[789]	dzelulu	[913]
cumaté	[1282]	dzepapa'i	[1022], [1025]
cumaté-rana	[1283], [1285]		
cupiúba	[535]		
cupuaçu sauvage	[1039]		
cupuaí	[1039]		
cupuí	[1039]		

Appendix 2. — Continuation.

E	
ébène rose.....	[160]
ébène soufrée.....	[160]
ébène verte	[162]
efea.....	[515]
ëhepuk	[106]
ehnaijap	[505]
ëhpaimë	[1027]
ei li.....	[1641], [1685]
eilowakulu.....	[1766]
ëkepüimë.....	[1440]
ekesima	[408]
ekesimaimë.....	[978]
ekesimë	[408]
ele'u.....	[399]
elekesi.....	[877]
elewu.....	[399]
eleyulu.....	[746], [788], [809], [811], [972]
eleyululan.....	[913], [939], [961]
embaúba.....	[1737], [1739]
embaúba-bengué	[1751]
enajop	[505], [508]
encens	[184], [187], [188], [191], [195], [209], [221]
encens blanc.....	[196]
enëmi'o	[1718], [1720]
envira-amarela	[56], [84], [91]
envira-amargosa.....	[49]
envira-atá	[27]
envira-branca.....	[22], [83], [91]
envira-caitetu.....	[43], [51]
envira-caju.....	[24]
envira-cheirosa	[71]
envira-da-mata	[41], [60]
envira-do-mangue	[1037]
envira-fofa	[58]
envira-jodó	[24]
envira-preta	[38], [47], [48], [53], [58], [67],[82], [85], [89]
envira-rolinha.....	[58]
envira-sapotinha.....	[1029]
envira-sarasará	[92]
envira-surucucu.....	[67], [79]
envira-surucucu-folha-miúda.....	[38]
envira-taia	[24], [48]
envira-vermelha.....	[91]
epalai.....	[99], [1442]
ëpalai.....	[1442]
epkui.....	[110]
erva-de-bugre	[1539]
erva-de-lagarto.....	[1528], [1539]
erva-de-rato	[1487]
escorrega-macaco.....	[919], [1440]
espadeira	[797], [800]
espinho-de-judeu.....	[1496]
esponjeira	[915], [916]
esporão-de-galo	[1496]
etekèle	[970]
etune	[1016], [1018], [1019]
etuwe	[321], [1016], [1018]
évéa	[515]
ëwa.....	[1034]
ëwok pokan.....	[67], [79]
F	
faeira	[1419], [1421], [1422]
falsa-quina.....	[1453]
falso-mangue	[399]
farinha-seca	[358]
faux gaïac blanc	[790]
faux simarouba.....	[164]
fava-amargosa.....	[972]
fava-ataná	[911]
fava-bolota	[913]
fava-de-rosca.....	[794]
fava-de-tambaqui	[879]
fava-marimari.....	[761]
fava-uim-amarela	[785]
faveira.....	[877], [969], [979]
faveira-benguê	[912]
faveira-camuzé.....	[936]
faveira-de-bolacha	[970]
faveira-de-chorão	[913]
faveira-de-empingem	[970]
faveirinha	[759]
faya pao	[1717]
fayaati	[164]
feijão-bravo-preto	[236]
feli kwali.....	[1786], [1788]
féréol	[952]
férole	[1152]
feuille d'argent.....	[637]
fève de tonka	[789]
feyati	[164]
féy-chasèr	[157]
féy-darjan	[637]
féy-sab	[438]
féy-tabak	[561], [562], [1719]
figuier	[1167], [1173]
figuier blanc	[1174]
figuier étrangleur	[1174]
fiman	[623]
fimel-manmanyawé	[82]
flambeau rouge.....	[96]
folha-de-couro.....	[1508]
folha-de-prata.....	[624], [637]

Appendix 2. — Continuation.

folha-de-serra [1154]
 freijó [416], [422]
 freijó-branco [416]
 frei-jorge [422]
 fromager [1008]
 fronmajé [1008]
 fruta-de-anel [1576]
 fruta-de-cutia [2], [3]
 fruta-de-jaboti [1249]
 fruta-de-pombo [15]
 frutinheira [1307]
 fruto-de-jacamin [1391]
 fruto-de-jacu [1080]
 fungutii kookoo [354], [356]

G

gaan bii udu [798]
 gaan busi moni [211]
 gaan busi mulunba [1206]
 gaan busi tobitutu [1509]
 gaan bwasiman weko [976]
 gaan dyaba udu [1015]
 gaan kaabasi ana [47], [50]
 gaan kinboto [1700]
 gaan maamadosu [1454]
 gaan moni [176], [219], [221]
 gaan pepee anga sawtu [43]
 gaan tatu tiki [1582], [1590]
 gaan uku tiki [43], [47]
 gaan wiwii bakuuman [1628]
 gaanman udu ana [1398]
 gaïac de Cayenne [789]
 galinha-choca [494]
 gambui [740]
 gameleira [1167]
 gameleira-branca [1169], [1173]
 gangi udu [965]
 gangouti [1565]
 gangouti-blan [1566]
 garrote [1152]
 gaulette [267], [326], [338]
 gaulette à petites feuilles [281]
 gaulette azon [340]
 gaulette blanche [286], [354]
 gaulette blanche à grandes feuilles [321]
 gaulette marécage [339]
 gaulette noire [264], [322], [339]
 gaulette rouge [303], [304]
 gawenti [1368]
 gayak [789]
 gayak-blan [790]
 gayak-rivié [967]

geebi udu [110], [112]
 geli apisi [608], [652], [666], [671]
 geli tyabisi [808], [809], [811], [812]
 geli udu [1385], [1569]
 génipa [1470]
 genipapo [1470], [1471]
 gibomasa [1252]
 giin ati [162]
 gógo purubumna [107]
 gogó-de-guariba [122], [1766]
 gógo-puvemna [108]
 goiaba-de-anta [1043]
 goiaba-do-mato [1315]
 goiabeira-preta [1437], [1438]
 goiabinha [1315]
 golèt [320], [326], [335], [338]
 golèt-azon [340]
 golèt-blan [286], [354]
 golèt-blan-gran-fey [321]
 golèt-endjen [303], [304]
 golèt-fronmi [295]
 golèt-grigri [333]
 golèt-koumaté [267], [268]
 golèt-marikaj [339]
 golèt-nwé [264], [322], [339]
 golèt-rouj [303], [304]
 golèt-ti-fey [279], [281], [282]
 gonfolo bleu [1793]
 gonfolo gris [1797]
 gonfolo kwali [1795], [1796], [1797]
 gonfolo rose [1795]
 goupi [535]
 gouyave-so [1318]
 gouyavié-montangn [1315]
 gouyavié-sovaj [1104]
 gouyav-kaka-poule [1434]
 gouyav-nwé [1434]
 graine merle [1043]
 graine roche [356]
 grão-de-galo [1686]
 graô-de-galo [427]
 graviola-da-mata [27]
 grènn-bich [114]
 grènn-koumarou [1099], [1490]
 grènn-kwata [1666], [1702]
 grènn-malenbé [1323], [1325]
 grènn-mèl [1043]
 grènn-oko [400], [402], [404], [407], [411], [1535]
 grènn-ròch [356]
 grènn-toti [1535]
 grignon fou [1788], [1791]
 grignon franc [673]
 grignon sainte marie [1799]
 grignon-boni [1797]

Appendix 2. — Continuation.

grignon-fou	[1786], [1788], [1791], [1795], [1797]	ihap-kamwi-purubumna	[463]
grignon-rouj	[673]	ihap-kamwi-seine	[454]
grignon-sentmarie	[1791]	ihap-kamwi-washiune	[468]
grigi	[322], [340]	ii	[184], [207], [216], [1350], [1352]
grigri-blan	[354]	iiru	[877]
grigri-rouj	[264], [309]	iiru-purubumna	[883]
griyō	[673]	ikalikanali	[603], [614]
gro-komou	[155]	ikun-kamwi	[1021]
guaçatunga	[1539]	ilakipté	[820]
guaiabinha	[1302]	ilakopi	[1718], [1795]
guaicá	[647]	iliwa	[27], [31]
guajará	[1597], [1598]	iliwaimé	[29], [43], [52], [1461]
guajara-do-igapó	[1706]	imavui	[1002]
guajuru	[262]	imavui-kamwi	[429], [1037]
guamirim	[1233], [1307], [1317]	imbaúba	[1737], [1739]
guamirim-branco	[1292]	imbaúba-branca	[1736]
guanambi	[222]	imbaúba-da-mata	[1740], [1752]
guapeva	[1686]	imbaúba-da-várzea	[1736]
guarabu	[918], [919], [920]	imbaúba-do-vinho	[1756]
guarapiranga	[925]	imbaúbarana	[1749], [1750], [1752]
guarecoló	[128]	imbaúba-vermelha	[1738], [1740]
guariuba	[1154]	imi'i	[1042], [1044]
guariúba-amarela	[1155]	impitit-ahavukune	[1630]
guariúba-branca	[1147]	impitit-washiunó	[1644]
guaxupita	[1506]	inajá	[147]
gwegwe	[928], [929]	inajai	[157]
gyebi udu	[110], [112]	inajarana	[1028]
		inajarana-envira	[1020]
		inam-etni	[1249], [1772]
		inam-etni-ayeweyo	[1766]
		inam-etni-duwo	[1252], [1285]
		inam-etni-kamwi	[1252], [1256]
		inam-etni-priyo	[1301]
		inam-etni-seine	[1233], [1456]
		inam-etni-seinó	[1215], [1637], [1681], [1702]
		inamkat-duwé	[1252]
		inamkat-seinó	[1215]
		inamkat-wahuyo	[1252]
		inámu pita	[49], [1307]
		inámu sí	[557], [1724], [1733], [1766]
		inámu sí sili	[1188]
		inámu sí wila	[1766]
		inámusi wila	[557], [1724], [1733]
		inaya	[147]
		ingá	[814], [815], [839], [844], [845], [857], [860]
		inga átā	[820]
		inga dju	[858]
		inga kala	[820], [844], [850]
		inga katapali	[866]
		inga láli	[791]
		inga láni	[791]
		inga mani'o	[857]
		inga masulapa	[815]
		inga mulua ya	[819], [820]

H

haka	[747]
halikuli	[157]
halikwana	[1157]
hapélek	[1010]
hapi	[1316]
hawahawaju	[934]
heli apotpé	[912]
hévéa	[515]
heymassoli	[1375]
houmiri baumier	[537]

I

iáyū	[1772], [1776], [1782], [1784]
ididiju	[933]
iduk-veiti	[936]
iduras-veiti	[522]
iduras-vey	[522]
idyakopi	[1718]
ihap-kamwi	[171], [454], [458], [460], [499], [501]
ihap-kamwi-duwé	[451]
ihap-kamwi-priye	[471]

Appendix 2. — Continuation.

inga pini	[814]	inharé.....	[1185]
inga piyã.....	[850]	inharé-folha-miúda.....	[1196]
inga sili.....	[817], [832]	inharé-paina	[1185]
inga sisi	[816], [818], [850]	inimopo'i piyãnga.....	[1323], [1325], [1328], [1329]
inga sisi pay	[843], [860]	inura	[309]
inga sówã.....	[837]	inutauviye	[296], [299], [301], [303], [306], [307]
inga takwá u	[824], [839]	inuuman	[246], [451], [458], [460]
inga tawa	[858]	inuva	[95]
inga tupe'it	[824]	inuva-kamwi	[1490]
inga tupewi.....	[824]	inyai	[859]
inga u	[816], [845]	inyamu bati	[1201]
inga ù	[818]	inyamu ekunali wewe	[1137]
inga u po'i	[814]	inyau	[859]
inga u sówã.....	[818], [860]	inyekulan	[765]
inga wasa	[825]	ipê	[162]
inga yiwa puku	[825]	ipê-branco-do-brejo.....	[169]
inga yowa	[842]	ipê-da-várzea	[770]
inga yowa puku	[825]	ipê-do-igapó	[879]
ingá-banana.....	[861], [866]	ipê-rana	[770]
ingá-branca	[816], [820], [823]	ipewi	[879]
ingá-cabeludo	[821]	ipewi pilã.....	[770]
ingá-caixão	[861]	ipo átã	[249]
ingá-canela	[846], [853]	irimwi	[692], [693]
ingá-chata.....	[858]	isao.....	[152]
ingá-cipó	[825]	isau.....	[152]
ingá-cipo-preta	[863]	isauminyo.....	[1124]
ingá-costella	[820]	isoimë watkï.....	[517], [521]
ingá-de-baixo.....	[858]	isuu-ã	[955]
ingá-de-leite	[853]	isuu-ára	[955]
ingá-de-macaco	[864]	isuu-ára-kamwi.....	[99], [102]
ingá-de-pelo	[856]	isuu-aveya.....	[1723]
ingá-de-veado	[820]	isuu-gaáha	[99]
ingá-do-baixo	[977]	isuu-vey	[1723]
ingá-do-brejo.....	[866]	itaúba	[614]
ingá-dura.....	[977]	itaúba-abacate	[982]
ingá-duro	[859]	itaúba-rana	[872]
ingá-facão	[820]	itey-avain.....	[531], [533]
ingá-feijão	[823]	itey-avan.....	[531], [533]
ingá-ferradura.....	[832]	itukuyen	[389]
ingá-ferro	[820], [843]	ityulanano anakoko	[898]
ingaí	[837]	ityulanano kupešini	[345]
ingá-peua	[842]	ityulu anakoko	[897]
ingarana	[824], [976]	ityulutano kuwasini	[1173]
ingarana-da-terra-firme.....	[978]	ityulutano yapopale	[301]
ingarana-vermelho.....	[879]	ityumban kalapa	[1112]
ingá-verdadeiro.....	[821]	ityutano kilikili maululu	[1009]
ingá-xixi	[818]	ityutano kwai	[400]
ingá-xixica	[816], [818], [850]	ivaí	[1141], [1724]
ingazinho	[832]	ivatingi	[1016]
ingi noto	[684]	ivey	[1762]
ingi pipa.....	[689], [690], [691], [692], [693], [695]	iwa pane	[591]
ingi udu	[505], [508]	iwa pe.....	[1185], [1193]
ingi waway	[365]	iwa pitã	[1249]
ingii pipa.....	[689], [690], [691], [692], [693], [695]	iwa tá'y	[321], [356]

Appendix 2. — Continuation.

iwa tā'iy sili.....	[358]	jawala.....	[141]
iwā ū.....	[1218], [1233]	jawala mili.....	[788]
iwa'e.....	[574], [643]	jawilëkë.....	[1718]
iwaka'i.....	[1360]	jenejapo.....	[958]
iwaka'i sili.....	[1357]	jenëkapo.....	[427]
ian-ethni.....	[5], [1439]	jénipa.....	[1470]
iwawo.....	[1136]	jenipapinho.....	[943]
iwawo sówī.....	[1129]	jenipapo.....	[1470]
iwē'i.....	[1714]	jenipapo-do-campo.....	[1503]
iwen-hetni.....	[5]	jenipaporana.....	[715], [716]
íwi.....	[61], [65], [67]	jinipa.....	[1470]
íwi li.....	[31]	jinipa-dilo.....	[1472]
íwi lo.....	[27]	jinipa-rivié.....	[1472]
íwi lū.....	[26], [29]	ító	[1121], [1127]
íwi lū sī.....	[67]	ító-branco	[1133]
íwi lū sili.....	[67]	ító-da-terra-firme.....	[1119], [1122]
íwi sī.....	[1031], [1032], [1034], [1036]	joão-mole.....	[1329]
íwi tay.....	[24], [27]	joérana	[913]
íwi tay sili.....	[24]	jonndèf.....	[1666], [1671], [1682]
íwidju.....	[1373]	juçara	[150]
íwi.....	[1621]	julu	[913]
íwiiyu.....	[1373]	juqueiri-guaçu	[932]
íwikwisí.....	[286], [1552]	jurema-branca	[740]
íwilá.....	[1621], [1622]	jurubeba.....	[1721]
íwilá pilá.....	[1622]	jutaí.....	[806]
íwilú e'e.....	[35]	jutaí-açu.....	[806]
		jutaí-mirim.....	[777]
		jutaí-rana.....	[774]

J

jaboti.....	[1786], [1788]
jaboticaba-rana.....	[1317]
jaboti-da-terra-firme.....	[1786], [1788]
jacareúba	[222]
jacaterão.....	[1070]
jalá	[1571], [1589]
jalëkë.....	[1718]
janitá.....	[1154]
japananim.....	[912]
jaraí.....	[1706]
jaraná-amarela	[726]
jaraná-mirim	[717]
jarandeu.....	[976]
jata.....	[157]
jatauba	[806]
jatereu	[722]
jatobá	[806]
jatobaí-do-igapó	[896]
jatuaúba	[1121]
jatuaúba-branca.....	[1120]
jatuaúba-preta	[1119], [1123]
jauari.....	[141]
jaune d'œuf	[1666], [1682]
jaune d'œuf à grandes feuilles	[1604]

K

ka'a ki.....	[3], [522]
ka'a pelele.....	[1410], [1460]
ka'a polopi.....	[1154]
ka'a sala	[1329]
ka'a sili	[436]
ka'a wewe	[964]
ka'i ákáni	[117]
ka'i ka'i nu'a.....	[695], [717]
ka'i ka'i nu'a sili	[698]
ka'i láká ni	[117]
ka'i nu'a	[729]
ka'i pimá	[1762]
ka'i tuli	[422], [631]
kaabasi ana	[42], [47], [51]
kaabasi bon	[696]
kaapa.....	[1111], [1112]
kabanafoo.....	[731], [805], [933]
kabuman	[1566]
kafé-lane.....	[1709]
kafé-sovaj	[1550]
kagegut.....	[522], [1257], [1282], [1307]
kagegut-ahavukune.....	[1282]

Appendix 2. — Continuation.

kagegut-puvemna	[1219], [1251], [1282]	kaloj ēwa	[63], [81], [91]
kagegut-puveyo	[1251]	kalojimë	[91]
kahambag	[13]	kalu melei	[1597], [1598]
kahikti	[147]	kamala kuyulu	[1079]
ka'i piye	[606]	kamata	[1070], [1079]
kaibune-ā	[978]	kamata-kamwi	[1064], [1070]
kaibune-āra	[978]	kameli	[1597], [1598]
kaikui amoman	[576], [1184], [1193]	kamina weko	[863]
kaikui wapa	[1566]	kana halale	[1178]
kaikushi anyali	[41]	kananbouli	[1712]
kaisu	[1114]	kananbuli	[1712]
kaisu pilā	[1114]	kanapali	[1712]
kaisu sī	[1114]	kanapupali	[1683]
kajou	[8]	kanari-makak	[729]
kajou-gran-bwa	[7], [9]	kanawa popalin	[1628], [1631]
kaju	[805]	kanawaimë	[1800]
kaka a buuku	[952]	kanegma	[1143], [1550], [1724]
kaka-chien	[806]	kaneli apisi	[603]
kaka-enrièt	[1053]	kanike	[1523]
kakao	[1038]	kankan udu	[1000], [1001], [1002]
kakao-gran-bwa	[1040]	kankantii	[1008]
kakao-larivyé	[1022]	kapasi amandaa	[114]
kakao-sovaj	[1022]	kapasi pomiidyi	[43]
kakaulan	[1769]	kapasi tuno	[1438]
kakaw	[1038]	kapasi wiwii	[1718]
kalai ka'a	[1225]	kapaya watı	[1490]
kalaipé'i	[733], [734], [736], [805], [877]	kapia'i	[1039]
kalaipé'u	[733], [805]	kapokier	[1008]
kalaipiu	[733]	kapuwa nana	[1051]
kalalawa akunepili	[402], [404]	karapa	[1111], [1112]
kalamüli	[1616], [1690]	karapa-oyak	[17], [18], [19]
kalamüli sili	[1616]	karavia	[1119]
kalamüli sili	[1597]	karavia-kamwi	[742], [1118]
kalapa	[1111], [1112]	karegut	[522], [1257], [1282], [1307]
kalapabosi	[789]	karegut-puvemna	[1219], [1251]
kalapaimë	[309], [1123]	karegut-seine	[933]
kalapa-oyak	[1120]	karevru	[805], [912]
kalapapo:si	[789], [790]	karevru-danō	[402]
kalapapolı	[356], [358]	kasa	[794]
kalape'u	[733], [805]	kasanyan	[247]
kalapi alekane	[1127]	kasigu	[770]
kalapi alikan	[1788]	kasila	[522], [1310]
kalapime	[1240]	kasili wewe	[1389]
kalate to tälawáwá	[1743]	kasilikita'i	[1550], [1553]
kalaujalan	[733], [975]	kasilikita'i	[1550], [1553]
kalawilulan	[1213]	kasilikita'i sili	[1549]
kaliaku la ka'a	[1122], [1124], [1125]	kasilikita'i sili	[1549]
kaliaku mili	[1102]	kasim	[33]
kaligni-blan	[331]	kasim-kamwi	[35]
kalima'i	[1565], [1566], [1567]	kasimya	[31]
kalima'i sili	[1565]	kasimyalan	[26]
kalima'i yowa	[1565]	kasiu-van	[427]
kalimai'sī	[1566], [1567]	kaston	[41]
kalipoimë	[339]	kasu	[8]

Appendix 2. — Continuation.

kasun.....	[8]	kokoepu	[148]
katanom.....	[1301]	kokonoto bon.....	[148]
katapali	[866]	kokoonto bon.....	[148]
katatru-avain	[1009], [1011]	kolokolo.....	[321]
katatu-avain.....	[515], [1009], [1011]	komou	[153]
katauma	[400]	komu	[153]
katsu	[1114]	konfitir-makak	[376], [1454], [1455]
katu.....	[1157], [1167], [1174]	kongo.....	[107]
katulimia.....	[790]	kongo-ama	[107]
katulimya	[400], [401], [408], [414]	konoimë.....	[324]
katumá.....	[400]	konolie.....	[234]
katun udu	[1009], [1010], [1011]	konomelulan	[930]
kaupi.....	[696]	konopo.....	[1430], [1431]
kaw udu	[1146]	konoto epi.....	[262]
kawap.....	[526], [1137], [1138]	kookoo.....	[281], [286], [296], [319], [321], [322], [326], [333], [334], [338], [339], [340]
kawap-kamwi	[211], [1124], [1133]	kooponyo.....	[816], [850]
kawe lá	[254]	kopaya.....	[164]
kawee babun udu	[1209]	kopi.....	[535]
kawee mulunba	[1209]	kopi kwali	[1805]
kawukwine-awak.....	[1745]	kopu.....	[696]
kayman buba boyti.....	[1623]	korosol-gran-bwa.....	[30]
kayman udu	[1529]	korosol-kamwi.....	[43], [241]
kee maka udu	[1522], [1523]	korosol-sovaj	[27], [31]
kee udu	[1522], [1523]	kotika uman gonbe.....	[1474]
kelekele	[1761]	kotkotoloju	[933]
kelekele sili	[1771]	koton-périch	[173]
kélépukaju.....	[1582]	koton-siam	[1009], [1011]
kenkii udu.....	[1500]	koubari.....	[806]
ki'elei.....	[778]	koubari-savann	[1708]
ki'elepun	[1710]	koumaté	[1282], [1291], [1292]
kiabisi.....	[770]	koupawa.....	[766]
kiapoku	[1504]	koupaya.....	[164]
kigiksau	[23], [54], [65], [67], [71], [78]	koupiton	[376]
kiiki matakí	[380]	koupou-asou-sovaj	[1039]
kiikii tiki	[1029]	kouy-serjan	[696]
kiinsikin banda.....	[1240]	kowai epi.....	[72]
kilikili amoti	[1036]	kuapitano	[1249]
kilikili maululu	[1010], [1024], [1034]	kuasi bita	[1710]
kinboto	[1702], [1703]	kuasimyan	[1161]
kinolo epi	[1080]	kuasini	[1174], [1175]
kinoto epi	[788]	kuata mopeli	[1387]
kinuwup.....	[1442], [1454], [1455], [1457]	kuata tumali	[729]
kiriksau	[23], [54], [65], [67], [71], [78]	kubaka	[1008]
kisipyulu ulemali	[692]	kubešini	[354], [356]
kisipyululan	[647]	kubi'i.....	[535]
kiwiliimë	[757]	kudagman	[574], [598]
kiyelapolan	[497]	kudagman-purubumna.....	[574]
kiyelewu	[778]	kudagman-wašiunó	[892]
koai	[152]	kudumig.....	[1086]
kobe	[1034], [1035]	kudumig-duwë.....	[1053]
kodya weko	[859]	kudumig-kamwi	[1042], [1049]
kofilan	[1452]	kudumig-seine.....	[1041]
koko.....	[148]	kuepi.....	[321]
kokobe weko	[952], [975], [977]		

Appendix 2. — Continuation.

kuipali	[12]	kuluwi	[1437], [1438]
kuje	[120]	kuma	[106]
kujeimë	[1500]	kumá	[106]
kukundiefu	[374], [376], [377]	kumak	[1008]
kula waju	[265]	kumaka	[1008]
kula:la	[222]	kumakalan	[516], [533]
kulali wepo	[1764]	kumak-kamwi	[245]
kulalu	[1802]	kumalaki	[774], [942]
kulani'i	[234]	kumalaki sili	[268]
kulata	[438]	kumanti udu	[96], [98]
kulataju	[1341]	kumati	[1216], [1240], [1248], [1254]
kulawa:i	[1790]	kumati sili	[1239], [1296]
kulawaja	[303]	kumati u	[1216], [1240], [1248]
kulawaju	[303]	kumboeta ši	[281], [283], [292], [293], [297]
kule kule	[1737]	kumboeta sipyoti	[283], [292], [293]
kule kule tapelelapöt	[1740]	kumboeta šipyoti	[302]
kule pi	[1752]	kumesi	[1315]
kule'i	[586], [746]	kumesi sili	[1282], [1283], [1305]
kule'i sī	[576]	kumete	[1076], [1307]
kulebogo	[1006]	kumeti	[303], [1076], [1216], [1307]
kulebogo tikalayi	[429]	kumetilan	[964], [1791], [1802]
kuleglī	[1737]	kumidan	[1527]
kulekle	[1737]	kumiklan	[234]
kuleku	[239]	kumin	[153]
kulekulan	[742]	kumu	[153]
kulekule ilebe	[1736]	kunapo	[1430], [1431]
kulekule ilupiwat	[1736]	kunapolan	[364], [369], [508], [982], [1417]
kulelu	[239]	kunatepi	[924]
kulikuli inga	[818]	kunau	[463], [1433]
kulisa	[120]	kunawa	[1373]
kulisili wokulu	[832]	kunawalu:i	[1433]
kulitu	[207]	kuni weju	[1087], [1070]
kului ahmit	[858]	kunui	[496]
kulukai	[650]	kunuliyé	[234]
kululu inga	[864], [975], [977]	kupa	[364]
kululu inga sili	[867], [975]	kupahja	[164]
kulumá kalate	[1751]	kupaiwa	[766]
kulumá pilā	[1756]	kupaiya	[164]
kulumá sī	[1751], [1753], [1758]	kupaja	[164]
kulumá tololo	[1749], [1750], [1752]	kupajwa	[766]
kulumnë	[293], [294]	kupali elepali	[1769]
kulumnë epí	[293], [297]	kupawa	[766]
kulumoto	[1086], [1616], [1666]	kupaya	[164]
kulunë epku	[293], [297]	kupayalan	[169]
kulunë ki	[287]	kupë	[1470]
kulupi'i	[984]	kupeïwa	[766]
kulupitá	[374], [376]	kupeïwa	[768], [769], [798]
kulupitá átā	[1242]	kupemi	[955]
kulupitá i	[374], [377]	kupešini	[354], [356]
kulupiyñ	[1201], [1203]	kupešinilan	[283], [292], [293], [295], [297]
kulupiyñ u	[1205]	kupi'i	[535]
kuluwa:i	[419], [429], [430], [431]	kupuh-ahavukune	[1039]
kuluwa:i sili	[,], [419]	kupuh-kamwi	[1039]
kuluwa:i u	[432]	kurip	[157]

Appendix 2. — Continuation.

kusali yepo	[1381], [1648]	kwadi	[1791]
kusangula	[522]	kwai	[400]
kusapoi.....	[363], [365], [371], [1417]	kwailan.....	[563]
kuše.....	[120]	kwairu.....	[247]
kusewelan.....	[451], [458], [460], [1019]	kwairu-kamwi	[567]
kusi wewe.....	[1500]	kwali [1788], [1800], [1802], [1806], [1808], [1810]	[1788], [1800], [1802], [1806], [1808], [1810]
kusi'í	[788]	kwali	[1791], [1802], [1810]
kusi'u së'ë	[257]	kwali e'e	[1810]
kusili ape	[150]	kwali pilá..... [1800], [1803], [1810]	[1800], [1803], [1810]
kusili palatali	[1599], [1628], [1646], [1651], [1656]	kwali pitá..... [1800], [1803], [1810]	[1800], [1803], [1810]
kusili sī ka'a	[1461]	kwali sī	[1799]
kusili yepo	[1788]	kwali sili	[1786], [1788], [1790], [1803]
kusisi	[501]	kwaluk	[778]
kutali.....	[751]	kwanal'í	[234]
kutsili palatahan	[1637]	kwanda	[720]
kuuku-ára-purubumna	[60]	kwapo'i..... [1159], [1178], [1183], [1742]	[1159], [1178], [1183], [1742]
kuuku-araybu-seine	[354]	kwapo'i	[1167]
kuuku-ariut	[1553]	kwapo'i átā	[1745]
kuuku-ariut-kamwi	[1029], [1772]	kwapo'i kalate	[1743]
kuuku-ariut-priye	[1527]	kwapo'i sili	[1171], [1182], [1183], [1330]
kuuku-ariut-seine	[1552]	kwapo'i sówî	[1157], [1178], [1183]
kuukumwi-priye.....	[53], [71], [72], [81], [82]	kwapo'u..... [363], [365], [371]	[363], [365], [371]
kuukumwi-purubumna	[60]	kwasi	[1710]
kuukumwi-seine.....	[74], [80]	kwasiilikita'i	[1550], [1553]
kuwai	[152]	kwasiilikita'i sili	[1549]
kuwajimë	[1503]	kwasii-muumuuwa	[145]
kuwali	[1791], [1810]	kwasikwasi tiki	[1535], [1539], [1550]
kuwali	[1802]	kwasini	[1169], [1173]
kuwalilan	[1786]	kwašini	[1174], [1175]
kuwaluk	[778]	kwasini	[1169], [1173]
kuwapityano	[1220], [1249]	kwasisi	[1721]
kuwapo'u	[363], [365], [371]	kwašiši	[1721]
kuwasí	[1710]	kwašvan	[1710]
kuwasimyan	[1178]	kwaswe-muumuuwa	[145]
kuwasini	[1169]	kwata bobí	[408], [1598], [1599]
kuwateli.....	[687], [688], [705], [709], [712], [719], [720]	kwata kama	[913]
kuwe enekali	[22]	kwata kaya	[401], [408], [414]
kuwepi	[263], [267], [312], [313], [317], [318]	kwata laway	[41], [733]
kuwepilan	[334]	kwata patu	[729]
kuwepilan tipulu	[330]	kwata pepe	[402]
kuya lá	[555], [1254]	kwata pepee	[400], [402]
kuya'í lá	[1421]	kwata pili	[602], [628], [666]
kuyaken elepali	[1075], [1489], [1494]	kwata poa	[692]
kuyali	[1712]	kwata weli	[978]
kuyali tapaiko	[768]	kwatan patu	[729]
kuyali tapityi	[915]	kwatili	[687], [688], [691], [705], [707], [712], [719], [720]
kuyali topaitye	[915]	kwatili tabulu	[701], [723]
kuyau-kamwi.....	[1250], [1456], [1664]	kwatili tamunen	[720]
kuyau-kamwi-duwë	[1652], [1678]	kwatri	[699], [1430], [1431]
kuyau-kamwi-duwō	[1682]	kwatri-duwë	[1361]
kuyau-kamwi-puvemna	[1659]	kwatri-duwō	[726]
kuyau-kamwi-seinó	[1652]	kwatri-purubumna	[726], [1717]
kwa'íyan	[563]	kwatri-puvemna	[687], [688]
kwachi	[1710]	kwatri-seinó	[721], [727]

Appendix 2. — Continuation.

kwatri-washiune	[383], [389], [726]	lébenn	[162]
kwatri-washiune-duwē	[382]	lébenn-ròz	[160]
kwatri-washiune-priye	[710]	lébenn-vert	[159]
kwatri-washiune-seine	[728]	lebi agi	[896]
kwatri-washiune-seinō	[723]	lebi giin ati	[160]
kwatri-washiunō	[1363]	lebi kookoo	[539]
kwatri-washiunō-duwē	[1363]	lebi luabi	[685], [687], [722], [723]
kwatri-washiunō-purubumna	[1360]	lebi mapa	[121]
kwatsindi	[1174]	lebi moni	[18], [184], [202], [209], [1099]
kwebi kookoo	[264], [267]	lebi musupu	[1041]
kwem	[1750], [1757], [1758]	lebi nenba	[757]
kwem purubumna	[1752]	lebi pindya udu	[548], [551]
kwem-duwō	[1754], [1757]	lebi sali	[181]
kwem-purubumna	[1749], [1750]	lebi tobitutu	[138]
kwem-puvemna	[1754]	lebi tongo	[1283], [1285], [1292]
kwem-seinō	[1753]	lebi tyabisi	[745], [746], [747]
kwem-tino	[1753]	lebi udu	[501], [1637]
kwep	[319], [321]	lebi weko	[816], [850]
kwepan	[1712]	leiteira	[126], [1150], [1153]
kwepan-washiunu	[1329], [1569], [1712]	lele anapamigín	[365]
kwepi	[263], [267], [312], [313], [317], [318]	lele elepali	[745], [747]
kwépi	[358]	lemu epi	[1642]
kwepilan	[291], [331], [335]	liba apisi	[1419], [1422]
kwep-puvemna	[267]	liba awaa	[141]
kwik	[162]	liba gobaya	[1318]
kwik-duwē	[1453]	likanau apisi	[626]
kwik-kamwi	[159], [169]	limãozinho	[1520], [1522]
kwik-marikasmatgene	[169]	lirio-do-mato	[1489]
kwik-washiune	[1515]	lisa pao	[1440]
kwip	[157]	lixeira	[438]

L

lace	[549], [551], [552]
lace-branco	[548], [1527]
lace-vermelho	[550]
lace-vermelho-do-mato	[554]
lamoussé	[74], [89]
lamoussé blanc	[91]
lamuci	[74]
lanjélik	[778]
lanmou-rouj	[1474]
lanmoussé	[74], [89]
lanmoussé-blan	[91]
lanmoussé-fronmi	[427]
lanmoussé-nwé	[85]
lansan-rouj	[181]
lansan-roz	[181]
lansan-ti-fey	[1136]
lapalapa	[96], [98], [110], [859]
lapo-sigar	[692]
laranja-do-mato	[1429]
laranjinha	[56], [1020]

lébenn	[162]
lébenn-ròz	[160]
lébenn-vert	[159]
lebi agi	[896]
lebi giin ati	[160]
lebi kookoo	[539]
lebi luabi	[685], [687], [722], [723]
lebi mapa	[121]
lebi moni	[18], [184], [202], [209], [1099]
lebi musupu	[1041]
lebi nenba	[757]
lebi pindya udu	[548], [551]
lebi sali	[181]
lebi tobitutu	[138]
lebi tongo	[1283], [1285], [1292]
lebi tyabisi	[745], [746], [747]
lebi udu	[501], [1637]
lebi weko	[816], [850]
leiteira	[126], [1150], [1153]
lele anapamigín	[365]
lele elepali	[745], [747]
lemu epi	[1642]
liba apisi	[1419], [1422]
liba awaa	[141]
liba gobaya	[1318]
likanau apisi	[626]
limãozinho	[1520], [1522]
lirio-do-mato	[1489]
lisa pao	[1440]
lixeira	[438]
loka	[806]
loka bii udu	[798]
louro	[598], [652]
louro-abacate	[586], [620]
louro-alho	[416]
louro-aritu	[607]
louro-babão	[616]
louro-branco	[637]
louro-capitiu	[576]
louro-chumbo	[603]
louro-cravo	[607]
louro-da-serra	[986]
louro-do-igapó	[615]
louro-faia	[1419], [1421], [1422]
louro-gamelá	[673]
louro-í	[1129]
louro-itaúba	[614]
louro-peludo	[593]
louro-pimenta	[625]
louro-pirarucu	[603]
louro-prata	[637]
louro-preto	[603], [623], [636]
louro-puxuri	[581]
louro-seda	[637]

Appendix 2. — Continuation.

louro-vermelho..... [673]
 loussé [15], [16]
 luabi [698], [703], [704], [709], [721], [722], [726]
 lunba [1207], [1210]
 luweluwé [1528], [1535]
 luwi [1178]

M

maamadosu [1454], [1456], [1457]
 mabee sii [1042], [1052], [1053]
 macacarecua [696]
 macacaúba [925]
 maçandarubinha [1084]
 maçarandúba [1621]
 maçarandúba-da-terra-firme [1623]
 maçarandúba-de-folha miuda [1624]
 maçarandúba-jacaré [1624]
 maçarandúbarana [1599], [1605]
 macaúba [140]
 macoubé [117]
 macoucou [130]
 macucá-murici [547]
 macucu-bobo [315]
 macucu-farinha-seca [322]
 macucu-fofo [303], [304]
 macucu-murici [273]
 macucu-sangue [306], [307]
 macucu-terra [309]
 madyo uman [1213]
 madyo uman wiwii [567]
 madyoma uwii [567]
 mahayapna [157]
 mahot cigare [690], [691], [692], [693]
 mahot cochon [1034], [1035], [1036]
 mahot coton [1010], [1011]
 mahot jaune [726]
 mahot marécage [725]
 mahot noir [695], [701], [707]
 mahot rouge [722], [723]
 mainyapo [1043], [1064], [1529]
 mainyapo weli [1066]
 maipuli apepiton [1379]
 maipuli elepali [1452], [1487]
 maipuli kiyelapoli [1078]
 maipuli solapali [850]
 maipyuli epu [1029]
 maipyuli italali [1036]
 maipyuli kiyelapole [1487]
 maipyuli omoseli [1034]
 maisyi [53]
 maka mapa [114]
 maka sii [246], [247], [248]

makalawalaimë [309]
 makamakalu [1740]
 makasu [379]
 makit [902], [903]
 makitwa [902], [903]
 makuk [1787]
 makwebo [529], [785]
 makweebo [785]
 makwelebo [943], [946], [948]
 mal bwadiwosu [608]
 malakaipyo [1203]
 malakana kuwa [1108]
 malakapuli'i [1799], [1802], [1808]
 mal-akasya [793], [912], [915]
 malakatano [1135]
 malako [774], [919]
 malaliapu [157]
 malaliapu lo [157]
 malamala [138]
 malay elepali [948]
 malay wokili [1585]
 mal-bwa-dérose [650]
 mal-bwa-kannon [1740], [1749], [1750], [1751],
 [1752], [1753], [1754], [1756], [1757], [1758]
 mal-bwa-makak [913]
 mal-bwa-pian [716]
 malihaliiämë [1515]
 malimali [759], [933]
 malimali sī [933]
 malimali u [758]
 malimiapo [281]
 malipa [147]
 malipa tay [147]
 malipa wewe [746], [788], [809], [972]
 malipaimë [1443]
 malmani [1345], [1349], [1354]
 malobi [1616], [1617], [1618]
 mal-simarouba [1708]
 malupete [74]
 mamanten [1627], [1631]
 mamanyawe [81], [82]
 mamao [1022]
 mamão-do-mato [245]
 mamão-rana [245], [1022]
 mamãozinho [1094]
 mamatki-arin [95]
 mameleiro [499]
 mamhalju [1070]
 maminha-de-porco [1520], [1522]
 mamiqueira [1523]
 mamorana [1022]
 mamorana-da-terra-firme [1010]
 mamorana-grande [1010]
 man agu siton [1494]

Appendix 2. — Continuation.

man musupu witi	[1079]	mao-kaka	[695]
man papay.....	[1740]	mao-kochon	[1031], [1032], [1034], [1035], [1036]
man pendyeku.....	[83]	mao-koton	[1009], [1010], [1011]
man pengeku.....	[83]	maoksi-adudu.....	[536]
man puyo.....	[1188]	mao-kwatarai	[717]
man tapupa	[715], [716]	maomao	[1022]
man tsaka	[775]	mao-marikaj	[725]
manaca-rana.....	[1769], [1770], [1771]	mao-nwé	[695], [701], [705], [707]
manche-hache	[225]	mao-pilon	[717]
mandapuçà.....	[1043]	mao-rouj	[720], [722], [723]
mandasabay.....	[1658]	mao-sigar.....	[690], [691], [692], [693]
mandioqueira	[1796]	mapa	[95], [106], [114], [117], [121], [1151]
mandioqueira-azul.....	[1791]	mapalaba	[236], [515], [1442]
mandioqueira-lisa.....	[1797]	maparajúba.....	[1621], [1623], [1628]
mandioqueira-preta	[1797]	maparajúba-branca	[1642]
mandioqueira-rosa.....	[1795]	maparajúba-vermelha	[1676]
mangabaraña	[1607], [1627]	mapa-ri	[877]
manga-brava.....	[365], [371]	mapatirana	[1750], [1753], [1757]
manga-de-anta.....	[1707]	mapiku.....	[789], [790]
mangu.....	[366], [389]	mapili poá	[164]
mangue-amarelo.....	[399]	mapiwalan.....	[501]
mangue-branco	[399]	mapuxiqui-vermelho	[740]
mangue-d'agua-doce.....	[1429]	marapuama	[1374]
mangue-da-mata	[393]	maras-gamana	[1438]
mangue-de-botão	[398]	maraaura	[766]
mangue-negro	[398]	maravuvuiá.....	[508]
manguerana.....	[382], [383], [388], [389], [393]	maraximbé	[1595]
mangue-vermelho.....	[1430], [1431]	maremi-etni.....	[234]
manhundi kwali	[1786], [1787], [1788]	maria congo	[107], [108]
mani.....	[378], [380]	marimari	[759]
máni.....	[378]	marinaiwa	[185], [188], [191], [216]
mani epu	[380]	marinaiwa-kamwi	[196]
mani'i.....	[703], [726]	marinaiwa-purubumna.....	[206]
manig.....	[30]	marinaiwa-puvemna	[205]
manig-ahavukune.....	[27]	marinaiwa-washiunó	[182]
manig-amutri	[31]	marindiba.....	[405]
manig-ihipiné.....	[37]	marinheiro	[1120]
manig-kamwi	[22], [37]	maripa	[147]
manig-retni	[26]	mari-pwèl.....	[1454], [1455]
maniimé.....	[380]	marirana	[263]
manil marécage	[380]	mariuvra	[992]
manil montagne	[378]	mariuvra-seine	[992]
manimé.....	[378]	mariuvra-washiunó	[508]
manmanyawé	[48], [53], [60], [79]	marmelada	[1434]
manmanyawé-piman	[43], [44], [46], [49], [51]	marmelo-bravo	[1433]
manmanyawé-yanman.....	[67]	marmita-de-macaco	[729]
manni-marikaj.....	[380]	marmite de singe	[729]
manni-montangn	[378]	marupa	[1714]
manni-rouj	[222]	maruparana	[1711]
mantiotio	[98]	marupazinho	[1713]
mantuga maka.....	[1721]	masalaipyo	[1547]
mao-blanc	[687], [688], [707], [717], [724]	masaláni	[1621], [1624]
mão-de-gato	[1185]	masulapa	[815]
mao-jònñ	[726]	masulapa pe u.....	[837], [850]

Appendix 2. — Continuation.

masulapa pilá.....	[837]	mekolo nohunu.....	[444]
mata-calado.....	[1553]	meku huhu.....	[1099]
mataki.....	[107], [108], [378]	meku kuwaili.....	[728]
matamatá	[705], [722], [723], [724]	meku susu	[374], [376], [377], [1489], [1490]
matamata wewe.....	[1616]	mekuhuluimé.....	[1371]
matamatá-azulada.....	[704]	mekulak	[1406]
matamatá-branco	[707]	mekwa.....	[1373]
matamatá-gameleira	[685]	mékwa.....	[1373]
matamataica	[716]	melamelulan.....	[1475], [1477]
matamatá-jibóia	[698]	melancieira	[741], [742]
matamatá-mirim	[712]	melekene	[110], [532]
matamatá-preto	[701]	melekene sili.....	[532]
matamatá-rósea	[703]	melekene u	[110]
matamatá-roxa	[707]	meli.....	[537], [726]
matamatá-verdadeira	[701]	meli lan	[1307]
mata-pasto-grande.....	[934]	meliju.....	[1421]
mata-pau	[365], [371]	meliki udu	[533]
matata	[802]	melimeli	[1141]
matau'í	[58]	mene	[963]
matawai gedu	[964]	mepu	[806]
matolosi	[583]	mëpu	[806]
matula.....	[1401], [1402], [1487]	mëpuimé	[885]
máu'í	[1022]	mésoupou.....	[1043]
máu'í sili.....	[173]	mëwá.....	[893], [951]
maumau	[1022]	miatoá	[1385]
maurissi	[992]	mifungafunga.....	[1002]
mavévé-chien	[1722], [1723]	migukat.....	[596], [608], [628], [663], [669]
mavévé-sikriyé	[1527]	migukat-kamwi	[645]
mavinbi	[1446], [1452]	mihitui	[8]
mavinbi-kamwi	[991], [1446], [1764]	mihitui-kamwi	[9]
mavinbi-priyu	[1770]	mihitui-tivarabuyené	[9]
mavinvi	[1446], [1452]	mihukat	[596], [608], [628], [663], [669]
mavinvi-kamwi.....	[991], [1446], [1459], [1489], [1764]	mihukat-kamwi	[645], [1328]
mavinvi-priyu.....	[1770]	mihukat-kamwi-wahuyo	[645]
maya-kongo	[107], [108]	mijelemili	[1529]
mayawa'í.....	[505], [508]	mijetaimé	[583], [603]
maypa	[147]	mïkotawa	[846]
mbagwi	[1565], [1566]	mïkotowa	[846]
mbagwi-purubumna.....	[1565]	miku ka'a.....	[1245], [1250]
mbagwi-puvemna.....	[927]	miku lapi'a sôwî	[447]
mbagwi-seine	[1566]	miku lapi'a u	[444], [1618]
mbarui	[1565], [1566]	mila kulan	[652]
mbarui-purubumna.....	[1565]	mila suku.....	[1715], [1717]
mbarui-puvemna.....	[927]	mila yowa	[1252]
mbarui-seine	[1566]	mile'i.....	[992], [993]
mbinbi-washiune.....	[1489]	miliki tiki	[129]
me:ku ubupo.....	[1442], [1455]	miliki udu	[106]
me:ku yongay	[1001]	milimi	[74], [90], [91]
mee'í	[164]	milisi	[152]
mejjú.....	[46]	mincouart	[1373]
meikolo muhunu.....	[444]	miniyu lá	[501]
meikolo muhupu.....	[444]	minua'í	[1389]
meikolo tamít.....	[1028]	miraceem	[1700]
meikolo wewe.....	[72]	miraúba.....	[1095], [1102], [1105]

Appendix 2. — Continuation.

miret	[58], [439], [440], [441], [443], [444]	mongii soke.....	[1604]
miret-kamwi.....	[22]	mongo babun udu.....	[1207]
miret-unikweune.....	[79]	mongo gwegwe.....	[909]
mirindiba	[414]	mongo mataki	[378]
mirindiba-branca.....	[400], [402]	mongo toso paasaa	[1203]
mirindiba-doce.....	[514]	moni	[189], [207], [209]
mirindiba-rosa.....	[986]	monokoe	[896]
mirukat	[596], [608], [628], [663], [669]	monokoe u.....	[889], [897], [898], [956]
mirukat-kamwi.....	[645], [1328]	mope.....	[13]
misobisobi.....	[234]	morcegueira.....	[745], [747], [809]
mitiku	[802]	morelia	[992]
mitū luway	[44], [49], [82]	morocototó	[499]
mitū pi.....	[1082]	morototó	[138]
mitū'ay	[1342]	morototó-branco	[137]
miumili	[928], [930]	moukaya	[140]
miumiu-ähinō	[856]	mouriri.....	[1095]
miumiu-akamá-arib.....	[814], [825], [858], [863]	mouroumourou.....	[145]
miumiu-akamá-aribinē	[814], [825], [858], [863]	mourousi.....	[992]
miumiu-asiru	[835], [849], [864], [960], [964]	mousigo	[1201]
miumiu-asiru-wašiunu	[1582]	mousigo-rouj.....	[1203]
miumiu-asukwinō	[846]	moussigot	[1201]
miumiu-avukun-kamwi.....	[833]	moussigot blanc.....	[1205]
miumiu-kasiuminio.....	[824]	moussigot rouge	[1203]
miumiu-maoksi-arib.....	[825], [844], [846]	moutouchi marécage	[928]
miumiu-maoksi-arib-duwē	[833]	moutouchi montagne	[909]
miumiu-maoksi-aribinē	[844], [846]	moutouchi noir	[943], [946], [948]
miumiu-platno	[820]	moutouchi rivière	[928]
miumiu-platno-kamwi	[874]	moutouchi rubané	[929]
miumiu-platno-mna.....	[841]	moutouchi savane	[929]
miumiu-platno-purubumna	[818]	moutouchi-marikaj.....	[791], [928]
miumiu-puveyo	[839]	moutouchi-montangn	[909], [945], [947]
miumiu-sababuyu	[833]	moutouchi-nwé	[943], [946]
miumiu-wašiunu	[859], [960]	moutouchi-ribanné	[929], [930]
miyū'i pepo	[1529]	moutouchi-rivié	[928]
miyuluwa	[195]	moutouchi-savann	[928]
mo:pe.....	[13]	moysi	[152]
moçataíba	[952]	moyu tasi.....	[1406]
moela-de-mutum	[1341], [1342]	moyu'i	[1217], [1250]
mojoi	[419], [505], [508]	mpitit-ahavukune	[1630]
mokaya.....	[140]	mpitit-kamwi	[1317]
molongó	[110]	mpitit-wašiune	[1188]
molongó-de-colher	[120]	mpitit-wašiunē	[222]
molototo	[137], [138]	mpitit-wašiunō	[1644]
molototo'u	[138]	mubulu	[145]
momaj.....	[690]	mucajá	[140]
mombin	[13]	mucuráo	[715]
mombin blanc.....	[15], [16]	muhu	[1159], [1162], [1169], [1173]
mombin fou	[15], [16]	muhu-ivine	[1174], [1178]
momow.....	[1022], [1023], [1025]	muhu-kamwi	[1159], [1162]
monben	[13]	muhu-puvemna	[1160]
monben-blan	[15], [16]	muhat	[928]
monben-fou	[15], [16], [17]	muhat-ahavukunō	[909]
monben-sovaj	[15], [16]	muhat-wašiunō	[1631], [1637]
mõngi.....	[1027]	muira-gonçalo	[1389]

Appendix 2. — Continuation.

muira-jibóia	[751], [752]	murta	[1251], [1252]
muira-pajé	[789]	murta-grande	[1307]
muirapenima	[1745]	murta-parida	[1307]
muira-pinima	[1149]	murtinha	[1282]
muirapiranga	[797], [800], [890]	muruci	[992]
muira-piranga	[1152]	muruci-vermelho	[993]
muirá-pixuna	[885]	murumuru	[143], [145]
muira-puama	[1374]	murupita	[532], [533], [534]
muira-pucu	[1529]	mururé	[1147], [1148]
muiraquatiara	[10]	mururérana	[1151]
muiratinga	[1150]	murusi	[995]
muiratinga-de-folha-miúda	[1188]	musansi	[955]
muiraúba	[1095], [1096], [1098]	musupu	[1044], [1045], [1054], [1070], [1091]
muiraximbé	[1142]	mutamba	[1014]
muiva	[1053]	mutamba-preta	[1016]
mukaya	[140]	mutata	[516]
mukulakula	[774]	mutende	[1440]
mukulap	[1406]	mutende kwali	[1802], [1808]
mukulupiȳ	[1201], [1203]	mutu pupot eimë	[376]
mukwalakwala	[774]	mutu'i	[1068], [1070], [1079], [1080]
mukwisi	[286], [1552]	mutuju	[1202], [1203], [1207]
mulateiro	[918], [919], [920]	mutupupot	[1454], [1457]
mulatorana	[1440]	mutusi	[924], [928], [930], [945]
mule'i	[992], [993]	mutusi e'e	[928]
mule'i ityudano	[995]	mutusi pilá	[928]
mule'ilan	[523]	mutusi piyū	[946]
mulei	[498], [617], [989], [994], [1731]	mutusi sī	[924], [930]
mulei pilá	[993], [1731]	mutusilan	[929]
mulei sī	[1724]	mututy	[928]
mulei sili	[558]	mututy-da-terra-firme	[929], [930]
mulei sówī	[993], [1731]	mutyengetyenge	[885]
mulekju	[1371]	muumu	[145]
mulewabo	[508]	muyū'i pepo	[1529]
mulewalan	[79]	mwanba	[72]
mulisi	[152]		
mulishi	[152]		
mulototou	[138]	N	
mulua ya	[819], [820]	nana polan	[1049]
mulumulu	[145]	néflier	[1043]
mulumulu sī	[143]	neko udu	[889], [924], [967]
mulunba	[1210]	neko udu gaan wiwi	[892]
mulungu	[802]	nēmī'o	[1718], [1720]
mulungu-brabo	[898]	nemo ákā	[200], [208], [209]
mulungu-rosa-verde	[1637]	nengee udu	[260], [286]
mumu	[145]	niaouli	[1277], [1278]
munguba	[1027]	nkasa	[972]
munguba-da-mata	[1015]	nkasa tyabisi	[969]
munguba-rana	[1022]	nkatu	[1157], [1167], [1174]
munpatutpē	[1490]	noix de cajou	[8]
munu'i e'e	[789]	noix de Pará	[684]
munu'i sī	[790]	noix du Brésil	[684]
murarena	[811]	noni	[1483]
murici	[992]	nopitya	[374]
muriri	[1095], [1098], [1099]		

Appendix 2. — Continuation.

noya elepali	[999]
nukunuku	[1025]
nyama suwi	[1059], [1080]
nyanboka	[1598], [1626], [1650], [1652], [1656], [1658], [1662], [1664], [1666], [1667], [1674], [1680], [1682], [1686], [1687], [1688]
nyon udu.....	[937], [938]

O

odun	[1146]
oticica	[331]
oka epu	[48], [63]
okajpu epeli.....	[490]
okalat	[691]
okalatimë	[695]
okoipyo	[1433]
okomë oki	[1389]
okopiaphawin.....	[1743]
olho-de-cabra	[897]
oloi.....	[8]
olojimë.....	[7], [9]
ololome	[377], [1664], [1703]
olome	[377], [1664], [1703]
olomelan	[1655], [1678]
olowaimë.....	[1236]
ombatapo	[242], [1185], [1192]
omole	[749]
omose.....	[83], [85]
omose tubulu	[20]
onohkoi.....	[889]
onohkoimë.....	[892]
onohkowe	[889], [890], [952]
onokoe	[896]
onokoe u	[889], [897], [898], [956]
orelha-de-burro	[527]
orelha-de-macaco	[794]
osi epít	[548], [550], [551], [554]
otono epityi.....	[1211]
oubarouna.....	[501]
ouregou.....	[65]

P

paa letowape.....	[952], [953]
paa lo'o.....	[436], [519]
paansu miti	[766]
paasaa.....	[156]
paata sii	[877]
paawisi besi	[1249]
paaya.....	[1149], [1152], [1193]

pacapuá	[943]
pacovi.....	[92]
pada	[164]
padula	[1487]
padyawa	[821], [825], [834], [850], [856], [860], [863]
pahe-avan	[129]
pahe-avan-kamwi	[1776]
pahe-avan-priyo.....	[119]
pahe-avan-puvemna	[124]
paidya	[1149]
paidya tabile.....	[1152]
paidya tubulu.....	[1149]
paidyawa	[825], [834], [856]
paig	[1149], [1199]
paig-priye	[1195]
paig-purubumna	[1152]
paig-puvemna.....	[1152]
paig-seine	[514], [943], [1149], [1152], [1185], [1196], [1198], [1199]
paila	[1149]
paila lā.....	[1150], [1152], [1196], [1198], [1199]
paila wili.....	[1149]
pailaimë.....	[1152], [1233]
pailawa	[825]
paipaiyo wokulu.....	[382], [389]
paipayo namipay	[1395], [1396], [1397]
pairi.....	[1149], [1199]
pairi-priye.....	[1149], [1195]
pairi-purubumna	[1152]
pairi-puvemna	[1152]
pairi-seine.....	[514], [943], [1185]
pairi-seinō	[1149], [1152], [1185], [1196], [1198], [1199]
paivan	[1627]
pajilé	[223], [224], [226], [227]
pajurá.....	[354], [1683]
pajurá-de-obidos	[1683]
pajurá-de-racha	[263]
pajurá-grande	[356]
pajurá-pedra	[356]
pajurá-verdadeiro	[263], [264], [267], [268]
pajurazinho	[273], [333]
pakad	[1146]
pakasa	[1146]
pakatsa	[1146]
pakea'i	[1285], [1307]
pakea'i sī	[1240], [1308]
pakeli	[1453]
pakih-em	[27], [48], [51]
pakih-em-priye	[44], [82]
pakih-em-purubumna	[41]
pakih-etni	[121]
pakila apisan	[82]
pakila eu	[1339]
pakila hapisan	[82]

Appendix 2. — Continuation.

pakila potay.....	[1044]	pali'i sī.....	[330], [335]
pakila siku.....	[195]	pali'i sili.....	[268], [333]
pakila sipyoli.....	[219], [982], [999]	pali'i towu.....	[320]
pakila šipyoli.....	[206], [207], [208], [209], [216]	pali'i wate'e.....	[354]
pakila udu.....	[563]	palipali.....	[295]
pakila wesopotali.....	[1044]	palipyali.....	[1379]
pakila yuyulu.....	[1049], [1053]	palipyo.....	[279], [324], [1771], [1776]
pakilemu.....	[129]	paliwi	[797], [800]
pakolo talalan.....	[101]	paliwu.....	[797], [800]
pakoweni.....	[1712]	palmié-bache.....	[152]
paku akayu.....	[1712]	paluke.....	[120]
paku inga.....	[837], [853], [862]	palulipo.....	[984]
paku-amana.....	[1226]	palulu ipyo.....	[1788]
pakuku.....	[1688]	palulumili.....	[1389]
pakuku.....	[1645], [1670]	palulumuli.....	[501]
pakulea.....	[1233]	paluwa.....	[1]
pakuli.....	[379]	pamā	[1196]
pakulilan.....	[374], [376]	pana akala	[1425]
pakumila.....	[1304]	paná miwi	[749]
pakumleimē.....	[1784]	pana'í.....	[808], [809], [875]
pakusin-amana.....	[1220], [1233], [1245], [1298]	pana'í wu.....	[17], [747]
pakutpē.....	[497], [498], [1504]	panacoco	[890], [952]
pala'i.....	[164]	panacoco blanc.....	[896]
pala'i wātā.....	[967]	panakoko.....	[889], [890], [891], [896], [897], [898], [952]
palaka pomiidyi.....	[1292]	panakoko-blan	[896]
palakasana.....	[254]	panakoko-marikaj	[875]
palakta.....	[1621]	panao	[809]
palakta jalan.....	[1628]	panapana.....	[1105]
palaktaimē.....	[1621]	panauna-kamwi.....	[570]
palaku'i.....	[101]	panauna-priye	[1605]
palakū'i.....	[99]	panauna-puvemna	[568]
palakusinian.....	[955]	panawilili	[863]
palakusinen	[102], [955]	panda	[169]
palakusinyan	[102]	pandalan	[168]
palakuta pilā.....	[102]	papaati	[918], [919], [920]
palakuta piyū.....	[101], [102]	papacuaió	[1712]
palakwa.....	[1662]	papa-terra	[1490]
palala.....	[442], [444], [447]	papay-bich.....	[95]
palálá ka'a.....	[992]	papaye biche.....	[95]
palalagwa.....	[1022]	papelilan.....	[1138]
palalapo.....	[933]	paracanaúba	[1733]
palalipanuimē.....	[1474]	paracutaca	[952]
palata	[1621]	paracutaca-da-terra-firme	[955]
palatabali.....	[1616]	paraiba	[1714]
palawakasi.....	[922]	parajuba	[777]
palemulu.....	[1721]	pará-pará	[164]
palétuvier blanc.....	[1]	parcourí	[379]
palétuvier gris.....	[399]	pareine-ára	[222]
palétuvier montagne.....	[386], [389]	parépou-djab	[157]
palétuvier rouge.....	[1430], [1431]	paricá	[744]
pali udu.....	[99], [101], [102], [1442]	paricá-branco	[926], [927], [932]
pali'i	[333], [334], [339]	paricá-de-esponja	[915]
pali'i lá	[279]	paricarana-de-espinho	[932]
pali'i pilá	[334]	paricazinho	[805]

Appendix 2. — Continuation.

parinari	[354]	pau-marfim-falso	[123]
parkouri	[379]	pau-mirim	[1722]
pasami	[1242]	pau-mulato	[1440]
pasi'i	[156]	pau-para-tudo	[1707]
pasi'i wapo	[361], [389], [393], [982]	pau-pereira	[107], [108]
pasi'i wapo sili	[382], [388], [389], [393], [982], [1732]	pau-pombo	[15], [16]
pasis	[535]	pau-rainha	[1152]
pasisi	[535]	pau-rainha-roxo	[1149]
pasiwet	[1283]	pau-rosa	[583]
pasiwi	[156]	pau-rosa-branca	[1136]
passuaré	[964]	pau-roxo	[918], [919], [920]
pasyolawa	[1252]	pau-tartaruga	[1149], [1152]
patakwik	[365], [372], [373]	pauteloso	[583]
patakwik-ahinē	[373]	pau-vermelho	[1368]
patakwik-kamwi	[371]	pawe	[919], [920]
patali	[144]	paxiúba	[156]
patau	[154]	paxiubarana	[382], [383], [388], [389], [393]
pataua	[154]	payakousa	[1581], [1582], [1583], [1584], [1587], [1589], [1590]
patauá	[154]	payawa	[821], [825], [834], [850], [856], [860], [863]
patawa	[154]	payawalu	[1418]
pativyé-blanc	[1]	payawalu'i	[1123]
pativyé-gran-bwa	[361]	payawalu'i sili	[1115], [1119], [1126]
pativyé-gri	[399], [1429]	paykuli	[379]
pativyé-janm	[1430], [1431]	payula	[321], [322], [336], [344]
pativyé-montagn	[382], [383], [386], [388], [389], [393]	payula piyū	[331]
pativyé-rouj	[1430], [1431]	payula sili	[333], [335]
patsi'i	[156]	payulilan	[61], [67], [69]
patsitsi	[535]	payuy	[1]
patula	[1401], [1402], [1487]	peigne macaque	[1000], [1001], [1002]
patulapo	[1036]	peilili	[786]
patumu	[1001]	pëinékë emnamotem	[1002]
pau-bicho	[733]	peke'a	[248]
pau-d'arco	[159]	peke'a lá	[246]
pau-d'arco-amarelo	[162]	pekea	[248]
pau-d'arco-rana	[567]	pékéya	[248]
pau-d'arco-roxo	[160]	pekui	[1710]
pau-de-arapuce	[1565], [1566]	pelepele	[365]
pau-de-cachimbo	[3]	pelepele apotokon	[1535], [1772]
pau-de-chicle	[114]	pelepele sili	[363], [368], [370], [372]
pau-de-colher	[114], [126]	pelo emulutano	[129]
pau-de-formiga	[427]	pelowi	[107]
pau-de-gafanhoto	[1576]	pelowi'u	[108]
pau-de-índio	[1189]	pelulu	[540], [989]
pau-de-jangada	[1000], [1002]	pëmu epit	[2]
pau-de-leite	[95]	pëmulimë	[553]
pau-de-polvora	[234]	pende paaya	[1149]
pau-de-remo	[955], [1442]	pendyeku	[74], [83], [85], [89]
pau-de-sangue	[951]	pëneju	[1316]
pau-de-sobre	[1142]	pengeku	[83], [85], [89]
pau-doce	[514], [1701]	pengn-makak	[1000], [1001], [1002]
pau-estálador	[1785]	pente-de-macaco	[1000], [1001], [1002]
pau-ferro	[777], [1731]	pëpë	[156]
pau-jacaré	[1529]	pepee anga sawtu	[42], [43], [44], [47], [49]
pau-lacre	[549], [551], [552]		

Appendix 2. — Continuation.

pepee boyti.....	[1605]	pilima'i piyū.....	[1792], [1795]
pepino-doce	[95]	pilima'i wu	[1792], [1795]
pepino-do-mato	[95]	pīlimokju	[1503]
péquéia.....	[248]	pilisi aminiyu.....	[1009], [1010]
pequiá	[248]	pilitalaipyo	[1528]
periquiteira-da-terra-firme	[173]	pilityalapo	[1384]
periquitinha	[234]	piliwa epulu.....	[297], [293], [1296]
perna-de-moça	[1440]	piliwa molototo'u	[137]
peruvia-kamwi.....	[1145]	piliwa mulototou	[137]
pesilan.....	[326]	piliwa pi	[293], [1296]
pésolowa.....	[1118]	pimenta-de-nambú.....	[484], [490]
pesupi.....	[496], [510]	pimenta-longa	[1398]
pet du diable	[516]	pina.....	[149], [150]
pëtëlimë	[1180]	pina'i	[43], [51], [79]
petit génipa	[1472]	pina'i piye.....	[20]
pëtpëjot	[1398]	pina'i piyū.....	[43], [60], [67]
petumo.....	[1002]	pina'i sī.....	[46], [49], [51]
petumo ka'apewat.....	[1000]	pina'i sili.....	[20], [46], [69]
petumo tsing	[1001]	pina'i tay.....	[43], [79]
petun.....	[1002]	pina'i teákā lopiwa	[48]
pëunpë	[1714]	pina'i to wi'i	[46], [51], [69]
pija píloman	[1504]	pina'i to wu	[49], [82]
pija upo.....	[963]	pina'i u	[44]
pikau ki'iy	[484], [1323]	pindaíba-de-folha-pequena	[85], [89]
pikau lemiú'i.....	[484]	pindo uhu	[153]
pikau meyu	[1323]	pindya	[1404]
pikau'i	[1016], [1018], [1019]	pindya udu.....	[548], [550], [551]
piki li.....	[1251]	pino	[149], [153]
piki li sili	[1279], [1282]	pinó.....	[1761]
pikia.....	[248]	pino akusiway	[153]
pikililan	[1310]	pino e'e	[153]
pikimisiki	[926], [927]	pino tamanuwa	[153]
pikin busi suku.....	[1357]	pintadinha.....	[309], [339]
pikin katu.....	[1171], [1182]	pipi	[1379]
pikin loka	[777]	pipya tamili	[1489]
pikin maamadosu	[1457]	piquiá.....	[248]
pikin mapa	[95]	piquiá-marfim	[96]
pikin paanga.....	[768], [772]	piquiárana	[246]
pikin pendyeku	[89]	piquiárana-do-igapó	[247]
pikyalaipyo	[1105]	pirayaura-kinha.....	[41]
pila lea.....	[1246], [1252], [1280], [1282], [1292], [1296], [1307], [1308], [1310]	pirigamepé	[715]
pila lea sili	[1297], [1310]	pirimavan	[1213], [1214]
pila lea u.....	[1307]	pirivri	[356]
pila lea wate'e	[1307]	pisat	[85], [89]
pila siay	[43], [51], [79], [82]	pisulu	[1043]
pila sila.....	[49], [82]	pisusuk	[1472]
pilate'i.....	[100]	pisusukimë	[715]
pilapisi.....	[522]	pitaguará-amarelo	[1506]
pilapuku'i	[576]	pitaica	[955]
pilika	[181], [187], [214], [221]	pitanga	[1233], [1249]
pilikolo'i.....	[1790]	pitanga-da-mata	[1096]
pililipyo.....	[72]	pitanga-miuda	[1283]
pilima'i	[1791]	pitangueira-de-cachorro	[1296]
		pitomba-brava	[1587]

Appendix 2. — Continuation.

pitomba-de-macaco.....	[1576]	puluma.....	[1743], [1751], [1752]
pitomba-do-norte.....	[1581], [1582], [1583], [1589]	pulumá.....	[1752]
pitombeira.....	[1581], [1582], [1583], [1589]	puluwi.....	[1437], [1438]
pitombinha	[249]	pune-etni-kamwi.....	[1410], [1489], [1499]
pitsiwit-mididju.....	[1011]	pune-etni-priye.....	[1418]
piyawapan wewe.....	[1717]	pune-etni-seine.....	[1492], [1494]
podo tiki	[67]	pun-kamwi.....	[1371]
pois sucré	[821], [825], [834], [858], [863]	pup	[156]
pois sucré blanc	[850]	pupunha-de-porco.....	[157]
pokata	[169]	puruhy	[1434]
pokololomilí.....	[745], [775]	purui-grande	[1437], [1438], [1454], [1455]
polata	[1621]	purui-pequeno	[1434]
pole	[934]	purumá	[1758]
polipyo.....	[279], [324], [1771], [1776]	purumaí	[1753]
polipyo tagulewe	[338]	putsulu.....	[1043]
polipyoli.....	[279], [324], [1771], [1776]	puyo.....	[1185], [1196]
polo ali	[458]	pwait	[99]
pololu peta	[856]	pwa-sab	[797]
pombinho	[1141], [1724]	pwa-sikré	[814], [815], [817], [821], [824], [825], [832], [834], [837], [839], [842], [845], [846], [853], [857], [863], [864]
pomiidyilan	[1225]	pwa-sikré-bakòv	[858]
pomme cajou	[8]	pwa-sikré-blan.....	[850]
pommier-cajou.....	[8]	pwa-sikré-gran-bwa	[860]
ponm-sousouri	[1721]	pwa-sikré-koubari.....	[859]
ponó.....	[691]	pwa-sikré-krapo.....	[820]
ponoimë.....	[705]	pwa-sikré-rouj	[844], [856]
popolika	[536]	pwé	[106]
pororoca.....	[777]	pwé-blan	[117]
posenti	[516]		
posindyo	[818], [846]		
postii	[516]		
potípotpiu	[1349]		
poto:no palemulu.....	[1721]		
pourouma	[1749], [1752]		
pracaxi.....	[922]		
pracuba-de-leite.....	[1701]		
préfontenn	[794]		
prin-lans.....	[262]		
prin-savann	[992]		
prin-zikak.....	[262]		
psuk-awak	[1345]		
psum	[537]		
puduku	[1721]		
puduku-wakáu	[1523]		
puduku-wakáu-purubumna.....	[1522]		
puhu	[1043]		
pukut	[701], [720], [722], [728]		
pukutsa'í.....	[1442]		
pukuu	[85], [89], [91], [867]		
pukuu-kamwi.....	[443]		
pukuu-seine.....	[91]		
pukuu-seinó	[91]		
pulelu inga	[975]		
pulo paaya.....	[1184], [1185], [1192], [1193]		
pulkuni	[816], [818], [850]		

Q

quariquara-branca	[1717]
quaruba.....	[1806], [1810]
quaruba-azul	[1791]
quaruba-branca	[1802]
quarubarana	[1788]
quaruba-rosa	[1802]
quaruba-vermelha.....	[1802], [1811]
quebra-machado.....	[547]
quina.....	[1710]
quina-de-Caiena.....	[1710]
quinaquina	[1453]
quinarana	[107], [108], [109], [1357], [1358], [1360], [1361], [1453]
rabo-de-mico.....	[825]
rabo-de-tucano	[1802]
radjé-chank	[522]
rainha.....	[1492]
rapari.....	[877]
rapé-de-índio.....	[1189]
remelo-de-cachorro	[1099]
rézen-bòdlanmé	[1403]
rim-de-paca	[770], [772]

Appendix 2. — Continuation.

ripeiro	[701], [702], [720]	sapupira-amarela	[812]
ripeiro-branco	[700]	sapupira-da-várzea	[787], [809]
roucou grand bois.....	[451], [458], [460], [463],[468], [471], [472]	sardinheiro	[496]
roukou-gran-bwa.....	[451], [454], [458], [460], [463],[468], [471], [472]	satiné rubané	[1152]
roukou-sovaj.....	[451], [454], [458], [460], [463],[468], [471], [472]	satiné-ribanné.....	[1152]
S		saulanani	[499]
sa'i melu ka'a	[22], [23]	sauwalai.....	[141]
sa'i melu ka'a wu.....	[41]	sawa	[246]
sabana foo	[933]	sawali	[246]
sabana gwegwe	[928]	sawu-anen	[1743]
sabana kwali	[1791]	sawu-anen-kamwi.....	[438]
sabana luabi.....	[725]	sèd.....	[574], [598], [647]
sabana matakí.....	[380]	sèd-akajou	[1114]
sabana toso paasaa	[1201]	sèd-bagas	[181], [636]
sablier.....	[516]	sèd-blanc	[643]
saboeiro	[733]	sèd-darjan.....	[637]
saboeiro-da-várzea	[805]	sede	[1114]
sacaca	[504]	sedee	[1114]
sadyabe.....	[1212], [1295]	sèd-gri	[635], [636], [643]
saint-martin blanc grandes feuilles.....	[892]	sèd-gro-lapo	[652]
saint-martin gris	[808]	sèd-jonn	[603], [652], [666], [671]
saint-martin jaune	[809], [811]	sèd-kalalou	[643]
saint-martin rouge.....	[745], [747]	sèd-kannel	[575], [603]
saipyalala	[551], [552]	sèd-marikaj.....	[576], [591]
sakaugi	[930]	sèd-nwé	[624], [652]
sakaugi-kamwi.....	[929]	sèd-rémi	[959], [961]
sakauri.....	[930]	sedri-eminyo	[583]
sakauri-kamwi	[929]	sedri-kamwi.....	[608]
sakeg	[809], [913]	sedri-kamwi-seine.....	[581]
sakeg-kamwi.....	[754], [794], [910], [912], [916]	sedri-priyo	[608]
sakeg-seinó	[882]	sedri-priyu	[649]
sakusaku.....	[1328], [1329]	sedri-puvemna.....	[669]
sala'i	[1520]	sedri-puvemna-priyo	[668]
sala'i sili	[1520]	sedri-seine	[176]
sala'i u	[1522]	sedri-wahuyo	[1114]
salala	[829]	sèd-rouj	[1114]
salasala.....	[1739]	sèd-sam	[416]
sali.....	[184], [216]	seityape	[1212], [1279], [1295], [1315]
samaati	[1133]	seivan	[1473]
samaliyapo	[1114]	sekeleu.....	[1425]
samaúma	[1008]	sen-marten-blan	[892]
sangue-de-galo.....	[929], [930]	sen-marten-gri.....	[808]
santi kookoo.....	[334], [340]	sen-marten-jonn.....	[809], [811]
santo	[297]	sen-marten-rouj.....	[745], [747]
sanvonié	[1567], [1576]	sen-martin-blan	[741], [742]
sapucaia.....	[729]	sepeipyo	[224]
sapululi.....	[155]	sepundi	[863]
sapupira.....	[788]	sepupi.....	[224], [1581]
		seringaí.....	[517], [519]
		seringueira-itaúba.....	[515]
		seweli	[247]
		siduwapali	[576], [578], [582], [585]
		sieuju.....	[1550]
		sihkëimë.....	[1052]

Appendix 2. — Continuation.

sihkëimëimë	[1568]	sisi	[816], [818], [850]
sihkëpani	[955]	sisi pay	[843], [860]
sii payopo	[1099]	sisi sī	[816]
siki	[1059]	sitaipyo	[603]
siki lā	[1055], [1075], [1084], [1087]	sitronél	[102]
siki sī	[1063], [1070]	sitronél-blan	[99], [101]
siklet	[114]	sitronél-rouj	[1442]
siktukahmit	[427]	sitru-kamwi	[463], [1371]
silimaipo	[982]	siwaluwa	[1119]
silityo	[872]	siwiman	[83]
siliwi	[711], [722]	siwinyani	[548], [550]
simajali	[181], [211]	sobreiro	[740]
simali	[1114]	soim	[1211]
simaliapo	[1114]	sokolo	[1712]
šimaluba	[1714]	sokosoko	[117], [126]
simalupa	[1714]	sokosoko'i	[129]
simarouba	[1714]	sokowe	[31]
simartē	[1127]	soloma	[338], [339]
simartē-puvemna	[857]	solosolo bakoloidyi	[345]
simatē	[1127]	sooke	[1671]
simig	[806]	sopu uwii	[949]
simii	[806]	sopu wiwii	[949]
simili	[806]	sorva	[106]
šimili	[806]	sorvão	[106]
simili tabile	[806]	sorvinha	[114]
simili tamune	[806]	sucupira-açu	[788]
simili tubulu	[806]	sucupira-amarela	[794]
šimililan	[919], [920], [926], [927]	sucupira-preta	[788]
simira	[1500]	sucuúba	[112]
simyali epi	[442], [444], [447]	sucuúbarana	[110]
sineki udu	[978], [980]	sulu iwi	[711], [722]
singabusu	[1565], [1566], [1567]	suluku'a	[1728]
singapeetu	[778]	sumaliba	[1714]
šipa yepo	[1092]	sumaúma	[1008]
sipi	[191], [196], [210], [218]	sumaúma-brava	[1011]
sipi lā	[205], [216]	sumaypa	[1714]
sipi laāngä	[205], [216]	supiárana	[496]
sipi sili	[200], [205], [216]	supun udu	[1039]
sipi towi'i	[205], [216]	supupi	[1581]
sipiki udu	[1381]	suwilani	[548], [549], [554]
sipilulan	[580], [647]	suwilani pilá	[551]
sipilupipyo	[576]	suwilani wu	[554], [1209]
šipyä	[188], [189], [196]	suwimba	[548], [549]
sipyalala	[1078], [1089]	suwimba-ahavukune	[554], [1727]
sipyö	[188], [189]	suwimba-kamwi	[555]
šipyö	[188], [189], [195], [196]	suwimba-purubumna	[552]
sirasira	[188], [210], [211], [213]	suwimpa	[548], [549]
sirasira-seine	[191]	suwimpa-ahauwukune	[554]
sirasra	[188], [210], [211], [213]	suwimpa-kamwi	[555]
sirisiri	[173]	suwimpa-purubumna	[552]
siri-siri	[173]	suwinyani	[550]
siriúba	[1]	suwinyani enekan	[548]
siriz-gran-bwa	[1249]	switi amini	[1606]
siriz-ronde	[1249]		

Appendix 2. — Continuation.

T

taapu tiki.....	[1141]	
tabuku.....	[1637]	
tabululi.....	[155]	
tabulupo.....	[1470]	
tacacazeiro.....	[1034], [1035], [1036]	
tachi.....	[959], [961], [963], [964], [965]	
tadju.....	[162]	
tagulewe.....	[26]	
tagulewe itudyano.....	[429]	
tahuma.....	[558], [1474], [1527], [1547], [1549]	
tahuma seine.....	[1551]	
tahuma-purubumna.....	[1535]	
tahuma-seine.....	[1535]	
taihem.....	[624]	
taitetu kasi ni.....	[18]	
taitetu kásí ni.....	[18]	
taíwí lenipi'á.....	[1399]	
taíwí lenipiá.....	[1057]	
taíwí lenipiá lá.....	[1087]	
taka.....	[1683]	
takalawelu.....	[1053], [1062], [1082]	
takalawelu ka'a lulu.....	[1090]	
takalawelu ki.....	[1078], [1089]	
takalawelu lá.....	[1050], [1070], [1078], [1080], [1086], [1090]	
takalawelu lá sili.....	[1086], [1090]	
takalawelu sí.....	[1090]	
takalawelu sili.....	[1080]	
takalawelu sówí.....	[1057]	
takamala.....	[1605]	
takamalaimë.....	[1318]	
takamalakma.....	[1316]	
takana.....	[642]	
takinan.....	[1148]	
takini.....	[1148]	
takulawa'i.....	[741]	
takulewe.....	[26]	
takulu welu.....	[1082]	
takulu wila.....	[777], [879]	
takweni.....	[1148]	
talaiwi.....	[720]	
talaiwi sí.....	[722], [723]	
talakua'i.....	[1102], [1105]	
talakua'i to wu.....	[1105]	
talakwa'i.....	[1105]	
talala.....	[442], [444], [447], [806], [944]	
tale'i li.....	[1529]	
talepêpê.....	[1494]	
talidi.....	[964]	
talili.....	[1395]	
talipi.....	[1287], [1440]	
talukuwa epi.....	[555], [1105]	
taluwaho	[1095], [1099], [1102]	
tamacoaré.....	[223], [224], [226], [227]	
tamalen	[733]	
tamaniwa.....	[1627], [1702], [1703]	
tamanokwale	[984]	
tamanqueira	[1520], [1522], [1523]	
tamanqueira-de-leite.....	[120]	
tamanuwa.....	[1616]	
tamaquaré	[223], [224], [226], [227]	
tamujom	[695]	
tamukwálê áká.....	[227], [1245], [1250]	
tamunen alaulama	[964]	
tamunen apukuitya	[101]	
tamunen ilakopi	[1791]	
tamunen isyano manduliyi	[51]	
tamunen kuwateli	[707]	
tamunen pilika	[206], [211]	
tamunen salasala	[1740]	
tamunen seityape	[1315]	
tamunen suwinyani	[550]	
tamunen waiki.....	[863]	
tamunen wonu	[584]	
tângala yiki	[1492]	
tângala yiki sili	[1492]	
tangaraca	[1070]	
tanimbuca	[401], [404], [405], [407], [408], [411]	
tanimbuca-do-igapó	[412]	
tanimbuca-rana	[412]	
tanimi	[863]	
tanmaren-gran-bwa	[805]	
tapaiúna	[778]	
tapaka	[797], [798], [800], [801]	
tapaka'i	[742], [797], [907]	
tapakadja'i	[715]	
tapalaimë	[3]	
tapatangman	[1184], [1193]	
tápë lemiú'i	[19], [1525], [1595]	
tapele dju'a	[114]	
tapeleyiwa	[114]	
tapeleyuwa	[114]	
tapeliwa	[13]	
taperebá	[13]	
taperibá	[13]	
tapi'i ka'a lilu	[1494]	
tapi'i ka'alulu	[1452], [1494]	
tapi'i ka'alulu sili	[1492]	
tapi'i lo wapa'a	[1560]	
tapi'i wapa'a	[1459]	
tapi'i wapa'a sili	[1461]	
tapi'ilo wapa'a	[463]	
tapi'ilupabi hun	[428]	
tapi'it lupabi	[1034]	
tapiá	[496], [497]	
tapichó	[501]	

Appendix 2. — Continuation.

tapilen katulimya.....	[1389]	tawali	[692]
tapilen kuwali.....	[1800]	tawali pilá.....	[692]
tapilen kuwatali.....	[720]	tawali sī	[692]
tapilen pipyo	[1385]	tawisowiso	[458]
tapilen salasala.....	[1737], [1739]	tawu.....	[155]
tapilen seityape.....	[1279]	taxi.....	[961], [963]
tapili.....	[1440]	taxi-branco	[964]
tapilili.....	[15], [16]	taxi-branco-da-terra-firme	[960]
tapilupami.....	[1035], [1036]	taxi-da-terra-firme	[964]
tapiseipyo	[78]	taxi-pitomba-preta.....	[965]
tapooka mapa.....	[1628]	taxi-preto.....	[958], [963]
tapouliman.....	[408], [414]	taxi-preto-da-várzea	[1406]
tapowonulen	[356], [358]	taxirana	[964]
tapuken nopitya	[377]	taxirana-branca.....	[938]
tapuliyuipo.....	[1283]	taxiseiro.....	[1406]
tapululi.....	[155]	tayau'i.....	[969], [979]
tapulupo.....	[1470]	tayi.....	[162]
taquari.....	[517], [519]	tayi lá	[566], [567]
tararanga	[1752], [1758]	tchenbyendan.....	[95]
tararanga-vermelha.....	[1754]	tchò-bèf.....	[28], [1010]
tarii	[1395]	tchò-dérò.....	[788]
tariri	[1395], [1396], [1397]	tembetaru.....	[1523]
tarit	[1395]	tento	[890], [897], [898]
tartaruginha	[257], [496]	tento-grande.....	[896]
tarumá.....	[567]	tento-miúdo	[748]
tasi	[505], [963], [1406]	tento-preto	[893]
tasi epi.....	[962]	tento-verdadeiro	[748]
tasi piyū.....	[963]	ten-wašiunó.....	[114]
tasi sī	[963]	tenway.....	[516]
tasiasiaiy	[1702]	tépepulu.....	[1233], [1239], [1240]
tata pilili.....	[15], [16]	tépepuluimë	[1237], [1240], [1604]
tata pota	[358]	teposi kili.....	[889], [897], [898], [956]
tatajuba	[246], [1146]	tepotsi átā.....	[544]
tatapililit.....	[15]	tépu.....	[1583], [1592]
tatapiririca.....	[15], [16]	tépuimë	[1589]
tatsi	[963], [964]	tépuklem	[1649]
tatsi inán	[961]	tété-kwata.....	[1599]
tatu	[1582], [1583], [1587]	tetey bii udu	[800]
tatu apite	[374], [377]	tetey weko	[825]
tatu kásı.....	[1419], [1422], [1423], [1770]	tewitulū u.....	[1487]
tatu mila.....	[249], [529]	tewitulū wate'e.....	[1487]
tatu tunu.....	[952]	tewitutū wate'e	[1487]
tatu udu	[1582], [1590]	ti-bwa-lélé	[1772], [1776], [1782], [1783], [1784]
tatu'i.....	[938]	ti-golet	[293]
tatu'i u.....	[875]	ti-golèt-rouj	[293]
tatulimá.....	[400], [415], [751]	tihsigli epít	[519]
tatulimá sili	[401], [408], [414]	ti-jinipa	[1472]
tatumila sili	[557]	tikalaye ulemali	[691]
tauarí.....	[693], [695]	tikalayen tonoloipyo	[1571]
tauarí-amarelo	[692]	tikilin wewe	[1385]
tauarí-folha-grande	[691]	ti-koubari	[777]
tauarí-folha-miuda	[692]	timaa-avain.....	[1713], [1714]
tauni.....	[1148]	timaa-avain-purubumna	[1707]
tavu.....	[155]	ti-marikasmatgene	[380]

Appendix 2. — Continuation.

timbaúba.....	[794], [939]	toulouri.....	[151]
timborana-foliolo-fina	[927]	tourém	[1753]
timborana-foliolo-graúdo	[926]	tourouri.....	[151]
timuvukti	[1105]	trapiá.....	[239]
timuvukti-duwē.....	[1102]	tsalikwanā'i.....	[370]
timuvuyen.....	[757]	tsibila lipi	[322]
timuvuyen-kamwi	[943]	tsibila lipikit	[335]
tingí moni	[187], [188], [191], [195], [196]	tsibit.....	[339]
tininipi	[10]	tsimini.....	[806]
tinteiro	[497]	tu pikin weko	[1368]
tipulu alaidya.....	[47]	tubá-pairi	[1152]
tipulu alaulama.....	[961]	tuchauá	[1529]
tipulu aloome	[523]	tucumá-branco	[144]
tipulu apukuitya.....	[99], [102]	tucumá-do-Pará.....	[146]
tipulu apukuityalan	[943], [955]	tucunaré-mereçá.....	[1099]
tipulu kiyele'u.....	[778]	tudjulata'i	[18]
tipulu sepeipyo	[938]	tuká nákū	[490], [1349]
titikö	[729]	tuká nákū sili	[484], [490]
titikö uhu	[684]	tuká nili.....	[1354]
titim batibataa	[793]	tukánákū sili	[983]
tiviru	[1111]	tukuke	[729]
tiviru-seinô	[1112]	tukuman udu	[245]
tivu-arigni	[1402]	tukusi pailali.....	[1152]
ti-washiunu	[378], [381]	tukuwi	[1737]
tiyapotano ilakopi.....	[1797]	tukuwi-awaig.....	[1740]
tiyapotano mutusi	[928]	tukuyuy	[1434], [1435]
tiyawasisyen.....	[1576]	tukuyuy-kamwi	[994], [1439], [1481], [1503], [1664], [1668]
tobitetu.....	[137]	tukwanru.....	[1185], [1191], [1196]
tohmopala	[1565], [1566]	tukwanru-kamwi	[1192]
toko	[693], [717], [722]	tula.....	[729]
tokolo abolimeli	[1149]	tulaimë	[729]
tokolo enulu	[536], [1389]	tuleke	[1737], [1739], [1740]
tokolo pailali	[1149]	tuli	[818], [820], [825], [1582], [1584], [1589], [1592]
tokolo-pipinatsi	[458]	tuli átā	[1581], [1583], [1587], [1592]
tokowe	[637]	tuli átā u	[1574], [1584], [1587]
tokuhsa	[463]	tuli átā yowa	[1587]
tokulojem	[859]	tuli sī	[1594]
tomoipo	[1552]	tuli ii	[191]
tona.....	[1783]	tulili	[832], [836], [960], [964]
tonawewe	[1783]	tulili wala	[959], [961]
tonka.....	[789], [790]	tulipalapo	[281], [282]
tonmat-sousouri	[1721]	tulisi	[1585], [1589], [1594]
tonolo ipyo.....	[1196], [1566], [1567], [1569], [1581]	tulisimë	[834]
tonolo polipyo.....	[1196], [1566], [1567], [1569]	tululi	[151]
tononipö	[1566], [1567]	tulumele	[1341]
topi	[1095], [1099]	tulumele u	[1342]
topuwunu	[962]	tulusili	[1585], [1589], [1594]
torém	[1738]	tumau-atamwa	[539], [540], [541]
toso paasaa	[1203], [1204]	tunu'i	[757]
tossopassa marécage.....	[1201]	tunu'i pilá.....	[751]
tossopassa montagne	[1203]	turamira	[537]
toto	[652], [666]	turury.....	[1153]
totoka.....	[684]	tutu	[1291]
touka.....	[684]		

Appendix 2. — Continuation.

tutuka	[684]	uluwu kásĩ	[1722], [1723]		
tutukẽ	[684]	uluyatu	[1762]		
tuu-ariut	[1576], [1582], [1709]	uman feyfi finga	[827]		
tuu-aška	[1585], [1592]	uman papay	[1737]		
tuu-kamwi	[1585], [1593], [1709]	uman tapupa	[1470], [1472]		
tuu-kamwi-seine	[1576]	umari-bravo	[1143]		
tuuvan	[151]	umeg	[537]		
tuwonule	[1667]	umi'í	[1042], [1044]		
tyabisi	[788]	umiri	[537]		
tyaka	[1683]	umpatapu	[1185], [1192]		
tyentyen udu	[1142]	urtiga-brava	[1761]		
tyoto amote	[797]	urubuzeiro	[978]		
U					
uaçima-do-campo	[1016]	urucurana	[451], [454], [458], [460], [463], [468], [471], [472], [499], [1389]		
ujara-branco	[1597]	urucurana-branca	[530]		
uarutama	[249]	uu	[1671]		
ubaia	[1249]	uu-ahavukune	[1666]		
ubuçú	[151]	uu-aška	[1665]		
uchirana	[273], [541], [544], [547], [550]	uu-kamwi	[1605], [1626], [1628], [1641], [1664], [1667], [1675], [1680], [1682], [1686]		
ucuquirana	[1606]	uu-kamwi-duwẽ	[1649], [1664], [1670]		
ucuúba	[1210]	uu-kamwi-purubumna	[1626]		
ucuúba-amarela	[1205]	uu-kamwi-puvemna	[1700]		
ucuúba-branca	[1210]	uu-kamwi-seine	[1666], [1680]		
ucuúba-chico-de-assis	[1205]	uu-kamwi-wahuyo	[1599], [1652], [1686], [1706]		
ucuúba-da-terra-firme	[1207], [1208]	uu-purubumna	[1670]		
ucuúba-do-igapó	[1210]	uva-de-macaco	[1754]		
ucuúba-preta	[1207]	uvayan-akig	[631]		
ucuúbarana	[1201], [1203]	uwa kaya	[1461]		
ucuúba-vermelha	[1203], [1209]	uwa ki'iy	[1492]		
ukalat	[691]	uwa pitág ndi	[1249]		
uki'í	[46], [51]	uwahe	[636]		
uku tiki	[63]	uwahe hun	[627]		
ukum	[106]	uwahe pitág	[574], [608]		
ukum-kamwi	[117], [1706]	uwahe tsing	[645], [652]		
ulana beta	[944], [945], [947], [952]	uwaitya-avain	[110]		
ulana elepali	[702], [709]	uwakaa	[1451]		
ulemali	[692], [695]	uwato eptyi	[250]		
ulimẽ	[247]	V			
ulu maululu	[1010]	vaquinha	[522]		
uluimẽ	[496]	vassourinha	[1219]		
uluku lá	[1037]	vénéré	[1718], [1720]		
uluku panali	[458], [468]	ventosa	[536]		
uluku panali sili	[454], [460]	vermelhão	[286]		
uluku tawa	[171]	viniré	[1718], [1720]		
ulukupalani	[458], [468]	violeta	[918], [919], [920]		
ulukupalani sili	[454], [460]	virarú	[1433]		
ulukupi	[1151], [1153]	viratay	[24]		
ulukupi sili	[1153]	virola	[1209]		
ululu'u	[912], [913]	visgueiro	[913]		
ulumapi	[1749], [1752]				
ulusulu	[30], [35]				

Appendix 2. — Continuation.

vouapa.....	[879]	waiyen.....	[1037]
vouapa-tabaca.....	[797]	waka'i.....	[1360]
		waka'i sili.....	[1357]
		waka'i u.....	[1360]
		wakaba.....	[415]
		wakaba-kaubimna.....	[100]
		wakali'i.....	[1373]
		wakap.....	[973]
		wakapou.....	[973]
		wakap-priye.....	[973]
		wakap-seine.....	[973]
		wakapu.....	[973]
		wakapu-giten.....	[931]
		wakáu.....	[1429], [1451]
		wakáu-dariu.....	[173]
		wakavu-kamwi.....	[2], [19], [496], [497], [498], [499], [501], [1437]
		wake.....	[248]
		wakukwa-adava.....	[715], [729]
		wakukwa-adava-kamwi.....	[716]
		wakukwa-adudu.....	[1599]
		wakukwa-ára.....	[1424]
		wakukwa-rimwa.....	[691], [693]
		wakukwa-tiranó.....	[369], [374], [376], [377]
		wakukwa-tiranó-kamwi.....	[379]
		wakukwa-tiranó-purubumna.....	[223], [226]
		wala ekone.....	[498]
		wala potili.....	[834], [866]
		walaku ki'iy.....	[1478]
		walakuseli.....	[191], [196]
		walalu'i.....	[1441], [1442]
		walapa kuwalili.....	[1788], [1810]
		walapulu.....	[1038]
		walapulu lá.....	[3]
		walatiwá.....	[378]
		walatiwá sili.....	[1379]
		walekone.....	[926], [927]
		waletsi.....	[1210]
		waletsi lawöt.....	[1209]
		waletsi pitág.....	[1206]
		waletsi tsing.....	[1207]
		wali lapi'a.....	[257]
		wali wowo.....	[536]
		walili pipyo.....	[636]
		walili tuli.....	[1130]
		walima.....	[535]
		walime.....	[1209]
		walime i.....	[701]
		walimei.....	[1202]
		wališi.....	[1210]
		walolo.....	[1207], [1209]
		walulan.....	[1143]
		walusi.....	[1210]
		waluši.....	[1210]

Appendix 2. — Continuation.

waluši ityudano	[1207], [1209]	wap-seinō	[798], [800]
walušilan	[1209]	wapu	[149], [150]
walusilan tamunen	[1207]	wapuimë	[150]
waluti	[1210]	wap-waikmwiunō	[798]
waluwa'i	[176], [202], [208], [209], [210], [216]	wara	[146]
waluwai	[184], [200], [202], [211], [215], [219]	warahu	[262]
waluwai pilá	[184], [202], [211], [215], [219]	waravru	[1038]
waluwai sī	[184], [187], [200]	waravru-ahavukune	[1031], [1039]
waluwai sili	[204]	waravru-kamwi	[2], [496], [497], [498], [499], [501], [1015], [1022], [1515]
waluwalu	[692]	waravru-kamwi-seine	[168]
wamisalu	[1628], [1686], [1688]	waravru-puvemna	[1730]
wamutu'í	[1068], [1070], [1079], [1080]	waravru-wašiunu	[1015]
wan ede	[1707]	warerey	[1442]
wana	[673]	waru	[970]
wana kwali	[1799], [1800], [1810]	waru-kamwi	[972]
wanaku-danō	[751], [952]	was	[149]
wanaku-divyo	[952]	wasakau	[436]
wanaku-duwē	[748]	wasaku	[516]
wanaku-kamwi	[890]	wasaku sili	[1711]
wanaku-marikasmatgene	[892]	wasay	[149]
wanaku-priyo	[897]	wasei	[149]
wanaku-umutinō	[748]	wasepuku	[1666], [1667], [1688]
wanani	[380]	wasey	[149]
wanāni	[380]	wasey iipi manu	[149]
wandekole	[514]	wasey iipip'sí	[150]
wan-édé	[1707]	wasey sī	[150]
wanetun	[745], [746], [747], [969]	wasipa	[877]
wap	[797]	wasiwasi uwii	[427]
wapa	[797], [800]	wassay	[149]
wapa courbaril	[798]	wata babun udu	[1210]
wapa marécage	[879]	wataa bii udu	[800]
wapa rivière	[800]	wataa kanu	[416]
wapa-charles	[922]	wataa matakí	[380]
wapa-dilos	[879]	wataki	[107], [108], [109]
wapa-gra	[797]	watala	[691], [806]
wapaaimë	[749], [907], [919]	watampana	[879], [880], [881]
wapa-koubari	[798]	watapa	[797], [800]
wapa-larivye	[800], [879]	watapalili	[15], [16]
wapa-marikaj	[879]	watapana	[879]
wapa-mèg	[907]	watau pokan	[603]
wapa-rouj	[800]	wateipa upadyali	[383], [389], [393]
wapa-sek	[879]	watili	[722]
wap-duwō	[797]	watili pitág	[726]
wapiju	[1120]	watili tsing	[685], [701], [723], [724]
wapiku	[751]	watoipyo	[1733]
wapitá	[1662]	watolo	[1201], [1203]
wapiu	[1120], [1124]	watoyati	[1733]
wap-kamwi	[770]	watsey	[149]
wap-kamwi-priyu	[942]	watuliya	[1628]
wap-kamwi-puvemna	[777]	watumo	[324]
wapolimë	[1598]	wau	[1666], [1686]
wapotoimë	[898], [952]	wau átā	[1669]
wap-purubumna	[879]		
wap-puvemna	[770]		

Appendix 2. — Continuation.

wawayu.....	[1666], [1667]	wila náná.....	[1655]
wayaka	[603], [627], [636]	wila náná kweli.....	[379]
wayamu sasamili.....	[124], [1323]	wila náná sili.....	[1655]
wayamu tapulu.....	[858], [864]	wila paye	[812], [972]
wayamu wolekotopo.....	[1455]	wila pele	[1599]
waye.....	[635], [650], [1037]	wila pikikit	[544]
wayilu	[61], [67], [69]	wila pilatá.....	[1374]
wayu	[1437]	wila pile lu.....	[223], [224], [227], [1368]
wayun	[1437]	wila pipé mu.....	[984]
weko	[814], [821], [834], [850], [857], [860]	wila pisí'u	[1139], [1571]
weli	[537]	wila pitág hun.....	[1680]
wen	[637]	wila pitág piñág	[1666]
wen-etni	[637]	wila pitág tsing	[1664]
wen-kamwi.....	[591]	wila poloelaa.....	[1769]
wesekapo eptyi	[380]	wila pulu'a	[1599]
wesepuku	[1628]	wila pulua.....	[281]
weti apisi	[624], [637]	wila suku	[1715], [1717]
weti baka.....	[333]	wila tai.....	[1374]
weti bee	[1082]	wila taiwa	[1366]
weti dondomisinga.....	[912]	wila takulu	[777]
weti kiikii	[1029]	wila takulu pilá.....	[918]
weti kookoo	[334], [1424], [1425]	wila tátá.....	[919]
weti mapa.....	[117]	wila tawa	[988], [1642]
weti musupu	[1070]	wila tí	[1386], [1387]
weti pendyeku	[91]	wila tipita toto	[1250]
weti pengeku	[91]	wila to pokو	[1426]
weti pindya udu	[550]	wila to tawa	[122], [1468]
weti sali	[211]	wila towakape sī.....	[436]
weti sikin apisi.....	[624], [637]	wila tuwi	[530]
weti tiki	[1349]	wila ū.....	[1218], [1233]
weti tobitutu	[137]	wila umí'i	[303], [304], [306], [311]
wewe lémopka.....	[1012]	wila umí'i wu.....	[308], [1307]
wewe pipyo	[919], [920]	wila wasey	[1294]
weweimë	[920]	wila yisí	[264]
weyu'i	[281], [282]	wila yowa	[1252]
widajimë	[144]	wila'a	[153]
wiipë	[377], [379]	wilakayulu	[472]
wila atsi	[1523]	wilapilátá.....	[1374]
wila è'è	[1702]	wilau piyú'á	[1754], [1757]
wila iki'ý	[1141]	wilimë	[567]
wila imí'i	[303], [304], [306], [311]	win udu	[1389]
wila imí'i wu.....	[308], [1307]	winámë'i	[943], [946]
wila ka'alulu wipë'i	[1415]	winámë'i piyú	[943]
wila kalayulu	[463], [472]	winámë'i sī	[946]
wila kayulu	[463]	wipitano ityumban wewe.....	[109]
wila kea	[1608], [1646]	wipitano kuwateli	[701]
wila ki	[875]	wipitano pilika.....	[214]
wila kitá.....	[1094], [1095], [1102]	wiratay	[51]
wila kitá u.....	[1104]	wítik.....	[95], [117]
wilá lë.....	[926], [927]	wiwis-ahavukunó	[429]
wilá lë pilá	[926], [927]	wiwis-kamwi	[1743]
wilá lë sī.....	[926]	wiwis-kasiuvan	[427]
wila lo	[122]	wiwis-priyo.....	[429]
wila munuwi	[1371]	wiwis-purubumna	[429]

Appendix 2. — Continuation.

wiwit hun.....	[67], [80], [81]	yanipa i	[1472]
wiyekane	[1381]	yanipa lali.....	[1434]
wiyu	[1373]	yanipa lali sili.....	[1463]
woki	[153]	yanipa lali u.....	[1503]
wokili apukuitya.....	[96]	yanipa u	[1454], [1455], [1457]
wokili besuli	[355]	yanita	[988], [1154]
wokili kisipyulu ulemali.....	[695]	yankomini.....	[1009], [1010], [1011]
wokili kupesini	[336], [338], [339]	yanu'í	[293]
woko isyale.....	[788]	yanu'í pilá.....	[292], [293], [295], [1188]
woko mapili	[598]	yanu'í u	[1198], [1199]
woko molokotoli	[1341], [1342]	yanu'í wate'e.....	[1198], [1199]
woko popi	[1099], [1384], [1637]	yanu'í yowa	[292], [295], [1188]
woko wokulu	[1628]	yapakaní'i	[959], [961]
wokunse	[1357], [1360]	yape'a pitá	[1535], [1539]
wololo	[1207]	yape'a pitá sili[557], [1527], [1528], [1539], [1540], [1547]	
wololo'i	[1207]	yape'a pitá u	[1535]
wololo'u	[1208]	yapopale	[35], [303]
wonu.....	[673]	yapopale membo	[336]
wopoli	[1631]	yapopali	[541]
wosiwosi.....	[1802], [1808]	yapu tulu'i	[1020], [1028], [1029]
wosiwosi tamunen.....	[1810]	yapu tulu'i sī.....	[1020], [1029]
woto kasimyate.....	[29]	yapu'í	[740]
woto kolalu	[167]	yapu'í pilá.....	[740]
woto kuwalili.....	[1809]	yapui	[649]
wulimë	[247]	yapukay	[729]
		yápkay	[729]
		yapukay sili	[687]
		yapukuliwa sili.....	[1515]
		yapukuliwa u.....	[874]
yá	[684]	yariwapna.....	[1718], [1720]
yá laánga	[1645], [1670], [1688]	yariwapna-kamwi	[1715]
yahiwemna	[1718], [1720]	yásí lea'iy	[497], [999]
yahiwemna-kamwi	[1715]	yasimanbo	[164]
yakale'i	[222]	yata'i.....	[157]
yakami ka'a.....	[570]	yatoa'i	[1121]
yakami'i.....	[570]	yatoa'i paye.....	[1124], [1125]
yalakalu pupo	[1454]	yatoa'i sī.....	[1118], [1120], [1124], [1137], [1138], [1567]
yalakalu tumali epi	[729]	yatoa'i sili	[1122], [1124], [1125]
yalakasi.....	[245]	yatoa'i u	[1138], [1139]
yalakasila	[522]	yauknabui.....	[1339], [1346], [1349], [1628], [1629], [1769]
yalakasilo.....	[522]	yauknabui-puvemna	[1358]
yalakopi.....	[1718]	yauknabwi-kiabumna	[490]
yalami	[1233]	yauknabwi-wahuye	[1626]
yalamilulan.....	[120]	yawa luway	[1547]
yalayala.....	[1737], [1738], [1751], [1752], [1756]	yawa poi li	[1287], [1429]
yalayala tupulu	[1740]	yawa poi li sī.....	[1248]
yali taku'a sili.....	[1399]	yawa popita	[41]
yalipi	[1080]	yawalelan	[501]
yango	[972]	yawalū lemo	[238]
yani	[1111]	yawasi pita	[1188], [1724]
yani pilá	[1111], [1112]	yawatai	[427]
yani sī.....	[1111]	yawayi lū.....	[1215], [1250], [1257], [1652]
yanipa	[1470]	yáwī inga	[846]
yanipa e'e	[1470]	yáwī kala	[53]

Appendix 2. — Continuation.

yáwí ki'íy	[1249]	yolokan topulukali.....	[649]
yáwí yiki	[53]	yolulu.....	[913]
yáwí'i	[85], [89], [94]	yongo.....	[972]
yáwí'i pilá	[85], [92]	yooka wiwii	[1366]
yáwí'i sí	[83], [91]	yū ásīsī	[1721]
yáwí'i sili	[89]	yu'i li	[1784]
yawu	[1373]	yu-kig.....	[120]
yawu-seine.....	[1550]	yuku'ipyo	[1198], [1199]
yaya.....	[214]	yukutuna.....	[443], [1119]
yaya'i	[181], [214]	yukuyapoi	[517]
yaya'i sí.....	[181], [214]	yululu.....	[913]
yayamadou marécage.....	[1210]	yuumwi.....	[922], [936]
yayamadou montagne.....	[1206], [1207]	yuumwi-seine	[793]
yemelu	[1124], [1125]	yuumwi-wašiunu	[733]
yemi'i	[691], [695]	yuwa	[106]
yenge pao	[1199]	yuwa piso	[1192]
yikisi'i	[1576]	zamād.....	[403]
yikisi'i lo	[1576]	zaman.....	[403]
yita'i	[806]	zanmann	[403]
yit-ára.....	[783], [1366]	zanmann-dan-bwa	[415]
yit-gataibi	[1032], [1035]	zanmann-sovaj.....	[401], [405], [408], [414]
yit-itaibi	[1002], [1032], [1035]	zépini-gran-fey	[1520], [1522]
yit-itaibi-ahawukune	[1031]	zépini-tífey	[1523]
yit-itaibi-priye	[1000]	zimortel.....	[802]
yit-itaibi-priyo	[1001]	zoliv	[1597], [1598], [1626], [1627], [1633], [1637], [1646], [1648], [1652], [1662], [1668], [1682], [1686]
yit-itaibi-purubumna.....	[1035]	zoliv-maypouri	[1631]
yit-itaibi-puvemna.....	[1000]	zongnon-danbwa	[363], [365], [371], [372]
yit-itaibi-seine	[1000]	zouti-montangn.....	[1761]
yiwa piso	[1192]	zoyó-kamwi.....	[365]
yoloka pomiydi	[1091]		
yolokan pomiydi	[43], [47], [608]		

APPENDIX 3. — Index of scientific names/Index des noms scientifiques.

A			
<i>Abarema</i>	503	<i>Achras guapeba</i>	695
<i>Abarema barbouriana</i>	503	<i>Achras mayana</i>	687
<i>Abarema brachystachya</i>	517	<i>Achras nitida</i>	690
<i>Abarema curvicarpa</i>	503	<i>Achrouteria durifructa</i>	686
<i>Abarema gallorum</i>	503	<i>Achrouteria pomifera</i>	687
<i>Abarema jupunba</i> var. <i>jupunba</i>	503	<i>Acia amara</i>	417
<i>Abarema jupunba</i> var. <i>trapezifolia</i>	504	<i>Acia dulcis</i>	416
<i>Abarema laeta</i>	504	<i>Acinodendron affine</i>	561
<i>Abarema mataybifolia</i>	504	<i>Acinodendron amplexans</i>	569
<i>Abarema racemosa</i>	546	<i>Acinodendron amplum</i>	562
<i>Abarema</i> sp. A	504	<i>Acinodendron argyrophyllum</i>	563
<i>Abarema trapezifolia</i>	504	<i>Acinodendron axilliflorum</i>	561
<i>Abbevillea martiana</i>	597	<i>Acinodendron beurlingii</i>	562
<i>Aberemoa guianensis</i>	373	<i>Acinodendron boissieranum</i>	569
<i>Aberemoa longifolia</i>	372	<i>Acinodendron cecidophorum</i>	562
<i>Ablania guianensis</i>	452	<i>Acinodendron chrysophylla</i>	563
<i>Ablania laurifolia</i>	452	<i>Acinodendron dispar</i>	563
<i>Abrophaes mirabilis</i>	565	<i>Acinodendron egense</i>	563
<i>Acacia</i>	504	<i>Acinodendron elatum</i>	563
<i>Acacia angustiloba</i>	505	<i>Acinodendron ellipticum</i>	569
<i>Acacia aspidioides</i>	535	<i>Acinodendron eriodontum</i>	563
<i>Acacia biceps</i>	527	<i>Acinodendron fandlerianum</i>	566
<i>Acacia catinga</i>	527	<i>Acinodendron floridum</i>	566
<i>Acacia frondosa</i>	527	<i>Acinodendron gratissimum</i>	564
<i>Acacia glomerosa</i>	537	<i>Acinodendron guianense</i>	566
<i>Acacia guacamayo</i>	504	<i>Acinodendron holosericeum</i>	564
<i>Acacia guianensis</i>	538	<i>Acinodendron hypoleucum</i>	564
<i>Acacia jupunba</i>	503	<i>Acinodendron kappleri</i>	564
<i>Acacia langlassei</i>	537	<i>Acinodendron lepidotum</i>	564
<i>Acacia leucocephala</i>	527	<i>Acinodendron longifolium</i>	565
<i>Acacia macroloba</i>	535	<i>Acinodendron longipedunculatum</i>	565
<i>Acacia mangium</i>	504	<i>Acinodendron longispicatum</i>	565
<i>Acacia microphylla</i>	505	<i>Acinodendron macrophyllum</i>	568
<i>Acacia niopo</i>	505	<i>Acinodendron matthaei</i>	565
<i>Acacia peregrina</i>	505	<i>Acinodendron melinonii</i>	565
<i>Acacia polyphylla</i>	537	<i>Acinodendron microcarpum</i>	562
<i>Acacia psilostachya</i>	535	<i>Acinodendron minutiflorum</i>	565
<i>Acacia pulcherrima</i>	538	<i>Acinodendron mucronatum</i>	564
<i>Acacia riparia</i>	537	<i>Acinodendron naudinii</i>	564
<i>Acajuba occidentalis</i>	363	<i>Acinodendron parviflorum</i>	567
ACANTHACEAE	362	<i>Acinodendron phaeophyllum</i>	566
<i>Acanthiphyllym ilicifolium</i>	585	<i>Acinodendron planinerve</i>	562
<i>Acanthiphyllym spruceanum</i>	585	<i>Acinodendron plukanetii</i>	566
<i>Acanthiphyllym strepitans</i>	585	<i>Acinodendron poeppigii</i>	566
<i>Acca glazioviana</i>	604	<i>Acinodendron prasinum</i>	567
ACHARIACEAE	362	<i>Acinodendron pteropodium</i>	567
<i>Achras balata</i>	689	<i>Acinodendron punctatum</i>	568
<i>Achras brasiliensis</i>	704	<i>Acinodendron pyrifolium</i>	568
<i>Achras buchananiifolia</i>	700	<i>Acinodendron robustum</i>	569
<i>Achras caimito</i>	694	<i>Acinodendron scorpioides</i>	569
<i>Achras glaucescens</i>	699	<i>Acinodendron seriale</i>	568
		<i>Acinodendron tomentellum</i>	568
		<i>Acinodendron tomentosum</i>	569

Appendix 3. — Continuation.

<i>Acinodendron trinervium</i>	569	<i>Acrodiclidium chrysophyllum</i>	480
<i>Acioa amara</i>	417	<i>Acrodiclidium debile</i>	480
<i>Acioa dulcis</i>	416	<i>Acrodiclidium guianense</i>	480, 481
<i>Acioa guianensis</i>	416	<i>Acrodiclidium guianense</i> var. <i>caudatum</i>	481
<i>Acioa somnolens</i>	416	<i>Acrodiclidium guianense</i> var. <i>oppositifolium</i>	480
<i>Acosmum nitens</i>	527	<i>Acrodiclidium guianense</i> var. <i>reticulatum</i>	481
<i>Acosmum paeclarum</i>	515	<i>Acrodiclidium itauba</i>	481
<i>Acrandra glazioviana</i>	604	<i>Acrodiclidium kunthianum</i>	491
<i>Acrocomia aculeata</i>	390	<i>Acrodiclidium martinianum</i>	481
<i>Acrocomia antiguana</i>	391	<i>Acrodiclidium meisneri</i>	481
<i>Acrocomia antioquiensis</i>	390	<i>Acrodiclidium pachycarpum</i>	481
<i>Acrocomia belizensis</i>	391	<i>Acrodiclidium rigidum</i>	480
<i>Acrocomia christopherensis</i>	391	<i>Acrodiclidium vernicosum</i>	481
<i>Acrocomia chunta</i>	391	<i>Actinostigma speciosum</i>	435
<i>Acrocomia cubensis</i>	390	<i>Adenobasium obtusifolium</i>	453
<i>Acrocomia erioacantha</i>	391	<i>Adenophaedra grandifolia</i>	457
<i>Acrocomia fusiformis</i>	390	<i>Adenostephanus guyanensis</i>	646
<i>Acrocomia glaucophylla</i>	390	<i>Aegiphila amazonica</i>	472
<i>Acrocomia globosa</i>	390	<i>Aegiphila arborea</i>	472
<i>Acrocomia grenadana</i>	391	<i>Aegiphila arborescens</i>	472
<i>Acrocomia guianensis</i>	390	<i>Aegiphila arborescens</i> var. <i>breviflora</i>	472
<i>Acrocomia horrida</i>	390	<i>Aegiphila arborescens</i> var. <i>longiflora</i>	472
<i>Acrocomia hospes</i>	391	<i>Aegiphila cowanii</i>	472
<i>Acrocomia ierensis</i>	391	<i>Aegiphila guianensis</i>	472
<i>Acrocomia intumescens</i>	390	<i>Aegiphila integrifolia</i>	472
<i>Acrocomia karukerana</i>	391	<i>Aegiphila integrifolia</i> var. <i>lopez-palacii</i>	472
<i>Acrocomia lasiospatha</i>	390	<i>Aegiphila parviflora</i>	472
<i>Acrocomia media</i>	391	<i>Aegiphila villosa</i>	472
<i>Acrocomia mexicana</i>	390	<i>Aegopricum betulinum</i>	463
<i>Acrocomia microcarpa</i>	390	<i>Agave foetida</i>	722
<i>Acrocomia minor</i>	390	<i>Agonandra silvatica</i>	635
<i>Acrocomia mokayayba</i>	390	<i>Aguava guianensis</i>	613
<i>Acrocomia odorata</i>	390	<i>Aguava tomentosa</i>	622
<i>Acrocomia panamensis</i>	391	<i>Aiouea demerarensis</i>	474
<i>Acrocomia pilosa</i>	391	<i>Aiouea guianensis</i>	474, 476
<i>Acrocomia quisqueyana</i>	391	<i>Aiouea impressa</i>	474
<i>Acrocomia sclerocarpa</i>	390	<i>Aiouea longipetiolata</i>	474
<i>Acrocomia sclerocarpa</i> var. <i>wallaceana</i>	390	<i>Aiouea minutiflora</i>	474
<i>Acrocomia sphaerocarpa</i>	390	<i>Aiouea montana</i>	474
<i>Acrocomia spinosa</i>	391	<i>Aiouea opaca</i>	475
<i>Acrocomia subinermis</i>	391	<i>Aiouea rubra</i>	474
<i>Acrocomia tenuifrons</i>	390	<i>Aiouea schomburgkii</i>	474
<i>Acrocomia totai</i>	390	<i>Aiouea tenella</i>	474
<i>Acrocomia ulei</i>	391	<i>Albizia</i>	504
<i>Acrocomia viegasii</i>	391	<i>Albizia caribaea</i>	504
<i>Acrocomia vinifera</i>	390	<i>Albizia corymbosa</i>	515
<i>Acrocomia wallaceana</i>	390	<i>Albizia dubia</i>	503
<i>Acrocomia zapotecis</i>	390, 391	<i>Albizia guacamayo</i>	504
<i>Acrodiclidium amarum</i>	481	<i>Albizia hassleri</i>	504
<i>Acrodiclidium anacardioides</i>	481	<i>Albizia niopoides</i>	504
<i>Acrodiclidium appelii</i>	480	<i>Albizia pedicellaris</i>	505
<i>Acrodiclidium aubletii</i>	480	<i>Alchornea acroneura</i>	458
<i>Acrodiclidium cannella</i>	480	<i>Alchornea brachygynne</i>	457
<i>Acrodiclidium cayennense</i>	480	<i>Alchornea brevistyla</i>	458

Appendix 3. — Continuation.

<i>Alchornea cordata</i>	458	<i>Alibertia latifolia</i> var. <i>parvifolia</i>	650
<i>Alchornea discolor</i>	457	<i>Alibertia myrciifolia</i>	652
<i>Alchornea gardneri</i>	457	<i>Alibertia myrciifolia</i> var. <i>tepuensis</i>	652
<i>Alchornea glandulosa</i> var. <i>floribunda</i>	458	<i>Alibertia oligantha</i>	650
<i>Alchornea glandulosa</i> var. <i>parvifolia</i>	458	<i>Alibertia steinbachii</i>	652
<i>Alchornea glaziovii</i>	457	<i>Alibertia tenuifolia</i>	652
<i>Alchornea intermedia</i>	458	<i>Alibertia triflora</i>	652
<i>Alchornea janeirensis</i>	458	<i>Alibertia triloba</i>	652
<i>Alchornea latifolia</i>	459	<i>Alibertia uniflora</i>	652
<i>Alchornea macrophylla</i>	458	<i>Alibertia utilis</i>	650
<i>Alchornea martiana</i>	459	<i>Alicastrum guianense</i>	583
<i>Alchornea nemoralis</i>	458	<i>Alicastrum rubescens</i>	584
<i>Alchornea nemoralis</i> var. <i>floribunda</i>	458	<i>Aliteria sagotii</i>	585
<i>Alchornea nemoralis</i> var. <i>intermedia</i>	458	<i>Allantoma fagifolia</i>	493
<i>Alchornea nemoralis</i> var. <i>janeirensis</i>	458	<i>Allantoma multiflora</i>	493
<i>Alchornea nemoralis</i> var. <i>lanceolata</i>	458	<i>Allantoma subramosa</i>	493
<i>Alchornea nemoralis</i> var. <i>parvifolia</i>	458	<i>Allophylus angustatus</i>	679
<i>Alchornea nemoralis</i> var. <i>psilorhachis</i>	458	<i>Allophylus edulis</i>	679
<i>Alchornea nemoralis</i> var. <i>rotundifolia</i>	458	<i>Allophylus edulis</i> var. <i>gracilis</i>	679
<i>Alchornea obovata</i>	458	<i>Allophylus edulis</i> var. <i>subsessilis</i>	679
<i>Alchornea orinocensis</i>	459	<i>Allophylus latifolius</i>	679
<i>Alchornea parvifolia</i>	458	<i>Allophylus leucoclados</i>	680
<i>Alchornea psilorhachis</i>	458	<i>Allophylus robustus</i>	680
<i>Alchornea rotundifolia</i>	458	<i>Alseis longifolia</i>	650
<i>Alchornea schomburgkii</i>	457	<i>Alseodaphne cujumary</i>	485
<i>Alchornea triplinervia</i>	458	<i>Alsodeia bahiensis</i>	720
<i>Alchornea triplinervia</i> f. <i>intermedia</i>	458	<i>Alsodeia brevipes</i>	720
<i>Alchornea triplinervia</i> f. <i>psilorhachis</i>	458	<i>Alsodeia falcata</i>	720, 721
<i>Alchornea triplinervia</i> var. <i>boliviiana</i>	458	<i>Alsodeia falcata</i> var. <i>grandifolia</i>	721
<i>Alchornea triplinervia</i> var. <i>crassifolia</i>	458	<i>Alsodeia flavescens</i>	720
<i>Alchornea triplinervia</i> var. <i>genuina</i>	458	<i>Alsodeia floribunda</i>	720
<i>Alchornea triplinervia</i> var. <i>iricuranoides</i>	458	<i>Alsodeia guianensis</i>	720, 721
<i>Alchornea triplinervia</i> var. <i>janeirensis</i>	458	<i>Alsodeia guianensis</i> var. <i>brevipes</i>	720
<i>Alchornea triplinervia</i> var. <i>laevigata</i>	458	<i>Alsodeia guianensis</i> var. <i>laxiflora</i>	720
<i>Alchornea triplinervia</i> var. <i>lanceolata</i>	458	<i>Alsodeia guianensis</i> var. <i>parviflora</i>	720
<i>Alchornea triplinervia</i> var. <i>nemoralis</i>	458	<i>Alsodeia laxiflora</i>	720
<i>Alchornea triplinervia</i> var. <i>parvifolia</i>	458	<i>Alsodeia macrocarpa</i>	720
<i>Alchornea triplinervia</i> var. <i>tomentella</i>	458	<i>Alsodeia martinii</i>	721
<i>Alchornea triplinervia</i> var. <i>trinitatis</i>	458	<i>Alsodeia paniculata</i>	721
<i>Alchorneopsis floribunda</i>	458	<i>Alsodeia piparea</i>	678
<i>Alchorneopsis trimera</i>	458	<i>Alsodeia prunifolia</i>	721
<i>Alcoceratothrix longibracteata</i>	550	<i>Alsodeia pubiflora</i>	721
<i>Alcoceratothrix rugosa</i>	550, 551	<i>Alsodeia riana</i>	721
<i>Alcoceratothrix rugosa</i> var. <i>minor</i>	551	<i>Alsodeia rinorea</i>	720
<i>Alcoceratothrix stipulacea</i>	550	<i>Amaioua affinis</i>	656
<i>Alexa</i>	505	<i>Amaioua corymbosa</i>	650
<i>Alexa grandiflora</i>	505	<i>Amaioua edulis</i>	650
<i>Alexa wachenheimii</i>	505	<i>Amaioua eriopila</i>	655
<i>Alibertia benensis</i>	652	<i>Amaioua fagifolia</i>	650
<i>Alibertia edulis</i>	650	<i>Amaioua glomerulata</i>	650
<i>Alibertia edulis</i> var. <i>edulis</i>	650	<i>Amaioua grandifolia</i>	656
<i>Alibertia granulosa</i>	650	<i>Amaioua guianensis</i>	650, 651
<i>Alibertia latifolia</i>	650	<i>Amaioua guianensis</i> var. <i>macrantha</i>	651
<i>Alibertia latifolia</i> var. <i>pargueniana</i>	650	<i>Amaioua surinamensis</i>	656

Appendix 3. — Continuation.

<i>Amaioua ursina</i>	656	<i>Anacardium spruceanum</i>	363
<i>Amaioua utilis</i>	650	<i>Anadenanthera</i>	503, 505
<i>Amanoa congesta</i>	636	<i>Anadenanthera peregrina</i>	505
<i>Amanoa divaricata</i>	638	<i>Anartia attenuata</i>	386
<i>Amanoa grandiflora</i>	636, 637	<i>Anartia flavicans</i>	386
<i>Amanoa guianensis</i>	636, 637	<i>Anartia glabrata</i>	386
<i>Amanoa guianensis</i> var. <i>grandiflora</i>	636	<i>Anartia meyeri</i>	387
<i>Amanoa macrocarpa</i>	636	<i>Anartia oblongifolia</i>	386
<i>Amanoa neglecta</i>	637	<i>Anartia olivacea</i>	386
<i>Amanoa potamophila</i>	636	<i>Anavinga parvifolia</i>	672
<i>Amanoa racemosa</i>	638	<i>Anavinga samyda</i>	675
<i>Amapa guianensis</i>	573	<i>Anaxagorea acuminata</i>	366
<i>Ambaiba humboldtiana</i>	711	<i>Anaxagorea brevipedicellata</i>	366
<i>Ambaiba latiloba</i>	711	<i>Anaxagorea costaricensis</i>	367
<i>Ambaiba obtusa</i>	711	<i>Anaxagorea dolichocarpa</i>	366
<i>Ambaiba palmata</i>	711	<i>Anaxagorea megalophylla</i>	366
<i>Ambaiba peltata</i>	711	<i>Anaxagorea multiflora</i>	366
<i>Ambaiba schiedeana</i>	711	<i>Anaxagorea mutica</i>	366
<i>Ambaiba sciadophylla</i>	712	<i>Anaxagorea phaeocarpa</i>	367
<i>Ambaiba surinamensis</i>	711	<i>Andira</i>	503
<i>Ambelania acida</i>	381	<i>Andira acuminata</i>	506
<i>Ambelania cucumerina</i>	381	<i>Andira amazonum</i>	544
<i>Ambelania sagotii</i>	381	<i>Andira aubletii</i>	544
<i>Ambelania tenuiflora</i>	381	<i>Andira bracteosa</i>	544
<i>Ambelania tenuiflora</i> var. <i>tenuiramea</i>	381	<i>Andira chiricana</i>	506
<i>Amerimnon affine</i>	506	<i>Andira coriacea</i>	506
<i>Amerimnon latifolium</i>	528	<i>Andira excelsa</i>	506
<i>Amerimnon pinnatum</i>	528, 535	<i>Andira grandiflora</i>	506
<i>Ampelocera edentula</i>	711	<i>Andira inermis</i>	506
<i>Ampelocera latifolia</i>	711	<i>Andira inermis</i> subsp. <i>grandiflora</i>	506
<i>Amphiodon</i>	505	<i>Andira inermis</i> var. <i>riedelii</i>	506
<i>Amphiodon effusus</i>	505	<i>Andira inermis</i> var. <i>sapindoides</i>	506
<i>Amphirrhox</i>	718	<i>Andira jamaicensis</i>	506
<i>Amphirrhox juruana</i>	718	<i>Andira jamaicensis</i> var. <i>sapindoides</i>	506
<i>Amphirrhox latifolia</i>	718	<i>Andira oblonga</i>	506
<i>Amphirrhox longifolia</i>	718	<i>Andira retusa</i>	506
<i>Amphirrhox surinamensis</i>	718	<i>Andira retusa</i> var. <i>oblonga</i>	506
<i>Amphymenium rohrrii</i>	536	<i>Andira retusa</i> var. <i>surinamensis</i>	506
<i>Amphymenium villosum</i>	536	<i>Andira riparia</i>	506
<i>Amyris acuminata</i>	402	<i>Andira sapindoides</i>	506
<i>Amyris altissima</i>	399	<i>Andira surinamensis</i>	506
<i>Amyris ambrosiaca</i>	402	<i>Andira surinamensis</i> var. <i>ovatifoliolata</i>	506
<i>Amyris arborescens</i>	634	<i>Andira wachenheimii</i>	506
<i>Amyris decandra</i>	400	<i>Andriapetalum cayennense</i>	647
<i>Amyris enneandra</i>	400	<i>Andriapetalum rubescens</i>	646
<i>Amyris guianensis</i>	402	<i>Andriapetalum rubescens</i> var. <i>acuminatum</i>	646
<i>Amyris heterophylla</i>	400	<i>Andriapetalum sessilifolium</i>	647
<i>Anacampa albescens</i>	387	<i>Andriapetalum sprucei</i>	646
ANACARDIACEAE	363	<i>Androstylium foceanum</i>	432
<i>Anacardium amapaense</i>	363	<i>Aneuriscus aubletii</i>	435
<i>Anacardium giganteum</i>	363	<i>Aneuriscus exserens</i>	435
<i>Anacardium microcarpum</i>	363	<i>Aniba affinis</i>	475
<i>Anacardium occidentale</i>	363	<i>Aniba anisosepala</i>	477
<i>Anacardium occidentale</i> var. <i>gardneri</i>	363	<i>Aniba castanea</i>	477

Appendix 3. — Continuation.

<i>Aniba citrifolia</i>	476	<i>Annona montana</i> f. <i>marcgravii</i>	369
<i>Aniba duckei</i>	478	<i>Annona mucosa</i>	369
<i>Aniba flexuosa</i>	479	<i>Annona muricata</i>	369
<i>Aniba gigantifolia</i>	477	<i>Annona neoelliptica</i>	369
<i>Aniba gonggrijpii</i>	477	<i>Annona obtusiflora</i>	369
<i>Aniba guianensis</i>	474, 476	<i>Annona obtusifolia</i>	369
<i>Aniba hostmanniana</i>	477	<i>Annona paludosa</i>	370
<i>Aniba jenmanii</i>	477	<i>Annona palustris</i>	367, 368
<i>Aniba kappleri</i>	477	<i>Annona palustris</i> var. <i>grandifolia</i>	368
<i>Aniba koumaroucapa</i>	477	<i>Annona parviflora</i>	369
<i>Aniba mas</i>	477	<i>Annona peduncularis</i>	378
<i>Aniba megacarpa</i>	480	<i>Annona perrottetii</i>	378
<i>Aniba megaphylla</i>	477	<i>Annona peruviana</i>	368
<i>Aniba muelleriana</i>	476	<i>Annona pisonis</i>	368, 369
<i>Aniba muricana</i>	478	<i>Annona poeppigiana</i>	370
<i>Aniba opaca</i>	476	<i>Annona prevostiae</i>	370
<i>Aniba ovalifolia</i>	485	<i>Annona pterocarpa</i>	369
<i>Aniba panurensis</i>	477	<i>Annona pteropetala</i>	369
<i>Aniba parviflora</i>	477	<i>Annona punctata</i>	374
<i>Aniba pittieri</i>	477	<i>Annona reticulata</i>	369
<i>Aniba reticulata</i>	479	<i>Annona reticulata</i> var. <i>mucosa</i>	369
<i>Aniba riparia</i>	477	<i>Annona rhombipetala</i>	372
<i>Aniba rosodora</i>	477	<i>Annona sericea</i>	370
<i>Aniba rosodora</i> var. <i>amazonica</i>	477	<i>Annona sessiliflora</i>	373
<i>Aniba salicifolia</i>	476	<i>Annona sphaerocarpa</i>	369
<i>Aniba simulans</i>	478	<i>Annona squamosa</i>	369
<i>Aniba taubertiana</i>	478	<i>Annona symphyocarpa</i>	370
<i>Aniba terminalis</i>	478	<i>Annona tenuiflora</i>	370
<i>Aniba tessmannii</i>	476	<i>Annona tenuipes</i>	370
<i>Aniba trinitatis</i>	476	<i>Annona tessmannii</i>	369
<i>Aniba williamsii</i>	478	<i>Annona trinitensis</i>	370
<i>Aniba</i> sp. A	478	<i>Annona trunciflora</i>	367
<i>Aniba</i> sp. B	478	<i>Annona uliginosa</i>	368
<i>Aniba</i> sp. C	478	<i>Anomalocalyx uleanus</i>	462
<i>Anneslia fasciculata</i>	507	<i>Anonocarpus amazonicus</i>	583
<i>Annona ambotay</i>	367	<i>Anthodon ellipticus</i>	415
<i>Annona australis</i>	368	<i>Anthodon glomeratus</i>	415
<i>Annona axilliflora</i>	374	<i>Anthodon grandiflorus</i>	416
<i>Annona biflora</i>	369	<i>Anthodon oblongifolius</i>	415
<i>Annona calycina</i>	371	<i>Antidesma triplinervium</i>	458
ANNONACEAE	366	<i>Antirhea guianensis</i>	666
<i>Annona chrysopetala</i>	375	<i>Antirhea panamensis</i>	666
<i>Annona cuspidata</i>	367	<i>Antirhea surinamensis</i>	666
<i>Annona exsucca</i>	367, 371, 372, 374, 379	<i>Antonia laurifolia</i>	657
<i>Annona foetida</i>	367	<i>Antonia ovata</i>	546
<i>Annona glabra</i>	367, 368	<i>Antonia ovata</i> var. <i>excelsa</i>	546
<i>Annona hostmannii</i>	376	<i>Antonia ovata</i> var. <i>pilosa</i>	546
<i>Annona humilis</i>	370	<i>Antonia pilosa</i>	546
<i>Annona hypoglauca</i>	369	<i>Antonia pubescens</i>	546
<i>Annona jenmanii</i>	370	<i>Apalatoa aromatica</i>	509
<i>Annona laurifolia</i>	368	<i>Apalatoa oblonga</i>	509
<i>Annona longifolia</i>	372	<i>Apalatoa pubescens</i>	509
<i>Annona marcgravii</i>	369	<i>Apalatoa spicata</i>	511
<i>Annona montana</i>	369	<i>Apalatoa tomentosa</i>	511

Appendix 3. — Continuation.

<i>Aparisthium cordatum</i>	458	<i>Artanthe celtidifolia</i>	640
<i>Aparisthium macrophyllum</i>	458	<i>Artanthe cernua</i>	642
<i>Aparisthium spruceanum</i>	459	<i>Artanthe elongata</i>	640
<i>Apatitia blakeoides</i>	560	<i>Artanthe eximia</i>	642
<i>Apeiba albiflora</i>	551	<i>Artanthe galleottii</i>	640
<i>Apeiba aspera</i>	551	<i>Artanthe richardifolia</i>	642
<i>Apeiba burchellii</i>	551	<i>Artanthe ruiziana</i>	642
<i>Apeiba echinata</i>	551	<i>Artanthe spectabilis</i>	642
<i>Apeiba glabra</i>	551	<i>Artanthe spectabilis</i> var. <i>glabrior</i>	642
<i>Apeiba hispida</i>	551	<i>Arthrosamanea corymbosa</i>	515
<i>Apeiba hypoleuca</i>	551	<i>Aruba cedron</i>	705
<i>Apeiba laevis</i>	551	<i>Aruba crustacea</i>	706
<i>Apeiba petoumo</i>	551	<i>Aruba guianensis</i>	705
<i>Apeiba tibourbou</i>	551	<i>Aruba orinocensis</i>	706
<i>Apeiba tibourbou</i> var. <i>krukoffii</i>	551	<i>Arum sagittaefolium</i>	722
<i>Apeiba tibourbou</i> var. <i>rugosa</i>	551	<i>Asimina arborea</i>	368
<i>Apeiba ulmifolia</i>	555	<i>Aspidosperma album</i>	381
<i>Apeiba</i> sp. A.	551	<i>Aspidosperma carapanauba</i>	381
APOCYNACEAE	381	<i>Aspidosperma centrale</i>	381
AQUIFOLIACEAE	388	<i>Aspidosperma chiapense</i>	381
<i>Arachis hypogaea</i>	722	<i>Aspidosperma chiapense</i> f. <i>tenax</i>	381
<i>Aralia capitata</i>	390	<i>Aspidosperma cruentum</i>	381
ARALIACEAE	388, 389	<i>Aspidosperma desmanthum</i>	381
<i>Aralia ovata</i>	390	<i>Aspidosperma excelsum</i>	381
<i>Arawakia</i>	438, 439	<i>Aspidosperma helstonei</i>	382
<i>Arawakia angustata</i>	439	<i>Aspidosperma kuhlmannii</i>	382
<i>Arawakia caputmonisia</i>	439	<i>Aspidosperma latisiliquum</i>	381
<i>Arawakia coriacea</i>	439	<i>Aspidosperma macrophyllum</i> subsp. <i>mori</i>	382
<i>Arawakia divesora</i>	439	<i>Aspidosperma marcgavianum</i>	382
<i>Arawakia glossophylla</i>	439	<i>Aspidosperma matudae</i>	381
<i>Arawakia lanceolata</i>	439	<i>Aspidosperma oblongum</i>	382
<i>Arawakia lingulata</i>	438	<i>Aspidosperma sandwithianum</i>	382
<i>Arawakia longicuneata</i>	438	<i>Aspidosperma sanguineum</i>	381
<i>Arawakia macrocarpa</i>	438	<i>Aspidosperma schultesii</i>	382
<i>Arawakia manchamancha</i>	439	<i>Aspidosperma</i> sp. A.	382
<i>Arawakia ob lanceolata</i>	438	<i>Aspidosperma spruceanum</i>	382
<i>Arawakia panamaea</i>	439	<i>Asterolepidion elatum</i>	582
<i>Arawakia parvifolia</i>	439	<i>Astrocaryum awarra</i>	392
<i>Arawakia pithecozia</i>	439	<i>Astrocaryum guara</i>	391
<i>Arawakia rhizophoroides</i>	438	<i>Astrocaryum guianense</i>	392
<i>Arawakia rileyi</i>	439	<i>Astrocaryum jauari</i>	391
<i>Arawakia sphenophylla</i>	438	<i>Astrocaryum minus</i>	391
<i>Arawakia weddelliana</i>	438	<i>Astrocaryum murumuru</i>	391
<i>Ardisia hostmannii</i>	646	<i>Astrocaryum rodriquesii</i>	391
<i>Ardisia orinocensis</i>	646	<i>Astrocaryum sciophilum</i>	391
<i>Ardisia schomburgkiana</i>	646	<i>Astrocaryum segregatum</i>	392
<i>Ardisia surinamensis</i>	646	<i>Astrocaryum tucumoides</i>	392
<i>Areca bacaba</i>	394	<i>Astrocaryum vulgare</i>	392
ARECACEAE	390	<i>Astrocaryum yauaperyense</i>	391
<i>Aristoclesia esculenta</i>	435	<i>Astronium lecointei</i>	363
<i>Arouna guianensis</i>	511	<i>Astronium obliquum</i>	363
<i>Arrudea purpurea</i>	433	<i>Astronium ulei</i>	363
<i>Artanthe adunca</i>	640	<i>Athenaea guianensis</i>	673
<i>Artanthe adunca</i> f. <i>angustifolia</i>	640	<i>Attalea cryptantha</i>	392

Appendix 3. — Continuation.

<i>Attalea macropetala</i>	392	<i>Aulomyrcia edulis</i>	617, 618
<i>Attalea maripa</i>	392	<i>Aulomyrcia elaeodendra</i>	613
<i>Attalea regia</i>	392	<i>Aulomyrcia emarginata</i>	614
<i>Aubletia laevis</i>	551	<i>Aulomyrcia exsucca</i>	613
<i>Aubletia petoumo</i>	551	<i>Aulomyrcia fragilis</i>	614
<i>Aubletia tibourbou</i>	551	<i>Aulomyrcia gardneriana</i>	614
<i>Aulomyrcia alloiota</i>	623	<i>Aulomyrcia gardneriana</i> var. <i>caerulescens</i>	614
<i>Aulomyrcia alloiota</i> var. <i>cuneata</i>	623	<i>Aulomyrcia gardneriana</i> var. <i>virescens</i>	614
<i>Aulomyrcia alloiota</i> var. <i>obovata</i>	623	<i>Aulomyrcia glandulosa</i>	614
<i>Aulomyrcia alloiota</i> var. <i>ovalis</i>	623	<i>Aulomyrcia glandulosa</i> var. <i>elliptica</i>	614
<i>Aulomyrcia alloiota</i> var. <i>pyramidalis</i>	623	<i>Aulomyrcia glandulosa</i> var. <i>longifolia</i>	614
<i>Aulomyrcia alloiota</i> var. <i>subcordata</i>	623	<i>Aulomyrcia glandulosa</i> var. <i>obovata</i>	614
<i>Aulomyrcia alternifolia</i>	613	<i>Aulomyrcia glaucescens</i>	616
<i>Aulomyrcia amazonica</i>	609	<i>Aulomyrcia glaucescens</i> var. <i>grandifolia</i>	616
<i>Aulomyrcia androsaemoides</i>	615	<i>Aulomyrcia glaucescens</i> var. <i>parvifolia</i>	616
<i>Aulomyrcia angustifolia</i>	614	<i>Aulomyrcia goyazensis</i>	616
<i>Aulomyrcia aureolanata</i>	623	<i>Aulomyrcia hepatica</i>	614
<i>Aulomyrcia biformis</i>	614	<i>Aulomyrcia hirtiflora</i>	622
<i>Aulomyrcia bimarginata</i>	614	<i>Aulomyrcia hostmanniana</i>	609
<i>Aulomyrcia botrys</i>	614	<i>Aulomyrcia hostmanniana</i> var. <i>gracilior</i>	609
<i>Aulomyrcia bracteata</i>	614	<i>Aulomyrcia hostmanniana</i> var. <i>robustior</i>	609
<i>Aulomyrcia buxifolia</i>	616	<i>Aulomyrcia inaequiloba</i>	615, 616
<i>Aulomyrcia buxifolia</i> var. <i>elliptica</i>	616	<i>Aulomyrcia inaequiloba</i> var. <i>nitida</i>	616
<i>Aulomyrcia buxifolia</i> var. <i>ovalis</i>	616	<i>Aulomyrcia inaequiloba</i> var. <i>paniculata</i>	615
<i>Aulomyrcia buxizans</i>	615	<i>Aulomyrcia jequitinhonhensis</i> var. <i>glauca</i>	615
<i>Aulomyrcia caeruleascens</i>	616	<i>Aulomyrcia lancea</i>	623
<i>Aulomyrcia caesia</i>	616	<i>Aulomyrcia lancifolia</i>	615
<i>Aulomyrcia camaraeana</i>	616	<i>Aulomyrcia lanuginosa</i>	623
<i>Aulomyrcia capivarhyensis</i>	623	<i>Aulomyrcia lanuginosa</i> var. <i>pyramidata</i>	623
<i>Aulomyrcia cassinioides</i>	613, 614	<i>Aulomyrcia laruotteana</i> var. <i>peruviana</i>	616
<i>Aulomyrcia cassinioides</i> var. <i>glabrata</i>	614	<i>Aulomyrcia lauriflora</i>	613
<i>Aulomyrcia cassinioides</i> var. <i>velutina</i>	614	<i>Aulomyrcia leptoclada</i>	609
<i>Aulomyrcia chrysophylla</i>	611	<i>Aulomyrcia leucadendron</i>	613
<i>Aulomyrcia clauseniana</i>	614	<i>Aulomyrcia leucantha</i>	623
<i>Aulomyrcia conduplicata</i>	613	<i>Aulomyrcia lingua</i>	614
<i>Aulomyrcia confusa</i>	623	<i>Aulomyrcia lingua</i> var. <i>glabrata</i>	614
<i>Aulomyrcia corymbosa</i>	609	<i>Aulomyrcia lingua</i> var. <i>rufa</i>	614
<i>Aulomyrcia costata</i>	620, 622	<i>Aulomyrcia longipes</i>	623
<i>Aulomyrcia coumete</i>	611	<i>Aulomyrcia longipes</i> var. <i>latifolia</i>	623
<i>Aulomyrcia crassicaulis</i>	613	<i>Aulomyrcia longipes</i> var. <i>obovata</i>	623
<i>Aulomyrcia cuneata</i>	613	<i>Aulomyrcia longipes</i> var. <i>spathulata</i>	623
<i>Aulomyrcia cuprea</i>	611	<i>Aulomyrcia mansoni</i>	614
<i>Aulomyrcia curatellifolia</i>	622, 623	<i>Aulomyrcia maritima</i>	614
<i>Aulomyrcia curatellifolia</i> var. <i>australis</i>	623	<i>Aulomyrcia micrantha</i>	623
<i>Aulomyrcia curatellifolia</i> var. <i>grandifolia</i>	623	<i>Aulomyrcia microcarpa</i>	613
<i>Aulomyrcia curatellifolia</i> var. <i>parvifolia</i>	623	<i>Aulomyrcia minutiflora</i>	616
<i>Aulomyrcia daphnoides</i>	613	<i>Aulomyrcia multiflora</i>	616
<i>Aulomyrcia decrescens</i>	614	<i>Aulomyrcia multiflora</i> var. <i>grandifolia</i>	616
<i>Aulomyrcia desertorum</i>	614	<i>Aulomyrcia nigropunctata</i>	614
<i>Aulomyrcia detergens</i>	609	<i>Aulomyrcia obscura</i>	614, 615
<i>Aulomyrcia detergens</i> var. <i>depauperata</i>	609	<i>Aulomyrcia obscura</i> var. <i>longipes</i>	615
<i>Aulomyrcia detergens</i> var. <i>dives</i>	609	<i>Aulomyrcia obtecta</i>	614
<i>Aulomyrcia dichroma</i>	613	<i>Aulomyrcia obtusa</i>	613
<i>Aulomyrcia dumosa</i>	617	<i>Aulomyrcia obtusa</i> var. <i>grandifolia</i>	613

Appendix 3. — Continuation.

<i>Aulomyrcia obtusa</i> var. <i>longipes</i>	613	<i>Aulomyrcia sphaerocarpa</i> var. <i>intermedia</i>	616
<i>Aulomyrcia obtusa</i> var. <i>panicularis</i>	613	<i>Aulomyrcia sphaerocarpa</i> var. <i>obtusata</i>	616
<i>Aulomyrcia obtusa</i> var. <i>pauciflora</i>	613	<i>Aulomyrcia sphaerocarpa</i> var. <i>ovata</i>	616
<i>Aulomyrcia obtusa</i> var. <i>schomburgkiana</i>	613	<i>Aulomyrcia sphaerocarpa</i> var. <i>pauciflora</i>	616
<i>Aulomyrcia obtusa</i> var. <i>surinamensis</i>	613	<i>Aulomyrcia spruceana</i>	609
<i>Aulomyrcia obtusa</i> var. <i>tenuifolia</i>	613	<i>Aulomyrcia suaveolens</i>	613
<i>Aulomyrcia ottonis</i>	623	<i>Aulomyrcia subobliqua</i>	622
<i>Aulomyrcia ovalifolia</i>	616	<i>Aulomyrcia suffruticosa</i>	614
<i>Aulomyrcia ovata</i>	618	<i>Aulomyrcia surinamensis</i>	613
<i>Aulomyrcia pallens</i>	613, 614	<i>Aulomyrcia tetramera</i>	611
<i>Aulomyrcia pallens</i> var. <i>ovalis</i>	614	<i>Aulomyrcia tobagensis</i>	611
<i>Aulomyrcia pallens</i> var. <i>ovata</i>	614	<i>Aulomyrcia tomentosa</i>	622
<i>Aulomyrcia pallens</i> var. <i>subcordata</i>	614	<i>Aulomyrcia uaupensis</i>	614
<i>Aulomyrcia pallida</i>	616	<i>Aulomyrcia vacciniifolia</i>	614
<i>Aulomyrcia paniculata</i>	615	<i>Aulomyrcia vauthieriana</i>	609
<i>Aulomyrcia paraensis</i>	609	<i>Aulomyrcia velhensis</i>	615
<i>Aulomyrcia perforata</i>	616	<i>Aulomyrcia vinacea</i>	616
<i>Aulomyrcia pirarensis</i>	615	<i>Aulomyrcia wullschlaegeliana</i>	619
<i>Aulomyrcia platyclada</i>	617	<i>Avicennia</i>	362
<i>Aulomyrcia platyclada</i> var. <i>kaieteurensis</i>	617	<i>Avicennia africana</i>	362
<i>Aulomyrcia plumbea</i>	614	<i>Avicennia elliptica</i>	362
<i>Aulomyrcia poeppigiana</i>	614	<i>Avicennia floridana</i>	362
<i>Aulomyrcia polymorpha</i>	611	<i>Avicennia germinans</i>	362
<i>Aulomyrcia polymorpha</i> var. <i>decorticans</i>	611	<i>Avicennia germinans</i> f. <i>aberrans</i>	362
<i>Aulomyrcia polymorpha</i> var. <i>duriuscula</i>	611	<i>Avicennia germinans</i> f. <i>brasiliensis</i>	362
<i>Aulomyrcia polymorpha</i> var. <i>lasiopus</i>	611	<i>Avicennia germinans</i> f. <i>venezuelensis</i>	362
<i>Aulomyrcia polymorpha</i> var. <i>leucophloea</i>	611	<i>Avicennia germinans</i> var. <i>cumanensis</i>	362
<i>Aulomyrcia pruinosa</i>	613	<i>Avicennia germinans</i> var. <i>guayaquilensis</i>	362
<i>Aulomyrcia prunifolia</i>	622, 623	<i>Avicennia lamarckiana</i>	362
<i>Aulomyrcia prunifolia</i> var. <i>brevipes</i>	623	<i>Avicennia meyeri</i>	362
<i>Aulomyrcia prunifolia</i> var. <i>longipes</i>	623	<i>Avicennia nitida</i>	362
<i>Aulomyrcia puberula</i>	623	<i>Avicennia nitida</i> var. <i>trinitensis</i>	362
<i>Aulomyrcia pusilla</i>	614	<i>Avicennia oblongifolia</i>	362
<i>Aulomyrcia pyrifolia</i>	618	<i>Avicennia officinalis</i> var. <i>nitida</i>	—
<i>Aulomyrcia pyrifolia</i> var. <i>gracilis</i>	618	<i>Avicennia tomentosa</i>	362
<i>Aulomyrcia pyrifolia</i> var. <i>robusta</i>	618	<i>Avicennia tomentosa</i> var. <i>campechensis</i>	362
<i>Aulomyrcia regeliana</i>	615	<i>Avicennia tomentosa</i> var. <i>cumanensis</i>	362
<i>Aulomyrcia regeliana</i> var. <i>oppositifolia</i>	615	<i>Avicennia tomentosa</i> var. <i>guayaquilensis</i>	362
<i>Aulomyrcia regeliana</i> var. <i>sparsifolia</i>	615	<i>Aydendron aciphyllum</i>	491
<i>Aulomyrcia roraimae</i>	615	<i>Aydendron affine</i>	475
<i>Aulomyrcia roraimensis</i>	613	<i>Aydendron cannella</i>	480
<i>Aulomyrcia rosulans</i>	623	<i>Aydendron cayennense</i>	480
<i>Aulomyrcia rotundifolia</i>	614	<i>Aydendron citrifolium</i>	476
<i>Aulomyrcia salzmannii</i>	611	<i>Aydendron cujumary</i>	485
<i>Aulomyrcia saxatilis</i>	618	<i>Aydendron hostmannianum</i>	477
<i>Aulomyrcia schaueriana</i>	612	<i>Aydendron pachycarpum</i>	481
<i>Aulomyrcia schomburgkiana</i>	613	<i>Aydendron panurensense</i>	477
<i>Aulomyrcia schrankiana</i>	613	<i>Aydendron parviflorum</i>	477
<i>Aulomyrcia scrobiculata</i>	615	<i>Aydendron riparium</i>	477
<i>Aulomyrcia spathulata</i>	623	<i>Aydendron salicifolium</i>	476
<i>Aulomyrcia sphaerocarpa</i>	616	<i>Aydendron trinitatis</i>	476
<i>Aulomyrcia sphaerocarpa</i> var. <i>arborescens</i>	616	<i>Ayenia morii</i>	551
<i>Aulomyrcia sphaerocarpa</i> var. <i>complicata</i>	616		
<i>Aulomyrcia sphaerocarpa</i> var. <i>gracilis</i>	616		

Appendix 3. — Continuation.

B	
<i>Bactris globosa</i>	390, 391
<i>Bactris minor</i>	390
<i>Bactris pavoniana</i>	390
<i>Bactris sciophila</i>	391
<i>Bactyriobium grande</i>	508
<i>Bactyriobium molle</i>	508
<i>Badamia commersonii</i>	441
<i>Badula schomburgkiana</i>	646
<i>Badula schomburgkiana</i> var. <i>brasiliensis</i>	646
<i>Bagassa guianensis</i>	583
<i>Bagassa sagotiana</i>	583
<i>Bagassa tiliifolia</i>	583
<i>Balanitium cordifolium</i>	430
<i>Balizia pedicellaris</i>	505
<i>Banara fagifolia</i>	671
<i>Banara glandulosa</i>	671
<i>Banara guianensis</i>	671
<i>Banara guianensis</i> var. <i>mollis</i>	671
<i>Banara guianensis</i> var. <i>spruceana</i>	671
<i>Banara laxiflora</i>	677
<i>Banara mollis</i>	671
<i>Banara pubescens</i>	671
<i>Banara pyramidata</i>	671
<i>Banara tulasnei</i>	671
<i>Barthollesia excelsa</i>	492
<i>Basanacantha phyllosepala</i>	665
<i>Basanacantha spinosa</i>	665
<i>Basanacantha spinosa</i> f. <i>grandiflora</i>	665
<i>Basanacantha spinosa</i> f. <i>puberula</i>	665
<i>Basanacantha spinosa</i> f. <i>pubiloba</i>	665
<i>Basanacantha spinosa</i> var. <i>longipedunculata</i>	665
<i>Basanacantha spinosa</i> var. <i>macrocalyx</i>	665
<i>Basanacantha spinosa</i> var. <i>polyantha</i>	665
<i>Basanacantha spinosa</i> var. <i>typica</i>	665
<i>Batesia</i>	506
<i>Batesia floribunda</i>	506
<i>Bathysa difformis</i>	652
<i>Batocarpus amazonicus</i>	583
<i>Batocarpus maranhensis</i>	583
<i>Bauhinia</i>	507
<i>Bauhinia cinnamomea</i>	507
<i>Bauhinia eilertssii</i>	507
<i>Bauhinia versteegii</i>	507
<i>Beilschmiedia hexanthera</i>	478
<i>Bellucia arborescens</i>	559
<i>Bellucia aubletii</i>	560
<i>Bellucia brasiliensis</i>	560
<i>Bellucia circumscissa</i>	560
<i>Bellucia egensis</i>	559
<i>Bellucia grossularioides</i>	560
<i>Bellucia hostmannii</i>	560
<i>Bellucia macrophylla</i>	560
<i>Bellucia mespiloides</i>	560
<i>Bellucia multiflora</i>	560
<i>Bellucia nervosa</i>	560
<i>Bellucia quinquenervia</i>	560
<i>Bellucia subrotundifolia</i>	560
<i>Bellucia superba</i>	560
<i>Bernardia denticulata</i>	457
<i>Bernardia grandifolia</i>	457
<i>Bernoullia swietenoides</i>	555
<i>Bertholletia excelsa</i>	492
<i>Bertholletia nobilis</i>	492
<i>Besleria caerulea</i>	722
<i>Bignonia aquatilis</i>	398
<i>Bignonia araliacea</i>	396
BIGNONIACEAE	395
<i>Bignonia conspicua</i>	396
<i>Bignonia copaia</i>	396
<i>Bignonia digitata</i>	398
<i>Bignonia dura</i>	398
<i>Bignonia filicifolia</i>	398
<i>Bignonia flavescens</i>	396
<i>Bignonia fluviatilis</i>	398
<i>Bignonia latisiliqua</i>	381
<i>Bignonia procera</i>	396
<i>Bignonia serratifolia</i>	396
<i>Bignonia triflora</i>	655
<i>Billardiera</i>	655
<i>Billardiera paniculata</i>	654
<i>Bixa arborea</i>	398
BIXACEAE	398
<i>Bixa</i> sp. A	398
<i>Blakea macrophylla</i>	560
<i>Blakea quinquenervia</i>	560
<i>Blakea triplinervia</i>	560
<i>Blochmannia weigeltiana</i>	644
<i>Blondea latifolia</i>	452, 453
<i>Bocagea asbeckii</i>	377
<i>Bocagea multiflora</i>	370
<i>Bocageopsis multiflora</i>	370
<i>Bocageopsis multiflora</i> var. <i>angustifolia</i>	370
<i>Bocoa</i>	503, 507
<i>Bocoa prouacensis</i>	507
<i>Bocoa viridiflora</i>	507
<i>Bombacopsis macrocalyx</i>	556
<i>Bombax aquaticum</i>	556
<i>Bombax crassum</i>	553
<i>Bombax cumanense</i>	552
<i>Bombax flaviflorum</i>	556
<i>Bombax globosum</i>	553
<i>Bombax guineense</i>	552
<i>Bombax macrocalyx</i>	556
<i>Bombax macrocarpum</i>	556
<i>Bombax mompxoxense</i>	552
<i>Bombax munguba</i>	556

Appendix 3. — Continuation.

<i>Bombax occidentale</i>	552	<i>Brosimum palmarum</i>	583
<i>Bombax orientale</i>	552	<i>Brosimum panamense</i>	583
<i>Bombax orinocense</i>	398	<i>Brosimum paraense</i>	584
<i>Bombax pentandrum</i>	552	<i>Brosimum parinarioides</i>	584
<i>Bombax rigidifolium</i>	556	<i>Brosimum platyneurum</i>	584
<i>Bombax sclerophyllum</i>	553	<i>Brosimum rigidum</i>	584
<i>Bombax spruceanum</i>	556	<i>Brosimum rotundatum</i>	583
<i>Bombax surinamense</i>	553	<i>Brosimum rubescens</i>	584
<i>Bonafousia attenuata</i>	386	<i>Brosimum tessmannii</i>	583
<i>Bonafousia latiflora</i>	386	<i>Brosimum utile</i> subsp. <i>ovatifolium</i>	584
<i>Bonafousia obliqua</i>	387	<i>Brosimum velutinum</i>	583
<i>Bonafousia oblongifolia</i>	386	<i>Brotera maritima</i>	555
<i>Bonafousia olivacea</i>	386	<i>Bruinsmania insertoides</i>	660
<i>Bonafousia perrottetii</i>	387	<i>Bubroma guazuma</i>	553
<i>Bonafousia sananho</i>	387	<i>Bubroma invira</i>	553
<i>Bonafousia undulata</i>	387	<i>Bubroma polybotryum</i>	553
<i>Bonafousia undulata</i> var. <i>ovalifolia</i>	387	<i>Bubroma tomentosum</i>	553
<i>Bonnetia meridionalis</i>	408	<i>Bubroma ulmifolia</i>	553
<i>Bonnetia palustris</i>	408	<i>Buceras catappa</i>	440
<i>Bontia</i>	362	<i>Buchenavia amazonia</i>	443
<i>Bontia germinans</i>	362	<i>Buchenavia capitata</i>	443
<i>Borellia aspera</i>	446	<i>Buchenavia discolor</i>	443
<i>Botryarrhena pendula</i>	651	<i>Buchenavia gracilis</i>	443
<i>Botryodendrum capitatum</i>	390	<i>Buchenavia grandis</i>	441
<i>Bourreria viridis</i>	446	<i>Buchenavia guianensis</i>	440
<i>Bowdichia brachypetala</i>	509	<i>Buchenavia huberi</i>	441
<i>Bowdichia guianensis</i>	513	<i>Buchenavia macahensis</i>	443
<i>Bowdichia martiusii</i>	512	<i>Buchenavia macrophylla</i>	442
<i>Braddleya legalis</i>	718	<i>Buchenavia megalophylla</i>	442
<i>Brignolia acuminata</i>	660	<i>Buchenavia nitidissima</i>	442
<i>Brignolia pubigera</i>	660	<i>Buchenavia ochroprumna</i>	443
<i>Britoa acida</i>	625	<i>Buchenavia parvifolia</i>	443
<i>Brosimopsis acutifolia</i>	583	<i>Buchenavia ptariensis</i>	443
<i>Brosimopsis amplifolia</i>	584	<i>Buchenavia stellae</i>	442
<i>Brosimopsis diandra</i>	584	<i>Buchenavia tetraphylla</i>	443
<i>Brosimopsis lactescens</i>	584	<i>Buchenavia vaupesana</i>	443
<i>Brosimopsis oblongifolia</i>	584	<i>Buchenavia viridiflora</i>	443
<i>Brosimum acutifolium</i>	583	<i>Bucida angustifolia</i>	440, 443
<i>Brosimum angustifolium</i>	584	<i>Bucida buceras</i> var. <i>angustifolia</i>	440
<i>Brosimum aubletii</i>	583	<i>Bucida capitata</i>	443
<i>Brosimum belicense</i>	584	<i>Buena lambertiana</i>	661
<i>Brosimum brevipedunculatum</i>	584	<i>Buena latifolia</i>	653
<i>Brosimum caloxylon</i>	584	<i>Buena obtusifolia</i>	653
<i>Brosimum discolor</i>	583	<i>Buena skinneri</i>	653
<i>Brosimum guianense</i>	583	<i>Buena triflora</i>	653
<i>Brosimum krukovi</i>	584	<i>Bumelia cuneifolia</i>	686
<i>Brosimum lactescens</i>	584	<i>Bumelia egensis</i>	691
<i>Brosimum lanciferum</i>	584	<i>Bumelia nervosa</i>	699, 700
<i>Brosimum lecointei</i>	583	<i>Bunchosia argentea</i>	547
<i>Brosimum lemeei</i>	583	<i>Bunchosia decussiflora</i>	547
<i>Brosimum longistipulatum</i>	584	<i>Burhardia aromatic</i>	597
<i>Brosimum ojoche</i>	584	<i>Burhardia grandiflora</i>	598
<i>Brosimum ovatifolium</i>	584	<i>Bursera altissima</i>	399
<i>Brosimum pallescens</i>	584	<i>Bursera aracouchini</i>	400

Appendix 3. — Continuation.

<i>Bursera caudata</i>	404	<i>Byrsonima rugosa</i>	550
BURSERACEAE	399	<i>Byrsonima sericea</i>	548
<i>Bursera decandra</i>	400	<i>Byrsonima sericea</i> f. <i>eglandulosa</i>	548
<i>Byrsonima</i>	551	<i>Byrsonima sericea</i> var. <i>angustifolia</i>	548
<i>Byrsonima aerugo</i>	547	<i>Byrsonima sericea</i> var. <i>eglandulosa</i>	548
<i>Byrsonima altissima</i>	547	<i>Byrsonima sericea</i> var. <i>pubescens</i>	548
<i>Byrsonima amazonica</i>	548	<i>Byrsonima spicata</i>	549
<i>Byrsonima amazonica</i> var. <i>lucidula</i>	548	<i>Byrsonima spicata</i> f. <i>propinqua</i>	549
<i>Byrsonima angustifolia</i>	549	<i>Byrsonima stipulacea</i>	550
<i>Byrsonima aubletii</i>	547	<i>Bytneria morii</i>	551
<i>Byrsonima berteroana</i>	549		
<i>Byrsonima biacuminata</i>	549	C	
<i>Byrsonima christianeae</i>	547	<i>Cabralea brachystachya</i>	572
<i>Byrsonima chrysophylla</i>	549	<i>Cabralea burchellii</i>	572
<i>Byrsonima chrysophylla</i> f. <i>glandulifera</i>	549	<i>Cabralea cangerana</i>	572
<i>Byrsonima chrysophylla</i> f. <i>kunthiana</i>	549	<i>Cabralea canjerana</i>	572
<i>Byrsonima coleostachya</i>	551	<i>Cabralea cauliflora</i>	572
<i>Byrsonima coriacea</i>	548-550	<i>Cabralea corcovadensis</i>	572
<i>Byrsonima coriacea</i> f. <i>angustifolia</i>	549	<i>Cabralea eichleriana</i>	572
<i>Byrsonima coriacea</i> f. <i>eglandulosa</i>	550	<i>Cabralea eichleriana</i> var. <i>macrantha</i>	572
<i>Byrsonima coriacea</i> f. <i>propinqua</i>	549	<i>Cabralea erismatica</i>	572
<i>Byrsonima coriacea</i> var. <i>spicata</i>	549	<i>Cabralea estrellensis</i>	572
<i>Byrsonima cotinifolia</i>	548	<i>Cabralea gaudichaudii</i>	572
<i>Byrsonima crassifolia</i>	547	<i>Cabralea glaberrima</i>	572
<i>Byrsonima cumingiana</i>	548	<i>Cabralea glaziovii</i>	572
<i>Byrsonima densa</i>	548	<i>Cabralea jussiaeana</i>	572
<i>Byrsonima densa</i> var. <i>emarginata</i>	548	<i>Cabralea lacaziana</i>	572
<i>Byrsonima discolor</i>	547	<i>Cabralea laevis</i>	572
<i>Byrsonima eglandulosa</i>	548	<i>Cabralea lagoensis</i>	572
<i>Byrsonima elegans</i>	551	<i>Cabralea lagoensis</i> var. <i>glabra</i>	572
<i>Byrsonima fagifolia</i>	548	<i>Cabralea lundii</i>	572
<i>Byrsonima fendleri</i>	548	<i>Cabralea macrantha</i>	572
<i>Byrsonima ferruginea</i>	548	<i>Cabralea macrophylla</i>	572
<i>Byrsonima guadalupensis</i>	549	<i>Cabralea macrophylla</i> var. <i>decomposita</i>	572
<i>Byrsonima herbert-smithii</i>	551	<i>Cabralea multijuga</i>	572
<i>Byrsonima horneana</i>	549	<i>Cabralea oblongifoliola</i>	572
<i>Byrsonima hostmannii</i>	549	<i>Cabralea pallescens</i>	572
<i>Byrsonima karwinskiana</i>	548	<i>Cabralea pedunculata</i>	572
<i>Byrsonima krukoffii</i>	548	<i>Cabralea pilosa</i>	572
<i>Byrsonima laevigata</i>	548	<i>Cabralea pilosa</i> var. <i>glabrior</i>	572
<i>Byrsonima lanceolata</i>	548	<i>Cabralea poeppigii</i>	572
<i>Byrsonima laurifolia</i>	548	<i>Cabralea riedelii</i>	572
<i>Byrsonima laurifolia</i> var. <i>guatemalensis</i>	548	<i>Cabralea rojasii</i>	572
<i>Byrsonima longibracteata</i>	550	<i>Cabralea schwackei</i>	572
<i>Byrsonima lucidula</i>	548	<i>Cabralea silvatica</i>	572
<i>Byrsonima moritziana</i>	548	<i>Cabralea sulcata</i>	572
<i>Byrsonima obversa</i>	548	<i>Cabralea villosa</i>	572
<i>Byrsonima ophiticola</i>	549	<i>Cabralea warmingiana</i>	572
<i>Byrsonima panamensis</i>	548	<i>Cabralea warmingiana</i> var. <i>coriacea</i>	572
<i>Byrsonima peruviana</i>	549	<i>Cacao guianensis</i>	559
<i>Byrsonima peruviana</i> var. <i>eglandulosa</i>	549	<i>Cacao minus</i>	558
<i>Byrsonima propinqua</i>	549	<i>Cacao sativa</i>	558
<i>Byrsonima pulchra</i>	548		
<i>Byrsonima rufescens</i>	548		

Appendix 3. — Continuation.

<i>Cacao sylvestris</i>	559	<i>Calycolpus glaber</i> var. <i>angustilobus</i>	597
<i>Cacao theobroma</i>	558	<i>Calycolpus goetheanus</i>	597
<i>Caesalpinioideae</i>	503	<i>Calycolpus gracilis</i>	597
<i>Cailliea macrostachya</i>	535	<i>Calycolpus kegelianus</i>	597
<i>Calanthea pulcherrima</i>	411	<i>Calycolpus kegelianus</i> var. <i>gracilis</i>	597
<i>Calappa nucifera</i>	392	<i>Calycolpus kegelianus</i> var. <i>robustus</i>	597
<i>Calappa speciosa</i>	395	<i>Calycolpus megalodon</i>	597
<i>Caleatia caimito</i>	694	<i>Calycolpus ovalifolius</i>	597
<i>Calliandra</i>	503, 507	<i>Calycolpus pyrifera</i>	597
<i>Calliandra coriacea</i>	525	<i>Calycolpus revolutus</i>	597
<i>Calliandra hymenaeodes</i>	507	<i>Calycolpus schomburgkianus</i>	597
<i>Calliandra latifolia</i>	545	<i>Calycolpus schomburgkianus</i> var. <i>recurvatus</i>	597
<i>Calliandra patrisii</i>	507	<i>Calycolpus schomburgkianus</i> var. <i>speciosus</i>	597
<i>Calliandra schwackeana</i>	545	<i>Calycorrectes</i>	598, 602
<i>Calliandra surinamensis</i>	507	<i>Calycorrectes bergii</i>	598
<i>Calliandra tenuiflora</i>	507	<i>Calycorrectes grandifolius</i>	605
<i>Calliandra trinervia</i>	507	<i>Calycorrectes guyanensis</i>	602
<i>Calliandra</i> sp. A	508	<i>Calycorrectes latifolius</i>	598, 603
<i>Callicarpa discolor</i>	472	<i>Calycorrectes macrocalyx</i>	607
<i>Callicarpa globiflora</i>	472	<i>Calycorrectes protractus</i>	626
<i>Callicarpa integrifolia</i>	472	<i>Calyptranthes</i>	612
<i>Calocarpum buchananiifolium</i>	700	<i>Calyptranthes amshoffae</i>	610
<i>Calocarpum cooperi</i>	702	<i>Calyptranthes eugenoides</i>	626
<i>Calocarpum procerum</i>	700	<i>Calyptranthes fasciculata</i>	612
<i>Calocarpum procerum</i> var. <i>cuspidatum</i>	700	<i>Calyptranthes fasciculata</i> var. <i>genuina</i>	612
<i>Calocarpum procerum</i> var. <i>oblongum</i>	700	<i>Calyptranthes fasciculata</i> var. <i>hahnii</i>	612
CALOPHYLLACEAE	407	<i>Calyptranthes floribunda</i>	625
<i>Calophyllum antillanum</i>	407	<i>Calyptranthes forsteri</i>	612
<i>Calophyllum brasiliense</i>	407	<i>Calyptranthes lucida</i>	617
<i>Calophyllum brasiliense</i> subsp. <i>longifolium</i>	407	<i>Calyptranthes obtusa</i>	617
<i>Calophyllum brasiliense</i> subsp. <i>mariae</i>	407	<i>Calyptranthes pohliana</i>	617
<i>Calophyllum brasiliense</i> subsp. <i>verum</i>	407	<i>Calyptranthes pulchella</i>	618
<i>Calophyllum brasiliense</i> subsp. <i>wrightii</i>	407	<i>Calyptranthes pulchella</i> var. <i>cuneata</i>	618
<i>Calophyllum brasiliense</i> var. <i>antillanum</i>	407	<i>Calyptranthes pulchella</i> var. <i>grandiflora</i>	618
<i>Calophyllum brasiliense</i> var. <i>burchellii</i>	407	<i>Calyptranthes pulchella</i> var. <i>latifolia</i>	618
<i>Calophyllum brasiliense</i> var. <i>elongatum</i>	407	<i>Calyptranthes pulchella</i> var. <i>parviflora</i>	618
<i>Calophyllum brasiliense</i> var. <i>gardneri</i>	407	<i>Calyptranthes pullei</i>	618
<i>Calophyllum brasiliense</i> var. <i>genuina</i>	407	<i>Calyptranthes pullei</i> var. <i>immaculata</i>	618
<i>Calophyllum brasiliense</i> var. <i>rekoi</i>	407	<i>Calyptranthes sericea</i>	612
<i>Calophyllum brasiliense</i> var. <i>spruceanum</i>	407	<i>Calyptranthes sericea</i> var. <i>hahnii</i>	612
<i>Calophyllum calaba</i>	407	<i>Calyptranthes speciosa</i>	617
<i>Calophyllum chiapense</i>	407	<i>Calyptranthes tobagensis</i>	611
<i>Calophyllum ellipticum</i>	407	<i>Calyptranthes tonduzii</i>	626
<i>Calophyllum jacquinii</i>	407	<i>Calyptromyrcia elegans</i>	613
<i>Calophyllum longifolium</i>	407	<i>Calyptromyrcia spixiana</i>	613
<i>Calophyllum lucidum</i>	407	<i>Calyptromyrcia venosa</i>	623
<i>Calophyllum madruno</i>	434	<i>Calyptropsidium sartorianum</i>	626
<i>Calophyllum piaroanum</i>	407	<i>Cameraria guianensis</i>	385
<i>Calophyllum rekoi</i>	407	<i>Cameraria lutea</i>	386
<i>Calophyllum revolutum</i>	407	<i>Cameraria tamaquarina</i>	386
<i>Calycolpus angustifolius</i>	597	<i>Camphoromoea surinamensis</i>	484
<i>Calycolpus chnoiophyllus</i>	597	<i>Camphoromoea tenuiflora</i>	484
<i>Calycolpus cordatus</i>	597	<i>Campomanesia aromatica</i>	597
<i>Calycolpus glaber</i>	597	<i>Campomanesia beaurepaireiana</i>	597

Appendix 3. — Continuation.

<i>Campomanesia ciliata</i>	597	<i>Capparis sola</i>	411
<i>Campomanesia coaetanea</i>	597	<i>Capparis sola</i> var. <i>longiracemosa</i>	411
<i>Campomanesia glabra</i>	597	<i>Capparis stenophylla</i>	411
<i>Campomanesia glazioviana</i>	597	<i>Capparis surinamensis</i>	412
<i>Campomanesia grandiflora</i>	597, 625	<i>Capparis ternata</i>	412
<i>Campomanesia poiteaui</i>	598	<i>Caraipa ampla</i>	407
<i>Campomanesia sparsiflora</i>	597	<i>Caraipa colombiana</i>	408
<i>Campomanesia synchrona</i>	597	<i>Caraipa densifolia</i>	408
<i>Campomanesia tenuifolia</i>	597	<i>Caraipa excelsa</i>	408
<i>Campomanesia tetramera</i>	607	<i>Caraipa fasciculata</i>	408
<i>Camptouratea leblondii</i>	630	<i>Caraipa fasciculata</i> var. <i>laxiflora</i>	408
<i>Camptouratea sagotii</i>	630	<i>Caraipa ferruginea</i>	408
<i>Cananga ouregou</i>	374	<i>Caraipa latifolia</i>	429
<i>Cananga schomburgkiana</i>	376	<i>Caraipa laurifolia</i>	408
<i>Candolleodendron</i>	503, 508	<i>Caraipa laxiflora</i>	408
<i>Candolleodendron brachystachyum</i>	508	<i>Caraipa longifolia</i>	426
CANELLACEAE	409	<i>Caraipa melhemiana</i>	408
CANNABACEAE	409	<i>Caraipa myrciifolia</i>	408
<i>Caopia acuminata</i>	470	<i>Caraipa parvifolia</i>	408
<i>Caopia cayennensis</i>	470	<i>Caraipa psidiifolia</i>	408
<i>Caopia gracilis</i>	470	<i>Caraipa punctulata</i>	408
<i>Caopia guianensis</i>	470	<i>Caraipa racemosa</i>	408
<i>Caopia latifolia</i>	470	<i>Caraipa richardiana</i>	408
<i>Caopia sessilifolia</i>	471	<i>Caraipa richardiana</i> var. <i>distorta</i>	408
<i>Caoutchoua guianensis</i>	462	<i>Caraipa variabilis</i>	407, 408
<i>Capirona decorticans</i>	651	<i>Caraipa variabilis</i> "var. β "	407
<i>Capirona duckei</i>	652	<i>Caraipa</i> sp. A	408
<i>Capirona huberiana</i>	652	<i>Caraipa</i> sp. B	408
<i>Capirona leiophloea</i>	652	<i>Carapa guianensis</i>	573
<i>Capirona macrophylla</i>	651	<i>Carapa latifolia</i>	573
<i>Capirona surinamensis</i>	652	<i>Carapa macrocarpa</i>	573
<i>Capirona wurdackii</i>	652	<i>Carapa sericea</i>	579
CAPPARACEAE	410	<i>Carapa surinamensis</i>	573
<i>Capparidastrum elegans</i>	411	<i>Cardiopetalum surinamense</i>	370
<i>Capparidastrum frondosum</i>	411	CARICACEAE	412
<i>Capparidastrum osmanthum</i>	411	<i>Carica dodecaphylla</i>	412
<i>Capparidastrum solum</i>	411	<i>Carica spinosa</i>	412
<i>Capparis</i>	410	<i>Cariniana paraensis</i>	493
<i>Capparis acutifolia</i>	411	<i>Carolinea insignis</i>	556
<i>Capparis crotonantha</i>	412	<i>Carolinea macrocarpa</i>	556
<i>Capparis elegans</i>	411	<i>Carolinea princeps</i>	556
<i>Capparis flexuosa</i> subsp. <i>polyantha</i>	412	<i>Carpiphea dentata</i>	444
<i>Capparis frondosa</i>	411	<i>Carpotroche crispidentata</i>	362
<i>Capparis guaguaensis</i>	411	<i>Carpotroche laxiflora</i>	363
<i>Capparis leprieurii</i>	412	<i>Carpotroche linguisfolia</i>	362
<i>Capparis maroniensis</i>	412	<i>Carpotroche longifolia</i>	362
<i>Capparis montana</i>	703	<i>Carpotroche longifolia</i> var. <i>heliocarpa</i>	362
<i>Capparis osmantha</i>	411	<i>Carpotroche longifolia</i> var. <i>phasmatocarpa</i>	362
<i>Capparis pittieri</i>	412	<i>Carpotroche paludosa</i>	363
<i>Capparis polyantha</i>	412	<i>Carpotroche surinamensis</i>	362
<i>Capparis pulcherrima</i>	411	CARYOCARACEAE	413
<i>Capparis radiatiflora</i>	412	<i>Caryocar butyrosum</i>	413
<i>Capparis schunkei</i>	412	<i>Caryocar coccineum</i>	413
<i>Capparis sect. Eucapparis</i>	410	<i>Caryocar edule</i>	413

Appendix 3. — Continuation.

<i>Caryocar glabrum</i>	413	<i>Casearia glaberrima</i>	672
<i>Caryocar glabrum</i> var. <i>edule</i>	413	<i>Casearia glaziovii</i>	672
<i>Caryocar glabrum</i> var. <i>pilosum</i>	413	<i>Casearia grandiflora</i>	673
<i>Caryocar microcarpum</i>	413	<i>Casearia grandiflora</i> var. <i>hypoleuca</i>	673
<i>Caryocar riparium</i>	413	<i>Casearia grandiflora</i> var. <i>obtusifolia</i>	673
<i>Caryocar tessmannii</i>	413	<i>Casearia grandiflora</i> var. <i>pauciflora</i>	673
<i>Caryocar toxiferum</i>	413	<i>Casearia guianensis</i>	673
<i>Caryocar villosum</i>	413	<i>Casearia herbert-smithii</i>	675
<i>Caryocar villosum</i> var. <i>aesculifolium</i>	413	<i>Casearia hostmanniana</i>	672
<i>Caryocar villosum</i> var. <i>macrophyllum</i>	413	<i>Casearia hypoleuca</i>	673
<i>Caryodendron amazonicum</i>	459	<i>Casearia incana</i>	672
<i>Caryophyllus floribundus</i>	625	<i>Casearia iquitosensis</i>	678
<i>Caryophyllus fruticosus</i>	598	<i>Casearia javitensis</i>	678
<i>Casarettoa diversifolia</i>	474	<i>Casearia javitensis</i> var. <i>myriantha</i>	678
<i>Cascarilla lambertiana</i>	661	<i>Casearia laevigata</i>	676
<i>Cascarilla schomburgkii</i>	661	<i>Casearia lanceolata</i>	672
<i>Casearia</i>	672	<i>Casearia lasiosperma</i>	678
<i>Casearia acuminata</i>	676	<i>Casearia laurifolia</i>	678
<i>Casearia adamantium</i>	673	<i>Casearia lindeniana</i>	675
<i>Casearia adstringens</i>	673	<i>Casearia macrophylla</i>	674
<i>Casearia affinis</i>	675	<i>Casearia macrophylla</i> var. <i>barbatula</i>	674
<i>Casearia albicaulis</i>	673	<i>Casearia mariquitensis</i>	673
<i>Casearia anavinga</i>	673	<i>Casearia matthewsii</i>	675
<i>Casearia arborea</i>	672	<i>Casearia maximilianii</i>	678
<i>Casearia attenuata</i>	675	<i>Casearia microphylla</i>	674, 675
<i>Casearia bangii</i>	672	<i>Casearia miradorensis</i>	678
<i>Casearia belizensis</i>	672	<i>Casearia myriantha</i>	678
<i>Casearia benthamiana</i>	675	<i>Casearia negrensis</i>	674
<i>Casearia bicolor</i>	672	<i>Casearia nitida</i>	673
<i>Casearia blanchetiana</i>	676	<i>Casearia oligantha</i>	672
<i>Casearia brasiliensis</i>	672	<i>Casearia onacaensis</i>	675
<i>Casearia brighamii</i>	678	<i>Casearia ovoidea</i>	676
<i>Casearia cambessedesii</i>	672, 676	<i>Casearia parviflora</i>	672, 673, 675
<i>Casearia cambessedesii</i> var. <i>angustifolia</i>	672	<i>Casearia parviflora</i> var. <i>microphylla</i>	675
<i>Casearia cambessedesii</i> var. <i>parvifolia</i>	672	<i>Casearia parvifolia</i>	672, 673
<i>Casearia camporum</i>	673	<i>Casearia parvifolia</i> var. <i>microcarpa</i>	673
<i>Casearia capitata</i>	672	<i>Casearia parvifolia</i> var. <i>paraguariensis</i>	673
<i>Casearia caudata</i>	675	<i>Casearia pavoniana</i>	673
<i>Casearia celastroides</i>	676	<i>Casearia petraea</i>	676
<i>Casearia celtidifolia</i>	676	<i>Casearia piparea</i>	678
<i>Casearia chlorophoroidea</i>	675	<i>Casearia pitumba</i>	674
<i>Casearia commersoniana</i>	678	<i>Casearia platyphylla</i>	673
<i>Casearia congestiflora</i>	678	<i>Casearia poeppigii</i>	672
<i>Casearia cotticensis</i>	676	<i>Casearia prunifolia</i>	675
<i>Casearia decandra</i>	672, 673	<i>Casearia pubiflora</i>	673
<i>Casearia densiflora</i>	678	<i>Casearia punctata</i>	675
<i>Casearia densiflora</i> var. <i>parviflora</i>	678	<i>Casearia ramiflora</i>	673
<i>Casearia dentata</i>	678	<i>Casearia reflexa</i>	673
<i>Casearia ekmanii</i>	676	<i>Casearia reginae</i>	676
<i>Casearia fallax</i>	673	<i>Casearia rusbyana</i>	675
<i>Casearia floribunda</i>	673	<i>Casearia samyda</i>	675
<i>Casearia fockeana</i>	673	<i>Casearia schulziana</i>	675
<i>Casearia formosa</i>	675	<i>Casearia serrata</i>	672
<i>Casearia gaertneri</i>	678	<i>Casearia serrulata</i>	673, 675

Appendix 3. — Continuation.

<i>Casearia singularis</i>	675	<i>Cassipourea broadwayi</i>	649
<i>Casearia stipularis</i>	672	<i>Cassipourea cubensis</i>	649
<i>Casearia subsessiliflora</i>	676	<i>Cassipourea elliptica</i>	648, 649
<i>Casearia sylvestris</i>	675, 676	<i>Cassipourea elliptica</i> var. <i>alba</i>	649
<i>Casearia sylvestris</i> var. <i>angustifolia</i>	675	<i>Cassipourea elliptica</i> var. <i>latifolia</i>	649
<i>Casearia sylvestris</i> var. <i>benthamiana</i>	675	<i>Cassipourea elliptica</i> var. <i>ovata</i>	649
<i>Casearia sylvestris</i> var. <i>chlorophoroidea</i>	675	<i>Cassipourea elliptica</i> var. <i>parvifolia</i>	649
<i>Casearia sylvestris</i> var. <i>eichleri</i>	675	<i>Cassipourea elliptica</i> var. <i>pauciserrata</i>	649
<i>Casearia sylvestris</i> var. <i>martinicensis</i>	676	<i>Cassipourea elliptica</i> var. <i>typica</i>	648
<i>Casearia sylvestris</i> var. <i>myricoides</i>	675	<i>Cassipourea guianensis</i>	648, 649
<i>Casearia sylvestris</i> var. <i>paraensis</i>	675	<i>Cassipourea guianensis</i> var. <i>elliptica</i>	648
<i>Casearia sylvestris</i> var. <i>platyphylla</i>	675	<i>Cassipourea guianensis</i> var. <i>serrata</i>	649
<i>Casearia sylvestris</i> var. <i>tomentella</i>	675	<i>Cassipourea guianensis</i> var. <i>trichopoda</i>	649
<i>Casearia sylvestris</i> var. <i>wydleri</i>	675	<i>Cassipourea guildingii</i>	649
<i>Casearia tarapotina</i>	673	<i>Cassipourea latifolia</i>	649
<i>Casearia timbuchi</i>	674	<i>Cassipourea macrodonta</i>	649
<i>Casearia tremifolia</i>	676	<i>Cassipourea macrophylla</i>	649
<i>Casearia ulmifolia</i>	676	<i>Cassipourea podantha</i>	649
<i>Casearia umbellifera</i>	672	<i>Cassipourea quadrilocularis</i>	649
<i>Casearia zizyphoides</i>	676	<i>Cassipourea serrata</i>	649
<i>Casearia</i> sp. A	676	<i>Cassupa scarlatina</i>	660
<i>Casearia</i> sp. B	676	<i>Cassuvium pomiferum</i>	363
<i>Casearia</i> sp. C	676	<i>Cassuvium reniforme</i>	363
<i>Casearia</i> sp. D	676	<i>Cassuvium solitarium</i>	363
<i>Casinga procera</i>	672	<i>Castilla australis</i>	593
<i>Cassia</i>	503, 508	<i>Catappa guianensis</i>	441
<i>Cassia acuminata</i>	509	<i>Cathartocarpus brasiliensis</i>	508
<i>Cassia ampliflora</i>	537	<i>Cathartocarpus erubescens</i>	508
<i>Cassia annunciatia</i>	538	<i>Cathartocarpus grandis</i>	508
<i>Cassia apoucouita</i>	509	<i>Cathedra acuminata</i>	633
<i>Cassia brasiliensis</i>	508	<i>Cathedra aestuaria</i>	633
<i>Cassia brasiliensis</i> var. <i>tomentosa</i>	508	<i>Cathedra caurensis</i>	633
<i>Cassia calliantha</i>	537	<i>Cathedra crassifolia</i>	633
<i>Cassia chamaecrista</i>	722	<i>Cathedra guianensis</i>	633
<i>Cassia cowanii</i>	508	<i>Cathedra inaequilatera</i>	633
<i>Cassia dumetorum</i>	538	<i>Cathedra oblonga</i>	633
<i>Cassia fastuosa</i>	508	<i>Cathormion corymbosum</i>	515
<i>Cassia fulgens</i>	537	<i>Catinga aromatica</i>	604
<i>Cassia grandis</i>	508	<i>Catinga moschata</i>	598, 604
<i>Cassia mollis</i>	508	<i>Catinga oblongifolia</i>	598
<i>Cassia multijuga</i>	537	<i>Catis martiana</i>	393
<i>Cassia pachycarpa</i>	508	<i>Catostemma commune</i>	552
<i>Cassia regia</i>	508	<i>Catostemma fragrans</i>	552
<i>Cassia reticulata</i>	538	<i>Catostemma grazielae</i>	552
<i>Cassia richardiana</i>	537	<i>Catostemma macrospermum</i>	552
<i>Cassia sagotiana</i>	508	<i>Catostemma</i> sp. A	552
<i>Cassia solimoesensis</i>	509	<i>Cecropia arachnoidea</i>	712
<i>Cassia spruceana</i>	508	<i>Cecropia asperrima</i>	712
<i>Cassia strobilacea</i>	538	<i>Cecropia bureauiana</i>	711
<i>Cassia tarantan</i>	538	<i>Cecropia dielsiana</i>	712
<i>Cassipourea alba</i>	649	<i>Cecropia digitata</i> var. <i>grisea</i>	711
<i>Cassipourea belizensis</i>	649	<i>Cecropia distachya</i>	711
<i>Cassipourea brittoniana</i>	649	<i>Cecropia goodspeedii</i>	712
		<i>Cecropia granvilleana</i>	711

Appendix 3. — Continuation.

<i>Cecropia hondurensis</i>	712	<i>Cedrela mourae</i>	575
<i>Cecropia humboldtiana</i>	711	<i>Cedrela occidentalis</i>	574
<i>Cecropia inchuensis</i>	712	<i>Cedrela odorata</i>	574, 575
<i>Cecropia juranyiana</i>	712, 713	<i>Cedrela odorata</i> var. <i>xerogeiton</i>	575
<i>Cecropia latiloba</i>	711	<i>Cedrela pachyrhachis</i>	574
<i>Cecropia lisboana</i>	711	<i>Cedrela palustris</i>	575
<i>Cecropia obtusa</i>	711	<i>Cedrela paraguariensis</i>	574
<i>Cecropia orinocensis</i>	711	<i>Cedrela paraguariensis</i> var. <i>brachystachya</i>	574
<i>Cecropia palmata</i>	711	<i>Cedrela paraguariensis</i> var. <i>hassleri</i>	574
<i>Cecropia paraensis</i>	711	<i>Cedrela paraguariensis</i> var. <i>multijuga</i>	574
<i>Cecropia peltata</i>	711	<i>Cedrela pilgeri</i>	574
<i>Cecropia propinqua</i>	712	<i>Cedrela regnellii</i>	574
<i>Cecropia richardii</i>	711	<i>Cedrela rotunda</i>	575
<i>Cecropia riparia</i>	711	<i>Cedrela sintenisii</i>	575
<i>Cecropia scabrifolia</i>	712	<i>Cedrela tubiflora</i>	574
<i>Cecropia schiedeana</i>	711	<i>Cedrela tubiflora</i> f. <i>angustifolia</i>	574
<i>Cecropia sciadophylla</i>	712	<i>Cedrela tubiflora</i> f. <i>grandifolia</i>	574
<i>Cecropia sciadophylla</i> var. <i>decurrens</i>	712	<i>Cedrela tubiflora</i> f. <i>parvifoliola</i>	574
<i>Cecropia sciadophylla</i> var. <i>guamuesensis</i>	712	<i>Cedrela tubiflora</i> subsp. <i>bertoniensis</i>	574
<i>Cecropia sciadophylla</i> var. <i>juranyiana</i>	712	<i>Cedrela tubiflora</i> var. <i>grandifolia</i>	574
<i>Cecropia sciadophylla</i> var. <i>pedroa</i>	712	<i>Cedrela tubiflora</i> var. <i>intermedia</i>	574
<i>Cecropia sciadophylla</i> var. <i>subsessilis</i>	712	<i>Cedrela tubiflora</i> var. <i>lagenaria</i>	574
<i>Cecropia silvae</i>	713	<i>Cedrela velloziana</i>	574
<i>Cecropia surinamensis</i>	711	<i>Cedrela whitfordii</i>	575
<i>Cedrela adenophylla</i>	574	<i>Cedrela yucatana</i>	575
<i>Cedrela alliacea</i>	574	<i>Cedrelinga</i>	503, 508
<i>Cedrela alternifolia</i>	555	<i>Cedrelinga cateniformis</i>	508
<i>Cedrela balansae</i>	574	<i>Cedrota guianensis</i>	476
<i>Cedrela barbata</i>	574	<i>Cedrota longifolia</i>	476
<i>Cedrela brachystachya</i>	574	<i>Cedrus alternifolia</i>	555
<i>Cedrela brasiliensis</i>	574	<i>Ceiba anfractuosa</i>	552
<i>Cedrela brasiliensis</i> var. <i>australis</i>	574	<i>Ceiba caribaea</i>	552
<i>Cedrela brownii</i>	574	<i>Ceiba casearia</i>	552
<i>Cedrela brunellioides</i>	574	<i>Ceiba guineensis</i>	552, 553
<i>Cedrela caldasana</i>	574	<i>Ceiba guineensis</i> var. <i>ampla</i>	553
<i>Cedrela ciliolata</i>	575	<i>Ceiba guineensis</i> var. <i>clausa</i>	552
<i>Cedrela cubensis</i>	575	<i>Ceiba occidentalis</i>	552
<i>Cedrela dugesii</i>	574	<i>Ceiba pentandra</i>	552, 553
<i>Cedrela elliptica</i>	574	<i>Ceiba pentandra</i> f. <i>albolana</i>	552
<i>Cedrela fissilis</i>	574	<i>Ceiba pentandra</i> f. <i>grisea</i>	552
<i>Cedrela fissilis</i> var. <i>glabrior</i>	574	<i>Ceiba pentandra</i> var. <i>caribaea</i>	552
<i>Cedrela fissilis</i> var. <i>macrocarpa</i>	574	<i>Ceiba pentandra</i> var. <i>clausa</i>	552
<i>Cedrela fuscata</i>	574	<i>Ceiba pentandra</i> var. <i>dehiscens</i>	553
<i>Cedrela glaziovii</i>	574	<i>Ceiba pentandra</i> var. <i>indica</i>	552
<i>Cedrela guianensis</i>	574, 575	<i>Ceiba thonningii</i>	553
<i>Cedrela hassleri</i>	574	CELASTRACEAE	413
<i>Cedrela hirsuta</i>	574	<i>Celastrus myrtifolius</i>	650
<i>Cedrela huberi</i>	574	<i>Celtis albicans</i>	409
<i>Cedrela longiflora</i>	574	<i>Celtis canescens</i>	409
<i>Cedrela longipes</i>	575	<i>Celtis chichilea</i>	409
<i>Cedrela longipetiolulata</i>	575	<i>Celtis curiandiuba</i>	409
<i>Cedrela macrocarpa</i>	574	<i>Celtis lima</i>	409
<i>Cedrela mexicana</i>	574	<i>Celtis macrophylla</i>	409
<i>Cedrela mexicana</i> var. <i>puberula</i>	574	<i>Celtis micrantha</i>	409

Appendix 3. — Continuation.

<i>Celtis microcarpa</i>	409	<i>Chrysobalanus icaco</i> subsp. <i>ellipticus</i>	416
<i>Celtis mollis</i>	409	<i>Chrysobalanus icaco</i> subsp. <i>orbicularis</i>	416
<i>Celtis riparia</i>	409	<i>Chrysobalanus icaco</i> subsp. <i>pellocarpus</i>	416
<i>Celtis rufescens</i>	409	<i>Chrysobalanus icaco</i> var. <i>ellipticus</i>	416
<i>Celtis rugosa</i>	409	<i>Chrysobalanus icaco</i> var. <i>genuinus</i>	417
<i>Celtis schiedeana</i>	409	<i>Chrysobalanus icaco</i> var. <i>luteus</i>	416
<i>Cerbera triphylla</i>	382	<i>Chrysobalanus icaco</i> var. <i>macrocarpus</i>	417
<i>Cercidioideae</i>	503	<i>Chrysobalanus icaco</i> var. <i>pellocarpus</i>	416
<i>Cercophora anomala</i>	500	<i>Chrysobalanus icaco</i> var. <i>roseus</i>	417
<i>Cercouratea melinonii</i>	630	<i>Chrysobalanus incanus</i>	428
<i>Cerdana alliodora</i>	444	<i>Chrysobalanus interior</i>	417
<i>Chaenopleura ferruginea</i>	564	<i>Chrysobalanus luteus</i>	416
<i>Chaenopleura hypoleuca</i>	564	<i>Chrysobalanus montanus</i>	431
<i>Chaenopleura longifolia</i>	565	<i>Chrysobalanus orbicularis</i>	416
<i>Chaetocarpus schomburgkianus</i>	635	<i>Chrysobalanus pellocarpus</i>	416
<i>Chaetocarpus williamsii</i>	636	<i>Chrysobalanus purpureus</i>	416
<i>Chaetocarpus</i> sp. A	636	<i>Chrysobalanus savannarum</i>	416
<i>Chaetocrater capitatum</i>	672	<i>Chrysobalanus triandra</i>	422
<i>Chaetocrater javitensis</i>	678	<i>Chrysochlamys membranacea</i>	432
<i>Chaetocrater reflexus</i>	673	<i>Chrysophyllum abbreviatum</i>	689
<i>Chailletia capitulifera</i>	448	<i>Chrysophyllum alnifolium</i>	697
<i>Chailletia sessiliflora</i>	448	<i>Chrysophyllum ambelaniifolium</i>	693
<i>Chamaecrista</i>	503, 509	<i>Chrysophyllum anomalum</i>	693
<i>Chamaecrista apoucouita</i>	509	<i>Chrysophyllum argenteum</i> subsp. <i>auratum</i>	685
<i>Chamaesenna multijuga</i>	537	<i>Chrysophyllum argenteum</i> subsp. <i>nitidum</i>	686
<i>Chamaesenna reticulata</i>	538	<i>Chrysophyllum auratum</i>	685, 686
<i>Chaunochiton breviflorum</i>	633	<i>Chrysophyllum auratum</i> var. <i>majus</i>	686
<i>Chaunochiton kappleri</i>	633	<i>Chrysophyllum brasiliense</i>	704
<i>Chaunochiton purpurascens</i>	633	<i>Chrysophyllum brasiliense</i> var. <i>minus</i>	704
<i>Cheiloclinium cognatum</i>	413	<i>Chrysophyllum cayennense</i>	695
<i>Cheiloclinium lineolatum</i>	413	<i>Chrysophyllum cochlearium</i>	703
<i>Cheiloclinium neglectum</i>	414	<i>Chrysophyllum cuneifolium</i>	686
<i>Chimarrhis clausicorollata</i>	666	<i>Chrysophyllum cuspidatum</i>	688
<i>Chimarrhis duckei</i>	652	<i>Chrysophyllum cyanogenum</i>	687
<i>Chimarrhis longistipulata</i>	652	<i>Chrysophyllum discolor</i>	692
<i>Chimarrhis microcarpa</i>	652	<i>Chrysophyllum durifruitum</i>	686
<i>Chimarrhis turbinata</i>	652	<i>Chrysophyllum excelsum</i>	687
<i>Chionanthus guianensis</i>	635	<i>Chrysophyllum eximium</i>	686
<i>Chionanthus mayepea</i>	635	<i>Chrysophyllum guianense</i>	688
<i>Chionanthus tetrandrus</i>	635	<i>Chrysophyllum guyanense</i>	686
<i>Chione chambersii</i>	666	<i>Chrysophyllum hostmannianum</i>	686
<i>Chitonia fothbergilla</i>	565	<i>Chrysophyllum lucentifolium</i> subsp. <i>pachycarpum</i>	686
<i>Chitonia macrophylla</i>	568	<i>Chrysophyllum macrophyllum</i>	699, 704
<i>Chloroleucon</i>	503, 509	<i>Chrysophyllum manaosense</i>	686
<i>Chloroleucon acaciaoides</i>	509	<i>Chrysophyllum melinonii</i>	691
<i>Chloromyron verticillatum</i>	434	<i>Chrysophyllum minutiflorum</i>	695
<i>Chomelia chambersii</i>	666	<i>Chrysophyllum nitidissimum</i>	686
<i>Chomelia panamensis</i>	666	<i>Chrysophyllum nitidum</i>	686
<i>Chromolucuma congestifolia</i>	685	<i>Chrysophyllum pomiferum</i>	687
CHRYSOBALANACEAE	416	<i>Chrysophyllum prieurii</i>	687
<i>Chrysobalanus ellipticus</i>	416	<i>Chrysophyllum psilosphyllum</i>	686
<i>Chrysobalanus guianensis</i>	416	<i>Chrysophyllum ramiflorum</i>	689
<i>Chrysobalanus icaco</i>	416, 417	<i>Chrysophyllum reticulatum</i>	700
<i>Chrysobalanus icaco</i> f. <i>albus</i>	417	<i>Chrysophyllum richardii</i>	685

Appendix 3. — Continuation.

<i>Chrysophyllum rufocupreum</i>	686	<i>Chytroma salebrosa</i>	501
<i>Chrysophyllum sanguinolentum</i>	687	<i>Chytroma simiorum</i>	502
<i>Chrysophyllum sericeum</i>	685, 686	<i>Chytroma simiorum</i> var. <i>latifolia</i>	502
<i>Chrysophyllum sericeum</i> var. <i>acutifolium</i>	686	<i>Chytroma spruceana</i>	500
<i>Chrysophyllum sericeum</i> var. <i>obtusifolium</i>	686	<i>Chytroma urceolata</i>	500
<i>Chrysophyllum sessiliflorum</i>	695	<i>Cicca antillana</i>	638
<i>Chrysophyllum sparsiflorum</i>	687	<i>Cicca antillana</i> var. <i>pedicellaris</i>	638
<i>Chrysophyllum sparsiflorum</i> var. <i>fagifolium</i>	687	<i>Cicca pavoniana</i>	638
<i>Chrysophyllum steyermarkii</i>	687	<i>Cicca surinamensis</i>	638
<i>Chrysophyllum venezuelanense</i>	687	<i>Cinchona china</i>	653
<i>Chrysophyllum wurdackii</i>	695	<i>Cinchona grandiflora</i>	653
<i>Chrysophyllum</i> sp. A.	687	<i>Cinchona lambertiana</i>	661
<i>Chrysophyllum</i> sp. B.	687	<i>Cinchona longiflora</i>	653
<i>Chrysophyllum</i> sp. C	688	<i>Cinchona obtusifolia</i>	653
<i>Chrysophyllum</i> sp. D	688	<i>Cinchona quina</i>	653
<i>Chrysophyllum</i> sp. E	688	<i>Cinchona souzana</i>	655
<i>Chrysophyllum</i> sp. F	688	<i>Cinnamodendron tenuifolium</i>	409
<i>Chuncoa amazonia</i>	440	<i>Cinnamomum australe</i>	475
<i>Chuncoa diptera</i>	443	<i>Cinnamomum bahiense</i>	485
<i>Chuncoa oblonga</i>	443	<i>Cinnamomum brasiliense</i>	475
<i>Chuncoa obovata</i>	440	<i>Cinnamomum chana</i>	475
<i>Chytraculia eugeniooides</i>	626	<i>Cinnamomum cinnamomifolium</i>	475
<i>Chytraculia fasciculata</i>	612	<i>Cinnamomum cubense</i>	475
<i>Chytraculia forsteri</i>	612	<i>Cinnamomum elongatum</i>	475
<i>Chytraculia gardneriana</i>	626	<i>Cinnamomum filamentosum</i>	475
<i>Chytraculia lucida</i>	617	<i>Cinnamomum fruticosum</i>	475
<i>Chytraculia pohliana</i>	617	<i>Cinnamomum grisebachii</i>	474
<i>Chytraculia pulchella</i>	618	<i>Cinnamomum heterotepalum</i>	475
<i>Chytraculia sartoriana</i>	626	<i>Cinnamomum impressum</i>	474
<i>Chytroma amara</i>	501	<i>Cinnamomum johnstonii</i>	475
<i>Chytroma apiculata</i>	494	<i>Cinnamomum maynense</i>	475
<i>Chytroma basilaris</i>	501	<i>Cinnamomum mexicanum</i>	475
<i>Chytroma chartacea</i>	500	<i>Cinnamomum montanum</i>	474
<i>Chytroma cincturata</i>	495	<i>Cinnamomum paraguariense</i>	475
<i>Chytroma cistella</i>	500	<i>Cinnamomum peruvianum</i>	475
<i>Chytroma collina</i>	495	<i>Cinnamomum phoebe</i>	475
<i>Chytroma congestiflora</i>	501	<i>Cinnamomum pichisense</i>	475
<i>Chytroma corrugata</i>	501	<i>Cinnamomum pickellii</i>	475
<i>Chytroma crenata</i>	502	<i>Cinnamomum portosecurianum</i>	475
<i>Chytroma cupularis</i>	500	<i>Cinnamomum triplinerve</i>	474
<i>Chytroma decolorans</i>	495	<i>Cinnamomum xinguense</i>	475
<i>Chytroma foetida</i>	502	<i>Ciponima guianensis</i>	709
<i>Chytroma grandifolia</i>	495	<i>Ciponima scabridula</i>	709
<i>Chytroma holcogyne</i>	501	<i>Citharexylum amazonicum</i>	717
<i>Chytroma idatimon</i>	501	<i>Citharexylum bahamense</i>	718
<i>Chytroma labriculata</i>	493	<i>Citharexylum broadwayi</i>	718
<i>Chytroma laevicula</i>	500	<i>Citharexylum caudatum</i>	718
<i>Chytroma langsdorffii</i>	500	<i>Citharexylum cinereum</i>	717, 718
<i>Chytroma marawynensis</i>	500	<i>Citharexylum coriaceum</i>	718
<i>Chytroma monosperma</i>	500	<i>Citharexylum fruticosum</i>	717, 718
<i>Chytroma perspicua</i>	497	<i>Citharexylum fruticosum</i> f. <i>bahamense</i>	718
<i>Chytroma pilacarpa</i>	500	<i>Citharexylum fruticosum</i> f. <i>subserratum</i>	718
<i>Chytroma rorida</i>	500	<i>Citharexylum fruticosum</i> f. <i>subvillosum</i>	718
<i>Chytroma rubriflora</i>	501	<i>Citharexylum fruticosum</i> var. <i>brittonii</i>	718

Appendix 3. — Continuation.

<i>Citharexylum fruticosum</i> var. <i>smallii</i>	718	<i>Cleidion denticulatum</i>	457
<i>Citharexylum fruticosum</i> var. <i>subserratum</i>	718	<i>Cleome arborea</i>	411
<i>Citharexylum fruticosum</i> var. <i>subvillosum</i>	718	<i>Clidemia trichodes</i>	569
<i>Citharexylum fruticosum</i> var. <i>villosum</i>	718	<i>Clompanus frondosa</i>	557
<i>Citharexylum hybridum</i>	718	<i>Clompanus pruriens</i>	557
<i>Citharexylum laevigatum</i>	718	<i>Clompanus speciosa</i>	558
<i>Citharexylum macrophyllum</i>	717	<i>Clusia cartilaginosa</i>	433
<i>Citharexylum molle</i>	718	<i>CLUSIACEAE</i>	432
<i>Citharexylum pentandrum</i>	718	<i>Clusia colorans</i>	434
<i>Citharexylum polystachyum</i>	718	<i>Clusia couletii</i>	433
<i>Citharexylum pulverulentum</i>	718	<i>Clusia cuneata</i>	432
<i>Citharexylum quadrangulare</i>	718	<i>Clusia flavida</i>	432
<i>Citharexylum spinosum</i>	717, 718	<i>Clusia fockeana</i>	432
<i>Citharexylum spinosum</i> f. <i>brittonii</i>	718	<i>Clusia galactodendron</i>	433
<i>Citharexylum spinosum</i> f. <i>smallii</i>	718	<i>Clusia gentlei</i>	432
<i>Citharexylum spinosum</i> f. <i>subserratum</i>	718	<i>Clusia grandiflora</i>	432
<i>Citharexylum spinosum</i> f. <i>subvillosum</i>	718	<i>Clusia leprantha</i>	433
<i>Citharexylum spinosum</i> f. <i>villosum</i>	718	<i>Clusia lhotzkyana</i>	433
<i>Citharexylum subserratum</i>	718	<i>Clusia lhotzkyana</i> var. <i>polygama</i>	433
<i>Citharexylum surrectum</i>	718	<i>Clusia longifolia</i>	437
<i>Citharexylum teres</i>	718	<i>Clusia mammosa</i>	433
<i>Citharexylum tomentosum</i>	718	<i>Clusia martinii</i>	433
<i>Citharexylum villosum</i>	718	<i>Clusia maxima</i>	432
<i>Citrosma camporum</i>	707	<i>Clusia melchiorii</i>	433
<i>Citrosma cristata</i>	706	<i>Clusia microphylla</i>	434
<i>Citrosma cuspidata</i>	707	<i>Clusia minor</i>	433
<i>Citrosma decipiens</i>	707	<i>Clusia nemorosa</i>	433
<i>Citrosma discolor</i>	707	<i>Clusia nemorosa</i> var. <i>lhotzkyana</i>	433
<i>Citrosma guianensis</i>	707	<i>Clusia oblanceolata</i>	438
<i>Citrosma guianensis</i> var. <i>divergentifolia</i>	707	<i>Clusia octandra</i>	433
<i>Citrosma guianensis</i> var. <i>nuda</i>	707	<i>Clusia palmicida</i>	433, 434
<i>Citrosma myristicoides</i>	707	<i>Clusia panapanari</i>	434
<i>Citrosma poeppigii</i>	708	<i>Clusia parvicaudula</i>	433
<i>Clarisia ilicifolia</i>	585	<i>Clusia petiolata</i>	432
<i>Clarisia ilicifolia</i> var. <i>cuyunensis</i>	585	<i>Clusia pithecobia</i>	439
<i>Clarisia ilicifolia</i> var. <i>guianensis</i>	585	<i>Clusia pratensis</i>	433
<i>Clarisia ilicifolia</i> var. <i>micranthera</i>	585	<i>Clusia purpurea</i>	433
<i>Clarisia ilicifolia</i> var. <i>paraensis</i>	585	<i>Clusia scrobiculata</i>	434
<i>Clarisia nitida</i>	585	<i>Clusia stahelii</i>	432
<i>Clarisia racemosa</i>	585	<i>Clusia utilis</i>	433
<i>Clarisia spruceana</i>	585	<i>Clusia venosa</i>	433
<i>Clarisia strepitans</i>	585	<i>Coccoloba dugandiana</i>	643
<i>Clarisia strepitans</i> var. <i>cuyunensis</i>	585	<i>Coccoloba grandis</i>	643
<i>Clarisia strepitans</i> var. <i>guianensis</i>	585	<i>Coccoloba latifolia</i>	642
<i>Clarisia strepitans</i> var. <i>micranthera</i>	585	<i>Coccoloba mollis</i>	643
<i>Clarisia strepitans</i> var. <i>paraensis</i>	585	<i>Coccoloba paniculata</i>	643
<i>Clathrotropis</i>	503, 509	<i>Coccoloba polystachya</i>	643
<i>Clathrotropis brachypetala</i>	509	<i>Coccoloba polystachya</i> var. <i>glabra</i>	643
<i>Clathrotropis flava</i>	531	<i>Coccoloba polystachya</i> var. <i>jamaicensis</i>	643
<i>Clathrotropis surinamensis</i>	531	<i>Coccoloba polystachya</i> var. <i>mollis</i>	643
<i>Clavapetalum elatum</i>	582	<i>Coccoloba polystachya</i> var. <i>pubescens</i>	643
<i>Clavapetalum surinamense</i>	582	<i>Coccoloba rheifolia</i>	642
<i>Clavija septentrionalis</i>	464	<i>Coccoloba standleyana</i>	643
<i>Clavija sparsifolia</i>	719	<i>Coccoloba uvifera</i>	643

Appendix 3. — Continuation.

<i>Cochlospermum orinocense</i>	398	<i>Conomorpha ampla</i>	644
<i>Cochlospermum parkeri</i>	399	<i>Conomorpha macrophylla</i>	644
<i>Cochlospermum parvifolium</i>	399	<i>Conomorpha multipunctata</i>	644
<i>Cochlospermum wentii</i>	399	<i>Conomorpha utiarityi</i>	644
<i>Cochlospermum williamsii</i>	399	<i>Conostegia acuminata</i>	561
<i>Cocos aculeata</i>	390	<i>Conostegia parviflora</i>	567
<i>Cocos aequatorialis</i>	395	<i>Conuleum guyannense</i>	707
<i>Cocos chavesiana</i>	395	<i>Convolvulus pes-caprae</i>	722
<i>Cocos fusiformis</i>	390	<i>Copaiba guyanensis</i>	509
<i>Cocos inajai</i>	395	<i>Copaifera</i>	503, 509
<i>Cocos indica</i>	392	<i>Copaifera guyanensis</i>	509
<i>Cocos mamillaris</i>	392	<i>Copaifera</i> sp. A	509
<i>Cocos nana</i>	392	<i>Copedesma nitens</i>	568
<i>Cocos nucifera</i>	392	<i>Corallodendron fuscum</i>	515
<i>Cocos nucifera</i> var. <i>synphyllica</i>	392	<i>Corallodendron glaucum</i>	515
<i>Cocos speciosa</i>	395	<i>Corallodendron ovalifolia</i>	515
<i>Coeloclina lucida</i>	379	<i>Corallodendron patens</i>	515
<i>Coffea crassiloba</i>	665	<i>Cordia alliodora</i>	444
<i>Coffea occidentalis</i>	657	<i>Cordia belizensis</i>	444
<i>Coffea umbellata</i>	657	<i>Cordia bicolor</i>	444
<i>Coilotapalus obtusa</i>	711	<i>Cordia calophylla</i>	445
<i>Colicodendron obliquifolium</i>	411	<i>Cordia calyprata</i>	444
<i>Colicodendron pulchellum</i>	411	<i>Cordia carnosa</i>	444
<i>Colythrum grandiflorum</i>	667	<i>CORDIACEAE</i>	444
<i>COMBRETACEAE</i>	439	<i>Cordia collococa</i>	445
<i>Comocladia tapaculo</i>	365	<i>Cordia cordifolia</i>	446
<i>Compsoneura ulei</i>	594	<i>Cordia coriacea</i>	446
<i>Conceveiba guianensis</i>	459	<i>Cordia dentata</i>	444
<i>Conceveiba krukoffii</i>	459	<i>Cordia exaltata</i>	444
<i>Conceveiba macrophylla</i>	458	<i>Cordia exaltata</i> var. <i>melanoneura</i>	444
<i>Conceveiba martiana</i>	459	<i>Cordia expansa</i>	447
<i>Conceveiba megalophylla</i>	459	<i>Cordia fallax</i>	444
<i>Conceveiba simulata</i>	459	<i>Cordia formicarum</i>	445
<i>Conceveibastrum martianum</i>	459	<i>Cordia fulva</i>	444
<i>Conceveiba trigonocarpa</i>	459	<i>Cordia goeldiana</i>	445
<i>Conceveibum cordatum</i>	458	<i>Cordia guianensis</i>	444
<i>Conceveibum ovatum</i>	459	<i>Cordia hebbecarpa</i>	447
<i>Condaminea macrophylla</i>	651	<i>Cordia heterophylla</i>	447
<i>Conocarpus</i>	439, 440	<i>Cordia hirsuta</i>	445
<i>Conocarpus acutifolius</i>	439	<i>Cordia hirta</i>	446
<i>Conocarpus erectus</i>	439	<i>Cordia hispidissima</i>	445
<i>Conocarpus erectus</i> var. <i>arboreus</i>	439	<i>Cordia ierensis</i>	446
<i>Conocarpus erectus</i> var. <i>procumbens</i>	439	<i>Cordia laevifrons</i>	445
<i>Conocarpus procumbens</i>	439	<i>Cordia leptopoda</i>	444
<i>Conocarpus pubescens</i>	439	<i>Cordia lockhartii</i>	444
<i>Conocarpus racemosus</i>	440	<i>Cordia lomatoloba</i>	445
<i>Conocarpus supinus</i>	439	<i>Cordia melanoneura</i>	444
<i>Conohoria aubletii</i>	678	<i>Cordia miranda</i>	445
<i>Conohoria brevipes</i>	720	<i>Cordia muneco</i>	446
<i>Conohoria flavescens</i>	720	<i>Cordia naidophila</i>	445
<i>Conohoria laxiflora</i>	720	<i>Cordia nervosa</i>	445
<i>Conohoria passoura</i>	721	<i>Cordia nodosa</i>	445
<i>Conohoria riana</i>	721	<i>Cordia nodosa</i> var. <i>angustifolia</i>	445
<i>Conohoria rinorea</i>	720	<i>Cordia nodosa</i> var. <i>glabrior</i>	445

Appendix 3. — Continuation.

<i>Cordia nodosa</i> var. <i>hispidissima</i>	445	<i>Couepia habrantha</i>	418
<i>Cordia opaca</i>	446	<i>Couepia joaquinæ</i>	418
<i>Cordia ovata</i>	444	<i>Couepia leptostachya</i>	417
<i>Cordia panicularis</i>	445	<i>Couepia magnoliifolia</i>	418
<i>Cordia pubescens</i>	447	<i>Couepia martinii</i>	418
<i>Cordia sagotii</i>	446	<i>Couepia myrtifolia</i>	417
<i>Cordia scabrida</i>	444	<i>Couepia obovata</i>	419
<i>Cordia sericicalyx</i>	446	<i>Couepia parillo</i>	419
<i>Cordia sericicalyx</i> var. <i>latifolia</i>	446	<i>Couepia pauciflora</i>	419
<i>Cordia sprucei</i>	446	<i>Couepia rankiniae</i>	419
<i>Cordia tenuifolia</i>	444	<i>Couepia</i> sp. A	419
<i>Cordia tetrandra</i>	446	<i>Couepia surinamensis</i>	417
<i>Cordia tetraphylla</i>	443	<i>Couepia thyrsiflora</i>	418
<i>Cordia toqueve</i>	446	<i>Couepia versicolor</i>	417
<i>Cordia trichostyla</i>	444	<i>Couepia villosa</i>	419
<i>Cordia umbraculifera</i>	446	<i>Couma fasciculata</i>	386
<i>Cordia umbrosa</i>	445	<i>Couma guianensis</i>	382
<i>Cordia viridis</i>	446	<i>Couma pentaphylla</i>	386
<i>Cordia volubilis</i>	445	<i>Coumarouna coriacea</i>	543
<i>Cordiera edulis</i>	650	<i>Coumarouna nudipes</i>	543
<i>Cordiera latifolia</i>	650	<i>Coumarouna odorata</i>	513
<i>Cordiera myrciifolia</i>	652	<i>Coumarouna odorata</i> var. <i>tetraphylla</i>	513
<i>Cordiera oligantha</i>	650	<i>Coumarouna oppositifolia</i>	543
<i>Cordiera triflora</i>	652	<i>Coumarouna punctata</i>	513
<i>Cordiera trifolia</i>	652	<i>Coumarouna speciosa</i>	544
<i>Cornuella venezuelanensis</i>	687	<i>Coumarouna tetraphylla</i>	513
<i>Coryphadenia abnormis</i>	572	<i>Coumarouna trifoliolata</i>	513
<i>Corythophora amapaensis</i>	492	<i>Coupoïa aquatica</i>	655
<i>Corythophora labriculata</i>	493	<i>Coupoïa martiniana</i>	656
<i>Corythophora rimosa</i> subsp. <i>rimosa</i>	493	<i>Coupoïa micrantha</i>	656
<i>Corythophora rimosa</i> subsp. <i>rubra</i>	493	<i>Couralia fluviatilis</i>	398
<i>Cosmibuena arborea</i>	653	<i>Couratari braganceae</i>	493
<i>Cosmibuena gardenioides</i>	653	<i>Couratari calycina</i>	493
<i>Cosmibuena grandiflora</i>	653	<i>Couratari coriacea</i>	493
<i>Cosmibuena grandiflora</i> var. <i>latifolia</i>	653	<i>Couratari fagifolia</i>	493
<i>Cosmibuena latifolia</i>	653	<i>Couratari gloriosa</i>	493
<i>Cosmibuena obtusifolia</i>	653	<i>Couratari guianensis</i>	493
<i>Cosmibuena obtusifolia</i> var. <i>latifolia</i>	653	<i>Couratari multiflora</i>	493
<i>Cosmibuena ovalis</i>	653	<i>Couratari oblongifolia</i>	494
<i>Cosmibuena quinqueflora</i>	653	<i>Couratari oligantha</i>	494
<i>Cosmibuena skinneri</i>	653	<i>Couratari panamensis</i>	493
<i>Cosmibuena triflora</i>	653	<i>Couratari paraensis</i>	493
<i>Couepia bracteosa</i>	417	<i>Couratari pulchra</i>	493
<i>Couepia bracteosa</i> var. <i>grandifolia</i>	417	<i>Couratari reticulata</i>	494
<i>Couepia bracteosa</i> var. <i>minor</i>	417	<i>Couratari stellata</i>	494
<i>Couepia canomensis</i>	419	<i>Couratari vriesii</i>	493
<i>Couepia caryophylloides</i>	417	<i>Couroupita acreensis</i>	494
<i>Couepia cognata</i>	—	<i>Couroupita antillana</i>	494
<i>Couepia elata</i>	419	<i>Couroupita froesii</i>	494
<i>Couepia excelsa</i>	417	<i>Couroupita guianensis</i>	494
<i>Couepia exflexa</i>	417	<i>Couroupita guianensis</i> var. <i>surinamensis</i>	494
<i>Couepia glandulosa</i>	417	<i>Couroupita idolica</i>	494
<i>Couepia guianensis</i> subsp. <i>glandulosa</i>	417	<i>Couroupita membranacea</i>	494
<i>Couepia guianensis</i> subsp. <i>guianensis</i>	—	<i>Couroupita pedicularis</i>	494

Appendix 3. — Continuation.

<i>Couroupita peruviana</i>	494	<i>Coutarea lindeniana</i>	655
<i>Couroupita saintcroixiana</i>	494	<i>Coutarea portlandia</i>	655
<i>Couroupita surinamensis</i>	494	<i>Coutarea pubescens</i>	655
<i>Couroupita venezuelensis</i>	494	<i>Coutarea scherffiana</i>	655
<i>Coussapoa angustifolia</i>	713	<i>Coutarea speciosa</i>	655
<i>Coussapoa asperifolia</i>	713	<i>Crateva</i>	410, 411
<i>Coussapoa cayennensis</i>	713	<i>Crateva acuminata</i>	411, 412
<i>Coussapoa cornifolia</i>	713	<i>Crateva bahiana</i>	411
<i>Coussapoa cuneata</i>	713	<i>Crateva benthamii</i>	411
<i>Coussapoa emarginata</i>	715	<i>Crateva benthamii</i> var. <i>leptopetala</i>	411
<i>Coussapoa fagifolia</i>	713	<i>Crateva coriacea</i>	411
<i>Coussapoa ferruginea</i>	713	<i>Crateva glauca</i>	411
<i>Coussapoa ficina</i>	713	<i>Crateva gynandra</i>	411
<i>Coussapoa froesii</i>	713	<i>Crateva radiatiflora</i>	411, 412
<i>Coussapoa latifolia</i>	713	<i>Crateva tapia</i>	411
<i>Coussapoa latifolia</i> var. <i>obovata</i>	713	<i>Crateva tapia</i> var. <i>glauca</i>	411
<i>Coussapoa leprieuri</i>	713	<i>Crateva tapioides</i>	411
<i>Coussapoa microcephala</i>	713	<i>Cremanium paniculatum</i>	565
<i>Coussapoa microcephala</i> subsp. <i>cornifolia</i>	713	<i>Cremanium trinitatis</i>	565
<i>Coussapoa obovata</i>	713	<i>Cremastosperma brevipes</i>	370
<i>Coussapoa parvifolia</i>	713	<i>Cremastosperma poiteaui</i>	370
<i>Coussarea</i>	653	<i>Crepidospermum goudotianum</i>	402
<i>Coussarea albescens</i>	653	<i>Crepidospermum rhoifolium</i>	404
<i>Coussarea darienensis</i>	654	<i>Crepidospermum sprucei</i>	402
<i>Coussarea froelichia</i>	654	<i>Crossopetalum myrsinoides</i>	415
<i>Coussarea grandis</i>	653	<i>Croton benthamianus</i>	460
<i>Coussarea granvillei</i>	653	<i>Croton bilocularis</i>	460
<i>Coussarea hallei</i>	653	<i>Croton cajucara</i>	459
<i>Coussarea hirticalyx</i>	654	<i>Croton caryophyllus</i>	460
<i>Coussarea lutea</i>	653	<i>Croton cuneatus</i>	459
<i>Coussarea machadoana</i>	654	<i>Croton draconoides</i>	460
<i>Coussarea macrocarpa</i>	654	<i>Croton icabarui</i>	460
<i>Coussarea mapourioides</i>	653	<i>Croton kaieteuri</i>	460
<i>Coussarea martinii</i>	654	<i>Croton kavanayensis</i>	460
<i>Coussarea morii</i>	654	<i>Croton lanjouwianus</i>	460
<i>Coussarea multiflora</i>	657	<i>Croton martii</i>	459, 460
<i>Coussarea odoratissima</i>	657	<i>Croton martii</i> var. <i>latifolius</i>	460
<i>Coussarea paniculata</i>	654, 655	<i>Croton martii</i> var. <i>longifolius</i>	460
<i>Coussarea racemosa</i>	653	<i>Croton matourensis</i>	460
<i>Coussarea sertulifera</i>	657	<i>Croton matourensis</i> var. <i>benthamianus</i>	460
<i>Coutarea campanilla</i>	655	<i>Croton matourensis</i> var. <i>poeppigianus</i>	460
<i>Coutarea flavescens</i>	655	<i>Croton matourensis</i> var. <i>sericeus</i>	460
<i>Coutarea hexandra</i>	655	<i>Croton mimeticus</i>	460
<i>Coutarea hexandra</i> f. <i>albiflora</i>	655	<i>Croton monachinoensis</i>	460
<i>Coutarea hexandra</i> f. <i>grandiflora</i>	655	<i>Croton motilonorum</i>	459
<i>Coutarea hexandra</i> f. <i>pubescens</i>	655	<i>Croton pakaraimae</i>	460
<i>Coutarea hexandra</i> f. <i>roseiflora</i>	655	<i>Croton palanostigma</i>	460
<i>Coutarea hexandra</i> f. <i>tarapotensis</i>	655	<i>Croton perobtusus</i>	460
<i>Coutarea hexandra</i> var. <i>amazonica</i>	655	<i>Croton schiedeanus</i>	460
<i>Coutarea hexandra</i> var. <i>calycina</i>	655	<i>Croton seputubensis</i>	459
<i>Coutarea hexandra</i> var. <i>campanilla</i>	655	<i>Croton sericeus</i>	460
<i>Coutarea hexandra</i> var. <i>fluminensis</i>	655	<i>Croton subcoriaceus</i>	460
<i>Coutarea hexandra</i> var. <i>pubescens</i>	655	<i>Croton surinamensis</i>	459
<i>Coutarea hexandra</i> var. <i>speciosa</i>	655	<i>Croton tricolor</i>	460

Appendix 3. — Continuation.

<i>Croton</i> sp. A	460	<i>Cupania velutina</i>	681
<i>Crudia</i>	503, 509	<i>Cupirana aubletiana</i>	655
<i>Crudia aromatica</i>	509	<i>Cupirana martiniana</i>	656
<i>Crudia bracteata</i>	509	<i>Curatella americana</i>	448
<i>Crudia oblonga</i>	509	<i>Curatella americana</i> var. <i>pentagyna</i>	448
<i>Crudia parivoa</i>	511	<i>Curatella cambaiba</i>	448
<i>Crudia pubescens</i>	509	<i>Curatella grisebachiana</i>	448
<i>Crudia spicata</i>	511	<i>Cybianthus amplius</i>	644
<i>Crudia tomentosa</i>	511	<i>Cybianthus brownii</i>	645
<i>Cryptocarya guianensis</i>	478	<i>Cybianthus comatus</i>	645
<i>Cryptocarya maroniensis</i>	478	<i>Cybianthus egensis</i>	645
<i>Cryptocarya mucronata</i>	487	<i>Cybianthus guyanensis</i> subsp. <i>multipunctatus</i>	644
<i>Cryptocarya pachycarpa</i>	476	<i>Cybianthus leprieurii</i>	644
<i>Cryptocarya pyriformis</i>	479	<i>Cybianthus microbotrys</i>	644
<i>Cubaea paniculata</i>	543	<i>Cybianthus multipunctatus</i>	644
<i>Cubaea trigona</i>	543	<i>Cybianthus myrianthos</i>	645
<i>Cucullaria excelsa</i>	724	<i>Cybianthus nitidus</i>	645
<i>Cucullaria tetraphylla</i>	725	<i>Cybianthus potiae</i>	644
<i>Cucullaria tomentosa</i>	725	<i>Cybianthus prieurii</i>	645
<i>Cumetea alba</i>	611	<i>Cybianthus resinosus</i>	645
<i>Cumetea divaricata</i>	618	<i>Cybianthus schomburgkianus</i>	645
<i>Cumetea mini</i>	598	<i>Cybianthus subspicatus</i>	645
<i>Cumetea montana</i>	617	<i>Cybianthus surinamensis</i>	645
<i>Cumetea multiflora</i>	616	<i>Cybianthus venezuelanus</i>	645
<i>Cumetea tomentosa</i>	622	<i>Cybianthus viridiflorus</i>	645
<i>Cunuria bracteosa</i>	463	<i>Cybianthus</i> sp. A	645
<i>Cunuria spruceana</i> var. <i>bracteosa</i>	463	<i>Cyclolobium hostmannii</i>	527
<i>Cunuria uleana</i>	462	<i>Cymbopetalum brasiliense</i>	371
<i>Cupania aubletii</i>	681	<i>Cymbopetalum brasiliense</i> f. <i>latifolia</i>	371
<i>Cupania auriculata</i>	681	<i>Cymbopetalum odoratissimum</i>	371
<i>Cupania diphylla</i>	680	<i>Cymbopetalum venosum</i>	501
<i>Cupania frondosa</i>	681	<i>Cynodendron auratum</i>	685
<i>Cupania geminata</i>	680	<i>Cynodendron nitidum</i>	686
<i>Cupania guianensis</i>	681	<i>Cynometra</i>	503, 511
<i>Cupania hirsuta</i>	680	<i>Cynometra bauhinifolia</i>	511
<i>Cupania inelegans</i>	682	<i>Cynometra crassifolia</i>	511
<i>Cupania laevigata</i>	681, 682	<i>Cynometra hostmanniana</i>	511
<i>Cupania lanuginosa</i>	681	<i>Cynometra marginata</i>	511
<i>Cupania longifolia</i>	684	<i>Cynometra marginata</i> var. <i>guianensis</i>	511
<i>Cupania macrophylla</i>	684	<i>Cynometra marginata</i> var. <i>laevis</i>	511
<i>Cupania macrostylis</i>	681	<i>Cynometra parvifolia</i>	511
<i>Cupania multiflora</i>	683	<i>Cynophalla</i>	410, 412
<i>Cupania obovata</i>	681	<i>Cynophalla polyantha</i>	412
<i>Cupania porosa</i>	681	<i>Cyphomandra artocarpophyllus</i>	708
<i>Cupania praealta</i>	684	<i>Cyphomandra endopogon</i> subsp. <i>guianensis</i>	708
<i>Cupania reticulata</i>	681	<i>Cyphomandra hartwegii</i>	708
<i>Cupania rubiginosa</i>	681	<i>Cyphomandra mollicella</i>	708
<i>Cupania scrobiculata</i>	681	<i>Cyphomandra splendens</i>	708
<i>Cupania scrobiculata</i> f. <i>frondosa</i>	681	<i>Cyrillopsis paraensis</i>	471
<i>Cupania scrobiculata</i> f. <i>guianensis</i>	681	<i>Cytisus membranaceus</i>	528
<i>Cupania scrobiculata</i> f. <i>reticulata</i>	681		
<i>Cupania scrobiculata</i> var. <i>frondosa</i>	681		
<i>Cupania scrobiculata</i> var. <i>reticulata</i>	681		
<i>Cupania subrepanda</i>	681		

Appendix 3. — Continuation.

D		
<i>Dacryodes cuspidata</i>	399	<i>Dimorphandra multiflora</i> 512
<i>Dacryodes decidua</i>	399	<i>Dimorphandra polyandra</i> 512
<i>Dacryodes nitens</i>	399	<i>Dimorphandra pullei</i> 512
<i>Dacryodes oblongipetala</i>	399	<i>Dinizia</i> 503, 512
<i>Dacryodes roraimensis</i>	399	<i>Dinizia excelsa</i> 512
<i>Dacryodes</i> sp. A	399	<i>Diospyros capimnensis</i> 448
<i>Dacryodes villosa</i>	399	<i>Diospyros capreifolia</i> 448
<i>Dalbergia heptaphylla</i>	528	<i>Diospyros carbonaria</i> 448
<i>Dalbergia pentaphylla</i>	528	<i>Diospyros cavalcantei</i> 448
<i>Damburneya purpurea</i>	478	<i>Diospyros cayennensis</i> 449
<i>Darluca prostrata</i>	657	<i>Diospyros dichroa</i> 449
<i>Dasynema alnifolium</i>	452	<i>Diospyros duckei</i> 448
<i>Dasynema cuneifolium</i>	452	<i>Diospyros guianensis</i> 449
<i>Dasynema obtusifolium</i>	453	<i>Diospyros ierensis</i> 449
<i>Dasynema obtusum</i>	453	<i>Diospyros martinii</i> 449
<i>Dasynema pubescens</i>	452, 453	<i>Diospyros melinonii</i> 448
<i>Debraea floribunda</i>	722	<i>Diospyros praetermissa</i> 449
<i>Decaraphe hostmannii</i>	568	<i>Diospyros sericea</i> 449
<i>Deguelia</i>	528	<i>Diospyros</i> sp. A 449
<i>Dendrobangia boliviiana</i>	582	<i>Diospyros tetrandra</i> 449
<i>Dendrobangia tenuis</i>	582	<i>Diospyros vestita</i> 449
<i>Derris hedyosma</i>	528	<i>Diploön cuspidatum</i> 688
<i>Derris latifolia</i>	528	<i>Diploön venezuelanum</i> 688
<i>Detarioideae</i>	503	<i>Diplochita bracteata</i> 564
<i>Dialioideae</i>	503	<i>Diplochita florida</i> 566
<i>Dialium</i>	503, 511	<i>Diplochita fothergilla</i> 565
<i>Dialium guianense</i>	511	<i>Diplochita leucocephala</i> 568
<i>Diasperus antillanus</i>	638	<i>Diplochita macrophylla</i> 568
<i>Diasperus guianensis</i>	638	<i>Diplochita mucronata</i> 564
<i>Dibrachion guianense</i>	513	<i>Diplochita parviflora</i> 567
<i>Dibrachion riparium</i>	512	<i>Diplochita rosea</i> 566
DICHAPETALACEAE	447	<i>Diplochita serrulata</i> 568
<i>Dichrostachys macrostachya</i>	535	<i>Diplochita serrulata</i> var. <i>latifolia</i> 568
<i>Dicorynia</i>	503, 511	<i>Diplochita sessilifolia</i> 566
<i>Dicorynia guianensis</i>	511	<i>Diplochita tomentosa</i> 569
<i>Didymandra</i>	471, 472	<i>Diplocrater acuminatus</i> 633
<i>Didymandra purpurea</i>	471	<i>Diplothemium henryanum</i> 392
<i>Didymopanax decaphyllus</i>	388	<i>Diplotropis</i> 503, 512
<i>Didymopanax morototoni</i>	389	<i>Diplotropis brachypetala</i> 509
<i>Didymopanax morototoni</i> var. <i>poeppigii</i>	389	<i>Diplotropis guianensis</i> 513
<i>Didymopanax morototoni</i> var. <i>sessiliflorus</i>	389	<i>Diplotropis martiusii</i> 512
<i>Didymopanax poeppigii</i>	389	<i>Diplotropis purpurea</i> 513
<i>Didymopanax splendens</i>	389	<i>Dipteryx</i> 503, 513
DILLENIACEAE	448	<i>Dipteryx applanata</i> 543
<i>Dimorpha falcata</i>	514	<i>Dipteryx nudipes</i> 543
<i>Dimorphandra</i>	503, 512	<i>Dipteryx odorata</i> 513
<i>Dimorphandra coccinea</i>	512	<i>Dipteryx oppositifolia</i> 543
<i>Dimorphandra cuprea</i>	512	<i>Dipteryx oppositifolia</i> var. <i>parviflora</i> 543
<i>Dimorphandra glabrifolia</i>	512	<i>Dipteryx parviflora</i> 544
<i>Dimorphandra hohenkerkii</i>	512	<i>Dipteryx punctata</i> 513
<i>Dimorphandra ignea</i>	512	<i>Dipteryx speciosa</i> 544
<i>Dimorphandra macrostachya</i> subsp. <i>glabrifolia</i>	—	<i>Dipteryx tetraphylla</i> 513
		<i>Dipteryx trifoliolata</i> 513
		<i>Discipiper reticulatum</i> 642

Appendix 3. — Continuation.

<i>Discocarpus mazarunensis</i>	636	E
<i>Discophora froesii</i>	708	
<i>Discophora guianensis</i>	708	EBENACEAE 448
<i>Discophora panamensis</i>	708	<i>Ecclinusa abbreviata</i> 689
<i>Ditmaria floribunda</i>	722	<i>Ecclinusa bacuri</i> 688
<i>Diuroglossum rufescens</i>	554	<i>Ecclinusa cochlearia</i> 703
<i>Dodecastigma integrifolium</i>	461	<i>Ecclinusa costata</i> 689
<i>Dodecastigma mazarunense</i>	461	<i>Ecclinusa cuneifolia</i> 686
<i>Dodecastigma uleanum</i>	462	<i>Ecclinusa cyanogena</i> 687
<i>Douglassia laurina</i>	474	<i>Ecclinusa eximia</i> 686
<i>Drypetes fanshawei</i>	648	<i>Ecclinusa guianensis</i> 688
<i>Drypetes maguireana</i>	636	<i>Ecclinusa lanceolata</i> 689
<i>Drypetes spruceana</i>	636	<i>Ecclinusa prieurii</i> 687
<i>Drypetes variabilis</i>	648	<i>Ecclinusa ramiflora</i> 689
<i>Duchassaingia glauca</i>	515	<i>Ecclinusa rufocuprea</i> 686
<i>Duchassaingia ovalifolia</i>	515	<i>Ecclinusa sanguinolenta</i> 687
<i>Duguettia adiscandra</i>	371	<i>Echites brasiliensis</i> 387
<i>Duguettia cadaverica</i>	371	<i>Eganthus poeppigii</i> 634
<i>Duguettia calycina</i>	371	<i>Elaeagia brasiliensis</i> 652
<i>Duguettia calycina</i> subsp. <i>jenmanii</i>	371	ELAEOCARPACEAE 449
<i>Duguettia calycina</i> subsp. <i>versteegii</i>	371	<i>Elaeodendron macrophyllum</i> 414
<i>Duguettia caudata</i>	372	<i>Elaeoluma nuda</i> 689
<i>Duguettia cuspidata</i>	371	<i>Elaeoluma</i> sp. A 689
<i>Duguettia elegans</i>	371	<i>Elaphrium altissimum</i> 399
<i>Duguettia eximia</i>	371	<i>Elaphrium decandrum</i> 400
<i>Duguettia friesii</i>	371	<i>Elaphrium enneandrum</i> 400
<i>Duguettia granvilleana</i>	371	<i>Elaphrium guianense</i> 402
<i>Duguettia insculpta</i>	372	<i>Elaphrium heptaphyllum</i> 402
<i>Duguettia longifolia</i>	372	<i>Elaphrium heterophyllum</i> 400
<i>Duguettia obovata</i>	372	<i>Elizabetha leiogyne</i> 533
<i>Duguettia paraensis</i>	371	<i>Elizabetha paraensis</i> 533
<i>Duguettia pycnastera</i>	372	<i>Elizabetha princeps</i> 533
<i>Duguettia riparia</i>	372	<i>Elvasia caurensis</i> 628
<i>Duguettia sandwithii</i>	372	<i>Elvasia elvasiooides</i> 628
<i>Duguettia surinamensis</i>	372	<i>Elvasia hostmannia</i> 628
<i>Duguettia trunciflora</i>	372	<i>Elvasia macrostipularis</i> 628
<i>Duguettia yeshidan</i>	372	<i>Elvasia sagotii</i> 628
<i>Duhamelia coccinea</i>	660	<i>Embothrium pinnatum</i> 647
<i>Duhamelia glabra</i>	650	<i>Emmotum fagifolium</i> 582
<i>Dulacia</i>	633	<i>Enckea lata</i> 642
<i>Dulacia guianensis</i>	633	<i>Enckea reticulata</i> 642
<i>Duroia aquatica</i>	655	<i>Enckea smilacifolia</i> 642
<i>Duroia eriopila</i>	655, 656	<i>Endandra itauba</i> 481
<i>Duroia eriopila</i> f. <i>glabra</i>	656	<i>Endlicheria bracteolata</i> 479
<i>Duroia longiflora</i>	656	<i>Endlicheria canescens</i> 479
<i>Duroia martiniana</i>	656	<i>Endlicheria chalisea</i> 479
<i>Duroia micrantha</i>	656	<i>Endlicheria endllicheriopsis</i> 485
<i>Duroia sprucei</i>	656	<i>Endlicheria glaberrima</i> 479
<i>Duroia surinamensis</i>	656	<i>Endlicheria glomerata</i> 480
<i>Dussia</i>	503, 513	<i>Endlicheria grandis</i> 490
<i>Dussia cayennensis</i>	513	<i>Endlicheria impressa</i> 474
<i>Dussia discolor</i>	513	<i>Endlicheria melinonii</i> 479
<i>Dussia micranthera</i>	513	<i>Endlicheria punctulata</i> 479
<i>Dussia tessmannii</i>	513	<i>Endlicheria pyriformis</i> 479

Appendix 3. — Continuation.

<i>Endlicheria rubriflora</i>	479	<i>Erisma nitidum</i>	722
<i>Endlicheria szyszlowiczii</i>	479	<i>Erisma pallidiflorum</i>	722
<i>Endlicheria trianae</i>	479	<i>Erisma parvifolium</i>	722
<i>Endlicheria wurdackiana</i>	479	<i>Erisma parvifolium</i> var. <i>pallidiflorum</i>	722
<i>Endlicheria</i> sp. A	480	<i>Erisma pulverulentum</i>	722
<i>Endosteira oppositifolia</i>	649	<i>Erisma uncinatum</i>	722
<i>Endusa punctata</i>	634	<i>Eryngium foetidum</i>	722
<i>Englerella macrocarpa</i>	701	<i>Erythrina</i>	503, 515
<i>Englerella speciosa</i>	701	<i>Erythrina amasina</i>	515
<i>Englerophoenix caribaeum</i>	392	<i>Erythrina atrosanguinea</i>	515
<i>Englerophoenix longirostrata</i>	392	<i>Erythrina caffra</i>	515
<i>Englerophoenix maripa</i>	392	<i>Erythrina darienensis</i>	515
<i>Englerophoenix regia</i>	392	<i>Erythrina fusca</i>	515
<i>Entada urbaeana</i>	536	<i>Erythrina fusca</i> var. <i>inermis</i>	515
<i>Enterolobium</i>	503, 513	<i>Erythrina glauca</i>	515
<i>Enterolobium oldemanii</i>	513	<i>Erythrina micropteryx</i>	515
<i>Enterolobium schomburgkii</i>	514	<i>Erythrina moelebei</i>	515
<i>Enterolobium</i> sp. A	514	<i>Erythrina ovalifolia</i>	515
<i>Eperua</i>	503, 514	<i>Erythrina patens</i>	515
<i>Eperua bijuga</i>	514	<i>Erythrina pisamo</i>	515
<i>Eperua bijuga</i> f. <i>typica</i>	514	<i>Erythrina poeppigiana</i>	515
<i>Eperua falcata</i>	514	<i>Erythrociton brasiliensis</i>	667
<i>Eperua grandiflora</i>	514	ERYTHROXYLACEAE	455
<i>Eperua jenmanii</i>	514	<i>Erythroxylum acutifolium</i>	455
<i>Eperua kourouensis</i>	514	<i>Erythroxylum albertianum</i>	457
<i>Eperua rubiginosa</i>	514	<i>Erythroxylum amazonicum</i>	455
<i>Eperua schomburgkiana</i>	514	<i>Erythroxylum areolatum</i>	457
<i>Eperua stipulata</i>	514	<i>Erythroxylum aristigerum</i>	457
<i>Ephedranthus guianensis</i>	372	<i>Erythroxylum aristigerum</i> var. <i>bahiense</i>	457
<i>Ephedranthus guianensis</i> var. <i>oligantha</i>	372	<i>Erythroxylum bahiense</i>	457
<i>Epigenia integerrima</i>	709	<i>Erythroxylum citrifolium</i>	455
<i>Epleienda sinemariensis</i>	606	<i>Erythroxylum citrifolium</i> var. <i>latifolium</i>	455
<i>Eremoluma krukoffii</i>	694	<i>Erythroxylum citrifolium</i> var. <i>minus</i>	455
<i>Eremoluma sagotiana</i>	701	<i>Erythroxylum coelophlebium</i> var. <i>grisebachii</i>	455
<i>Eremoluma williamii</i>	702	<i>Erythroxylum comosum</i>	457
<i>Eremoluma wurdackii</i>	694	<i>Erythroxylum costaricense</i>	456
<i>Eriodendron anfractuosum</i>	552	<i>Erythroxylum duckei</i>	455
<i>Eriodendron anfractuosum</i> var. <i>africanum</i>	552	<i>Erythroxylum ellipticum</i>	456
<i>Eriodendron anfractuosum</i> var. <i>caribaeum</i>	552	<i>Erythroxylum filipes</i>	456
<i>Eriodendron anfractuosum</i> var. <i>guianense</i>	552	<i>Erythroxylum fimbriatum</i>	455
<i>Eriodendron anfractuosum</i> var. <i>indicum</i>	552	<i>Erythroxylum floribundum</i>	456
<i>Eriodendron caribaeum</i>	552	<i>Erythroxylum gomphiooides</i>	455
<i>Eriodendron guineense</i>	552	<i>Erythroxylum grisebachii</i>	455
<i>Eriodendron inerme</i>	553	<i>Erythroxylum kapplerianum</i>	455
<i>Eriodendron occidentale</i>	552	<i>Erythroxylum kirkianum</i>	457
<i>Eriodendron orientale</i>	552	<i>Erythroxylum laurinum</i>	456
<i>Eriodendron pentandrum</i>	552	<i>Erythroxylum leptoneurum</i>	455
<i>Eriotheca crassa</i>	553	<i>Erythroxylum ligustrinum</i>	455
<i>Eriotheca globosa</i>	553	<i>Erythroxylum lineolatum</i>	455
<i>Eriotheca longitubulosa</i>	553	<i>Erythroxylum lucidum</i>	456
<i>Eriotheca macrophylla</i> subsp. <i>sclerophylla</i>	553	<i>Erythroxylum lucidum</i> var. <i>costaricense</i>	456
<i>Eriotheca</i> sp. A	553	<i>Erythroxylum macrophyllum</i>	456
<i>Eriotheca surinamensis</i>	553	<i>Erythroxylum mapuerae</i>	455
<i>Erisma floribundum</i>	722	<i>Erythroxylum micranthum</i>	455

Appendix 3. — Continuation.

<i>Erythroxylum mucronatum</i>	455, 457	<i>Eschweilera montana</i>	497
<i>Erythroxylum mucronatum</i> var. <i>majus</i>	455	<i>Eschweilera odora</i>	495
<i>Erythroxylum paraense</i>	455	<i>Eschweilera ovalis</i>	501
<i>Erythroxylum roraimae</i>	457	<i>Eschweilera pallida</i>	495
<i>Erythroxylum squamatum</i>	457	<i>Eschweilera parviflora</i>	497
<i>Erythroxylum squamatum</i> var. <i>microcarpum</i>	457	<i>Eschweilera patrisii</i>	501
<i>Erythroxylum suberosum</i>	457	<i>Eschweilera pedicellata</i>	497
<i>Erythroxylum suberosum</i> f. <i>brevipetiolatum</i>	457	<i>Eschweilera pilosa</i>	497
<i>Erythroxylum tabascense</i>	456	<i>Eschweilera piresii</i> subsp. <i>viridipetala</i>	497
<i>Erythroxylum testaceum</i>	457	<i>Eschweilera platycarpa</i>	497
<i>Erythroxylum trinerve</i>	457	<i>Eschweilera poiteaui</i>	502
<i>Erythroxylum venezuelense</i>	457	<i>Eschweilera polyantha</i>	496
<i>Eschweilera</i>	494	<i>Eschweilera praeclera</i>	502
<i>Eschweilera acuminatissima</i>	495	<i>Eschweilera retroflexa</i>	495
<i>Eschweilera alata</i>	494	<i>Eschweilera rigida</i>	500
<i>Eschweilera alba</i>	495	<i>Eschweilera rorida</i>	500
<i>Eschweilera amara</i>	501	<i>Eschweilera sagotiana</i>	497
<i>Eschweilera apiculata</i>	494	<i>Eschweilera salebrosa</i>	501
<i>Eschweilera brancoensis</i>	500	<i>Eschweilera sandwithiana</i>	495
<i>Eschweilera carrii</i>	500	<i>Eschweilera simiorum</i>	502
<i>Eschweilera chartacea</i>	500	<i>Eschweilera simiorum</i> var. <i>latifolia</i>	502
<i>Eschweilera chartaceifolia</i>	495	<i>Eschweilera simplex</i>	371
<i>Eschweilera collina</i>	495	<i>Eschweilera spruceana</i>	500
<i>Eschweilera confertiflora</i>	500	<i>Eschweilera squamata</i>	498
<i>Eschweilera congestiflora</i>	501	<i>Eschweilera subglandulosa</i>	498
<i>Eschweilera coriacea</i>	495	<i>Eschweilera tapuya</i>	497
<i>Eschweilera corrugata</i>	501	<i>Eschweilera vageleri</i>	495
<i>Eschweilera decolorans</i>	495	<i>Eschweilera wachenheimii</i>	499
<i>Eschweilera eymaana</i>	495	<i>Eschweilera</i> sp. A	499
<i>Eschweilera flaccida</i>	497	<i>Eschweilera</i> sp. B	499
<i>Eschweilera floribunda</i>	496	<i>Esenbeckia attenuata</i>	667
<i>Eschweilera fracta</i>	495	<i>Esenbeckia cowanii</i>	667
<i>Eschweilera fractiflexa</i>	495	<i>Esenbeckia fasciculata</i>	667
<i>Eschweilera gracilipes</i>	496	<i>Esenbeckia grandiflora</i>	667
<i>Eschweilera grandiflora</i>	495, 499	<i>Esenbeckia grandiflora</i> var. <i>macrophylla</i>	667
<i>Eschweilera grandifolia</i>	495	<i>Esenbeckia grandiflora</i> var. <i>peruviana</i>	667
<i>Eschweilera grata</i>	497	<i>Esenbeckia obovalifolia</i>	667
<i>Eschweilera holcogyme</i>	501	<i>Esenbeckia rigida</i>	667
<i>Eschweilera idatimon</i>	501	<i>Ethnora maripa</i>	392
<i>Eschweilera idatimonoides</i>	497	<i>Eugenia</i>	598
<i>Eschweilera jenmanii</i>	501	<i>Eugenia acuminatissima</i>	599
<i>Eschweilera jucunda</i>	497	<i>Eugenia acutiloba</i>	599
<i>Eschweilera labriculata</i>	493	<i>Eugenia adenocalyx</i>	601
<i>Eschweilera laevicarpa</i>	496	<i>Eugenia albicans</i>	598
<i>Eschweilera laevisfolia</i>	499	<i>Eugenia alexandri</i>	599
<i>Eschweilera longipes</i>	497	<i>Eugenia alfaroana</i>	599
<i>Eschweilera longipes</i> f. <i>genuina</i>	497	<i>Eugenia amanuensis</i>	599
<i>Eschweilera longipes</i> f. <i>platycarpa</i>	497	<i>Eugenia amazonica</i>	604
<i>Eschweilera lutea</i>	501	<i>Eugenia anastomosans</i>	598
<i>Eschweilera macrophylla</i>	497	<i>Eugenia arivoa</i>	606
<i>Eschweilera marawynensis</i>	500	<i>Eugenia armeniaca</i>	598
<i>Eschweilera matamata</i>	495	<i>Eugenia atropunctata</i>	602
<i>Eschweilera melinonii</i>	497	<i>Eugenia atropunctata</i> var. <i>gracilis</i>	602
<i>Eschweilera micrantha</i>	496	<i>Eugenia atropunctata</i> var. <i>robusta</i>	602

Appendix 3. — Continuation.

<i>Eugenia aubletiana</i>	598	<i>Eugenia eschholtziana</i>	606
<i>Eugenia australis</i>	599	<i>Eugenia eschholtziana</i> var. <i>angustifolia</i>	606
<i>Eugenia berlynensis</i>	605	<i>Eugenia eschholtziana</i> var. <i>latifolia</i>	606
<i>Eugenia berteroana</i>	599	<i>Eugenia exaltata</i>	601
<i>Eugenia biflora</i>	598, 599	<i>Eugenia excelsa</i>	601
<i>Eugenia biflora</i> var. <i>hoffmannseggii</i>	599	<i>Eugenia fallax</i>	618, 622
<i>Eugenia biflora</i> var. <i>lancea</i>	599	<i>Eugenia fasciculiflora</i>	604
<i>Eugenia biflora</i> var. <i>ludibunda</i>	599	<i>Eugenia feijoi</i>	604
<i>Eugenia biflora</i> var. <i>mini</i>	598	<i>Eugenia fenzliana</i>	604
<i>Eugenia biflora</i> var. <i>myriostigma</i>	599	<i>Eugenia ferreiraeanae</i>	602
<i>Eugenia biflora</i> var. <i>pallens</i>	598	<i>Eugenia ferruginea</i>	612
<i>Eugenia biflora</i> var. <i>virgultosa</i>	598	<i>Eugenia fieldingii</i>	599
<i>Eugenia biflora</i> var. <i>wallenii</i>	599	<i>Eugenia flavesrens</i>	602
<i>Eugenia brachybotrya</i>	571	<i>Eugenia flavesens</i> var. <i>guyanensis</i>	602
<i>Eugenia brachythrix</i>	599	<i>Eugenia flavesens</i> var. <i>longifolia</i>	602
<i>Eugenia bracteata</i>	610	<i>Eugenia flavesens</i> var. <i>parvifolia</i>	602
<i>Eugenia brownsbergii</i>	600	<i>Eugenia flavonigra</i>	603
<i>Eugenia brownsbergii</i> var. <i>glauca</i>	600	<i>Eugenia flavonigra</i> var. <i>guadalupensis</i>	603
<i>Eugenia buxizans</i>	615	<i>Eugenia flavonigra</i> var. <i>martinicensis</i>	603
<i>Eugenia calothysrsa</i>	604	<i>Eugenia floribunda</i>	624
<i>Eugenia casarettiana</i>	604	<i>Eugenia florida</i>	602
<i>Eugenia catinga</i>	604	<i>Eugenia forsteri</i>	605
<i>Eugenia caurensis</i>	599	<i>Eugenia freireana</i>	599
<i>Eugenia chrysophylloides</i>	600	<i>Eugenia freireana</i> var. <i>angustifolia</i>	599
<i>Eugenia chrysophyllum</i>	600	<i>Eugenia freireana</i> var. <i>latifolia</i>	599
<i>Eugenia ciliolata</i>	624	<i>Eugenia galbaoensis</i>	602
<i>Eugenia citrifolia</i>	600	<i>Eugenia gardneriana</i>	602
<i>Eugenia coffeifolia</i>	606	<i>Eugenia gardneriana</i> var. <i>depauperata</i>	602
<i>Eugenia coffeifolia</i> var. <i>grandifolia</i>	606	<i>Eugenia gardneriana</i> var. <i>dives</i>	602
<i>Eugenia coffeifolia</i> var. <i>parvifolia</i>	606	<i>Eugenia gardneriana</i> var. <i>ovata</i>	602
<i>Eugenia coloradoensis</i>	602	<i>Eugenia gardneriana</i> var. <i>rigida</i>	602
<i>Eugenia compta</i>	605	<i>Eugenia gerdae</i>	603
<i>Eugenia correae</i>	603	<i>Eugenia gomesiana</i>	603
<i>Eugenia costata</i>	604	<i>Eugenia gongylocarpa</i>	603
<i>Eugenia coumetae</i>	611	<i>Eugenia griseiflora</i>	603
<i>Eugenia cowanii</i>	601	<i>Eugenia guianensis</i>	613
<i>Eugenia cryptadena</i>	601, 606	<i>Eugenia hagendorffii</i>	625
<i>Eugenia cryptadena</i> var. <i>gracilis</i>	601	<i>Eugenia hartii</i>	599
<i>Eugenia cucullata</i>	601	<i>Eugenia hirsuta</i>	611
<i>Eugenia cupulata</i>	601	<i>Eugenia hoffmannseggii</i>	599
<i>Eugenia cuspidata</i>	601	<i>Eugenia hoffmannseggii</i> var. <i>grandifolia</i>	599
<i>Eugenia deflexa</i>	611	<i>Eugenia hoffmannseggii</i> var. <i>parvifolia</i>	599
<i>Eugenia denigrata</i>	601	<i>Eugenia inaequiloba</i>	615
<i>Eugenia densiracemosa</i>	601	<i>Eugenia inocarpa</i>	605
<i>Eugenia desvauxiana</i>	597	<i>Eugenia inundata</i> var. <i>acutifolia</i>	599
<i>Eugenia divaricata</i>	618	<i>Eugenia jamaicana</i>	599
<i>Eugenia doniana</i>	606	<i>Eugenia jamaicensis</i>	599
<i>Eugenia egensis</i>	601	<i>Eugenia kiaerskoviana</i>	604
<i>Eugenia egensis</i> var. <i>angustifolia</i>	601	<i>Eugenia lambertiana</i>	603
<i>Eugenia egensis</i> var. <i>bimarginata</i>	601	<i>Eugenia lambertiana</i> var. <i>bispidula</i>	603
<i>Eugenia egensis</i> var. <i>grandifolia</i>	601	<i>Eugenia lancea</i>	599
<i>Eugenia egensis</i> var. <i>latifolia</i>	601	<i>Eugenia lasiopus</i>	611
<i>Eugenia egensis</i> var. <i>parvifolia</i>	601	<i>Eugenia latifolia</i>	603
<i>Eugenia egensis</i> var. <i>tenuiramis</i>	601	<i>Eugenia laxiflora</i>	618

Appendix 3. — Continuation.

<i>Eugenia leptophlebia</i>	599	<i>Eugenia paniculiflora</i>	619
<i>Eugenia leucanthera</i>	604	<i>Eugenia paraensis</i>	604
<i>Eugenia leucophloea</i>	624, 625	<i>Eugenia parodiana</i>	601
<i>Eugenia leucophloea</i> var. <i>warmingiana</i>	625	<i>Eugenia patens</i>	605
<i>Eugenia llewelynii</i>	600	<i>Eugenia patrisii</i>	605
<i>Eugenia loretensis</i>	599	<i>Eugenia patula</i>	602
<i>Eugenia luciae</i>	603	<i>Eugenia perforata</i>	601
<i>Eugenia ludibunda</i>	599	<i>Eugenia periplocifolia</i>	618
<i>Eugenia lugens</i>	604	<i>Eugenia perorebi</i>	602
<i>Eugenia macarensis</i>	599	<i>Eugenia piauhiensis</i>	606
<i>Eugenia macrocalyx</i>	607	<i>Eugenia pisonis</i>	604
<i>Eugenia maculata</i>	601	<i>Eugenia pleurosiiphonea</i>	604
<i>Eugenia maranhensis</i>	624	<i>Eugenia polyantha</i>	615
<i>Eugenia marowynensis</i>	603	<i>Eugenia polyneura</i>	625
<i>Eugenia martiusiana</i>	606	<i>Eugenia polystachya</i>	605
<i>Eugenia maximiliana</i>	624	<i>Eugenia portoricensis</i>	605
<i>Eugenia megalocarpa</i>	606	<i>Eugenia portoricensis</i> var. <i>brevipes</i>	605
<i>Eugenia melanosticta</i>	602	<i>Eugenia prieurii</i>	605, 606
<i>Eugenia melinonii</i>	606	<i>Eugenia prieurii</i> var. <i>robusta</i>	605, 606
<i>Eugenia membranacea</i>	602	<i>Eugenia prieurii</i> var. <i>tenuiramis</i>	605
<i>Eugenia meyeriana</i>	599	<i>Eugenia prosoneura</i>	603
<i>Eugenia meyeriana</i> var. <i>depauperata</i>	599	<i>Eugenia protracta</i>	624
<i>Eugenia meyeriana</i> var. <i>dives</i>	599	<i>Eugenia pseudopsidium</i>	605
<i>Eugenia microcarpos</i>	598	<i>Eugenia pseudopsidium</i> var. <i>genuina</i>	605
<i>Eugenia microcarpos</i> var. <i>mini</i>	598	<i>Eugenia pseudopsidium</i> var. <i>portoricensis</i>	605
<i>Eugenia mikaniana</i>	619	<i>Eugenia psidiooides</i>	605
<i>Eugenia mimus</i>	604	<i>Eugenia ptariensis</i>	598
<i>Eugenia mini</i>	598	<i>Eugenia pycnoneura</i>	625
<i>Eugenia mini</i> var. <i>microcarpos</i>	598	<i>Eugenia pyrifolia</i>	618
<i>Eugenia modesta</i> var. <i>jamaicensis</i>	599	<i>Eugenia racemifera</i>	602, 604
<i>Eugenia mollis</i>	620	<i>Eugenia racemosa</i>	599
<i>Eugenia montana</i>	617	<i>Eugenia ramiflora</i> var. <i>montana</i>	—
<i>Eugenia monticola</i>	603, 624	<i>Eugenia richardiana</i>	599
<i>Eugenia monticola</i> var. <i>racemosa</i>	603	<i>Eugenia riparia</i>	604
<i>Eugenia moraviana</i> var. <i>gardneriana</i>	602	<i>Eugenia ripidocarpa</i>	604
<i>Eugenia morii</i>	604	<i>Eugenia rivularis</i>	625
<i>Eugenia moschata</i>	604	<i>Eugenia roraimana</i>	606
<i>Eugenia multiflora</i>	616	<i>Eugenia rugosa</i>	604
<i>Eugenia muricata</i>	604	<i>Eugenia rutidocarpa</i>	604
<i>Eugenia muricata</i> var. <i>guyanensis</i>	604	<i>Eugenia salicifolia</i>	599
<i>Eugenia myriostigma</i>	599	<i>Eugenia salzmannii</i>	624
<i>Eugenia nemoralis</i>	602	<i>Eugenia schaueriana</i>	612
<i>Eugenia neograndifolia</i>	605	<i>Eugenia schlechtendaliana</i>	605
<i>Eugenia nicholsii</i>	599	<i>Eugenia schomburgkii</i>	603
<i>Eugenia nigra</i>	571	<i>Eugenia seriatoracemosa</i>	602
<i>Eugenia nitida</i>	615	<i>Eugenia sericiflora</i>	599
<i>Eugenia oblongifolia</i>	598	<i>Eugenia sinemariensis</i>	606
<i>Eugenia obovata</i>	602	<i>Eugenia smaragdina</i>	603
<i>Eugenia ochra</i>	606	<i>Eugenia smaragdina</i> var. <i>angustifolia</i>	603
<i>Eugenia oligoneura</i>	602	<i>Eugenia smaragdina</i> var. <i>brevipes</i>	603
<i>Eugenia oligophylla</i>	603	<i>Eugenia smaragdina</i> var. <i>rigida</i>	603
<i>Eugenia omissa</i>	605	<i>Eugenia sparsiflora</i>	597
<i>Eugenia oneillii</i>	625	<i>Eugenia sphaerosperma</i>	601
<i>Eugenia pallens</i>	598	<i>Eugenia spicata</i>	604
<i>Eugenia paniculata</i>	611	<i>Eugenia</i> sp. A	607

Appendix 3. — Continuation.

<i>Eugenia</i> sp. B	607, 608	<i>Euplassa meridionalis</i>	646
<i>Eugenia</i> sp. C	607	<i>Euplassa pinnata</i>	646, 647
<i>Eugenia</i> sp. D	607	<i>Eurychaenia punctata</i>	563, 568
<i>Eugenia</i> sp. E	607	<i>Eurychaenia punctata</i> var. <i>latifolia</i>	563
<i>Eugenia</i> sp. F	607	<i>Euterpe badiocarpa</i>	393
<i>Eugenia</i> sp. G	607	<i>Euterpe beardii</i>	393
<i>Eugenia</i> sp. H	608	<i>Euterpe brasiliiana</i>	393
<i>Eugenia</i> sp. I	608	<i>Euterpe confertiflora</i>	393
<i>Eugenia</i> sp. J	608	<i>Euterpe cuatrecasana</i>	393
<i>Eugenia</i> sp. K	608	<i>Euterpe jataipuensis</i>	393
<i>Eugenia</i> sp. L	608	<i>Euterpe langloisii</i>	393
<i>Eugenia</i> sp. M	608	<i>Euterpe oleracea</i>	393
<i>Eugenia</i> sp. N	608	<i>Euterpe petiolata</i>	393
<i>Eugenia</i> sp. O	608	<i>Euterpe precatoria</i>	393
<i>Eugenia</i> sp. P	608	<i>Euterpe stenophylla</i>	393
<i>Eugenia</i> sp. Q	608	<i>Euterpe subruminata</i>	393
<i>Eugenia</i> sp. R	608	<i>Excoecaria aerea</i>	465
<i>Eugenia spruceana</i>	606	<i>Excoecaria arguta</i>	464
<i>Eugenia stictopetala</i>	606	<i>Excoecaria biglandulosa</i>	464-467
<i>Eugenia subobliqua</i>	622	<i>Excoecaria biglandulosa</i> f. <i>oblongata</i>	465
<i>Eugenia sulcata</i>	612	<i>Excoecaria biglandulosa</i> f. <i>obovata</i>	465
<i>Eugenia sylvatica</i>	602	<i>Excoecaria biglandulosa</i> var. <i>aubletiana</i>	465
<i>Eugenia tapacumensis</i>	606	<i>Excoecaria biglandulosa</i> var. <i>aucuparia</i>	464
<i>Eugenia tapacumensis</i> var. <i>angustifolia</i>	606	<i>Excoecaria biglandulosa</i> var. <i>clauseniana</i>	466
<i>Eugenia tapacumensis</i> var. <i>latifolia</i>	606	<i>Excoecaria biglandulosa</i> var. <i>cuneata</i>	465
<i>Eugenia teffensis</i>	605	<i>Excoecaria biglandulosa</i> var. <i>dracunculoides</i>	466
<i>Eugenia teffensis</i> var. <i>subcordata</i>	605	<i>Excoecaria biglandulosa</i> var. <i>glandulata</i>	465
<i>Eugenia teffensis</i> var. <i>truncata</i>	605	<i>Excoecaria biglandulosa</i> var. <i>grandifolia</i>	466
<i>Eugenia tenuiflora</i>	606	<i>Excoecaria biglandulosa</i> var. <i>hamata</i>	465
<i>Eugenia tenuiramis</i>	601	<i>Excoecaria biglandulosa</i> var. <i>hippomane</i>	466
<i>Eugenia tetramera</i>	607	<i>Excoecaria biglandulosa</i> var. <i>intercedens</i>	466
<i>Eugenia tinge-lingua</i>	602	<i>Excoecaria biglandulosa</i> var. <i>klotzschiana</i>	465
<i>Eugenia tomentosa</i>	622	<i>Excoecaria biglandulosa</i> var. <i>lanceolata</i>	465
<i>Eugenia vallis</i>	623	<i>Excoecaria biglandulosa</i> var. <i>leptadenia</i>	466
<i>Eugenia variifolia</i>	625	<i>Excoecaria biglandulosa</i> var. <i>montevidensis</i>	466
<i>Eugenia vellozoi</i>	605	<i>Excoecaria biglandulosa</i> var. <i>moritziana</i>	465
<i>Eugenia verruculosa</i>	605	<i>Excoecaria biglandulosa</i> var. <i>pavoniana</i>	465
<i>Eugenia virgata</i>	599	<i>Excoecaria biglandulosa</i> var. <i>petiolaris</i>	466
<i>Eugenia virgultosa</i>	598, 599	<i>Excoecaria biglandulosa</i> var. <i>prunifolia</i>	465
<i>Eugenia virgultosa</i> var. <i>jamaicensis</i>	599	<i>Excoecaria biglandulosa</i> var. <i>salicifolia</i>	465
<i>Eugenia vismiifolia</i>	625	<i>Excoecaria biglandulosa</i> var. <i>serrata</i>	465
<i>Eugenia wallenii</i>	599	<i>Excoecaria guianensis</i>	463
<i>Eugenia wentii</i>	607	<i>Excoecaria ilicifolia</i>	585
<i>Eugenia willdenowii</i>	605	<i>Excoecaria marginata</i>	465, 466
<i>Eugenia wullschlaegeliana</i>	607	<i>Excoecaria marginata</i> f. <i>major</i>	465
<i>Eugenia xylopifolia</i>	599	<i>Excoecaria marginata</i> var. <i>conjugens</i>	466
<i>Eugenia xylopifolia</i> var. <i>brevipes</i>	599	<i>Excoecaria marginata</i> var. <i>grandifolia</i>	466
<i>Eugeniodes guianense</i>	709	<i>Excoecaria marginata</i> var. <i>lanceolata</i>	465
<i>Eugeniodes martinicense</i>	710	<i>Excoecaria marginata</i> var. <i>longifolia</i>	466
<i>Eugeniodes martinicense</i> var. <i>paraguayense</i>	710	<i>Excoecaria marginata</i> var. <i>spathulata</i>	465
<i>Eugeniodes paraense</i>	709	<i>Excoecaria obtusiloba</i>	465
<i>Eugeniopsis richardiana</i>	612	<i>Excoecaria occidentalis</i>	466
<i>Eumachia boliviiana</i>	656	<i>Excoecaria suberosa</i>	466
<i>Euosma latifolia</i>	713	<i>Exelloidendron barbatum</i>	419
EUPHORBIACEAE	457	<i>Exostema souzanum</i>	655

Appendix 3. — Continuation.

F			
FABACEAE	503	Faramea egensis	657
<i>Fagara acreana</i>	668	<i>Faramea hondureae</i>	657
<i>Fagara acutifolia</i>	670	<i>Faramea laxula</i>	657
<i>Fagara amapaense</i>	668	<i>Faramea longifolia</i>	658
<i>Fagara apiculata</i>	669	<i>Faramea longifolia</i> var. <i>petiolaris</i>	658
<i>Fagara astrigera</i>	671	<i>Faramea martinii</i>	654
<i>Fagara caribaea</i>	669	<i>Faramea maynensis</i>	657
<i>Fagara coco</i> var. <i>formosana</i>	—	<i>Faramea multiflora</i>	657
<i>Fagara dellomei</i>	669	<i>Faramea multiflora</i> var. <i>amazonica</i>	657
<i>Fagara ekmanii</i>	669	<i>Faramea multiflora</i> var. <i>benensis</i>	657
<i>Fagara elephantiasis</i>	669	<i>Faramea multiflora</i> var. <i>epedunculata</i>	657
<i>Fagara guianensis</i>	669	<i>Faramea multiflora</i> var. <i>maynensis</i>	657
<i>Fagara idae</i>	668	<i>Faramea multiflora</i> var. <i>salicifolia</i>	657
<i>Fagara juniperina</i>	668	<i>Faramea occidentalis</i>	657
<i>Fagara luizii</i>	669	<i>Faramea occidentalis</i> subsp. <i>lonchocalyx</i>	657
<i>Fagara machadoi</i>	668	<i>Faramea occidentalis</i> var. <i>brachycalyx</i>	657
<i>Fagara microcarpa</i>	670	<i>Faramea occidentalis</i> var. <i>meridionalis</i>	657
<i>Fagara minutiflora</i>	668	<i>Faramea odoratissima</i>	657
<i>Fagara obscura</i>	670	<i>Faramea parviflora</i>	657
<i>Fagara occidentalis</i>	669	<i>Faramea pedunculata</i>	658
<i>Fagara ocumarensis</i>	668	<i>Faramea pendula</i>	656
<i>Fagara paulae</i>	669	<i>Faramea planitiarum</i>	658
<i>Fagara pentandra</i>	669	<i>Faramea polytriadophora</i>	658
<i>Fagara perrottetii</i>	669	<i>Faramea salicifolia</i>	657
<i>Fagara pringlei</i>	668	<i>Faramea salicifolia</i> f. <i>paniculata</i>	657
<i>Fagara procera</i>	668	<i>Faramea salicifolia</i> f. <i>subumbellata</i>	657
<i>Fagara pubescens</i>	670	<i>Faramea sertulifera</i>	657
<i>Fagara regnelliana</i>	671	<i>Faramea sessiliflora</i> var. <i>pedunculata</i>	658
<i>Fagara regnelliana</i> var. <i>calvata</i>	671	<i>Faramea sessiliflora</i>	658
<i>Fagara rhoifolia</i>	669-671	<i>Faramea standleyana</i>	657
<i>Fagara rhoifolia</i> f. <i>angustifolia</i>	671	<i>Faramea stenura</i>	657
<i>Fagara rhoifolia</i> f. <i>intermedia</i>	671	<i>Faramea talamancarum</i>	657
<i>Fagara rhoifolia</i> f. <i>latifolia</i>	671	<i>Faramea talamancarum</i> f. <i>acutifolia</i>	657
<i>Fagara rhoifolia</i> f. <i>paucijuga</i>	671	<i>Faramea tenuifolia</i>	657
<i>Fagara rhoifolia</i> subsp. <i>pubescens</i>	670	<i>Faramea truncata</i>	656
<i>Fagara rhoifolia</i> var. <i>intermedia</i>	671	<i>Faramea zetekii</i>	657
<i>Fagara rhoifolia</i> var. <i>peltophora</i>	670	<i>Faramea</i> sp. A	659
<i>Fagara rhoifolia</i> var. <i>petiolulata</i>	670	<i>Ferdinandusa goudotiana</i>	659
<i>Fagara rhoifolia</i> var. <i>surparanaensis</i>	671	<i>Ferdinandusa goudotiana</i> var. <i>eciliata</i>	659
<i>Fagara rothschuhii</i>	671	<i>Ferdinandusa goudotiana</i> var. <i>psilocarpa</i>	659
<i>Fagara ruiziana</i>	670	<i>Ferdinandusa paraensis</i>	659
<i>Fagara trinitensis</i>	669	<i>Ferdinandusa paraensis</i> var. <i>palustris</i>	659
<i>Fagara warmingiana</i>	668	<i>Ferdinandusa</i> sp. A	659
<i>Faramea albescens</i>	653	<i>Ferolia amazonica</i>	431
<i>Faramea amazonica</i>	657	<i>Ferolia campestris</i>	430
<i>Faramea belizensis</i>	657	<i>Ferolia excelsa</i>	431
<i>Faramea benensis</i>	657	<i>Ferolia guianensis</i>	584
<i>Faramea coarinensis</i>	658	<i>Ferolia montana</i>	431
<i>Faramea corymbosa</i>	656	<i>Feuilleea acrocephala</i>	516
<i>Faramea costata</i>	658	<i>Feuilleea affinis</i>	526
<i>Faramea cuatrecasasii</i>	657	<i>Feuilleea aggregata</i>	517
<i>Faramea cuencana</i>	657	<i>Feuilleea alba</i>	517
		<i>Feuilleea bahiensis</i>	520
		<i>Feuilleea bourgonii</i>	517

Appendix 3. — Continuation.

<i>Feuilleea brachyptera</i>	521	<i>Feuilleea stipularis</i>	525
<i>Feuilleea bracteosa</i>	521	<i>Feuilleea strigillosa</i>	524
<i>Feuilleea capitata</i>	517	<i>Feuilleea tenuiflora</i>	507
<i>Feuilleea cataractae</i>	545	<i>Feuilleea tenuifolia</i>	518
<i>Feuilleea catharinæ</i>	525	<i>Feuilleea terminalis</i>	505
<i>Feuilleea cayennensis</i>	518	<i>Feuilleea thibaudiana</i>	526
<i>Feuilleea conferta</i>	519	<i>Feuilleea trinervia</i>	507
<i>Feuilleea coriacea</i>	525	<i>Feuilleea umbellifera</i>	526
<i>Feuilleea corymbosa</i>	515	<i>Feuilleea umbratrica</i>	526
<i>Feuilleea cylindrica</i>	518	<i>Feuilleea uraguensis</i>	526
<i>Feuilleea disticha</i>	518	<i>Feuilleea venosa</i>	519
<i>Feuilleea divaricata</i>	545	<i>Feuilleea virgultosa</i>	527
<i>Feuilleea edulis</i>	518	<i>Ficus acarouaniensis</i>	587
<i>Feuilleea fasciculata</i>	507	<i>Ficus ajahuensis</i>	591
<i>Feuilleea fastuosa</i>	519	<i>Ficus albert-smithii</i>	585
<i>Feuilleea flagelliformis</i>	519	<i>Ficus alvareziana</i>	591
<i>Feuilleea graciliflora</i>	519	<i>Ficus amazonica</i>	585
<i>Feuilleea heterophylla</i>	520	<i>Ficus americana</i> subsp. <i>greiffiana</i>	585
<i>Feuilleea ignota</i>	521	<i>Ficus americana</i> subsp. <i>guianensis</i>	586
<i>Feuilleea inaequalis</i>	545	<i>Ficus americana</i> subsp. <i>subapiculata</i>	586
<i>Feuilleea ingoides</i>	520	<i>Ficus anacardiifolia</i>	586
<i>Feuilleea jupunba</i>	503	<i>Ficus anguina</i>	589
<i>Feuilleea laeta</i>	504	<i>Ficus angustifolia</i>	585
<i>Feuilleea lateriflora</i>	520	<i>Ficus anthelmintica</i>	588
<i>Feuilleea latifolia</i>	545	<i>Ficus arctocarpa</i>	589
<i>Feuilleea leiocalycina</i>	520	<i>Ficus arpazusa</i>	590
<i>Feuilleea leptoloba</i>	524	<i>Ficus arukensis</i>	590
<i>Feuilleea longiflora</i>	529	<i>Ficus baccata</i>	590
<i>Feuilleea macrophylla</i>	521	<i>Ficus bopiana</i>	588
<i>Feuilleea marginata</i>	522	<i>Ficus broadwayi</i>	590
<i>Feuilleea mathewsoniana</i>	522	<i>Ficus caballina</i>	586
<i>Feuilleea meisneriana</i>	526	<i>Ficus cabusana</i>	589
<i>Feuilleea micradenia</i>	503	<i>Ficus catappifolia</i>	586
<i>Feuilleea myriantha</i>	526	<i>Ficus chaconiana</i>	589
<i>Feuilleea niopoides</i>	504	<i>Ficus chiribiquetensis</i>	586
<i>Feuilleea nobilis</i>	522	<i>Ficus citrifolia</i>	588
<i>Feuilleea nuda</i>	525	<i>Ficus clusiifolia</i>	586
<i>Feuilleea pedicellaris</i>	505	<i>Ficus complicata</i>	590
<i>Feuilleea pezizifera</i>	523	<i>Ficus corpulenta</i>	586
<i>Feuilleea pilosula</i>	523	<i>Ficus coybana</i>	588
<i>Feuilleea poeppigiana</i>	524	<i>Ficus cremersii</i>	586
<i>Feuilleea punctata</i>	524	<i>Ficus cyclophylla</i>	589
<i>Feuilleea quassiifolia</i>	523	<i>Ficus daphniphylla</i>	590
<i>Feuilleea rubiginosa</i>	524	<i>Ficus doliaria</i>	587
<i>Feuilleea rufinervis</i>	524	<i>Ficus donnell-smithii</i>	587
<i>Feuilleea salzmanniana</i>	525	<i>Ficus duartei</i>	587
<i>Feuilleea scabriuscula</i>	519	<i>Ficus duckeana</i>	587
<i>Feuilleea schomburgkii</i>	514	<i>Ficus duquei</i>	589
<i>Feuilleea sciadion</i>	526	<i>Ficus duquei</i> var. <i>obtusiloba</i>	589
<i>Feuilleea sessiliflora</i>	518	<i>Ficus elliptica</i>	590
<i>Feuilleea sesuya</i>	520	<i>Ficus erratica</i>	586
<i>Feuilleea setifera</i>	523	<i>Ficus erythrosticta</i>	590
<i>Feuilleea speciosa</i>	521	<i>Ficus euomphala</i>	591
<i>Feuilleea splendens</i>	525	<i>Ficus fagifolia</i>	591

Appendix 3. — Continuation.

<i>Ficus fanshawei</i>	591	<i>Ficus nymphoides</i>	589
<i>Ficus fasciculata</i>	590	<i>Ficus ob lanceolata</i>	586
<i>Ficus florenciana</i>	587	<i>Ficus ochroleuca</i>	590
<i>Ficus frondosa</i>	585	<i>Ficus orinocensis</i>	590
<i>Ficus garcesii</i>	591	<i>Ficus padifolia</i>	590
<i>Ficus gemina</i>	590	<i>Ficus pakkensis</i>	589
<i>Ficus glaucescens</i>	588	<i>Ficus palmicida</i>	591
<i>Ficus gleasonii</i>	586	<i>Ficus paludica</i>	589
<i>Ficus gomelleira</i>	587	<i>Ficus panamensis</i>	589
<i>Ficus grabhamii</i>	590	<i>Ficus panurensis</i>	589
<i>Ficus grandaeva</i>	588	<i>Ficus paraensis</i>	589
<i>Ficus greiffiana</i>	585	<i>Ficus parkeri</i>	588
<i>Ficus grenadensis</i>	586	<i>Ficus parkeriana</i>	586
<i>Ficus guadalajarana</i>	588	<i>Ficus peroblonga</i>	590
<i>Ficus guianensis</i>	586	<i>Ficus pertusa</i>	590
<i>Ficus halliana</i>	590	<i>Ficus peruviana</i>	590
<i>Ficus haughtii</i>	589	<i>Ficus picardae</i>	588
<i>Ficus hebetifolia</i>	588	<i>Ficus piresiana</i>	591
<i>Ficus hernandezii</i>	588	<i>Ficus planicostata</i>	590
<i>Ficus holosericea</i>	587	<i>Ficus plicato-ostiolata</i>	591
<i>Ficus hydrophila</i>	590	<i>Ficus plumieri</i>	588
<i>Ficus hypochrysea</i>	585	<i>Ficus popenoei</i> subsp. <i>malacocarpa</i>	591
<i>Ficus ierensis</i>	589	<i>Ficus populnea</i> f. <i>citrifolia</i>	588
<i>Ficus insipida</i> subsp. <i>scabra</i>	588	<i>Ficus populnea</i> f. <i>planicostata</i>	590
<i>Ficus involuta</i> var. <i>urbaniana</i>	589	<i>Ficus prinoides</i> var. <i>subtriplinervia</i>	590
<i>Ficus juruensis</i>	591	<i>Ficus protensa</i>	588
<i>Ficus kanukuensis</i>	591	<i>Ficus pseudoradula</i>	588
<i>Ficus krukovii</i>	588	<i>Ficus putumayonis</i>	589
<i>Ficus lancifolia</i>	590	<i>Ficus radicans</i>	590
<i>Ficus lanjouwii</i>	588	<i>Ficus radula</i>	588
<i>Ficus laurifolia</i>	588	<i>Ficus ramiflora</i>	586
<i>Ficus leiophylla</i>	588	<i>Ficus regularis</i>	591
<i>Ficus leucosticta</i>	589	<i>Ficus rolanderi</i>	590
<i>Ficus llanensis</i>	591	<i>Ficus rubricosta</i>	588
<i>Ficus llewelynii</i>	591	<i>Ficus sapida</i>	590
<i>Ficus maguirei</i>	589	<i>Ficus savannarum</i>	591
<i>Ficus malacocarpa</i>	591	<i>Ficus scabrida</i>	591
<i>Ficus maroana</i>	586	<i>Ficus schumacheri</i>	591
<i>Ficus maroniensis</i>	588	<i>Ficus sodiroi</i>	589
<i>Ficus martinii</i>	586	<i>Ficus sonorae</i>	590
<i>Ficus mathewsi</i>	586	<i>Ficus splendens</i>	586
<i>Ficus maxima</i>	588	<i>Ficus sprucei</i>	586
<i>Ficus mendelsonii</i>	591	<i>Ficus subapiculata</i>	586
<i>Ficus mensalis</i>	586	<i>Ficus subscabrida</i>	588
<i>Ficus metensis</i>	586	<i>Ficus subtriplinervia</i>	590
<i>Ficus mexicana</i>	588	<i>Ficus suffocans</i>	590
<i>Ficus morantensis</i>	590	<i>Ficus sulcipes</i>	590
<i>Ficus murilloi</i>	589	<i>Ficus surinamensis</i>	585
<i>Ficus murilloi</i> var. <i>cajambrensis</i>	589	<i>Ficus tamatamae</i>	586
<i>Ficus myriasycea</i>	586	<i>Ficus tapajozensis</i>	591
<i>Ficus myrmecophila</i>	589	<i>Ficus thelephora</i>	589
<i>Ficus myrtifolia</i>	590	<i>Ficus trachelosyce</i>	591
<i>Ficus niceforoi</i>	586	<i>Ficus trigona</i>	591
<i>Ficus nymphaeifolia</i>	589	<i>Ficus turbinata</i>	590

Appendix 3. — Continuation.

<i>Ficus uberrima</i>	589	<i>Gardenia oblongifolia</i>	659
<i>Ficus ulei</i>	589	<i>Gardenia suaveolens</i>	663
<i>Ficus umbonigera</i>	586	<i>Gardenia tetricantha</i>	665
<i>Ficus urbaniana</i>	589	<i>Garuga gigantea</i>	366
<i>Ficus vaupesana</i>	586	<i>Garuga guianensis</i>	366
<i>Ficus vicencionis</i>	589	<i>Garuga schomburgkiana</i>	366
<i>Ficus vulpina</i>	591	<i>Garuga schomburgkiana</i> var. <i>salzmanniana</i>	366
<i>Ficus weberbaueri</i>	591	<i>Garuga spruceana</i>	366
<i>Ficus williamsii</i>	589	<i>Gaulettia canomensis</i>	419
<i>Ficus wuliana</i>	591	<i>Gaulettia elata</i>	419
<i>Firensia hirsuta</i>	445	<i>Gaulettia parillo</i>	419
<i>Firensia lutea</i>	443	<i>Geanthemum cadavericum</i>	371
<i>Flacourtie benthamii</i>	679	<i>Geissospermum argenteum</i>	382
<i>Flacourtie digyna</i>	679	<i>Geissospermum excelsum</i>	381
<i>Flacourtie nitida</i>	679	<i>Geissospermum fuscum</i>	384
<i>Folianthera guianensis</i>	538	<i>Geissospermum laeve</i>	383
<i>Fothergilla mirabilis</i>	565, 566	<i>Geissospermum sericeum</i>	384
<i>Franchetella anibifolia</i>	701	<i>Geissospermum vellozii</i>	383
<i>Franchetella gonggrijpii</i>	698	<i>Gelsemium araliaceum</i>	396
<i>Franchetella platyphylla</i>	700	<i>Gelsemium avellaneda</i>	395
<i>Franchetella reticulata</i>	700	<i>Gelsemium insigne</i>	398
<i>Franchetella unilocularis</i>	700	<i>Gelsemium speciosum</i>	396
<i>Froelichia</i>	655	<i>Genipa americana</i> f. <i>grandifolia</i>	659
<i>Froelichia paniculata</i>	654	<i>Genipa americana</i> f. <i>jorgensenii</i>	659
<i>Froesiodendron surinamense</i>	370	<i>Genipa americana</i> f. <i>parvifolia</i>	659
<i>Fusaea decurrents</i>	372	<i>Genipa americana</i> var. <i>americana</i>	659
<i>Fusaea longifolia</i>	372	<i>Genipa americana</i> var. <i>caruto</i>	659
<i>Fusaea rhombipetala</i>	372	<i>Genipa barbata</i>	659
		<i>Genipa caruto</i>	659
		<i>Genipa codonocalyx</i>	659
		<i>Genipa edulis</i>	650
		<i>Genipa eriopila</i>	655
		<i>Genipa excelsa</i>	659
		<i>Genipa grandifolia</i>	659
		<i>Genipa humilis</i>	659
		<i>Genipa merianae</i>	655
		<i>Genipa oblongifolia</i>	659
		<i>Genipa pubescens</i>	659
		<i>Genipa spruceana</i>	659
		<i>Geoffroea acutifolia</i>	506
		<i>Geoffroea discolor</i>	513
		<i>Geoffroea inermis</i>	506
		<i>Geoffroea obtusifolia</i>	506
		<i>Geoffroea pubescens</i>	506
		<i>Geoffroea retusa</i>	506
		<i>Geoffroea surinamensis</i>	506
		<i>Geonoma baculifera</i>	353
		<i>Gerascanthus alliodorus</i>	444
		<i>Gerascanthus bicolor</i>	444
		<i>Gerascanthus calophyllus</i>	445
		<i>Gerascanthus cordifolius</i>	446
		<i>Gerascanthus exaltatus</i>	444
		<i>Gerascanthus goeldianus</i>	445
		<i>Gerascanthus heterophyllus</i>	447

Appendix 3. — Continuation.

<i>Gerascanthus hirtus</i>	446	<i>Guapeba laevigata</i>	699
<i>Gerascanthus lomatolobus</i>	445	<i>Guapeba lasiocarpa</i>	695
<i>Gerascanthus muneco</i>	446	<i>Guapeba laurifolia</i>	694
<i>Gerascanthus naidophilus</i>	445	<i>Guapeba parinariooides</i>	698
<i>Gerascanthus opacus</i>	446	<i>Guapira broadwayana</i>	627
<i>Gerascanthus panicularis</i>	445	<i>Guapira eggersiana</i>	627
<i>Gerascanthus pubescens</i>	447	<i>Guapira guianensis</i>	627
<i>Gerascanthus sagotii</i>	446	<i>Guapiraheimeriana</i>	627
<i>Gerascanthus scabridus</i>	444	<i>Guapira salicifolia</i>	627
<i>Gerascanthus sericalyx</i>	446	<i>Guapira</i> sp. A	627
<i>Gerascanthus sprucei</i>	446	<i>Guarania laurifolia</i>	638
<i>Gerascanthus tetrandrus</i>	446	<i>Guarania ramiflora</i>	638
<i>Gerascanthus tetraphyllus</i>	443	<i>Guarea</i>	575
<i>Gerascanthus toqueve</i>	447	<i>Guarea affinis</i>	578
<i>Gerascanthus viridis</i>	446	<i>Guarea alba</i>	576
<i>Gimbernacia oblonga</i>	443	<i>Guarea alternans</i>	576
<i>Gimbernacia obovata</i>	440	<i>Guarea andreana</i>	576
<i>Gloeospermum</i>	718	<i>Guarea aubletii</i>	575, 576
<i>Gloeospermum sphaerocarpum</i>	718, 719	<i>Guarea bahiensis</i>	575
<i>Gloeospermum sphaerocarpum</i> var. <i>latifolium</i>	719	<i>Guarea bangii</i>	577
<i>Gloeospermum ulei</i>	721	<i>Guarea bilibil</i>	576
<i>Glossocentrum collinum</i>	565	<i>Guarea bilocularis</i>	579
<i>Glossopetalum glabrum</i>	467	<i>Guarea borisii</i>	578
<i>Glossopetalum tomentosum</i>	467	<i>Guarea brachystachya</i>	575
<i>Glycoxylon huberi</i>	703	<i>Guarea cabirme</i>	576
<i>Glycydendron amazonicum</i>	462	<i>Guarea campestris</i>	576
<i>Glycyrrhiza undulata</i>	506	<i>Guarea carinata</i>	575
<i>Goeppertia sericea</i> var. <i>bracteolata</i>	479	<i>Guarea chichon</i>	578
<i>Gomidesia minutiflora</i>	609, 610	<i>Guarea cinnamomea</i>	575
<i>Gomphia candollei</i>	629	<i>Guarea concinna</i>	578
<i>Gordonia fruticosa</i>	710	<i>Guarea convergens</i>	575
<i>Gordonia humboldtii</i>	710	<i>Guarea costata</i>	575
<i>Gordonia planchonii</i>	710	<i>Guarea costulata</i>	577
<i>Gordonia semiserrata</i>	710	<i>Guarea culebrana</i>	578
<i>Gossampinus alba</i>	552	<i>Guarea davisii</i>	578
<i>Gossampinus rumphii</i>	552	<i>Guarea densiflora</i>	576
<i>GOUPIACEAE</i>	467	<i>Guarea depauperata</i>	576
<i>Gouphia glabra</i>	467	<i>Guarea duckei</i>	579
<i>Gouphia paraensis</i>	467	<i>Guarea eggersii</i>	576
<i>Gouphia tomentosa</i>	467	<i>Guarea francavillana</i>	576
<i>Granatum guianense</i>	573	<i>Guarea gigantea</i>	578
<i>Grangeria brasiliensis</i>	420	<i>Guarea glauca</i>	576
<i>Greeneina affinis</i>	592	<i>Guarea gomma</i>	575
<i>Greeneina poeppigiana</i>	592	<i>Guarea grandiflora</i>	578
<i>Grias tetrapetala</i>	499	<i>Guarea grandifolia</i>	578
<i>Guaiabara uvifera</i>	643	<i>Guarea grandifoliola</i>	576
<i>Guajava acutangula</i>	625	<i>Guarea guara</i>	575
<i>Guajava oligosperma</i>	626	<i>Guarea guedesii</i>	577
<i>Guanabanus palustris</i>	367	<i>Guarea guidonia</i>	575
<i>Guapeba brasiliensis</i>	695	<i>Guarea huberi</i>	578
<i>Guapeba caimito</i>	694	<i>Guarea huberi</i> var. <i>peruviana</i>	578
<i>Guapeba coriacea</i>	695	<i>Guarea klugii</i>	579
<i>Guapeba glazioviana</i>	698	<i>Guarea kunthiana</i>	576
<i>Guapeba glomerata</i>	698	<i>Guarea kunthiana</i> var. <i>densiflora</i>	576

Appendix 3. — Continuation.

<i>Guarea kunthiana</i> var. <i>hahniana</i>	576	<i>Guarea trichiliooides</i> var. <i>colombiana</i>	576
<i>Guarea kunthiana</i> var. <i>hahnii</i>	576	<i>Guarea trichiliooides</i> var. <i>decandra</i>	576
<i>Guarea leticiana</i>	576	<i>Guarea trichiliooides</i> var. <i>pachycarpa</i>	577
<i>Guarea longipetiolata</i>	578	<i>Guarea trichiliooides</i> var. <i>pallida</i>	576
<i>Guarea macrantha</i>	577	<i>Guarea trompillo</i>	578
<i>Guarea macrophylla</i>	577	<i>Guarea trunciflora</i>	579
<i>Guarea macrophylla</i> subsp. <i>pachycarpa</i>	577	<i>Guarea tuberculata</i> var. <i>purgans</i>	575
<i>Guarea mancharra</i>	578	<i>Guarea ulei</i>	578
<i>Guarea megalantha</i>	578	<i>Guarea williamsii</i>	576
<i>Guarea megantha</i>	578	<i>Guarea xiorensana</i>	576
<i>Guarea megaphylla</i>	577	<i>Guatteria aberemoa</i>	373
<i>Guarea membranacea</i>	576	<i>Guatteria aberemoa</i> var. <i>microcarpa</i>	373
<i>Guarea microsepala</i>	578	<i>Guatteria acutissima</i>	376
<i>Guarea mucronulata</i>	573	<i>Guatteria anteridifera</i>	373
<i>Guarea multiflora</i>	575	<i>Guatteria anthracina</i>	374
<i>Guarea multijuga</i>	575	<i>Guatteria asplundiana</i>	376
<i>Guarea oblongiflora</i>	576	<i>Guatteria atra</i>	376
<i>Guarea obtusifolia</i>	580	<i>Guatteria axilliflora</i>	374
<i>Guarea paraensis</i>	577	<i>Guatteria blepharophylla</i>	373
<i>Guarea parva</i>	576	<i>Guatteria brevipes</i>	370, 371
<i>Guarea pedicellata</i>	579	<i>Guatteria buchtienii</i>	376
<i>Guarea pittieri</i>	578	<i>Guatteria calimensis</i>	376
<i>Guarea poeppigii</i>	576	<i>Guatteria calliantha</i>	376
<i>Guarea pohlii</i>	576	<i>Guatteria calophylla</i>	373
<i>Guarea pohlii</i> var. <i>glabra</i>	576	<i>Guatteria caniflora</i>	374, 375
<i>Guarea puberula</i>	576	<i>Guatteria caniflora</i> var. <i>angustifolia</i>	375
<i>Guarea pubescens</i> subsp. <i>pubescens</i>	578	<i>Guatteria caniflora</i> var. <i>latifolia</i>	375
<i>Guarea pubescens</i> subsp. <i>pubiflora</i>	578	<i>Guatteria cargadero</i>	375
<i>Guarea pubiflora</i>	578	<i>Guatteria chrysopetala</i>	375, 376
<i>Guarea pubiflora</i> var. <i>angustifoliola</i>	578	<i>Guatteria chrysopetala</i> var. <i>major</i>	376
<i>Guarea pubiflora</i> var. <i>parvifolia</i>	578	<i>Guatteria chrysopetala</i> var. <i>tenuipeps</i>	376
<i>Guarea punctata</i>	577	<i>Guatteria citriodora</i>	373
<i>Guarea purgans</i>	575	<i>Guatteria coeloneura</i>	376
<i>Guarea quadrangularis</i>	575	<i>Guatteria collina</i>	376
<i>Guarea racemiformis</i>	576	<i>Guatteria conspicua</i>	373
<i>Guarea rhabdotocarpa</i>	578	<i>Guatteria crassipetala</i>	374
<i>Guarea richardiana</i>	578	<i>Guatteria cylindrocarpa</i>	373
<i>Guarea rosea</i>	577	<i>Guatteria dielsiana</i>	373
<i>Guarea rubescens</i>	576	<i>Guatteria discolor</i>	373
<i>Guarea rubricalyx</i>	576	<i>Guatteria duckeana</i> var. <i>subcordata</i>	373
<i>Guarea rubrisepala</i>	576	<i>Guatteria elliptica</i>	376
<i>Guarea rusbyi</i>	576	<i>Guatteria excellens</i>	373
<i>Guarea scabra</i>	578	<i>Guatteria flavovirens</i>	377
<i>Guarea silvatica</i>	579	<i>Guatteria foliosa</i>	373
<i>Guarea simplicifolia</i>	576	<i>Guatteria gamosepala</i>	376
<i>Guarea spicata</i>	578	<i>Guatteria glauca</i>	374
<i>Guarea steinbachii</i>	576	<i>Guatteria gracilipes</i>	376
<i>Guarea subsessiliflora</i>	577	<i>Guatteria guentheri</i>	376
<i>Guarea subsessiliflora</i> var. <i>polyphyllaria</i>	577	<i>Guatteria guianensis</i>	373
<i>Guarea surinamensis</i>	576	<i>Guatteria hyposericea</i>	377
<i>Guarea sylvestris</i>	576	<i>Guatteria insignis</i>	377
<i>Guarea trichiliooides</i>	575-577	<i>Guatteria intermedia</i>	374
<i>Guarea trichiliooides</i> var. <i>brachystachya</i>	576	<i>Guatteria juninensis</i>	376

Appendix 3. — Continuation.

<i>Guatteria krukoffii</i>	377	<i>Guatteria vestita</i>	376
<i>Guatteria lanceolata</i>	376	<i>Guatteria vestita</i> var. <i>angustifolia</i>	376
<i>Guatteria lasiocalyx</i>	376	<i>Guatteria vestita</i> var. <i>latifolia</i>	376
<i>Guatteria latipetala</i>	376	<i>Guatteria wachenheimii</i>	377
<i>Guatteria leiocarpa</i>	376	<i>Guatteria wessels-boerii</i>	376
<i>Guatteria leucotricha</i>	374	<i>Guatteriopsis blepharophylla</i>	373
<i>Guatteria liesneri</i>	374	<i>Guatteriopsis sessiliflora</i>	373
<i>Guatteria longestipitata</i>	376	<i>Guazuma blumei</i>	554
<i>Guatteria macrocalyx</i>	376	<i>Guazuma bubroma</i>	554
<i>Guatteria micans</i>	377	<i>Guazuma coriacea</i>	554
<i>Guatteria microsperma</i>	377	<i>Guazuma guazuma</i>	553
<i>Guatteria montis-trinitatis</i>	376	<i>Guazuma guazuma</i> var. <i>tomentosa</i>	554
<i>Guatteria multiflora</i>	370	<i>Guazuma guazuma</i> var. <i>ulmifolia</i>	553
<i>Guatteria multivenia</i>	373	<i>Guazuma invira</i>	553
<i>Guatteria obliqua</i>	376	<i>Guazuma parvifolia</i>	554
<i>Guatteria oblonga</i>	374	<i>Guazuma polybotrya</i>	553
<i>Guatteria occidentalis</i>	376	<i>Guazuma tomentosa</i>	553, 554
<i>Guatteria olivacea</i>	376	<i>Guazuma tomentosa</i> var. <i>cumanensis</i>	554
<i>Guatteria ouregou</i>	374	<i>Guazuma tomentosa</i> var. <i>monpoxensis</i>	554
<i>Guatteria ouregou</i> var. <i>latifolia</i>	374	<i>Guazuma tomentosa</i> var. <i>parvifolia</i>	554
<i>Guatteria ovalifolia</i>	376	<i>Guazuma ulmifolia</i>	553, 554
<i>Guatteria oxycarpa</i>	373	<i>Guazuma ulmifolia</i> var. <i>glabra</i>	554
<i>Guatteria pannosa</i>	374	<i>Guazuma ulmifolia</i> var. <i>tomentella</i>	554
<i>Guatteria paraensis</i>	373	<i>Guazuma ulmifolia</i> var. <i>tomentosa</i>	553
<i>Guatteria parviflora</i>	376	<i>Guazuma ulmifolia</i> var. <i>trianae</i>	554
<i>Guatteria peduncularis</i>	378	<i>Guazuma ulmifolia</i> var. <i>velutina</i>	554
<i>Guatteria platyphylla</i>	375	<i>Guazuma utilis</i>	554
<i>Guatteria pleiocarpa</i>	376	<i>Guenetia macrosperma</i>	552
<i>Guatteria podocarpa</i>	374	<i>Guettarda acreana</i>	666
<i>Guatteria podocarpa</i> var. <i>oligocarpa</i>	374	<i>Guettarda argentea</i>	660
<i>Guatteria podocarpa</i> var. <i>polycarpa</i>	374	<i>Guettarda argentea</i> var. <i>glabrata</i>	660
<i>Guatteria poeppigiana</i>	374	<i>Guettarda coccinea</i>	660
<i>Guatteria poiteaui</i>	370, 371	<i>Guettarda fanshawei</i>	666
<i>Guatteria pteropus</i>	375, 376	<i>Guettarda leiantha</i>	666
<i>Guatteria pteropus</i> var. <i>angustior</i>	376	<i>Guianodendron</i>	503, 515
<i>Guatteria pteropus</i> var. <i>cinerea</i>	376	<i>Guianodendron praeclarum</i>	515
<i>Guatteria punctata</i>	374	<i>Guidonia adstringens</i>	673
<i>Guatteria rhamnoides</i>	376	<i>Guidonia parvifolia</i>	672
<i>Guatteria richardii</i>	376	<i>Guidonia procera</i>	672
<i>Guatteria robusta</i>	373	<i>Guidonia ramiflora</i>	673
<i>Guatteria sagotiana</i>	376	<i>Guidonia stipularis</i>	672
<i>Guatteria sagotiana</i> var. <i>gracilior</i>	376	<i>Guidonia sylvestris</i>	675
<i>Guatteria sandwithii</i>	376	<i>Guidonia ulmifolia</i>	676
<i>Guatteria schomburgkiana</i>	376	<i>Gustavia angusta</i>	499
<i>Guatteria schomburgkiana</i> var. <i>angustifolia</i>	376	<i>Gustavia antillana</i>	499
<i>Guatteria schomburgkiana</i> var. <i>holosericea</i>	376	<i>Gustavia augusta</i>	499
<i>Guatteria schomburgkiana</i> var. <i>latifolia</i>	376	<i>Gustavia augusta</i> var. <i>brasiliensis</i>	499
<i>Guatteria scytophylla</i>	377	<i>Gustavia augusta</i> var. <i>calycaris</i>	499
<i>Guatteria sessiliflora</i>	373	<i>Gustavia augusta</i> var. <i>guianensis</i>	499
<i>Guatteria sessilis</i>	376	<i>Gustavia augusta</i> var. <i>verrucosa</i>	499
<i>Guatteria spruceana</i>	376	<i>Gustavia brasiliiana</i>	500
<i>Guatteria sylvicola</i>	376	<i>Gustavia brasiliiana</i> var. <i>minor</i>	500
<i>Guatteria ucayaliana</i>	373	<i>Gustavia brasiliensis</i>	499
<i>Guatteria umbonata</i>	376	<i>Gustavia calycaris</i>	499

Appendix 3. — Continuation.

<i>Gustavia eximia</i>	500	<i>Hasseltia pubescens</i>	677
<i>Gustavia fastuosa</i>	499, 500	<i>Hasseltia rigida</i>	677
<i>Gustavia fastuosa</i> var. <i>angustisepala</i>	500	<i>Hasseltia tomentulosa</i>	677
<i>Gustavia fastuosa</i> var. <i>latisepala</i>	500	<i>Havetia flava</i>	432
<i>Gustavia fustis-mortui</i>	500	<i>Havetia flexilis</i>	432
<i>Gustavia hexapetala</i>	499	<i>Havetia octandra</i>	433
<i>Gustavia insignis</i>	499	<i>Havetiopsis</i> sect. <i>Havetiopsis</i>	432
<i>Gustavia laciniosa</i>	499	<i>Havetiopsis flava</i>	432
<i>Gustavia marcgraaviana</i>	499	<i>Havetiopsis flexilis</i>	432
<i>Gustavia meizocarpa</i>	499	<i>Hebepeplatum humiriifolium</i>	546
<i>Gustavia membrillo</i>	499	<i>Hedera capitata</i>	390
<i>Gustavia mexicana</i>	499	<i>Hedera frondosa</i>	390
<i>Gustavia microcarpa</i>	500	<i>Hedwigia hostmannii</i>	405
<i>Gustavia poeppigiana</i> var. <i>rigida</i>	499	<i>Hedwigia panamensis</i>	404
<i>Gustavia pterocarpa</i>	500	<i>Hedwigia rhoifolia</i>	404, 407
<i>Gustavia tetrapetala</i>	499	<i>Hedwigia simplicifolia</i>	580
<i>Gustavia theophrasta</i>	499	<i>Hedycrea incana</i>	428
<i>Gustavia urceolata</i>	499	<i>Heisteria barbata</i>	633
<i>Gymnobalanus fendleri</i>	485	<i>Heisteria densifrons</i>	633
 H			
<i>Habzelia aromatic</i> a	379	<i>Heisteria flexuosa</i>	634
<i>Habzelia cubensis</i>	379	<i>Heisteria kappleri</i>	633
<i>Habzelia discreta</i>	380	<i>Heisteria krukovi</i>	634
<i>Haemocharis camellioides</i>	710	<i>Heisteria micrantha</i>	634
<i>Haemocharis caracasana</i>	710	<i>Heisteria microcalyx</i>	633, 634
<i>Haemocharis intermedia</i>	710	<i>Heisteria microcarpa</i>	633
<i>Haemocharis parviflora</i>	710	<i>Heisteria nitida</i>	634
<i>Haemocharis praemorsa</i>	710	<i>Heisteria ovata</i>	634
<i>Haemocharis pubescens</i>	710	<i>Heisteria parvicalyx</i>	634
<i>Haemocharis semiserrata</i>	710	<i>Heisteria rubricalyx</i>	634
<i>Haemocharis symplocoidea</i>	710	<i>Heisteria sessilis</i>	634
<i>Hamelia glabra</i>	650	<i>Heisteria subsessilis</i>	634
<i>Hamelia sessiliflora</i>	650	<i>Heisteria surinamensis</i>	634
<i>Hancornia amapa</i>	386	<i>Heisteria vageleri</i>	634
<i>Handroanthus araliaceus</i>	396	<i>Hekkingia bordenavei</i>	719
<i>Handroanthus atractocarpus</i>	396	<i>Helicosstylis affinis</i>	592
<i>Handroanthus avellanedae</i>	395	<i>Helicosstylis duckei</i>	592
<i>Handroanthus capitatus</i>	395	<i>Helicosstylis elegans</i>	592
<i>Handroanthus durus</i>	398	<i>Helicosstylis obtusifolia</i>	592
<i>Handroanthus flavescens</i>	396	<i>Helicosstylis pedunculata</i>	591
<i>Handroanthus impetiginosus</i>	395	<i>Helicosstylis podogyne</i>	592
<i>Handroanthus obscurus</i>	395	<i>Helicosstylis poeppigiana</i>	592
<i>Handroanthus serratifolius</i>	396	<i>Helicosstylis poeppigiana</i> var. <i>macrophylla</i>	592
<i>Handroanthus subtilis</i>	396	<i>Helicosstylis tormentosa</i>	592
<i>Hasseltia floribunda</i>	676	<i>Helicosstylis</i> sp. A	592
<i>Hasseltia laxiflora</i>	677	<i>Hemicrepidiospermum rhoifolium</i>	404
<i>Hasseltia micrantha</i>	677	<i>Henriettea brasiliensis</i>	561
<i>Hasseltia monagensis</i>	677	<i>Henriettea brunnescens</i>	560
<i>Hasseltia panamensis</i>	677	<i>Henriettea duckeana</i>	560
<i>Hasseltia peruviana</i>	677	<i>Henriettea flavescentia</i>	560
<i>Hasseltia psittacarum</i>	677	<i>Henriettea maroniensis</i>	561
		<i>Henriettea multiflora</i>	561
		<i>Henriettea parviflora</i>	561
		<i>Henriettea patrisiana</i>	561
		<i>Henriettea ramiflora</i>	561

Appendix 3. — Continuation.

<i>Henriettea stellaris</i>	561	<i>Himatanthus phagedaenicus</i>	384
<i>Henriettea succosa</i>	561	<i>Himatanthus rigidus</i>	384
<i>Henriettea succosa</i> var. <i>guianensis</i>	561	<i>Himatanthus speciosus</i>	384
<i>Henriettea surinamensis</i>	561	<i>Himatanthus sucuuba</i>	384
<i>Henriettea trinervia</i>	561	<i>Himatanthus tarapotensis</i>	384
<i>Henriettea williamii</i>	561	<i>Hippomane aucuparia</i>	464
<i>Henriettella duckeana</i>	560	<i>Hippomane biglandulosa</i>	464, 467
<i>Henriettella flavescens</i>	560	<i>Hippomane glandulosa</i>	464
<i>Henriettella parviflora</i>	561	<i>Hippomane zeocca</i>	466
<i>Henriettella patrisiana</i>	561	<i>Hirtella acayacensis</i>	421
<i>Henriettella ramiflora</i>	561	<i>Hirtella aggregata</i>	421
HERNANDIACEAE	467	<i>Hirtella americana</i> var. <i>hexandra</i>	421
<i>Hernandia guianensis</i>	467	<i>Hirtella americana</i> var. <i>oblongifolia</i>	421
<i>Herrania guianensis</i>	559	<i>Hirtella apetala</i>	425
<i>Hevea guianensis</i>	462	<i>Hirtella araguariensis</i>	419
<i>Hevea peruviana</i>	462	<i>Hirtella bicornis</i> var. <i>bicornis</i>	419
<i>Hexactina corymbosa</i>	650	<i>Hirtella bicornis</i> var. <i>pubescens</i>	420
<i>Heymassoli inermis</i>	634, 635	<i>Hirtella bracteata</i>	422
<i>Heymassoli spinosa</i>	634, 635	<i>Hirtella bracteosa</i>	421
<i>Hibiscus arboreus</i>	558	<i>Hirtella canomensis</i>	419
<i>Hibiscus bracteosus</i>	558	<i>Hirtella castanea</i>	422
<i>Hibiscus fragrantissimus</i>	558	<i>Hirtella caudata</i>	420
<i>Hibiscus pernambucensis</i>	558	<i>Hirtella ciliata</i>	420
<i>Hibiscus tiliaceus</i> f. <i>immaculatus</i>	558	<i>Hirtella coppenamensis</i>	421
<i>Hibiscus tiliaceus</i> subsp. <i>pernambucensis</i>	558	<i>Hirtella coriacea</i>	421
<i>Hibiscus tiliaceus</i> var. <i>pernambucensis</i>	558	<i>Hirtella cosmibuena</i>	422
<i>Hieronyma alchorneoides</i>	637	<i>Hirtella damaziana</i>	420
<i>Hieronyma alchorneoides</i> var. <i>stipulosa</i>	637	<i>Hirtella davisii</i>	420
<i>Hieronyma andina</i>	637	<i>Hirtella egensis</i>	420
<i>Hieronyma blanchetiana</i>	637	<i>Hirtella filiformis</i>	421
<i>Hieronyma caribaea</i>	637	<i>Hirtella glandulostipula</i>	420
<i>Hieronyma choocoensis</i>	637	<i>Hirtella glandulosa</i>	420
<i>Hieronyma ferruginea</i>	637	<i>Hirtella guyanensis</i>	421
<i>Hieronyma gentlei</i>	638	<i>Hirtella hexandra</i>	421
<i>Hieronyma guatemalensis</i>	637	<i>Hirtella hirsuta</i>	421
<i>Hieronyma heterotricha</i>	637	<i>Hirtella hispidula</i>	420
<i>Hieronyma laxiflora</i>	637	<i>Hirtella hookeri</i>	420
<i>Hieronyma mattogrossensis</i>	637	<i>Hirtella indecora</i>	421
<i>Hieronyma mollis</i>	637	<i>Hirtella jamaicensis</i>	422
<i>Hieronyma oblonga</i>	637	<i>Hirtella lanceolata</i>	421
<i>Hieronyma oblonga</i> f. <i>glabra</i>	637	<i>Hirtella macrophylla</i>	420
<i>Hieronyma oblonga</i> var. <i>benthamii</i>	637	<i>Hirtella macrosepala</i>	420
<i>Hieronyma oblonga</i> var. <i>blanchetiana</i>	637	<i>Hirtella manigera</i>	422
<i>Hieronyma oblonga</i> var. <i>crassifolia</i>	637	<i>Hirtella margae</i>	421
<i>Hieronyma oblonga</i> var. <i>nervata</i>	637	<i>Hirtella melinonii</i>	422
<i>Hieronyma oblonga</i> var. <i>obtusa</i>	637	<i>Hirtella multiflora</i>	422
<i>Hieronyma ovatifolia</i>	637	<i>Hirtella nemorosa</i>	421
<i>Hieronyma poasana</i>	637	<i>Hirtella nitida</i>	421
<i>Hieronyma tectissima</i>	637	<i>Hirtella obidensis</i>	421
<i>Hilairanthus nitidus</i>	362	<i>Hirtella oblongifolia</i>	421
<i>Hilairanthus tomentosus</i>	362	<i>Hirtella octandra</i>	426
<i>Himatanthus articulatus</i>	384	<i>Hirtella paniculata</i>	421, 422
<i>Himatanthus bracteatus</i>	384	<i>Hirtella peruviana</i>	422
<i>Himatanthus lancifolius</i>	384	<i>Hirtella praealta</i>	419, 420

Appendix 3. — Continuation.

<i>Hirtella racemosa</i>	421, 422	<i>Horau racemosus</i>	440
<i>Hirtella racemosa</i> var. <i>hexandra</i>	421	<i>Hornschuchia caudata</i>	377
<i>Hirtella racemosa</i> var. <i>metallica</i>	421	<i>Hortia excelsa</i>	667
<i>Hirtella racemosa</i> var. <i>oblongifolia</i>	421	<i>Hortia superba</i>	667
<i>Hirtella racemosa</i> var. <i>racemosa</i>	421	<i>Hostmannia elvasiooides</i>	628
<i>Hirtella rosea</i>	421	<i>Hostmannia sagotii</i>	628
<i>Hirtella rotundata</i>	420	<i>Huberodendron swieteniooides</i>	555
<i>Hirtella rubra</i>	420	<i>Hudsonia arborea</i>	443
<i>Hirtella scandens</i>	421	<i>Humiria amplexicaulis</i>	467
<i>Hirtella silicea</i>	422	<i>Humiria arenaria</i>	467
<i>Hirtella subsetosa</i>	420	<i>Humiria balsamifera</i>	467
<i>Hirtella suffulta</i>	422	<i>Humiria balsamifera</i> f. <i>attenuata</i>	467
<i>Hirtella tenuifolia</i>	422	HUMIRIACEAE	467
<i>Hirtella triandra</i>	422	<i>Humiriastrum excelsum</i>	468
<i>Hirtella velutina</i>	420	<i>Humiriastrum subcrenatum</i>	468
<i>Hirtella violacea</i>	421	<i>Humiria subcrenata</i>	468
<i>Hirtella wachenheimii</i>	420	<i>Hura brasiliensis</i>	462
<i>Hisingera benthamii</i>	679	<i>Hura crepitans</i>	462
<i>Hisingera ciliatifolia</i>	679	<i>Hura crepitans</i> f. <i>oblongifolia</i>	462
<i>Hisingera lucens</i>	679	<i>Hura crepitans</i> f. <i>orbicularis</i>	462
<i>Homalium anzoateguiense</i>	677	<i>Hura crepitans</i> f. <i>ovata</i>	462
<i>Homalium choocoense</i>	677	<i>Hura crepitans</i> var. <i>genuina</i>	462
<i>Homalium columbianum</i>	677	<i>Hura crepitans</i> var. <i>membranacea</i>	462
<i>Homalium densiflorum</i>	677	<i>Hura crepitans</i> var. <i>senegalensis</i>	462
<i>Homalium eleutherostylum</i>	677	<i>Hura crepitans</i> var. <i>strepens</i>	462
<i>Homalium euryptetalum</i>	677	<i>Hura senegalensis</i>	462
<i>Homalium guianense</i>	677	<i>Hura strepens</i>	462
<i>Homalium hemistylum</i>	677	<i>Hybanthus surinamensis</i>	718
<i>Homalium hondurens</i> e	677	<i>Hydrochorea</i>	503, 515
<i>Homalium integrifolium</i>	677	<i>Hydrochorea corymbosa</i>	515
<i>Homalium leiogynum</i>	677	<i>Hymenaea</i>	503, 516
<i>Homalium mattogrossense</i>	677	<i>Hymenaea courbaril</i>	516
<i>Homalium mituense</i>	677	<i>Hymenaea venosa</i>	535
<i>Homalium mollicellum</i>	677	<i>Hymenaea</i> sp. A	516
<i>Homalium napimoga</i>	677	<i>Hymenolobium</i>	503, 516
<i>Homalium nicaraguense</i>	677	<i>Hymenolobium excelsum</i>	516
<i>Homalium obtusatum</i>	677	<i>Hymenolobium flavum</i>	516
<i>Homalium pedicellatum</i>	677	<i>Hymenolobium heterocarpum</i>	516
<i>Homalium pittieri</i>	677	<i>Hymenolobium petraeum</i>	516
<i>Homalium pleiandrum</i>	677	<i>Hymenolobium pulcherrimum</i>	516
<i>Homalium puberulum</i>	677	<i>Hymenopus amapaensis</i>	422
<i>Homalium racemosum</i>	677	<i>Hymenopus caudatus</i>	422
<i>Homalium racemosum</i> subsp. <i>barbellatum</i>	677	<i>Hymenopus divaricatus</i>	423
<i>Homalium racoubea</i>	677	<i>Hymenopus glabriflorus</i>	424
<i>Homalium riparium</i>	677	<i>Hymenopus heteromorphus</i> var. <i>glabrus</i>	424
<i>Homalium schippiae</i>	677	<i>Hymenopus heteromorphus</i> var. <i>heteromorphus</i>	424
<i>Homalium senarium</i>	677	<i>Hymenopus intrapetiolaris</i>	424
<i>Homalium spicatum</i>	677	<i>Hymenopus laevigatus</i>	424
<i>Homalium stenosepalum</i>	677	<i>Hymenopus latifolius</i>	424
<i>Homalium surinamense</i>	677	<i>Hymenopus latistipulus</i>	424
<i>Homalium trichocladium</i>	677	<i>Hymenopus macrophyllus</i>	425
<i>Homalium trichostemon</i>	677	<i>Hymenopus occultans</i>	425
<i>Homalolepis cedron</i>	705	<i>Hymenopus reticulatus</i>	425
<i>Homalolepis morettii</i>	705	HYPERICACEAE	470

Appendix 3. — Continuation.

<i>Hypericum acuminatum</i>	470	<i>Inga aggregata</i>	517
<i>Hypericum cayennense</i>	470	<i>Inga alata</i>	517
<i>Hypericum eugeniifolium</i>	470	<i>Inga alatocarpa</i>	521
<i>Hypericum guianense</i>	470	<i>Inga alba</i>	517, 522
<i>Hypericum latifolium</i>	470	<i>Inga albicans</i>	518
<i>Hypericum reticulatum</i>	470	<i>Inga albicoria</i>	518
<i>Hypericum rufescens</i>	471	<i>Inga altissima</i>	517
<i>Hypericum sessilifolium</i>	471	<i>Inga amazonica</i>	521
		<i>Inga amazonica</i> var. <i>bracteifera</i>	521
		<i>Inga amazonica</i> var. <i>lomatophylla</i>	521
		<i>Inga amazonica</i> var. <i>membranacea</i>	521
		<i>Inga apta</i>	517
		<i>Inga aria</i>	518
<i>Icica acuminata</i>	402	<i>Inga arinensis</i>	526
<i>Icica altissima</i>	399	<i>Inga arrabidae</i>	525
<i>Icica aracouchini</i>	400	<i>Inga assimilis</i>	517
<i>Icica decandra</i>	400	<i>Inga auristellae</i>	517
<i>Icica enneandra</i>	400, 401	<i>Inga bahiensis</i>	520
<i>Icica goudotiana</i>	402	<i>Inga benthamiana</i>	519
<i>Icica guianensis</i>	402	<i>Inga bourgoni</i>	517
<i>Icica heptaphylla</i>	402	<i>Inga brachyptera</i>	521
<i>Icica heterophylla</i>	400	<i>Inga brachystachya</i>	517
<i>Icica hostmannii</i>	402	<i>Inga brachystachys</i>	517
<i>Icica insignis</i>	403	<i>Inga bracteosa</i>	521
<i>Icica polybotrya</i>	404	<i>Inga brevialata</i>	526
<i>Icica spruceana</i>	404	<i>Inga brevipedicellata</i>	516
<i>Icica surinamensis</i>	402	<i>Inga calocephala</i>	521
<i>Icica viridiflora</i>	402	<i>Inga calycina</i>	518
<i>Icicopsis caudata</i>	404	<i>Inga canaminensis</i>	525
<i>Icicopsis insignis</i>	403	<i>Inga capitata</i>	517, 518, 524
<i>Icicopsis subserrata</i>	405	<i>Inga capitata</i> var. <i>brevicalyx</i>	518
<i>Icicopsis tenuifolia</i>	405	<i>Inga capitata</i> var. <i>latifolia</i>	524
<i>Ilex acuminata</i>	388	<i>Inga capitata</i> var. <i>tenuior</i>	518
<i>Ilex celastroides</i>	388	<i>Inga capuchoi</i>	518
<i>Ilex gentlei</i>	388	<i>Inga carachensis</i>	517
<i>Ilex guianensis</i>	388	<i>Inga cataractae</i>	545
<i>Ilex inundata</i>	388	<i>Inga catharinae</i>	525
<i>Ilex jenmanii</i>	388	<i>Inga cayennensis</i>	518
<i>Ilex laureola</i>	388	<i>Inga cayennensis</i> var. <i>sessiliflora</i>	518
<i>Ilex laureola</i> var. <i>genuina</i>	388	<i>Inga chorrorana</i>	519, 521
<i>Ilex laureola</i> var. <i>neglecta</i>	388	<i>Inga ciliata</i>	524
<i>Ilex macoucoua</i>	388	<i>Inga comewynensis</i>	525
<i>Ilex macrolaurus</i>	388	<i>Inga complanata</i>	519
<i>Ilex martiniana</i>	388	<i>Inga conferta</i>	519
<i>Ilex occidentalis</i>	388	<i>Inga conglomerata</i>	522, 523
<i>Ilex panamensis</i>	388	<i>Inga cordatoalata</i>	518
<i>Ilex paraguensis</i>	388	<i>Inga coriacea</i>	525
<i>Ilex riparia</i>	388	<i>Inga corymbifera</i>	522
<i>Ilex</i> sp. A	388	<i>Inga corymbifera</i> var. <i>brasiliensis</i>	522
<i>Ilex</i> sp. B	388	<i>Inga crassiflora</i>	519
<i>Inga</i>	503, 516	<i>Inga crevauxii</i>	518
<i>Inga acreana</i>	516	<i>Inga cycladenia</i>	524
<i>Inga acrocephala</i>	516	<i>Inga cyclocarpa</i>	522
<i>Inga acutifolia</i>	526	<i>Inga cylindrica</i>	518
<i>Inga affinis</i>	523, 526		

Appendix 3. — Continuation.

<i>Inga disticha</i>	518	<i>Inga loubryana</i>	521
<i>Inga divaricata</i>	545	<i>Inga macradenia</i>	526
<i>Inga dulcis</i>	525	<i>Inga macrophylla</i>	521, 523
<i>Inga dysantha</i>	518	<i>Inga macrophylla</i> var. <i>stenoptera</i>	521
<i>Inga edulis</i>	518, 519	<i>Inga mapiriensis</i>	520
<i>Inga edulis</i> var. <i>parviflora</i>	519	<i>Inga marginata</i>	522
<i>Inga ellsworthiana</i>	525	<i>Inga marginata</i> var. <i>itayensis</i>	522
<i>Inga excelsa</i>	522	<i>Inga mathewsiana</i>	522
<i>Inga fagifolia</i> f. <i>genuina</i>	522	<i>Inga meisneriana</i>	526
<i>Inga fagifolia</i> f. <i>pedicellaris</i>	522	<i>Inga melinonii</i>	522
<i>Inga fagifolia</i> var. <i>intermedia</i>	522	<i>Inga merianae</i>	520
<i>Inga fagifolia</i> var. <i>marginata</i>	522	<i>Inga microstachya</i>	523
<i>Inga falcistipula</i>	518	<i>Inga mitaraka</i>	522
<i>Inga fanchoniana</i>	519	<i>Inga multiflora</i>	521
<i>Inga fasciculata</i>	507	<i>Inga myriantha</i>	526
<i>Inga fastuosa</i>	519	<i>Inga myriocephala</i>	516
<i>Inga flagelliformis</i>	519	<i>Inga niopo</i>	505
<i>Inga floribunda</i>	525	<i>Inga nitida</i>	523
<i>Inga fraxinea</i>	517	<i>Inga nobilis</i>	522
<i>Inga galibica</i>	520	<i>Inga nobilis</i> var. <i>pavoniana</i>	522
<i>Inga gladiata</i>	526	<i>Inga nouragensis</i>	523
<i>Inga glomerata</i>	545	<i>Inga nubium</i>	523
<i>Inga graciliflora</i>	519	<i>Inga nuda</i>	525
<i>Inga graciliflora</i> var. <i>peruviana</i>	519	<i>Inga nuda</i> var. <i>longiflora</i>	525
<i>Inga gracilifolia</i>	519	<i>Inga odorata</i>	522
<i>Inga gracilipes</i>	526	<i>Inga ouraphylla</i>	521
<i>Inga grandiflora</i>	519	<i>Inga paraensis</i>	523
<i>Inga guaremalensis</i>	519	<i>Inga parviflora</i>	517, 520
<i>Inga guayaquilensis</i>	522	<i>Inga pavoniana</i>	522
<i>Inga heterophylla</i>	520	<i>Inga pedicellaris</i>	505
<i>Inga hostmannii</i>	525	<i>Inga peduncularis</i>	518
<i>Inga huberi</i>	520	<i>Inga pendula</i>	534
<i>Inga humboldtiana</i>	522	<i>Inga perrottetii</i>	525
<i>Inga hymenaeodes</i>	507	<i>Inga pezizifera</i>	523
<i>Inga ierensis</i>	524	<i>Inga pilosiuscula</i>	523
<i>Inga inaequalis</i>	545	<i>Inga pilosula</i>	523
<i>Inga ingoides</i>	520	<i>Inga platycarpa</i>	523
<i>Inga jenmanii</i>	520	<i>Inga poeppigiana</i>	524
<i>Inga jucunda</i>	522	<i>Inga polystachya</i>	518
<i>Inga laeta</i>	504	<i>Inga popayanensis</i>	524
<i>Inga lateriflora</i>	520	<i>Inga prieurii</i>	525
<i>Inga lateriflora</i> var. <i>latior</i>	520	<i>Inga protracta</i>	520
<i>Inga latifolia</i>	545	<i>Inga puberula</i>	522
<i>Inga lawranceana</i>	526	<i>Inga pubiramea</i>	529
<i>Inga laxiflora</i>	520	<i>Inga punctata</i>	524
<i>Inga leiocalycina</i>	520, 521	<i>Inga punctata</i> subsp. <i>chagrensis</i>	524
<i>Inga leptoloba</i>	524	<i>Inga punctata</i> var. <i>elongata</i>	524
<i>Inga leptostachya</i>	522	<i>Inga punctata</i> var. <i>panamensis</i>	524
<i>Inga lomatophylla</i>	521	<i>Inga pycnostachya</i>	522
<i>Inga lomatophylla</i> var. <i>bracteifera</i>	521	<i>Inga quadrangularis</i>	521
<i>Inga longiflora</i>	521	<i>Inga quassiaifolia</i>	523
<i>Inga longipedunculata</i>	521	<i>Inga radiata</i>	526
<i>Inga longituba</i>	519	<i>Inga ramiflora</i>	545
<i>Inga loretana</i>	522	<i>Inga recordii</i>	526

Appendix 3. — Continuation.

<i>Inga retinocarpa</i>	524	<i>Inga uraguensis</i> f. <i>tomentosula</i>	526
<i>Inga rhynchocalyx</i>	524	<i>Inga uraguensis</i> var. <i>parvifolia</i>	526
<i>Inga riedeliana</i>	522	<i>Inga urnifera</i>	523
<i>Inga riedeliana</i> var. <i>surinamensis</i>	522	<i>Inga velloziana</i>	526
<i>Inga riopalenquensis</i>	523	<i>Inga venosa</i>	519
<i>Inga rubiginosa</i>	524	<i>Inga vera</i> subsp. <i>affinis</i>	525
<i>Inga rufinervis</i>	524	<i>Inga versicolor</i>	523
<i>Inga rutilans</i>	526	<i>Inga virgultosa</i>	527, 530
<i>Inga salzmanniana</i>	525	<i>Inga vouapifolia</i>	520
<i>Inga sanctae-annae</i>	523	<i>Inga yngá</i>	519
<i>Inga sapida</i>	522	<i>Inga yunckeri</i>	521
<i>Inga sarmientosa</i>	524	<i>Inga</i> sp. A	527
<i>Inga scabriuscula</i>	519	<i>Inga</i> sp. B	527
<i>Inga scabriuscula</i> var. <i>villosior</i>	519	<i>Inga</i> sp. C	527
<i>Inga sciadion</i>	526	<i>Inga</i> sp. D	527
<i>Inga semialata</i>	522	<i>Iriartea durissima</i>	394
<i>Inga semialata</i> var. <i>latifolia</i>	522	<i>Iriartea exorrhiza</i>	394
<i>Inga sericantha</i>	522	<i>Iriartea exorrhiza</i> var. <i>elegans</i>	394
<i>Inga sertulifera</i>	520, 525	<i>Iriartea exorrhiza</i> var. <i>orbigniana</i>	394
<i>Inga sertulifera</i> subsp. <i>leptopus</i>	520	<i>Iriartea orbigniana</i>	394
<i>Inga sertulifera</i> var. <i>minor</i>	520	<i>Iriartea philonotia</i>	394
<i>Inga setifera</i>	523	<i>Iroucana guianensis</i>	673
<i>Inga soluta</i>	526	<i>Iryanthera elongata</i>	594
<i>Inga speciosa</i>	521	<i>Iryanthera hostmannii</i>	594
<i>Inga speciosa</i> var. <i>bracteifera</i>	521	<i>Iryanthera krukovi</i>	595
<i>Inga speciosa</i> var. <i>lomatophylla</i>	521	<i>Iryanthera microcarpa</i>	595
<i>Inga speciosa</i> var. <i>membranacea</i>	521	<i>Iryanthera paraensis</i>	594
<i>Inga splendens</i>	525	<i>Iryanthera sagotiana</i>	594
<i>Inga splendens</i> var. <i>hostmannii</i>	525	<i>Iryanthera sessilis</i>	594
<i>Inga splendens</i> var. <i>superba</i>	525	<i>Iryanthera tessmannii</i>	595
<i>Inga spruceana</i>	517	<i>Ischyranthera laevigata</i>	560
<i>Inga spuria</i> var. <i>sordida</i>	526	<i>Isertia breviflora</i>	660
<i>Inga standleyana</i>	524	<i>Isertia bullata</i>	660
<i>Inga stenocarpa</i>	520	<i>Isertia coccinea</i>	660
<i>Inga stipularis</i>	525	<i>Isertia coccinea</i> var. <i>hypoleuca</i>	660
<i>Inga striata</i>	525	<i>Isertia coccinea</i> var. <i>pentamera</i>	660
<i>Inga strigillosa</i>	524	<i>Isertia commutata</i>	660
<i>Inga suaveolens</i>	525	<i>Isertia flava</i>	660
<i>Inga subsericantha</i>	523	<i>Isertia glabra</i>	660
<i>Inga superba</i>	525	<i>Isertia hoehnei</i>	660
<i>Inga suturalis</i>	526	<i>Isertia hypoleuca</i>	660
<i>Inga tapajozensis</i>	517	<i>Isertia parviflora</i>	660
<i>Inga tenuiflora</i>	526	<i>Isertia parviflora</i> var. <i>hirta</i>	660
<i>Inga tenuiflora</i> var. <i>glabrior</i>	526	<i>Isertia pterantha</i>	660
<i>Inga tenuifolia</i>	518	<i>Isertia scarlatina</i>	660
<i>Inga thibaudiana</i>	526	<i>Isertia spiciformis</i>	660
<i>Inga thyrsoides</i>	517	<i>Isertia viscosa</i>	660
<i>Inga trapezifolia</i>	504	<i>Ivira pruriens</i>	557
<i>Inga tubiformis</i>	521	IXONANTHACEAE	471
<i>Inga tysonii</i>	522	<i>Ixora ferrea</i>	660
<i>Inga umbellata</i>	520	<i>Ixora gleasonii</i>	661
<i>Inga umbellifera</i>	526	<i>Ixora occidentalis</i>	657
<i>Inga umbratrica</i>	526	<i>Ixora panurensis</i>	661
<i>Inga uncinata</i>	519	<i>Ixora schomburgkiana</i>	661
<i>Inga uraguensis</i>	526	<i>Ixora sessilifolia</i>	658

Appendix 3. — Continuation.

<i>Ixora ulei</i>	661	<i>Klugiodendron laetum</i>	504
		<i>Klugiodendron umbrianum</i>	504
J		<i>Krugella hartii</i>	698, 699
<i>Jacaranda amazonensis</i>	396	<i>Krugia elliptica</i>	612
<i>Jacaranda copaia</i> subsp. <i>copaia</i>	396	<i>Krugia ferruginea</i>	612
<i>Jacaranda copaia</i> subsp. <i>spectabilis</i>	396	<i>Kubitzkia mezii</i>	480
<i>Jacaranda copaia</i> var. <i>paraensis</i>	396	<i>Kuhlia mollis</i>	671
<i>Jacaranda copaia</i> var. <i>spectabilis</i>	396	<i>Kummeria brasiliensis</i>	708
<i>Jacaranda duckei</i>	397	<i>Kutchubaea insignis</i>	661
<i>Jacaranda filicifolia</i>	397, 398	<i>Kutchubaea laxiflora</i>	661
<i>Jacaranda filicifolia</i> var. <i>puberula</i>	398	<i>Kyrtanthus longiflorus</i>	663
<i>Jacaranda obtusifolia</i> subsp. <i>rhombifolia</i>	397	L	
<i>Jacaranda obtusifolia</i> var. <i>rhombifolia</i>	398	<i>Labatia bilocularis</i>	694
<i>Jacaranda paraensis</i>	396	<i>Labatia caimito</i>	694
<i>Jacaranda procera</i>	396	<i>Labatia glomerata</i>	698
<i>Jacaranda rhombifolia</i>	397	<i>Labatia laevigata</i>	699
<i>Jacaranda spectabilis</i>	396	<i>Labatia lasiocarpa</i>	695
<i>Jacaranda superba</i>	396	<i>Labatia macrocarpa</i>	699
<i>Jacaratia actinophylla</i>	412	<i>Labatia parinarioides</i>	698
<i>Jacaratia costaricensis</i>	413	<i>Labatia parviflora</i>	698
<i>Jacaratia dodecaphylla</i>	412, 413	<i>Labatia pedunculata</i>	698
<i>Jacaratia dodecaphylla</i> f. <i>longiflora</i>	412	<i>Labatia reticulata</i>	694, 695
<i>Jacaratia dodecaphylla</i> var. <i>lucida</i>	413	<i>Labatia tovarensis</i>	698
<i>Jacaratia spinosa</i>	412	<i>Lacistema aggregatum</i>	471
<i>Japarandiba antillana</i>	499	<i>Lacistema aggregatum</i> var. <i>elongatum</i>	471
<i>Japarandiba augusta</i>	499	<i>Lacistema angustum</i>	471
<i>Japarandiba brasiliiana</i>	500	<i>Lacistema boliviianum</i>	471
<i>Japarandiba fastuosa</i>	499, 500	<i>Lacistema coriaceum</i>	471
<i>Japarandiba hexapetala</i>	500	<i>Lacistema curtum</i>	471
<i>Japarandiba marcgraaviana</i>	499	<i>Lacistema ellipticum</i>	472
<i>Japarandiba pterocarpa</i>	500	<i>Lacistema elongatum</i>	471
<i>Jessenia bataua</i>	394	<i>Lacistema floribundum</i>	471
<i>Jessenia bataua</i> subsp. <i>oligocarpa</i>	394	<i>Lacistema grandifolium</i>	472
<i>Jessenia oligocarpa</i>	394	<i>Lacistema guyanense</i>	471
<i>Jessenia polycarpa</i>	394	<i>Lacistema myricoides</i>	471
<i>Jessenia repanda</i>	394	<i>Lacistema myricoides</i> var. <i>stipitatum</i>	471
<i>Jessenia weberbaueri</i>	394	<i>Lacistema occidentale</i>	472
<i>Joncquetia paniculata</i>	365	<i>Lacistema orinocense</i>	471
<i>Jucunda tomentosa</i>	569	<i>Lacistema pacificum</i>	472
<i>Jugastrum coriaceum</i>	495	<i>Lacistema poeppigii</i>	471
<i>Jugastrum poiteaui</i>	502	<i>Lacistema polystachyum</i>	472
<i>Juglans catappa</i>	440	<i>Lacistema pubescens</i>	472
<i>Jupunba jupunba</i>	503	<i>Lacistema purpureum</i>	471
<i>Jupunba trapezifolia</i>	504	<i>Lacistema recurvum</i>	471
K		<i>Lacistema rosidiscum</i>	471
<i>Kajuputi leucadendra</i>	608, 609	<i>LACISTEMATACEAE</i>	471
<i>Kajuputi leucadendra</i> var. <i>angustifolia</i>	609	<i>Lacistema weberbaueri</i>	471
<i>Kaukenia surinamensis</i>	690	<i>Lacmellea aculeata</i>	385
<i>Kippistia cognata</i>	413	<i>Lacmellea guyanensis</i>	385
		<i>Lacunaria colonensis</i>	630
		<i>Lacunaria crenata</i>	628

Appendix 3. — Continuation.

<i>Lacunaria grandiflora</i>	629	<i>Lauraceae</i> sp. D	492
<i>Lacunaria jenmanii</i> subsp. <i>jenmanii</i>	629	<i>Lauraceae</i> sp. E	492
<i>Lacunaria jenmanii</i> subsp. <i>subsessilis</i>	629	<i>Lauraceae</i> sp. G	492
<i>Lacunaria panamensis</i>	629	<i>Lauraceae</i> sp. H	492
<i>Lacunaria pauciflora</i>	629	<i>Lauraceae</i> sp. I	492
<i>Lacunaria silvatica</i>	629	<i>Lauraceae</i> sp. J	492
<i>Lacunaria umbonata</i>	629	<i>Laurea tiliifolia</i>	583
<i>Ladenbergia lambertiana</i>	661	<i>Lauro-cerasus myrtifolia</i>	650
<i>Ladenbergia lucens</i>	661	<i>Laurus aestivalis</i>	483
<i>Ladenbergia puberula</i>	661	<i>Laurus canaliculata</i>	484
<i>Ladenbergia schomburgkii</i>	661	<i>Laurus cerifera</i>	485
<i>Ladenbergia venamoensis</i>	661	<i>Laurus crassifolia</i>	480
<i>Laetia</i>	671	<i>Laurus difformis</i>	474
<i>Laetia caseariooides</i>	672	<i>Laurus divaricata</i>	485
<i>Laetia glandulosa</i>	671	<i>Laurus elongata</i>	475
<i>Laetia obtusifolia</i>	672	<i>Laurus exaltata</i> var. <i>lancifolia</i>	486
<i>Laetia procera</i>	672	<i>Laurus floribunda</i>	485
<i>Lafoensia replicata</i>	547	<i>Laurus globosa</i>	482
<i>Lafoensia vandelliana</i>	547	<i>Laurus hexandra</i>	474
<i>Lafoensia vandelliana</i> subsp. <i>replicata</i>	547	<i>Laurus hibua</i>	482
<i>Laguncularia</i>	440	<i>Laurus koumaroucupa</i>	477
<i>Laguncularia obovata</i>	440	<i>Laurus leptobotra</i>	484
<i>Laguncularia racemosa</i>	440	<i>Laurus leucoxylon</i>	486
<i>Laguncularia racemosa</i> f. <i>longifolia</i>	440	<i>Laurus montana</i>	474
LAMIACEAE	472	<i>Laurus mucronata</i>	487
<i>Lanessania oligandra</i>	594	<i>Laurus nitida</i>	484
<i>Lanessania turbinata</i>	594	<i>Laurus ocotea</i>	486
<i>Langsdorfia instrumentaria</i>	669	<i>Laurus parviflora</i>	486
<i>Laplacea camelliifolia</i>	710	<i>Laurus puberula</i>	488
<i>Laplacea camelliooides</i>	710	<i>Laurus purpurea</i>	478
<i>Laplacea caracasana</i>	710	<i>Laurus reticulata</i>	483
<i>Laplacea fruticosa</i>	710	<i>Laurus retroflexa</i>	485
<i>Laplacea fruticosa</i> var. <i>chimantae</i>	710	<i>Laurus salicifolia</i>	476
<i>Laplacea fruticosa</i> var. <i>pulcherrima</i>	710	<i>Laurus triplinervis</i>	474
<i>Laplacea fruticosa</i> var. <i>symplocooides</i>	710	<i>Laxoplumeria baehniana</i>	385
<i>Laplacea intermedia</i>	710	<i>Lecostemon amazonicum</i>	648
<i>Laplacea parviflora</i>	710	<i>Lecostemon crassipes</i>	648
<i>Laplacea praemorsa</i>	710	<i>Lecostemon crassipes</i> var. <i>cayennense</i>	648
<i>Laplacea pubescens</i>	710	<i>Lecostemon sylvestre</i>	648
<i>Laplacea pubescens</i> var. <i>camelliifolia</i>	710	LECYTHIDACEAE	492
<i>Laplacea pubescens</i> var. <i>minor</i>	710	<i>Lecythis</i>	500
<i>Laplacea pubescens</i> var. <i>subcaudata</i>	710	<i>Lecythis acuminatissima</i>	495
<i>Laplacea pulcherrima</i>	710	<i>Lecythis alba</i>	501
<i>Laplacea quinoderma</i>	710	<i>Lecythis amara</i>	501
<i>Laplacea raimondiana</i>	710	<i>Lecythis basizone</i>	497
<i>Laplacea semiserrata</i>	710	<i>Lecythis bracteata</i>	494
<i>Laplacea speciosa</i>	710	<i>Lecythis chartacea</i>	500
<i>Laplacea speciosa</i> var. <i>intermedia</i>	710	<i>Lecythis cognata</i>	501
<i>Laplacea symplocooides</i>	710	<i>Lecythis confertiflora</i>	500
<i>Lasianthera amazonica</i>	708	<i>Lecythis congestiflora</i>	501
LAURACEAE	474, 492	<i>Lecythis coriacea</i>	495
Lauraceae sp. A	492	<i>Lecythis corrugata</i>	501
Lauraceae sp. B	492	<i>Lecythis couratari</i>	493
Lauraceae sp. C	492	<i>Lecythis crassinoda</i>	503

Appendix 3. — Continuation.

<i>Lecythis cupularis</i>	500	<i>Lecythis wullschlaegeliana</i>	497
<i>Lecythis davisii</i>	503	<i>Lecythis zabucajo</i>	502
<i>Lecythis davisii</i> var. <i>gracilipes</i>	503	<i>Legnotis cassipourea</i>	649
<i>Lecythis fagifolia</i>	493	<i>Legnotis elliptica</i>	648
<i>Lecythis gracilipes</i>	496, 497	<i>Legnotis macrophylla</i>	649
<i>Lecythis grandiflora</i>	495	<i>Leguminosae</i>	503, 530
<i>Lecythis grandifolia</i>	495	<i>Lemniscia floribunda</i>	468
<i>Lecythis hians</i>	503	<i>Lemniscia guianensis</i>	468
<i>Lecythis holcogyne</i>	501	<i>Leonia</i>	718
<i>Lecythis idatimon</i>	501	<i>Leonia glycycarpa</i>	719
<i>Lecythis idatimonoidea</i>	497	<i>Leonia glycycarpa</i> var. <i>racemosa</i>	719
<i>Lecythis jucunda</i>	497	<i>Leonia melinoniana</i>	719
<i>Lecythis laevicula</i>	500	<i>Leonia racemosa</i>	719
<i>Lecythis laevifolia</i>	499	<i>Leonia sphaerocarpum</i>	718
<i>Lecythis langsdorffii</i>	500	<i>Leonia surinamensis</i>	718
<i>Lecythis lecomtei</i>	503	<i>Leonia ulei</i>	721
<i>Lecythis longipes</i>	497	<i>LEPIDOBOTRYACEAE</i>	546
<i>Lecythis longipes</i> f. <i>genuina</i>	497	<i>Leptobalanus apetalus</i> var. <i>apertus</i>	425
<i>Lecythis longipes</i> f. <i>platycarpa</i>	497	<i>Leptobalanus apetalus</i> var. <i>apetalus</i>	425
<i>Lecythis longipes</i> var. <i>platycarpa</i>	497	<i>Leptobalanus granvillei</i>	425
<i>Lecythis lutea</i>	501	<i>Leptobalanus latus</i>	425
<i>Lecythis macrophylla</i>	497	<i>Leptobalanus longistylus</i>	425
<i>Lecythis marawynensis</i>	500	<i>Leptobalanus octandrus</i>	426
<i>Lecythis martinii</i>	501	<i>Leptobalanus persaudii</i>	426
<i>Lecythis melinonii</i>	497	<i>Leptobalanus sprucei</i>	426
<i>Lecythis micrantha</i>	496	<i>Leptodaphne subalpina</i>	484
<i>Lecythis monosperma</i>	500	<i>Leptodaphne tenuiflora</i>	484
<i>Lecythis multiflora</i>	493	<i>Leptolobium</i>	503, 527
<i>Lecythis odora</i>	495	<i>Leptolobium costulatum</i>	531
<i>Lecythis parviflora</i>	497	<i>Leptolobium nitens</i>	527
<i>Lecythis pedicellata</i>	497	<i>Leptolobium nitidulum</i>	527
<i>Lecythis persistens</i> subsp. <i>aurantiaca</i>	502	<i>Leptospermum leucadendrum</i>	608
<i>Lecythis persistens</i> subsp. <i>persistens</i>	501	<i>Leucaena</i>	503, 527
<i>Lecythis peruviana</i>	495	<i>Leucaena boliviiana</i>	537
<i>Lecythis pilosa</i>	497	<i>Leucaena glauca</i>	527
<i>Lecythis platycarpa</i>	497	<i>Leucaena leucocephala</i>	527
<i>Lecythis pneumatophora</i>	502	<i>Licania</i>	416
<i>Lecythis poiteaui</i>	493, 502	<i>Licania affinis</i>	425, 426
<i>Lecythis praeclera</i>	502	<i>Licania alba</i>	426
<i>Lecythis racemiflora</i>	502	<i>Licania amapaensis</i>	422
<i>Lecythis retroflexa</i>	495	<i>Licania anisophylla</i>	429
<i>Lecythis rigida</i>	500	<i>Licania aperta</i>	425
<i>Lecythis rorida</i>	500	<i>Licania apetala</i>	425
<i>Lecythis rubicunda</i>	501	<i>Licania apetala</i> var. <i>aperta</i>	425
<i>Lecythis sagotiana</i>	497	<i>Licania apetala</i> var. <i>pendula</i>	425
<i>Lecythis salebrosa</i>	501	<i>Licania aubletiana</i>	416
<i>Lecythis simiorum</i>	502	<i>Licania axilliflora</i>	428
<i>Lecythis spruceana</i>	500	<i>Licania benthamii</i>	424
<i>Lecythis subglandulosa</i>	498	<i>Licania biglandulosa</i>	424
<i>Lecythis tapuya</i>	497	<i>Licania bothynophylla</i>	426
<i>Lecythis tumefacta</i>	503	<i>Licania bracteosa</i>	432
<i>Lecythis validissima</i>	503	<i>Licania canescens</i>	427
<i>Lecythis venusta</i>	501	<i>Licania capinensis</i>	432
<i>Lecythis wachenheimii</i>	499	<i>Licania caracasana</i>	425

Appendix 3. — Continuation.

<i>Licania caudata</i>	422	<i>Licania microcarpa</i>	428
<i>Licania coriacea</i>	427	<i>Licania minutiflora</i>	430
<i>Licania costata</i>	424	<i>Licania obovata</i>	424
<i>Licania crassifolia</i>	428	<i>Licania obovatifolia</i>	432
<i>Licania cyathodes</i>	427	<i>Licania obtusifolia</i>	432
<i>Licania dahlgrenii</i>	425	<i>Licania occultans</i>	425
<i>Licania davillifolia</i>	427	<i>Licania octandra</i>	426
<i>Licania densiflora</i>	427	<i>Licania ovalifolia</i>	429
<i>Licania discolor</i>	427	<i>Licania pachystachya</i>	430
<i>Licania divaricata</i>	423	<i>Licania pallida</i>	429
<i>Licania elliptica</i>	427	<i>Licania pallidula</i>	427
<i>Licania fanshawei</i>	427	<i>Licania paniculata</i>	427
<i>Licania floribunda</i>	425	<i>Licania parinariooides</i>	432
<i>Licania galibica</i>	429	<i>Licania parinariooides</i> var. <i>latifolia</i>	432
<i>Licania glabra</i>	424	<i>Licania parviflora</i>	428, 429, 432
<i>Licania glabriflora</i>	424	<i>Licania parviflora</i> f. <i>brevifolia</i>	429
<i>Licania gracilis</i>	428	<i>Licania parviflora</i> f. <i>longifolia</i>	429
<i>Licania granvillei</i>	425	<i>Licania parviflora</i> var. <i>conduplicata</i>	428
<i>Licania grisea</i>	428	<i>Licania parviflora</i> var. <i>pallida</i>	429
<i>Licania guianensis</i>	424, 430	<i>Licania parviflora</i> var. <i>subfalcata</i>	429
<i>Licania helvola</i>	429	<i>Licania parviflora</i> var. <i>submembranacea</i>	428
<i>Licania heteromorpha</i>	423, 424	<i>Licania parvifolia</i>	428
<i>Licania heteromorpha</i> var. <i>glabra</i>	424	<i>Licania parvifructa</i>	429
<i>Licania heteromorpha</i> var. <i>grandifolia</i>	424	<i>Licania pendula</i>	425
<i>Licania hookeri</i> var. <i>obtusa</i>	426	<i>Licania persaudii</i>	426
<i>Licania hostmannii</i>	428	<i>Licania pruinosa</i>	429
<i>Licania huberiana</i>	432	<i>Licania pubiflora</i>	425
<i>Licania hylaea</i>	425	<i>Licania reticulata</i>	425
<i>Licania hypargyrea</i>	428	<i>Licania riparia</i>	430
<i>Licania hypoleuca</i>	427	<i>Licania robusta</i>	430
<i>Licania incana</i>	428	<i>Licania rodriguesii</i>	430
<i>Licania incana</i> var. <i>axilliflora</i>	428	<i>Licania rondonii</i>	430
<i>Licania intrapetiolaris</i>	424	<i>Licania schomburgkiana</i>	426
<i>Licania intrapetiolaris</i> var. <i>brevis</i>	424	<i>Licania sellowiana</i>	426
<i>Licania irwinii</i>	428	<i>Licania silvae</i>	430
<i>Licania kanukuensis</i>	427	<i>Licania sprucei</i>	426
<i>Licania kunthiana</i>	428	<i>Licania stahelii</i>	429
<i>Licania kuntzeana</i>	425	<i>Licania takutuensis</i>	426
<i>Licania laevigata</i>	424	<i>Licania turiuva</i>	426
<i>Licania lata</i>	425	<i>Licania utilis</i>	426
<i>Licania latifolia</i>	424	<i>Licania venosa</i>	426
<i>Licania latistipula</i>	424	<i>Licania wilson-browniei</i>	432
<i>Licania laxiflora</i>	428	<i>Licania</i> sp. A	430
<i>Licania leptostachya</i>	428	<i>Licania</i> sp. B	430
<i>Licania leptostachya</i> var. <i>axilliflora</i>	428	<i>Licania</i> sp. C	430
<i>Licania leptostachya</i> var. <i>crassifolia</i>	428	<i>Licaria amara</i>	481
<i>Licania licaniiiflora</i>	432	<i>Licaria appellii</i>	480
<i>Licania longifolia</i>	426	<i>Licaria cannella</i>	480
<i>Licania longistyla</i>	426	<i>Licaria cayennensis</i>	480
<i>Licania macrophylla</i>	425, 428	<i>Licaria chrysophylla</i>	480
<i>Licania majuscula</i>	428	<i>Licaria crassifolia</i>	480
<i>Licania maxima</i>	428	<i>Licaria debilis</i>	480
<i>Licania membranacea</i>	429	<i>Licaria foveolata</i>	481
<i>Licania micrantha</i>	429	<i>Licaria guianensis</i>	480

Appendix 3. — Continuation.

<i>Licaria martiniana</i>	481	<i>Lonchocarpus</i>	503, 528
<i>Licaria meisneri</i>	481	<i>Lonchocarpus amerimnum</i>	535
<i>Licaria pachycarpa</i>	481	<i>Lonchocarpus discolor</i>	528
<i>Licaria parvifolia</i>	476	<i>Lonchocarpus hedyosmus</i>	528
<i>Licaria polyphylla</i>	481	<i>Lonchocarpus heptaphyllus</i>	528
<i>Licaria rigida</i>	480	<i>Lonchocarpus latifolius</i>	528
<i>Licaria rufotomentosa</i>	481	<i>Lonchocarpus macrocarpus</i> var. <i>sericophyllus</i>	528
<i>Licaria simulans</i>	481	<i>Lonchocarpus paniculatus</i>	528
<i>Licaria subbullata</i>	481	<i>Lonchocarpus staudtii</i>	506
<i>Licaria vernicosa</i>	481	<i>Lonchocarpus swartzii</i>	528
<i>Licaria wilhelminensis</i>	481	<i>Loretoa peruviana</i>	652
<i>Lieutautia mirabilis</i>	565	<i>Loreya acutifolia</i>	559
<i>Lignonia monodynama</i>	719	<i>Loreya arborescens</i>	559
<i>Limadendron</i>	503, 527	<i>Loreya brunnescens</i>	560
<i>Limadendron hostmannii</i>	527, 530	<i>Loreya flavescens</i>	560
LINACEAE	546	<i>Loreya maguirei</i>	559
<i>Lindackeria maynensis</i>	363	<i>Loreya mespilooides</i>	560
<i>Lindackeria maynensis</i> var. <i>laxiflora</i>	363	<i>Loreya subrotundifolia</i>	560
<i>Lindackeria paludosa</i>	363	<i>Loxopterygium sagotii</i>	364
<i>Lindleya semiserrata</i>	710	<i>Lucuma acreana</i>	699
<i>Lingoum esculentum</i>	537	<i>Lucuma anibifolia</i>	701
<i>Lingoum officinale</i>	536	<i>Lucuma buchananiifolia</i>	700
<i>Lingoum rohri</i>	536	<i>Lucuma caimito</i>	694
<i>Lingoum rufescens</i>	536	<i>Lucuma dissepala</i>	702
<i>Lingoum villosum</i>	536	<i>Lucuma dolichophylla</i>	702
<i>Lingoum violaceum</i>	536	<i>Lucuma duckei</i>	702
<i>Linnaeobreynia pulcherrima</i>	411	<i>Lucuma durlandii</i>	697
<i>Linociera tetrandra</i>	635	<i>Lucuma ephedrantha</i>	697
<i>Liriosma cerifera</i>	633	<i>Lucuma excelsa</i>	700
<i>Liriosma guianensis</i>	633	<i>Lucuma glazioviana</i>	698
<i>Lisanthius caerulescens</i>	722	<i>Lucuma glomerata</i>	698
<i>Lithocardium alliodorum</i>	444	<i>Lucuma gutta</i>	702
<i>Lithocardium bicolor</i>	444	<i>Lucuma hartii</i>	698, 699
<i>Lithocardium calophyllum</i>	445	<i>Lucuma huallagae</i>	699
<i>Lithocardium cordifolium</i>	446	<i>Lucuma hypoglauca</i>	698
<i>Lithocardium exaltatum</i>	444	<i>Lucuma jenmanii</i>	702
<i>Lithocardium hebecarpum</i>	447	<i>Lucuma laevigata</i>	699
<i>Lithocardium heterophyllum</i>	447	<i>Lucuma lasiocarpa</i>	695
<i>Lithocardium hispidissimum</i>	445	<i>Lucuma laurifolia</i>	694
<i>Lithocardium lockhartii</i>	444	<i>Lucuma laurifolia</i> var. <i>reticulata</i>	694
<i>Lithocardium mirandum</i>	445	<i>Lucuma littoralis</i>	702
<i>Lithocardium muneco</i>	446	<i>Lucuma lucentifolia</i>	687
<i>Lithocardium nervosum</i>	445	<i>Lucuma melinonii</i>	694
<i>Lithocardium nodosum</i>	445	<i>Lucuma minutiflora</i>	700
<i>Lithocardium paniculare</i>	445	<i>Lucuma oblonga</i>	700
<i>Lithocardium pubescens</i>	447	<i>Lucuma paraensis</i>	699
<i>Lithocardium scabridum</i>	444	<i>Lucuma pentasperma</i>	687
<i>Lithocardium sericalyx</i>	446	<i>Lucuma platyphylla</i>	700
<i>Lithocardium sprucei</i>	446	<i>Lucuma procera</i>	700
<i>Lithocardium tenuifolium</i>	444	<i>Lucuma procera</i> var. <i>cuspidata</i>	700
<i>Lithocardium tetrandrum</i>	446	<i>Lucuma psammophila</i> var. <i>macrophylla</i>	698
<i>Lithocardium tetraphyllum</i>	443	<i>Lucuma pulverulenta</i>	695
<i>Lithocardium toqueve</i>	446	<i>Lucuma rivicoa</i>	699
LOGANIACEAE	546	<i>Lucuma rivicoa</i> var. <i>glaucophylla</i>	699

Appendix 3. — Continuation.

<i>Lucuma sagotiana</i>	701	<i>Macrolobium campestre</i>	529
<i>Lucuma speciosa</i>	701	<i>Macrolobium chrysostachyum</i>	528
<i>Lucuma temare</i>	694	<i>Macrolobium chrysostachyum</i> var. <i>parviflorum</i>	528
<i>Lucuma tomentosa</i>	689	<i>Macrolobium elegans</i>	529
<i>Lucuma tuberculata</i>	702	<i>Macrolobium guianense</i>	529
<i>Lucuma venulosa</i>	693	<i>Macrolobium huberianum</i>	529
<i>Luehea alternifolia</i>	555	<i>Macrolobium hymenaeoides</i>	529
<i>Luehea althaeiflora</i>	555	<i>Macrolobium hymeneifolium</i>	528
<i>Luehea ferruginea</i>	555	<i>Macrolobium multijugum</i>	529
<i>Luehea flavescens</i>	555	<i>Macrolobium outea</i>	529
<i>Luehea maroniensis</i>	555	<i>Macrolobium pinnatum</i>	529
<i>Luehea rosea</i>	555	<i>Macrolobium stamineum</i>	529
<i>Luehea rugosa</i>	555	<i>Macrolobium utea</i>	529
<i>Luehea speciosa</i>	555	<i>Macrolobium vuapa</i>	529
<i>Luehea tarapotina</i>	555	<i>Macropiper latum</i>	642
<i>Lueheopsis althaeiflora</i>	555	<i>Macrosamanea</i>	503, 529
<i>Lueheopsis flavescens</i>	555	<i>Macrosamanea longiflora</i>	529
<i>Lueheopsis rosea</i>	555	<i>Macrosamanea pedicellaris</i>	505
<i>Lueheopsis rugosa</i>	555	<i>Macrosamanea pubiramea</i>	529
<i>Luma bergii</i>	604	<i>Macrostegia ruiziana</i>	474
LYTHRACEAE	547	<i>Macroule coutinhoi</i>	531
M			
<i>Maba melinonii</i>	448	<i>Magonia amentacea</i>	643
<i>Maba sericea</i>	449	<i>Magonia brachysepala</i>	643
<i>Mabea argutissima</i>	463	<i>Mahurea palustris</i>	408
<i>Mabea caudata</i>	463	<i>Malouetia duckei</i>	385
<i>Mabea maynensis</i>	462	<i>Malouetia guianensis</i>	385
<i>Mabea piriri</i>	462	<i>Malouetia tamaquarina</i>	386
<i>Mabea salicoides</i>	462	<i>Malpighia altissima</i>	547
<i>Mabea schomburgkii</i>	463	<i>Malpighia argentea</i>	547
<i>Mabea speciosa</i>	462	MALPIGHIACEAE	547
<i>Mabea subsessilis</i>	463	<i>Malpighia coriacea</i>	548
<i>Mabea taquari</i>	463	<i>Malpighia crassifolia</i>	547
<i>Mabea taquari</i> var. <i>angustifolia</i>	463	<i>Malpighia densa</i>	548
<i>Macaglia alba</i>	381	<i>Malpighia elegans</i>	551
<i>Macaglia excelsa</i>	381	<i>Malpighia emarginata</i>	606
<i>Macaglia oblonga</i>	382	<i>Malpighia guadalupensis</i>	549
<i>Macoubea fasciculata</i>	386	<i>Malpighia laevigata</i>	548
<i>Macoubea guianensis</i>	385	<i>Malpighia moureila</i>	547
<i>Macoubea guianensis</i> var. <i>reticulata</i>	385	<i>Malpighia pulchra</i>	548
<i>Macoubea paucifolia</i>	385	<i>Malpighia spicata</i>	549, 550
<i>Macoubea reticulata</i>	385	MALVACEAE	551
<i>Macoubea sinuosa</i>	385	<i>Manabea arborescens</i>	472
<i>Macoubea sprucei</i> var. <i>paucifolia</i>	385	<i>Manabea laevis</i>	722
<i>Macoucoua guianensis</i>	388	<i>Manabea villosa</i>	472
<i>Macrocteron cuneatus</i>	459	<i>Manicaria atricha</i>	393
<i>Macrolobium</i>	503, 528	<i>Manicaria pluknetii</i>	393
<i>Macrolobium acaciifolium</i>	528	<i>Manicaria saccifera</i>	393
<i>Macrolobium acaciifolium</i> var. <i>vestitum</i>	528	<i>Manicaria saccifera</i> var. <i>mediterranea</i>	393
<i>Macrolobium angustifolium</i>	528	<i>Manicaria saccifera</i> var. <i>pluknetii</i>	393
<i>Macrolobium bifolium</i>	529	<i>Manilkara amazonica</i>	690
		<i>Manilkara balata</i>	689, 690
		<i>Manilkara balata</i> var. <i>cruegeri</i>	689
		<i>Manilkara balata</i> var. <i>domingensis</i>	690
		<i>Manilkara balata</i> var. <i>gutta</i>	689

Appendix 3. — Continuation.

<i>Manilkara balata</i> var. <i>hartii</i>	689	<i>Matayba arborescens</i>	681, 683
<i>Manilkara balata</i> var. <i>melinonii</i>	689, 690	<i>Matayba chimantensis</i>	682
<i>Manilkara balata</i> var. <i>schomburgkii</i>	689	<i>Matayba fallax</i>	682
<i>Manilkara balata</i> var. <i>sieberi</i>	689	<i>Matayba guianensis</i>	682
<i>Manilkara bidentata</i> subsp. <i>bidentata</i>	689	<i>Matayba guianensis</i> f. <i>genuina</i>	682
<i>Manilkara bidentata</i> subsp. <i>surinamensis</i>	690	<i>Matayba guianensis</i> subf. <i>subovalis</i>	682
<i>Manilkara bidentata</i> var. <i>cruegeri</i>	689	<i>Matayba inelegans</i>	682
<i>Manilkara darienensis</i>	689	<i>Matayba jauaensis</i>	682
<i>Manilkara huberi</i>	690	<i>Matayba laevigata</i>	682
<i>Manilkara longiciliata</i>	690	<i>Matayba macrolepis</i>	681
<i>Manilkara nitida</i>	690	<i>Matayba macrostylis</i>	681
<i>Manilkara paraensis</i>	690	<i>Matayba miquelii</i>	682
<i>Manilkara riedleana</i>	690	<i>Matayba oligandra</i>	682
<i>Manilkara siqueiraei</i>	690	<i>Matayba oligandra</i> var. <i>occidentalis</i>	682
<i>Manilkara surinamensis</i>	690	<i>Matayba oligandra</i> var. <i>ptariana</i>	682
<i>Manilkara williamsii</i>	689	<i>Matayba opaca</i>	682
<i>Mapouria biacuminata</i>	664	<i>Matayba opaca</i> var. <i>fallax</i>	682
<i>Mapouria chionantha</i>	665	<i>Matayba peruviana</i> subsp. <i>oligandra</i>	682
<i>Mapouria corumbensis</i>	664	<i>Matayba reducta</i>	681
<i>Mapouria corymbifera</i>	664	<i>Matayba sororopaniana</i>	682
<i>Mapouria ficigemma</i>	664	<i>Matayba stenodictya</i>	682
<i>Mapouria guianensis</i>	664	<i>Matisia apaporiensis</i>	556
<i>Mapouria myriantha</i>	664	<i>Matisia lasiocalyx</i>	555
<i>Mapouria ornithophila</i>	664	<i>Matisia muricata</i>	555
<i>Mapouria patrisii</i>	664	<i>Matisia ochrocalyx</i>	555
<i>Mapouria sambucina</i>	664	<i>Matthiola argentea</i>	660
<i>Mapouria simira</i>	666	<i>Mauria multiflora</i>	365
<i>Maprounea guianensis</i>	463	<i>Mauria obtusa</i>	365
<i>Maquira calophylla</i>	592	<i>Mauria subbijuga</i>	365
<i>Maquira guianensis</i>	592	<i>Mauritia flexuosa</i>	393
<i>Maquira sclerophylla</i>	592	<i>Mauritia flexuosa</i> var. <i>venezuelana</i>	393
<i>Maquira</i> sp. A	592	<i>Mauritia minor</i>	393
<i>Margaritaria adeliooides</i>	638	<i>Mauritia sagus</i>	393
<i>Margaritaria nobilis</i>	638	<i>Mauritia setigera</i>	393
<i>Margaritaria nobilis</i> var. <i>antillana</i>	638	<i>Mauritia sphaerocarpa</i>	393
<i>Margaritaria nobilis</i> var. <i>hypomalaca</i>	638	<i>Mauritia vinifera</i>	393
<i>Margaritopsis boliviiana</i>	656	<i>Maximiliana caribaea</i>	392
<i>Marialva guianensis</i>	437	<i>Maximiliana elegans</i>	392
<i>Marlierea acuminata</i>	612	<i>Maximiliana inajai</i>	395
<i>Marlierea brachymischa</i>	625	<i>Maximiliana longirostrata</i>	392
<i>Marlierea elliptica</i>	612	<i>Maximiliana macrogyne</i>	392
<i>Marlierea ferruginea</i>	612	<i>Maximiliana macropetala</i>	392
<i>Marlierea gleasonii</i>	611	<i>Maximiliana mariipa</i>	392
<i>Marlierea guyanensis</i>	612	<i>Maximiliana martiana</i>	392
<i>Marlierea montana</i>	617	<i>Maximiliana regia</i>	392
<i>Marlierea multiglomerata</i>	617	<i>Maximiliana stenocarpa</i>	392
<i>Marlierea richardiana</i>	612, 617	<i>Maximiliana tetrasticha</i>	392
<i>Marlierea suffruticosa</i>	617	<i>Maximilanea orinocensis</i>	398
<i>Marmaroxylon racemosum</i>	546	<i>Maximilanea parkeri</i>	399
<i>Martiodendron</i>	503, 530	<i>Maximilanea parvifolia</i>	399
<i>Martiodendron parviflorum</i>	530	<i>Mayepia guianensis</i>	635
<i>Martiusia parviflora</i>	530	<i>Mayna laxiflora</i>	363
<i>Mastichodendron williamsii</i>	692	<i>Mayna linguifolia</i>	362
<i>Matayba affinis</i>	682	<i>Mayna longifolia</i>	362

Appendix 3. — Continuation.

<i>Mayna longifolia</i> var. <i>heliocarpa</i>	362	<i>Melastoma grossularioides</i>	560
<i>Mayna longifolia</i> var. <i>phasmatocarpa</i>	362	<i>Melastoma holosericeum</i>	564
<i>Mayna muricida</i>	362	<i>Melastoma laevigata</i>	722
<i>Mayna pacifica</i> var. <i>pusilla</i>	362	<i>Melastoma lasiopetalum</i>	568
<i>Mayna paludosa</i>	363	<i>Melastoma lepidotum</i>	564
<i>Maytenus</i>	414, 415	<i>Melastoma leucanthum</i>	567
<i>Maytenus cardenasi</i>	415	<i>Melastoma longifolium</i>	565
<i>Maytenus erythrocarpa</i>	415	<i>Melastoma macrophyllum</i>	568
<i>Maytenus floribunda</i>	415	<i>Melastoma megalophyllum</i>	569
<i>Maytenus guianensis</i>	414	<i>Melastoma minutiflorum</i>	565
<i>Maytenus guyanensis</i>	414	<i>Melastoma montanum</i>	567
<i>Maytenus guyanensis</i> f. <i>crenulata</i>	414	<i>Melastoma mucronatum</i>	564
<i>Maytenus myrsinoides</i>	415	<i>Melastoma parviflorum</i>	567
<i>Maytenus oblongata</i>	415	<i>Melastoma patens</i>	568
<i>Maytenus reissekii</i>	415	<i>Melastoma paulense</i>	559
<i>Maytenus</i> sp. A	415	<i>Melastoma pendulifolium</i>	567
<i>Maytenus</i> sp. B	415	<i>Melastoma pendulum</i>	565
<i>Meborea guianensis</i>	638	<i>Melastoma prasinum</i>	567
<i>Meioloma guianensis</i>	693	<i>Melastoma pulchrum</i>	567
<i>Meladendron leucocladum</i>	608	<i>Melastoma punctatum</i>	568
<i>Melaleuca amboinensis</i>	609	<i>Melastoma quinquenervium</i>	567
<i>Melaleuca leucadendra</i>	608, 609	<i>Melastoma ramiflorum</i>	561
<i>Melaleuca leucadendra</i> var. <i>albida</i>	609	<i>Melastoma scorpioides</i>	569
<i>Melaleuca leucadendra</i> var. <i>angusta</i>	609	<i>Melastoma sepiarium</i>	567
<i>Melaleuca leucadendra</i> var. <i>angustifolia</i>	609	<i>Melastoma serrulatum</i>	568
<i>Melaleuca leucadendra</i> var. <i>coriacea</i>	609	<i>Melastoma splendens</i>	569
<i>Melaleuca leucadendra</i> var. <i>cunninghamii</i>	609	<i>Melastoma strangulatum</i>	568
<i>Melaleuca leucadendra</i> var. <i>lancifolia</i>	609	<i>Melastoma suaveolens</i>	567
<i>Melaleuca leucadendra</i> var. <i>mimosoides</i>	609	<i>Melastoma succosum</i>	561
<i>Melaleuca maidenii</i>	609	MELASTOMATACEAE	559, 566
<i>Melaleuca mimosoides</i>	609	<i>Melastoma tamonea</i>	566
<i>Melaleuca quinquenervia</i>	609	<i>Melastoma tomentosum</i>	569
<i>Melaleuca rigida</i>	609	<i>Melastoma trinervium</i>	567, 569
<i>Melaleuca rubriflora</i>	609	MELIACEAE	572
<i>Melaleuca smithii</i>	609	<i>Melia guara</i>	575, 578
<i>Melaleuca viridiflora</i> var. <i>angustifolia</i>	609	<i>Melicoccus pedicellaris</i>	682
<i>Melaleuca viridiflora</i> var. <i>rubriflora</i>	609	<i>Meliosma</i> sp. A	671
<i>Melanoxyylon amazonicum</i>	537	<i>Merizadenia amplifolia</i>	385
<i>Melanoxyylon speciosum</i>	537	<i>Merizadenia sananho</i>	387
<i>Melastoma acuminatum</i>	567	<i>Mesopanax capitatus</i>	390
<i>Melastoma arborescens</i>	559	<i>Mespilodaphne ceanothifolia</i>	484
<i>Melastoma argyrophyllum</i>	563	<i>Mespilodaphne complicata</i>	485
<i>Melastoma balbisianum</i>	568	<i>Mespilodaphne fasciculata</i>	485
<i>Melastoma bracteatum</i>	564	<i>Mespilodaphne laxiflora</i>	487
<i>Melastoma chrysophyllum</i>	563	<i>Mespilodaphne nitida</i>	484
<i>Melastoma compressum</i>	566	<i>Mespilodaphne oblonga</i>	487
<i>Melastoma egense</i>	559	<i>Mespilodaphne opifera</i>	487
<i>Melastoma elatum</i>	563	<i>Mespilodaphne pyriformis</i>	479
<i>Melastoma flavescens</i>	560	<i>Metrosideros albida</i>	609
<i>Melastoma fothergilla</i>	565, 566	<i>Metrosideros coriacea</i>	609
<i>Melastoma fothergilla</i> var. <i>lanceolata</i>	566	<i>Metrosideros quinquenervia</i>	609
<i>Melastoma fothergilla</i> var. <i>ovalis</i>	566	METTENIUSACEAE	582
<i>Melastoma fulvum</i>	563	<i>Mezia anacardioides</i>	481
<i>Melastoma glabrum</i>	567	<i>Mezia itauba</i>	481

Appendix 3. — Continuation.

<i>Mezilaurus anacardiooides</i>	481	<i>Miconia guianensis</i> var. <i>vulgaris</i>	566
<i>Mezilaurus itauba</i>	481	<i>Miconia habrolepis</i>	568
<i>Miconia</i>	561	<i>Miconia holosericea</i>	564
<i>Miconia acuminata</i>	561	<i>Miconia holosericea</i> var. <i>bracteata</i>	564
<i>Miconia affinis</i>	561	<i>Miconia holosericea</i> var. <i>mucronata</i>	564
<i>Miconia amacurensis</i>	562	<i>Miconia holosericea</i> var. <i>subquintuplinervia</i>	564
<i>Miconia ampla</i>	562	<i>Miconia hypoleuca</i>	564
<i>Miconia amplexans</i>	569	<i>Miconia involucrata</i>	562
<i>Miconia anceps</i>	569	<i>Miconia kappleri</i>	564
<i>Miconia argyrophylla</i>	563	<i>Miconia lambertiana</i>	565
<i>Miconia argyrophylla</i> subsp. <i>gracilis</i>	563	<i>Miconia lepidota</i>	564, 568
<i>Miconia argyrophylla</i> var. <i>attenuata</i>	563	<i>Miconia lepidota</i> var. <i>grandifolia</i>	568
<i>Miconia attenuata</i>	567	<i>Miconia leucocephala</i>	568
<i>Miconia attenuata</i> var. <i>subquintuplinervia</i>	567	<i>Miconia longifolia</i>	563, 565
<i>Miconia aubletiana</i>	567	<i>Miconia longifolia</i> var. <i>aubletiana</i>	563
<i>Miconia axilliflora</i>	561	<i>Miconia longipedunculata</i>	565
<i>Miconia beurlingii</i>	562	<i>Miconia longispicata</i>	565
<i>Miconia bifrons</i>	564	<i>Miconia longispicata</i> var. <i>minor</i>	565
<i>Miconia blanchetiana</i>	564	<i>Miconia longistyla</i>	563
<i>Miconia boissieriana</i>	569	<i>Miconia macrophylla</i>	567, 568
<i>Miconia borealis</i>	565	<i>Miconia macrophylla</i> var. <i>hostmannii</i>	568
<i>Miconia bracteolaris</i>	563	<i>Miconia macrophylla</i> var. <i>latifolia</i>	568
<i>Miconia bracteolaris</i> var. <i>cardiophora</i>	563	<i>Miconia macrophylla</i> var. <i>leucocephala</i>	568
<i>Miconia cayumbensis</i>	562	<i>Miconia macrophylla</i> var. <i>serrulata</i>	568
<i>Miconia cecidophora</i>	562	<i>Miconia matthaei</i>	565
<i>Miconia chrysophylla</i>	563	<i>Miconia matthaei</i> var. <i>undulata</i>	565
<i>Miconia collina</i>	567	<i>Miconia megaphylla</i>	562
<i>Miconia congesta</i>	566	<i>Miconia melanodendron</i>	565
<i>Miconia crispula</i>	567	<i>Miconia melinonii</i>	565
<i>Miconia cristulata</i>	567	<i>Miconia microcarpa</i>	562
<i>Miconia cruegeriana</i>	564	<i>Miconia minutiflora</i>	565
<i>Miconia darienensis</i>	566	<i>Miconia minutiflora</i> var. <i>latifolia</i>	565
<i>Miconia discolor</i>	568	<i>Miconia mirabilis</i>	565
<i>Miconia dispar</i>	563	<i>Miconia mucronata</i>	564
<i>Miconia egensis</i>	563	<i>Miconia mucronulata</i>	567
<i>Miconia elata</i>	563	<i>Miconia myriantha</i>	566
<i>Miconia elliptica</i>	569	<i>Miconia naudiniana</i>	564
<i>Miconia eriodonta</i>	563	<i>Miconia nemoralis</i>	567
<i>Miconia eriodonta</i> var. <i>oblongifolia</i>	563	<i>Miconia obovalis</i>	569
<i>Miconia eurychaeniooides</i>	563	<i>Miconia palustris</i>	567
<i>Miconia fendleriana</i>	566	<i>Miconia panicularis</i>	563
<i>Miconia fleischeriana</i>	567	<i>Miconia parviflora</i>	567
<i>Miconia florida</i>	566	<i>Miconia phaeophylla</i>	566
<i>Miconia fothergilla</i>	565	<i>Miconia planinervia</i>	562
<i>Miconia fragilis</i>	564	<i>Miconia platyhedra</i>	568
<i>Miconia fulva</i>	563	<i>Miconia pluknetii</i>	566
<i>Miconia fulva</i> var. <i>angustifolia</i>	563	<i>Miconia poeppigii</i>	566
<i>Miconia fulva</i> var. <i>aubletiana</i>	563	<i>Miconia prasina</i>	567
<i>Miconia fulva</i> var. <i>poeppigii</i>	563	<i>Miconia prasina</i> var. <i>angustifolia</i>	567
<i>Miconia fulva</i> var. <i>tinctoria</i>	563	<i>Miconia prasina</i> var. <i>attenuata</i>	567
<i>Miconia glossocentra</i>	565	<i>Miconia prasina</i> var. <i>collina</i>	567
<i>Miconia gratissima</i>	564	<i>Miconia prasina</i> var. <i>crispula</i>	567
<i>Miconia guianensis</i>	566	<i>Miconia prasina</i> var. <i>denticulata</i>	567
<i>Miconia guianensis</i> var. <i>ovalis</i>	566	<i>Miconia pteropoda</i>	567

Appendix 3. — Continuation.

<i>Miconia pubipetala</i>	567	<i>Micropholis grandifolia</i>	692
<i>Miconia punctata</i>	563, 568	<i>Micropholis guatemalensis</i>	692
<i>Miconia punctata</i> var. <i>brevifolia</i>	568	<i>Micropholis guyanensis</i> subsp. <i>duckeana</i>	692
<i>Miconia punctata</i> var. <i>latifolia</i>	563	<i>Micropholis guyanensis</i> subsp. <i>guyanensis</i>	691
<i>Miconia pusilliflora</i>	562	<i>Micropholis imrayana</i>	691
<i>Miconia pyrifolia</i>	568	<i>Micropholis longipedicellata</i>	692
<i>Miconia repandocrenata</i>	567	<i>Micropholis martiana</i>	691
<i>Miconia revoluta</i>	567	<i>Micropholis melinoniana</i>	692
<i>Miconia robusta</i>	569	<i>Micropholis mensalis</i>	692
<i>Miconia ruficalyx</i>	568	<i>Micropholis mexicana</i>	692
<i>Miconia schomburgkii</i>	564	<i>Micropholis mucronata</i>	693
<i>Miconia scorpioides</i>	569	<i>Micropholis obscura</i>	693
<i>Miconia sepiaria</i>	567	<i>Micropholis paraensis</i>	691
<i>Miconia serialis</i>	568	<i>Micropholis porphyrocarpa</i>	693
<i>Miconia serrulata</i>	568	<i>Micropholis portoricensis</i>	691
<i>Miconia splendens</i>	569	<i>Micropholis portoricensis</i> var. <i>curvata</i>	691
<i>Miconia surinamensis</i>	566	<i>Micropholis portoricensis</i> var. <i>mesuifolia</i>	691
<i>Miconia symplectocaulos</i>	569	<i>Micropholis rigida</i>	693
<i>Miconia tetraspermoides</i>	569	<i>Micropholis rosadinha-brava</i>	692
<i>Miconia tomentella</i>	568	<i>Micropholis rufa</i>	691
<i>Miconia tomentosa</i>	569	<i>Micropholis sanctae-rosae</i>	693
<i>Miconia tomentosa</i> var. <i>ovata</i>	569	<i>Micropholis</i> sp. A	693
<i>Miconia trichotoma</i>	567	<i>Micropholis steyermarkii</i>	697
<i>Miconia trinervia</i>	569	<i>Micropholis truncata</i>	691
<i>Miconia trinitatis</i>	565	<i>Micropholis trunciflora</i>	693
<i>Miconia tschudyoides</i>	569	<i>Micropholis ulei</i>	691
<i>Miconia umbrifera</i>	569	<i>Micropholis venulosa</i>	693
<i>Miconia verticilliflora</i>	567	<i>Micropholis wurdackii</i>	691
<i>Miconia wilsonii</i>	565	<i>Micropteryx poeppigiana</i>	515
<i>Micrandra bracteosa</i>	463	<i>Mimosa acacioides</i>	505
<i>Micrandra brownsbergensis</i>	463	<i>Mimosa alba</i>	517
<i>Micrandra elata</i>	463	<i>Mimosa atakta</i>	504
<i>Micrandra glaziovii</i>	463	<i>Mimosa biceps</i>	527
<i>Micrandra rossiana</i>	463	<i>Mimosa bourgoni</i>	517
<i>Micrandra santanderensis</i>	463	<i>Mimosa coriacea</i>	525
<i>Micrandra spruceana</i>	463	<i>Mimosa corymbosa</i>	515
<i>Micranthera clusiifolia</i>	437	<i>Mimosa cylindrica</i>	518
<i>Micranthera clusiooides</i>	437	<i>Mimosa dulcis</i>	525
<i>Micropholis achradiformis</i>	691	<i>Mimosa fastuosa</i>	519
<i>Micropholis acutangula</i>	691	<i>Mimosa flagelliformis</i>	519
<i>Micropholis balata</i>	692	<i>Mimosa fraxinea</i>	517
<i>Micropholis calophylloides</i>	693	<i>Mimosa frondosa</i>	527
<i>Micropholis cayennensis</i>	691	<i>Mimosa glandulosa</i>	527
<i>Micropholis chrysophylloides</i>	691	<i>Mimosa guianensis</i>	538
<i>Micropholis chrysophylloides</i> var. <i>truncata</i>	691	<i>Mimosa hymenaeodes</i>	507
<i>Micropholis cowanii</i>	692	<i>Mimosa inaequalis</i>	545
<i>Micropholis cruegeriana</i>	691	<i>Mimosa inga</i>	518, 519
<i>Micropholis curvata</i>	691, 692	<i>Mimosa ingoides</i>	520
<i>Micropholis cyrtobotrya</i>	691	<i>Mimosa latifolia</i>	545
<i>Micropholis discolor</i>	692	<i>Mimosa leucocephala</i>	527
<i>Micropholis dominicensis</i>	692	<i>Mimosa lucida</i>	523
<i>Micropholis egensis</i>	691	<i>Mimosa macroloba</i>	535
<i>Micropholis eggersiana</i>	691	<i>Mimosa macrophylla</i>	521
<i>Micropholis eugenifolia</i>	697	<i>Mimosa niopo</i>	505

Appendix 3. — Continuation.

<i>Mimosa nitida</i>	523	<i>Misanteca appelli</i>	480
<i>Mimosa pacay</i>	517, 518	<i>Misanteca cannella</i>	480
<i>Mimosa parae</i>	520	<i>Misanteca cayennensis</i>	480
<i>Mimosa parvifolia</i>	505	<i>Misanteca chrysophylla</i>	480
<i>Mimosa pendula</i>	534	<i>Misanteca debilis</i>	480
<i>Mimosa peregrina</i>	505	<i>Misanteca martiniana</i>	481
<i>Mimosa pilosula</i>	523	<i>Misanteca meisneri</i>	481
<i>Mimosa polyphylla</i>	537	<i>Misanteca pachycarpa</i>	481
<i>Mimosa pulcherrima</i>	538	<i>Misanteca polyphylla</i>	481
<i>Mimosa quassiiifolia</i>	523	<i>Misanteca rigida</i>	480
<i>Mimosa ramiflora</i>	545	<i>Misanteca subbullata</i>	481
<i>Mimosa rubiginosa</i>	524	<i>Misanteca vernicosa</i>	481
<i>Mimosa semialata</i>	522	<i>Misanteca wilhelminensis</i>	481
<i>Mimosa sericea</i>	524	<i>Mitranthes eugeniooides</i>	626
<i>Mimosa splendens</i>	525	<i>Mitranthes eugeniooides</i> var. <i>oblongifolia</i>	626
<i>Mimosa terminalis</i>	505	<i>Mitranthes eugeniooides</i> var. <i>ovata</i>	626
<i>Mimosa trapezifolia</i>	503, 504	<i>Mitranthes gardneriana</i>	626
<i>Mimosa umbellata</i>	526	<i>Mitranthes sartoriana</i>	626
<i>Mimosa umbellifera</i>	526	<i>Mitropsidium eugeniooides</i>	626
<i>Mimosa virgultosa</i>	527	<i>Mitropsidium gardnerianum</i>	626
<i>Mimosa wilsonii</i>	514	<i>Mitropsidium ob lanceolatum</i>	626
<i>Mimosa yng'a</i>	518, 519	<i>Mitropsidium oligospermum</i>	626
<i>Mimosa zygia</i>	545	<i>Mitropsidium pittieri</i>	626
<i>Mimusops</i>	689, 690	<i>Mitropsidium sartorianum</i>	626
<i>Mimusops amazonica</i>	690	<i>Mollinedia casca</i>	582
<i>Mimusops balata</i>	689, 690	<i>Mollinedia glabricalis</i>	583
<i>Mimusops balata</i> var. <i>cruegeri</i>	689	<i>Mollinedia grandifolia</i>	582
<i>Mimusops balata</i> var. <i>domingensis</i>	690	<i>Mollinedia grazielae</i>	582
<i>Mimusops balata</i> var. <i>gutta</i>	689	<i>Mollinedia grosseserrata</i>	582
<i>Mimusops balata</i> var. <i>hartii</i>	689	<i>Mollinedia krukovii</i>	582
<i>Mimusops balata</i> var. <i>melinonii</i>	689, 690	<i>Mollinedia latifolia</i>	582
<i>Mimusops balata</i> var. <i>schomburgkii</i>	689	<i>Mollinedia laurina</i>	582
<i>Mimusops balata</i> var. <i>sieberi</i>	689	<i>Mollinedia neblinensis</i>	583
<i>Mimusops bidentata</i>	689, 690	<i>Mollinedia ovata</i>	582
<i>Mimusops darienensis</i>	689	<i>Mollinedia ptariensis</i>	582
<i>Mimusops domingensis</i>	690	<i>Mollinedia rusbyana</i>	582
<i>Mimusops huberi</i>	690	<i>Mollinedia tessmannii</i>	582
<i>Mimusops longiciliata</i>	690	<i>Monadelphanthus floridus</i>	651
<i>Mimusops maparajuba</i>	690	MONIMIACEAE	582
<i>Mimusops nitida</i>	690	<i>Monopteryx</i>	503, 530
<i>Mimusops paraensis</i>	690	<i>Monopteryx inpa</i>	530, 531
<i>Mimusops paraensis</i> var. <i>densiflora</i>	690	<i>Monteverdia floribunda</i>	415
<i>Mimusops paraensis</i> var. <i>discolor</i>	690	<i>Monteverdia myrsinoides</i>	415
<i>Mimusops riedeliana</i>	690	<i>Monteverdia oblongata</i>	415
<i>Mimusops riedleana</i>	690	<i>Moquilea Aubletiana</i>	416
<i>Mimusops sieberi</i>	689	<i>Moquilea bothynophylla</i>	426
<i>Mimusops siqueiraei</i>	690	<i>Moquilea bracteosa</i>	417
<i>Mimusops surinamensis</i>	690	<i>Moquilea canomensis</i>	419
<i>Minquartia guianensis</i>	634	<i>Moquilea couepia</i>	417
<i>Minquartia macrophylla</i>	634	<i>Moquilea floribunda</i>	425
<i>Minquartia parvifolia</i>	634	<i>Moquilea glandulosa</i>	417
<i>Minquartia punctata</i>	634	<i>Moquilea guianensis</i>	416, 430
<i>Misanteca amara</i>	481	<i>Moquilea kuntzeana</i>	425
<i>Misanteca anacardioides</i>	481	<i>Moquilea licaniiiflora</i>	432

Appendix 3. — Continuation.

<i>Moquilea longistyla</i>	425	<i>Mouriri crassifolia</i>	570
<i>Moquilea minutiflora</i>	430	<i>Mouriri duckeana</i>	570
<i>Moquilea orinocensis</i>	425	<i>Mouriri dumetosa</i>	570
<i>Moquilea parillo</i>	419	<i>Mouriri francavilliana</i>	570
<i>Moquilea parviflora</i>	432	<i>Mouriri grandiflora</i>	570
<i>Moquilea pendula</i>	425	<i>Mouriri guianensis</i>	570
<i>Moquilea riparia</i>	430	<i>Mouriri huberi</i>	571
<i>Moquilea</i> sp. A	430	<i>Mouriri macrophylla</i>	570
<i>Moquilea sprucei</i>	426	<i>Mouriri nervosa</i>	571
<i>Moquilea turiuva</i>	426	<i>Mouriri nigra</i>	571
<i>Moquilea utilis</i>	426	<i>Mouriri oligantha</i>	571
MORACEAE	583	<i>Mouriri pendulifolia</i>	570
<i>Morinda angustifolia</i>	661	<i>Mouriri plasschaertii</i>	571
<i>Morinda aspera</i>	662	<i>Mouriri polyantha</i>	571
<i>Morinda chachuca</i>	661	<i>Mouriri princeps</i>	570
<i>Morinda citrifolia</i>	661, 662	<i>Mouriri sagotiana</i>	571, 606
<i>Morinda citrifolia</i> f. <i>potteri</i>	662	<i>Mouriri sideroxylon</i>	571
<i>Morinda coreia</i> var. <i>stenophylla</i>	661	<i>Mouriri subumbellata</i>	571
<i>Morinda ligulata</i>	662	<i>Mouriri ulei</i>	571
<i>Morinda littoralis</i>	662	<i>Mouriri vernicosa</i>	571
<i>Morinda macrophylla</i>	661	<i>Mouriri weddelli</i>	571
<i>Morinda mudia</i>	661	<i>Moutouchi crispata</i>	536
<i>Morinda multiflora</i>	661	<i>Moutouchi draco</i>	536
<i>Morinda nodosa</i>	661	<i>Moutouchi suberosa</i>	536
<i>Morinda pubescens</i> var. <i>aspera</i>	662	<i>Mussaenda glomerulata</i>	650
<i>Morinda quadrangularis</i>	662	<i>Mussaenda spinosa</i>	665
<i>Morinda stenophylla</i>	661	<i>Mutisiopersea nivea</i>	490
<i>Morinda teysmanniana</i>	662	<i>Myginda myrsinoides</i>	415
<i>Morinda tinctoria</i>	661	<i>Myrcia</i>	609
<i>Morinda tomentosa</i>	661	<i>Myrcia acuminata</i>	619
<i>Morinda zollingeriana</i>	662	<i>Myrcia acuminata</i> var. <i>bullata</i>	619
<i>Morisonia</i>	411, 412	<i>Myrcia acuminata</i> var. <i>genuina</i>	619
<i>Morisonia frondosa</i>	411	<i>Myrcia acuminata</i> var. <i>meridensis</i>	619
<i>Morisonia pittieri</i>	412	<i>Myrcia acuminata</i> var. <i>peruviana</i>	619
<i>Morisonia polyantha</i>	412	<i>Myrcia acuminata</i> var. <i>tovarensis</i>	619
<i>Morisonia pulcherrima</i>	411	<i>Myrcia acutata</i>	620
<i>Morisonia sola</i>	411	<i>Myrcia acutiloba</i>	621
<i>Moronobea coccinea</i>	435	<i>Myrcia adpresepilosa</i>	615
<i>Moronobea esculenta</i>	435	<i>Myrcia aguitensis</i>	622
<i>Moronobea globulifera</i>	435	<i>Myrcia alagoensis</i>	620
<i>Moronobea grandiflora</i>	435	<i>Myrcia alagoensis</i> var. <i>intermedia</i>	620
<i>Moronobea macoubea</i>	435	<i>Myrcia alagoensis</i> var. <i>oblongata</i>	620
<i>Moronobea montana</i>	435	<i>Myrcia alagoensis</i> var. <i>ovata</i>	620
<i>Moschoxylum cipo</i>	579	<i>Myrcia alloiota</i>	623
<i>Moschoxylum pentandrum</i>	581	<i>Myrcia alloiota</i> var. <i>obovata</i>	623
<i>Moschoxylum propinquum</i>	581	<i>Myrcia alloiota</i> var. <i>subcordata</i>	623
<i>Moschoxylum surinamense</i>	582	<i>Myrcia alternifolia</i>	613
<i>Mouriri abnormis</i>	572	<i>Myrcia amazonica</i>	609
<i>Mouriri acutiflora</i>	570	<i>Myrcia ambigua</i>	622
<i>Mouriri acutiflora</i> var. <i>amblyodon</i>	570	<i>Myrcia ambigua</i> var. <i>dives</i>	622
<i>Mouriri acutiflora</i> var. <i>oligantha</i>	570	<i>Myrcia ambigua</i> var. <i>latifolia</i>	622
<i>Mouriri angulicosta</i>	570	<i>Myrcia ambigua</i> var. <i>multiflora</i>	622
<i>Mouriri anomala</i>	570	<i>Myrcia ambigua</i> var. <i>pauciflora</i>	622
<i>Mouriri collocarpa</i>	570	<i>Myrcia ambigua</i> var. <i>rostrata</i>	622

Appendix 3. — Continuation.

<i>Myrcia ambigua</i> var. <i>sylvatica</i>	622	<i>Myrcia daphnoides</i> var. <i>nervosa</i>	615
<i>Myrcia amshoffae</i>	610	<i>Myrcia decorticans</i>	611
<i>Myrcia androsaemoides</i>	615	<i>Myrcia decrescens</i>	614
<i>Myrcia androsaemoides</i> var. <i>parvifolia</i>	615	<i>Myrcia deflexa</i>	611, 612
<i>Myrcia angustifolia</i>	614	<i>Myrcia deflexa</i> var. <i>dussii</i>	612
<i>Myrcia arimensis</i>	615	<i>Myrcia desertorum</i>	614
<i>Myrcia attenuata</i>	610	<i>Myrcia detergens</i>	609
<i>Myrcia augustana</i>	621	<i>Myrcia diaphanosticta</i>	615
<i>Myrcia ayresiana</i>	619	<i>Myrcia dictyoneura</i>	621
<i>Myrcia barrensis</i>	621	<i>Myrcia didrichseniana</i>	615
<i>Myrcia belizensis</i>	622	<i>Myrcia discolor</i>	619
<i>Myrcia berberis</i>	619, 620	<i>Myrcia divaricata</i>	618
<i>Myrcia berberis</i> var. <i>angustifolia</i>	620	<i>Myrcia divergens</i>	618
<i>Myrcia berberis</i> var. <i>latifolia</i>	620	<i>Myrcia doloresensis</i>	602
<i>Myrcia botrys</i>	614	<i>Myrcia doniana</i>	606
<i>Myrcia brachylopadia</i>	621	<i>Myrcia duchassaingiana</i>	611
<i>Myrcia bracteata</i>	610	<i>Myrcia dumosa</i>	617
<i>Myrcia bracteolaris</i>	618	<i>Myrcia duriuscula</i>	611
<i>Myrcia brandamii</i>	620	<i>Myrcia dussii</i>	612
<i>Myrcia caerulescens</i>	616	<i>Myrcia edulis</i>	617
<i>Myrcia camapuana</i>	615	<i>Myrcia elaeodendra</i>	613
<i>Myrcia camaraeana</i>	616	<i>Myrcia elegans</i>	613
<i>Myrcia capivarhyensis</i>	623	<i>Myrcia ellipticifolia</i>	616
<i>Myrcia carnea</i>	611	<i>Myrcia elongata</i>	620
<i>Myrcia cassinioides</i>	613	<i>Myrcia elongata</i> var. <i>brunnea</i>	620
<i>Myrcia catharinæ</i>	621	<i>Myrcia elongata</i> var. <i>grandifolia</i>	620
<i>Myrcia chilensis</i>	619	<i>Myrcia elongata</i> var. <i>ochracea</i>	620
<i>Myrcia chrysophylla</i>	611	<i>Myrcia emarginata</i>	614
<i>Myrcia ciarensis</i>	621	<i>Myrcia erythroxylon</i>	599, 620
<i>Myrcia collina</i>	615	<i>Myrcia erythroxylon</i> var. <i>caeruleascens</i>	620
<i>Myrcia communis</i>	621	<i>Myrcia erythroxylon</i> var. <i>virescens</i>	599
<i>Myrcia communis</i> var. <i>glabrata</i>	621	<i>Myrcia exsucca</i>	613
<i>Myrcia communis</i> var. <i>latifolia</i>	621	<i>Myrcia fallax</i>	618
<i>Myrcia complicata</i>	618	<i>Myrcia fasciculata</i>	612
<i>Myrcia compressa</i>	622	<i>Myrcia fastigiata</i>	615
<i>Myrcia corcovadensis</i>	621	<i>Myrcia fastigiata</i> var. <i>coriacea</i>	615
<i>Myrcia coroicensis</i>	621	<i>Myrcia ferruginea</i>	611, 612
<i>Myrcia corymbosa</i>	609	<i>Myrcia ferruginea</i> var. <i>domingensis</i>	611
<i>Myrcia costaricensis</i>	619	<i>Myrcia floribunda</i>	623
<i>Myrcia costata</i>	619, 620, 622	<i>Myrcia formosiana</i>	619
<i>Myrcia costata</i> var. <i>bahiensis</i>	620	<i>Myrcia forsteri</i>	612
<i>Myrcia costata</i> var. <i>minensis</i>	620	<i>Myrcia fosteri</i>	612
<i>Myrcia coumete</i>	611	<i>Myrcia friburgensis</i>	620
<i>Myrcia coumetoides</i>	619	<i>Myrcia frontinensis</i>	617
<i>Myrcia crassicaulis</i>	613	<i>Myrcia frontinensis</i> var. <i>gamaeana</i>	617
<i>Myrcia crassinervia</i>	611	<i>Myrcia gamaeana</i>	617
<i>Myrcia cucullata</i>	619	<i>Myrcia gardneriana</i>	621
<i>Myrcia cuneata</i>	613	<i>Myrcia gentlei</i>	609
<i>Myrcia cuprea</i>	611	<i>Myrcia gigas</i>	612
<i>Myrcia curatellifolia</i>	622, 623	<i>Myrcia glaberrima</i>	616
<i>Myrcia curatellifolia</i> var. <i>grandifolia</i>	623	<i>Myrcia glandulosa</i>	614
<i>Myrcia curatellifolia</i> var. <i>parvifolia</i>	623	<i>Myrcia glandulosa</i> var. <i>obovata</i>	614
<i>Myrcia cymosopaniculata</i>	615	<i>Myrcia glaucescens</i>	616
<i>Myrcia daphnoides</i>	613, 615	<i>Myrcia graciliflora</i>	612

Appendix 3. — Continuation.

<i>Myrcia gracilis</i>	620, 621	<i>Myrcia leptoclada</i>	609
<i>Myrcia gracilis</i> var. <i>opaca</i>	620	<i>Myrcia leptoclada</i> var. <i>glazioviana</i>	609
<i>Myrcia gracilis</i> var. <i>prasina</i>	620	<i>Myrcia leucadendron</i>	613
<i>Myrcia gracilis</i> var. <i>sessiliflora</i>	621	<i>Myrcia leucantha</i>	623
<i>Myrcia granulata</i>	625	<i>Myrcia leucophlaea</i>	611
<i>Myrcia guajavifolia</i>	620, 621	<i>Myrcia lindeniana</i>	619
<i>Myrcia guajavifolia</i> f. <i>grandifolia</i>	621	<i>Myrcia lingua</i>	614
<i>Myrcia guajavifolia</i> var. <i>bullata</i>	620	<i>Myrcia lingua</i> var. <i>rufa</i>	614
<i>Myrcia guajavifolia</i> var. <i>impunctata</i>	620	<i>Myrcia longicaudata</i>	622
<i>Myrcia guajavifolia</i> var. <i>perforata</i>	620	<i>Myrcia longipes</i>	623
<i>Myrcia guianensis</i>	613	<i>Myrcia longipes</i> f. <i>obovata</i>	623
<i>Myrcia guianensis</i> var. <i>cuneata</i>	613	<i>Myrcia lucida</i>	616
<i>Myrcia hayneana</i>	619, 621	<i>Myrcia luetzelburgii</i>	621
<i>Myrcia hayneana</i> var. <i>paraensis</i>	621	<i>Myrcia lundiana</i>	609
<i>Myrcia hepatica</i>	614	<i>Myrcia macrophylla</i>	619
<i>Myrcia heringeriana</i>	616	<i>Myrcia magnoliifolia</i>	619, 620
<i>Myrcia hiemalis</i>	613	<i>Myrcia magnoliifolia</i> var. <i>angustifolia</i>	620
<i>Myrcia hirtellifolia</i>	611	<i>Myrcia magnoliifolia</i> var. <i>latifolia</i>	620
<i>Myrcia hirtiflora</i>	622	<i>Myrcia magnoliifolia</i> var. <i>parvifolia</i>	620
<i>Myrcia hostmanniana</i>	609	<i>Myrcia mansoni</i>	614
<i>Myrcia hostmanniana</i> var. <i>robustior</i>	609	<i>Myrcia marahanensis</i>	623
<i>Myrcia humboldtiana</i>	611, 612	<i>Myrcia martiana</i>	620
<i>Myrcia humboldtiana</i> var. <i>caribaea</i>	612	<i>Myrcia martinicensis</i>	621
<i>Myrcia humboldtiana</i> var. <i>orinocensis</i>	612	<i>Myrcia melanoclada</i>	619
<i>Myrcia impressa</i>	621	<i>Myrcia melanosticta</i>	621
<i>Myrcia inaequiloba</i>	615	<i>Myrcia membranacea</i>	623
<i>Myrcia incisa</i>	615	<i>Myrcia micrantha</i>	620
<i>Myrcia kegeliana</i>	620	<i>Myrcia microcarpa</i>	613
<i>Myrcia kegeliana</i> var. <i>angustifolia</i>	620	<i>Myrcia mikaniiana</i>	619, 621
<i>Myrcia kegeliana</i> var. <i>latifolia</i>	620	<i>Myrcia mikaniiana</i> var. <i>angustifolia</i>	621
<i>Myrcia kegeliana</i> var. <i>longifolia</i>	620	<i>Myrcia mikaniiana</i> var. <i>latifolia</i>	621
<i>Myrcia kegeliana</i> var. <i>pendula</i>	620	<i>Myrcia mini</i>	598
<i>Myrcia kegeliana</i> var. <i>vulgaris</i>	620	<i>Myrcia minoriflora</i>	609
<i>Myrcia klotzschiana</i>	620, 621	<i>Myrcia minutiflora</i>	616
<i>Myrcia klotzschiana</i> var. <i>impellucida</i>	621	<i>Myrcia multiflora</i>	616
<i>Myrcia kunthiana</i>	619	<i>Myrcia multiflora</i> f. <i>ovalifolia</i>	616
<i>Myrcia laevigata</i>	620, 621	<i>Myrcia multiflora</i> var. <i>glaucescens</i>	616
<i>Myrcia laevigata</i> var. <i>brunnea</i>	620	<i>Myrcia multiflora</i> var. <i>ramulosa</i>	616
<i>Myrcia laevigata</i> var. <i>canescens</i>	621	<i>Myrcia multiglomerata</i>	617
<i>Myrcia laevis</i>	616	<i>Myrcia myoporina</i>	613
<i>Myrcia lamprosericea</i>	621	<i>Myrcia negrensis</i>	621
<i>Myrcia lancea</i>	623	<i>Myrcia neoforsteri</i>	612
<i>Myrcia lanceolata</i>	610, 611	<i>Myrcia neolucida</i>	617
<i>Myrcia lanceolata</i> var. <i>angustifolia</i>	610	<i>Myrcia neomontana</i>	617
<i>Myrcia lanceolata</i> var. <i>avenia</i>	611	<i>Myrcia neospeciosa</i>	617
<i>Myrcia lanceolata</i> var. <i>grandifolia</i>	610	<i>Myrcia nigropunctata</i>	614
<i>Myrcia lanceolata</i> var. <i>latifolia</i>	611	<i>Myrcia nitens</i>	621
<i>Myrcia lanceolata</i> var. <i>racemosa</i>	611	<i>Myrcianthes elegans</i>	613
<i>Myrcia langsdorffii</i>	621	<i>Myrcianthes montana</i>	617
<i>Myrcia lanuginosa</i>	623	<i>Myrcianthes prodigiosa</i>	624
<i>Myrcia lasiopus</i>	611	<i>Myrcianthes puberula</i>	623
<i>Myrcia latifolia</i>	620	<i>Myrcianthes spixiana</i>	613
<i>Myrcia lauriflora</i>	613	<i>Myrcianthes terminalis</i>	615
<i>Myrcia lehmannii</i>	615	<i>Myrcia obscura</i>	614, 615

Appendix 3. — Continuation.

<i>Myrcia obscura</i> var. <i>longipes</i>	615	<i>Myrciaria ferruginea</i>	624
<i>Myrcia obtecta</i>	613, 614	<i>Myrciaria floribunda</i>	624
<i>Myrcia obtecta</i> var. <i>alternifolia</i>	613	<i>Myrciaria hagendorffii</i>	625
<i>Myrcia obtusa</i>	613	<i>Myrciaria leucadendron</i>	624
<i>Myrcia orstediana</i>	619	<i>Myrciaria leucophloea</i>	624, 625
<i>Myrcia oitchi</i>	609	<i>Myrciaria leucophloea</i> var. <i>conferta</i>	624
<i>Myrcia oocarpa</i>	619	<i>Myrciaria leucophloea</i> var. <i>laxa</i>	624
<i>Myrcia opaca</i>	621	<i>Myrciaria leucophloea</i> var. <i>warmingiana</i>	625
<i>Myrcia opaca</i> var. <i>angustifolia</i>	621	<i>Myrciaria longicaudata</i>	625
<i>Myrcia opaca</i> var. <i>latifolia</i>	621	<i>Myrciaria longipes</i>	624
<i>Myrcia ovalifolia</i>	616	<i>Myrciaria longipes</i> var. <i>opaca</i>	624
<i>Myrcia oxyoentophylla</i>	621	<i>Myrciaria longipes</i> var. <i>pellucida</i>	624
<i>Myrcia paivae</i>	617	<i>Myrciaria maragnanensis</i>	624
<i>Myrcia paivae</i> var. <i>gracilis</i>	617	<i>Myrciaria maranhensis</i>	624
<i>Myrcia pallens</i>	613, 614	<i>Myrciaria marowynensis</i>	603
<i>Myrcia pallens</i> var. <i>ovalis</i>	614	<i>Myrciaria maximiliana</i>	624
<i>Myrcia pallens</i> var. <i>subcordata</i>	614	<i>Myrciaria mexicana</i>	625
<i>Myrcia pallida</i>	616	<i>Myrciaria nitida</i>	615
<i>Myrcia paracatuensis</i>	615	<i>Myrciaria nitida</i> var. <i>chartacea</i>	615
<i>Myrcia paraensis</i>	609	<i>Myrciaria nitida</i> var. <i>coriacea</i>	615
<i>Myrcia patula</i>	602	<i>Myrciaria nitida</i> var. <i>dives</i>	615
<i>Myrcia pellucida</i>	620	<i>Myrciaria oneillii</i>	625
<i>Myrcia phaeoclada</i>	620	<i>Myrciaria polyantha</i>	615
<i>Myrcia phaeoclada</i> var. <i>alagoensis</i>	620	<i>Myrciaria prasina</i>	624
<i>Myrcia phaeoclada</i> var. <i>guyanensis</i>	620	<i>Myrciaria protracta</i>	624
<i>Myrcia piauhiensis</i>	623	<i>Myrciaria ramiflora</i>	606
<i>Myrcia pilosa</i>	622	<i>Myrciaria rivularis</i>	625
<i>Myrcia platyclada</i>	617	<i>Myrciaria rivularis</i> var. <i>baporeti</i>	625
<i>Myrcia plicatocostata</i>	620	<i>Myrciaria salzmannii</i>	624
<i>Myrcia plumbea</i>	614	<i>Myrciaria schaueriana</i>	612
<i>Myrcia poeppigiana</i>	614	<i>Myrciaria schuechiana</i>	624
<i>Myrcia polyantha</i>	618	<i>Myrciaria schuechiana</i> var. <i>deflexa</i>	624
<i>Myrcia prunifolia</i>	622, 623	<i>Myrciaria schuechiana</i> var. <i>latifolia</i>	624
<i>Myrcia prunifolia</i> var. <i>angustior</i>	622	<i>Myrciaria sellowiana</i>	624
<i>Myrcia prunifolia</i> var. <i>obovata</i>	623	<i>Myrciaria splendens</i>	624
<i>Myrcia prunifolia</i> var. <i>ovata</i>	623	<i>Myrciaria tenuiramis</i>	624
<i>Myrcia pseudomini</i>	619	<i>Myrciaria tolypantha</i>	624
<i>Myrcia puberula</i>	623	<i>Myrciaria tolypantha</i> var. <i>angustifolia</i>	624
<i>Myrcia pulchella</i>	618	<i>Myrciaria tolypantha</i> var. <i>latifolia</i>	624
<i>Myrcia pullei</i>	618	<i>Myrciaria uliginosa</i>	624
<i>Myrcia pusilla</i>	614	<i>Myrciaria verticillata</i>	624
<i>Myrcia pyrifolia</i>	618	<i>Myrciaria vismiifolia</i>	625
<i>Myrcia queimadensis</i>	613	<i>Myrcia riedeliana</i>	621
<i>Myrcia renatoana</i>	615	<i>Myrciariopsis baporeti</i>	625
<i>Myrcia reticulata</i>	619	<i>Myrcia riparia</i>	621
<i>Myrcia rhabdoides</i>	615	<i>Myrcia roraimae</i>	615
<i>Myrcia rhodeosepala</i>	623	<i>Myrcia roraimensis</i>	613
<i>Myrciaria amazonica</i>	624	<i>Myrcia rostrata</i>	619-622
<i>Myrciaria arborea</i>	625	<i>Myrcia rostrata</i> f. <i>communis</i>	621
<i>Myrciaria arborea</i> var. <i>rostrata</i>	625	<i>Myrcia rostrata</i> f. <i>flexuosa</i>	622
<i>Myrciaria axillaris</i>	624	<i>Myrcia rostrata</i> f. <i>gracilis</i>	620
<i>Myrciaria baporeti</i>	625	<i>Myrcia rostrata</i> f. <i>pseudomini</i>	619
<i>Myrciaria ciliolata</i>	624, 625	<i>Myrcia rostrata</i> f. <i>sericiflora</i>	621
<i>Myrciaria ciliolata</i> var. <i>warmingiana</i>	625	<i>Myrcia rostrata</i> var. <i>brunea</i>	619

Appendix 3. — Continuation.

<i>Myrcia rosulans</i>	623	<i>Myrcia torta</i>	613, 615
<i>Myrcia rotundifolia</i>	614	<i>Myrcia torta</i> f. <i>glaucia</i>	615
<i>Myrcia rufidula</i>	619	<i>Myrcia vacciniifolia</i>	614
<i>Myrcia rufula</i>	619, 620	<i>Myrcia velhensis</i>	615
<i>Myrcia rufula</i> var. <i>martiana</i>	620	<i>Myrcia velutina</i>	621
<i>Myrcia sagotii</i>	609	<i>Myrcia velutina</i> var. <i>canescens</i>	621
<i>Myrcia sartoriana</i>	621	<i>Myrcia velutina</i> var. <i>ochracea</i>	621
<i>Myrcia saxatilis</i>	618	<i>Myrcia venezuelensis</i>	619
<i>Myrcia saxicola</i>	619, 621	<i>Myrcia venosa</i>	623
<i>Myrcia saxicola</i> var. <i>grandifolia</i>	621	<i>Myrcia verruculata</i>	623
<i>Myrcia schippii</i>	622	<i>Myrcia ypanemensis</i>	621
<i>Myrcia schomburgkiana</i>	598	<i>Myrcia yungasensis</i>	615
<i>Myrcia schrankiana</i>	613	<i>Myrcia</i> sp. A	624
<i>Myrcia schuechiana</i>	621	<i>Myrcia</i> sp. B	624
<i>Myrcia scrobiculata</i>	615	<i>Myriaspore decipiens</i>	560
<i>Myrcia sellowiana</i>	621	<i>Myriaspore egensis</i>	559
<i>Myrcia sellowiana</i> var. <i>bullata</i>	621	<i>Myriaspore integrifolia</i>	560
<i>Myrcia sellowiana</i> var. <i>costata</i>	621	<i>Myriaspore paulensis</i>	559
<i>Myrcia sepiaria</i>	619	<i>Myriaspore surinamensis</i>	559
<i>Myrcia sericea</i>	620	<i>Myristica americana</i>	596
<i>Myrcia sericiflora</i>	621	<i>MYRISTICACEAE</i>	594
<i>Myrcia servata</i>	618	<i>Myristica cordifolia</i>	595
<i>Myrcia silvatica</i>	625	<i>Myristica fatua</i>	596
<i>Myrcia sororia</i>	619	<i>Myristica fulva</i>	596
<i>Myrcia spathulata</i>	623	<i>Myristica hostmannii</i>	594
<i>Myrcia sphaerocarpa</i>	616	<i>Myristica melinonii</i>	595
<i>Myrcia spixiana</i>	613	<i>Myristica mocoia</i>	595
<i>Myrcia splendens</i>	598, 618, 619, 621, 622	<i>Myristica mouchico</i>	594
<i>Myrcia splendens</i> var. <i>chrysocoma</i>	622	<i>Myristica mouchigo</i>	594
<i>Myrcia splendens</i> var. <i>genuina</i>	619	<i>Myristica panamensis</i>	595
<i>Myrcia splendens</i> var. <i>guantanamana</i>	622	<i>Myristica platysperma</i>	595
<i>Myrcia splendens</i> var. <i>micropora</i>	619	<i>Myristica sagotiana</i>	594
<i>Myrcia splendens</i> var. <i>mini</i>	598	<i>Myristica sebifera</i>	595, 596
<i>Myrcia splendens</i> var. <i>obscura</i>	619	<i>Myristica sebifera</i> var. <i>cordifolia</i>	595
<i>Myrcia splendens</i> var. <i>robustior</i>	621	<i>Myristica sebifera</i> var. <i>curvinervia</i>	595
<i>Myrcia spruceana</i>	620	<i>Myristica sebifera</i> var. <i>longifolia</i>	596
<i>Myrcia sprucei</i>	609	<i>Myristica surinamensis</i>	596
<i>Myrcia stemmeriana</i>	615	<i>Myristica virola</i>	595
<i>Myrcia suaveolens</i>	613	<i>Myrobalanus badamia</i>	441
<i>Myrcia subobliqua</i>	622	<i>Myrobalanus catappa</i>	440
<i>Myrcia subsessilis</i>	622	<i>Myrobalanus commersonii</i>	441
<i>Myrcia subsessilis</i> var. <i>ovalis</i>	622	<i>Myrobalanus eichleriana</i>	441
<i>Myrcia subsessilis</i> var. <i>subcordata</i>	622	<i>Myrobalanus excelsa</i>	440
<i>Myrcia sulcata</i>	612	<i>Myrobalanus guianensis</i>	441
<i>Myrcia superba</i>	621	<i>Myrobalanus lucida</i>	441
<i>Myrcia surinamensis</i>	613	<i>Myrobalanus nitidissima</i>	442
<i>Myrcia sylvatica</i>	622	<i>Myrobalanus oblonga</i>	443
<i>Myrcia taubatensis</i>	616	<i>Myrobalanus obovata</i>	440
<i>Myrcia taubatensis</i> var. <i>ovalis</i>	616	<i>Myrobalanus pamea</i>	440
<i>Myrcia terebinthacea</i>	614	<i>Myrobalanus rubrigemmis</i>	441
<i>Myrcia tetramera</i>	611	<i>Myrobalanus scutifera</i>	441
<i>Myrcia tingens</i>	621	<i>Myrobalanus terminalia</i>	441
<i>Myrcia tobagensis</i>	611	<i>Myrodendrum amplexicaule</i>	467
<i>Myrcia tomentosa</i>	622	<i>Myrodendrum balsamiferum</i>	467

Appendix 3. — Continuation.

<i>Myroxylon benthamii</i>	679	<i>Myrtus latifolia</i>	603
<i>Myroxylon ciliatifolium</i>	679	<i>Myrtus lauriflora</i>	613
<i>Myroxylon digynum</i>	679	<i>Myrtus leucadendra</i>	608
<i>Myroxylon lucens</i>	679	<i>Myrtus lucida</i>	622
<i>Myrsine guianensis</i>	645	<i>Myrtus lugens</i>	604
<i>Myrsine ovalifolia</i>	645	<i>Myrtus micrantha</i>	624
<i>Myrsine rapanea</i>	645, 646	<i>Myrtus mikaniiana</i>	619
<i>Myrsine rapanea</i> f. <i>angustata</i>	645	<i>Myrtus mini</i>	598
<i>Myrsine rapanea</i> f. <i>communis</i>	645	<i>Myrtus multiflora</i>	616
<i>Myrsine rapanea</i> f. <i>paviflora</i>	645	<i>Myrtus muricata</i>	604
<i>Myrsine rapanea</i> f. <i>robusta</i>	645	<i>Myrtus pallens</i>	598
<i>Myrsine rapanea</i> var. <i>major</i>	645	<i>Myrtus patrisii</i>	605
<i>Myrsine rapanea</i> var. <i>minor</i>	645	<i>Myrtus paucinervia</i>	626
<i>Myrsine rapanea</i> var. <i>subacuta</i>	645	<i>Myrtus pilosa</i>	622
MYRTACEAE	597, 626, 627	<i>Myrtus polyantha</i>	618, 619
<i>Myrtaceae</i> sp. A	626	<i>Myrtus pseudopsidium</i>	605
<i>Myrtaceae</i> sp. B	626	<i>Myrtus psidioides</i>	597
<i>Myrtaceae</i> sp. C	627	<i>Myrtus pyrifolia</i>	613
<i>Myrtilluma eugeniifolia</i>	697	<i>Myrtus revoluta</i>	597
<i>Myrtus acuminata</i>	619	<i>Myrtus riparia</i>	604
<i>Myrtus alba</i>	609	<i>Myrtus rostrata</i>	619
<i>Myrtus amazonica</i>	609	<i>Myrtus ruizii</i>	619
<i>Myrtus aubletii</i>	622	<i>Myrtus saligna</i>	608
<i>Myrtus berberis</i>	619	<i>Myrtus saxicola</i>	619
<i>Myrtus berteroana</i>	599	<i>Myrtus sepiaria</i>	619
<i>Myrtus biflora</i>	598, 599	<i>Myrtus sinemariensis</i>	606
<i>Myrtus biflora</i> f. <i>subsericea</i>	599	<i>Myrtus sparsiflora</i>	597
<i>Myrtus biflora</i> var. <i>salicifolia</i>	599	<i>Myrtus splendens</i>	618
<i>Myrtus biflora</i> var. <i>yapacani</i>	599	<i>Myrtus stoupyi</i>	618, 622
<i>Myrtus bracteata</i>	610	<i>Myrtus sylvatica</i>	622
<i>Myrtus bracteolaris</i>	618, 622	<i>Myrtus torta</i>	613
<i>Myrtus camaraeana</i>	616	<i>Myrtus umbellata</i>	571
<i>Myrtus carnea</i>	611	<i>Myrtus verruculosa</i>	605
<i>Myrtus cayennensis</i>	600	<i>Myrtus verticillata</i>	624
<i>Myrtus claraensis</i>	626	<i>Myrtus virgultosa</i>	598
<i>Myrtus complicata</i>	618	<i>Myrtus willdenowii</i> var. <i>portoricensis</i>	605
<i>Myrtus costata</i>	619		
<i>Myrtus coumete</i>	611		
<i>Myrtus curatellifolia</i>	622		
<i>Myrtus deflexa</i>	611, 612		
<i>Myrtus duartii</i>	613	<i>Napimoga guianensis</i>	677
<i>Myrtus egensis</i>	601	<i>Naucleopsis guianensis</i>	592
<i>Myrtus elegans</i>	613	<i>Nectandra acutangula</i>	486
<i>Myrtus exsucca</i>	613	<i>Nectandra albiflora</i>	483
<i>Myrtus fascicularis</i>	597	<i>Nectandra amazonum</i>	482
<i>Myrtus ferruginea</i>	612	<i>Nectandra ambigua</i>	482
<i>Myrtus flavescens</i>	602	<i>Nectandra antillana</i>	483
<i>Myrtus floribunda</i>	624	<i>Nectandra bombycina</i>	482
<i>Myrtus florida</i>	602	<i>Nectandra bredemeyeriana</i>	482
<i>Myrtus goetheana</i>	597	<i>Nectandra capanahuensis</i>	482
<i>Myrtus guianensis</i>	613	<i>Nectandra caudata</i>	481
<i>Myrtus hayneana</i>	619	<i>Nectandra cissiflora</i>	482
<i>Myrtus inocarpa</i>	605	<i>Nectandra discolor</i> var. <i>subvenosa</i>	483
<i>Myrtus lancea</i>	599	<i>Nectandra floribunda</i>	485

Appendix 3. — Continuation.

<i>Nectandra glabrescens</i>	482	<i>Nectandra willdenoviana</i>	482
<i>Nectandra globosa</i>	482	<i>Neea caulinflora</i>	627
<i>Nectandra globosa</i> var. <i>barbeyana</i>	482	<i>Neea constricta</i>	627
<i>Nectandra grandiflora</i> var. <i>latifolia</i>	483	<i>Neea floribunda</i>	627
<i>Nectandra grandis</i>	490	<i>Neea mollis</i>	627
<i>Nectandra guanaiensis</i>	483	<i>Neea ovalifolia</i>	627
<i>Nectandra hihua</i>	482	<i>Neea</i> sp. A	628
<i>Nectandra kunthiana</i>	491	<i>Neea</i> sp. B	628
<i>Nectandra kuntzeana</i>	482	<i>Neea</i> sp. C	628
<i>Nectandra latifolia</i>	478	<i>Neea spruceana</i>	628
<i>Nectandra laurel</i> var. <i>triquetra</i>	483	<i>Neea weberbaueri</i>	628
<i>Nectandra leucantha</i> var. <i>attenuata</i>	483	<i>Neleixa prostrata</i>	657
<i>Nectandra leucantha</i> var. <i>guianensis</i>	483	<i>Nemaluma engleri</i>	697
<i>Nectandra leucantha</i> var. <i>peruviana</i>	483	<i>Nematospermum laevigatum</i>	471
<i>Nectandra longifolia</i>	486	<i>Neocalyptrocalyx leprieurii</i>	412
<i>Nectandra lucida</i>	482	<i>Neocalyptrocalyx maroniensis</i>	412
<i>Nectandra magnoliifolia</i>	482	<i>Neocalyptrocalyx morii</i>	412
<i>Nectandra maranonensis</i>	483	<i>Neocalyptrocalyx surinamensis</i>	412
<i>Nectandra matogrossensis</i>	483	<i>Neohuberia matamata</i>	495
<i>Nectandra megaphylla</i>	483	<i>Neolabatia glomerata</i>	698
<i>Nectandra meyeriana</i>	491	<i>Neolabatia hypoglauca</i>	698
<i>Nectandra mollis</i>	483	<i>Neolabatia macrocarpa</i>	699
<i>Nectandra mollis</i> var. <i>attenuata</i>	483	<i>Neolabatia parviflora</i>	698
<i>Nectandra mollis</i> var. <i>intermedia</i>	483	<i>Neopometia ptychandra</i>	703
<i>Nectandra mollis</i> var. <i>venosa</i>	483	<i>Neoraputia cowanii</i>	668
<i>Nectandra mollis</i> var. <i>villosa</i>	483	<i>Neoraputia paraensis</i>	667, 668
<i>Nectandra myriantha</i>	482	<i>Neoxythecia cladantha</i>	695
<i>Nectandra myriantha</i> var. <i>attenuata</i>	482	<i>Neoxythecia dura</i>	695
<i>Nectandra myriantha</i> var. <i>glabrata</i>	482	<i>Neoxythecia robusta</i>	696
<i>Nectandra pallida</i>	482	<i>Neoxythecia schulzii</i>	695
<i>Nectandra paulii</i>	482	<i>Neoxythecia wurdackii</i>	695
<i>Nectandra pisi</i>	482	<i>Newtonia psilostachya</i>	535
<i>Nectandra pittieri</i>	483	<i>Newtonia suaveolens</i>	536
<i>Nectandra polita</i>	479	<i>Niopa peregrina</i>	505
<i>Nectandra polita</i> var. <i>oerstedii</i>	479	<i>Nonatelia cataractarum</i>	662
<i>Nectandra polyphylla</i>	481	<i>Nonatelia lutea</i>	653
<i>Nectandra pulverulenta</i>	483	<i>Nonatelia sessiliflora</i>	652
<i>Nectandra purpurea</i>	478	<i>Noyeria rubra</i>	593
<i>Nectandra purusensis</i>	483	NYCTAGINACEAE	627, 628
<i>Nectandra reticulata</i>	483	<i>Nyctaginaceae</i> sp. A	628
<i>Nectandra revoluta</i>	484	<i>Nyctaginaceae</i> sp. B	628
<i>Nectandra rubra</i>	491	<i>Nyctaginaceae</i> sp. C	628
<i>Nectandra schomburgkii</i>	483	<i>Nyctaginaceae</i> sp. D	628
<i>Nectandra</i> sp. A	483	O	
<i>Nectandra steinbachii</i>	482	OCHNACEAE	628
<i>Nectandra surinamensis</i>	482, 486	<i>Ocotea</i>	483
<i>Nectandra tessmannii</i>	483	<i>Ocotea acutangula</i>	486
<i>Nectandra urophylla</i>	482	<i>Ocotea amazonica</i>	483
<i>Nectandra vaga</i>	482	<i>Ocotea arenaensis</i>	486
<i>Nectandra vaga</i> var. <i>major</i>	482	<i>Ocotea argyrophylla</i>	484
<i>Nectandra vaga</i> var. <i>sprucei</i>	482	<i>Ocotea baturitensis</i>	488
<i>Nectandra vaga</i> var. <i>vulgaris</i>	482		
<i>Nectandra villosa</i>	483		
<i>Nectandra villosa</i> var. <i>venosa</i>	483		

Appendix 3. — Continuation.

<i>Ocotea botryophylla</i>	486	<i>Ocotea pauciflora</i>	487
<i>Ocotea canaliculata</i>	484	<i>Ocotea percurrens</i>	487
<i>Ocotea castanea</i>	489	<i>Ocotea persulcata</i>	487
<i>Ocotea caudata</i>	484	<i>Ocotea puberula</i>	488
<i>Ocotea ceanothifolia</i>	484	<i>Ocotea puberula</i> var. <i>truncata</i>	488
<i>Ocotea cernua</i>	484	<i>Ocotea punctulata</i>	479
<i>Ocotea cinerea</i>	484	<i>Ocotea pyramidata</i>	488
<i>Ocotea citrifolia</i>	489	<i>Ocotea rubra</i>	491
<i>Ocotea commutata</i>	485	<i>Ocotea rusbyana</i>	487
<i>Ocotea complicata</i>	485	<i>Ocotea scabrella</i>	489
<i>Ocotea cooperi</i>	491	<i>Ocotea schomburgkiana</i>	489
<i>Ocotea cujumary</i>	485	<i>Ocotea scrobiculata</i>	484
<i>Ocotea dielsiana</i>	487	<i>Ocotea scrobiculifera</i>	485
<i>Ocotea diffusa</i>	485	<i>Ocotea splendens</i>	489
<i>Ocotea dissimilis</i>	489	<i>Ocotea subalpina</i>	484
<i>Ocotea duckei</i>	485	<i>Ocotea subglabra</i>	488
<i>Ocotea duotincta</i>	486	<i>Ocotea subracemosa</i>	486
<i>Ocotea endlicheriopsis</i>	485	<i>Ocotea subsericea</i>	486
<i>Ocotea fasciculata</i>	485	<i>Ocotea subterminalis</i>	489
<i>Ocotea fendleri</i>	485	<i>Ocotea tomentella</i>	489
<i>Ocotea flavescens</i>	475	<i>Ocotea trianae</i>	487
<i>Ocotea flexuosa</i>	479	<i>Ocotea ucayalensis</i>	488
<i>Ocotea floribunda</i>	485	<i>Ocotea urophylla</i>	484
<i>Ocotea froesii</i>	486	<i>Ocotea vernicosa</i>	481
<i>Ocotea glandulosa</i>	485	<i>Ocotea wachenheimii</i>	486
<i>Ocotea globifera</i>	489	<i>Ocotea</i> sp. A	489
<i>Ocotea globosa</i>	482	<i>Ocotea</i> sp. B	489
<i>Ocotea glomerata</i>	486	<i>Ocotea</i> sp. C	489
<i>Ocotea grandifolia</i>	487	<i>Ocotea</i> sp. D	489
<i>Ocotea guianensis</i>	486	<i>Ocotea</i> sp. E	489
<i>Ocotea guianensis</i> var. <i>aurea</i>	486	<i>Ocotea</i> sp. F	490
<i>Ocotea guianensis</i> var. <i>subsericea</i>	486	<i>Ocotea</i> sp. G	490
<i>Ocotea kunthiana</i>	491	<i>Ocotea</i> sp. H	490
<i>Ocotea lasseriana</i>	486	<i>Ocotea</i> sp. I	490
<i>Ocotea latifolia</i>	478	<i>Ocotea</i> sp. J	490
<i>Ocotea laxiflora</i>	487	<i>Octavia sessiliflora</i>	652
<i>Ocotea lenticellata</i>	486	<i>Odina francoana</i>	365
<i>Ocotea leptobotra</i>	484	<i>Odontandra quadrijuga</i>	581
<i>Ocotea leucoxylon</i>	486	<i>Oedematopus octandrus</i>	433
<i>Ocotea leucoxylon</i> var. <i>elongata</i>	486	<i>Oenocarpus bacaba</i>	394
<i>Ocotea longifolia</i>	486	<i>Oenocarpus bacaba</i> var. <i>grandis</i>	394
<i>Ocotea macropoda</i>	488	<i>Oenocarpus bataua</i> var. <i>bataua</i>	394
<i>Ocotea marmellensis</i> var. <i>acrensis</i>	484	<i>Oenocarpus bataua</i> var. <i>oligocarpus</i>	394
<i>Ocotea marowynensis</i>	484	<i>Oenocarpus grandis</i>	394
<i>Ocotea martiniana</i>	488	<i>Oenocarpus hoppii</i>	394
<i>Ocotea mollis</i>	483	<i>Oenocarpus oligocarpa</i>	394
<i>Ocotea montis-insulae</i>	487	<i>Oenocarpus seje</i>	394
<i>Ocotea mucronata</i>	487	<i>Ogcodeia guianensis</i>	592
<i>Ocotea nigra</i>	487	OLACACEAE	633
<i>Ocotea nunesii</i>	485	<i>Olax</i>	633
<i>Ocotea oblonga</i>	487	<i>Olax guianensis</i>	633
<i>Ocotea opifera</i>	487	<i>Olax schomburgkii</i>	633
<i>Ocotea paraensis</i>	488	OLEACEAE	635
<i>Ocotea paranapiacabensis</i>	488	<i>Olmedia affinis</i>	592

Appendix 3. — Continuation.

<i>Olmedia asperula</i>	592	<i>Oreodaphne pauciflora</i>	487
<i>Olmedia calophylla</i>	592	<i>Oreodaphne retroflexa</i>	485
<i>Olmedia caurensis</i>	593	<i>Oreodaphne schomburgkiana</i>	489
<i>Olmedia ferruginea</i>	593	<i>Oreodaphne splendens</i>	489
<i>Olmedia grandifolia</i>	593	<i>Oreodaphne subalpina</i>	484
<i>Olmedia guianensis</i>	592	<i>Oreodaphne urophylla</i>	484
<i>Olmedia habas</i>	593	<i>Oreodaphne willdenoviana</i>	485
<i>Olmedia laevis</i>	593	<i>Oreopanax capitatus</i>	390
<i>Olmedia maquira</i>	592	<i>Oreopanax capitatus</i> var. <i>minor</i>	390
<i>Olmedia poeppigiana</i>	592	<i>Oreopanax destructor</i>	390
<i>Olmedia polycephala</i>	592	<i>Oreopanax liebmannii</i>	390
<i>Olmedia tomentosa</i>	592	<i>Oreopanax meiocephalus</i>	390
<i>Olmedioperebea calophylla</i>	592	<i>Oreopanax morototoni</i>	389
<i>Olmedioperebea sclerophylla</i>	592	<i>Oreopanax oligocarpus</i>	390
<i>Olmediopsis lanceolata</i>	593	<i>Ormosia</i>	503, 531
<i>Olmediopsis obliqua</i>	593	<i>Ormosia bolivarensis</i>	531
<i>Oluntos trigona</i>	591	<i>Ormosia cinerea</i>	531
<i>Omphalea glandulata</i>	465	<i>Ormosia coarctata</i>	531
<i>Oncoba maynensis</i>	363	<i>Ormosia coccinea</i>	531
<i>Oncoba maynensis</i> var. <i>laxiflora</i>	363	<i>Ormosia costulata</i>	531
<i>Ophiocaryon chironectes</i>	671	<i>Ormosia costulata</i> var. <i>trifoliolata</i>	531
<i>Ophiocaryon paradoxum</i>	671	<i>Ormosia coutinhoi</i>	531
OPILIACEAE	635	<i>Ormosia crassicarpa</i>	532
<i>Oreodaphne acutangula</i>	486	<i>Ormosia cuneata</i>	531
<i>Oreodaphne acutifolia</i> var. <i>latifolia</i>	488	<i>Ormosia flava</i>	531
<i>Oreodaphne alba</i>	475	<i>Ormosia heterophylla</i>	532
<i>Oreodaphne amazonica</i>	483	<i>Ormosia lignivalvis</i>	532
<i>Oreodaphne caudata</i>	484	<i>Ormosia melanocarpa</i>	532
<i>Oreodaphne cayennensis</i>	484	<i>Ormosia nobilis</i>	531, 532
<i>Oreodaphne cernua</i>	484	<i>Ormosia nobilis</i> var. <i>boliviensis</i>	531
<i>Oreodaphne commutata</i>	485	<i>Ormosia pacimonensis</i>	544
<i>Oreodaphne diospyrifolia</i> var. <i>incompta</i>	487	<i>Ormosia paraensis</i>	532
<i>Oreodaphne divaricata</i>	485	<i>Ormosia stipularis</i>	532
<i>Oreodaphne domingensis</i>	485	<i>Ormosia trifoliolata</i>	531
<i>Oreodaphne fasciculata</i>	485	<i>Ormosia</i> sp. A	532
<i>Oreodaphne glomerata</i>	486	<i>Ormosia</i> sp. B	532
<i>Oreodaphne grandifolia</i>	487	<i>Ormosia</i> sp. C	532
<i>Oreodaphne guianensis</i>	486	<i>Ormosiopsis flava</i>	531
<i>Oreodaphne guianensis</i> var. <i>argentea</i>	486	<i>Ossaea flavescens</i>	560
<i>Oreodaphne guianensis</i> var. <i>aurea</i>	486	<i>Osteophloeum platyspermum</i>	595
<i>Oreodaphne hookeriana</i>	482	<i>Ouratea candellei</i>	629
<i>Oreodaphne hostmanniana</i>	488	<i>Ouratea cerebroidea</i>	629
<i>Oreodaphne leptobotra</i>	484	<i>Ouratea decagyna</i>	629
<i>Oreodaphne leucoxylon</i>	486	<i>Ouratea francinae</i>	629
<i>Oreodaphne leucoxylon</i> var. <i>elongata</i>	486	<i>Ouratea guianensis</i>	629
<i>Oreodaphne lindeniana</i>	486	<i>Ouratea leblondii</i>	630
<i>Oreodaphne longifolia</i>	487	<i>Ouratea macrocarpa</i>	630
<i>Oreodaphne macrothyrsus</i>	485	<i>Ouratea melinonii</i>	630
<i>Oreodaphne marowynensis</i>	484	<i>Ouratea retrorsa</i>	630
<i>Oreodaphne martiniana</i>	488	<i>Ouratea sagotii</i>	630
<i>Oreodaphne moritziana</i>	486	<i>Ouratea scottii</i>	630
<i>Oreodaphne opifera</i>	487	<i>Outea acaciifolia</i>	528
<i>Oreodaphne paraensis</i>	487	<i>Outea guianensis</i>	529
<i>Oreodaphne parviflora</i>	486	<i>Outea multijuga</i>	529

Appendix 3. — Continuation.

<i>Oxandra asbeckii</i>	377	<i>Palicourea pedunculosa</i>	665
<i>Oxandra nervosa</i>	380	<i>Palma cocos</i>	392
<i>Oxandra xylopioides</i>	377	<i>Palma maripa</i>	392
<i>Oxisma arborescens</i>	559	<i>Palma moeia</i>	390
<i>Oxyanthus isthmia</i>	663	<i>Palma spinosa</i>	390
<i>Oxydectes cajucara</i>	459	<i>Paloue</i>	503, 530, 532, 533
<i>Oxydectes caryophyllus</i>	460	<i>Paloue brasiliensis</i>	532
<i>Oxydectes cuneatus</i>	459	<i>Paloue guianensis</i>	532
<i>Oxydectes draconoides</i>	460	<i>Paloue induta</i>	530, 532
<i>Oxydectes martii</i>	459	<i>Paloue leiogyne</i>	533
<i>Oxydectes matourensis</i>	460	<i>Paloue paraensis</i>	533
<i>Oxydectes palanostigma</i>	460	<i>Paloue princeps</i>	533
<i>Oxydectes schiedeana</i>	460	<i>Paloue riparia</i>	533
<i>Oxydectes surinamensis</i>	459	<i>Pamea guianensis</i>	440
<i>Oxymeris furfuracea</i>	566	<i>Panax decaphyllus</i>	388
<i>Oxythece ambelanifolia</i>	693	<i>Panax morototoni</i>	389
<i>Oxythece robusta</i>	696	<i>Panax splendens</i>	389
<i>Oxythece robusta</i> var. <i>brevipetiolata</i>	696	<i>Panax undulatus</i>	389
 P			
<i>Pachira aquatica</i>	556	<i>Pancratium caribaeum</i>	722
<i>Pachira aquatica</i> var. <i>occidentalis</i>	556	<i>Panopsis acuminata</i>	646
<i>Pachira aquatica</i> var. <i>surinamensis</i>	556	<i>Panopsis cuaensis</i>	646
<i>Pachira dolichocalyx</i>	556	<i>Panopsis hameliifolia</i>	647
<i>Pachira flaviflora</i>	556	<i>Panopsis rubescens</i>	646
<i>Pachira insignis</i>	556	<i>Panopsis rubescens</i> var. <i>simulans</i>	646
<i>Pachira macrocalyx</i>	556	<i>Panopsis rubescens</i> var. <i>sprucei</i>	646
<i>Pachira macrocarpa</i>	556	<i>Panopsis sessilifolia</i>	647
<i>Pachira pustulifera</i>	556	<i>Panopsis sprucei</i>	646
<i>Pachira spruceana</i>	556	<i>Panzera falcata</i>	514
<i>Pachira villosula</i>	556	<i>Papaya spinosa</i>	412
<i>Pagamea guianensis</i>	662	<i>Papilioideae</i>	503
<i>Pagamea guianensis</i> var. <i>angustifolia</i>	662	<i>Parahancornia amapa</i>	386
<i>Pagamea guianensis</i> var. <i>parviflora</i>	662	<i>Parahancornia fasciculata</i>	386
<i>Pagamea thrysiflora</i>	662	<i>Parahancornia tabernaemontana</i>	385
<i>Palala hostmannii</i>	594	<i>Paralabatia acutangula</i>	691
<i>Palala mocoa</i>	595	<i>Paralabatia belizensis</i>	697
<i>Palala panamensis</i>	595	<i>Paralabatia durlandii</i>	697
<i>Palala platysperma</i>	595	<i>Paralabatia potosina</i>	697
<i>Palala sebifera</i>	595	<i>Paralea guianensis</i>	449
<i>Palala surinamensis</i>	596	<i>Paramachaerium</i>	503, 533
<i>Palicourea barbinervia</i>	662	<i>Paramachaerium ormosioides</i>	533
<i>Palicourea brachyloba</i>	662	<i>Paramicropholis acutangula</i>	691
<i>Palicourea chionantha</i>	665	<i>Paramyriaria ciliolata</i>	624
<i>Palicourea elliptica</i>	664	<i>Paraprotium amazonicum</i>	400
<i>Palicourea guianensis</i>	662	<i>Parinari amazonica</i>	431
<i>Palicourea guianensis</i> f. <i>glabra</i>	662	<i>Parinari barbata</i>	419
<i>Palicourea guianensis</i> subsp. <i>barbinervia</i>	662	<i>Parinari brachystachya</i>	431
<i>Palicourea guianensis</i> subsp. <i>occidentalis</i>	662	<i>Parinari campestris</i>	430
<i>Palicourea guianensis</i> var. <i>glabrescens</i>	662	<i>Parinari elliottii</i>	431
<i>Palicourea guianensis</i> var. <i>tetramera</i>	662	<i>Parinari excelsa</i>	431
<i>Palicourea guianensis</i> var. <i>trimera</i>	662	<i>Parinari excelsa</i> subsp. <i>holstii</i>	431
		<i>Parinari excelsa</i> var. <i>fulvescens</i>	431
		<i>Parinari glazioviana</i>	431
		<i>Parinari guianensis</i>	584
		<i>Parinari guyanensis</i>	584

Appendix 3. — Continuation.

<i>Parinari holstii</i>	431	<i>Pausandra flagellarhachis</i>	463
<i>Parinari holstii</i> var. <i>longifolia</i>	431	<i>Pausandra fordii</i>	463
<i>Parinari laxiflora</i>	431	<i>Pausandra integrifolia</i>	461
<i>Parinari laxiflora</i> var. <i>lata</i>	431	<i>Pausandra martinii</i>	463
<i>Parinari lucidissima</i>	431	<i>Pausandra quadriglandulosa</i>	464
<i>Parinari mildbraedii</i>	431	<i>Pausandra sericea</i>	464
<i>Parinari montana</i>	431	<i>Pausandra trianae</i>	464
<i>Parinari nalaensis</i>	431	<i>Paypayrola bordenavei</i>	719
<i>Parinariopsis licaniflora</i>	432	<i>Paypayrola grandiflora</i>	719
<i>Parinari pajura</i>	431	<i>Paypayrola guianensis</i>	719
<i>Parinari parvifolia</i>	431	<i>Paypayrola hulkiana</i>	719
<i>Parinari pilosa</i>	419	<i>Paypayrola longifolia</i>	719
<i>Parinari riparia</i>	431	<i>Paypayrola ventricosa</i>	719
<i>Parinari rodolphii</i>	431	<i>Payrola guianensis</i>	719
<i>Parinari salicifolia</i>	431	<i>Peckia microbotrys</i>	644
<i>Parinari</i> sp. A	431	<i>Peckia nitida</i>	645
<i>Parinari sylvestris</i>	431	<i>Peckia prieurii</i>	645
<i>Parinari tenuifolia</i>	431	<i>Peckia purpurea</i>	645
<i>Parinari verdickii</i>	431	<i>Peckia subspicata</i>	645
<i>Parinari whytei</i>	431	<i>Peckia surinamensis</i>	645
<i>Pariti tiliaceum</i> f. <i>immaculatum</i>	558	<i>Peiranisia aristulata</i>	537
<i>Paritium pernambucense</i>	558	<i>Peiranisia multijuga</i>	537
<i>Parivoa grandiflora</i>	514	<i>Pekea butyrosa</i>	413
<i>Parivoa tomentosa</i>	511	<i>Pekea couroupita</i>	494
<i>Parkia</i>	503, 530	<i>Pekea ternata</i>	413
<i>Parkia alliodora</i>	534	<i>Pekea villosa</i>	413
<i>Parkia arborea</i>	533	<i>Peltogyne</i>	503, 534
<i>Parkia decussata</i>	533	<i>Peltogyne amplissima</i>	534
<i>Parkia gigantocarpa</i>	530, 533	<i>Peltogyne lecointei</i>	534
<i>Parkia ingens</i>	534	<i>Peltogyne paniculata</i>	534
<i>Parkia inundabilis</i>	534	<i>Peltogyne paniculata</i> subsp. <i>pubescens</i>	534
<i>Parkia microcephala</i>	534	<i>Peltogyne pubescens</i>	534
<i>Parkia nitida</i>	533	<i>Peltogyne</i> sp. A	535
<i>Parkia oppositifolia</i>	533	<i>Peltogyne venosa</i>	535
<i>Parkia paryphosphaera</i>	533	<i>Peltospermum latisiliquum</i>	381
<i>Parkia pendula</i>	534	<i>Peltospermum patrisii</i>	381
<i>Parkia reticulata</i>	534	<i>Pentaclethra</i>	503, 535
<i>Parkia</i> sp. A	534	<i>Pentaclethra brevifila</i>	535
<i>Parkia sylvatica</i>	534	<i>Pentaclethra filamentosa</i>	535
<i>Parkia ulei</i> var. <i>surinamensis</i>	534	<i>Pentaclethra macroloba</i>	535
<i>Parkia velutina</i>	534	<i>Pentamorpha graveolens</i>	667
<i>Paryphosphaera arborea</i>	533	<i>PENTAPHYLACACEAE</i>	635
<i>Passaveria lanceolata</i>	689	<i>Pentascyphus thrysiflorus</i>	683
<i>Passaveria obovata</i>	689	<i>Pera arborea</i>	636
<i>Passaveria obovata</i> var. <i>tomentosa</i>	689	<i>Pera bicolor</i>	636
<i>Passiflora foetida</i>	722	<i>PERACEAE</i>	635
<i>Passoura guianensis</i>	721	<i>Pera ferruginea</i>	636
<i>Patrisia bicolor</i>	678	<i>Pera glabrata</i>	636
<i>Patrisia parviflora</i>	679	<i>Pera schomburgkiana</i>	636
<i>Patrisia pyrifera</i>	678	<i>Perebea australis</i>	593
<i>Patrisia sagotiana</i>	678	<i>Perebea calophylla</i>	592
<i>Patrisia tomentosa</i>	679	<i>Perebea guianensis</i>	593
<i>Pausandra densiflora</i>	464	<i>Perebea laurifolia</i>	592
<i>Pausandra extorris</i>	464	<i>Perebea mollis</i> subsp. <i>rubra</i>	593

Appendix 3. — Continuation.

<i>Perebea rubra</i>	593	<i>Pharmacosycea rigida</i>	588
<i>Perebea xinguana</i>	592	<i>Phellocarpus floridus</i>	536
<i>Periclistia latifolia</i>	719	<i>Phellocarpus laxiflorus</i>	536
<i>Peridium bicolor</i>	636	<i>Phoebe antillana</i> var. <i>cubensis</i>	475
<i>Peridium ferrugineum</i>	636	<i>Phoebe brasiliensis</i>	475
<i>Peridium glabratum</i>	636	<i>Phoebe cinnamomifolia</i>	475
<i>Peridium oblongifolium</i>	636	<i>Phoebe cubensis</i>	475
<i>Peridium schomburgkianum</i>	636	<i>Phoebe elongata</i>	475
<i>Peridium schomburgkii</i>	636	<i>Phoebe filamentosa</i>	475
<i>Persea ceanothifolia</i>	484	<i>Phoebe fruticosa</i>	475
<i>Persea cinnamomifolia</i>	475	<i>Phoebe granatensis</i>	475
<i>Persea globosa</i>	482	<i>Phoebe heterotepala</i>	475
<i>Persea incana</i>	483	<i>Phoebe impressa</i>	474
<i>Persea latifolia</i>	478	<i>Phoebe johnstonii</i>	475
<i>Persea leucoxylon</i>	486	<i>Phoebe maynensis</i>	475
<i>Persea longifolia</i>	486	<i>Phoebe mexicana</i>	475
<i>Persea macropoda</i>	488	<i>Phoebe mexicana</i> var. <i>bourgeauana</i>	475
<i>Persea mexicana</i>	475	<i>Phoebe montana</i>	474
<i>Persea mollis</i>	483	<i>Phoebe paraguariensis</i>	475
<i>Persea montana</i>	474	<i>Phoebe peruviana</i>	474, 475
<i>Persea nivea</i>	490	<i>Phoebe peruviana</i> var. <i>glabriflora</i>	475
<i>Persea parviflora</i>	486	<i>Phoebe pichisensis</i>	475
<i>Persea retroflexa</i>	485	<i>Phoebe pickellii</i>	475
<i>Persea richardiana</i>	488	<i>Phoebe poeppigii</i>	475
<i>Persea tenuiflora</i>	484	<i>Phoebe triplinervis</i>	474, 475
<i>Persea triplinervis</i> var. <i>valenzuelana</i>	475	<i>Phoebe triplinervis</i> var. <i>cubensis</i>	475
<i>Persea</i> sp. A	490	<i>Phoebe valenzuelana</i>	475
<i>Persoonia guareoides</i>	573	<i>Pholacilia diversifolia</i>	580
<i>Peschiera benthamiana</i>	387	<i>Pholacilia trinitensis</i>	580
<i>Peschiera benthamiana</i> var. <i>myriantha</i>	387	<i>Phosanthus coccineus</i>	660
<i>Peschiera benthamiana</i> var. <i>stenantha</i>	387	PHYLLANTHACEAE	636
<i>Peschiera linkii</i>	387	<i>Phyllanthus antillanus</i>	638
<i>Peschiera multiflora</i>	387	<i>Phyllanthus antillanus</i> var. <i>hypomalacus</i>	638
<i>Peschiera myriantha</i>	387	<i>Phyllanthus antillanus</i> var. <i>pedicularis</i>	638
<i>Peschiera ochracea</i>	387	<i>Phyllanthus attenuatus</i>	638
<i>Peschiera stenantha</i>	387	<i>Phyllanthus guianensis</i>	638
<i>Peschiera surinamensis</i>	387	<i>Phyllanthus juglandifolius</i>	638
<i>Petaloma mouriri</i>	571	<i>Phyllanthus nobilis</i>	638
<i>Peteniodendron belicense</i>	697	<i>Phyllanthus nobilis</i> var. <i>antillanus</i>	638
<i>Peteniodendron durlandii</i>	697	<i>Phyllanthus nobilis</i> var. <i>brasiliensis</i>	638
<i>Peteniodendron potosinum</i>	697	<i>Phyllanthus nobilis</i> var. <i>guyanensis</i>	638
<i>Petrocarya campestris</i>	430	<i>Phyllanthus nobilis</i> var. <i>hypomalacus</i>	638
<i>Petrocarya excelsa</i>	431	<i>Phyllanthus nobilis</i> var. <i>martii</i>	638
<i>Petrocarya montana</i>	431	<i>Phyllanthus nobilis</i> var. <i>panamensis</i>	638
<i>Pharmacosycea glaucescens</i>	588	<i>Phyllanthus nobilis</i> var. <i>pavonianus</i>	638
<i>Pharmacosycea grandaeva</i>	588	<i>Phyllanthus nobilis</i> var. <i>peruvianus</i>	638
<i>Pharmacosycea guyanensis</i>	588	<i>Phyllanthus nobilis</i> var. <i>riedelianus</i>	638
<i>Pharmacosycea hernandezii</i>	588	<i>Phyllocalyx wentii</i>	607
<i>Pharmacosycea laurifolia</i>	590	<i>Phytolacca javanica</i>	440
<i>Pharmacosycea mexicana</i>	588	PICRAMNIACEAE	639
<i>Pharmacosycea parkeriana</i>	586	<i>Picramnia cooperi</i>	640
<i>Pharmacosycea peruviana</i>	590	<i>Picramnia eosina</i>	640
<i>Pharmacosycea pseudoradula</i>	588	<i>Picramnia guianensis</i>	639
<i>Pharmacosycea radula</i>	588	<i>Picramnia krukovii</i>	640

Appendix 3. — Continuation.

<i>Picramnia latifolia</i>	639	<i>Piper fasciculare</i>	471
<i>Picramnia longissima</i>	639	<i>Piper fatoanum</i>	640
<i>Picramnia macrostachys</i>	639	<i>Piper flavescentia</i>	640
<i>Picramnia sellowii</i> subsp. <i>spruceana</i>	640	<i>Piper gigantifolium</i>	642
<i>Picramnia spruceana</i>	640	<i>Piper guanaianum</i>	640
<i>Picramnia tariri</i>	639	<i>Piper herzogii</i>	640
<i>Picramnia tenuis</i>	640	<i>Piper illudens</i>	641
<i>Picramnia umbrosa</i>	639	<i>Piper intersitum</i>	641
<i>Picrolemma pseudocoffea</i>	705	<i>Piper intersitum</i> f. <i>porcecitense</i>	641
<i>Picrolemma sprucei</i>	705	<i>Piper kuntzei</i>	640
<i>Pilocarpus latifolius</i>	668	<i>Piper latum</i>	642
<i>Pilocarpus racemosus</i>	668	<i>Piper lineatum</i> var. <i>hirtipetiolatum</i>	640
<i>Pilophora saccifera</i>	393	<i>Piper multinervium</i>	640, 641
<i>Pilophora testicularis</i>	393	<i>Piper multinervium</i> var. <i>amplum</i>	641
<i>Pimecaria odorata</i>	634	<i>Piper multinervium</i> var. <i>cayoense</i>	641
<i>Pionandra hartwegii</i>	708	<i>Piper multinervium</i> var. <i>flavicans</i>	641
<i>Piparea</i>	678	<i>Piper multinervium</i> var. <i>hirsuticaule</i>	641
<i>Piparea dentata</i>	678	<i>Piper multinervium</i> var. <i>kantelolense</i>	641
<i>Piparea multiflora</i>	678	<i>Piper multinervium</i> var. <i>paralense</i>	641
PIPERACEAE	640	<i>Piper multinervium</i> var. <i>peracutum</i>	641
<i>Piper aduncifolium</i>	640	<i>Piper multinervium</i> var. <i>productipes</i>	641
<i>Piper aduncum</i>	640	<i>Piper multinervium</i> var. <i>protractifolium</i>	641
<i>Piper aduncum</i> var. <i>brachyarthrum</i>	640	<i>Piper multinervium</i> var. <i>puberulipedunculum</i>	641
<i>Piper aduncum</i> var. <i>cordulatum</i>	640	<i>Piper multinervium</i> var. <i>pubescenticaule</i>	641
<i>Piper aduncum</i> var. <i>laevifolium</i>	640	<i>Piper multinervium</i> var. <i>skutchii</i>	641
<i>Piper aggregatum</i>	471	<i>Piper multinervium</i> var. <i>tamashense</i>	641
<i>Piper anguillispicum</i>	640	<i>Piper multinervium</i> var. <i>telanum</i>	641
<i>Piper angustifolium</i>	640	<i>Piper nonconformans</i>	640
<i>Piper angustifolium</i> var. <i>cordulatum</i>	640	<i>Piper ob lanceolatum</i> var. <i>fragilicaule</i>	640
<i>Piper araguense</i>	642	<i>Piper obliquum</i> var. <i>eximium</i>	642
<i>Piper arbelaezii</i> var. <i>subglabrirameum</i>	642	<i>Piper ovalifolium</i>	642
<i>Piper boliviianum</i>	642	<i>Piper pangoense</i>	642
<i>Piper cabellense</i>	642	<i>Piper perlongispicum</i>	642
<i>Piper cardenasii</i>	640	<i>Piper pseudovelutinum</i> var. <i>flavescens</i>	640
<i>Piper celtidifolium</i>	640	<i>Piper purpurascens</i>	640
<i>Piper cernuum</i>	642	<i>Piper reciprocum</i>	641
<i>Piper cernuum</i> var. <i>araguense</i>	642	<i>Piper reticulatum</i>	642
<i>Piper cernuum</i> var. <i>biformipilum</i>	642	<i>Piper reticulatum</i> var. <i>santarosanum</i>	642
<i>Piper cernuum</i> var. <i>glabricaule</i>	642	<i>Piper richardiifolium</i>	642
<i>Piper cernuum</i> var. <i>perlongispicum</i>	642	<i>Piper smilacifolium</i>	642
<i>Piper coccoboides</i>	642	<i>Piper stevensonii</i>	640
<i>Piper consanguineum</i>	640	<i>Piper submolle</i>	640
<i>Piper cuatrecasasii</i>	641	<i>Piper tarapotianum</i>	642
<i>Piper cumbricola</i>	641	<i>Piper tiliifolium</i>	583
<i>Piper discophorum</i>	642	<i>Piptadenia cateniformis</i>	508
<i>Piper disparispicum</i>	640	<i>Piptadenia cobi</i>	538
<i>Piper duchassaingii</i>	642	<i>Piptadenia guianensis</i>	538
<i>Piper elongatifolium</i>	640	<i>Piptadenia niopo</i>	505
<i>Piper elongatum</i>	640	<i>Piptadenia peregrina</i>	505
<i>Piper elongatum</i> var. <i>brachyarthrum</i>	640	<i>Piptadenia polystachya</i>	538
<i>Piper elongatum</i> var. <i>cordulatum</i>	640	<i>Piptadenia psilospatha</i>	535
<i>Piper elongatum</i> var. <i>laevifolium</i>	640	<i>Piptadenia suaveolens</i>	536
<i>Piper elongatum</i> var. <i>pampayacusum</i>	640	<i>Piratinera acutifolia</i>	583
<i>Piper eximium</i>	642	<i>Piratinera discolor</i>	583

Appendix 3. — Continuation.

<i>Piratinera guianensis</i>	583	<i>Pithecellobium subcorymbosum</i>	515
<i>Piratinera lancifera</i>	584	<i>Pithecellobium trapezifolium</i>	504
<i>Piratinera lemeei</i>	583	<i>Pitumba edulis</i>	674
<i>Piratinera mollis</i>	583	<i>Pitumba guianensis</i>	674
<i>Piratinera panamensis</i>	583	<i>Planchonella guianensis</i>	687
<i>Piratinera paraensis</i>	584	<i>Planchonella melinonii</i>	691
<i>Piratinera rubescens</i>	584	<i>Platanocephalus orientalis</i>	661
<i>Piratinera scabridula</i>	583	<i>Platonia esculenta</i>	435
<i>Piratinera velutina</i>	583	<i>Platonia insignis</i>	435
<i>Pirigara hexapetala</i>	499	<i>Platyluma calophylloides</i>	693
<i>Pirigara tetrapetala</i>	499	<i>Platymiscium</i>	503, 528
<i>Piscidia florida</i>	536	<i>Platymiscium dubium</i>	535
<i>Pisonia albiflora</i>	627	<i>Platymiscium duckei</i>	535
<i>Pisonia broadwayana</i>	627	<i>Platymiscium duckei</i> var. <i>nigrum</i>	535
<i>Pisonia eggersiana</i>	627	<i>Platymiscium filipes</i>	535
<i>Pisonia guianensis</i>	627	<i>Platymiscium nigrum</i>	535
<i>Pisonia salicifolia</i>	627	<i>Platymiscium pinnatum</i>	528, 535
<i>Pithecellobium acacioides</i>	509	<i>Platymiscium pinnatum</i> subsp. <i>polystachyum</i>	535
<i>Pithecellobium adiantifolium</i> var. <i>multipinnatum</i>	529	<i>Platymiscium polystachyum</i>	535
<i>Pithecellobium barbourianum</i>	503	<i>Platymiscium polystachyum</i> var. <i>fendleri</i>	535
<i>Pithecellobium benthamianum</i>	504	<i>Platymiscium trinitatis</i>	530, 535
<i>Pithecellobium bicolor</i>	545	<i>Platymiscium trinitatis</i> var. <i>duckei</i>	535
<i>Pithecellobium brongniartii</i>	503	<i>Platymiscium trinitatis</i> var. <i>nigrum</i>	535
<i>Pithecellobium caribaeum</i>	504	<i>Pleroma holosericea</i>	564
<i>Pithecellobium cataractae</i>	545	<i>Pleurothyrium chrysothyrsus</i>	491
<i>Pithecellobium cateniformis</i>	508	<i>Pleurothyrium cowanianum</i>	491
<i>Pithecellobium cauliflorum</i> f. <i>niveum</i>	545	<i>Pleuteron frondosa</i>	411
<i>Pithecellobium chagrense</i>	545	<i>Plinia acutissima</i>	625
<i>Pithecellobium corymbosum</i>	515	<i>Plinia baporeti</i>	625
<i>Pithecellobium corymbosum</i> var. <i>longipes</i>	515	<i>Plinia rivularis</i>	625
<i>Pithecellobium curvicarpum</i>	503	<i>Plumeria ambigua</i>	384
<i>Pithecellobium divaricatum</i>	545	<i>Plumeria articulata</i>	384
<i>Pithecellobium fanshawei</i>	503	<i>Plumeria bracteata</i>	384
<i>Pithecellobium foreroi</i>	545	<i>Plumeria floribunda</i>	384
<i>Pithecellobium glomeratum</i>	545	<i>Plumeria floribunda</i> var. <i>acutifolia</i>	384
<i>Pithecellobium hassleri</i>	504	<i>Plumeria floribunda</i> var. <i>calycina</i>	384
<i>Pithecellobium huberi</i>	545	<i>Plumeria floribunda</i> var. <i>crassipes</i>	384
<i>Pithecellobium inaequale</i>	545	<i>Plumeria lancifolia</i>	384
<i>Pithecellobium jupunba</i>	503	<i>Plumeria lancifolia</i> var. <i>major</i>	384
<i>Pithecellobium laetum</i>	504	<i>Plumeria lancifolia</i> var. <i>microphylla</i>	384
<i>Pithecellobium latifolium</i>	545	<i>Plumeria martii</i>	384
<i>Pithecellobium latifolium</i> subsp. <i>tayronense</i>	545	<i>Plumeria microcalyx</i>	384
<i>Pithecellobium longiflorum</i>	529	<i>Plumeria phagedaenica</i>	384
<i>Pithecellobium longiramosum</i>	545	<i>Plumeria speciosa</i>	384
<i>Pithecellobium mataybifolium</i>	504	<i>Plumeria sucuuba</i>	384
<i>Pithecellobium micradenium</i>	503	<i>Plumeria tarapotensis</i>	384
<i>Pithecellobium miquelianum</i>	529	<i>Pochota flaviflora</i>	556
<i>Pithecellobium niopoides</i>	504	<i>Podoluma benai</i>	694
<i>Pithecellobium pedicellare</i>	505	<i>Podoluma glaziovii</i>	694
<i>Pithecellobium pilosulum</i>	545	<i>Poecilanthe effusa</i>	505
<i>Pithecellobium polycarpum</i>	504	<i>Poecilanthe hostmannii</i>	527
<i>Pithecellobium racemiflorum</i>	546	<i>Poecilanthe ovalifolia</i>	505
<i>Pithecellobium racemosum</i>	546	<i>Pogonophora schomburgkiana</i>	636
<i>Pithecellobium schomburgkii</i>	514	<i>Pogonophora schomburgkiana</i> f. <i>elliptica</i>	636

Appendix 3. — Continuation.

<i>Pogonophora schomburgkiana</i> var. <i>longifolia</i>	636	<i>Pourouma cinerascens</i>	715
<i>Pogonophora trianae</i>	464	<i>Pourouma crassivenosa</i>	714
<i>Pogonophyllum elatum</i>	463	<i>Pourouma cuatrecasasii</i>	715
<i>Pogonorhynchus amplexans</i>	569	<i>Pourouma digitata</i>	714
<i>Pogonorhynchus sessillifolius</i>	569	<i>Pourouma folleata</i>	715
<i>Pogopetalum acutum</i>	582	<i>Pourouma fuliginea</i>	715
<i>Pohlana instrumentaria</i>	669	<i>Pourouma guianensis</i>	714
<i>Pohlana langsdorffii</i>	669	<i>Pourouma heterophylla</i>	715
<i>Polembryum castanocarpum</i>	667	<i>Pourouma isophlebia</i>	715
<i>Polembryum jussieui</i>	667	<i>Pourouma laevis</i>	716
POLYGONACEAE	642	<i>Pourouma lawrancei</i>	714
<i>Polygonum uvifera</i>	643	<i>Pourouma maroniensis</i>	716
<i>Poraqueiba guianensis</i>	582	<i>Pourouma melinonii</i>	715
<i>Poraqueiba surinamensis</i>	582	<i>Pourouma mildbraediana</i>	715
<i>Poraresia anomala</i>	636	<i>Pourouma minor</i>	715
<i>Portesia diversifolia</i>	580	<i>Pourouma mollis</i>	715
<i>Portesia echinocarpa</i>	580	<i>Pourouma palmata</i>	714
<i>Portesia montana</i>	580	<i>Pourouma radula</i>	715
<i>Portesia ovata</i>	580	<i>Pourouma saulensis</i>	715
<i>Portesia simplicifolia</i>	580	<i>Pourouma scabra</i>	715
<i>Portesia trinitensis</i>	580	<i>Pourouma schultesii</i>	714
<i>Portlandia acuminata</i>	655	<i>Pourouma steyermarkii</i>	716
<i>Portlandia hexandra</i>	655	<i>Pourouma subplicata</i>	715
<i>Portlandia speciosa</i>	655	<i>Pourouma substrigosa</i>	715
<i>Posoqueria acuminata</i>	663	<i>Pourouma subtriloba</i>	715
<i>Posoqueria decora</i>	663	<i>Pourouma tomentosa</i> subsp. <i>maroniensis</i>	—
<i>Posoqueria gracilis</i>	663	<i>Pourouma umbellata</i>	715
<i>Posoqueria insignis</i>	663	<i>Pourouma umbellifera</i>	715
<i>Posoqueria latifolia</i>	663	<i>Pourouma velutina</i>	716
<i>Posoqueria latifolia</i> subsp. <i>gracilis</i>	663	<i>Pourouma villosa</i>	716
<i>Posoqueria longiflora</i>	663	<i>Pourouma</i> sp. A	716
<i>Posoqueria lucida</i>	663	<i>Pourouma</i> sp. B	716
<i>Posoqueria macrophylla</i>	663	<i>Pouteria</i>	693
<i>Posoqueria macropus</i>	663	<i>Pouteria achradiformis</i>	691
<i>Posoqueria metensis</i>	663	<i>Pouteria acutangula</i>	691
<i>Posoqueria multiflora</i>	663	<i>Pouteria aeranea</i>	692
<i>Posoqueria mutisii</i>	663	<i>Pouteria ambelanifolia</i>	693
<i>Posoqueria panamensis</i>	663	<i>Pouteria anibifolia</i>	701
<i>Posoqueria panamensis</i> subsp. <i>grandiflora</i>	663	<i>Pouteria anomala</i>	693
<i>Posoqueria revoluta</i>	663	<i>Pouteria aubrevillei</i>	694
<i>Posoqueria spraguei</i>	663	<i>Pouteria auyantepuiensis</i>	689
<i>Possira arborescens</i>	539	<i>Pouteria balata</i>	692
<i>Possira dodecandra</i>	539	<i>Pouteria bangii</i>	694
<i>Possira triphylla</i>	539	<i>Pouteria benai</i>	694
<i>Potamoxylon fluviatile</i>	398	<i>Pouteria benoitii</i>	699
<i>Poulsenia</i> sp. A	593	<i>Pouteria bilocularis</i>	694
<i>Pourouma acutiflora</i>	714	<i>Pouteria bopiensis</i>	692
<i>Pourouma apaporiensis</i>	715	<i>Pouteria bouffardiana</i>	699
<i>Pourouma apaporiensis</i> var. <i>macrophylla</i>	715	<i>Pouteria brachyandra</i>	694
<i>Pourouma aspera</i>	714	<i>Pouteria caimito</i>	694, 699
<i>Pourouma aurea</i>	715	<i>Pouteria caimito</i> var. <i>laurifolia</i>	694
<i>Pourouma bicolor</i> subsp. <i>bicolor</i>	714	<i>Pouteria caimito</i> var. <i>strigosa</i>	699
<i>Pourouma bicolor</i> subsp. <i>digitata</i>	714	<i>Pouteria cataractae</i>	693
<i>Pourouma camaratana</i>	714	<i>Pouteria cayennensis</i>	695

Appendix 3. — Continuation.

<i>Pouteria cearensis</i>	701	<i>Pouteria leucophaea</i>	695
<i>Pouteria chrysophylloides</i>	691	<i>Pouteria loretensis</i>	695
<i>Pouteria cladantha</i>	695	<i>Pouteria lucentifolia</i>	687
<i>Pouteria congestifolia</i>	685	<i>Pouteria macrocarpa</i>	699
<i>Pouteria cooperi</i>	702	<i>Pouteria macrophylla</i>	699
<i>Pouteria coriacea</i>	695	<i>Pouteria marginata</i> var. <i>impressa</i>	—
<i>Pouteria cruegeriana</i>	691	<i>Pouteria maxima</i>	700
<i>Pouteria cuatrecasasii</i>	699	<i>Pouteria mayana</i>	687
<i>Pouteria cuspidata</i>	695, 696	<i>Pouteria melanopoda</i>	700
<i>Pouteria cuspidata</i> subsp. <i>dura</i>	695	<i>Pouteria melinoniana</i>	692
<i>Pouteria cuspidata</i> subsp. <i>robusta</i>	696	<i>Pouteria melinonii</i>	691
<i>Pouteria cyrtobotrya</i>	691	<i>Pouteria mensalis</i>	692
<i>Pouteria dasystyla</i>	702	<i>Pouteria meyeri</i>	701
<i>Pouteria dasystyla</i> var. <i>abaetensis</i>	702	<i>Pouteria micans</i>	694
<i>Pouteria decorticans</i>	696	<i>Pouteria minutiflora</i>	695
<i>Pouteria deliciosa</i>	696	<i>Pouteria neglecta</i>	702
<i>Pouteria demerarae</i>	698	<i>Pouteria nuda</i>	689
<i>Pouteria dibrachiata</i>	687	<i>Pouteria obidensis</i>	698
<i>Pouteria discolor</i>	692	<i>Pouteria oblanceolata</i>	700
<i>Pouteria dissepala</i>	702	<i>Pouteria paraensis</i>	699
<i>Pouteria dominicensis</i>	692	<i>Pouteria pedunculata</i>	698
<i>Pouteria duckeana</i>	692	<i>Pouteria pentasperma</i>	687
<i>Pouteria dura</i>	695	<i>Pouteria petenensis</i>	687
<i>Pouteria durlandii</i>	697	<i>Pouteria piresii</i>	692
<i>Pouteria echinocarpa</i>	702	<i>Pouteria pittieri</i>	698
<i>Pouteria egensis</i>	691	<i>Pouteria platyphylla</i>	700
<i>Pouteria egregia</i>	697	<i>Pouteria polyneura</i>	693
<i>Pouteria engleri</i>	697	<i>Pouteria pomifera</i>	687
<i>Pouteria ephedrantha</i>	697	<i>Pouteria porphyrocarpa</i>	693
<i>Pouteria eugeniifolia</i>	697	<i>Pouteria potosina</i>	697
<i>Pouteria excelsa</i>	700	<i>Pouteria procera</i>	700
<i>Pouteria filipes</i>	697	<i>Pouteria ptychandra</i>	703
<i>Pouteria fimbriata</i>	697	<i>Pouteria putamen-ovi</i>	700
<i>Pouteria flava</i>	693	<i>Pouteria reticulata</i>	700
<i>Pouteria flavilatex</i>	697	<i>Pouteria retinervis</i>	701
<i>Pouteria franciscana</i>	698	<i>Pouteria rivularis</i>	693
<i>Pouteria glazioviana</i>	698	<i>Pouteria robusta</i>	696
<i>Pouteria glomerata</i>	698	<i>Pouteria robusta</i> var. <i>longifolia</i>	696
<i>Pouteria glomerata</i> var. <i>glabrescens</i>	698	<i>Pouteria robusta</i> var. <i>typica</i>	696
<i>Pouteria glomerata</i> var. <i>typica</i>	698	<i>Pouteria rodriguesiana</i>	701
<i>Pouteria gonggrijpii</i>	698	<i>Pouteria rufa</i>	691
<i>Pouteria grandis</i>	698	<i>Pouteria sagotiana</i>	701
<i>Pouteria guianensis</i>	698	<i>Pouteria sanctae-rosae</i>	693
<i>Pouteria gutta</i>	702	<i>Pouteria scytalophora</i>	694
<i>Pouteria hartii</i>	698, 699	<i>Pouteria setosa</i>	691
<i>Pouteria hispida</i>	699	<i>Pouteria silvicola</i>	692
<i>Pouteria hondurensis</i>	701	<i>Pouteria singularis</i>	701
<i>Pouteria hypoglauca</i>	698	<i>Pouteria solimoesensis</i>	699
<i>Pouteria jariensis</i>	699	<i>Pouteria speciosa</i>	701
<i>Pouteria jenmanii</i>	702	<i>Pouteria steyermarkii</i>	697
<i>Pouteria klugii</i>	693	<i>Pouteria stipulifera</i>	701
<i>Pouteria laevigata</i>	699	<i>Pouteria suffulta</i>	697
<i>Pouteria lasiocarpa</i>	695	<i>Pouteria surinamensis</i>	704
<i>Pouteria laurifolia</i>	694	<i>Pouteria temare</i>	694

Appendix 3. — Continuation.

<i>Pouteria tenuisepala</i>	701	<i>Protium excelsior</i>	399
<i>Pouteria torta</i> subsp. <i>glabra</i>	—	<i>Protium fimbriatum</i>	400
<i>Pouteria torta</i> subsp. <i>tuberculata</i>	702	<i>Protium firmum</i>	400
<i>Pouteria tovarensis</i>	698	<i>Protium gallicum</i>	402
<i>Pouteria trichopoda</i>	699	<i>Protium giganteum</i> var. <i>crassifolium</i>	402
<i>Pouteria truncata</i>	691	<i>Protium giganteum</i> var. <i>giganteum</i>	402
<i>Pouteria ulei</i>	691	<i>Protium goudotianum</i>	402
<i>Pouteria unilocularis</i>	700	<i>Protium guianense</i>	402
<i>Pouteria venosa</i> subsp. <i>amazonica</i>	702	<i>Protium heptaphyllum</i>	402, 403
<i>Pouteria venulosa</i>	693	<i>Protium heptaphyllum</i> var. <i>floribundum</i>	403
<i>Pouteria virescens</i>	702	<i>Protium heptaphyllum</i> var. <i>multiflorum</i>	402
<i>Pouteria weddelliana</i>	698	<i>Protium heptaphyllum</i> var. <i>puberulum</i>	402
<i>Pouteria williamii</i>	702	<i>Protium heptaphyllum</i> var. <i>surinamense</i>	402
<i>Pouteria wurdackii</i>	702	<i>Protium heptaphyllum</i> var. <i>unifoliolatum</i>	403
<i>Pouteria</i> sp. A	702	<i>Protium hostmannii</i>	402, 403
<i>Pouteria</i> sp. B	703	<i>Protium hostmannii</i> var. <i>brasiliense</i>	403
<i>Pouteria</i> sp. C	703	<i>Protium inodorum</i>	403
<i>Pouteria</i> sp. D	703	<i>Protium insigne</i>	403
<i>Pouteria</i> sp. E	703	<i>Protium joannis</i>	405
<i>Pouteria</i> sp. F	703	<i>Protium martianum</i>	405
<i>Pouteria</i> sp. G	703	<i>Protium melinonii</i>	403
<i>Pouteria</i> sp. H	703	<i>Protium morii</i>	403
<i>Pouteria</i> sp. I	703	<i>Protium multiflorum</i>	402
<i>Pradosia cochlearia</i>	703	<i>Protium neglectum</i>	404, 405
<i>Pradosia huberi</i>	703	<i>Protium neglectum</i> var. <i>robustum</i>	404
<i>Pradosia ptychandra</i>	703	<i>Protium occultum</i>	403
<i>Pradosia subverticillata</i>	704	<i>Protium octandrum</i>	403
<i>Pradosia surinamensis</i>	704	<i>Protium opacum</i> subsp. <i>rabelianum</i>	403
<i>Pradosia verticillata</i>	704	<i>Protium orinocense</i>	401
<i>Pradosia</i> sp. A	704	<i>Protium pallidum</i>	403
<i>Preslianthes pittieri</i>	412	<i>Protium paraense</i>	400
<i>Prieurella cuneifolia</i>	686	<i>Protium pauciflorum</i>	405
<i>Prieurella manaosensis</i>	686	<i>Protium pernervatum</i>	402
<i>Prieurella prieurii</i>	687	<i>Protium picramnioides</i>	404
PRIMULACEAE	644	<i>Protium pilosum</i>	403
PROTEACEAE	646	<i>Protium plagiocarpium</i>	404
<i>Protium</i>	399	<i>Protium polybotryum</i>	404
<i>Protium almecega</i>	404	<i>Protium puberulentum</i>	400
<i>Protium altissimum</i>	399	<i>Protium rhoifolium</i>	404
<i>Protium altsonii</i>	400	<i>Protium robustum</i>	404
<i>Protium amazonicum</i>	400	<i>Protium sagotianum</i>	404
<i>Protium angustifolium</i>	403	<i>Protium schomburgkianum</i>	401
<i>Protium apiculatum</i>	400	<i>Protium spruceanum</i>	404
<i>Protium aracouchini</i>	400	<i>Protium stevensonii</i>	404
<i>Protium aracouchini</i> var. <i>angustifolium</i>	400	<i>Protium strumosum</i>	405
<i>Protium calendulinum</i>	400	<i>Protium subserratum</i>	405
<i>Protium crassifolium</i>	402	<i>Protium surinamense</i>	405
<i>Protium cuneatum</i>	400	<i>Protium tenuifolium</i>	405
<i>Protium cupreatum</i>	402	<i>Protium titubans</i>	405
<i>Protium decandrum</i>	400	<i>Protium trifoliolatum</i>	405
<i>Protium demerarensis</i>	401	<i>Prunus accumulans</i>	649
<i>Protium divaricatum</i> subsp. <i>fumarium</i>	402	<i>Prunus myrtifolia</i>	649, 650
<i>Protium divaricatum</i> var. <i>intermedium</i>	400	<i>Prunus myrtifolia</i> var. <i>accumulans</i>	649
<i>Protium duckei</i>	402	<i>Prunus sphaerocarpa</i>	650

Appendix 3. — Continuation.

<i>Prunus tikalana</i>	650	<i>Psidium persoonii</i>	625
<i>Pseudima frutescens</i>	681, 683	<i>Psidium protractum</i>	626
<i>Pseudobombax amapaense</i>	556	<i>Psidium salutare</i>	626
<i>Pseudobombax munguba</i>	556	<i>Psidium sartorianum</i>	626
<i>Pseudochimarrhis difformis</i>	652	<i>Psidium sartorianum</i> var. <i>yucatanense</i>	626
<i>Pseudochimarrhis turbinata</i>	652	<i>Psidium socorrense</i>	626
<i>Pseudocladia colombiana</i>	694	<i>Psidium solisii</i>	626
<i>Pseudocladia melinonii</i>	694	<i>Psidium tenuifolium</i>	597
<i>Pseudocladia minutiflora</i>	695	<i>Psidium yucatanense</i>	626
<i>Pseudocladia neblinaensis</i>	694	<i>Psychotria anceps</i>	664
<i>Pseudocladia scytalophora</i>	694	<i>Psychotria anceps</i> var. <i>robustior</i>	664
<i>Pseudolabatia filipes</i>	697	<i>Psychotria ardisiifolia</i>	664
<i>Pseudolabatia raoulantonia</i>	697	<i>Psychotria boliviiana</i>	656
<i>Pseudolmedia alnifolia</i>	593	<i>Psychotria brachyloba</i>	662
<i>Pseudolmedia brosimifolia</i>	593	<i>Psychotria carthagagenensis</i>	664
<i>Pseudolmedia bucidifolia</i>	443	<i>Psychotria cataractarum</i>	662
<i>Pseudolmedia ferruginea</i>	593	<i>Psychotria chionantha</i>	665
<i>Pseudolmedia guaranitica</i>	593	<i>Psychotria chlorantha</i>	664
<i>Pseudolmedia hirsuta</i>	593	<i>Psychotria corumbensis</i>	664
<i>Pseudolmedia hirtellifolia</i>	593	<i>Psychotria cyanea</i>	657
<i>Pseudolmedia laevigata</i>	593	<i>Psychotria elliptica</i>	664
<i>Pseudolmedia laevis</i>	593	<i>Psychotria ernestii</i>	665
<i>Pseudolmedia mildbraedii</i>	593	<i>Psychotria familiarifolia</i>	664
<i>Pseudolmedia multinervis</i>	593	<i>Psychotria ficigemma</i>	664
<i>Pseudolmedia obliqua</i>	593	<i>Psychotria floribunda</i>	664
<i>Pseudolmedia sagotii</i>	593	<i>Psychotria foetidiflora</i>	656
<i>Pseudopiptadenia</i>	503, 535	<i>Psychotria guianensis</i>	664
<i>Pseudopiptadenia psilostachya</i>	535	<i>Psychotria kukenanensis</i>	656
<i>Pseudopiptadenia suaveolens</i>	536	<i>Psychotria lawrancei</i>	656
<i>Pseudosorocea poeppigii</i>	585	<i>Psychotria lucida</i>	664
<i>Pseudosorocea uaupensis</i>	594	<i>Psychotria lutea</i>	653, 662
<i>Pseudoxandra cuspidata</i>	377	<i>Psychotria magnoliifolia</i>	664
<i>Pseudoxandra lucida</i>	377	<i>Psychotria mapouria</i>	664
<i>Pseudoxandra</i> sp. A	377	<i>Psychotria mapouriodes</i>	665
<i>Pseudoxythece ambelaniifolia</i>	693	<i>Psychotria mapouriodes</i> var. <i>chionantha</i>	665
<i>Psidium acidum</i>	625	<i>Psychotria matthewsii</i>	665
<i>Psidium acutangulum</i>	625	<i>Psychotria nitida</i>	664
<i>Psidium acutangulum</i> var. <i>acidum</i>	625	<i>Psychotria palicourea</i>	662
<i>Psidium acutangulum</i> var. <i>crassirame</i>	625	<i>Psychotria paracatuensis</i>	664
<i>Psidium acutangulum</i> var. <i>oblongatum</i>	625	<i>Psychotria parviflora</i>	666
<i>Psidium acutangulum</i> var. <i>tenuirame</i>	625	<i>Psychotria patrisii</i>	664
<i>Psidium aromaticum</i>	597, 598	<i>Psychotria pedunculosa</i>	665
<i>Psidium aromaticum</i> var. <i>grandiflorum</i>	598	<i>Psychotria plowmanii</i>	656
<i>Psidium ciliatum</i>	626	<i>Psychotria puberulenta</i>	656
<i>Psidium claraense</i>	626	<i>Psychotria ronaldii</i>	656
<i>Psidium eugenoides</i>	626	<i>Psychotria salicifolia</i>	664
<i>Psidium galapageium</i>	626	<i>Psychotria sambucina</i>	664
<i>Psidium galapageium</i> var. <i>howellii</i>	626	<i>Psychotria simira</i>	666
<i>Psidium grandiflorum</i>	597, 625	<i>Psychotria tinctoria</i>	666
<i>Psidium microphyllum</i>	626	<i>Psychotria turboensis</i>	656
<i>Psidium minutiflorum</i>	626	<i>Psychotria viburnoides</i>	664
<i>Psidium molinae</i>	626	<i>Psychotria willdenowii</i>	664
<i>Psidium oligospermum</i>	626	<i>Pterocarpus</i>	503, 536
<i>Psidium paucinerve</i>	626	<i>Pterocarpus amazonicus</i>	537

Appendix 3. — Continuation.

<i>Pterocarpus apalatoa</i>	536	<i>Quararibea amazonica</i>	557
<i>Pterocarpus belizensis</i>	536	<i>Quararibea apaporiensis</i>	556
<i>Pterocarpus crispatus</i>	536	<i>Quararibea duckei</i>	557
<i>Pterocarpus draco</i>	536	<i>Quararibea guianensis</i>	557
<i>Pterocarpus esculentus</i>	537	<i>Quararibea lasiocalyx</i>	555
<i>Pterocarpus floribundus</i>	536	<i>Quararibea machin</i>	557
<i>Pterocarpus grandis</i>	537	<i>Quararibea muricata</i>	555
<i>Pterocarpus hayesii</i>	536	<i>Quararibea ochrocalyx</i>	555
<i>Pterocarpus hemipterus</i>	536	<i>Quararibea spatulata</i>	557
<i>Pterocarpus magnicarpus</i>	536	<i>Quassia alatifolia</i>	706
<i>Pterocarpus michelii</i>	537	<i>Quassia amara</i>	705
<i>Pterocarpus moutouchi</i>	536	<i>Quassia amara</i> f. <i>paniculata</i>	705
<i>Pterocarpus officinalis</i>	536	<i>Quassia amara</i> var. <i>paniculata</i>	705
<i>Pterocarpus ormosioides</i>	533	<i>Quassia aruba</i>	705
<i>Pterocarpus reticulatus</i>	536	<i>Quassia cedron</i>	705
<i>Pterocarpus rohrii</i>	536	<i>Quassia crocea</i>	705
<i>Pterocarpus rohrii</i> var. <i>rubiginosus</i>	536	<i>Quassia crustacea</i>	706
<i>Pterocarpus rufescens</i>	536	<i>Quassia cuspidata</i>	705
<i>Pterocarpus rupestris</i>	536	<i>Quassia dioica</i>	706
<i>Pterocarpus santalinoides</i>	537	<i>Quassia glauca</i>	706
<i>Pterocarpus sapindoides</i>	506	<i>Quassia guianensis</i>	705
<i>Pterocarpus steinbachianus</i>	536	<i>Quassia multiflora</i>	706
<i>Pterocarpus suberosus</i>	536	<i>Quassia officinalis</i>	706
<i>Pterocarpus villosus</i>	536	<i>Quassia orinocensis</i>	706
<i>Pterocarpus violaceus</i>	536	<i>Quassia simaruba</i>	706
<i>Pterocarpus violaceus</i> var. <i>angustifolia</i>	536	<i>Quiina acutangula</i>	632
<i>Ptychopetalum olacoides</i>	634	<i>Quiina albiflora</i>	631
PUTRANJIVACEAE	648	<i>Quiina berryi</i>	630
<i>Pyrostoma ternatum</i>	473	<i>Quiina cajambrensis</i>	630
		<i>Quiina colonensis</i>	630
		<i>Quiina congesta</i>	630
		<i>Quiina crenata</i>	628
		<i>Quiina cruegeriana</i>	630
		<i>Quiina decaisneana</i>	631
		<i>Quiina guaporensis</i>	630
		<i>Quiina guianensis</i>	630
		<i>Quiina integrifolia</i>	631
		<i>Quiina leptoclada</i>	631
		<i>Quiina oblanceolata</i>	631
		<i>Quiina obovata</i>	631
		<i>Quiina oiapicensis</i>	631
		<i>Quiina panamensis</i>	629
		<i>Quiina peruviana</i>	630
		<i>Quiina pteridophylla</i>	632
		<i>Quiina rigidifolia</i>	631
		<i>Quiina sessilis</i>	632
		<i>Quiina silvatica</i>	629
		<i>Quiina tessmannii</i>	630
		<i>Quiina yatuensis</i>	632
		<i>Quiina</i> sp. A	632

Q

<i>Qualea acuminata</i>	722
<i>Qualea albiflora</i>	724
<i>Qualea amapaensis</i>	722
<i>Qualea apodocarpa</i>	724
<i>Qualea caerulea</i>	722
<i>Qualea dinizii</i>	723
<i>Qualea ferruginea</i>	724
<i>Qualea glaberrima</i>	724
<i>Qualea lutea</i>	722
<i>Qualea melinonii</i>	724
<i>Qualea moriboomiorum</i>	723
<i>Qualea polychroma</i>	723
<i>Qualea rosea</i>	724
<i>Qualea rubiginosa</i>	724
<i>Qualea rubiginosa</i> var. <i>angustior</i>	724
<i>Qualea speciosa</i>	722
<i>Qualea tricolor</i>	724
<i>Qualea violacea</i>	724
<i>Quapoya colorans</i>	434
<i>Quapoya microphylla</i>	434
<i>Quapoya panapanari</i>	434

Appendix 3. — Continuation.

R			
<i>Racaria sylvatica</i>	685	<i>Rheedia macrantha</i>	434
<i>Racosperma mangium</i>	504	<i>Rheedia macrophylla</i>	434
<i>Racoubea guianensis</i>	677	<i>Rheedia macrophylla</i> var. <i>benthamiana</i>	434
<i>Raddia elliptica</i>	415	<i>Rheedia madruno</i>	434
<i>Raddia fasciculata</i>	415	<i>Rheedia rostrata</i>	435
<i>Raddia firmifolia</i>	415	<i>Rheedia sagotiana</i>	434
<i>Raddia glomerata</i>	415	<i>Rhigospira paucifolia</i>	385
<i>Raddia impressifolia</i>	416	<i>Rhigospira reticulata</i>	385
<i>Raddia lacunosa</i>	415	<i>Rhigospira sinuosa</i>	385
<i>Raddia oblongifolia</i>	415	<i>Rhizaeris alba</i>	440
<i>Raddia pachyphylla</i>	415	<i>Rhizobolus butyrosus</i>	413
<i>Radlkoferella brachyandra</i>	694	<i>Rhizobolus glaber</i>	413
<i>Radlkoferella grandis</i>	698	<i>Rhizobolus saouvari</i>	413
<i>Radlkoferella littoralis</i>	702	<i>Rhizobolus souari</i>	413
<i>Ragala sanguinolenta</i>	687	<i>Rhizophora americana</i>	649
<i>Raimondia tenuiflora</i>	370	RHIZOPHORACEAE	648
<i>Randia armata</i>	665	<i>Rhizophora mangle</i>	649
<i>Randia guianensis</i>	665	<i>Rhizophora mangle</i> var. <i>racemosa</i>	649
<i>Randia ovata</i>	665	<i>Rhizophora mangle</i> var. <i>samoensis</i>	649
<i>Randia spinosa</i>	665	<i>Rhizophora racemosa</i>	649
<i>Rapanea</i>	645	<i>Rhizophora samoensis</i>	649
<i>Rapanea guianensis</i>	645	<i>Rhodognaphalopsis flaviflora</i>	556
<i>Rapanea guianensis</i> var. <i>andicola</i>	645	<i>Rhodostemonodaphne elephantopus</i>	490
<i>Rapanea guianensis</i> var. <i>bogotensis</i>	645	<i>Rhodostemonodaphne grandis</i>	490
<i>Rapanea guianensis</i> var. <i>calensis</i>	645	<i>Rhodostemonodaphne kunthiana</i>	491
<i>Rapanea oblonga</i>	645	<i>Rhodostemonodaphne leptoclada</i>	491
<i>Raputia paraensis</i>	667	<i>Rhodostemonodaphne morii</i>	491
<i>Raputia</i> sp. A	668	<i>Rhodostemonodaphne revolutifolia</i>	491
<i>Rauvolfia duckei</i>	386	<i>Rhodostemonodaphne rufovirgata</i>	491
<i>Rauvolfia paraensis</i>	386	<i>Rhodostemonodaphne saulensis</i>	491
<i>Rauvolfia pentaphylla</i>	386	<i>Riana guianensis</i>	721
<i>Recordoxylon</i>	503, 537	<i>Richardella cayennensis</i>	695
<i>Recordoxylon amazonicum</i>	537	<i>Richardella cladantha</i>	695
<i>Recordoxylon speciosum</i>	537	<i>Richardella glomerata</i>	698
<i>Remijia lambertiana</i>	661	<i>Richardella hypoglauca</i>	698
RHABDODENDRACEAE	648	<i>Richardella macrocarpa</i>	699
<i>Rhabdodendron amazonicum</i>	648	<i>Richardella macrophylla</i>	699
<i>Rhabdodendron arirambae</i>	648	<i>Richardella rivicoa</i>	699
<i>Rhabdodendron crassipes</i>	648	<i>Richardella speciosa</i>	701
<i>Rhabdodendron duckei</i>	648	<i>Richardella temare</i>	694
<i>Rhabdodendron longifolium</i>	648	<i>Richeria australis</i>	638
<i>Rhabdodendron paniculatum</i>	648	<i>Richeria grandis</i>	638
<i>Rhabdodendron sylvestre</i>	648	<i>Richeria grandis</i> var. <i>divaricata</i>	638
RHAMNACEAE	648	<i>Richeria grandis</i> var. <i>genuina</i>	638
<i>Rhamnus micrantha</i>	409	<i>Richeria grandis</i> var. <i>laurifolia</i>	638
<i>Rhamnus micranthus</i>	409	<i>Richeria grandis</i> var. <i>obovata</i>	638
<i>Rheedia acuminata</i>	434, 435	<i>Richeria grandis</i> var. <i>racemosa</i>	638
<i>Rheedia acuminata</i> var. <i>floribunda</i>	435	<i>Richeria laurifolia</i>	638
<i>Rheedia benthamiana</i>	434	<i>Richeria obovata</i>	638
<i>Rheedia brasiliensis</i>	434	<i>Richeria racemosa</i>	638
<i>Rheedia floribunda</i>	434, 435	<i>Richeria submembranacea</i>	639
<i>Rheedia kappleri</i>	435	<i>Rinorea amapensis</i>	720
		<i>Rinorea bahiensis</i>	720
		<i>Rinorea brevipes</i>	720

Appendix 3. — Continuation.

<i>Rinorea falcata</i>	720	<i>Rollinia jimenezii</i>	369
<i>Rinorea flavescens</i>	720	<i>Rollinia jimenezii</i> var. <i>nelsonii</i>	369
<i>Rinorea guianensis</i>	720	<i>Rollinia micrantha</i>	369
<i>Rinorea laxiflora</i>	720	<i>Rollinia mucosa</i>	369
<i>Rinorea macrocarpa</i>	720	<i>Rollinia mucosa</i> subsp. <i>aequatorialis</i>	369
<i>Rinorea martinii</i>	721	<i>Rollinia mucosa</i> subsp. <i>portoricensis</i>	369
<i>Rinorea neglecta</i>	720	<i>Rollinia mucosa</i> var. <i>macropoda</i>	369
<i>Rinorea paniculata</i>	721	<i>Rollinia mucosa</i> var. <i>neglecta</i>	369
<i>Rinorea passoura</i>	721	<i>Rollinia multiflora</i>	367
<i>Rinorea passoura</i> f. <i>andersonii</i>	721	<i>Rollinia neglecta</i>	369
<i>Rinorea passoura</i> f. <i>grandifolia</i>	721	<i>Rollinia orthopetala</i>	369
<i>Rinorea passoura</i> f. <i>leiosperma</i>	721	<i>Rollinia pachyptera</i>	367
<i>Rinorea passoura</i> var. <i>andersonii</i>	721	<i>Rollinia permensis</i>	369
<i>Rinorea passoura</i> var. <i>grandifolia</i>	721	<i>Rollinia procera</i>	367
<i>Rinorea pectinosquamata</i>	721	<i>Rollinia pterocarpa</i>	369
<i>Rinorea pubiflora</i> f. <i>andersonii</i>	721	<i>Rollinia puberula</i>	367
<i>Rinorea pubiflora</i> f. <i>grandifolia</i>	721	<i>Rollinia pulchrinervia</i>	369
<i>Rinorea pubiflora</i> var. <i>grandifolia</i>	721	<i>Rollinia resinosa</i>	367
<i>Rinorea pubiflora</i> var. <i>pubiflora</i>	721	<i>Rollinia sieberi</i>	369
<i>Rinorea pulleana</i>	721	<i>Rollinia sphaerantha</i>	367
<i>Rinorea riana</i>	721	<i>Rollinia surinamensis</i>	367
<i>Rinorea scandens</i>	721	<i>Rollinia tinifolia</i>	367
<i>Rinorea surinamensis</i>	720	<i>Rollinia uniflora</i>	367
<i>Rinoreocarpus</i>	721	<i>Rondeletia glomerulata</i>	650
<i>Rinoreocarpus salmoneus</i>	721	ROSACEAE	649
<i>Rinoreocarpus ulei</i>	721	<i>Roucheria angulata</i>	546
<i>Rittera dodecandra</i>	539	<i>Roucheria humiriifolia</i>	546
<i>Rittera triphylla</i>	539	<i>Roucheria laxiflora</i>	546
<i>Robinia coccinea</i>	531	<i>Roupala acuminata</i>	647
<i>Robinia nicou</i>	528	<i>Roupala affinis</i>	647
<i>Robinia panacoco</i>	541	<i>Roupala boissieriana</i>	647
<i>Robinia rubiginosa</i>	681	<i>Roupala borealis</i>	647
<i>Robinia sepium</i>	528	<i>Roupala brasiliensis</i> var. <i>macropoda</i>	647
<i>Robinia tomentosa</i>	541	<i>Roupala complicata</i>	647
<i>Robinsonia guianensis</i>	633	<i>Roupala darienensis</i>	647
<i>Robinsonia melianthifolia</i>	633	<i>Roupala dentata</i>	647
<i>Rohria petioliflora</i>	448	<i>Roupala discolor</i>	647
<i>Rohria schreberi</i>	448	<i>Roupala dissimilis</i>	647
<i>Rohria tapura</i>	448	<i>Roupala diversifolia</i>	647
<i>Rollinia biflora</i>	369	<i>Roupala gardneri</i>	647
<i>Rollinia brevipes</i>	367	<i>Roupala gardneri</i> var. <i>angustata</i>	647
<i>Rollinia broadwayi</i>	367	<i>Roupala gardneri</i> var. <i>dentata</i>	647
<i>Rollinia broadwayi</i> var. <i>cuneata</i>	367	<i>Roupala gardneri</i> var. <i>integrifolia</i>	647
<i>Rollinia cardiantha</i>	367	<i>Roupala glabrata</i>	647
<i>Rollinia curvipetala</i>	369	<i>Roupala hameliifolia</i>	647
<i>Rollinia cuspidata</i>	367	<i>Roupala laurifolia</i>	648
<i>Rollinia deliciosa</i>	369	<i>Roupala macropoda</i>	647
<i>Rollinia elliptica</i>	369	<i>Roupala martii</i>	647
<i>Rollinia exsucca</i>	367	<i>Roupala martii</i> var. <i>pinnata</i>	647
<i>Rollinia exsucca</i> subsp. <i>elongata</i>	367	<i>Roupala martii</i> var. <i>simplicifolia</i>	647
<i>Rollinia exsucca</i> subsp. <i>resinosa</i>	367	<i>Roupala mayana</i>	647
<i>Rollinia gardneri</i>	367	<i>Roupala media</i>	647
<i>Rollinia glaucescens</i>	367	<i>Roupala montana</i>	647
<i>Rollinia incurva</i>	367	<i>Roupala montana</i> var. <i>complicata</i>	647

Appendix 3. — Continuation.

<i>Roupala montana</i> var. <i>dentata</i>	647	<i>Ryania speciosa</i>	678, 679
<i>Roupala montana</i> var. <i>heterophylla</i>	647	<i>Ryania speciosa</i> var. <i>bicolor</i>	678
<i>Roupala nitida</i>	648	<i>Ryania speciosa</i> var. <i>subuliflora</i>	678
<i>Roupala ovalis</i>	647	<i>Ryania speciosa</i> var. <i>tomentosa</i>	679
<i>Roupala panamensis</i>	647	<i>Ryania tomentosa</i>	679
<i>Roupala pinnata</i>	646, 647	S	
<i>Roupala pyrifolia</i>	647	SABIACEAE	671
<i>Roupala repanda</i>	647	<i>Sabicea edulis</i>	650
<i>Roupala sessilifolia</i>	647	<i>Sacoglottis amazonica</i>	468
<i>Roupala sphenophylla</i>	647	<i>Sacoglottis cydonioides</i>	468
<i>Roupala tomentosa</i> var. <i>dentata</i>	648	<i>Sacoglottis dichotoma</i>	468
<i>Roupala tomentosa</i> var. <i>sellowii</i>	647	<i>Sacoglottis excelsa</i>	468
<i>Roupala veraguensis</i>	647	<i>Sacoglottis guianensis</i>	468
<i>Roupala yauaperyensis</i>	646	<i>Sacoglottis guianensis</i> f. <i>dolichocarpa</i>	468
<i>Rourea bakeriana</i>	581	<i>Sacoglottis guianensis</i> var. <i>hispidula</i>	468
<i>Ruagea microsepala</i>	578	<i>Sacoglottis guianensis</i> var. <i>maiор</i>	468
RUBIACEAE	650	<i>Sacoglottis subcrenata</i>	468
<i>Rudgea carolina</i>	666	<i>Sagotanthus kappleri</i>	633
<i>Rudgea crassiloba</i>	665	<i>Sagotia brachysepala</i>	464
<i>Rudgea dasyantha</i>	666	<i>Sagotia racemosa</i>	464
<i>Rudgea duckei</i>	665	<i>Sagotia racemosa</i> var. <i>brachysepala</i>	464
<i>Rudgea fissistipula</i>	666	<i>Sagotia racemosa</i> var. <i>genuina</i>	464
<i>Rudgea graciliflora</i>	665	<i>Sagotia racemosa</i> var. <i>ligularis</i>	464
<i>Rudgea lanceifolia</i>	666	<i>Sagotia racemosa</i> var. <i>microsepala</i>	464
<i>Rudgea oldemanii</i>	666	<i>Sagotia tafelbergii</i>	464
<i>Rudgea ovalifolia</i>	666	<i>Saguerus americanus</i>	393
<i>Rudgea prancei</i>	666	<i>Sagus americana</i>	393
<i>Rudgea scandens</i>	657	<i>Sahagunia racemifera</i>	585
<i>Rudgea schomburgkiana</i>	665	<i>Sahagunia strepitans</i>	585
<i>Rudgea sipapoensis</i>	666	<i>Salacia affinis</i>	416
<i>Rudgea standleyana</i>	665	<i>Salacia amygdalina</i>	416
<i>Ruellia macrocalyx</i>	474	<i>Salacia cognata</i>	413
<i>Ruizterania albiflora</i>	724	<i>Salacia cognata</i> var. <i>egensis</i>	413
<i>Ruizterania apodocarpa</i>	724	<i>Salacia cognata</i> var. <i>genuina</i>	413
<i>Ruizterania ferruginea</i>	724	<i>Salacia duckei</i>	416
<i>Ruizterania rubiginosa</i>	724	<i>Salacia elliptica</i>	415
<i>Ruizterania rubiginosa</i> var. <i>angustior</i>	724	<i>Salacia elliptica</i> var. <i>apiculata</i>	415
<i>Ruprechtia amentacea</i>	643	<i>Salacia elliptica</i> var. <i>oblongifolia</i>	415
<i>Ruprechtia brachysepala</i>	643	<i>Salacia erythroxyloides</i>	415
<i>Ruprechtia latifolia</i>	643	<i>Salacia glomerata</i>	415
<i>Ruprechtia marowynensis</i>	643	<i>Salacia grandiflora</i>	416
<i>Ruprechtia martii</i>	643	<i>Salacia guianensis</i>	415
<i>Ruprechtia nitida</i>	643	<i>Salacia impressifolia</i>	416
<i>Rupiliocarpon</i> sp. A	546	<i>Salacia induta</i>	416
RUTACEAE	667	<i>Salacia juruana</i>	416
<i>Ryania bicolor</i>	678	<i>Salacia lacunosa</i>	415
<i>Ryania candollei</i>	678	<i>Salacia lineolata</i>	414
<i>Ryania casiquiarensis</i>	679	<i>Salacia oblongifolia</i>	415
<i>Ryania parviflora</i>	679	<i>Salacia obtusifolia</i>	415
<i>Ryania patrisii</i>	679	<i>Salacia obtusifolia</i> var. <i>parviflora</i>	415
<i>Ryania pyrifera</i>	678, 679	<i>Salacia pachyphylla</i>	415
<i>Ryania pyrifera</i> var. <i>subuliflora</i>	678		
<i>Ryania pyrifera</i> var. <i>tomentosa</i>	679		
<i>Ryania sagotiana</i>	678		

Appendix 3. — Continuation.

<i>Salacia polyanthomaniaca</i>	416	<i>Sapium biglandulosum</i> var. <i>meyerianum</i>	465
<i>Salacia sphaerocarpa</i>	414	<i>Sapium biglandulosum</i> var. <i>moritzianum</i>	465
SALICACEAE	671	<i>Sapium biglandulosum</i> var. <i>oligoneurum</i>	466
<i>Salmasia guianensis</i>	421	<i>Sapium biglandulosum</i> var. <i>salicifolium</i>	465
<i>Salmasia racemosa</i>	421	<i>Sapium biglandulosum</i> var. <i>serratum</i>	465
<i>Salvadora surinamensis</i>	645	<i>Sapium biglandulosum</i> var. <i>sulciferum</i>	466
<i>Samama citrifolia</i>	661	<i>Sapium bogotense</i>	466
<i>Samanea corymbosa</i>	515	<i>Sapium caribaeum</i>	466
<i>Samanea pedicellaris</i>	505	<i>Sapium caudatum</i>	466
<i>Samara floribunda</i>	646	<i>Sapium ciliatum</i>	464
<i>Samyda arborea</i>	672	<i>Sapium clausenianum</i>	466
<i>Samyda guidonia</i>	575	<i>Sapium contortum</i>	467
<i>Samyda iroucana</i>	673	<i>Sapium cremostachyum</i>	465
<i>Samyda lancifolia</i>	673	<i>Sapium endlicherianum</i>	466
<i>Samyda niviana</i>	672	<i>Sapium fendleri</i>	466
<i>Samyda obtusifolia</i>	672	<i>Sapium fragile</i>	466
<i>Samyda octandra</i>	673	<i>Sapium giganteum</i>	466
<i>Samyda parviflora</i>	673, 675, 676	<i>Sapium glandulatum</i>	465
<i>Samyda parvifolia</i>	672	<i>Sapium glandulosum</i>	464
<i>Samyda pitumba</i>	674	<i>Sapium guaricense</i>	466
<i>Samyda procera</i>	672	<i>Sapium haematospermum</i> var. <i>saltense</i>	467
<i>Samyda stipularis</i>	672	<i>Sapium hamatum</i>	465
<i>Samyda sylvestris</i>	675	<i>Sapium hemsleyanum</i>	466
<i>Samyda virgata</i>	672	<i>Sapium hippomane</i>	464
<i>Samyda viridiflora</i>	672	<i>Sapium integrifolium</i>	466
<i>Sandwithia guyanensis</i>	464	<i>Sapium intercedens</i>	466
<i>Sandwithiodoxa egregia</i>	697	<i>Sapium itzananum</i>	467
<i>Saouari glabra</i>	413	<i>Sapium ixiamasense</i>	467
<i>Saouari villosa</i>	413	<i>Sapium izabalense</i>	467
SAPINDACEAE	679	<i>Sapium jamaicense</i>	464
<i>Sapindus arborescens</i>	681, 683	<i>Sapium klotzschianum</i>	465, 466
<i>Sapindus frutescens</i>	681, 683	<i>Sapium klotzschianum</i> var. <i>glaziovii</i>	466
<i>Sapium aereum</i>	465	<i>Sapium lanceolatum</i>	465
<i>Sapium alainianum</i>	467	<i>Sapium lateriflorum</i>	467
<i>Sapium albomarginatum</i>	466	<i>Sapium leptadenium</i>	466
<i>Sapium argutum</i>	464	<i>Sapium marginatum</i>	465, 466
<i>Sapium aubletianum</i>	465	<i>Sapium marginatum</i> f. <i>majus</i>	465
<i>Sapium aucuparium</i>	464-466	<i>Sapium marginatum</i> var. <i>conjugens</i>	466
<i>Sapium aucuparium</i> subsp. <i>moritzianum</i>	465	<i>Sapium marginatum</i> var. <i>grandifolium</i>	466
<i>Sapium aucuparium</i> var. <i>hippomane</i>	465	<i>Sapium marginatum</i> var. <i>lanceolatum</i>	465
<i>Sapium aucuparium</i> var. <i>petiolare</i>	466	<i>Sapium marginatum</i> var. <i>longifolium</i>	466
<i>Sapium aureum</i>	465	<i>Sapium marginatum</i> var. <i>spathulatum</i>	465
<i>Sapium biglandulosum</i>	464-467	<i>Sapium microdentatum</i>	467
<i>Sapium biglandulosum</i> f. <i>minus</i>	465	<i>Sapium moaense</i>	467
<i>Sapium biglandulosum</i> f. <i>oblongatum</i>	465	<i>Sapium montanum</i>	464
<i>Sapium biglandulosum</i> f. <i>ovovata</i>	465	<i>Sapium montevidense</i>	465
<i>Sapium biglandulosum</i> f. <i>pavonianum</i>	465	<i>Sapium moritzianum</i>	465
<i>Sapium biglandulosum</i> var. <i>aubletianum</i>	465	<i>Sapium muelleri</i>	466
<i>Sapium biglandulosum</i> var. <i>aucuparium</i>	464	<i>Sapium naiguatense</i>	466
<i>Sapium biglandulosum</i> var. <i>bogotense</i>	466	<i>Sapium nitidum</i>	467
<i>Sapium biglandulosum</i> var. <i>cuneatum</i>	465	<i>Sapium obtusatum</i>	466
<i>Sapium biglandulosum</i> var. <i>hamatum</i>	465	<i>Sapium obtusilobum</i>	465
<i>Sapium biglandulosum</i> var. <i>klotzschianum</i>	465	<i>Sapium occidentale</i>	466
<i>Sapium biglandulosum</i> var. <i>lanceolatum</i>	465	<i>Sapium oligoneurum</i>	466

Appendix 3. — Continuation.

<i>Sapium paranaense</i>	466	<i>Scyphonymchium multiflorum</i>	683
<i>Sapium paucinervium</i>	467	<i>Secretania loranthacea</i>	634
<i>Sapium paucistamineum</i>	466	<i>Senegalia</i>	503, 537
<i>Sapium pavonianum</i>	465	<i>Senegalia glomerosa</i>	537
<i>Sapium petiolare</i>	466	<i>Senegalia guacamayo</i>	504
<i>Sapium pittieri</i>	466	<i>Senegalia langlassei</i>	537
<i>Sapium poeppigii</i>	466	<i>Senegalia liebmannii</i>	504
<i>Sapium pohlianum</i>	466	<i>Senegalia lobana</i>	537
<i>Sapium prunifolium</i>	465	<i>Senegalia polyphylla</i>	537
<i>Sapium punctatum</i>	466	<i>Senegalia tomentella</i>	537
<i>Sapium pycnostachys</i>	466	<i>Senna</i>	503, 537
<i>Sapium salicifolium</i>	465	<i>Senna marilandica</i>	509
<i>Sapium saltense</i>	467	<i>Senna multijuga</i>	537
<i>Sapium schippii</i>	467	<i>Senna reticulata</i>	538
<i>Sapium serratum</i>	465	<i>Sextonia rubra</i>	491
<i>Sapium suberosum</i>	466	<i>Sickingia tinctoria</i>	666
<i>Sapium submarginatum</i>	466	<i>Siderodendrum ferreum</i>	660
<i>Sapium subserratum</i>	466	<i>Siderodendrum macrophyllum</i>	661
<i>Sapium sulciferum</i>	466	<i>Siderodendrum triflorum</i>	660
<i>Sapium taburu</i>	466	<i>Sideroxyloides ferreum</i>	660
SAPOTACEAE	685, 686, 695, 696	<i>Sideroxylon acutangulum</i>	691
<i>Sapota mulleri</i>	689, 690	<i>Sideroxylon bangii</i>	694
<i>Sarcaulus brasiliensis</i>	704	<i>Sideroxylon calophyloides</i>	692, 693
<i>Sarcaulus macrophyllus</i>	705	<i>Sideroxylon cuspidatum</i> var. <i>crassifolium</i>	695
<i>Sarcocephalus leichhardtii</i>	662	<i>Sideroxylon cyrtobotryum</i>	691
<i>Sarcomphalus cinnamomum</i>	648	<i>Sideroxylon durum</i>	695
<i>Sassafridium macrophyllum</i>	483	<i>Sideroxylon egense</i>	691
<i>Scheelea maripa</i>	392	<i>Sideroxylon eugeniiifolium</i>	697
<i>Scheelea tetrasticha</i>	392	<i>Sideroxylon guianense</i>	693
<i>Schefflera decaphylla</i>	388	<i>Sideroxylon guyanense</i>	691
<i>Schefflera morototoni</i>	389	<i>Sideroxylon hondurensse</i>	701
<i>Schefflera morototoni</i> var. <i>sessiliflorus</i>	389	<i>Sideroxylon meyeri</i>	701
<i>Schefflera paraensis</i>	388	<i>Sideroxylon paraense</i>	691
<i>Schefflera splendens</i>	389	<i>Sideroxylon quinilla</i>	691
<i>Schinus pubescens</i>	669	<i>Sideroxylon robustum</i>	696
<i>Schistostemon dichotomus</i>	468	<i>Sideroxylon rufum</i>	691
<i>Schistostemon sylvaticus</i>	468	<i>Sideroxylon rugosum</i>	691, 692
<i>Schmidelia edulis</i>	679	<i>Sideroxylon rugosum</i> var. <i>egense</i>	691
<i>Schousboea commutata</i>	440	<i>Sideroxylon rugosum</i> var. <i>parviflorum</i>	692
<i>Sciadaphyllum capitatum</i>	390	<i>Sideroxylon ulei</i>	691
<i>Sciadaphyllum decaphyllum</i>	388	<i>Sideroxylon uniloculare</i>	700
<i>Sclerolobium albiflorum</i>	543	<i>Sideroxylon venulosum</i>	693
<i>Sclerolobium amplifolium</i>	542	<i>Sideroxylon williamsii</i>	692
<i>Sclerolobium guianense</i>	542	<i>Silvia anacardioides</i>	481
<i>Sclerolobium guianense</i> var. <i>radlkoferi</i>	542	<i>Silvia itauba</i>	481
<i>Sclerolobium melanocarpum</i>	542	<i>Simaba alata</i>	706
<i>Sclerolobium melinonii</i>	542	<i>Simaba angustifolia</i>	706
<i>Sclerolobium micropetalum</i>	543	<i>Simaba cedron</i>	705
<i>Sclerolobium myrmecophilum</i>	542	<i>Simaba crustacea</i>	706
<i>Sclerolobium paraense</i>	543	<i>Simaba cuspidata</i>	705
<i>Sclerolobium radlkoferi</i>	542	<i>Simaba foetida</i>	706
<i>Sclerolobium subbullatum</i>	542	<i>Simaba guianensis</i>	705, 706
<i>Sclerolobium tinctorium</i> var. <i>uleanum</i>	542	<i>Simaba guianensis</i> subsp. <i>polyphylla</i>	706
<i>Sclerolobium uleanum</i>	542	<i>Simaba morettii</i>	705

Appendix 3. — Continuation.

<i>Simaba multiflora</i>	706	<i>Siphonia guianensis</i>	462
<i>Simaba orinocensis</i>	706	<i>Sloanea</i>	449-455
<i>Simaba polyphylla</i>	706	<i>Sloanea acutiflora</i>	449
<i>Simarouba amara</i>	705, 706	<i>Sloanea alnifolia</i>	452
<i>Simarouba amara</i> var. <i>opaca</i>	706	<i>Sloanea alnifolia</i> var. <i>lancea</i>	452
<i>Simarouba amara</i> var. <i>puberula</i>	706	<i>Sloanea alnifolia</i> var. <i>ovalis</i>	452
<i>Simarouba amara</i> var. <i>typica</i>	706	<i>Sloanea amplifrons</i>	452
<i>SIMAROUBACEAE</i>	705	<i>Sloanea aubletii</i>	454
<i>Simarouba glauca</i>	706	<i>Sloanea brachytepala</i>	450
<i>Simarouba glauca</i> var. <i>latifolia</i>	706	<i>Sloanea bracteosa</i>	453
<i>Simarouba glauca</i> var. <i>typica</i>	706	<i>Sloanea brevipes</i>	450
<i>Simarouba opaca</i>	706	<i>Sloanea breviseta</i>	452
<i>Simira nitida</i>	664	<i>Sloanea calva</i>	450
<i>Simira palicourea</i>	662	<i>Sloanea caribaea</i>	454
<i>Simira tinctoria</i>	666	<i>Sloanea conferta</i>	450
<i>Siparuna amazonica</i>	708	<i>Sloanea corymbiflora</i>	453
<i>Siparuna archeri</i>	707	<i>Sloanea cuneifolia</i>	452
<i>Siparuna argyrochrysea</i>	707	<i>Sloanea echinocarpa</i>	450
<i>Siparuna arianeae</i>	707	<i>Sloanea egensis</i>	453
<i>Siparuna camporum</i>	707	<i>Sloanea eichleri</i>	450
<i>Siparuna cavalcantei</i>	707	<i>Sloanea erythrocarpa</i>	451
<i>SIPARUNACEAE</i>	706	<i>Sloanea fernando-costae</i>	453
<i>Siparuna crassiflora</i>	707	<i>Sloanea ferruginea</i>	455
<i>Siparuna cristata</i>	706	<i>Sloanea filiformis</i>	454
<i>Siparuna cristata</i> var. <i>macrophylla</i>	706	<i>Sloanea floribunda</i>	451
<i>Siparuna cristata</i> var. <i>petiolaris</i>	706	<i>Sloanea garckeana</i>	451
<i>Siparuna cuspidata</i>	707	<i>Sloanea gentryi</i>	453
<i>Siparuna decipiens</i>	707	<i>Sloanea gracilis</i>	453
<i>Siparuna discolor</i>	707	<i>Sloanea grandiflora</i>	452
<i>Siparuna duckeana</i>	707	<i>Sloanea granulosa</i>	452
<i>Siparuna emarginata</i>	707	<i>Sloanea guianensis</i> subsp. <i>guianensis</i>	452
<i>Siparuna espinhicensis</i>	708	<i>Sloanea guianensis</i> subsp. <i>purdieei</i>	452
<i>Siparuna foetida</i>	707	<i>Sloanea guianensis</i> subsp. <i>stipitata</i>	452
<i>Siparuna guianensis</i>	707	<i>Sloanea guianensis</i> var. <i>microcarpa</i>	452
<i>Siparuna guianensis</i> var. <i>divergentifolia</i>	707	<i>Sloanea inermis</i>	453
<i>Siparuna guianensis</i> var. <i>glabrescens</i>	707	<i>Sloanea kappleriana</i>	453
<i>Siparuna guianensis</i> var. <i>longifolia</i>	707	<i>Sloanea latifolia</i>	452
<i>Siparuna guianensis</i> var. <i>nitens</i>	707	<i>Sloanea lindneri</i>	452
<i>Siparuna itacaiunensis</i>	707	<i>Sloanea longicaudata</i>	454
<i>Siparuna lepidantha</i>	707	<i>Sloanea longipes</i>	451
<i>Siparuna lepidiflora</i>	707	<i>Sloanea macrantha</i>	454
<i>Siparuna monogyna</i>	706	<i>Sloanea maroana</i>	451
<i>Siparuna obconica</i>	706	<i>Sloanea maximowicziana</i>	452
<i>Siparuna pachyantha</i>	707	<i>Sloanea megacarpa</i>	454
<i>Siparuna panamensis</i>	707	<i>Sloanea microcarpa</i>	452
<i>Siparuna poeppigii</i>	708	<i>Sloanea morii</i>	453
<i>Siparuna savanicola</i>	707	<i>Sloanea nitida</i>	453
<i>Siparuna sprucei</i>	708	<i>Sloanea obidensis</i>	453
<i>Siparuna ucayaliensis</i>	707	<i>Sloanea obtusa</i>	453
<i>Siparuna williamsii</i>	706	<i>Sloanea obtusifolia</i>	453
<i>Siphoneugena baporeti</i>	625	<i>Sloanea paniculata</i>	451
<i>Siphoneugena cantareirae</i>	625	<i>Sloanea parviflora</i>	453
<i>Siphoneugena legrandii</i>	625	<i>Sloanea parviflora</i> var. <i>pedicellata</i>	453
<i>Siphoneugena micrantha</i>	625	<i>Sloanea pseudodentata</i>	453

Appendix 3. — Continuation.

<i>Sloanea ptariana</i>	452	<i>Spachea elegans</i>	551
<i>Sloanea pubescens</i>	452, 453	<i>Spachea herbert-smithii</i>	551
<i>Sloanea purdiei</i>	452	<i>Spathularia longifolia</i>	718
<i>Sloanea regelii</i>	452	<i>Spermacoce caerulescens</i>	722
<i>Sloanea rojasiae</i>	454	<i>Sphenista peruviana</i>	422
<i>Sloanea rufa</i>	454	<i>Spirotropis</i>	503, 538
<i>Sloanea sinemariensis</i>	450, 454	<i>Spirotropis candellei</i>	538
<i>Sloanea sinemariensis</i> var. <i>melinonii</i>	450	<i>Spirotropis longifolia</i>	538
<i>Sloanea sipapoana</i>	451	<i>Spondias aurantiaca</i>	364
<i>Sloanea stipitata</i>	452	<i>Spondias axillaris</i>	365
<i>Sloanea synandra</i>	454	<i>Spondias cytherea</i>	364
<i>Sloanea trichosticha</i>	454	<i>Spondias dubia</i>	365
<i>Sloanea usurpatrix</i>	449	<i>Spondias graveolens</i>	365
<i>Sloanea wurdackii</i>	454	<i>Spondias lucida</i>	364
<i>Sloanea</i> sp. A	454	<i>Spondias lutea</i>	364, 365
<i>Sloanea</i> sp. B	454	<i>Spondias lutea</i> var. <i>glabra</i>	365
<i>Sloanea</i> sp. C	454	<i>Spondias lutea</i> var. <i>maxima</i>	365
<i>Sloanea</i> sp. D	454	<i>Spondias mombin</i>	364
<i>Sloanea</i> sp. E	454	<i>Spondias myrobalanus</i>	364
<i>Sloanea</i> sp. F	455	<i>Spondias nigrescens</i>	365
<i>Sloanea</i> sp. G	455	<i>Spondias oghigee</i>	365
<i>Sloanea</i> sp. H	455	<i>Spondias pseudomyrobalanus</i>	364
<i>Soaresia nitida</i>	585	<i>Spondias purpurea</i> var. <i>venulosa</i>	365
<i>Socratea albolineata</i>	394	<i>Spondias radlkoferi</i>	365
<i>Socratea durissima</i>	394	<i>Spondias venulosa</i>	365
<i>Socratea elegans</i>	394	<i>Spondias zanzee</i>	365
<i>Socratea exorrhiza</i>	394	<i>Sponia</i>	409
<i>Socratea gracilis</i>	394	<i>Sponia canescens</i>	409
<i>Socratea hoppii</i>	394	<i>Sponia chichilea</i>	409
<i>Socratea macrochlamys</i>	394	<i>Sponia crassifolia</i>	409
<i>Socratea microchlamys</i>	394	<i>Sponia grisea</i>	409
<i>Socratea orbigniana</i>	394	<i>Sponia integerrima</i>	409
<i>Socratea philonotia</i>	394	<i>Sponia lima</i>	409
SOLANACEAE	708	<i>Sponia macrophylla</i>	409
<i>Solanum circinatum</i>	708	<i>Sponia micrantha</i>	409
<i>Solanum crinitum</i>	708	<i>Sponia mollis</i>	409
<i>Solanum cyananthum</i>	708	<i>Sponia peruviana</i>	409
<i>Solanum cyananthum</i> var. <i>jubatum</i>	708	<i>Sponia riparia</i>	409
<i>Solanum endopogon</i> subsp. <i>guianense</i>	708	<i>Sponia rufescens</i>	409
<i>Solanum formosum</i>	708	<i>Sponia schiedeana</i>	409
<i>Solanum jubatum</i>	708	<i>Sprucella cyrtobotrya</i>	691
<i>Solanum splendens</i>	708	<i>Stannia grandiflora</i>	663
<i>Solena armata</i>	665	<i>Stannia metensis</i>	663
<i>Solena decora</i>	663	<i>Stannia panamensis</i>	663
<i>Solena gracilis</i>	663	<i>Steffensia adunca</i>	640
<i>Solena latifolia</i>	663	<i>Steffensia celtidifolia</i>	640
<i>Solena longiflora</i>	663	<i>Steffensia cocoloboides</i>	642
<i>Solena revoluta</i>	663	<i>Steffensia elongata</i>	640
<i>Sophia carolina</i>	556	<i>Steffensia eximia</i>	642
<i>Sorocea guayanensis</i>	594	<i>Steffensia richardifolia</i>	642
<i>Sorocea muriculata</i> subsp. <i>uaupensis</i>	594	<i>Stemmadenia nervosa</i>	387
<i>Sorocea nitida</i>	585	STEMONURACEAE	708
<i>Sorocea stenophylla</i>	585	<i>Stenocalyx albicans</i>	598
<i>Sorocea uaupensis</i>	594	<i>Stenocalyx patrisii</i>	605

Appendix 3. — Continuation.

<i>Stenocalyx patrisii</i> var. <i>grandifolius</i>	605	<i>Stryphnodendron guianense</i> f. <i>floribundum</i>	538
<i>Stenocalyx patrisii</i> var. <i>parvifolius</i>	605	<i>Stryphnodendron melinonii</i>	538
<i>Stenocalyx portoricensis</i>	605	<i>Stryphnodendron moricolor</i>	538
<i>Stenocalyx pseudopsidium</i>	605, 606	<i>Stryphnodendron polystachyum</i>	538
<i>Stenostomum acreanum</i>	666	<i>Stryphnodendron pulcherrimum</i>	538
<i>Stenostomum guianense</i>	666	<i>Stryphnodendron purpureum</i>	538
<i>Stephanium guianense</i>	662	<i>Stryphnodendron</i> sp. A	539
<i>Stephanoluma rugosa</i>	692	<i>Styloceras macrostachyum</i>	459
<i>Sterculia frondosa</i>	557	<i>Stylogyne amazonica</i>	646
<i>Sterculia glabrifolia</i>	557	<i>Stylogyne brasiliensis</i>	646
<i>Sterculia ivira</i>	557	<i>Stylogyne kappleri</i>	646
<i>Sterculia kayae</i>	557	<i>Stylogyne micans</i>	646
<i>Sterculia multiovula</i>	557	<i>Stylogyne orinocensis</i>	646
<i>Sterculia parviflora</i>	557	<i>Stylogyne poeppigii</i>	646
<i>Sterculia pilosa</i>	558	<i>Stylogyne schomburgkiana</i>	646
<i>Sterculia pruriens</i>	557	<i>Stylogyne surinamensis</i>	646
<i>Sterculia pruriens</i> var. <i>grandiflora</i>	557	STYRACACEAE	709
<i>Sterculia pruriens</i> var. <i>parviflora</i>	557	<i>Styrax burchellii</i>	709
<i>Sterculia roseiflora</i>	557	<i>Styrax erymophyllum</i>	709
<i>Sterculia speciosa</i>	558	<i>Styrax glabratus</i>	709
<i>Sterculia surinamensis</i>	558	<i>Styrax guyanensis</i>	709
<i>Sterculia villifera</i>	558	<i>Styrax guyanensis</i> var. <i>japurensis</i>	709
<i>Steudelia racemosa</i>	719	<i>Styrax heteroclitus</i>	709
<i>Stilaginella amazonica</i>	637	<i>Styrax lauraceus</i>	709
<i>Stilaginella benthamii</i>	637	<i>Styrax leiophyllum</i>	709
<i>Stilaginella blanchetiana</i>	637	<i>Styrax longifolius</i>	709
<i>Stilaginella ferruginea</i>	637	<i>Styrax macrophyllum</i>	709
<i>Stilaginella laxiflora</i>	637	<i>Styrax pallidus</i>	709
<i>Stilaginella oblonga</i>	637	<i>Styrax pearcei</i>	709
<i>Stillingia aucuparia</i>	464	<i>Styrax pearcei</i> var. <i>boliviensis</i>	709
<i>Stillingia biglandulosa</i>	465, 467	<i>Styrax psilophyllum</i>	709
<i>Stillingia cremostachya</i>	465	<i>Styrax sieberi</i>	709
<i>Stillingia dracunculoides</i>	466	<i>Styrax squamulosus</i>	709
<i>Stillingia guianensis</i>	463	<i>Styrax tessmannii</i>	709
<i>Stillingia haematantha</i>	466	<i>Surenus brownii</i>	574
<i>Stillingia hippomane</i>	465	<i>Surenus fissilis</i>	574
<i>Stillingia marginata</i>	465	<i>Surenus glaziovii</i>	574
<i>Stillingia prunifolia</i>	465	<i>Surenus guianensis</i>	574
<i>Stillingia salicifolia</i>	465	<i>Surenus mexicana</i>	574
<i>Strigilia glabrata</i>	709	<i>Surenus velloziana</i>	574
<i>Strigilia guyanensis</i>	709	<i>Swartzia</i>	503, 538
<i>Strigilia leiophylla</i>	709	<i>Swartzia acuminata</i>	541
<i>Strigilia macrophylla</i>	709	<i>Swartzia acuminata</i> var. <i>platygynae</i>	541
<i>Strigilia pallida</i>	709	<i>Swartzia acuminata</i> var. <i>puberula</i>	541
<i>Strigilia psilophylla</i>	709	<i>Swartzia acuminata</i> var. <i>tridynamia</i>	541
<i>Strychnodaphne floribunda</i>	485	<i>Swartzia alata</i>	540
<i>Strychnodaphne puberula</i>	488	<i>Swartzia amshoffiana</i>	539
<i>Strychnodaphne puberula</i> var. <i>angustata</i>	488	<i>Swartzia apetala</i> var. <i>acuminata</i>	539
<i>Strychnodaphne puberula</i> var. <i>truncata</i>	488	<i>Swartzia aptera</i>	539
<i>Strychnos cayennensis</i>	547	<i>Swartzia arborescens</i>	539
<i>Stryphnodendron</i>	503, 538	<i>Swartzia bannia</i>	539
<i>Stryphnodendron angustum</i>	538	<i>Swartzia benthamiana</i>	539
<i>Stryphnodendron floribundum</i>	538	<i>Swartzia bifida</i>	539
<i>Stryphnodendron guianense</i>	538	<i>Swartzia brachyrachis</i> var. <i>peruviana</i>	541

Appendix 3. — Continuation.

<i>Swartzia brachystachya</i>	508	<i>Synzyganthera</i>	471, 472
<i>Swartzia canescens</i>	539	<i>Synzyganthera purpurea</i>	471
<i>Swartzia coriacea</i>	543	<i>Systemonodaphne mezii</i>	480
<i>Swartzia dodecandra</i>	539		
<i>Swartzia erythrocarpa</i>	542		
<i>Swartzia grandifolia</i>	540		
<i>Swartzia guianensis</i>	540		
<i>Swartzia hostmannii</i>	541		
<i>Swartzia leblondii</i>	540, 541	<i>Tabebuia aquatilis</i>	398
<i>Swartzia longicarpa</i>	541	<i>Tabebuia araliacea</i>	396
<i>Swartzia longifolia</i>	538	<i>Tabebuia avellanedae</i>	395
<i>Swartzia minutiflora</i>	507	<i>Tabebuia capitata</i>	395
<i>Swartzia ob lanceolata</i>	541	<i>Tabebuia dugandii</i>	395
<i>Swartzia opacifolia</i>	542	<i>Tabebuia dura</i>	398
<i>Swartzia panacoco</i> var. <i>panacoco</i>	541	<i>Tabebuia fluviatilis</i>	398
<i>Swartzia panacoco</i> var. <i>sagotii</i>	541	<i>Tabebuia glomerata</i>	395
<i>Swartzia parviflora</i>	539	<i>Tabebuia hypolepra</i>	395
<i>Swartzia peruviana</i>	541	<i>Tabebuia impetiginosa</i>	395
<i>Swartzia platygyné</i>	541	<i>Tabebuia insignis</i>	398
<i>Swartzia polyphylla</i>	541	<i>Tabebuia ipe</i> var. <i>integra</i>	395
<i>Swartzia prouacensis</i>	507	<i>Tabebuia monticola</i>	396
<i>Swartzia rariflora</i>	539	<i>Tabebuia nicaraguensis</i>	395
<i>Swartzia rosea</i>	539	<i>Tabebuia obscura</i>	396
<i>Swartzia sericea</i>	542	<i>Tabebuia obscura</i> var. <i>schultesiana</i>	396
<i>Swartzia similis</i>	541	<i>Tabebuia palmeri</i>	395
<i>Swartzia tomentosa</i>	541	<i>Tabebuia roraimae</i>	398
<i>Swartzia tomentosa</i> var. <i>sagotii</i>	541	<i>Tabebuia schunkevigoi</i>	395
<i>Swartzia triphylla</i>	539	<i>Tabebuia serratifolia</i>	396
<i>Swartzia urubuensis</i>	542	<i>Tabebuia stenocalyx</i>	398
<i>Swartzia viridiflora</i>	507	<i>Tabebuia subtilis</i>	396
<i>Sweetia nitens</i>	527	<i>Tabebuia subtilis</i> var. <i>schultesiana</i>	396
<i>Sweetia praeclará</i>	515	<i>Tabernaemontana disparifolia</i>	386
<i>Swietenia alternifolia</i>	555	<i>Tabernaemontana albescens</i>	387
<i>Syagrus chavesiana</i>	395	<i>Tabernaemontana attenuata</i>	386
<i>Syagrus inajai</i>	395	<i>Tabernaemontana aubletii</i>	385
<i>Syagrus speciosa</i>	395	<i>Tabernaemontana benthamiana</i>	387
<i>Syagrus stratincola</i>	395	<i>Tabernaemontana fasciculata</i>	386
<i>Sycocarpus rusbyi</i>	576	<i>Tabernaemontana flavicans</i>	386
<i>Symphonia coccinea</i>	435	<i>Tabernaemontana laevis</i>	383
<i>Symphonia esculenta</i>	435	<i>Tabernaemontana lagenaria</i>	387
<i>Symphonia gabonensis</i>	435	<i>Tabernaemontana linkii</i>	387
<i>Symphonia globulifera</i>	435	<i>Tabernaemontana macrophylla</i>	385
<i>Symphonia globulifera</i> var. <i>gabonensis</i>	435	<i>Tabernaemontana meyeri</i>	387
<i>Symphonia globulifera</i> var. <i>macoubea</i>	435	<i>Tabernaemontana multiflora</i>	387
<i>Symphonia grandiflora</i>	435	<i>Tabernaemontana muricata</i>	387
<i>Symphonia microphylla</i>	435	<i>Tabernaemontana myriantha</i>	387
<i>Symphonia</i> sp. A	436	<i>Tabernaemontana obliqua</i>	387
<i>Symphonia utilissima</i>	435	<i>Tabernaemontana oblongifolia</i>	386
SYMPLOCACEAE	709	<i>Tabernaemontana olivacea</i>	386
<i>Symplocos ciponima</i>	709	<i>Tabernaemontana paucifolia</i>	385
<i>Symplocos guianensis</i>	709	<i>Tabernaemontana perrottetii</i>	387
<i>Symplocos guianensis</i> var. <i>paraensis</i>	709	<i>Tabernaemontana poeppigii</i>	387
<i>Symplocos martinicensis</i>	710	<i>Tabernaemontana reticulata</i>	385
<i>Symplocos paraensis</i>	709	<i>Tabernaemontana sananho</i>	387
		<i>Tabernaemontana stenantha</i>	387

Appendix 3. — Continuation.

<i>Tabernaemontana undulata</i>	387	<i>Talisia praealta</i>	684
<i>Taberna poeppigii</i>	387	<i>Talisia pulverulenta</i>	682
<i>Tachibota guianensis</i>	421	<i>Talisia reticulata</i>	685
<i>Tachigali</i>	503, 542	<i>Talisia rosea</i>	683
<i>Tachigali albiflora</i>	543	<i>Talisia sancarlosiana</i>	684
<i>Tachigali amplifolia</i>	542	<i>Talisia simaboides</i>	685
<i>Tachigali angustifolia</i>	543	<i>Talisia squarrosa</i>	685
<i>Tachigali bracteolata</i>	543	<i>Talisia sylvatica</i>	685
<i>Tachigali eriocalyx</i>	543	<i>Tamonea</i>	566
<i>Tachigali glauca</i>	542	<i>Tamonea fothergilla</i>	565
<i>Tachigali grandiflora</i>	543	<i>Tamonea guianensis</i>	566
<i>Tachigali guianensis</i>	542	<i>Tamonea lasiopetala</i>	568
<i>Tachigali melanocarpa</i>	542	<i>Tamonea tomentosa</i> var. <i>auriculata</i>	569
<i>Tachigali melinonii</i>	542	<i>Tanibouca guianensis</i>	440, 441
<i>Tachigali micropetala</i>	543	<i>Taonabo dentata</i>	635
<i>Tachigali myrmecophila</i>	542	<i>Taonabo punctata</i>	635
<i>Tachigali paniculata</i>	543	<i>Tapirira</i>	365
<i>Tachigali paniculata</i> var. <i>angustifolia</i>	543	<i>Tapirira bethanniana</i>	365
<i>Tachigali paniculata</i> var. <i>comosa</i>	543	<i>Tapirira bijuga</i>	365
<i>Tachigali paraensis</i>	543	<i>Tapirira fanshawei</i>	365
<i>Tachigali pulchra</i>	543	<i>Tapirira guianensis</i>	365
<i>Tachigali purpurea</i>	513	<i>Tapirira guianensis</i> subsp. <i>subandina</i>	365
<i>Tachigali richardiana</i>	543	<i>Tapirira guianensis</i> var. <i>cuneata</i>	365
<i>Tachigali rusbyi</i>	543	<i>Tapirira guianensis</i> var. <i>elliptica</i>	365
<i>Tachigali sericea</i>	543	<i>Tapirira marchandii</i>	365
<i>Tachigali</i> sp. A	543	<i>Tapirira myriantha</i>	365
<i>Tachigali subbullata</i>	542	<i>Tapirira obtusa</i>	365
<i>Tachigali sulcata</i>	543	<i>Tapirira pao-pombo</i>	365
<i>Tachigali trigona</i>	543	<i>Tapirira pao-pombo</i> var. <i>major</i>	365
<i>Tachigali uleana</i>	542	<i>Tapirira pearcei</i>	365
<i>Tachigali ulei</i>	543	<i>Tapirira peckoltiana</i>	365
<i>Talipariti pernambucense</i>	558	<i>Tapirocarpus talisia</i>	684
<i>Talipariti tiliaceum</i> var. <i>pernambucense</i>	558	<i>Tapura amazonica</i>	447
<i>Talisia allenii</i>	684	<i>Tapura amazonica</i> var. <i>ciliata</i>	447
<i>Talisia cararensis</i>	684	<i>Tapura amazonica</i> var. <i>cuspidata</i>	447
<i>Talisia carinata</i> f. <i>acutisepala</i>	685	<i>Tapura amazonica</i> var. <i>dasyphylla</i>	447
<i>Talisia clathrata</i> subsp. <i>canescens</i>	683	<i>Tapura amazonica</i> var. <i>sublanceolata</i>	447
<i>Talisia elephantipes</i>	684	<i>Tapura capitulifera</i>	448
<i>Talisia furfuracea</i>	683	<i>Tapura ciliata</i>	447
<i>Talisia glabra</i>	683	<i>Tapura cucullata</i>	448
<i>Talisia glandulifera</i>	683	<i>Tapura guianensis</i>	448
<i>Talisia guianensis</i>	683	<i>Tapura negrensis</i>	448
<i>Talisia hemidasya</i>	683	<i>Tapura singularis</i>	448
<i>Talisia hexaphylla</i>	684	<i>Taralea</i>	503, 543
<i>Talisia longifolia</i>	684	<i>Taralea nudipes</i>	543
<i>Talisia macrophylla</i>	684	<i>Taralea oppositifolia</i>	543
<i>Talisia megaphylla</i>	684	<i>Taralea oppositifolia</i> var. <i>parviflora</i>	543
<i>Talisia micrantha</i>	685	<i>Taralea</i> sp. A	544
<i>Talisia microphylla</i>	684	<i>Tariri guianensis</i>	639
<i>Talisia mollis</i>	684	<i>Tassia ovata</i>	543
<i>Talisia pachycarpa</i>	684	<i>Tecoma adenophylla</i>	395
<i>Talisia panamensis</i>	684	<i>Tecoma albiflora</i>	398
<i>Talisia pedicellaris</i>	682, 683	<i>Tecoma aquatilis</i>	398
<i>Talisia pentantha</i>	684	<i>Tecoma araliacea</i>	396

Appendix 3. — Continuation.

<i>Tecoma atractocarpa</i>	396	<i>Terminalia nyssifolia</i>	441
<i>Tecoma avellanedae</i>	395	<i>Terminalia obidensis</i>	443
<i>Tecoma avellanedae</i> var. <i>alba</i>	395	<i>Terminalia oblonga</i>	443
<i>Tecoma capitata</i>	395	<i>Terminalia obovata</i>	440, 443
<i>Tecoma conspicua</i>	396	<i>Terminalia ochroprumna</i>	443
<i>Tecoma dura</i>	398	<i>Terminalia odontoptera</i>	440
<i>Tecoma flavescens</i>	396	<i>Terminalia ovatifolia</i>	441
<i>Tecoma fluviatilis</i>	398	<i>Terminalia pamea</i>	440
<i>Tecoma impetiginosa</i>	395	<i>Terminalia paraensis</i>	441
<i>Tecoma insignis</i>	398	<i>Terminalia parvifolia</i>	443
<i>Tecoma integra</i>	395	<i>Terminalia rubrigemmis</i>	441
<i>Tecoma ipe</i> f. <i>leucotricha</i>	395	<i>Terminalia scutifera</i>	441
<i>Tecoma ipe</i> var. <i>integra</i>	395	<i>Terminalia subcordata</i>	441
<i>Tecoma ipe</i> var. <i>integrifolia</i>	395	<i>Terminalia tanibouca</i>	441
<i>Tecoma leucoxylon</i> var. <i>miquelii</i>	395	<i>Terminalia tarapotensis</i>	443
<i>Tecoma leucoxylon</i> var. <i>salpingophora</i>	398	<i>Terminalia tetraphylla</i>	443
<i>Tecoma meyeriana</i>	398	<i>Terminalia viridiflora</i>	443
<i>Tecoma nigricans</i>	396	<i>Ternstroemia delicatula</i>	635
<i>Tecoma obscura</i>	395	<i>Ternstroemia dentata</i>	635
<i>Tecoma patrisiana</i>	396	<i>Ternstroemia punctata</i>	635
<i>Tecoma roraimae</i>	398	<i>Ternstroemia revoluta</i>	635
<i>Tecoma speciosa</i>	396	<i>Tetragastris altissima</i>	399
<i>Temenia regia</i>	392	<i>Tetragastris hostmannii</i>	405
<i>Tephrosia barclayana</i>	527	<i>Tetragastris panamensis</i>	404
<i>Tephrosia mimosoides</i>	527	<i>Tetragastris panamensis</i> var. <i>grandifolia</i>	404
<i>Terminalia amazonia</i>	440	<i>Tetragastris panamensis</i> var. <i>hirtella</i>	404
<i>Terminalia aubletii</i>	440	<i>Tetragastris paraensis</i>	404
<i>Terminalia badamia</i>	441	<i>Tetragastris phanerosepala</i>	399
<i>Terminalia capitata</i>	443	<i>Tetragastris pilosa</i>	403
<i>Terminalia carinata</i>	440	<i>Tetragastris stevensonii</i>	404
<i>Terminalia catappa</i>	440, 441	<i>Tetragastris trifoliolata</i>	405
<i>Terminalia catappa</i> var. <i>subcordata</i>	441	<i>Tetrameranthus guianensis</i>	377
<i>Terminalia chiriquensis</i>	443	<i>Tetramerium occidentale</i>	657
<i>Terminalia dichotoma</i>	440, 441	<i>Tetramerium odoratissimum</i>	657
<i>Terminalia erecta</i>	439	<i>Tetramerium sessilifolium</i>	658
<i>Terminalia erecta</i> var. <i>procumbens</i>	439	<i>Tetratome latifolia</i>	582
<i>Terminalia eriantha</i>	441	<i>Tetratome lepidota</i>	708
<i>Terminalia excelsa</i>	440	THEACEAE	710
<i>Terminalia firma</i>	441	<i>Theobroma album</i>	426
<i>Terminalia grandis</i>	441	<i>Theobroma cacao</i>	558, 559
<i>Terminalia guyanensis</i>	441	<i>Theobroma cacao</i> f. <i>leiocarpum</i>	558
<i>Terminalia hayesii</i>	441	<i>Theobroma cacao</i> f. <i>pentagonum</i>	558
<i>Terminalia hilariana</i>	443	<i>Theobroma cacao</i> subsp. <i>leiocarpum</i>	558
<i>Terminalia intermedia</i>	441	<i>Theobroma cacao</i> subsp. <i>pentagonum</i>	558
<i>Terminalia kydiana</i>	441	<i>Theobroma cacao</i> subsp. <i>sativum</i>	558
<i>Terminalia latifolia</i>	441	<i>Theobroma cacao</i> subsp. <i>sphaerocarpum</i>	559
<i>Terminalia latifolia</i> var. <i>dichotoma</i>	441	<i>Theobroma cacao</i> var. <i>leiocarpum</i>	558
<i>Terminalia lucida</i>	441	<i>Theobroma cacao</i> var. <i>typicum</i>	559
<i>Terminalia macrophylla</i>	442	<i>Theobroma caribaeum</i>	558
<i>Terminalia mauritiana</i>	441	<i>Theobroma ferrugineum</i>	559
<i>Terminalia megalophylla</i>	442	<i>Theobroma guazuma</i>	553
<i>Terminalia moluccana</i>	441	<i>Theobroma guianense</i>	559
<i>Terminalia myrobalana</i>	441	<i>Theobroma integrerrimum</i>	558
<i>Terminalia nitidissima</i>	442	<i>Theobroma kalagua</i>	559

Appendix 3. — Continuation.

<i>Theobroma leiocarpum</i>	558	<i>Tocoyena latifolia</i>	663
<i>Theobroma minus</i>	558	<i>Tocoyena longifolia</i>	663
<i>Theobroma pentagonum</i>	558	<i>Tocoyena macrophylla</i>	663
<i>Theobroma saltzmannianum</i>	558	<i>Tocoyena mutisii</i>	663
<i>Theobroma sapidum</i>	559	<i>Tocoyena speciosa</i>	663
<i>Theobroma sativum</i>	558, 559	<i>Tocoyena undulatifolia</i>	663
<i>Theobroma sativum</i> var. <i>leucospermum</i>	559	<i>Tonsella elliptica</i>	415
<i>Theobroma sativum</i> var. <i>melanospermum</i>	559	<i>Tonsella glomerata</i>	415
<i>Theobroma sphaerocarpum</i>	559	<i>Tonsella oblongifolia</i>	415
<i>Theobroma subincanum</i>	559	<i>Tontelea egensis</i>	413
<i>Theobroma sylvestre</i>	559	<i>Tontelea erythroxyloides</i>	415
<i>Theobroma tomentosum</i>	553	<i>Tontelea guianensis</i>	415
<i>Theobroma velutinum</i>	559	<i>Tontelea obtusifolia</i>	415
<i>Theophrasta glycyarpa</i>	719	<i>Toquera tormentosa</i>	447
<i>Theophrasta sparsifolia</i>	719	<i>Torrubia broadwayana</i>	627
<i>Thyrsanthus fasciculatus</i>	386	<i>Torrubia eggarsiana</i>	627
<i>Thrysodium dasytrichum</i>	366	<i>Torrubiaheimeriana</i>	627
<i>Thrysodium giganteum</i>	366	<i>Torrubia salicifolia</i>	627
<i>Thrysodium guianense</i>	366	<i>Touchiroa aromatica</i>	509
<i>Thrysodium paraense</i>	366	<i>Touchiroa oblonga</i>	509
<i>Thrysodium puberulum</i>	366	<i>Touchiroa parivoa</i>	511
<i>Thrysodium salzmannianum</i>	366	<i>Touchiroa tomentosa</i>	511
<i>Thrysodium schomburgkianum</i>	366	<i>Toulicia elliptica</i>	685
<i>Thrysodium schomburgkianum</i> var. <i>salzmannianum</i>	366	<i>Toulicia guianensis</i>	685
<i>Thrysodium spruceanum</i>	366	<i>Toulicia pulvinata</i>	685
<i>Ticorea foetida</i>	668, 722	<i>Tounatea acuminata</i>	541
<i>Ticorea longiflora</i>	668	<i>Tounatea acuminata</i> var. <i>puberula</i>	541
<i>Tingulonga almacega</i>	404	<i>Tounatea aptera</i>	539
<i>Tingulonga altissima</i>	399	<i>Tounatea arborescens</i>	539
<i>Tingulonga aracouchini</i>	400	<i>Tounatea benthamiana</i>	539
<i>Tingulonga caudata</i>	404	<i>Tounatea brachystachya</i>	508
<i>Tingulonga enneandra</i>	401	<i>Tounatea dodecandra</i>	539
<i>Tingulonga gigantea</i>	402	<i>Tounatea grandifolia</i>	540
<i>Tingulonga guianensis</i>	402	<i>Tounatea guianensis</i>	540
<i>Tingulonga heptaphylla</i>	402	<i>Tounatea hostmannii</i>	541
<i>Tingulonga hostmannii</i>	402	<i>Tounatea longifolia</i>	538
<i>Tingulonga insignis</i>	403	<i>Tounatea panacoco</i>	541
<i>Tingulonga martiana</i>	405	<i>Tounatea polyphylla</i>	541
<i>Tingulonga multiflora</i>	402	<i>Tounatea rosea</i>	539
<i>Tingulonga polybotrya</i>	404	<i>Tounatea sericea</i>	542
<i>Tingulonga spruceana</i>	404	<i>Tournefortia foetidissima</i>	722
<i>Tingulonga subserrata</i>	405	<i>Touroulia guianensis</i>	633
<i>Tingulonga tenuifolia</i>	405	<i>Touroulia jenmanii</i>	629
<i>Tingulonga trifoliolata</i>	405	<i>Touroulia pteridophylla</i>	632
<i>Tinus hostmannii</i>	646	<i>Touroulia solitaria</i>	633
<i>Tinus orinocensis</i>	646	<i>Tovomita</i>	436
<i>Tinus schomburgkiana</i>	646	<i>Tovomita angustata</i>	439
<i>Tinus surinamensis</i>	646	<i>Tovomita bahiensis</i>	437
<i>Tipuana erythrocarpa</i>	544	<i>Tovomita brevistaminea</i>	436
<i>Tocoyena gracilis</i>	663	<i>Tovomita calodictyos</i>	436
<i>Tocoyena guianensis</i>	666	<i>Tovomita caloneura</i>	437
<i>Tocoyena guianensis</i> var. <i>communis</i>	666	<i>Tovomita caputmonstia</i>	439
<i>Tocoyena guianensis</i> var. <i>glabriuscula</i>	666	<i>Tovomita cephalostigma</i>	438
<i>Tocoyena insignis</i>	663	<i>Tovomita choisyana</i>	437

Appendix 3. — Continuation.

<i>Tovomita clusiifolia</i>	437	<i>Trattinnickia schwackeana</i>	407
<i>Tovomita coriacea</i>	439	<i>Trattinnickia subchoripetala</i>	406
<i>Tovomita divesora</i>	439	<i>Trema</i>	409
<i>Tovomita excelsa</i>	437	<i>Trema canescens</i>	409
<i>Tovomita fanshawei</i>	437	<i>Trema chichilea</i>	409
<i>Tovomita gazelii</i>	437	<i>Trema floridana</i>	409
<i>Tovomita glossophylla</i>	439	<i>Trema integririma</i>	409
<i>Tovomita grata</i>	437	<i>Trema laxiflora</i>	409
<i>Tovomita guianensis</i>	437	<i>Trema lima</i>	409
<i>Tovomita hameliifolia</i>	438	<i>Trema macrophylla</i>	409
<i>Tovomita jenmanii</i>	436	<i>Trema melinonii</i>	409
<i>Tovomita lanceolata</i>	439	<i>Trema micrantha</i>	409
<i>Tovomita lingulata</i>	438	<i>Trema micrantha</i> var. <i>floridana</i>	409
<i>Tovomita longicuneata</i>	438	<i>Trema micrantha</i> var. <i>obtusata</i>	409
<i>Tovomita longifolia</i>	437	<i>Trema micrantha</i> var. <i>strigillosa</i>	409
<i>Tovomita macrocarpa</i>	438	<i>Trema mollis</i>	409
<i>Tovomita manchamancha</i>	439	<i>Trema riparia</i>	409
<i>Tovomita mangle</i>	438	<i>Trema rufescens</i>	409
<i>Tovomita maxima</i>	438	<i>Trema schiedeana</i>	409
<i>Tovomita melinonii</i>	437	<i>Trema strigillosa</i>	409
<i>Tovomita nigrescens</i>	438	<i>Trichilia acariaeantha</i>	580
<i>Tovomita panamaea</i>	439	<i>Trichilia acutantha</i>	580
<i>Tovomita parvifolia</i>	439	<i>Trichilia alta</i>	581
<i>Tovomita pithecobia</i>	439	<i>Trichilia Aubletii</i>	576
<i>Tovomita rhizophoroides</i>	438	<i>Trichilia biolleyi</i>	580
<i>Tovomita richardiana</i>	437	<i>Trichilia biolleyi</i> var. <i>nicoiensis</i>	580
<i>Tovomita rileyi</i>	439	<i>Trichilia brachystachya</i>	580
<i>Tovomita saulensis</i>	438	<i>Trichilia buchtienii</i>	580
<i>Tovomita schomburgkiana</i>	438	<i>Trichilia canjerana</i>	572
<i>Tovomita schomburgkii</i>	438	<i>Trichilia caucana</i>	579
<i>Tovomita sphenophylla</i>	438	<i>Trichilia caudata</i>	681
<i>Tovomita stigmatosa</i>	438	<i>Trichilia chiriquina</i>	580
<i>Tovomita tenuiflora</i>	438	<i>Trichilia cipo</i>	579
<i>Tovomita triflora</i>	436	<i>Trichilia colombiana</i>	580
<i>Tovomita umbellata</i>	438	<i>Trichilia compacta</i>	581
<i>Tovomita weddelliana</i>	438	<i>Trichilia cuneata</i>	579
<i>Tovomitopsis membranacea</i>	432	<i>Trichilia cuneifolia</i>	581
<i>Tovomita</i> sp. A	439	<i>Trichilia curranii</i>	581
<i>Tragia grandifolia</i>	457	<i>Trichilia davisii</i>	581
<i>Trattinnickia boliviiana</i>	405	<i>Trichilia diversifolia</i>	580
<i>Trattinnickia burserifolia</i>	405, 406	<i>Trichilia echinocarpa</i>	580
<i>Trattinnickia burserifolia</i> var. <i>obtusa</i>	405	<i>Trichilia euneura</i>	579
<i>Trattinnickia burserifolia</i> var. <i>quinquejuga</i>	406	<i>Trichilia eurysepala</i>	581
<i>Trattinnickia cuspidata</i>	399	<i>Trichilia excelsa</i>	580
<i>Trattinnickia demerarae</i>	406	<i>Trichilia flava</i>	580
<i>Trattinnickia glaziovii</i>	407	<i>Trichilia froesii</i>	579
<i>Trattinnickia guianensis</i>	405	<i>Trichilia fuscescens</i>	579
<i>Trattinnickia lawrancei</i> var. <i>boliviiana</i>	405	<i>Trichilia gaudichaudii</i>	579
<i>Trattinnickia multiflora</i>	406	<i>Trichilia gigantophylla</i>	581
<i>Trattinnickia rhoifolia</i>	407	<i>Trichilia goudotiana</i>	580
<i>Trattinnickia rhoifolia</i> var. <i>sprucei</i>	407	<i>Trichilia grandis</i>	581
<i>Trattinnickia rhoifolia</i> var. <i>willdenowii</i>	407	<i>Trichilia guara</i>	575, 578
<i>Trattinnickia ryanii</i>	407	<i>Trichilia harmsii</i>	580
<i>Trattinnickia schomburgkii</i>	405	<i>Trichilia heydeana</i>	580

Appendix 3. — Continuation.

<i>Trichilia insignis</i>	579	<i>Trichocarpus guianensis</i>	452
<i>Trichilia izabalana</i>	580	<i>Trichocarpus laurifolius</i>	452
<i>Trichilia krukovi</i>	581	<i>Trigonia laevis</i>	722
<i>Trichilia laminensis</i>	581	<i>Trigynaea anastomosans</i>	371
<i>Trichilia lanceolata</i>	581	<i>Trigynaea angustifolia</i>	378
<i>Trichilia lepidota</i> subsp. <i>leucastera</i>	579	<i>Trigynaea antillana</i>	378
<i>Trichilia leucastera</i>	579	<i>Trigynaea boliviensis</i>	378
<i>Trichilia lobulata</i>	581	<i>Trigynaea caudata</i>	377
<i>Trichilia longeracemosa</i>	579	<i>Trigynaea grandis</i>	378
<i>Trichilia macrophylla</i>	580	<i>Trigynaea matthewsii</i>	378
<i>Trichilia magnifica</i>	581	<i>Trigynaea perrottetii</i>	378
<i>Trichilia martiana</i>	579	<i>Trigynaea perrottetii</i> var. <i>lanceolata</i>	378
<i>Trichilia micrantha</i>	580	<i>Trigynaea rufescens</i>	378
<i>Trichilia micropetala</i>	580	<i>Trilix glandulosa</i>	671
<i>Trichilia mollis</i>	580	<i>Trilix macrobotrys</i>	671
<i>Trichilia montana</i>	580, 581	<i>Triplaris americana</i>	643
<i>Trichilia montana</i> var. <i>acutivalvis</i>	581	<i>Triplaris boliviiana</i>	643
<i>Trichilia montana</i> var. <i>fendleriana</i>	580	<i>Triplaris brasiliiana</i>	643
<i>Trichilia moritzii</i>	581	<i>Triplaris estriata</i>	643
<i>Trichilia oaxacana</i>	580	<i>Triplaris euryphylla</i>	643
<i>Trichilia obtusanthera</i>	580	<i>Triplaris felipensis</i>	643
<i>Trichilia pallida</i>	580	<i>Triplaris formicosa</i>	643
<i>Trichilia paracaimana</i>	581	<i>Triplaris guanaiensis</i>	643
<i>Trichilia paraensis</i>	581	<i>Triplaris laxa</i>	644
<i>Trichilia pauciflora</i>	581	<i>Triplaris martiana</i> var. <i>oblongifolia</i>	644
<i>Trichilia peruviana</i>	580	<i>Triplaris noli-tangere</i>	643
<i>Trichilia polyantha</i>	576	<i>Triplaris pavonii</i>	643
<i>Trichilia polyneura</i>	581	<i>Triplaris pyramidalis</i>	643
<i>Trichilia portoricensis</i>	580	<i>Triplaris schomburgkiana</i>	643
<i>Trichilia propinqua</i>	581	<i>Triplaris siphonopetala</i>	644
<i>Trichilia ptarihana</i>	682	<i>Triplaris surinamensis</i>	644
<i>Trichilia pubescens</i>	578	<i>Triplaris surinamensis</i> var. <i>benthamiana</i>	644
<i>Trichilia quadrijuga</i>	581	<i>Triplaris surinamensis</i> var. <i>chamisoana</i>	644
<i>Trichilia riedelii</i>	580	<i>Triplaris surinamensis</i> var. <i>crassifolia</i>	644
<i>Trichilia roraimana</i>	580	<i>Triplaris vahliana</i>	644
<i>Trichilia sampaioana</i>	579	<i>Triplaris weigeltiana</i>	644
<i>Trichilia schomburgkii</i>	581	<i>Triplaris williamsii</i>	644
<i>Trichilia septentrionalis</i>	581	<i>Trymatococcus amazonicus</i>	594
<i>Trichilia simplicifolia</i>	580	<i>Trymatococcus guanabarinus</i>	592
<i>Trichilia siqueiraei</i>	581	<i>Trymatococcus oligandrus</i>	594
<i>Trichilia skutchii</i>	581	<i>Trymatococcus paraensis</i>	594
<i>Trichilia</i> sp. A	582	<i>Trymatococcus turbinatus</i>	594
<i>Trichilia stelligera</i>	579	<i>Tschudyia robusta</i>	569
<i>Trichilia subsessilifolia</i>	581		
<i>Trichilia subsimplex</i>	581		
<i>Trichilia surinamensis</i>	582		
<i>Trichilia tocacheana</i>	581		
<i>Trichilia tonduzii</i>	580	<i>Ucriana humboldtii</i>	663
<i>Trichilia trinitensis</i>	580	<i>Ucriana insignis</i>	663
<i>Trichilia triphylla</i>	581	<i>Ucriana macrophylla</i>	663
<i>Trichilia truncata</i>	581	<i>Ucriana mutisii</i>	663
<i>Trichilia weddellii</i>	580	ULMACEAE	711
<i>Trichilia weddellii</i> var. <i>parvifolia</i>	580	<i>Unona aromatica</i>	379
<i>Trichilia weddellii</i> var. <i>stylosa</i>	580	<i>Unona crassipetala</i>	374

U

Appendix 3. — Continuation.

<i>Unona discreta</i>	380	<i>Urbanella oblonga</i>	700
<i>Unona fuscata</i>	374	<i>Urbanella procera</i>	700
<i>Unona lucida</i>	379	<i>Urera acuminata</i>	716
<i>Unona pachypetala</i>	374	<i>Urera alceifolia</i>	716
<i>Unona viridiiflora</i>	371	<i>Urera caracasana</i>	716
<i>Unona xylopioides</i>	379	<i>Urera caracasana</i> var. <i>miquelii</i>	716
<i>Unonopsis andersonii</i>	378	<i>Urera caracasana</i> var. <i>mitis</i>	716
<i>Unonopsis angustifolia</i>	378	<i>Urera caracasana</i> var. <i>subpeltata</i>	716
<i>Unonopsis antillana</i>	378	<i>Urera caracasana</i> var. <i>tomentosa</i>	716
<i>Unonopsis boliviensis</i>	378	<i>Urera corallina</i>	716
<i>Unonopsis buchtienii</i>	378	<i>Urera jacquinii</i>	716
<i>Unonopsis glaucopetala</i>	378	<i>Urera jacquinii</i> var. <i>corallina</i>	716
<i>Unonopsis gracilis</i>	378	<i>Urera jacquinii</i> var. <i>miquelii</i>	716
<i>Unonopsis grandis</i>	378	<i>Urera jacquinii</i> var. <i>subpeltata</i>	716
<i>Unonopsis guaraya</i>	378	<i>Urera jacquinii</i> var. <i>ulmifolia</i>	716
<i>Unonopsis guatterioides</i>	378	<i>Urera mitis</i>	716
<i>Unonopsis guatterioides</i> f. <i>elongata</i>	378	<i>Urera subpeltata</i>	716
<i>Unonopsis lindmanii</i>	378	<i>Urera subpeltata</i> var. <i>morifolia</i>	716
<i>Unonopsis matthewsii</i>	378	<i>Urostigma amazonica</i>	585
<i>Unonopsis oblanceolata</i>	379	<i>Urostigma anacardiifolium</i>	586
<i>Unonopsis obovata</i>	378	<i>Urostigma angustifolium</i>	585
<i>Unonopsis perrottetii</i>	378	<i>Urostigma baccatum</i>	590
<i>Unonopsis perrottetii</i> var. <i>lanceolata</i>	378	<i>Urostigma catappifolium</i>	586
<i>Unonopsis rigida</i>	379	<i>Urostigma cestrifolium</i> f. <i>major</i>	590
<i>Unonopsis rufescens</i>	378	<i>Urostigma clusiifolium</i>	586
<i>Unonopsis stipitata</i>	378	<i>Urostigma complicatum</i>	590
<i>Unonopsis williamsii</i>	378	<i>Urostigma costatum</i>	591
<i>Uragoga anceps</i>	664	<i>Urostigma cyclophyllum</i>	589
<i>Uragoga barbinervia</i>	662	<i>Urostigma doliarium</i>	587
<i>Uragoga brachyloba</i>	662	<i>Urostigma erythrostictum</i>	590
<i>Uragoga carthagensis</i>	664	<i>Urostigma euomphalum</i>	591
<i>Uragoga chionantha</i>	665	<i>Urostigma fagifolium</i>	591
<i>Uragoga chlorantha</i>	664	<i>Urostigma geminum</i>	590
<i>Uragoga corymbifera</i>	664	<i>Urostigma gomelleira</i>	587
<i>Uragoga elliptica</i>	664	<i>Urostigma lancifolium</i>	590
<i>Uragoga ficigemma</i>	664	<i>Urostigma laurifolium</i>	588
<i>Uragoga fissistipula</i>	666	<i>Urostigma leucostictum</i>	589
<i>Uragoga floribunda</i>	664	<i>Urostigma mattheusii</i>	586
<i>Uragoga lanceifolia</i>	666	<i>Urostigma nymphaeifolium</i>	589
<i>Uragoga magnoliifolia</i>	664	<i>Urostigma padifolium</i>	590
<i>Uragoga mapouria</i>	664	<i>Urostigma paraense</i>	589
<i>Uragoga myriantha</i>	664	<i>Urostigma pertusum</i>	590
<i>Uragoga ovalifolia</i>	666	<i>Urostigma populneum</i> f. <i>mexicanum</i>	590
<i>Uragoga palicourea</i>	662	<i>Urostigma protensum</i>	588
<i>Uragoga patrisii</i>	664	<i>Urostigma rolanderi</i>	590
<i>Uragoga pedunculosa</i>	665	<i>Urostigma sapidum</i>	590
<i>Uragoga salicifolia</i>	664	<i>Urostigma schiedeanum</i>	590
<i>Uragoga sambucina</i>	664	<i>Urostigma schumacheri</i>	591
<i>Uragoga simira</i>	666	<i>Urostigma splendens</i>	586
<i>Uragoga tinctoria</i>	666	<i>Urostigma subapiculatum</i>	586
<i>Uragoga viburnoides</i>	664	<i>Urostigma subtriplinervium</i>	590
<i>Urbanella buchananitifolia</i>	700	<i>Urostigma subtriplinervium</i> f. <i>ellipticum</i>	590
<i>Urbanella cuspidata</i>	700	<i>Urostigma subtriplinervium</i> f. <i>major</i>	590
<i>Urbanella excelsa</i>	700	<i>Urostigma sulcipes</i>	590

Appendix 3. — Continuation.

<i>Urostigma trigonum</i>	591	<i>Virola michelii</i>	595
<i>Urostigma turbinatum</i>	590	<i>Virola mocoia</i>	595
<i>Urostigma williamsii</i>	589	<i>Virola multicostata</i>	595
<i>Urtica acuminata</i>	716	<i>Virola myctis</i>	596
<i>Urtica aestuans</i>	722	<i>Virola panamensis</i>	595
<i>Urtica alceifolia</i>	716	<i>Virola peruviana</i> var. <i>tomentosa</i>	—
<i>Urtica alnifolia</i>	409	<i>Virola sebifera</i>	595
<i>Urtica caracasana</i>	716	<i>Virola sebifera</i> var. <i>curvinervia</i>	595
URTICACEAE	711	<i>Virola surinamensis</i>	596
<i>Urtica corallina</i>	716	<i>Virola venezuelensis</i>	596
<i>Urtica mitis</i>	716	<i>Virola warburgii</i>	596
<i>Urtica tiliifolia</i>	716	<i>Vismia</i>	470, 471, 625
<i>Urtica ulmifolia</i>	716	<i>Vismia acuminata</i>	470
<i>Utea guyannensis</i>	529	<i>Vismia amazonica</i>	470
<i>Uterveria frondosa</i>	411	<i>Vismia angusta</i>	470
<i>Uva brasiliensis</i>	371	<i>Vismia buchtienii</i>	470
<i>Uvaria aromatica</i>	379	<i>Vismia caparosa</i>	470
<i>Uvaria brasiliensis</i>	371	<i>Vismia cayennensis</i>	470, 471
<i>Uvaria febrifuga</i>	379	<i>Vismia cayennensis</i> var. <i>sessilifolia</i>	471
<i>Uvaria guatterioides</i>	378	<i>Vismia ferruginea</i>	470
<i>Uvaria monosperma</i>	374	<i>Vismia floribunda</i>	470
<i>Uvaria ouregou</i>	374	<i>Vismia glaziovii</i>	470
<i>Uvaria spectabilis</i>	372	<i>Vismia gracilis</i>	470
<i>Uvaria viridiflora</i>	371	<i>Vismia guianensis</i>	470
<i>Uvifera latifolia</i>	642	<i>Vismia guianensis</i> var. <i>acuminata</i>	470
<i>Uvifera polystachya</i>	643	<i>Vismia guttifera</i> var. <i>guianensis</i>	470
V			
<i>Vantanea cupularis</i>	470	<i>Vismia latifolia</i>	470
<i>Vantanea guianensis</i>	468	<i>Vismia macrophylla</i>	470
<i>Vantanea maculicarpa</i>	469	<i>Vismia macrophylla</i> var. <i>glabrescens</i>	470
<i>Vantanea ovicarpa</i>	469	<i>Vismia ramuliflora</i>	471
<i>Vantanea parviflora</i>	470	<i>Vismia reticulata</i>	470
<i>Varronia calyptata</i>	444	<i>Vismia rufescens</i>	471
<i>Varronia calyptata</i> var. <i>hartweggi</i>	444	<i>Vismia sandwithii</i>	471
<i>Vatairea</i>	503, 544	<i>Vismia sessilifolia</i>	471
<i>Vatairea erythrocarpa</i>	544	<i>Vitellaria dissepala</i>	702
<i>Vatairea guianensis</i>	544	<i>Vitellaria glaucophylla</i>	699
<i>Vatairea paraensis</i>	544	<i>Vitellaria littoralis</i>	702
<i>Vatairea surinamensis</i>	544	<i>Vitellaria procera</i>	700
<i>Vataireopsis</i>	503, 544	<i>Vitellaria pulverulenta</i>	695
<i>Vataireopsis surinamensis</i>	544	<i>Vitellaria rivicoa</i>	699
VERBENACEAE	566, 717	<i>Vitex brittoniana</i>	472
<i>Verbena stoechadifolia</i>	722	<i>Vitex compressa</i>	472
<i>Verticillaria acuminata</i>	434	<i>Vitex compressa</i> f. <i>angustifolia</i>	472
<i>Verticillaria madruno</i>	434	<i>Vitex guianensis</i>	473
<i>Vexillifera micranthera</i>	513	<i>Vitex multiflora</i>	473
VIOLACEAE	718	<i>Vitex orinocensis</i> var. <i>multiflora</i>	473
<i>Virola boliviensis</i>	596	<i>Vitex stahelii</i>	473
<i>Virola glaziovii</i>	596	<i>Vitex triflora</i>	473, 474
<i>Virola kwatae</i>	595	<i>Vitex triflora</i> f. <i>quinquefoliolata</i>	474
<i>Virola melinonii</i>	595	<i>Vitex triflora</i> var. <i>angustiloba</i>	474
		<i>Vitex triflora</i> var. <i>coriacea</i>	474
		<i>Vitex triflora</i> var. <i>floribunda</i>	474
		<i>Vitex triflora</i> var. <i>hirsuta</i>	474
		<i>Vitex triflora</i> var. <i>kraatzii</i>	474

Appendix 3. — Continuation.

<i>Vitex triflora</i> var. <i>quinquefoliolata</i>	474	<i>Wikstroemia camelliifolia</i>	710
<i>Vitex triflora</i> var. <i>tenuifolia</i>	474	<i>Wikstroemia fruticosa</i>	710
<i>Vochysia arcuata</i>	725	<i>Wikstroemia intermedia</i>	710
<i>Vochysia cayennensis</i>	724	<i>Wikstroemia parviflora</i>	710
VOCHysiACEAE	722	<i>Wikstroemia symplocoides</i>	710
<i>Vochysia densiflora</i>	724	<i>Willdenovia schreberi</i>	663
<i>Vochysia excelsa</i>	724	<i>Willughbeia acida</i>	381
<i>Vochysia glaberrima</i>	724	<i>Wittelsbachia orinocensis</i>	398
<i>Vochysia guianensis</i>	724		
<i>Vochysia lucida</i>	724		
<i>Vochysia melinonii</i>	724	X	
<i>Vochysia neyratii</i>	725		
<i>Vochysia paraensis</i>	725	<i>Xantolis tuberculata</i>	702
<i>Vochysia rufescens</i>	725	<i>Xantolis venulosa</i>	693
<i>Vochysia sabatieri</i>	725	<i>Ximenia aculeata</i>	634
<i>Vochysia sofiae</i>	725	<i>Ximenia americana</i>	634, 635
<i>Vochysia speciosa</i>	725	<i>Ximenia americana</i> f. <i>inermis</i>	634, 635
<i>Vochysia surinamensis</i>	725	<i>Ximenia americana</i> var. <i>oblonga</i>	634
<i>Vochysia tetraphylla</i>	725	<i>Ximenia americana</i> var. <i>ovata</i>	635
<i>Vochysia tomentosa</i>	725	<i>Ximenia arborescens</i>	634
<i>Vochysia vismiifolia</i>	725	<i>Ximenia elliptica</i>	634, 635
<i>Votomita guianensis</i>	572	<i>Ximenia fluminensis</i>	635
<i>Vouacapoua</i>	503, 506, 544	<i>Ximenia inermis</i>	634
<i>Vouacapoua amazonum</i>	544	<i>Ximenia laurina</i>	635
<i>Vouacapoua americana</i>	544	<i>Ximenia loranthifolia</i>	635
<i>Vouacapoua aubletti</i>	544	<i>Ximenia montana</i>	634
<i>Vouacapoua inermis</i>	506	<i>Ximenia multiflora</i>	635
<i>Vouacapoua retusa</i>	506	<i>Ximenia oblonga</i>	635
<i>Vouacapoua sapindoides</i>	506	<i>Ximenia spinosa</i>	634
<i>Vouacapoua surinamensis</i>	506	<i>Ximenia verrucosa</i>	634
<i>Vouapa acaciifolia</i>	528	<i>Xiphidium caeruleum</i>	722
<i>Vouapa angustifolia</i>	528	<i>Xyladenius glandulosus</i>	671
<i>Vouapa bifolia</i>	529	<i>Xylocarpus carapa</i>	573
<i>Vouapa chrysostachya</i>	528	<i>Xylon pentandrum</i>	552
<i>Vouapa guianensis</i>	529	<i>Xylopia acuminata</i>	366
<i>Vouapa multijuga</i>	529	<i>Xylopia aromatica</i>	379
<i>Vouapa staminea</i>	529	<i>Xylopia benthamii</i>	379
<i>Vouarana guianensis</i>	685	<i>Xylopia benthamii</i> var. <i>dolichopetala</i>	379
<i>Voyara montana</i>	703	<i>Xylopia benthamii</i> var. <i>subnuda</i>	379
<i>Voyria caerulea</i>	722	<i>Xylopia cayennensis</i>	379
		<i>Xylopia cinerea</i>	380
		<i>Xylopia crinita</i>	379
		<i>Xylopia cubensis</i>	379
		<i>Xylopia discreta</i>	380
		<i>Xylopia dunaliana</i>	379
		<i>Xylopia excellens</i>	380
		<i>Xylopia frutescens</i>	380
		<i>Xylopia grandiflora</i>	379
		<i>Xylopia longifolia</i>	379
		<i>Xylopia lucida</i>	379
		<i>Xylopia meridensis</i>	380
		<i>Xylopia neglecta</i>	380
		<i>Xylopia nervosa</i>	380
		<i>Xylopia nitida</i>	379, 380

W

<i>Wallenia myrianthos</i>	645
<i>Webera quinquenervia</i>	560
<i>Weigeltia capitellata</i>	645
<i>Weigeltia microbotrys</i>	644
<i>Weigeltia myrianthos</i>	645
<i>Weigeltia parviflora</i>	644
<i>Weigeltia potiae</i>	644
<i>Weigeltia schomburgkiana</i>	645
<i>Weigeltia surinamensis</i>	645
<i>Wibelia guianensis</i>	719

Appendix 3. — Continuation.

<i>Xylopia nitida</i> var. <i>longifolia</i>	379	<i>Zanthoxylum luizii</i>	669
<i>Xylopia parviflora</i>	380	<i>Zanthoxylum machadoi</i>	668
<i>Xylopia pulcherrima</i>	380	<i>Zanthoxylum microcarpum</i>	670
<i>Xylopia salicifolia</i>	380	<i>Zanthoxylum minutiflorum</i>	668
<i>Xylopia setosa</i>	380	<i>Zanthoxylum obscurum</i>	670
<i>Xylopia surinamensis</i>	380	<i>Zanthoxylum obscurum</i> var. <i>ruizianum</i>	670
<i>Xylopia ulei</i>	379	<i>Zanthoxylum occidentale</i>	669
<i>Xylopia xylopioides</i>	379	<i>Zanthoxylum ocumarensse</i>	668
<i>Xylocarpum acuminatum</i>	366	<i>Zanthoxylum paulae</i>	669
<i>Xylocarpum aromaticum</i>	379	<i>Zanthoxylum peltophorum</i>	670
<i>Xylocarpum frutescens</i>	380	<i>Zanthoxylum pentandrum</i>	669
<i>Xylocarpum grandiflorum</i>	379	<i>Zanthoxylum perrottetii</i>	669
<i>Xylocarpum longifolium</i>	379	<i>Zanthoxylum pringlei</i>	668
<i>Xylocarpum neglectum</i>	380	<i>Zanthoxylum procerum</i>	668
<i>Xylocarpum nitidum</i>	380	<i>Zanthoxylum pubescens</i>	669
<i>Xylosma armata</i>	679	<i>Zanthoxylum regnellianum</i>	671
<i>Xylosma benthamii</i>	679	<i>Zanthoxylum rhoifolium</i>	669-671
<i>Xylosma ciliatifolia</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>formosanum</i>	671
<i>Xylosma digyna</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>peltophorum</i>	670
<i>Xylosma pallidifolia</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>petiolulatum</i>	671
<i>Xylosma pilosa</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>pubescens</i>	670
<i>Xylosma pou-orfilae</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>sessilifolium</i>	671
<i>Xylosma tessmannii</i>	679	<i>Zanthoxylum rhoifolium</i> var. <i>surparanaense</i>	671
<i>Xylosterculia pilosa</i>	558	<i>Zanthoxylum ruizianum</i>	670
 Y		<i>Zanthoxylum sobrevielae</i>	669
<i>Yutajea liesneri</i>	660	<i>Zanthoxylum sorbifolium</i>	669
 Z		<i>Zanthoxylum trinitense</i>	669
<i>Zamzela racemosa</i>	422	<i>Zeyheria digitata</i>	398
<i>Zanthoxylum achidek</i>	669	<i>Zeyheria fluviatilis</i>	398
<i>Zanthoxylum acreanum</i>	668	<i>Zinowiewia aymardii</i>	416
<i>Zanthoxylum acuminatum</i> subsp. <i>juniperinum</i>	668	<i>Ziziphus cinnamomum</i>	648
<i>Zanthoxylum acutifolium</i>	670	<i>Ziziphus itacaiunensis</i>	648
<i>Zanthoxylum acutifolium</i> var. <i>petiolulatum</i>	670	<i>Zollernia</i>	503, 544
<i>Zanthoxylum amapaense</i>	668	<i>Zollernia surinamensis</i>	544
<i>Zanthoxylum apiculatum</i>	669	<i>Zschokkea aculeata</i>	385
<i>Zanthoxylum astrigerum</i>	671	<i>Zschokkea guyanensis</i>	385
<i>Zanthoxylum belizense</i>	669	<i>Zwingera amara</i>	705
<i>Zanthoxylum caribaeum</i>	669	<i>Zwingera aruba</i>	705
<i>Zanthoxylum dellomei</i>	669	<i>Zwingera orinocensis</i>	706
<i>Zanthoxylum ekmanii</i>	669	<i>Zygia</i>	503, 545
<i>Zanthoxylum elephantiasis</i>	669	<i>Zygia arborescens</i>	545
<i>Zanthoxylum gentlei</i>	669	<i>Zygia cataractae</i>	545
<i>Zanthoxylum hermaphroditum</i>	669	<i>Zygia divaricata</i>	545
<i>Zanthoxylum horridum</i>	669	<i>Zygia foreroi</i>	545
<i>Zanthoxylum idae</i>	668	<i>Zygia glomerata</i>	545
<i>Zanthoxylum juniperinum</i>	668	<i>Zygia huberi</i>	545
<i>Zanthoxylum langsdownii</i>	669	<i>Zygia inaequalis</i>	545

APPENDIX 4. — List of cited exsiccata held at CAY, with the corresponding barcode(s) and identifier in the CAY database/Liste des exsiccata cités conservés à CAY, avec le(s) code(s)-barres et l'identifiant correspondants dans la base de données de CAY.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
A. Le Goff 201	CAY002062	49848	C. Moretti 1027	CAY024960	54949
A.H. Gentry & C. Feuillet 63218	CAY057498	125443	C. Moretti 1027	CAY024961	54949
A.H. Gentry & E.M. Zardini 50232	CAY117685	26300	C. Moretti 1027	CAY024962	54949
A.H. Gentry & E.M. Zardini 50232	CAY117686	26300	C. Moretti 1035	CAY097284	54956
A.H. Gentry & E.M. Zardini 50232	CAY117687	26300	C. Moretti 354	CAY031793	54631
A.H. Gentry et al. 63044	CAY107936	26408	C. Moretti 354	CAY031794	54631
A.H. Gentry et al. 63129	CAY065839	26428	C. Moretti 354	CAY076072	54631
A.J.M. Leeuwenberg 11653	CAY084756	50267	C. Moretti 354	CAY076073	54631
B. Bordenave & Bétian 7946	CAY052938	128460	C. Moretti 354	CAY076074	54631
B. Bordenave 5040	CAY031363	120284	C. Moretti 358	CAY024909	54634
B. Bordenave 5040	CAY037898	120284	C. Moretti 571	CAY090070	54724
B. Bordenave 887	CAY035965	118859	C. Moretti 606	CAY017475	113993
B. Dutrèvre & F. Crozier 19	CAY014118	110320	C. Moretti 606	CAY017476	113993
B. Dutrèvre & F. Crozier 7	CAY014160	109706	C. Moretti 626	CAY084776	54760
B. Dutrèvre & F. Crozier 7	CAY014161	109706	C. Moretti 626	CAY084777	54760
B. Dutrèvre 388	CAY046121	129711	C. Moretti 626	CAY084778	54760
B. Dutrèvre 399	CAY046135	129722	C. Moretti 626	CAY084779	54760
B. Dutrèvre 489	CAY046165	129747	C. Moretti 979	CAY189796	54930
B. Dutrèvre 489	CAY046166	129747	C. Moretti 979	CAY189797	54930
B. Dutrèvre 569	CAY046265	129826	C. Moretti 979	CAY189798	54930
B. Dutrèvre 569	CAY046266	129826	C. Moretti 979	CAY189799	54930
B. Riéra & D. Sabatier 1282	CAY111689	71097	C. Persson et al. 2069	CAY074182	65513
B. Riéra & D. Sabatier 1808	CAY184383	71366	C. Sastre & C. Moretti 3852	CAY031333	78155
B. Riéra & D. Sabatier 1904	CAY080261	71471	C. Sastre & C. Moretti 4020	CAY081340	78284
B. Riéra & D. Sabatier 1925	CAY177421	71498	C. Sastre & D.A. Bell 8125	CAY092717	101602
B. Riéra 1092	CAY082971	70926	C. Sastre 1621	CAY006060	101468
B. Riéra 1257	CAY089645	71079	C. Sastre 265	CAY039442	121682
B. Riéra 470	CAY024975	70367	C. Sastre 4009	CAY029953	78277
B. Riéra 470	CAY024976	70367	C. Sastre 4146	CAY087851	78386
B. Riéra 470	CAY024977	70367	C. Sastre 4516	CAY047646	78586
B. Riéra 470	CAY024978	70367	C. Sastre 4704	CAY111999	78744
B. Riéra 470	CAY024979	70367	C. Sastre 6178	CAY195078	79381
B. Riéra 715	CAY047253	70587	C. Sastre 6387	CAY030370	79528
B. Torke & S. Gonzalez 191	CAY017584	114080	C. Sastre 6387	CAY206007	79528
B.M. Boom & S.A. Mori 1795	CAY007760	5647	C. Sastre 6405	CAY024890	79543
B.M. Boom & S.A. Mori 1805	CAY108143	5657	C. Sastre et al. 4125	CAY002146	78367
B.M. Boom & S.A. Mori 1811	CAY027866	5661	D. Larpin 1021	CAY079504	49627
B.M. Boom & S.A. Mori 1885	CAY081790	5727	D. Larpin 670	CAY178249	49285
B.M. Boom & S.A. Mori 2116		5951	D. Larpin 687	CAY090052	49302
B.M. Boom & S.A. Mori 2222	CAY065795	6056	D. Larpin 687	CAY090053	49302
B.M. Boom & S.A. Mori 2344	CAY003294	6175	D. Larpin 702	CAY093451	49317
B.M. Boom & S.A. Mori 2362	CAY107512	6192	D. Larpin 846	CAY090420	49461
B.M. Boom & S.A. Mori 2415	CAY204020	6243	D. Larpin 882	CAY007947	49497
C. Baraloto et al. 3500	CAY100717	303164	D. Larpin 885	CAY100031	49500
C. Baraloto et al. 8056	CAY111112	309085	D. Larpin 946	CAY178975	49561
C. Delnatte 1445	CAY080555	135143	D. Larpin 946	CAY178976	49561
C. Delnatte 1549	CAY083525	135993	D. Loubray 1122	CAY028739	52181
C. Delnatte 1556	CAY083808	136195	D. Loubray 1122	CAY028740	52181
C. Feuillet 10338		23966	D. Loubray 1122	CAY028741	52181
C. Feuillet 1122	CAY208440	21457	D. Loubray 1135	CAY086128	52194
C. Feuillet 1122	CAY208441	21457	D. Loubray 1579	CAY191201	52634
C. Feuillet 2263	CAY082560	22031	D. Loubray 1579	CAY191202	52634
C. Feuillet 2961	CAY086169	22347	D. Loubray 1626	CAY064032	52681
C. Feuillet 3936	CAY098062	22778	D. Loubray 1626	CAY064033	52681
C. Feuillet 616	CAY026346	20987	D. Loubray 1698	CAY162707	52756
C. Feuillet 616	CAY026347	20987	D. Loubray 1774	CAY189288	52835
C. Feuillet 715	CAY195226	21082	D. Loubray 1892	CAY001594	52955
C. Feuillet et al. 10067		23699	D. Loubray 1892	CAY001595	52955
C. Feuillet et al. 10083		23715	D. Loubray 1892	CAY001596	52955
C. Feuillet et al. 10115	CAY027154	23746	D. Loubray 1892	CAY001597	52955
C. Feuillet et al. 10319	CAY057803	23947	D. Loubray 1934	CAY010151	52988
C. Feuillet et al. 10319	CAY057803	23947	D. Loubray 2000	CAY214652	53054
C. Feuillet et al. 10354		23982	D. Loubray 2097	CAY166396	53151

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
D. Loubry 2339	CAY001299	53379	D. Sabatier & J.-F. Molino 5135	CAY104867	306670
D. Loubry 2392	CAY097482	53432	D. Sabatier & J.-F. Molino 5135	CAY104868	306670
D. Loubry 2413	CAY079895	53453	D. Sabatier & J.-F. Molino 5136	CAY103757	305608
D. Loubry 2413	CAY079896	53453	D. Sabatier & J.-F. Molino 5137	CAY068500	133559
D. Loubry 55	CAY178198	51470	D. Sabatier & J.-F. Molino 5137	CAY068501	133559
D. Loubry 621	CAY109565	51819	D. Sabatier & J.-F. Molino 5187	CAY077841	134265
D. Loubry 621	CAY109566	51819	D. Sabatier & J.-F. Molino 5194	CAY077848	134271
D. Loubry 833	CAY089648	52026	D. Sabatier & J.-F. Molino 5194	CAY077849	134271
D. Paget 201	CAY019505	109889	D. Sabatier & J.-F. Molino 5198	CAY081047	135560
D. Paget 204	CAY019507	109887	D. Sabatier & J.-F. Molino 5202	CAY081048	135561
D. Sabatier & B. Riéra 1741	CAY024999	73036	D. Sabatier & J.-F. Molino 5204	CAY077877	134292
D. Sabatier & B. Riéra 3113		74341	D. Sabatier & J.-F. Molino 5205	CAY077878	134293
D. Sabatier & D.-Y. Alexandre 889	CAY002037	72313	D. Sabatier & J.-F. Molino 5300	CAY109448	307861
D. Sabatier & E. Fonty 5589	CAY111701	307863	D. Sabatier & J.-F. Molino 5300	CAY111699	307861
D. Sabatier & E. Fonty 5589	CAY111702	307863	D. Sabatier & J.-F. Molino 5304	CAY083534	135997
D. Sabatier & E. Fonty 5590	CAY109367	308430	D. Sabatier & J.-F. Molino 5321	CAY083538	136000
D. Sabatier & E. Fonty 5590	CAY109368	308430	D. Sabatier & J.-F. Molino 5322bis	CAY083618	136063
D. Sabatier & E. Fonty 5591	CAY111636	307812	D. Sabatier & J.-F. Molino 5332	CAY084961	136466
D. Sabatier & E. Fonty 5597	CAY108458	307929	D. Sabatier & J.-F. Molino 5334	CAY109142	308351
D. Sabatier & E. Fonty 5662	CAY108064	307251	D. Sabatier & J.-F. Molino 5334	CAY109143	308351
D. Sabatier & E. Fonty 5662	CAY108065	307251	D. Sabatier & J.-F. Molino 5340	CAY083542	136003
D. Sabatier & E. Fonty 5760	CAY108076	307262	D. Sabatier & J.-F. Molino 5340	CAY083543	136003
D. Sabatier & E. Fonty 5760	CAY108077	307262	D. Sabatier & J.-F. Molino 5341	CAY083625	136069
D. Sabatier & E. Fonty 5762	CAY111808	308679	D. Sabatier & J.-F. Molino 5347	CAY083544	136004
D. Sabatier & J.-F. Molino 4820	CAY055483	122728	D. Sabatier & J.-F. Molino 5530	CAY110483	302615
D. Sabatier & J.-F. Molino 4834	CAY108332	307580	D. Sabatier & J.-F. Molino 5530	CAY110484	302615
D. Sabatier & J.-F. Molino 4834	CAY108333	307580	D. Sabatier & J.-F. Molino 5550	CAY110476	302605
D. Sabatier & J.-F. Molino 4836	CAY065673	180231	D. Sabatier & J.-F. Molino 5552	CAY112311	309008
D. Sabatier & J.-F. Molino 4836	CAY065674	180231	D. Sabatier & J.-F. Molino 5552	CAY112312	309008
D. Sabatier & J.-F. Molino 4836	CAY065675	180231	D. Sabatier & J.-F. Molino 5554	CAY110485	302616
D. Sabatier & J.-F. Molino 4945	CAY072480	131724	D. Sabatier & J.-F. Molino 5555	CAY112313	309009
D. Sabatier & J.-F. Molino 4945	CAY072481	131724	D. Sabatier & J.-F. Molino 5555	CAY112314	309009
D. Sabatier & J.-F. Molino 4951	CAY072512	131751	D. Sabatier & J.-F. Molino 5556	CAY101324	303983
D. Sabatier & J.-F. Molino 4958	CAY047409	130061	D. Sabatier & J.-F. Molino 5556	CAY101325	303983
D. Sabatier & J.-F. Molino 4958	CAY047410	130061	D. Sabatier & J.-F. Molino 5620	CAY111731	308503
D. Sabatier & J.-F. Molino 4977	CAY072482	131725	D. Sabatier & J.-F. Molino 5620	CAY111732	308503
D. Sabatier & J.-F. Molino 5005	CAY073264	132362	D. Sabatier & J.-F. Molino 5625	CAY111938	308821
D. Sabatier & J.-F. Molino 5006bis	CAY072487	131730	D. Sabatier & J.-F. Molino 5628	CAY108476	307642
D. Sabatier & J.-F. Molino 5012	CAY073265	132363	D. Sabatier & J.-F. Molino 5629	CAY108286	307529
D. Sabatier & J.-F. Molino 5016	CAY072491	131734	D. Sabatier & J.-F. Molino 5629	CAY108287	307529
D. Sabatier & J.-F. Molino 5017	CAY072532	131765	D. Sabatier & J.-F. Molino 5634	CAY108327	307576
D. Sabatier & J.-F. Molino 5025	CAY072494	131736	D. Sabatier & J.-F. Molino 5634	CAY108328	307576
D. Sabatier & J.-F. Molino 5026	CAY072495	131737	D. Sabatier & J.-F. Molino 5642	CAY111829	308693
D. Sabatier & J.-F. Molino 5026	CAY072496	131737	D. Sabatier & J.-F. Molino 5647	CAY110480	302611
D. Sabatier & J.-F. Molino 5027	CAY072537	131770	D. Sabatier & J.-F. Molino 5648	CAY108460	307931
D. Sabatier & J.-F. Molino 5029	CAY072538	131771	D. Sabatier & J.-F. Molino 5649	CAY111827	308692
D. Sabatier & J.-F. Molino 5070	CAY068572	133594	D. Sabatier & J.-F. Molino 5649	CAY111828	308692
D. Sabatier & J.-F. Molino 5070	CAY068573	133594	D. Sabatier & J.-F. Molino 5651	CAY108383	307637
D. Sabatier & J.-F. Molino 5072	CAY104862	306667	D. Sabatier & J.-F. Molino 5653	CAY103390	305358
D. Sabatier & J.-F. Molino 5072	CAY104863	306667	D. Sabatier & J.-F. Molino 5653	CAY103391	305358
D. Sabatier & J.-F. Molino 5076	CAY070542	134070	D. Sabatier & J.-F. Molino 5671	CAY104439	306530
D. Sabatier & J.-F. Molino 5076	CAY070543	134070	D. Sabatier & J.-F. Molino 5671	CAY104440	306530
D. Sabatier & J.-F. Molino 5079	CAY078252	200672	D. Sabatier & J.-F. Molino 5672	CAY108082	307267
D. Sabatier & J.-F. Molino 5082	CAY111930	308816	D. Sabatier & J.-F. Molino 5672	CAY108083	307267
D. Sabatier & J.-F. Molino 5082	CAY111931	308816	D. Sabatier & J.-F. Molino 5683	CAY103758	305609
D. Sabatier & J.-F. Molino 5082	CAY111932	308816	D. Sabatier & J.-F. Molino 5683	CAY103759	305609
D. Sabatier & J.-F. Molino 5086	CAY068717	133647	D. Sabatier & J.-F. Molino 5692	CAY108088	307270
D. Sabatier & J.-F. Molino 5086	CAY068718	133647	D. Sabatier & J.-F. Molino 5692	CAY108089	307270
D. Sabatier & J.-F. Molino 5099	CAY104857	306663	D. Sabatier & J.-F. Molino 5705	CAY108095	307275
D. Sabatier & J.-F. Molino 5101	CAY069672	133839	D. Sabatier & J.-F. Molino 5707	CAY103444	305366
D. Sabatier & J.-F. Molino 5117	CAY069682	133849	D. Sabatier & J.-F. Molino 5708	CAY103794	305702
D. Sabatier & J.-F. Molino 5127	CAY104860	306666	D. Sabatier & J.-F. Molino 5717	CAY111651	307826
D. Sabatier & J.-F. Molino 5127	CAY104861	306666	D. Sabatier & J.-F. Molino 5717	CAY111652	307826
D. Sabatier & J.-F. Molino 5128	CAY069684	133851	D. Sabatier & J.-F. Molino 5722	CAY108450	307921
D. Sabatier & J.-F. Molino 5130	CAY110516	302651	D. Sabatier & J.-F. Molino 5722	CAY192120	307921
D. Sabatier & J.-F. Molino 5134	CAY108479	307645	D. Sabatier & J.-F. Molino 5727	CAY108504	307666
			D. Sabatier & J.-F. Molino 5728	CAY103772	305620

Appendix 4. — Continuation.

Collecte	CB	id_echantillon	Collecte	CB	id_echantillon
D. Sabatier & J.-F. Molino 5728	CAY103773	305620	D. Sabatier & M.-F. Prévost 2559	CAY042696	73809
D. Sabatier & J.-F. Molino 5738	CAY108086	307269	D. Sabatier & M.-F. Prévost 2560	CAY033960	73810
D. Sabatier & J.-F. Molino 5738	CAY108087	307269	D. Sabatier & M.-F. Prévost 2561		73811
D. Sabatier & J.-F. Molino 5750	CAY108581	307953	D. Sabatier & M.-F. Prévost 2568	CAY097787	73818
D. Sabatier & J.-F. Molino 5750	CAY165648	307953	D. Sabatier & M.-F. Prévost 2569	CAY000778	73819
D. Sabatier & J.-F. Molino 5794	CAY109144	308352	D. Sabatier & M.-F. Prévost 2575	CAY168700	73823
D. Sabatier & J.-F. Molino 5794	CAY112385	308352	D. Sabatier & M.-F. Prévost 2578	CAY008653	73825
D. Sabatier & J.-F. Molino 5794	CAY112386	308352	D. Sabatier & M.-F. Prévost 2578	CAY008654	73825
D. Sabatier & J.-L. Smock 5782	CAY108457	307928	D. Sabatier & M.-F. Prévost 2585	CAY107805	73833
D. Sabatier & M.-F. Prévost 1266	CAY100045	72614	D. Sabatier & M.-F. Prévost 2602	CAY031108	73850
D. Sabatier & M.-F. Prévost 1266	CAY100046	72614	D. Sabatier & M.-F. Prévost 2611	CAY192105	73860
D. Sabatier & M.-F. Prévost 1345	CAY107354	72705	D. Sabatier & M.-F. Prévost 2617	CAY099164	73866
D. Sabatier & M.-F. Prévost 1345	CAY107355	72705	D. Sabatier & M.-F. Prévost 2645	CAY099881	73896
D. Sabatier & M.-F. Prévost 1351	CAY100022	72713	D. Sabatier & M.-F. Prévost 2696	CAY097849	73949
D. Sabatier & M.-F. Prévost 1354	CAY082906	72717	D. Sabatier & M.-F. Prévost 2703	CAY049672	73956
D. Sabatier & M.-F. Prévost 1354	CAY082907	72717	D. Sabatier & M.-F. Prévost 2716	CAY189806	73971
D. Sabatier & M.-F. Prévost 1363	CAY156799	72726	D. Sabatier & M.-F. Prévost 2718	CAY192141	73973
D. Sabatier & M.-F. Prévost 1363	CAY156800	72726	D. Sabatier & M.-F. Prévost 2751	CAY043034	73996
D. Sabatier & M.-F. Prévost 1559	CAY097591	72902	D. Sabatier & M.-F. Prévost 2753	CAY026799	73998
D. Sabatier & M.-F. Prévost 1637	CAY099288	72958	D. Sabatier & M.-F. Prévost 2760	CAY031023	74006
D. Sabatier & M.-F. Prévost 1671	CAY193432	72985	D. Sabatier & M.-F. Prévost 2780		86961
D. Sabatier & M.-F. Prévost 1671	CAY193433	72985	D. Sabatier & M.-F. Prévost 2800	CAY195214	74044
D. Sabatier & M.-F. Prévost 1672	CAY199260	72986	D. Sabatier & M.-F. Prévost 2831	CAY177097	74073
D. Sabatier & M.-F. Prévost 1684	CAY102052	72995	D. Sabatier & M.-F. Prévost 2842	CAY042966	74084
D. Sabatier & M.-F. Prévost 1836	CAY082642	73126	D. Sabatier & M.-F. Prévost 2872	CAY007061	74107
D. Sabatier & M.-F. Prévost 1843	CAY099804	73132	D. Sabatier & M.-F. Prévost 2876	CAY001845	74111
D. Sabatier & M.-F. Prévost 1853	CAY098820	73143	D. Sabatier & M.-F. Prévost 2882	CAY009447	74117
D. Sabatier & M.-F. Prévost 1856	CAY084813	73146	D. Sabatier & M.-F. Prévost 2889	CAY088924	74125
D. Sabatier & M.-F. Prévost 1877	CAY031026	73170	D. Sabatier & M.-F. Prévost 2894		74130
D. Sabatier & M.-F. Prévost 1917	CAY043030	73210	D. Sabatier & M.-F. Prévost 2895	CAY100190	74131
D. Sabatier & M.-F. Prévost 1973	CAY043076	73261	D. Sabatier & M.-F. Prévost 2898	CAY167670	74134
D. Sabatier & M.-F. Prévost 2116	CAY042266	73388	D. Sabatier & M.-F. Prévost 2901	CAY075801	74135
D. Sabatier & M.-F. Prévost 2121	CAY193730	73393	D. Sabatier & M.-F. Prévost 2927	CAY043774	74166
D. Sabatier & M.-F. Prévost 2147	CAY185133	73420	D. Sabatier & M.-F. Prévost 2930	CAY165217	74171
D. Sabatier & M.-F. Prévost 2152	CAY183282	73425	D. Sabatier & M.-F. Prévost 2942		74182
D. Sabatier & M.-F. Prévost 2157	CAY191411	73431	D. Sabatier & M.-F. Prévost 2944	CAY080172	74185
D. Sabatier & M.-F. Prévost 2157	CAY191412	73431	D. Sabatier & M.-F. Prévost 2947	CAY185764	74189
D. Sabatier & M.-F. Prévost 2159	CAY190613	73433	D. Sabatier & M.-F. Prévost 2956		74197
D. Sabatier & M.-F. Prévost 2173	CAY030032	73445	D. Sabatier & M.-F. Prévost 2958	CAY160216	74199
D. Sabatier & M.-F. Prévost 2173	CAY030033	73445	D. Sabatier & M.-F. Prévost 2970	CAY165157	74212
D. Sabatier & M.-F. Prévost 2173	CAY080685	73445	D. Sabatier & M.-F. Prévost 2973	CAY098899	74216
D. Sabatier & M.-F. Prévost 2173	CAY080686	73445	D. Sabatier & M.-F. Prévost 2979	CAY042066	74222
D. Sabatier & M.-F. Prévost 2185	CAY192317	73456	D. Sabatier & M.-F. Prévost 3016	CAY003325	74250
D. Sabatier & M.-F. Prévost 2189	CAY116863	73463	D. Sabatier & M.-F. Prévost 3022	CAY214371	74256
D. Sabatier & M.-F. Prévost 2194	CAY043877	73468	D. Sabatier & M.-F. Prévost 3035	CAY165798	74269
D. Sabatier & M.-F. Prévost 2198	CAY081991	73473	D. Sabatier & M.-F. Prévost 3064	CAY087415	74293
D. Sabatier & M.-F. Prévost 2203	CAY027253	73478	D. Sabatier & M.-F. Prévost 3081	CAY111686	74310
D. Sabatier & M.-F. Prévost 2233	CAY048406	73507	D. Sabatier & M.-F. Prévost 3086	CAY191286	74315
D. Sabatier & M.-F. Prévost 2248	CAY178538	73518	D. Sabatier & M.-F. Prévost 3090	CAY189896	74319
D. Sabatier & M.-F. Prévost 2249	CAY096542	73519	D. Sabatier & M.-F. Prévost 3102	CAY114685	74330
D. Sabatier & M.-F. Prévost 2255	CAY178805	73524	D. Sabatier & M.-F. Prévost 3117	CAY167030	74345
D. Sabatier & M.-F. Prévost 2255	CAY178806	73524	D. Sabatier & M.-F. Prévost 3117	CAY167031	74345
D. Sabatier & M.-F. Prévost 2263		73531	D. Sabatier & M.-F. Prévost 3125	CAY162725	74353
D. Sabatier & M.-F. Prévost 2267	CAY200963	73535	D. Sabatier & M.-F. Prévost 3161	CAY029984	74388
D. Sabatier & M.-F. Prévost 2268	CAY089646	73537	D. Sabatier & M.-F. Prévost 3165	CAY076943	74392
D. Sabatier & M.-F. Prévost 2270	CAY165155	73540	D. Sabatier & M.-F. Prévost 3169	CAY031029	74396
D. Sabatier & M.-F. Prévost 2278	CAY043876	73548	D. Sabatier & M.-F. Prévost 3209	CAY082410	74436
D. Sabatier & M.-F. Prévost 2279	CAY158812	73549	D. Sabatier & M.-F. Prévost 3215	CAY162616	74442
D. Sabatier & M.-F. Prévost 2281	CAY196811	73551	D. Sabatier & M.-F. Prévost 3219		74445
D. Sabatier & M.-F. Prévost 2341	CAY196731	73603	D. Sabatier & M.-F. Prévost 3243	CAY165393	74466
D. Sabatier & M.-F. Prévost 2343	CAY109649	73605	D. Sabatier & M.-F. Prévost 3253	CAY006187	74476
D. Sabatier & M.-F. Prévost 2344	CAY042014	73606	D. Sabatier & M.-F. Prévost 3262	CAY197999	74485
D. Sabatier & M.-F. Prévost 2417	CAY165272	73681	D. Sabatier & M.-F. Prévost 3296	CAY163440	74519
D. Sabatier & M.-F. Prévost 2495	CAY043096	73753	D. Sabatier & M.-F. Prévost 3298	CAY088902	74521
D. Sabatier & M.-F. Prévost 2537	CAY165235	73788	D. Sabatier & M.-F. Prévost 3306	CAY033552	74529
D. Sabatier & M.-F. Prévost 2538	CAY214488	73789	D. Sabatier & M.-F. Prévost 3308	CAY034501	74532
D. Sabatier & M.-F. Prévost 2541	CAY043746	73791	D. Sabatier & M.-F. Prévost 3308	CAY034502	74532
D. Sabatier & M.-F. Prévost 2541	CAY043747	73791	D. Sabatier & M.-F. Prévost 3319	CAY100306	74543

Appendix 4. — Continuation.

Collecte	CB	id_echantillon	Collecte	CB	id_echantillon
D. Sabatier & M.-F. Prévost 3333	CAY189775	74556	D. Sabatier & M.-F. Prévost 4341	CAY206212	88502
D. Sabatier & M.-F. Prévost 3346	CAY001999	74568	D. Sabatier & M.-F. Prévost 4359	CAY017468	113987
D. Sabatier & M.-F. Prévost 3348		74570	D. Sabatier & M.-F. Prévost 4359	CAY200803	113987
D. Sabatier & M.-F. Prévost 3354	CAY107358	74576	D. Sabatier & M.-F. Prévost 4361	CAY192538	320838
D. Sabatier & M.-F. Prévost 3370	CAY028646	74593	D. Sabatier & M.-F. Prévost 4385	CAY198563	89253
D. Sabatier & M.-F. Prévost 3379	CAY163881	74603	D. Sabatier & M.-F. Prévost 4403	CAY215462	89260
D. Sabatier & M.-F. Prévost 3382	CAY081853	74605	D. Sabatier & M.-F. Prévost 4419	CAY108566	307898
D. Sabatier & M.-F. Prévost 3391	CAY029186	74612	D. Sabatier & M.-F. Prévost 4421	CAY089644	200750
D. Sabatier & M.-F. Prévost 3394	CAY199131	74614	D. Sabatier & M.-F. Prévost 4423	CAY000185	107662
D. Sabatier & M.-F. Prévost 3400	CAY079378	74619	D. Sabatier & M.-F. Prévost 4424	CAY085370	136540
D. Sabatier & M.-F. Prévost 3407	CAY099816	74623	D. Sabatier & M.-F. Prévost 4425	CAY107529	307016
D. Sabatier & M.-F. Prévost 3414	CAY195010	74630	D. Sabatier & M.-F. Prévost 4426	CAY001097	108122
D. Sabatier & M.-F. Prévost 3420	CAY030040	74636	D. Sabatier & M.-F. Prévost 4426	CAY001098	108122
D. Sabatier & M.-F. Prévost 3422	CAY098399	74638	D. Sabatier & M.-F. Prévost 4427	CAY016723	110526
D. Sabatier & M.-F. Prévost 3426	CAY034924	74642	D. Sabatier & M.-F. Prévost 4427	CAY016724	110526
D. Sabatier & M.-F. Prévost 3550	CAY197874	74706	D. Sabatier & M.-F. Prévost 4427	CAY016725	110526
D. Sabatier & M.-F. Prévost 3562	CAY165792	74716	D. Sabatier & M.-F. Prévost 4429	CAY206224	313288
D. Sabatier & M.-F. Prévost 3564	CAY198650	74718	D. Sabatier & M.-F. Prévost 4429	CAY206225	313288
D. Sabatier & M.-F. Prévost 3568	CAY089965	74722	D. Sabatier & M.-F. Prévost 4432	CAY004660	108813
D. Sabatier & M.-F. Prévost 3589	CAY178561	74744	D. Sabatier & M.-F. Prévost 4432	CAY004661	108813
D. Sabatier & M.-F. Prévost 3593	CAY185126	74748	D. Sabatier & M.-F. Prévost 4434	CAY004662	108814
D. Sabatier & M.-F. Prévost 3640	CAY001864	74799	D. Sabatier & M.-F. Prévost 4434	CAY004663	108814
D. Sabatier & M.-F. Prévost 3648	CAY089709	74806	D. Sabatier & M.-F. Prévost 4631	CAY018341	109274
D. Sabatier & M.-F. Prévost 3651	CAY109612	74808	D. Sabatier & M.-F. Prévost 4675	CAY193839	321169
D. Sabatier & M.-F. Prévost 3657	CAY165639	74814	D. Sabatier & M.-F. Prévost 4826	CAY055784	123501
D. Sabatier & M.-F. Prévost 3658	CAY007148	74815	D. Sabatier & M.-F. Prévost 4826	CAY055785	123501
D. Sabatier & M.-F. Prévost 3658	CAY007149	74815	D. Sabatier & M.-F. Prévost 4829	CAY065661	180223
D. Sabatier & M.-F. Prévost 3660	CAY034219	74817	D. Sabatier & M.-F. Prévost 4830	CAY055795	123917
D. Sabatier & M.-F. Prévost 3660	CAY034220	74817	D. Sabatier & M.-F. Prévost 4830	CAY055796	123917
D. Sabatier & M.-F. Prévost 3663	CAY007037	74820	D. Sabatier & M.-F. Prévost 4848	CAY164787	317540
D. Sabatier & M.-F. Prévost 3670	CAY114692	74827	D. Sabatier & M.-F. Prévost 4857	CAY058938	126741
D. Sabatier & M.-F. Prévost 3678	CAY193511	74836	D. Sabatier & M.-F. Prévost 4857	CAY058939	126741
D. Sabatier & M.-F. Prévost 3692	CAY003899	74850	D. Sabatier & M.-F. Prévost 4910	CAY047412	130063
D. Sabatier & M.-F. Prévost 3758	CAY193781	74918	D. Sabatier & M.-F. Prévost 4910	CAY047413	130063
D. Sabatier & M.-F. Prévost 3762	CAY096960	74922	D. Sabatier & M.-F. Prévost 4914	CAY047415	130065
D. Sabatier & M.-F. Prévost 3766	CAY199246	74925	D. Sabatier & M.-F. Prévost 4921	CAY065641	180210
D. Sabatier & M.-F. Prévost 3780	CAY043050	74939	D. Sabatier & M.-F. Prévost 4921	CAY065642	180210
D. Sabatier & M.-F. Prévost 3797	CAY028845	74955	D. Sabatier & M.-F. Prévost 4921	CAY065643	180210
D. Sabatier & M.-F. Prévost 3797	CAY028846	74955	D. Sabatier & M.-F. Prévost 4926	CAY047418	130067
D. Sabatier & M.-F. Prévost 3833	CAY031036	74989	D. Sabatier & M.-F. Prévost 4926	CAY047419	130067
D. Sabatier & M.-F. Prévost 3850	CAY166404	75005	D. Sabatier & M.-F. Prévost 4928	CAY072506	131746
D. Sabatier & M.-F. Prévost 3855	CAY007081	75010	D. Sabatier & M.-F. Prévost 4928	CAY072507	131746
D. Sabatier & M.-F. Prévost 3857	CAY215784	75012	D. Sabatier & M.-F. Prévost 4962	CAY065656	180219
D. Sabatier & M.-F. Prévost 3886	CAY109661	75039	D. Sabatier & M.-F. Prévost 4962	CAY065657	180219
D. Sabatier & M.-F. Prévost 3896	CAY097639	75052	D. Sabatier & M.-F. Prévost 5046	CAY073268	132366
D. Sabatier & M.-F. Prévost 3937	CAY034915	75090	D. Sabatier & M.-F. Prévost 5046	CAY073269	132366
D. Sabatier & M.-F. Prévost 3944	CAY194933	75097	D. Sabatier & M.-F. Prévost 5047	CAY073329	132395
D. Sabatier & M.-F. Prévost 3946	CAY041950	75099	D. Sabatier & M.-F. Prévost 5047	CAY073330	132395
D. Sabatier & M.-F. Prévost 3946	CAY041950	75099	D. Sabatier & M.-F. Prévost 5049	CAY073270	132367
D. Sabatier & M.-F. Prévost 3953	CAY163640	75106	D. Sabatier & M.-F. Prévost 5055	CAY073318	132387
D. Sabatier & M.-F. Prévost 3967	CAY198628	75115	D. Sabatier & M.-F. Prévost 5055	CAY073319	132387
D. Sabatier & M.-F. Prévost 3983	CAY081846	75132	D. Sabatier & M.-F. Prévost 5276	CAY093832	300365
D. Sabatier & M.-F. Prévost 4064	CAY166403	75201	D. Sabatier & M.-F. Prévost 5402	CAY089482	137854
D. Sabatier & M.-F. Prévost 4066	CAY184139	75203	D. Sabatier & M.-F. Prévost 5403	CAY086411	136681
D. Sabatier & M.-F. Prévost 4068	CAY169307	75205	D. Sabatier & M.-F. Prévost 5411	CAY088389	137359
D. Sabatier & M.-F. Prévost 4072	CAY043897	75207	D. Sabatier & M.-F. Prévost 5421	CAY110488	302620
D. Sabatier & M.-F. Prévost 4087	CAY008073	75221	D. Sabatier & M.-F. Prévost 5421	CAY110489	302620
D. Sabatier & M.-F. Prévost 4124	CAY115324	75249	D. Sabatier & M.-F. Prévost 5595	CAY112114	308827
D. Sabatier & M.-F. Prévost 4126	CAY206134	75251	D. Sabatier & P. Birnbaum 4451	CAY004655	108809
D. Sabatier & M.-F. Prévost 4137	CAY198606	75259	D. Sabatier & P. Birnbaum 4608	CAY032504	122343
D. Sabatier & M.-F. Prévost 4170		75282	D. Sabatier & S. Gonzalez 5370	CAY088478	137377
D. Sabatier & M.-F. Prévost 4177	CAY024594	75288	D. Sabatier & S. Gonzalez 5371	CAY089473	137845
D. Sabatier & M.-F. Prévost 4200	CAY042105	75301	D. Sabatier & S. Gonzalez 5381	CAY088483	137382
D. Sabatier & M.-F. Prévost 4201	CAY074911	75302	D. Sabatier & S. Gonzalez 5385	CAY110349	302445
D. Sabatier & M.-F. Prévost 4221	CAY178911	318524	D. Sabatier & S. Gonzalez 5385	CAY110350	302445
D. Sabatier & M.-F. Prévost 4226	CAY090187	75307	D. Sabatier & S. Gonzalez 5387	CAY109443	307864
D. Sabatier & M.-F. Prévost 4234	CAY165256	75309	D. Sabatier & S. Gonzalez 5387	CAY111703	307864
D. Sabatier & M.-F. Prévost 4282	CAY191746	320771	D. Sabatier & S. Gonzalez 5387	CAY111704	307864
			D. Sabatier & S. Gonzalez 5399	CAY088490	137387

Appendix 4. — Continuation.

Collecte	CB	id_echantillon	Collecte	CB	id_echantillon
D. Sabatier 1001	CAY081771	72387	D. Sabatier 3146	CAY002075	74374
D. Sabatier 1009	CAY157743	72395	D. Sabatier 3500	CAY027874	74658
D. Sabatier 1018	CAY158435	72404	D. Sabatier 3516	CAY049349	74673
D. Sabatier 1018	CAY158436	72404	D. Sabatier 3519	CAY098228	74675
D. Sabatier 1027	CAY098263	72413	D. Sabatier 3519	CAY098229	74675
D. Sabatier 1027	CAY098264	72413	D. Sabatier 3521	CAY028874	74677
D. Sabatier 1031	CAY193899	72417	D. Sabatier 3524	CAY109616	74680
D. Sabatier 1051	CAY216065	72435	D. Sabatier 3526	CAY003186	74681
D. Sabatier 1052	CAY191524	72436	D. Sabatier 3528	CAY033408	74683
D. Sabatier 1055	CAY075808	72439	D. Sabatier 3535	CAY089738	74693
D. Sabatier 1098	CAY099540	72476	D. Sabatier 3536	CAY191122	74695
D. Sabatier 1106	CAY099808	88434	D. Sabatier 3540	CAY208014	74699
D. Sabatier 1170	CAY024988	72540	D. Sabatier 3547	CAY021939	115835
D. Sabatier 1170	CAY024989	72540	D. Sabatier 3556	CAY178426	74710
D. Sabatier 1196	CAY081839	72563	D. Sabatier 3571	CAY194240	74726
D. Sabatier 1213	CAY024995	88445	D. Sabatier 3573	CAY028735	74729
D. Sabatier 1219	CAY099184	72578	D. Sabatier 3573	CAY028736	74729
D. Sabatier 1220		72579	D. Sabatier 3574		100683
D. Sabatier 1234	CAY160526	315544	D. Sabatier 3575	CAY158548	74730
D. Sabatier 1237	CAY089915	72587	D. Sabatier 4370	CAY059003	126801
D. Sabatier 1242	CAY157848	72592	D. Sabatier 4462	CAY018329	109260
D. Sabatier 1249	CAY062800	72599	D. Sabatier 4462	CAY018330	109260
D. Sabatier 1267	CAY098464	72615	D. Sabatier 4462	CAY018331	109260
D. Sabatier 1277	CAY028804	72625	D. Sabatier 4682	CAY056511	124542
D. Sabatier 1423	CAY214275	72785	D. Sabatier 4707	CAY037573	120032
D. Sabatier 1430	CAY099105	72792	D. Sabatier 4815	CAY032512	122350
D. Sabatier 1433	CAY007948	72795	D. Sabatier 4853	CAY112115	308828
D. Sabatier 1462	CAY107609	72817	D. Sabatier 4861	CAY065604	180175
D. Sabatier 1464	CAY097878	72819	D. Sabatier 4861	CAY065605	180175
D. Sabatier 1488	CAY089687	72843	D. Sabatier 4862	CAY072503	131743
D. Sabatier 1512	CAY042000	72867	D. Sabatier 4863	CAY072504	131744
D. Sabatier 1516		72871	D. Sabatier 4872	CAY065607	180177
D. Sabatier 1528	CAY100055	72881	D. Sabatier 4873	CAY065608	180178
D. Sabatier 1539	CAY165265	72891	D. Sabatier 4874	CAY065609	180180
D. Sabatier 1541	CAY214307	72893	D. Sabatier 4896	CAY175566	319490
D. Sabatier 1546	CAY030251	72896	D. Sabatier 4896	CAY175567	319490
D. Sabatier 1691	CAY088543	73001	D. Sabatier 4897	CAY065610	180181
D. Sabatier 1772	CAY089724	73064	D. Sabatier 4897	CAY213525	180181
D. Sabatier 2100	CAY049582	73372	D. Sabatier 4897	CAY213526	180181
D. Sabatier 2102	CAY016579	109948	D. Sabatier 5139	CAY088462	137363
D. Sabatier 2102	CAY016580	109948	D. Sabatier 5571	CAY104864	306668
D. Sabatier 2241	CAY098713	73512	D. Sabatier 5571	CAY104865	306668
D. Sabatier 2246	CAY194601	73516	D. Sabatier 5572	CAY108066	307252
D. Sabatier 2259	CAY010165	73528	D. Sabatier 5573	CAY104787	306623
D. Sabatier 2294	CAY043878	73561	D. Sabatier 5574	CAY109439	307855
D. Sabatier 2300	CAY010148	73566	D. Sabatier 5574	CAY111685	307855
D. Sabatier 2302	CAY007810	73568	D. Sabatier 5778	CAY108070	307257
D. Sabatier 2306	CAY042703	73572	D. Sabatier 5785	CAY112111	308825
D. Sabatier 2308	CAY091640	73574	D. Sabatier 5785	CAY112112	308825
D. Sabatier 2312		73577	D. Sabatier 5792	CAY111681	307853
D. Sabatier 2316	CAY191365	73580	D. Sabatier 5792	CAY111682	307853
D. Sabatier 2317	CAY033078	73581	D. Sabatier 5799	CAY112116	308829
D. Sabatier 2321	CAY032799	73585	D. Sabatier 5843	CAY164785	317538
D. Sabatier 2329	CAY176772	73590	D. Sabatier 824	CAY009805	72262
D. Sabatier 2338	CAY100320	73600	D. Sabatier 824	CAY009806	72262
D. Sabatier 2339	CAY098596	73601	D. Sabatier 824	CAY009807	72262
D. Sabatier 2346		73608	D. Sabatier 828	CAY114668	72265
D. Sabatier 2347	CAY165574	73609	D. Sabatier 839	CAY009621	72276
D. Sabatier 2372	CAY192736	73637	D. Sabatier 839	CAY009622	72276
D. Sabatier 2447	CAY158896	73707	D. Sabatier 839	CAY009623	72276
D. Sabatier 2448	CAY158810	73708	D. Sabatier 842		72279
D. Sabatier 2450	CAY099386	73710	D. Sabatier 844	CAY160615	72281
D. Sabatier 2529	CAY023804	73781	D. Sabatier 844	CAY160769	72281
D. Sabatier 2530	CAY174605	73782	D. Sabatier 854	CAY104025	72290
D. Sabatier 28	CAY107778	71638	D. Sabatier 854	CAY104026	72290
D. Sabatier 28	CAY107779	71638	D. Sabatier 858	CAY097377	72293
D. Sabatier 3005	CAY165795	74237	D. Sabatier 865	CAY086653	72300
D. Sabatier 3012	CAY046448	74246	D. Sabatier 865	CAY086654	72300

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
D. Sabatier 876	CAY183724	72310	G. Cremers & M. Hoff 11240	CAY046830	16510
D. Sabatier 876	CAY183725	72310	G. Cremers & P. Petronelli 11776	CAY090377	17038
D. Sabatier 922	CAY034012	72333	G. Cremers 14582	CAY157875	86731
D. Sabatier 925	CAY163835	72336	G. Cremers 14582	CAY157876	86731
D. Sabatier 931	CAY024880	200021	G. Cremers 6509	CAY043457	11766
D. Sabatier 931	CAY024881	200021	G. Cremers 6509	CAY043458	11766
D. Sabatier 960	CAY190805	320674	G. Cremers 7287	CAY030116	12561
D. Sabatier 965	CAY180659	72360	G. Cremers 7287	CAY074017	12561
D. Sabatier 965	CAY180660	72360	G. Cremers 8233		13527
D. Sabatier 981	CAY084858	72369	G. Cremers 8385	CAY178924	13678
D. Sabatier 984	CAY097873	72370	G. Cremers 8385	CAY178925	13678
D. Sabatier 984	CAY097874	72370	G. Cremers 8397	CAY030810	13690
D. Sabatier 990		72376	G. Cremers 8397	CAY030811	13690
D. Sabatier 996	CAY029806	72382	G. Cremers 8448	CAY074712	13741
D. Sabatier et al. 4393	CAY028795	89250	G. Cremers 8448	CAY074713	13741
D. Sabatier et al. 4457	CAY018320	109264	G. Cremers 8448	CAY074714	13741
D. Sabatier et al. 4457	CAY018321	109264	H. de Foresta 125ter	CAY028226	116329
D. Sabatier et al. 4604	CAY004963	109255	H. de Foresta 125ter	CAY028227	116329
D. Sabatier et al. 4622	CAY018373	109315	H. de Foresta 490	CAY062806	25083
D. Sabatier et al. 4623	CAY018363	109312	H. de Foresta 760	CAY213342	25338
D. Sabatier et al. 4623	CAY018364	109312	H. de Foresta 760	CAY213343	25338
D. Sabatier et al. 4712	CAY028312	116348	H. Jacquemin 1763	CAY027031	47219
D. Sabatier et al. 4712	CAY028313	116348	H. Jacquemin 1763	CAY027032	47219
D. Sabatier et al. 4780	CAY032505	122344	H. Jacquemin 1903	CAY084951	47350
D. Sabatier et al. 4781	CAY036008	118893	H. Jacquemin 1903	CAY084952	47350
D. Sabatier et al. 4786	CAY036005	118891	H. Jacquemin 1903	CAY084953	47350
D. Sabatier et al. 4794	CAY037587	120045	H. Jacquemin 2435	CAY178197	47821
D. Sabatier et al. 4880	CAY112647	309162	H. Puig 12055	CAY040283	68911
D. Sabatier et al. 4891	CAY182887	322615	H. Richard & A. Mambe 622	CAY113931	310344
D. Sabatier et al. 5161	CAY069651	133820	H. Richard & Bagadi 674	CAY113954	310464
D. Sabatier et al. 5250	CAY089467	137841	H. Richard & J. Ateni 432	CAY113910	310325
D. Sabatier et al. 5252	CAY088465	137366	H. Richard 70	CAY084997	136501
D. Sabatier et al. 5252	CAY088466	137366	H. Richard 70	CAY084998	136501
D. Sabatier et al. 5262	CAY088473	137372	H. van der Werff et al. 12951	CAY184925	86206
D. Sabatier et al. 5802	CAY109471	308551	H.D. Clarke 3146	CAY177774	318415
D. Sabatier et al. 5802	CAY109472	308551	H.D. Clarke 5485	CAY167091	316444
D. Sabatier et al. 5803	CAY109473	308552	I. Petrov 185	CAY024986	65590
D. Sabatier et al. 5805	CAY109478	308558	I. Petrov 185	CAY024987	65590
D. Sabatier et al. 5805	CAY109479	308558	J. Engel & P. Petronelli 19		324443
D. Sabatier et al. 5806	CAY111815	308684	J. Engel et al. 14		324140
D. Sabatier et al. 5806	CAY111816	308684	J.-F. Molino & D. Sabatier 1703	CAY101028	123719
D. Sabatier et al. 5812	CAY175547	319483	J.-F. Molino & D. Sabatier 1725	CAY029130	108271
D. Sabatier et al. 5812	CAY175548	319483	J.-F. Molino & D. Sabatier 1948	CAY004673	108822
D. Sabatier et al. 6003	CAY175584	319504	J.-F. Molino & D. Sabatier 1968	CAY004968	112287
D. Sabatier et al. 6009		324134	J.-F. Molino & D. Sabatier 1987	CAY037600	120056
D. Sabatier et al. 6043		324133	J.-F. Molino & D. Sabatier 1991	CAY037604	120060
D. Sabatier et al. 6122	CAY175563	319489	J.-F. Molino & D. Sabatier 1993		123908
D. Sabatier et al. 6122	CAY175564	319489	J.-F. Molino & D. Sabatier 1994	CAY037607	120062
D. Sabatier et al. 6122	CAY175565	319489	J.-F. Molino & D. Sabatier 1994	CAY037608	120062
D. Sabatier et al. 6161		324135	J.-F. Molino & D. Sabatier 1995	CAY037609	120063
D. Sabatier et al. 6272	CAY175612	319540	J.-F. Molino & D. Sabatier 1995	CAY037610	120063
F. Billiet & B. Jadin 1232	CAY203839	3797	J.-F. Molino & D. Sabatier 2096	CAY056355	124406
F. Billiet & B. Jadin 1232	CAY203840	3797	J.-F. Molino & D. Sabatier 2099	CAY056358	124409
F. Billiet & B. Jadin 1431		3875	J.-F. Molino & D. Sabatier 2133	CAY071558	131545
F. Billiet 1286	CAY188245	3834	J.-F. Molino & D. Sabatier 2147	CAY071566	131551
F. Hallé 1018	CAY028108	43133	J.-F. Molino & D. Sabatier 2158	CAY071577	131560
F. Hallé 1018	CAY028109	43133	J.-F. Molino & D. Sabatier 2158	CAY071578	131560
F. Hallé 626	CAY009779	43056	J.-F. Molino & D. Sabatier 2158	CAY071579	131560
F. Hallé 629	CAY030373	43057	J.-F. Molino & D. Sabatier 2164	CAY073354	132410
G. Angel 29	CAY107617	87030	J.-F. Molino & D. Sabatier 2164	CAY073355	132410
G. Bourdy 2941	CAY057276	125237	J.-F. Molino & D. Sabatier 2173	CAY073213	132320
G. Cremers & F. Crozier 15215	CAY101097	303911	J.-F. Molino & D. Sabatier 2174	CAY073372	132424
G. Cremers & J.-J. de Granville 13176	CAY042621	18399			
G. Cremers & J.-J. de Granville 13176	CAY042622	18399			
G. Cremers & J.-J. de Granville 13176	CAY101109	18399			
G. Cremers & J.-J. de Granville 13176	CAY101110	18399			

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
J.-F. Molino & D. Sabatier 2174	CAY073373	132424	J.-F. Molino & D. Sabatier 2773	CAY111910	308774
J.-F. Molino & D. Sabatier 2218	CAY073393	132439	J.-F. Molino & D. Sabatier 2788	CAY112310	309007
J.-F. Molino & D. Sabatier 2230	CAY073402	132448	J.-F. Molino & D. Sabatier 2797	CAY104841	306649
J.-F. Molino & D. Sabatier 2230	CAY073403	132448	J.-F. Molino & D. Sabatier 2803	CAY111676	307849
J.-F. Molino & D. Sabatier 2246	CAY073228	132332	J.-F. Molino & D. Sabatier 2810	CAY104628	306572
J.-F. Molino & D. Sabatier 2248	CAY073420	132462	J.-F. Molino & D. Sabatier 2815	CAY104634	306578
J.-F. Molino & D. Sabatier 2249	CAY073421	132463	J.-F. Molino & D. Sabatier 2816	CAY111653	307828
J.-F. Molino & D. Sabatier 2249	CAY073422	132463	J.-F. Molino & D. Sabatier 2834	CAY111678	307851
J.-F. Molino & D. Sabatier 2250	CAY073423	132464	J.-F. Molino & D. Sabatier 2834	CAY111679	307851
J.-F. Molino & D. Sabatier 2254	CAY073424	132465	J.-F. Molino & D. Sabatier 2844	CAY104840	306648
J.-F. Molino & D. Sabatier 2254	CAY073425	132465	J.-F. Molino & D. Sabatier 2846	CAY104847	306653
J.-F. Molino & D. Sabatier 2255	CAY073350	132408	J.-F. Molino & D. Sabatier 2850	CAY164794	317545
J.-F. Molino & D. Sabatier 2255	CAY073351	132408	J.-F. Molino & D. Sabatier 2869	CAY164543	317086
J.-F. Molino & D. Sabatier 2262	CAY073235	132338	J.-F. Molino & D. Sabatier 2871	CAY164544	317087
J.-F. Molino & D. Sabatier 2262	CAY073236	132338	J.-F. Molino & D. Sabatier 3313	CAY164504	317046
J.-F. Molino & D. Sabatier 2267	CAY073429	132469	J.-F. Molino & D. Sabatier 3313	CAY164505	317046
J.-F. Molino & D. Sabatier 2270	CAY073357	132412	J.-F. Molino & M.-F. Prévost 2531	CAY104426	306499
J.-F. Molino & D. Sabatier 2270	CAY073358	132412	J.-F. Molino 731	CAY043008	122879
J.-F. Molino & D. Sabatier 2284	CAY081074	135581	J.-F. Molino 907	CAY185009	123010
J.-F. Molino & D. Sabatier 2287	CAY077768	134209	J.-F. Molino 907	CAY185010	123010
J.-F. Molino & D. Sabatier 2287	CAY077769	134209	J.-F. Molino 921	CAY165928	123016
J.-F. Molino & D. Sabatier 2290	CAY077772	134211	J.-F. Molino 921	CAY165929	123016
J.-F. Molino & D. Sabatier 2307	CAY081084	135591	J.-F. Molino 936	CAY043011	123024
J.-F. Molino & D. Sabatier 2309	CAY081085	135592	J.-F. Molino 938	CAY048618	123026
J.-F. Molino & D. Sabatier 2319	CAY093843	300376	J.-F. Molino 1107	CAY191831	123150
J.-F. Molino & D. Sabatier 2329	CAY081089	135596	J.-F. Molino 1155	CAY201026	123188
J.-F. Molino & D. Sabatier 2332	CAY077786	134223	J.-F. Molino 1181	CAY017300	123207
J.-F. Molino & D. Sabatier 2332	CAY077787	134223	J.-F. Molino 1181	CAY117276	123207
J.-F. Molino & D. Sabatier 2335	CAY077789	134225	J.-F. Molino 1197	CAY034281	123228
J.-F. Molino & D. Sabatier 2337	CAY077791	134227	J.-F. Molino 1197	CAY034282	123228
J.-F. Molino & D. Sabatier 2349	CAY077796	134232	J.-F. Molino 1198	CAY099521	107184
J.-F. Molino & D. Sabatier 2365	CAY081096	135603	J.-F. Molino 1363	CAY093089	123434
J.-F. Molino & D. Sabatier 2367	CAY081098	135605	J.-F. Molino 1556	CAY042995	123599
J.-F. Molino & D. Sabatier 2399	CAY081104	135611	J.-F. Molino 1556	CAY042996	123599
J.-F. Molino & D. Sabatier 2401	CAY080998	135515	J.-F. Molino 1617	CAY097555	123648
J.-F. Molino & D. Sabatier 2403	CAY081000	135517	J.-F. Molino 1766	107190	
J.-F. Molino & D. Sabatier 2417	CAY093847	300380	J.-F. Molino 1769	CAY177161	123762
J.-F. Molino & D. Sabatier 2419	CAY081018	135535	J.-F. Molino 1769	CAY177162	123762
J.-F. Molino & D. Sabatier 2436	CAY081027	135543	J.-F. Molino 1769	CAY080698	123886
J.-F. Molino & D. Sabatier 2437	CAY089462	137836	J.-F. Molino 1924	CAY071196	131444
J.-F. Molino & D. Sabatier 2438	CAY093794	300327	J.-F. Molino 2144	CAY071197	131444
J.-F. Molino & D. Sabatier 2442	CAY081032	135547	J.-F. Molino 2144	CAY073409	132453
J.-F. Molino & D. Sabatier 2448	CAY081109	135616	J.-F. Molino 2235	CAY173359	317945
J.-F. Molino & D. Sabatier 2461	CAY083920	136296	J.-F. Molino 3328	CAY220112	323767
J.-F. Molino & D. Sabatier 2464	CAY109508	308460	J.-F. Molino 3429	CAY220109	323762
J.-F. Molino & D. Sabatier 2464	CAY109509	308460	J.-F. Molino 3430		
J.-F. Molino & D. Sabatier 2484	CAY093799	300332	J.-F. Molino et al. 1661	CAY162966	123683
J.-F. Molino & D. Sabatier 2469	CAY083639	136082	J.-F. Molino et al. 1661	CAY162967	123683
J.-F. Molino & D. Sabatier 2496	CAY084969	136474	J.-F. Molino et al. 1852	CAY181740	123825
J.-F. Molino & D. Sabatier 2505	CAY084972	136477	J.-F. Molino et al. 1981	CAY004960	112292
J.-F. Molino & D. Sabatier 2657	CAY104420	306484	J.-F. Molino et al. 2007	CAY035622	118650
J.-F. Molino & D. Sabatier 2661	CAY104416	306481	J.-F. Molino et al. 2041	CAY035638	118666
J.-F. Molino & D. Sabatier 2664	CAY109464	307843	J.-F. Molino et al. 2041	CAY035639	118666
J.-F. Molino & D. Sabatier 2664	CAY111670	307843	J.-F. Molino et al. 2070	CAY056330	124386
J.-F. Molino & D. Sabatier 2673	CAY104854	306660	J.-F. Molino et al. 2070	CAY056331	124386
J.-F. Molino & D. Sabatier 2679	CAY220162	323823	J.-F. Molino et al. 2071	CAY056332	124387
J.-F. Molino & D. Sabatier 2679	CAY220163	323823	J.-F. Molino et al. 2075	CAY056338	124391
J.-F. Molino & D. Sabatier 2692	CAY108505	307667	J.-F. Molino et al. 2076	CAY056339	124392
J.-F. Molino & D. Sabatier 2692	CAY109460	307667	J.-F. Molino et al. 2083	CAY071165	131426
J.-F. Molino & D. Sabatier 2693	CAY104869	306671	J.-F. Molino et al. 2083	CAY071166	131426
J.-F. Molino & D. Sabatier 2696	CAY104853	306659	J.-F. Molino et al. 2086	CAY056348	124399
J.-F. Molino & D. Sabatier 2708	CAY111648	307823	J.-F. Molino et al. 2108	CAY056367	124417
J.-F. Molino & D. Sabatier 2713	CAY111909	308771	J.-F. Molino et al. 2112	CAY056372	124421
J.-F. Molino & D. Sabatier 2714	CAY108330	307578	J.-F. Molino et al. 2112	CAY056373	124421
J.-F. Molino & D. Sabatier 2716	CAY111640	307816	J.-F. Molino et al. 2116	CAY056376	124424
J.-F. Molino & D. Sabatier 2719	CAY111677	307850	J.-F. Molino et al. 2124	CAY071173	131430
J.-F. Molino & D. Sabatier 2736	CAY111907	308769	J.-F. Molino et al. 2130	CAY071184	131437
J.-F. Molino & D. Sabatier 2751	CAY111649	307824	J.-F. Molino et al. 2143	CAY071564	131549

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
J.-F. Molino et al. 3305	CAY161999	317040	J.-J. de Granville 17659	CAY111313	304502
J.-F. Molino et al. 3309	CAY164501	317043	J.-J. de Granville 17659	CAY111314	304502
J.-F. Molino et al. 3315	CAY220157	323815	J.-J. de Granville 17659	CAY111322	304502
J.-F. Molino et al. 3325	CAY164512	317052	J.-J. de Granville 17664	CAY111335	304599
J.-F. Molino et al. 3329	CAY164793	317544	J.-J. de Granville 17664	CAY111336	304599
J.-F. Molino et al. 3333	CAY164515	317056	J.-J. de Granville 17664	CAY111337	304599
J.-F. Molino et al. 3352	CAY164530	317073	J.-J. de Granville 1869	CAY097293	28657
J.-F. Molino et al. 3366	CAY164532	317075	J.-J. de Granville 2200	CAY167222	28938
J.-F. Molino et al. 3371	CAY164541	317084	J.-J. de Granville 2200	CAY167223	28938
J.-F. Molino et al. 3372	CAY164542	317085	J.-J. de Granville 2902	CAY104000	29618
J.-F. Molino et al. 3375	CAY164535	317078	J.-J. de Granville 2902	CAY104001	29618
J.-F. Molino et al. 3382	CAY220124	323783	J.-J. de Granville 3327	CAY097273	30033
J.-F. Molino et al. 3397	CAY164520	317061	J.-J. de Granville 3411	CAY203825	30115
J.-F. Molino et al. 3406	CAY164526	317069	J.-J. de Granville 3411	CAY203826	30115
J.-F. Molino et al. 3423	CAY164517	317058	J.-J. de Granville 3411	CAY203827	30115
J.-F. Villiers & C. Feuillet 1974	CAY115300	85196	J.-J. de Granville 3411	CAY203828	30115
J.-F. Villiers & C. Feuillet 2086	CAY028096	85307	J.-J. de Granville 3750	CAY203890	30581
J.-F. Villiers 1873	CAY090056	85093	J.-J. de Granville 3750	CAY203891	30581
J.-F. Villiers 1873	CAY090057	85093	J.-J. de Granville 44	CAY024895	26602
J.-J. de Granville & F. Crozier 17010	CAY045625	129332	J.-J. de Granville 44	CAY024896	26602
J.-J. de Granville & F. Crozier 17010	CAY045626	129332	J.-J. de Granville 4866	CAY157732	32310
J.-J. de Granville & F. Kahn 11201	CAY201174	108516	J.-J. de Granville 4866	CAY157733	32310
J.-J. de Granville & F. Kahn 11201	CAY201175	108516	J.-J. de Granville 488	CAY163694	27222
J.-J. de Granville & F. Kahn 11201	CAY201176	108516	J.-J. de Granville 488	CAY163695	27222
J.-J. de Granville & F. Kahn 11201	CAY201177	108516	J.-J. de Granville 5550	CAY161483	33674
J.-J. de Granville & F. Kahn 11201	CAY201178	108516	J.-J. de Granville 5550	CAY161484	33674
J.-J. de Granville & F. Kahn 11201	CAY201179	108516	J.-J. de Granville 5909	CAY003766	34003
J.-J. de Granville & F. Kahn 11201	CAY201180	108516	J.-J. de Granville 638	CAY074838	27367
J.-J. de Granville & F. Kahn 11201	CAY201181	108516	J.-J. de Granville 652	CAY196158	27381
J.-J. de Granville & F. Kahn 11201	CAY201182	108516	J.-J. de Granville 652	CAY196159	27381
J.-J. de Granville & F. Kahn 5410	CAY024862	33420	J.-J. de Granville 7000	CAY088444	35006
J.-J. de Granville & F. Kahn 5410	CAY024863	33420	J.-J. de Granville 7000	CAY088445	35006
J.-J. de Granville 10319	CAY203865	38339	J.-J. de Granville 7000	CAY088446	35006
J.-J. de Granville 10319	CAY203866	38339	J.-J. de Granville 7000	CAY088447	35006
J.-J. de Granville 10319	CAY203867	38339	J.-J. de Granville 7203	CAY202720	35200
J.-J. de Granville 10319	CAY203868	38339	J.-J. de Granville 7203	CAY202721	35200
J.-J. de Granville 10319	CAY203869	38339	J.-J. de Granville 7203	CAY202722	35200
J.-J. de Granville 10319	CAY203870	38339	J.-J. de Granville 7203	CAY202723	35200
J.-J. de Granville 10319	CAY203871	38339	J.-J. de Granville 7203	CAY202724	35200
J.-J. de Granville 10319	CAY027241	91843	J.-J. de Granville 7203	CAY202725	35200
J.-J. de Granville 1169	CAY027242	91843	J.-J. de Granville 7734	CAY043476	35717
J.-J. de Granville 1169	CAY028081	91852	J.-J. de Granville 7734	CAY210504	35717
J.-J. de Granville 1197	CAY028082	91852	J.-J. de Granville 7755	CAY115173	35738
J.-J. de Granville 1197	CAY063800	91870	J.-J. de Granville 8340	CAY117750	36327
J.-J. de Granville 1220	CAY176702	88381	J.-J. de Granville 8340	CAY117751	36327
J.-J. de Granville 13403	CAY019012	108321	J.-J. de Granville 8340	CAY117752	36327
J.-J. de Granville 13833	CAY019013	108321	J.-J. de Granville 8340	CAY117753	36327
J.-J. de Granville 14611	CAY025248	115336	J.-J. de Granville B-3724	CAY194665	30527
J.-J. de Granville 1475	CAY028088	28273	J.-J. de Granville B-3724	CAY194666	30527
J.-J. de Granville 1475	CAY028089	28273	J.-J. de Granville B-3833	CAY026682	30723
J.-J. de Granville 1475	CAY028090	28273	J.-J. de Granville B-3833	CAY026683	30723
J.-J. de Granville 16824	CAY052495	128058	J.-J. de Granville B-4467	CAY007959	31550
J.-J. de Granville 17272	CAY047585	130226	J.-J. de Granville B-4467	CAY007960	31550
J.-J. de Granville 17272	CAY047586	130226	J.-J. de Granville B-4482	CAY046717	31578
J.-J. de Granville 17651	CAY098123	301315	J.-J. de Granville B-4488	CAY088442	31590
J.-J. de Granville 17651	CAY098124	301315	J.-J. de Granville B-4488	CAY088443	31590
J.-J. de Granville 17651	CAY098125	301315	J.-J. de Granville B-4549	CAY090068	91964
J.-J. de Granville 17651	CAY098126	301315	J.-J. de Granville B-4549	CAY090069	91964
J.-J. de Granville 17651	CAY098127	301315	J.-J. de Granville B-4563	CAY007736	31707
J.-J. de Granville 17651	CAY098128	301315	J.-J. de Granville B-4563	CAY007737	31707
J.-J. de Granville 17651	CAY098129	301315	J.-J. de Granville B-4741	CAY007576	32066
J.-J. de Granville 17659	CAY111306	304502	J.-J. de Granville B-4741	CAY007577	32066
J.-J. de Granville 17659	CAY111308	304502	J.-J. de Granville B-4744	CAY007384	32072
J.-J. de Granville 17659	CAY111309	304502	J.-J. de Granville B-4760	CAY046478	32105
J.-J. de Granville 17659	CAY111311	304502	J.-J. de Granville B-4760	CAY046479	32105
J.-J. de Granville 17659	CAY111312	304502	J.-J. de Granville B-4792	CAY169834	32167

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
J.-J. de Granville B-4989	CAY048597	32551	J.W. Grimes & P. Acevedo-Rodríguez 3322	CAY106937	89306
J.-J. de Granville B-5195	CAY027845	32979	J.W. Grimes & P. Acevedo-Rodríguez 3322	CAY106938	89306
J.-J. de Granville B-5195	CAY171665	32979	J.W. Grimes & P. Acevedo-Rodríguez 3322	CAY106939	89306
J.-J. de Granville B-5195	CAY171666	32979	J.W. Grimes & P. Acevedo-Rodríguez 3322	CAY106940	89306
J.-J. de Granville B-5356	CAY007638	33310	J.W. Grimes et al. 3312	CAY191022	42702
J.-J. de Granville B-5356	CAY007639	33310	K.M. Redden et al. 3306	CAY110361	302458
J.-J. de Granville B-5392	CAY160514	33382	L. Andersson 1993	CAY075547	1284
J.-J. de Granville B-5392	CAY160515	33382	L. Barrabé 232	CAY058760	126582
J.-J. de Granville B-5401	CAY167206	33401	L. Barrabé 232	CAY058761	126582
J.-J. de Granville B-5401	CAY167207	33401	L.E. Skog & C. Feuillet 7440	CAY075446	82206
J.-J. de Granville B-5436	CAY020414	33473	L.R. Phillippe et al. 26984	CAY167623	316578
J.-J. de Granville B-5436	CAY020415	33473	M. Blanc 128	CAY000097	107207
J.-J. de Granville B-5486	CAY010453	33574	M. Fleury 1348	CAY020646	107484
J.-J. de Granville B-5486	CAY010454	33574	M. Fleury 1348	CAY020647	107484
J.-J. de Granville et al. 10656	CAY171867	38682	M. Fleury 2053	CAY108609	307982
J.-J. de Granville et al. 10765	CAY214685	38792	M. Fleury 2166	CAY108718	308083
J.-J. de Granville et al. 10796	CAY184787	38823	M. Fleury 2176	CAY108711	308076
J.-J. de Granville et al. 10907		38935	M. Fleury 238		24136
J.-J. de Granville et al. 15000	CAY039089	121353	M. Fleury 567		24402
J.-J. de Granville et al. 15000	CAY039090	121353	M. Fleury 911	CAY198646	322153
J.-J. de Granville et al. 15181	CAY039292	121543	M. Hoff 5980	CAY165586	44749
J.-J. de Granville et al. 16138	CAY058515	126378	M.-F. Prévost & C. Feuillet 3964	CAY016768	110873
J.-J. de Granville et al. 16705	CAY050221	127939	M.-F. Prévost & C. Feuillet 3964	CAY016769	110873
J.-J. de Granville et al. 6000	CAY010617	34088	M.-F. Prévost & C. Feuillet 3968	CAY014333	110867
J.-J. de Granville et al. 6121	CAY092504	34199	M.-F. Prévost & C. Feuillet 3968	CAY014334	110867
J.-J. de Granville et al. 6268	CAY075704	34327	M.-F. Prévost & C. Proisy 4550	CAY035533	118581
J.-J. de Granville et al. 6330	CAY003561	34381	M.-F. Prévost & D. Barthélémy 3687	CAY018142	109289
J.-J. de Granville et al. 6480	CAY176089	34518	M.-F. Prévost & D. Barthélémy 3687	CAY018143	109289
J.-J. de Granville et al. 6480	CAY176090	34518	M.-F. Prévost & D. Sabatier 2193	CAY031106	68100
J.-J. de Granville et al. 6503	CAY097245	34541	M.-F. Prévost & D. Sabatier 2193	CAY031107	68100
J.-J. de Granville et al. 7632	CAY089669	35615	M.-F. Prévost & D. Sabatier 2249	CAY109502	68143
J.-J. de Granville et al. 7911	CAY198417	35894	M.-F. Prévost & D. Sabatier 2280	CAY114014	68170
J.-J. de Granville et al. 8057	CAY203092	36039	M.-F. Prévost & D. Sabatier 2356	CAY006020	68227
J.-J. de Granville et al. 8057	CAY203093	36039	M.-F. Prévost & D. Sabatier 2360	CAY114695	310536
J.-J. de Granville et al. 8057	CAY203094	36039	M.-F. Prévost & D. Sabatier 2377	CAY087604	68243
J.-J. de Granville et al. 8057	CAY203095	36039	M.-F. Prévost & D. Sabatier 2378	CAY086802	68244
J.-J. de Granville et al. 8057	CAY203096	36039	M.-F. Prévost & D. Sabatier 2464	CAY191557	68303
J.-J. de Granville et al. 8057	CAY203097	36039	M.-F. Prévost & D. Sabatier 2564	CAY088947	68395
J.-J. de Granville et al. 8099	CAY089805	36081	M.-F. Prévost & D. Sabatier 2610	CAY043767	68440
J.-J. de Granville et al. 8601	CAY007476	36595	M.-F. Prévost & D. Sabatier 2625	CAY081821	68452
J.-J. de Granville et al. 8765	CAY039581	36759	M.-F. Prévost & D. Sabatier 2680	CAY087447	68496
J.-J. de Granville et al. 8915	CAY009689	36912	M.-F. Prévost & D. Sabatier 2705	CAY189764	68829
J.-J. de Granville et al. 9032	CAY075775	37030	M.-F. Prévost & D. Sabatier 2742	CAY167649	68545
J.-J. de Granville et al. 9032	CAY075776	37030	M.-F. Prévost & D. Sabatier 2744	CAY162831	68547
J.-J. de Granville et al. 9032	CAY075777	37030	M.-F. Prévost & D. Sabatier 2755	CAY078270	68558
J.-J. de Granville et al. 9032	CAY075778	37030	M.-F. Prévost & D. Sabatier 2755	CAY078272	68558
J.-J. de Granville et al. 9229A	CAY027380	37232	M.-F. Prévost & D. Sabatier 2763	CAY199220	68565
J.-J. de Granville et al. 9455	CAY095706	37466	M.-F. Prévost & D. Sabatier 2763	CAY199221	68565
J.-J. de Granville et al. 9913	CAY204075	37933	M.-F. Prévost & D. Sabatier 2767	CAY185274	68569
J.-J. de Granville et al. 9913	CAY204076	37933	M.-F. Prévost & D. Sabatier 2773	CAY034138	68575
J.-J. de Granville et al. 9913	CAY204077	37933	M.-F. Prévost & D. Sabatier 2773	CAY034139	68575
J.-J. de Granville et al. 9913	CAY204078	37933	M.-F. Prévost & D. Sabatier 2780	CAY049606	68582
J.-J. de Granville et al. 9913	CAY204079	37933	M.-F. Prévost & D. Sabatier 2787	CAY100077	68589
J.-J. de Granville et al. 9913	CAY204079	37933	M.-F. Prévost & D. Sabatier 2789	CAY107595	68591
J.-J. Pipoly 10414	CAY167895	99924	M.-F. Prévost & D. Sabatier 2792	CAY006326	68594
J.-P. Lescure 708	CAY162898	51129	M.-F. Prévost & D. Sabatier 2805	CAY191048	68607
J.-P. Lescure 708	CAY162899	51129	M.-F. Prévost & D. Sabatier 2815	CAY098056	68616
J.-P. Lescure 723	CAY026991	51144	M.-F. Prévost & D. Sabatier 2817	CAY167376	68618
J.-P. Lescure 723	CAY026992	51144	M.-F. Prévost & D. Sabatier 2817	CAY167377	68618
J.-P. Lescure 816	CAY062769	51238			
J.-P. Lescure 816	CAY062770	51238			
J.-P. Lescure 816	CAY062771	51238			
J.-P. Lescure 839	CAY167890	51262			
J.-P. Lescure 839	CAY167891	51262			
J.-P. Lescure 839	CAY167892	51262			
J.-P. Lescure 877	CAY028065	51301			
J.-P. Lescure 877	CAY028066	51301			
J.-P. Lescure 943	CAY046542	51365			
J.-P. Lescure 943	CAY046543	51365			

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
M.-F. Prévost & D. Sabatier 2818	CAY192272	320807	M.-F. Prévost & P. Grenand 885	CAY189873	66918
M.-F. Prévost & D. Sabatier 2839	CAY161492	68638	M.-F. Prévost & P. Grenand 885	CAY189874	66918
M.-F. Prévost & D. Sabatier 2848	CAY158591	68646	M.-F. Prévost & P. Grenand 890	CAY169449	66923
M.-F. Prévost & D. Sabatier 2912	CAY193961	68660	M.-F. Prévost & P. Grenand 893	CAY027427	66926
M.-F. Prévost & D. Sabatier 2946	CAY092152	68691	M.-F. Prévost & P. Grenand 893	CAY027428	66926
M.-F. Prévost & D. Sabatier 2950	CAY049352	68695	M.-F. Prévost & P. Grenand 904	CAY189577	66934
M.-F. Prévost & D. Sabatier 2956	CAY033896	68701	M.-F. Prévost & P. Grenand 904	CAY189578	66934
M.-F. Prévost & D. Sabatier 2961	CAY028654	68706	M.-F. Prévost & P. Grenand 904	CAY189579	66934
M.-F. Prévost & D. Sabatier 2961	CAY028655	68706	M.-F. Prévost & P. Grenand 904	CAY189580	66934
M.-F. Prévost & D. Sabatier 2970	CAY178628	68716	M.-F. Prévost & P. Grenand 910	CAY162365	66940
M.-F. Prévost & D. Sabatier 2973	CAY098605	68719	M.-F. Prévost & P. Grenand 910	CAY162366	66940
M.-F. Prévost & D. Sabatier 2990	CAY206150	68735	M.-F. Prévost & P. Grenand 910	CAY162367	66940
M.-F. Prévost & D. Sabatier 2990	CAY206151	68735	M.-F. Prévost & P. Grenand 930	CAY095715	66957
M.-F. Prévost & D. Sabatier 2994	CAY163016	315579	M.-F. Prévost & P. Grenand 934	CAY186809	66961
M.-F. Prévost & D. Sabatier 2996	CAY028340	68740	M.-F. Prévost & P. Grenand 943	CAY049826	66970
M.-F. Prévost & D. Sabatier 3007	CAY204426	68751	M.-F. Prévost & P. Grenand 943	CAY049827	66970
M.-F. Prévost & D. Sabatier 3010	CAY213274	68753	M.-F. Prévost & P. Grenand 943	CAY049828	66970
M.-F. Prévost & D. Sabatier 3385	CAY196073	321625	M.-F. Prévost & P. Grenand 944	CAY215805	323571
M.-F. Prévost & D. Sabatier 3479	CAY016640	110074	M.-F. Prévost & P. Grenand 944	CAY215806	323571
M.-F. Prévost & D. Sabatier 3986	CAY047382	130038	M.-F. Prévost & P. Grenand 946	CAY184316	66972
M.-F. Prévost & D. Sabatier 3988	CAY017302	113265	M.-F. Prévost & P. Grenand 946	CAY184317	66972
M.-F. Prévost & D. Sabatier 3990	CAY017305	113267	M.-F. Prévost & P. Grenand 947	CAY084887	66973
M.-F. Prévost & D. Sabatier 3990	CAY017306	113267	M.-F. Prévost & P. Grenand 947	CAY084888	66973
M.-F. Prévost & D. Sabatier 3996	CAY022605	116236	M.-F. Prévost & P. Grenand 947	CAY084889	66973
M.-F. Prévost & D. Sabatier 4457	CAY037497	119976	M.-F. Prévost & P. Grenand 947	CAY084890	66973
M.-F. Prévost & D. Sabatier 4580	CAY035541	118586	M.-F. Prévost & P. Grenand 956	CAY085966	66980
M.-F. Prévost & D. Sabatier 4580	CAY035542	118586	M.-F. Prévost & P. Grenand 962	CAY112096	66985
M.-F. Prévost & D. Sabatier 4623	CAY035431	118492	M.-F. Prévost & P. Grenand 962	CAY112097	66985
M.-F. Prévost & D. Sabatier 4647	CAY035043	200243			
M.-F. Prévost & D. Sabatier 4660	CAY035828	118798	M.-F. Prévost & P. Grenand 970		66992
M.-F. Prévost & D. Sabatier 4662	CAY037512	119989	M.-F. Prévost & P. Grenand 976	CAY194684	66997
M.-F. Prévost & D. Sabatier 4664	CAY035420	118481	M.-F. Prévost & P. Grenand 976	CAY194685	66997
M.-F. Prévost & D. Sabatier 4720	CAY037535	120003	M.-F. Prévost & P. Grenand 992	CAY086074	67013
M.-F. Prévost & D. Sabatier 4766	CAY055517	122757	M.-F. Prévost & P. Grenand 995	CAY095744	67016
M.-F. Prévost & D. Sabatier 4766	CAY055518	122757	M.-F. Prévost & P. Grenand 995	CAY095745	67016
M.-F. Prévost & D. Sabatier 4766	CAY091966	122757	M.-F. Prévost & P. Grenand 995	CAY095746	67016
M.-F. Prévost & D. Sabatier 4792	CAY065618	180190	M.-F. Prévost & P. Grenand 996	CAY032977	67017
M.-F. Prévost & D. Sabatier 4818	CAY055776	123495	M.-F. Prévost & P. Grenand 996	CAY115404	67017
M.-F. Prévost & D. Sabatier 4819	CAY055777	123496			
M.-F. Prévost & D. Sabatier 4822	CAY056566	124584	M.-F. Prévost 1076	CAY032111	67090
M.-F. Prévost & D. Sabatier 4891	CAY065627	180196	M.-F. Prévost 1082	CAY017053	112488
M.-F. Prévost & D. Sabatier 4895	CAY065629	180198	M.-F. Prévost 1082	CAY017054	112488
M.-F. Prévost & D. Sabatier 4936	CAY073332	132397	M.-F. Prévost 1106	CAY069181	67111
M.-F. Prévost & D. Sabatier 4939	CAY080192	200687	M.-F. Prévost 1106	CAY069182	67111
M.-F. Prévost & D. Sabatier 4952	CAY076261	200661	M.-F. Prévost 1110	CAY034825	67115
M.-F. Prévost & D. Sabatier 4976	CAY082055	135784	M.-F. Prévost 1119	CAY205954	67123
M.-F. Prévost & P. Grenand 1000	CAY032110	67021	M.-F. Prévost 1135	CAY040240	67138
M.-F. Prévost & P. Grenand 1000	CAY185117	67021	M.-F. Prévost 1135	CAY040241	67138
M.-F. Prévost & P. Grenand 1005	CAY046932	67025	M.-F. Prévost 1137	CAY003896	67140
M.-F. Prévost & P. Grenand 1005	CAY046933	67025	M.-F. Prévost 1137	CAY003897	67140
M.-F. Prévost & P. Grenand 1005	CAY046934	67025	M.-F. Prévost 1146	CAY115500	67149
M.-F. Prévost & P. Grenand 1005	CAY046934	67025	M.-F. Prévost 1148	CAY192292	67151
M.-F. Prévost & P. Grenand 1042	CAY191332	67056	M.-F. Prévost 1148	CAY192293	67151
M.-F. Prévost & P. Grenand 1064	CAY003905	67078	M.-F. Prévost 1205	CAY187198	67197
M.-F. Prévost & P. Grenand 1964		67902	M.-F. Prévost 1205	CAY187199	67197
M.-F. Prévost & P. Grenand 2027	CAY162666	67955	M.-F. Prévost 1206	CAY047013	67198
M.-F. Prévost & P. Grenand 2032	CAY075007	67960	M.-F. Prévost 1206	CAY047014	67198
M.-F. Prévost & P. Grenand 2032	CAY075008	67960	M.-F. Prévost 1207	CAY197953	67199
M.-F. Prévost & P. Grenand 2057	CAY198921	67975	M.-F. Prévost 1207	CAY197954	67199
M.-F. Prévost & P. Grenand 2057	CAY198922	67975	M.-F. Prévost 1207	CAY197955	67199
M.-F. Prévost & P. Grenand 2059	CAY003956	67977	M.-F. Prévost 1221	CAY170325	67213
M.-F. Prévost & P. Grenand 3753	CAY021967	115857	M.-F. Prévost 1221	CAY170326	67213
M.-F. Prévost & P. Grenand 3753	CAY021968	115857	M.-F. Prévost 1271	CAY192614	67262
M.-F. Prévost & P. Grenand 3778	CAY019286	109631	M.-F. Prévost 1271	CAY192615	67262
M.-F. Prévost & P. Grenand 3778	CAY019287	109631	M.-F. Prévost 1281	CAY194218	67272
M.-F. Prévost & P. Grenand 4274	CAY022008	115898	M.-F. Prévost 1281	CAY194219	67272
M.-F. Prévost & P. Grenand 4304	CAY022635	116258	M.-F. Prévost 1291	CAY024874	67282
M.-F. Prévost & P. Grenand 4324	CAY022650	116268	M.-F. Prévost 1301	CAY165338	67292
M.-F. Prévost & P. Grenand 885	CAY189872	66918	M.-F. Prévost 1301	CAY165339	67292
			M.-F. Prévost 1361	CAY162093	67351

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
M.-F. Prévost 1361	CAY162094	67351	M.-F. Prévost 3742	CAY018189	109284
M.-F. Prévost 1364	CAY165432	67354	M.-F. Prévost 3756	CAY004937	109030
M.-F. Prévost 1364	CAY165433	67354	M.-F. Prévost 3768	CAY019216	109588
M.-F. Prévost 1460		67446	M.-F. Prévost 3768	CAY019217	109588
M.-F. Prévost 1472	CAY194491	67457	M.-F. Prévost 3775	CAY019288	109630
M.-F. Prévost 1472	CAY194492	67457	M.-F. Prévost 3775	CAY019289	109630
M.-F. Prévost 1506	CAY047772	67487	M.-F. Prévost 3781	CAY018053	109110
M.-F. Prévost 1506	CAY047773	67487	M.-F. Prévost 3840	CAY019222	109592
M.-F. Prévost 1579	CAY031957	67556	M.-F. Prévost 3840	CAY019223	109592
M.-F. Prévost 1619	CAY043575	67594	M.-F. Prévost 3845	CAY019265	109638
M.-F. Prévost 1619	CAY043576	67594	M.-F. Prévost 3847	CAY018311	109472
M.-F. Prévost 1619	CAY043577	67594	M.-F. Prévost 3856	CAY018450	109431
M.-F. Prévost 1628	CAY048355	67603	M.-F. Prévost 3900	CAY014019	110845
M.-F. Prévost 1628	CAY048356	67603	M.-F. Prévost 3900	CAY014020	110845
M.-F. Prévost 1636	CAY174617	67611	M.-F. Prévost 4014	CAY045804	129427
M.-F. Prévost 165	CAY215194	66229	M.-F. Prévost 4023	CAY017303	113266
M.-F. Prévost 165	CAY215195	66229	M.-F. Prévost 4023	CAY017304	113266
M.-F. Prévost 169	CAY164494	66233	M.-F. Prévost 4056	CAY166247	316826
M.-F. Prévost 169	CAY164495	66233	M.-F. Prévost 4056	CAY166248	316826
M.-F. Prévost 169	CAY164496	66233	M.-F. Prévost 4059	CAY017088	112515
M.-F. Prévost 1693	CAY115338	67665	M.-F. Prévost 4059	CAY017088	112515
M.-F. Prévost 1693	CAY115339	67665	M.-F. Prévost 4059	CAY017089	112515
M.-F. Prévost 1693	CAY115340	67665	M.-F. Prévost 4059	CAY017089	112515
M.-F. Prévost 1718	CAY033105	67689	M.-F. Prévost 4065	CAY015618	112538
M.-F. Prévost 1718	CAY033106	67689	M.-F. Prévost 4065	CAY015619	112538
M.-F. Prévost 178	CAY007294	66242	M.-F. Prévost 4166	CAY022009	115899
M.-F. Prévost 1810	CAY041533	67772	M.-F. Prévost 4166	CAY022010	115899
M.-F. Prévost 1810	CAY041534	67772	M.-F. Prévost 4194	CAY016006	113000
M.-F. Prévost 185	CAY176761	66249	M.-F. Prévost 4262	CAY035563	118600
M.-F. Prévost 185	CAY176762	66249	M.-F. Prévost 4262	CAY035564	118600
M.-F. Prévost 1858	CAY046758	67816	M.-F. Prévost 4476	CAY022401	116108
M.-F. Prévost 1858	CAY046759	67816	M.-F. Prévost 4523	CAY022620	116246
M.-F. Prévost 1873	CAY034904	67830	M.-F. Prévost 4523	CAY022621	116246
M.-F. Prévost 1873	CAY034905	67830	M.-F. Prévost 4566	CAY035556	118596
M.-F. Prévost 1893	CAY033226	67841	M.-F. Prévost 4566	CAY035557	118596
M.-F. Prévost 1910	CAY078356	67858	M.-F. Prévost 4684	CAY035551	118592
M.-F. Prévost 204	CAY187171	66268	M.-F. Prévost 4684	CAY035552	118592
M.-F. Prévost 204	CAY187172	66268	M.-F. Prévost 4686	CAY037524	119996
M.-F. Prévost 2071	CAY215263	67988	M.-F. Prévost 4686	CAY037525	119996
M.-F. Prévost 2071	CAY215264	67988	M.-F. Prévost 478	CAY028077	66533
M.-F. Prévost 2161	CAY117327	68069	M.-F. Prévost 478	CAY028078	66533
M.-F. Prévost 2161	CAY117328	68069	M.-F. Prévost 478	CAY028079	66533
M.-F. Prévost 2169	CAY167581	68077	M.-F. Prévost 478	CAY028080	66533
M.-F. Prévost 2170	CAY030208	68078	M.-F. Prévost 649	CAY178208	66699
M.-F. Prévost 2170	CAY030209	68078	M.-F. Prévost 709	CAY096769	66754
M.-F. Prévost 2719	CAY013727	68525	M.-F. Prévost 750	CAY213505	66794
M.-F. Prévost 2966	CAY043781	68712	M.-F. Prévost 750	CAY213506	66794
M.-F. Prévost 3064	CAY181018	68790	M.-F. Prévost 752	CAY198923	66796
M.-F. Prévost 3075	CAY027098	88498	M.-F. Prévost 752	CAY198924	66796
M.-F. Prévost 3127	CAY033816	200579	M.-F. Prévost 761	CAY191860	66804
M.-F. Prévost 3198	CAY177045	107047	M.-F. Prévost 761	CAY191861	66804
M.-F. Prévost 3207	CAY159811	107046	M.-F. Prévost 761	CAY191862	66804
M.-F. Prévost 3278	CAY165959	316382	M.-F. Prévost 791	CAY194328	66831
M.-F. Prévost 3301	CAY172179	319660	M.-F. Prévost 791	CAY194329	66831
M.-F. Prévost 3302	CAY033231	89235	M.-F. Prévost 840		66877
M.-F. Prévost 3329	CAY050029	127759	M.-F. Prévost 840		66889
M.-F. Prévost 3330	CAY117322	312115	M.-F. Prévost 854	CAY194660	
M.-F. Prévost 3379	CAY056844	124839	M.-F. Prévost et al. 4440	CAY022769	116398
M.-F. Prévost 3423	CAY068411	133490	M.-F. Prévost et al. 4450	CAY028307	116344
M.-F. Prévost 3442	CAY035913	118818	M.-F. Prévost et al. 4465	CAY035203	200383
M.-F. Prévost 3442	CAY035914	118818	M.-F. Prévost et al. 4470	CAY028308	116345
M.-F. Prévost 3456	CAY176448	318162	M.-F. Prévost et al. 4470	CAY028309	116345
M.-F. Prévost 3527	CAY168912	316876	M.-F. Prévost et al. 4604	CAY035500	118553
M.-F. Prévost 3567	CAY000004	107577	M.-F. Prévost et al. 4604	CAY035501	118553
M.-F. Prévost 3655	CAY002610	108943	M.-F. Prévost et al. 4604	CAY035502	118553
M.-F. Prévost 3655	CAY002611	108943	M.-F. Prévost et al. 4671	CAY035519	118569
M.-F. Prévost 3667	CAY018274	109309	M.-F. Prévost et al. 4695	CAY037526	119997
M.-F. Prévost 3667	CAY018275	109309	M.-F. Prévost et al. 4695	CAY037527	119997
			M.-F. Prévost et al. 5014	CAY110474	302602

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
M.J. Jansen-Jacobs et al. 230	CAY180746	94573	P. Grenand 1503	CAY089768	41997
M.J. Jansen-Jacobs et al. 2658	CAY115377	95950	P. Grenand 1503	CAY089769	41997
M.J. Jansen-Jacobs et al. 78	CAY074927	94485	P. Grenand 1503	CAY089770	41997
N. Marshall & J. Rombold 146		53747	P. Grenand 1510	CAY062790	42004
O. Poncy & J. Munzinger 1637	CAY114414	310431	P. Grenand 1510	CAY062791	42004
O. Poncy 2866	CAY156945	315004	P. Grenand 1525	CAY030390	42017
O. Poncy 901	CAY004773	107242	P. Grenand 1525	CAY030391	42017
O. Poncy et al. 1430	CAY021960	115850	P. Grenand 1525	CAY030393	42017
O. Poncy et al. 1430	CAY021961	115850	P. Grenand 1527	CAY189082	42019
O. Poncy et al. 1680	CAY056663	124671	P. Grenand 1527	CAY189083	42019
O. Poncy et al. 1746	CAY059163	126951	P. Grenand 1543	CAY112401	42035
O. Poncy et al. 1748	CAY059164	126952	P. Grenand 1543	CAY112402	42035
O. Poncy et al. 1830	CAY059234	127019	P. Grenand 1543	CAY112403	42035
O. Tostain 2818	CAY111559	307743	P. Grenand 1543	CAY112404	42035
O. Tostain 2818	CAY111560	307743	P. Grenand 1594		42087
O. Tostain et al. 1499	CAY086490	136757	P. Grenand 1776	CAY169823	42262
O. Tostain et al. 2725	CAY101513	304295	P. Grenand 1776	CAY169824	42262
O. Tostain et al. 2725	CAY101514	304295	P. Grenand 1776	CAY169825	42262
O. Tostain et al. 2725	CAY101515	304295	P. Grenand 1808	CAY003139	42297
P. Acevedo-Rodríguez et al. 4871	CAY007030	364	P. Grenand 1808	CAY003140	42297
P. Acevedo-Rodríguez et al. 4871	CAY007031	364	P. Grenand 1845	CAY033940	42334
P. Acevedo-Rodríguez et al. 4911	CAY007157	402	P. Grenand 1845	CAY033941	42334
P. Acevedo-Rodríguez et al. 5012	CAY001980	479	P. Grenand 1845	CAY117367	42334
P. Acevedo-Rodríguez et al. 5012	CAY001981	479	P. Grenand 1902	CAY049851	42396
P. Acevedo-Rodríguez et al. 6130	CAY201024	101126	P. Grenand 1902	CAY115098	42396
P. Acevedo-Rodríguez et al. 6130	CAY201025	101126	P. Grenand 1902	CAY115099	42396
P. Béna 1232	CAY005436	113911	P. Grenand 1933	CAY112101	42428
P. Béna 1232	CAY005437	113911	P. Grenand 1933	CAY112102	42428
P. Béna 1232	CAY005438	113911	P. Grenand 2107	CAY196014	42601
P. Béna 1313	CAY169866	3413	P. Grenand 2107	CAY196015	42601
P. Béna 1313	CAY169867	3413	P. Grenand 2118	CAY083302	42612
P. Béna 1313	CAY169868	3413	P. Grenand 2118	CAY083303	42612
P. Chareyre 30D	CAY019730	110212	P. Grenand 2119	CAY097324	42613
P. Grenand & M.-F. Prévost 1959	CAY181806	42454	P. Grenand 237	CAY179001	40766
P. Grenand & M.-F. Prévost 1959	CAY181807	42454	P. Grenand 237	CAY179002	40766
P. Grenand & M.-F. Prévost 2017	CAY083291	42512	P. Grenand 2868	CAY095068	42663
P. Grenand & M.-F. Prévost 2040		42535	P. Grenand 3131	CAY000352	107789
P. Grenand 1013	CAY095057	41575	P. Grenand 3502	CAY109168	308374
P. Grenand 1013	CAY095058	41575	P. Grenand 3502	CAY109169	308374
P. Grenand 1013	CAY095059	41575	P. Grenand 356	CAY096533	40889
P. Grenand 1044	CAY162020	41606	P. Grenand 510	CAY024893	41057
P. Grenand 1044	CAY162021	41606	P. Grenand 510	CAY024894	41057
P. Grenand 1116	CAY157851	41679	P. Grenand 510	CAY081233	41057
P. Grenand 1116	CAY157852	41679	P. Grenand 510	CAY081234	41057
P. Grenand 1126	CAY193940	41689	P. Grenand 543	CAY083285	41094
P. Grenand 1126	CAY193941	41689	P. Grenand 543	CAY083286	41094
P. Grenand 1126	CAY193942	41689	P. Grenand 574	CAY034859	41127
P. Grenand 1126	CAY193943	41689	P. Grenand 574	CAY034860	41127
P. Grenand 1222	CAY109273	41739	P. Grenand 574	CAY200809	41127
P. Grenand 1222	CAY109274	41739	P. Grenand 574	CAY200810	41127
P. Grenand 1291	CAY083390	41810	P. Grenand 626	CAY031015	41181
P. Grenand 1291	CAY083391	41810	P. Grenand 626	CAY031016	41181
P. Grenand 1303	CAY170332	41821	P. Grenand 626	CAY214258	41181
P. Grenand 1303	CAY170333	41821	P. Grenand 626	CAY214259	41181
P. Grenand 1303	CAY170334	41821	P. Grenand 637	CAY165874	41192
P. Grenand 1328	CAY075402	41847	P. Grenand 637	CAY165875	41192
P. Grenand 1328	CAY075403	41847	P. Grenand 642	CAY107505	41197
P. Grenand 1328	CAY075404	41847	P. Grenand 642	CAY107506	41197
P. Grenand 1488	CAY098286	41982	P. Grenand 642	CAY107507	41197
P. Grenand 1488	CAY098287	41982	P. Grenand 642	CAY107508	41197
P. Grenand 1488	CAY098364	41982	P. Grenand 645	CAY180721	41200
P. Grenand 1488	CAY098365	41982	P. Grenand 645	CAY180722	41200
P. Grenand 15	CAY071711	40535	P. Grenand 649	CAY007416	41204
			P. Grenand 649	CAY007417	41204
			P. Grenand 649	CAY007418	41204
			P. Grenand 649	CAY007419	41204
			P. Grenand 677	CAY155295	41232
			P. Grenand 677	CAY155296	41232
			P. Grenand 699	CAY089665	41255

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
P. Grenand 699	CAY089666	41255	R.A.A. Oldeman 2627		63180
P. Grenand 699	CAY089667	41255	R.A.A. Oldeman 2672	CAY196899	63265
P. Grenand 699	CAY089668	41255	R.A.A. Oldeman 2673	CAY064785	63268
P. Grenand 701	CAY006784	41257	R.A.A. Oldeman 2673	CAY064786	63268
P. Grenand 701	CAY006785	41257	R.A.A. Oldeman 2756	CAY079748	63430
P. Grenand 715	CAY074719	41271	R.A.A. Oldeman 2756	CAY079749	63430
P. Grenand 715	CAY074720	41271	R.A.A. Oldeman 3060	CAY032624	63990
P. Grenand 738	CAY048125	41294	R.A.A. Oldeman 3060	CAY032625	63990
P. Grenand 738	CAY048126	41294	R.A.A. Oldeman 3064	CAY093904	63998
P. Grenand 738	CAY048127	41294	R.A.A. Oldeman 3064	CAY093905	63998
P. Grenand 738	CAY048128	41294	R.A.A. Oldeman 3066	CAY056165	124240
P. Grenand 756	CAY048790	41312	R.A.A. Oldeman 3066	CAY056166	124240
P. Grenand 756	CAY048791	41312	R.A.A. Oldeman 3066	CAY200632	124240
P. Grenand 756	CAY048792	41312	R.A.A. Oldeman 3188	CAY198002	64244
P. Grenand 774	CAY196777	41330	R.A.A. Oldeman 3191	CAY032737	64249
P. Grenand 774	CAY196778	41330	R.A.A. Oldeman 3191	CAY032738	64249
P. Grenand 774	CAY196779	41330	R.A.A. Oldeman B-1017	CAY010019	60032
P. Grenand 774	CAY196780	41330	R.A.A. Oldeman B-1017	CAY010020	60032
P. Grenand et al. 3380	CAY098141	301327	R.A.A. Oldeman B-1122	CAY047845	60230
P. Grenand et al. 3425	CAY108763	308133	R.A.A. Oldeman B-1205	CAY111153	304123
P. Grenand et al. 3442	CAY093693	300234	R.A.A. Oldeman B-1205	CAY111154	304123
P. Grenand et al. 3521	CAY108467	307938	R.A.A. Oldeman B-1338	CAY116262	60650
P. Mutchnick 773	CAY161614	315466	R.A.A. Oldeman B-1441	CAY198164	60854
P. Mutchnick et al. 117	CAY162659	105540	R.A.A. Oldeman B-1441	CAY198165	60854
P. Petronelli 277	CAY015654	112560	R.A.A. Oldeman B-1467	CAY165590	60916
P. Petronelli 300	CAY089483	137855	R.A.A. Oldeman B-1467	CAY165591	60916
P. Petronelli 300	CAY089484	137855	R.A.A. Oldeman B-1468	CAY165292	60919
P. Petronelli 300	CAY089485	137855	R.A.A. Oldeman B-1645	CAY167084	61251
P.G. Delprete & F. Crozier 7127	CAY022363	116079	R.A.A. Oldeman B-1645	CAY167085	61251
P.G. Delprete & F. Crozier 7150	CAY022338	116057	R.A.A. Oldeman B-1825	CAY090339	61602
P.J.M. Maas et al. 8062	CAY200779	53695	R.A.A. Oldeman B-1825	CAY090340	61602
P.J.M. Maas et al. 9351	CAY039532	121758	R.A.A. Oldeman B-1913	CAY097321	61766
P.J.M. Maas et al. 9635	CAY070183	133946	R.A.A. Oldeman B-1936	CAY075203	61810
R.A.A. Oldeman & C. Sastre 156	CAY095820	58381	R.A.A. Oldeman B-1936	CAY075204	61810
R.A.A. Oldeman & C. Sastre 163	CAY049496	58395	R.A.A. Oldeman B-1946	CAY116311	61830
R.A.A. Oldeman & C. Sastre 167	CAY009778	58405	R.A.A. Oldeman B-2026	CAY090151	61993
R.A.A. Oldeman & C. Sastre 77	CAY049495	58210	R.A.A. Oldeman B-2028	CAY183737	61997
R.A.A. Oldeman 1221	CAY189304	60423	R.A.A. Oldeman B-2084	CAY027273	62120
R.A.A. Oldeman 1221	CAY189305	60423	R.A.A. Oldeman B-2101	CAY048587	62155
R.A.A. Oldeman 1314	CAY191357	60601	R.A.A. Oldeman B-2312	CAY101119	62543
R.A.A. Oldeman 1862	CAY008581	61672	R.A.A. Oldeman B-2317	CAY194287	62554
R.A.A. Oldeman 1862	CAY008582	61672	R.A.A. Oldeman B-2398	CAY096048	62712
R.A.A. Oldeman 1862	CAY008583	61672	R.A.A. Oldeman B-2398	CAY096049	62712
R.A.A. Oldeman 1897	CAY033263	61736	R.A.A. Oldeman B-2424	CAY176060	62764
R.A.A. Oldeman 1897	CAY033264	61736	R.A.A. Oldeman B-2424	CAY176061	62764
R.A.A. Oldeman 1908	CAY064796	61758	R.A.A. Oldeman B-2687	CAY194502	63293
R.A.A. Oldeman 1908	CAY064797	61758	R.A.A. Oldeman B-2784	CAY189138	63486
R.A.A. Oldeman 1912	CAY085710	61764	R.A.A. Oldeman B-2784	CAY189139	63486
R.A.A. Oldeman 1912	CAY085711	61764	R.A.A. Oldeman B-2825	CAY090210	63568
R.A.A. Oldeman 1912	CAY085712	61764	R.A.A. Oldeman B-2825	CAY090211	63568
R.A.A. Oldeman 1916	CAY109520	61773	R.A.A. Oldeman B-3045	CAY195727	63960
R.A.A. Oldeman 1916	CAY109521	61773	R.A.A. Oldeman B-3045	CAY195728	63960
R.A.A. Oldeman 2001		61942	R.A.A. Oldeman B-3097	CAY030485	64062
R.A.A. Oldeman 2106	CAY065392	62164	R.A.A. Oldeman B-3097	CAY030486	64062
R.A.A. Oldeman 2106	CAY065393	62164	R.A.A. Oldeman B-3110	CAY026857	64089
R.A.A. Oldeman 2106	CAY065396	62164	R.A.A. Oldeman B-3110	CAY026858	64089
R.A.A. Oldeman 2193	CAY180743	62325	R.A.A. Oldeman B-3139	CAY020459	64144
R.A.A. Oldeman 2238	CAY176237	62408	R.A.A. Oldeman B-3139	CAY020460	64144
R.A.A. Oldeman 2238	CAY176238	62408	R.A.A. Oldeman B-3228	CAY090435	64322
R.A.A. Oldeman 2242	CAY028067	62412	R.A.A. Oldeman B-3261	CAY178283	64387
R.A.A. Oldeman 2242	CAY028068	62412	R.A.A. Oldeman B-3288	CAY194388	64438
R.A.A. Oldeman 2386	CAY181760	62690	R.A.A. Oldeman B-3358		64556
R.A.A. Oldeman 2386	CAY181761	62690	R.A.A. Oldeman B-3538	CAY163912	64739
R.A.A. Oldeman 2386	CAY181762	62690	R.A.A. Oldeman B-3553	CAY010466	64753
R.A.A. Oldeman 2386	CAY181763	62690	R.A.A. Oldeman B-3579	CAY046739	64779
R.A.A. Oldeman 2386			R.A.A. Oldeman B-3580	CAY184971	64780
R.A.A. Oldeman 2386			R.A.A. Oldeman B-3602	CAY159683	64803
R.A.A. Oldeman 2386			R.A.A. Oldeman B-4007	CAY001986	64984

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
R.A.A. Oldeman B-4007	CAY001987	64984	S.A. Mori & C. Gracie 23929	CAY030129	57889
R.A.A. Oldeman B-4125	CAY100867	65097	S.A. Mori & C. Gracie 24205	CAY196036	57978
R.A.A. Oldeman B-4125	CAY100868	65097	S.A. Mori & C. Gracie 24212	CAY076795	57984
R.A.A. Oldeman B-4132	CAY010613	65105	S.A. Mori & J.J. Pipoly 15425	CAY214546	55888
R.A.A. Oldeman B-4183	CAY021241	65156	S.A. Mori & J.J. Pipoly 15459	CAY078062	55909
R.A.A. Oldeman B-4183	CAY021242	65156	S.A. Mori & J.J. Pipoly 15475	CAY195224	55920
R.A.A. Oldeman B-4183	CAY021243	65156	S.A. Mori & J.J. Pipoly 15498	CAY160589	55930
R.A.A. Oldeman B-521	CAY195533	59035	S.A. Mori & J.J. Pipoly 15515	CAY099204	55941
R.A.A. Oldeman B-521	CAY195534	59035	S.A. Mori & J.J. Pipoly 15515	CAY099205	55941
R.A.A. Oldeman B-652	CAY087527	59306	S.A. Mori & J.J. Pipoly 15528	CAY195632	55949
R.A.A. Oldeman B-652	CAY087528	59306	S.A. Mori & J.J. Pipoly 15534	CAY183773	55955
R.A.A. Oldeman B-688	CAY096501	59377	S.A. Mori & J.J. Pipoly 15554	CAY15970	55970
R.A.A. Oldeman B-723	CAY192676	59446	S.A. Mori & J.J. Pipoly 15562	CAY042235	55978
R.A.A. Oldeman B-813	CAY027149	59621	S.A. Mori & J.J. Pipoly 15603	CAY075992	56009
R.A.A. Oldeman B-813	CAY027150	59621	S.A. Mori & J.J. Pipoly 15603	CAY075993	56009
R.A.A. Oldeman B-849A	CAY097307	59702	S.A. Mori & N.P. Smith 25047	CAY015333	111604
R.A.A. Oldeman B-884	CAY208397	59772	S.A. Mori & T.D. Pennington 17937	CAY001970	56086
R.A.A. Oldeman T-321	CAY220404	58730	S.A. Mori & T.D. Pennington 17983	CAY030470	56117
R.A.A. Oldeman T-390	CAY048625	58800	S.A. Mori & T.D. Pennington 18001	CAY084878	56125
R.A.A. Oldeman T-390	CAY048626	58800	S.A. Mori & T.D. Pennington 18034	CAY166399	56134
R.A.A. Oldeman T-650	CAY097373	59302	S.A. Mori & T.D. Pennington 18052	CAY050679	56151
R.A.A. Oldeman T-663	CAY176074	59329	S.A. Mori & T.D. Pennington 18056	CAY032735	56190
R.A.A. Oldeman T-734	CAY189767	59465	S.A. Mori & Y. Veyret 8972	CAY160549	55310
R.A.A. Oldeman T-737	CAY169726	59471	S.A. Mori 14824	CAY030473	55410
R.A.A. Oldeman T-737	CAY169727	59471	S.A. Mori 15537	CAY087682	55958
R.A.A. Oldeman T-762	CAY098221	59523	S.A. Mori 22252	CAY043765	57074
R.A.A. Oldeman T-763	CAY064194	59524	S.A. Mori 25053	CAY015344	111593
R.A.A. Oldeman T-763	CAY064195	59524	S.A. Mori et al. 14899	CAY027852	55466
R.A.A. Oldeman T-909	CAY195460	59822	S.A. Mori et al. 14900	CAY024885	55467
R.A.A. Oldeman T-909	CAY195461	59822	S.A. Mori et al. 14900	CAY024886	55467
R.C. Ek 1745	CAY070249	134009	S.A. Mori et al. 14927	CAY002334	55486
R.C. Ek 1835	CAY070281	134041	S.A. Mori et al. 15014	CAY027877	55559
R.S. Cowan 38757		8784	S.A. Mori et al. 15024	CAY066071	55567
S. Barrier & C. Feuillet 2633	CAY104592	1809	S.A. Mori et al. 15027	CAY006115	55570
S. Barrier 4026	CAY076916	2117	S.A. Mori et al. 15028	CAY070881	55571
S. Barrier 5018	CAY106194	2648	S.A. Mori et al. 15075	CAY027885	55609
S. Barrier 5018	CAY206287	2648	S.A. Mori et al. 15115	CAY174957	55867
S. Barrier 5018	CAY206288	2648	S.A. Mori et al. 15394	CAY028666	56066
S. Gonzalez 1093	CAY108491	307655	S.A. Mori et al. 15690	CAY024585	56075
S. Gonzalez 1093	CAY108492	307655	S.A. Mori et al. 15701	CAY023778	56076
S. Gonzalez 1093	CAY116659	307655	S.A. Mori et al. 15702	CAY019160	56422
S. Gonzalez 1093	CAY116660	307655	S.A. Mori et al. 19167	CAY027837	56424
S. Gonzalez 2334	CAY168495	316669	S.A. Mori et al. 20758	CAY027849	56468
S. Gonzalez 2334	CAY168496	316669	S.A. Mori et al. 20774	CAY165775	56494
S.A. Mori & B.M. Boom 14691	CAY075173	55318	S.A. Mori et al. 20790	CAY027880	56505
S.A. Mori & B.M. Boom 14709	CAY070876	55332	S.A. Mori et al. 20800	CAY163914	56511
S.A. Mori & B.M. Boom 14721	CAY032732	55341	S.A. Mori et al. 20808	CAY027861	56545
S.A. Mori & B.M. Boom 14746	CAY027879	55358	S.A. Mori et al. 20850	CAY032144	56552
S.A. Mori & B.M. Boom 14752	CAY005994	55363	S.A. Mori et al. 20862	CAY032145	56552
S.A. Mori & B.M. Boom 14764	CAY010621	55372	S.A. Mori et al. 20862	CAY027842	56588
S.A. Mori & B.M. Boom 14784	CAY098781	55382	S.A. Mori et al. 20928	CAY096704	56604
S.A. Mori & B.M. Boom 15146	CAY177609	55665	S.A. Mori et al. 20960	CAY030998	56610
S.A. Mori & B.M. Boom 15162	CAY194077	55681	S.A. Mori et al. 20968	CAY028674	56611
S.A. Mori & B.M. Boom 15192	CAY090007	55702	S.A. Mori et al. 20973	CAY001603	56639
S.A. Mori & B.M. Boom 15225	CAY024985	55727	S.A. Mori et al. 21018	CAY007334	56802
S.A. Mori & B.M. Boom 15232	CAY165309	55734	S.A. Mori et al. 21520	CAY076393	56918
S.A. Mori & B.M. Boom 15236		55737	S.A. Mori et al. 21661	CAY157843	56942
S.A. Mori & B.M. Boom 15239	CAY044794	55740	S.A. Mori et al. 21691	CAY157844	56942
S.A. Mori & B.M. Boom 15253	CAY005527	55754	S.A. Mori et al. 21691	CAY031535	56992
S.A. Mori & B.M. Boom 15288	CAY027882	55782	S.A. Mori et al. 22034	CAY065474	57126
S.A. Mori & B.M. Boom 15302	CAY034925	55794	S.A. Mori et al. 22719	CAY112106	57168
S.A. Mori & B.M. Boom 15333	CAY196068	55819	S.A. Mori et al. 22781	CAY065484	57183
S.A. Mori & B.M. Boom 15360	CAY034217	55840	S.A. Mori et al. 22811	CAY177077	57184

Appendix 4. — Continuation.

Collecte	CB	id_échantillon	Collecte	CB	id_échantillon
S.A. Mori et al. 23207	CAY115073	57378	Service Forestier 4005	CAY192123	80409
S.A. Mori et al. 23276	CAY024335	57406	Service Forestier 4005	CAY192124	80409
S.A. Mori et al. 23307	CAY009838	57420	Service Forestier 4436	CAY157745	314891
S.A. Mori et al. 23320	CAY087072	57431	Service Forestier 4436	CAY157746	314891
S.A. Mori et al. 23340	CAY190586	57441	Service Forestier 5118	CAY093537	300099
S.A. Mori et al. 23344	CAY032771	57444	Service Forestier 5147	CAY093548	300108
S.A. Mori et al. 23417	CAY155254	57496	Service Forestier 6096	CAY010398	80743
S.A. Mori et al. 23427	CAY048485	57506	Service Forestier 6096	CAY010399	80743
S.A. Mori et al. 23464	CAY162797	57543	Service Forestier 7075	CAY092950	80892
S.A. Mori et al. 23532	CAY167283	57611	Service Forestier 7075	CAY092951	80892
S.A. Mori et al. 23682	CAY116373	57762	Service Forestier 7237	CAY163015	81011
S.A. Mori et al. 23694	CAY206070	57774	Service Forestier 7445	CAY112105	81116
S.A. Mori et al. 23792	CAY009462	57823	Service Forestier 7816	CAY101549	81321
S.A. Mori et al. 23841	CAY163902	57842	Service Forestier 7816	CAY101550	81321
S.A. Mori et al. 23898	CAY076794	57870	Service Forestier 7855	CAY167196	81351
S.A. Mori et al. 23938	CAY191099	57896	Service Forestier 7855	CAY167197	81351
S.A. Mori et al. 24002	CAY167624	57935	Service Forestier 7911	CAY097016	81400
S.A. Mori et al. 24011	CAY065537	57942	Service Forestier 7911	CAY097017	81400
S.A. Mori et al. 24021	CAY074676	57949	Service Forestier 7911	CAY097018	81400
S.A. Mori et al. 24136	CAY091925	138263	Service Forestier M-23	CAY187648	80098
S.A. Mori et al. 24258	CAY066418	131253	Service Forestier M-23	CAY187649	80098
S.A. Mori et al. 24370	CAY165615	316336	Service Forestier M-63	CAY085057	6915
S.A. Mori et al. 24757	CAY000383	107818	Service Forestier M-63	CAY085058	6915
S.A. Mori et al. 24778	CAY000379	107814	T.D. Pennington & S.A. Mori 12134	CAY100315	65402
S.A. Mori et al. 24798	CAY002718	109238	T.D. Pennington & S.A. Mori 12137	CAY088949	65405
S.A. Mori et al. 24828	CAY015406	111776	T.D. Pennington & S.A. Mori 12177	CAY089800	65446
S.A. Mori et al. 24980	CAY019864	111654	T.D. Pennington et al. 13840	CAY097565	65466
S.A. Mori et al. 24995	CAY035293	200456	T.D. Pennington et al. 13840	CAY097566	65466
S.A. Mori et al. 25034	CAY015359	111582	T.D. Pennington et al. 13867	CAY192014	65493
S.A. Mori et al. 25174	CAY015318	111566	T.W. Henkel et al. 3133	CAY191514	105317
S.A. Mori et al. 25177	CAY015315	111569	V. Hequet 159	CAY030889	88648
S.A. Mori et al. 25303	CAY015677	113251	V. Hequet 3003	CAY073057	132183
S.A. Mori et al. 25427	CAY023010	117161	V. Hequet 3018bis	CAY073070	132196
S.A. Mori et al. 25434	CAY023028	117178	V. Hequet 3018bis	CAY073114	132196
S.A. Mori et al. 25515	CAY022666	116284	W.J. Hahn 3534	CAY169018	42803
S.A. Mori et al. 25536	CAY036865	119417	X. Cuniberti 22	CAY109376	308440
S.A. Mori et al. 25554	CAY036885	119436	Y. Caraglio 508	CAY073077	132203
S.A. Mori et al. 25564	CAY036895	119447	Y. Caraglio 802	CAY019094	109542
S.A. Mori et al. 25667	CAY057038	125019			
S.A. Mori et al. 25669	CAY057040	125021			
S.A. Mori et al. 25711	CAY057079	125060			
S.A. Mori et al. 25729	CAY057094	125075			
S.A. Mori et al. 26472	CAY077948	134355			
S.A. Mori et al. 26487	CAY077959	134365			
S.A. Mori et al. 26487	CAY077960	134365			
S.A. Mori et al. 26487	CAY077961	134365			
S.A. Mori et al. 26487	CAY077962	134365			
S.A. Mori et al. 26542	CAY080517	135106			
S.A. Mori et al. 26555	CAY080521	135110			
S.A. Mori et al. 26555	CAY080522	135110			
S.A. Mori et al. 26565	CAY077995	134397			
S.A. Mori et al. 8864	CAY116315	311203			

APPENDIX 5. — Full names of abbreviated references in the checklist/Noms complets des références abrégées du catalogue.

Titre_abr	[Auteurs]	Titre
Abh. Königl. Böhm. Ges. Wiss.		Abhandlungen der Königlichen Böhmischen Gesellschaft der Wissenschaften
Abh. Königl. Ges. Wiss. Göttingen		Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen
Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss.		Abhandlungen der Mathematisch-Physikalischen Klasse der Königlich Bayerischen Akademie der Wissenschaften
Acta Amazonica Supl.		Acta Amazonica. Suplemento
Acta Biol. Venez.		Acta Biologica Venezuelica
Acta Bot. Hung.		Acta Botanica Hungarica
Acta Bot. Neerl.		Acta Botanica Neerlandica
Acta Bot. Venez.		Acta Botánica Venezolana
Acta Helv. Phys.-Math.		Acta Helvetica Physico-Mathematico-Anatomico-Botanico-Medica
Acta Horti Berg.		Acta Horti Bergiani
Actes Soc. Hist. Nat. Paris		Actes de la Société d'Histoire Naturelle de Paris
Adansonia [Baillon]	Baillon	Adansonia
Adnot. Bot.		Adnotationes botanicae
Advances Econ. Bot.		Advances in Economic Botany
Agric. Colon.		Agricoltura Colonial
Allg. Med.-Pharm. Fl.		Allgemeine medizinisch-pharmazeutische Flora
Allg. Naturgesch.		Allgemeine Naturgeschichte für alle Stände
Alsoogr. Amer.		Alsographia Americana
Amer. J. Bot.		American Journal of Botany
Amer. Midl. Naturalist		American Midland Naturalist
Anais 15 Congr. Soc. Bot. Brasil		Anais do XV Congresso da Sociedade de Botânica do Brasil
Anais Acad. Brasil. Ci.		Anais da Academia Brasileira de Ciências
Anais Reunião Sul-Amer. Bot.		Anais Reunião Sul-Americana de Botânica
Anal. Soc. Rural Argent. (Cat. Descr. Maderas)		Catalogo descriptivo de las maderas
Anales Acad. Ci. Med. Habana		Anales de la Academia de Ciencias Medicas
Anales Ci. Parag.		Anales Científicos Paraguayos
Anales Inst. Biol. Univ. Nac. México		Anales del Instituto de Biología de la Universidad Nacional de México
Anales Inst. Bot. Cavanilles		Anales del Instituto Botanico A. J. Cavanilles
Anales Inst. Méd.-Nac. México		Anales del Instituto Médico Nacional
Anales Jard. Bot. Madrid		Anales del Jardín Botánico de Madrid
Anales Mus. Hist. Nat. Montevideo		Anales del Museo de Historia Natural de Montevideo
Anales Soc. Esp. Hist. Nat.		Anales de la Sociedad Española de Historia Natural
Anexos Mem. Inst. Butantan, Secc. Bot.		Anexos das Memórias do Instituto de Butantan, Secção de Botânica
Ann. Acad. Brasil Sci.		Annaes da Academia Brasileira de Ciencias
Ann. Bot. Syst. [Walpers]	Walpers	Annales botanices systematicae
Ann. Carnegie Mus.		Annals of the Carnegie Museum
Ann. Gen. Sci. Phys.		Annales Générales des Sciences Physiques
Ann. Hort. Belge Étrangère		Annales d'Horticulture Belge et Étrangère
Ann. Inst. Colon. Marseille		Annales de l'Institut Colonial de Marseille
Ann. Jard. Bot. Buitenzorg, Suppl.		Annales du Jardin Botanique de Buitenzorg, Supplément
Ann. K. K. Naturhist. Hofmus.		Annalen des K.K. Naturhistorischen Hofmuseums
Ann. Mag. Nat. Hist.		The Annals and Magazine of Natural History
Ann. Missouri Bot. Gard.		Annals of the Missouri Botanical Garden
Ann. Mus. Bot. Lugduno-Batavi		Annales Musei Botanici Lugduno-Batavi
Ann. Mus. Colon. Marseille		Annales du Musée Colonial de Marseille
Ann. Mus. Congo		Annales du Musée du Congo Belge
Ann. Mus. Natl. Hist. Nat.		Annales du Muséum national d'Histoire Naturelle
Ann. Naturhist. Mus. Wien		Annalen des Naturhistorischen Museums in Wien
Ann. New York Acad. Sci.		Annals of the New York Academy of Sciences
Ann. Sci. Nat. (Paris)		Annales des Sciences Naturelles
Ann. Sci. Nat., Bot.		Annales des Sciences Naturelles, Botanique
Ann. Soc. Bot. Lyon		Annales de la Société Botanique de Lyon
Ann. Univ. Abidjan, C.		Annales de l'Université d'Abidjan, série C
Annuaire Conserv. Jard. Bot. Genève		Annuaire du Conservatoire et du Jardin Botaniques de Genève
Apocyn. S. Am.		On the Apocynaceae of South America
Apont.		Apontamentos phytogeographicos
Arb. Arbust. Venez.		Arboles y arbustos nuevos de Venezuela
Arbeiten Königl. Bot. Gart. Breslau		Arbeiten aus dem Kgl. Botanischen Garten zu Breslau
Arch. Bot.		Archives de Botanique (Caen)
Arch. Bot. Bull. Mens.		Archives de Botanique, Bulletin Mensuel
Arch. Bot. São Paulo		Archivos de Botânica do São Paulo
Arch. Inst. Biol. Veg.		Archivos do Instituto de Biologia Vegetal
Arch. Jard. Bot. Rio de Janeiro		Arquivos do Jardim Botânico do Rio de Janeiro
Arch. Med. Bras.		Archivo Medico Brasileiro
Arch. Mus. Hist. Nat.		Archives du Muséum d'Histoire Naturelle
Arch. Mus. Nac. Rio de Janeiro		Arquivos do Museu Nacional do Rio de Janeiro
Ark. Bot.		Arkiv för Botanik

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Ark. Bot., n.s.		<i>Arkiv für Botanik, Andra Serien</i>
Arq. Bot. Estado São Paulo		<i>Arquivos de Botânica do Estado de São Paulo</i>
Arq. Serv. Florest.		<i>Arquivos do Servico Florestal</i>
Atas Simp. Biota Amazônica		<i>Atas do Simpósio sóbre a Biota Amazônica</i>
Atlantic J.		<i>Atlantic Journal, and Friend of Knowledge</i>
Atti Riunione Sci. Ital.		<i>Atti della Riunione degli Scienziati Italiani</i>
Autik. Bot.		<i>Autikon Botanikon</i>
Beih. Bot. Centralbl.		<i>Beihefte zum Botanischen Centralblatt</i>
Beitr. Erythroxylon		<i>Beiträge zur Kenntniss der Gattung Erythroxylum</i>
Ber. Deutsch. Bot. Ges.		<i>Berichte der Deutschen Botanischen Gesellschaft</i>
Beskr. Guin. Pl.		<i>Beskrivelse af Guineiske Planter</i>
Biblioth. Bot.		<i>Bibliotheca Botanica</i>
Bih. Kongl. Svenska Vetensk.-Akad. Handl.		<i>Bihang til Kongliga Svenska Vetenskaps-Akademiens Handlingar</i>
Biol. Cent.-Amer.		<i>Biología Centrali-Americana</i>
BioLlania, ed. espec.		<i>BioLlania, edición especial</i>
Bol. Bot. Univ. São Paulo		<i>Boletim de Botânica</i>
Bol. Centro Invest. Biol. Univ. Zulia		<i>Boletín del Centro de Investigaciones Biológicas, Universidad de Zulia, Maracaibo, Venezuela.</i>
Bol. Ci. Técn. Mus. Comercial Venezuela		<i>Boletín Científico y Técnico del Museo Comercial de Venezuela</i>
Bol. Inst. Bot. (São Paulo)		<i>Boletim do Instituto de Botânica</i>
Bol. Mens. Mus. Prod. Argent.		<i>Boletín Mensual del Museo de Productos Argentinos</i>
Bol. Mus. Goeldi Hist. Nat. Ethnogr.		<i>Boletim do Museu Goeldi de Historia Natural e Ethnographia</i>
Bol. Mus. Nac. Rio de Janeiro, Bot.		<i>Boletim do Museu Nacional de Rio de Janeiro, Botânica</i>
Bol. Mus. Paraense Emílio Goeldi, N.S., Bot.		<i>Boletim do Museu Paraense Emílio Goeldi, nova série, Botânica</i>
Bol. Mus. Paraense Hist. Nat. Ethnogr.		<i>Boletim do Museu Paraense de Historia Natural e Ethnographia</i>
Bol. Soc. Argent. Bot.		<i>Boletín de la Sociedad Argentina de Botánica</i>
Bol. Soc. Venez. Ci. Nat.		<i>Boletín de la Sociedad Venezolana de Ciencias Naturales</i>
Bol. Técn. Inst. Agron. N.		<i>Boletim Técnico do Instituto Agronômico de Norte</i>
Bol. Técn. Minist. Agric.		<i>Boletín Técnico. Ministerio de Agricultura y Cria</i>
Bot. Beechey Voy.		<i>The Botany of Captain Beechey's Voyage</i>
Bot. Centralbl.		<i>Botanisches Centralblatt</i>
Bot. Cult.		<i>Le Botaniste Cultivateur</i>
Bot. Gaz.		<i>Botanical Gazette</i>
Bot. J. Linn. Soc.		<i>Botanical Journal of the Linnean Society</i>
Bot. Jahrb. Syst.		<i>Botanische Jahrbücher für Systematik</i>
Bot. Közlem.		<i>Botanikae Közlemények</i>
Bot. Mag.		<i>Botanical Magazine</i>
Bot. Mat. Med.		<i>A Botanical Materia Medica</i>
Bot. Misc.		<i>Botanical Miscellany</i>
Bot. Mus. Leafl.		<i>Botanical Museum Leaflets</i>
Bot. Nomencl.		<i>A Botanical Nomenclator</i>
Bot. Sam. Venez		<i>Über die Behandlung von Sämereien und Pflanzen des tropischen Süd-Amerika, besonders Venezuela's</i>
Bot. Tidsskr.		<i>Botanisk Tidsskrift</i>
Bot. Voy. Herald [Seemann]	Seemann	<i>The Botany of the Voyage of H.M.S. Herald</i>
Bot. Voy. Sulphur [Bentham]	Bentham	<i>The Botany of the Voyage of H.M.S. Sulphur</i>
Bull. Acad. Roy. Sci. Belgique		<i>Bulletin de l'Academie Royale des Sciences et Belles-lettres de Belgique</i>
Bull. Acad. Roy. Sci. Bruxelles		<i>Bulletin de l'Academie Royale des Sciences et Belles-lettres de Bruxelles</i>
Bull. Bernice P. Bishop Mus.		<i>Bulletin of the Bernice P. Bishop Museum. Honolulu</i>
Bull. Bur. Pl. Industr. U.S.D.A.		<i>U.S. Department of Agriculture Bureau of Plant Industry Bulletin</i>
Bull. Herb. Boissier		<i>Bulletin de l'Herbier Boissier</i>
Bull. Jard. Bot. Buitenzorg		<i>Bulletin du Jardin Botanique de Buitenzorg</i>
Bull. Jard. Bot. État Bruxelles		<i>Bulletin du Jardin Botanique de l'État à Bruxelles</i>
Bull. Jard. Bot. Natl. Belg.		<i>Bulletin du Jardin Botanique National de Belgique</i>
Bull. Mens. Soc. Linn. Paris		<i>Bulletin Mensuel de la Société Linnéenne de Paris</i>
Bull. Misc. Inform. Kew		<i>Bulletin of Miscellaneous Information</i>
Bull. Mus. Natl. Hist. Nat.		<i>Bulletin du Muséum National d'Histoire Naturelle</i>
Bull. Mus. Natl. Hist. Nat., B, Adansonia		<i>Bulletin du Muséum National d'Histoire Naturelle, série B, Adansonia</i>
Bull. New York Bot. Gard.		<i>Bulletin of the New York Botanical Garden</i>
Bull. Soc. Bot. France		<i>Bulletin de la Société Botanique de France</i>
Bull. Soc. Bot. France, Lett. Bot.		<i>Bulletin de la Société Botanique de France, Lettres botaniques</i>
Bull. Soc. Bot. Genève		<i>Bulletin de la Société Botanique de Genève</i>
Bull. Soc. Imp. Naturalistes Moscou		<i>Bulletin de la Société Impériale des Naturalistes de Moscou</i>
Bull. Soc. Linn. Normandie		<i>Bulletin de la Société Linnéenne de Normandie</i>
Bull. Soc. Natl. Acclim. France		<i>Bulletin de la Société Nationale d'Acclimatation de France</i>
Bull. Soc. Roy. Bot. Belgique		<i>Bulletin de la Société Royale de Botanique de Belgique</i>
Bull. Torrey Bot. Club		<i>Bulletin of the Torrey Botanical Club</i>
Bull. U.S. Natl. Mus.		<i>Bulletin of the United States National Museum</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Byrsonima		<i>Arbeiten aus dem Botanischen Institut des Kgl. Lyceum Hosianum in Braunsberg</i>
Cacao Manual [F. Hardy] Caribbean Forest. Cat. Fl. Domingensis Cat. Fl. Venez. [Pittier] Cat. Hist. Nat. Corrientes Cat. Horti Vindob. Cat. Pl. Cub. [Grisebach] Cat. Pl. Horti Monsp. Char. Gen. Pl. Choix Pl. Ci. & Cult. Civ. Nat. Hist. Jamaica. Comm. Lin. Telegr., Bot.	Hardy Pittier Grisebach	Hardy's Cacao Manual Caribbean Forester <i>Catalogus florae domingensis</i> <i>Catálogo de la Flora Venezolana</i> <i>Catálogo Historia Natural de Corrientes</i> <i>Catalogus horti academicii vindobonensis</i> <i>Catalogus plantarum cubensium</i> <i>Catalogus plantarum Horti Botanici Monspeliensis</i> <i>Characteres generum plantarum</i> <i>Choix de plantes</i> <i>Ciencia e Cultura</i> <i>The Civil and Natural History of Jamaica in Three Parts</i> <i>Comissão de Linhas Telegraphicas estratégicas de Matto Grosso ao Amazonia. Anexo no. 5 Historia natural. Botanica</i> <i>Commentarii Phytographicici</i> <i>Commentationes de leguminosarum generibus</i> <i>Comptes Rendus des Séances de la Société de biogéographie</i> <i>Comunicaciones Botánicas del Museo de Historia Natural de Montevideo</i> <i>Contributions from the Arnold Arboretum of Harvard University</i> <i>Contribucion al Conocimiento de los Áboles de la Argentina</i> <i>Contributions from the Gray Herbarium of Harvard University</i> <i>Contribuciones a la historia natural Colombiana</i> <i>Contributions du Jardin Botanique de Rio de Janeiro</i> <i>Contribuciones Ocasionales del Museo de Historia Natural del Colegio De La Salle</i> <i>Contributions from the United States National Herbarium</i> <i>Contributions from the University of Michigan Herbarium</i> <i>Dissertatio de Cortice Geoffraeae Surinamensis</i> <i>On the cultivation of the plants belonging to the natural order of Proteaceae</i> <i>The Cyclopædia</i> <i>Dagbok öfwer en Ostindisk resa</i> <i>Denkschriften der Königlichen Akademie der Wissenschaften zu München</i> <i>Description des Guttiferae de l'Inde</i> <i>Description of the genus Pinus</i> <i>Description des plantes nouvelles</i> <i>Descriptionis rariorum plantarum</i> <i>Descriptions of three hundred new species of South American Plants</i> <i>Diagnoses plantarum novarum Mexicanarum</i> <i>Diccionario botánico de los nombres vulgares cubanos y puerto-riqueños</i> <i>Dictionnaire des Sciences Naturelles</i> <i>Monadelphiae Classis Dissertationes Decem</i> <i>Dissertaçao sobre as plantas do Brazil</i> <i>The Druggists' Circular</i> <i>In genus Echitis observationes</i> <i>Eclogae Americanae</i> <i>The Edinburgh New Philosophical journal</i> <i>Edinburgh Philosophical Journal</i> <i>Empire Forestry Journal</i> <i>Encyclopédie Méthodique</i> <i>Encyclopédie Méthodique. Supplément</i> <i>Enumeratio plantarum Horti regii botanici Berolinensis altera</i> <i>Enumeratio Myrtacearum Brasiliensium</i> <i>Enumeratio Palmarum Novarum</i> <i>Enumeratio Plantarum</i> <i>Enumeratio Plantarum Horti Regii Berolinensis</i> <i>Enumeratio Plantarum Guatemalensis</i> <i>Enumeratio Plantarum Horti Regii Berolinensis. Supplementum</i> <i>Enumeração das substâncias brasileiras</i> <i>Enumeratio Systematica Plantarum</i> <i>An enumeration of the vascular plants known from Surinam</i> <i>Essai sur les propriétés médicales des plantes</i> <i>Essay on the plants collected by Mr. Eugene Fitzalan during Lieut. Smith's expedition to the estuary of the Burdekin (1860)</i> <i>Estudo botânico do Nordeste</i> <i>Estudios científicos del doctor Andres Posada</i> <i>Étude générale du groupe des Euphorbiacées</i> <i>De Euphorbiacearum Generibus</i>
Comm. Phytogr. Commentat. Legum. Gen. Compt. Rend. Séances Soc. Biogéogr. Comun. Bot. Mus. Hist. Nat. Montevideo Contr. Arnold Arbor. Contr. Conoc. Arb. Argent. Contr. Gray Herb. Contr. Hist. Nat. Colomb. Contr. Jard. Bot. Rio de Janeiro Contr. Ocas. Mus. Hist. Nat. Colegio De La Salle Contr. U.S. Natl. Herb. Contr. Univ. Michigan Herb. Cort. Geogr. Surinam. Cult. Prot. Cycl. [Rees] Dagb. Ostind. Resa Denkschr. Königl. Akad. Wiss. München Descr. Guttif. Inde Descr. Pinus Descr. Pl. Nouv. Descr. Rar. Pl. Surin. [Rottbøll] Descr. S. Amer. Pl. Diagn. Pl. Nov. Mexic. Dicc. Bot. Nombres Vulg. Cub. Puerto-Riq. Dict. Sci. Nat. Diss. Diss. Pl. Brazil Druggists' Circ. Echitis Eclog. Amer. Edinburgh New Philos. J. Edinburgh Philos. J. Empire For. J. Encycl. [J. Lamarck & al.] Encycl. [J. Lamarck & al.] Suppl. Enum. Hort. Berol. Alt. Enum. Myrt. Bras. Enum. Palm. Nov. Enum. Pl. [Vahl] Enum. Pl. [Willdenow] Enum. Pl. Guatem. Enum. Pl. Suppl. [Willdenow] Enum. Subst. Braz. Enum. Syst. Pl. Enum. Vasc. Pl. Surinam Essai Propr. Méd. Pl. Essay Pl. Burdekin Estud. Bot. Nordeste Braz. Estudios Científicos Étude Euphorb. Euphorb. Gen.	Rees Rottbøll	

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Euphorb. Surinam		<i>The Euphorbiaceae of Surinam</i>
Exped. Roosevelt-Rondon. Bot.		<i>Expedição Scientifica Roosevelt-Rondon. Annexo n. 2. Botanica</i>
Expl. Bot. Cuenca Maracaibo		<i>Exploraciones Botánicas y otras en la Cuenca de Maracaibo</i>
Explor. Bot. Afrique Occ. Franç.		<i>Exploration botanique de l'Afrique occidentale française</i>
Expos. Fam. Nat.		<i>Exposition des familles naturelles et de la germination des plantes</i>
Fam. Nat. Syn. Monogr.		<i>Familiarum Naturalium Regni Vegetabilis Synopses Monographicæ</i>
Feddes Repert.		<i>Feddes Repertorium</i>
Festschr. Ascherson		<i>Festschrift zur Feier des siebzigsten Geburtstages des Herrn Professor Dr. Paul Ascherson</i>
Ficus		<i>Ficus Genus</i>
Field & Lab.		<i>Field & Laboratory</i>
Fieldiana, Bot.		<i>Fieldiana, Botany</i>
Fitologiya		<i>Fitologija</i>
Fl. Antill.		<i>Flore des Antilles</i>
Fl. Avila		<i>Flora del Avila</i>
Fl. Bras. [Martius]	Martius	<i>Flora Brasiliensis</i>
Fl. Bras. Merid. [A. St.-Hil.] (quarto ed.)	Saint-Hilaire	<i>Flora Brasiliæ meridionalis (quarto edition)</i>
Fl. Brit. W.I. [Grisebach]	Grisebach	<i>Flora of the British West Indian islands</i>
Fl. Cochinch.		<i>Flora Cochinchinensis</i>
Fl. Coimbatore [M. Chandrabose & N.C. Nair]	Nair	<i>Chandrabose & Flora of Coimbatore</i>
Fl. Columb. [H. Karsten]	Karsten	<i>Florae Columbiae</i>
Fl. Cuba [Gómez & Roig]	Gómez	<i>Flora de Cuba</i>
Fl. Cuba [Gómez]	Gómez & Roig	<i>Flora de Cuba</i>
Fl. Ecuador		<i>Flora of Ecuador</i>
Fl. Filip. [F.M. Blanco]	Blanco	<i>Flora de Filipinas</i>
Fl. Flumin.		<i>Florae Fluminensis</i>
Fl. Flumin. Icon.		<i>Florae Fluminensis Icones</i>
Fl. Guadeloupe [Stehlé]	Stehlé	<i>La Flore de la Guadeloupe et Dépendance</i>
Fl. Guianas		<i>Flora of the Guianas</i>
Fl. Guyane Franç.		<i>Flore de la Guyane française</i>
Fl. Habanera		<i>Flora Habanera</i>
Fl. Hawaii. Fam.		<i>Flora Hawaiensis</i>
Fl. Huayaquilensis		<i>Flora Huayaquilensis</i>
Fl. Ilustr. Catar.		<i>Flora Ilustrada Catarinense</i>
Fl. Ind.		<i>Flora Indica</i>
Fl. Ind. Occid.		<i>Flora Indiae Occidentalis</i>
Fl. Indica [N.L. Burman]	Burman	<i>Flora Indica</i>
Fl. Ins. Austr.		<i>Florulae Insularum Australium Prodromus</i>
Fl. Jamaic. [Linnaeus]	Linnaeus	<i>Flora Jamaicensis</i>
Fl. Jamaica [Fawcett & Rendle]	Fawcett & Rendle	<i>Flora of Jamaica</i>
Fl. Jamaica [Macfadyen]	Macfadyen	<i>Flora of Jamaica</i>
Fl. Mexic.		<i>Flora mexicana</i>
Fl. N. Territory		<i>The flora of the Northern Territory</i>
Fl. Ned. Ind.		<i>Flora van Nederlandsch Indie</i>
Fl. Neotrop. Monogr.		<i>Flora Neotropica Monographs</i>
Fl. Oware		<i>Flore d'Oware</i>
Fl. Peruv. [Ruiz & Pavón]	Ruiz & Pavón	<i>Flora Peruiana</i>
Fl. Peruv. Prodr.		<i>Flora Peruviana, et Chilensis prodromus</i>
Fl. Phan. Antill. Franç.		<i>Flore phanérogamique des Antilles françaises</i>
Fl. S.E. U.S. [Small]	Small	<i>Flora of the Southeastern United States</i>
Fl. Seneg. Tent.		<i>Florae Senegambiae Tentamen</i>
Fl. Serres Jard. Eur.		<i>Flore des serres et des jardins de l'Europe</i>
Fl. South. U.S.		<i>Flora of the southern United States</i>
Fl. Suriname		<i>Flora of Suriname</i>
Fl. Trinidad & Tobago		<i>Flora of Trinidad and Tobago</i>
Fl. Trop. Afr.		<i>Flora of Tropical Africa</i>
Fl. Venez.		<i>Flora de Venezuela</i>
Fl. Venez. Guayana		<i>Flora of the Venezuelan Guayana</i>
Fragm. Bot.		<i>Fragmenta Botanica</i>
Fruct. Sem. Pl.		<i>De fructibus et seminibus plantarum</i>
Gard. Bull. Singapore		<i>Gardens' Bulletin Singapore</i>
Gard. Dict.		<i>The Gardeners Dictionary</i>
Gen. Cass. Syn.		<i>Synopsis generis Cassiae</i>
Gen. Fl. Pl.		<i>The Genera of Flowering Plants</i>
Gen. Hist.		<i>A General History of the Dichlamydeous Plants</i>
Gen. Inga, Bot.		<i>The genus Inga, Botany</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Gen. Pl. [Bentham & Hooker f.]	Bentham & Hooker	<i>Genera Plantarum</i>
Gen. Sp. Synon. Cand.		<i>Genera, species et synonyma Candolleana</i>
Gentes Herb.		<i>Gentes Herbarum</i>
Ges. Naturf. Freunde Berlin Neue Schriften		<i>Der Gesellschaft Naturforschender Freunde zu Berlin Neue Schriften</i>
Getreue Darstell. Gew.		<i>Getreue Darstellung und Beschreibung der in der Arzneykunde Gebräuchlichen Gewächse</i>
Global Fl.		<i>Plant Gateway's the Global Flora</i>
Gött. Gel. Anz.		<i>Göttingische Gelehrte Anzeigen</i>
Harvard Pap. Bot.		<i>Harvard Papers in Botany</i>
Herb. Amb.		<i>Herbarium Amboinense</i>
Het Instituut		<i>Het Instituut van Verslagen en Mededeelingen</i>
Hieronima alchorneoides		<i>Euphorbiaceae. Hieronima (Gen. nov.) alchorneoides (Sp. nov.)</i>
Hist. Fís. Cuba, Bot.		<i>Historia física, política y natural de la Isla de Cuba. Botanica</i>
Hist. Nat. Palm.		<i>Historia naturalis Palmarum</i>
Hist. Nat. Quinquinas		<i>Histoire naturelle des Quinquinas</i>
Hist. Phys. Cuba, Pl. Vasc.		<i>Histoire physique, politique et naturelle de l'ile de Cuba. Botanique. Plantes vasculaires</i>
Hist. Pl. [Baillon]	Baillon	<i>Histoire des Plantes</i>
Hist. Pl. Guiane		<i>Histoire des plantes de la Guiane Française</i>
Hist. Univ. Règne Vég., Planches		<i>Histoire Universelle du Règne Végétal, Planches</i>
Hooker's Icon. Pl.		<i>Hooker's Icônes Plantarum; or figures, with brief descriptive characters and remarks of new or rare plants</i>
Hooker's J. Bot. Kew Gard. Misc.		<i>Hooker's Journal of Botany and Kew Garden Miscellany</i>
Hort. Brit. [Loudon]	Loudon	<i>Hortus Britannicus</i>
Hort. Brit. [Sweet]	Sweet	<i>Hortus Britannicus</i>
Hort. Jamaic.		<i>Hortus Jamaicensis</i>
Hort. Nymphenb.		<i>Hortus Nymphenburgensis</i>
Hort. Suburb. Calcutt.		<i>Hortus suburbanus Calcuttensis</i>
Horti Bononi. Pl. Nov.		<i>Horti botanici Bononiensis plantae novae vel minus cognitae</i>
Hufeland. III.		<i>Hufelandiae Illustratio</i>
Icon. [Cavanilles]	Cavanilles	<i>Icônes et descriptions plantarum</i>
Icon. Pict. Pl. Rar.		<i>Icônes pictae plantarum rariorum</i>
Icon. Pl.		<i>Icônes plantarum</i>
Icon. Pl. Rar. [Jacquin]	Jacquin	<i>Icônes plantarum rariorum</i>
Icon. Sel. Pl.		<i>Icônes selectae plantarum</i>
III. Bot. Himal. Mts.		<i>Illustrations of the botany and other branches of the natural history of the Himalayan mountains</i>
III. Dict. Gard.		<i>The Illustrated Dictionary of Gardening</i>
III. Hort.		<i>L'Illustration Horticole</i>
Ind. Bibl. Pl. Col. Com. Rondon		<i>Indice bibliográfico e numérico das plantas colhidas pela Comissão Rondon</i>
Index Kew.		<i>Index Kewensis plantarum phanerogamarum</i>
Index Kew. Suppl.		<i>Index Kewensis plantarum phanerogamarum; Supplementum</i>
Index Linn. Herb.		<i>Index to the Linnean Herbarium</i>
Index Palm.		<i>Index Palmarum</i>
Index Seminum (Berlin)		<i>Index seminum in Horti Botanici Berolinensis</i>
Indig. Palms Surin.		<i>The Indigenous Palms of Suriname</i>
Inst. Rei Herb.		<i>Institutiones Rei Herbariae</i>
Isis [Oken]	Oken	<i>Isis oder Encyclopädische Zeitung.</i>
Iter Hispan.		<i>Iter Hispanicum</i>
J. Arnold Arbor.		<i>Journal of the Arnold Arboretum</i>
J. Board Agric. British Guiana		<i>Journal of the Board of Agriculture of British Guiana</i>
J. Bot.	Hooker	<i>Journal of Botany</i>
J. Bot. [Hooker]	Hooker	<i>Journal of Botany, British and Foreign</i>
J. Bot. [Morot]	Morot	<i>Journal de Botanique</i>
J. Bot. Agric.		<i>Journal de Botanique appliquée à l'Agriculture</i>
J. Bot. Res. Inst. Texas		<i>Journal of the Botanical Research Institute of Texas</i>
J. Dept. Agric. Porto Rico		<i>The journal of the Department of Agriculture of Porto Rico</i>
J. Hist. Nat.		<i>Journal d'Histoire Naturelle</i>
J. Linn. Soc., Bot.		<i>Journal of the Linnean Society, Botany</i>
J. Proc. Linn. Soc., Bot.		<i>Journal of the Proceedings of the Linnean Society, Botany</i>
J. Straits Branch Roy. Asiat. Soc.		<i>Journal of the Straits Branch of the Royal Asiatic Society</i>
J. Wash. Acad. Sci.		<i>Journal of the Washington Academy of Sciences</i>
Jaarb. Kon. Ned. Maatsch. Aanm. Tuinb.		<i>Jaarboek van de Koninklijke Nederlandse Maatschappij tot Aanmoediging van den Tuinbouw</i>
Jahrb. Gewächsk.		<i>Jahrbücher der Gewächskunde</i>
Jahrb. Königl. Bot. Gart. Berlin		<i>Jahrbuch des Königlichen Botanischen Gartens und des Botanischen Museums zu Berlin</i>
Jour. As. Soc. Bengal		<i>Journal of the Asiatic Society of Bengal</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Kew Bull.		<i>Kew Bulletin</i>
Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd.		<i>Kongelige Danske videnskabernes Selskabs Skrifter, Naturvidenskabeli Mathematiske Afdeling</i>
Kongl. Svenska Vetensk.-Akad. Handl.		<i>Kongliga Svenska Vetenskaps-Akademiens Handlingar</i>
Kongl. Vetensk.-Akad. Handl.		<i>Kongliga Vetenskaps-Akademiens Handlingar</i>
Kungl. Svenska Vetenskapsakad. Handl.		<i>Kungliga Svenska Vetenskapsakademiens Handlingar</i>
Legum. Argent.		<i>Las Leguminosas Argentinas</i>
Legum. Pl. Hawaii		<i>The Leguminous Plants of Hawaii</i>
Legumes Ilha de Maracá		<i>Legumes of the Ilha de Maracá</i>
Lista Pl. Salvador [Standley & S. Calderon]	Standley & Calderon	<i>Tropical Woods</i>
London J. Bot.		<i>London Journal of Botany</i>
London Med. J.		<i>The London Medical Journal</i>
Mag. Neuesten Entdeck. Gesammten Naturk. Ges. Naturf. Freunde Berlin		<i>Magazin für die neuesten Entdeckungen in der gesammten Naturkunde, Gesellschaft Naturforschender Freunde zu Berlin</i>
Malvenfam.		<i>Ueber einige künstliche Geschlechter aus der Malven-Familie</i>
Man. S.E. Fl.		<i>Manual of the Southeastern Flora</i>
Mant. 1 [Schultes]	Schultes	<i>Mantissa in volumen primum</i>
Mant. 3 [Schultes] Add.	Schultes	<i>Mantissa in volumen tertium. Additamentum</i>
Mant. Pl.		<i>Mantissa Plantarum</i>
Mater. Med.		<i>Materia Medica</i>
Meded. Bot. Mus. Herb. Rijks Univ. Utrecht		<i>Mededeelingen van het Botanisch Museum en Herbarium van de Rijks Universiteit te Utrecht</i>
Meded. Landbouwhoogeschool Wageningen		<i>Mededeelingen van de Landbouwhoogeschool te Wageningen</i>
Meded. Rijks-Herb.		<i>Mededeelingen van 's Rijks Herbarium</i>
Melast. Bras.		<i>Melastome Brasiliane</i>
Melet. Bot.		<i>Meletemata Botanica</i>
Mém. Acad. Imp. Sci. Saint-Pétersbourg, sér. 6, Sci. Math., Seconde Pt. Sci. Nat.		<i>Mémoires de l'Académie Impériale des Sciences de Saint-Pétersbourg</i>
Mém. Acad. Sci. Toulouse		<i>Mémoires de l'Académie des Sciences, Inscriptions et Belles-Lettres de Toulouse</i>
Mem. Cl. Sci. Fis. Mat. Nat. Reale Accad. Italia		<i>Memorie della Classe di Scienze Fisiche, Matematiche e Naturali. Reale Accademia d'Italia</i>
Mem. Mat. Fis. Soc. Ital. Sci. Modena		<i>Memorie di Matematica e di Fisica della Società Italiana delle Scienze Residente in Modena</i>
Mem. Math. Phis. Acad. Real Sci. Lisboa		<i>Memorias da Academia Real das Ciencias de Lisboa</i>
Mém. Mus. Hist. Nat.		<i>Mémoires du Muséum d'Histoire Naturelle</i>
Mém. Mus. Natl. Hist. Nat., sér. B, Bot.		<i>Mémoires du Muséum National d'Histoire Naturelle, série B, Botanique</i>
Mem. New York Bot. Gard.		<i>Memoirs of the New York Botanical Garden</i>
Mem. Pacific Trop. Bot. Gard.		<i>Memoirs of Pacific Tropical Botanical Garden</i>
Mém. Rubiac.		<i>Mémoire sur la famille des Rubiacées</i>
Mem. Soc. Ci. Nat. La Salle		<i>Memoria de la Sociedad de Ciencias Naturales La Salle</i>
Mem. Soc. Cub. Hist. Nat. «Felipe Poey»		<i>Memorias de la Sociedad Cubana de Historia Natural «Felipe Poey»</i>
Mém. Soc. Hist. Nat. Paris		<i>Mémoires de la Société d'Histoire Naturelle de Paris</i>
Mém. Soc. Phys. Genève		<i>Mémoires de la Société de Physique et d'Histoire Naturelle de Genève</i>
Mem. Torrey Bot. Club		<i>Memoirs of the Torrey Botanical Club</i>
Mem. Wern. Nat. Hist. Soc.		<i>Memoirs of the Wernerian Natural History Society</i>
Mimos.		<i>Mimoses</i>
Misc. Bot. [Bertol.]	Bertoloni	<i>Miscellanea Botanica</i>
Mitt. Bot. Vereins Kreis Freiburg		<i>Mitteilungen des Botanischen Vereins für den Kreis Freiburg und das Land Baden</i>
Mitt. Inst. Allg. Bot. Hamburg		<i>Mitteilungen aus dem Institut für Allgemeine Botanik in Hamburg</i>
Molec. Phylogen. Evol.		<i>Molecular Phylogenetics and Evolution</i>
Monog. Bux.		<i>Monographie des Buxacees et des Stylocérées</i>
Monogr. Anonac.		<i>Monographie de la famille des Anonacées</i>
Monogr. Melast.		<i>Monographia Melastomacearum</i>
Monogr. Paullinia		<i>Monographie der Sapindaceengattung Paullinia</i>
Monogr. Phan. [A.DC. & C.DC.]	Candolle & Candolle	<i>Monographiae Phanerogamarum</i>
Monogr. Syst. Bot. Missouri Bot. Gard.		<i>Monographs in Systematic Botany from the Missouri Botanical Garden</i>
Mus. Bot.		<i>Museum Botanicum</i>
N. Amer. Fl.		<i>North American Flora</i>
N. Amer. Sylv.		<i>The North American sylva</i>
N. Amer. Trees		<i>North American Trees</i>
Nachtr. Vollst. Lex. Gärtn.		<i>Nachtrag zum Vollständigen Lexicon der Gärtnerei und Botanik</i>
Narr. Exped. Zaire		<i>Narrative of an expedition to explore the river Zaire, usually called the Congo, in South Africa, in 1816</i>
Nat. Appl. Sci. Bull. Univ. Philipp.		<i>Natural and Applied Science Bulletin</i>
Nat. Hist. Misc.		<i>Natural History Miscellanea</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Nat. Pflanzenfam. [Engler & Prantl]	Engler & Prantl	<i>Die Natürlichen Pflanzenfamilien</i>
Nat. Pflanzenfam. [Engler & Prantl] Nachtr.	Engler & Prantl	<i>Die Natürlichen Pflanzenfamilien Nachträge</i>
Natl. Hort. Mag.		<i>National Horticultural Magazine</i>
Natuurk. Verh. Wet. Haarlem		<i>Natuurkundige verhandelingen van de Hollandsche Maatschappij der Wetenschappen te Haarlem</i>
Natuurw. Stud. Suriname & Curacao		<i>Uitgaven Natuurwetenschappelijke Studiekring voor Suriname en de Nederlandse Antillen</i>
Ned. Kruidk. Arch.		<i>Nederlandsch Kruidkundig Archief</i>
Neue Denkschr. Schw. Naturf. Ges.		<i>Neue Denkschriften der Allgemeinen Schweizerischen Gesellschaft für die Gesammten Naturwissenschaften</i>
Neue Entdeck. Pflanzenk.		<i>Neue Entdeckungen im Ganzen Umfang der Pflanzenkunde</i>
Neue Schriften Ges. Naturf. Freunde Berlin		<i>Der Gesellsschaft Naturforschender Freunde zu Berlin</i>
New Fl. [Rafinesque]	Rafinesque	<i>New Flora and Botany of North America</i>
Nomencl. Bot. [Dennstedt]	Dennstedt	<i>Nomenclator Botanicus</i>
Nomencl. Bot. [Raeusch.]	Raeuschel	<i>Nomenclator Botanicus</i>
Nomencl. Bot. [Steudel]	Steudel	<i>Nomenclator Botanicus</i>
Nordic J. Bot.		<i>Nordic Journal of Botany</i>
Not. Pl. Asiat.		<i>Notulae ad Plantas Asiaticas</i>
Notas Fl. Colombia		<i>Notas a la Flora de Colombia</i>
Notes Bot. Amaz.		<i>Notes of a Botanist on the Amazon & Andes</i>
Notes Bot. Sapot.		<i>Notes botaniques. Sapotacées</i>
Notes Roy. Bot. Gard. Edinburgh		<i>Notes from the Royal Botanic Garden</i>
Notizbl. Bot. Gart. Berlin-Dahlem		<i>Notizblatt des Botanischen Gartens und Museums zu Berlin-Dahlem</i>
Notizbl. Königl. Bot. Gart. Berlin		<i>Notizblatt des Königlicher Botanischen Gartens und Museums zu Dahlem bei Steglitz (Berlin)</i>
Notul. Syst. (Paris)		<i>Notulae Systematae</i>
Nouv. Ann. Mus. Hist. Nat.		<i>Nouvelles Annales du Muséum d'Histoire Naturelle</i>
Nouv. Bull. Sci. Soc. Philom. Paris		<i>Nouveau Bulletin des Sciences, publié par la Société Philomatique de Paris</i>
Nov. Gen. Sp. [H.B.K.]	Humboldt, Bonpland & Kunth	<i>Nova genera et species plantarum</i>
Nov. Gen. Sp. Pl. [Martius]	Martius	<i>Nova genera et species plantarum</i>
Nov. Gen. Sp. Pl. [Poeppig & Endlicher]	Poeppig & Endlicher	<i>Nova genera ac species plantarum</i>
Nov. Pl. Sp.		<i>Novae plantarum species praeserti Indiae Orientalis</i>
Nov. Stirp. Bras.		<i>Novarum stirpium : Brasiliensium decades</i>
Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.		<i>Nova Acta Academiae Caesareae Leopoldino-Carolinae Germanicae Naturae Curiosorum</i>
Nova Acta Phys.-Med. Acad. Caes. Leop.- Carol. Nat. Cur.		<i>Nova Acta Physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum</i>
Nuevo Cat. Fl. Vasc. Venezuela		<i>Nuevo Catálogo de la Flora Vascular de Venezuela</i>
Numer. List		<i>A numerical list of dried specimens of plants, in the East India Companys Museum collected under the superintendence of Dr. Wallich of the Company's botanic garden at Calcutta</i>
Nuovi Ann. Sci. Nat.	Jacquin	<i>Nuovi annali delle scienze naturali</i>
Observ. Bot. [Jacquin]	Jacquin	<i>Observationum botanicarum</i>
Observ. Bot. [Swartz]	Swartz	<i>Observationes botanicae</i>
Observ. Bot. [Van Heurck]	Van Heurck	<i>Observationes botanicae</i>
Occas. Pap. Bernice Pauahi Bishop Mus.		<i>Occasional Papers of the Bernice Pauahi Bishop Museum of Polynesian Ethology and Natural History</i>
Opusc. Neotrop.	Kerchove	<i>Opuscula Neotropicana</i>
Palm. Mattogross.		<i>Palmae Mattogrossenses</i>
Palmiers [Kerchove]	Kerchove	<i>Les palmiers</i>
Palms Amazon		<i>Palms of the Amazon</i>
Pflanzenr. [Engler]	Engler	<i>Das Pflanzenreich</i>
Pflanzenw. Ost-Afrikas		<i>Die Pflanzenwelt Ost-Afrikas</i>
Pharm. J. Trans.		<i>The Pharmaceutical Journal and Transactions</i>
Physis (Buenos Aires)		<i>Physis : Revista de la Sociedad Argentina de Ciencias Naturales</i>
Piperac. N. South Amer.		<i>The Piperaceae of Northern South America</i>
Pl. Aequinoct.		<i>Plantae Aequinoctiales</i>
Pl. Bequaert.		<i>Plantae Bequaertiae</i>
Pl. Bras.		<i>Plantarum Brasiliensium</i>
Pl. Bras. Icon. Descr.		<i>Plantarum Brasiliæ icones et descriptiones hactenus ineditæ</i>
Pl. Columb. [Linden]	Linden	<i>Plantæ Columbianæ</i>
Pl. Guian. [Rudge]	Rudge	<i>Plantarum Guianæ rariorum icones et descriptiones</i>
Pl. Hartw. [Bentham]	Bentham	<i>Plantas Hartwegianas</i>
Pl. Jard. Rio de Janeiro		<i>Plantas novas cultivadas no Jardim Botanico do Rio de Janeiro</i>
Pl. Jav. Rar. [Bennett]	Bennett	<i>Plantae Javanicae Rariores</i>
Pl. Nam.		<i>Plant names</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Pl. Nouv. Amér.		Plantes nouvelles d'Amérique
Pl. Nouv. Découv.		Plantes nouvellement découvertes récemment dénommées et classées
Pl. Nov. Hisp.		Plantae Novae Hispaniae
Pl. Rar. Hort. Schoenbr.		Plantarum rariorum horti caesarei Schoenbrunnensis descriptiones et icones
Pl. Surin.		Plantæ Surinamenses
Pl. Syst. Evol.		Plant Systematics and Evolution
Pl. Usuel. Bras.		Plantes usuelles des Brésiliens
Pl. Util. Col. Franç.		Les plantes utiles des colonies françaises
Pl. Wright. [Grisebach]	Grisebach	Plantæ Wrightianæ e Cuba orientali
Polygon. Guttif. Lecythid. Surinam		The Polygonaceae, Guttiferae and Lecythidaceae of Surinam
Pomona Coll. J. Econ. Bot.		Pomona College Journal of Economic Botany and Subtropical Horticulture
Prim. Fl. Esseq.		Primitiae Floræ Essequeboensis
P. ir. Rostlin		O Pirozeností Rostlin
Proc. Amer. Acad. Arts		Proceedings of the American Academy of Arts and Sciences
Proc. Biol. Soc. Washington		Proceedings of the Biological Society of Washington
Proc. Calif. Acad. Sci.		Proceedings of the California Academy of Sciences
Proc. Kon. Ned. Akad. Wetensch. C		Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen
Proc. Linn. Soc. New South Wales		Proceedings of the Linnean Society of New South Wales
Proc. Roy. Hort. Soc. London		Proceedings of the Royal Horticultural Society of London
Proc. Roy. Soc. Queensland	Candolle	Proceedings of the Royal Society of Queensland
Prodr. [A. P. de Candolle]		Prodromus systematis regni vegetabilis
Prodr. [Swartz]	Swartz	Prodromus descriptionum vegetabilium
Prodr. Fl. Ind. Orient.		Prodromus llorae peninsulae Indiae Orientalis
Prodr. Fl. Nov. Holland.		Prodromus florae Novae Hollandiae
Prodr. Monogr. Hypéric.		Prodromus d'une monographie de la famille des Hypéricinées
Prodr. Pl. Ind. Occid. [Hamilton]	Hamilton	Prodromus plantarum Indiae Occidentalis
Prodr. Stirp. Chap. Allerton		Prodromus stirpium in horto ad Chapel Allerton vigentium
Prot.-App. Enum. Palm. Nov.		Protesto Appendix ao Enumeratio Palmarum Novarum
Publ. Avulsas Mus. Paraense Emílio Goeldi		Publicações Avulsas do Museu Goeldi
Publ. Carnegie Inst. Wash.		Publications of the Carnegie Institution of Washington
Publ. Field Mus. Nat. Hist., Bot. Ser.		Publications of the Field Museum of Natural History, Botanical Series
Publ. Inst. Nac. Pesq. Amaz., Bot.		Publicações Instituto Nacional de Pesquisas de Amazônia Botânica
Pug. Pl. Amer. Centr.		Pugillus plantarum novarum Americae Centralis et Meridionalis
Rain Forests Golfo Dulce		The Rain Forests of Golfo Dulce
Recueil Trav. Bot. Néerl.		Recueil des Travaux Botaniques Néerlandais
Reis. Br.-Guiana [Ri. Schomburgk]	Schomburgk	Reisen in Britisch-Guiana in den Jahren 1840-1844
Reise Bras. [Spix & Mart.]	Spix & Martius	Reise in Brasilien
Rep. (Annual) Missouri Bot. Gard.		Annual Report of the Missouri Botanical Garden
Repert. Bot. Syst. [Walpers]	Walpers	Repertorium botanices systematicae
Repert. Pharm. [J.A. Buchner]	Buchner	Repertorium für die Pharmacie
Repert. Spec. Nov. Regni Veg.		Repertorium specierum novarum regni vegetabilis
Repert. Spec. Nov. Regni Veg. Beih.		Repertorium specierum novarum regni vegetabilis, Beihefte
Rev. Anacard.		Révision du groupe des Anacardiacées
Rev. Bot. Appl. Agric. Colon.		Revue de botanique appliquée et d'agriculture coloniale
Rev. Bot. Appl. Agric. Trop.		Revue de botanique appliquée et d'agriculture tropicale
Rev. Engenh.		Revista Engenharia
Rev. Hort. (Paris)		Revue Horticole - Journal d'horticulture pratique
Revis. Gen. Pl.		Revisio generum plantarum
Revis. Tabernaemontana		A Revision of Tabernaemontana
Revista Acad. Colomb. Ci. Exact.		Revista de la Academia Colombiana de Ciencias Exactas
Revista Argent. Agron.		Revista Argentina de Agronomía
Revista Argent. Bot.		Revista Argentina de Botánica
Revista Brasil. Biol.		Revista Brasileira de Biología
Revista Braz.		Revista Brasileira
Revista Fac. Agron. (Maracay)		Revista de la Facultad de Agronomía
Revista Inst. Parag.		Revista del Instituto Paraguayo
Revista Soc. Cub. Bot.		Revista de la Sociedad Cubana de Botánica
Revista Sudamer. Bot.		Revista Sudamericana de Botánica
Rutac.		Rutaceae, fragmenta botanica
Sci. Surv. Porto Rico & Virgin Islands		Scientific Survey of Porto Rico and the Virgin Islands
Seg. Contr. Conoc. Arbol. Argent.		Segunda contribución al conocimiento de los árboles de la Argentina
Select. Stirp. Amer. Hist.		Selectarum stirpium americanarum historia
Sert. Palm. Brasil.		Sertum Palmarum Brasiliensium
Sitzungsber. Ges. Naturf. Freunde Berlin		Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin
Sitzungsber. Math.-Phys. Cl. Königl. Bayer.		Sitzungsberichte der Mathematisch-Physikalischen Classe (Klasse) der K. B.
Akad. Wiss. München		Akademie der Wissenschaften zu München
Skr. Naturhist.-Selsk.		Skrivter af Naturhistorie-Selskabet
Smithsonian Contr. Bot.		Smithsonian Contributions to Botany
Smithsonian Misc. Collect.		Smithsonian Miscellaneous Collections

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
Solan. Syn.		<i>Solanorum generumque affinium synopsis</i>
Sp. Pl.		<i>Species Plantarum</i>
Sp. Pl., ed. 4 [Willdenow]	Willdenow	<i>Species Plantarum, edition 4</i>
Stand. Cycl. Hortic.		<i>The Standard Cyclopedia of Horticulture</i>
Stirp. Surinam. Select.		<i>Stirpes Surinamenses selectae</i>
Suppl. Carp.		<i>Supplementum carpologiae</i>
Suppl. Hort. Bot. Hafn.		<i>Supplementum horti botanici Hafniensis</i>
Suppl. Pl.		<i>Supplementum Plantarum</i>
Surinam Exsic. [Weigelt]	Weigelt	<i>Surinam exsiccatae</i>
Sylva Tellur.		<i>Sylva Telluriana</i>
Symb. Antill. [Urban]	Urban	<i>Symbolae Antillanae seu fundamenta florae Indiae occidentalis</i>
Symb. Bot. [C. Presl]	Presl	<i>Symbolae botanicae</i>
Symb. Bot. [Vahl]	Vahl	<i>Symbolae botanicae</i>
Symb. Fl. Argent.		<i>Symbolae ad floram Argentinam</i>
Syn. Pl. [D. Dietrich]	Dietrich	<i>Synopsis plantarum</i>
Syn. Pl. [Persoon]	Persoon	<i>Synopsis plantarum</i>
Syn. Queensl. Fl.		<i>A synopsis of the Queensland flora</i>
Syst. Bot.		<i>Systematic Botany</i>
Syst. Geogr. Pl.		<i>Systematics and Geography of Plants</i>
Syst. Laur.		<i>Systema Laurinarum</i>
Syst. Mat. Med. Veg. Bras.		<i>Systema materiae medicae vegetabilis Brasiliensis</i>
Syst. Nat. [A. P. de Candolle]	Candolle	<i>Regni vegetabilis systema naturale</i>
Syst. Piperac. [F.A.W. Miquel]	Miquel	<i>Systema Piperacearum</i>
Syst. Veg. [Roemer & Schultes]	Roemer & Schultes	<i>Systema vegetabilium</i>
Syst. Veg. [Sprengel]	Sprengel	<i>Systema vegetabilium</i>
Syst. Veg. Fl. Peruv. Chil.		<i>Systema vegetabilium florae Peruviana et Chilensis</i>
Tabl. École Bot.		<i>Tableau de l'Ecole de Botanique du Jardin du Roi</i>
Tabl. Encycl.		<i>Tableau Encyclopédique et Méthodique des trois Règnes de la Nature</i>
Tent. Suppl.		<i>Tentamen supplementi ad systematis vegetabilium</i>
Third Conf. Interamer. Agric. Caracas		<i>Third Inter-American Conference on Agriculture Caracas</i>
Tijdschr. Nat. Geschied. Physiol.		<i>Tijdschrift voor natuurlijke geschiedenis en physiologie</i>
Trab. Comm. Sci. Expl., Bot., Rio de Janeiro		<i>Trabalhos da Comissão Científica de Exploração</i>
Trab. Mus. Comercial Venezuela		<i>Trabajos del Museo Comercial de Venezuela</i>
Trab. Mus. Nac. Ci. Nat., Ser. Bot.		<i>Trabajos del Museo Nacional de Ciencias Naturales y Jardín Botánico, Serie Botánica</i>
Trab. Soc. Velllos.		<i>Trabalhos da Sociedade Vellosoiana</i>
Traité Arbr. Arbust.		<i>Traité des arbres et arbustes que l'on cultive en France</i>
Traité Bot. Méd. Phan.		<i>Traité de botanique médicale phanérogamique</i>
Trans. & Proc. Bot. Soc. Edinburgh		<i>Transactions and Proceedings of the Botanical Society of Edinburgh</i>
Trans. Cambridge Philos. Soc.		<i>Transactions of the Cambridge Philosophical Society</i>
Trans. Hort. Soc. London		<i>Transactions of the Horticultural Society of London</i>
Trans. Linn. Soc. London		<i>Transactions of the Linnean Society of London</i>
Trans. Linn. Soc. London, Bot.		<i>Transactions of the Linnean Society of London, 2nd series: Botany</i>
Trans. Soc. London Encour. Arts		<i>Transactions of the Society (London) for the Encouragement of Arts</i>
Trat. Elem. Bot.		<i>Tratado elemental de botánica</i>
Trav. Brazil [H. Koster]	Koster	<i>Travels in Brazil</i>
Trop. Woods		<i>Tropical Woods</i>
U.S.D.A. Div. Bot. Bull.		<i>U.S. Department of Agriculture. Division of Botany Bulletin</i>
Uebersicht Theobroma		<i>Übersicht der bis jetzt bekannten Arten von Theobroma</i>
Univ. Calif. Publ. Bot.		<i>University of California Publications in Botany</i>
Veg. Kingd.		<i>The Vegetable Kingdom</i>
Vég. Ut. Afr. Trop. Franç.		<i>Les végétaux utiles de l'Afrique tropicale française</i>
Vegetationsbilder [G. Karsten & H. Schenck]	Karsten & Schenck	<i>Vegetationsbilder</i>
Verh. Batav. Genootsch. Kunst.		<i>Verhandelingen van het Bataviaansch Genootschap van Kunsten en Wetenschappen</i>
Verh. Bot. Vereins Prov. Brandenburg		<i>Verhandlungen des Botanischen Vereins für die Provinz Brandenburg</i>
Verh. Kon. Ned. Akad. Wetensch., Afd.		<i>Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen</i>
Natuurk.		
Verslagen Meded. Afd. Natuurk. Kon. Akad.		<i>Verslagen en Mededeelingen van de Afdeeling Natuurkunde</i>
Wetensch.		
Verz. Landwirtsch. Gartn. Kulturpfl.		<i>Verzeichnis Landwirtschaftlicher und Gärtnerischer Kulturpflanzen</i>
Vidensk. Meddel. Dansk Naturhist. Foren.		<i>Videnskabelige meddelelser fra den Naturhistoriske forening i Kjøbenhavn</i>
Kjøbenhavn		
Voy. Amér. Mér.		<i>Voyage dans l'Amérique Méridionale</i>
Voy. Uranie, Bot.		<i>Voyage autour du monde</i>
Wiss. Erg. Deut. Zentr.-Afr. Exped., Bot.		<i>Wissenschaftliche Ergebnisse der Deutschen Zentral-Afrika-Expedition 1907-1908</i>

Appendix 5. — Continuation.

Titre_abr	[Auteurs]	Titre
<i>Wiss. Z. Friedrich-Schiller-Univ. Jena, Math.-Naturwiss. Reihe (Beitr. Phytotax., 6)</i> <i>World Checkl. & Bibliogr. Araliaceae</i>		<i>Wissenschaftliche Zeitschrift der Friedrich-Schiller-Universität Jena</i> <i>World Checklist and Bibliography of Araliaceae</i>

APPENDIX 6. — List of the 143 unnamed species/Liste des 143 espèces non nommées.

<i>Abarema</i> sp. A.....	[737]	<i>Helicostylis</i> sp. A.....	[1186]
<i>Aniba</i> sp. A.....	[587]	<i>Hymenaea</i> sp. A	[807]
<i>Aniba</i> sp. B.....	[588]	<i>Ilex</i> sp. A	[135]
<i>Aniba</i> sp. C	[589]	<i>Ilex</i> sp. B.....	[136]
<i>Apeiba</i> sp. A.....	[1003]	<i>Inga</i> sp. A	[868]
<i>Aspidosperma</i> sp. A.....	[105]	<i>Inga</i> sp. B	[869]
<i>Bixa</i> sp. A.....	[172]	<i>Inga</i> sp. C.....	[870]
<i>Calliandra</i> sp. A.....	[756]	<i>Inga</i> sp. D.....	[871]
<i>Caraipa</i> sp. A.....	[229]	<i>Lauraceae</i> sp. A	[674]
<i>Caraipa</i> sp. B.....	[230]	<i>Lauraceae</i> sp. B	[675]
<i>Casearia</i> sp. A.....	[1542]	<i>Lauraceae</i> sp. C.....	[676]
<i>Casearia</i> sp. B	[1543]	<i>Lauraceae</i> sp. D	[677]
<i>Casearia</i> sp. C.....	[1544]	<i>Lauraceae</i> sp. E	[678]
<i>Casearia</i> sp. D	[1545]	<i>Lauraceae</i> sp. F	[679]
<i>Catostemma</i> sp. A.....	[1007]	<i>Lauraceae</i> sp. G	[680]
<i>Chaetocarpus</i> sp. A.....	[1382]	<i>Lauraceae</i> sp. H	[681]
<i>Chrysophyllum</i> sp. A.....	[1609]	<i>Lauraceae</i> sp. I	[682]
<i>Chrysophyllum</i> sp. B.....	[1610]	<i>Lauraceae</i> sp. J	[683]
<i>Chrysophyllum</i> sp. C	[1611]	<i>Licania</i> sp. A.....	[348]
<i>Chrysophyllum</i> sp. D	[1612]	<i>Licania</i> sp. B.....	[349]
<i>Chrysophyllum</i> sp. E.....	[1613]	<i>Licania</i> sp. C	[350]
<i>Chrysophyllum</i> sp. F	[1614]	<i>Maquira</i> sp. A.....	[1190]
<i>Copaifera</i> sp. A	[767]	<i>Maytenus</i> sp. A.....	[251]
<i>Couepia</i> sp. A.....	[275]	<i>Maytenus</i> sp. B.....	[252]
<i>Croton</i> sp. A.....	[511]	<i>Meliosma</i> sp. A.....	[1524]
<i>Cybianthus</i> sp. A.....	[1416]	<i>Micropolis</i> sp. A	[1638]
<i>Dacryodes</i> sp. A.....	[180]	<i>Moquilea</i> sp. A.....	[353]
<i>Diospyros</i> sp. A.....	[448]	<i>Myrcia</i> sp. A	[1312]
<i>Elaeoluma</i> sp. A.....	[1620]	<i>Myrcia</i> sp. B	[1313]
<i>Endlicheria</i> sp. A.....	[601]	<i>Myrtaceae</i> sp. A	[1320]
<i>Enterolobium</i> sp. A.....	[795]	<i>Myrtaceae</i> sp. B	[1321]
<i>Eriotheca</i> sp. A.....	[1013]	<i>Myrtaceae</i> sp. C	[1322]
<i>Eschweilera</i> sp. A.....	[713]	<i>Nectandra</i> sp. A	[622]
<i>Eschweilera</i> sp. B.....	[714]	<i>Neea</i> sp. A	[1332]
<i>Eugenia</i> sp. A.....	[1259]	<i>Neea</i> sp. B	[1333]
<i>Eugenia</i> sp. B	[1260]	<i>Neea</i> sp. C	[1334]
<i>Eugenia</i> sp. C.....	[1261]	<i>Nyctaginaceae</i> sp. A	[1335]
<i>Eugenia</i> sp. D	[1262]	<i>Nyctaginaceae</i> sp. B	[1336]
<i>Eugenia</i> sp. E	[1263]	<i>Nyctaginaceae</i> sp. C	[1337]
<i>Eugenia</i> sp. F	[1264]	<i>Nyctaginaceae</i> sp. D	[1338]
<i>Eugenia</i> sp. G	[1265]	<i>Ocotea</i> sp. A.....	[653]
<i>Eugenia</i> sp. H	[1266]	<i>Ocotea</i> sp. B	[654]
<i>Eugenia</i> sp. I.....	[1267]	<i>Ocotea</i> sp. C	[655]
<i>Eugenia</i> sp. J	[1268]	<i>Ocotea</i> sp. D	[656]
<i>Eugenia</i> sp. K.....	[1269]	<i>Ocotea</i> sp. E	[657]
<i>Eugenia</i> sp. L	[1270]	<i>Ocotea</i> sp. F	[658]
<i>Eugenia</i> sp. M.....	[1271]	<i>Ocotea</i> sp. G	[659]
<i>Eugenia</i> sp. N	[1272]	<i>Ocotea</i> sp. H	[660]
<i>Eugenia</i> sp. O	[1273]	<i>Ocotea</i> sp. I	[661]
<i>Eugenia</i> sp. P	[1274]	<i>Ocotea</i> sp. J	[662]
<i>Eugenia</i> sp. Q	[1275]	<i>Ormosia</i> sp. A	[899]
<i>Eugenia</i> sp. R	[1276]	<i>Ormosia</i> sp. B	[900]
<i>Faramea</i> sp. A.....	[1466]	<i>Ormosia</i> sp. C	[901]
<i>Ferdinandusa</i> sp. A.....	[1469]	<i>Parinari</i> sp. A	[359]
<i>Guapira</i> sp. A	[1326]	<i>Parkia</i> sp. A.....	[917]

Appendix 6. — Continuation.

<i>Peltogyne</i> sp. A	[921]	<i>Raputia</i> sp. A	[1513]
<i>Persea</i> sp. A	[664]	<i>Ruptiliocarpon</i> sp. A	[981]
<i>Poulsenia</i> sp. A	[1194]	<i>Sloanea</i> sp. A	[475]
<i>Pourouma</i> sp. A	[1759]	<i>Sloanea</i> sp. B	[476]
<i>Pourouma</i> sp. B	[1760]	<i>Sloanea</i> sp. C	[477]
<i>Pouteria</i> sp. A	[1691]	<i>Sloanea</i> sp. D	[478]
<i>Pouteria</i> sp. A	[1705]	<i>Sloanea</i> sp. E	[479]
<i>Pouteria</i> sp. B	[1692]	<i>Sloanea</i> sp. F	[480]
<i>Pouteria</i> sp. C	[1693]	<i>Sloanea</i> sp. G	[481]
<i>Pouteria</i> sp. D	[1694]	<i>Sloanea</i> sp. H	[482]
<i>Pouteria</i> sp. E	[1695]	<i>Stryphnodendron</i> sp. A	[940]
<i>Pouteria</i> sp. F	[1696]	<i>Sympmania</i> sp. A	[381]
<i>Pouteria</i> sp. G	[1697]	<i>Tachigali</i> sp. A	[966]
<i>Pouteria</i> sp. H	[1698]	<i>Taralea</i> sp. A	[968]
<i>Pouteria</i> sp. I	[1699]	<i>Tovomita</i> sp. A	[397]
<i>Pseudoxandra</i> sp. A	[75]	<i>Trichilia</i> sp. A	[1140]
<i>Quiina</i> sp. A	[1365]		