

# First record of *Cocculinella* (Mollusca, Gastropoda, Cocculiniformia) from the Lower Pleistocene of Southern Italy with the description of two new species

Bruno DELL'ANGELO

Via Santelia 55/12A, I-16153 Genova (Italy)  
bruno.dellangelo@chitons.it

Maurizio SOSSO

Via Bengasi 4/4, I-16153 Genova (Italy)  
maurizio.soso@tele2.it

Antonio BONFITTO

Dipartimento di Biologia evoluzionistica e sperimentale,  
Via Selmi 3, I-40126 Bologna (Italy)  
antonio.bonfitto@unibo.it

---

Dell'Angelo B., Sosso M. & Bonfitto A. 2011. — First record of *Cocculinella* (Mollusca, Gastropoda, Cocculiniformia) from the Lower Pleistocene of Southern Italy with the description of two new species. *Geodiversitas* 33 (4): 739-746. DOI: 10.5252/g2011n4a9.

## ABSTRACT

*Cocculinella freti* n. sp., from Archi and Valanidi (Reggio Calabria) and *Cocculinella bertolasoi* n. sp., from Valanidi (Reggio Calabria) and Capo Milazzo (Messina, Sicily), are described from three southern Italian Lower Pleistocene outcrops (bathyal assemblages). No species of the genus *Cocculinella* Thiele, 1909 is known from the Atlantic and the Mediterranean, whereas three living species are known from the Indo-Pacific. Very little, mostly questionable information is known on fossil species of the genus. The present report is the first unequivocal fossil record of *Cocculinella*, allowing the stratigraphic and biogeographic range of the genus to be extended to the Lower Pleistocene of the Mediterranean.

## KEY WORDS

Mollusca,  
Gastropoda,  
Cocculiniformia,  
Cocculinellidae,  
Pleistocene,  
Mediterranean basin,  
Italy,  
new species.

## RÉSUMÉ

Première occurrence de *Cocculinella* (Mollusca, Gastropoda, Cocculiniformia) du Pléistocene inférieur du sud de l'Italie, avec la description de deux nouvelles espèces.

*Cocculinella freti* n. sp. et *Cocculinella bertolasoi* n. sp. sont décrites de trois affleurements du Pléistocène inférieur (biocénoses bathyale) du sud de l'Italie.

**MOTS CLÉS**  
 Mollusca,  
 Gastropoda,  
*Cocculiniformia*,  
*Cocculinellidae*,  
 Pléistocene,  
 bassin Méditerranéen,  
 Italie,  
 espèces nouvelles.

La première espèce a été trouvée dans des échantillons d'Archi et Valanidi (province de Reggio Calabria), la deuxième dans des échantillons de Valanidi et de Capo Milazzo (province de Messina, Sicile). Le genre *Cocculinella* Thiele, 1909 n'est pas connu dans les faunes modernes de la mer Méditerranée et de l'océan Atlantique, alors que trois espèces vivantes sont signalées dans les océans Indien et Pacifique. Dans le même temps, les informations sur les espèces fossiles sont rares et incertaines. Les deux espèces nouvelles représentent donc la première occurrence de ce genre dans les faunes fossiles et nous permettent d'étendre sa présence au Pléistocène inférieur du sud de l'Italie.

## INTRODUCTION

The *Cocculinellidae* Moskalev, 1971 are a family of symmetrical limpets, with a narrowly elliptical aperture and subcentral apex (Haszprunar 1998). The radula is highly modified and highly diagnostic (Moskalev 1976; Marshall 1983, 1985).

A single genus, *Cocculinella* Thiele, 1909, is known for the family, including three species: *Cocculinella coercita* Hedley, 1907, from Australia, *C. osteophila* Marshall, 1983, from New Zealand and *C. minutissima* (Smith, 1904), type species of the genus, from the Andaman Islands (Indian Ocean) (Marshall 1983, 1985; Haszprunar 1988, 1998). Another species, *C. kopua* Marshall, 1983, described from a single dead shell from New Zealand, was re-considered, by Marshall (1985), as a possible synonym of *C. osteophila*. Two other Australian species, *C. tasmanica* (May, 1919) and *C. mayi* Finlay, 1926 are recorded in some web-databases, such as "A Guide to the Marine Molluscs of Tasmania" (<http://www.molluscssoftasmania.net> [2010]) or OBIS (Ocean Biogeographic Information System); these species are not considered belonging to *Cocculinella* by Haszprunar (1998), and are referred to *Tectisumen* Finlay, 1926, a genus attributed to family Lepetellidae Dall, 1882 by other authors (e.g., Keen 1960; Moskalev 1978).

All cocculinellid species live and feed on fish bones.

This paper focuses on the finding of four fossil shells of *Cocculinella* in three Lower Pleistocene out-

crops in southern Italy, attributed to two different species, described as new. No living species of this genus are known in the Mediterranean Sea and in the Atlantic Ocean, and there is very little information on fossil species, and so the description of these new species is particularly important for the knowledge of the stratigraphic and biogeographic distribution of the genus.

## MATERIAL AND METHODS

The shells were found by Stefano Palazzi in two Lower Pleistocene outcrops in Southern Italy, Archi (Reggio Calabria, 1 shell) and Valanidi (Reggio Calabria, 2 specimens), and by Luca Bertolaso at Capo Milazzo, Sicily (Fig. 1).

Archi is a clay quarry, locally known as "Fornace Aloï", situated in the northern suburbs of Reggio Calabria, with a rich lower Pleistocene fauna, that can be attributed to a bathyal paleoenvironment, 500 to 1000 m deep (Guadagno *et al.* 1979; Di Geronimo *et al.* 1997). The molluscs have been listed by Micali & Villari (1986), Rindone & Vazzana (1989) and Di Geronimo *et al.* (1997).

Valanidi is a clay quarry, near Bovetto (Reggio Calabria), with a rich lower Pleistocene fauna, that can be attributed to a bathyal biocoenoses. This outcrop has scarcely been reported in the literature (Seguenza 1879; Barrier 1987), and there is no published list of the molluscs occurring there, apart from isolated reports of some rare species (e.g., Vazzana 1996).




FIG. 1. – Location of the outcrops in Southern Italy, where *Cocculinella* Thiele, 1909 species were found.

The Neogene sedimentary sequence at Capo Milazzo (Messina, NE Sicily) is well known (Ruggieri & Greco 1965; Fois 1990). The *Cocculinella* shell was found in bathyal deposits of Lower Pleistocene age outcropping at Punta Lazzi, which can be attributed to a bathyal biocoenoses (Palazzi & Villari 1996). There have been several papers on Capo Milazzo molluscs (Gignoux 1913; Mars 1956), and the Plio-Pleistocene bathyal molluscan assemblages are well documented (Palazzi & Villari 1996).

SEM micrographs were taken using a JEOL model JSM-5200.

#### ABBREVIATIONS

AL	anterior length (distance from the posterior extremity of protoconch to the anterior end of teleoconch parallel to apertural plain);
H	shell height;

L shell length;  
W shell width.

#### Repository

BD	Bruno Dell'Angelo collection, Genova, Italy;
MZB	Museo di Zoologia dell'Università di Bologna, Italy.

#### SYSTEMATICS

##### Superorder COCCULINIFORMIA

##### Superfamily LEPETELLOIDEA

##### Family COCCULINELLIDAE Moskalev, 1971

##### Genus *Cocculinella* Thiele, 1909

TYPE SPECIES. — *Acmaea minutissima* Smith, 1904; Recent, Andaman Islands, Indian Ocean, 238–457 m.

TABLE 1. — Measurements (in mm) of *Cocculinella freti* n. sp. shells.

<b><i>Cocculinella freti</i> n. sp.</b>	<b>L</b>	<b>W</b>	<b>H</b>	<b>AL</b>	<b>L/W</b>	<b>L/H</b>	<b>L/AL</b>	<b>W/H</b>
Holotype (Archi)	3.03	1.04	1.38	1.77	2.91	2.20	1.71	0.75
Paratype (Valanidi)	5.21	1.39	2.41	2.68	3.75	2.16	1.94	0.58

TABLE 2. — Measurements (in mm) of *Cocculinella bertolasoi* n. sp. shells.

<b><i>Cocculinella bertolasoi</i> n. sp.</b>	<b>L</b>	<b>W</b>	<b>H</b>	<b>AL</b>	<b>L/W</b>	<b>L/H</b>	<b>L/AL</b>	<b>W/H</b>
Holotype (Valanidi)	3.87	1.62	1.92	1.95	2.39	2.02	1.98	0.84
Paratype (Capo Milazzo)	2.03	1	1.19	1.22	2.03	1.71	1.66	0.84

*Cocculinella freti* n. sp.  
(Fig. 2A-F)

TYPE MATERIAL. — Holotype: MZB 45691, shell from Archi (Fig. 2A, B, D-F). Paratype: BD, shell from Valanidi (Fig. 2C).

ETYMOLOGY. — From the latin word “*fretum*”, “strait” (of Messina), alluding to the occurrence at two outcrops near the Calabrian coast of the Messina Strait.

TYPE LOCALITY AND HORIZON. — Archi, Reggio Calabria Province, Southern Italy, Lower Pleistocene.

MEASUREMENTS. — see Table 1.

DESCRIPTION

Shell patelliform, up to 5.21 mm long (3.03 mm the holotype), thin and fragile. Aperture narrowly elliptical, ends evenly rounded, sides subparallel and very shallowly rounded, slightly convex from side to side so that only the center contacts a flat surface. Spire depressed-conical, apex slightly behind the centre, anterior slope profile slightly convex, posterior slope profile shallowly concave, anterior and posterior slope areas regularly connected with lateral areas, posterior outline sub-angular. Protoconch clearly demarcated, 285 µm long and 188 µm wide in the holotype, of about one whorl, tip of apical fold perfectly fused behind the apertural rim. Teleoconch with only fine collateral growth lines.

REMARKS

The paratype from Valanidi has the same narrow and elongate general shape as the holotype. The

protoconch is partially eroded, so it was possible only to measure the length, 310 µm, which is closely accordant with that of the holotype (285 µm).

DISTRIBUTION

*Cocculinella freti* n. sp. is only known from the lower Pleistocene of Archi and Valanidi, Reggio Calabria Province, Southern Italy.

*Cocculinella bertolasoi* n. sp.  
(Fig. 2G-L)

TYPE MATERIAL. — Holotype: MZB 45692, shell from Valanidi (Figs 2G, H, J-L). Paratype: BD, shell from Capo Milazzo (Fig. 2I).

ETYMOLOGY. — The specific name honours our friend Luca Bertolaso (Reggio Emilia), who collected part of the material here presented and made it available to the authors.

TYPE LOCALITY. — Valanidi, Reggio Calabria Province, Southern Italy.

TYPE STAGE. — Lower Pleistocene.

MEASUREMENTS. — see Table 2.

DESCRIPTION

Shell patelliform, up to 3.87 mm long (the holotype), thin and fragile. Aperture narrowly elliptical, ends evenly rounded, sides subparallel and very shallowly rounded, slightly convex from side to side so that only the center contacts a flat surface. Spire depressed-conical, apex slightly behind the




FIG. 2. — Lower Pleistocene *Cocculinella* Thiele, 1909 species: A-F, *Cocculinella freti* n. sp.; A, B, D-F, holotype (MZB 45691) from Archi (Reggio Calabria Province) in dorsal (A) and lateral (B) views; D, E, protoconch; F, teleoconch; C, paratype from Valanidi (Reggio Calabria Province), dorsal view, not coated; G-L, *Cocculinella bertolstoi* n. sp.; G, H, J-L, holotype (MZB 45692), from Valanidi (Reggio Calabria Province) in dorsal (G) and lateral (H) views; J, K, protoconch; L, teleoconch; I, paratype from Capo Milazzo (Messina Province, Sicily), not coated. Scale bars: A, B, I, 500 µm; C, G, H, 1 mm; D-F, J-L, 100 µm.

centre, anterior slope profile slightly convex, posterior slope profile practically straight or shallowly concave, anterior and posterior slope areas regularly connected with lateral areas, posterior outline regularly rounded. Protoconch clearly demarcated, 335 µm long and 220 µm wide in the holotype, of about one whorl, tip of apical fold perfectly fused behind the apertural rim. Teleoconch with only fine collabral growth lines.

#### REMARKS

The paratype from Capo Milazzo has the same general shape as the holotype. *Cocculinella bertolstoi* n. sp.

differs from *C. freti* n. sp. in general shape, more narrow and elongate in *C. freti* n. sp. (L/W 2.91-3.75 in *C. freti* n. sp. vs 2.03-2.39 in *C. bertolstoi* n. sp.; W/H 0.58-0.75 in *C. freti* n. sp. vs 0.84 in *C. bertolstoi* n. sp.) and in having a more globose and larger protoconch (285 × 188 µm in *C. freti* n. sp. vs 335 × 220 µm in *C. bertolstoi* n. sp.).

#### DISTRIBUTION

*Cocculinella bertolstoi* n. sp. is only known from the lower Pleistocene of Valanidi (Reggio Calabria Province, Calabria) and Capo Milazzo (Messina Province, Sicily), Southern Italy.

TABLE 3. — *Cocculinella* Thiele, 1909 species, published data.

	shells	Protoconch size (µm)	L (mm)	L/W	L/H	L/AL	W/H
<i>C. minutissima</i> (Smith, 1904)	1	—	3.5	2.63	3.5	—	1.33
<i>C. osteophila</i> Marshall, 1983	6	233 × 150	1.13-2.30	1.82-2.33	3.12-4.83	1.41-1.64	1.41-1.72
<i>C. coercita</i> Hedley, 1907	6	—	2.60-6.60	2.20-4.12	2.60-3.77	1.57-1.94	0.80-1.24
<i>C. freti</i> n. sp.	2	285 × 188	3.03-5.21	2.91-3.75	2.16-2.20	1.71-1.94	0.58-0.75
<i>C. bertolasoi</i> n. sp.	2	335 × 220	2.03-3.87	2.03-2.39	1.71-2.02	1.66-1.98	0.84

## DISCUSSION

There are few taxonomically useful shell characters that can be used to separate the species of the family Cocculinellidae. All the known species (1.13-6.6 mm long) have a narrowly elliptical aperture and subcentral apex, and exhibit a weak, concentric sculpture. The protoconch has a fused tip and has no significant sculpture. The radula is greatly modified and highly diagnostic (Haszprunar 1998). The present species are attributed to the genus *Cocculinella* on the basis of the close similarity of the shell to those of the known Recent *Cocculinella* species, *C. minutissima* from the Indian Ocean, and *C. coercita* and *C. osteophila* from the south-western Pacific Ocean. Very few data are available on the variability of these species. All available morphometric data is tabulated here (Table 3).

Two *Cocculinella* species were reported by Vera-Peláez et al. (1995) from the Pliocene of Estepona: *C. cf. minutissima* (Smith, 1904) and *C. compressiuscula* (Karsten, 1849), described from the Miocene of the North Sea basin. These were simply names in a list of molluscs, with no other data or figures, and were not reported by Landau et al. (2003) in their paper on Estepona Vertigastropoda. *Patella compressiuscula* Karsten, 1849 is probably referable to *Lepetella*. B. Landau (pers. com.) has informed us that no species referable to *Cocculinella* has been found in the Estepona Pliocene, so it is better to ignore these reports.

*Cocculinella salisburyensis* Ludbrook, 1956 was described from the Pliocene of South Australia, but the only available specimen (holotype) has a much more strongly posterior apex than any Recent *Cocculinella* and is very doubtfully referable to this genus (Marshall 1983: 142).

The present report is thus the first fossil record of *Cocculinella*, allowing the stratigraphic and biogeographic range of the genus to be extended to the Lower Pleistocene of the Mediterranean.

An Indo-Pacific affinity for some molluscan taxa from the Plio-Pleistocene bathyal deposits of the Mediterranean basin is fairly well known (e.g., La Perna et al. [2004] for the nuculanid genus *Jupiteria* Bellardi, 1875 and *Zealeda* Marwick, 1924; La Perna [2007] for the nuculanid genus *Deminucula* Iredale, 1931). This affinity can be explained by the faunal continuity between the two areas, as part of the Tethys Realm, until the Early Miocene (Popov 1993; Harzhauser et al. 2002, 2007), when the formation of a landbridge between the Eurasia and Arabian plates, in the late Burdigalian (approximately 19 Ma), separated western (Proto Mediterranean Atlantic Region) and eastern (Proto Indo-West-Pacific Region) parts (Rögl 1998, 1999; Harzhauser et al. 2007). However, as regard to the genus *Cocculinella*, the hypothesis of a fragmentation of the area of origin can only be considered assuming that the taxon is much older and with a wider distribution than the one appearing in the fossil record.

Other members of the superorder Cocculiniformia have a fossil distribution not so erratic as suggested by the occurrence of *Cocculinella*; in the genus *Cocculina* Dall, 1882 (family Cocculinidae, superfamily Cocculinoidea), *Cocculina miocaenica* Boettger, 1901 is known from the Paratethys (Romania, Poland) and North Sea basin (The Netherlands) Miocene, *Cocculina adunca* (Jeffreys, 1883) from the Spanish and Italian Early Pliocene, *Cocculina dittmeri* (Anderson, 1964) from German Miocene and *Cocculina unica* Bałuk, 2006 from the Paratethys (Poland) Miocene (Anderson 1964; Janssen 1984; Landau et al. 2003; Bałuk 2006).

## Acknowledgements

We wish to thank Luca Bertolaso (Reggio Emilia, Italy) and Stefano Palazzi (Modena, Italy) for the material studied here, Angelo Vazzana (Reggio Emilia) for information on the Valanidi outcrop, Bernard Landau (Albufeira, Portugal) for his help regarding the supposed fossil *Cocculinella* reports from Spain. Karen Gowlett-Holmes (Tasmania, Australia), Bruce A. Marshall (Museum of New Zealand Te Papa Tongarewa, Wellington, New Zealand), Robert Hamilton-Bruce, Thierry Laperousaz and Ben McHenry (South Australian Museum, Adelaide, Australia), Lisa-ann Gershwin (Queen Victoria Museum and Art Gallery, Tasmania, Australia) helped us with bibliography and information on living Australian and New Zealand species. We thank Rafael La Perna (Università di Bari, Italy), Mathias Harzhauser (Natural History Museum, Vienna) and Annemarie Ohler (Muséum national d'Histoire naturelle, Paris) for their useful comments and constructive criticism on an earlier version of the manuscript.

## REFERENCES

- ANDERSON H. J. 1964. — Die miocene Reinbek-Stufe in Nord- und Westdeutschland und ihre Mollusken-Fauna. *Fortschritte in der Geologie von Rheinland und Westfalen* 14: 31-368.
- BAŁUK W. 2006. — Middle Miocene (Badenian) gastropods from Korytnica, Poland; Part V Addenda et Corrigenda ad Prosobranchia. *Acta Geologica Polonica* 56: 177-220.
- BARRIER P. 1987. — Stratigraphie des dépôts pliocènes et quaternaires du Détriot de Messine, in BARRIER P., DI GERONIMO I. & MONTENANT C. (eds), Le Détriot de Messine (Italie). Évolution tectono-sédimentaire récente (Pliocène et Quaternaire) et environnement actuel. *Documents et travaux de l'Institut Géologique Albert de Lapparent* 11: 59-81.
- DI GERONIMO I., D'ATRI A., LA Perna R., ROSSO A., SANFILIPPO R. & VIOLANTI D. 1997. — The Pleistocene bathyal section of Archi (Southern Italy). *Bollettino della Società Paleontologica Italiana* 36: 189-212.
- FOIS E. 1990. — La successione neogenica di Capo Milazzo (Sicilia Nord-Orientale). *Rivista Italiana di Paleontologia e Stratigrafia* 95: 397-440.
- GIGNOUX M. 1913. — *Les formations marines pliocènes et quaternaires de l'Italie du Sud et de la Sicile*. Thèse, faculté des Sciences de l'Université de Lyon, Lyon: xxiv+694, 21 pls.
- GROVE S. 2010. — *A Guide to the Marine Molluscs of Tasmania*. <http://www.molluscssoftasmania.net/Genus%20pages/Cocculinella.html> (last access 3/07/2010).
- GUADAGNO M., TADDEI RUGGIERO E., DE BLASIO I., PLACELLA B. & SGARRELLA F. 1979. — La sezione pleistocenica di Archi (RC). *Bollettino della Società dei Naturalisti in Napoli* 88: 1-29.
- HARZHAUSER M., PILLER W. E. & STEININGER F. F. 2002. — Circum-Mediterranean Oligo-Miocene biogeographic evolution – the gastropods' point of view. *Palaeogeography, Palaeoclimatology, Palaeoecology* 183: 103-133.
- HARZHAUSER M., KROH A., MANDIC O., PILLER W. E., GÖHLICH U., REUTER M. & BERNING B. 2007. — Biogeographic responses to geodynamics: A key study all around the Oligo-Miocene Tethyan Seaway. *Zoologischer Anzeiger-Journal of Comparative Zoology* 246: 241-256.
- HASZPRUNAR G. 1988. — Anatomy and relationships of the Boone-feeding limpets, *Cocculinella minutissima* (Smith) and *Osteopelta mirabilis* Marshall (Archaeogastropoda). *Journal of Molluscan Studies* 54: 1-20.
- HASZPRUNAR G. 1998. — Superorder Cocculiniformia, in BEESLEY P. L., ROSS G. J. B. & WELLS A. (eds), *Mollusca: The Southern Synthesis. Fauna of Australia*. Vol. 5. CSIRO Publishing, Melbourne, Part B: 653-664.
- JANSSEN A. W. 1984. — *Mollusken uit het Mioceen van Winterswijk-Miste. Een inventarisatie met beschrijvingen en afbeeldingen van alle aangetroffen soorten*. Koninklijke Nederlandse Natuurhistorische Vereniging, Amsterdam, 451 p.
- LA Perna R. 2007. — The deep-water protobranch *Deminucula* (Bivalvia) in the Mediterranean Plio-Pleistocene and the contribution of palaeobiogeography to taxonomy. *Bollettino Malacologico* 43: 3-12.
- LA Perna R., CEREGATO A. & TABANELLI C. 2004. — Mediterranean Pliocene protobranchs: the genera *Jupiteria* Bellardi, 1875, *Ledella* Verrill & Bush, 1897 and *Zealeda* Marwick, 1924 (Mollusca, Bivalvia). *Bollettino Malacologico* 40: 25-36.
- LANDAU B., MARQUET R. & GRIGIS M. 2003. — The Early Pliocene Gastropoda (Mollusca) of Estepona, southern Spain. *Palaeontos* 3: 1-87.
- MARS M. 1956. — Faunes malacologiques du Pliocène et du Quaternaire de Milazzo (Sicile). *Bulletin du Muséum d'Histoire naturelle de Marseille* 16: 33-52.
- MARSHALL B. A. 1983. — The family Cocculinellidae (Mollusca: Gastropoda) in New Zealand. *National Museum of New Zealand Records* 2: 139-143.
- MARSHALL B. A. 1985. — Recent and Tertiary Cocculinidae and Pseudococculinidae (Mollusca: Gastropoda) from New Zealand and New South Wales. *New Zealand Journal of Zoology* 12: 505-546.
- MICALI P. & VILLARI A. 1986. — Riscoperta di *Calliostoma sayanus* (G. Seguenza, 1873) e considerazioni

- sul Pleistocene di Archi (RC). *Bollettino Malacologico* 22: 267-272.
- MOSKALEV L. I. 1976. — [On the generic classification in Cocculinidae (Gastropoda, Prosobranchia)]. *Trudy Instituta Okeanologii, Akademiya Nauk SSSR* 99: 59-70 (in Russian).
- MOSKALEV L. I. 1978. — [Lepetellidae (Gastropoda, Prosobranchia) and related forms]. *Trudy Instituta Okeanologii, Akademiya Nauk SSSR* 113: 132-146 (in Russian).
- OBIS INDO-PACIFIC MOLLUSCAN DATABASE. — <http://clade.ansp.org/obis/> (last access 03/07/2010).
- PALAZZI S. & VILLARI A. 1996. — Malacofaune batiali Plio-Pleistoceniche del Messinese. 2: Capo Milazzo. *Naturalista siciliano* (4) 20: 237-279.
- POPOV S. V. 1993. — Zoogeography of the Late Eocene basins of western Eurasia based on bivalve mollusks. *Stratigraphy and Geological Correlation* 2/6: 103-118.
- RINDONE V. & VAZZANA A. 1989. — Alcune specie di molluschi delle argille batiali del piano Siciliano (Pleistocene inf.) della cava di Archi (Reggio Calabria). *Bollettino Malacologico* 25: 233-240.
- RÖGL F. 1998. — Palaeogeographic considerations for Mediterranean and Paratethys seaways (Oligocene to Miocene). *Annalen des Naturhistorischen Museums in Wien* 99: 279-310.
- RÖGL F. 1999. — Circum-Mediterranean Miocene paleogeography, in RÖSSNER G. & HEISSIG K. (eds), *The Miocene Land Mammals of Europe*. Dr Fritz Pfeil Verlag, Munich: 39-48.
- RUGGIERI G. & GRECO A. 1965. — Studi geologici e paleontologici su Capo Milazzo con particolare riguardo al Milazziano. *Geologica Romana* 4: 41-48, pls 1-11.
- SEGUENZA G. 1879. — Le formazioni terziarie nella provincia di Reggio (Calabria). *Reale Accademia dei Lincei, Memorie della Classe di scienze fisiche, matematiche e naturali* 6: 1-446.
- VAZZANA A. 1996. — Malacofauna batiale del Pleistocene inferiore del Vallone Catrica (Reggio Calabria, Italia). *Bollettino Malacologico* 31: 143-162 (dated 1995, published 1996).
- VERA-PELÁEZ J. L., LOZANO-FRANCISCO M. C., MUÑIZ-SOLÍS R., GILI C., MARTINELL J., DOMÈNECH R., PALMQUIST P. & GUERRA-MERCHÁN A. 1995. — Estudio preliminar de la malacofauna del Plioceno de Estepona (Málaga, España). *Iberus* 13: 93-117.

Submitted on 30 March 2010;  
accepted on 17 December 2010.