

***In situ* foraminifera in grounding zone diamict: a working hypothesis**

PHILIP J. BART¹, LAURA COQUEREAU^{1,2}, SOPHIE WARNY¹ and WOJCIECH MAJEWSKI³

¹*Department of Geology and Geophysics and Museum of Natural Science, E235 Howe-Russell, Louisiana State University, Baton Rouge, LA 70803, USA*

²*Department of Oceanography and Marine Environments, 4 Place Jussieu, Université Pierre et Marie Curie, 75252 Paris Cedex 05, France*

³*Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, 00-818 Warsaw, Poland
pbart@lsu.edu*

Abstract: The ice-proximal diamict sediment deposited on the foreset of a grounding zone wedge in Glomar Challenger Basin on the eastern Ross Sea continental shelf yielded a low abundance assemblage of foraminifera at two piston core sites. We found 302 small well-preserved specimens representing 18 species of benthic foraminifera from 825 ml of sediment. Only three poorly preserved specimens of the planktonic foraminifera *Neogloboquadrina pachyderma* (sinistral) were found. Our combined analyses of preservation state, assemblage composition and stable isotopes suggest that the benthic foraminifera may be *in situ*. This possibility is of interest to palaeoclimatologists who use ice-proximal sediments on the Antarctic continental shelves to radiocarbon date the post-glacial retreat history.

Received 19 May 2015, accepted 26 January 2016

Key words: continental shelf, grounding zone wedge, ice sheet, Last Glacial Maximum, Ross Sea, West Antarctica

Introduction

Geophysical data (i.e. seismic and multibeam) collected on the Antarctic continental shelves show that large-scale glacial troughs were eroded by ice streams when grounded ice advanced to the outer shelf during the Last Glacial Maximum (LGM) (e.g. Jakobsson *et al.* 2011, O'Brien *et al.* 2015). Since the LGM, grounded ice has retreated to the inner continental shelf. During the retreat, some palaeo-ice-stream troughs were partially filled with a backstepping succession of grounding zone wedges (GZWs), i.e. subaqueous terminal moraines (e.g. Graham *et al.* 2010). The GZWs are of interest because they provide unequivocal evidence showing the locations at which the retreat of grounded ice was temporarily interrupted by a pause and slight re-advance.

Piston cores collected on the outer continental shelf always show that ice sheet retreat is recorded by an overall upcore lithologic transition from diamict that is tens of metres thick to a condensed section of diatom ooze (e.g. Domack *et al.* 1999). The diamict was deposited by or in proximity to grounded ice whereas ooze has accumulated in an ice-distal open-marine setting since ice shelf retreat. The challenge has been to find *in situ* material to radiocarbon date the retreat history with all its discrete pauses. The task is complicated because ice-contact and ice-proximal sediments contain much recycled material.

During recent years, denser seismic grids and large-area multibeam surveys (e.g. Graham *et al.* 2010) reveal that

the 3D morphology of GZWs includes eroded topset morphology, which is a surface over which grounded ice streamed. The eroded sediment is transported, in this case northward, to a depositional foreset surface, formed beyond the limit of grounded ice. The GZW foreset received sediment delivered by ice streaming over the topset. The topset and foreset zones are distinct and are clearly separated by an ice sheet grounding line. Megascale glacial lineations (MSGs, i.e. subglacial bedforms) are not found on the GZW foreset surface. Furthermore, given that the GZW foreset probably was a site of rapid sedimentation, long cores through the foreset lessen the possibility that benthic organisms have created a post-depositional bioturbation disturbance possibly reworking younger post-glacial material downward into the underlying marine diamict.

In a recent study of a GZW on the middle continental shelf of the Glomar Challenger Basin palaeo-ice stream trough in eastern Ross Sea, Bart & Cone (2012) emphasized the existence of a GZW topset with MSGs that can be distinguished from a GZW foreset surface. The foreset has tens of metres of relief and is a few kilometres wide (Fig. 1). The existence of an undisturbed GZW foreset opens the possibility that *in situ* foraminifera and other life assemblages inhabited this setting and that the stratigraphic record has not experienced post-depositional disturbance. Video images acquired by the Whillans Ice Stream Subglacial Access Research Drilling (WISSARD) project show that an




Fig. 1a. Global Challenger Basin (GCB), eastern Ross Sea, Antarctica. Dark yellow indicates the drainage basin for the palaeo Whillans Ice Stream when the West Antarctic Ice Sheet was grounded at the middle continental shelf of GCB. Light yellow indicates the modern drainage, where the WISSARD project sampled the grounding zone wedge (GZW) environment. **b.** Multibeam bathymetry of the GZW foreset. **c.** Cross section of the GZW at PC12 and PC5 in grey. White dashed lines show the inferred time-transgressive GZW progradation. M = McMurdo Sound, MSGLs = megascale glacial lineations, WA = West Antarctic.

ecosystem, including fish and invertebrates, exists at the modern marine termination of a GZW. This discovery of surprisingly rich life at this site is remarkable because this unique setting is below a 1000 metre thick floating ice shelf, and is more than 500 km from the nearest open-water environment. The existence of fauna in this environment is of special interest to palaeoclimatologists who use the backstepping succession of ice-proximal deposits and submarine landforms on the Antarctic continental shelf to reconstruct the ice sheet's post-glacial retreat (e.g. Post *et al.* 2014). An *in situ* fossil assemblage in ice-proximal marine sediments could be used to i) reconstruct the changing environmental conditions at the grounding line environment (e.g. Mackensen 2012), and ii) radiocarbon date the various positions occupied by grounded ice during the post-glacial retreat of ice from the continental shelf (e.g. Bart & Cone 2012).

Here, we show that ice-proximal diamict sediment deposited on the foreset of a GZW in the Glomar




Fig. 2. Foraminifera recovered from NBP0802-PC12 and NBP0803-PC5 cores. Scale bar indicates 80 μ m. **a.** *Astrononion echolsi* Kennett, **b.** *Bulimina aculeata* d'Orbigny, **c.** *Cassidulina teretis* Tappan, **d.** *Cibicides* sp., **e.** *Ehrenbergina glabra* Heron-Allen & Earland, **f.** *Epistominella* spp., **g.** *Fissurina* sp., **h.** *Stainforthia concava* (Höglund), **i.** *Globocassidulina subglobosa* (Brady), **j.** *Homalohedra acuticosta* (Reuss), **k.** *Lagena* sp., **l.** *Melonis* sp. (broken), **m.** unidentified Miliolid, **n.** *Nonionella* sp., **o.** *Oolina* spp., **p.** *Oolina* spp., **q.** *Trifarina earlandi* (Parr), **r.** *Textularia* sp. (agglutinated), **s.** *Trochammina* sp. (agglutinated), **t.** *Neogloboquadrina pachyderma* (Ehrenberg) (planktonic).

Challenger Basin palaeo-ice-stream trough of the eastern Ross Sea contains 18 species of well-preserved extant benthic foraminifera. In combination with the ecosystem discovered by the WISSARD team, our results open the possibility that an *in situ* ice-proximal sub-ice shelf assemblage lived on the GZW foreset prior to decoupling retreat of the West Antarctic Ice Sheet (WAIS) and is preserved in marine diamict deposited on the middle continental shelf.

Methods

Our study focused on isolating foraminifera from ice-proximal marine diamict that was deposited at a GZW when the WAIS was grounded on the middle continental shelf within the axis of the Glomar Challenger

Basin palaeo-ice-stream trough in eastern Ross Sea. Two cores collected during an RVIB *Nathaniel B. Palmer* 2008 cruise (NBP0802) were selected based on the morphology of the GZW as interpreted from a seismic grid and large-area multibeam survey of the trough. Both cores penetrated olive-green diatom-rich ooze overlying homogenous grey diamict. PC12 was selected because it penetrates the GZW foreset at a location corresponding to a zone where the last interval of marine diamict deposition occurred prior to the onset of lift-off and retreat of grounded ice (Fig. 1b). PC5 penetrated GZW diamict at an interior location on the eroded scarp of the GZW topset (Fig. 1c).

Fifty-five 15 ml samples of marine diamict and two 15 ml core-top samples of post-glacial diatom ooze were processed individually following the techniques described by Bart & Cone (2012), i.e. before picking, foraminifera were concentrated by floating air-dried > 45 µm residue. This procedure removed potentially reworked specimens with sediment filled tests. Distinction between *in situ* and reworked foraminifera was made on the basis of preservation state (e.g. Majewski & Anderson 2009). Foraminifera tests (i.e. the hard shells) that appeared to be whole and devoid of physical/chemical damage in visible light were picked and analysed by scanning

electron microscope (SEM) to confirm whether or not the tests were indeed damage free. The SEM images were also used for taxonomic identification. The core-top samples of open-marine diatom ooze sediment from PC12 and PC5 were processed for comparison with the foraminiferal assemblages isolated from glacial-age diamict. Foraminiferal abundances from the marine diamict and ooze sediment were calculated.

Results

Foraminiferal preservation state

A total of 302 whole foraminifera were picked from 55 samples of diamict from PC12 and PC5. In visible light, 98% of tests (296 specimens) are white and translucent to opaque, whereas the remaining 2% are discoloured yellow-brown. All tests are small and thin, i.e. there are no heavily calcified or sediment filled tests. Among the 40 tests selected for SEM analysis, 32 (80%) show no surface damage (Fig. 2). A detailed visual examination of the SEM images showed that the majority of tests possess well-defined pores and sharp edges. Two tests show signs of minor chemical dissolution such as etching and/or slightly enlarged pores (e.g. Fig. 2a). Five show physical damage, which appears




Fig. 3. Distribution of foraminiferal species abundances (i.e. number of specimens) in **a.** NBP0802-PC12, and **b.** NBP0803-PC5 on the middle shelf grounding zone wedge diamict of Global Challenger Basin, eastern Ross Sea. Names highlighted are those of agglutinated species.

Table I. Distribution of foraminiferal species in NBP0802-PC12 and NBP0803-PC5 from core-top open-marine ooze overlying the middle shelf grounding zone wedge of Global Challenger Basin, eastern Ross Sea.

Species	NBP0802-PC12	NBP0803-PC5
<i>Trochammina</i> spp.	54	122
<i>Pseudobolivina</i> sp.*	8	18
<i>Miliolinella</i> sp.	0	15
<i>Ammodiscus</i> sp.	1	6
<i>Spirillina</i> sp.	0	2
<i>Pelosina</i> sp.	0	2
<i>Melonis</i> sp.	2	1
<i>Stainforthia</i> sp.	1	1
<i>Miliammina</i> sp.	7	1

*Agglutinated species.

as fractures or holes on the test walls (e.g. Fig. 2l & m) and one (a planktonic) shows an eroded surface (Fig. 2t). Aside from the damage to the planktonic foraminifera, no significant differences in the degree of physical or chemical damage of the tests were observed between the two core sites.

Assemblage diversity: the grounding zone wedge marine diamict

We isolated 302 foraminifera, averaging less than six minute specimens per 15 ml sample (i.e. 0.3 specimens cm⁻³). No extinct taxa were found in the processed picked samples. Nineteen taxa (18 benthics and the planktonic *Neogloboquadrina pachyderma* (Ehrenberg)) were identified from the diamict samples from PC12 and PC5 (Fig. 3). The assemblage is dominated by *Globocassidulina subglobosa* (Brady) and *Epistominella* spp. All other species generally constitute < 5% of the assemblages. We found only two agglutinated genera, *Textularia* and *Trochammina*, in our samples at 2.3% and 8.8% in PC12 and PC5, respectively (e.g. Fig. 2r & s).

Assemblage diversity: the interglacial-age ooze

Foraminiferal abundance was 20 times higher in the core-top diatom ooze samples with an average of 120 tests per 15 ml sample. The assemblage of the core-top ooze samples includes mainly agglutinated forms (Table I). Among the nine genera found in the surface samples, *Trochammina* is the most common (> 70% in both PC5 and PC12) and is associated mainly with species of *Pseudobolivina*, *Miliolinella* and *Miliammina*. Six of the nine genera are the same in the two core-top diatom ooze samples (Table I). Four of the nine genera in the core-top sample appear in the GZW diamict assemblage.

Stable isotopic data

Small size and low abundances of foraminifera precluded obtaining geochemical data for individual specimens. For

Table II. $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ data from benthic foraminifera.

Core	Core depth (in cm)	$\delta^{18}\text{O}$ (in ‰)	$\delta^{13}\text{C}$ (in ‰)
NBP0802-PC12	184–194	4.52	-0.77
	194–198	4.34	-0.37
	198–206	3.42	-0.74
	206–214	-	-
	214–222	4.57	-0.12
NBP0803-PC5	222–232	3.99	-0.48
	240–250	4.30	-0.73
	250–264	4.28	-0.77
	264–274	-	-
	274–282	4.31	-0.74
	282–288	4.36	-0.84

this reason, isotopic results were obtained on bulk foraminiferal samples. The low abundance of calcareous species in the core-top ooze precluded obtaining any stable isotopic data on core-top foraminifera. Results of the oxygen and carbon isotope analyses are given in Table II. The $\delta^{18}\text{O}$ values were highly positive ranging from 3.42‰ to 4.57‰, with mean $\delta^{18}\text{O}$ values of $4.17 \pm 0.47\text{‰}$ and $4.31 \pm 0.04\text{‰}$ for PC12 and PC5, respectively. Values of $\delta^{13}\text{C}$ measured on foraminiferal tests range from -0.84‰ to -0.12‰.

Discussion

Preservation of minute benthic foraminifera in grounding zone wedge diamict

Discolouration and other preservation issues may be the way to distinguish between *in situ* and reworked foraminiferal tests. Foraminiferal biostratigraphical analyses of Cape Roberts Project core data (e.g. Webb & Strong 2006) show that a low diversity and low abundance assemblage similar to that found in our piston core diamict samples has existed, more or less continuously in the Ross Sea region since at least the early Oligocene. In other words, the diamict assemblage that was isolated in piston cores might be recycled from pre-LGM or even much older strata.

A good example of reworked calcareous benthic foraminifera from Antarctic Holocene sediment was illustrated in Majewski & Anderson (2009). They showed three populations of thickly testate *Globocassidulina biora* (Crespin) from a single sample. Some of the worst preserved specimens were clearly compacted with poorly preserved walls and no signs of pores. The less altered were light-to-dark orange in colour with a rough surface and partly or completely obscured pores, and finally the best preserved were white specimens with smooth walls and well-preserved pores. This example shows what kind of alterations could be expected for reworked foraminiferal tests.

The SEM images confirm that the majority (80%) of foraminiferal tests that were isolated from diamict are devoid of significant damage (Fig. 2). A majority of our

specimens from GZW diamict show smooth test walls with well-preserved pores. They are all small and hollow, with thin walls that would seemingly make them susceptible to breakage during compaction or transport. When damage is present, it is mostly found on either the last chamber or along test margins. The slight physical breakage observed on some foraminifera may be due to *in situ* predation or sample processing.

Highly eroded and discoloured surfaces are observed on only the three planktonic foraminifera (e.g. Fig. 2t), which suggests that those three specimens may be reworked from older sediment. Conversely, these specimens may have been exposed to a shallow Carbon Compensation Depth (CCD). All other tests are benthic foraminifera and these exhibit at most only minor effects of dissolution manifested as microscopic surface etching or slightly enlarged pores (Fig. 2). This relatively slight chemical damage might have been due to less prolonged exposure to corrosive waters in the Ross Sea where the average depth of the CCD is *c.* 550 m (Osterman & Kellogg 1979).

The assemblage also includes a few agglutinated forms. Most of them tend to disintegrate after death because of the fragile nature of their tests (Corliss 1985). Such paucity of agglutinated forms in sediments older than a few thousands of years is not uncommon (e.g. Majewski & Anderson 2009) and it may obscure reconstructions of fossil assemblages. In this case, it seems to provide no arguments for either a reworked or *in situ* character of the GZW assemblage, as the original ratio of calcareous forms is unknown.

In summary, the overall excellent preservation state of the foraminifera is consistent with an assemblage that is *in situ*. However, this physical criterion cannot be considered definitive proof of the *in situ* assemblage because it cannot be proved that recycled foraminifera cannot survive physical and chemical damage intact.

Difference between grounding zone wedge diamict and open-marine benthic foraminiferal assemblages

We assumed that the modern ooze assemblage can be taken as an approximation to the foraminiferal assemblage that inhabited the open-marine assemblage during pre-LGM interglacials. In this line of reasoning, if the foraminifera isolated from GZW diamict were reworked from an ooze deposited during a previous interglacial, then the diamict assemblage should be composed of a subset of the ooze assemblage.

The high abundance of foraminifera in the core-top oozes analysed is due at least in part to the higher productivity and extremely low influx of terrigenous sediment to the open water continental shelf. The dominance of agglutinated foraminifera in our core-top assemblage (Table I) is typical for the modern eastern Ross Sea open-marine conditions (McKnight 1962, Pflum 1966, Osterman & Kellogg 1979). If this modern open-marine assemblage is a potential proxy for pre-LGM open-marine assemblages in the eastern Ross Sea, then it can be expected that the reworked fraction would be dominated by the most abundant robust tests from the modern-like ooze that, following Schröder (1988), should




Fig. 4. Environmental tolerances for foraminiferal species found in PC12 and PC5 of the grounding zone wedge diamict. Grey shading indicates a common range of environmental conditions that all species in the diamict share.

be the agglutinated *Trochammina*. Its occurrence in the diamict permits the possibility that it was reworked from pre-LGM interglacial ooze, but alternately, its very low abundance may indicate the opposite, i.e. that its occurrences are primarily *in situ* and it belongs to a very different assemblage dominated by calcareous forms.

The diamict assemblage is dominated by two calcareous taxa, i.e. *G. subglobosa* and *Epistominella* spp. (Fig. 3), both of which are also known from the Ross Sea (Osterman & Kellogg 1979, Ward *et al.* 1987, Melis & Salvi 2009) and other modern Antarctic continental shelf areas (Pudsey *et al.* 2006, Majewski 2013). Although neither of these species is found in the modern diatom ooze from our two core-top samples (Table I), it cannot be ruled out that pre-LGM open-marine conditions in the eastern Ross Sea were different than Recent, facilitating much different foraminiferal assemblages.

The environmental affinities of the well-preserved benthic foraminifera

The *in situ* assemblage should include only those species that could tolerate the expected conditions for an ice-proximal environment. Benthic foraminiferal assemblages are controlled by many factors including food availability and its seasonal delivery, sediment input, substrate type, water mass properties and bathymetry, temperature, salinity and oxygen content (e.g. Osterman & Kellogg 1979, Majewski 2005, Melis & Salvi 2009, Gooday *et al.* 2014). The dominant foraminifera in the diamict assemblage (*G. subglobosa*, *Epistominella* spp. and *Bulimina aculeata* d'Orbigny) are typical of cold-water benthic assemblages (Nelson *et al.* 1993, Polyak & Solheim 1994, Ishman & Foley 1996). Figure 4 summarizes the ranges for three environmental parameters for the benthic species that were isolated in the diamict based on data from McKnight (1962), Pflum (1966), Ward *et al.* (1987), Mackensen & Hald (1988), Asioli (1995), Harloff & Mackensen (1997), Majewski (2005) and other summary data (Encyclopedia of Life 2009, <http://www.eol.org>). The grey-shaded area shows that there is a common range of environmental conditions that all species in the diamict share (Fig. 4). This overlap is a minimum requirement for the GZW assemblage to be considered *in situ*.

Reports on foraminiferal communities from under ice sheets around Antarctica are not uncommon. The best studied are fossil assemblages from the area of the Larsen Ice Shelf collapse aimed at developing foraminiferal proxies for tracking past ice shelf retreats (Murray & Pudsey 2004). Abundant planktonic foraminifera were described from beneath Amery Ice Shelf reflecting sub-ice shelf circulation (Hemer *et al.* 2007). Foraminifera were also encountered from below the Ross Ice Shelf. Lipps *et al.* (1979) reported empty foraminiferal tests from

under 420 m of ice as far south as 430 km from open sea. More recently, Pawlowski *et al.* (2005) described allogromiid foraminifera and gromiids along with empty tests from the ANDRILL site HWD-2, 12 km east of the ice edge. They interpret the presence of the rather diverse living protist community as the result of advected phytodetritus that was apparently lacking at the site investigated by Lipps *et al.* (1979).

The question to be answered is whether our GZW diamict assemblage is similar to what could be hypothetically anticipated in habitats located many tens of kilometres from open ocean as its potential food source. Korsun *et al.* (1995) showed that ice-proximal Arctic fjord settings lack large tests but have abundant juveniles, which they argued is an opportunistic response to environmental stress of glacier proximity. These glacier-proximal settings were also dominated by abundant monothalamous allogromiid species that lack calcified tests (Korsun & Hald 2000, Sabbatini *et al.* 2007). The allogromiid foraminifera are also prominent in marginal settings of the South Shetland Islands (Majewski *et al.* 2007) and in McMurdo Sound (Gooday *et al.* 1996), and were reported from below the Ross Ice Shelf (Pawlowski *et al.* 2005). Because they do not possess robust tests, they are not present in the fossil record and are also commonly overlooked in studies of Recent assemblages, if sediment samples are washed and dried prior to picking. The minute size and scarcity of the predominantly calcareous benthic foraminifera found in GZW diamict seems to coincide with an assemblage originally dominated by allogromiid foraminifera with only a minor share of calcareous forms. Their small size and thin walls suggest that this could have been an element of an opportunistic assemblage strongly affected by scarce and/or seasonal delivery of phytodetritus to the sea floor (Gooday 1993).

It is true, however, that similar suites of minute calcareous benthic foraminifera were commonly described also from open-sea environments where food is limited and/or strongly seasonal, for example, the *Epistominella* trough assemblage found off the Larsen A Ice Shelf (Ishman & Szymcek 2003) or the *Epistominella* spp. dominated assemblage from central Pine Island Bay (Majewski 2013), where persistent sea ice often hinders primary production. A widespread assemblage dominated by *G. subglobosa* was also described from surface sediments of the western Ross Sea (Assemblage 3 from Osterman & Kellogg 1979). Thus, one could argue that the major components of our GZW assemblage are actually more characteristic of open-water conditions.

On the other hand, marginal, especially sub- or near-glacier environments, are definitely under-investigated. There are simply hardly any modern analogue data. Sub-ice shelf facies with abundant calcareous foraminifera were reported from Marguerite Trough (Kilfeather *et al.* 2011). The most

glacier-proximal assemblage of the Holocene record from Firth of Tay (Majewski & Anderson 2009) corresponds in terms of species composition and minute size of tests with the GZW assemblage. Sediments interpreted to represent sub-ice shelf environments from Pine Island Bay contain benthic foraminifera that represent roughly similar assemblages dominated by sparse and minute species (unpublished data).

Following these arguments, species composition of the GZW fossil assemblage seems to be coherent and to represent opportunistic, food-limited communities. This assemblage seems to correspond well with conditions of scarce phytodetritus only occasionally delivered by subglacial currents, but on the other hand could also be associated with an open-sea environment with primary production restricted by persistent sea ice and/or oligotrophic waters.

Evaluation of stable isotopic data

Both $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values may differ significantly between different benthic foraminiferal species from a single site (e.g. Majewski *et al.* 2012). Because, due to the minute size of the specimens, our measurements were performed on bulk samples grouping together specimens belonging to different species, it is difficult to unequivocally evaluate these data. At face value, the oxygen and carbon stable isotopic values are reasonable and consistent between different samples (Table II). They do not have values that would suggest possible diverse provenances of analysed foraminifera, in turn supporting the hypothesis that at least a majority of analysed specimens are of similar origin.

Implications for dating of ice sheet grounding events

Throughout the earlier discussion, we showed that diamict deposited on a GZW foreset contains a low abundance of minute, well-preserved, extant foraminifera. Our analyses of the preservation states, assemblage compositions, as well as $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values, failed to provide conclusive evidence that the benthic foraminifera are *in situ*. Nonetheless, nothing in our data requires that the *in situ* assumption be abandoned as a working hypothesis. This finding may have a pronounced significance for radiocarbon dating deglaciation events after the LGM.

So far, two basic strategies have been used to directly date the marine record of Antarctic ice sheet retreat. The most widely used approach is to constrain the time corresponding to the resumption of marine sedimentation that followed ice sheet retreat. In this approach, dateable materials, either foraminifera and/or acid insoluble organic matter (AIOM), are isolated from the basal-most section of post-glacial diatom ooze that overlies the

glacial-age diamict. This approach has its weaknesses, however. It assumes that ooze sedimentation began immediately after ice sheet retreat and has since been continuous and undisturbed. If this assumption is not correct, it may lead to significant underestimating of deglaciation ages.

The second strategy is to date the grounding event itself. This relies on isolating dateable material from glacial-age diamict. Licht & Andrews (2002) reasoned that the advance of grounded ice eroded and recycled proglacial sediment that was originally deposited on the previously open-water continental shelf. Given that proglacial sediments (and the dateable material it contains) become recycled and incorporated into the diamict, the youngest radiocarbon dates of either AIOM or foraminifera in diamict provide a maximum age of the grounding event.

In a departure from the strategy employed by Licht & Andrews (2002), we suggest that well-preserved foraminifera from the marine diamict deposited on the middle continental shelf GZW foreset are *in situ*. This assumption has already been employed by Bart & Cone (2012). Their radiocarbon dates from two widely separated core sites are tightly clustered and stratigraphically ordered, providing support for the *in situ* nature of the well-preserved foraminiferal assemblages. Still, those dates could also be considered suspect because i) many specimens had to be combined to produce a single radiocarbon date, and ii) the dates suggest that the WAIS retreated significantly earlier than is deduced from other marine and terrestrial locations.

Our present study does not show unequivocally that the *in situ* assumption of Bart & Cone (2012) is correct, but nothing in the nature of the well-preserved minute calcareous foraminifera assemblages found in GZWs suggests that they are reworked. The possibility that the best preserved fraction is *in situ* is important because radiocarbon dates of carefully isolated *in situ* foraminifera can be used to date the timing and durations of grounding events in the many places where GZWs are found on the Antarctic continental shelves (e.g. Graham *et al.* 2010) more precisely than by dating post-glacial diatom ooze. This new approach can significantly improve our knowledge of Antarctic deglaciation history and environmental conditions existing where grounded ice streams meet the marine settings, providing better data for climatic modelling.

Conclusions

To our knowledge, foraminifera have not yet been observed from samples and observations made by the WISSARD team, thus this is the first analysis of GZW foreset ecology. As for our analyses of the GZW preserved on the eastern Ross Sea middle continental

shelf, we propose that further work is needed before the *in situ* hypothesis is confirmed because the data do not convincingly require that the foraminifera that were isolated from the diamict are *in situ*. Conversely, nothing in our new data requires that the *in situ* assumption be abandoned as a working hypothesis.

Acknowledgements

The research was supported by NSF OPP grant #1246357 to Bart. Thanks are extended to the ECO Captain, ship crew and marine/electronic support crews of the R/VIB *Nathaniel B. Palmer* for assistance acquiring core and geophysical data during difficult field conditions during the NBP0802 and 0803 cruises to Glomar Challenger Basin. Thanks are also extended to the curatorial staff of the Antarctic Marine Geology Research Facilities in Tallahassee for providing the core samples requested from the archive. We thank Patrick Quilty and the journal editor for suggestions that improved the manuscript.

Author contribution

Coquereau sampled, processed and analysed the samples for foraminifera under the supervision of Bart and Warny. Foraminifera identifications were confirmed by Majewski and Barun Sen Gupta. All authors participated in the data interpretation and writing of the manuscript.

References

- ASIOLI, A. 1995. Living (stained) benthic foraminifera distribution in the western Ross Sea (Antarctica). *Palaeopelagos*, **5**, 201–214.
- BART, P.J. & CONE, A.N. 2012. Early stall of West Antarctic Ice Sheet advance on the eastern Ross Sea middle shelf followed by retreat at 27,500 ¹⁴C yr bp. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **335**, 10.1016/j.palaeo.2011.08.007.
- CORLISS, B.H. 1985. Microhabitats of benthic foraminifera within deep-sea sediments. *Nature*, **314**, 10.1038/314435a0.
- DOMACK, E.W., JACOBSEN, E.A., SHIPP, S. & ANDERSON, J.B. 1999. Late Pleistocene-Holocene retreat of the West Antarctic Ice Sheet system in the Ross Sea: part 2—sedimentologic and stratigraphic signature. *Geological Society of America Bulletin*, **111**, 1517–1536.
- GOODAY, A.J. 1993. Deep-sea benthic foraminiferal species which exploit phytodetritus – characteristic features and controls on distribution. *Marine Micropaleontology*, **22**, 187–205.
- GOODAY, A.J., BOWSER, S.S. & BERNHARD, J.M. 1996. Benthic foraminiferal assemblages in Explorers Cove, Antarctica: a shallow-water site with deep-sea characteristics. *Progress in Oceanography*, **37**, 117–166.
- GOODAY, A.J., ROTHE, N., BOWSER, S.S. & PAWLOWSKI, J. 2014. Benthic foraminifera. In DE BROYER, C., KOUUBI, P., GRIFFITHS, H.J., *et al.*, eds. *Biogeographic atlas of the Southern Ocean*. Cambridge: Scientific Committee on Antarctic Research, 74–82.
- GRAHAM, A.G.C., LARTER, R.D., GOHL, K., DOWDESHELL, J.A., HILLENBRAND, C.D., SMITH, J.A., EVANS, J., KUHN, G. & DEEN, T. 2010. Flow and retreat of the Late Quaternary Pine Island-Thwaites palaeo-ice stream, West Antarctica. *Journal of Geophysical Research—Earth Surface*, **115**, 10.1029/2009JF001482.
- HARLOFF, J. & MACKENSEN, A. 1997. Recent benthic foraminiferal associations and ecology of the Scotia Sea and Argentine Basin. *Marine Micropaleontology*, **31**, 1–29.
- HEMER, M.A., POST, A.L., O'BRIEN, P.E., CRAVEN, M., TRUSWELL, E.M., ROBERTS, D. & HARRIS, P.T. 2007. Sedimentological signatures of the sub-Amery Ice Shelf circulation. *Antarctic Science*, **19**, 497–506.
- ISHMAN, S.E. & FOLEY, K.M. 1996. Modern benthic foraminifer distribution in the Amerasian Basin, Arctic Ocean. *Micropaleontology*, **42**, 206–220.
- ISHMAN, S.E. & SZYMCEK, P. 2003. Foraminiferal distributions in the former Larsen-A Ice Shelf and Prince Gustav Channel region, eastern Antarctic Peninsula margin: a baseline for Holocene paleoenvironment interpretation. *Antarctic Research Series*, **79**, 239–260.
- JAKOBSSON, M., ANDERSON, J.B., NITSCHKE, F.O., DOWDESHELL, J.A., GYLLENCREUTZ, R., KIRCHNER, N., MOHAMMAD, R., O'REGAN, M., ALLEY, R.B., ANANDAKRISHNAN, S., ERIKSSON, B., KIRSHNER, A., FERNANDEZ, R., STOLLDOERF, T., MINZONI, R. & MAJEWSKI, W. 2011. Geological record of ice shelf break-up and grounding line retreat, Pine Island Bay, West Antarctica. *Geology*, **39**, 10.1130/G32153.1.
- KILFEATHER, A.A., COFAIGH, C.O., LLOYD, J.M., DOWDESHELL, J.A., XU, S. & MORETON, S.G. 2011. Ice-stream retreat and ice-shelf history in Marguerite Trough, Antarctic Peninsula: sedimentological and foraminiferal signatures. *Geological Society of America Bulletin*, **123**, 997–1015.
- KORSUN, S. & HALD, M. 2000. Seasonal dynamics of benthic foraminifera in a glacially fed fjord of Svalbard, European Arctic. *Journal of Foraminiferal Research*, **30**, 251–271.
- KORSUN, S.A., POGODINA, I.A., FORMAN, S.L. & LUBINSKI, D.J. 1995. Recent foraminifera in glaciomarine sediments from three arctic fjords of Novaja Zemlja and Svalbard. *Polar Research*, **14**, 15–32.
- LICHT, K.J. & ANDREWS, J.T. 2002. The ¹⁴C record of Late Pleistocene ice advance and retreat in the central Ross Sea, Antarctica. *Arctic, Antarctic, and Alpine Research*, **34**, 324–333.
- LIPPS, J.H., RONAN, T.E. & DELACA, T.E. 1979. Life below the Ross Ice Shelf, Antarctica. *Science*, **203**, 447–449.
- MACKENSEN, A. & HALD, M. 1988. *Cassidulina teretis* Tappan and *Cassidulina laevigata* d'Orbigny: their modern and late Quaternary distribution in northern seas. *Journal of Foraminiferal Research*, **18**, 16–24.
- MACKENSEN, A. 2012. Strong thermodynamic imprint on Recent bottom-water and epibenthic $\delta^{13}\text{C}$ in the Weddell Sea revealed: implications for glacial Southern Ocean ventilation. *Earth and Planetary Science Letters*, **317**, 10.1016/j.epsl.2011.11.030.
- MAJEWSKI, W. 2005. Benthic foraminiferal communities: distribution and ecology in Admiralty Bay, King George Island, West Antarctica. *Polish Polar Research*, **26**, 159–214.
- MAJEWSKI, W. 2013. Benthic foraminifera from Pine Island and Ferrero bays, Amundsen Sea. *Polish Polar Research*, **34**, 169–200.
- MAJEWSKI, W. & ANDERSON, J.B. 2009. Holocene foraminiferal assemblages from Firth of Tay, Antarctic Peninsula: paleoclimate implications. *Marine Micropaleontology*, **73**, 135–247.
- MAJEWSKI, W., LECROQ, B., SINNIGER, F. & PAWLOWSKI, J. 2007. Monothalamous foraminifera from Admiralty Bay, King George Island, West Antarctica. *Polish Polar Research*, **28**, 187–210.
- MAJEWSKI, W., WELLNER, J.S., SZCZUCIŃSKI, W. & ANDERSON, J.B. 2012. Holocene oceanographic and glacial changes recorded in Maxwell Bay, West Antarctica. *Marine Geology*, **326**, 67–79.
- McKNIGHT Jr, W.M. 1962. The distribution of foraminifera off parts of the Antarctic coast. *Bulletin of American Paleontology*, **44**, 65–158.
- MELIS, R. & SALVI, G. 2009. Late Quaternary foraminiferal assemblages from western Ross Sea (Antarctica) in relation to the main glacial and marine lithofacies. *Marine Micropaleontology*, **70**, 10.1016/j.marmicro.2008.10.003.

- MURRAY, J.W. & PUDSEY, C.J. 2004. Living (stained) and dead foraminifera from the newly ice-free Larsen Ice Shelf, Weddell Sea, Antarctica: ecology and taphonomy. *Marine Micropaleontology*, **53**, 10.1016/j.marmicro.2004.04.001.
- NELSON, C.S., COOK, P.J., HENDY, C.H. & CUTHBERTSON, A.M. 1993. Oceanographic and climatic changes over the past 160,000 years at Deep Sea Drilling Project Site 594 off southeastern New Zealand, southwest Pacific Ocean. *Paleoceanography*, **8**, 435–458.
- O'BRIEN, P.E., SMITH, J., STARK, J.S., JOHNSTON, G., RIDDLE, M. & FRANKLIN, D. 2015. Submarine geomorphology and sea floor processes along the coast of Vestfold Hills, East Antarctica, from multibeam bathymetry and video data. *Antarctic Science*, **27**, 10.1017/S095410201500371.
- OSTERMAN, L.E. & KELLOGG, T.B. 1979. Recent benthic foraminiferal distribution from the Ross Sea, Antarctica: relation to ecologic and oceanographic conditions. *Journal of Foraminiferal Research*, **9**, 250–269.
- PAWLOWSKI, J., FAHRNI, J.F., GUIARD, J., CONLAN, K., HARDECKER, J., HABURA, A. & BOWSER, S.S. 2005. Allogromiid foraminifera and gromiids from under the Ross Ice Shelf: morphological and molecular diversity. *Polar Biology*, **28**, 514–522.
- PFLUM, C.E. 1966. The distribution of foraminifera in the eastern Ross Sea, Amundsen Sea, and Bellingshausen, Antarctica. *Bulletin of American Paleontology*, **50**, 151–209.
- POLYAK, L. & SOLHEIM, A. 1994. Late-glacial and postglacial environments in the northern Barents Sea west of Franz Josef Land. *Polar Research*, **13**, 197–207.
- POST, A.L., GALTON-FENZI, B.K., RIDDLE, M.J., HERRAIZ-BORREGUERO, L., O'BRIEN, P.E., HEMER, M.A., MCMINN, A., RASCH, D. & CRAVEN, M. 2014. Modern sedimentation, circulation and life beneath the Amery Ice Shelf, East Antarctica. *Continental Shelf Research*, **74**, 77–87.
- PUDSEY, C.J., MURRAY, J.W., APPLEBY, P. & EVANS, J. 2006. Ice shelf history from petrographic and foraminiferal evidence, northeast Antarctic Peninsula. *Quaternary Science Reviews*, **25**, 2357–2379.
- SABBATINI, A., MORIGI, C., NEGRI, A. & GOODAY, A.J. 2007. Distribution and biodiversity of stained monothalamous foraminifera from Tempelfjord, Svalbard. *Journal of Foraminiferal Research*, **37**, 93–106.
- SCHRÖDER, C.J. 1988. Subsurface preservation of agglutinated foraminifera in the northwest Atlantic Ocean. *Abhandlungen der geologischen Bundesanstalt*, **41**, 325–336.
- WARD, B.L., BARRETT, P.J. & VELLA, P. 1987. Distribution and ecology of benthic foraminifera in McMurdo Sound, Antarctica. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **58**, 139–153.
- WEBB, P.N. & STRONG, C.P. 2006. Foraminiferal biostratigraphy and paleoecology in upper Oligocene–lower Miocene glacial marine sequences 9, 10, and 11, CRP-2/2A drill hole, Victoria Land Basin, Antarctica. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **231**, 71–100.