

Staying alive on an active volcano: 80 years population dynamics of *Cytisus aeolicus* (Fabaceae) from Stromboli (Aeolian Islands, Italy)

Journal: Ecological Processes

Zaia R., Pasta S., Di Rita F., Laudicina V.A., Lo Cascio P., Magri D., Troia A., Guarino R.^{1*}

¹University of Palermo, Department STEBICEF, via Archirafi 38, 90123 Palermo, Italy. Email: riccardo.guarino@unipa.it

**Appendix 1 - photo collection of the growing site of
Cytisus aeolicus from Stromboli Island**



Fig. A1.1: view of the sand deposit of Rina Grande, seen from the s-pop PV (see Fig. 1 in the main text). On the other side of Rina Grande, the s-pops SS (left) and SC (right). (Photo: R. Guarino).



Fig. A1.2: contact between the s-pop VB (see Fig. 1 in the main text) and the neighbouring *Spartium junceum* shrubland, in the area where soil samples were collected (Photo: R. Guarino).



Fig. A1.3: young individuals of *C. aeolicus* in the area named Guardiani, slightly above the upper limit of the s-pop VB (see Fig. 1 in the main text) (photo: R. Guarino).



Fig. A1.4: view of the s-pop SS (see Fig. 1 in the main text) (photo: R. Guarino).