

COMPARATIVE DISTRIBUTION OF MOLLUSKS IN DREDGED AND UN-DREDGED PORTIONS OF AN ESTUARY, WITH A SYSTEMATIC LIST OF SPECIES¹

BY JAMES E. SYKES AND JOHN R. HALL, *FISHERY BIOLOGISTS*
BUREAU OF COMMERCIAL FISHERIES BIOLOGICAL LABORATORY
ST. PETERSBURG BEACH, FLA. 33706

ABSTRACT

A survey of benthic mollusks in Boca Ciega Bay, Fla., showed a much smaller number and variety of species in the soft sediments in dredged canals than in the predominantly sand and shell sediments in undredged

areas. Samples contained an average of 60.5 live mollusks and 3.8 species in undredged areas and 1.1 individuals and 0.6 species in dredged canals. A list of mollusks collected in this survey and in past studies is appended.

This report compares the numbers and varieties of mollusks in fine sediments of dredged canals with those found in undisturbed bottoms of sand and shell in Boca Ciega Bay, Fla. The bay is a shallow coastal lagoon of about 70 km.² which connects with Tampa Bay at its southern end (fig. 1). Some of the previous investigations in the lagoon included studies of sediments (Goodell and Gorsline, 1961; Taylor and Saloman, 1969); hydrology (Saloman and Taylor, 1968); submerged vegetation (Pomeroy, 1960; Phillips, 1960); fishes (Springer and Woodburn, 1960; Sykes and Finucane, 1964); and benthic invertebrates (Hutton, Eldred, Woodburn, and Ingle, 1956; Bullock and Boss²).

A recent evaluation of the effects of dredging and filling has documented a large loss of estuarine resources in Boca Ciega Bay (Taylor and Saloman, 1968). It was here that scientists and conservationists were finally successful in suppressing a dredge-fill proposal of 202 ha. (Sykes, 1967). This is also the bay in which the U.S. Army Corps of Engineers denied a dredge-fill application for the first time on the basis of fish and wildlife values, thus providing a stimulus for more comprehensive assessments of the

biological and recreational aspects involved in future bayfill developments.

Thorson (1956) and others have concluded that sediment composition is a cardinal factor in controlling the settlement and viability of many marine invertebrates. The distribution of sessile benthic mollusks indicates to the marine ecologist the ability of the environment to support life. Marked deficiencies in abundance and variety indicate abnormality of the environment, and the degree of deficiency is roughly proportional to the degree of abnormality.

PROCEDURES

Between September 1963 and August 1964, we took 107 bottom samples at 31 stations in Boca Ciega Bay (figs. 1 and 2). Seven stations were in canals between finger fills (1-7), and the other 24 (8-31) were in relatively undisturbed areas of the bay. We collected algae, sea grasses, and benthic animals with a bucket dredge and bottom drag (Taylor, 1965). In water less than 1 m. deep, three shovelfuls of bay bottom (about 15 l.) were substituted for the dredge haul. One station sample consisted of the combined catch from one bucket dredge (or three shovelfuls) and one bottom drag. At each station a subsample of sediment was taken from the dredge or shovel and was later analyzed at Florida State University.

¹ Contribution No. 57, Bureau of Commercial Fisheries Biological Laboratory, St. Petersburg Beach, Fla. 33706.

² Bullock, R., and O. Boss. 1963. Ecological distribution of marine mollusks in Boca Ciega Bay, Florida. Winter term project. Mimeographed report on file at Florida Presbyterian College, St. Petersburg, Fla. 33733.




FIGURE 1.—Collecting stations 20 to 31 and area of stations 1 to 19 (see fig. 2), Boca Ciega Bay, Fla.



FIGURE 2.—Collecting stations 1 to 19, Boca Ciega Bay, Fla., between Johns Pass and Corey Causeway.

We washed samples for benthic organisms on a 24-mesh sieve which had an opening of 0.701 mm. and fixed the material retained by the sieve in a 10 percent sea-water Formalin³ mixture. A protein stain (rose bengal) was added to facilitate the separation of small organisms from debris. Identified animals were preserved in 70 percent isopropanol.

³ Trade names referred to in this publication do not imply endorsement of commercial products.

We identified 168 species of mollusks representing 69 families; of these, representatives of 156 species were collected live. We based determinations on standard taxonomic works (Clench, 1941-69; McLean, 1951; Olsson, Harbison, Fargo, and Pilsbry, 1953; Abbott, 1954, 1968; Perry and Schwengel, 1955; Warmke and Abbott, 1962; Keen, 1963; Wagner and Abbott, 1967; an unpublished report by Bullock and Boss (see footnote 2), and collections at the University of

South Florida ⁴ and the U.S. National Museum. Specimens from this study were deposited in the invertebrate reference collection of the BCF (Bureau of Commercial Fisheries) Biological Laboratory, St. Petersburg Beach, Fla.

MOLLUSK-SEDIMENT RELATIONS

Comparison of mollusks and bottom types showed that species and individuals were much less numerous in soft sediments of canals than in sandy sediments in undredged areas of Boca Ciega Bay (tables 1 and 2). Canal sediments, which averaged 85 percent silt and clay, had 16 live mollusks in 14 samples. Living specimens collected at the seven canal stations were the gastropods *Nassarius vibex* and *Haminoea antillarum*, and the pelecypods *Brachidontes exustus*, *Anomalocardia cuneimeris*, and *Mercenaria campechiensis*. These species and 151 others were collected live from the 24 stations in undredged areas of the bay. Sediments from natural bottom, which averaged 91 percent sand and shell, yielded 5,631 live mollusks in 93 samples.

Pratt (1953) suggested that soft sediments and associated hydrological conditions may be limiting because (1) rapid deposition has a smothering effect, (2) high organic content of soft sediments depletes dissolved oxygen, and (3) weak currents in areas of deposition are insufficient for the removal of toxic metabolic wastes. Comparisons of sediments and environmental factors (Taylor and Saloman, 1968) in dredged and undredged areas at sampling stations lead us to conclude that the soft sediment is the principal factor limiting the abundance and diversity of benthic mollusks in bayfill canals of Boca Ciega Bay. Such sediments are as thick as 4 m. in waterways that were dredged 15 years ago.

ACKNOWLEDGMENTS

The authors gratefully acknowledge confirmations and corrections of identifications made by Harry W. Wells, Department of Biology, University of Delaware, Newark, Del.; Joseph Rosewater, Division of Mollusks, U.S. National Museum, Washington, D.C.; and George Radwin, San Diego Natural History Museum, San Diego, Calif.

⁴ Hillman Collection, University of South Florida, Tampa, Fla. 33620.

TABLE 1.—Depth, bottom type, and number of live mollusks collected at stations in dredged canals in Boca Ciega Bay, Fla., 1963-64

Canal station	Depth	Bottom type		Species per sample	Individual per sample
		Sand size and larger	Silt and clay		
1.....	M. 4	Percent 6	Percent 94	Number 1.0	Number 1.0
2.....	3	7	93	1.0	1.5
3.....	4	5	95	0	0
4.....	3	60	40	1.0	2.0
5.....	6	7	93	0	0
6.....	4	6	94	.5	1.0
7.....	4	14	86	.5	2.5
Average.....	4	15	85	.6	1.1

TABLE 2.—Depth, bottom type, and number of live mollusks collected at stations in undredged areas of Boca Ciega Bay, Fla., 1963-64

Stations in undredged areas	Depth	Bottom type		Species per sample	Individual per sample
		Sand size and larger	Silt and clay		
8.....	M. 2.5	Percent 98	Percent 2	Number 3.5	Number 8.0
9.....	1.0	99	1	3.0	7.3
10.....	1.5	89	11	.5	.5
11.....	1.5	97	3	4.0	25.0
12.....	2.5	96	4	.5	2.5
13.....	2.0	96	4	3.5	18.0
14.....	1.7	88	12	.5	2.0
15.....	.5	92	8	2.6	20.6
16.....	.7	99	1	.3	.7
17.....	5.0	83	17	1.0	2.5
18.....	1.5	90	4	6.5	58.0
19.....	1.0	84	16	1.6	24.0
20.....	1.5	99	1	5.6	140.0
21.....	1.0	98	2	3.4	67.8
22.....	2.0	98	2	2.8	71.8
23.....	7.0	98	2	4.3	86.6
24.....	2.0	79	19	11.7	316.0
25.....	.7	97	3	3.2	8.4
26.....	7.0	99	1	4.0	16.7
27.....	.7	97	3	5.0	11.0
28.....	3.0	12	88	4.7	56.3
29.....	2.0	87	12	6.7	89.0
30.....	.3	98	2	6.2	78.0
31.....	2.0	95	5	4.0	86.5
Average.....	1.9	91	9	3.8	60.5

LITERATURE CITED

- ABBOTT, R. TUCKER.
1954. American seashells. D. Van Nostrand Co., Princeton, N.J., 541 pp.
1968. Seashells of North America. Golden Press, N.Y., 280 pp.
- CLENCH, WILLIAM J. (EDITOR).
1941-69. Monographs of the marine mollusks of the Western Atlantic. Johnsonia, vols. I-IV. Department of Mollusks, Mus. Comp. Zool., Harvard Univ., Cambridge, Mass.
- DRAGOVICH, ALEXANDER, and JOHN A. KELLY, JR.
1964. Ecological observations of macro-invertebrates in Tampa Bay, Florida, 1961-1962. Bull. Mar. Sci. Gulf Carib. 14: 74-102.
- GOODELL, H. G., and D. S. GORSLINE.
1961. A sedimentologic study of Tampa Bay, Florida. Rep. Int. Geol. Congr., 21 Sess., Pt. 23, pp. 75-88. Int. Ass. Sediment., Copenhagen, 1961.

- HUTTON, ROBERT F., BONNIE ELDRED, KENNETH D. WOODBURN, and ROBERT INGLE.
1956. The ecology of Boca Ciega Bay with special reference to dredging and filling operations. Fla. State Bd. Conserv., Tech. Ser. 17(1), 86 pp.
- KEEN, A. MYRA.
1963. Marine molluscan genera of western North America. Stanford Univ. Press, Stanford, Calif., 126 pp.
- MCLEAN, RICHARD A.
1951. The Pelecypoda or bivalve mollusks of Porto Rico and the Virgin Islands. Scientific Survey of Porto Rico and the Virgin Islands. N.Y. Acad. Sci. 17, 183 pp.
- OLSSON, AXEL A., ANNE HARBISON, WILLIAM G. FARGO, and HENRY A. PILSBRY.
1953. Pliocene Mollusca of southern Florida. Monogr. Acad. Nat. Sci. Philadelphia 8, 458 pp.
- PERRY, LOUISE M., and JEANNE S. SCHWENGEL.
1955. Marine shells of the western coast of Florida. Palaeontological Research Institute, Ithaca, N.Y., 318 pp.
- PHILLIPS, RONALD C.
1960. Ecology and distribution of marine algae found in Tampa Bay, Boca Ciega Bay, and at Tarpon Springs, Florida. Quart. J. Fla. Acad. Sci. 23: 222-260.
- POMEROY, LAWRENCE R.
1960. Primary productivity of Boca Ciega Bay, Florida. Bull. Mar. Sci. Gulf Carib. 10: 1-10.
- PRATT, DAVID M.
1953. Abundance and growth of *Venus mercenaria* and *Callocardia morrhua* in relation to the character of bottom sediments. J. Mar. Res. 12: 60-74.
- SALOMAN, CARL H.
1965. Bait shrimp (*Penaeus duorarum*) in Tampa Bay, Florida—biology, fishery economics, and changing habitat. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 52C, iii + 16 pp.
- SALOMAN, CARL H., and JOHN L. TAYLOR.
1968. Hydrographic observations in Tampa Bay, Florida, and the adjacent Gulf of Mexico—1965-1966. U.S. Fish Wildl. Serv., Data Rep. 24, 39 pp. on 6 microfiches.
- SPRINGER, VICTOR G., and KENNETH D. WOODBURN.
1960. An ecological study of the fishes of the Tampa Bay area. Fla. State Bd. Conserv., Prof. Pap. Ser. 1, 104 pp.
- SYKES, JAMES E.
1967. The role of research in the preservation of estuaries. Trans. 32d N. Amer. Wildl. Natur. Resour. Conf., pp. 150-160.
- SYKES, JAMES E., and JOHN H. FINUCANE.
1964. Occurrence in Tampa Bay, Florida, of immature species dominant in Gulf of Mexico commercial fisheries. U.S. Fish Wildl. Serv., Fish. Bull. 65: 369-379.
- TAYLOR, JOHN L.
1965. Bottom samplers for estuarine research. Chesapeake Sci. 6: 233-234.
- TAYLOR, JOHN L., and CARL H. SALOMAN.
1968. Some effects of hydraulic dredging and coastal development in Boca Ciega Bay, Florida. U.S. Fish Wildl. Serv., Fish. Bull. 67: 213-241.
1969. Sediments, oceanographic observations, and floristic data from Tampa Bay, Florida, and adjacent waters, 1961-65. U.S. Fish Wildl. Serv., Data Rep. 34, i + 561 pp. on 9 microfiches.
- THORSON, GUNNAR.
1956. Marine level-bottom communities of recent seas, their temperature adaptation and their "balance" between predators and food animals. Trans. N.Y. Acad. Sci., Sect. 2, 18: 693-700.
- WAGNER, ROBERT J. L., and R. TUCKER ABBOTT (EDITORS).
1967. Van Nostrand's standard catalog of shells. 2d ed. D. Van Nostrand Co., Inc., Princeton, N.J., 303 pp.
- WARMKE, GERMAINE L., and R. TUCKER ABBOTT.
1962. Caribbean seashells. Livingston Publishing Co., Narberth, Penn., 348 pp.

APPENDIX

A Checklist Of Mollusks From Boca Ciega Bay, Florida

We identified 168 species of mollusks representing 69 families from BCF collections in Boca Ciega Bay. Of these, members of 156 species were alive. The number was increased to 72 families and 188 species by including mollusks recorded in other studies. The additions are coded within the list by surname initials of the authors who reported them: B. and B.—Bullock and Boss (see footnote 2); H.—Hutton et al. (1956); D. and K.—Dragovich and Kelly (1964). Mollusks not collected alive in this investigation are denoted by an asterisk (*). We identified some specimens only after comparison with specimens in the U.S. National Museum (+). Classifications are based on Abbott (1954, 1968) and Warmke and Abbott (1962).

CLASS GASTROPODA

Fissurellidae

Diodora cayenensis (Lamarck)

Trochidae

**Calliostoma jujubinum tampaense* (Conrad)

Turbinidae

Arene tricarinata (Stearns)

Turbo castaneus (Gmelin)

Neritidae

**Neritina reclinata* (Say)

Melanellidae

- Melanella bilineata* (Alder)
- Melanella intermedia* (Cantraine)

Epitonidae

- Epitonium angulatum* (Say)
- Epitonium hypnypresi* (Kiener)
- Epitonium rupicola* (Kurts)

Rissoidae

- Rissoina chesneli* (Michaud)

Vitrinellidae

- Cyclostremiscus beawi* Fisher—further verification pending.
- Cyclostremiscus suppressus* Dall—further verification pending.
- + *Teinostoma cryptospira* (no author on specimen)—further verification pending.

Truncatellidae

- **Truncatella pulchella* Pfeiffer

Turritellidae

- Vermicularia fargoii* Olsson

Caecidae

- Caecum cooperi* S. Smith
- Caecum pulchellum* Stimpson
- Meioceras nitidum* (Stimpson)

Modulidae

- Modulus modulus* (Linne)

Cerithiidae

- Bitium varium* (Pfeiffer)
- Cerithiopsis emersoni* (C. B. Adams)—B. and B.
- Cerithiopsis greeni* (C. B. Adams)
- Cerithium muscarum* Say
- Cerithium floridanum* Mörch
- Seila adamsi* (H. C. Lea)

Triphoridae

- Triphora nigrocincta* (C. B. Adams)

Potamididae

- Batillaria minima* (Gmelin)

Calyptraeidae

- Calyptraea centralis* Conrad
- Crepidula aculeata* (Gmelin)
- Crepidula fornicata* (Linne)
- Crepidula maculosa* Conrad
- Crepidula plana* Say

Strombidae

- Strombus alatus* Gmelin—B. and B.

Naticidae

- Natica pusilla* Say
- Polinices duplicatus* (Say)
- Sinum perspectivum* (Say)—B. and B.

Muricidae

- Eupleura sulcidentata* Dall
- Murex cellulusus* Conrad
- Murex pomum* Gmelin
- Thais haemastoma floridana* (Conrad)—B. and B.
- Urosalpinx perrugata* (Conrad)

Buccinidae

- Busycon contrarium* (Conrad)
- Busycon spiratum* (Lamarek)

Columbellidae

- + *Anachis semiplicata* Stearns
- Anachis obesa* (C. B. Adams)
- Anachis ostreicola* Sowerby
- Columbella rusticooides* Heilprin—B. and B.
- Mitrella lunata* (Say)

Melongenidae

- Melongena corona* (Gmelin)

Nassaridae

- Nassarius vibex* (Say)

Fasciolaridae

- Fasciolaria hunteria* (Perry)
- Fasciolaria tulipa* (Linne)
- Pleuroploca gigantea* (Kiener)—H.

Olividae

- Oliva sayana* Ravenel
- Olivella perplexa* Olsson
- Olivella mutica* (Say)
- Olivella floralia* Duclou

Marginellidae

- Bullata ovuliformis* (Orbigny)
- Hyalina avenacea* (Deshayes)
- Marginella aureocincta* Stearns
- Persicula lavalleana* (Orbigny)
- Prunum apicinum* (Menke)

Conidae

- Conus floridanus* Gabb—H.
- Conus jaspideus* Gmelin—H.
- **Conus stearnsi* Conrad

Terebridae

- Terebra concava vinosa* Dall
- Terebra dislocata* Say
- Terebra proteata* Conrad

Turridae

- **Glyphoturris rugirima* (Dall)
- **Monilispira leucocyma* (Dall)
- Pyrgocythara hemphilli* Bartsch and Rehder
- Stellatoma stellata* (Stearns)

Bullidae

- Bulla striata* Bruguière

Atyidae

- Haminoea antillarum* (Orbigny)
- Haminoea succinea* (Conrad)

Retusidae*Retusa canaliculata* (Say)**Pyramidellidae***Odostomia acutidens* Dall*Odostomia impressa* (Say)*Odostomia producta* Dall*Odostomia seminuda* C. B. Adams*Odostomia* sp.*Pyramidella crenulata* (Holmes)+ *Sayella hemphilli* (Dall)*Turbonilla conradi* Bush (Dall)*Turbonilla dalli* Bush**Acteocinidae***Cylichna bidentata* (Orbigny)**Acteonidae***Acteon punctostriatus* (C. B. Adams)**Aplysidae***Bursatella leachi plei* Rang**Ellobiidae***Melampus coffeus* (Linné)**CLASS AMPHINEURA****Ischnochitonidae***Chaetopleura apiculata* (Say)*Ischnochiton papillosus* (C. B. Adams)**Chitonidae***Chiton tuberculatus* Linné—H.**CLASS SCAPHOPODA****Dentaliidae***Dentalium eboreum* Conrad*Dentalium antillarum* Orbigny*Dentalium* sp. (resembles *D. texasianum* Philippi)**CLASS PELECYPODA****Solemyacidae***Solemya occidentalis* Deshayes**Nuculidae***Nucula proxima* Say**Nuculanidae***Nuculana acuta* Conrad**Arcidae***Anadara transversa* (Say)* *Arca zebra* Swainson*Barbatia cancellaria* (Lamarek)*Barbatia candida* (Helbling)—D. and K.*Noetia ponderosa* (Say)**Glycymerididae***Glycymeris pectinata* (Gmelin)**Pinnidae***Atrina rigida* Lightfoot**Mytilidae***Amygdalum papyria* (Conrad)*Brachidontes exustus* (Linné)*Lioberus castaneus* (Say)*Lithophaga bisulcata* (Orbigny)*Modiolus demissus* (Dillwyn)—H.*Modiolus demissus granosissimus* (Sowerby)—B. and B.*Modiolus americanus* (Leach)*Musculus lateralis* (Say)**Plicatulidae***Plicatula gibbosa* Lamarek**Ostreidae***Crassostrea virginica* (Gmelin)*Ostrea equestris* Say*Ostrea frons* Linné**Pectinidae***Aequipeecten irradians concentricus* (Say)**Anomidae***Anomia simplex* Orbigny**Crassatellidae***Crassinella lunulata* Conrad**Carditidae***Cardita floridana* Conrad*Venericardia perplana* (Conrad)*Venericardia tridentata* (Say)**Corbiculidae***Pseudocyrena floridana* (Conrad)—B. and B.**Erycinidae***Erycina floridana* Vanatta*Mysella planulata* (Stimpson)**Montacutidae***Montacuta floridana* Dall**Lucinidae*** *Anodontia alba* Link*Anodontia philippiana* (Reeve)*Codakia orbiculata* (Montagu)—B. and B.*Lucina multilineata* (Tuomey and Holmes)*Phacoides nassula* (Conrad)**Diplodontidae***Diplodonta punctata* (Say)**Chamidae*** *Chama congregata* Conrad**Cardiidae***Dinocardium robustum vanhyningi* (Clench and L. C. Smith)*Laevicardium mortoni* (Conrad)*Trachycardium egmontianum* (Shuttleworth)*Trachycardium muricatum* (Linné)

Veneridae

Anomalocardia cuneimeris (Conrad)
Chione cancellata (Linné)
Chione grus (Holmes)
Chione intapurpurea Conrad—B. and B.
Cyclinella tenuis (Recluz)
Dosinia discus (Reeve)
Dosinia elegans Conrad—B. and B.
Macrocallista nimboza Lightfoot
Mercenaria campechiensis (Gmelin)
Parastarte triquetra (Conrad)
Pitar fulminata (Menke)
Pitar simpsoni (Dall)
**Transennella conradina* Dall
**Transennella simpsoni* Dall

Petricolidae

Petricola pholadiformis Lamarck

Mactridae

Anatina plicatella (Lamarck)
Mactra fragilis Gmelin
Mulinia lateralis (Say)
Spisula solidissima (Dillwyn)—B. and B.

Tellinidae

Psanmotreta intastriata (Say)—B. and B.
Macoma constricta (Bruguère)
Tellidora cristata Recluz
Tellina aequistriata Say
Tellina alternata Say
Tellina lineata Turton
Tellina mera Say
Tellina tampaensis Conrad
Tellina sp. (resembles *T. texana* Dall)
Tellina versicolor DeKay

Semellidae

Abra aequalis (Say)
Cumingia tellinoides (Conrad)
Semele bellastrata (Conrad)
Semele proficua (Pulteney)
Semele purpurascens (Gmelin)

Donacidae

Donax variabilis Say

Sanguinolaridae

Tagelus divisus Spengler
Tagelus plebeius (Lightfoot)

Solenidae

Ensis minor Dall

Corbulidae

Corbula barrattiana C. B. Adams
Corbula caribaea Orbigny
Corbula contracta Say

Pholadidae

Cyrtoplcura costata (Linné)
Martesia cuneiformis Say

Pandoridae

**Pandora trilineata* Say
Pandora bushiana Dall

Lyonsiidae

Lyonsia hyalina floridana (Conrad)

Periplomatidae

Periploma sp.