

HELMINTH PARASITES OF PORPOISES OF THE GENUS  
*STENELLA* IN THE EASTERN TROPICAL PACIFIC,  
WITH DESCRIPTIONS OF TWO NEW SPECIES:  
*MASTIGONEMA STENELLAE* GEN. ET SP. N.  
(NEMATODA: SPIRUROIDEA) AND *ZALOPHOTREMA*  
*PACIFICUM* SP. N. (TREMATODA: DIGENEA)

MURRAY D. DAILEY<sup>1</sup> AND WILLIAM F. PERRIN<sup>2</sup>

ABSTRACT

Parasite frequencies in 72 spotted porpoise, *Stenella graffmani* (Lönningberg, 1934), and 19 spinner porpoise, *S. cf. S. longirostris* (Gray, 1828), are reported and analyzed with respect to age of host. In addition to the new nematode *Mastigonema stenellae* gen. et sp. n. from the fore stomach of *S. graffmani* and the new fluke *Zalophotrema pacificum* sp. n. from the pancreatic duct of *S. graffmani* and *S. cf. S. longirostris*, parasites reported include *Anisakis simplex*, *Halocercus delphini*, *Crassicauda* sp., *Campula rochebruni*, *Tetrabothrium forsteri*, *Strobilocephalus triangularis*, *Phyllobothrium delphini*, and *Mono-rygma grimaldii*. Irreversible lesions in the ventral region of the skull due to infection of the air sinuses by a *Crassicauda*-like nematode were more frequent in skulls of calves of *S. graffmani* than in skulls of subadults and adults (129 specimens total), indicating that the parasite, or a condition correlated with its occurrence, is a significant factor in natural mortality of the spotted porpoise in the eastern tropical Pacific.

Porpoises of the genus *Stenella* are of great importance to the U.S. tropical tuna fishery (Perrin, 1969, 1970a). Yellowfin tuna associate with porpoise schools in the eastern Pacific, and the fishermen make use of this association in locating and capturing the tuna. Despite this economic importance, little is known of the ecology of the porpoises. Parasites have been previously reported from *Stenella longirostris*, *S. roseiventris*, *S. coeruleoalba*, *S. attenuata*, and *S. graffmani* (Dailey and Brownell, 1972). These reports have usually been from a small sample and did not include numbers of parasites per animal or percentage of sample infected. This report deals with two members of the genus in the eastern tropical Pacific: the spotted porpoise and the spinner porpoise. The taxonomy of the genus is currently in a chaotic state. Spinner porpoises from the eastern Pacific have been referred by recent authors to *Stenella microps* (Gray, 1846) and to *S.*

*longirostris* (Gray, 1828)—Perrin and Hunter (1972). The spotted porpoise *S. graffmani* (Lönningberg, 1934) was described from Acapulco, but Rice and Scheffer (1968) recently referred all spotted porpoises to *S. dubia* (G. Cuvier, 1812), a decision that does not reflect a consensus of current opinion (Perrin, 1970b; Mitchell, 1970; Caldwell et al., 1971). We here refer the spotted porpoise examined to *S. graffmani* and the spinner porpoise tentatively to *S. longirostris*. The "eastern" and "whitebelly" spinner porpoises (Perrin, 1972) are considered to be geographical variants within one species and are not treated separately. This usage is provisional pending the results of current taxonomic investigations.

MATERIALS AND METHODS

We examined 72 specimens of *S. graffmani* and 19 of *S. cf. S. longirostris* (11 eastern and 8 whitebelly). These were collected during the course of investigations of morphology and reproduction. We collected parasites as we prepared skeletons of the animals for use

<sup>1</sup> Department of Biology, California State University at Long Beach, Long Beach, CA 90840.

<sup>2</sup> Southwest Fisheries Center, National Marine Fisheries Service, NOAA, La Jolla, CA 92037.

as museum specimens. Capture data and museum numbers are presented in Appendix Tables 1 and 2. Also, we examined 67 skulls of spotted porpoise and 29 of spinner porpoise from museum collections, for bone damage caused by nematode infection. The museum specimens were from the eastern tropical Pacific and are listed below. Abbreviations used are: AMNH, American Museum of Natural History, New York; LACM, Los Angeles County Museum; SDMNH, San Diego Museum of Natural History; USNM, United States National Museum, Washington, D.C.

*S. graffmani*: AMNH130216, LACM27412, 27413, 27419, 27420, 27421, 27422, 27425, 27426, 27427, 27428, 27429, 27431, 27434, 31362, 31431; SDMNH20637, 21199; USNM-261427, 261428, 261429, 261430, 261431, 261432, 261433, 261434, 395264, 395265, 395266, 395267, 395268, 395615, 395616, 395617, 395618, 395926, 395927, 395928, 395929, 395935, 395936, 395937, 395938, 396027, 396029, 396034, 396037. The following LACM specimens, identified by field numbers, had no museum numbers when examined: GDF42, GDF57, MI563, RLB404, RLB500, WAW42, WAW51, WAW55. The following specimens, identified by field numbers are in the research collection at the Southwest Fisheries Center (SWFC below), La Jolla, Calif: CV160, JSL88, JSL89, JSL90, JSL95, RICE65-1, RICE66-3, WFP7, WPSG-01, WPSG02, WPSG03.

*S. cf. S. longirostris*: LACM27414, 27423, 27424, 27428, 27430, 27432, 27433; SDMNH-21200, 21427; USNM395269, 395270, 395271, 395272, 395273, 395274, 395275, 395930, 395931, 395932, 395933, 395934, 396030, 396032, 396035; LACM field numbers GDF56, RLB238, RLB405, WAW43, WAW50, WAW-52, WAW53, WAW54, WAW58, WAW60; SWFC field numbers CV240, CV241, CV245, RICE65-149, RICE67-102.

While rough-cleaning the porpoise carcasses for skeletal preparation, we examined the blubber, mesenteries, peritoneum, musculature, blowhole region, maxillary sinuses, mammary glands, and viscera (esophagus, stomachs, heart, lungs, liver, spleen, kidneys, gonads, intestine)

for larval and adult parasites. We searched 3-m lengths of the intestine at the duodenal end, mid-length, and posterior end. Counts of cysts in the blubber should be considered as relative indices; we did not examine all the blubber carefully. We filled out a data form for each carcass indicating precisely which areas had been checked for parasites. All worms and cysts were fixed in Lavdowsky's (Formalin<sup>3</sup>-Acetic acid-Alcohol) or Bouin's fluid and stored in 70% ethanol. Cestode, trematode, and acanthocephalan whole mounts were stained with celestine blue B or Semichon's carmine. Specimens were sectioned singly or in situ at 6, 8, and 10  $\mu$  and stained with Mayer's hematoxylin and eosin Y or Mallory's trichrome. All material except nematode whole mounts were dehydrated in ethanol, cleared in xylene, and mounted in Piccolyte. Nematodes were cleared and mounted in glycerine. Drawings were made with the aid of a drawing tube. Measurements are given in millimeters unless otherwise stated. Average measurements are given with ranges in parentheses.

## THE PARASITES

Counts of parasites collected are tabulated in Appendix Tables 1 and 2.

### Nematoda

*Anisakis simplex* (Rudolphi, 1809, det. Krabbe, 1878)

This nematode was the most frequent parasite collected during this study. The taxonomic criteria used in identification of these specimens were those presented by Davey (1971) in his revision of the genus *Anisakis*. *A. simplex* was found in the stomachs of both host species and represents a new host report for both *S. graffmani* and *S. longirostris*. Anisakids have been previously reported from other species of *Stenella* (Delyamure, 1955; Kagei et al., 1967; Zam, Caldwell, and Caldwell, 1971; Dailey and Brownell, 1972).

<sup>3</sup>Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

*Halocercus delphini* Baylis and  
Daubney, 1925

This parasite has been reported previously from the genus *Stenella* by Zam et al. (1971). They found three of 17 *S. plagiodon* (Cope, 1866) taken off the Florida coast infected with *H. delphini*. This report represents a new host record from *S. graffmani* and *S. longirostris*. Infections occurred in bunches with the anterior ends of the parasites embedded in capsules found outside of the bronchial walls (Figure 1). The body of each parasite lies in the bronchiole oriented in an anterior direction, never towards the apex of the lobe, similar to the orientation described for *Skrjabinalius cryptocephalus* (see Delyamure, 1955).

*Mastigonema stenellae* gen. et sp. n.  
(Nematoda: Spiruridae)

This nematode is described from three female specimens taken from the forestomachs of two *S. graffmani* (both recovered prior to present study) and one *S. longirostris*.

*Mastigonema* gen. n. (Ascaropsinae)

Mouth with two unlobed lips. Cervical cuticle not inflated or flanged. Buccal capsule cylindrical, with few connecting, annular, thickenings. Esophagus long, not divided into anterior and posterior portions; valve of typical spirurid form. Females prodelphic, ovaries post-esophageal. Vulva in middle third of body. Cuticle transversely striated. Oviparous. Parasites of Cetacea.

*Mastigonema stenellae* sp. n. (Figures 2-6)

With characters of the genus: body whip-shaped, with thin anterior and enlarged posterior portions. Stoma triangular; two large lips. Buccal capsule reinforced with connecting rings, cervical papillae present, eggs with enlarged cap.

Female: 25.5 (23.5-28.5) long by 1.2 in maximum width, 7.0 from posterior end. Buccal capsule 0.054 long, tapering from 0.045 wide at anterior to 0.032 at esophageal junction.





FIGURE 1.—*Halocercus delphini* with anterior end embedded in capsule. (X7)

Nerve ring 0.281 from anterior end, surrounding esophagus. Esophagus 8.5-11.7 long, widening at approximately 2.59 from anterior end. Ovaries directed anteriorly then reflexing sharply. Vulva muscular, 16.64 from anterior, vagina vera 0.13 long by 0.086 wide, vagina uterina 0.81 long by 0.11 wide. Eggs 0.034 long by 0.032 wide, with swollen cap slightly larger than bottom portion. Anal lip large, anus 0.22 from posterior tip. Tail conical with large mucron.

Hosts: *Stenella graffmani*, *S. longirostris*.  
Location: Forestomach.

Locality: Eastern tropical Pacific Ocean;  
holotype from *S. longirostris* collected  
at approximately lat 8°N, long 107°W.  
Type specimens: USNM Helminth Collection  
holotype female No. 72159, paratype  
No. 72160.

Remarks: A search of the literature indicates  
this is the first report of a spirurid



FIGURES 2-6.—*Mastigonema stenellae* sp. n.: 2. Entire worm. 3. Cephalic region, showing annular thickenings. 4. Esophageal-intestinal valve. 5. Caudal region. 6. Eggs, showing enlarged caps.

nematode parasitizing a marine mammal. The general body shape of this nematode is similar to that seen in *Vasorhabdochona cablei* Martin and Zam, 1967, *Trichospirura leptostoma* Smith and Chitwood, 1967, and *Freitasia teixeirai* Baruš and Coy, 1965. However, the structure of lips and buccal capsule place it in the subfamily Ascaropsinae rather than with the rhabdochonids. Of the genera in this subfamily it most closely resembles the genus *Pereirai* Cuocolo, 1943. It differs from *Pereirai* in the following characteristics; lips are unlobed, cuticular flange is lacking, buccal capsule has

few annular thickenings, esophagus not divided into anterior and posterior portions, and vulva in posterior half of body.

#### Crassicaudidae Skrjabin et Andrews, 1934

*Crassicauda* sp. A single infection of this tissue nematode was found in the 78 animals examined (1.3%). The worms were in the abdominal muscle adjacent to the mammary gland of *S. longirostris*. A large abdominal cyst was also found in the same area. Sections revealed large numbers of *Crassicauda* sp. eggs as well as portions of the female worm (Figure 7). Entire worms were not recovered so specific identification was not possible. However, egg




FIGURE 7.—Section through abdominal cyst with *Crassicauda* eggs. (X150)

measurements resemble those given for *C. grampicola* by Johnston and Mawson (1941) from Risso's porpoise, *Grampus griseus* (Cuvier, 1812), taken at New South Wales, Australia.

Another single specimen with *Crassicauda* affinities was recovered by James G. Mead, University of Chicago, Department of Anatomy, while dissecting the head of a *S. graffmani*. The worm was taken from the pterygoid air sinus. The pterygoid bone was eroded in the region in which the specimen was found. We assume that extensive bone damage found by us in other specimens we examined (Figure 8) was also caused by infestations of this nematode, although, judging from the published literature, the possibility cannot be dismissed that more than one species is involved. Reysenbach de Haan (1957) reported *Stenurus globicephalae* Baylis and Daubney, 1952, in the sinuses and cava tympani of the pilot whale, *Globicephala melaena* (Traill, 1809). Norris and Prescott (1961) collected "large quantities of a nematode (*Stenurus* sp.)" from the lateral air sinus of a dall porpoise, *Phocoenoides dalli* (True, 1885). Tomilin and Smyshlyayev (1968) reported that a "nematode *Stenurus minor* attacks the hearing organs of the Black Sea porpoise *Phocaena* and by clogging the cavity around the earbone may kill its host or severely complicate its existence." Brodie (1971) reported *Crassicauda* sp. as a middle ear parasite of the white whale, *Delphinapterus leucas* Pallas, 1776, in Cumberland Sound, Baffin Island. Yamada (1956) examined skulls of approximately 80 false killer whales, *Pseudorca crassidens*, from a stranded group of approximately 150 and found "basket-like" lesions in the pterygoid region of 40 skulls. He ascribed the damage to parasitic infection, "most probably by nematodes," and stated that such damage is a common occurrence in *Grampus* and *Globicephala* in Japanese waters.

## Trematoda

The trematodes recovered during this study are represented by two genera known from marine mammals and three specimens of one genus that has been found in teleost hosts.

### *Zalophotrema pacificum* sp. n. (Figures 9 and 10)

Description: Based on measurements from 26 specimens. With characteristics of the genus. Body elongate. Dorsoventrally flattened, 10 mm (8.3-12.6) long by 1.7 mm (1.4-2.0) wide. Cuticle spinose. Oral sucker 535  $\mu$  (398-592) long by 575  $\mu$  (439-673) wide; opening slightly sub-terminal. Pharynx pyriform, 494  $\mu$  (398-541) long by 314  $\mu$  (255-367) wide. Intestinal cecum bifurcates immediately caudad to pharynx with each lateral branch dividing into short anterior and long posterior portions. Both anterior and posterior cecal branches contain numerous diverticula; anterior branch terminating lateral to posterior edge of oral sucker, posterior branch terminating near posterior end of body. Excretory pore terminal. Acetabulum 600  $\mu$  (520-710) in diameter, located approximately mid-body. Cirrus unarmed. Testes large, lobed, in tandem and extending posterior to equatorial third of body. Ovary pretesticular, lobed, 333  $\mu$  (235-418) long by 477  $\mu$  (310-765) wide. Mehlis's gland posterior and dorsal to ovary. Uterus coiled, preovarian. Genital pore just preacetabular. Vitellaria follicular, occupying entire anterior half of body from level of genital pore to shoulder region of ceca, extending posterior from genital pore in lateral bands to posterior end of body. Eggs, oblong, operculate, round in cross-section, mean measurements of 20 eggs, 61  $\mu$  long by 42  $\mu$  wide.

Hosts: *Stenella longirostris*, *Stenella graffmani*.

Location: Pancreatic duct.

Locality: Eastern tropical Pacific Ocean; holotype from *S. graffmani* collected at lat 12°5'N, long 93°18'W.

Type specimen: USNM Helm. Coll. No. 72158.

Remarks: Currently there are three species in the genus *Zalophotrema* (Stunkard and Alvey, 1929): *Z. hepaticum* Stunkard and Alvey, 1929, described from the bile ducts of a California sea lion, *Zalophus californianus* (Lesson, 1828), at the New York Aquarium; *Z. curilensis* Gubanov, 1952, from the bile ducts of





FIGURE 8.—Ventral view of skull of spotted porpoise, *Stenella graffmani*, showing bone lesions (arrows).



a sperm whale, *Physeter catodon* L; and *Z. lubimowi* Petrov and Chertkova, 1963, from the liver of a southern sea lion, *Otaria byronia* (Blainville, 1820) at the Moscow Zoo. *Zalophotrema pacificum* sp. n. can be easily separated from the three previously described species. *Z. lubimowi* differs from all other species in the genus in that it does not have confluent vitellarian fields anteriorly. *Z. curilensis* differs from *Z. pacificum* by larger body size (18-22.5 mm long by 5-7 mm wide) as well as the acetabulum being larger than the oral sucker. *Z. pacificum* sp. n. is most similar to *Z. hepaticum* in general body shape but differs in the following characteristics: body width (*Z. hepaticum* 3.0-3.6 mm wide), sucker ratio (in *Z. hepaticum* oral sucker almost twice as large as acetabulum while in *Z. pacificum* sp. n. suckers are approximately same size), ratio of distance between suckers to body length (*Z. hepaticum* 1:13, *Z. pacificum* sp. n. 1:2.5), distribution (percentage) of preacetabular vitellaria (*Z. hepaticum* nearly twice as large as those of *Z. pacificum* sp. n.), and egg size (75  $\mu$  long by 50  $\mu$  wide in *Z. hepaticum*).

*Campula rochebruni* (Poirier, 1886)  
Bittner et Sprehn, 1928

This parasite was recovered from the stomachs and hepatopancreatic ducts of three (two *S. graffmani* and one *S. longirostris*) of 78 (3.9%) animals examined. This is a new host record for this parasite and represents the first report from the genus *Stenella*. *C. rochebruni* was originally described from *Delphinus delphis* L. in European waters.

Incidental Trematodes

Three specimens of the genus *Hirudinella*

FIGURES 9-10.—*Zalophotrema pacificum* sp. n. 9. Entire worm, ventral view; 10. egg. Abbreviations used: a - acetabulum; gp - genital pore; o - ovary; os - oral sucker; p - pharynx; t - testis; u - uterus.



Garsin, 1730, were recovered from the fore-stomachs of three hosts (two *S. longirostris* and one *S. graffmani*). Members of this genus are commonly parasites of teleost fishes and were probably ingested with a meal of fish and not established parasites of these hosts.

### Cestoda

*Tetrabotrium forsteri* (Kreff, 1871)  
Fuhrmann, 1904

This cestode was recovered from the intestine of over 50% of the examined hosts and occurred in both *S. graffmani* and *S. longirostris*. This is the first report of this parasite from either of these hosts. *T. forsteri* (Kreff, 1871) was first described by Krefft (1871) under the name of *Taenia forsteri* from *Delphinus delphis* L. (= *D. forsteri*) caught off the coast of New Zealand and later transferred to the genus *Tetrabotrium* by Fuhrmann (1904). A complete review of the literature on this species is given by Delyamure (1955).

*Strobilocephalus triangularis* (Diesing, 1850)  
Baer, 1932

This tetrabotriid has a complex taxonomic history that is covered by Delyamure (1955). The parasite apparently anchors its scolex in the colon wall when small and grows inside a fibrotic capsule formed by the host (Figures 11 and 12). This is the first report of this parasite from the genus *Stenella*. It has been reported from the Pacific Ocean by Baer (1955), who described material taken from an unidentified dolphin captured in the waters of the Marshall Islands. There are numerous reports of the parasite from Atlantic Ocean hosts (*Hyperoodon ampullatus*) (Forster, 1770) (= *H. rostratus*), *Mesoplodon bidens* (Sowerby, 1804), *Lagenorhynchus acutus* (Gray, 1828), *Steno bredanensis* Lesson, 1828 (= *S. frontatus*), *Delphinus* sp. (Delyamure, 1955).

*Phyllobotrium delphini* (Bosc, 1802) and  
*Monorygma grimaldii* (Moniez, 1881) Baylis, 1919

These larval cestodes were recovered from




FIGURE 11.—Section through the capsule and scolex of *Strobilocephalus triangularis* from the rectum of *S. graffmani*. (X10.5)




FIGURE 12.—*Strobilocephalus triangularis* attached in rectum of *Stenella graffmani*. (X1)

a large proportion of the animals examined. The history of these worms has been discussed by Delyamure (1955), Dollfus (1964), Williams (1968), and Dailey and Brownell (1972). *P. delphini* was always found in the blubber and *M. grimaldii* in the mesenteries.

### PARASITE LOAD RELATIVE TO AGE OF HOST

The availability of data for a large sample of porpoise specimens, especially of *S. graffmani*, dissected in relatively uniform manner provides an unique opportunity to examine infection relative to age of the host for certain of the more frequently encountered parasite species. We discuss only the parasite species for which patterns of infestation are apparent in the data.

#### Spotted Porpoise, *Stenella graffmani*

The sample of *S. graffmani* examined may be

divided into four age classes on the basis of total length, the developmental pattern categories defined by Perrin (1970b) and degree of sexual maturity (Figure 13). Females were classified as adult by the presence of corpora of ovulation in the ovaries and males by presence of spermatogenesis in the seminiferous tubules, as determined by histological examination. Sixty-one of the 72 specimens were collected during a 1-month period, and the first three modes probably represent discrete reproductive cohorts. Analysis below of parasitism relative to age is in terms of the four age classes, except as noted in the case of the *Crassicauda*-like nematode in the air sinuses.

The percent infection for *Anisakis* in *S. graffmani* shows increase with age of the host (Figure 14), but very large numbers of worms were found only in subadults, and the parasite number frequency distribution for that class is obviously skewed to the right. The inference to be drawn is that either some reduction in number of worms occurs in a heavily infected porpoise as it matures, due to the life cycle




FIGURE 13.—Frequency distributions of total length, by age class, of spotted porpoise, *Stenella graffmani*, included in sample examined for parasites.

of the parasite and/or to possibly differential feeding habits of young and mature porpoise, or heavily infected individuals die before attaining maturity.

The case of the *Crassicauda*-like parasite in the air sinuses of *S. graffmani* certainly provides the most dramatic instance of infection varying with age. As stated above, the sinuses were not searched during dissection of most of the porpoise specimens, and only one specimen of the *Crassicauda*-type was recovered. Direct evidence, however, is provided of heavy infection by the areas of bone erosion on the lower surfaces of the skull (as illustrated in Figure 8). Given this circumstance, we expanded the series of porpoise specimens examined to include skeletal specimens in various museum collections. In all, 129 skulls of spotted porpoise from the eastern tropical Pacific were examined (see Materials and Methods). Since correlated data on size, reproductive condition, and color pattern were not available for most of the museum specimens examined, it was necessary to use other criteria of age for those specimens. The fetal and neonatal skulls examined were all from specimens we dissected.




FIGURE 14.—Frequency distributions of levels of infection by *Anisakis* in four age classes of *Stenella graffmani*.

Those of the remaining skulls that were museum specimens were relegated to the three categories of calves, subadults, and adults by criteria of cranial features, using the sample of skulls from specimens we dissected to correlate these features with total length, sexual maturity, and color pattern. Museum skulls were assigned to the calf class when in condylobasal length, telescoping of facial elements, and closure of occipital sutures, they had not reached the adult condition. Skulls exhibiting adult length, telescoping, and occipital closure were divided into adults and subadults by the criterion of presence or absence of distal fusion of the premaxillary and maxillary bones of the rostrum. Rough correlation of the onset of fusion with attainment of sexual maturity was found among the skulls from specimens for which external data were also available (Table 1. Errors in classifying subadult skulls as adult and vice versa are approximately equal. Results of examination of the 129 skulls for bone damage (Table 2) demonstrate clearly that the rate of infection is very much higher in

TABLE 1.—Incidence of distal fusion of pmx (pre-maxillary) and mx (maxillary) bones in specimens of *Stenella graffmani*.

| Age class <sup>1</sup> | Examined | Pmx and mx fused | Error in classification |
|------------------------|----------|------------------|-------------------------|
| Subadults              | 31       | 6                | 19.4%                   |
| Adult                  | 21       | 18               | 14.3%                   |

<sup>1</sup> As determined by criterion of total length, color pattern, and sexual maturity.

TABLE 2.—Lesions of the ventral side of the skull in *Stenella graffmani*, presumably due to infection by a *Crassicauda*-like nematode.

| Age class                      | Range of condylobasal length (mm) | Examined | Lesions present |
|--------------------------------|-----------------------------------|----------|-----------------|
| Nearterm fetuses and neonatals | 209-236                           | 10       | 0               |
| Calves                         | 302-359                           | 18       | 8 (44.4%)       |
| Subadults                      | 356-446                           | 45       | 5 (11.1%)       |
| Adults                         | 360-466                           | 56       | 4 (7.1%)        |
| Total                          |                                   | 129      | 17 (13.2%)      |

calves than in subadults or adults. What is more, the most severely damaged skulls were those of calves. Such extensive damage is certainly irreversible, in that if repaired, traces would still be apparent in the form of bone callouses, depressions, or altered shape of bone elements. No such traces were found in any of the adult skulls examined, although we searched for them carefully. We conclude, therefore, that heavy damage is indicative of a probably morbid condition and that parasitism or a correlated condition is a significant factor in natural mortality of the spotted porpoise in the eastern tropical Pacific.

The patterns of infection relative to age by cysts of the cestodes *Monorygma* and *Phyllobothrium* (Figure 15) closely resemble each other, with little or no occurrence before subadulthood and 100% occurrence and greatest numbers in adults. As these are larval forms, the cysts presumably would have to be ingested by predators of the porpoise in order for the life cycles of the parasites to progress. Killer whales, *Orcinus orca*, have been observed by Perrin to prey on spotted porpoise, and sharks of several species have been observed to feed on dead and injured spotted porpoise during tuna fishing operations (Leatherwood, Perrin,

Garvie, and La Grange, manuscript).<sup>4</sup> Killer whales and/or sharks are therefore possible candidates for terminal hosts of these two cestodes, although adult Tetracystidae have only been found in elasmobranchs. However, a life cycle involving a warm blooded intermediate host serving a cold blooded definitive host has not been previously demonstrated.

The two nonlarval cestodes, *Tetrabothrium forsteri* (Figure 16) and *Strobilocephalus triangularis* (Table 3), exhibit different patterns of occurrence, more like those of the nematodes, occurring with greatest numbers and frequencies in calves in the case of *T. forsteri* and in calves and subadults in the case of *S. triangularis*. This was especially striking for *S. triangularis*. In the two calves and two subadults with extremely heavy infections, the last meter of the intestine was all but blocked and swollen to three or four times its normal diameter. No adult examined was found to be infected with *S. triangularis*.

### Spinner Porpoise, *Stenella* cf. *S. longirostris*

The smaller sample of spinner porpoise was divided into three age classes on the basis of length and sexual maturity, with subadults again defined as immature animals of adult size. Despite the small number of data for the spinner (Appendix Table 2), some differences are apparent between the patterns of infection for this species and for *S. graffmani*. For *Anisakis*, the specimen with the greatest number of worms (CV285) was a large adult female, which contrasts with the finding for *S. graffmani*, in which heaviest infections occurred in subadults. Bone erosion suggesting severe infection of the ventral air sinuses by a nematode was encountered in only 1 skull (of a subadult) of 53 examined. The parasite specimen was not recovered, precluding identification, but the damage was similar to that caused by the *Crassicauda*-like nematode in *S. graffmani*.

<sup>4</sup> Leatherwood, J. S., W. F. Perrin, R. L. Garvie, and J. C. La Grange. Observations of sharks attacking porpoise (*Stenella* spp.). (Unpubl. manusc.)




FIGURE 15.—Frequency distributions of levels of infection by cysts of *Monorygma grimaldii* and *Phyllobothrium delphini* in four age classes of *Stenella graffmani*.

Infection by cestode cysts showed the same pattern as in *S. graffmani*, except that one adult specimen was found to be free of *Phyllobothrium* cysts. Also, as in *S. graffmani*, calves were most heavily infected with *T. forsteri*, and *Strobilocephalus triangularis* was encountered only in an immature specimen, a calf.

### ACKNOWLEDGMENTS

The authors would like to express their appreciation to Gerald Schmidt, University of Northern Colorado; John Schacher, University of California, Los Angeles; Mary Pritchard, University of Nebraska; May Belle Chitwood,

University of California, Davis; Kenneth S. Norris, University of California, Los Angeles; and Leo Margolis, Fisheries Research Board of Canada, Nanaimo, for their help and advice during the study. We would also like to give special thanks to Barry L. Hill, Lorraine G. Peterson, and Kenneth S. Raymond, for their valuable technical contributions. Randolph Jose, Ralston-Purina Corporation; Craig J. Orange, Inter-American Tropical Tuna Commission; and the captains and crews of the tuna vessels *Carol Virginia*, *Pacific Queen*, *Conquest*, *Marietta*, and *Queen Mary* helped us obtain porpoise for dissection. Richard G. Van Gelder, American Museum of Natural History; Donald R. Patten,




FIGURE 16.—Frequency distributions of levels of infection by *Tetrabothrium forsteri* in four age classes of *Stenella graffmani*.

TABLE 3.—*Strobilocephalus triangularis* in *Stenella graffmani*.

| Age class | Examined | Infected  |
|-----------|----------|-----------|
| Neontals  | 5        | 0         |
| Calves    | 14       | 3 (21.4%) |
| Subadults | 26       | 3 (11.5%) |
| Adults    | 23       | 0         |
| Total     | 68       | 6 ( 8.8%) |

Los Angeles County Museum; Joseph R. Jehl, Jr., San Diego Museum of Natural History; and Charles O. Handley, Jr., U.S. National Museum, kindly made materials in their charge available for study.

## LITERATURE CITED

- BAER, J. G.  
1955. Cestodes d'un dauphin de L'océan pacifique. Bull. Soc. Neuchatel. Sci. Nat., Ser. 3, 78:33-36.
- BARUŠ, V., AND O. COY.  
1965. *Freitasia teixeirai* gen. n. et sp. n. and other nematodes parasitizing *Anolis equestris* (Squamata: Iguanidae). Folia Parasitol. (Praha) 15:41-54.
- BRODIE, P. F.  
1971. A reconsideration of aspects of growth, reproduction, and behavior of the white whale (*Delphinapterus leucas*), with reference to the Cumberland Sound, Baffin Island, population. J. Fish. Res. Board Can. 28:1309-1318.
- CALDWELL, D. K., M. C. CALDWELL, W. F. RATHJEN, AND J. R. SULLIVAN.  
1971. Cetaceans from the Lesser Antillean Island of St. Vincent. Fish. Bull., U.S. 69:303-312.
- DAILEY, M. D., AND R. L. BROWNELL.  
1972. A checklist of marine mammal parasites. In Mammals of the sea: Biology and medicine, p. 528-589. Charles C. Thomas, Springfield, Ill.
- DAVEY, J. T.  
1971. A revision of the genus *Anisakis* Dujardin, 1845 (Nematoda: Ascaridata). J. Helminthol. 45:51-72.
- DELYAMURE, S. L.  
1955. [The helminth fauna of marine mammals in the light of their ecology and phylogeny.] Izd. Akad. Nauk SSSR, Mosk., 517 p. (Translated by Israel Program Sci. Transl., 1968, 522 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, Va., as TT 67-51202).
- DOLLFUS, R. P.  
1964. A propos de la récolte, à Banyuls d'un cystique de Cestode chez *Tursiops truncatus* (Montagu, 1821). Les cystiques de Cestodes chez les Cétacés et Pinnipèdes. Vie Milieu, Suppl. 17:177-204.
- FUHRMANN, O.  
1904. Die Tetrabothrien der Säugetiere. Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Abt. I Orig. 35:744-752.
- JOHNSTON, T. H., AND P. M. MAWSON.  
1941. Nematodes from Australian marine mammals. Rec. S. Aust. Mus. Adelaide 6:429-434.
- KAGEL, N., T. OSHIMA, A. KOBAYASHI, M. KUMADA, T. KOYAMA, Y. KOMIYA, AND A. TAKEMURA.  
1967. Survey of *Anisakis* spp. (Anisakinae, Nematoda) on marine mammals in the coast of Japan. [In Japanese, English summ.] Jap. J. Parasitol. 16:427-435.
- KREFFT, G.  
1871. On Australian Entozoa, with descriptions of new species. Trans. Entom. Soc. N. S. W. 2:206-232.
- MARTIN, W., AND S. ZAM.  
1967. *Vasorhabdochona cablei*, gen. et sp. n. (Nematoda) from blood vessels of the marine fish, *Gillichthys mirabilis* Cooper. J. Parasitol. 53:389-391.
- MITCHELL, E.  
1970. Pigmentation pattern evolution in delphinid cetaceans: An essay in adaptive coloration. Can. J. Zool. 48:717-740.

- NORRIS, K. S., AND J. H. PRESCOTT.  
 1961. Observations on Pacific cetaceans of California and Mexican waters. Univ. Calif. Publ. Zool. 63:291-370, pl. 27-41.
- PERRIN, W. F.  
 1969. Using porpoise to catch tuna. World Fishing 18(6):42-45.  
 1970a. The problem of porpoise mortality in the U.S. tropical tuna fishery. Proc. Sixth Annu. Conf. Biol. Sonar Diving Mammals, p. 45-48.  
 1970b. Color pattern of the eastern Pacific spotted porpoise *Stenella graffmani* Lönnberg (Cetacea, Delphinidae). Zoologica (N.Y.) 54:135-149.  
 1972. Color patterns of spinner porpoises (*Stenella* cf. *S. longirostris*) of the eastern Pacific and Hawaii, with comments on delphinid pigmentation. Fish. Bull., U.S. 70:983-1003.
- PERRIN, W. F., AND J. R. HUNTER.  
 1972. Escape behavior of the Hawaiian spinner porpoise (*Stenella* cf. *S. longirostris*). Fish. Bull., U.S. 70:49-60.
- REYSENBACH DE HAAN, F. W.  
 1957. Hearing in whales. Acta Otolaryngol., Suppl. 134, 114 p.
- RICE, D. W., AND V. B. SCHEFFER.  
 1968. A list of the marine mammals of the world. U.S. Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 579, 16 p.
- SMITH, W., AND M. B. CHITWOOD.  
 1967. *Trichospirura leptostoma* gen. et sp. n. (Nematoda: Thelazioidea) from the pancreatic ducts of the white-eared marmoset *Callithrix jacchus*. J. Parasitol. 53:1270-1272.
- TOMILIN, A. G., AND M. I. SMYSHLYAYEV.  
 1968. [Some factors affecting whale mortality (diseases of cetaceans).] Byull. Mosk. O-va. Ispyt. Prir. Otd. Biol. 3:5-12. Transl. in Soviet Studies on Cetaceans, Joint Publ. Res. Serv. Transl. No. 49777, 1970, 19 p. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, Va.
- WILLIAMS, H. H.  
 1968. The taxonomy, ecology and host-specificity of some phyllobothriidae (Cestoda: Tetraphyllidae), a critical revision of *Phyllobothrium* Beneden, 1849 and comments on some allied genera. Philos. Trans. R. Soc. Lond., Ser. B, Biol. Sci. 253:231-307.
- YAMADA, M.  
 1956. An analysis in mass osteology of the false killer whale, *Pseudorca crassidens* (Owen). Part 1. Okajimas Folia Anat. Jap. 28:451-463.
- ZAM, S. G., D. K. CALDWELL, AND M. C. CALDWELL.  
 1971. Some endoparasites from small odontocete cetaceans collected in Florida and Georgia. Cetology 2:1-11.

APPENDIX TABLE 1.—Parasites in *Stenella graffmani*.

| Host         |   |  |             |     |                        | Parasites noted         |                            |                                     |                                      |                           |                               |                                      |                                |                            |                             |                          |
|--------------|---|--|-------------|-----|------------------------|-------------------------|----------------------------|-------------------------------------|--------------------------------------|---------------------------|-------------------------------|--------------------------------------|--------------------------------|----------------------------|-----------------------------|--------------------------|
| Field number | Museum number of skeletal specimen <sup>1</sup> | Collection number or locality <sup>2</sup> | Length (cm) | Sex | Age class <sup>3</sup> | <i>Anisakis simplex</i> | <i>Halocercus delphini</i> | <i>Mastigonema stenellae</i> sp. n. | <i>Zalophotrema pacificum</i> sp. n. | <i>Campula rochebruni</i> | <i>Tetrabothrium forsteri</i> | <i>Strobilocephalus triangulatus</i> | <i>Phyllobothrium delphini</i> | <i>Monoryzma grimaldii</i> | <i>Bolbosoma vasculosum</i> | <i>Bolbosoma balanae</i> |
| CV259        | USNM395276                                      | 6  | 1,676       | M   | 3                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 2                             | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV260        | USNM395277                                      | 6  | 1,750       | F   | 4                      | 1                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 1                              | 3                          | 0                           | 0                        |
| CV261        | USNM395278                                      | 2  | 1,657       | F   | 3                      | 1                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 0                              | 2                          | 1                           | 0                        |
| CV262        | USNM395279                                      | 2  | 1,646       | F   | 3                      | 20                      | —                          | 0                                   | 0                                    | 0                         | 1                             | 1                                    | 0                              | 0                          | 0                           | 0                        |
| CV263        | USNM395328                                      | 6  | 1,759       | M   | 3                      | 35                      | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 0                              | 5                          | 0                           | 0                        |
| CV264        | USNM395329                                      | 6  | 1,716       | M   | 3                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 2                             | 0                                    | 18                             | 7                          | 0                           | 0                        |
| CV265        | USNM395330                                      | 2  | 1,954       | M   | 4                      | 1                       | —                          | 0                                   | 0                                    | 0                         | 4                             | 0                                    | 20                             | 26                         | 0                           | 2                        |
| CV266        | USNM395331                                      | 2  | 1,681       | M   | 3                      | 1                       | —                          | 0                                   | 0                                    | 0                         | 0                             | +                                    | 0                              | 2                          | 0                           | 0                        |
| CV267        | USNM395332                                      | 2  | 1,787       | M   | 4                      | 2                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 18                             | 23                         | 0                           | 0                        |
| CV268        | USNM395333                                      | 2  | 1,900       | F   | 4                      | 4                       | —                          | 0                                   | 0                                    | 0                         | 2                             | 0                                    | —                              | 27                         | 0                           | 0                        |
| CV269        | USNM395334                                      | 2  | 1,837       | F   | 4                      | 2                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | —                              | 14                         | 0                           | 0                        |
| CV270        | USNM395336                                      | 2  | 1,815       | F   | 4                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 1                             | 0                                    | 5                              | 52                         | 0                           | 1                        |
| CV271        | USNM395337                                      | 2  | 1,821       | F   | 4                      | 8                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | —                              | 13                         | 0                           | 0                        |
| CV272        | USNM395338                                      | 2  | 1,850       | F   | 4                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 2                             | 0                                    | 25                             | 45                         | 0                           | 1                        |
| CV273        | USNM395339                                      | 2  | 1,630       | M   | 3                      | 61                      | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 0                              | 1                          | 0                           | 1                        |
| CV274        | USNM395385                                      | 6  | 1,782       | M   | 3                      | 0                       | —                          | 0                                   | 0                                    | 0                         | +                             | 0                                    | 0                              | 5                          | 0                           | 0                        |
| CV275        | USNM395386                                      | 6  | 1,570       | M   | 3                      | 2                       | —                          | 0                                   | 0                                    | 0                         | 1                             | 0                                    | 8                              | 3                          | 0                           | 0                        |
| CV276        | USNM395387                                      | 6  | 1,578       | M   | 3                      | 16                      | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 0                              | 1                          | 0                           | 0                        |
| CV277        | USNM395388                                      | 2  | 1,816       | F   | 4                      | 8                       | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 5                              | 2                          | 0                           | 1                        |
| CV278        | USNM395389                                      | 6  | 1,632       | M   | 3                      | 18                      | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                    | 2                              | 0                          | 0                           | 0                        |
| CV279        | USNM395390                                      | 6  | 2,183       | M   | 4                      | 8                       | —                          | 0                                   | 0                                    | 0                         | 5                             | 0                                    | 27                             | 50                         | 2                           | 0                        |
| CV280        | USNM395391                                      | 6  | 2,032       | M   | 4                      | 14                      | —                          | 0                                   | 0                                    | 0                         | 5                             | 0                                    | 32                             | 14                         | 0                           | 0                        |

APPENDIX TABLE 1.—Parasites in *Stenella graffmani*.—Continued.

| Host         |   |  |             |     |                        | Parasites noted         |                            |                                     |                                      |                           |                                  |                                      |                                |                            |                             |                          |
|--------------|---|--|-------------|-----|------------------------|-------------------------|----------------------------|-------------------------------------|--------------------------------------|---------------------------|----------------------------------|--------------------------------------|--------------------------------|----------------------------|-----------------------------|--------------------------|
| Field number | Museum number of skeletal specimen <sup>1</sup> | Collection number or locality <sup>2</sup> | Length (cm) | Sex | Age class <sup>3</sup> | <i>Anisakis simplex</i> | <i>Halocercus delphini</i> | <i>Mastigonema stenellae</i> sp. n. | <i>Zalophotrema pacificum</i> sp. n. | <i>Campula rochebruni</i> | <i>Tetrahelminthium forsteri</i> | <i>Sirobilocephalus triangulatus</i> | <i>Phyllobothrium delphini</i> | <i>Monorygma grimaldii</i> | <i>Bolbosoma vasculosum</i> | <i>Bolbosoma batanae</i> |
| CV281        | USNM395392                                      | 6  | 1,739       | F   | 3                      | 27                      | —                          | 0                                   | 0                                    | 0                         | +                                | 0                                    | 1                              | 1                          | 0                           | 0                        |
| CV282        | USNM395393                                      | 6  | 1,740       | F   | 3                      | 2                       | —                          | 0                                   | 0                                    | 0                         | 2                                | 0                                    | 6                              | 4                          | 0                           | 0                        |
| CV283        | USNM395394                                      | 6  | 2,021       | M   | 4                      | 5                       | —                          | 0                                   | 0                                    | 0                         | 13                               | 0                                    | 90                             | 28                         | 0                           | 0                        |
| CV284        | USNM395395                                      | 2  | 1,927       | F   | 4                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 0                                | —                                    | —                              | —                          | 0                           | 1                        |
| CV286        | USNM395397                                      | 2  | 1,647       | M   | 3                      | 0                       | —                          | 0                                   | 0                                    | 100+                      | 0                                | 0                                    | 3                              | 2                          | 2                           | 0                        |
| CV287        | USNM395417                                      | 2  | 1,732       | M   | 3                      | 0                       | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 4                              | 7                          | 0                           | 0                        |
| CV289        | USNM395594                                      | 2  | 1,767       | M   | 4                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 2                                | 0                                    | 3                              | 4                          | 0                           | 0                        |
| CV290        | USNM395595                                      | 2  | 1,627       | F   | 3                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 6                                | 0                                    | 0                              | 1                          | 0                           | 0                        |
| CV291        | USNM395596                                      | 6  | 1,927       | F   | 4                      | 1                       | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 24                             | 9                          | 0                           | 0                        |
| CV292        | USNM395597                                      | 2  | 1,895       | M   | 4                      | 5                       | 0                          | 0                                   | 34                                   | 5                         | 15                               | 0                                    | —                              | —                          | 0                           | 0                        |
| CV293        | USNM395598                                      | 6  | 1,996       | M   | 4                      | 2                       | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 5                              | 12                         | 0                           | 0                        |
| CV295        | USNM395527                                      | 6  | 1,390       | M   | 2                      | 9                       | 0                          | 0                                   | 0                                    | 0                         | 19                               | 0                                    | 0                              | 0                          | 1                           | 2                        |
| CV296        | USNM395528                                      | 2  | 1,298       | M   | 2                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | +                                    | 0                              | 0                          | 0                           | 0                        |
| CV297        | USNM395529                                      | 6  | 1,429       | F   | 2                      | 2                       | 0                          | 0                                   | 0                                    | 0                         | 1                                | 0                                    | 0                              | 0                          | 0                           | 2                        |
| CV298        | USNM395530                                      | 2  | 1,540       | M   | 3                      | 4                       | 0                          | 0                                   | 0                                    | 0                         | 5                                | +                                    | 0                              | 0                          | 0                           | 0                        |
| CV300        | USNM395532                                      | 6  | 1,438       | M   | 2                      | 11                      | 0                          | 0                                   | 0                                    | 0                         | 20+                              | 0                                    | 0                              | 0                          | 2                           | 2                        |
| CV303        | USNM395535                                      | 6  | 1,433       | M   | 2                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 4                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV304        | USNM395458                                      | 2  | 862         | M   | 1                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV305        | USNM395459                                      | 2  | 780         | F   | 1                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV306        | USNM395460                                      | 6  | 913         | M   | 1                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV307        | USNM395461                                      | 2  | 803         | F   | 1                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV308        | USNM395466                                      | 6  | 793         | F   | 1                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV309        | USNM395463                                      | 6  | 1,878       | F   | 4                      | 9                       | 0                          | 0                                   | 0                                    | 0                         | 9                                | 0                                    | 21                             | 26                         | 0                           | 0                        |
| CV310        | USNM395464                                      | 2  | 1,801       | M   | 4                      | 3                       | 0                          | 0                                   | 0                                    | 0                         | 1                                | 0                                    | 21                             | 17                         | 0                           | 0                        |
| CV311        | USNM395465                                      | 6  | 1,995       | M   | 4                      | 4                       | 0                          | 0                                   | 0                                    | 0                         | 3                                | 0                                    | 18                             | 23                         | 0                           | 0                        |
| CV312        | USNM395462                                      | 2  | 1,640       | F   | 3                      | 7                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 1                              | 0                          | 0                           | 0                        |
| CV313        | USNM395467                                      | 2  | 1,601       | M   | 3                      | 8                       | 0                          | 0                                   | 0                                    | 0                         | 5                                | 0                                    | 0                              | 2                          | 0                           | 0                        |
| CV314        | USNM395468                                      | 2  | 1,842       | F   | 4                      | 1                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | +                              | +                          | 0                           | 0                        |
| CV315        | USNM395603                                      | 2  | 1,621       | M   | 3                      | 7                       | +                          | 0                                   | 0                                    | 0                         | 12                               | 0                                    | 3                              | 5                          | 0                           | 0                        |
| CV316        | USNM395604                                      | 2  | 1,679       | M   | 3                      | 12                      | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 1                              | 4                          | 0                           | 0                        |
| CV317        | USNM395605                                      | 2  | 1,375       | M   | 2                      | 7                       | 0                          | 0                                   | 0                                    | 0                         | 5                                | 0                                    | 0                              | 0                          | 0                           | 2                        |
| CV318        | USNM395606                                      | 2  | 1,444       | F   | 2                      | 8                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV319        | USNM395607                                      | 6  | 1,681       | F   | 3                      | 2                       | 0                          | 0                                   | 0                                    | 0                         | 1                                | 0                                    | 5                              | 0                          | 0                           | 0                        |
| CV320        | USNM395608                                      | 2  | 1,670       | M   | 3                      | 20                      | +                          | 0                                   | 0                                    | 0                         | 13                               | 0                                    | 7                              | 0                          | 0                           | 0                        |
| CV321        | USNM395609                                      | 2  | 1,525       | M   | 3                      | 1                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 1                              | 0                          | 0                           | 0                        |
| CV322        | USNM395610                                      | 6  | 1,423       | F   | 2                      | 9                       | 0                          | 0                                   | 0                                    | 0                         | 6                                | 0                                    | 0                              | 0                          | 0                           | 0                        |
| CV323        | USNM395611                                      | 2  | 1,617       | M   | 3                      | 5                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 4                              | 2                          | 0                           | 0                        |
| CV324        | USNM395612                                      | 6  | 1,356       | F   | 2                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 13                               | 0                                    | +                              | 0                          | 0                           | 0                        |
| CV325        | USNM395613                                      | 6  | 1,348       | M   | 2                      | 4                       | +                          | 0                                   | 0                                    | 0                         | 8                                | 0                                    | 1                              | 0                          | 2                           | 2                        |
| PQ01         | USNM395407                                      | lat -8°N,<br>long 107°W                    | 1,978       | M   | 4                      | 3                       | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 5                              | 18                         | 0                           | 0                        |
| PQ02         | USNM395408                                      | lat -8°N,<br>long 107°W                    | 1,590       | M   | 2                      | 7                       | —                          | 0                                   | 0                                    | 0                         | 0                                | +                                    | 0                              | 0                          | 0                           | 0                        |
| 4PQ04        | USNM395410                                      | lat -8°N,<br>long 107°W                    | 1,677       | F   | 3                      | 3                       | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 2                              | 8                          | 1                           | 0                        |
| WFP10        | CAS15662  | lat -14°N,<br>long 100°W                   | 1,865       | F   | 4                      | 11                      | —                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | —                              | —                          | 0                           | 0                        |
| WFP11        | UCMP86284                                       | lat -14°N,<br>long 99-108°W                | 1,304       | M   | 2                      | —                       | +                          | —                                   | —                                    | —                         | —                                | +                                    | —                              | —                          | —                           | —                        |
| WFP12        | MVZ140640                                       | lat -14°N,<br>long 100°W                   | 1,671       | M   | 3                      | —                       | —                          | —                                   | —                                    | —                         | —                                | 0                                    | —                              | —                          | —                           | —                        |
| WFP13        | UCMP23071                                       | lat -10°N,<br>long 100°W                   | 1,572       | M   | 3                      | —                       | 0                          | —                                   | —                                    | —                         | —                                | —                                    | —                              | —                          | —                           | —                        |
| WFP19        | CAS15663  | lat -10°N,<br>long 100°W                   | 1,536       | M   | 3                      | —                       | 0                          | —                                   | —                                    | —                         | —                                | —                                    | —                              | —                          | —                           | —                        |
| WFP20        | MVZ140643                                       | lat -10°N,<br>long 100°W                   | 1,708       | F   | 3                      | —                       | 0                          | —                                   | —                                    | —                         | —                                | —                                    | —                              | —                          | —                           | —                        |
| WFP3         | LACM37549                                       | lat 14°30'N,<br>long 99°10'W               | 1,252       | F   | 2                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 2                                | 0                                    | 0                              | 0                          | 0                           | 2                        |
| WFP4         | LACM37550                                       | lat 14°30'N,<br>long 99°10'W               | 1,260       | M   | 2                      | 0                       | 0                          | 0                                   | 0                                    | 0                         | 0                                | 0                                    | 0                              | 0                          | 0                           | 0                        |

<sup>1</sup> CAS = California Academy of Sciences, San Francisco  
 LACM = Los Angeles County Museum  
 MVZ = Museum of Vertebrate Zoology, University of California, Berkeley  
 USNM = United States National Museum, Washington, D.C.

<sup>2</sup> Locality for collection number 2: lat 12°51'N, long 93°18'W  
 Locality for collection number 6: lat 7°11'N, long 90°32'W

<sup>3</sup> 1 = near term fetus or neonatal  
 2 = calf  
 3 = subadult  
 4 = adult

<sup>4</sup> This specimen also infected with *Crassicauda*.



APPENDIX TABLE 2.—Parasites in *Stenella* cf. *S. longirostris*.

| Host         |   |  |             |     |                        | Parasites noted          |                            |                                     |                                      |                           |                               |                                       |                               |                           |                             |                          |
|--------------|---|--|-------------|-----|------------------------|--------------------------|----------------------------|-------------------------------------|--------------------------------------|---------------------------|-------------------------------|---------------------------------------|-------------------------------|---------------------------|-----------------------------|--------------------------|
| Field number | Museum number of skeletal specimen <sup>1</sup> | Collection number or locality <sup>2</sup> | Length (cm) | Sex | Age class <sup>3</sup> | <i>Antisakis simplex</i> | <i>Halocercus delphini</i> | <i>Mastigonema stenellae</i> sp. n. | <i>Zalophotrema pacificum</i> sp. n. | <i>Campyla rochebruni</i> | <i>Terrabothrium forsteri</i> | <i>Sirobiloccephalus triangulatus</i> | <i>Phyllbothrium delphini</i> | <i>Monorygia grimaldi</i> | <i>Bolbosoma vasculosum</i> | <i>Bolbosoma batanae</i> |
| CV285        | USNM395396                                      | 2  | 1,769       | F   | 4                      | 56                       | —                          | 0                                   | 50+                                  | 50+                       | 3                             | 0                                     | 27                            | 13                        | 2                           | 0                        |
| CV288        | USNM395593                                      | 6  | 1,726       | F   | 4                      | 2                        | —                          | 0                                   | —                                    | —                         | —                             | —                                     | +                             | +                         | —                           | —                        |
| CV294        | USNM395526                                      | 6  | 1,494       | F   | 2                      | 1                        | +                          | 0                                   | 0                                    | 0                         | 10                            | 0                                     | 8                             | 0                         | 1                           | 0                        |
| CV299        | USNM395531                                      | 2  | 1,189       | M   | 2                      | 4                        | 0                          | 0                                   | 0                                    | 0                         | 5                             | 0                                     | 0                             | 0                         | 0                           | 4                        |
| CV301        | USNM395533                                      | 2  | 1,396       | F   | 2                      | 0                        | 0                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 0                             | 0                         | 2                           | 0                        |
| CV302        | USNM395534                                      | 2  | 1,184       | M   | 2                      | 9                        | 0                          | 0                                   | 0                                    | 0                         | 20+                           | 0                                     | 0                             | 0                         | 0                           | 1                        |
| PQ03         | USNM395409                                      | lat -8°N,<br>long 107°W                    | 1,571       | F   | 3                      | 8                        | 0                          | 0                                   | 0                                    | 0                         | 8                             | 0                                     | 8                             | 2                         | 2                           | 6                        |
| PQ05         | USNM395411                                      | lat -8°N,<br>long 107°W                    | 1,680       | F   | 4                      | 5                        | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 13                            | 13                        | 0                           | 0                        |
| PQ06         | USNM395412                                      | lat -8°N,<br>long 107°W                    | 1,729       | F   | 4                      | 9                        | —                          | 1                                   | 0                                    | 0                         | 0                             | 0                                     | 0                             | 4                         | 1                           | 0                        |
| PQ07         | USNM395413                                      | lat -8°N,<br>long 107°W                    | 1,761       | F   | 4                      | 9                        | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 8                             | 6                         | 1                           | 0                        |
| PQ08         | USNM395414                                      | lat -8°N,<br>long 107°W                    | 1,734       | M   | 4                      | 1                        | —                          | 0                                   | 0                                    | 0                         | 2                             | 0                                     | 7                             | 17                        | 0                           | 0                        |
| PQ09         | USNM395599                                      | lat -8°N,<br>long 109°45'W                 | 1,727       | M   | 4                      | 1                        | 0                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 21                            | 8                         | 1                           | 2                        |
| PQ10         | USNM395600                                      | lat -8°N,<br>long 107°W                    | 1,086       | F   | 2                      | 7                        | —                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 0                             | 0                         | 0                           | 1                        |
| PQ11         | USNM395601                                      | lat 8°N,<br>long 109°45'W                  | 1,054       | F   | 2                      | 0                        | —                          | 0                                   | 0                                    | 0                         | 6                             | +                                     | 0                             | 0                         | 0                           | 1                        |
| WFP14        | CAS15664  | lat -14°N,<br>long 99-108°W                | 1,618       | F   | 4                      | —                        | +                          | —                                   | —                                    | —                         | —                             | —                                     | —                             | —                         | —                           | —                        |
| WFP15        | CAS15665  | lat -14°N,<br>long 100°W                   | 1,637       | M   | 4                      | —                        | +                          | —                                   | —                                    | —                         | —                             | —                                     | —                             | —                         | —                           | —                        |
| WFP21        | MVZ140641                                       | lat -14°N,<br>long 100°W                   | 1,651       | M   | 3                      | —                        | +                          | —                                   | —                                    | —                         | —                             | —                                     | —                             | —                         | —                           | —                        |
| WFP22        | CAS15668  | lat -10°N,<br>long 100°W                   | 1,750       | M   | 4                      | —                        | —                          | —                                   | —                                    | —                         | —                             | —                                     | —                             | —                         | —                           | —                        |
| WFP6         | LACM37551                                       | lat 10°40'N,<br>long 108°40'W              | 896         | F   | 2                      | 0                        | 0                          | 0                                   | 0                                    | 0                         | 0                             | 0                                     | 0                             | 0                         | 0                           | 0                        |

<sup>1</sup> CAS = California Academy of Sciences, San Francisco  
 LACM = Los Angeles County Museum  
 MVZ = Museum of Vertebrate Zoology, University of California, Berkeley  
 USNM = United States National Museum, Washington, D.C.  
<sup>2</sup> Locality for collection number 2: lat 12°51'N, long 93°18'W  
 Locality for collection number 6: lat 7°11'N, long 90°32'W

<sup>3</sup> 1 = near term fetus or neonatal  
 2 = calf  
 3 = subadult  
 4 = adult