



American  
Bryological and  
Lichenological  
Society

---

## New Localities for Hawaiian Mosses

Author(s): Douglas R. Smith

Source: *The Bryologist*, Summer, 1967, Vol. 70, No. 2 (Summer, 1967), pp. 237-245

Published by: American Bryological and Lichenological Society

Stable URL: <https://www.jstor.org/stable/3240950>

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



American Bryological and Lichenological Society is collaborating with JSTOR to digitize, preserve and extend access to *The Bryologist*

rows, about 1 mm long; stomata superficial, at extreme base of the urn; peristome teeth rusty or red-brown, split to a low basal membrane into 32 long, filiform, densely spiculose, anastomosed divisions spirally wound together, about 1-1.3 mm long. Spores spherical, 11-15 $\mu$ , smooth or nearly so. Calyptae cucullate, smooth, naked.

PERU. PROV. LIMA: Depto. Lima, Cascadas de Barranco, terreno húmedo, afloramiento de agua dulce, ambiente calcáreo, alt. 5 m, A. Malaga, 4 Oct. 1964 (MICH—holotype).

## New Localities for Hawaiian Mosses

DOUGLAS R. SMITH<sup>1</sup>

**Abstract.** One hundred twelve mosses are reported from Hawaii with 78 as new additions to the individual floras of the six larger islands. Eighteen species previously known from a single island have been found on other islands. *Glossadelphus*, previously known only from Maui, is reported from Kauai, Oahu, and Molokai.

Recent contributions to the Hawaiian moss flora (Crosby, 1965; Hörmann, 1964; Schultze-Motel, 1962) have supplemented known distributions within the islands since Bartram's publications (1933, 1939, 1942, 1952). The few new additions to the individual island floras indicate that mosses from easily accessible localities are now fairly well known. This report is based primarily upon collections made by Dr. Otto Degener in somewhat isolated parts of the islands of Kauai, Oahu, Molokai, Lanai, Maui, and Hawaii during 1950-1955. Degener was assisted in much of this work by Miss Amy B. H. Greenwell and others. Because of the abundance of duplicate gatherings in Degener's material, only a few representative collections are cited from each station. These and portions of all other collections cited have been deposited in the Miami University Herbarium, with a representative series in the New York Botanical Garden, the University of Massachusetts, and the Bernice P. Bishop Museum, as requested by the collector. Some private collections were made available to supplement this report and permission to cite these is acknowledged, for they fill distributional gaps.

A careful search for small species in *Fissidens* resulted in seven new island records for this genus. Degener's collections of *Glossadelphus*, ". . . confined to the island of Maui" (Bartram, 1933), extend the range of this genus to three other islands — Kauai, Oahu, and Molokai. The number of additions to individual island bryofloras is: Kauai (20), Oahu (16), Molokai (13), Lanai (2), Maui (2), and Hawaii (25). A summary of the apparent total number of species reported for each island to date is: Kauai (100), Oahu (118), Molokai (67), Lanai (31), Maui (176), and Hawaii (138). It is clear that additional collections from more remote areas will expose new interisland range extensions as well as an occasional addition to the bryoflora of the state.

<sup>1</sup> Department of Botany, Miami University, Oxford, Ohio 45056.

In general, the sequence and nomenclature follow Bartram (1933) as has been customary in recent treatments of the Hawaiian moss flora. The "manual name" is given in parentheses where another name is now used as a result of recent taxonomic work, new interpretations of the Code, or where I have returned to the original spelling.

- Fissidens hawaiicus* Bartr. KAUAI: Alakai Swamp, *Degener* 22301E. OAHU: Sacred Falls, *Miller* 3331. MOLOKAI: Waikolu Valley, 4,000 ft., *Miller* 3825. MAUI: Haleakala, Koolau Gap, 7,500 ft., *Miller* 4263. HAWAII: 12 mi. below Kulani Prison, *Miller* 5222. New to Molokai and Hawaii.
- Fissidens oahuensis* Bartr. KAUAI: Halemanu Cliff lookout, 3,000 ft., *Miller* 3019. OAHU: Poamoho Gulch, 500 ft., *Degener* 21199. MAUI: Hanakalau Trail, 1,500 ft., *Miller* & *Lamberton* 3919. New to Kauai. Additionally reported from Hawaii.
- Fissidens baldwinii* Broth. KAUAI: Waioli Stream, *Degener* 22304H. OAHU: Kawaiiki Ditch Trail 1,000 ft., *Miller* 2313. MOLOKAI: Pipe-line Trail, upper Waikolu Valley, 4,000 ft., *Miller* 3824. New to Molokai. Additionally reported from Maui.
- Fissidens delicatulus* Angst. KAUAI: Moloaa Forest Reserve, *Degener* 22309D; Alakai Swamp, *Degener* 22301E. OAHU: E spur of E ridge, W branch of East Makaleha Valley, 1,800 ft., *Degener* 20877B. Additionally reported from Lanai, Maui, and Hawaii.
- Fissidens pacificus* Angst. KAUAI: Wainiha Valley, 750 ft., *Miller* 3247. OAHU: Kaipapau Valley, 1,100 ft., c. fr., *Miller* 2386. HAWAII: Honokanenui Valley, Kohala, *Degener* 22356F. New to Kauai and Hawaii. Additionally reported from Maui. This is the first reported discovery of a sporophyte.
- Fissidens mauiensis* C. Müll. KAUAI: S of Nawiliwili Bay, *Degener* 22291B. OAHU: E spur of E ridge, W branch of East Makaleha Valley, 1,800 ft., *Degener* 20876C. New to Kauai and Oahu. Previously known only from Maui.
- Fissidens lancifolius* Bartar. OAHU: Makaleha Valley, 1,600 ft., *Miller* 2674. MOLOKAI: W rim of Waikolu Canyon, 4,000 ft., *Miller* 3759. MAUI: Haleakala, 9 mi. W of Hana, 1,500 ft., *Miller* 4528. New to Molokai.
- Ceratodon purpureus* (Hedw.) Brid. KAUAI: Moloaa Forest Reserve, *Degener* & *Greenwell* 22309F. New to Kauai. Additionally reported from Maui and Hawaii.
- Trematodon latinervis* C. Müll. HAWAII: Near Kulani Prison, 4,000 ft., *Degener* 22281B. Additionally reported from Kauai, Oahu, Molokai, and Maui.
- Dicranella hochreutineri* Card. KAUAI: Mohihi Road, *Degener* & *Greenwell* 22287C. HAWAII: Kohala, W of Halona cone, 3,750 ft., *Lohman* H521. Additionally reported from Oahu and Maui.
- Dicranella hillebrandii* (C. Müll.) Broth. MOLOKAI: Cliffs overlooking Kalaupapa, *Degener* 24384.
- Dicranella integrifolia* (Broth.) Bartr. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21755B. Additionally reported from Oahu, Molokai and Maui.
- Campylospodium euphorocladum* (C. Müll.) Besch. KAUAI: Alakai Swamp, *Degener* & *Greenwell* 22301G. HAWAII: Kohala, Pololu-Honokanenui region, *Degener* 22341G. New to Kauai. Additionally reported from Oahu, Molokai, and Maui.
- Campylopus skottsbergii* Broth. HAWAII: Pololu Valley, Kohala, *Degener* & *Greenwell* 22354C; Waipio Valley, *Degener* 22278. New to Hawaii. Additionally reported from Kauai, Oahu and Maui.
- Campylopus introflexus* (Hedw.) Brid. KAUAI: S of Nawiliwili Bay, *Degener* 22288B, 22289. HAWAII: Above Kulani Prison, *Degener* 21933D, 22335D, 22338A. Additionally reported from Oahu.
- Campylopus densifolius* Angst. KAUAI: Kopiwi Trail, Kokee region, *Degener* & *Greenwell* 21621AA; along Mohihi Road, *Degener* & *Greenwell* 22285A; Alakai Swamp, *Degener* & *Greenwell* 22302, 22307B, 22346B. MOLOKAI: Cliffs above Kalaupapa, *Degener* 24382; head of Waikolu Valley, rainy region, *Degener* & *Tousley* 24403. HAWAII: Above Kulani Prison, *Degener* & *Greenwell* 21933E; Pololu-Honokanenui region, Kohala, *Degener* & *Greenwell* 22343C, 22348A; Pololu Valley, Kohala, *Degener* & *Greenwell* 22354D. Additionally reported from Oahu and Maui.

The following two species are mostly separable on the basis of Bartram's key, but further studies are needed to show whether they are variants of the same species. For now it seems reasonable to retain both species.

- Campylopus hawaiioco-flexuosus* (C. Müll.) Par. KAUAI: E of Kalalau Lookout, *Degener & Greenwell* 21620. S of Nawiliwili Bay, *Degener* 22291; Kalalau rim near lookout, O. & I. *Degener* 23887; Alakai Swamp, *Degener & Greenwell* 22295E, 22296. MOLOKAI: Head of Waikolu Valley, *Degener & Tousley*, 24402, 24452. MAUI: Above McGregor, W Maui, 2,500-4,000 ft., *Degener* 24531. HAWAII: In *kipuka* on S side of Saddle Road near mile-post 16, *Degener* 22277D; above Kulani Prison, *Degener & Greenwell* 22358. Additionally reported from Oahu.
- Campylopus purpureo-flavescens* (C. Müll.) Par. HAWAII: Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22343C; above Kulani Prison, *Degener* 22363B. Additionally reported from Kauai, Oahu, Molokai and Maui.
- Campylopus wheeleri* (C. Müll.) Hampe ex Par. KAUAI: S of Nawiliwili Bay, *Degener* 22288A; Alakai Swamp, *Degener & Greenwell* 22346A. HAWAII: Makaha Falls, Akaka, *Kikudome* 36B. New to Hawaii. Additionally reported from Maui.
- Campylopus boswellii* (C. Müll.) Par. KAUAI: Alakai Swamp, *Degener & Greenwell* 22297. New to Kauai. Additionally reported from Molokai and Maui.
- Campylopus richardii* Brid. [= *C. umbellatus* (Arnott) Par.] OAHU: SW side of Poamoho Gulch, SW of Brodie Camp, 500 ft., *Degener* 21197, 21204. HAWAII: Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22355B, 22356F; above Kulani Prison, *Degener & Greenwell* 22359. Additionally reported from Kauai, Lanai and Maui.
- Campylopus exasperatus* (Nees & Blum.) Brid. KAUAI: NE of Kapalikea, Hanalei, *Degener* 20448. HAWAII: Along Saddle Road, 21 mi. from Hilo, *Degener* 19924. New to Kauai. Additionally reported from Oahu and Maui.
- Campylopus tubulosus* Bartr. KAUAI: S of Nawiliwili Bay, *Degener* 22288F. HAWAII: Waipio Valley, *Degener*, 22278C; Keauohana Forest Reserve, near Pahoa, Puna, *Degener* 22282. New to Hawaii. Additionally reported from Maui.
- Dicranodontium falcatum* Broth. OAHU: S Kipapa Ridge Trail, Koolau Mts., 2,000 ft., *Miller* 2100, 2133; Wahiaawa, central ridge to Mt. Kaala, 3,900 ft., *Chock & Lamberton* 477M. New to Oahu. Additionally reported from Kauai, Maui and Hawaii.
- Dicranum speirophyllum* Mont. KAUAI: E of Kalalau Lookout, *Degener & Greenwell* 21620C; Kalalau Lookout, *Degener* 22284. MAUI: Koolau Gap, Haleakala, 7,500 ft., *Lohman* 4252A. HAWAII: Above Kulani Prison, *Degener & Greenwell* 22335, 22338. Additionally reported from Oahu and Molokai.
- Leucoloma molle* (C. Müll.) Mitt. HAWAII: Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22356A, 22357A. New to Hawaii. Additionally reported from Oahu and Maui.
- Octoblepharum albidum* Hedw. KAUAI: Second point E of Haupu, Kipu, 1,000-1,700 ft., *Miller* 3097. New to Kauai. Additionally reported from Oahu and Maui.
- Leucobryum gracile* Sull. KAUAI: E of Kohala Lookout, *Degener & Greenwell* 21620B; along Mohihi Road, *Degener & Greenwell* 22285, 22287A; S of Nawiliwili Bay, *Degener* 22286B, 22289; ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22309, 22310; Alakai Swamp, *Degener & Greenwell* 22295, 22301, 22307, 22346E. OAHU: S Haleauau Valley, below Kaala summit, *Degener* 21217E. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21755F; near Pahoa, Puna, in woods, *Degener* 21758; Keauohana Forest Reserve near Pahoa, Puna, *Degener* 22282E; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22348. Additionally reported from Molokai, Lanai, and Maui.
- Leucobryum solfatarae* C. Müll. [= *L. hawaiiense* var. *solfatarae* (C. Müll.) Bartr.] MOLOKAI: Head of Waikolu Valley, *Degener & Tousley* 24446. MAUI: Above McGregor, 2,500-4,000 ft., *Degener* 24536. Additionally reported from Kauai and Hawaii.
- Leucobryum solfatarae* var. *hawaiense* (Reichardt) Wijk & Marg. [= *L. hawaiense* (Reichardt) Bartr.] KAUAI: S of Nawiliwili Bay, *Degener* 22286D. MOLOKAI: Head of Waikolu Valley, rainy region, *Degener & Tousley* 24404; Nualele Valley, rain forest E of Waikolu, *Degener et al.* 24422, 24425, 24430. Additionally reported from Oahu, Maui, and Hawaii. Reichardt's original spelling of the name for this taxon was "hawaiensis" (Reichardt, 1877), not "hawaiense" (Wijk & Margadant, 1962).
- Leucobryum pachyphyllum* C. Müll. KAUAI: Alakai Swamp, *Degener* 22301B, *Degener & Greenwell* 22303; S of Nawiliwili Bay, *Degener* 22286C. Additionally reported from Oahu, Molokai, Lanai, and Maui.
- Syrrhopodon oahuensis* Broth. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21755C. MAUI: 9 mi. W of Hana, 1,500 ft., Haleakala, *Miller* 4547. New to Hawaii and Maui. Additionally reported only from Oahu.

- Syrrhopodon hawaiicus* C. Müll. MOLOKAI: Pipe-line trail, head of Waikolu Valley, 4,000 ft., Miller 3830. New to Molokai. Additionally reported from Oahu, Maui, and Hawaii.
- Calymperes tenerum* C. Müll. OAHU: Papali Gulch, Hauula, Koolau Range, 1,500 ft., Miller 2600. HAWAII: Keauohana Forest Reserve near Kalapana, Miller & Lamberton 5080; Kalapana Beach, Miller & Lamberton 5085. New to Oahu and Hawaii. Additionally reported from Kauai and Maui.
- Merceyopsis crassinervis* (Broth.) Bartr. HAWAII: Above Kulani Prison, Mauna Loa, Degener & Greenwell 21933A; summit of Hualalai, Greenwell 185. New to Hawaii. Additionally reported from Oahu and Maui.
- Anoectangium euchloron* (Schwaegr.) Mitt. KAUAI: Ridge behind Papaa, Moloaa Forest Reserve, Degener & Greenwell 22309. OAHU: SW side of Poamoho Gulch, SW of Brodie Camp, 500 ft., Degener 21198. HAWAII: Above Kulani Prison, Mauna Loa, Degener & Greenwell 21933B; Waipio Valley, Degener 22278E; Pololu-Honokanenui Valley, Kohala, Degener & Greenwell 22335A, 22341C, 22353C, 22360A. New to Kauai. Additionally reported from Molokai, Lanai, and Maui.
- Anoectangium haleakalae* (C. Müll.) Par. KAUAI: Ridge behind Papaa, Moloaa Forest Reserve, Degener & Greenwell 22309C. MOLOKAI: Cliffs and ledges overlooking Kalaupapa, Degener et al. 24387. HAWAII: Pololu Valley, Kohala, Degener & Greenwell 22340. New to Kauai and Molokai. Additionally reported from Oahu and Maui.
- Weissia viridula* Hedw. OAHU: SW side of Poamoho Gulch, SW of Brodie Camp, 500 ft., Degener 21201, 21207. HAWAII: Above Kulani Prison, Mauna Loa, Degener & Greenwell 21933C. Additionally reported from Kauai.
- Weissia ovalis* (Williams) Bartr. KAUAI: S of Nawiliwili Bay, Degener 22288C. Additionally reported from Oahu, Molokai, Lanai, Maui, and Hawaii.
- Trichostomum mauiense* Broth. KAUAI: S of Nawiliwili Bay, Degener 22288D. MOLOKAI: Cliffs and ledges overlooking Kalaupapa, Degener et al. 24383. New to Molokai. Additionally reported from Oahu, Maui, and Hawaii.
- Trichostomum oblongifolium* Bartr. OAHU: E spur of E ridge, W branch of East Makaleha Valley, Degener 20878D. HAWAII: Honokanenui Valley, Kohala, Degener & Greenwell 22342C, 22353B. Additionally reported from Maui.
- Pseudosymblepharis mauiensis* (C. Müll.) Broth. OAHU: E spur of E ridge. W branch of East Makaleha Valley, Degener 20878B. HAWAII: Pololu-Honokanenui region, Kohala, Degener & Greenwell 22350, 22351B. New to Hawaii. Additionally reported from Maui.
- Tortella caespitosa* (Schwaegr.) Limp. OAHU: Koko Head, 100 ft., in gulches, Miller & Lohman 2026. New to Oahu. Additionally reported from Lanai, Maui, and Hawaii.
- Barbula vinealis* ssp. *cylindrica* (Tayl.) Podp. OAHU: DuPont Trail, N slope of Kaala, 3,000 ft., Degener 20530. New to Oahu. Additionally reported only from Maui.
- Tortula princeps* De Not. HAWAII: Edges of caves and surrounding area, Mauna Kea, 13,000 ft., Greenwell B146. New to Hawaii. Additionally reported from Maui.
- Tortula alpina* var. *inermis* (Milde) De Not. HAWAII: NE of Halepohaku, Mauna Kea, on exposed lava, 9,500 ft., Degener 20425A, 20427. Additionally reported from Maui.
- Rhacomitrium lanuginosum* var. *pruininosum* Hook. f. & Wils. OAHU: Waianae Mts., Mt. Kalona, E ridge on exposed rocks, 950 m, Fosberg 13004. HAWAII: In *kipuka* on S side of Saddle Road near Milepost 16, Degener 22277B. New to Oahu. Additionally reported from Kauai, Molokai and Maui.
- Funaria subintegra* Broth. KAUAI: Kokee Camp, 3,700 ft., Miller 3007. HAWAII: Greenwell Ranch above Captain Cook, 2,500-5,000 ft., Miller & Lamberton 5380. Additionally reported from Oahu, Molokai, Lanai, and Maui.
- Pohlia mauiensis* (Broth.) Schultze-Mot. HAWAII: Honokanenui Valley, Kohala, Degener & Greenwell 23356C. New to Hawaii. Additionally reported from Maui.
- Bryum vino-viride* Bartr. HAWAII: Papaloa, above Captain Cook, Kona, 5,000 ft., Degener 20423. New to Hawaii. Additionally reported from Kauai and Oahu.
- Bryum megalostegium* Sull. KAUAI: E of Hanalei, road bank, 400 ft., Degener & Hatheway 20466. Additionally reported from Oahu, Maui and Hawaii.
- Bryum atrovirens* Vill. ex Brid. [=B. erythrocarpum Schwaegr.] MAUI: Near Hana, Degener 21928. HAWAII: Kilaea region, on bare pahoehoe lava, O. & I. Degener 23817. Additionally reported from Oahu.
- Bryum caespiticium* Hedw. HAWAII: NW of Halepohaku, Mauna Kea, in crevices of lava, 9,500 ft., Degener 20425B. New to Hawaii. Additionally reported from Maui.

- Bryum decaisnei* Dozy & Molk. HAWAII: Above Kulani Prison, *Degener & Greenwell* 22338B. New to Hawaii. Additionally reported from Maui.
- Bryum argenteum* var. *lanatum* (P. Beauv.) B.S.G. OAHU: Kealia Trail, Kealia, Waianae Mts., 1,700 ft., *Miller* 2554; summit of Mt. Kaala, 4,000 ft., *Miller* 1431. New to Oahu. Additionally reported from Kauai, Maui and Hawaii.
- Rhodobryum giganteum* (Hook.) Schimp. MAUI: Above McGregor, 2,500-4,000 ft., *c. fr.*, *Degener* 24533. Additionally reported from Kauai and Oahu. This is the first report of a sporophyte from Hawaii.
- Mnium rostratum* Schrad. KAUAI: Waioli Stream, *Degener & Greenwell* 22304F. OAHU: Kaipapau Valley, 1,100-1,500 ft., *Miller* 2362; E West Manoa Falls, Manoa Valley, *Miller* 2574; Papali Gulch, Hauula, Koolau Range, 1,500 ft., *Miller* 2629. New to Kauai and Oahu. Additionally reported from Maui.
- Rhizogonium pungens* Sull. KAUAI: S of Nawiliwili Bay, *Degener* 22286A; Waioli Stream, *Degener & Greenwell* 22298D; Alakai Swamp, *Degener & Greenwell* 22301. HAWAII: Keauhona Forest Reserve near Pahoa, Puna, *Degener* 22282A; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22343A. Additionally reported from Oahu, Molokai, Lanai, and Maui.
- Rhizogonium spiniforme* (Hedw.) Bruch. KAUAI: Waioli Stream, *Degener & Greenwell* 22304A; Alakai Swamp, *Degener & Greenwell* 22346. OAHU: Near Summit Ridge overlooking head of Makua Valley, *Degener* 21164; S Haleauau Valley below Kaala summit, *Degener* 21217. HAWAII: Honokanenui Valley, Kohala, *Degener & Greenwell* 22356E. Additionally reported from Molokai, Lanai, and Maui.
- Sciaromium tricostata* (Sull.) Mitt. [= *Limbella tricostata* (Sull.) C. Müll.] KAUAI: Along Mohihi Road, *Degener & Greenwell* 22285, 22287; Alakai Swamp, *Degener & Greenwell* 22302A. OAHU: Central Haleauau Gulch, S of Kaala, partly and completely submerged in rock stream, 3,100 ft., *Degener & Cainbec* 20679; S Haleauau Valley below Kaala summit, *Degener* 21217A. MOLOKAI: Nualele Valley, rain forest, *Degener et al.* 24369. MAUI: Above McGregor, 2,500-4,000 ft., *Degener* 24526, 24529, 24542. Additionally reported from Hawaii. This is the oldest legitimate name for the taxon according to the *Index Muscorum*.
- Philonotis hawaiica* (C. Müll.) Broth. KAUAI: S of Nawiliwili Bay, *Degener* 22288E. LANAI: NE summit, *Degener* 21929D. MAUI: Keanae, dripping rocks, *O. & I. Degener* 2382I. HAWAII: Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22335, 22356B. New to Lanai. Additionally reported from Oahu and Molokai.
- Philonotis falcata* (Hook.) Mitt. OAHU: S Haleauau Valley below Kaala summit, *Degener* 21217B. Additionally reported from Kauai, Molokai, and Maui.
- Philonotis turneriana* (Schwaegr.) Mitt. OAHU: E spur of E ridge, W branch of East Makaleha Valley, *Degener* 20878E. MOLOKAI: Nualele rain forest, *Degener et al.* 24371, 24372; head of Waikolu Valley, rainy region, *Degener & Tousley* 24367. HAWAII: Waipio Valley, *Degener* 22278D; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22262A, 22335C, 22351, 22362A; Below Kulani Prison, dense forest, 4,000 ft., *Degener* 22283; E of Honokanenui Valley, Kohala, *Degener & Greenwell* 22342. Additionally reported from Kauai, Lanai, and Maui.
- Zygodon tetragonostomus* A. Br. HAWAII: Above Kulani Prison, Mauna Loa, *Degener & Greenwell* 21933, 22364. Additionally reported from Oahu and Maui.
- Orthotrichum hillebrandii* C. Müll. OAHU: S Kipapa Ridge Trail, Koolau Mts., 1,200-2,500 ft., *Miller* 2101, 2108. New to Oahu. Additionally reported from Maui and Hawaii.
- Macromitrium brevisetum* Mitt. KAUAI: Waioli Stream, *Degener & Greenwell* 22305A. MOLOKAI: Mapulehu Valley, *Degener & Tam* 22467. New to Molokai. Additionally reported from Oahu, Maui, and Hawaii.
- Macromitrium emersulum* C. Müll. MOLOKAI: Mapulehu Valley, *Degener & Tam* 24473. New to Molokai. Additionally reported from Kauai, Oahu, Maui, and Hawaii.
- Macromitrium piliferum* Schwaegr. KAUAI: Waioli Stream, *Degener & Greenwell* 22304D; Ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22309B; Alakai Swamp, *Degener & Greenwell* 22300B, 22346C. MOLOKAI: Nualele Valley, rain forest, *Degener et al.* 24392, 24420; Waikolu Valley, rainy region, *Degener & Tousley* 24407. LANAI: NE summit, *Degener* 21929C. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21754B, 21755D; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22343, 22353C. Additionally reported from Oahu and Maui.

*Macromitrium owahense* C. Müll. KAUAI: Mohihi Road, *Degener & Greenwell* 22287D; ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22309B. MOLOKAI: Head of Waikolu Valley, epiphytic, *O. & I. Degener* 23837. Additionally reported from Oahu, Lanai, Maui and Hawaii.

*Rhacopilum cuspidigerum* (Schwaegr.) Mitt. KAUAI: E side of Hanalei Valley, road cut, 400 ft., *Degener & Hatheway* 20465, 20471; S of Nawiliwili Bay, *Degener* 22288, 22294; Waioli Stream, *Degener & Greenwell* 22298; ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22301B, 22309. OAHU: E spur of E ridge, W branch of East Makaleha Valley, *Degener* 20876A; Kahana Valley, *Degener & Silva* 21080; SW side of Poamoho Gulch, SW of Brodie Camp, 500 ft., *Degener* 21200, 21206; S. Haleauau Valley, below Kaala summit, *Degener* 21217C. MOLOKAI: Mapulehu Valley, *Degener & Tam* 24373, 24468, 24472; cliffs and ledges overlooking Kalaupapa, *Degener* 24378; Waikolu Valley, *Degener & Tousley* 24411, 24444, 24445; Nualele Valley, *Degener et al.* 24459. HAWAII: Near Pahoa, Puna, *Degener* 21758A; Honokanenui Valley, Kohala, *Degener & Greenwell* 22353; Waipio Valley, *Degener* 22278A; E of Honokanenui Valley, Kohala, *Degener & Greenwell* 22360; Pololu Valley, Kohala, *Degener & Greenwell* 22339, 22340, 22354B; Kaumana Caves, near Hilo, *Degener* 22276C. Additionally reported from Lanai and Maui. Reported as "Rhamomitrium cuspidigerum" by Crosby (1965).

*Trachypus bicolor* Reinw. & Hornsch. HAWAII: Hualalai, Mt. Laalaau, *Kondo & Miller s.n.* New to Hawaii. Additionally reported from Maui.

*Aerobryopsis longissima* (Dozy & Molk.) Fleisch. KAUAI: Waioli Stream, *Degener & Greenwell* 22298C; ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22309E. OAHU: S Haleauau Valley below Kaala summit, *Degener* 21217D. MOLOKAI: Nualele Valley, rain forest, E of Waikolu, *Degener et al.* 24417, 24418. HAWAII: Near Pahoa, Puna in woods, *Degener* 21758D; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22335B, 22334. Additionally reported from Lanai and Maui.

*Barbella trichophora* (Mont.) Fleisch. in Broth. HAWAII: E of Honokanenui Valley, Kohala, *Degener & Greenwell* 22342A; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22355D, 22362B. Additionally reported from Kauai, Oahu, and Molokai.

*Floribundaria floribunda* (Dozy & Molk.) Fleisch. OAHU: S Haleauau Valley below Kaala summit, *Degener* 21217E. HAWAII: Dry water course, Holokaole, Kona, *Greenwell* B109. New to Hawaii. Additionally reported from Maui.

*Baldwinella kealeensis* (Reichardt) Bartr. KAUAI: S of Nawiliwili Bay, *Degener* 22291C. OAHU: Narrow gulch SW of Dillingham Ranch, Mokuleia, dark forest, 2,500 ft., *Degener* 21189; S Haleauau Valley below Kaala summit, *Degener* 21217B. MOLOKAI: Head of Waikolu Valley, *Degener & Tousley* 24449, 24451. HAWAII: Pololu-Honokanenui region, Kohala, *Degener* 22337. Additionally reported from Maui.

*Homaliodendron flabellatum* (Sm.) Fleisch. KAUAI: Kopiwi Trail, Kokee region, *Degener & Greenwell* 21621A; S of Nawiliwili Bay, *Degener* 22291A; Alakai Swamp, *Degener & Greenwell* 22295; Waioli Stream, *Degener & Greenwell* 22298, 22304B, 22308; Mohihi Road, *Degener & Greenwell* 22311. OAHU: E spur of E ridge, W branch of East Makaleha Valley, dark gulch, 1,800 ft., *Degener* 20876B, 20878F. MOLOKAI: Nualele Valley, *Degener et al.*, 24458. HAWAII: Below Kulani Prison, dense forest, 4,000 ft., *Degener* 22275. Additionally reported from Lanai and Maui.

*Thamnium speciosum* Broth. HAWAII: Greenwell Ranch, Kealakekua, 4,200-5,000 ft., *Greenwell* B-229. New to Hawaii. Additionally reported from Maui.

*Daltonia contorta* C. Müll. KAUAI: Kalalau Lookout, 4,000 ft., *Miller* 2818; Waimea, Alakai Swamp Trail, 4,000 ft., *Miller* 2883. MOLOKAI: NW of Mt. Kolekole, Kawela, 3,800 ft., *Miller & Lamberton* 3432; W rim of Waikolu Valley, 4,000 ft., *Miller & Lamberton* 3737, 3746; Mt. Kolekole, 3,700 ft., and below to 3,000 ft., Kawela, *Miller & Lamberton* 3681. HAWAII: Kohala Mts., Waimea Forest Reserve, 4,000 ft., *Miller & Lamberton* 4877; in kipuka in 1843 lava flow, Saddle Road, 15 mi. from Hilo, *Miller & Lamberton* 5047. New to Kauai, Molokai, and Hawaii. Previously known only from Maui.

*Daltonia rufescens* Broth. OAHU: Halawa Trail, Koolau Mts., 2,300 ft., *Miller* 2781F. HAWAII: Near Halepohaku, Mauna Kea, 8,000 ft., *Lamberton* 1345; Kohala Mts., forest gate above Waimea 3,000 ft., *Miller & Lamberton* 4819A. New to Oahu and Hawaii. Previously known only from Maui.

*Daltonia pseudostenophylla* Bartr. OAHU: Halawa Trail, Koolau Mts., 2,300 ft., *Miller* 2786A. HAWAII: 6 mi. below Kulani Prison, *Miller & Lamberton* 5262B; in kipuka in 1843 lava

- flow, Saddle Road, 15 mi. from Hilo, *Miller & Lamberton* 5047 (mixed with *D. contorta*). New to Oahu and Hawaii. Previously known only from Maui.
- Distichophyllum paradoxum* (Mont.) Mitt. KAUAI: Waioli Stream, *Degener & Greenwell* 22299B. MOLOKAI: Head of Waikolu Valley, rainy region, *Degener & Tousley* 24366; Nualele Valley, *Degener et al.* 24397, 24456. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21754C, 21755; near Pahoa, Puna, in woods, *Degener* 21758C, 22282C; below Kulani Prison, dense forest, 4,000 ft., *Degener* 22281A. Additionally reported from Oahu and Maui.
- Distichophyllum freycinetii* (Schwaegr.) Mitt. MOLOKAI: Waikolu Valley, rainy region, *Degener & Tousley* 24406. HAWAII: Below Kulani Prison, dense forest, 4,000 ft., *Degener* 22279; Keauohana Forest Reserve, near Pahoa, Puna, *Degener* 22282C. Additionally reported from Kauai, Oahu, and Lanai.
- Hookeria acutifolia* Hook. & Grev. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21754A. New to Hawaii. Additionally reported from Oahu and Maui.
- Hookeriopsis purpurea* (C. Müll.) Broth. HAWAII: Pauahi Clinkers, in old *aa* lava flows, 4,000 ft., *Greenwell* B7A. Additionally reported from Kauai, Oahu, Molokai, and Maui.
- Fabronia degeneri* Bartr. OAHU: Kealia Trail, Kealia, Waianae Mts., 1,700 ft., *Miller* 2555. HAWAII: Christ Church yard, 1,300 ft., *Greenwell* B-34. New to Oahu and Hawaii. Previously reported only from Molokai.
- Claopodium whippleanum* (Sull.) Ren. & Card. [= *Claopodium hawaiiense* Williams.] KAUAI: W side, Nonou, Wailua, St. John & Bishop 24942, 24946. New to Kauai. Additionally reported from Oahu and Maui.
- Thuidium plicatum* Mitt. KAUAI: Kopiwai Trail, Kokee region, *Degener & Greenwell* 21621A; Waioli Stream, *Degener & Greenwell* 22305; ridge behind Papaa, Moloaa Forest Reserve, *Degener & Greenwell* 22310A; Mohihi Road, *Degener & Greenwell* 22311A. HAWAII: Puu Oo Trail, 6,000 ft., *Degener* 20713; in *kipuka*, S side of Saddle Road near milepost 16, *Degener* 22277C; Pololu-Honokanenui region, Kohala, *Degener* 22337A, 22354A. Additionally reported from Oahu, Molokai, and Maui.
- Thuidium havaiense* Reichardt. [= *T. hawaiiense* Reichardt.] KAUAI: Alakai Swamp, *Degener & Greenwell* 22295C; Waioli Stream, *Degener & Greenwell* 22304E. OAHU: E spur of E ridge, W branch of East Makaleha Valley, dark gulch, 1,800 ft., *Degener* 20876, 20877. MOLOKAI: Nualele Valley rain forest, *Degener et al.* 24393; head of Waikolu Valley, *Degener & Tousley* 24443. Additionally reported from Lanai, Maui, and Hawaii. The original spelling of the name for this taxon was "havaiense" (Reichardt, 1877).
- Pleuropus wilkesianus* (Sull.) Broth. KAUAI: Waioli Stream, *Degener & Greenwell* 22298B. MOLOKAI: Cliffs and ledges overlooking Kalaupapa, *Degener et al.* 24380. HAWAII: Kau-mana Caves, near Hilo, *Degener* 22276; above Kulani Prison, *Degener & Greenwell* 22338C; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22355C. Additionally reported from Oahu, Lanai, and Maui.
- Pleuropus wilkesianus* var. *sciurooides* (C. Müll.) Bartr. LANAI: NE summit of Lanai, *Degener* 21929. HAWAII: Honokanenui Valley, Kohala, *Degener & Greenwell* 22353A. Additionally reported from Kauai, Oahu, Molokai, and Maui.
- Brachythecium rutabulum* (Hedw.) B.S.G. MAUI: Above McGregor, W Maui, 2,500-4,000 ft., *Degener* 24532, 24537. Known only from Maui.
- Brachythecium hawaiiicum* Bartr. HAWAII: Edges of caves and surrounding area, Mauna Kea, 13,000 ft., *Greenwell* B149. New to this altitude on Mauna Kea. Additionally reported from Maui.
- Eurhynchium selaginellifolium* (C. Müll.) Bartr. KAUAI: Waioli Stream, *Degener & Greenwell* 22299B, 22305. OAHU: S Haleauau Valley below Kaala summit, *Degener* 21237, 21217F; E spur of E ridge, W branch of East Makaleha Valley, dark gulch, 1,800 ft., *Degener* 20878. MOLOKAI: Mapulehu Valley, *Degener & Tam* 24472, 24475, 24476; Nualele Valley rain forest, *Degener et al.* 24394. HAWAII: Kaumana Caves, near Hilo, *Degener* 22276A; Pololu-Honokanenui region, Kohala, *Degener & Greenwell* 22351A, 22339A. New to Molokai. Additionally reported from Maui.
- Rhynchosstiella hawaiica* (C. Müll.) Broth. OAHU: S Haleauau Valley, below Kaala summit, *Degener* 21217G. Known only from Oahu.
- Entodon solanderi* (Angst.) Jaeg. KAUAI: Wainiha Valley, below Honoona Pali, 750 ft., *Miller* 3181. New to Kauai. Additionally reported from Oahu and Maui.
- Entodon subcuspidatus* (Hampe ex C. Müll.) Bartr. KAUAI: Wainiha Valley, 0.5 mile below

- water intake, 750 ft., *Miller* 3225. New to Kauai. Previously reported only from Maui. *Plagiothecium draytonii* (Sull.) Bartr. OAHU: Summit of Mt. Kaala, 4,000 ft., *Miller* 1409A. New to Oahu. Additionally reported from Kauai, Molokai, Maui, and Hawaii.
- Heterophyllum subauriculatum* (Hampe ex C. Müll.) Broth. HAWAII: Kohala Mts., Waimea Forest Reserve, 4,000 ft., *Miller* & *Lamberton* 4853; open *koa*-*Myoporum* forest area, Puulehua, South Kona, 5,000 ft., *Greenwell* B-80. New to Hawaii. Previously reported only from Maui.
- Brotherella opaeodon* (Sull.) Broth. KAUAI: Mohihi Road, *Degener* & *Greenwell* 22285B; E of Kalalau Lookout, *Degener* & *Greenwell* 21620A. OAHU: S Haleauau Valley below Kaala summit, *Degener* 21217H. MAUI: Gulch above The Needle, Iao Valley, *Degener* 21930. HAWAII: Keauohana Forest Reserve, near Pahoa, Puna, *Degener* 22282B.
- Sematophyllum hawaiiense* (Broth.) Broth. KAUAI: Nualolo Trail, Kokee to Na Pali coast, 2,800-3,500 ft., *Miller* 2962; Halemanu Cliff Lookout, Waiahulu Stream, Waimea, 3,000 ft., *Miller* 3024, 3025. New to Kauai. Additionally reported from Oahu, Maui, and Hawaii.
- Acroporium fuscus-flavum* (C. Müll.) Broth. KAUAI: Alakai Swamp, *Degener* & *Greenwell* 22297, 22300; Waioli Stream, *Degener* & *Greenwell* 22304C, 22306. MOLOKAI: Nualele Valley rain forest, E of Waikolu, *Degener* et al. 24421. HAWAII: Near Pahoa, Puna, in woods, *Degener* 21758B; Kulani Prison Road, 4,000 ft., *Degener* 21755A; below Kulani Prison, dense forest, 4,000 ft., *Degener* 22280A, 22281; Keauohana Forest Reserve, near Pahoa, Puna, *Degener* 22282D; Pololu-Honokanenui region, Kohala, *Degener* & *Greenwell* 22335, 22352, 22362; above Kulani Prison, *Degener* & *Greenwell* 22358A. Additionally reported from Oahu, Lanai, and Maui.
- Trichosteleum hamatum* (Dozy & Molk.) Jaeg. KAUAI: Alakai Swamp, *Degener* & *Greenwell* 22297A. MAUI: E branch, Makapipi Gulch, 9 mi. W of Hana, 1,500-3,500 ft., *Miller* & *Lamberton* 4597. HAWAII: Kulani Prison Road, 4,000 ft., *Degener* 21754, 22280; in kipuka, S side of Saddle Road, near milepost 16, *Degener* 22277.
- Taxithelium mundulum* (Sull.) Bartr. HAWAII: Pololu-Honokanenui region, Kohala, *Degener* & *Greenwell* 22343D. Additionally reported from Kauai, Oahu, and Lanai.
- Glossadelphus baldwinii* Broth. KAUAI: Second point E of Haupu, 1,000-1,400 ft., *Miller* 3086. OAHU: S Haleauau Valley, below Kaala summit, *Degener* 21217H; narrow Gulch SW of Dillingham Ranch, Mokuleia, dark forest, 2,000 ft., *Degener* 21189. MOLOKAI: Upper Waikolu Valley, Pipe-line Trail, 4,000 ft., *Miller* 3785. New to Kauai, Oahu, and Molokai. Previously reported only from Maui.
- Glossadelphus zollingeri* var. *filicaulis* (Fleisch.) Fleisch. KAUAI: Waioli Stream, *Degener* & *Greenwell* 22305AA; ridge behind Papaa, Moloaa Forest Reserve, *Degener* & *Greenwell* 22310C. MOLOKAI: Head of Waikolu Valley, aquatic in rivulet, *O.* & *I.* *Degener* 23838. New to Kauai and Molokai.
- Ectropothecium sandwichense* (Hook. & Arnott) Mitt. KAUAI: Wainiha Valley, 0.5 mi. below water intake, 750 ft., *Miller* 3237. OAHU: Manoa Valley, E of West Manoa Falls, Koolau Mts., 800 ft., *Miller* 2074, 2075. HAWAII: Pololu-Honokanenui region, Kohala, *Degener* & *Greenwell* 22355A. New to Kauai and Oahu. Additionally reported from Maui.
- Ectropothecium viridifolium* Bartr. MAUI: Along Makamakaole Stream, W Maui, *Degener* 21151. New to Maui. Additionally reported from Oahu and Lanai.
- Isopterygium albescens* (Hook.) Jaeg. KAUAI: Alakai Swamp, *Degener* & *Greenwell* 22301C. Additionally reported from Oahu, Molokai, Lanai, Maui, and Hawaii.
- Vesicularia periridis* (Ångst.) C. Müll. [= *V. graminicolor* (Ångst.) Broth.] KAUAI: Ridge behind Papaa, Moloaa Forest Reserve, *Degener* & *Greenwell* 22309G. OAHU: Near summit ridge overlooking head of Makua Valley, *Degener* 21164; S Haleauau Valley below Kaala summit, *Degener* 21217I. MOLOKAI: Mapulehu Valley, *Degener* & *Tam* 24374. LANAI: NE summit of Lanai, *Degener* 21929B; near wharf, S side of Lanai, dripping cliffs, 20 ft. from ocean, *Degener* 21931. MAUI: Makamakaole Stream, *Degener* 21151, 21153; Maliku, Maui, spring-fed ledges at ocean, *Degener* 21932. HAWAII: Kaumana Caves, near Hilo, *Degener* 22276B; Keauohana Forest Reserve, near Pahoa, Puna, *Degener* 22282C; Pololu-Honokanenui region, Kohala, *Degener* & *Greenwell* 22349, 22351C. New to Kauai, Molokai, and Lanai.
- Ctenidium decurrens* (Sull.) Broth. OAHU: E spur of E ridge, W branch of East Makaleha Valley, dark gulch, 1,800 ft., *Degener* 20878A. MAUI: Koolau Gap, Haleakala, 7,500-8,000 ft., *Miller* & *Lamberton* 4283. HAWAII: Above Kulani Prison, Mauna Loa, *Degener* & *Greenwell* 21933F, 22338C, 22361. Additionally reported from Kauai and Molokai.

*Ctenidium elegantulum* Broth. OAHU: Kaaawa Gulch, Mt. Kaala, Kamananui, 1,300 ft., Miller 2500. HAWAII: Honokanenui Valley, Kohala, Degener & Greenwell 22356. New to Oahu and Hawaii. Previously known only from Maui.

*Pogonatum baldwinii* (C. Müll.) Par. KAUAI: Near Kokee, road cut, Degener 20471; Mohihi Road, Degener & Greenwell 22287B. MOLOKAI: Head of Waikolu Valley, vertical bank, O. & I. Degener 23836. HAWAII: Below Kulani Prison, dense forest, 4,000 ft., Degener 22281. Additionally reported from Oahu, Lanai, and Maui.

#### LITERATURE CITED

- BARTRAM, E. B. 1933. Manual of Hawaiian mosses. Bernice P. Bishop Mus. Bull. 101: 1-275.  
\_\_\_\_\_. 1939. Supplement to the manual of Hawaiian mosses. Bernice P. Bishop Mus., Occ. Pap. 15: 93-108.  
\_\_\_\_\_. 1942. Mosses collected by Hawaiian Bog Survey of 1938. *Ibid.* 16: 321-336.  
\_\_\_\_\_. 1952. High altitude mosses from Mauna Kea, Island of Hawaii. *Ibid.* 20: 297-300.  
CROSBY, M. R. 1965. New records for Hawaiian Island mosses. THE BRYOLOGIST 68: 457-462.  
HÖRMANN, H. 1964. Contributions to two Hawaiian mosses: *Aerobryopsis longissima* (Dozy et Molk.) Fleisch. and *A. scariosa* Bartr. Advancing Frontiers of Plant Sci. 8: 33-40.  
\_\_\_\_\_. 1964. Preliminary contributions to Dr. Degener's survey of Hawaiian mosses. Rev. Bryol. & Lichénol. 33: 550-554.  
REICHARDT, H. W. 1877. Beitrag zur Kryptogamenflora der Hawaiischen Inseln. Sitzber. Akad. Wiss. Wien. 75: 553-582.  
SCHULTZE-MOTEL, W. 1962. Beitrag zur Kenntnis der Laubmose der Hawaii-Inseln. Willdenowia 3: 97-107. (1963).  
WIJK, R. VAN DER, W. D. MARGADANT & P. A. FLORSCHÜTZ. 1959, 1962, 1964. Index Muscorum. Vol. 1-3. Utrecht.  
\_\_\_\_\_. & W. D. MARGADANT. 1962. New combinations in mosses VII. Taxon 11: 221-223.