by Sue Bottom, sbottom15@hotmail.com

The Royal Horticultural Society is the administrator of the International Orchid Register for new orchid hybrids. The RHS had adopted changes to orchid nomenclature that affect the names we call our orchids. The orchid taxonomists who have proposed these changes are referred to somewhat derisively as the lumpers and the splitters, because they are busy lumping some species together into an existing genus or splitting species apart into separate sometimes new genera. I have happily continued to use the old Sanders names with which I am familiar and comfortable. There are many new hybrids being referred to by the new nomenclature so I decided it was time for this old dog to learn some new tricks.

Central American Cattleyas Split into Guarianthes. The splitters pulled the Central American bifoliate species out of the cattleya genus in 2003 and created the new genus Guarianthe (Gur.) to house them. Dressler and Higgins felt this group was out of place in the Cattleya genus, partly on the basis of DNA analysis, and proposed the new generic name based on Guaria, a Costa Rican word for orchid, and the Greek term for flower, anthe.




Photo Credits: Gur. aurantiaca by Gene Crocker of Carter and Holmes, Gur. bowringiana by Jean Wilson, Gur. deckeri by Michael Blietz of Exotic Orchids, Gur. patinii by Jean Wilson, Gur. skinneri by Jean Wilson

This has created the need for new intergeneric names to describe the hybrids with Guarianthes. The table below lists the more common intergenerics that have been registered. The combinations with various genera are listed, along with a note about what Sanders would have called it. The Sanders information assumes none of the reorganizations in the cattleya alliance occurred.

Guarianthe Combinations with Other Genera in the Cattleya Alliance					
Combine Guarianthe with:		To Create This Intergeneric:		Sanders Would Have Called It:	
B.	Brassavola (B)	Bsn.	Brassanthe	Bc.	Brassocattleya
Bc.	Brassocattleya (B x C)	Bct.	Brassocatanthe	Bc.	Brassocattleya
Bro.	Broughtonia (Bro)	Grt.	Guaritonia	Ctna.	Cattleytonia



by Sue Bottom, sbottom15@hotmail.com

Guarianthe Combinations with Other Genera in the Cattleya Alliance						
Combine Guarianthe with:		To Create This Intergeneric:		Sanders Would Have Called It:		
C.	Cattleya (C)	Ctt.	Cattlianthe	C.	Cattleya	
Ctyh.	Cattleychea (C x Psh)	Pgy.	Proguarleya	Ерс.	Epicattleya	
Ctna.	Cattleytonia (Bro x C)	Gct.	Guaricattonia	Ctna.	Cattleytonia	
Cty.	Catyclia (C x E)	Eny.	Enanthleya	Ерс.	Epicattleya	
Clty.	Caulocattleya (C x Cau)	Gty.	Guarthroleya	Clty.	Caulocattleya	
E.	Encyclia (E)	Gcy.	Guaricyclia	Epc.	Epicattleya	
Ерс.	Epicattleya (C x Epi)	Ett.	Epicatanthe	Ерс.	Epicattleya	
Epi.	Epidendrum (Epi)	Gdd.	Guaridendrum	Ерс.	Epicattleya	
L.	Laelia (L)	Lnt.	Laelianthe	Lc.	Laeliocattleya	
Lc.	Laeliocattleya (C x L)	Lcn.	Laeliocatanthe	Lc.	Laeliocattleya	
Mel.	Meloara (C x Cau x L)	Jkf.	Jackfowlieara	Lcr.	Laeliocatarthron	
Мус.	Myrmecattleya (C x Mcp)	Сср.	Guaricatophila	Smbc.	Schombocattleya	
Psh.	Prosthechea (Psh)	Grc.	Guarechea	Epc.	Epicattleya	
Rcc.	Rhyncatclia (C x E)	Bul.	Bullara	Epc.	Epicattleya	
Ryc.	Rhyncatlaelia (C x L x RI)	Rchg.	Rechingerara	Blc.	Brassolaeliocattleya	
Rby	Rhynchobrassoleya (B x C x RI)	Chz.	Cahuzacara	Bc.	Brassocattleya	
RI.	Rhyncholaelia (RI)	Ryn.	Rhyncanthe	Bc.	Brassocattleya	
Rlc.	Rhyncholaeliocattleya (C x RI)	Rth.	Rhyncattleanthe	Bc.	Brassocattleya	
Rly	Rhyntonleya (Bro x C x RI)	Vkt.	Volkertara	Stlma.	Stellamizutaara	

Some Brassavolas Split into Rhyncholaelias. The splitters pulled two species out of the Brassavolas and created the new genus Rhyncholaelia to house them. Though this change was made prior to the turn of the century, many growers still refer to them as Brassavolas. The fimbriated lip of the digbyana is very popular with hybridizers and has been incorporated into many hybrids such that the B in the majority of the old Brassocattleyas and Brassolaeliocattleyas is from B. digbyana.

by Sue Bottom, sbottom15@hotmail.com




Photo Credits: RI. digbyana by David Genovese and RI. glauca by P. Nelson

The addition of the Rhyncholaelia genus created the need for new genus names to describe its hybrids.

Rhyncholaelia Combinations with Other Genera in the Cattleya Alliance						
Combine Rhyncholaelia with:		To Create This Intergeneric:		Sanders Would Have Called It:		
B.	Brassavola (B)	Rcv.	Rhynchovola	B.	Brassavola	
Bct.	Brassocatanthe (B x C x Gur)	Chz.	Cahuzacara	Вс	Brassocattleya	
Bc.	Brassocattleya (B x C)	Rby.	Rhynchobrassoleya	Вс	Brassocattleya	
Ctyl.	Catcylaelia (C x E x L)	Aea.	Appletonara	Yhra.	Yahiroara	
C.	Cattleya (C)	Rlc.	Rhyncholaeliocattleya	Bc.	Brassocattleya	
Ctyh.	Cattleychea (C x Psh)	Pry.	Prosrhyncholeya	Vnra.	Vaughnara	
Ctna.	Cattleytonia (Bro x C)	Rly.	Rhyntonleya	Stlma.	Stellamizutaara	
Ctt.	Cattlianthe (C x Gur)	Rth.	Rhyncattleanthe	Вс	Brassocattleya	
Cty.	Catyclia (C x E)	Rcc.	Rhyncatclia	Vnra.	Vaughnara	



by Sue Bottom, sbottom15@hotmail.com

Rhyncholaelia Combinations with Other Genera in the Cattleya Alliance						
Combine Rhyncholaelia with:		To Create This Intergeneric:		Sanders Would Have Called It:		
Clty.	Caulocattleya (C x Cau)	Rry.	Rhynarthrolyea	Hook.	Hookerara	
Eny.	Enanthleya (C x E x Gur)	Bul.	Bullara	Vnra.	Vaughnara	
Epc.	Epicattlya (C x Epi)	Rnd.	Rhycatdendrum	Vnra.	Vaughnara	
Epi.	Epidendrum (Epi)	Rdd.	Rhynchodendrum	Вері.	Brassoepidendrum	
Gur.	Guarianthe (Gur)	Rhy.	Rhyncanthe	Bc.	Brassocattleya	
Gct.	Guaricattonia (Bro x C x Gur)	Vkt.	Volkertara	Stlma.	Stellamizutaara	
L.	Laelia (L)	Lrn.	Laelirhynchos	BI.	Brassolaelia	
Lcn.	Laeliocatanthe (C x Gur x L)	Rchg.	Rechingerara	Blc.	Brassolaeliocattleya	
Lcr.	Laeliocatarthron (C x Cau x L)	Mel.	Meloara	Col.	Collierara	
Lc.	Laeliocattleya (C x L)	Ryc.	Rhyncatlaelia	Blc.	Brassolaeliocattleya	
Led.	Ledienara (C x Cau x Gur x L)	Jkf.	Jackfowlieara	Col.	Collierara	
Мус.	Myrmecocattleya (C x Mcp)	Rmy.	Rhynchomyremeleya	Recc.	Recchara	

Cattleyode Laelias Lumped into Cattleyas. The lumpers moved the large flowered Cattleyode Brazilian Laelias into the Cattleya genus. This did not result in any need for new intergeneric names, it just changed the name of many familiar orchids. L. purpurata has been in the parentage of perhaps 90% of the previously called laeliocattleya orchids that are now known as cattleyas.




Photo Credits: C. crispa by Dalton Baptista, C. grandis by Mauro Rosin, C. lobata by Mauro Rosin, C. purpurata by Woolf Orchid Culture and C. tenebrosa by Mauro Rosin



by Sue Bottom, sbottom15@hotmail.com

Other Brazilian Laelias Lumped into Cattleyas. The lumpers moved the rest of the Brazilian laelias into the Cattleya genus, including the colorful Brazilian rupicolous laelias. Only Laelias from Mexico and Central America are now still considered to be Laelias.

Other Brazilian Laelias Moved Into the Genus Cattleya							
			1				
C. alaorii	C. jongheana	C. perrinii	C. pumila	C. sincorana			
		Sendanta Res Spins					
C. bradei	C. briegeri	C. cinnabarina	C. harpophylla	C. longipes			

Photo Credits: C. alaorii , C.jongheana, C. perrinii, C. pumila, C.sincorana, C. bradei by Francisco Miranda, C. briegeri by Fred Clarke, C. cinnabarina by Mauro Rosin, C. harpophylla by Maureen Puligano, C. longipes by Francisco Miranda

Schomburgkias Eliminated, Plants Split into Myrmecophilas and Laelias. The hollow pseudobulb Schomburgkias were moved into the Myrmecophila genus. The genus name Schomburgkia, which meant ant lover and referred to the symbiotic relationship these plants had with ants living within the hollow pseudobulbs, is no more. This also eliminated the intergeneric Schombocattleyas (Smbc.), etc. A partial listing includes the following:




Photo Credits: Mcp. albopurpurea by Judy Cook, Mcp. brysiana by Jean Wilson, Mcp. exaltata by James Jeansonne, Mcp. thomsoniana by Fred Clarke and Mcp. tibicinis by Stephen William Swan of Burbank O. Nursery

Page 5 of 6

by Sue Bottom, sbottom15@hotmail.com

The solid pseudobulb Schomburgkias are supposedly more closely related to the Mexican laelias such as L. anceps based on DNA sequencing. Some of the Schomburgkias including the South American Schomburgkias moved into the Laelia genus include:




Photo Credits: L. Iyonsii by Jean Wilson, L. rosea by Weyman Bussey, L. splendid by Jose Portilla of Ecuagenera, L. superbiens by Patricia Harding, and L. undulata by Mauro Rosin

Sophronitis Eliminated, Plants Lumped into Cattleyas. The lumpers moved all the Sophronitis species into the Cattleya genus. These mostly cool growing orchids have been used to impart their intense red coloration to their progeny. Lumping the Sophronitis in with the Cattleyas did not result in any new intergeneric names, rather it eliminated the need for many intergeneric names like Sophrocattleya (Sc.), Sophrolaelia (Sl.), Sophrolaeliocattleya (Slc.) and Potinara (Pot.). Some notable renamed Sophronitis include:




Photo Credits: C. brevipedunculata, C. cernua and C. coccinea by John Varigos, C. wittigiana by Mauro Rosin

The reclassifications of the various genera align the orchid groups more geographically. Laelias are found from Mexico through central America into Northern South America. Cattleyas are primarily a Brazilian genus with some unifoliate species extending into the northern Andes and Central America. The Guarianthe are found in Central America. The negative consequence of this new nomenclature is the renaming of so many of the hybrids in the Cattleya alliance.

Having said all this, I haven't changed any of the labels on my plants.

Page 6 of 6

All contents copyright © 2013-2017 by Sue and Terry Bottom. All rights reserved. No part of this document or the related photos may be reproduced or transmitted in any form, by any means without the prior written permission of the author.